The fluctuations, under time reversal, of the natural time and the entropy distinguish similar looking electric signals of different dynamics

P. A. Varotsos,^{1,*} N. V. Sarlis,¹ E. S. Skordas,¹ and M. S. Lazaridou¹

¹Solid State Section and Solid Earth Physics Institute, Physics Department, University of Athens, Panepistimiopolis, Zografos 157 84, Athens, Greece

We show that the scale dependence of the fluctuations of the natural time itself under time reversal provides a useful tool for the discrimination of seismic electric signals (*critical* dynamics) from noises emitted from manmade sources as well as for the determination of the scaling exponent. We present recent data of electric signals detected at the Earth's surface, which confirm that the value of the entropy in natural time as well as its value under time reversal are smaller than that of the entropy of a "uniform" distribution.

PACS numbers: 91.25.Qi, 91.30.Px, 05.45.Tp

I. INTRODUCTION

In a time series comprising N events, the natural time $\chi_k = k/N$ serves as an index^{1,2,3} for the occurrence of the k-th event. In natural time analysis, the time evolution of the pair of the two quantities (χ_k, Q_k) is considered, where Q_k denotes in general a quantity proportional to the energy released during the k-th event. In the case of dichotomous electric signals (e.g., seismic electric signal (SES) activities, i.e., low frequency \leq 1Hz electric signals that precede earthquakes, e.g., see Refs. 4, 5, 6, 7, 8, 9, 10, 11, 12, 13) Q_k stands for the duration of the k-th pulse (cf. The SES activities should not be confused with pulses of very short durations observed some minutes before earthquakes¹⁴). It has been shown¹⁵ that natural time domain is optimal for enhancing the signals' localization in the time-frequency space, thus conforming to the desire to reduce uncertainty and extract signal information as much as possible. The entropy S in natural time is defined¹⁶ as the derivative with respect to q of the fluctuation function $\langle \chi^q \rangle - \langle \chi \rangle^q$ at q = 1:

$$S \equiv \langle \chi \ln \chi \rangle - \langle \chi \rangle \ln \langle \chi \rangle \tag{1}$$

where $\langle f(\chi) \rangle = \sum_{k=1}^{N} p_k f(\chi_k)$ and $p_k = Q_k / \sum_{n=1}^{N} Q_n$. It is dynamic entropy^{17,18} and exhibits¹⁹ concavity, positivity and Lesche^{20,21} stability. Note that S should not be confused with $\operatorname{Cov}[\chi, \ln \chi] \equiv \langle \chi \ln \chi \rangle - \langle \chi \rangle \langle \ln \chi \rangle$ since in general $\langle \ln \chi \rangle \neq \ln \langle \chi \rangle$. The value of the entropy upon considering the time reversal \mathcal{T} , i.e., $\mathcal{T}p_k = p_{N-k+1}$, is labelled by S_- . The value of S_- is^{19,22,23}, in general, different from S, and thus S does satisfy the conditions to be "causal" in the following sense (see Ref.19 and references therein): When studying a dynamical system evolving in time, the "causality" of an operator describing this evolution assures that the values assumed by the operator, at each time instant, depends solely on the past values of the system. Hence, a "causal" operator should be able to represent the evolution of the system according to the (true) time arrow, thus the operator can represent a real physical system evolving in time and reveal the differences arising upon time-reversal of the series.

The statistical properties of S and S_{-} have been stud-

ied in a variety of models $^{22,23}.$ In the case of a "uniform" distribution $S = S_{-} = S_{u} (= \ln 2/2 - 1/4 \approx 0.0966).$ The "uniform" distribution (defined in Refs.1,24) has been analytically studied in Ref.17 and corresponds to the case when Q_k are independent and identically distributed (IID) positive random variables of finite variance including the case of Markovian dichotomous electric signals studied in Ref.16. The "uniform" distribution corresponds to $p(\chi) = 1$, where $p(\chi)$ is a continuous probability density function (PDF) corresponding to the point probabilities p_k used so far. When Q_k of a "uniform" distribution are perturbed by a small linear trend, we find²² (see also Eq.(4), below) that $(S - S_u)(S_- - S_u) < 0$ (cf. this simple example, which shows that S captures the effect of a linear trend, may be considered as clarifying the meaning of S, see Section V of Ref.22). Another model studied is when the increments of Q_k are positive IID, in this case we find²³ that $S \approx 0.048$ and $S_{-} \approx 0.088$ which are both smaller than S_u . The same holds, i.e., that both S and S_{-} are smaller than S_{u} , in the examples of an on-off intermittency model discussed in Ref.22 as well as for a multiplicative cascades $model^{23}$ adjusted to describe turbulence data.

A case of practical importance is that of the SES activities. SES activities (*critical* dynamics) exhibit infinitely ranged long-range temporal correlations^{22,23,24} which are destroyed²³ after shuffling the durations Q_k randomly. An interesting property emerged from the data analysis of several SES activities refers to the fact²² that both S and S_- values are smaller than the value of S_u , i.e.,

$$S, S_{-} < S_{u}, \tag{2}$$

in addition to the fact that for SES activities 1,2,3 the variance

$$\kappa_1 \equiv \langle \chi^2 \rangle - \langle \chi \rangle^2 \approx 0.070. \tag{3}$$

These findings -which do not hold¹⁹ for "artificial" noises (AN) (i.e., electric signals emitted from manmade sources)- have been supported by numerical simulations in fractional Brownian motion (fBm) time series^{22,23} that have an exponent α_{DFA} , resulted from the Detrended Fluctuation Analysis (DFA)^{25,26}, close to unity. This

model have been applied since fBm (with a self-similarity index $H \approx 1$) has been found²⁷ as an appropriate type of modeling process for the SES activities. These simulations resulted in values of S and S_{-} that do obey relation (2) (see Fig.4 of Ref.22) and $\kappa_1 \approx 0.070$ (see Fig.3 of Ref.23). It was then conjectured²² that the validity of the relation (2) stems from infinitely ranged long-range temporal correlations (cf. $H \approx 1$). On the other hand, for short-range temporal correlations (e.g. when modeling Q_k by an autoregressive process $Q_k =$ $aQ_{k-1} + g_k + c, |a| < 1$ and c stands for an appropriate constant to ensure positivity of Q_k or $Q_k = |aQ_{k-1} + g_k|$ where g_k is Gaussian IID variables) the values of both S and S_{-} approach (see Appendix A) that of S_{u} and $\kappa_1 \rightarrow \kappa_u$, where $\kappa_u = 1/12$ denotes the corresponding value of the "uniform" distribution 23 .

The scope of this paper is twofold: First, in Section II, we point out the usefulness of the study of the fluctuations of the natural time itself under time reversal. In particular, it enables the determination of the scaling exponent, thus allowing the distinction of SES activities from similar looking AN. Second, in Section III, we provide the most recent experimental data that strengthen the validity of the relations (2) and (3) for SES activities. The earthquakes that followed the latter SES activities are described in Section IV. Section V, summarizes our conclusions.

II. THE FLUCTUATIONS OF NATURAL TIME UNDER TIME REVERSAL

The way through which the entropy in natural time captures the influence of the effect of a small linear trend has been studied, as mentioned, in Ref.22 on the basis of the parametric family of PDFs: $p(\chi; \epsilon) = 1 + \epsilon(\chi - 1/2)$, where ϵ measures the extent of the linear trend. Such a family of PDFs shares the interesting property $\mathcal{T}p(\chi; \epsilon) = p(\chi; -\epsilon)$, i.e, the action of the time reversal is obtained by simply changing the sign of ϵ . It has been shown²² that the entropy $S(\epsilon) \equiv S[p(\chi; \epsilon)]$, as well as that of the entropy under time reversal $S_{-}(\epsilon) \equiv$ $S[\mathcal{T}p(\chi; \epsilon)], S_{-}(\epsilon) = S(-\epsilon)$, depend *non*-linearly on the trend parameter ϵ :

$$S(\epsilon) = -\frac{1}{4} + \frac{\epsilon}{72} - \left(\frac{1}{2} + \frac{\epsilon}{12}\right) \ln\left(\frac{1}{2} + \frac{\epsilon}{12}\right). \quad (4)$$

However, it would be extremely useful to obtain a *linear* measure of ϵ in natural time. Actually, this is simply the average of the natural time itself:

$$\langle \chi \rangle = \int_0^1 \chi p(\chi; \epsilon) d\chi = \frac{1}{2} + \frac{\epsilon}{12}.$$
 (5)

If we consider the fluctuations of this simple measure upon time-reversal, we can obtain information on the long-range dependence of Q_k . We shall show that a measure of the long-range dependence emerges in natural time if we study the dependence of its fluctuations under time-reversal $\Delta \chi_l^2 \equiv \mathbb{E}[(\langle \chi \rangle - \langle \mathcal{T} \chi \rangle)^2]$ on the window length l that is used for the calculation. Since $\mathcal{T}p_k = p_{l-k+1}$, we have

$$\Delta \chi_l^2 \equiv \mathrm{E}[(\langle \chi \rangle - \langle \mathcal{T} \chi \rangle)^2] = \mathrm{E}\left\{ \left[\sum_{k=1}^l \frac{k}{l} \left(p_k - p_{l-k+1} \right) \right]^2 \right\},\tag{6}$$

where the symbol E[...] denotes the expectation value obtained when a window of length l is sliding through the time series Q_k . E[...] is well defined when all the $\{p_k\}_{k=1,2,...l}$ involved in its argument are also well defined. The evaluation of E[...] can be carried out either by full or by Monte Carlo calculation. In order to achieve this goal, from the original time-series $\{Q_k\}_{k=1,2,...L}$, we select segments $\{Q_{m_0+i-1}\}_{i=1,2,...l}$ of length l, and the argument of E[...] is computed by substituting $p_k = Q_{m_0+k-1} / \sum_{i=1}^l Q_{m_0+i-1}$. The sum of the resulting values over the number of the selected segments (different m_0) is assigned to E[...]. The full calculation refers to the case when m_0 takes all the $L - l + 1(m_0 = 1, 2, \ldots L - l + 1)$ possible values, whereas the Monte Carlo when m_0 is selected randomly.

By expanding the square in the last part of Eq.(6), we obtain

$$\Delta \chi_l^2 = \sum_{k=1}^l \left(\frac{k}{l}\right)^2 \mathbf{E}[(p_k - p_{l-k+1})^2] + \sum_{k \neq m} \frac{km}{l^2} \mathbf{E}[(p_k - p_{l-k+1})(p_m - p_{l-m+1})].$$
(7)

The basic relation¹⁷ that interrelates p_k is $\sum_{k=1}^{l} p_k = 1$ or equivalently $p_k = 1 - \sum_{m \neq k} p_m$. By subtracting from the

last expression its value for k = l - k + 1, we obtain $p_k - p_{l-k+1} = -\sum_{m \neq k} (p_m - p_{l-m+1})$, and thus

$$(p_k - p_{l-k+1})^2 = -\sum_{m \neq k} (p_k - p_{l-k+1})(p_m - p_{l-m+1}).$$
(8)

By substituting Eq.(8) into Eq.(7), we obtain

$$\Delta \chi_l^2 = -\sum_{k=1}^l \left(\frac{k}{l}\right)^2 \sum_{m \neq k} \operatorname{E}[(p_k - p_{l-k+1})(p_m - p_{l-m+1})] + \sum_{k \neq m} \frac{km}{l^2} \operatorname{E}[(p_k - p_{l-k+1})(p_m - p_{l-m+1})]$$
(9)

which simplifies to

$$\Delta \chi_l^2 = -\sum_{k,m} \frac{(k-m)^2}{l^2} \mathbb{E}[(p_k - p_{l-k+1})(p_m - p_{l-m+1})]$$
(10)

The negative sign appears because $(p_k - p_{l-k+1})$ and $(p_m - p_{l-m+1})$ are in general anti-correlated due to Eq.(8). Equation (10) implies that $\Delta \chi_l^2$ measures the long-range correlations in Q_k : If we assume that $-\mathrm{E}[(p_k - p_{l-k+1})(p_m - p_{l-m+1})] \propto (k-m)^{2\chi_H}/l^2$ (cf. p_k scales as 1/l, e.g. see¹⁷), we have that

$$\Delta \chi_l^2 \propto l^{4+2\chi_H}/l^4 \tag{11}$$

so that

$$\Delta \chi_l (\equiv \sqrt{\Delta \chi_l^2}) \propto l^{\chi_H}, \qquad (12)$$

where χ_H is a scaling exponent.

A. Fractional Brownian motion and fractional Gaussian noise time series

In order to examine the validity of the above result Eq.(12) when Q_k are coming from fBm or fractional Gaussian noise (fGn), we employed the following procedure: First, we generated fBm (or fGn) time-series X_k (consisting of 2×10^4 points) for a given value of ${\cal H}$ using the Mandelbrot-Weierstrass function 28,29,30 as described in Ref.22. Second, since Q_k should be positive, we normalized the resulting X_k time-series to zero mean and unit standard deviation and then added to the normalized time-series N_k a constant factor c to ensure the positivity of $Q_k = N_k + c$ (for the purpose of the present study we used c = 10). The resulting Q_k timeseries were then analyzed and the fluctuations of $\Delta \chi_l$ versus the scale l are shown in Figs. 1(a) and 1(d) for fGn and fBm, respectively. The upper three panels of Fig.1 correspond to fGn while the lower three to fBm. We observe (see Fig.1(b)) that for fGn we have the interconnection: $\chi_H \approx H - 1$ corresponding to descending curves(see Fig.1(a)), whereas for fBm the interconnection turns (see Fig.1(e)) to: $\chi_H \approx H$ corresponding to ascending curves (see Fig.1(d)).

In order to judge the merits or demerits of the procedure proposed here for the determination of the scaling exponent, we compare Figs.1(b) and 1(e) with Figs.1(c) and 1(f), respectively, that have been obtained by the well-established DFA method^{25,26}. This comparison reveals that the results are more or less comparable for fGn, while for fBm the exponent χ_H deviates less from the behavior of an ideal estimator of the true scaling exponent (drawn in dashed green) compared to α_{DFA} , especially for the largest H values.

B. The fluctuations of the natural time to distinguish seismic electric signal activities from similar looking AN

The physical meaning of the present analysis was further investigated by performing the same procedure in the time-series of the durations of those signals analyzed in Ref.19 that have enough number of pulses e.g. $\approx 10^2$ (cf. the signals depicted in Fig.3 could not be analyzed in view of the small number of pulses). The relevant results are shown in Fig.2. Their inspection interestingly indicates that all seven AN correspond to descending $\Delta \chi_l$ curves versus the scale l, while the three SES activities to ascending curves (in a similar fashion as in Figs.1(a) and 1(d), respectively) as expected from the fact that the latter exhibit²⁴ infinitely ranged long-ranged temporal correlations (having H close to unity), while the former do not. Hence, the method proposed here enables the detection of long-range correlations even for datasets of small size ($\approx 10^2$), thus allowing the distinction of SES activities from AN.

III. RECENT DATA OF SEISMIC ELECTRIC SIGNALS ACTIVITIES

First, Fig.3(a) depicts an electric signal, consisting of a number of pulses, that has been recorded on November 14, 2006 at a station labelled³¹ PIR lying in western Greece (close to Pirgos city). This signal has been clearly collected at eleven measuring electric dipoles with electrodes installed at sites that are depicted in a map given in Ref.31. The signal is presented (continuous line in red) in Fig. 3(a) in normalized units, i.e., by subtracting the mean value and dividing by the standard deviation. For the reader's convenience, the corresponding dichotomous representation is also drawn in Fig. 3(a) with a dotted (blue) line, while in Fig. 3(c) we show (in red crosses) how the signal is read in natural time. The computation of S and S_{-} leads to the following values: $S = 0.070 \pm 0.012$, $S_{-} = 0.051 \pm 0.010$. As for the variance κ_1 , the resulting value is $\kappa_1 = 0.062 \pm 0.010$. These values more or less obey the conditions (2) and (3) that have been found to hold for other SES activities²². Note that the feature of this SES activity, it is similar to the one observed at the same station before the magnitude $M \approx 6.7$ earthquake that occurred on Jan 8, 2006, see Ref.32.

A closer inspection of Fig. 3(a) reveals the following experimental fact: An additional electric signal has been also detected (in the gray shaded area of Fig. 3(a)), which consists of pulses with markedly smaller amplitude than those of the SES activity discussed in the previous paragraph. This is reproduced (continuous line in red) in Fig. 3(b) in an expanded time scale and for the sake of the reader's convenience its dichotomous representation is also marked by the dotted (blue) line, which leads to the natural time representation shown (dotted blue) in Fig. 3(c). The computation of S and S_{-} gives $S = 0.077 \pm 0.004$, $S_{-} = 0.082 \pm 0.004$, while κ_1 is found to be $\kappa_1 = 0.076 \pm 0.005$. Hence, these values also obey the conditions (2) and (3) for the classification of this signal as an SES activity.

The two aforementioned signals have been followed by two significant earthquakes as described in Section IV. This conforms to their classification as SES activities, which has been completed in an early version of this paper³³ on November 16, 2006.

Second, very recently, i.e., on July 2, 2007 and July 10,2007, two separate electric signals were recorded at a station labelled PAT lying in central Greece (close to Patras city) at 38.32°N21.90°E. The signals are presented (continuous line in red) in Fig. 3 (d) and 3 (e) in normalized units in a similar fashion as in Figs.3(a),(b). Their corresponding dichotomous representation are also drawn with dotted (blue) lines, while in Fig.3(f) we show (in red crosses and blue asterisks, respectively) how the signals are read in natural time. The computation of κ_1 , S and S_{-} leads to the following values: For the signal on July 2, 2007: $\kappa_1 = 0.072 \pm 0.005$, $S = 0.073 \pm 0.007$, $S_{-} = 0.081 \pm 0.006$, for the signal on July 10, 2007: $\kappa_1 = 0.073 \pm 0.004, S = 0.085 \pm 0.005, S_- = 0.080 \pm 0.004.$ An inspection of these values reveals that they obey the conditions (2) and (3) and hence both signals can be classified as SES activities. The procedure for the current study of the subsequent seismicity that occurred after these SES activities is described in the next Section.

IV. THE SEISMIC ACTIVITY THAT FOLLOWED THE SES ACTIVITIES

We discriminate that during the last decade SES activities are publicized *only* when their amplitude indicates that the impending earthquake has an expected^{13,23} magnitude comparable to 6.0 unit or larger.

A. The case of the SES activities of Figs.3(a),(b)

According to the Athens observatory (the data of which will be used here), a strong earthquake (EQ) with magnitude 5.8-units occurred at 13:43 UT on February 3, 2007, with epicenter at 35.8° N 22.6°E, i.e., almost 80 km to the southwest of the 6.9 EQ of January 8, 2006, (cf. the magnitude announced from Athens observatory is equal to ML+0.5, where ML stands for the local magnitude). This was preceded by a 5.2-units EQ that occurred at 22:25 UT on January 18, 2007 at 34.8°N 22.7°E. The occurrence of these two EQs confirm the classification as SES activities of the signals depicted in Figs.3(a) and 3(b). (Note that preseismic information based on SES activities is issued *only* when the magnitude of the strongest EQ of the impending EQ activity is estimated -by means of the SES amplitude³⁴- to be comparable to 6.0 units or $larger^{13}$.)

Here, we show that the occurrence times of the aforementioned two EQs can be estimated by following the procedure described in Refs.1,22,23,35 and using the order parameter of seismicity proposed in Ref.35, i.e, the normalized power spectrum in natural time $\Pi(\phi)$ as $\phi \rightarrow 0$ (see also below). We study how the seismicity evolved after the recording of the SES activities on November 14, 2006, at PIR station (which were classified as SES activities in the initially submitted version of the present paper on November 16, 2006). The study is made either in the area $A:N_{34.45}^{37.54}E_{20.95}^{24.54}$ or in the area $B:N_{34.65}^{37.24}E_{21.55}^{24.44}$ (see Fig.4), by considering three magnitude thresholds $M_{thres} = 3.2, 3.4$ and 3.6 (hence six combinations were studied in total). If we set the natural time for seismicity zero at the initiation of the SES activity at 17:19 UT on November 14, 2006, we form time series of seismic events³⁶ in natural time for various time windows as the number N of consecutive (small) EQs increases. We then compute the normalized power spectrum^{1,22,23,35} in natural time $\Pi(\phi)$ for each of the time windows. Excerpts of these results, which refer to the values during the periods: (a) December 25, 2006, to January 17, 2007, and (b): January 18 to January 31, 2007 are depicted with red crosses in Fig.5, respectively. This figure corresponds to the small area B with $M_{thres} = 3.4$. In the same figure, we plot in blue the normalized power spectrum obeying the relation 1,2,3,35

$$\Pi(\omega) = \frac{18}{5\omega^2} - \frac{6\cos\omega}{5\omega^2} - \frac{12\sin\omega}{5\omega^3}$$
(13)

which holds when the system enters the critical

stage ($\omega = 2\pi\phi$, where ϕ stands for the natural frequency 1,2,3,13). The date and the time of the occurrence of each small earthquake (with $M \geq 3.4$) that occurred in the area B, is written in green in each panel (see also Table I). An inspection of Fig.5(a) reveals that the red line approaches the blue line as N increases and a coincidence occurs at the small event of magnitude 3.7 that occurred at 03:22 UT on January 17, 2007, i.e., roughly two days before the 5.2-units EQ at 22:25 UT on January 18,2007. A similar behavior is observed in Fig.5(b) in which we see that a *coincidence* occurs at the small event of magnitude 3.6 at 18:40 UT on January 31, 2007, i.e., roughly three days before the strong EQ of magnitude 5.8-units that occurred at 13:43 UT on February 3, 2007. To ensure that these two coincidences in Figs.5(a)and (b) are true ones^{1,3,13,31,35} (see also below) we also calculate the evolution of the quantities κ_1, S and S_- and the results are depicted in Fig.6 for the three magnitude thresholds for each of the aforementioned two areas A and B.

The conditions for a coincidence to be considered as true are the following (e.g., see Ref.1, see $also^{3,13,31,35}$): First, the 'average' distance $\langle D \rangle$ between the empirical and the theoretical $\Pi(\phi)$ (i.e., the red and the blue line, respectively, in Fig.5) should be 1,2,13,31,35 smaller than or equal to 10^{-2} . See Fig.7, where we plot $\langle D \rangle$ versus the conventional time during the whole period after the recording of the SES activities on November 14, 2006, for both areas, i.e., the large one (area A) and the small (area B) and the three magnitude thresholds. For the sake of the readers convenience, the mean value of the results obtained for the three thresholds is also shown in black. Second, in the examples observed to date $^{1,3,13,31,34,35}.$ a few events *before* the coincidence leading to the strong EQ, the evolving $\Pi(\phi)$ has been found to approach that of Eq.(1), i.e., the blue one in Fig.5, from below (cf. this reflects that during this approach the κ_1 -value decreases as the number of events increases). In addition, both values S and S_{-} should be smaller than S_{u} at the coincidence. Finally, since the process concerned is selfsimilar (*critical* dynamics), the time of the occurrence of the (true) coincidence should not change, in principle, upon changing either the (surrounding) area or the magnitude threshold used in the calculation. Note that in Fig.7, at the last small events , i.e., the rightmost in Figs.5(a) and 5(b), respectively (i.e., the magnitude 3.7 event on January 17, 2007 and the second event of magnitude 3.6 on January 31, 2007) just before the occurrences of the 5.2-units and 5.8-units EQs, in both areas A and B, the mean value (see the black thick lines in Fig.7) of $\langle D \rangle$ obtained from the three magnitude thresholds become smaller than or equal to 10^{-2} . Hence, these two coincidences can be considered as true.

In summary, the SES activities recorded on November 14, 2006, at PIR station (presented in Figs.3(a),(b)) have been followed by two EQs with magnitudes 5.2-units and 5.8-units that occurred on January 18 and February 3, 2007. The time of the occurrences of these two EQs are determined within a narrow range of a few days upon analyzing, in natural time, the seismicity subsequent to the SES activities.

B. The case of the SES activities of Figs.3(d),(e)

The actual amplitude (in mV) of the most recent SES activities recorded at PAT on July 2, 2007 and July 10, 2007 (see Fig.3(d) and (e), respectively) can be visualized in Figs.8(a) and 8(b) where the original recordings of a measuring electric dipole (with length $L \approx 5$ km) are reproduced. For the sake of comparison, in Figs.8(c),(d) we also present the corresponding SES activities at the same station, i.e., PAT, that preceded^{23,31} the magnitude 6.0-class earthquakes that occurred with epicenters at 37.6°N20.9°E on April 11 and 12,2006. Furthermore, in Fig.8(e), we show the SES activity³⁷ at PAT on February 8, 2007, which was followed by a magnitude class 6.0 earthquake at 38.3°N20.4°E that occurred on March 25, 2007.

In order to determine the occurrence time of the impending EQs, we currently apply the procedure explained in the previous subsection by studying the seismicity in the areas A, B, C (see Fig.9). Since the result should exhibit *spatial* scale invariance, the epicenter(s) will lie either in the area B or in C depending on whether the areas A and B or A and C show *true* coincidence.



FIG. 1: (color online) Examples of log-log plot of the fluctuations $\Delta \chi_l$ of the natural time under time reversal versus the scale l for fGn (a) and fBm (d). (b) and (e) depict the values of the scaling exponent χ_H , introduced here, versus the true exponent H for fGn and fBm, respectively. For the sake of comparison, (c) and (f) are similar to (b) and (e), respectively, but for the DFA exponent α_{DFA} . The (blue) dotted curves show the $\pm \sigma$ deviation from the average value (obtained after 10² runs) depicted by the (red) solid curves. The (green) dashed straight lines correspond to the ideal behavior of each exponent and have been drawn as a guide to the eye.



FIG. 2: (color online) The log-log plot of $\Delta \chi_l$ versus the scale *l* for three SES activities (K1, K2 and U) and seven AN (n1-n5, n7 and n10) treated in Ref.19 (cf. these signals have enough number of pulses in order to apply the present analysis). The values of $\Delta \chi_l$ are divided by the corresponding values $\Delta \chi_5$ at the scale l = 5.

V. CONCLUSIONS

First, the scale dependence of the fluctuations of the natural time under time reversal distinguish similar looking electric signals emitted from systems of different dynamics providing a useful tool for the determination of the scaling exponent. In particular, SES activities (*critical* dynamics) are distinguished from noises emitted from man-made electrical sources.

Second, recent data of SES activities are presented

which confirm that the value of the entropy in natural time as well as its value under time reversal are smaller than that of a "uniform" distribution.



FIG. 3: (color online) (a) The electric signal recorded on November 14, 2006 at PIR station (sampling rate $f_{exp} = 1$ Hz). The signal is presented here in normalized units (see the text). The corresponding dichotomous representation is shown with the dotted (blue) line. The gray shaded area shows an additional signal (consisting of pulses of smaller duration) superimposed on the previously mentioned signal. (b) The signal belonging to the gray shaded area in (a) is given here in an expanded time scale, while its dichotomous representation is marked by the dotted (blue) line. (c) How the signals in (a) and (b) are read in natural time: the continuous (red crosses) and dotted (blue asterisks) bars correspond to the durations of the dichotomous representations marked in (a) and (b), respectively.

APPENDIX A: THE CASE OF SIGNALS THAT EXHIBIT SHORT-RANGE TEMPORAL CORRELATIONS

Here, we present results of modeling Q_k by shortranged temporal correlated time-series. Two examples were treated by numerical simulation: (i) A stationary autoregressive process $Q_k = aQ_{k-1} + g_k + c$, |a| < 1, where g_k are Gaussian IID random variables, and c stands for an appropriate constant to ensure positivity of Q_k . (ii) $Q_k = |aQ_{k-1} + g_k|$. Figure 10(a) depicts the results for S, S_- and κ_1 for the first example versus the number of Q_k , whereas Fig.10(b) refers to the second example. In both cases S and S_- converge to S_u whereas κ_1 to the value $\kappa_u = 1/12$ corresponding to the "uniform" distribution. For the reader's convenience, the values of



FIG. 4: Map of the area surrounding the station PIR (solid dot) and the epicenters of the EQs (stars) with magnitude 5.2-units and 5.8-units that occurred on January 18 and February 3, 2007, respectively. The seismicity subsequent to the SES activities on November 14, 2006, has been studied in the gray shaded areas (the large and the small area are designated A and B, respectively).



FIG. 5: (color online) The normalized power spectrum(red) $\Pi(\phi)$ of the seismicity as it evolves event by event (whose date and time of occurrence are written in green in each panel) after the initiation of the SES activity on November 14, 2006. The two excerpts presented here, refer to the periods: (a) December 25, 2006, to January 17, 2007, and (b): January 18 to January 31, 2007, and correspond to the area B with $M_{thres} = 3.4$. In each panel only the spectrum in the range $\phi \in [0, 0.5]$ (for the reasons discussed in Refs.1,35) is depicted (separated by the vertical lines), whereas the $\Pi(\phi)$ of Eq.(13) is depicted by blue color. The minor horizontal ticks for ϕ are marked every 0.1.



FIG. 6: (color online) Evolution of the quantities κ_1 , S and S_- ; they are shown in (a), (b) and (c), respectively, for three magnitude thresholds, i.e., $M \ge 3.2, 3.4$ and $M \ge 3.6$, for both areas, i.e., the large (area A) and the small (area B). The solid horizontal lines correspond to $\kappa_1 = 0.07$ and $S_u = \ln 2/2 - 1/4$.

 S_u and κ_u are designated by the horizontal solid black lines.

APPENDIX B: THE SEISMIC ACTIVITY THAT FOLLOWED THE RECENT SES ACTIVITIES OF FIGS.8(A),(B)

Considering the Athens observatory preliminary catalogue, the seismic activity $(M_{thres} \geq 3.0)$ that occurred



FIG. 7: (color online) The average distance $\langle D \rangle$ between the calculated and the theoretical $\Pi(\phi)$ curves versus the conventional time. The calculation of $\langle D \rangle$ is made upon the occurrence of every consecutive earthquake when starting the calculation after the initiation of the SES activities on November 14, 2006 (depicted in Figs.3(a),(b)) for the large (area A) and the small (area B) for the three magnitude thresholds $M_{thres} = 3.2$, 3.4 and 3.6. The black solid line corresponds to the mean value obtained from the three thresholds.

in area A (see Fig.9) after the initiation of the SES activity on July 2, 2007 (Fig.8(a)) until 03:27 UT of September 25, 2007 is shown in Fig.11(a). The evolution of the corresponding parameters κ_1 , S, S_- and $\langle D \rangle$ calculated for three magnitude thresholds, i.e., $M_{thres} = 3.0, 3.1$ and 3.2 are shown in Figs.11(b), (c) and (d) respectively. To investigate the spatial invariance, the computation was repeated for several smaller areas, three of which are shown in Figs.12,13 and 14 (which are *different* from the areas B and C of Fig.9) along with the evolution of the corresponding parameters. The same was repeated for an area (see Fig.15) somewhat larger than A. An inspection of all these figures, i.e., Figs.11 to15, suggests that presumably a true coincidence has just been approached, thus being very close to the critical point.

APPENDIX C: THE SEISMIC ACTIVITY THAT FOLLOWED THE MOST RECENT SES ACTIVITIES AT PAT AND PIR

Here, we report the update results of the seismic activity that followed the SES activity at PAT on October 9, 2008³⁸ and the SES activity at PIR on December 12, 2008³⁹ by following the procedure described by Sarlis et al.⁴⁰. The subsequent seismicity of the former SES activity was studied in the area $N_{37.5}^{38.6}E_{19.8}^{23.3}$ while that of the latter in the selectivity map of PIR depicted in Fig. 16. The results, when considering the seismicity until early in the morning of February 2, 2009, for magnitude threshold $M_{thres}=3.3$, are shown in Figs. 17 and 18 for the former and the latter SES activities at PAT and PIR, respectively. An inspection of these figures reveals that in both areas the probability $Prob(\kappa_1)$ versus κ_1 -calculated in all the possible regions of each area as described in Ref.⁴⁰- maximizes at $\kappa_1 \approx 0.070$ upon the occurrence of



FIG. 8: Recent SES activities recorded at PAT.

the events marked with arrows, thus probably indicating the approach to the critical point.

Note added on February 20, 2009: Actually, at 23:16 UT on February 16, 2009 a strong earthquake with magnitude Ms(ATH)=6.0 (i.e., $M_L(ATH)=5.5$) occurred with an epicenter at 37.1°N20.8°E, which clearly lies inside the selectivity map of PIR depicted in Fig.16.

We now present the results, when considering the seismicity until early in the morning of February 19, 2009. Figure 19, depicts the results for $M_{thres}=3.2$ in the selectivity map of PAT (i.e., in the area $N_{37.5}^{38.6}E_{19.8}^{23.3}$), which show that the aforementioned probability $Prob(\kappa_1)$ versus κ_1 maximizes at $\kappa_1 \approx 0.070$ upon the occurrence of a $M_L=3.2$ event at 00:35 UT on February 19, 2009 with epicenter at 37.8°N21.2°E. As for the results in the selectivity map of PIR (i.e., the shaded area in Fig.16), they are shown in Fig.20 for $M_{thres}=3.4$ and reveal the following: After the aforementioned strong earthquake on February 16, 2009, the bimodal curve $Prob(\kappa_1)$ versus κ_1 shows a clear local maximum at $\kappa_1 \approx 0.070$ upon the oc-



FIG. 9: A map showing the areas A, B and C at which the analysis of the seismicity in natural time is currently carried out.



FIG. 10: (color online) The thick colored lines in (a) and (b) depict the average value of S (red), S_{-} (blue) and κ_1 (green) versus the number of Q_k for the two examples (i) and (ii), respectively, mentioned in the Appendix. The thinner colored lines refer to the $\pm \sigma$ deviation from the average value.

currence of the $M_L=3.5$ event at 00:30 UT on February 18, 2009 with an epicenter at $34.9^{\circ}N23.4^{\circ}E$. This local maximum does exhibit magnitude threshold invariance (since it also appears when investigating $M_{thres}=3.3$ and $M_{thres}=3.5$).

Updated note on March 13, 2009: Several hours before the occurrence of the aforementioned magnitude 6.0 earthquake on February 16, 2009, an electrical anomaly of significant amplitude, see Fig.21, appeared at PIR. It basically consists of three bay like transient changes, thus having a form essentially different than the SES activities that preceded the major earthquakes on February 14, 2008 and June 8, 2008. This electrical anomaly could be in principle attributed to the 6.0 earthquake on February 16, 2009 that occurred almost 11 hours later, but this earthquake was also preceded by the SES activity at PIR on December 12, 2008. Alternatively, this anomaly might be related to a new impending earthquake. For this reason, a natural time analysis of the seismicity subsequent to the electrical anomaly of February 16, 2009 was carried out. In this calculation we considered the events that occurred in the PIR selectivity map shown in Fig.22 (cf. this, which is slightly different than that depicted in Fig.16, has been drawn by taking into account the whole region to the west of the Hellenides). The results of the analysis shown in Fig.23 indicate that for $M_{thres}=3.6$ and 3.5 (see a and b, respectively) the probability $Prob(\kappa_1)$ maximizes at $\kappa_1 \approx 0.070$ upon the occurrence of a $M_L=3.6$ event at 02:48 UT on March 9, 2009, while for $M_{thres}=3.3$ the maximization occurs upon the occurrence of a $M_L=3.3$ event on March 10, 2009. Interestingly, a similar analysis (subsequent to the SES activity at PAT) in the selectivity map of PAT, i.e., in the

- * Electronic address: pvaro@otenet.gr
- ¹ P. A. Varotsos, N. V. Sarlis, and E. S. Skordas, Practica of Athens Academy **76**, 294 (2001).
- ² P. A. Varotsos, N. V. Sarlis, and E. S. Skordas, Phys. Rev. E **66**, 011902 (2002).
- ³ P. Varotsos, N. Sarlis, and E. Skordas, Acta Geophys. Pol. 50, 337 (2002).
- ⁴ P. Varotsos and K. Alexopoulos, *Thermodynamics of Point Defects and their Relation with Bulk Properties* (North Holland, Amsterdam, 1986).
- ⁵ P. Varotsos, K. Alexopoulos, K. Nomicos, and M. Lazaridou, Nature (London) **322**, 120 (1986).
- ⁶ P. Varotsos, K. Alexopoulos, K. Nomicos, and M. Lazaridou, Tectonophysics **152**, 193 (1988).
- ⁷ P. Varotsos, N. Sarlis, and E. Skordas, Phys. Rev. Lett. 91, 148501 (2003).
- ⁸ P. Varotsos, N. Sarlis, and E. Skordas, Appl. Phys. Lett. 86, 194101 (2005).
- ⁹ P. Varotsos, N. Sarlis, and E. Skordas, Proc. Jpn. Acad. Ser. B **77**, 93 (2001).
- ¹⁰ P. Varotsos, N. Sarlis, and M. Lazaridou, Phys. Rev. B **59**, 24 (1999).
- ¹¹ N. Sarlis, M. Lazaridou, P. Kapiris, and P. Varotsos, Geophys. Res. Lett. **26**, 3245 (1999).
- ¹² P. Varotsos, N. Sarlis, M. Lazaridou, and P. Kapiris, J. Appl. Phys. 83, 60 (1998).
- ¹³ P. Varotsos, *The Physics of Seismic Electric Signals* (TER-RAPUB, Tokyo, 2005).
- ¹⁴ P. Varotsos, N. Sarlis, E. Skordas, and M. Lazaridou, Appl. Phys. Lett. **90**, 064104 (2007).
- ¹⁵ S. Abe, N. V. Sarlis, E. S. Skordas, H. K. Tanaka, and P. A. Varotsos, Phys. Rev. Lett. **94**, 170601 (2005).
- ¹⁶ P. A. Varotsos, N. V. Sarlis, and E. S. Skordas, Phys. Rev. E 68, 031106 (2003).
- ¹⁷ P. A. Varotsos, N. V. Sarlis, E. S. Skordas, and M. S. Lazaridou, Phys. Rev. E **70**, 011106 (2004).
- ¹⁸ P. A. Varotsos, N. V. Sarlis, E. S. Skordas, and M. S. Lazaridou, Phys. Rev. E **71**, 011110 (2005).
- ¹⁹ P. A. Varotsos, N. V. Sarlis, H. K. Tanaka, and E. S. Skordas, Phys. Rev. E **71**, 032102 (2005).
- ²⁰ B. Lesche, J. Stat. Phys. **27**, 419 (1982).
- ²¹ B. Lesche, Phys. Rev. E **70**, 017102 (2004).
- ²² P. A. Varotsos, N. V. Sarlis, E. S. Skordas, H. K. Tanaka, and M. S. Lazaridou, Phys. Rev. E **73**, 031114 (2006).
- ²³ P. A. Varotsos, N. V. Sarlis, E. S. Skordas, H. K. Tanaka, and M. S. Lazaridou, Phys. Rev. E **74**, 021123 (2006).
- ²⁴ P. A. Varotsos, N. V. Sarlis, and E. S. Skordas, Phys. Rev. E 67, 021109 (2003).
- ²⁵ C.-K. Peng, S. V. Buldyrev, S. Havlin, M. Simons, H. E. Stanley, and A. L. Goldberger, Phys. Rev. E **49**, 1685 (1994).
- ²⁶ S. V. Buldyrev, A. L. Goldberger, S. Havlin, R. N. Man-

area $N_{37.5}^{38.6} E_{19.8}^{23.3}$, reveals that a maximization of $\text{Prob}(\kappa_1)$ at $\kappa_1 \approx 0.070$ also occurs on March 10, 2009 (see Fig.24). We are currently investigating the spatial and magnitude threshold invariance of the aforementioned maxima to examine whether they actually point to an approach to the critical point.

tegna, M. E. Matsa, C.-K. Peng, M. Simons, and H. E. Stanley, Phys. Rev. E **51**, 5084 (1995).

- ²⁷ A. Weron, K. Burnecki, S. Mercik, and K. Weron, Phys. Rev. E **71**, 016113 (2005).
- ²⁸ B. Mandelbrot and J. R. Wallis, Water Resources Research 5, 243 (1969).
- ²⁹ J. Szulga and F. Molz, J. Stat. Phys. **104**, 1317 (2001).
- ³⁰ M. Frame. Mandelbrot. В. and N. Neger, fractal Geometry, Yale University, available from http://classes.yale.edu/fractals/, see http://classes.yale.edu/Fractals/RandFrac/fBm/fBm4.html.
- ³¹ See EPAPS document No. E-PLEEE8-74-190608 for additional information, originally from P. Varotsos, N. Sarlis, E. Skordas, H. Tanaka, and M. Lazaridou, Phys. Rev. E 74, 021123 (2006). For more information on EPAPS, see http:// www.aip.org/pubservs/epaps.html.
- ³² P. A. Varotsos, Proc. Japan Acad., Ser. B **82**, 86 (2006).
- ³³ P. Varotsos, N. Sarlis, E. Skordas, and M. Lazaridou (2006), arXiv:cond-mat/0611437v1 [cond-mat.stat-mech].
- ³⁴ The time of the impending earthquake can be determined by means of the procedure described in EPAPS Document No. E-PLEEE8-73-134603 for additional information, originally from P.A. Varotsos, N.V. Sarlis, E.S. Skordas, H.K. Tanaka and M.S. Lazaridou, Phys. Rev. E 73, 031114 (2006). For more information on EPAPS, see http://www.aip.org/pubservs/ epaps.html.
- ³⁵ P. Varotsos, N. Sarlis, H. Tanaka, and E. Skordas, Phys. Rev. E **72**, 041103 (2005).
- ³⁶ In the natural time analysis of the seismicity we study the evolution of the pair (χ_k, Q_k), where Q_k is proportional to the seismic moment M_0 . Following Refs.[1-4,9], for the reasons of this calculation the formulae $M_0 = 10^{M_L+10.5}$ (Nm), for $M_L \leq 3.6$ and $M_0 = 10^{1.5M_L+8.7}$ (Nm), for $3.6 < M_L \leq$ 5.0 proposed by the Global Seismological Services were used. Then, the normalized power spectrum is given[1-4,9] by

$$\Pi(\phi) \equiv \left| \sum_{k=1}^{N} p_k \exp\left(i2\pi\phi\chi_k\right) \right|^2,$$

where $p_k = (M_0)_k / \sum_{n=1}^N (M_0)_n$.

- ³⁷ P. Varotsos, N. Sarlis, E. Skordas, and M. Lazaridou (2007), arXiv:cond-mat/0703683v5 [cond-mat.stat-mech].
- ³⁸ P. A. Varotsos, N. V. Sarlis, and E. S. Skordas (2008), arXiv:cond-mat/0711.3766v5 [cond-mat.stat-mech].
- ³⁹ P. A. Varotsos, N. V. Sarlis, and E. S. Skordas, Submitted to Chaos (2009).
- ⁴⁰ N. V. Sarlis, E. S. Skordas, M. S. Lazaridou, and P. A. Varotsos, Proc. Japan Acad., Ser. B 84, 331 (2008).



FIG. 11: (color online) Results of the analysis of seismicity in the dark gray shaded area of (a) after the SES initiation on July 2, 2007.



FIG. 12: (color online) Results of the analysis of seismicity in the dark gray shaded area of (a) after the SES initiation on July 2, 2007.



FIG. 13: (color online) Results of the analysis of seismicity in the dark gray shaded area of (a) after the SES initiation on July 2, 2007.



FIG. 14: (color online) Results of the analysis of seismicity in the dark gray shaded area of (a) after the SES initiation on July 2, 2007.



FIG. 15: (color online) Results of the analysis of seismicity in the dark gray shaded area of (a) after the SES initiation on July 2, 2007.



FIG. 16: The selectivity map of PIR (shaded area) along with the selectivity map of PAT (rectangular area)



FIG. 17: The $\operatorname{Prob}(\kappa_1)$ versus κ_1 of the seismicity for $M_{thres}=3.3$ subsequent to the SES activity recorded at PAT on October 9, 2008 within the selectivity map of PAT shown in Fig. 16. For the sake of clarity, only the last 7 events are depicted. The arrows mark the maxima at $\kappa_1=0.070$ (see the text)



FIG. 18: The $\operatorname{Prob}(\kappa_1)$ versus κ_1 of the seismicity for $M_{thres}=3.3$ subsequent to the SES activity recorded at PIR on December 12, 2008 within the selectivity map of PIR shown in Fig. 16. For the sake of clarity, only the last 12 events are depicted. The arrows mark the maxima at $\kappa_1=0.070$ (see the text).



FIG. 19: $\operatorname{Prob}(\kappa_1)$ versus κ_1 of the seismicity for $M_{thres}=3.2$ subsequent to the SES activity recorded at PAT on October 9, 2008 within the area $N_{37.5}^{38.6} E_{19.8}^{23.3}$. For the sake of clarity, only the last 6 events are depicted. $\operatorname{Prob}(\kappa_1)$ exhibits a maximum at $\kappa_1=0.070$ upon the occurrence of the 3.2 event on February 19, 2009 (see the text).



FIG. 20: $\operatorname{Prob}(\kappa_1)$ versus κ_1 of the seismicity for $M_{thres}=3.4$ subsequent to the SES activity recorded at PIR on December 12, 2008 within the selectivity map of PIR shown in Fig. 16. For the sake of clarity, only the last 17 events are depicted. The arrows mark the maxima of $\operatorname{Prob}(\kappa_1)$ at $\kappa_1=0.070$ (see the text).



FIG. 21: Electrical recordings from PIR; for the sites of the electrodes see ${\rm Ref.}^{31}.$



FIG. 22: The selectivity map of PIR when including the whole region to the west of Hellenides (shaded area). The rectangular area corresponds to the selectivity map of PAT.



FIG. 23: $\operatorname{Prob}(\kappa_1)$ versus κ_1 of the seismicity in the shaded region of Fig.22 for $M_{thres}=3.6$ (a), 3.5 (b) and 3.3 (c) subse-



FIG. 24: Prob(κ_1) versus κ_1 of the seismicity for M_{thres}=3.3 subsequent to the SES activity recorded at PAT on October 9, 2008 within the area N^{38.6}_{37.5}E^{23.3}_{19.8}. For the sake of clarity, only the last 6 events are depicted. Prob(κ_1) exhibits a maximum at κ_1 =0.070 upon the occurrence of the event at 15:22 UT on March 10, 2009.

TABLE I: The catalogue ($M_{thres} = 3.2$) of the Institute of Geodynamics of the National Observatory of Athens (GI-NOA) (available from http://www.gein.noa.gr/services/monthly-list.html on February 5, 2007) for the Large area (area A) during the following period: From the initiation of the SES activity almost at 17:00 UT on November 14, 2006 (Fig.1 of the main text) until the occurrence of the 5.8-units earthquake on February 3, 2007. Note that Ms(ATH)=M+0.5, where M stands for the local magnitude ML or the "duration" magnitude MD. Two events reported by GI-NOA: at 17:32:23.7 on January 28, 2007, and at 5:11:06.5 on January 30, 2007, were omitted since they could not be independently confirmed by other seismic networks operating at that time.

N	Year	Month	Day	UT	$Lat.(^{o}N)$	$\operatorname{Lon.}(^{o}\mathrm{E})$	depth(km)	М
1	2006	11	16	4:20:40.0	35.66	23.26	54	3.5
2	2006	11	16	12:30:50.3	35.44	23.42	5	3.7
3	2006	11	16	18:50:02.5	36.97	23.11	25	3.7
4	2006	11	16	21:53:12.5	35.72	24.33	10	3.4
5	2006	11	17	16:31:26.4	36.69	23.32	24	3.2
6	2006	11	18	3:48:58.3	34.79	24.07	16	3.6
7	2006	11	18	21:51:21.8	34.78	24.36	34	3.2
8	2006	11	19	9:31:09.1	35.73	21.95	19	3.6
9	2006	11	19	10:54:52.3	35.99	23.65	10	3.2
10	2006	11	19	14:11:45.4	35.75	22.55	24	3.3
11	2006	11	21	5:57:05.5	34.81	24.28	54	3.3
12	2006	11	21	8:42:06.0	37.42	21.96	19	3.2
13	2006	11	21	12:50:34.0	35.45	23.90	10	3.3
14	2006	11	24	17:16:58.4	37.16	21.58	12	4.1
15	2006	11	24	17:42:56.3	35.60	23.50	7	3.3
16	2006	11	24	20:26:02.9	37.12	21.56	11	3.5
17	2006	11	24	21:41:44.1	37.21	21.63	8	3.3
18	2006	11	25	3:14:31.2	34.94	23.91	5	3.2
19	2006	11	25	4:55:01.3	37.15	21.86	9	3.3
20	2006	11	25	9:30:18.1	36.75	21.73	12	3.6
21	2006	11	26	17:58:05.5	35.45	23.30	11	4.1
22	2006	11	26	19:45:54.7	35.40	23.27	2	3.3
23	2006	11	26	23:57:49.1	35.47	23.36	5	3.4
24	2006	11	27	3:12:49.8	35.42	23.31	9	3.3
25	2006	11	27	11:58:03.7	35.53	23.43	24	3.2
26	2006	11	28	9:07:38.6	36.07	22.35	10	3.3
27	2006	11	28	15:07:43.1	36.06	22.42	43	3.6
28	2006	11	28	15:42:59.4	35.71	22.11	10	3.4
29	2006	11	28	22:54:24.0	37.06	21.51	12	3.2
30	2006	11	29	14:11:02.8	35.97	23.11	19	3.5
31	2006	11	30	23:30:58.6	35.43	23.32	3	3.2
32	2006	12	1	2:37:02.1	35.58	23.60	10	3.5

N	Year	Month	Day	UT	$Lat.(^{o}N)$	$\operatorname{Lon.}(^{o}\mathrm{E})$	depth(km)	М
33	2006	12	1	19:57:55.4	37.07	21.47	3	3.2
34	2006	12	1	22:49:35.6	35.72	22.52	114	3.7
35	2006	12	2	0:29:20.8	34.74	22.87	45	3.2
36	2006	12	2	10:30:35.3	37.42	22.04	4	3.3
37	2006	12	3	1:53:22.4	36.72	21.81	24	3.5
38	2006	12	3	2:37:36.7	36.40	21.66	10	3.2
39	2006	12	4	19:32:24.0	34.99	23.44	14	3.4
40	2006	12	7	22:22:49.4	35.73	23.17	32	3.2
41	2006	12	8	10:54:40.0	34.95	23.42	5	3.3
42	2006	12	8	12:48:15.5	37.02	21.93	10	3.2
43	2006	12	8	19:57:27.2	35.06	23.50	7	3.6
44	2006	12	9	22:45:56.5	35.89	23.38	24	3.4
45	2006	12	10	3:29:46.1	34.84	24.48	10	3.2
46	2006	12	10	12:41:27.6	35.80	23.04	59	3.2
47	2006	12	11	3:36:22.0	35.74	24.08	16	3.4
48	2006	12	11	19:11:29.4	35.93	23.40	15	3.2
49	2006	12	12	3:40:55.0	35.54	22.81	17	3.7
50	2006	12	13	9:45:40.0	37.25	21.94	20	3.2
51	2006	12	14	16:44:02.0	35.95	23.46	10	3.3
52^{-1}	2006	12	15	4:40:12.5	35.01	23.37	80	3.4
53	2006	12	15	10:23:32.1	36.09	22.25	45	3.7
54	2006	12	16	10:43:15.4	34.85	24.33	6	3.6
55	2006	12	16	15:23:33.2	34.92	23.45	$\overset{\circ}{5}$	3.7
56	2006	12	17	4:44:38.7	34.73	24.00	30	3.5
57	2006	12	17	$4 \cdot 44 \cdot 42.7$	35.15	24.12	59	3.5
58	2006	12	17	10:20:49.7	36.06	21.76	37	3.3
59	2006	12	17	14.13.23.4	36.21	21.70	10	34
60	2006	12	17	20: 0:20.6	34.84	24.21	34	4.0
61	2006	12	17	22.38.375	36.65	21.00	10	3.2
62	2006	12	20	4.34.09.2	37.48	21.00 21.50	10	3.2
63	2006	12	20	4:45:15.0	35.42	21.00	10	3.5
64	2006	12	20	19.27.325	34.60	23 79	11	3.8
65	2006	12	20	20.17.02.7	36.68	21.50	4	3.2
66	2006	12	20	6:58:02.3	34 94	24.03	10	3.6
67	2000	12	24	7.12.114	36.29	22.00	24	3.8
68	2000	12	25	14.08.594	34.83	22.20	39	4.4
69	2000	12	25	14.15.501	34 99	22.00 23.04	39	4.0
70	2000	12	25	14.18.50.7	35.09	23.04	27	1.0
71	2000	12	25	14.10.00.7 14.22.30.3	35.05	25.00 22.95	18	3.9
72	2000	12	25	14:57:00 5	35.04	22.55	10	4.1
73	2000	12	25	17.20.17.0	35.02	22.04	5	3.0
74	2000	12	25	18.13.44 5	35.00	23.02	10	3.2 3.7
75	2006	12	20	6.39.10 /	37 53	20.29	10	3.7 3.9
76	2000	12	20	2.12.13.4	36 77	21.01	5	0.2 २२
70	2000	12	29 30	2.00.04.9 99.51.38 0	35.30	21.01	12	 ვე
11	2000	12	3 U	22:01:00.0	əə.ə9	20.04	19	3.2

TABLE I: Continued

N	Year	Month	Day	UT	$Lat.(^{o}N)$	$Lon.(^{o}E)$	depth(km)	Μ
78	2006	12	31	7:00:51.9	35.85	22.08	10	3.2
79	2006	12	31	18: 6:27.5	35.14	22.76	12	3.7
80	2007	1	1	15:27:48.7	34.72	24.12	23	3.5
81	2007	1	3	15:04:04.9	35.37	23.29	10	3.4
82	2007	1	3	18:18:34.5	36.54	21.74	26	3.8
83	2007	1	4	7:57:08.3	37.37	21.56	15	3.3
84	2007	1	4	14:55:24.6	36.29	21.92	10	3.2
85	2007	1	4	17:42:54.7	34.91	23.61	45	3.3
86	2007	1	4	20:41:40.9	34.89	24.13	48	3.2
87	2007	1	4	22:47:39.1	36.96	21.07	25	3.2
88	2007	1	7	14:07:11.5	37.10	21.93	26	3.2
89	2007	1	8	16:13:02.8	35.10	23.03	10	3.4
90	2007	1	9	9:55:29.2	36.70	21.55	31	3.4
91	2007	1	9	15:54:47.3	35.89	23.61	3	3.2
92	2007	1	9	23:09:20.0	36.21	22.70	116	3.6
93	2007	1	11	5:50:39.5	35.02	22.48	38	3.8
94	2007	1	13	11:12:43.3	35.48	23.51	8	3.4
95	2007	1	14	9:09:23.0	35.30	23.38	3	3.8
96	2007	1	14	16:43:01.6	35.06	23.20	85	4.1
97	2007	1	15	0:55:18.8	37.47	21.07	15	3.2
98	2007	1	15	3:13:45.8	36.06	22.40	13	3.4
99	2007	1	15	6:56:47.0	35.42	23.56	33	3.3
100	2007	1	15	17:50:48.7	36.56	21.65	5	3.4
101	2007	1	17	1:52:18.8	36.20	21.58	34	3.7
102	2007	1	17	3:22:53.9	35.34	23.53	12	3.7
103	2007	1	18	22:25:23.0	34.84	22.67	39	4.7
104	2007	1	18	23:31:32.7	34.76	21.36	10	3.4
105	2007	1	19^{-3}	8:42:22.8	34.72	22.57	38	4.4
106	2007	1	20	7:05:37.0	34.95	24.09	19	3.5
107	2007	1	20	7:24:24.6	34.86	23.96	19	3.4
108	2007	1	20	10:24:55.4	36.88	22.10	10	3.2
109	2007	1	20	11:57:23.7	36.69	22.47	10	3.5
110	2007	1	20	23:50:04.1	34.93	24.40	28	3.3
111	2007	1	21	17.52.349	35.46	23.46	11	34
112	2007	1	25	10.24.47.3	35.33	23.40	2	3.3
113	2007	1	25	21.09.457	35.09	23.25	22	4.3
114	2007	1	28	15.19.23.6	36 29	23.23	25	3.4
115	2007	1	20	10.12.16 /	36.95	20.11	10	3.1
116	2007	1	30	5.11.08.9	35.94	21.15	28	0. 1 २२
117	2007	1	30	10.38.42 7	36.98	20.41	12	0.0 3 5
118	2007	1	30	10.30.42.7 21.11.43.0	36.13	21.10 22.17	30	0.0 3 3
110	2007	1	21	4.56.50 1	35.40	22.17	36	0.0 3.4
119	2007	1	21	4.50.59.1	34.67	22.10	20	3.4
120	2007	1	01 21	14.40.09.7	96.96	22.42 22.48	- <u>∠</u> ∂ 22	0.0 3 G
121	2007	1	01 1	16.45.95 9	30.20 34.60	22.40 22.46	∠ວ ງ1	0.0 2 0
122	2007	2	1	10.40.50.5 91.06-54.7	04.09 26.06	22.40 01.60	∠1 40	ა.ყ ეი
123 194	2007	2	1	21:00:04.7 6:30:05 6	30.00 25.91	21.02 22.25	42 22	3.2 9.9
124 195	2007	2	2	0:39:00.0 19:07:59 1	00.41 27 00	20.20 01 69	ى م	ე.ე ე∡
120	2007	2	2	13:27:33.1	31.22	21.03	4	3.4

TABLE I: Continued