Created-by-current states in long Josephson junctions

4 Sep 2008 T.L. BOYADJIEV^{1,2}, O.YU. ANDREEVA³, E.G. SEMERDJIEVA^{1,4} and YU.M. SHUKRINOV^{1,5}

JINR, Dubna, Moscow Region, 141980, Russia

Sofia University, Sofia, 1164, Bulgaria

Tumen Thermal Networks OAO TGK 10, Tobolsk, 626150, Russia

University of Plovdiv, Plovdiv, 4000, Bulgaria

Physical Technical Institute, Dushanbe, 734063, Tajikistan

PACS 74.50.+r - Tunneling phenomena; point contacts, weak links, Josephson effects

 ⁵ Physical Technical Institute, Dushanbe, 734063, Tajikista
 PACS 74.50.+r – Tunneling phenomena; point contact PACS 05.45.-a – Nonlinear dynamics and chaos
 Abstract. - Critical curves "critical current - external with inhomogeneity and variable width are studied. No of magnetic field where some fluxon states are stable junction is different from zero. Position and size of such its geometry, parameters of inhomogeneity and form right) pure fluxon states are appeared in the inhom in the junction length. We demonstrate new bifurch inhomogeneity and amplitude of the Josephson current of Owen and Scalapino, [1] the long Josephson junctions (LJJ) are attracting attention both experimentalists and theoreticians. They are good candidates for a wide range of applications such as superconducting quantum interfer-Abstract. - Critical curves "critical current - external magnetic field" of long Josephson junctions with inhomogeneity and variable width are studied. We demonstrate the existence of the regions of magnetic field where some fluxon states are stable only, if the external current through the junction is different from zero. Position and size of such regions depend on length of the junction, its geometry, parameters of inhomogeneity and form of the junction. The noncentral (left and right) pure fluxon states are appeared in the inhomogeneous Josephson junction with increase in the junction length. We demonstrate new bifurcation points with change in width of the inhomogeneity and amplitude of the Josephson current through the inhomogeneity.

 \bigcirc of applications such as superconducting quantum interfer- \bigcirc ence devices, Josephson voltage standards, logic elements, and Josephson flux-flow oscillators because of their non-linear behavior and quantum effects. [2–4] From the other \geq side, the nonlinearity makes the LJJ a very complex system, and some aspects of its properties are not investigated up to now. Particularly, the experimental methods rightarrow to determine the vortex structure under the critical curve are not developed well vet. But there are different kind of stable states in the LJJ, which bifurcation curves are close to the critical curve of the junction and manifest themselves in the experiments. [2, 5] Particularly, some "metastable" states (experimental points under overlap curve) were observed in Ref. [6]. Static bound states of fluxons in LJJ with artificial inhomogeneities were experimentally found in Ref. [5]. Experimental and theoretical investigation of such states is an interesting and actual problem.

An important step of the sample characterization in the experiment with LJJs is the measurement of its static properties, i.e. the dependence of the critical current I_c on external magnetic field H. This dependence provides

a technique to evaluate the several important parameters of LJJ from the experiment, particularly, a critical current density, a magnetic flux penetration field, an effective magnetic thickness. [7] Since the solution of the corresponding boundary value problem (BVP) cannot be performed analytically in general case, the most straightforward way to study the dependence $I_c(H)$ is its numerical simulation.

In our previous study we investigated the stability of the vortices in LJJ with inhomogeneities (LJJI) [8,9] and vortex structure in exponentially shaped Josephson junctions (ESJJ). [10–12] The obtained bifurcation curves of the mixed fluxon-antifluxon states in LJJI [9] and pure fluxon states in ESJJ demonstrate the intervals of magnetic field, where these states are stable only, if the current through the junction is not equal to zero. These peculiarities were not mention at that time. We call these states as the CbC-states (created-by-current), and the corresponding regions of the external magnetic field as CbC-regions.

In this Letter we study the CbC-states of mixed fluxon states $\Phi^{-1}\Phi^1$, $\Phi^1\Phi^{-1}$ and pure fluxon states Φ^n . We demonstrate new bifurcation points with change in width of the inhomogeneity and amplitude of the Josephson current through the inhomogeneity. The influence of the model parameters on CbC regions in the LJJI and ESJJ is investigated.

Method of calculation. – First we start with LJJI case. We consider the Josephson junctions in the overlap geometry. [2] In order to study a linear stability of the static distribution $\varphi(x)$ of magnetic flux $\phi(t, x)$, we write the perturbed expression for $\phi(t, x)$ as $\phi(t, x) = \varphi(x) + \varepsilon \xi(t)\psi(x)$, where ε is a parameter [13, 14]. Then in a first approximation with respect to ε , we get the following boundary value problem (BVP) for $\varphi(x)$

$$-\varphi_{xx} + j_C(x)\sin\varphi + \gamma = 0, \quad \varphi(-l) = \varphi(l) = h_e \quad (1)$$

and the corresponding Sturm-Liouville problem (SLP)

$$-\psi_{xx} + \sigma(x)\psi_x + q(x)\psi = \lambda\psi \quad \psi_x(-l) = \psi_x(l) = 0 \quad (2)$$

Here all the quantities are in dimensionless form [3], h_e is the external magnetic field, γ is the external current, and λ is an eigenvalue of the SLP, 2*l* is a length of the junction. The function $j_C(x)$ represents the Josephson current amplitude, so $|j_C(x)| \leq 1$. Because of the nonlinearity the BVP, eq.(1) has more then one solution for given parameters (counted set in case $h_e = 0, \gamma = 0, l \to \infty$) [15]. The potential $q(x) = j_D(x) \cos \varphi(x)$ is defined by the concrete static solution $\varphi(x)$, which is asymptotically stable with respect to small perturbations, if $\lim_{t\to\infty} |\xi(t)| \to 0$. As the junction's length $l < \infty$ in case under consideration and the potential $|q(x)| \leq 1$, the SLP eq.(2) is regular, so there exists a discrete low-bounded set of real eigenvalues $\lambda_n, n = 0, 1, 2, \dots$ and the minimal one $\lambda_0 \ge -1$. [16] The corresponding eigenfunctions satisfy the norm condition $\int \psi^2(x) \, dx - 1 = 0$. In case $\lambda_0 < 0$ the static solution

 $\varphi(x)$ is unstable, i.e. $|\xi(t)| \to \infty$, when $t \to \infty$. Solutions of the BVP eq.(1), eq.(2) describe the different fluxon states in LJJ. Numerical simulation simplifies the study and makes it possible to estimate the range of variation of the parameters in which one can expect stability or instability of the magnetic flux distributions in the Josephson junction.

The models of LJJ depend on m parameters, such as junction's length 2l, external magnetic field h_e , external current γ and so on. Let us denote the vector of all the parameters by p. In all cases we presume that the possible solutions of non-linear BVP eq.(1) continuously depend on the set p, i.e. $\varphi = \varphi(x, p)$. It follows that the potential of SLP eq.(2) generated by the solution $\varphi(x, p)$, depends on the model parameters p as well, i.e., q = q(x, p). Finally, the corresponding eigenvalues and eigenfunctions of eq.(2) are continuous functions of p, i.e. $\lambda_n = \lambda_n(p)$, $\psi_n = \psi_n(x, p)$.

Hence, the static solution $\varphi(x, p)$ is asymptotically (exponentially) stable with respect to small space-time perturbations in some bounded subset of the parameter region, if the minimal eigenvalue satisfies $\lambda_0(p) > 0$. [14] Every point on the hypersurface $\lambda_0(p) = 0$ is the bifurcation point for the solution under consideration. The values of the parameters which satisfy $\lambda_0(p) = 0$, are called the bifurcation (critical) values for this solution. The cross section of this hypersurface by a hyperplane, which correspond to fixed values of m-2 parameters, determine the bifurcation curve for the rest two parameters. From the experimental point of view, the most important are the bifurcation curves of kind "critical current-magnetic field"

$$\lambda_0(\gamma, h_e) = 0. \tag{3}$$

The numerical algorithm for the determination of the bifurcation curves is proposed in [17] and was successfully applied to various physical problems [9]

Bifurcation curves in LJJ with inhomogeneity. The static configurations of finite-length junction in the presence of an external magnetic field were studied by many authors. [1,8,9,14,18–21]. Particularly, S.Pagano et al. [18] for arbitrary junction lengths demonstrated the bifurcation curves of the fluxon states in the three different regimes typically observed: small, intermediate, and long junction. We consider here the LJJ which barrier layer contains one resistive rectangular inhomogeneity, characterized by its width Δ , the position of the center of the inhomogeneity ζ and the portion of Josephson current k_J through it. The existence of the inhomogeneity leads to the local change of the Josephson current through the junction, which can be modeled by $j_C(x) = 1 + k_J$ inside of the inhomogeneity and $j_C(x) = 1$ outside of it. At $k_J > 0$ the value of the current through the inhomogeneity exceeds the value of the current in other parts of the junction and such inhomogeneity are considered as a shunt. [14] At $k_J \in [-1, 0)$ the value of the current through the inhomogeneity is less than in other parts of the junction and such inhomogeneity represents a microresistor. Changing the thickness of the barrier layer inside of the inhomogeneity, we can model the transformation from the "shunt" to the "microresistor" inhomogeneity. The value $k_J = 0$ corresponds to a homogeneous junction. Results



Fig. 1: Critical curve of the LJJ with length 2l = 7 and width of inhomogeneity $\Delta = 0.7$. The insert shows the bifurcation curves in the marked by circle region.

of numerical solution of non-linear eigenvalue problem [9] eq.(1), eq.(2) are presented in fig. 1, where the curve





Fig. 2: (a) The distribution of the magnetic field $\varphi_x(x)$ along the junction for bifurcation states M, Φ^1 , and $\Phi^{-1}\Phi^1$ in LJJ with length 2l = 7, width of inhomogeneity $\Delta = 0.7$ at $h_e =$ 1.9; (b) The distribution of the magnetic field for LJJ with 2l = 12 for Φ^1 , Φ^1_l and Φ^1_r at $h_e = 1.5$ and $\gamma = 0$.

"critical current - external magnetic field" of the junction with 2l = 7 and the inhomogeneity with width $\Delta = 0.7$ in the center of the junction is shown. This curve is "constructed" as envelope of bifurcation curves eq.(3), which belong to different possible bound states [9]. The inhomogeneity leads to the appearance on the critical curves the fragments of bifurcation curves of the stable mixed states like $\Phi^n \Phi^{-n}$ and $\Phi^{-n} \Phi^n$, $n = 1, 2, \ldots$, which are not stable in homogeneous case. In one's turn this leads to the non-monotonic decrease of maximums of critical curve, when the magnetic field increases. As we can see in fig. 1, in the investigated region of magnetic field the critical curve is an envelope of the critical curves for the Meissner state M, mixed distributions $\Phi^1 \Phi^{-1}$, $\Phi^{-2} \Phi^2$, $\Phi^{-3} \Phi^3$ and pure fluxon states Φ^1 , Φ^3 , Φ^5 , Φ^7 . All "even" pure states like Φ^2 , Φ^4 , etc., are unstable, so their bifurcation curves eq.(3) do not share for critical curve of junction as whole.

The insert to the fig. 1 shows the details of formation of the critical curve in the marked region. In all our figures in this paper the solid lines correspond to the positive direction of current, the dashed lines correspond to the negative direction of the current. The critical curve of the junction consists of pieces of the bifurcation curves for Φ^1 , $\Phi^1 \Phi^{-1}$ and Φ^3 states in this region.

Table 1: Physical parameters for bifurcation states $M, \ \Phi^1, \ \Phi^{-1} \Phi^1$ and $\Phi^1 \Phi^{-1}$

Type	γ_{cr}	Ν	$\Delta \varphi / 2\pi$	$\varphi(0)/\pi$
Φ^1	0.228	1.317	1.69	1.438
M	0.065	0.073	0.847	0.048
$\Phi^{-1}\Phi^{1}$	0.034	1.654	2.28	1.58
$\Phi^{-1}\Phi^{1}$	0.017	1.43	2.19	1.329
$\Phi^1 \Phi^{-1}$	-0.017	2.57	2.19	2.671
$\Phi^1 \Phi^{-1}$	-0.034	2.346	2.28	2.42
M	-0.065	-0.073	0.847	-0.048
Φ^1	-0.228	0.683	1.69	0.562

The bifurcation curves for mixed $\Phi^1 \Phi^{-1}$ and $\Phi^{-1} \Phi^1$ states demonstrate an interesting peculiarity. As we can see, in the intervals of magnetic field (h_1, h_2) and (h_3, h_4) these CbC-states are stable only if the current through the junction is not equal to zero. A value of the current, which is needed for the creation of stable mixed distributions $\Phi^1 \Phi^{-1}$ and $\Phi^{-1} \Phi^1$ in CbC-regions, depends on the value of the external magnetic field. The table presents the values of the physical parameters for bifurcation states $\Phi^{-1}\Phi^{1}$ and $\Phi^{1}\Phi^{-1}$ at h = 1.9 in comparison with M and Φ^1 . Here the average value of the magnetic flux (number of fluxons) is defined as a functional $N[\varphi] = \frac{1}{2\pi l} \int_{-\infty}^{t} \varphi(x) dx$ and the full magnetic flux through the junction is $\Delta \varphi = \varphi(l) - \varphi(-l)$. [9] For nonbifurcation pure fluxon states at $\gamma = 0$ the number of vortices equals to the integer number: $N[\Phi^{\pm n}] = n$, while for non-bifurcation mixed fluxon states at $\gamma = 0$ we obtain $N[\Phi^n \Phi^{-n}] + N[\Phi^{-n} \Phi^n] = 2n$. [9] For non-bifurcation stable Meissner solution N[M] = 0. As we can see from the Table, the found bifurcation values satisfy the following relations. For Φ^n states the half of sum of N for positive and negative currents equal to n. For CbC states we should take the sum of $N[\Phi^n \Phi^{-n}]$ and $N[\Phi^{-n} \Phi^n]$ for opposite current directions to have integer number 2n. The magnetic flux in the center of junction $\varphi(0)/2\pi$ fulfil the

The distribution of the internal magnetic field $\varphi_x(x)$ along the junction for bifurcation states M, Φ^1 , $\Phi^{-1}\Phi^1$, and $\Phi^{-1}\Phi^1$ at the different values of γ_{cr} , presented in the Table is shown in fig. 2(a) in case of LJJ with 2l = 7and $\Delta = 0.7$ at h = 1.9. The inhomogeneity in LJJ attracts the bound states and it leads to the deformation of the distributions of the magnetic field for these states. In fig. 3 we present the bifurcation curves for vortices in LJJI with length 2l = 12 and the same width of inhomogeneity $\Delta = 0.7$ in the center of junction and the zero amplitude of Josephson current through inhomogeneity $j_C = 0$. It shows clear the tendency of changing of the different bifurcation curves with increase in magnetic field. The main features of this critical curve coincide with the results for

similar relationships.



Fig. 3: Bifurcation curves for LJJI at 2l = 12 with the inhomogeneity in the center with $\Delta = 0.7$

LJJ with 2l = 7. Particularly, we observe here the mixed states as well, and as a result, the non-monotonic decrease of maximums of the junction's critical curve with magnetic field.

But in contrast to the case 2l = 7, the new "noncentral" fluxon states are appeared here in addition to the pinned to the inhomogeneity "central" ones. We found that in case under consideration the noncentral bound states Φ_{I}^{1} , Φ_r^1, Φ_l^2 and Φ_r^{-2} are stable only. The distribution of the internal magnetic field along the junction for Φ^1 , Φ^1_i and Φ^1_r is shown in fig. 2(b). The new moment in case of 2l = 12is a fact, that the states Φ_l^1 and Φ_r^1 are not stable without electric current, at any value of external magnetic field. We have not found the stable $\Phi^1 \Phi^{-1}$ and $\Phi^{-1} \Phi^1$ states in this case also, so we consider that they are replaced with Φ_l^2 and Φ_r^{-2} when the length of LJJ is increased. Or we can say, that for LJJ with 2l = 12 the width's value $\Delta = 0.7$ is small enough to stabilize the $\Phi^1 \Phi^{-1}$ and $\Phi^{-1} \Phi^1$ states. We can see below in fig. 5(a) the transformation of the Φ^2 state into $\Phi^1 \Phi^{-1}$ and $\Phi^{-1} \Phi^1$ with the increase in Δ . The Φ_l^2 and Φ_r^{-2} demonstrate the CbC-regions which look like CbC-regions for mixed states $\Phi^1 \Phi^{-1}$ and $\Phi^{-1} \Phi^1$ for small junction length. Fig. 3 allows us to represent from the unified point of view the influence of the inhomogeneity on the bifurcation curves of the fluxon states in LJJ. It leads to: a)mixed fluxon-antifluxon states $\Phi^{-n}\Phi^{n}$; b)non-monotonic decrease of maximums of the junction's critical curve with magnetic field; c) right and left fluxon states; d)different position of the CbC-states of the different fluxon states.

The influence of the inhomogeneity width. – The influence of the inhomogeneity width on the CbCregions for mixed state $\Phi^1 \Phi^{-1}$ in the case of 2l = 7 is demonstrated by fig. 4. It shows the dependence of the minimal eigenvalue λ_0 on the external current γ for Φ^1 and $\Phi^1 \Phi^{-1}$ for the LJJ with two values of width of inhomogeneity $\Delta = 0.5$ and $\Delta = 1$ at $h_e = 2.2$. The zeroes of λ_0 determine the critical currents for corresponding states. The two values of current γ_{cr} determine the down and upper critical currents for the CbC states at fixed value of the external magnetic field $h_e = 2.2$. We can see that with increase in Δ the interval of the CbC-region for $\Phi^1 \Phi^{-1}$ is increased. The influence of the width of the inho-



Fig. 4: Dependence of the minimal eigenvalue λ_{min} on the external current γ for Φ^1 and $\Phi^1 \Phi^{-1}$ at two different values of the width of inhomogeneity $\Delta = 0.1$ and $\Delta = 0.5$.

mogeneity Δ on the bifurcation curve for Φ^2 is demonstrated in fig. 5(a). We show here the bifurcation curves at $\Delta = 0.3; 0.5; 1$. This figure demonstrates the transformation $\Phi^2 \rightarrow (\Phi^1 \Phi^{-1}, \Phi^{-1} \Phi^1)$ with increase in Δ . There is a bifurcation region in the interval $0.3 < \Delta < 0.5$, where the state Φ^2 is getting unstable, but $\Phi^1 \Phi^{-1}$ and $\Phi^{-1} \Phi^1$ are stable. The details of such transformation are shown in fig. 5(b). As we can see, the state Φ^2 is stable yet at $\Delta = 0.45$, but the corresponding region of magnetic field decreases with Δ . This region disappears completely in the interval $0.45 < \Delta < 0.5$.

Actually, fig. 5 shows in the current versus magnetic field plane typical features of the catastrophe theory. [22]This correspondence is stressed by fig. 5(c), where the dependence $F(\gamma)$ of the total energy on γ is shown for stable distributions Φ^2 , $\Phi^{-1}\Phi^1$ and related to them unstable ones at $h_e = 1.85$ (see fig. 5b). The cusp in the B_1 point is a bifurcation point for Φ^2 with change in the external current. Correspondingly, the points B_2 and B_3 are the cusps of the bifurcation points for mixed distribution $\Phi^{-1}\Phi^{1}$. Note, that the interval along the γ -axis which corresponds to the points B_2 and B_3 has not the point $\gamma = 0$, i.e. the current in B_2 point is a "creation current", and in the point B_3 - "annihilation current" of the distribution $\Phi^{-1}\Phi^{1}$. The states the two horns (in the interval of magnetic field $h_1 < h < h_2$ in the insert to fig. 1) reflect the fact that these states can not be stable without current through the junction. The detailed study of the observed features in the correspondence of the catastrophe theory will be done somewhere else.

Shifting of the inhomogeneity. – Fig. 6 shows the bifurcation curves for Φ^1 in LJJI with 2l = 7, $\Delta = 0.7$ and $k_J = -1$ at different position of the inhomogeneity $\zeta = 0; 0.5; 1; 1.5; 2$ and 2.5. We found that when the inhomogeneity is shifted from the center of the junction, the



Fig. 5: (a) The transformation of the critical curve for Φ^2 with the value of inhomogeneity's width; (b) The details of the transformation of Φ^2 at $\Delta = 0.45$; (c) The dependence of the total energy F on external current γ for distributions Φ^2 , $\Phi^{-1}\Phi^1$ at $h_e = 1.85$



Fig. 6: Bifurcation curves for Φ^1 at different position of the inhomogeneity.

pure fluxon states have the CbC-regions as well. The case $\zeta = 0$ correspond to the position of the inhomogeneity in the center of the junction.

The influence of the current value through the inhomogeneity. – The decrease in the value of the parameter k_J , which characterizes the amplitude of Josephson current through the inhomogeneity, leads to the same transformation $\Phi^2 \rightarrow (\Phi^1 \Phi^{-1}, \Phi^{-1} \Phi^1)$ as we have observed with increase in Δ . We demonstrate it for LJJI with 2l = 7 in fig. 7(a), where the transformation of the dependence $N(h_e)$ for Φ^2 with the value of the amplitude of Josephson current through the inhomogeneity is shown. As we can see from this figure, a bifurcation of the states exists in the interval $-1 < k_J < -0.735$. We found that a new bifurcation point exists at $k_J \approx -0.77$. The dependence $N(h_e)$ in LJJ with 2l = 7, $\Delta = 0.7$, $\zeta = 0$ and $k_J = -1$ is shown in fig. 7(b). It demonstrates clear the relations for N for the pure and mixed bifurcation states we mentioned above. Particularly, for pure fluxon state Φ^3 sum of the $N(h_e)$ for positive and negative current equals to 3, and for states $\Phi^2 \Phi^{-2}$, $\Phi^{-2} \Phi^2$ we get $N[\Phi^2 \Phi^{-2}] +$ $N[\Phi^{-2}\Phi^2] = 4.$



Fig. 7: a) Transformation of the dependence $N(h_e)$ for Φ^2 with the amplitude of Josephson current through the inhomogeneity; (b) The dependence $N(h_e)$ in LJJ with 2l = 7, $\Delta = 0.7$, $\zeta = 0$ and $k_J = -1$.

Exponentially shaped LJJ. – As a second example, we consider an exponentially shaped LJJ [10]- [12]. In Ref [12] we demonstrated that the ESJJ is equivalent to



Fig. 8: (a) Bifurcation curves for ESJJ at 2l = 7 and $\sigma = 0.07$; (b) Bifurcation curves for Φ^1 at different values of σ .

the Josephson junction with distributed inhomogeneity. So we may expect the CbC regions in the critical curves of the ESJJ as well. The corresponding BVP for the ESJJ has the form:

$$-\varphi_{xx} + \sin\varphi + \sigma(\varphi_x - h_e) - \gamma = 0, \varphi_x(-l) = h_e, \varphi_x(l) = h_e$$
(4)

where $0 \leq \sigma \ll 1$ is the form parameter.

Results of numerical solution of this BVP combined with the corresponding SLP [10] are presented in fig. 8(a), where the bifurcation curves for Meissner and four first fluxon states are shown for l = 7 and $\sigma = 0.07$. As we can see, the bifurcation curves of pure fluxon states in ESJJ demonstrate the CbC-regions. The CbC-states are realized at positive current in low magnetic fields and negative current in high magnetic fields. The exponential shape of the junction leads to the appearance of the distributed "geometrical" current $\sigma (\varphi_x - h_e)$, which modifies the bifurcation curves of the states in compare with the rectangular homogeneous junction. [12] Bifurcation curves for Φ^1 at l = 7 and different values of σ are shown in fig. 8(b). The increase in σ leads to the increase of the CbC-region. With increase in σ a situation might be realized, when the dependence of the upper critical current on the external magnetic field in CbC regions is getting nonmonotonic. We found that such non-monotonic behavior of the bifurcation curve Φ^1 in low magnetic fields appears in ESJJ with increase in junction's length as well.

In summary, we performed the numerical simulation of the critical curves of long Josephson junctions with inhomogeneity and variable width. In both cases we demonstrate the CbC-regions of magnetic field, where fluxon states are stable only if the current through the junction is different from zero. We showed that the position and size of these regions depend on length of the junction, its geometry, parameters of inhomogeneity and form of the junction. We consider that development of the experimental methods for detection of CbC-states will open a perspective for their applications.

* * *

We thank I.V. Puzynin and N.M. Plakida for useful discussion and cooperation. This work is partially supported by Sofia University Scientific foundation under Grant No 135/2008 and RFBR grant 08-02-00520-a.

REFERENCES

- [1] OWEN C.S., SCALAPINO D.J., Phys. Rev., 164 (1967) 538.
- [2] BARONE A. and PATTERNO J., Physics and Applications of the Josephson effect (John Wiley and Sons) 1982.
- [3] LICHAREV K.K., Dynamics of Josephson Junctions and Circuits (Gordon and Breach, New York) 1986.
- [4] PETRAGLIA A. and PEDERSEN N. F. CHRISTIANSEN P. L. USTINOV A. V., *Phys. Rev.*, B55 (1997) 8490.
- [5] VISTAVKIN A.N., DRACHEVSKI YU.F., COSHELETZ V.P., SERPUCHENKO I.L., Low Temp. Phys., 14 (1988) 646.
- [6] SCHWIDTAL K., Phys. Rev., **B2** (1970) 2526.
- [7] GOLDOBIN E., USTINOV A. V., Phys. Rev., 59 (1999) 11532.
- [8] FILIPPOV A.T., GAL'PERN YU.S., BOYADJIEV T.L. and PUZYNIN I.V., Phys. Lett. A, 120 (1987) 47.
- [9] PUZYNIN I.V. ET AL, , Particles & Nucley (Dubna), 38 (2007) 70.
- [10] SEMERDJIEVA E.G., BOYADJIEV T.L. and SHUKRINOV YU.M., Low Temp. Phys., 30 (2004) 610.
- [11] SHUKRINOV YU.M., SEMERDJIEVA E.G. and BOYADJIEV T.L., J. of Low Temp. Phys., 139 (299 (2005)).
- [12] SEMERDJIEVA E.G., BOYADJIEV T.L. and SHUKRINOV YU.M., Low Temp. Phys., 31 (2005) 1110.
- [13] FOGEL ET AL M.B., Phys. Rev., 15 (1977) 1578.
- [14] GAL'PERN YU.S. and FILIPPOV A.T., Sov. Phys. JETP, 59 (1984) 894.
- [15] ILIEV I.D., KHRISTOV E.KH. and KIRCHEV K.P., Spectral methods in soliton equations (Longman Sci. & Techn., Wiley) 1994.
- [16] LEVITAN B.M., SARGSJAN I.S., Introduction to Spectral Theory (Transl. Math. Monographs, AMS, Providence, RI) 1975.
- [17] BOYADJIEV T.L. ET AL., Comm. JINR P11-88-409 (Dubna) 1988.
- [18] PAGANO S. ET AL., Phys. Rev., B43 (1991) 5364.
- [19] CAPUTO J.-G., FLYTZANIS N., GAIDIDEI Y., STE-FANAKIS N., VAVALIS E., Supercond. Sci. Technol., 13 (2000) 423.
- [20] STEFANAKIS N. AND FLYTZANIS. N., Supercond. Sci. Technol., 14 (2001) 16.

- [21] USTINOV A.V., Long Josephson junctions and stacks, In: Superconductivity in Networks and Mesoscopic Structures, Eds. C. Giovannella & C. Lambert, (AIP) 1998.
- [22] TIM POSTON and IAN STEWART., Catastrophe Theory and its Applications (Dover) 1996.