On minimal actions of finite simple groups on homology spheres and Euclidean spaces

Bruno P. Zimmermann

Università degli Studi di Trieste Dipartimento di Matematica e Informatica 34100 Trieste, Italy

Abstract. We consider the following problem: for which classes of finite groups, and in particular finite simple groups, does the minimal dimension of a faithful, smooth action on a homology sphere coincide with the minimal dimension of a faithful, linear action on a sphere? We prove that the two minimal dimensions coincide for the linear fractional groups PSL(2, p) as well as for various classes of alternating and symmetric groups. We prove analogous results also for actions on Euclidean spaces.

1. Introduction

We consider actions of finite groups on spheres and homology spheres; if not stated otherwise, all actions will be assumed to be smooth (or locally linear) and faithful.

In a program to classify the finite, nonsolvable groups which admit an action on a sphere or homology sphere in dimension three or four, it is shown in [16-18], [20, 21] that the only finite (nonabelian) simple group acting on a homology 3-sphere is the alternating or linear fractional group $\mathbb{A}_5 \cong \mathrm{PSL}(2,5)$, and the only finite simple groups acting on a homology 4-sphere are the groups $\mathbb{A}_5 \cong \mathrm{PSL}(2,5)$ and $\mathbb{A}_6 \cong \mathrm{PSL}(2,9)$; a corresponding classification in dimension five is still open. This naturally motivated the following problems:

- For a given finite simple group G, determine the minimal dimension of a sphere or homology sphere on which G acts faithfully (does it coincide with the minimal dimension of a linear action on a sphere?).

- Show that there are only finitely many finite simple groups which admit an action on a homology sphere of a given dimension d.

Some partial results for linear fractional and alternating groups have been obtained in [22]. Concerning other classes of groups, it is shown in [9] that for finite p-groups the two minimal dimensions of an action on a homology sphere and of a linear action on a sphere coincide, and by [8] also for orientation-preserving actions of finite abelian groups. However, there are finite solvable groups for which the two minimal dimensions

do not coincide; specifically, the Milnor groups Q(8a, b, c) ([15]) do not admit faithful, orientation-preserving, linear actions on S^3 (neither free nor nonfree), but by [14] some of them admit a free action on a homology 3-sphere (see also section 4).

In the present paper, we determine the minimal dimensions for the linear fractional groups PSL(2, p) as well as for certain alternating and symmetric groups. In the following, a mod p homology sphere is a closed n-manifold with the mod p homology of the n-sphere (i.e., homology with coefficients in the integers $\mathbb{Z}_p \mod p$). Our main result is the following (where p denotes a prime number and q a positive integer):

Theorem 1. For the following finite groups, the minimal dimension of a faithful, smooth (orientation-preserving) action on a mod p homology sphere coincides with the minimal dimension of a faithful, linear (orientation-preserving) action on a sphere:

i) a semidirect product (metacyclic group) $\mathbb{Z}_p \rtimes \mathbb{Z}_q$, with an effective action of \mathbb{Z}_q on the normal subgroup \mathbb{Z}_p ;

- ii) a linear fractional group PSL(2, p);
- iii) a symmetric group \mathbb{S}_p ;
- iv) an alternating groups \mathbb{A}_p such that $p \equiv 3 \mod 4$.

See section 3 for the various minimal dimensions. An analogous result holds for faithful, smooth actions on Euclidean spaces:

Theorem 2. For the groups listed in Theorem 1, the minimal dimension of a faithful, smooth (orientation-preserving) action on a Euclidean space coincides with the minimal dimension of a faithful, linear (orientation-preserving) action.

Note that this is clear for actions with a global fixed point, so more generally one may ask: what is the minimal dimension of a fixed point-free action of a given finite group on a Euclidean space? See [7] for a survey on actions of finite groups on disks and Euclidean spaces.

2. The Borel spectral sequence associated to a group action

Let G be a finite group acting on a space X. Let EG denote a contractible space on which G acts freely, and BG = EG/G. We consider the twisted product $X_G = EG \times_G X = (EG \times X)/G$. The "Borel fibering" $X_G \to BG$, with fiber X, is induced by the projection $EG \times X \to EG$, and the equivariant cohomology of the G-space X is defined as $H^*(X_G; K)$. Our main tool will be the Leray-Serre spectral sequence E(X)associated to the Borel fibration $X_G \to BG$,

$$E_2^{i,j} = E_2^{i,j}(X) = H^i(BG; H^j(X; K)) = H^i(G; H^j(X; K)) \quad \Rightarrow \quad H^{i+j}(X_G; K),$$

i.e. converging to the graded group associated to a filtration of $H^*(X_G; K)$ ("Borel spectral sequence", see e.g. [2]); here K denotes any abelian coefficient group or commutative ring.

Now suppose that G acts on a (open or closed) n-manifold M; we denote by Σ the singular set of the G-action (all points in M with nontrivial stabilizer). Crucial for the proofs of Theorems 1 and 2 is the following Proposition 1, see [13], [10, 11] for a proof (see also [4, Proposition VII.10.1] for a Tate cohomology version).

Proposition 1. In dimensions greater than n, inclusion induces isomorphisms

$$H^*(M_G; K) \cong H^*(\Sigma_G; K).$$

We use this to prove the following:

Proposition 2. For an odd prime p and a positive integer q, let $G = \mathbb{Z}_p \rtimes \mathbb{Z}_q$ be a semidirect product with an effective action of \mathbb{Z}_q on the normal subgroup \mathbb{Z}_p . Suppose that G admits a faithful action on a manifold M with the mod p homology of the n-sphere, and that the subgroup \mathbb{Z}_p of G acts freely. Then n + 1 is a multiple of 2q if all elements of G act as the identity on $H^n(M; \mathbb{Z}_p) \cong \mathbb{Z}_p$ (the "orientation-preserving case"), and an odd multiple of q if some element of G acts as the minus identity (the "orientation-reversing case").

Proof. We consider first the Borel spectral sequence $E(\Sigma)$ converging to the cohomology of Σ_G , with $E_2^{i,j} = H^i(G; H^j(\Sigma; \mathbb{Z}_p))$. Let Σ_q denote the singular set of the subgroup \mathbb{Z}_q of G. The singular set Σ of G is the disjoint union of the singular sets of the p conjugates of \mathbb{Z}_q in G; these fixed point sets are all disjoint since the action of \mathbb{Z}_q on \mathbb{Z}_p is effective and the action of \mathbb{Z}_p is free, by assumption. Now the action of G on the cohomology $H^j(\Sigma; \mathbb{Z}_p)$ of the orbit $\mathbb{Z}_p(\Sigma_q) = \Sigma$ is induced (or co-induced) from the action of \mathbb{Z}_q on $H^j(\Sigma_q; \mathbb{Z}_p)$, and by Shapiro's Lemma ([4, Proposition III.6.2]), $H^i(G; H^j(\Sigma; \mathbb{Z}_p)$ is isomorphic to $H^i(\mathbb{Z}_q; H^j(\Sigma_q; \mathbb{Z}_p)$ and hence trivial, for i > 0. So also $H^*(\Sigma_G; \mathbb{Z}_p)$ is trivial, in positive dimensions. By Proposition 1, also $H^*(M_G; \mathbb{Z}_p) \cong H^*(\Sigma_G; \mathbb{Z}_p)$ is trivial, in sufficiently large dimensions.

Next we analyze the spectral sequence E(M) converging to the cohomology of M_G . The E_2 -terms $E_2^{i,j} = H^i(G; H^j(M; \mathbb{Z}_p))$ are concentrated in the two rows j = 0 and j = n where they are equal to \mathbb{Z}_p , with a possibly twisted action of G on $H^n(M; \mathbb{Z}_p) \cong \mathbb{Z}_p$. In particular, the only possibly nontrivial differentials of E(M) are $d_{n+1}^{i,j} : E_{n+1}^{i,n} \to E_{n+1}^{i+n+1,0}$, of bidegree (n+1, -n).

By [4, Theorem III.10.3], for i > 0

$$H^i(G; H^j(M; \mathbb{Z}_p)) \cong H^i(\mathbb{Z}_p; H^j(M; \mathbb{Z}_p))^{\mathbb{Z}_q}.$$

The cohomology ring $H^*(\mathbb{Z}_p;\mathbb{Z}_p)$ is the tensor product of a polynomial algebra $\mathbb{Z}_p[t]$ on a 2-dimensional generator t and an exterior algebra E(s) on a 1-dimensional generator s (see [1, Corollary II.4.2]); also, t is the image of s under the mod p Bockstein homomorphism.

Suppose first that \mathbb{Z}_q acts trivially on $H^n(M; \mathbb{Z}_p) \cong \mathbb{Z}_p$. Since the action of \mathbb{Z}_q on \mathbb{Z}_p is effective, also the action of \mathbb{Z}_q on $H^1(\mathbb{Z}_p; \mathbb{Z}_p)$ and $H^2(\mathbb{Z}_p; \mathbb{Z}_p)$ is effective: denoting by σ the action of a generator of \mathbb{Z}_q on the cohomology $H^i(\mathbb{Z}_p; \mathbb{Z}_p)$, one has $\sigma(s) = ks$ and $\sigma(t) = kt$, for some integer k representing an element of order q in \mathbb{Z}_p . Hence $\sigma(t^a) = k^a t^a$ and the only powers of t fixed by σ are those divisible by q (in dimensions i which are even multiples of q); similarly, $\sigma(st^a) = k^{a+1}t^{a+1}$, so the only elements in odd dimensions fixed by σ are the products st^a such that a + 1 is a multiple of q (in dimensions i such that i + 1 is an even multiple of q). Consequently, $H^i(G; H^j(M; \mathbb{Z}_p))$ is nontrivial exactly for j = 0 and n and in dimensions i such that either i or i + 1 is a multiple of 2q.

Since $H^*(M_G; \mathbb{Z}_p) \cong H^*(\Sigma_G; \mathbb{Z}_p)$ is trivial in sufficiently large dimensions, the differentials d_{n+1} of the spectral sequence $(E_2^{i,j}(M) = H^i(G; H^j(M; \mathbb{Z}_p))$ (concentrated in the rows j = 0 and n) have to be isomorphisms in large dimensions. This can happen only if n + 1 is a multiple of 2q, which completes the proof of Proposition 2 in the orientation-preserving case.

Now suppose that a generator of \mathbb{Z}_q acts as minus identity on $H^n(M;\mathbb{Z}_p) \cong \mathbb{Z}_p$, in particular q is even. Denoting by $\tilde{\sigma}$ the action of a generator of \mathbb{Z}_q on the cohomology $H^i(\mathbb{Z}_p; H^n(M;\mathbb{Z}_p))$, we now have that $\tilde{\sigma}(t^a) = -\sigma(t^a) = -k^a t^a$, $\tilde{\sigma}(st^a) = -\sigma(st^a) =$ $-k^{a+1}t^{a+1}$. Now the elements of $H^i(\mathbb{Z}_p; H^n(M;\mathbb{Z}_p)) = H^i(\mathbb{Z}_p;\mathbb{Z}_p)$ invariant under $\tilde{\sigma}$ are the powers of t by odd multiples of q/2 (in dimensions i which are odd multiples of q), and the products st^a such that a + 1 is an odd multiple of q/2 (in dimensions isuch that i + 1 is an odd multiple of q). Hence $H^i(G; H^j(M;\mathbb{Z}_p))$ is nontrivial exactly in the following situations: either j = 0 and i or i + 1 is an even multiple of q (since the action of G on $H^0(M;\mathbb{Z}_p)$ is trivial), or if j = n and i or i + 1 is an odd multiple of q(with the twisted action of G on $H^n(M;\mathbb{Z}_p)$). Since again $H^*(M_G;\mathbb{Z}_p) \cong H^*(\Sigma_G;\mathbb{Z}_p)$ is trivial in large dimensions, the differential d_{n+1} has to be an isomorphism and n + 1an odd multiple of q. (Note that, in order to obtain just the lower bound $n \ge q - 1$, one may apply the orientation-preserving case to the subgroup $\mathbb{Z}_p \rtimes \mathbb{Z}_{q/2}$ of $\mathbb{Z}_p \rtimes \mathbb{Z}_{q,2}$.)

This completes the proof of Proposition 2.

3. Proofs of Theorems 1 and 2

Proof of Theorem 1.

i) It is easy to see that a group $G = \mathbb{Z}_p \rtimes \mathbb{Z}_q$ as in Theorem 1 i) admits a faithful, linear action (a faithful, real, linear representation) on $S^{2q-1} \subset \mathbb{R}^{2q}$ if q is odd, on $S^q \subset \mathbb{R}^{q+1}$

if q is even and the action is orientation-preserving, and on $S^{q-1} \subset \mathbb{R}^q$ if some element of G reverses the orientation (see e.g. [4, Example 9.2.3, p.155]). We will show that the dimensions 2q - 1, q and q - 1 give lower bounds for the dimensions of actions of G on mod p homology spheres, in the respective cases.

Suppose that $G = \mathbb{Z}_p \rtimes \mathbb{Z}_q$ admits a faithful, smooth action on a mod p homology n-sphere M. If the subgroup \mathbb{Z}_p of G acts freely, Proposition 2 implies that n + 1 is a multiple of 2q if G acts orientation-preservingly, and an odd multiple of q otherwise. In particular, the minimal possibilities are n = 2q - 1 and n = q - 1, respectively, which coincide with the minimal dimension of a faithful, linear action on a sphere.

Suppose then that \mathbb{Z}_p has nonempty fixed point set F. Since the action is assumed to be smooth, F is a smooth submanifold; moreover, by Smith fixed point theory, Fis a mod p homology sphere of some dimension d, $0 \leq d < n$. Then, by Lefschetzduality with coefficients in \mathbb{Z}_p , the complement M - F is a G-invariant manifold with the mod p homology of a sphere of dimension n - d - 1. Proposition 2 implies now that $n - d - 1 \geq 2q - 1$, $n \geq 2q$ if q is odd, $n - d - 1 \geq q - 1$, $n \geq q$ if q is even.

This completes the proof of part i) of Theorem 1.

ii) The representation theory of the linear fractional groups PSL(2, p) is well-known, in particular PSL(2, p) admits a faithful, linear action (a faithful, real, linear representation) on $S^{p-2} \subset \mathbb{R}^{p-1}$ if $p \equiv 3 \mod 4$, and on $S^{(p-1)/2} \subset \mathbb{R}^{(p+1)/2}$ if $p \equiv 1 \mod 4$ (see e.g. [12], section 5.2, in particular Exercise 5.10).

The subgroup of PSL(2, p) represented by all upper triangular matrices is a semidirect product $\mathbb{Z}_p \rtimes \mathbb{Z}_q$, with q = (p-1)/2, where \mathbb{Z}_q is the subgroup represented by all diagonal matrices and \mathbb{Z}_p by all matrices with both diagonal entries equal to one; also, \mathbb{Z}_q acts effectively on the normal subgroup \mathbb{Z}_p . By i), the minimal dimension of a faithful, smooth, orientation-preserving action of $\mathbb{Z}_p \rtimes \mathbb{Z}_q$ on a mod p homology sphere is 2q - 1 = p - 2 if q is odd, and to q = (p - 1)/2 if q is even. Since PSL(2, p) admits linear actions in these dimensions, this proves Theorem 1 for the groups of type ii).

iii) and iv) Again the representation theory of the symmetric and alternating groups is well-known (see e.g. [12, section 5.1]), in particular the symmetric group \mathbb{S}_p and the alternating group \mathbb{A}_p admit a faithful, linear action (a faithful, real, linear representation) on $S^{p-2} \subset \mathbb{R}^{p-1}$, orientation-reversing and for an arbitrary positive integer p in the case of the symmetric group, for integers p > 5 in the case of the alternating group.

For an odd prime p, consider the semidirect product $G = \mathbb{Z}_p \rtimes \mathbb{Z}_{p-1}$, with an effective action of \mathbb{Z}_{p-1} on the normal subgroup \mathbb{Z}_p . The action by left-multiplication of G on the p left cosets of the subgroup \mathbb{Z}_{p-1} of G realizes G as a subgroup of the symmetric group \mathbb{S}_p . By i), the minimal dimension of a faithful, smooth action of G on a mod phomology sphere is p-2; since \mathbb{S}_p admits a faithful, linear action on S^{p-2} this proves part iii) of Theorem 1. For groups of type iv), we consider the subgroup $\mathbb{Z}_p \rtimes \mathbb{Z}_{(p-1)/2}$ of index two of $\mathbb{Z}_p \rtimes \mathbb{Z}_{p-1}$ which is realized as a subgroup of the alternating group \mathbb{A}_p . The alternating group \mathbb{A}_p admits a faithful action on S^{p-2} ; if (p-1)/2 is odd, by part i) of Theorem 1 this coincides with the minimal dimension of a faithful, smooth action of $\mathbb{Z}_p \rtimes \mathbb{Z}_{(p-1)/2}$ on a mod p homology sphere.

This completes the proof of Theorem 1.

Proof of Theorem 2.

As noted in the proof of part i) of Theorem 1, the group $G = \mathbb{Z}_p \rtimes \mathbb{Z}_q$ admits a faithful, linear action on \mathbb{R}^{2q} if q is odd, on \mathbb{R}^{q+1} if q is even and the action is orientationpreserving, and on \mathbb{R}^q if some element of G reverses the orientation.

Suppose that $G = \mathbb{Z}_p \rtimes \mathbb{Z}_q$ admits a faithful, smooth action on Euclidean space \mathbb{R}^n . By Smith fixed point theory (see [2]), the fixed point set F of the subgroup \mathbb{Z}_p of G is a \mathbb{Z}_p -acyclic manifold of some dimension $d \ge 0$, in particular non-empty. Note that the action of G extends to a continuous action of G on the sphere S^n and, again by Smith fixed point theory, the fixed point set \hat{F} of the subgroup \mathbb{Z}_p on S^n has the mod p homology of a sphere of dimension d. Now by Lefschetz-duality with coefficients in \mathbb{Z}_p , applied to the pair (S^n, \hat{F}) , the complement $\mathbb{R}^n - F = S^n - \hat{F}$ has the mod p homology of a sphere of dimension n - d - 1. Hence G admits a faithful, smooth action on the manifold $\mathbb{R}^n - F$, with the mod p homology of a sphere of dimension 2p of G acts freely. Now Proposition 2 implies that $n - d - 1 \ge 2q - 1$, $n \ge 2q$ if q is odd, and $n - d - 1 \ge q - 1$, $n \ge q$ if q is even.

Suppose that q is even and that G acts orientation-preservingly on \mathbb{R}^n . If the fixed point set F of \mathbb{Z}_p has dimension d = 0 then F is a single point which hence is a global fixed point of G. Now G acts orientation-preservingly on the boundary of an invariant neighbourhood of the fixed point which is a sphere of dimension n-1, and Proposition 2 implies that $n \ge 2q$. On the other hand, if d > 0 then, as noted above, $n-d-1 \ge q-1$ hence $n \ge q+d \ge q+1$.

Since these lower bounds for the dimension of a faithful, smooth action of $G = \mathbb{Z}_p \rtimes \mathbb{Z}_q$ on a Euclidean space can be realized by a faithful, linear action of the group, this completes the proof of Theorem 2 for the groups of type i). For all other groups, similar as in the proof of Theorem 1, it is a consequence of the case of $\mathbb{Z}_p \rtimes \mathbb{Z}_q$.

4. Remarks on continuous actions of some other finite groups

We present some other finite groups for which the two minimal dimensions for spheres and homology spheres coincide, even for the case of arbitrary continuous actions.

Proposition 3. For the following groups, the minimal dimension of a faithful, continuous action on a homology sphere coincides with the minimal dimension of a faithful, linear action on a sphere:

the unitary and symplectic groups $PSU(4,2) \cong PSp(4,3)$ and PSp(6,2);

the Weyl or Coxeter groups E_6 , E_7 and E_8 of the corresponding exceptional Lie algebras.

This is a consequence of the following well-known result from Smith fixed point theory ([19], see also [3]; iii) follows from the fact that the fixed point set of an orientationpreserving involution on a manifold is a mod 2 homology manifold of codimension least two, then one applies ii)).

Proposition 4.

i) For an odd prime number p, the minimal dimension of a faithful, continuous action of the elementary abelian p-group $(\mathbb{Z}_p)^k$ on a mod p homology sphere is 2k - 1 (and 2kfor an action on a mod p acyclic manifold).

ii) The minimal dimension of a faithful, continuous action of the elementary abelian 2-group $(\mathbb{Z}_2)^k$ on a mod 2 homology sphere is k-1 (and k for an action on a mod 2 acyclic manifold).

iii) The minimal dimension of a faithful, continuous, orientation-preserving action of the elementary abelian 2-group $(\mathbb{Z}_2)^k$ on a mod 2 homology sphere is k (and k + 1 for an action on a mod 2 acyclic manifold).

Proof of Proposition 3. We refer to [6] and its references for information about the subgroup structure and the character tables of the finite simple groups. The unitary group PSU(4,2) has a maximal subgroup $(\mathbb{Z}_3)^3$ and a faithful, linear action on \mathbb{S}^5 , so Proposition 4 i) implies that the two minimal dimensions coincide. Also, PSU(4,2) is a subgroup of index two in the Weyl group of type E_6 which has also a linear action on \mathbb{S}^5 .

The Weyl group of type E_8 has a subgroup $(\mathbb{Z}_3)^4$ and a linear action on \mathbb{S}^7 (see the closely related orthogonal group $O_8^+(2)$ in [6]), so the result follows again from Proposition 4 i).

The symplectic group PSp(6, 2) is a subgroup of index 2 in the Weyl group of type E_7 , and both act linearly on \mathbb{S}^6 . Since PSp(6, 2) has a subgroup $(\mathbb{Z}_2)^6$, Proposition 4 iii) applies. Alternatively, the group PSp(6, 2) has the alternating group \mathbb{A}_8 and the linear fractional group PSL(2, 8) as subgroups, and it is shown in [22, Corollary 2] and [17, Proposition 1] as an application of the Borel formula that the minimal dimension of a continuous action of \mathbb{A}_8 and PSL(2, 8) on a homology sphere is six (the results in [22] and [17] are formulated for smooth actions; since the Borel formula holds for continuous actions, the two specific results cited remain true in this more general setting).

This finishes the proof of Proposition 4.

As noted in the introduction, for some of the Milnor groups Q(8a, b, c) the minimal dimension of an action on a homology sphere is strictly smaller than the minimal dimension of a linear action on a sphere. We do not know similar examples for smooth or continuous actions on Euclidean space. Interesting examples of continuous actions not conjugate to smooth actions can be obtained as follows.

We consider again the Milnor groups Q(8a, b, c) ([15]); these groups have periodic cohomology of period four but do not admit a faithful, free, linear action on the 3-sphere. For odd, coprime integers $a > b > c \ge 1$, the Milnor group Q(8a, b, c) is a semidirect product $\mathbb{Z}_{abc} \ltimes Q_8$ of a normal cyclic subgroup $\mathbb{Z}_a \times \mathbb{Z}_b \times \mathbb{Z}_c \cong \mathbb{Z}_{abc}$ by the quaternion group $Q_8 = \{\pm 1, \pm i, \pm j, \pm k\}$ of order eight, where i, j and k act trivially on $\mathbb{Z}_a, \mathbb{Z}_b$ and \mathbb{Z}_c , respectively, and in a dihedral way on the other two.

It has been shown by Milgram [14] that some of the Milnor groups Q(8a, b, c), for odd, coprime integers $a > b > c \ge 1$, admit a faithful, free action on a homology 3-sphere; let Q be a Milnor group which admits such an action on a homology 3-sphere M. By the double suspension theorem (see e.g. [5]), the double suspension $M * S^1$ of M (or join with the 1-sphere) is homeomorphic to S^5 . Letting Q act trivially on S^1 , the actions of Q on M and S^1 induce a faithful, continuous, orientation-preserving action of Q on S^5 with fixed point set S^1 , and hence also on \mathbb{R}^5 (the complement of a fixed point).

References

- A.Adem, R.J.Milgram, Cohomology of finite groups. Grundlehren der math. Wissenschaften 309, Springer-Verlag 1994
- 2. G.Bredon, Introduction to compact Transformation Groups. Academic Press, New York 1972
- 3. M.R.Bridson, K.Vogtmann, Actions of automorphism groups of free groups on homology spheres and acyclic manifolds. arXiv:0803.2062
- K.S.Brown, Cohomology of Groups. Graduate Texts in Mathematics 87, Springer-Verlag 1982
- J.W.Cannon, The recognition problem: what is a topological manifold. Bull. Amer. Math. Soc., 832-866 (1978)
- J.H.Conway, R.T.Curtis, S.P.Norton, R.A.Parker, R.A.Wilson, Atlas of Finite Groups. Oxford University Press 1985
- M.W.Davis, A survey of results in higher dimensions. In: The Smith Conjecture (eds. J.W.Morgan, H.Bass), Academic Press, 227-240 (1984)
- R.M.Dotzel, Orientation preserving actions of finite abelian groups on spheres. Proc. Amer. Math. Soc. 100, 159-163 (1987)
- R.M.Dotzel, G.C.Hamrick, *p*-group actions on homology spheres. Invent. math. 62, 437-442 (1981)

- 10 A.E.Edmonds, Aspects of group actions on four-manifolds. Top. Appl. 31, 109-124 (1989)
- 11. A.E.Edmonds, *Homologically trivial group actions on 4-manifolds*. Electronic version available at arXiv:math.GT/9809055
- 12. W.Fulton, J.Harris, *Representation Theory: A First Course*. Graduate Texts in Mathematics 129, Springer-Verlag 1991
- 13. M.P.McCooey, Symmetry groups of 4-manifolds. Topology 41, 835-851 (2002)
- 14. R.J.Milgram, *Evaluating the Swan finiteness obstruction for finite groups*. Algebraic and Geometric Topology. Lecture Notes in Math. 1126 (Springer 1985), 127-158
- 15. J.Milnor, Groups which act on S^n without fixed points. Amer. J. Math. 79, 623-630 (1957)
- M.Mecchia, B.Zimmermann, On finite groups acting on Z₂-homology 3-spheres. Math. Z. 248, 675-693 (2004)
- M.Mecchia, B.Zimmermann, On finite simple groups acting on integer and mod 2 homology 3-spheres. J. Algebra 298, 460-467 (2006)
- M.Mecchia, B.Zimmermann, On finite simple and nonsolvable groups acting on homology 4-spheres. Top. Appl. 2006, 2933-2942
- P.A.Smith, Permutable periodic transformations. Proc. Nat. Acad. Sci. U.S.A. 30, 105
 108 (1944)
- 20. B.Zimmermann, On finite simple groups acting on homology 3-spheres. Top. Appl. 125, 199-202 (2002)
- B.Zimmermann, On the classification of finite groups acting on homology 3-spheres. Pacific J. Math. 217, 387-395 (2004)
- 22. B.Zimmermann, On the minimal dimension of a homology sphere on which a finite group acts. Math. Proc. Camb. Phil. Soc. 144, 397-401 (2008)