

Stability of a functional equation deriving from cubic and quartic functions

M. Eshaghi Gordji

Department of Mathematics, Semnan University,
P. O. Box 35195-363, Semnan, Iran
e-mail: madjid.eshaghi@gmail.com

A. Ebadian

Department of Mathematics, Urmia University, Urmia, Iran
e-mail: ebadian.ali@gmail.com

S. Zolfaghari

Department of Mathematics, Semnan University,
P. O. Box 35195-363, Semnan, Iran
e-mail: zolfagharys@yahoo.com

Abstract. In this paper, we obtain the general solution and the generalized Ulam-Hyers stability of the cubic and quartic functional equation

$$\begin{aligned} 4(f(3x+y) + f(3x-y)) &= -12(f(x+y) + f(x-y)) \\ &+ 12(f(2x+y) + f(2x-y)) - 8f(y) - 192f(x) + f(2y) + 30f(2x). \end{aligned}$$

1. INTRODUCTION

The stability problem of functional equations originated from a question of Ulam [34] in 1940, concerning the stability of group homomorphisms. Let $(G_1, .)$ be a group and let $(G_2, *)$ be a metric group with the metric $d(., .)$. Given $\epsilon > 0$, does there exist a $\delta > 0$, such that if a mapping $h : G_1 \rightarrow G_2$ satisfies the inequality $d(h(x.y), h(x) * h(y)) < \delta$ for all $x, y \in G_1$, then there exists a homomorphism $H : G_1 \rightarrow G_2$ with $d(h(x), H(x)) < \epsilon$ for all $x \in G_1$? In the other words, Under what condition does there exists a homomorphism near an approximate homomorphism? The concept of stability for functional equation arises when we replace the functional equation by an inequality which acts as a perturbation of the equation. In 1941, D. H. Hyers [9] gave the first affirmative answer to the question of Ulam for Banach spaces. Let $f : E \rightarrow E'$ be a mapping between Banach spaces such that

$$\|f(x+y) - f(x) - f(y)\| \leq \delta$$

for all $x, y \in E$, and for some $\delta > 0$. Then there exists a unique additive mapping $T : E \rightarrow E'$ such that

$$\|f(x) - T(x)\| \leq \delta$$

for all $x \in E$. Moreover if $f(tx)$ is continuous in $t \in \mathbb{R}$ for each fixed $x \in E$, then T is linear. Finally in 1978, Th. M. Rassias [31] proved the following theorem.

Theorem 1.1. *Let $f : E \rightarrow E'$ be a mapping from a normed vector space E into a Banach space E' subject to the inequality*

$$\|f(x+y) - f(x) - f(y)\| \leq \epsilon(\|x\|^p + \|y\|^p) \quad (1.1)$$

for all $x, y \in E$, where ϵ and p are constants with $\epsilon > 0$ and $p < 1$. Then there exists a unique additive mapping $T : E \rightarrow E'$ such that

$$\|f(x) - T(x)\| \leq \frac{2\epsilon}{2-2^p} \|x\|^p \quad (1.2)$$

for all $x \in E$. If $p < 0$ then inequality (1.1) holds for all $x, y \neq 0$, and (1.2) for $x \neq 0$. Also, if the function $t \mapsto f(tx)$ from \mathbb{R} into E' is continuous in real t for each fixed $x \in E$, then T is linear.

In 1991, Z. Gajda [5] answered the question for the case $p > 1$, which was raised by Rassias. This new concept is known as Hyers-Ulam-Rassias stability of functional equations (see [1-2], [5-10], [28-30]). On the other hand J. M. Rassias [25-27], generalized the Hyers stability result by presenting a weaker condition controlled by a product of different powers of norms. According to J. M. Rassias Theorem:

Theorem 1.2. *If it is assumed that there exist constants $\Theta \geq 0$ and $p_1, p_2 \in \mathbb{R}$ such that $p = p_1 + p_2 \neq 1$, and $f : E \rightarrow E'$ is a map from a norm space E into a Banach space E' such that the inequality*

$$\|f(x+y) - f(x) - f(y)\| \leq \epsilon \|x\|^{p_1} \|y\|^{p_2} \quad (1.1p)$$

for all $x, y \in E$, then there exists a unique additive mapping $T : E \rightarrow E'$ such that

$$\|f(x) - T(x)\| \leq \frac{\Theta}{2-2^p} \|x\|^p,$$

for all $x \in E$. If in addition for every $x \in E$, $f(tx)$ is continuous in real t for each fixed x , then T is linear (see [19-26]).

The oldest cubic functional equation, and was introduced by J. M. Rassias (in 2000-2001): [17-18], as follows:

$$f(x+2y) + 3f(x) = 3f(x+y) + f(x-y) + 6f(y).$$

Jun and Kim [11] introduced the following cubic functional equation

$$f(2x+y) + f(2x-y) = 2f(x+y) + 2f(x-y) + 12f(x) \quad (1.3)$$

and they established the general solution and the generalized Hyers-Ulam-Rassias stability for the functional equation (1.3). The function $f(x) = x^3$ satisfies the functional equation (1.3), which is thus called a cubic functional equation. Every solution of the cubic functional equation is said to be a cubic function. Jun and Kim proved that a function f between real vector spaces X and Y is a solution of (1.3) if and only if there exists a unique function $C : X \times X \times X \rightarrow Y$ such that $f(x) = C(x, x, x)$ for all $x \in X$, and C is symmetric for each fixed one variable and is additive for fixed two variables. The oldest quartic functional equation, and was introduced by J. M. Rassias (in 1999-2000): [16], [27], and then (in 2005) was employed by Won-Gil Park [15] and others, such that:

$$f(x+2y) + f(x-2y) = 4(f(x+y) + f(x-y)) + 24f(y) - 6f(x). \quad (1.4)$$

In fact they proved that a function f between real vector spaces X and Y is a solution of (1.4) if and only if there exists a unique symmetric multi-additive function $Q : X \times X \times X \times X \rightarrow Y$ such that $f(x) = Q(x, x, x, x)$ for all x (see also [3,4], [12-15], [33]). It is easy to show that the function $f(x) = x^4$ satisfies the functional equation (1.4), which is called a quartic functional equation and every solution of the quartic functional equation is said to be a quartic function.

We deal with the following functional equation deriving from quartic and cubic functions:

$$\begin{aligned} 4(f(3x+y) + f(3x-y)) &= -12(f(x+y) + f(x-y)) + 12(f(2x+y) + f(2x-y)) \\ &\quad - 8f(y) - 192f(x) + f(2y) + 30f(2x). \end{aligned} \quad (1.5)$$

It is easy to see that the function $f(x) = ax^4 + bx^3$ is a solution of the functional equation (1.5). In the present paper we investigate the general solution and the generalized Hyers-Ulam-Rassias stability of the functional equation (1.5).

2. GENERAL SOLUTION

In this section we establish the general solution of functional equation (1.5).

Theorem 2.1. *Let X, Y be vector spaces, and let $f : X \rightarrow Y$ be a function. Then f satisfies (1.5) if and only if there exists a unique symmetric multi-additive function $Q : X \times X \times X \times X \rightarrow Y$ and a unique function $C : X \times X \times X \rightarrow Y$ such that C is symmetric for each fixed one variable and is additive for fixed two variables, and that $f(x) = Q(x, x, x, x) + C(x, x, x)$ for all $x \in X$.*

Proof. Suppose there exists a symmetric multi-additive function $Q : X \times X \times X \times X \rightarrow Y$ and a function $C : X \times X \times X \rightarrow Y$ such that C is symmetric for each fixed one variable and is additive for fixed two variables, and that $f(x) = Q(x, x, x, x) + C(x, x, x)$ for all $x \in X$. Then it is easy to see that f satisfies (1.5). For the converse let f satisfy (1.5). We decompose f into the even part and odd part by setting

$$f_e(x) = \frac{1}{2}(f(x) + f(-x)), \quad f_o(x) = \frac{1}{2}(f(x) - f(-x)),$$

for all $x \in X$. By (1.5), we have

$$\begin{aligned} 4f_e(3x+y) + 4f_e(3x-y) &= \frac{1}{2}[4f(3x+y) + 4f(-3x-y) + 4f(3x-y) + 4f(-3x+y)] \\ &= \frac{1}{2}[4f(3x+y) + 4f(3x-y)] + \frac{1}{2}[4f((-3x) + (-y)) + 4f((-3x) - (-y))] \\ &= \frac{1}{2}[12f(2x+y) + 12f(2x-y) - 12f(x+y) - 12f(x-y) \\ &\quad - 8f(y) - 192f(x) + f(2y) + 30f(2x)] \\ &\quad + \frac{1}{2}[12f(-2x-y) + 12f((-2x) + y) - 12f(-x-y) - 12f(-x+y) \\ &\quad - 8f(-y) - 192f(-x) + f(-2y) + 30f(-2x)] \\ &= 12[\frac{1}{2}(f(2x+y) + f(-(2x+y))) + 12[\frac{1}{2}(f(2x-y) + f(-(2x-y)))] \\ &\quad - 12[\frac{1}{2}(f(x+y) + f(-(x+y)))] - 12[\frac{1}{2}(f(x-y) + f(-(x-y)))] \\ &\quad - 8[\frac{1}{2}(f(y) + f(-y))] - 192[\frac{1}{2}(f(x) + f(-x))] \\ &\quad + \frac{1}{2}[f(2y) + f(-2y)] + 30[\frac{1}{2}(f(2x) + f(-2x))] \\ &= 12(f_e(2x+y) + f_e(2x-y)) - 12(f_e(x+y) + f_e(x-y)) \\ &\quad - 8f_e(y) - 192f_e(x) + f_e(2y) + 30f_e(2x) \end{aligned}$$

for all $x, y \in X$. This means that f_e satisfies (1.5), or

$$4(f_e(3x+y) + f_e(3x-y)) = -12(f_e(x+y) + f_e(x-y)) + 12(f_e(2x+y) + f_e(2x-y)) \\ - 8f_e(y) - 192f_e(x) + f_e(2y) + 30f_e(2x). \quad (1.5(e))$$

Now putting $x = y = 0$ in (1.5(e)), we get $f_e(0) = 0$. Setting $x = 0$ in (1.5(e)), by evenness of f_e we obtain

$$f_e(2y) = 16f_e(y) \quad (2.1)$$

for all $y \in X$. Hence (1.5(e)) can be written as

$$f_e(3x+y) + f_e(3x-y) + 3(f_e(x+y) + f_e(x-y)) = 3(f_e(2x+y) + f_e(2x-y)) + 72f_e(x) + 2f_e(y) \quad (2.2)$$

for all $x, y \in X$. With the substitution $y := 2y$ in (2.2), we have

$$f_e(3x+2y) + f_e(3x-2y) + 3f_e(x+2y) + 3f_e(x-2y) = 48f_e(x+y) + 48f_e(x-y) + 72f_e(x) + 32f_e(y). \quad (2.3)$$

Replacing y by $x+2y$ in (2.2), we obtain

$$16f_e(2x+y) + 16f_e(2x-y) + 48f_e(x+y) + 48f_e(x-y) = 3f_e(3x+2y) + 3f_e(3x-2y) + 2f_e(x+2y) + 72f_e(x). \quad (2.4)$$

Substituting $-y$ for y in (2.4) gives

$$16f_e(2x-y) + 16f_e(2x+y) + 48f_e(x-y) + 48f_e(x+y) = 3f_e(3x-2y) + 3f_e(3x+2y) + 2f_e(x-2y) + 72f_e(x). \quad (2.5)$$

By utilizing equations (2.3), (2.4) and (2.5), we obtain

$$4f_e(2x+y) + 4f_e(2x-y) + f_e(x+2y) + f_e(x-2y) = 20f_e(x+y) + 20f_e(x-y) + 90f_e(x). \quad (2.6)$$

Interchanging x and y in (2.3), we get

$$f_e(2x+3y) + f_e(2x-3y) + 3f_e(2x+y) + 3f_e(2x-y) = 48f_e(x+y) + 48f_e(x-y) + 32f_e(x) + 72f_e(y). \quad (2.7)$$

If we add (2.3) to (2.7), we have

$$f_e(2x+3y) + f_e(3x+2y) + f_e(2x-3y) + f_e(3x-2y) + 3f_e(2x+y) + 3f_e(2x-y) + 3f_e(x+2y) + 3f_e(x-2y) \\ = 96f_e(x+y) + 96f_e(x-y) + 104f_e(x) + 104f_e(y). \quad (2.8)$$

And by utilizing equations (2.4), (2.5) and (2.8), we arrive at

$$\begin{aligned}
3f_e(2x+3y) + 3f_e(2x-3y) &= -25f_e(2x+y) - 25f_e(2x-y) \\
&- 4f_e(x-2y) - 4f_e(x+2y) + 224f_e(x+y) \\
&+ 224f_e(x-y) + 456f_e(x) + 216f_e(y).
\end{aligned} \tag{2.9}$$

Let us interchange x and y in (2.9). Then we see that

$$\begin{aligned}
3f_e(3x+2y) + 3f_e(3x-2y) &= -25f_e(x+2y) - 25f_e(x-2y) \\
&- 4f_e(2x-y) - 4f_e(2x+y) + 224f_e(x+y) \\
&+ 224f_e(x-y) + 456f_e(y) + 216f_e(x).
\end{aligned} \tag{2.10}$$

Comparing (2.10) with (2.3), we get

$$\begin{aligned}
4f_e(2x-y) + 4f_e(2x+y) &= -16f_e(x+2y) \\
&- 16f_e(x-2y) + 80f_e(x+y) + 80f_e(x-y) + 360f_e(y).
\end{aligned} \tag{2.11}$$

If we compare (2.11) and (2.6), we conclude that

$$f_e(x+2y) + f_e(x-2y) + 6f_e(x) = 4f_e(x+y) + 4f_e(x-y) + 24f_e(y).$$

This means that f_e is quartic function. Thus there exists a unique symmetric multi-additive function $Q : X \times X \times X \times X \rightarrow Y$ such that $f_e(x) = Q(x, x, x, x)$ for all $x \in X$. On the other hand we can show that f_o satisfies (1.5), or

$$\begin{aligned}
4(f_o(3x+y) + f_o(3x-y)) &= -12(f_o(x+y) + f_o(x-y)) + 12(f_o(2x+y) + f_o(2x-y)) \\
&- 8f_o(y) - 192f_o(x) + f_o(2y) + 30f_o(2x).
\end{aligned} \tag{1.5(o)}$$

Now setting $x = y = 0$ in (1.5(o)) gives $f_o(0) = 0$. Putting $x = 0$ in (1.5(o)), then by oddness of f_o , we have

$$f_o(2y) = 8f_o(y). \tag{2.12}$$

Hence (1.5(o)) can be written as

$$\begin{aligned}
f_o(3x+y) + f_o(3x-y) + 3f_o(x+y) \\
+ 3f_o(x-y) &= 3f_o(2x+y) + 3f_o(2x-y) + 12f_o(x)
\end{aligned} \tag{2.13}$$

for all $x, y \in X$. Replacing x by $x+y$, and y by $x-y$ in (2.13) we have

$$\begin{aligned}
8f_o(2x+y) + 8f_o(x+2y) + 24f_o(x) \\
+ 24f_o(y) &= 3f_o(3x+y) + 3f_o(x+3y) + 12f_o(x+y)
\end{aligned} \tag{2.14}$$

and interchanging x and y in (2.13) yields

$$\begin{aligned}
f_o(x+3y) - f_o(x-3y) + 3f_o(x+y) \\
- 3f_o(x-y) &= 3f_o(x+2y) - 3f_o(x-2y) + 12f_o(y).
\end{aligned} \tag{2.15}$$

Which on substitution of $-y$ for y in (2.13) gives

$$\begin{aligned}
& f_o(3x - y) + f_o(3x + y) + 3f_o(x - y) \\
& + 3f_o(x + y) = 3f_o(2x - y) + 3f_o(2x + y) + 12f_o(x).
\end{aligned} \tag{2.16}$$

Replace y by $x + 2y$ in (2.13). Then we have

$$\begin{aligned}
& 8f_o(2x + y) + 8f_o(x - y) + 24f_o(x + y) \\
& - 24f_o(y) = 3f_o(3x + 2y) + 3f_o(x - 2y) + 12f_o(x).
\end{aligned} \tag{2.17}$$

From the substitution $y := -y$ in (2.17) it follows that

$$\begin{aligned}
& 8f_o(2x - y) + 8f_o(x + y) + 24f_o(x - y) \\
& + 24f_o(y) = 3f_o(3x - 2y) + 3f_o(x + 2y) + 12f_o(x).
\end{aligned} \tag{2.18}$$

If we add (2.17) to (2.18), we have

$$\begin{aligned}
& 3f_o(3x - 2y) + 3f_o(3x + 2y) = 8f_o(2x + y) + 8f_o(2x - y) \\
& - 3f_o(x + 2y) - 3f_o(x - 2y) + 32f_o(x - y) \\
& + 32f_o(x + y) - 24f_o(x).
\end{aligned} \tag{2.19}$$

Let us interchange x and y in (2.19). Then we see that

$$\begin{aligned}
& 3f_o(2x + 3y) - 3f_o(2x - 3y) = 8f_o(x + 2y) - 8f_o(x - 2y) \\
& - 3f_o(2x + y) + 3f_o(2x - y) + 32f_o(x + y) \\
& - 32f_o(x - y) - 24f_o(y).
\end{aligned} \tag{2.20}$$

With the substitution $y := x + y$ in (2.13), we have

$$\begin{aligned}
& f_o(4x + y) + f_o(2x - y) + 3f_o(2x + y) \\
& - 3f_o(y) = 3f_o(3x + y) + 3f_o(x - y) + 12f_o(x)
\end{aligned} \tag{2.21}$$

and replacing $-y$ by y gives

$$\begin{aligned}
& f_o(4x - y) + f_o(2x + y) + 3f_o(2x - y) \\
& + 3f_o(y) = 3f_o(3x - y) + 3f_o(x + y) + 12f_o(x).
\end{aligned} \tag{2.22}$$

If we add (2.21) to (2.22), we have

$$\begin{aligned}
& f_o(4x + y) + f_o(4x - y) = 3f_o(3x + y) + 3f_o(3x - y) \\
& - 4f_o(2x - y) - 4f_o(2x + y) + 3f_o(x - y) \\
& + 3f_o(x + y) + 24f_o(x).
\end{aligned} \tag{2.23}$$

By comparing (2.16) with (2.23), we arrive at

$$\begin{aligned}
& f_o(4x + y) + f_o(4x - y) = 5f_o(2x + y) + 5f_o(2x - y) \\
& - 6f_o(x + y) - 6f_o(x - y) + 60f_o(x)
\end{aligned} \tag{2.24}$$

and replacing y by $2y$ in (2.13) gives

$$\begin{aligned} f_o(3x+2y) + f_o(3x-2y) &= 24f_o(x+y) + 24f_o(x-y) \\ &\quad - 3f_o(x+2y) - 3f_o(x-2y) + 12f_o(x). \end{aligned} \quad (2.25)$$

By comparing (2.25) with (2.19), we arrive at

$$\begin{aligned} 3f_o(x+2y) + 3f_o(x-2y) &= 20f_o(x+y) + 20f_o(x-y) \\ &\quad - 4f_o(2x+y) - 4f_o(2x-y) + 30f_o(x). \end{aligned} \quad (2.26)$$

Let us interchange x and y in (2.25). Then we see that

$$\begin{aligned} f_o(2x+3y) - f_o(2x-3y) &= 24f_o(x+y) - 24f_o(x-y) \\ &\quad - 3f_o(2x+y) + 3f_o(2x-y) + 12f_o(y). \end{aligned} \quad (2.27)$$

Thus combining (2.27) with (2.20) yields

$$\begin{aligned} 4f_o(x+2y) - 4f_o(x-2y) &= 3f_o(2x-y) - 3f_o(2x+y) \\ &\quad + 20f_o(x+y) - 20f_o(x-y) + 30f_o(y). \end{aligned} \quad (2.28)$$

By comparing (2.28) with (2.15), we arrive at

$$\begin{aligned} 4f_o(x+3y) - 4f_o(x-3y) &= 9f_o(2x-y) - 9f_o(2x+y) \\ &\quad + 48f_o(x+y) - 48f_o(x-y) + 138f_o(y). \end{aligned} \quad (2.29)$$

Which, by putting $y := 2y$ in (2.14), leads to

$$\begin{aligned} 64f_o(x+y) + 8f_o(x+4y) + 24f_o(x) \\ + 192f_o(y) &= 3f_o(3x+2y) + 3f_o(x+6y) + 12f_o(x+2y). \end{aligned} \quad (2.30)$$

Replacing y by $-y$ in (2.30) gives

$$\begin{aligned} 64f_o(x-y) + 8f_o(x-4y) + 24f_o(x) \\ - 192f_o(y) &= 3f_o(3x-2y) + 3f_o(x-6y) + 12f_o(x-2y). \end{aligned} \quad (2.31)$$

If we subtract (2.30) from (2.31), we obtain

$$\begin{aligned} 8f_o(x+4y) - 8f_o(x-4y) &= 3f_o(3x+2y) - 3f_o(3x-2y) \\ &\quad + 3f_o(x+6y) - 3f_o(x-6y) + 12f_o(x+2y) - 12f_o(x-2y) \\ &\quad + 64f_o(x-y) - 64f_o(x+y) - 384f_o(y). \end{aligned} \quad (2.32)$$

Setting x instead of y and y instead of x in (2.24), we get

$$\begin{aligned} f_o(x+4y) - f_o(x-4y) &= 5f_o(x+2y) - 5f_o(x-2y) \\ &\quad + 6f_o(x-y) - 6f_o(x+y) + 60f_o(y). \end{aligned} \quad (2.33)$$

Combining (2.32) and (2.33) yields

$$\begin{aligned}
3f_o(3x+2y) - 3f_o(3x-2y) &= 28f_o(x+2y) - 28f_o(x-2y) \\
&+ 3f_o(x-6y) - 3f_o(x+6y) + 16f_o(x+y) \\
&- 16f_o(x-y) + 864f_o(y)
\end{aligned} \tag{2.34}$$

and subtracting (2.18) from (2.17), we obtain

$$\begin{aligned}
3f_o(3x+2y) - 3f_o(3x-2y) &= 3f_o(x+2y) - 3f_o(x-2y) \\
&+ 8f_o(2x+y) - 8f_o(2x-y) + 16f_o(x+y) \\
&- 16f_o(x-y) - 48f_o(y).
\end{aligned} \tag{2.35}$$

By comparing (2.34) with (2.35), we arrive at

$$\begin{aligned}
3f_o(x+6y) - 3f_o(x-6y) &= 25f_o(x+2y) - 25f_o(x-2y) \\
&+ 8f_o(2x-y) - 8f_o(2x+y) + 912f_o(y).
\end{aligned} \tag{2.36}$$

Interchanging y with $2y$ in (2.29) gives the equation

$$\begin{aligned}
4f_o(x+6y) - 4f_o(x-6y) &= 48f_o(x+2y) - 48f_o(x-2y) \\
&+ 72f_o(x-y) - 72f_o(x+y) + 1104f_o(y).
\end{aligned} \tag{2.37}$$

We obtain from (2.36) and (2.37)

$$\begin{aligned}
44f_o(x+2y) - 44f_o(x-2y) &= 32f_o(2x-y) - 32f_o(2x+y) \\
&+ 216f_o(x+y) - 216f_o(x-y) + 336f_o(y).
\end{aligned} \tag{2.38}$$

By using (2.28) and (2.38), we lead to

$$f_o(2x+y) - f_o(2x-y) = 4f_o(x+y) - 4f_o(x-y) - 6f_o(y). \tag{2.39}$$

And interchanging x with y in (2.39) gives

$$f_o(x+2y) + f_o(x-2y) = 4f_o(x+y) + 4f_o(x-y) - 6f_o(x). \tag{2.40}$$

If we compare (2.40) and (2.26), we conclude that

$$8f_o(x+y) + 8f_o(x-y) + 48f_o(x) = 4f_o(2x+y) + 4f_o(2x-y).$$

This means that f_o is cubic function and that there exists a unique function $C : X \times X \times X \rightarrow Y$ such that $f_o(x) = C(x, x, x)$ for all $x \in X$, and C is symmetric for each fixed one variable and is additive for fixed two variables. Thus for all $x \in X$, we have

$$f(x) = f_e(x) + f_o(x) = C(x, x, x) + Q(x, x, x).$$

This completes the proof of Theorem. \square

The following Corollary is an alternative result of above Theorem 2.1.

Corollary 2.2. *Let X, Y be vector spaces, and let $f : X \rightarrow Y$ be a function satisfying (1.5). Then the following assertions hold.*

- a) *If f is even function, then f is quartic.*
- b) *If f is odd function, then f is cubic.*

3. STABILITY

We now investigate the generalized Hyers-Ulam-Rassias stability problem for functional equation (1.5). From now on, let X be a real vector space and let Y be a Banach space. Now before taking up the main subject, given $f : X \rightarrow Y$, we define the difference operator $D_f : X \times X \rightarrow Y$ by

$$D_f(x, y) = 4[f(3x + y) + f(3x - y)] - 12[f(2x + y) + f(2x - y)] + 12[f(x + y) + f(x - y)] - f(2y) + 8f(y) - 30f(2x) + 192f(x)$$

for all $x, y \in X$. We consider the following functional inequality :

$$\|D_f(x, y)\| \leq \phi(x, y)$$

for an upper bound $\phi : X \times X \rightarrow [0, \infty)$.

Theorem 3.1. *Let $s \in \{1, -1\}$ be fixed. Suppose that an even mapping $f : X \rightarrow Y$ satisfies $f(0) = 0$, and*

$$\|D_f(x, y)\| \leq \phi(x, y) \quad (3.1)$$

for all $x, y \in X$. If the upper bound $\phi : X \times X \rightarrow [0, \infty)$ is a mapping such that the series $\sum_{i=0}^{\infty} 2^{4si} \phi(0, \frac{x}{2^{si}})$ converges, and that $\lim_{n \rightarrow \infty} 2^{4sn} \phi(\frac{x}{2^{sn}}, \frac{y}{2^{sn}}) = 0$ for all $x, y \in X$, then the limit $Q(x) = \lim_{n \rightarrow \infty} 2^{4sn} f(\frac{x}{2^{sn}})$ exists for all $x \in X$, and $Q : X \rightarrow Y$ is a unique quartic function satisfying (1.5), and

$$\|f(x) - Q(x)\| \leq \frac{1}{16} \sum_{i=\frac{s-1}{2}}^{\infty} 2^{4s(i+1)} \phi(0, \frac{x}{2^{s(i+1)}}) \quad (3.2)$$

for all $x \in X$.

Proof. Let $s = 1$. Putting $x = 0$ in (3.1), we get

$$\|f(2y) - 16f(y)\| \leq \phi(0, y). \quad (3.3)$$

Replacing y by $\frac{x}{2}$ in (3.3), yields

$$\|f(x) - 16f(\frac{x}{2})\| \leq \phi(0, \frac{x}{2}). \quad (3.4)$$

Interchanging x with $\frac{x}{2}$ in (3.4), and multiplying by 16 it follows that

$$\|16f(\frac{x}{2}) - 16^2 f(\frac{x}{4})\| \leq 16\phi(0, \frac{x}{4}). \quad (3.5)$$

Combining (3.4) and (3.5), we lead to

$$\|16^2 f(\frac{x}{4}) - f(x)\| \leq \phi(0, \frac{x}{2}) + 16\phi(0, \frac{x}{4}). \quad (3.6)$$

From the inequality (3.4) we use iterative methods and induction on n to prove our next relation:

$$\|16^n f(\frac{x}{2^n}) - f(x)\| \leq \frac{1}{16} \sum_{i=0}^{n-1} 16^{i+1} \phi(0, \frac{x}{2^{i+1}}). \quad (3.7)$$

We multiply (3.7) by 16^m and replace x by $\frac{x}{2^m}$ to obtain that

$$\| 16^{m+n} f(\frac{x}{2^{m+n}}) - 16^m f(\frac{x}{2^m}) \| \leq \sum_{i=0}^{n-1} 16^{m+i} \phi(0, \frac{x}{2^{i+m+1}}).$$

This shows that $\{16^n f(\frac{x}{2^n})\}$ is a Cauchy sequence in Y by taking the limit $m \rightarrow \infty$. Since Y is a Banach space, it follows that the sequence $\{16^n f(\frac{x}{2^n})\}$ converges. We define $Q : X \rightarrow Y$ by $Q(x) = \lim_{n \rightarrow \infty} 2^{4n} f(\frac{x}{2^n})$ for all $x \in X$. It is clear that $Q(-x) = Q(x)$ for all $x \in X$, and it follows from (3.1) that

$$\| D_Q(x, y) \| = \lim_{n \rightarrow \infty} 16^n \| D_f(\frac{x}{2^n}, \frac{y}{2^n}) \| \leq \lim_{n \rightarrow \infty} 16^n \phi(\frac{x}{2^n}, \frac{y}{2^n}) = 0$$

for all $x, y \in X$. Hence by Corollary 2.2, Q is quartic. It remains to show that Q is unique. Suppose that there exists another quartic function $Q' : X \rightarrow Y$ which satisfies (1.5) and (3.2). Since $Q(2^n x) = 16^n Q(x)$, and $Q'(2^n x) = 16^n Q'(x)$ for all $x \in X$, we conclude that

$$\begin{aligned} \| Q(x) - Q'(x) \| &= 16^n \| Q(\frac{x}{2^n}) - Q'(\frac{x}{2^n}) \| \\ &\leq 16^n \| Q(\frac{x}{2^n}) - f(\frac{x}{2^n}) \| + 16^n \| Q'(\frac{x}{2^n}) - f(\frac{x}{2^n}) \| \\ &\leq 2 \sum_{i=0}^{\infty} 16^{n+i} \phi(0, \frac{x}{2^{n+i+1}}) \end{aligned}$$

for all $x \in X$. By letting $n \rightarrow \infty$ in this inequality, it follows that $Q(x) = Q'(x)$ for all $x \in X$, which gives the conclusion. For $s = -1$, we obtain

$$\| \frac{f(2^m x)}{16^m} - f(x) \| \leq \frac{1}{16} \sum_{i=-1}^{n-2} \frac{\phi(0, 2^{i+1} x)}{16^{i+1}},$$

from which one can prove the result by a similar technique. \square

Theorem 3.2. *Let $s \in \{1, -1\}$ be fixed. Suppose that an odd mapping $f : X \rightarrow Y$ satisfies*

$$\| D_f(x, y) \| \leq \phi(x, y) \tag{3.8}$$

for all $x, y \in X$. If the upper bound $\phi : X \times X \rightarrow [0, \infty)$ is a mapping such that $\sum_{i=0}^{\infty} 2^{3si} \phi(0, \frac{x}{2^{si}})$ converges, and that $\lim_{n \rightarrow \infty} 2^{3si} \phi(\frac{x}{2^{si}}, \frac{y}{2^{si}}) = 0$ for all $x, y \in X$, then the limit $C(x) = \lim_{n \rightarrow \infty} 2^{3sn} f(\frac{x}{2^{sn}})$ exists for all $x \in X$, and $C : X \rightarrow Y$ is a unique cubic function satisfying (1.5), and

$$\| f(x) - C(x) \| \leq \frac{1}{8} \sum_{i=\frac{s-1}{2}}^{\infty} 2^{3s(i+1)} \phi(0, \frac{x}{2^{s(i+1)}}) \tag{3.9}$$

for all $x \in X$.

Proof. Let $s = 1$. Set $x = 0$ in (3.8). We obtain

$$\| 8f(y) - f(2y) \| \leq \phi(0, y). \tag{3.10}$$

Replacing y by $\frac{x}{2}$ in (3.10) to get

$$\| 8f(\frac{x}{2}) - f(x) \| \leq \phi(0, \frac{x}{2}). \tag{3.11}$$

An induction argument now implies

$$\| 8^n f(\frac{x}{2^n}) - f(x) \| \leq \frac{1}{8} \sum_{i=0}^{n-1} 8^{i+1} \phi(0, \frac{x}{2^{i+1}}). \quad (3.12)$$

Multiply (3.12) by 8^m and replace x by $\frac{x}{2^m}$, we obtain that

$$\| 8^{m+n} f(\frac{x}{2^{m+n}}) - 8^m f(\frac{x}{2^m}) \| \leq \sum_{i=0}^{n-1} 8^{m+i} \phi(0, \frac{x}{2^{m+i+1}}). \quad (3.13)$$

The right hand side of the inequality (3.13) tends to 0 as $m \rightarrow \infty$ because of

$$\sum_{i=0}^{\infty} 8^i \phi(0, \frac{x}{2^{i+1}}) < \infty$$

by assumption, and thus the sequence $\{2^{3n} f(\frac{x}{2^n})\}$ is Cauchy in Y , as desired. Therefore we may define a mapping $C : X \rightarrow Y$ as $C(x) = \lim_{n \rightarrow \infty} 2^{3n} f(\frac{x}{2^n})$. The rest of proof is similar to the proof of Theorem 3.1. \square

Theorem 3.3. *Let $s \in \{1, -1\}$ be fixed. Suppose a mapping $f : X \rightarrow Y$ satisfies $f(0) = 0$, and $\|D_f(x, y)\| \leq \phi(x, y)$ for all $x, y \in X$. If the upper bound $\phi : X \times X \rightarrow [0, \infty)$ is a mapping such that*

$$\sum_{i=0}^{\infty} [(|s| + s) 2^{4si} \phi(0, \frac{x}{2^{si-1}}) + (|s| - s) 2^{3si} \phi(0, \frac{x}{2^{si-1}})] < \infty, \quad (3.14)$$

and

$$\lim_{n \rightarrow \infty} [(|s| + s) 2^{(4sn-1)} \phi(\frac{x}{2^{sn}}, \frac{y}{2^{sn}}) + (|s| - s) 2^{3sn} \phi(\frac{x}{2^{sn}}, \frac{y}{2^{sn}})] = 0, \quad (3.15)$$

for all $x, y \in X$. Then there exists a unique quartic function $Q : X \rightarrow Y$ and a unique cubic function $C : X \rightarrow Y$ satisfying

$$\| f(x) - Q(x) - C(x) \| \leq \sum_{i=\frac{s-1}{2}}^{\infty} \{ (\frac{2^{4s(i+1)}}{32} + \frac{2^{3s(i+1)}}{16}) [\phi(0, \frac{x}{2^{s(i+1)}}) + \phi(0, \frac{-x}{2^{s(i+1)}})] \}$$

for all $x \in X$.

Proof. Let $f_e(x) = \frac{1}{2}(f(x) + f(-x))$ for all $x \in X$. Then $f_e(0) = 0$ and f_e is even function satisfying $\|D_{f_e}(x, y)\| \leq \frac{1}{2}[\phi(x, y) + \phi(-x, -y)]$ for all $x, y \in X$. From Theorem 3.1, it follows that there exists a unique quartic function $Q : X \rightarrow Y$ satisfies

$$\| f_e(x) - Q(x) \| \leq \frac{1}{32} \sum_{i=\frac{s-1}{2}}^{\infty} \{ 2^{4s(i+1)} \phi(0, \frac{x}{2^{s(i+1)}}) + 2^{4s(i+1)} \phi(0, \frac{-x}{2^{s(i+1)}}) \} \quad (3.16)$$

for all $x \in X$. Let now $f_o(x) = \frac{1}{2}(f(x) - f(-x))$ for all $x \in X$. Then f_o is odd function satisfying

$$\|D_{f_o}(x, y)\| \leq \frac{1}{2}[\phi(x, y) + \phi(-x, -y)]$$

for all $x, y \in X$. Hence in view of Theorem 3.2, it follows that there exists a unique cubic function $C : X \rightarrow Y$ such that

$$\|f_o(x) - C(x)\| \leq \frac{1}{16} \sum_{i=\frac{s-1}{2}}^{\infty} \{2^{3s(i+1)} \phi(0, \frac{x}{2^{s(i+1)}}) + 2^{3s(i+1)} \phi(0, \frac{-x}{2^{s(i+1)}})\} \quad (3.17)$$

for all $x \in X$. On the other hand we have $f(x) = f_e(x) + f_o(x)$ for all $x \in X$. Then by combining (3.16) and (3.17), it follows that

$$\begin{aligned} \|f(x) - C(x) - Q(x)\| &\leq \|f_e(x) - Q(x)\| + \|f_o(x) - C(x)\| \\ &\leq \sum_{i=\frac{s-1}{2}}^{\infty} \{(\frac{2^{4s(i+1)}}{32} + \frac{2^{3s(i+1)}}{16})[\phi(0, \frac{x}{2^{s(i+1)}}) + \phi(0, \frac{-x}{2^{s(i+1)}})]\} \end{aligned}$$

for all $x \in X$, and the proof of Theorem is complete. \square

We are going to investigate the Hyers-Ulam -Rassias stability problem for functional equation (1.5).

Corollary 3.4. *Let $p \in (-\infty, 3) \cup (4, +\infty)$, $\theta > 0$. Suppose $f : X \rightarrow Y$ satisfies $f(0) = 0$, and inequality*

$$\|D_f(x, y)\| \leq \theta(\|x\|^p + \|y\|^p),$$

for all $x, y \in X$. Then there exists a unique quartic function $Q : X \rightarrow Y$, and a unique cubic function $C : X \rightarrow Y$ satisfying

$$\|f(x) - Q(x) - C(x)\| \leq \begin{cases} \theta\|x\|^p(\frac{1}{2^p-2^4} + \frac{1}{2^p-2^3}), & p > 4, \\ \theta\|x\|^p(\frac{1}{2^4-2^p} + \frac{1}{2^3-2^p}), & p < 3 \end{cases}$$

for all $x \in X$.

Proof. Let $s = 1$ in Theorem 3.3. Then by taking $\phi(x, y) = \theta(\|x\|^p + \|y\|^p)$ for all $x, y \in X$, the relations (3.14) and (3.15) hold for $p > 4$. Then there exists a unique quartic function $Q : X \rightarrow Y$ and a unique cubic function $C : X \rightarrow Y$ satisfying

$$\|f(x) - Q(x) - C(x)\| \leq \theta\|\frac{x}{2}\|^p(\frac{1}{1-2^{4-p}} + \frac{1}{1-2^{3-p}})$$

for all $x \in X$. Let now $s = -1$ in Theorem 3.3 and put $\phi(x, y) = \theta(\|x\|^p + \|y\|^p)$ for all $x, y \in X$. Then the relations (3.14) and (3.15) hold for $p < 3$. Then there exists a unique quartic function $Q : X \rightarrow Y$ and a unique cubic function $C : X \rightarrow Y$ satisfying

$$\|f(x) - Q(x) - C(x)\| \leq \theta\|x\|^p(\frac{1}{2^4-2^p} + \frac{1}{2^3-2^p})$$

for all $x \in X$. \square

Similarly, we can prove the following Ulam stability problem for functional equation (1.5) controlled by the mixed type product-sum function

$$(x, y) \mapsto \theta(\|x\|_X^u \|y\|_X^v + \|x\|^p + \|y\|^p)$$

introduced by J. M. Rassias (see for example [32]).

Corollary 3.5. *Let u, v, p be real numbers such that $u + v, p \in (-\infty, 3) \cup (4, +\infty)$, and let $\theta > 0$. Suppose $f : X \rightarrow Y$ satisfies $f(0) = 0$, and inequality*

$$\|D_f(x, y)\| \leq \theta(\|x\|_X^u \|y\|_X^v + \|x\|^p + \|y\|^p),$$

for all $x, y \in X$. Then there exists a unique quartic function $Q : X \rightarrow Y$, and a unique cubic function $C : X \rightarrow Y$ satisfying

$$\|f(x) - Q(x) - C(x)\| \leq \begin{cases} \theta \|x\|^p (\frac{1}{2^p-2^4} + \frac{1}{2^p-2^3}), & p > 4, \\ \theta \|x\|^p (\frac{1}{2^4-2^p} + \frac{1}{2^3-2^p}), & p < 3 \end{cases}$$

for all $x \in X$.

By Corollary 3.4, we solve the following Hyers-Ulam stability problem for functional equation (1.5).

Corollary 3.6. *Let ϵ be a positive real number. Suppose $f : X \rightarrow Y$ satisfies $f(0) = 0$, and $\|D_f(x, y)\| \leq \epsilon$, for all $x, y \in X$. Then there exists a unique quartic function $Q : X \rightarrow Y$, and a unique cubic function $C : X \rightarrow Y$ satisfying*

$$\|f(x) - Q(x) - C(x)\| \leq \frac{22}{105} \epsilon$$

for all $x \in X$.

Acknowledgement. The authors would like to express their sincere thanks to referee for his invaluable comments. The first author would like to thank the Semnan University for its financial support. Also, The third author would like to thank the office of gifted students at Semnan University for its financial support.

REFERENCES

- [1] T. Aoki, On the stability of the linear transformation in Banach spaces, J. Math. Soc. Japan. 2(1950), 64-66.
- [2] P. W. Cholewa, Remarks on the stability of functional equations, Aequationes Math. 27 (1984) 76-86.
- [3] Chung, Jukang K. and Sahoo, Prasanna K., On the general solution of a quartic functional equation, Bull. Korean Math. Soc. 40 (2003), no. 4, 565-576.
- [4] Cădariu, Liviu, Fixed points in generalized metric space and the stability of a quartic functional equation Bul. Ştiinţ. Univ. Politeh. Timiş. Ser. Mat. Fiz. 50(64) (2005), no. 2, 25-34.
- [5] Z. Gajda, On stability of additive mappings, Internat. J. Math. Math. Sci. 14(1991) 431-434.
- [6] P. Găvruta, A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings, J. Math. Anal. Appl. 184 (1994) 431-436.
- [7] A. Grabiec, The generalized Hyers-Ulam stability of a class of functional equations, Publ. Math. Debrecen 48 (1996) 217-235.
- [8] D. H. Hyers, G. Isac and Th. M. Rassias, Stability of Functional Equations in Several Variables, Birkhäuser, Basel, 1998.
- [9] D. H. Hyers, On the stability of the linear functional equation, Proc. Natl. Acad. Sci. 27 (1941) 222-224.
- [10] G. Isac and Th. M. Rassias, On the Hyers-Ulam stability of ψ -additive mappings, J. Approx. Theory 72(1993), 131-137.

- [11] K. W. Jung and H. M. Kim, The generalized Hyers-Ulam-Rassias stability of a cubic functional equation, *J. Math. Anal. Appl.* 274, (2002), no. 2, 267–278.
- [12] Lee, Sang Han, Im, Sung Mo and Hwang, In Sung, Quartic functional equations *J. Math. Anal. Appl.* 307 (2005), no. 2, 387–394.
- [13] Najati, Abbas, On the stability of a quartic functional equation *J. Math. Anal. Appl.* 340 (2008), no. 1, 569–574.
- [14] Park, Chun-Gil, On the stability of the orthogonally quartic functional equation *Bull. Iranian Math. Soc.* 31 (2005), no. 1, 63–70.
- [15] Park, Won-Gil and Bae, Jae-Hyeong, On a bi-quadratic functional equation and its stability *Nonlinear Anal.* 62 (2005), no. 4, 643–654.
- [16] Rassias, John Michael, Solution of the Ulam stability problem for quartic mappings, *J. Indian Math. Soc. (N.S.)* 67 (2000), no. 1-4, 169–178.
- [17] Rassias, John Michael, Solution of the Ulam stability problem for cubic mappings. *Glas. Mat. Ser. III* 36(56) (2001), no. 1, 63–72.
- [18] Rassias, John Michael, Solution of the Ulam stability problem for cubic mappings. *An. Univ. Timi c soara Ser. Mat.-Inform.* 38(2000), no. 1, 121–132.
- [19] J. M. Rassias, Complete solution of the multi-dimensional problem of Ulam. *Discuss. Math.* 14 (1994), 101–107.
- [20] J. M. Rassias, On the stability of a multi-dimensional Cauchy type functional equation. *Geometry, analysis and mechanics*, 365–375, World Sci.Publ., River Edge, NJ, 1994.
- [21] J. M. Rassias, Solution of a stability problem of Ulam. *Functional analysis, approximation theory and numerical analysis*, 241–249, World Sci.Publ., River Edge, NJ, 1994.
- [22] J. M. Rassias, Solution of a stability problem of Ulam. *Discuss. Math.* 12 (1992), 95–103 (1993).
- [23] J. M. Rassias, Solution of a problem of Ulam. *J. Approx. Theory* 57 (1989), no. 3, 268–273.
- [24] J. M. Rassias, On a new approximation of approximately linear mappings by linear mappings. *Discuss. Math.* 7 (1985), 193–196.
- [25] J. M. Rassias, On approximation of approximately linear mappings by linear mappings. *Bull. Sci. Math. (2)* 108 (1984), no. 4, 445–446.
- [26] J. M. Rassias, On approximation of approximately linear mappings by linear mappings. *J.Funct. Anal.* 46 (1982), no. 1, 126–130.
- [27] Rassias, John Michael, Solution of the Ulam stability problem for quartic mappings, *Glas. Mat. Ser. III* 34(54) (1999), no. 2, 243–252.
- [28] Th. M. Rassias (Ed.), *Functional Equations and Inequalities*, Kluwer Academic, Dordrecht, 2000.
- [29] Th. M. Rassias, On the stability of functional equations and a problem of Ulam, *Acta Math. Appl.* 62 (2000) 23–130.
- [30] Th. M. Rassias, On the stability of functional equations in Banach spaces, *J. Math. Anal. Appl.* 251 (2000) 264–284.
- [31] Th. M. Rassias, On the stability of the linear mapping in Banach spaces, *Proc. Amer. Math. Soc.* 72 (1978) 297–300.
- [32] K. Ravi, M. Arunkumar and J. M. Rassias, Ulam stability for the orthogonally general Euler-Lagrange type functional equation, *Int. J. Math. Stat.* 3 (2008), A08, 36–46. 39B55 (39B82)
- [33] E. Thandapani, K. Ravi and M. Arunkumar, On the solution of the generalized quartic functional equation *Far East J. Appl. Math.* 24 (2006), no. 3, 297–312.
- [34] S. M. Ulam, *Problems in Modern Mathematics*, Chapter VI, science ed., Wiley, New York, 1940.