# Circuit partitions and #P-complete products of inner products

Cristopher Moore Computer Science Department, University of New Mexico and the Santa Fe Institute Alexander Russell
Department of Computer Science & Engineering
University of Connecticut

November 9, 2021

#### Abstract

We present a simple, natural #P-complete problem. Let G be a directed graph, and let k be a positive integer. We define q(G;k) as follows. At each vertex v, we place a k-dimensional complex vector  $x_v$ . We take the product, over all edges (u,v), of the inner product  $\langle x_u, x_v \rangle$ . Finally, q(G;k) is the expectation of this product, where the  $x_v$  are chosen uniformly and independently from all vectors of norm 1 (or, alternately, from the Gaussian distribution). We show that q(G;k) is proportional to G's cycle partition polynomial, and therefore that it is #P-complete for any k > 1.

### 1 Introduction

Let  $x, y \in \mathbb{C}^k$  be k-dimensional complex-valued vectors. We denote their inner product as

$$\langle x, y \rangle = \sum_{i=1}^k x_i^* y_i.$$

Now suppose we have a directed graph G=(V,E). Let us associate a vector  $x_v \in \mathbb{C}^k$  with each vertex v, and consider the product over all edges (u,v) of the inner products of the corresponding vectors:

$$\prod_{(u,v)\in E} \langle x_u, x_v \rangle \ . \tag{1}$$

For instance, for the graph in Figure 1 this product is

$$\prod_{(u,v)\in E} \langle x_u, x_v \rangle = \langle x_1, x_2 \rangle \langle x_2, x_3 \rangle \langle x_3, x_1 \rangle |\langle x_3, x_4 \rangle|^2 . \tag{2}$$

The expectation of this product, where each  $x_v$  is chosen independently and uniformly from the set of vectors in  $\mathbb{C}^k$  of norm 1, is a type of moment, where each  $x_v$  appears with order  $d_v^{\text{in}} + d_v^{\text{out}}$ . It is a function of the graph G and the dimension k, which we denote as follows:

$$q(G;k) = \underset{\{x_v\}}{\text{Exp}} \prod_{(u,v)\in E} \langle x_u, x_v \rangle .$$




Figure 1: A little directed graph. We use the edge labels in the proof of Theorem 1.

A simple observation is that q(G;k) is zero unless G is Eulerian—that is, unless  $d_v^{\text{in}} = d_v^{\text{out}}$  for each vertex v. Since  $x_v$  appears in the product  $d_v^{\text{in}}$  times unconjugated and  $d_v^{\text{out}}$  times conjugated, multiplying  $x_v$  by  $e^{i\theta}$  multiplies q(G;k) by  $e^{i\theta(d_v^{\text{in}}-d_v^{\text{out}})}$ . But multiplying by a phase preserves the uniform measure, so the expectation is zero if  $d_v^{\text{in}} \neq d_v^{\text{out}}$  for any v.

So, let us suppose that G is Eulerian. In that case, what is q(G;k)? Does it have a combinatorial interpretation? And how difficult is it to calculate? Our main result is this:

**Theorem 1.** For any  $k \geq 2$ , computing q(G;k), given G as input, is #P-hard under Turing reductions.

If we extend #P to rational functions in the natural way, then we can replace #P-hardness in this theorem with #P-completeness.

Our proof is very simple; we show that q(G;k) is essentially identical to an existing graph polynomial, which is known to be #P-hard to compute. Along the way, we will meet some nice combinatorics, and glancingly employ the representation theory of the unitary and orthogonal groups.

## 2 The circuit partition polynomial

A circuit partition of G is a partition of G's edges into circuits. Let  $r_t$  denote the number of circuit partitions containing t circuits; for instance,  $r_1$  is the number of Eulerian circuits. The circuit partition polynomial j(G; z) is the generating function

$$j(G;z) = \sum_{t=1}^{\infty} r_t z^t.$$
 (3)

For instance, for the graph in Figure 1 we have  $j(G; z) = z + z^2$ . This polynomial was first studied by Martin [10], with a slightly different parametrization; see also [1, 3, 4, 6, 9, 11, 12].

Now consider the following theorem.

**Theorem 2.** For any Eulerian directed graph G = (V, E),

$$q(G;k) = \left(\prod_{v \in V} \frac{(k-1)!}{(k+d_v-1)!}\right) j(G;k),$$
(4)

where  $d_v$  denotes  $d_v^{\text{in}} = d_v^{\text{out}}$ .

*Proof.* Given a vector  $x \in \mathbb{C}^k$  and an integer d, the outer product of  $x^{\otimes d} = \underbrace{x \otimes \cdots \otimes x}_{d \text{ times}}$  with itself

is a tensor of rank 2d, or equivalently a linear operator on  $(\mathbb{C}^k)^{\otimes d}$ :

$$\left|x^{\otimes d}\right\rangle\left\langle x^{\otimes d}\right| = \left(\left|x\right\rangle\left\langle x\right|\right)^{\otimes d}.$$

In terms of indices, we can write

$$\left| x^{\otimes d} \right\rangle \left\langle x^{\otimes d} \right|_{\beta_1 \beta_2 \cdots \beta_d}^{\alpha_1 \alpha_2 \cdots \alpha_d} = \prod_{\ell=1}^d x_{\alpha_\ell} x_{\beta_\ell}^*.$$

Then  $\prod_{(u,v)\in E} \langle x_u, x_v \rangle$  is a contraction of the product of these tensors, where upper and lower indices correspond to incoming and outgoing edges respectively. For instance, for the graph in Figure 1 we can rewrite the product 2 as

$$\prod_{(u,v)\in E} \langle x_u, x_v \rangle = |x_1\rangle \langle x_1|_{\alpha}^{\gamma} |x_2\rangle \langle x_2|_{\beta}^{\alpha} |x_3 \otimes x_3\rangle \langle x_3 \otimes x_3|_{\gamma\delta}^{\beta\eta} |x_4\rangle \langle x_4|_{\eta}^{\delta}.$$

Here we use the Einstein summation convention, where any index which appears once above and once below is automatically summed from 1 to k. Now, since the  $x_v$  are independent for different v, we can compute q(G;k) by taking the expectation over each  $x_v$  separately. This gives a contraction of the tensors

$$X_d = \operatorname{Exp}_{x} \left| x^{\otimes d} \right\rangle \left\langle x^{\otimes d} \right| \,, \tag{5}$$

where  $d = d_v$ , over all v.

In order to calculate  $X_d$ , we introduce some notation. Let  $S_d$  denote the symmetric group on d elements. We identify a permutation  $\pi \in S_d$  with the linear operator on  $(\mathbb{C}^k)^{\otimes d}$  which permutes the d factors in the tensor product. That is,

$$\pi (x_1 \otimes x_2 \otimes \cdots \otimes x_d) = x_{\pi(1)} \otimes x_{\pi(2)} \otimes \cdots \otimes x_{\pi(d)},$$

or, using indices,

$$\pi_{\beta_1\beta_2\cdots\beta_d}^{\alpha_1\alpha_2\cdots\alpha_d} = \prod_{\ell=1}^d \delta_{\beta_\ell}^{\alpha_{\pi(\ell)}},$$

where  $\delta_j^i$  is the Kronecker delta operator,  $\delta_j^i = 1$  if i = j and 0 if  $i \neq j$ . Diagrammatically,  $\pi$  is a gadget with d incoming edges and d outgoing edges, wired to each other according to the permutation  $\pi$ .

We have the following lemma:

**Lemma 1.** With  $X_d$  defined as in (5), if x is uniform in the set of vectors in  $\mathbb{C}^k$  of norm 1, then

$$X_d = \frac{(k-1)!}{(k+d-1)!} \sum_{\pi \in S_d} \pi.$$
 (6)

*Proof.* First,  $X_d$  is a member of the commutant of the group U(k) of  $k \times k$  unitary matrices, since these preserve the uniform measure. That is  $X_d$  commutes with  $U^{\otimes d}$  for any  $U \in U(k)$ . By Schur duality, the commutant is a quotient of the group algebra  $\mathbb{C}[S_d]$ ; namely, the image of  $\mathbb{C}[S_d]$  under the identification above. Thus  $X_d$  is a superposition of permutations,  $\sum_{\pi \in S_d} a_{\pi}\pi$ .

We also have  $X_d\pi = \pi X_d = X_d$  for any  $\pi$ . Thus  $X_d$  is proportional to the uniform superposition on  $S_d$ , or equivalently the projection operator  $\Pi_{\text{sym}} = (1/d!) \sum_{\pi} \pi$  onto the totally symmetric subspace  $V_{\text{sym}}$  of  $(\mathbb{C}^k)^{\otimes d}$ . Since  $\operatorname{tr} X_d = \operatorname{Exp}_x |x|^{2d} = 1$  while  $\operatorname{tr} \Pi_{\text{sym}} = \dim V_{\text{sym}}$ , we have  $X_d = (1/\dim V_{\text{sym}}) \Pi_{\text{sym}}$ .

Finally, dim  $V_{\text{sym}}$  is the number of ways to label the d factors of the tensor product with basis vectors  $\{e_1, \ldots, e_k\}$  in nondecreasing order—or, for afficient ados, the number of semistandard tableaux with one row of length d and content ranging from 1 to k. This gives dim  $V_{\text{sym}} = \binom{k+d-1}{d}$ .

To illustrate some ideas that will recur in the next section, we give an alternate proof. First, note that  $\operatorname{tr} \pi$  is the number of ways to label each of  $\pi$ 's cycles with a basis vector ranging from 1 to k, or  $k^{c(\pi)}$  where  $c(\pi)$  denotes the number of cycles (including fixed points). Thus

$$\operatorname{tr} \sum_{\pi \in S_d} \pi = \sum_{\pi \in S_d} k^{c(\pi)} \,. \tag{7}$$

To compute this generating function, we use the fact that each permutation  $\pi \in S_d$  appears once in the following product, where 1 denotes the identity permutation, and  $\tau_{ij}$  denotes the transposition of the *i*th and *j*th object:

$$\sum_{\pi \in S_d} \pi = 1(1 + \tau_{1,2})(1 + \tau_{1,3} + \tau_{2,3}) \cdots (1 + \tau_{1,d} + \tau_{2,d} + \cdots + \tau_{d-1,d}). \tag{8}$$

This product works by describing a permutation  $\pi_t$  of t objects inductively as a permutation  $\pi_{t-1}$  of the first t-1 objects, composed either with the identity or with a transposition swapping the tth object with one of the previous t-1. If we apply the identity, then the tth object is a fixed point, and  $c(\pi_t) = c(\pi_{t-1}) + 1$ , gaining a factor of k in (7); but if we apply a transposition, then  $c(\pi_t) = c(\pi_{t-1})$ . Thus (8) becomes

$$\sum_{\pi \in S_d} k^{c(\pi)} = k(k+1)(k+2)\cdots(k+d-1) = \frac{(k+d-1)!}{(k-1)!}.$$

Comparing traces again gives (6).

All that remains is to interpret the operators  $X_{d_v}$ , and their contraction, diagrammatically. Lemma 1 tells us that, for each vertex v of G, taking the expectation over  $x_v$  converts it to a sum over all  $d_v$ ! ways to wire the incoming edges to the outgoing edges. But doing this at each vertex gives us a sum over all cycle partitions of G. Contracting these tensors gives the number of ways to label each cycle in a each partition with a basis vector ranging from 1 to k, so each cycle contributes a factor of k. Along with the scaling factor in (6), this completes the proof.

Next we show that the cycle partition polynomial is #P-hard. To our knowledge, the following theorem first appeared in [7]; we prove it here for completeness.

**Theorem 3.** For any fixed z > 1, computing j(G; z) from G is #P-hard under Turing reductions.




Figure 2: A planar graph G (black) and its oriented medial graph  $G_m$  (gray).

*Proof.* Recall that the *Tutte polynomial* of an undirected graph G = (V, E) can be written as a sum over all subsets S of E,

$$T(G; x, y) = \sum_{S \subseteq E} (x - 1)^{c(S) - c(G)} (y - 1)^{c(S) + |S| - n}.$$
 (9)

Here c(G) denotes the number of connected components in G. Similarly, c(S) denotes the number of connected components in the spanning subgraph (V, S), including isolated vertices. When x = y, we have

$$T(G; x, x) = \sum_{S \subseteq E} (x - 1)^{c(S) + \ell(S) - c(G)},$$
(10)

where  $\ell(S) = c(s) + |S| - n$  is the total excess of the components of S, i.e., the number of edges that would have to be removed to make each one a tree.

If G is planar, then we can define a directed medial graph  $G_m$  as in Figure 2. Each vertex of  $G_m$  corresponds to an edge of G, edges of  $G_m$  correspond to shared vertices in G, and we orient the edges of  $G_m$  so that they go counterclockwise around the faces of G. Each vertex of  $G_m$  has  $d^{\text{in}} = d^{\text{out}} = 2$ , so  $G_m$  is Eulerian.

The following identity is due to Martin [10]; see also [11], or [2] for a review.

$$j(G_m; z) = z^{c(G)} T(G; z+1, z+1).$$
(11)

We prove this using a one-to-one correspondence between subsets  $S \subseteq E$  and circuit partitions of  $G_m$ . Let v be a vertex of  $G_m$ , corresponding to an edge e of G. Then the circuit partition connects each of v's incoming edge to the outgoing edge on the same side of e if  $e \in S$ , and crosses over to the other side if  $e \notin S$ . It is easy to prove by induction that the number of circuits is then  $c(S) + \ell(S)$ , in which case (10) yields (11).

The theorem then follows from the fact, proven by Vertigan [13], that the Tutte polynomial for planar graphs is #P-hard under Turing reductions, except on the hyperbolas  $(x-1)(y-1) \in \{1,2\}$  or when  $(x,y) \in \{(1,1),(-1,-1),(\omega,\omega^*),(\omega^*,\omega)\}$  where  $\omega = \mathrm{e}^{2\pi i/3}$ . Thus computing j(G;z) for any z>1 is #P-hard, even in the special case where G is planar and where every vertex has  $d^{\mathrm{in}}=d^{\mathrm{out}}=2$ .

### 3 Real-valued vectors

We can also consider the case where the  $x_v$  are real-valued, and are chosen uniformly from the set of vectors in  $\mathbb{R}^k$  of norm 1. In this case, the inner product  $\langle x_u, x_v \rangle$  becomes symmetric, so the graph G becomes undirected. We might then expect q(G;k) to be related to the circuit partition polynomial for undirected circuits, and indeed this is the case.

We again wish to compute the tensor  $X_d = \operatorname{Exp}_x |x^{\otimes d}\rangle\langle x^{\otimes d}|$ . First, let  $M_d$  denote the set of perfect matchings of 2d objects; note that

$$|M_d| = (2d-1)!! = (2d-1)(2d-3)\cdots 5\cdot 3\cdot 1 = \frac{(2d)!}{2^d d!}.$$

We can identify each matching  $\mu \in M_d$  with a linear operator on  $(\mathbb{R}^k)^{\otimes d}$ , where the first d objects correspond to upper indices, and the last d correspond to lower indices. However, in addition to permutations that wire upper indices to lower ones with a bipartite matching, we now also have "cups" and "caps" that wire two upper indices, or two lower indices, to each other. For instance, if d = 2 then  $M_d$  includes three operators, corresponding to the three perfect matchings of 4 objects:

$$\delta_{\beta_1}^{\alpha_1} \delta_{\beta_2}^{\alpha_2} = \left| \begin{array}{cc} & & \\ &$$

The first two of these operators correspond to the identity permutation and the transposition  $\tau_{1,2}$  respectively, as in the previous section. The third one is a *cupcap*; it is the outer product of the vector  $\sum_{i=1}^k e_i \otimes e_i$  with itself, where  $e_i$  denotes the *i*th basis vector in  $\mathbb{R}^k$ . We denote it  $\gamma_{1,2}$ , and more generally  $\gamma_{ij} = \delta^{\alpha_i,\alpha_j}\delta_{\beta_i,\beta_j}$ .

Now, in the real-valued case, Lemma 1 becomes the following:

**Lemma 2.** If x is uniform in the set of vectors in  $\mathbb{R}^k$  of norm 1, then

$$X_d = \frac{1}{k(k+2)(k+4)\cdots(k+2d-2)} \sum_{\mu \in M_d} \mu = \frac{(k-2)!!}{(k+2d-2)!!} \sum_{\mu \in M_d} \mu,$$
 (13)

where  $n!! = n(n-2)(n-4)\cdots 6\cdot 4\cdot 2$  if n is even, and  $n(n-2)(n-4)\cdots 5\cdot 3\cdot 1$  if n is odd.

*Proof.* Analogous to the complex case,  $X_d$  is a member of the commutant of the group O(k) of  $k \times k$  orthogonal matrices, since these preserve the uniform measure. That is,  $X_d$  commutes with  $O^{\otimes d}$  for any  $O \in O(k)$ . The commutant of O(k) is the *Brauer algebra*; namely, the algebra consisting of linear combinations of the operators  $\mu \in M_d$ . Thus  $X_d$  is of the form  $\sum_{\mu \in M_d} a_{\mu}\mu$ .

In addition to being fixed under permutations as in the complex case,  $X_d$  is also fixed under partial transposes, which switch some upper indices with some lower ones. Thus  $X_d$  is proportional to the uniform superposition  $\sum_{\mu \in M_d} \mu$ . To find the constant of proportionality, we again compare traces.

As in the case of permutations, the trace of an operator  $\mu \in M_d$  is  $k^{c(\mu)}$ , where  $c(\mu)$  is the number of loops in the diagram resulting from joining the upper indices to the lower ones. For instance, for the operators in (12), we have  $\operatorname{tr} 1 = k^2$ ,  $\operatorname{tr} \tau_{1,2} = k$ , and  $\operatorname{tr} \gamma_{1,2} = k$ . Thus we wish to calculate

$$\operatorname{tr} \sum_{\mu \in M_d} \mu = \sum_{\mu \in M_d} k^{c(\mu)} \,. \tag{14}$$

We can write  $\sum_{\mu \in M_d}$  as a product, analogous to (8):

$$\sum_{\pi \in S_d} \pi = 1(1 + \tau_{1,2} + \gamma_{1,2})(1 + \tau_{1,3} + \gamma_{1,3} + \tau_{2,3} + \gamma_{2,3}) \cdots (1 + \tau_{1,d} + \gamma_{1,d} + \cdots + \tau_{d-1,d} + \gamma_{d-1,d}).$$

This product describes a matching  $\mu_t$  of 2t objects inductively as a matching  $\mu_{t-1}$  of the first 2(t-1) objects, composed either with the identity, or with a transposition or cupcap connecting the tth upper object with the ith lower one and the tth lower object with the ith upper one, or vice versa. If we apply the identity, then the tth upper object is matched to the tth lower one, and  $c(\mu_t) = c(\mu_{t-1}) + 1$ , gaining a factor of k in (14); but if we apply a transposition or cupcap, then  $c(\mu_t) = c(\mu_{t-1})$ . Thus (14) becomes

$$\sum_{\pi \in S_d} k^{c(\pi)} = k(k+2)(k+4)\cdots(k+2d-2) = \frac{(k+2d-2)!!}{(k-2)!!}.$$

We again have  $\operatorname{tr} X_d = \operatorname{Exp}_x |x|^{2d} = 1$ , and comparing traces gives (13).

As before, q(G;k) is a contraction of the tensors  $X_d$ . However, now G is undirected, with no distinction between incoming and outgoing edges, so at each vertex of degree  $d_v$  the appropriate tensor is  $X_{d_v/2}$ . Applying Lemma 2 to each v sums over all the ways to match v's edges with each other, and hence sums over all possible partitions of G's edges into undirected cycles. The trace of the resulting diagram is again the number of ways to label each cycle with a basis vector. So, if define a polynomial  $j_{\text{undirected}}(G;z)$  as  $\sum_{t=1}^{\infty} r_t z^t$ , where  $r_t$  is the number of partitions with t cycles, then Theorem 2 becomes

**Theorem 4.** For any undirected graph G = (V, E) where every vertex has even degree, if we define q(G; k) by selecting the  $x_v$  independently and uniformly from the set of vectors in  $\mathbb{R}^k$  with norm 1, then

$$q(G;k) = \left(\prod_{v \in V} \frac{(k-2)!!}{(k+d_v-2)!!}\right) j_{\text{undirected}}(G;k).$$

$$(15)$$

To our knowledge, the computational complexity of  $j_{\text{undirected}}(G; z)$  is open, although it seems likely that it is also #P-hard.

### 4 The Gaussian distribution

Our results above assume that each  $x_v$  is chosen uniformly from the set of vectors in  $\mathbb{C}^k$  or  $\mathbb{R}^k$  of norm 1. Another natural measure would be to choose each component of  $x_v$  independently from the Gaussian distribution with variance 1/k, so that  $\text{Exp}[|x_v|^2] = 1$ .

For vectors in  $\mathbb{C}^k$ , the probability density of the norm |x| is then

$$p(|x|) = \frac{2k^{k+1}}{k!} |x|^{2k-1} e^{-k|x|^2},$$
(16)

Compared to the case where  $|x_v| = 1$ , each  $x_v$  contributes scaling factor of  $|x_v|^{2d}$  to the product (1). The even moments of (16) are

$$\operatorname{Exp}\left[|x|^{2d}\right] = \frac{(d+k-1)!}{k^d(k-1)!},$$

so in the Gaussian distribution (4) becomes

$$q(G;k) = \left(\prod_{v \in V} \frac{1}{k^{d_v}}\right) j(G;k) = \frac{1}{k^m} j(G;k),$$
(17)

where m denotes the number of edges.

We could also have derived this directly from the Gaussian analog of Lemma 1. If x is chosen according to the Gaussian distribution on  $\mathbb{C}^k$ , and we again let  $X_d$  denote  $\operatorname{Exp}_x |x^{\otimes d}\rangle\langle x^{\otimes d}|$ , then

$$X_d = \frac{1}{k^d} \sum_{\pi \in S_d} \pi. \tag{18}$$

Similarly, in the real-valued case, if we choose each component of  $x \in \mathbb{R}^k$  from the Gaussian distribution on  $\mathbb{R}$  with variance 1/k, then (15) becomes

$$q(G;k) = \frac{1}{k^m} j_{\text{undirected}}(G;k), \qquad (19)$$

since

$$X_d = \frac{1}{k^d} \sum_{\mu \in M_d} \mu. \tag{20}$$

Both (18) and (20) are consequences of Wick's Theorem [8, 15], that if  $x_1, \ldots, x_{2t}$  obey a multi-variate Gaussian distribution with mean zero, then

$$\operatorname{Exp}\left[\prod_{i=1}^{2t} x_i\right] = \sum_{\mu \in M_t} \prod_{(i,j) \in \mu} \operatorname{Exp}[x_i x_j].$$

**Acknowledgments.** We are grateful to Piotr Śniady for teaching us the sum (8), and to Jon Yard for introducing us to the Brauer algebra. This work was supported by the NSF under grant CCF-0829931, and by the DTO under contract W911NF-04-R-0009.

### References

- [1] R. Arratia, B. Bollobás, and G. Sorkin, The interlace polynomial: A new graph polynomial. Proc. 11th Annual ACM-SIAM Symposium on Discrete Algorithms 237–245 (2000).
- [2] Andrea Austin, The Circuit Partition Polynomial with Applications and Relation to the Tutte and Interlace Polynomials. *Rose-Hulman Undergraduate Mathematics Journal*, 8(2) (2007).
- [3] Béla Bollobás, Evaluations of the Circuit Partition Polynomial. *Journal of Combinatorial Theory, Series B* 85, 261–268 (2002)
- [4] André Bouchet, Tutte-Martin polynomials and orienting vectors of isotropic systems. *Graphs Combin.* 7(3) 235–252 (1991).
- [5] Richard Brauer, On Algebras Which are Connected with the Semisimple Continuous Groups. *Annals of Mathematics*, 38(4) 857–872 (1937).

- [6] Joanna A. Ellis-Monaghan, New results for the Martin polynomial. *Journal of Combinatorial Theory*, Series B 74, 326–352 (1998).
- [7] Joanna A. Ellis-Monaghan and Irasema Sarmiento, Distance hereditary graphs and the interlace polynomial. *Combinatorics, Probability and Computing* 16(6) 947–973 (2007).
- [8] L. Isserlis, On a formula for the product-moment coefficient of any order of a normal frequency distribution in any number of variables. *Biometrika* 12: 134–139 (1918).
- [9] F. Jaeger, On Tutte polynomials and cycles of plane graphs. *Journal of Combinatorial Theory*, Series B 44, 127–146 (1988).
- [10] P. Martin, Enumérations eulériennes dans les multigraphes et invariants de Tutte-Grothendieck. Thesis, Grenoble 1977.
- [11] Michel Las Vergnas, On Eulerian partitions of graphs. Research Notes in Mathematics 34, 62–75 (1979).
- [12] Michel Las Vergnas, On the evaluation at (3,3) of the Tutte polynomial of a graph. *Journal of Combinatorial Theory, Series B* 44, 367–372 (1988).
- [13] Dirk Vertigan, The Computational Complexity of Tutte Invariants for Planar Graphs. SIAM J. Comput., 35(3) 690–712 (2006).
- [14] Hans Wenzl, On the Structure of Brauer's Centralizer Algebras. *Annals of Mathematics* 128(1) 173–193 (1988).
- [15] Gian-Carlo Wick, The evaluation of the collision matrix. *Physical Review* 80(2): 268–272 (1950).