Confidence Statements for Ordering Quantiles

Carlos A. de B. Pereira Institute of Mathematics and Statistics University of São Paulo, São Paulo, Brazil cpereira@ime.usp.br

Cassio P. de Campos
Dalle Molle Institute for Artificial Intelligence
Manno, Switzerland
cassio@idsia.ch

Adriano Polpo Federal University of São Carlos São Carlos, Brazil polpo@ufscar.br

June, 2014

Abstract

This work proposes *Quor*, a simple yet effective nonparametric method to compare independent samples with respect to corresponding quantiles of their populations. The method is solely based on the order statistics of the samples, and independence is its only requirement. All computations are performed using exact distributions with no need for any asymptotic considerations, and yet can be run using a fast quadratic-time dynamic programming idea. Computational performance is essential in high-dimensional domains, such as gene expression data. We describe the approach and discuss on the most important assumptions, building a parallel with assumptions and properties of widely used techniques for the same problem. Experiments using real data from biomedical studies are performed to empirically compare Quor and other methods in a classification task over a selection of high-dimensional data sets.

1 Introduction

A common procedure in Statistics and Machine Learning when dealing with data sets of thousands of variables is to *sort* all these variables according to some measure that identifies how important they are to predict and/or retrospectively understand a certain target variable (or equivalently an indicator that tells in which group or *population* belongs each sample). Classical examples of such a procedure are the Student's *t*-test and the Wilcoxon's rank-sum *u*-test [7, 8, 15], whose statistics are often used to sort variables into some order of importance. Arguably, they represent the most commonly used methods for this problem in biomedical applications, in part because of their prompt availability and easiness of use. A typical scenario is to have gene expression data of cancer patients, and a *class* variable that identifies whether the patient *relapsed* or not (in other word, whether the cancer came back after treatment/surgery or not). The ability to sort variables in some meaningful order has a range of applications in many fields, and can also be seen as means of performing *feature selection* [16, 26].

This work describes a simple yet effective approach, named *Quor*, to sort variables according to the order relationship of arbitrary quantiles of the variable's distributions under different groups. The method computes a value that indicates the confidence that such quantiles of these distributions are sorted in some pre-defined way. As example, suppose we have two populations and are interested in the median values of a variable representing the level of expression of a particular gene. The goal is to obtain the confidence that "the median expression of that gene in the first population is strictly smaller (or greater) than the median expression of the same gene in the second population". The comparison of medians might suggest that

the gene is under-expressed, over-expressed, or simply that there is no significant difference of expression between the populations. As other methods with similar purpose, Quor can be used as a first aid for the later application of other sophisticated statistical or biological procedures: Its simplicity may avoids expensive and time-consuming analyses of uninteresting variables, or at least may help to prioritize further analyses in order to save valuable time and biological materials/machinery.

Methods for this problem may rely on eventually unrealistic hypotheses, such as normality of samples, asymptotic behavior of some statistics, reasonably large-sized samples, approximate computations, comparisons of only two populations, need of equivalent number of samples in the groups (across distinct variables), necessity of multiple-test corrections (to avoid encountering significant results *by chance*), no evidence for the null hypothesis, among others [18]. These issues might be aggravated by having data sets with only a few patients and a large number of genes. On the other hand, Quor is nonparametric and assumes nothing but independence of samples. It can deal with different number of samples and missing data, and yet can properly compare these variables; all computations are performed exactly, without any asymptotic assumption. Moreover, its computations can be carried on in quadratic time in the number of samples, which is (roughly) as fast as other methods based on *t*-tests and *u*-tests. Other approaches for median test do exist (we refer to Chapter 6 of [9] for some examples), but they either fail in at least one of the issues mentioned previously in this paragraph, or are not able to order arbitrary quantiles of (arbitrary) many populations.

This paper is divided as follows. Section 2 describes the approach in details and presents its computational complexity. Sections 3 and 4 analyze real biomedical data sets and compare the empirical performance of the methods. Section 5 concludes the paper and discuss on future work.

2 Unveiling Quor

We describe here the details of Quor and present an efficient algorithm for its computation. The method is built on the ideas of confidence statements developed long ago [3, 13] and revisited more recently [27]. The proposed method uses nonparametric confidence intervals for quantiles based on the binomial distribution [6]. Its goal is to compute a confidence value indicating how much one believes that quantile parameters of different populations/groups are ordered among themselves. We do not assume any particular quantile nor a specific number of populations, even though the case of comparing medians of two populations is arguably the most common scenario for its application.

The problem is defined as follows. Let Q_1,\ldots,Q_n represent the quantiles at arbitrary percentages q_1,\ldots,q_n , respectively, for n populations, and let $\mathbf{x}_i=(x_i^{(1)},\ldots,x_i^{(m_i)})$, for $i=1,\ldots,n$, be data samples from those populations, where the sample from population i has size m_i . The goal is to produce a confidence value in [0,1] that indicates how much we believe in the statement $Q_{i_1}< Q_{i_2}<\ldots< Q_{i_n}$, where (i_1,\ldots,i_n) is a permutation of $(1,\ldots,n)$. As in other nonparametric methods [24], the order and the values of the tuples of order statistics will be the solely observations needed.

Consider an event A related to a random variable X_i whose quantile at q_i is Q_i . $\Pr(A \mid Q_i)$ indicates the probability of the event A with Q_i known. The quantile Q_i of X_i is a population parameter that satisfies the following inequality:

$$\Pr(\{X_i \le Q_i\} \mid Q_i) \ge q_i,\tag{1}$$

while in the continuous case, this inequality is tight.

Let $(X_i^{(1)}, X_i^{(2)}, \dots, X_i^{(m_i)})$ be the ordered vector of m_i independent and identically distributed continuous random variables. Since the probability of one observation being smaller than the population quantile Q_i is q_i , it is straightforward that for the jth $(j=1,2,\dots,m_i)$ order statistics, $X_i^{(j)}$, the following probability expression holds:

$$\Pr(\{X_i^{(j)} \le Q_i\} \mid Q_i) = \sum_{k=j}^{m_i} {m_i \choose k} q_i^k (1 - q_i)^{m_i - k}, \tag{2}$$

and symmetrically

$$\Pr(\{X_i^{(j)} \ge Q_i\} \mid Q_i) = \sum_{k=0}^{j-1} {m_i \choose k} q_i^k (1 - q_i)^{m_i - k}.$$
(3)

These equalities come from probabilities obtained with a binomial distribution with sample size m_i and probability of success q_i . Consider a sequence of pairs of order statistics $((X_1^{(j_1)}, X_1^{(j_1')}), (X_2^{(j_2)}; X_2^{(j_2')}), \ldots, (X_n^{(j_n)}; X_n^{(j_n')}))$, each of them chosen from the ith group (with abuse of notation, let first and last groups have $X_1^{(j_1)} = -\infty$ and $X_n^{(j_n')} = \infty$, respectively), and consider the event E as follows.

$$\mathbf{E} = \bigcap_{i=1}^{n} \{X_{i}^{(j_{i})} \leq Q_{i} \leq X_{i}^{(j'_{i})}\} = \bigcap_{i=1}^{n} \left(\{X_{i}^{(j'_{i})} \geq Q_{i}\} \setminus \{X_{i}^{(j_{i})} > Q_{i}\} \right).$$

Given the independence among the samples, one can compute $Pr(E \mid Q_1, \dots, Q_n) =$

$$\prod_{i=1}^{n} \max \left\{ 0; \left[\Pr(\{X_i^{(j_i')} \ge Q_i\} \mid Q_i) - \left(1 - \Pr(\{X_i^{(j_i)} \le Q_i\} \mid Q_i)\right) \right] \right\},\tag{4}$$

using the product of binomial probabilities from Equations (2) and (3).

Let \mathbf{x}_i be sorted, for every i. After these samples are observed, the only unknown quantities of interest are the quantiles Q_i of the populations being studied. By replacing random variables with their observations, we create the statement \mathbf{e} :

$$e = \bigcap_{i=1}^{n} \{x_i^{(j_i)} \le Q_i \le x_i^{(j_i')}\},\tag{5}$$

which has confidence given by Expression (4). Note that we only need the orders j_i and j_i' for every i, and not the actual observed values of X_i , for computing with Expression (4), that is, the confidence of e could be equivalently defined as the confidence of $((0;j_1');(j_2;j_2');\ldots;(j_n;1+m_n))$, where we conveniently define that, for every $i=1,\ldots,n,$ $x_i^{(j)}=-\infty$ for every j<1 and $x_i^{(j)}=\infty$ for every $j>m_i$. Hence, we might say that e is represented by the list $((0;j_1');(j_2;j_2');\ldots;(j_n;1+m_n))$. At this point after sampling, we call the value of Expression (4) a confidence value instead of probability value, in order to keep terminology precise. This confidence regards the unknown quantities of interest, in our case the parameters Q_i . The idea is to look for statements e that are able to tell us something about Q_i . Without loss of generality, assume that we take a list of pairs of order statistics $((0;j_1');(j_2;j_2');\ldots;(j_n;1+m_n))$ such that, in the observed sets, we have $x_i^{(j_i)} < x_{i+1}^{(j_{i'})} < x_{i+1}^{(j_{i+1})}$, for every $1 \le i < n$ (we do not lose generality because in case we want to sort these observations in some other order, we simply rename the variables). With this fact and a quick analysis of Expression (5), we derive

$$\left(\forall_{1 \le i < n} : x_i^{(j_i)} < x_i^{(j_i')} < x_{i+1}^{(j_{i+1})}\right) \land \mathbf{e} \Rightarrow \bigcap_{i=1}^{n-1} \{Q_i < Q_{i+1}\},\tag{6}$$

that is, the assertion in the left-hand side of Expression (6) implies an order for the quantiles, so its confidence is a lower bound for the confidence of the right-hand side. Because we know how to compute the confidence value of e through Expression (4), and any time the assertion $x_i^{(j_i)} < x_i^{(j_i')} < x_{i+1}^{(j_{i+1})}$ is false for any i the result of Expression (4) becomes zero, we have the following relation.

$$\operatorname{Conf}(\mathbf{e}) \le \operatorname{Conf}\left(\bigcap_{i=1}^{n-1} \{Q_i < Q_{i+1}\}\right). \tag{7}$$

In order to compute the best possible lower bound for the confidence value in the right-hand side of Expression (7), we run over all tuples $((0; j'_1); (j_2; j'_2); \ldots; (j_n; 1 + m_n))$ of orders that comply with the linear order

$$x_1^{(0)} < x_1^{(j_1')} < x_2^{(j_2)} < x_2^{(j_2')} < \ldots < x_n^{(j_n)} < x_n^{(1+m_n)}$$

and keep the maximum confidence value of the e statements built from these tuples as our estimation for the confidence that $Q_1 < Q_2 < \ldots < Q_n$ holds true. Because we want to maximize such confidence, we will always choose j_i such that $x_i^{(j_i)}$ is the smallest possible value greater than $x_{i-1}^{(j'_{i-1})}$, that is, the value of j_i , in order to maximize the confidence, is uniquely computable from the value of j'_{i-1} (that is because there is no reason to leave a larger gap between them if a smaller gap is possible, as smaller gaps will yield higher confidence values). Hence, we can ease on the terminology and compute the confidence of e by $\operatorname{Conf}(j'_1,\ldots,j'_{n-1})$, as these values are enough to find all j_2,\ldots,j_n (that lead to the highest possible confidence) and then to use Expression (4) to obtain the confidence value. For easy of computation, we define the confidence using a sum of logarithms.

$$\operatorname{Conf}_{i}(j'_{i-1}, j'_{i}) = \log \left(\max \left\{ 0; \Pr(\{X_{i}^{(j'_{i})} \ge Q_{i}\} \mid Q_{i}) + \Pr(\{X_{i}^{(j_{i})} \le Q_{i}\} \mid Q_{i}) - 1 \right\} \right), \tag{8}$$

where j_i is obtained from j'_{i-1} (by looking the data) as just explained, and $\log 0$ is $-\infty$. Note that the value of $\mathrm{Conf}_i(j'_{i-1},j'_i)$ will equal to $-\infty$ whenever the values j'_{i-1},j'_i do not induce a "valid" order in the observations, that is, whenever there is no element $x_i^{(j_i)}$ strictly inside $[x_{i-1}^{(j'_{i-1})},x_i^{(j'_i)}]$. We are interested in $\exp(\sum_{i=1}^n\mathrm{Conf}_i(j'_{i-1},j'_i))$, which is our statistical conclusion based on the observed samples that the quantiles (of the populations) are ordered according to the permutation $(1,\ldots,n)$. This procedure is presented in Algorithm 1, which is explained in the continuation. We recall that if one wants to compute the confidence of some other order, for instance $Q_{i_1} < Q_{i_2} < \ldots < Q_{i_n}$, where (i_1,\ldots,i_n) is a permutation of $(1,\ldots,n)$, then they simply need to rename the variables accordingly before invoking the algorithm, while if one wants to check for every possible order of the quantiles, there would be n! permutations to check, which is fast for small values of n (the vast majority of biomedical studies has $n \leq 6$ groups).

Algorithm 1 (Quor Core) Finding the confidence value of a statement about the ordering of a quantile parameter among n populations.

Input a data set with samples $\mathbf{x}_i = (x_i^{(1)}, \dots, x_i^{(m_i)})$, for $i = 1, \dots, n$, and the quantiles of interest q_i , for each i

Output the log-confidence value that the statement $Q_1 < Q_2 < \ldots < Q_n$ holds.

- 1 For every i in $1, \ldots, n$, sort \mathbf{x}_i .
- 2 Pre-compute the values that appear in Equations (2) and (3) by making a cache to be used in the computation of Equation (8), for every $i=1,\ldots,n$: cache $(i,0)\leftarrow(1-q_i)^{m_i}$ and for $j=1,\ldots,m_i$:

$$\mathrm{cache}(i,j) \leftarrow \mathrm{cache}(i,j-1) + \binom{m_i}{j} q_i^j (1-q_i)^{m_i-j}.$$

- 3 Let D_i be a vector of size m_i (defined from 1 to m_i), for each $i=1,\ldots,n-1$. Initialize $D_1[\ell'_1] \leftarrow \operatorname{Conf}_1(0,\ell'_1)$, for every $1 \leq \ell'_1 \leq m_1$. If n=2, then go to line 5.
- 4 For i = 2, ..., n 1, do:

For $\ell'_i = 1, \ldots, m_i$, do:

$$D_i[\ell_i'] = \max_{1 \le \ell_{i-1}' \le m_{i-1}} (D_{i-1}[\ell_{i-1}'] + \mathsf{Conf}_i(\ell_{i-1}', \ell_i')).$$

5 Return $\max_{1 \le \ell'_{n-1} \le m_{n-1}} (D_{n-1}[\ell'_{n-1}] + \operatorname{Conf}_n(\ell'_{n-1}, 1 + m_n)).$

Theorem 1 Algorithm 1 uses space O(n+m) and takes time $O(m \log m)$ if n=2, and $O(m \max_i m_i)$ otherwise.

Proof. Step 1 needs to sort each of the samples x_i , which takes $O(m_i \log m_i)$ for each i, so in total $O(m \log \max_i m_i)$. Step 2 pre-computes the partial binomial sums. By doing it in a proper order, this

can be accomplished in constant time for each $\operatorname{cache}(i,j)$, and the loop will execute $O(\sum_i m_i) = O(m)$ times. Step 3 initializes the data structures D_i for the dynamic programming. Summed altogether, they spend O(m) space and O(m) time to initialize. Step 4 performs the dynamic programming. The number of times both loops are executed altogether is O(m). The internal maximization takes less than $O(\max_i m_i)$, so this step has time $O(m \max_i m_i)$ in the worst case (and it is not run for n=2). Finally, Step 5 takes time O(m).

Algorithm 1 is very fast. First, many common practical cases have n=2. In this case, the Step 4 of the algorithm is skipped and the whole algorithm runs in linear time (except for the sorting of Step 1). Second, this worst-case time complexity is pessimistic, and the algorithm will usually run in sub-quadratic time. Third, m tends to be a reasonable small number (for example, biomedical data sets hardly contain more than one hundred patients).

The correctness of Algorithm 1 comes from its simple dynamic programming formulation. At each main loop i of Step 4, we have already solved the problem up to i-1 groups in the best possible way, for each one of the positions one could choose for the last placeholder ℓ'_{i-1} , which we saved in $D_{i-1}[\ell'_{i-1}]$. The choices of the yet-to-decide values ℓ'_i, \ldots, ℓ'_n do not depend on the positions of the placeholders before ℓ'_{i-1} , only on ℓ'_{i-1} itself. Thus, the dynamic programming does the job of increasingly building new solutions. This step is inspired by other dynamic programming algorithms, such as those for k-means in one dimensional vectors [23], yet different because of the nature of our confidence function to be optimized. It is worth noting that some computations in Algorithm 1 may suffer from numerical problems. We have implemented it using incomplete Beta functions and arbitrary precision when needed; this is usually the case if m is reasonably large (greater than some hundreds).

The confidence value obtained with Quor can be used in procedures that want to optimize their utility functions, because they can provide information about differences in quantiles as well as similarities in quantiles (in the case that confidences are low in both directions). This is in contrast to usual hypothesis testing, where no evidence in favor of the null hypothesis can be obtained. A possible negative characteristic of Quor is that its computations may often lead to ties in the results, which is an intrinsic situation of exact computations with the binomial distribution. However, if computations are carried on with a clever implementation that avoids numerical issues, then ties should arguably be considered as effective ties, and other procedures/ideas could be devised to break the ties. For our current needs, ties have not constituted a problem, mainly because we are usually interested in the top confidence values, where the ties seem to be less often.

3 Schizophrenia data

We show a practical scenario by analyzing the Schizophrenia data set from the Stanley Medical Research Institute's online genomics database [12, 21]. The sample sizes are $m_1=34$ individuals in the control group and $m_2=34$ patients with schizophrenia. A total of 20992 microarray probes were obtained. The usual goal in such an analysis is to find the most differentially expressed genes. For that purpose, we decided to evaluate the confidence of the statements $\{Q_1 < Q_2\}$ and $\{Q_1 > Q_2\}$, where $q_1 = q_2 = 1/2$, that is, we compared the medians of the populations. This is performed for each gene in the data set. We argue that the difference in the populations' medians indicate that genes might be differentially expressed (at least it shows that the distributions are not equal). Not surprisingly, there are **no** significant genes if we perform either t-test or u-test with multiple-test correction (we have used the Holm-Bonferroni correction). On the other hand, Quor measure has a clear interpretation as the confidence of the difference in medians, so if one chooses genes with, for instance, confidence above 95%, then there are 56 genes in this situation, of which 14 cases suggest that ill patients (group 2) have under-expressed genes ($\{Q_1 > Q_2\}$) and 42 cases suggest over-expressed genes ($\{Q_1 < Q_2\}$).

Because of its characteristic, one might see a single Quor confidence as a more conservative approach than other tests, as it has fewer assumptions (for instance, all but one high-confidence genes from Quor also had u-test pvalue < 5%), but this vanishes if multiple test correction is performed. Table 1 shows the

Table 1: Number of rejected and non-rejected genes among those with Quor confidence $\geq 95\%$ in the Schizophrenia data. First line is performed without using multiple test correction for t and u tests. Second does correction assuming that only 56 hypothesis tests are executed. If one corrects for all the genes, no significant gene is found by t or u tests.

	t-test		<i>u</i> -test	
	Non-Reject	Reject	Non-Reject	Reject
no correction	18	38	1	55
with correction	48	8	43	13

number of null hypothesis (that is, no difference) rejections with t-test and u-test assuming that only 56 tests were performed (exactly for those genes with high Quor confidence). This illustrates a hypothetical procedure where Quor is firstly executed, followed by the hypothesis testing (we decided to show this view otherwise those tests would not be able to reject the null hypothesis for any gene). In the table we see that t and u tests turn out to be very conservative because of multiple test correction, and the prior use of Quor can alleviate this situation and consequently reduce an excessive amount of Type II errors.

4 Evaluation as Feature Selection

In this section we apply different hypothesis testing methods in the literature and Quor to the task of feature selection, with the ultimate goal of building a classifer with a subset of all variables. We do not compare these approaches with yet other methods for feature selection, for example those which consider correlations between variables and information measures, because they have very different goals, while ours is to compare methods with similar purpose, such as *t*-test, *u*-test, *ks*-test, and Quor itself. For that purpose, we try to predict a yet unseen class/group given the observations of the model covariates. In each data set, the class has a particular meaning, and the number of samples and variables vary. Table 2 shows the main characteristics of the data sets with which we work. Please refer to the corresponding citation for additional information on the data [1, 2, 10, 12, 17, 20, 22, 25]. These data were obtained from internet repositories [14, 19, 21].

Table 2: Data set characteristics. m is the number of patients, shown by the amount in each group. PS stands for $Proteomic\ spectra$, GEP for $Gene\ Expression\ Profiling$.

Data set	m	# Feat.	Type	Class
Prostate C.	8+13	12600	GEP	Relapse/Not
Schizophr.	34+32	20992	GEP	Disease/Not
Lung C.	24+15	2880	GEP	Relapse/Not
Breast C.	12+7	24481	GEP	Relapse/Not
Colon C.	22+40	2000	GEP	Disease/Not
Lymphoma	22+23	4026	GEP	Dis. Subtype
Leukemia	27+11	7129	GEP	Dis. Subtype
Ovarian C.	162+91	15154	PS	Disease/Not

The idea is to fix the classifier while varying the feature selection approach among Quor (median as the chosen quantile), t-test, u-test and ks-test (Kolmogorov-Smirnov test [4]). To avoid a result that is specific to a single classifier, we perform the experiment with two very distinct classifiers: a Bayesian network and a C4.5 decision tree (both inferred from data), implemented in the Weka package [11]. For each data set, we run a five-fold cross-validation procedure repeated 20 times (so as the test set is never available during

training; five folds are chosen because some of the data sets contain very few patients). Results are shown in Tables 3 and 4 for the Bayesian network and the decision tree, respectively. Numbers represent average accuracy over $5 \cdot 20 = 100$ runs. Quor, as a feature selection procedure, has different characteristics from the t-test, and each of them seems to perform better in distinct data sets; on the other hand, when compared to the u-test, Quor seems to show a mild improvement in accuracy. This might be explained by the greater proximity of ideas, in some sense, of the u-test and Quor, with the latter demonstrating a slight better performance in these specific data sets.

Table 3: Average accuracy (over 100 runs) of a Bayesian network classifier using the 20 best ranked features for each selection method.

Data set	Quor	t-test	u-test	ks-test
Prostate C.	47.85	43.55	42.45	42.60
Schizophrenia	59.08	57.99	58.98	60.09
Lung C.	66.39	65.54	65.45	66.52
Breast C.	69.50	74.75	64.08	66.67
Colon C.	82.87	76.60	83.37	81.84
Lymphoma	91.33	92.44	92.33	91.67
Leukemia	96.23	91.54	96.07	92.48
Ovarian C.	97.41	98.00	97.23	97.35
Average	76.02	75.52	75.56	75.55

Table 4: Average accuracy (over 100 runs) of a C4.5 decision tree classifier using the 20 best ranked features for each selection method.

Data set	Quor	t-test	u-test	ks-test
Prostate C.	40.45	40.70	39.55	39.90
Schizophrenia	55.91	55.24	56.81	56.87
Lung C.	60.02	59.07	63.09	64.27
Breast C.	50.67	59.50	51.67	58.33
Colon C.	83.65	77.40	80.57	81.01
Lymphoma	76.89	77.44	77.11	77.44
Leukemia	87.66	84.70	86.55	86.11
Ovarian C.	96.68	97.21	96.66	96.58
Average	68.27	68.23	68.74	69.76

5 Conclusions

This study proposes a method called Quor to compute the confidence of a statement about the order of arbitrary quantiles of many populations. Its only assumption is independence of samples, which makes it applicable to many domains. Among Quor properties, we highlight its nonparametric nature, the possibility of processing data sets with missing data and varying number of samples from each population (across multiple tests), the confidence interpretation that might avoid the need of multiple-test correction, and its efficient exact computations. Quor will be made available under an open source license both as an easy-to-install R package and as a plugin for the Weka Data Mining package [11]. Besides confidences on the order of quantiles of the populations, the package also computes confidence intervals based on order statistics (we refrained from detailing on them here; see Chapter 7 of [5]).

Our empirical analysis has indicated that Quor results are in line with the most used techniques for hypothesis testing, which are also often used to order genes according to their statistics/p-values. Quor also performs well as a feature selection procedure (when compared to hypothesis testing methods) in a benchmark of high-dimensional data sets. As future work, we will study the occurrence of ties in the results of Quor, and means to break such ties. While they have not been a major issue so far in our analyses, we consider important to have a sound procedure to resolve them, especially to deal with discrete ordinal data. We are also pursuing applications of the method in recent data sets from the latest microarray technologies. As the number of covariates increases as Quor shall become important for its computational performance and minimal assumptions.

Acknowledgements

This work was partially supported by the Brazilian agencies FAPESP, CNPq and CAPES.

References

- [1] A. A. Alizadeh, M. B. Eisen, R. E. Davis, C. Ma, I. S. Lossos, A. Rosenwald, J. C. Boldrick, H. Sabet, T. Tran, X. Yu, J. I. Powell, Liming Yang, Gerald E. Marti, Troy Moore, James Hudson Jr, Lisheng Lu, David B. Lewis, Robert Tibshirani, Gavin Sherlock, Wing C. Chan, Timothy C. Greiner, Dennis D. Weisenburger, James O. Armitage, Roger Warnke, Ronald Levy, Wyndham Wilson, Michael R. Grever, John C. Byrd, David Botstein, Patrick O. Brown, and Louis M. Staudt. Distinct types of diffuse large B-cell lymphoma identified by gene expression profiling. *Nature*, 403(3):503–511, 2000.
- [2] U. Alon, N. Barkai, D. A. Notterman, K. Gish, S. Ybarra, D. Mack, and A. J. Levine. Broad Patterns of Gene Expression Revealed by Clustering Analysis of Tumor and Normal Colon Tissues Probed by Oligonucleotide Arrays. *PNAS*, 96:6745–6750, 1999.
- [3] D. Basu. *On ancillary statistics, pivotal quantities and confidence statements*, pages 1–29. Concordia University Publication, 1981.
- [4] I. M. Chakravarti, R. G. Laha, and J. Roy. *Handbook of Methods of Applied Statistics*, volume I. John Wiley and Sons, 1967.
- [5] H. A. David and H. N. Nagaraja. Order Statistics. Wiley-Interscience, 2003. 3rd edition.
- [6] M. DeGroot. Probability and Statistics. Addison-Wesley, New York, 1975. 2nd edition.
- [7] J. Demšar. Statistical comparisons of classifiers over multiple data sets. *Journal of Machine Learning Research*, 7:1–30, 2006.
- [8] M. P. Fay and M. A. Proschan. Wilcoxon-Mann-Whitney or t-test? On assumptions for hypothesis tests and multiple interpretations of decision rules. *Statistics Surveys*, 4:1–39, 2010.
- [9] J. K. Gibbons and S. Chakraborti. *Nonparametric Statistical Inference*. Marcel Dekker Inc., 4th edition, 2003.
- [10] T. R. Golub, D. K. Slonim, P. Tamayo, C. Huard, M. Gaasenbeek, J. P. Mesirov, H. Coller, M. L. Loh, J. R. Downing, M. A. Caligiuri, C. D. Bloomfield, and E. S. Lander. Molecular Classification of Cancer: Class Discovery and Class Prediction by Gene Expression Monitoring. *Science*, 286:531–537, 1999.
- [11] Mark Hall, Eibe Frank, Geoffrey Holmes, Bernhard Pfahringer, Peter Reutemann, and Ian H. Witten. The weka data mining software: An update. *SIGKDD Explorations*, 11(1), 2009.

- [12] B. Higgs, M. Elashoff, S. Richman, and B. Barci. An online database for brain disease research. BMC Genomics, 7:70, 2006.
- [13] J. Kiefer. Conditional confidence statements and confidence estimators. *Journal of American Statistical Association*, 72(360):789–808, 1977.
- [14] Jinyan Li, Huiqing Liu, and Limsoon Wong. Mean-entropy discretized features are effective for classifying high-dimensional biomedical data. In *3rd ACM SIGKDD Workshop on Data Mining in Bioinformatics*, pages 17–24, 2003. Kent Ridge Bio-medical Dataset.
- [15] H. B. Mann and D. R. Whitney. On a test of whether one of two random variables is stochastically larger than the other. *Annals of Mathematical Statistics*, 18(1):50–60, 1947.
- [16] T. M. Mitchell. *Machine Learning*. McGraw-Hill, 1997.
- [17] Emanuel F. Petricoin III, Ali M Ardekani, Ben A. Hitt, Peter J. Levine, Vincent A. Fusaro, Seth M. Steinberg, Gordon B. Mills, Charles Simone, David A. Fishman, Elise C. Kohn, and Lance A. Liotta. Use of proteomic patterns in serum to identify ovarian cancer. *The Lancet*, 359, 2002.
- [18] A. E Raftery. Bayesian model selection in social research. *Sociological methodology*, 25:111–164, 1995.
- [19] BioInformatics Group Seville. Bioinformatics research group web page, 2014. http://www.upo.es/eps/bigs, Accessed on Jan/2014.
- [20] Dinesh Singh, Phillip G. Febbo, Kenneth Ross, Donald G. Jackson, Judith Manola, Christine Ladd, Pablo Tamayo, Andrew A. Renshaw, Anthony V. D'Amico, Jerome P. Richie, Eric S. Lander, Massimo Loda, Philip W. Kantoff, Todd R. Golub, and William R. Sellers. Gene expression correlates of clinical prostate cancer behavior. *Cancer Cell*, 1(2):203–209, 2002.
- [21] The Stanley Medical Research Institute. Stanley Medical Research Institute online genomics database: Altar A, 2012. Available at http://www.stanleygenomics.org; accessed on Jan/2014.
- [22] L. J. van't Veer, H. Dai, M. J. van de Vijver, Y. D. He, A. A. Hart, M. Mao, H. L. Peterse, K. van der Kooy, M. J. Marton, A. T. Witteveen, G. J. Schreiber, R. M. Kerkhoven, C. Roberts, P. S. Linsley, R. Bernards, and S. H. Friend. Gene expression profiling predicts clinical outcome of breast cancer. *Nature*, 415(6871):530–536, 2002.
- [23] Haizhou Wang and Mingzhou Song. Ckmeans.1d.dp: Optimal k-means Clustering in One Dimension by Dynamic Programming. *The R Journal*, 3/2:29–33, 2011.
- [24] L. Wasserman. All of Nonparametric Statistics. Springer, New York, 2006.
- [25] Dennis A. Wigle, Igor Jurisica, Niki Radulovich, Melania Pintilie, Janet Rossant, Ni Liu, Chao Lu, James Woodgett, Isolde Seiden, Michael Johnston, Shaf Keshavjee, Gail Darling, Timothy Winton, Bobby-Joe Breitkreutz, Paul Jorgenson, Mike Tyers, Frances A. Shepherd, and Ming Sound Tsao. Molecular Profiling of Non-Small Cell Lung Cancer and Correlation with Disease-free Survival. Cancer Research, 62:3005–3008, 2002.
- [26] Ian Witten, Eibe Frank, and Mark Hall. *Data Mining: Practical Machine Learning Tools and Techniques*. Morgan Kaufmann, 2011.
- [27] A. Zellner, H. Keuzenkamp, and M. McAleer. Simplicity, Inference and Modeling: Keeping it Sophisticatedly Simple. Cambridge University Press, 2004.




Figure 1: Data of four transcripts from the Schizophrenia data set. In (a), Quor Conf=0.43, but t and u reject the null hypothesis at 0.002 and 0.02 (that is, they consider the difference between groups to be quite significant). In (b), t cannot reject the null hypothesis, and in (c) neither t nor u can (in both cases Quor Conf is at least 0.95). In (d), both tests and Quor strongly agree (Quor Conf=0.99, t-test pvalue=8e-4, u-test pvalue=2e-4).