# NOETHERIANITY FOR INFINITE-DIMENSIONAL TORIC VARIETIES

JAN DRAISMA, ROB H. EGGERMONT, ROBERT KRONE, AND ANTON LEYKIN

ABSTRACT. We consider a large class of monomial maps respecting an action of the infinite symmetric group, and prove that the toric ideals arising as their kernels are finitely generated up to symmetry. Our class includes many important examples where Noetherianity was recently proved or conjectured. In particular, our results imply Hillar-Sullivant's Independent Set Theorem and settle several finiteness conjectures due to Aschenbrenner, Martin del Campo, Hillar, and Sullivant.

We introduce a *matching monoid* and show that its monoid ring is Noetherian up to symmetry. Our approach is then to factorize a more general equivariant monomial map into two parts going through this monoid. The kernels of both parts are finitely generated up to symmetry: recent work by Yamaguchi-Ogawa-Takemura on the (generalized) Birkhoff model provides an explicit degree bound for the kernel of the first part, while for the second part the finiteness follows from the Noetherianity of the matching monoid ring.

### 1. INTRODUCTION AND MAIN RESULT

Families of algebraic varieties parameterized by combinatorial data arise in various areas of mathematics, such as statistics (e.g., phylogenetic models parameterized by trees [AR08, DK09, DE15, PS05] or the relations among path probabilities in Markov chains parameterized by path length [HdTY14, Nor15]), commutative algebra (e.g., Segre powers of a fixed vector space parameterized by the exponent [Sno13] or Laurent lattice ideals [HdC13]), and combinatorics (e.g., algebraic matroids arising from determinantal ideals parameterized by matrix sizes [KR13] or edge ideals of hypergraphs parameterized by the number of vertices [GP13]). A natural question is whether such families *stabilize* as some of the combinatorial data tend to infinity. A recently established technique for proving such stabilization is passing to an infinite-dimensional limit of the family, giving some equations for that limit, and showing that those equations cut out a suitably Noetherian space. This then implies that the limit itself is given by finitely many further equations, and that the family stabilizes. This technique is applied, for instance, in the proof of the Independent Set Theorem [HS12], and in the first author's work on the Gaussian k-factor model, chirality varieties, and tensors of bounded rank [Dra10, DK14].

In the present paper, we follow a similar approach, utilizing the new concept of a *matching monoid* to prove that stabilization happens for a large class of toric varieties. Our Main Theorem provides one-step proofs for several existing results that were established in a rather less general context; and it settles conjectures and questions from [AH07, HS12, HdC13]. There is a list of three such consequences

The first and second author are supported by a Vidi grant from the Netherlands Organisation for Scientific Research (NWO).

The third and fourth author are supported in part by the NSF under grant DMS 1151297.

at the end of this Introduction. Moreover, we show Noetherianity in a constructive manner by complementing the Main Theorem with an algorithm that produces a finite set of equations whose orbits define the infinite-dimensional toric variety under consideration.

Instead of working with inverse systems of affine varieties, we work directly with direct limits of their coordinate rings. In fact, we formulate our Main Theorem directly in the infinite-dimensional setting, as going back to families of finitedimensional coordinate rings of toric varieties is fairly straightforward. Throughout,  $\mathbb{N}$  denotes  $\{0, 1, 2, 3, \ldots\}$ , and for  $k \in \mathbb{N}$  we write  $[k] := \{0, \ldots, k-1\}$ . We write  $\operatorname{Sym}(\mathbb{N})$  for the group of all bijections  $\mathbb{N} \to \mathbb{N}$ , and  $\operatorname{Inc}(\mathbb{N})$  for the monoid of all strictly increasing maps  $\mathbb{N} \to \mathbb{N}$ . Let Y be a set equipped with an action of Sym( $\mathbb{N}$ ). We require that the action has the following property: for each  $y \in Y$  there exists a  $k_y \in \mathbb{N}$  such that y is fixed by all of  $\operatorname{Sym}(\mathbb{N} \setminus [k_y])$ , i.e., by all elements of  $Sym(\mathbb{N})$  that fix  $[k_y]$  element-wise. In this setting,  $Inc(\mathbb{N})$  also acts on Y, as follows: for  $\pi \in \text{Inc}(\mathbb{N})$  and  $y \in Y$ , choose a  $\pi' \in \text{Sym}(\mathbb{N})$  that agrees with  $\pi$  on  $[k_y]$ , set  $\pi y := \pi' y$ , and observe that this does not depend on the choice of  $\pi'$ . Observe that for each  $y \in Y$  the  $\operatorname{Inc}(\mathbb{N})$ -orbit  $\operatorname{Inc}(\mathbb{N})y$  is contained in  $\operatorname{Sym}(\mathbb{N})y$ , and that the latter is in fact equal to the orbit of y under the countable subgroup of  $Sym(\mathbb{N})$ consisting of permutations fixing all but finitely many natural numbers. See also [HS12, Section 5].

Let R be a Noetherian ring (commutative, with 1), and let R[Y] be the commutative R-algebra of polynomials in which the elements of Y are the variables and the coefficients come from R. The group  $Sym(\mathbb{N})$  acts by R-algebra automorphisms on R[Y] by permuting the variables. Furthermore, let k be a natural number, and let  $Z = \{z_{ij} \mid i \in [k], j \in \mathbb{N}\}$  be a second set of variables, with a  $Sym(\mathbb{N})$ -action given by  $\pi z_{ij} = z_{i\pi(j)}$ . Extend this action to an action by R-algebra automorphisms of R[Z]. Note that the  $Sym(\mathbb{N})$ -actions on R[Y], Z, R[Z] all have the property required of the action on Y. Hence they also yield  $Inc(\mathbb{N})$ -actions, by means of injective Ralgebra endomorphisms in the case of R[Y] and R[Z]. In general, when a monoid II acts on a ring S by means of endomorphisms, S is called  $\Pi$ -Noetherian if every II-stable ideal in S is generated by the union of finitely many  $\Pi$ -orbits of elements, i.e., if S is Noetherian as a module under the skew monoid ring  $S * \Pi$ ; see [HS12].

**Theorem 1.1** (Main Theorem). Assume that  $\operatorname{Sym}(\mathbb{N})$  has only finitely many orbits on Y. Let  $\varphi : R[Y] \to R[Z]$  be a  $\operatorname{Sym}(\mathbb{N})$ -equivariant homomorphism that maps each  $y \in Y$  to a monomial in the  $z_{ij}$ . Then ker  $\varphi$  is generated by finitely many  $\operatorname{Inc}(\mathbb{N})$ -orbits of binomials, and im  $\varphi \cong R[Y]/\ker \varphi$  is an  $\operatorname{Inc}(\mathbb{N})$ -Noetherian ring.

If an ideal is  $\operatorname{Sym}(\mathbb{N})$ -stable, then it is certainly  $\operatorname{Inc}(\mathbb{N})$ -stable, so the last statement implies that  $R[Y]/\ker \varphi$  is  $\operatorname{Sym}(\mathbb{N})$ -Noetherian. The conditions in the theorem are sharp in the following senses.

(1) The ring R[Y] itself is typically not  $Sym(\mathbb{N})$ -Noetherian, let alone  $Inc(\mathbb{N})$ -Noetherian. Take, for instance,  $Y = \{y_{ij} \mid i, j \in \mathbb{N}\}$  with  $Sym(\mathbb{N})$  acting diagonally on both indices, and take any R with  $1 \neq 0$ . Then the  $Sym(\mathbb{N})$ orbits of the monomials

### $y_{12}y_{21}, y_{12}y_{23}y_{31}, y_{12}y_{23}y_{34}y_{41}, \dots$

generate a Sym( $\mathbb{N}$ )-stable ideal that is not generated by any finite union of orbits (see [AH07, Proposition 5.2]).

- (2) The *R*-algebra *R*[*Z*] is Sym(N)-Noetherian, and even Inc(N)-Noetherian [Coh87, HS12]—this is the special case of our theorem where *Y* = *Z* and *φ* is the identity—but Sym(N)-stable subalgebras of *R*[*Z*] need not be, even when generated by finitely many Sym(N)-orbits of polynomials. For instance, an (as yet) unpublished theorem due to Krasilnikov says that in characteristic 2, the ring generated by all 2 × 2-minors of a 2 × N-matrix of variables is not Sym(N)-Noetherian. Put differently, we do not know if the finite generatedness of ker *φ* in the Main Theorem continues to hold if *φ* is an arbitrary Sym(N)-equivariant homomorphism, but certainly the quotient is not, in general, Sym(N)-Noetherian.
- (3) Moreover, subalgebras of R[Z] generated by finitely many Inc(N)-orbits of monomials need not be Inc(N)-Noetherian; see Krasilnikov's example in [HS12]. However, our Main Theorem implies that subalgebras of R[Z] generated by finitely many Sym(N)-orbits of monomials are Inc(N)-Noetherian.

Our Main Theorem applies to many problems on Markov bases of families of point sets. In such applications, the following strengthening is sometimes useful.

**Corollary 1.2.** Assume that  $\operatorname{Sym}(\mathbb{N})$  has only finitely many orbits on Y, and let S be an R-algebra with trivial  $\operatorname{Sym}(\mathbb{N})$ -action. Let  $\varphi : R[Y] \to S[Z]$  be a  $\operatorname{Sym}(\mathbb{N})$ -equivariant R-algebra homomorphism that maps each  $y \in Y$  to an element of S times a monomial in the  $z_{ij}$ . Then ker  $\varphi$  is generated by finitely many  $\operatorname{Inc}(\mathbb{N})$ -orbits of binomials, and im  $\varphi \cong R[Y]/\ker \varphi$  is an  $\operatorname{Inc}(\mathbb{N})$ -Noetherian ring.

Proof of the Corollary given the Main Theorem. Let  $y_p, p \in [N]$  be representatives of the Sym(N)-orbits on Y. Then for all  $p \in [N]$  and  $\pi \in$  Sym(N) we have  $\varphi(\pi y_p) = s_p \pi u_p$  for some monomial  $u_p$  in the  $z_{ij}$  and some  $s_p$  in S. Apply the Main Theorem to  $Y' := Y \times \mathbb{N}$  and  $Z \cup Z'$  with  $Z' := \{z'_{p,j} \mid p \in [N], j \in \mathbb{N}\}$  and  $\varphi'$  the map that sends the variable  $(\pi y_p, j)$  to  $z'_{p,j} \pi u_p$ . Consider the commutative diagram

$$\begin{array}{ccc} R[Y'] & \stackrel{\varphi'}{\longrightarrow} R[Z \cup Z'] \\ & & & & \downarrow \\ & & & \downarrow \\ \rho:(y,j) \mapsto y & & & \downarrow \\ \psi:z'_{jj} \mapsto s_p \\ & & & R[Y] & \stackrel{\varphi}{\longrightarrow} S[Z] \end{array}$$

of Sym( $\mathbb{N}$ )-equivariant *R*-algebra homomorphisms. By the Main Theorem, im  $\varphi'$  is Inc( $\mathbb{N}$ )-Noetherian, hence so is its image under  $\psi$ ; and this image equals im  $\varphi$  because  $\rho$  is surjective. Similarly, ker( $\psi \circ \varphi'$ ) is generated by finitely many Inc( $\mathbb{N}$ )-orbits (because this is the case for both ker  $\varphi'$  and ker  $\psi|_{\operatorname{im} \varphi'}$ ), hence so is its image under  $\rho$ ; and this image is ker  $\varphi$  because  $\rho$  is surjective.  $\Box$ 

Here are some consequences of our Main Theorem.

- (1) Our Main Theorem implies [AH07, Conjecture 5.10] that chains of ideals arising as kernels of monomial maps of the form  $y_{i_1,\ldots,i_k} \mapsto z_{i_1}^{a_1} \cdots z_{i_k}^{a_k}$ , where the indices  $i_1,\ldots,i_k$  are required to be distinct, stabilize. In [AH07] this is proved in the squarefree case, where the  $a_j$  are equal to 1. In the Laurent polynomial setting more is known [HdC13].
- (2) A consequence of [dLST95] is that for any  $n \ge 4$  the vertex set  $\{v_{ij} := e_i + e_j \mid i \ne j\} \subseteq \mathbb{R}^n$  of the (n-1)-dimensional second hypersimplex has a Markov basis corresponding to the relations  $v_{ij} = v_{ji}$  and  $v_{ij} + v_{kl} = v_{il} + v_{kj}$ . Here is a qualitative generalisation of this fact. Let m and k be fixed

natural numbers. For every  $n \in \mathbb{N}$  consider a finite set  $P_n \subseteq \mathbb{Z}^m \times \mathbb{Z}^{k \times n}$ . Let  $\operatorname{Sym}(n)$  act trivially on  $\mathbb{Z}^m$  and by permuting columns on  $\mathbb{Z}^{k \times n}$ . Assume that there exists an  $n_0$  such that  $\operatorname{Sym}(n)P_{n_0} = P_n$  for  $n \ge n_0$ ; here we think of  $\mathbb{Z}^{k \times n_0}$  as the subset of  $\mathbb{Z}^{k \times n}$  where the last  $n - n_0$  columns are zero. Then Corollary 1.2 implies that there exists an  $n_1$  such that for any Markov basis  $M_{n_1}$  for the relations among the points in  $P_{n_1}$ ,  $\operatorname{Sym}(n)M_{n_1}$  is a Markov basis for  $P_n$  for all  $n \ge n_1$ . For the second hypersimplex,  $n_0$  equals 2 and  $n_1$  equals 4.

(3) A special case of the previous consequence is the Independent Set Theorem of [HS12]. We briefly illustrate how to derive it directly from Corollary 1.2. Let m be a natural number and let  $\mathcal{F}$  be a family of subsets of a finite set [m]. Let T be a subset of [m] and assume that each  $F \in \mathcal{F}$  contains at most one element of T. In other words, T is an independent set in the hypergraph determined by  $\mathcal{F}$ . For  $t \in [m] \setminus T$  let  $r_t$  be a natural number. Set  $Y := \{y_{\alpha} \mid \alpha \in \mathbb{N}^T \times \prod_{t \in [m] \setminus T} [r_t]\}$  and  $Z := \{z_{F,\alpha} \mid F \in \mathcal{F}, \alpha \in \mathbb{N}^T \mid T \in \mathbb{N}\}$  $\mathbb{N}^{F\cap T} \times \prod_{F\setminus T} [r_t]$ , and let  $\varphi$  be the homomorphism  $\mathbb{Z}[Y] \to \mathbb{Z}[Z]$  that maps  $y_{\alpha}$  to  $\prod_{F \in \mathcal{F}} z_{F,\alpha|_F}$ , where  $\alpha|_F$  is the restriction of  $\alpha$  from [m] to F. Then  $\varphi$  is equivariant with respect to the action of  $Sym(\mathbb{N})$  on the variables induced by the diagonal action of  $\text{Sym}(\mathbb{N})$  on  $\mathbb{N}^T$ , and (a strong form of) the Independent Set Theorem boils down to the statement that ker  $\varphi$  is generated by finitely many  $Sym(\mathbb{N})$ -orbits of binomials. By the condition that T is an independent set, each z-variable has at most one index running through all of N. Setting S to be  $\mathbb{Z}[z_{F,\alpha} \mid F \cap T = \emptyset]$ , we find that Y, S, the remaining  $z_{F,\alpha}$ -variables, with  $|F \cap T| = 1$ , and the map  $\varphi$  satisfy the conditions of the corollary. The conclusion of the corollary now implies the Independent Set Theorem.

The remainder of the paper is organized as follows: In Section 2 we reduce the Main Theorem to a particular class of maps  $\varphi$  related to matching monoids of complete bipartite graphs. For these maps, finite generation of the kernel follows from recent results on the Birkhoff model [YOT14]; see Section 3, where we also describe the image of  $\varphi$ . In Section 4 we prove Noetherianity of im $\varphi$ , still for our special  $\varphi$ . As in [Coh87, HS12], the strategy in Section 4 is to prove that a partial order on certain monoids is a well-partial-order. In our case, these are said matching monoids, and the proof that they are well-partially-ordered is quite subtle. In Section 5 we establish that a finite Inc(N)-generating set of ker  $\varphi$  is (at least theoretically) computable. The last Section describes a simpler procedure that one can attempt in order to obtain a generating set; at the moment, we do not know if this procedure is guaranteed to terminate. We conclude the paper with a computational example for which termination does occur.

Acknowledgments. We thank a referee for pointing out the recent results [YOT14] on the Markov degree of the (generalized) Birkhoff model—which are stronger than a bound that we had in the original version of this paper—and for the remark that the integral polytopes capturing this model are not only normal but even compressed (see Remark 3.2).

### 2. Reduction to matching monoids

In this section we reduce the Main Theorem to a special case to be treated in the next two sections. To formulate this special case, let  $N \in \mathbb{N}$  and for each  $p \in [N]$  let  $k_p \in \mathbb{N}$ . First, introduce a set Y' of variables  $y'_{p,J}$  where  $p \in [N]$  and  $J = (j_l)_{l \in [k_p]} \in \mathbb{N}^{[k_p]}$  is a  $k_p$ -tuple of *distinct* natural numbers. The group Sym( $\mathbb{N}$ ) acts on Y' by  $\pi y'_{p,J} = y'_{p,\pi(J)}$  where  $\pi(J) = (\pi(j_l))_{l \in [k_p]}$ . This action has finitely many orbits and satisfies the condition preceding the Main Theorem. Second, let X be a set of variables  $x_{p,l,j}$  with  $p \in [N], l \in [k_p], j \in \mathbb{N}$  and let Sym( $\mathbb{N}$ ) act on Xby its action on the last index.

**Proposition 2.1.** Let  $\varphi' : R[Y'] \to R[X]$  be the *R*-algebra homomorphism sending  $y'_{p,J}$  to  $\prod_{l \in [k_p]} x_{p,l,j_l}$ . Then the Main Theorem implies that ker  $\varphi'$  is generated by finitely many  $\operatorname{Inc}(\mathbb{N})$ -orbits of binomials, and that im  $\varphi'$  is an  $\operatorname{Inc}(\mathbb{N})$ -Noetherian ring. Conversely, if these two statements hold for all choices of  $N, k_1, \ldots, k_N \in \mathbb{N}$ , then the Main Theorem holds.

*Proof.* The first statement is immediate—note that the pair (p, l) comprising the first two indices of the variables  $x_{p,l,j}$  takes on finitely many, namely,  $\sum_{p} k_p$  values.

For the second statement, consider a monomial map  $\varphi : R[Y] \to R[Z]$  with  $Z = \{z_{i,j} \mid i \in [k], j \in \mathbb{N}\}$  as in the Main Theorem. Let N be the number of Sym( $\mathbb{N}$ )-orbits on Y and let  $y_p, p \in [N]$  be representatives of the orbits. Set  $k_p := k_{y_p}$  for  $p \in [N]$ , so that  $\pi y_p$  depends only on the restriction of  $\pi \in \text{Sym}(\mathbb{N})$  to  $[k_p]$ . We have thus determined the values of N and the  $k_p$ , and we let Y', X be as above.

Let  $\psi : R[Y'] \to R[Y]$  be the *R*-algebra homomorphism defined by sending  $y'_{p,J}$  to  $\pi y_p$  for any  $\pi \in \text{Sym}(\mathbb{N})$  satisfying  $\pi(l) = j_l$ ,  $l \in [k_p]$ . This homomorphism is  $\text{Sym}(\mathbb{N})$ -equivariant. The composition  $\varphi'' := \varphi \circ \psi : R[Y'] \to R[Z]$  satisfies the conditions of the Main Theorem. Since  $\psi$  is surjective, it maps any generating set for ker  $\varphi''$  onto a generating set for ker  $\varphi$ ; moreover, we have im  $\varphi'' = \text{im } \varphi$ . Hence the conclusions of the Main Theorem for  $\varphi''$  imply those for  $\varphi$ .

Next write  $\varphi''(y_{p,J}) = \prod_{i \in [k], j \in \mathbb{N}} z_{i,j}^{d_{p,i,j}}$ . Observe that  $d_{p,i,j} = 0$  whenever  $j \notin J$ , using the fact that any permutation that fixes J also fixes  $y_{p,J}$ , and hence must also fix  $\varphi''(y_{p,J})$  by Sym(N)-equivariance. Now let  $\varphi' : K[Y'] \to K[X]$  be as above and define  $\rho : R[X] \to R[Z]$  by  $\rho(x_{p,l,j}) = \prod_{i \in [k]} z_{i,j}^{d_{p,i,j}}$ . By construction, we have  $\rho \circ \varphi' = \varphi''$ .

Now im  $\varphi''$  is a quotient of im  $\varphi'$  and ker  $\varphi''$  is generated by ker  $\varphi'$  together with pre-images of generators of ker $(\rho|_{\mathrm{im} \varphi'})$ , hence the conclusions of the Main Theorem for  $\varphi'$  imply those for  $\varphi''$ , as desired.

In what follows, we will drop the accents on the *y*-variables and write Y for the set of variables  $y_{p,J}$ , X for the set of variables  $x_{p,l,j}$ , and  $\varphi$  for the *R*-algebra homomorphism

(1) 
$$\varphi: R[Y] \to R[X], \ y_{p,J} \mapsto \prod_{l \in [k_p]} x_{p,l,j_l}.$$

Monomials in the  $x_{p,l,j}$  will be denoted  $x^A$  where  $A \in \prod_{p \in [N]} \mathbb{N}^{[k_p] \times \mathbb{N}}$  is an [N]-tuple of finite-by-infinite matrices  $A_p$ . Note that  $\varphi(y_{p,J})$  equals  $x^A$  where only the p-th component  $A_p$  of A is non-zero and in fact has all row sums equal to 1, all

column sums labelled by J equal to 1, and all other column sums equal to 0. Thus  $A_p$  can be thought of as the adjacency matrix of a matching of the maximal size  $k_p$  in the complete bipartite graph with bipartition  $[k_p] \bigsqcup \mathbb{N}$ . Thus the monomials in  $m \varphi$  form the Abelian monoid generated by such matchings (with p varying). We call a monoid like this a *matching monoid*. In the next section we characterize these monomials among all monomials in the  $x_{p,l,j}$ , and find a bound on the relations among the  $\varphi(y_{p,J})$ .



FIGURE 1. A bipartite graph on  $[2] \bigsqcup \mathbb{N}$  and its corresponding monomial  $x_p^{A_p}$  (top). This graph can be decomposed into matchings in two different ways (middle and bottom). Each decomposition represents a monomial in the preimage  $\varphi^{-1}(x_p^{A_p})$ .

### 3. Relations among matchings

We retain the setting at the end of the previous section: Y is the set of variables  $y_{p,J}$  with p running through [N] and  $J \in \mathbb{N}^{[k_p]}$  running through the  $[k_p]$ -tuples of distinct natural numbers; X is the set of variables  $x_{p,l,j}$  with  $p \in [N], l \in [k_p], j \in \mathbb{N}$ , and  $\varphi$  is the map in (1). In this section we describe both the kernel and the image of  $\varphi$ . Note that if some  $k_p$  is zero, then the corresponding (single) variable  $y_{p,()}$  is mapped by  $\varphi$  to 1. The image of  $\varphi$  does not change if we disregard those p, and the kernel changes only in that we forget about the generators  $y_{p,()} - 1$ . Hence we may and will assume that all  $k_p$  are strictly positive. The following lemma gives a complete characterization of the  $x^A$  in the image of  $\varphi$ .

**Proposition 3.1.** For an [N]-tuple  $A \in \prod_{p \in [N]} \mathbb{N}^{[k_p] \times \mathbb{N}}$  the monomial  $x^A$  lies in the image of  $\varphi$  if and only if for all  $p \in [N]$  the matrix  $A_p \in \mathbb{N}^{[k_p] \times \mathbb{N}}$  has all row sums equal to a number  $d_p \in \mathbb{N}$  and all column sums less than or equal to  $d_p$ .

We call such A good. Note that  $d_p$  is unique since all  $k_p$  are strictly positive. We call the vector  $(d_p)_p$  the *multi-degree* of A and of  $x^A$ .

**Remark 3.2.** By replacing  $\mathbb{N}$  with [n] for some natural number n greater than or equal to the maximum of the  $k_p$ , the proposition boils down to the statement that for each p the lattice polytope in  $\mathbb{R}^{[k_p] \times [n]}$  with defining inequalities  $\forall_{ij} a_{ij} \geq 0$ ,

 $\forall_i \sum_j a_{ij} = 1$ , and  $\forall_j \sum_i a_{ij} \leq 1$  is normal (in the case where  $n = k_p$  this is the celebrated *Birkhoff polytope*). This is a not new result; in fact, this polytope satisfies a stronger property, namely, it is *compressed*. This follows, for instance, from [Sul06, Theorem 2.4] or from the main theorem of [OH01]; see also [YOT14, Section 4.2]. For completeness, we include a proof of the proposition using elementary properties of matchings in bipartite graphs.

Proof. Let  $x_p$  denote the vector of variables  $x_{p,l,j}$  for  $l \in [k_p]$  and  $j \in \mathbb{N}$ . By definition of  $\varphi$ , the monomial  $x^A$  lies in im  $\varphi$  if and only if the monomial  $x_p^{A_p}$  lies in im  $\varphi$  for all  $p \in [N]$ . Thus it suffices to prove that  $x_p^{A_p}$  lies in im  $\varphi$  if and only if all row sums of  $A_p$  are equal, say to  $d \in \mathbb{N}$ , and all column sums of  $A_p$  are at most d. The "only if" part is clear, since every variable  $y_{p,J}$  is mapped to a monomial  $x_p^B$  where  $B \in \mathbb{N}^{[k_p] \times \mathbb{N}}$  has all row sums 1 and all column sums at most 1. For the "if" part we proceed by induction on d: assume that the statement holds for d-1, and consider a matrix  $A_p$  with row sums d and column sums  $\leq d$ , where d is at least 1. Clearly, the "if" part is true in the case d = 0.

Think of  $A_p$  as the adjacency matrix of a bipartite graph  $\Gamma$  (with multiple edges) with bipartition  $[k_p] \bigsqcup \mathbb{N}$  (see Figure 1). With this viewpoint in mind, we will invoke some standard results from combinatorics, and refer to [Sch03, Chapter 16]. The first observation is that  $\Gamma$  contains a matching that covers all vertices in  $[k_p]$ . Indeed, otherwise, by Hall's marriage theorem, after permuting rows and columns,  $A_p$  has the block structure

$$A_p = \begin{bmatrix} A_{11} & 0\\ A_{12} & A_{22} \end{bmatrix}$$

with  $A_{11} \in \mathbb{N}^{[l] \times [l-1]}$  for some  $l, 1 \leq l \leq k_p$ . But then the entries of  $A_{11}$  added rowwise add up to ld, and added column-wise add up to at most (l-1)d, a contradiction. Hence  $\Gamma$  contains a matching that covers all of  $[k_p]$ . Next, let  $S \subseteq \mathbb{N}$  be the set of column indices where  $A_p$  has column sum equal to the upper bound d. We claim that  $\Gamma$  contains a matching that covers all of S. Indeed, otherwise, again by Hall's theorem, after permuting rows and columns  $A_p$  has the structure

$$A_p = \begin{bmatrix} A_{11} & A_{12} \\ 0 & A_{22} \end{bmatrix}$$

with  $A_{11} \in \mathbb{N}^{[l-1] \times [l]}$  for some  $l, 1 \leq l \leq |S|$ ; here the first l columns correspond to a subset of the original S. Now the entries of  $A_{11}$  added columnwise yield ld, while the entries of  $A_{11}$  added rowwise yield at most (l-1)d, a contradiction.

Finally, we invoke a standard result in matching theory (see [Sch03, Theorem 16.8]), namely that since  $\Gamma$  contains a matching that covers all of  $[k_p]$  and a matching that covers all of S, it also contains a matching that covers both. Let B be the adjacency matrix of this matching, so that B has all row sums 1 and all column sums  $\leq 1$ , with equality at least in the columns labelled by S. Then  $A'_p := A_p - B$  satisfies the induction hypothesis for d-1, so  $x_p^{A'_p} \in \operatorname{im} \varphi$ . Also,  $x_p^B = \varphi(y_{p,J})$ , where  $j_a \in \mathbb{N}$  is the neighbour of  $a \in [k_p]$  in the matching given by B. Hence,  $x_p^{A_p} = x_p^{A'_p} x_p^B \in \operatorname{im} \varphi$  as claimed.

This concludes the description of the image of  $\varphi$ . For the kernel, we quote the following result.

**Theorem 3.3** (Theorem 2.1 of [YOT14]). The kernel of  $\varphi$  from (1) is generated by binomials in the  $y_{p,J}$  of degree at most 3.

Indeed, for each fixed p, and replacing  $\mathbb{N}$  by some [n] with  $n \geq k_p$ , the monomial map (1) captures precisely the generalization of the Birkhoff model studied in [YOT14], where each voter choses  $k_p$  among n candidates. Then their Theorem 2.1 yields that the kernel is generated in degrees 2 and 3. Since this holds for each  $n \geq k_p$ , it also holds for  $\mathbb{N}$  instead of [n]. Moreover, taking the union over all p of sets of generators for each individual p yields a set of generators for the kernel of  $\varphi$ . A straightforward consequence of the theorem is the following.

**Corollary 3.4.** The kernel of  $\varphi$  from (1) is generated by finitely many  $\text{Inc}(\mathbb{N})$ -orbits of binomials.

### 4. Noetherianity of matching monoid rings

By Corollary 3.4 and Proposition 2.1, Main Theorem follows from the following proposition.

**Proposition 4.1.** The ring  $S = R[x^A \mid A \in \prod_{p \in [N]} \mathbb{N}^{[k_p] \times \mathbb{N}} \text{ good}]$  is  $\operatorname{Inc}(\mathbb{N})$ -Noetherian.

Let  $G \subset \prod_{p \in [N]} \mathbb{N}^{[k_p] \times \mathbb{N}}$  be the set of good (N-tuples of) matrices, so the monomials of S are precisely  $x^A$  for  $A \in G$ . The monoids  $\operatorname{Sym}(\mathbb{N})$  and  $\operatorname{Inc}(\mathbb{N})$  act on G by permuting or shifting columns, so we have  $\pi x^A = x^{\pi A}$ , where the  $\pi(j)$ -th column of the matrix  $(\pi A)_p$  equals the *j*-th column of  $A_p$ . Let  $d_A = (d_{A,p})_p \in \mathbb{N}^{[N]}$ denote the multi-degree of A; recall that this means that all row sums of  $A_p$  are equal to  $d_{A,p}$ . To prove Noetherianity we will define a partial order  $\preceq$  on G and prove that  $\preceq$  is a well-partial-order. Thus we need some basic results from order theory.

A partial order  $\leq$  on a set P is a *well-partial-order* (or *wpo*) if for every infinite sequence  $p_1, p_2, \ldots$  in P, there is some i < j such that  $p_i \leq p_j$ ; see [Kru72] for alternative characterisations. For instance, the natural numbers with the usual total order  $\leq$  is a well-partial-order, and so is the componentwise partial order on the Cartesian product of any finite number of well-partially-ordered sets. Combining these statements yields Dickson's Lemma [Dic13] that  $\mathbb{N}^k$  is well-partially-ordered. This can be seen as a special case of Higman's Lemma [Hig52], for a beautiful proof of which we refer to [NW63].

**Lemma 4.2** (Higman's Lemma). Let  $(P, \preceq)$  be a well-partial-order and let  $P^* := \bigcup_{l=0}^{\infty} P^l$ , the set of all finite sequences of elements of P. Define the partial order  $\preceq'$  on  $P^*$  by  $(a_0, \ldots, a_{l-1}) \preceq' (b_0, \ldots, b_{m-1})$  if and only if there exists a strictly increasing function  $\rho : [l] \to [m]$  such that  $a_j \preceq b_{\rho(j)}$  for all  $j \in [l]$ . Then  $\preceq'$  is a well-partial-order.

Our interest in well-partial-orders stems from the following application. Consider a commutative monoid M with an action of a (typically non-commutative) monoid II by means of monoid endomorphisms. We suggestively call the elements of M monomials. Assume that we have a  $\Pi$ -compatible monomial order  $\leq$  on M, i.e., a well-order that satisfies  $a < b \Rightarrow ac < bc$  and  $a < b \Rightarrow \pi a < \pi b$  for all  $a, b, c \in M$ and  $\pi \in \Pi$ . Then it follows that the divisibility relation | defined by a|b if there exists a  $c \in M$  with ac = b is a partial order, and also that  $a \leq \pi a$  for all  $a \in M$ . Define a third partial order, the  $\Pi$ -divisibility order,  $\leq$  on M by  $a \leq b$  if there exists a  $\pi \in \Pi$  and a  $c \in M$  such that  $c\pi a = b$ . A straightforward computation shows that  $\leq$  is, indeed, a partial order—antisymmetry follows using  $a \leq \pi a$ .

**Proposition 4.3.** If  $\leq$  is a well-partial-order, then for any Noetherian ring R, the R-algebra R[M] is  $\Pi$ -Noetherian.

*Proof.* This statement was proved in [HS12] for the case where R is a field. The more general case can be proved with the same argument by incorporating work done in [AH07].

Note that the monoid  $\{x^A \mid A \in G\}$  that we are considering here can be given a monomial order which respects the  $\text{Inc}(\mathbb{N})$ -action. For example, take the lexicographic order, where the variables  $x_{p,i,j}$  are ordered by their indices:  $x_{p,i,j} < x_{p',i',j'}$ if and only if p < p'; or p = p' and j < j'; or p = p', j = j', and i < i'.

The Inc( $\mathbb{N}$ )-divisibility order gives a partial order  $\leq$  on the set G of good (N-tuples of) matrices by  $A \leq B$  if and only if there is a monomial  $x^C \in S$  and  $\pi \in \text{Inc}(\mathbb{N})$  such that  $x^C \pi(x^A) = x^B$ , or equivalently there is  $\pi \in \text{Inc}(\mathbb{N})$  such that  $B - \pi A \in G$ . Note that  $A \leq B$  not only implies there is some  $\pi \in \text{Inc}(\mathbb{N})$  such that all  $A_{p,i,j} \leq B_{p,i,\pi(j)}$ , but additionally that all (N-tuples of) column sums of  $B - \pi A$  are at most  $d_B - d_A \in \mathbb{N}^{[N]}$ . This prevents us from applying Higman's Lemma directly to  $(G, \leq)$ . To encode this condition on column sums, for any  $A \in G$ , let  $\tilde{A} \in \prod_{p \in [N]} \mathbb{N}^{[k_p+1] \times \mathbb{N}}$  be the N-tuple of matrices such that for all  $p \in [N]$ , the first  $k_p$  rows of  $\tilde{A}_p$  are equal to  $A_p$ , and the last row of  $\tilde{A}_p$  is such that all column sums

$$\tilde{A}_{p,i,j} = \begin{cases} A_{p,i,j} & \text{for } i < k_p, \text{ and} \\ d_{A,p} - \sum_{l=0}^{k_p - 1} A_{p,l,j} & \text{for } i = k_p. \end{cases}$$

We let  $\tilde{G}$  be the set of *N*-tuples of matrices of the form  $\tilde{A}$  with  $A \in G$ . It is precisely the set of *N*-tuples of matrices of the form  $\tilde{A} \in \prod_{p \in [N]} \mathbb{N}^{[k_p+1] \times \mathbb{N}}$  with the property that there exists a  $d_A \in \mathbb{N}^{[N]}$  such that for each  $p \in [N]$  the first  $k_p$  row sums of  $A_p$  are equal to  $d_{A,p}$  and all column sums of  $A_p$  are equal to  $d_{A,p}$ . Since  $A \in G$  has only finitely many *N*-tuples of non-zero columns,  $\tilde{A}$  will have all but finitely many *N*-tuples of columns equal to  $((0, \ldots, 0, d_{A,p})^T)_{p \in [N]}$ . Such *N*-tuples of columns will be called *trivial* (of degree  $d_A$ ). The *N*-tuple of *j*th columns of  $\tilde{A}$ will be denoted  $\tilde{A}_{..j}$ . We define the action of  $\text{Inc}(\mathbb{N})$  on  $\tilde{G}$  as  $\pi(\tilde{A}) = \widetilde{\pi(A)}$ . Note that for any  $j \notin \text{im}(\pi)$ , the column  $(\pi \tilde{A})_{..j}$  is trivial of degree  $d_A$ , rather than uniformly zero.

**Proposition 4.4.** For  $A, B \in G$ ,  $A \preceq B$  if and only if there is  $\pi \in \text{Inc}(\mathbb{N})$  such that  $\pi \tilde{A} \leq \tilde{B}$  entry-wise.

*Proof.* The condition that  $(\pi \tilde{A})_{p,i,j} \leq \tilde{B}_{p,i,j}$  for all  $p \in [N]$ , all  $i < k_p$ , and all  $j \in \mathbb{N}$  is equivalent to the condition that  $B - \pi A$  is non-negative. Using the fact that

$$\tilde{B}_{p,k_p,j} - (\pi \tilde{A})_{p,k_p,j} = (d_{B,p} - d_{A,p}) - \sum_{i=0}^{\kappa_p - 1} (B_p - \pi A_p)_{i,j},$$

the condition that  $\tilde{B}_{p,k_p,j} - (\pi \tilde{A})_{p,k_p,j} \ge 0$  for all  $p \in [N]$  and all  $j \in \mathbb{N}$  is equivalent to the condition that every N-tuple of column sums of  $B - \pi A$  is less than or equal to  $d_B - d_A$ . Therefore  $\pi \tilde{A} \le \tilde{B}$  if and only if  $B - \pi A \in G$ . **Example 4.5.** Let A and B be the following good matrices in  $\mathbb{N}^{[2] \times \mathbb{N}}$ :

<i>A</i> =	$=\begin{bmatrix}3\\0\end{bmatrix}$	$\begin{array}{c} 0 \\ 1 \end{array}$	$\begin{array}{c} 0 \\ 1 \end{array}$	$\begin{array}{c} 0 \\ 1 \end{array}$	$\begin{array}{c} 0 \\ 0 \end{array}$	  ],	$B = \left[ \right]$	3 0	$\frac{1}{2}$	$\begin{array}{c} 0 \\ 1 \end{array}$	$\begin{array}{c} 0 \\ 1 \end{array}$	$\begin{array}{c} 0 \\ 0 \end{array}$	  ].

Note that  $\pi A \leq B$  when  $\pi$  is the identity, however  $A \not\preceq B$ . Consider

	3	0	0	0	0	• • • •			3	1	0	0	0	• • • •	1
$\tilde{A} =$	0	1	1	1	0		,	$\tilde{B} =$	0	2	1	1	0		
	0	2	2	2	3	••••			1	1	3	3	4	•••	

and note that there is no  $\pi \in \text{Inc}(\mathbb{N})$  such that  $\pi A \leq B$ .

We will work with finite truncations of N-tuples of matrices in  $\tilde{G}$ . Let H be the set of N-tuples of matrices  $A \in \bigcup_{\ell=0}^{\infty} \prod_{p \in [N]} \mathbb{N}^{[k_p+1] \times [\ell]}$  such that there exists  $d_A \in \mathbb{N}^{[N]}$  such that for all p, all column sums of  $A_p$  are equal to  $d_{A,p}$  and the first  $k_p$ row sums are at most  $d_{A,p}$ ; we call  $d_A$  the multi-degree of A. Note that the condition on row sums is relaxed, which will allow us to freely remove columns from matrices while still remaining in the set H. For  $A \in H$  the number of columns of A is called the *length* of A and denoted  $\ell_A$ . We give H the partial order  $\preceq$  defined as follows. For  $A, B \in H, A \preceq B$  if and only if there is a strictly increasing map  $\rho : [\ell_A] \rightarrow [\ell_B]$ such that  $\rho A \leq B$ . Just as in  $\tilde{G}$ , here  $\rho A$  is defined by  $(\rho A)_{..j} = A_{..\rho^{-1}(j)}$  for  $j \in im(\rho)$ , and  $(\rho A)_{..j}$  trivial (of degree  $d_A$ ) for  $j \in [\ell_B] \setminus im(\rho)$ . For an N-tuple of matrices A and a set  $J \subset \mathbb{N}$ , let  $A_{..J}$  denote the N-tuple of matrices obtained from A by taking only the columns  $A_{..j}$  with  $j \in J$ .

Some care must be taken in the definition of H since we allow matrices with no columns. In all other cases, the degree of  $A \in H$  is uniquely determined by its entries. However for the length 0 case the degree is arbitrary, so we will consider H as having a distinct length 0 element  $Z^d$  with degree d for each  $d \in \mathbb{N}^{[N]}$ , and we define  $Z^d \leq A$  if and only if  $d \leq d_A$ . Additionally, define  $A_{..\emptyset} = Z^{d_A}$ .

**Definition 4.6.** For  $A \in H$ , the *N*-tuple of *j*th columns of *A* is *bad* if for some  $p \in [N]$ , we have  $A_{p,k_p,j} < d_{A,p}/2$ . If  $A_{p,k_p,j} < d_{A,p}/2$ , we will call *j* a *bad index* of *A* (with respect to *p*). Let  $H_t$  denote the set of *N*-tuples of matrices in H with exactly *t* bad indices.

We will use induction on t to show that  $(H_t, \preceq)$  is well-partially ordered for all  $t \in \mathbb{N}$ . This will in turn be used to prove that  $(H, \preceq)$  and then  $(\tilde{G}, \preceq)$  are well-partially ordered. First we prove the base case:

## **Proposition 4.7.** $(H_0, \preceq)$ is well-partially ordered.

*Proof.* Let  $A^{(1)}, A^{(2)}, \ldots$  be any infinite sequence in  $H_0$ . We will show that there are r < s such that  $A^{(r)} \preceq A^{(s)}$ .

Fix  $p \in [N]$ . There are now two possibilities: either the degrees of the elements of the sequence  $A_p^{(1)}, A_p^{(2)}, \ldots$  are bounded by some  $d_p \in \mathbb{N}$ , or they are not. In the former case, it follows that the number of non-trivial columns in any  $A_p^{(r)}$  is bounded by  $d_pk_p$ . Then there is a subsequence  $B_p^{(1)}, B_p^{(2)}, \ldots$  of  $A_p^{(1)}, A_p^{(2)}, \ldots$  such that every element has the same degree and same number of non-trivial columns. In the latter case,  $A_p^{(1)}, A_p^{(2)}, \ldots$  has a subsequence with strictly increasing degree and moreover a subsequence  $B_p^{(1)}, B_p^{(2)}, \ldots$  with the property that  $d_{B^{(s+1)},p} \geq 2d_{B^{(s)},p}$ for all  $s \in \mathbb{N}$ . In either case, without loss of generality, we replace  $A^{(1)}, A^{(2)}, \ldots$  by  $B^{(1)}, B^{(2)}, \ldots$ . We repeat this procedure for all  $p \in [N]$ , and we find that  $A^{(1)}, A^{(2)}, \ldots$  contains a subsequence  $B^{(1)}, B^{(2)}, \ldots$  such that for all  $p \in [N]$ , one of the following two statements holds.

1: Both  $d_{B^{(t)},p}$  and the number of non-trivial columns in  $B_p$  are constant. 2: We have  $d_{B^{(t+1)},p} \ge 2d_{B^{(t)},p}$  for all t.

It now suffices to show that there are r < s such that  $B^{(r)} \leq B^{(s)}$  for all r < s. Define the partial order  $\sqsubseteq$  on  $H_0$  by  $A \sqsubseteq B$  if and only if there exists strictly increasing  $\rho : [\ell_A] \rightarrow [\ell_B]$  such that  $A_{\cdot,j} \leq B_{\cdot,\rho(j)}$  for all  $j \in [\ell_A]$ . By Higman's Lemma (Lemma 4.2),  $\sqsubseteq$  is a wpo. This means that there exist r < s such that  $B^{(r)} \sqsubseteq B^{(s)}$ . Fix such a pair r < s. We will show that  $B^{(r)} \preceq B^{(s)}$ .

Let  $\rho : [\ell_{B^{(r)}}] \to [\ell_{B^{(s)}}]$  be a strictly increasing map that witnesses  $B^{(r)} \sqsubseteq B^{(s)}$ . We claim that it also witnesses  $B^{(r)} \preceq B^{(s)}$ . For this, we have to show that  $\rho B^{(r)} \leq B^{(s)}$ . By the properties of  $\sqsubseteq$ , we already have  $(\rho B^{(r)})_{..\rho(j)} \leq B^{(s)}_{..\rho(j)}$ , which is to say that it suffices to show that for all  $j \notin \operatorname{im}(\rho)$ , we have  $d_{B^{(r)}} \leq (B^{(s)}_{p,k_p,j})_{p \in [N]}$ .

Let  $p \in [N]$ . Suppose we are in the case that both  $d_{B^{(t)},p}$  and the number of non-trivial columns in  $B_p$  are constant. Since  $\rho$  must map non-trivial columns of  $B_p^{(r)}$  to non-trivial columns of  $B_p^{(s)}$ , we conclude that if  $j \notin \operatorname{in}(\rho)$ , then the *j*-th column of  $B_p^{(s)}$  is trivial, and hence  $(B_{p,k_p,j}^{(s)}) = d_{B^{(s)},p}$ . But the latter equals  $d_{B^{(r)},p}$ , so certainly  $d_{B^{(r)},p} \leq (B_{p,k_p,j}^{(s)})$ .

Alternatively, suppose we have  $d_{B^{(t+1)},p} \ge 2d_{B^{(t)},p}$  for all t. Since  $B_p^{(s)}$  has no bad columns, we have

$$B_{p,k_p,j}^{(s)} \ge \frac{1}{2} d_{B^{(s)},p} \ge d_{B^{(r)},p}.$$

This is exactly what we wanted to show.

So in both cases, we find that  $d_{B^{(r)},p} \leq B_{p,k_p,j}^{(s)}$  for all  $j \notin \operatorname{im}(\rho)$ . This is true for all p, so we have  $d_{B^{(r)}} \leq (B_{p,k_p,j}^{(s)})_{p \in [N]}$ . We conclude that  $B^{(r)} \preceq B^{(s)}$ , as we wanted to show.

### **Proposition 4.8.** $(H_t, \preceq)$ is well-partially ordered for all $t \in \mathbb{N}$ .

*Proof.* The base case, t = 0, is given by Proposition 4.7. For t > 0, assume by induction that  $(H_{t-1}, \preceq)$  is well-partially ordered. For any  $A \in H_t$ , let  $j_A$  be the largest bad index of A. Then A can be decomposed into three parts: the N-tuple of matrices of all N-tuples of columns before  $j_A$ ,  $A_{..j_A}$  itself, and the N-tuple of matrices of all N-tuples of columns after  $j_A$ . This decomposition is represented by the map

$$\delta: \mathbf{H}_t \to \mathbf{H}_{t-1} \times \prod_{p \in [N]} \mathbb{N}^{[k_p+1]} \times \mathbf{H}_0$$
$$A \mapsto (A_{\dots\{0,\dots,j_A-1\}}, A_{\dots j_A}, A_{\dots\{j_A+1,\dots,\ell_A-1\}}).$$

Let the partial order  $\sqsubseteq$  on  $\mathcal{H}_{t-1} \times \prod_{p \in [N]} \mathbb{N}^{[k_p+1]} \times \mathcal{H}_0$  be the product order of the wpos  $(\mathcal{H}_{t-1}, \preceq)$ ,  $(\mathbb{N}^{[k+1]}, \leq)$  and  $(\mathcal{H}_0, \preceq)$ . Note that the product order of any finite number of wpos is also a wpo. Suppose for some  $A, B \in \mathcal{H}_t$  that  $\delta(A) \sqsubseteq \delta(B)$ . This implies that  $A_{\cdot,j_A} \leq B_{\cdot,j_B}$  and that there exist strictly increasing maps  $\rho$  and

 $\sigma$  such that  $\rho(A_{\cdot\cdot [j_A]}) \leq B_{\cdot\cdot [j_B]}$  and  $\sigma(A_{\cdot\cdot \{j_A+1,\ldots,\ell_A-1\}}) \leq B_{\cdot\cdot \{j_B+1,\ldots,\ell_B-1\}}$ . We combine these into a single strictly increasing map  $\tau: [\ell_A] \to [\ell_B]$  defined by

$$\tau(j) = \begin{cases} \rho(j) & \text{for } 0 \le j < j_A \\ j_B & \text{for } j = j_A \\ \sigma(j - j_A - 1) + j_B + 1 & \text{for } j_A < j < \ell_A \end{cases}$$

illustrated in Figure 2. Then  $\tau A \leq B$  so  $A \leq B$ . Since  $\sqsubseteq$  is a wpo,  $(\mathbf{H}_t, \preceq)$  is also a wpo.



FIGURE 2.  $\delta(A) \sqsubseteq \delta(B)$  implies  $A \preceq B$ .

**Proposition 4.9.**  $(H, \preceq)$  is well-partially ordered.

*Proof.* For any  $A \in H$ , if j is a bad index of A, then for some  $p \in [N]$ , we have  $d_{A,p}/2 > \sum_{i \in [k_p]} A_{p,i,j}$ . Letting  $J_p \subset \mathbb{N}$  be the set of bad indices of A with respect to p and let  $J \subset \mathbb{N}$  be the union of the  $J_p$ . Then

$$|J_p|\frac{d_{A,p}}{2} < \sum_{j \in J_p} \sum_{i \in [k_p]} A_{p,i,j} \le \sum_{i \in [k_p]} \sum_{j \in \mathbb{N}} A_{p,i,j} \le k_p d_A$$

with the last inequality due to the row sum condition on  $A_p$ . Therefore  $|J_p| \le 2k_p - 1$ , and hence  $|J| \le 2\sum_{p \in [N]} k_p - N$ .

Let  $A^{(1)}, A^{(2)}, \ldots$  be any infinite sequence in H. Since the numbers of bad Ntuples of columns of elements of H are bounded by  $2\sum_{p\in[N]} k_p - N$  there exists a subsequence which is contained in  $H_t$  for some  $0 \le t \le 2\sum_{p\in[N]} k_p - N$ . By Proposition 4.8 there is r < s with  $A^{(r)} \preceq A^{(s)}$ .

## **Proposition 4.10.** $(G, \preceq)$ is well-partially ordered.

Proof. Let  $A^{(1)}, A^{(2)}, \ldots$  be any infinite sequence in G. Each  $A^{(r)}$  has some  $j_r > 0$  such that all N-tuples of columns  $A^{(r)}_{\cdots m}$  are zero for  $m \ge j_r$ . Consider the sequence  $\tilde{A}^{(1)}_{\cdots [j_1]}, \tilde{A}^{(2)}_{\cdots [j_2]}, \ldots$  in H obtained by truncating each  $\tilde{A}^{(r)}$  to the first  $j_r$  N-tuples of columns. By Proposition 4.9 there is some r < s and  $\rho : [j_r] \to [j_s]$  such that  $\rho \tilde{A}^{(r)}_{\cdots [j_r]} \le \tilde{A}^{(s)}_{\cdots [j_s]}$ . Note that this implies  $d_{A^{(r)}} \le d_{A^{(s)}}$ . Extend  $\rho$  to some  $\pi \in \operatorname{Inc}(\mathbb{N})$  so then

$$(\pi \tilde{A}^{(r)})_{\cdot \cdot [j_s]} = \rho(\tilde{A}^{(r)}_{\cdot \cdot [j_r]}) \le \tilde{A}^{(s)}_{\cdot \cdot [j_s]}.$$

The remaining N-tuples of columns of  $\pi \tilde{A}^{(r)}$  and  $\tilde{A}^{(s)}$  are trivial, so  $\pi \tilde{A}^{(r)} \leq \tilde{A}^{(s)}$  follows from the fact that  $d_{A^{(r)}} \leq d_{A^{(s)}}$ . Therefore  $A^{(r)} \leq A^{(s)}$  by Proposition 4.4.

Now we can apply Proposition 4.3 to the monoid  $\{x^A \mid A \in G\}$  which proves that the ring  $R[x^A \mid A \in G]$  is  $Inc(\mathbb{N})$ -Noetherian. This concludes the proof of Proposition 4.1.

### 5. Buchberger's algorithm for matching monoid algebras

Assume the general setting of Proposition 4.3: M is a monoid with  $\Pi$ -action and  $\Pi$ -compatible monomial order  $\leq$ . For a polynomial f and an ideal I in K[M], we can define  $\operatorname{Im}(f)$ ,  $\operatorname{lc}(f)$ ,  $\operatorname{in}(I)$ , division with remainder, and the concept of equivariant Gröbner basis from [BD11]; all relative to the monomial order  $\leq$ . We now derive a version of Buchberger's algorithm for computing such a Gröbner basis, under an additional assumption. For  $a, b \in M$  we define the set of *least common multiples* 

 $lcm(a,b) = \{ l \in M : a|l, b|l \text{ and } (a|l', b|l', l'|l \Rightarrow l' = l) \}.$ 

We require the following variant of conditions EGB3 and EGB4 from [BD11]:

EGB34. For all  $f, g \in K[M]$ , the set of triples in  $M \times \Pi f \times \Pi g$ defined by

$$T_{f,g} = \{ (l', f', g') \mid f' \in \Pi f, g' \in \Pi g, l' \in \operatorname{lcm}(\operatorname{lm}(f'), \operatorname{lm}(g')) \},\$$

is a union of a finite number of  $\Pi$ -orbits:

$$T_{f,g} = \bigcup_{i} \Pi(l_i, f_i, g_i), \quad i \in [r].$$

In particular, EGB34 implies that for all  $a, b \in M$  and  $\pi \in \Pi$ , we have  $\pi \operatorname{lcm}(a, b) \subseteq \operatorname{lcm}(\pi a, \pi b)$ . (This is what condition EGB3 of [BD11] looks like when least common multiples are not unique.)

If EGB34 is fulfilled, then there is a unique *inclusion-minimal* finite set of orbit generators as above, which we denote

$$O_{f,g} = \{ (l_i, f_i, g_i) \mid i \in [r] \}.$$

Indeed, suppose that O and O' are both inclusion-minimal sets of orbit generators for  $T_{f,g}$ . For any triple  $t \in O$ , there are  $t' \in O', \pi \in \Pi$  such that  $\pi t' = t$ , and similarly  $t'' \in O, \tau \in \Pi$  such that  $\tau t'' = t'$ . Now  $t = \pi \tau t''$  and since O is minimal, t = t''. But since  $\Pi$  is compatible with a monomial order,  $\pi \tau t'' = t''$  implies that also the intermediate expression  $t' = \tau t''$  equals t''. Hence  $O \subseteq O'$  and equality holds by minimality of O'.

**Definition 5.1.** For monic  $f, g \in K[M]$  define the set of S-polynomials to be

$$S_{f,g} = \{af' - bg' \mid (l', f', g') \in O_{f,g}; a, b \in M; \text{ and } a \operatorname{Im}(f') = b \operatorname{Im}(g') = l'\}.$$

Furthermore, define  $\Pi$ -reduction of a polynomial f with respect to a set  $G \subseteq K[M]$  as follows: while there exist  $g \in G$  and  $\pi \in \Pi$  with  $\pi \operatorname{Im}(g)|\operatorname{Im}(f)$ , replace f by

$$f' := f - \frac{\operatorname{lc}(f)\operatorname{lm}(f)}{\operatorname{lc}(g)\pi\operatorname{lm}(g)}\pi g;$$

and when no such g and  $\pi$  exist, return the *remainder* f'.

One can generalize Gröbner theory to our equivariant setting for a monoid algebra satisfying EGB34. In particular, Buchberger's criterion holds, and the following procedure produces an equivariant Gröbner basis *if* it terminates.

Algorithm 5.2. G = BUCHBERGER(F)

**Require:** F is a finite set of monic elements in K[M], the algebra of a monoid M equipped with a  $\Pi$ -action, satisfying the assumptions above and the condition EGB34.

**Ensure:** G is an equivariant Gröbner basis of  $\langle F \rangle$ .

1:  $G \leftarrow \overline{F}$ 2:  $S \leftarrow \bigcup_{f,g \in G} S_{f,g}$ 3: while  $S \neq \emptyset$  do -- in particular, compute  $O_{f,g}$  needed in Definition 5.1 4: pick  $f \in S$  $S \leftarrow S \setminus \{f\}$ 5: $h \leftarrow$  the  $\Pi$ -reduction of f with respect to G 6: if  $h \neq 0$  then 7:  $G \leftarrow G \cup \{h\}$ 8:  $S \leftarrow S \cup \left(\bigcup_{g \in G} S_{g,h}\right)$ 9: 10: end if 11: end while

This algorithm has been implemented for the particular case where K[M] is a polynomial ring and  $\Pi = \text{Inc}(\mathbb{N})$  (i.e. the algorithm described in [BD11]) in the package EquivariantGB [Kro] for the computer algebra system Macaulay2 [GS02]. When the algebra K[M] is  $\Pi$ -Noetherian, termination of Algorithm 5.2 is guaranteed, but in general we cannot make this claim.

We now turn our attention to the task of computing a finite  $\text{Inc}(\mathbb{N})$ -generating set of binomials of a general toric map as in the Main Theorem. By the proof of Proposition 2.1 we may assume that Y is as in (1), i.e., it consists of variables  $y_{p,J}$  where p runs through [N] and J runs through all  $k_p$ -tuples of distinct natural numbers. Section 2 then leads to the following analysis of this task.

**Problem 5.3.** Fix the names of algebras and maps in the following diagram:

$$R[Y] \xrightarrow{\varphi} R[X] \xrightarrow{\psi} R[Z].$$

Here  $\varphi$  is the map defined by (1), whose image is the R-algebra spanned by the matching monoid, and  $\psi$  is any Sym(N)-equivariant monomial map from R[X] to  $R[z_{ij} \mid i \in [k], j \in \mathbb{N}]$ . For ker( $\psi \circ \varphi$ ), how does one compute

- (a) a finite set of generators up to  $Inc(\mathbb{N})$ -symmetry?
- (b) a finite Inc(N)-Gröbner basis with respect to a given Inc(N)-compatible monomial order on K[Y]?

The algorithm we are about to construct solves Problem 5.3(a); indeed, we do not know whether a finite  $\text{Inc}(\mathbb{N})$ -Gröbner basis as in part (b) exists! Our algorithm relies on the fact that we may replace R[X] above by the matching monoid algebra im  $\varphi = R[x^A : A \text{ good}]$ , so as to get the sequence

(2) 
$$R[Y] \xrightarrow{\varphi} R[x^A \mid A \text{ good}] \xrightarrow{\psi} R[Z].$$

Most of our computations will take place in the ring  $R[x^A \mid A \text{ good}][Z]$ , which is itself a matching monoid with N replaced by N+k and  $k_p = 1$  for  $p \in [N+k] \setminus [N]$ . This monoid is Gröbner friendly by the following proposition.

**Proposition 5.4.** Let M be a submonoid of  $\mathbb{N}^{[k] \times \mathbb{N}}$  that is generated by the Sym $(\mathbb{N})$ orbits of a finite number of matrices. For  $\Pi = \text{Inc}(\mathbb{N})$ , the monoid algebra K[M]satisfies EGB34.

*Proof.* Any such K[M] is the image of some map  $\varphi$  as in the Main Theorem (with R = K), and so is  $\operatorname{Inc}(\mathbb{N})$ -Noetherian. Similarly  $K[M^3] = K[M]^{\otimes 3}$  is  $\operatorname{Inc}(\mathbb{N})$ -Noetherian. For any  $a, b \in M$ , the monomial ideal  $\langle T_{a,b} \rangle \subseteq K[M^3]$  is  $\operatorname{Inc}(\mathbb{N})$ -stable. Let  $L \subseteq T_{a,b}$  be a minimal finite  $\operatorname{Inc}(\mathbb{N})$ -generating set of  $\langle T_{a,b} \rangle$ .

For any  $(l, \pi a, \sigma b) \in T_{a,b}$ , there is some  $(m, a', b') \in L$  and  $\tau \in \text{Inc}(\mathbb{N})$  such that  $\tau(m, a', b')|(l, \pi a, \sigma b)$ . It is clear that  $\tau a' = \pi a$  and  $\tau b' = \sigma b$ . Since a' and b' divide m,  $\pi a$  and  $\sigma b$  must divide  $\tau m$ , and in turn  $\tau m$  divides l. But  $l \in \text{lcm}(\pi a, \sigma b)$  by assumption, so  $l = \tau m$ . Therefore  $(l, \pi a, \sigma b) = \tau(m, a', b')$ . This shows that  $T_{a,b}$  is the union of the Inc( $\mathbb{N}$ )-orbits of the elements of L, and then  $L = O_{a,b}$ .

To establish the same fact for a general pair  $f, g \in K[M]$  we first determine  $O_{a,b}$ where  $a = \operatorname{Im}(f)$  and  $b = \operatorname{Im}(g)$ . For any  $(l, \pi f, \sigma g) \in T_{f,g}$ , the triple  $(l, \pi a, \sigma b) \in T_{a,b}$  is in the orbit of some  $(m, a', b') \in O_{a,b}$ . This implies  $a' = \tau a$  for some  $\tau \in \operatorname{Inc}(\mathbb{N})$ , but  $\tau$  is not unique. Define

$$\Lambda_{a,a'} := \{ \tau \in \operatorname{Inc}(\mathbb{N}) \mid a' = \tau a; \text{ and } n \in \operatorname{im} \tau \text{ for all } n > \ell_{a'} \}.$$

Here  $\ell_{a'}$  denotes the *length* of a' as in Section 4, the maximum index value among all non-zero columns of a'. Note that  $\Lambda_{a,a'}$  is a finite set.

Since  $\pi a$  is in the orbit of a',  $\pi$  factors through some  $\tau \in \Lambda_{a,a'}$ . So  $(l, \pi f, \sigma g) = (\gamma m, \alpha \tau_1 f, \beta \tau_2 g)$  for some  $\gamma, \alpha, \beta \in \text{Inc}(\mathbb{N}), \tau_1 \in \Lambda_{a,a'}$  and  $\tau_2 \in \Lambda_{b,b'}$ . Therefore

$$T_{f,g} \subseteq \bigcup_{(m,f',g') \in U_{f,g}} \Pi m \times \Pi f' \times \Pi g'$$

where

$$U_{f,g} = \bigcup_{(m,a',b')\in T_{a,b}} \{ (m,\tau_1 f, \tau_2 g) \mid \tau_1 \in \Lambda_{a,a'}, \tau_2 \in \Lambda_{b,b'} \}.$$

For each (m, f', g'), the set  $\Pi m \times \Pi f' \times \Pi g'$  is the union of a finite number of  $\operatorname{Inc}(\mathbb{N})$ -orbits. To prove this one can follow closely the proof of [BD11] Lemma 3.4. From the finite set of generators we select only those  $(\gamma m, \alpha f', \beta g')$  with  $\gamma m \in \operatorname{lcm}(\alpha f', \beta g')$ , and call this set  $O_{(m, f', g')}$ . Then

$$O_{f,g} = \bigcup_{(m,f',g') \in U_{f,g}} O_{(m,f',g')}$$

is as desired.

Algorithm 5.5.  $T = \text{TORICIDEAL}(\varphi)$ Require:  $\varphi : R[Y] \to R[Z]$  is a monomial map as in the Main Theorem. Ensure: T is a finite set of generators of ker  $\varphi$  as  $\text{Inc}(\mathbb{N})$ -stable ideal.

<sup>1:</sup> Replace Y by the set of variables  $\{y_{p,J}\}_{p,J}$  as in the proof of Proposition 2.1.

<sup>2:</sup> Decompose  $\varphi$  with the composition of two maps  $\varphi$  and  $\psi$  as in diagram (2).

3: Consider the ideal  $I_{\psi} \subset R[x^A \mid A \text{ good}][Z]$  generated by the finite set F of binomials  $\psi(x^A) - x^A$ , where  $A \in \prod_{p \in [N]} \mathbb{N}^{[k_p] \times \mathbb{N}}$  is good of multi-degree

 $d \in \{(1, 0, \dots, 0), \dots, (0, 0, 0, \dots, 1)\} \subseteq \mathbb{N}^{[N]}$ 

and  $\leq$ -minimal; the Inc( $\mathbb{N}$ )-orbits of such monomials  $x^A$  generate  $R[x^A \mid A \text{ good}]$ . 4: Run Algorithm 5.2 for the input F with respect to a monomial order that eleminates the variables Z. Since  $R[x^A \mid A \text{ good}][Z]$  is the monoid algebra

of a monoid where  $\text{Inc}(\mathbb{N})$ -divisibility is a w.p.o., the algorithm terminates. Standard elimination theory implies that  $G' = G \cap R[x^A \mid A \text{ good}]$  generates

$$I_{\psi} \cap R[x^A \mid A \text{ good}] = \ker \psi \cap \operatorname{im} \varphi.$$

5: Let T consist of preimages of elements in G' (one per element) and a finite number of binomials whose orbits generating ker  $\varphi$  (see Corollary 3.4).

**Remark 5.6.** We can execute Algorithm 5.5 for any coefficient ring R (not necessarily a field), since all polynomials that appear in computation are binomials with coefficients  $\pm 1$ .

In the following two remarks we comment on two major subroutines not spelled out in the sketch of the algorithm above.

**Remark 5.7.** Unlike in the usual Buchberger algorithm, the task of computing S-polynomials in Algorithm 5.2 is far from being trivial. To accomplish that, one needs to compute the set  $O_{f,g}$ , which can be done following the lines of the proof of Proposition 5.4. While this procedure is *effective*, by no means it is *efficient*.

**Remark 5.8.** In the last step of Algorithm 5.5 a preimage  $\varphi^{-1}(g)$  of an element  $g \in G$  can be computed by reducing the problem to one of computing maximal matchings of bipartite graphs, a well studied problem in combinatorics. Any monomial  $x^A \in \operatorname{im} \varphi$  can be considered as a collection of N bipartite graphs with adjacency matrices  $A_0, \ldots, A_{N-1}$  as in Section 3, where each  $A_p$  has bipartition  $[k_p] \sqcup \mathbb{N}$ . Fixing  $A_p$ , let  $S \subset \mathbb{N}$  be the set of vertices in the second partition with degree  $d_{A_p}$  (i.e. the indices of the columns of  $A_p$  with column sum equal to  $d_{A_p}$ ). A matching B covering  $[k_p]$  and S can be computed using the Hungarian method or other algorithms for computing weighted bipartite matchings (see [Sch03] Chapter 17 for more details). The matching B directly corresponds to a variable  $y_{p,J} \in Y$  with  $\varphi(y_{p,J}) = x^B$ . Since B covers S, it follows that  $A_p - B$  is a good matrix. Therefore  $x^{A_p}/\varphi(y_{p,J})$  is also in im  $\varphi$  and can be decomposed further by repeating the process.

Algorithm 5.5 has an important theoretical consequence, a solution to Problem 5.3(a): a finite  $Inc(\mathbb{N})$ -generating set of the toric ideals in the Main Theorem is computable. However, in view of Remark 5.7 and a more elementary procedure (albeit with no termination guarantee) given in the following section that solves a harder Problem 5.3(b) for a small example, we postpone a practical implementation of Algorithm 5.5.

### 6. An example, and a more naïve implementation

A more elementary approach to Problem 5.3—indeed, to the hardest variant is, for a given order on [Y, Z], to directly apply the algorithm of [BD11] to the graph of the entire map  $\psi \circ \varphi$ , rather than computing generators for the kernels

of  $\psi$  and  $\varphi$  separately as in Algorithm 5.5. The advantages of this approach are that it is simpler to implement, and that it produces not just a generating set, but an  $\text{Inc}(\mathbb{N})$ -equivariant Gröbner basis. The disadvantage is that we do not know whether the procedure is guaranteed to terminate. We now set up a version of the usual equivariant Buchberger algorithm that is particularly easy to implement, and conclude with one nontrivial computational example.

For convenience let  $\omega = \psi \circ \varphi$ . Let  $I_{\omega} \subset R[Y, Z]$  be the ideal corresponding to the graph of  $\omega$ , so  $I_{\omega}$  is generated by the binomials of the form  $y - \omega(y)$  for each variable  $y \in Y$ . Choosing a representative  $y_p = y_{p,(0,\ldots,k_p-1)}$  of each Sym(N)-orbit in Y, the ideal is Inc(N)-generated by the finite set

$$F := \{ \sigma y_p - \omega(\sigma y_p) \mid p \in [N], \ \sigma \in \operatorname{Sym}([k_p]) \}.$$

Choose an  $\operatorname{Inc}(\mathbb{N})$ -compatible monomial order  $\leq$  on R[Y, Z] that eliminates Z. Then apply to F the equivariant Gröbner basis algorithm from [BD11] (which is essentially Algorithm 5.2). Note that since we are working in a polynomial ring R[Y, Z], rather than a more complicated monoid ring  $R[X \mid X \text{ good}][Z]$ , every pair of monomials has only one lcm, which is is straightforward to compute. If the procedure terminates with output G, then  $G \cap R[Y]$  is an  $\operatorname{Inc}(\mathbb{N})$ -equivariant Gröbner basis of  $I_{\omega} \cap R[Y] = \ker \omega$ .

This procedure can be adapted to make use of existing, fast implementations of traditional Gröbner basis algorithms. For each  $n \in \mathbb{N}$  truncate to the first n index values by defining  $Y_n := \{y_{p,J} \mid J \in [n]^{k_p}\}, Z_n := \{z_{i,j} \in Z \mid j \in [n]\}$ , and  $F_n := \{y - \omega(y) \mid y \in Y_n\}$ . Let  $I_n$  be the ideal in  $R[Y_n, Z_n]$  generated by  $F_n$ . Note that each  $I_n$  is Sym([n])-stable and that  $\bigcup_{n \in \mathbb{N}} I_n = I_{\omega}$ . Let  $\operatorname{Inc}(m, n)$  denote the set of all strictly increasing maps  $[m] \to [n]$ , and equip  $K[Y_n, Z_n]$  with the restriction of the Inc( $\mathbb{N}$ )-monomial order  $\leq$ .

Algorithm 6.1.  $G = \text{TRUNCATEDBUCHBERGER}(\omega)$ 

**Require:**  $\varphi : R[Y] \to R[Z]$  is a monomial map in the Main Theorem. **Ensure:** G is an Inc(N)-equivariant Gröbner basis of ker  $\varphi$ .

 $\begin{array}{l} n \leftarrow \max_{p \in [N]} k_p \\ \textbf{while true do} \\ F_n \leftarrow \{y - \omega(y) \mid y \in Y_n\} \\ G_n \leftarrow \text{GRÖBNERBASIS}(F_n) \\ m \leftarrow \lfloor (n+1)/2 \rfloor \\ \textbf{if } m \ge \max_{p \in [N]} k_p \text{ and } G_n = \text{Inc}(m,n)G_m \textbf{ then} \\ G \leftarrow G_m \cap R[Y] \\ \text{return } G \\ \textbf{end if} \\ n \leftarrow n+1 \\ \textbf{end while} \end{array}$ 

Here GRÖBNERBASIS denotes any algorithm to compute a traditional Gröbner basis. If TRUNCATEDBUCHBERGER( $\omega$ ) terminates, this implies that there is some  $m \geq \max_{p \in [N]} k_p$  such that  $\operatorname{Inc}(m, n)G_m$  satisfies Buchberger's criterion for some  $n \geq 2m - 1$ . Then  $G_m$  satisfies the equivariant Buchberger criterion, so  $G_m$  is an equivariant Gröbner basis. Because we require that  $m \geq \max_{p \in [N]} k_p$ , the set  $G_m$ generates  $I_{\omega}$  up to  $\operatorname{Inc}(\mathbb{N})$ -action. Finally  $G = G_m \cap R[Y]$  is an equivariant Gröbner basis for ker  $\omega$ .

 $\begin{array}{c|c} \hline & & \\ \hline & & \\ \triangleleft y_{1,2}y_{0,1}^2 - y_{1,0}^2y_{0,2} \triangleright \\ \triangleleft y_{2,0}y_{0,1}^2 - y_{1,0}y_{0,2}^2 \triangleright \end{array}$ degree 2  $\triangleleft y_{1,3}y_{0,2} - y_{1,2}y_{0,3} \triangleright$  $\triangleleft y_{2,0}y_{1,0} - y_{1,2}y_{0,2} \triangleright$  $y_{2,1}y_{0,2}^2 - y_{2,0}^2y_{0,1}$  $y_{2,1}y_{0,1} - y_{1,2}y_{0,2}$  $y_{2,3}y_{0,1} - y_{2,1}y_{0,3}$  $y_{2,3}y_{1,0} - y_{2,0}y_{1,3}$  $y_{3,1}y_{2,0} - y_{3,0}y_{2,1}$  $y_{3,2}y_{0,1} - y_{3,1}y_{0,2}$  $\begin{array}{c} y_{2,1}y_{1,0}y_{0,3} - y_{2,0}y_{1,3}y_{0,1} \\ y_{2,1}^2y_{0,3} - y_{2,0}^2y_{1,3} \\ y_{2,3}y_{1,2}y_{0,2} - y_{2,0}^2y_{1,3} \end{array}$  $y_{3,2}y_{1,0} - y_{3,0}y_{1,2}$ degree 4  $\begin{array}{c} y_{3,0}y_{1,2}y_{0,2} - y_{2,0}y_{1,3}y_{0,3} \\ y_{3,0}y_{1,2}^2 - y_{2,0}y_{1,3}^2 \\ y_{3,0}y_{2,1}^2 - y_{2,3}y_{2,0}y_{1,3} \\ y_{3,0}y_{2,1}^2 - y_{2,3}y_{2,0}y_{1,3} \\ y_{3,1}y_{0,2}^2 - y_{2,1}y_{0,3}^2 \end{array}$  $\begin{array}{c} y_{2,1}y_{1,2}y_{0,3}y_{0,2}-\overrightarrow{y_{2,0}^{2}}y_{1,3}y_{0,1}\\ y_{3,1}y_{2,3}y_{1,3}y_{0,4}-\overrightarrow{y_{3,0}^{2}}y_{2,1}y_{1,4}\end{array}$  $y_{3,1}y_{2,3}^2y_{0,4} - y_{3,0}^2y_{2,4}y_{2,1}$  $y_{3,2}y_{2,3}y_{1,3}y_{0,4} - y_{3,0}^2y_{2,4}y_{1,2}$  $y_{3,1}y_{1,0}y_{0,2} - y_{3,0}y_{1,2}y_{0,1}$  $\begin{array}{l} y_{4,1}y_{2,3}y_{1,4}y_{0,4}-y_{4,0}^2y_{2,1}y_{1,3}\\ y_{4,1}y_{3,2}y_{1,4}y_{0,4}-y_{4,0}^2y_{3,1}y_{1,2}\end{array}$  $y_{3,1}y_{1,2}y_{0,2} - y_{2,1}y_{1,3}y_{0,3}$  $\begin{array}{c} y_{3,1}y_{2,3}y_{0,3} - y_{3,0}^2y_{2,1} \\ y_{3,1}^2y_{0,2} - y_{3,0}^2y_{1,2} \end{array}$  $y_{4,1}y_{3,4}y_{2,4}y_{0,5} - y_{4,0}^2y_{3,1}y_{2,5}$  $y_{3,2}y_{1,3}y_{0,3} - y_{3,0}^2y_{1,2}$ degree 5  $y_{3,2}y_{2,0}y_{1,3} - y_{3,0}y_{2,3}y_{1,2}$  $y_{3,2}y_{2,0}y_{1,4} - y_{3,0}y_{2,4}y_{1,2}$  $y_{3,2}y_{2,1}y_{0,3} - y_{3,1}y_{2,3}y_{0,2}$  $y_{3,2}y_{2,1}y_{0,4} - y_{3,1}y_{2,4}y_{0,2}$  $y_{4,0}y_{2,3}y_{1,3} - y_{3,0}y_{2,4}y_{1,4}$  $y_{4,1}y_{2,3}y_{0,3} - y_{3,1}y_{2,4}y_{0,4}$  $y_{4,2}y_{1,3}y_{0,3} - y_{3,2}y_{1,4}y_{0,4}$  $y_{4,2}y_{2,0}y_{1,3} - y_{4,0}y_{2,3}y_{1,2}$  $y_{4,2}y_{2,1}y_{0,3} - y_{4,1}y_{2,3}y_{0,2}$ 

FIGURE 3. An Inc( $\mathbb{N}$ )-equivariant Gröbner basis for ker  $\omega$  in Example 6.2. The five highlighted binomials form a Sym( $\mathbb{N}$ )-equivariant Markov basis according to [KKL14].

**Example 6.2.** Set  $Y := \{y_{j_0,j_1} \mid j_0, j_1 \in \mathbb{N}, j_0 \neq j_1\}$  and  $Z := \{z_i \mid i \in \mathbb{N}\}$ , each consisting of a single Sym( $\mathbb{N}$ )-orbit, and define the monomial map  $\omega : R[Y] \to R[Z]$  by

$$\omega: y_{j_0,j_1} \mapsto z_{j_0}^2 z_{j_1}.$$

Whether ker  $\omega$  is finitely generated was posed as an open question in [HdC13] (Remark 1.6). This is answered in the affirmative by Theorem 1.1, but by applying Algorithm 6.1 we have also explicitly computed an Inc(N)-equivariant Gröbner basis. The Gröbner basis computations were carried out using the software package 4ti2 [HHM08], which features algorithms specifically designed for computing Gröbner bases of toric ideals. The monomial order on Y is lexicographic, where variables are ordered by  $y_{i,j} < y_{i',j'}$  if i < i', or i = i' and j < j'.

The result displayed in Figure 3 consists of 51 generators with indices from  $\{0, 1, 2, 3, 4, 5\}$  and degrees up to 5. Note that a minimal generating set resulting

from a study of the family of equivariant toric maps of the form

$$y_{ij} \mapsto z_i^a z_j^b, \quad i, j \in \mathbb{N}, i \neq j,$$

for fixed  $a, b \in \mathbb{N}$  in [KKL14] is much smaller.

**Remark 6.3.** As pointed out in the Introduction, the technique laid out in this article does not settle the question whether the finite generatedness of ker  $\varphi$  in the Main Theorem persists when  $\text{Inc}(\mathbb{N})$  acts with finitely many orbits on Y and the monomial map  $\varphi$  is required to be merely  $\text{Inc}(\mathbb{N})$ -equivariant (though we do know that im  $\varphi$  needs not be  $\text{Inc}(\mathbb{N})$ -Noetherian in this case).

However, a naïve elimination procedure terminates, for instance, for the  $\text{Inc}(\mathbb{N})$ analogue of Example 6.2, i.e., for the same map, but with the smaller set of variables

$$Y := \{ y_{j_0, j_1} \mid j_0, j_1 \in \mathbb{N}, \ j_0 > j_1 \}$$

A computation that can be carried out with EquivariantGB [Kro] produces a finite number of generators of the kernel:

$$\{y_{3,1}y_{2,0} - y_{3,0}y_{2,1}, y_{3,2}^2y_{1,0} - y_{3,1}y_{3,0}y_{2,1}, y_{4,2}y_{3,2}y_{1,0} - y_{4,0}y_{3,1}y_{2,1}\}.$$

### References

- [AH07] Matthias Aschenbrenner and Christopher J. Hillar. Finite generation of symmetric ideals. Trans. Am. Math. Soc., 359(11):5171–5192, 2007.
- [AR08] Elizabeth S. Allman and John A. Rhodes. Phylogenetic ideals and varieties for the general Markov model. *Adv. Appl. Math.*, 40(2):127–148, 2008.
- [BD11] Andries E. Brouwer and Jan Draisma. Equivariant Gröbner bases and the two-factor model. Math. Comput., 80:1123–1133, 2011.
- [Coh87] Daniel E. Cohen. Closure relations, Buchberger's algorithm, and polynomials in infinitely many variables. In *Computation theory and logic*, volume 270 of *Lect. Notes Comput. Sci.*, pages 78–87, 1987.
- [DE15] Jan Draisma and Rob H. Eggermont. Finiteness results for Abelian tree models. J. Eur. Math. Soc., 17:711–738, 2015.
- [Dic13] Leonard E. Dickson. Finiteness of the odd perfect and primitive abundant numbers with n distinct prime factors. Amer. J. Math., 35(4):413–422, 1913.
- [DK09] Jan Draisma and Jochen Kuttler. On the ideals of equivariant tree models. *Math. Ann.*, 344(3):619–644, 2009.
- [DK14] Jan Draisma and Jochen Kuttler. Bounded-rank tensors are defined in bounded degree. Duke Math. J., 163(1):35–63, 2014.
- [dLST95] Jesús A. de Loera, Bernd Sturmfels, and Rekha R. Thomas. Gröbner bases and triangulations of the second hypersimplex. *Combinatorica*, 15:409–424, 1995.
- [Dra10] Jan Draisma. Finiteness for the k-factor model and chirality varieties. Adv. Math., 223:243–256, 2010.
- [GP13] Elizabeth Gross and Sonja Petrović. Combinatorial degree bound for toric ideals of hypergraphs. Int. J. Algebra Comput., 23(6):1503–1520, 2013.
- [GS02] Daniel R Grayson and Michael E Stillman. Macaulay 2, a software system for research in algebraic geometry. Available at http://www.math.illinois.edu/Macaulay2/, 2002.
- [HdC13] Christopher J. Hillar and Abraham Martín del Campo. Finiteness theorems and algorithms for permutation invariant chains of Laurent lattice ideals. J. Symb. Comput., 50:314–334, 2013.
- [HdTY14] David Haws, Abraham Martín del Campo, Akimichi Takemura, and Ruriko Yoshida. Markov degree of the three-state toric homogeneous Markov chain model. *Beitr. Algebra Geom.*, 55(1):161–188, 2014.
- [HHM08] Raymond Hemmecke, Ralf Hemmecke, and Peter Malkin. 4ti2 version 1.3.2– computation of hilbert bases, graver bases, toric gröbner bases, and more, 2008.
- [Hig52] Graham Higman. Ordering by divisibility in abstract algebras. Proc. Lond. Math. Soc., III. Ser., 2:326–336, 1952.

- [HS12] Christopher J. Hillar and Seth Sullivant. Finite groebner bases in infinite dimensional polynomial rings and applications. *Advances in Mathematics*, 229(1):1–25, 2012.
- [KKL14] Thomas Kahle, Robert Krone, and Anton Leykin. Equivariant lattice generators and markov bases. In International Symposium on Symbolic and Algebraic Computation, 2014.
- [KR13] Franz J. Kiraly and Zvi Rosen. Symmetries and finiteness for algebraic matroids. poster at MEGA 2013, 2013.
- [Kro] Robert Krone. Equivariantgb: equivariant gröbner bases in macaulay2. Available at http://people.math.gatech.edu/~rkrone3/EquivariantGB.html.
- [Kru72] Joseph B Kruskal. The theory of well-quasi-ordering: A frequently discovered concept. Journal of Combinatorial Theory, Series A, 13(3):297–305, 1972.
- [Nor15] Patrik Norén. The three-state toric homogeneous Markov chain model has Markov degree two. J. Symb. Comput., 68:285–296, 2015.
- [NW63] C Nash-Williams. On well-quasi-ordering finite trees. Mathematical Proceedings of the Cambridge Philosophical Society, 59(04):833–835, 1963.
- [OH01] Hidefumi Ohsugi and Takayuki Hibi. Convex polytopes all of whose reverse lexicographic initial ideals are squarefree. *Proc. Am. Math. Soc.*, 129(9):2541–2546, 2001.
- [PS05] Lior Pachter and Bernd Sturmfels, editors. Algebraic Statistics for Computational Biology, Cambridge, 2005. Cambridge University Press.
- [Sch03] Alexander Schrijver. Combinatorial optimization: polyhedra and efficiency, volume 24. Springer Verlag, 2003.
- [Sno13] Andrew Snowden. Syzygies of Segre embeddings and  $\Delta$ -modules. Duke Math. J., 162(2):225–277, 2013.
- [Sul06] Seth Sullivant. Compressed polytopes and statistical disclosure limitation. Tohoku Math. J. (2), 58(3):433-445, 2006.
- [YOT14] Takashi Yamaguchi, Mitsunori Ogawa, and Akimichi Takemura. Markov degree of the Birkhoff model. J. Algebr. Comb., 40(1):293–311, 2014.

(Jan Draisma) DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, TECHNISCHE UNI-VERSITEIT EINDHOVEN, P.O. BOX 513, 5600 MB EINDHOVEN, THE NETHERLANDS, AND VRIJE UNIVERSITEIT AND CENTRUM VOOR WISKUNDE EN INFORMATICA, AMSTERDAM, THE NETHERLANDS *E-mail address*: j.draisma@tue.nl

(Rob H. Eggermont) DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, TECHNISCHE UNIVERSITEIT EINDHOVEN, P.O. BOX 513, 5600 MB EINDHOVEN, THE NETHERLANDS *E-mail address*: r.h.eggermont@tue.nl

(Robert Krone) School of Mathematics, Georgia Institute of Technology, 686 Cherry Street, Atlanta, GA 30332-0160 USA

E-mail address: krone@math.gatech.edu

(Anton Leykin) School of Mathematics, Georgia Institute of Technology, 686 Cherry Street, Atlanta, GA 30332-0160 USA

E-mail address: leykin@math.gatech.edu