Expansion Trees with Cut

Stefan Hetzl and Daniel Weller

April 15, 2013

Abstract

Herbrand's theorem is one of the most fundamental insights in logic. From the syntactic point of view it suggests a compact representation of proofs in classical first- and higher-order logic by recording the information which instances have been chosen for which quantifiers, known in the literature as expansion trees.

Such a representation is inherently analytic and hence corresponds to a cut-free sequent calculus proof. Recently several extensions of such proof representations to proofs with cut have been proposed. These extensions are based on graphical formalisms similar to proof nets and are limited to prenex formulas.

In this paper we present a new approach that directly extends expansion trees by cuts and covers also non-prenex formulas. We describe a cut-elimination procedure for our expansion trees with cut that is based on the natural reduction steps. We prove that it is weakly normalizing using methods from the ε -calculus.

1 Introduction

Herbrand's theorem [16, 8], one of the most fundamental insights of logic, characterizes the validity of a formula in classical first-order logic by the existence of a propositional tautology composed of instances of that formula.

From the syntactic point of view this theorem induces a way of describing proofs: by recording which instances have been picked for which quantifiers we obtain a description of a proof up to its propositional part, a part we often want to abstract from. An example for a formalism that carries out this abstraction are Herbrand proofs [8]. This generalizes nicely to most classical systems with quantifiers, for example to simple type theory as in the expansion tree proofs of [23]. Such formalisms are compact and useful proof certificates in many situations; they are for example produced naturally by methods of automated deduction such as instantiation-based reasoning [21].

These formalisms consider only instances of the formula that has been proved and hence are *analytic* proof formalisms (corresponding to cut-free proofs in the sequent calculus). Considering an expansion tree to be a compact representation of a proof, it is thus natural to ask about the possibility of extending this kind of representation to *non-analytic* proofs (corresponding to proofs with cut in the sequent calculus).

In addition to enlarging the scope of instance-based proof representations, the addition of cuts to expansion trees also sheds more light on the computational content of classical logic. This is a central topic of proof theory and has therefore attracted considerable attention, see [25, 13, 12], [6], [27, 28], [7], [20], or [5], for different investigations in this direction and [1] for a survey covering classical arithmetic.

Two instance-based proof formalisms incorporating a notion of cut have recently been proposed: proof forests [15] and Herbrand nets [22]. While proof forests are motivated by the game semantics for classical arithmetic of [11], Herbrand nets are based on methods for proof nets [14]. These two formalisms share a number of properties: both of them work in a graphical notation for proofs, both work on prenex formulas only, for both weak but no strong normalization results are known.

In this paper we present a new approach which works directly in the formalism of expansion tree proofs and hence naturally extends the existing literature in this tradition. As [15, 22] we define a cut-elimination procedure and prove it weakly normalizing but in contrast to [15, 22] we also treat non-prenex formulas, therefore avoiding the distortion of the intuitive meaning of a formula by prenexification.

We describe expansion trees with cuts for non-prenex end-sequents and cuts, including their correctness criterion and how to translate from and to sequent calculus. We describe natural cut-reduction steps and show that they are weakly normalizing. A technical key for proving weak normalization is to use methods of Hilbert's ε -calculus which is a formalism for representing non-analytic firstorder proofs modulo propositional logic. The reader is invited to compare our treatment, in particular the termination measure, with the proof of the first ε -theorem in [19], see [24] for an exposition in English.

2 Expansion Trees

In this whole paper we work with classical first-order logic. Formulas and terms are defined as usual. In order to simplify the exposition, we restrict our attention to formulas in negation normal form (NNF). Mutatis mutandis all notions and results of this paper generalize to arbitrary formulas. We write \overline{A} for the de Morgan dual of a formula A. A *literal* is an atom $P(t_1, \ldots, t_n)$ or a negated atom $\overline{P}(t_1, \ldots, t_n)$.

Definition 1. Expansion trees and a function $Sh(\cdot)$ (for *shallow*) that maps an expansion tree to a formula are defined inductively as follows:

- 1. A literal L is an expansion tree with Sh(L) = L.
- 2. If E_1 and E_2 are expansion trees and $\circ \in \{\land,\lor\}$, then $E_1 \circ E_2$ is an expansion tree with $\operatorname{Sh}(E_1 \circ E_2) = \operatorname{Sh}(E_1) \circ \operatorname{Sh}(E_2)$.
- 3. If $\{t_1, \ldots, t_n\}$ is a set of terms and E_1, \ldots, E_n are expansion trees with $\operatorname{Sh}(E_i) = A[x \setminus t_i]$ for $i = 1, \ldots, n$, then $E = \exists x A + t_1 E_1 \cdots + t_n E_n$ is an expansion tree with $\operatorname{Sh}(E) = \exists x A$.
- 4. If E_0 is an expansion tree with $\operatorname{Sh}(E_0) = A[x \setminus y]$, then $E = \forall x A + {}^y E_0$ is an expansion tree with $\operatorname{Sh}(E) = \forall x A$.

The $+^{t_i}$ are called \exists -expansions and the $+^{\alpha} \forall$ -expansions, and both \forall - and \exists -expansions are called expansions. The variable y of a \forall -expansion $+^y$ is called

eigenvariable of this expansion. We say that $+^{t_i}$ dominates all the expansions in E_i . Similarly, $+^{\alpha}$ dominates all the expansions in E_0 .

Definition 2. We define the function $Dp(\cdot)$ (for *deep*) that maps an expansion tree to a formula as follows:

$$Dp(L) = L \text{ for a literal } L,$$

$$Dp(E_1 \circ E_2) = Dp(E_1) \circ Dp(E_2) \text{ for } \circ \in \{\land, \lor\},$$

$$Dp(\exists x A + {}^{t_1} E_1 \dots + {}^{t_n} E_n) = \bigvee_{i=1}^n Dp(E_i), \text{ and}$$

$$Dp(\forall x A + {}^y E_0) = Dp(E_0).$$

We also say that E is an expansion tree of Sh(E).

Definition 3. A *cut* is a set $C = \{E_1, E_2\}$ of two expansion trees s.t. $Sh(E_1) = Sh(E_2)$.

A formula is called positive if its top connective is \lor or \exists or a positive literal. An expansion tree E is called positive if $\operatorname{Sh}(E)$ is positive. It will sometimes be useful to consider a cut as an ordered pair: to that aim we will write a cut as $C = (E_1, E_2)$ with parentheses instead of curly braces with the convention that E_1 is the positive expansion tree. For a cut $C = (E_1, E_2)$, we define $\operatorname{Sh}(C) = \operatorname{Sh}(E_1)$ which is also called *cut-formula* of C. We define $\operatorname{Dp}(C) = \operatorname{Dp}(E_1) \land \operatorname{Dp}(E_2)$

Definition 4. Let C be a set of cuts with pairwise different cut-formulas and let \mathcal{E} be a set of expansion trees of pairwise different formulas. Then $\mathcal{P} = C, \mathcal{E}$ is called *expansion pre-proof* if each two \forall -expansions in \mathcal{P} have different eigenvariables (*regularity*), and if Sh(\mathcal{P}) does not contain free variables.

For an expansion pre-proof $\mathcal{P} = \mathcal{C}, \mathcal{E}$ we define $\operatorname{Sh}(\mathcal{P}) = \operatorname{Sh}(\mathcal{E})$, which corresponds to the end-sequent of a sequent calculus proof, and $\operatorname{Dp}(\mathcal{P}) =$ $\operatorname{Dp}(\mathcal{E}), \operatorname{Dp}(\mathcal{C})$ (which is a sequent of quantifier-free formulas). For an eigenvariable α in \mathcal{P} , define $q(\alpha)$ to be the \forall -expansion whose eigenvariable it is.

Example 1. Consider the straightforward proof of $P(a) \to \exists z Q(z) \text{ from } \exists y \forall x (P(x) \to Q(f(y)))$ via a cut on $\forall x \exists y (P(x) \to Q(f(y)))$. In negation normal formal these formulas are $\overline{P}(a) \lor \exists z Q(z), \exists y \forall x (\overline{P}(x) \lor Q(f(y)))$, and $\forall x \exists y (\overline{P}(x) \lor Q(f(y)))$. The proof will be represented by the expansion pre-proof $\mathcal{P} = \{E^+, E^-\}, E_1, E_2$ where

$$\begin{split} E^{+} &= \exists x \forall y \left(P(x) \land \overline{Q}(f(y)) \right) +^{a} \left(\forall y \left(P(a) \land \overline{Q}(f(y)) \right) +^{\gamma} P(a) \land \overline{Q}(f(\gamma)) \right) \\ E^{-} &= \forall x \exists y \left(\overline{P}(x) \lor Q(f(y)) \right) +^{\beta} \left(\exists y \left(\overline{P}(\beta) \lor Q(f(y)) \right) +^{\alpha} \left(\overline{P}(\beta) \lor Q(f(\alpha)) \right) \right) \\ E_{1} &= \forall y \exists x \left(P(x) \land \overline{Q}(f(y)) \right) +^{\alpha} \left(\exists x \left(P(x) \land \overline{Q}(f(\alpha)) \right) +^{\beta} P(\beta) \land \overline{Q}(f(\alpha)) \right) \\ E_{2} &= \overline{P}(a) \lor \left(\exists z Q(z) +^{f(\gamma)} Q(f(\gamma)) \right) \end{split}$$

We have $\operatorname{Sh}(\mathcal{P}) = \operatorname{Sh}(E_1, E_2) = \forall y \exists x (P(x) \land \overline{Q}(f(y))), \overline{P}(a) \lor \exists z Q(z) \text{ and}$

$$Dp(\mathcal{P}) = Dp(E^+) \wedge Dp(E^-), Dp(E_1), Dp(E_2)$$

= $(P(a) \wedge \overline{Q}(f(\gamma))) \wedge (\overline{P}(\beta) \vee Q(f(\alpha))), P(\beta) \wedge \overline{Q}(f(\alpha)), \overline{P}(a) \vee Q(f(\gamma)))$

As in [15, 22] it would also be possible in our setting to use a graphical notation. However, we refrain from doing so in order to avoid the parallel use of two different notations: a graphical for examples and a more abstract notation for carrying out proofs.

Let us now move on to isolating the proofs in the set of pre-proofs. The correctness criterion of expansion tree proofs [23], but also those of proof forests [15] and Herbrand nets [22], has two (main) components: 1. a tautology-condition on one or more quantifier-free formulas and 2. an acyclicity condition on one or more orderings. While the tautology condition of [23] generalizes to the setting of cuts in a straightforward way, the acyclicity condition needs a bit more work: in the setting of cut-free expansion trees it is enough to require the acyclicity of an order on the \exists -expansions. In our setting that includes cuts we also have to speak about the order of cuts (w.r.t. each other and w.r.t. \exists -expansions). To simplify our treatment of this order we also include \forall -expansions. Together this leads to the following inference ordering constraints in expansion proofs.

Definition 5. Let $\mathcal{P} = \mathcal{C}, \mathcal{E}$ be an expansion pre-proof. We will define the *dependency relation* $<_{\mathcal{P}}$, which is a binary relation on the set of expansions and cuts in \mathcal{P} . First, we define the binary relation $<_{\mathcal{P}}^{0}$ (writing $<^{0}$ if \mathcal{P} is clear from the context) as the least relation satisfying (C being a cut in \mathcal{P}):

- 1. $v <^0 w$ if w is an \exists -expansion in \mathcal{P} whose term contains the eigenvariable of the \forall -expansion v
- 2. $v <^0 w$ if v is an expansion in \mathcal{P} that dominates the expansion w
- 3. $C <^0 v$ if v is an expansion in C
- 4. $v < {}^0 C$ if Sh(C) contains the eigenvariable of the \forall -expansion v

 $<_{\mathcal{P}}$ is then defined to be the transitive closure of $<^0$. Again, we write < for $<_{\mathcal{P}}$ if \mathcal{P} is clear from the context.

Definition 6. An *expansion proof* is an expansion pre-proof \mathcal{P} that satisfies the following conditions:

- 1. $<_{\mathcal{P}}$ is acyclic (i.e. $x <_{\mathcal{P}} x$ holds for no x),
- 2. $Dp(\mathcal{P})$ is a tautology.

As there is no cycle containing cuts only, $<_{\mathcal{P}}$ is cyclic iff $w <_{\mathcal{P}} w$ for an expansion w, and we will make use of this property without further mention.

Example 2. Coming back to the expansion pre-proof \mathcal{P} of Example 1, note that $\operatorname{Dp}(\mathcal{P}) = (P(a) \land \overline{Q}(f(\gamma))) \land (\overline{P}(\beta) \lor Q(f(\alpha))), P(\beta) \land \overline{Q}(f(\alpha)), \overline{P}(a) \lor Q(f(\gamma)))$ is a tautology (of the form $A \land B, \overline{B}, \overline{A}$). Let us now consider the theory induced by \mathcal{P} : in \mathcal{P} each term belongs to at most one \exists - and at most one \forall -expansion In such a situation we can uniformly notate all expansions as Qt for some term t and $Q \in \{\exists, \forall\}$. The expansions of \mathcal{P} are then written as $\exists a, \forall \gamma, \forall \beta, \exists \alpha, \forall \alpha, \exists \beta, \text{ and } \exists f(\gamma)$. Furthermore, \mathcal{P} contains a single cut C. Then $<^0$ is exactly:

- 1. $\forall \gamma <^0 \exists f(\gamma), \forall \beta <^0 \exists \beta, \forall \alpha <^0 \exists \alpha,$
- 2. $\exists a <^0 \forall \gamma, \forall \beta <^0 \exists \alpha, \forall \alpha <^0 \exists \beta,$

3. $C <^0 \exists a, C <^0 \forall \gamma, C <^0 \forall \beta, C <^0 \exists \alpha,$

4. there is no $v <^0 C$ as the cut formula of C is variable-free.

As the reader is invited to verify, < is acyclic.

3 Basic Operations on Expansion Proofs

Our cut-elimination algorithm, described in Section 5, will be based on natural rewrite rules of expansion proofs. In order to fully specify those, we first need to clarify some basic operations on expansion proofs.

3.1 Expansion Trees with Merges

One on these basic operations is the *merge* of expansion pre-proofs. If we have two expansion pre-proofs E_1 and E_2 with $\text{Sh}(E_1) = \text{Sh}(E_2)$ we want to define a new expansion pre-proof $E_1 \cup E_2$ which merges E_1 and E_2 . For example

$$(\exists x P(x) + {}^{a} P(a)) \cup (\exists x P(x) + {}^{b} P(b)) = \exists x P(x) + {}^{a} P(a) + {}^{b} P(b).$$

In general however, this operation can be considerably more complicated.

Example 3. Consider the following merge operation in an expansion pre-proof:

$$(\forall x A + {}^{u} E_{1}) \cup (\forall x A + {}^{v} E_{2}), \exists x B + {}^{f(u)} F_{1} + {}^{f(v)} F_{2}.$$

When propagating the merge node into the subtrees of the two trees being merged, the two eigenvariables u and v will need to be unified, say by globally applying the substitution $[v \setminus u]$. As eigenvariables are global, the result of this unification is that the two \exists -expansions $+^{f(u)}$ and $+^{f(v)}$ in the expansion tree of $\exists x B$ will also be identified, violating the set-nature of the expansions of an existential formula. Globally applying the substitution $[v \setminus u]$ therefore requires merging the two trees $F_1[v \setminus u]$ and $F_2[v \setminus u]$.

We hence see that carrying out a merge operation does not only induce other merge operations on subtrees but also substitutions and vice versa: carrying out a substitution may induce additional merge operations. In order to give a clear formal definition of these operations we will consider *expansion pre-proofs with merges*: a data structure of expansion pre-proofs which, in addition, contains an object-level merge-operation \sqcup .

Definition 7. An *expansion tree with merges* is defined by the same inductive definition as expansion trees in Definition 1 to which we add the following clause:

5. If E_1 and E_2 are expansion trees with merges s.t. $\operatorname{Sh}(E_1) = \operatorname{Sh}(E_2)$, then $E_1 \sqcup E_2$ is an expansion tree with merges and $\operatorname{Sh}(E_1 \sqcup E_2) = \operatorname{Sh}(E_1) = \operatorname{Sh}(E_2)$.

We also extend $Dp(\cdot)$ to expansion trees with merges by setting $Dp(E_1 \sqcup E_2) = Dp(E_1) \lor Dp(E_2)$. Expansion (pre-)proofs with merges are defined analogously to expansion (pre-)proofs (without merge).

3.2 Substitution

We now develop the definition of substitution via expansion trees with merges indicated in the beginning of this section. In the following, for a formula or term F we denote by V(F) the set of variables free in F. To make sure that the application of a substitution transforms expansion trees (with merges) into expansion trees (with merges) we have to restrict the set of permitted substitutions: a substitution σ can only be applied to an expansion tree (with merges) E if it is a renaming on the eigenvariables of E, more precisely: if $\alpha \in EV(E)$ implies that $\alpha\sigma$ is a variable. Otherwise it would destroy the \forall -expansions. Furthermore, to ensure no cycles are introduced in the dependency relation, we have to impose an additional restriction on the eigenvariables introduced by σ : $\beta \in V(\alpha\sigma)$ implies that for all \exists -expansions w in \mathcal{P} with an expansion term tsuch that $\alpha \in V(t)$, we have $w \notin q(\beta)$. A substitution fulfilling these conditions will be called *admissible for* \mathcal{P} .

Later we will give an operational meaning to the merge by means of a reduction system. This will allow us to define a notion of substitution for expansion tree proofs *without merge*.

Definition 8. Let *E* be an expansion tree with merges and let σ be a substitution.

- 1. For a literal $L, L\sigma$ is defined as for formulas.
- 2. $(E_1 \circ E_2)\sigma = E_1\sigma \circ E_2\sigma$ for $\circ \in \{\land,\lor\}$.
- 3. Let $E = \exists x A + t_1 E_1 \cdots + t_n E_n$, let $\{s_1, \ldots, s_k\}$ be $\{t_1\sigma, \ldots, t_n\sigma\}$ and define

$$E\sigma = \exists x \, A\sigma + {}^{s_1} \bigsqcup_{\substack{i \in \{1, \dots, n\} \\ t_i\sigma = s_1}} E_i \sigma \dots + {}^{s_k} \bigsqcup_{\substack{i \in \{1, \dots, n\} \\ t_i\sigma = s_k}} E_i \sigma \dots$$

4. $(\forall x A + {}^{\alpha} E)\sigma = \forall x A\sigma + {}^{\alpha\sigma} E\sigma.$

5.
$$(E_1 \sqcup E_2)\sigma = E_1\sigma \sqcup E_2\sigma$$
.

For an expansion pre-proof $\mathcal{P} = C_1, \ldots, C_k, E_1, \ldots, E_n$ and a substitution σ s.t. $\alpha \in EV(\mathcal{P})$ implies that $\alpha \sigma$ is a variable we define $\mathcal{P}\sigma = C_1\sigma, \ldots, C_k\sigma, E_1\sigma, \ldots, E_n\sigma$.

To every expansion w in $\mathcal{P}\sigma$ we can naturally associate a non-empty set of predecessors w.r.t. substitution $\operatorname{pred}_s(w)$ in \mathcal{P} (note that $\operatorname{pred}_s(w)$ is always a singleton, except in case 3 of the above definition). As usual in the term rewriting literature, $\mathcal{P}[]$ denotes an expansion pre-proof context, i.e. an expansion pre-proof with a hole and $\mathcal{P}[E]$ denotes the expansion pre-proof obtained from filling this hole with the expansion tree E.

Lemma 1. Let $\mathcal{P} = \mathcal{P}'[E]$ be an expansion proof with merges and σ a substitution admissible for \mathcal{P} . Then $\mathcal{Q} = \mathcal{P}'[E\sigma]$ is an expansion proof with merges, and $\operatorname{Sh}(\mathcal{P}) = \operatorname{Sh}(\mathcal{Q})$.

Proof. The existence of a cycle in Q implies that of one in \mathcal{P} , see Appendix for details.

3.3 Merge

As we have seen in Example 3, carrying out a merge operation may require to identify two eigenvariables globally, i.e. on the level of the expansion preproof. The object-level merge operations are hence executed by the following reduction system which, in addition to local term rewriting, includes global variable renaming.

Definition 9. We define a reduction system $\stackrel{\sqcup}{\mapsto}$ on expansion pre-proofs with merges.

- 1. $\mathcal{P}[L \sqcup L] \xrightarrow{\sqcup} \mathcal{P}[L]$ for a literal L.
- 2. $\mathcal{P}[(E'_1 \circ E''_1) \sqcup (E'_2 \circ E''_2)] \xrightarrow{\sqcup} \mathcal{P}[(E'_1 \sqcup E'_2) \circ (E''_1 \sqcup E''_2)]$ for $\circ \in \{\land, \lor\}$.
- 3. $\mathcal{P}[(\forall x A + \alpha_1 E_1) \sqcup (\forall x A + \alpha_2 E_2)] \xrightarrow{\sqcup} \mathcal{P}[\forall x A + \alpha_1 (E_1 \sqcup E_2)][\alpha_2 \setminus \alpha_1].$
- 4. If $E_1 = \exists x A + {}^{r_1} E_{1,1} \dots + {}^{r_k} E_{1,k} + {}^{s_1} F_1 \dots + {}^{s_l} F_l$ and $E_2 = \exists x A + {}^{r_1} E_{2,1} \dots + {}^{r_k} E_{2,k} + {}^{t_1} G_1 \dots + {}^{t_m} G_m$ where $\{s_1, \dots, s_l\} \cap \{t_1, \dots, t_m\} = \emptyset$, then

$$\mathcal{P}[E_1 \sqcup E_2] \stackrel{\sqcup}{\mapsto} \mathcal{P}[\exists x A + {}^{r_1}(E_{1,1} \sqcup E_{2,1}) \ldots + {}^{r_k}(E_{1,k} \sqcup E_{2,k}) + {}^{s_1}F_1 \ldots + {}^{s_l}F_l + {}^{t_1}G_1 \ldots + {}^{t_m}G_m]$$

Write $\xrightarrow{\sqcup}$ for the reflexive and transitive closure of $\stackrel{\sqcup}{\mapsto}$.

As with substitution, for $\mathcal{P} \stackrel{\sqcup}{\mapsto} \mathcal{P}'$ we associate to every expansion n in \mathcal{P}' a non-empty set $\operatorname{pred}_{\sqcup}^{0}(n)$ of predecessor expansions from \mathcal{P} in the natural way, noting that $\operatorname{pred}_{\sqcup}^{0}(n)$ is a singleton in cases 1, 2 of the definition, and contains at most 2 elements in cases 3, 4. We extend $\operatorname{pred}_{\sqcup}^{0}$ to $\stackrel{\sqcup}{\to}$ by denoting the reflexive and transitive closure of $\operatorname{pred}_{\sqcup}^{0}$ by $\operatorname{pred}_{\sqcup}$.

Lemma 2. The relation $\stackrel{\sqcup}{\rightarrow}$ is confluent and strongly normalizing. Its normal forms have no merge nodes.

Proof. See Appendix.

By $\mathcal{P}\downarrow$ we denote the normal form of \mathcal{P} under $\stackrel{\sqcup}{\rightarrow}$. We now use the above reduction system on object-level merge nodes for defining the actual merge operation on expansion trees without merge nodes.

Definition 10. Let E_1, E_2 be expansion trees with $Sh(E_1) = Sh(E_2)$, then $E_1 \cup E_2$ is defined as $(E_1 \sqcup E_2) \downarrow$.

The merge operation is extended to expansion pre-proofs in the natural way: expansion trees and cuts with the same shallow formula are merged, the others are combined by set-theoretic union, where merging of a cut is defined as follows: for cuts $C_1 = (E_1^+, E_1^-)$ and $C_2 = (E_2^+, E_2^-)$ with $\operatorname{Sh}(C_1) = \operatorname{Sh}(C_2)$ we define $C_1 \cup C_2$ as $(E_1^+ \cup E_2^+, E_1^- \cup E_2^-)$.

Lemma 3. If $\mathcal{P}_1 \sqcup \mathcal{P}_2$ is an expansion proof with merge such that $\operatorname{Sh}(\mathcal{P}_1) = \operatorname{Sh}(\mathcal{P}_2)$, then $\mathcal{P}_1 \cup \mathcal{P}_2$ is an expansion proof and $\operatorname{Sh}(\mathcal{P}_1 \cup \mathcal{P}_2) = \operatorname{Sh}(\mathcal{P}_1) = \operatorname{Sh}(\mathcal{P}_2)$.

Proof. See Appendix.

The role of the merge operation is to recursively identify such variables that denote the same value. For the purpose of cut-elimination, its principal use consists in defining which parts of an expansion tree are to be duplicated by a reduction. It is not surprising that this is technically involved as it is also the case in other comparable formalisms. Indeed, it is maybe in the technical details of *how* the decision what to duplicate is taken where the existing formalisms differ most: in the ε -calculus [19], the object-level syntax of ε -terms ensures maximal identifications, in proof forests [15], the reduction steps duplicate too much and are hence interleaved with pruning steps and in Herbrand nets [22] the notion of kingdom from the literature on proof nets is used for determining what to duplicate.

4 Expansion Proofs and Sequent Calculus

In this section we will clarify the relationship between our expansion proofs and the sequent calculus. The concrete version of sequent calculus is of no significance to the results presented here, they hold mutatis mutandis for every version that is common in the literature. For technical convenience we choose a calculus where a sequent is a set of formulas and all rules are invertible.

Definition 11. The calculus **LK** is defined as follows: initial sequents are of the form Γ , A, \overline{A} for an atom A. The inference rules are

$$\frac{\Gamma, A[x \setminus \alpha]}{\Gamma, \forall x A} \ \forall \quad \frac{\Gamma, \exists x A, A[x \setminus t]}{\Gamma, \exists x A} \ \exists \quad \frac{\Gamma, A \quad \Gamma, B}{\Gamma, A \wedge B} \ \land \quad \frac{\Gamma, A, B}{\Gamma, A \vee B} \ \lor \quad \frac{\Gamma, A \quad \overline{A}, \Gamma}{\Gamma} \ \text{cut}$$

with the usual side conditions: α must not appear in Γ , $\forall x A$ and t must not contain a variable which is bound in A.

Due to the global nature of expansion proofs, they correspond to regular **LK**-proofs. An **LK**-proof is called *regular* if each two \forall -inferences have different eigenvariables. From now on we assume w.l.o.g. that all **LK**-proofs are regular.

4.1 From Sequent Calculus to Expansion Proofs

In this section we describe how to read off expansion trees from **LK**-proofs which leads to a completeness theorem for expansion proofs. For representing a formula A that is introduced by (implicit) weakening we use the natural coercion of A into an expansion tree, denoted by $A^{\rm E}$. For a sequent $\Gamma = A_1, \ldots, A_n$ we define $\Gamma^{\rm E} = A_1^{\rm E}, \ldots, A_n^{\rm E}$.

Definition 12. For an **LK**-proof π define the expansion proof $\text{Exp}(\pi)$ by induction on π :

- 1. If π is an initial sequent Γ, A, \overline{A} , then $\text{Exp}(\pi) = \Gamma^{\text{E}}, A, \overline{A}$
- 2. If $\pi = \frac{(\pi_A)}{\Gamma, A \land B} \land$ with $\operatorname{Exp}(\pi_A) = \mathcal{P}_A, E_A$ and $\operatorname{Exp}(\pi_B) = \mathcal{P}_B, E_B$ where $\operatorname{Sh}(E) \land A$ and $\operatorname{Sh}(E) \land B$, then $\operatorname{Exp}(\pi) = \mathcal{P}_B, E_B$.

where $\operatorname{Sh}(E_A) = A$ and $\operatorname{Sh}(E_B) = B$, then $\operatorname{Exp}(\pi) = \mathcal{P}_A \cup \mathcal{P}_B, E_A \wedge E_B$.

3. If
$$\pi = \frac{(\pi')}{\Gamma, A, B}$$
 with $\operatorname{Exp}(\pi') = \mathcal{P}, E_A, E_B$ where $\operatorname{Sh}(E_A) = A$ and $\operatorname{Sh}(E_B) = B$, then $\operatorname{Exp}(\pi) = \mathcal{P}, E_A \vee E_B$.
 (π_A)

- 4. If $\pi = \frac{\Gamma, A[x \setminus \alpha]}{\Gamma, \forall x A} \forall$ with $\operatorname{Exp}(\pi_A) = \mathcal{P}, E$ where $\operatorname{Sh}(E) = A[x \setminus \alpha]$, then $\operatorname{Exp}(\pi) = \mathcal{P}, \forall x A + {}^{\alpha} E.$
- 5. If $\pi = \frac{(\pi_A)}{\Gamma, \exists x A, A[x \setminus t]} \exists$ with $\operatorname{Exp}(\pi_A) = \mathcal{P}, E, E_t$ where $\operatorname{Sh}(E) = \exists x A$

and
$$\operatorname{Sh}(E_t) = A[x \setminus t]$$
, then $\operatorname{Exp}(\pi) = \mathcal{P}, E \cup \exists x A + {}^t E_t$

6. If $\pi = \frac{(\pi^+)}{\Gamma} \frac{(\pi^-)}{\Gamma}$ for A positive with $\operatorname{Exp}(\pi^+) = \mathcal{P}^+, E^+$ and $\operatorname{Exp}(\pi^-) = \mathcal{P}^-, E^-$ where $\operatorname{Sh}(E^+) = A$ and $\operatorname{Sh}(E^-) = \overline{A}$, then $\operatorname{Exp}(\pi) = (E^+, E^-), \mathcal{P}^+ \cup \mathcal{P}^-$.

Note that the behavior of the above definition of $\text{Exp}(\cdot)$ on binary rules is to merge expansions of both subproofs (including cuts). This is the reason for the relationship between sequent calculus proofs and expansion proofs which on the one hand are strongly connected structurally [9, 10] but at the same time have different complexity [4].

Theorem 1 (completeness). If π is an **LK**-proof of a sequent Γ , then $\text{Exp}(\pi)$ is an expansion proof of Γ . If π is cut-free then so is $\text{Exp}(\pi)$.

Proof. That $\text{Exp}(\pi)$ is an expansion pre-proof follows directly from the definitions as we are dealing with regular **LK**-proofs only. By a straightforward induction on π one shows that $\text{Dp}(\text{Exp}(\pi))$ is a tautology. Acyclicity is also shown inductively by observing that if α is a free variable in the end-sequent of π , then α is not an eigenvariable in $\text{Exp}(\pi)$. This implies that if w is the new expansion introduced in the construction of $\text{Exp}(\pi)$, and v is an old expansion in $\text{Exp}(\pi)$, then $w \ge v$, which in turn yields acyclicity.

4.2 From Expansion Proofs to Sequent Calculus

In this section we show how to construct an **LK**-proof from a given expansion proof. To this aim we introduce a calculus **LKE** that works on expansion preproofs instead of sequents (of formulas) following the treatment in [23].

Definition 13. The axioms of **LKE** are of the form $\mathcal{P}, A, \overline{A}$ for an atom A. The inference rules are

$$\frac{\mathcal{P}, E_0}{\mathcal{P}, \forall x A +^{\alpha} E_0} \forall \frac{\mathcal{P}, \exists x A +^{t_1} E_1 \cdots +^{t_{n-1}} E_{n-1}, E_n}{\mathcal{P}, \exists x A +^{t_1} E_1 \cdots +^{t_n} E_n} \exists \frac{\mathcal{P}, E_1 - \mathcal{P}, E_2}{\mathcal{P}, E_1 \wedge E_2} \land \frac{\mathcal{P}, E_1, E_2}{\mathcal{P}, E_1 \vee E_2} \lor \frac{\mathcal{P}, E_1 - E_2, \mathcal{P}}{\{E_1, E_2\}, \mathcal{P}} \text{ cut}$$

with the following side conditions: $\operatorname{Sh}(E_1) = \overline{\operatorname{Sh}(E_2)}$ for the cut and the eigenvariable condition for \forall : α must not occur in $\operatorname{Sh}(\mathcal{P}, \forall x A + {}^x E_0)$.

The reader is invited to note that $\operatorname{Sh}(\mathcal{P}, \forall x A + {}^{x}E_{0})$ does *not* include the cut formulas of \mathcal{P} , they may – and indeed often have to – contain the eigenvariable α . Furthermore, it should be kept in mind that the expansion terms at the \exists -rule form a *set*, i.e. the above rule allows to take any instance as there is no such thing as a last or rightmost instance. An important feature of the above calculus, which is easily verified, is that if π is an **LKE**-proof, then $\operatorname{Sh}(\pi)$ is an **LK**-proof. In the following proof we describe how to transform expansion proofs to **LK**-proofs.

Theorem 2 (soundness). If \mathcal{P} is an expansion proof of a sequent Γ , then there is an **LK**-proof of Γ . If \mathcal{P} is cut-free, then so is the **LK**-proof.

Proof. It is enough to construct an **LKE**-proof π of \mathcal{P} , as then $\operatorname{Sh}(\pi)$ is a proof of $\operatorname{Sh}(\mathcal{P}) = \Gamma$. The construction will be carried out by induction on the number of nodes in \mathcal{P} .

If $\mathcal{P} = \mathcal{P}', E_1 \lor E_2$ for some \mathcal{P}', E_1 and E_2 , then both \mathcal{P}', E_1, E_2 is a strictly smaller expansion proof. By the induction hypothesis we obtain an **LKE**-proofs π' of \mathcal{P}', E_1, E_2 from which a proof of \mathcal{P} is obtained by an \lor -inference. For $\mathcal{P} = \mathcal{P}', E_1 \land E_2$, proceed analogously.

If there are no top-level conjunctions or disjunctions, then by the acyclicity of $<_{\mathcal{P}}$ there must be a $<_{\mathcal{P}}$ -minimal top-level quantifier or cut. For the case of cut proceed as follows: let $\mathcal{P} = C, \mathcal{P}'$ for some \mathcal{P}' and a $<_{\mathcal{P}}$ -minimal cut $C = \{E_1, E_2\}$. Then both E_1, \mathcal{P}' and E_2, \mathcal{P}' are strictly smaller expansion proofs because $Dp(E_i, \mathcal{P}')$ is a tautology as $Dp(\mathcal{P})$ is one and the orderings are suborderings of \mathcal{P} hence also acyclic. By the induction hypothesis we obtain **LKE**-proofs π_1, π_2 of E_1, \mathcal{P}' and E_2, \mathcal{P}' respectively from which a proof of \mathcal{P} is obtained by a cut.

For the case of the minimal node being a quantifier, proceed analogously. As in the cut-free case the eigenvariable condition of the \forall -rule is ensured by the acyclicity of the dependency relation.

Definition 14. The **LK**-proof constructed in the above proof will be called $Seq(\mathcal{P})$.

5 Cut-Elimination

In this section we define a natural reduction system for expansion proofs whose normal forms are cut-free expansion proofs. We prove weak normalization and discuss the status of other properties such as strong normalization and confluence in comparison to other systems from the literature.

5.1 Cut-Reduction Steps

Before we present our cut-reduction steps, we have to discuss regularity: in contrast to the operations we have defined so far, cut-reduction will *duplicate* sub-proofs, making it necessary to discuss the renaming of variables (as in the case of the sequent calculus). We will carefully indicate, in the case of a duplication, which subtrees should be subjected to a variable renaming, and which variables are to be renamed.

The cut-reduction steps, relating expansion proofs $\mathcal{P}, \mathcal{P}'$ and written $\mathcal{P} \mapsto \mathcal{P}'$, are

$$\{ \exists x \ A + {}^{t_1} E_1 \dots + {}^{t_n} E_n, \forall x \ \bar{A} + {}^{\alpha} E \}, \mathcal{P}$$

$$\mathcal{P} \cup \{ E_1 \vee \dots \vee E_n, E\eta_1[\alpha \backslash t_1] \wedge \dots \wedge E\eta_n[\alpha \backslash t_n] \} \cup \bigcup_{i=1}^n \mathcal{P}\eta_i[\alpha \backslash t_i]$$

$$\{ E_1 \vee E_2, E_1' \wedge E_2' \}, \mathcal{P} \mapsto \{ E_1, E_1' \} \cup \{ E_2, E_2' \} \cup \mathcal{P}$$

$$\{ A, \overline{A} \}, \mathcal{P} \mapsto \mathcal{P} \quad \text{for an atom } A.$$

where η_i are renamings of the eigenvariables of \mathcal{P}, E to fresh variables.

These reduction rules are very natural: an atomic cut is simply removed and a propositional cut is decomposed. The reduction of a quantified cut is, when thinking about cut-elimination in the sequent calculus, intuitively immediately appealing: An existential cut is replaced by a cut on a disjunction of the instances. We emphasize here that due to the eigenvariable condition in the sequent calculus, such a rule cannot directly be stated with such formal clarity and elegance. Note that the rule makes use of the merge operation which, as will become clear in the following sections, will prevent redundancies that would be introduced by using the set-union \cup .

One surprising aspect of the quantifier-reduction rule is the presence of \mathcal{P} , without a substitution applied, on the rhs of the rule: in general, \mathcal{P} will contain α , and one would expect that occurrences of α are redundant (since α is "eliminated" by the rule). The reason why this occurrence of \mathcal{P} must be present is that α is not, in fact, eliminated since some t_i might contain it. This situation occurs, for example, when translating from a regular **LK**-proof where an \exists -quantifier may be instantiated by any term, and we happen to choose an eigenvariable from a different branch of the proof. In the sequent calculus, this situation can in principle be avoided by using a different witness for the \exists -quantifier, but realizing such a renaming in expansion proofs is technically non-trivial due to the global nature of eigenvariables. For simplicity of exposition, we therefore allow this somewhat unnatural situation and leave a more detailed analysis for future work.

Remark 1. We note that this phenomenon also occurs in the proof forests of [15], where it is called *bridge*. There, bridges are dealt with by a *pruning reduction*, and the weak normalization proof of that system depends on this pruning. In our setting, we do not need additional machinery for proving weak normalization (see Section 5.4). Furthermore, the counterexample to strong normalization from [15] also contains a bridge; we investigate (a translation of) this counterexample in Section 5.5 and find that it is not a counterexample for our reduction.

As before, if $\mathcal{P} \mapsto \mathcal{P}'$ we can associate in a natural way (formally, using the pred_{\u0354} and pred_s functions defined before) to every expansion w in \mathcal{P}' a unique predecessor (w.r.t. cut-reduction) in \mathcal{P} . This predecessor is denoted by pred_c(w). Note that pred_c(w) is a single expansion, while pred_{\u0354}(w) is a set of expansions; this is explained by the fact that all expansions in pred_{\u0354}(w) are ,,copies" of pred_c(w).

Lemma 4. If $\mathcal{P} \mapsto \mathcal{P}'$ and \mathcal{P} is an expansion proof, then \mathcal{P}' is an expansion proof. Furthermore, $\operatorname{Sh}(\mathcal{P}) = \operatorname{Sh}(\mathcal{P}')$.

Proof. See Appendix.

Example 4. For the sake of conciseness, we use the notation $E(\alpha)$ for an expansion tree with an indicated variable α , and E(t) for the expansion tree obtained from $E(\alpha)$ by (syntactically) substituting t for α . We will also identify formulas and quantifier-node-free expansion trees. With this in mind, consider the expansion proof $\mathcal{P} = -P0$, $Pf^{4}0$, $E(\alpha)$, $\{C^+, C^-\}$ with

$$\begin{split} E(\alpha) &= \exists x F(x) + {}^{\alpha}F(\alpha) + {}^{f\alpha}F(f\alpha) \\ C^+ &= \exists x G(x) + {}^{0}G(0) + {}^{f^{2}0}G(f^{2}0) \\ C^- &= \forall x \overline{G(x)} + {}^{\alpha}\overline{G(\alpha)}, \end{split}$$

where $F(x) = Px \land \neg Pfx$ and $G(x) = Px \land \neg Pf^2x$. Then, since in this case substitution does not introduce any merge nodes and no eigenvariable renaming is necessary,

$$\mathcal{P} \mapsto \neg P0, Pf^40, E(\alpha) \cup E(0) \cup E(f^{2}0), \{G(0) \lor G(f^{2}0), \overline{G(0)} \land \overline{G(f^{2}0)}\}$$

where the substitutions $[\alpha \setminus 0], [\alpha \setminus f^2 0]$ were applied and

$$E(\alpha) \cup E(0) \cup E(f^{2}0) = \exists x F(x) + {}^{\alpha}F(\alpha) + {}^{f\alpha}F(f\alpha) + {}^{0}F(0) + {}^{f0}F(f0) + {}^{f^{2}0}F(f^{2}0) + {}^{f^{3}0}F(f^{3}0)$$

Finally, this proof reduces to

$$\neg P0, Pf^40, E(\alpha) \cup E(0) \cup E(f^{2}0)$$

by the propositional cut-reduction rules. The reader is invited to verify that tautology-hood of $Dp(\mathcal{P})$ is preserved (the α -instances are redundant in this case). The final expansion proof does not contain any \forall -nodes, so acyclicity of the dependency relation is trivial.

In the sequel, by \rightarrow we denote the reflexive, transitive closure of the mapping \mapsto .

5.2 Complexity Measures

Our next aim is to prove weak normalization of our reduction system \rightarrow . It turns out that the strategy of the proof of the *first* ε -theorem can be applied to expansion trees. For simplicity, we just state the second ε -theorem, which is a consequence of the first: for every proof of an ε -free formula in the ε -calculus, there exists a proof of the same formula in which no ε 's occur. It is known that proofs in the ε -calculus can be translated to **LK**-proofs with cut, and vice-versa. This translation shows us that closed ε -terms correspond to eigenvariables in the sequent calculus, which in turn correspond to \forall -expansions in expansion proofs. Equipped with this observation, we can find suitable versions of the notions of rank and degree which in turn will allow us to prove weak normalization. In fact, these notions can be formulated in a natural way using the language of expansion trees we have introduced so far. In the following, we fix max $\emptyset = 0$.

Definition 15. Let w be a \forall -expansion in \mathcal{P} , and let > be its dependency relation. A sequence of \forall -expansions w, w_1, \ldots, w_k of \mathcal{P} such that $w > w_1 > w_1$

 $\cdots > w_k$ is called a >-chain descending from w of length k. We now define the rank $\operatorname{rk}(w)$ for expansions w and the degree $\operatorname{deg}(w)$ for \forall -expansions w:

 $rk(w) = \max\{rk(u) \mid w \text{ dominates } u\} + 1,$ $deg(w) = \max\{length \text{ of } c \mid c \text{ } i\text{-chain descending from } w\}.$

A trivial but crucial property of deg is that it is order-preserving w.r.t. the dependency relation, i.e. v > w implies $\deg(v) > \deg(w)$. For use in our weak normalization proof, we extend the notion of rank to expansion proofs, calling expansions w occurring in a cut *critical*.

Definition 16. For an expansion proof \mathcal{P} and $r \in \mathbb{N}$, the rank $\operatorname{rk}(\mathcal{P})$ and the order with respect to $r \ o(\mathcal{P}, r)$ are defined as

 $\begin{aligned} \mathrm{rk}(\mathcal{P}) &= \max\{\mathrm{rk}(w) \mid w \text{ critical}\},\\ o(\mathcal{P}, r) &= \#\{w \mid w \text{ critical } \forall \text{-expansion } \wedge \mathrm{rk}(w) = r\}. \end{aligned}$

5.3 Elimination of Propositional Connectives

Since expansion proofs work modulo propositional validity, it can be expected that the elimination of propositional parts of cuts is simple. This is indeed the case: from our cut-reduction steps, it is immediately clear that purely propositional cuts can be eliminated in linear time (since each propositional connective and each atom in a cut-formula induces a single cut-reduction step). In fact, it is easy to see that if \mathcal{P}, \mathcal{C} is an expansion proof where \mathcal{C} contains only propositional cuts, then \mathcal{P} is also an expansion proof. Hence purely propositional cuts can simply be dropped. This is in line with the results of [29], where it is shown that quantifier-free cuts can be eliminated from **LK**-proofs at the cost of propositional proof search.

The following result builds on these observations, showing that propositional parts of cuts can be eliminated while preserving the complexity measures we have defined in the previous section. This will yield a convenient ,,intermediate normal form" that will be used in the proof of weak normalization.

Definition 17. An expansion proof $\mathcal{P} = \mathcal{C}, \mathcal{E}$ is $\vee \wedge$ -normal if no $(E_1, E_2) \in \mathcal{C}$ is of the form $E_1 = E_l \vee E_r$.

In particular, if \mathcal{P} is $\vee \wedge$ -normal and no cut in \mathcal{P} contains a quantifier, then \mathcal{P} contains only atomic cuts.

Lemma 5. For every expansion proof \mathcal{P} there is a $\vee \wedge$ -normal expansion proof \mathcal{P}^* such that $\mathcal{P} \to \mathcal{P}^*$, $\operatorname{Sh}(\mathcal{P}^*) = \operatorname{Sh}(\mathcal{P})$, $\operatorname{rk}(\mathcal{P}^*) = \operatorname{rk}(\mathcal{P})$ and $o(\mathcal{P}^*, r) = o(\mathcal{P}, r)$ for all r.

Proof. We proceed by induction on the number of $\vee \wedge$ -cuts in \mathcal{P} , showing by induction on the structure of \mathcal{P} that rk is preserved. See the appendix for details.

5.4 Weak Normalization

This section is dedicated to proving that there exists a terminating strategy for the application of the cut-reduction rules. Given an expansion proof \mathcal{P} , our

reduction strategy will be based on picking a degree-maximal \forall -expansion from the set $M(\mathcal{P}) = \{w \mid w \text{ critical and } \operatorname{rk}(w) = \operatorname{rk}(\mathcal{P})\}$. The following results establish some invariances of rank and domination under substitution and $\xrightarrow{\sqcup}$ reduction, which are crucial for the weak normalization proof.

Lemma 6. Let v, w be expansions in $\mathcal{P}[\alpha \setminus t] \downarrow$ such that α does not occur in any cut-formula in \mathcal{P} , and let $v' \in \operatorname{pred}_{\sqcup}(v)$ and $w' \in \operatorname{pred}_{\sqcup}(w)$. Then v dominates w if and only if v' dominates w'. Furthermore, $\operatorname{rk}(v') = \operatorname{rk}(v)$.

Proof. By induction on the definition of $P\sigma$ and $\mathcal{P}_1 \stackrel{\sqcup}{\mapsto} \mathcal{P}_2$. See the appendix for details.

Lemma 7. Let $\mathcal{P} \xrightarrow{\sqcup} \mathcal{P}'$ and v be an expansion in \mathcal{P}' and $v' \in \operatorname{pred}_{\sqcup}(v)$. Then $\operatorname{rk}(v') = \operatorname{rk}(v)$.

Proof. By induction on a $\xrightarrow{\sqcup}$ -sequence of \mathcal{P} .

Lemma 8. Let $r = \operatorname{rk}(\mathcal{P})$ and σ a substitution. Then $o(\mathcal{P}\sigma, r) = o(\mathcal{P}, r)$. Furthermore, let E_1, E_2 be expansion trees and $E = (E_1 \sqcup E_2) \downarrow$, then $o(E, r) = o(E_1, r) = o(E_2, r)$.

Proof. Using Lemma 6 for substitution, and for merge the fact that expansions of rank r are uppermost and hence merged. See Appendix for details.

We are ready to state the main tool of the termination proof. It shows that when reducing an appropriate quantified cut, the number of \forall -expansions with maximal rank decreases, while the maximal rank does not increase. The difficulty lies in showing that while the expansion proof with merge that is constructed by the cut-reduction rule may, in fact, contain more \forall -expansions of maximal rank, this increase will be eliminated by the merge-normalization.

Lemma 9. Let $\mathcal{P}_1 \mapsto \mathcal{P}_2$ by the quantifier reduction-rule, let $r = \operatorname{rk}(\mathcal{P}_1)$ and denote the reduced \forall -expansion by w. If $w \in M(\mathcal{P})$ and $\operatorname{deg}(w)$ is maximal in $M(\mathcal{P}_1)$, then $\operatorname{rk}(\mathcal{P}_2) \leq r$ and $o(\mathcal{P}_2, r) = o(\mathcal{P}_1, r) - 1$.

Proof. We have $\operatorname{rk}(\mathcal{P}_2) \leq r$ since the rank changes for no expansions by Lemmas 6 and 7. To show that $o(\mathcal{P}_2, r) = o(\mathcal{P}_1, r) - 1$, it suffices to show that for all non-reduced cuts $G \in \mathcal{P}_1$ containing a \forall -expansion of rank $r, \alpha \notin \operatorname{V}(\operatorname{Sh}(G))$ and if $\beta \in \operatorname{V}(\operatorname{Sh}(G))$ then $w \notin q(\beta)$: If this is so, then $\operatorname{Sh}(G)[\alpha \setminus t_i] = \operatorname{Sh}(G)$ and (as can be checked by induction) the regularization η_i is reversed w.r.t. $\operatorname{Sh}(G)$ by the merge, and hence the cuts are merged. Therefore, by Lemma 8, their order stays the same. Furthermore, E_i, E do not contain expansions of maximal rank (since w has maximal rank), and w does not have a successor in \mathcal{P}_2 , hence $o(\mathcal{P}_2, r) = o(\mathcal{P}_1, r) - 1$.

To show the claim, consider a \forall -expansion v of rank r in G. If $\alpha \in V(Sh(G))$, then $v > q(\alpha) = w$ and therefore $\deg(v) > \deg(w)$, which contradicts maximality of $\deg(w)$. Similarly, $\beta \in V(Sh(G))$ implies $q(\beta) < w$. Assuming $w < q(\beta)$ yields w < v and again the contradictory $\deg(w) < \deg(v)$.

Theorem 3 (Weak Normalization). For every expansion proof \mathcal{P} there is a cut-free expansion proof \mathcal{P}^* with $\operatorname{Sh}(\mathcal{P}) = \operatorname{Sh}(\mathcal{P}^*)$ and $\mathcal{P} \to \mathcal{P}^*$.

Proof. First, we apply the propositional cut-reduction rules exhaustively to \mathcal{P} to obtain an $\vee \wedge$ -normal expansion proof \mathcal{P}^* (Lemma 5). If \mathcal{P}^* is cut-free, we are done. Otherwise, $M(\mathcal{P})$ contains a \forall -expansion. Let $n \in M(\mathcal{P})$ be a \forall -expansion such that deg(n) is maximal in $M(\mathcal{P})$. Since deg(n) is maximal and \mathcal{P}^* is $\vee \wedge$ -normal, no node dominates n. Hence we may apply the quantifier-reduction rule to n, which decreases $o(\mathcal{P}^*, r)$ by Lemma 9. At some point, $o(\mathcal{P}^*, r) = 0$, and the next cut-reduction will be applied to a \forall -expansion of rank < r. Since, by Lemmas 5 and 9, $\operatorname{rk}(\mathcal{P}^*)$ never increases, we conclude termination of the strategy by double induction. Finally, $\operatorname{Sh}(\mathcal{P}) = \operatorname{Sh}(\mathcal{P}^*)$ by Lemma 4.

5.5 Strong Normalization

Having shown weak normalization of the cut-reduction rules in the previous section, it is important to turn to the question of strong normalization, i.e. whether *all* reduction sequences are of finite length. We conjecture that our cut-reduction rules are indeed strongly normalizing, and present some evidence for this claim by discussing how our reduction rules behave on a translation of the example [15, Figure 14], which causes a failure of strong normalization in the setting of proof forests.

This example can be translated as an expansion proof of the form $\mathcal{P} = (C_1^+, C_1^-), (C_2^+, C_2^-), \mathcal{P}'$ (where \mathcal{P}' is cut-free) with

$$\begin{array}{rcl} C_1^+ &=& \exists x \,\forall y \,\overline{P(x,y)} &+ {}^c \forall y \,\overline{P(c,y)} + {}^\gamma \overline{P(c,\gamma)} \\ &+ {}^\gamma \forall y \,\overline{P(\gamma,y)} + {}^\delta \,\overline{P(\gamma,\delta)} \\ C_1^- &=& \forall x \,\exists y \,P(x,y) &+ {}^\alpha \exists y \,P(\alpha,y) + {}^\beta \,P(\alpha,\beta) \\ C_2^+ &=& \exists x \,\forall y \,\overline{Q(x,y)} &+ {}^c \forall y \,\overline{Q(c,y)} + {}^\epsilon \overline{Q(c,\epsilon)} \\ &+ {}^\epsilon \forall y \,\overline{Q(\epsilon,y)} + {}^\iota \,\overline{Q(\epsilon,\iota)} \\ C_2^- &=& \forall x \,\exists y \,Q(x,y) &+ {}^\beta \exists y \,Q(\beta,y) + {}^\alpha \,Q(\beta,\alpha). \end{array}$$

It can be checked that any application of our cut-reduction rules to such a proof terminates. This is essentially due to the different treatment of bridges (i.e. dependencies between different sides of a cut, see Section 5.1) in our formalism: at the core of the non-termination of [15, Figure 14] lies a single bridge [15, Figure 16] which induces a cycle. In our setting, if \mathcal{P} is an expansion proof containing a single cut, and $\mathcal{P} \mapsto \mathcal{P}'$ via a quantifier reduction rule, then \mathcal{P}' still contains only a single cut.

Indeed, a reduction sequence similar to the non-terminating one described in [15, Figure 17] exists, and it ends in such an expansion proof containing only a single cut which is, also in our setting, a bridge. The cut reduces then to a single propositional cut, the elimination of which is easily seen to be strongly normalizing.

In the setting of proof forests, the non-termination due to bridges is handled by adding a pruning reduction. One explanation for the fact that in our setting, we are able to get by without such a reduction, is the use of the merge in the definition of the cut-reduction rules. The merge has the advantage that it is very natural, it is an extension of the merge for cut-free expansion proofs from [23], and it is useful also in applications not related to cut-elimination, as in the proof of Theorem 1.

5.6 Confluence

It is well-known that cut-elimination (and similar procedures) in classical logic are typically not confluent, see e.g. [27, 26, 3] for case studies and [2, 17] for asymptotic results. Neither the proof forests of [15] nor the Herbrand nets of [22] have a confluent reduction. The situation is analogous in our formalism: the reduction is not confluent. In fact, one can use the same example to demonstrate this; let

$$\mathcal{P} = \{\exists x A + {}^{s} A[x \backslash s] + {}^{t} A[x \backslash t], \forall x \overline{A} + {}^{\alpha} \overline{A}[x \backslash \alpha]\}, \\ \{\exists x B + {}^{\alpha} B[x \backslash \alpha], \forall x \overline{B} + {}^{\beta} \overline{B}[x \backslash \beta]\}, \\ \exists x \exists y C + {}^{\alpha} (\exists y C[x \backslash \alpha] + {}^{\beta} C[x \backslash \alpha, y \backslash \beta]).$$

which is the translation of [15, Figure 12] into an expansion proof with cut. Then it can be verified by a quick calculation that the choice of reducing either the cut on A or that on B first determines which of two normal forms is obtained.

However cut-elimination in classical logic can be shown confluent on the level of the (cut-free) expansion tree on a certain class of proofs [18]. For future work we hope to use such techniques for describing a confluent reduction in expansion proofs whose normal form is unique and most general in the sense that it contains all other normal forms as sub-expansions.

6 Conclusion

In this paper we have presented expansion proofs with cut for full first-order logic including non-prenex formulas. Our definitions extend the existing notion of cut-free expansion proofs in a natural way. We have given a cut-elimination procedure and proved weak normalization; strong normalization remains an open problem. Our proof of weak normalization is inspired by the ε -calculus which allowed to cover also the non-prenex case without technical difficulties. The complex object-level syntax of the ε -calculus is avoided in our work by taking care of the mutual dependencies of variables by the merge operation of expansion trees.

It should be noted that the ε -calculus is, in a sense, more general than expansion-proofs since there are formulas in the ε -language which do not arise by translation from usual formulas. But of course, our objective is not to create a general formalism, but rather to find a good model of cut-elimination for the classical first-order sequent calculus! For this purpose, we believe expansion proofs with cut are very promising, as they are compact, focus on the firstorder level of proofs, and admit natural cut-reduction rules which are weakly normalizing — and perhaps even strongly normalizing.

Acknowledgements. The authors would like to thank D. Miller and K. Chaudhuri, M. Baaz, W. Heijltjes and R. McKinley for many helpful discussions about expansion trees, the ε -calculus, proof forests and Herbrand nets respectively.

References

- Jeremy Avigad. The computational content of classical arithmetic. In Solomon Feferman and Wilfried Sieg, editors, *Proofs, Categories, and Computations: Essays in Honor of Grigori Mints*, pages 15–30. College Publications, 2010.
- [2] Matthias Baaz and Stefan Hetzl. On the non-confluence of cut-elimination. Journal of Symbolic Logic, 76(1):313–340, 2011.
- [3] Matthias Baaz, Stefan Hetzl, Alexander Leitsch, Clemens Richter, and Hendrik Spohr. Cut-Elimination: Experiments with CERES. In Franz Baader and Andrei Voronkov, editors, *Logic for Programming, Artificial Intelligence, and Reasoning (LPAR) 2004*, volume 3452 of *Lecture Notes in Computer Science*, pages 481–495. Springer, 2005.
- [4] Matthias Baaz, Stefan Hetzl, and Daniel Weller. On the complexity of proof deskolemization. *Journal of Symbolic Logic*, 77(2):669–686, 2012.
- [5] Matthias Baaz and Alexander Leitsch. Cut-elimination and Redundancyelimination by Resolution. *Journal of Symbolic Computation*, 29(2):149– 176, 2000.
- [6] Franco Barbanera and Stefano Berardi. A Symmetric Lambda Calculus for Classical Program Extraction. Information and Computation, 125(2):103– 117, 1996.
- [7] Ulrich Berger, Wilfried Buchholz, and Helmut Schwichtenberg. Refined Program Extraction from Classical Proofs. Annals of Pure and Applied Logic, 114:3–25, 2002.
- [8] Samuel R. Buss. On Herbrand's Theorem. In Logic and Computational Complexity, volume 960 of Lecture Notes in Computer Science, pages 195– 209. Springer, 1995.
- [9] Kaustuv Chaudhuri, Stefan Hetzl, and Dale Miller. A Systematic Approach to Canonicity in the Classical Sequent Calculus. In Patrick Cégielski and Arnaud Durand, editors, *Computer Science Logic (CSL) 2012*, volume 16 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 183–197. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, 2012.
- [10] Kaustuv Chaudhuri, Stefan Hetzl, and Dale Miller. The Isomorphism Between Expansion Proofs and Multi-Focused Sequent Proofs. submitted, 2012.
- [11] Thierry Coquand. A semantics of evidence for classical arithmetic. Journal of Symbolic Logic, 60(1):325–337, 1995.
- [12] Pierre-Louis Curien and Hugo Herbelin. The Duality of Computation. In Proceedings of the Fifth ACM SIGPLAN International Conference on Functional Programming (ICFP '00), pages 233–243. ACM, 2000.
- [13] Vincent Danos, Jean-Baptiste Joinet, and Harold Schellinx. A New Deconstructive Logic: Linear Logic. Journal of Symbolic Logic, 62(3):755–807, 1997.

- [14] Jean-Yves Girard. Linear logic. Theoretical Computer Science, 50(1):1–101, 1987.
- [15] Willem Heijltjes. Classical proof forestry. Annals of Pure and Applied Logic, 161(11):1346–1366, 2010.
- [16] Jacques Herbrand. Recherches sur la théorie de la démonstration. PhD thesis, Université de Paris, 1930.
- [17] Stefan Hetzl. The Computational Content of Arithmetical Proofs. Notre Dame Journal of Formal Logic, 53(3):289–296, 2012.
- [18] Stefan Hetzl and Lutz Straßburger. Herbrand-Confluence for Cut-Elimination in Classical First-Order Logic. In Patrick Cégielski and Arnaud Durand, editors, Computer Science Logic (CSL) 2012, volume 16 of Leibniz International Proceedings in Informatics (LIPIcs), pages 320–334. Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik, 2012.
- [19] David Hilbert and Paul Bernays. Grundlagen der Mathematik II. Springer, 1939.
- [20] Ulrich Kohlenbach. Applied Proof Theory: Proof Interpretations and their Use in Mathematics. Springer, 2008.
- [21] Konstantin Korovin. Instantiation-Based Automated Reasoning: From Theory to Practice. In Renate A. Schmidt, editor, 22nd International Conference on Automated Deduction (CADE), volume 5663 of Lecture Notes in Computer Science, pages 163–166. Springer, 2009.
- [22] Richard McKinley. Proof nets for Herbrand's Theorem. ACM Transactions on Computational Logic, 14(1), 2013.
- [23] Dale Miller. A Compact Representation of Proofs. Studia Logica, 46(4):347– 370, 1987.
- [24] Georg Moser and Richard Zach. The Epsilon Calculus and Herbrand Complexity. Studia Logica, 82(1):133–155, 2006.
- [25] Michel Parigot. λμ-Calculus: An Algorithmic Interpretation of Classical Natural Deduction. In Andrei Voronkov, editor, Logic Programming and Automated Reasoning, International Conference LPAR'92, Proceedings, volume 624 of Lecture Notes in Computer Science, pages 190–201. Springer, 1992.
- [26] Diana Ratiu and Trifon Trifonov. Exploring the Computational Content of the Infinite Pigeonhole Principle. 22(2):329–350, 2012. Journal of Logic and Computation.
- [27] Christian Urban. Classical Logic and Computation. PhD thesis, University of Cambridge, October 2000.
- [28] Christian Urban and Gavin Bierman. Strong Normalization of Cut-Elimination in Classical Logic. Fundamenta Informaticae, 45:123–155, 2000.

[29] Daniel Weller. On the Elimination of Quantifier-Free Cuts. Theoretical Computer Science, 412(49):6843–6854, 2011.

7 Appendix

In this appendix we describe the technical details that have been omitted from the main paper. Note that the numbering of the results is non-monotonic: For those results that are stated in the main text, we have retained the numbers, while we introduce new ones for intermediate results presented exclusively in the appendix. We also include some examples here that did not fit into the main paper.

7.1 Basic Operations on Expansion Proofs

Example 5. Consider the following expansion tree E and $E\sigma$ with $\sigma = [\alpha \backslash c]$.

$$E = \exists x \forall y R(x, y) + {}^{c} (\forall y R(c, y) + {}^{\beta} R(c, \beta)) + {}^{\alpha} (\forall y R(\alpha, y) + {}^{\gamma} R(\alpha, \gamma))$$
$$E\sigma = \exists x \forall y R(x, y) + {}^{c} ((\forall y R(c, y) + {}^{\beta} R(c, \beta)) \sqcup (\forall y R(c, y) + {}^{\gamma} R(c, \gamma)))$$

Note that the only expansion w in E that has no w' in $E\sigma$ such that $\operatorname{pred}_s(w') = w$ is the $+^{\alpha}$ expansion — it is replaced by a \sqcup -node.

By induction on the definition, it is easy to see that if σ is a renaming and \mathcal{P} an expansion tree (without merge), then $\mathcal{P}\sigma$ is an expansion tree without merge. An important property is that substitution commutes with $Dp(\cdot)$ and $Sh(\cdot)$.

Lemma 10. Let \mathcal{P} be an expansion pre-proof with merges and σ a substitution. Then

- $\operatorname{Dp}(\mathcal{P}\sigma)$ is logically equivalent to $\operatorname{Dp}(\mathcal{P})\sigma$, and
- $\operatorname{Sh}(\mathcal{P}\sigma) = \operatorname{Sh}(\mathcal{P})\sigma$.

Proof. By induction on the structure of \mathcal{P} .

Lemma 11. Let $\mathcal{P} = \mathcal{P}'[E]$ be an expansion proof with merges and σ a substitution admissible for \mathcal{P} . Then $\mathcal{Q} = \mathcal{P}'[E\sigma]$ is an expansion proof with merges, and $\operatorname{Sh}(\mathcal{P}) = \operatorname{Sh}(\mathcal{Q})$.

Proof. Lemma 10 implies the latter claim since $\operatorname{Sh}(\mathcal{P})$ does not contain free variables. The same Lemma also implies that $\operatorname{Dp}(\mathcal{Q})$ is a tautology since propositional tautology-hood is preserved under substitution. Regularity is preserved since every subtree of E is "copied" exactly once to create $E\sigma$. To show that $<_{\mathcal{Q}}$ is acyclic, we show that $w <_{\mathcal{Q}} w$ implies $\operatorname{pred}_s(w) <_{\mathcal{P}} \operatorname{pred}_s(w)$. The only non-trivial case is that there exist an \exists -expansion v such that $\operatorname{pred}_s(v)$ has an expansion term containing α , and a \forall -expansion u with eigenvariable β , such that $w >_{\mathcal{Q}} v >_{\mathcal{Q}} u >_{\mathcal{Q}} w$, and $\beta \in \operatorname{V}(\alpha\sigma)$. But this implies $\operatorname{pred}_s(u) >_{\mathcal{P}} \operatorname{pred}_s(w) >_{\mathcal{P}} \operatorname{pred}_s(v) = q(\beta)$, which contradicts admissibility of σ .

Example 6. Continuing Example 5, we have $E\sigma \stackrel{\sqcup}{\mapsto} E' \stackrel{\sqcup}{\mapsto} E''$ where

$$E' = \exists x \forall y R(x, y) + {}^{c} (\forall y R(c, y) + {}^{\beta} R(c, \beta) \sqcup R(c, \beta))$$
$$E'' = \exists x \forall y R(x, y) + {}^{c} (\forall y R(c, y) + {}^{\beta} R(c, \beta))$$

The only expansion w in E' with a non-trivial set $\operatorname{pred}_{\sqcup}^{0}(w)$ is the $+^{\beta}$ expansion: it has $\operatorname{pred}_{\sqcup}^{0}(w) = \{+^{\beta}, +^{\gamma}\}$. Similarly, the only w in E'' with non-trivial $\operatorname{pred}_{\sqcup}^{0}(w)$ is $R(c,\beta)$: here, $\operatorname{pred}_{\sqcup}^{0}(w)$ consists of the two occurrences of $R(c,\beta)$ in E'.

Lemma 12. The relation $\stackrel{\square}{\rightarrow}$ is confluent and strongly normalizing. Its normal forms have no merge nodes.

Proof. Local confluence follows immediately from the absence of critical pairs.

Let *m* be a merge node in an expansion pre-proof with merges \mathcal{P} , then the weight of this node w(m) is the number of nodes below it in \mathcal{P} . Let m_1, \ldots, m_l be the merge nodes in an expansion pre-proof with merges \mathcal{P} , then the weight of \mathcal{P} is defined as $w(\mathcal{P}) = \sum_{i=1}^{l} w(m_i)$. The application of eigenvariable renamings and merges decreases the lexicographic ordering $\langle |\text{EV}(\mathcal{P})|, w(\mathcal{P}) \rangle$, hence $\stackrel{\sqcup}{\rightarrow}$ is strongly normalizing and we can conclude confluence.

That its normal forms have no more merge nodes is immediate.

Example 7. Continuing Example 6, we have $E'' = E\sigma \downarrow$. There is exactly one expansion w in E'' with a non-trivial set $\operatorname{pred}_{\sqcup}(w)$ (containing expansions from $E\sigma$): namely, $\operatorname{pred}_{\sqcup}(+^{\beta}) = \{+^{\beta}, +^{\gamma}\}$.

An important property is that $\stackrel{\sqcup}{\rightarrow}$ preserves the proof properties of \mathcal{P} .

Lemma 13. If $\mathcal{P} \stackrel{\sqcup}{\mapsto} \mathcal{P}'$ then

- if $Dp(\mathcal{P})$ is valid then $Dp(\mathcal{P}')$ is valid, and
- if $<_{\mathcal{P}}$ is acyclic then $<_{\mathcal{P}'}$ is acyclic, and
- if \mathcal{P} is regular than so is \mathcal{P}'

Proof. By inspection of the definition, we show that there exists a variable renaming σ such that Dp(\mathcal{P}) σ → Dp(\mathcal{P}'). The variable renaming is used in case 3. Note that logical equivalence is not preserved due to case 2. Acyclicity is shown by verifying that for all expansions v, w from \mathcal{P}' , if $w >_{\mathcal{P}}^{0} v$ then there exist $w' \in \operatorname{pred}_{\sqcup}^{0}(w)$ such that for all $v' \in \operatorname{pred}_{\sqcup}^{0}(v)$ we have $w' >_{\mathcal{P}}^{0} v'$. This allows to translate a cycle from $>_{\mathcal{P}'}$ to $>_{\mathcal{P}}$. Finally, regularity of \mathcal{P}' follows since no new \forall -expansions are introduced.

Lemma 14. If $\mathcal{P}_1 \sqcup \mathcal{P}_2$ is an expansion proof with merge such that $\operatorname{Sh}(\mathcal{P}_1) = \operatorname{Sh}(\mathcal{P}_2)$, then $\mathcal{P}_1 \cup \mathcal{P}_2$ is an expansion proof and $\operatorname{Sh}(\mathcal{P}_1 \cup \mathcal{P}_2) = \operatorname{Sh}(\mathcal{P}_1) = \operatorname{Sh}(\mathcal{P}_2)$.

Proof. Since $\operatorname{Sh}(\mathcal{P}_1) = \operatorname{Sh}(\mathcal{P}_2)$, all non-cut expansion-trees are merged and we have $\operatorname{Sh}(\mathcal{P}_1 \cup \mathcal{P}_2) = \operatorname{Sh}(\mathcal{P}_1) = \operatorname{Sh}(\mathcal{P}_2)$ by definition. The proof-properties of $\mathcal{P}_1 \sqcup \mathcal{P}_2$ are carried over to $\mathcal{P}_1 \cup \mathcal{P}_2$ by Lemma 13 and Lemma 1: in case 3 of Definition 9, the eigenvariable renaming is admissible since $q(\alpha_1)$ and $q(\alpha_2)$ are dominated by the same expansions, and are contained in the same cut (if any). Finally, $\mathcal{P}_1 \cup \mathcal{P}_2$ does not contain merge nodes by Lemma 12.

7.2 Cut-Elimination

Towards verifying that \mapsto is really a binary relation on expansion proofs, as claimed, we have to prove a technical result on the behavior of the merge w.r.t. cut-reduction, namely that those \forall -expansions that do not depend upon the reduced \forall -expansion are merged.

Lemma 15. Let $\mathcal{P} \mapsto \mathcal{P}'$ by the quantifier-reduction rule. We write $\mathcal{P}' = \mathcal{P}'' \downarrow$ where \mathcal{P}'' is the expansion tree with merge constructed by the reduction rule. Let w be the \forall -expansion indicated by the rule, and let v be a \forall -expansion in \mathcal{P} with $w \leqslant v$. Then there exists a \forall -expansion u in \mathcal{P}' s.t. $v' \in \operatorname{pred}_{\sqcup}(u)$ for all copies v' of v in \mathcal{P}'' .

Proof. By induction on the merge-reduction sequence. The assumption $w \leq v$ ensures that in case 4 of Definition 9, the subtrees containing copies of u will be merged since the \exists -expansions dominating them belong to the $+^{r_i}$ -part. \Box

Lemma 16. If $\mathcal{P}_1 \mapsto \mathcal{P}_2$ and \mathcal{P}_1 is an expansion proof, then \mathcal{P}_2 is an expansion proof. Furthermore, $\operatorname{Sh}(\mathcal{P}_1) = \operatorname{Sh}(\mathcal{P}_2)$.

Proof. We only give the proof for the quantifier cut-reduction step; the proof for the other reduction steps is analogous and simpler. Let $\sigma_i = [\alpha \setminus t_i]$ and assume

$$\mathcal{P}_1 = \{ \exists x \, A + {}^{t_1} E_1 \cdots + {}^{t_n} E_n, \forall x \, \bar{A} + {}^{\alpha} E \}, \mathcal{P} \\ \mapsto \mathcal{P} \cup \{ E_1 \lor \cdots \lor E_n, E\eta_1 \sigma_1 \land \cdots \land E\eta_n \sigma_n \} \cup \bigcup_{i=1}^n \mathcal{P}\eta_i \sigma_i = \mathcal{P}_2,$$

where η_i are renamings establishing regularity. First, note that the η_i are trivially admissible for \mathcal{P}_1 since only new variables are introduced. Next, we show that the σ_i are admissible for $\mathcal{P} \cup \{E_1 \lor \cdots \lor E_n, E\eta_1 \land \cdots \land E\eta_n\} \cup \bigcup_{j=1}^n \mathcal{P}\eta_j$. Hence assume $\beta \in V(t_i)$ and that there exists an \exists -expansion w in $\mathcal{P}\eta_j$ or $E\eta_j$ with expansion term t such that $\alpha \in V(t)$ and that $w < q(\beta)$. This is only possible if there is a \forall -expansion v with eigenvariable γ such that $v < q(\beta)$ and w dominates v. But since γ is a fresh variable introduced by η_j , this implies that v dominates $q(\beta)$, hence β is also a fresh variable introduced by η_j , which contradicts $\beta \in V(t_i)$.

Towards showing that \mathcal{P}_2 is an expansion proof, we have to make the definition of expansion proof with merge slightly more liberal: we allow cuts $\{C^+, C^-\}$ such that $\operatorname{Sh}(C^+) = \overline{\operatorname{Sh}(C^-)}\eta$, where η is a renaming (the usual definition requires η to be the identity renaming). The results from Section 3.1, hold as well for this definition. η will be chosen such that after merge-normalization, all cuts will be syntactically correct.

Now we show that \mathcal{P}_2 is an expansion proof: writing $\mathcal{P}_2 = \mathcal{P}'_2 \downarrow$, by Lemma 1 and Lemma 3 it suffices to show that \mathcal{P}'_2 is regular (which it is by construction), that its dependency relation is acyclic, that $Dp(\mathcal{P}'_2)$ is a tautology, and that $Sh(\mathcal{P}'_2) = Sh(\mathcal{P}_1)$ (which holds by construction as well).

To show that $\operatorname{Dp}(\mathcal{P}'_2)$ is valid, we start by reducing the problem: It can be checked (using the propositional tautology $(A \wedge A') \vee (B \wedge B') \to (A \vee B) \wedge (A' \vee B')$) that $\operatorname{Dp}(\mathcal{P}'_2)$ is implied by

$$F = \operatorname{Dp}(\mathcal{P}) \vee (\bigvee_{i=1}^{n} \operatorname{Dp}(E_{i})) \wedge (\bigwedge_{i=1}^{n} \operatorname{Dp}(E\eta_{i}\sigma_{i})) \vee \bigvee_{i=1}^{n} \operatorname{Dp}(\mathcal{P}\eta_{i}\sigma_{i}).$$

By Lemma 10, F is logically equivalent to

$$F' = \operatorname{Dp}(\mathcal{P}) \vee (\bigvee_{i=1}^{n} \operatorname{Dp}(E_{i})) \wedge (\bigwedge_{i=1}^{n} \operatorname{Dp}(E)\eta_{i}\sigma_{i}) \vee \bigvee_{i=1}^{n} \operatorname{Dp}(\mathcal{P})\eta_{i}\sigma_{i}.$$

Hence it suffices to show that F' is valid.

Note that $\operatorname{Dp}(\mathcal{P}_1) = \operatorname{Dp}(\mathcal{P}) \lor (\bigvee_{i=1}^n \operatorname{Dp}(E_i) \land \operatorname{Dp}(E))$. Since $\operatorname{Dp}(\mathcal{P}_1)$ is valid, the formulas $\operatorname{Dp}(\mathcal{P})\eta_i\sigma_i \lor \operatorname{Dp}(E)\eta_i\sigma_i$ and $\operatorname{Dp}(\mathcal{P})\lor \bigvee_{i=1}^n \operatorname{Dp}(E_i)$ are valid. Using propositional reasoning, in particular validity of $(A \lor B) \land (C \lor D) \to A \lor (B \land C) \lor D$, we obtain validity of F'.

Next, we show that acyclicity of $<_{\mathcal{P}_1}$ implies acyclicity of $<_{\mathcal{P}_2}$. This follows from the fact that if $x <_{\mathcal{P}_2} y$ implies that there exist $x' \in \operatorname{pred}_c(x), y' \in \operatorname{pred}_c(y)$ such that $x' <_{\mathcal{P}_1} y'$. Hence a cycle in $<_{\mathcal{P}_2}$ gives rise to a cycle in $<_{\mathcal{P}_1}$.

Finally, we have to show that all cuts in \mathcal{P}_2 are syntactically correct. By construction, the only "incorrect" cut in \mathcal{P}'_2 is the indicated one. We have $\operatorname{Sh}(E\sigma_i) = \operatorname{Sh}(E\eta_i\eta_i^{-1}\sigma_i) = \operatorname{Sh}(E\eta_i\sigma_i)\eta_i^{-1}$ since η_i is a renaming to fresh variables. Since $\operatorname{Sh}(E)\sigma_i = \operatorname{Sh}(E_i)$, this yields $\overline{\operatorname{Sh}(E_i)}\eta_i = \operatorname{Sh}(E\eta_i\sigma_i)$, hence the "incorrect" cut fulfills the liberalized definition. We even have $\operatorname{Sh}(E_i)\eta_i = \operatorname{Sh}(E_i)\eta'_i$ for a variable renaming η'_i such that if $\beta \in \operatorname{dom}(\eta'_i)$ then $q(\alpha) \notin q(\beta)$. For if $\beta \in \operatorname{V}(\operatorname{Sh}(E_i))$ and $\beta \notin \operatorname{V}(\operatorname{Sh}(E\eta_i\sigma_i))$ then $\beta \in \operatorname{V}(\operatorname{Sh}(c))$ where c is the indicated cut in \mathcal{P}_1 , and therefore $q(\beta) <_{\mathcal{P}_1} q(\alpha)$. Since $<_{\mathcal{P}_1}$ is acyclic, we have $q(\alpha) \notin q(\beta)$. Hence we can take for η'_i just η_i where these β are not renamed. Finally, Lemma 15 implies that the copies of the variables in dom(η'_i) are identified by the merge, which yields correctness of the cuts in \mathcal{P}_2 .

Example 8. Consider an expansion proof with three cuts $\mathcal{P} = C_1, C_2, C_3, \mathcal{E}$ where $C_i = \{C_i^+, C_i^-\}$ for $1 \le i \le 3$ where

$$\begin{split} C_1^+ &= \exists x \forall y \, P(x, y) +^c \left(\forall y \, P(c, y) +^{\gamma} P(c, \gamma) \right) \\ C_1^- &= \forall x \exists y \, \overline{P}(x, y) +^{\alpha} \left(\exists y \, \overline{P}(\alpha, y) +^c \overline{P}(\alpha, c) \right) \\ C_2^+ &= \exists x Q(\alpha, x) +^{\alpha} Q(\alpha, \alpha) +^c Q(\alpha, c) \\ C_2^- &= \forall x Q(\alpha, x) +^{\beta} \overline{Q(\alpha, \beta)} \\ C_3^+ &= \exists x \forall y \, R(x, y) +^{\beta} \left(\forall y \, P(\beta, y) +^{\lambda} P(\beta, \lambda) \right) \\ C_3^- &= \forall x \exists y \, \overline{R}(x, y) +^{\delta} \left(\exists y \, \overline{R}(\delta, y) +^c \overline{R}(\delta, c) +^{\alpha} \overline{R}(\delta, \alpha) \right) \end{split}$$

Assuming that \mathcal{E} is cut-free and contains no \forall -nodes, it is of no importance in this context, and so we do not give its definition. Denote the expansions in these trees from left to right, top to bottom, by w_1, \ldots, w_{12} (i.e. w_1 is the $+^c$ expansion in C_1^+, w_{12} is the $+^\alpha$ expansion in C_3^- , etc.). Then the maximal >-chain descending from w_9 is $w_9 > w_8 > w_7 > w_3$, yielding deg $(w_9) = 3$, deg $(w_8) = 2$, deg $(w_7) = 1$, deg $(w_3) = 0$. In fact, w_9 is the node of maximal degree in \mathcal{P} . Furthermore, $\mathrm{rk}(w_i) = 1$ and $\mathrm{rk}(w_j) = 2$ for $i \in \{2, 4, 5, 6, 7, 9, 11, 12\}$ and $j \in \{1, 3, 8, 10\}$.

Lemma 17. For every expansion proof \mathcal{P} there is a $\vee \wedge$ -normal expansion proof \mathcal{P}^* such that $\mathcal{P} \to \mathcal{P}^*$, $\operatorname{Sh}(\mathcal{P}^*) = \operatorname{Sh}(\mathcal{P})$, $\operatorname{rk}(\mathcal{P}^*) = \operatorname{rk}(\mathcal{P})$ and $o(\mathcal{P}^*, r) = o(\mathcal{P}, r)$ for all r.

Proof. If \mathcal{P} is not $\vee \wedge$ -normal, then $\mathcal{P} = (E_1 \vee E_2, E'_1 \wedge E'_2), \mathcal{P}'$ and hence

$$\mathcal{P} \mapsto (E_1, E_1'), (E_2, E_2'), \mathcal{P}' = \mathcal{P}^*.$$

Since the number of $\vee \wedge$ -cuts in \mathcal{P}^* is strictly smaller than the number of $\vee \wedge$ -cuts in \mathcal{P} , and $\operatorname{rk}(w) = \operatorname{rk}(\operatorname{pred}_c(w))$ for all expansions w in \mathcal{P}^* , we conclude by induction.

Let (E_1, E_2) be a cut in an expansion proof. Since $Sh(E_1) = \overline{Sh(E_2)}$ we can associate in a natural way to every expansion w in E_1 a non-empty set of

dual expansions in E_2 , the set of *dual expansions* dl(w). This association is symmetric, i.e. $v \in dl(w)$ exactly if $w \in dl(v)$.

Example 9. We continue Example 8, giving the sets of dual expansions for C_3 : dl(w_8) = { w_{10} }, dl(w_9) = { w_{11} , w_{12} }, dl(w_{10}) = { w_8 }, dl(w_{11}) = { w_9 }, dl(w_{12}) = { w_9 }.

Lemma 18. Let w be a critical expansion. Then rk(w) = rk(w') for all $w' \in dl(w)$.

Proof. By structural induction on the cut-formula, noting that since w, w' occur in the same cut, they have the same cut-formula.

Lemma 19. If \mathcal{P} contains a critical expansion, then $M(\mathcal{P})$ contains a \forall -expansion.

Proof. Since $M(\mathcal{P})$ is non-empty, the result follows from Lemma 18.

The following result is a trivial consequence of the definition, and will ensure that a cut-reduction rule is applicable to expansions in $M(\mathcal{P})$.

Lemma 20. Let \mathcal{P} be an expansion proof and $w \in M(\mathcal{P})$. Then no expansion dominates w.

Lemma 21. Let w, v be quantifier nodes in $\mathcal{P}[\alpha \setminus t] \downarrow$ such that α does not occur in any cut-formula in \mathcal{P} , and let $w' \in \operatorname{pred}_{\sqcup}(w)$ and $v' \in \operatorname{pred}_{\sqcup}(v)$. Then wdominates v if and only if w' dominates v'. Furthermore, $\operatorname{rk}(w') = \operatorname{rk}(w)$.

Proof. By induction on the definition of $\mathcal{P}\sigma$, it is easy to show that w dominates v in $\mathcal{P}\sigma$ iff $\operatorname{pred}_s(w)$ dominates $\operatorname{pred}_s(v)$. Next, we show that if $\mathcal{P}_1 \stackrel{\sqcup}{\mapsto} \mathcal{P}_2$ then for w, v nodes in \mathcal{P}_2 and $w' \in \operatorname{pred}^0_{\sqcup}(w), v' \in \operatorname{pred}^0_{\sqcup}(v)$, w dominates v iff w' dominates v. This is obvious in case 2 of the definition. In case 1, we have $E[L \sqcup L] \stackrel{\sqcup}{\mapsto} E[L]$, and it suffices to observe that a node w dominates L in E[L] iff all $w' \in \operatorname{pred}^0_{\sqcup}(w)$ dominate both occurrences of L in $E[L \sqcup L]$. In cases 3 and 4, we reason analogously, using the result for pred_s we just proved for case 3. Finally, we extend the result to $\operatorname{pred}_{\sqcup}$ by induction on its definition. $\operatorname{rk}(w') = \operatorname{rk}(w)$ follows immediately from the first statement.

Lemma 22. Let $r = \operatorname{rk}(\mathcal{P})$ and σ a substitution. Then $o(\mathcal{P}\sigma, r) = o(\mathcal{P}, r)$. Furthermore, let E_1, E_2 be expansion trees and $E = E_1 \sqcup E_2 \downarrow$, then $o(E, r) = o(E_1, r) = o(E_2, r)$.

Proof. $o(\mathcal{P}\sigma, r) = o(\mathcal{P}, r)$ holds for all r by Lemma 6. Let v be a \forall -node of rank r in E. Then there is a $w \in \operatorname{pred}_{\sqcup}(v)$ such that $\operatorname{rk}(w) = r$, and since $\operatorname{Sh}(E_1) = \operatorname{Sh}(E_2)$ and by Lemma 20, there is a unique w' corresponding to w in E_2 . It is then easy to see by induction on an appropriate $\xrightarrow{\sqcup}$ -sequence that $\operatorname{pred}_{\sqcup}(v) = \{w, w'\}$. From this, the claim follows. \Box

We conclude by giving a more detailed proof of the weak-normalization result.

Lemma 23. Let $\mathcal{P}_1 \mapsto \mathcal{P}_2$ by the quantifier reduction-rule, let $r = \operatorname{rk}(\mathcal{P}_1)$ and denote the reduced \forall -expansion by w. If $w \in M(\mathcal{P})$ and $\operatorname{deg}(w)$ is maximal in $M(\mathcal{P}_1)$, then $\operatorname{rk}(\mathcal{P}_2) \leq r$ and $o(\mathcal{P}_2, r) = o(\mathcal{P}_1, r) - 1$.

Proof. We have $\operatorname{rk}(\mathcal{P}_2) \leq r$ since the rank changes for no expansions by Lemmas 6 and 7. To show that $o(\mathcal{P}_2, r) = o(\mathcal{P}_1, r) - 1$, it suffices to show that for all non-reduced cuts $G \in \mathcal{P}_1$ containing a \forall -expansion of rank $r, \alpha \notin \operatorname{V}(\operatorname{Sh}(G))$ and if $\beta \in \operatorname{V}(\operatorname{Sh}(G))$ then $w \leq q(\beta)$: If this is so, then $\operatorname{Sh}(G)[\alpha \setminus t_i] = \operatorname{Sh}(G)$ and by Lemma 15, the regularization η_i is reversed w.r.t. $\operatorname{Sh}(G)$ by the merge, and hence the cuts are merged. Therefore, by Lemma 8, their order stays the same. Furthermore, E_i, E do not contain expansions of maximal rank (since w has maximal rank), and w does not have a successor in \mathcal{P}_2 , hence $o(\mathcal{P}_2, r) = o(\mathcal{P}_1, r) - 1$.

To show the claim, consider a \forall -expansion v of rank r in G. If $\alpha \in V(Sh(G))$, then $v > q(\alpha) = w$ and therefore $\deg(v) > \deg(w)$, which contradicts maximality of $\deg(w)$. Similarly, $\beta \in V(Sh(G))$ implies $q(\beta) < w$. Assuming $w < q(\beta)$ yields w < v and again the contradictory $\deg(w) < \deg(v)$.

Theorem 33 (Weak Normalization). For every expansion proof \mathcal{P} there is a cut-free expansion proof \mathcal{P}^* with $\operatorname{Sh}(\mathcal{P}) = \operatorname{Sh}(\mathcal{P}^*)$ and $\mathcal{P} \to \mathcal{P}^*$.

Proof. First, we apply the propositional cut-reduction rules exhaustively to \mathcal{P} to obtain an ∨∧-normal expansion proof \mathcal{P}^* (Lemma 5). If \mathcal{P}^* is cut-free, we are done. Otherwise, $M(\mathcal{P})$ contains a \forall -expansion by Lemma 19. Let $n \in M(\mathcal{P})$ be a \forall -expansion such that deg(n) is maximal in $M(\mathcal{P})$. Since deg(n) is maximal and \mathcal{P}^* is ∨∧-normal, no node dominates n by Lemma 20. Hence we may apply the quantifier-reduction rule to n, which decreases $o(\mathcal{P}^*, r)$ by Lemma 9. At some point, $o(\mathcal{P}^*, r) = 0$, and the next cut-reduction will be applied to a \forall -expansion of rank < r. Since, by Lemmas 5 and 9, rk (\mathcal{P}^*) never increases, we conclude termination of the strategy by double induction. Finally, Sh $(\mathcal{P}) = \operatorname{Sh}(\mathcal{P}^*)$ by Lemma 4.