

On the arithmetic of crossratios and generalised Mertens' formulas

Jouni Parkkonen

Frédéric Paulin

March 3, 2018

Abstract

We develop the relation between hyperbolic geometry and arithmetic equidistribution problems that arises from the action of arithmetic groups on real hyperbolic spaces, especially in dimension ≤ 5 . We prove generalisations of Mertens' formula for quadratic imaginary number fields and definite quaternion algebras over \mathbb{Q} , counting results of quadratic irrationals with respect to two different natural complexities, and counting results of representations of (algebraic) integers by binary quadratic, Hermitian and Hamiltonian forms with error bounds. For each such statement, we prove an equidistribution result of the corresponding arithmetically defined points. Furthermore, we study the asymptotic properties of crossratios of such points, and expand Pollicott's recent results on the Schottky-Klein prime functions.¹

1 Introduction

The aim of this paper is to prove various equidistributions results (and their related asymptotic counting results) of arithmetically defined points in tori (quotient of the real or complex line \mathbb{R} or \mathbb{C} , or of Hamilton's quaternion algebra \mathbb{H} , by arithmetic lattices), organised using appropriate complexities (see for instance, amongst the numerous literature, [Oh, Har, Ser, BQ] and their references for other types of results).

We denote by Δ_x the unit Dirac mass at a point x , by $\xrightarrow{*}$ the weak-star convergence of measures, and by Leb_E the standard Lebesgue measure on a Euclidean space E . Given an imaginary quadratic number field K , we will denote by \mathcal{O}_K its ring of integers, by D_K its discriminant, by ζ_K its zeta function and by \mathfrak{n} its norm.

The first two statements are equidistribution results of rational points (satisfying congruence properties) in the complex field and Hamilton's quaternion algebra, analogous to the equidistribution result of Farey fractions in the real field, where the complexity is the norm of the denominator.

¹**Keywords:** counting, equidistribution, Mertens formula, quadratic irrational, binary Hermitian form, binary Hamiltonian form, norm form, representation of integers, group of automorphs, crossratio, Bianchi group, Hamilton-Bianchi group, common perpendicular, hyperbolic geometry. **AMS codes:** 11D85, 11E39, 11F06, 11N45, 11R52, 20H10, 20G20, 30F40, 53A35, 53C22

Theorem 1 *Let \mathfrak{m} be a (nonzero) fractional ideal of the ring of integers of an imaginary quadratic number field K , with norm $\mathfrak{n}(\mathfrak{m})$. As $s \rightarrow +\infty$,*

$$\frac{|D_K| \zeta_K(2)}{2 \pi s^2} \sum_{\substack{(u,v) \in \mathfrak{m} \times \mathfrak{m} \\ \mathfrak{n}(\mathfrak{m})^{-1} \mathfrak{n}(v) \leq s, \ \theta_K u + \theta_K v = \mathfrak{m}}} \Delta_{\frac{u}{v}} \xrightarrow{*} \text{Leb}_{\mathbb{C}} .$$

Let \mathbb{H} be Hamilton's quaternion algebra over \mathbb{R} , with reduced norm $\mathbb{N} : x \mapsto x\bar{x}$. Let A be a quaternion algebra over \mathbb{Q} , which is definite ($A \otimes_{\mathbb{Q}} \mathbb{R} = \mathbb{H}$), with reduced discriminant D_A . Let \mathcal{O} be a maximal order in A , and let \mathfrak{m} be a (nonzero) left ideal of \mathcal{O} , with reduced norm $\mathbb{N}(\mathfrak{m})$ (see [Vig] for definitions). We refer to Equation (14) for a more general result.

Theorem 2 *As s tends to $+\infty$, we have*

$$\frac{\pi^2 \zeta(3) \prod_{p|D_A} (p^3 - 1)}{360 D_A s^4} \sum_{\substack{(u,v) \in \mathfrak{m} \times \mathfrak{m} \\ \mathbb{N}(\mathfrak{m})^{-1} \mathbb{N}(v) \leq s, \ \mathcal{O}u + \mathcal{O}v = \mathfrak{m}}} \Delta_{uv^{-1}} \xrightarrow{*} \text{Leb}_{\mathbb{H}} .$$

The next three statements are equidistribution results on \mathbb{R} or \mathbb{C} of points arithmetically constructed by using quadratic irrationals.

We first prove that the set of traces $\text{tr } \alpha$ of the real quadratic irrationals α over \mathbb{Q} at distance at least $s > 0$ from their Galois conjugate α^σ , in a given orbit $G \cdot \alpha_0$ by homographies of a finite index subgroup G of the modular group $\text{PSL}_2(\mathbb{Z})$, equidistributes to the Lebesgue measure on \mathbb{R} , as $s \rightarrow 0$, using as complexity (the inverse of) the distance to their Galois conjugate (see [PP2, PP4] for background on this height). Given a real integral quadratic irrational α over \mathbb{Q} , we denote by R_α the regulator of $\mathbb{Z} + \alpha\mathbb{Z}$, and by $Q_\alpha(t) = t^2 - (\text{tr } \alpha)t + \mathfrak{n}(\alpha)$ its associated monic quadratic polynomial (the integrality assumption is only used in this introduction, to simplify the notation). We denote by H_x the stabiliser of $x \in \mathbb{P}_1(\mathbb{R}) = \mathbb{R} \cup \{\infty\}$ in a subgroup H of $\text{PSL}_2(\mathbb{R})$.

Theorem 3 *Let α_0 be a real integral quadratic irrational over \mathbb{Q} and let G be a finite index subgroup of $\text{PSL}_2(\mathbb{Z})$. Then, as $\epsilon \rightarrow 0$,*

$$\frac{\pi^2 [\text{PSL}_2(\mathbb{Z}) : G] \epsilon}{6 [\text{PSL}_2(\mathbb{Z})_{\alpha_0} : G_{\alpha_0}] R_{\alpha_0}} \sum_{\alpha \in G \cdot \alpha_0 : |\alpha - \alpha^\sigma| \geq \epsilon} \Delta_{\text{tr } \alpha} \xrightarrow{*} \text{Leb}_{\mathbb{R}} .$$

We refer to Theorems 13 and 15 for extensions of the above result to quadratic irrationals over an imaginary quadratic number field (using relative traces) or over a rational quaternion algebra, and we only quote in this introduction the following special case of Theorem 13.

Corollary 4 *Let $\phi = \frac{1+\sqrt{5}}{2}$ be the Golden Ratio, let K be an imaginary quadratic number field with $D_K \neq -4$, let \mathfrak{c} be a nonzero ideal in \mathcal{O}_K , and let $\Gamma_0(\mathfrak{c})$ be the Hecke congruence subgroup $\left\{ \pm \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{PSL}_2(\mathcal{O}_K) : c \in \mathfrak{c} \right\}$. Then, as $\epsilon \rightarrow 0$,*

$$\frac{|D_K|^{\frac{3}{2}} \zeta_K(2) \mathfrak{n}(\mathfrak{c}) \prod_{p|\mathfrak{c}} \left(1 + \frac{1}{\mathfrak{n}(\mathfrak{p})}\right) \epsilon^2}{4 \pi^2 k_{\mathfrak{c}} \ln \phi} \sum_{\alpha \in \Gamma_0(\mathfrak{c}) \cdot \phi, |\alpha - \alpha^\sigma| \geq \epsilon} \Delta_{\text{tr } \alpha} \xrightarrow{*} \text{Leb}_{\mathbb{C}} .$$

where $k_{\mathfrak{c}}$ is the smallest $k \in \mathbb{N} - \{0\}$ such that the $2k$ -th term of Fibonacci's sequence belongs to \mathfrak{c} .

Given two real quadratic irrationals α, β over \mathbb{Q} , we introduce the *relative height* $h_\alpha(\beta)$ of β with respect to α , measuring how close the (unordered) pair $\{\beta, \beta^\sigma\}$ is to the pair $\{\alpha, \alpha^\sigma\}$, by

$$h_\alpha(\beta) = \frac{\min\{|\beta - \alpha| |\beta^\sigma - \alpha^\sigma|, |\beta - \alpha^\sigma| |\beta^\sigma - \alpha|\}}{|\beta - \beta^\sigma|}.$$

See Equation (25) for an expression of $h_\alpha(\beta)$ using crossratios of $\alpha, \beta, \alpha^\sigma, \beta^\sigma$. Consider the points

$$x_\alpha^\pm(\beta) = \frac{\mathbf{n}(\beta) - \mathbf{n}(\alpha) \pm ((\mathbf{n}(\alpha) - \mathbf{n}(\beta))^2 + (\mathbf{tr} \beta - \mathbf{tr} \alpha)(\mathbf{tr} \beta \mathbf{n}(\alpha) - \mathbf{tr} \alpha \mathbf{n}(\beta)))^{\frac{1}{2}}}{\mathbf{tr} \beta - \mathbf{tr} \alpha}. \quad (1)$$

We will prove that when β varies in an orbit of the modular group $\mathrm{PSL}_2(\mathbb{Z})$, the relative height $h_\alpha(\beta)$ is a well defined complexity modulo the stabiliser of α , and, except finitely many orbits under this stabiliser, the points $x_\alpha^\pm(\beta)$ are well defined and real. The following results says that these points, when β has relative height at most s tending to ∞ , equidistribute to the (unique up to scalar) measure on $\mathbb{R} - \{\alpha_0, \alpha_0^\sigma\}$ which is absolutely continuous with respect to the Lebesgue measure and invariant under the stabiliser of $\{\alpha_0, \alpha_0^\sigma\}$ under $\mathrm{PSL}_2(\mathbb{R})$.

Theorem 5 *Let α, β be real integral quadratic irrationals and let G be a finite index subgroup of $\mathrm{PSL}_2(\mathbb{Z})$. Then, as $s \rightarrow +\infty$,*

$$\frac{\pi^2 [\mathrm{PSL}_2(\mathbb{Z}) : G]}{24 [\mathrm{PSL}_2(\mathbb{Z})_\beta : G_\beta] R_\beta s} \sum_{\beta' \in G \cdot \beta, h_\alpha(\beta') \leq s} \Delta_{x_\alpha^-(\beta')} + \Delta_{x_\alpha^+(\beta')} \xrightarrow{*} \frac{dt}{|Q_\alpha(t)|}.$$

This result, whose proof uses asymptotic properties of crossratios, implies that

$$\mathrm{Card}\{\beta' \in \mathrm{SO}_{Q_\alpha}(\mathbb{Z}) \setminus \mathrm{PSL}_2(\mathbb{Z}) \cdot \beta, h_\alpha(\beta') \leq s\} \sim \frac{48 R_\alpha R_\beta}{\pi^2 |\alpha - \alpha^\sigma|} s.$$

We refer to Theorem 20 for a version with congruences, and to Theorem 21 for an extension of this result to quadratic irrationals over an imaginary quadratic extension of \mathbb{Q} .

Towards the same measure, we also have the following equidistribution result of integral representations of quadratic norm forms (see Section 5.3 for generalisations).

Theorem 6 *If α is a real quadratic irrational over \mathbb{Q} , then, as $s \rightarrow +\infty$,*

$$\frac{\pi^2}{12 s} \sum_{(u,v) \in \mathbb{Z}^2, (u,v)=1, |u^2 - \mathbf{tr} \alpha uv + \mathbf{n}(\alpha) v^2| \leq s} \Delta_{\frac{u}{v}} \xrightarrow{*} \frac{dt}{|Q_\alpha(t)|},$$

Our final equidistribution result for this introduction is the following equidistribution of coefficients of binary Hermitian forms in an orbit under the Bianchi group $\mathrm{SL}_2(\mathcal{O}_K)$, using as complexity their first coefficient (see Subsection 5.1 for extensions, in particular to binary Hamiltonian forms). Given an imaginary quadratic number field K , let $f(u, v) = a|u|^2 + 2\mathrm{Re}(b\bar{u}v) + c|v|^2$ be a binary Hermitian form, which is integral over \mathcal{O}_K (that is $a = a(f), c = c(f) \in \mathbb{Z}$ and $b = b(f) \in \mathcal{O}_K$), and indefinite (that is $\mathrm{Disc}(f) = |b|^2 - ac > 0$). We denote by \cdot the action of $\mathrm{SL}_2(\mathbb{C})$ on the set of binary Hermitian forms by precomposition, and by $\mathrm{SU}_f(\mathcal{O}_K)$ the stabiliser of f in $\mathrm{SL}_2(\mathcal{O}_K)$.

Theorem 7 *Let G be a finite index subgroup of $\mathrm{SL}_2(\mathcal{O}_K)$. As $s \rightarrow +\infty$,*

$$\frac{[\mathrm{SL}_2(\mathcal{O}_K) : G] |D_K|^{\frac{3}{2}} \zeta_K(2) \mathrm{Disc}(f)}{\iota_G \pi^2 \mathrm{Covol}(\mathrm{SU}_f(\mathcal{O}_K) \cap G) s^2} \sum_{f' \in G \cdot f, 0 < |a(f')| \leq s} \Delta_{\frac{b(f')}{a(f')}} \xrightarrow{*} \mathrm{Leb}_{\mathbb{C}} .$$

Identifying \mathcal{O}_K with the upper triangular unipotent subgroup of $\mathrm{SL}_2(\mathcal{O}_K)$, this theorem implies the following counting result of integral binary Hermitian forms (ordered by their first coefficient) in an orbit of the Bianchi group $\mathrm{SL}_2(\mathcal{O}_K)$: as $s \rightarrow +\infty$,

$$\mathrm{Card}\{f' \in \mathcal{O}_K \backslash \mathrm{SL}_2(\mathcal{O}_K) \cdot f : 0 < |a(f')| \leq s\} \sim \frac{\pi^2 \iota_G \mathrm{Covol}(\mathrm{SU}_f(\mathcal{O}_K))}{2 |D_K|^{\frac{1}{2}} \zeta_K(2) \mathrm{Disc}(f)} s^2 .$$

All these theorems come with an error term, and will be (not immediate) consequences of the geometric equidistribution results of [PP8]. The particular case of the result of loc. cit. that we will use, stated in Section 2, is the following one (see also [OhS1, OhS2] for related counting and equidistribution results in real hyperbolic spaces and [Kim, PP9] in negatively curved symmetric spaces) : given a lattice Γ in the isometry group of the real n -dimensional hyperbolic space $\mathbb{H}_{\mathbb{R}}^n$, and two horoballs or totally geodesic subspaces D^-, D^+ whose stabilisers in Γ have cofinite volume, the initial tangent vectors of the common perpendiculars between D_- and the images under Γ of D^+ equidistribute in the unit normal bundle of D^- .

The paper is organised according to the arithmetic applications: In Section 3, we apply Section 2 with both D^- and D^+ horoballs to prove generalisations of the classical Mertens formula, describing the asymptotic behaviour of the average order of Euler's function, for the rings of integers of quadratic imaginary number fields and maximal orders in definite quaternion algebras over \mathbb{Q} . In Section 4, we consider counting and equidistribution of quadratic irrationals in terms of two complexities, the height introduced in [PP2] and the relative height mentioned above. In Subsection 4.1, the geometric result is applied when D^- is a horoball and D^+ is a geodesic line, and in Subsection 4.3 and 4.4 (where we extend Pollicott's result on the asymptotic of crossratios to prove the convergence of generalised Schottky-Klein functions) when both D^- and D^+ are geodesic lines. In the final section, we consider representations of integers by binary quadratic, Hermitian and Hamiltonian forms, applying the geometric result when D^- is a horoball and D^+ is either a geodesic line or a totally geodesic hyperplane or (when considering positive definite forms) a point.

Acknowledgement: The first author thanks the Université Paris-Sud (Orsay) for a visit of a month and a half which allowed most of the writing of this paper, under the financial support of the ERC grant GADA 208091.

2 Geometric counting and equidistribution

In this section, we briefly review a simplified version of the geometric counting and equidistribution results proved in [PP8], whose arithmetic applications will be considered in the other parts of this paper (see also [PP7] for related references and [PP9] for the case of locally symmetric spaces).

Let $n \geq 2$, let Γ be a discrete nonelementary group of isometries of the n -dimensional real hyperbolic space $\mathbb{H}_{\mathbb{R}}^n$, and let $M = \Gamma \backslash \mathbb{H}_{\mathbb{R}}^n$ be the quotient orbifold. Let D^- and D^+

be nonempty proper closed convex subsets in $\mathbb{H}_{\mathbb{R}}^n$, such that the families $(\gamma D^-)_{\gamma \in \Gamma}$ and $(\gamma D^+)_{\gamma \in \Gamma}$ are locally finite in $\mathbb{H}_{\mathbb{R}}^n$ (see [PP8, §3.3] for more general families in any simply connected complete Riemannian manifold with pinched negative curvature).

We denote by $\partial_{\infty} \mathbb{H}_{\mathbb{R}}^n$ the boundary at infinity of $\mathbb{H}_{\mathbb{R}}^n$, by $\Lambda \Gamma$ the limit set of Γ and by $(\xi, x, y) \mapsto \beta_{\xi}(x, y)$ the Busemann cocycle on $\partial_{\infty} \mathbb{H}_{\mathbb{R}}^n \times \mathbb{H}_{\mathbb{R}}^n \times \mathbb{H}_{\mathbb{R}}^n$ defined by

$$(x, y, \xi) \mapsto \beta_{\xi}(x, y) = \lim_{t \rightarrow +\infty} d(\rho_t, x) - d(\rho_t, y) ,$$

where $\rho : t \mapsto \rho_t$ is any geodesic ray with point at infinity ξ and d is the hyperbolic distance.

For every $v \in T^1 \mathbb{H}_{\mathbb{R}}^n$, let $\pi(v) \in \mathbb{H}_{\mathbb{R}}^n$ be its origin, and let v_-, v_+ be the points at infinity of the geodesic line in $\mathbb{H}_{\mathbb{R}}^n$ whose tangent vector at time $t = 0$ is v . We denote by $\partial_{\pm}^1 D^{\mp}$ the *outer/inner unit normal bundle* of ∂D^{\mp} , that is, the set of $v \in T^1 \mathbb{H}_{\mathbb{R}}^n$ such that $\pi(v) \in \partial D^{\mp}$ and the closest point projection on D^{\mp} of $v_{\pm} \in \partial_{\infty} \widetilde{M} - \partial_{\infty} D^{\mp}$ is $\pi(x)$. For every γ, γ' in Γ such that γD^- and $\gamma' D^+$ have a common perpendicular (that is, if the closures $\overline{\gamma D^-}$ and $\overline{\gamma' D^+}$ in $\mathbb{H}_{\mathbb{R}}^n \cup \partial_{\infty} \mathbb{H}_{\mathbb{R}}^n$ are disjoint), we denote by $\alpha_{\gamma, \gamma'}$ this common perpendicular (starting from γD^- at time $t = 0$), by $\ell(\alpha_{\gamma, \gamma'})$ its length, by $v_{\gamma, \gamma'}^- \in \gamma \partial_+^1 D^-$ its initial tangent vector and by $v_{\gamma, \gamma'}^+ \in \gamma' \partial_-^1 D^+$ its terminal tangent vector. Let Γ_{D^-} and Γ_{D^+} be the stabilisers in Γ of the subsets D^- and D^+ , respectively. The *multiplicity* of $\alpha_{\gamma, \gamma'}$ is

$$m_{\gamma, \gamma'} = \frac{1}{\text{Card}(\gamma \Gamma_{D^-} \gamma^{-1} \cap \gamma' \Gamma_{D^+} \gamma'^{-1})} ,$$

which equals 1 when Γ acts freely on $T^1 \mathbb{H}_{\mathbb{R}}^n$ (for instance when Γ is torsion-free). Let

$$\mathcal{N}_{D^-, D^+}(t) = \sum_{\substack{(\gamma, \gamma') \in \Gamma \setminus ((\Gamma/\Gamma_{D^-}) \times (\Gamma/\Gamma_{D^+})) \\ \overline{\gamma D^-} \cap \overline{\gamma' D^+} = \emptyset, \ell(\alpha_{\gamma, \gamma'}) \leq t}} m_{\gamma, \gamma'} = \sum_{\substack{[\gamma] \in \Gamma_{D^-} \setminus \Gamma/\Gamma_{D^+} \\ \overline{D^-} \cap \overline{\gamma D^+} = \emptyset, \ell(\alpha_{e, \gamma}) \leq t}} m_{e, \gamma} ,$$

where Γ acts diagonally on $\Gamma \times \Gamma$. When Γ has no torsion, $\mathcal{N}_{D^-, D^+}(t)$ is the number (with multiplicities coming from the fact that $\Gamma_{D^{\pm}} \setminus D^{\pm}$ is not assumed to be embedded in M) of the common perpendiculars of length at most t between the images of D^- and D^+ in M . We refer to [PP8, §4] for the use of Hölder potentials on $T^1 \mathbb{H}_{\mathbb{R}}^n$ to modify this counting function by adding weights, which could be useful for some further arithmetic applications.

Recall the following notions (see for instance [Rob]). The *critical exponent* of Γ is

$$\delta_{\Gamma} = \limsup_{N \rightarrow +\infty} \frac{1}{N} \ln \text{Card}\{\gamma \in \Gamma : d(x_0, \gamma x_0) \leq N\} ,$$

which is positive, finite and independent of the base point $x_0 \in \mathbb{H}_{\mathbb{R}}^n$. Let $(\mu_x)_{x \in \mathbb{H}_{\mathbb{R}}^n}$ be a *Patterson density* for Γ , that is, a family $(\mu_x)_{x \in \mathbb{H}_{\mathbb{R}}^n}$ of nonzero finite measures on $\partial_{\infty} \mathbb{H}_{\mathbb{R}}^n$ whose support is $\Lambda \Gamma$, such that $\gamma_* \mu_x = \mu_{\gamma x}$ and

$$\frac{d\mu_x}{d\mu_y}(\xi) = e^{-\delta_{\Gamma} \beta_{\xi}(x, y)}$$

for all $\gamma \in \Gamma$, $x, y \in \mathbb{H}_{\mathbb{R}}^n$ and $\xi \in \partial_{\infty} \mathbb{H}_{\mathbb{R}}^n$. The *Bowen-Margulis measure* \tilde{m}_{BM} for Γ on $T^1 \mathbb{H}_{\mathbb{R}}^n$ is defined, using Hopf's parametrisation $v \mapsto (v_-, v_+, \beta_{v_+}(x_0, \pi(v)))$ of $T^1 \mathbb{H}_{\mathbb{R}}^n$, by

$$d\tilde{m}_{\text{BM}}(v) = e^{-\delta_{\Gamma} (\beta_{v_-}(\pi(v), x_0) + \beta_{v_+}(\pi(v), x_0))} d\mu_{x_0}(v_-) d\mu_{x_0}(v_+) dt .$$

The measure \tilde{m}_{BM} is nonzero and independent of $x_0 \in \mathbb{H}_{\mathbb{R}}^n$. It is invariant under the geodesic flow, the antipodal map $v \mapsto -v$ and the action of Γ , and thus defines a nonzero measure m_{BM} on T^1M , called the *Bowen-Margulis measure* on $M = \Gamma \backslash \mathbb{H}_{\mathbb{R}}^n$, which is invariant under the geodesic flow of M and the antipodal map. When m_{BM} is finite, denoting the total mass of a measure m by $\|m\|$, the probability measure $\frac{m_{\text{BM}}}{\|m_{\text{BM}}\|}$ is then uniquely defined, and it is the unique probability measure of maximal entropy for the geodesic flow (see [OtP]). This holds for instance when M has finite volume or when Γ is geometrically finite.

Using the endpoint homeomorphism $v \mapsto v_+$ from $\partial_+^1 D^-$ to $\partial_{\infty} \mathbb{H}_{\mathbb{R}}^n - \partial_{\infty} D^-$, the *skinning measure* $\tilde{\sigma}_{D^-}$ of Γ on $\partial_+^1 D^-$ is defined by

$$d\tilde{\sigma}_{D^-}(v) = e^{-\delta \beta_{v_+}(\pi(v), x_0)} d\mu_{x_0}(v_+),$$

see [OhS1, §1.2] when D^- is a horoball or a totally geodesic subspace in $\mathbb{H}_{\mathbb{R}}^n$ and [PP6], [PP8] for the general case of convex subsets in variable curvature and with a potential.

The measure $\tilde{\sigma}_{D^-}$ is independent of $x_0 \in \mathbb{H}_{\mathbb{R}}^n$, it is nonzero if $\Lambda\Gamma$ is not contained in $\partial_{\infty} D^-$, and satisfies $\tilde{\sigma}_{\gamma D^-} = \gamma_* \tilde{\sigma}_{D^-}$ for every $\gamma \in \Gamma$. Since the family $(\gamma D^-)_{\gamma \in \Gamma}$ is locally finite in $\mathbb{H}_{\mathbb{R}}^n$, the measure $\sum_{\gamma \in \Gamma/\Gamma_{D^-}} \gamma_* \tilde{\sigma}_{D^-}$ is a well defined Γ -invariant locally finite (Borel nonnegative) measure on $T^1 \mathbb{H}_{\mathbb{R}}^n$, hence induces a locally finite measure σ_{D^-} on $T^1 M = \Gamma \backslash T^1 \mathbb{H}_{\mathbb{R}}^n$, called the *skinning measure* of D^- in $T^1 M$. We refer to [OhS2, §5] and [PP6, Theo. 9] for finiteness criteria of the skinning measure σ_{D^-} , in particular satisfied when M has finite volume and if either D^- is a horoball centred at a parabolic fixed point of Γ or if D^- is a totally geodesic subspace.

The following result on the asymptotic behaviour of the counting function \mathcal{N}_{D^-, D^+} in real hyperbolic space is a special case of much more general results [PP8, Coro. 20, 21, Theo. 28]. Furthermore, the equidistribution result of the initial and terminal tangent vectors of the common perpendiculars holds simultaneously in the outer and inner tangent bundles of D^- and D^+ . We refer to [PP7] for a survey of the particular cases known before [PP8] due to Huber, Margulis, Herrmann, Cosentino, Roblin, Oh-Shah, Martin-McKee-Wambach, Pollicott, and the authors for instance.

For every $t \geq 0$, let

$$m_t(x) = \sum_{\gamma \in \Gamma/\Gamma_{D^+} : \overline{D^-} \cap \overline{\gamma D^+} = \emptyset, \alpha_{e, \gamma}(0) = x, \ell(\alpha_{e, \gamma}) \leq t} m_{e, \gamma}$$

be the multiplicity of a point $x \in \partial D^-$ as the origin of common perpendiculars with length at most t from D^- to the elements of the Γ -orbit of D^+ . We denote by Δ_x the unit Dirac mass at a point x .

Theorem 8 *Let Γ, D^-, D^+ be as above. Assume that the measures $m_{\text{BM}}, \sigma_{D^-}, \sigma_{D^+}$ are nonzero and finite. Then*

$$\mathcal{N}_{D^-, D^+}(t) \sim \frac{\|\sigma_{D^-}\| \|\sigma_{D^+}\|}{\delta_{\Gamma} \|m_{\text{BM}}\|} e^{\delta_{\Gamma} t},$$

as $t \rightarrow +\infty$. If Γ is arithmetic or if M is compact, then the error term is $O(e^{(\delta_{\Gamma} - \kappa)t})$ for some $\kappa > 0$. Furthermore, the origins of the common perpendiculars equidistribute in the

boundary of D^- :

$$\lim_{t \rightarrow +\infty} \frac{\delta_\Gamma \|m_{\text{BM}}\|}{\|\sigma_{D^-}\| \|\sigma_{D^+}\|} e^{-\delta_\Gamma t} \sum_{x \in \partial D^-} m_t(x) \Delta_x = \frac{\pi_* \tilde{\sigma}_{D^-}}{\|\sigma_{D^-}\|} \quad (2)$$

for the weak-star convergence of measures on the locally compact space $T^1 \mathbb{H}_{\mathbb{R}}^n$. \square

For smooth functions ψ with compact support on ∂D^- , there is an error term in the above equidistribution claim, of the form $O(s^{\delta_\Gamma - \kappa} \|\psi\|_\ell)$ where $\kappa > 0$ and $\|\psi\|_\ell$ is the Sobolev norm of ψ for some $\ell \in \mathbb{N}$, as proved in [PP8, Theo. 28].

When M has finite volume, we have $\delta_\Gamma = n - 1$, the Bowen-Margulis measure m_{BM} coincides up to a multiplicative constant with the Liouville measure on $T^1 M$, and the skinning measures of points, horoballs and totally geodesic subspaces D^\pm coincide again up to a multiplicative constant with the (homogeneous) Riemannian measures on $\partial_\pm^1 D^\mp$ induced by the Riemannian metric of $T^1 \mathbb{H}_{\mathbb{R}}^n$. These proportionality constants were computed in [PP7, §7] and [PP8, Prop. 29]:

$$\|m_{\text{BM}}\| = 2^{n-1} (n-1) \text{Vol}(\mathbb{S}^{n-1}) \text{Vol}(M),$$

if D^- is a horoball, then $\|\sigma_{D^-}\| = 2^{n-1} (n-1) \text{Vol}(\Gamma_{D^-} \setminus D^-)$, and if D^- is a totally geodesic submanifold in $\mathbb{H}_{\mathbb{R}}^n$ of dimension $k^- \in \{0, \dots, n-1\}$ with pointwise stabiliser of order m^- , then $\tilde{\sigma}_{D^-} = \text{Vol}_{\partial_+^1 D^-}$, so that $\|\sigma_{D^-}\| = \frac{\text{Vol}(\mathbb{S}^{n-k^- - 1})}{m^-} \text{Vol}(\Gamma_{D^-} \setminus D^-)$. See [PP9, §3] for the computation of the proportionality constants in the complex hyperbolic case.

Using these explicit expressions, we now reformulate Theorem 8 using these explicit expressions, considering the following cases.

(1) If D^- and D^+ are totally geodesic submanifolds in $\mathbb{H}_{\mathbb{R}}^n$ of dimensions k^- and k^+ in $\{1, \dots, n-1\}$, respectively, such that $\text{Vol}(\Gamma_{D^-} \setminus D^-)$ and $\text{Vol}(\Gamma_{D^+} \setminus D^+)$ are finite, let

$$c(D^-, D^+) = \frac{\text{Vol}(\mathbb{S}^{n-k^- - 1}) \text{Vol}(\mathbb{S}^{n-k^+ - 1}) \text{Vol}(\Gamma_{D^-} \setminus D^-) \text{Vol}(\Gamma_{D^+} \setminus D^+)}{2^{n-1} (n-1) \text{Vol}(\mathbb{S}^{n-1}) \text{Vol}(M)}.$$

(2) If D^- and D^+ are horoballs in $\mathbb{H}_{\mathbb{R}}^n$ centred at parabolic fixed points of Γ , let

$$c(D^-, D^+) = \frac{2^{n-1} (n-1) \text{Vol}(\Gamma_{D^-} \setminus D^-) \text{Vol}(\Gamma_{D^+} \setminus D^+)}{\text{Vol}(\mathbb{S}^{n-1}) \text{Vol}(M)}.$$

(3) If D^- is a horoball in $\mathbb{H}_{\mathbb{R}}^n$ centred at a parabolic fixed point of Γ and D^+ is a totally geodesic subspace in $\mathbb{H}_{\mathbb{R}}^n$ of dimension $k^+ \in \{1, \dots, n-1\}$ such that $\text{Vol}(\Gamma_{D^+} \setminus D^+)$ is finite, let

$$c(D^-, D^+) = \frac{\text{Vol}(\mathbb{S}^{n-k^+ - 1}) \text{Vol}(\Gamma_{D^-} \setminus D^-) \text{Vol}(\Gamma_{D^+} \setminus D^+)}{\text{Vol}(\mathbb{S}^{n-1}) \text{Vol}(M)}.$$

(4) If D^- is a horoball in $\mathbb{H}_{\mathbb{R}}^n$ centred at a parabolic fixed point of Γ and D^+ is a point of $\mathbb{H}_{\mathbb{R}}^n$, let

$$c(D^-, D^+) = \frac{\text{Vol}(\Gamma_{D^-} \setminus D^-)}{\text{Vol}(M)}.$$

Corollary 9 *Let Γ be a discrete group of isometries of $\mathbb{H}_{\mathbb{R}}^n$ such that the orbifold $M = \Gamma \backslash \mathbb{H}_{\mathbb{R}}^n$ has finite volume. In each of the cases (1) to (4) above, if m^{\pm} is the cardinality of the pointwise fixator of D^{\pm} , then*

$$\mathcal{N}_{D^-, D^+}(t) \sim \frac{c(D^-, D^+)}{m^- m^+} e^{(n-1)t}.$$

If Γ is arithmetic or if M is compact, then there exists $\kappa > 0$ such that, as $t \rightarrow +\infty$,

$$\mathcal{N}_{D^-, D^+}(t) = \frac{c(D^-, D^+)}{m^- m^+} e^{(n-1)t} (1 + O(e^{-\kappa t})).$$

Furthermore, the origins of the common perpendiculars equidistribute in ∂D^- to the induced Riemannian measure: if D^- is a horoball centred at a parabolic fixed point of Γ , then

$$\frac{m^+ (n-1) \text{Vol}(\Gamma_{D^-} \backslash D^-)}{c(D^-, D^+)} e^{-(n-1)t} \sum_{x \in \partial D^-} m_t(x) \Delta_x \xrightarrow{*} \text{Vol}_{\partial D^-}, \quad (3)$$

as $t \rightarrow +\infty$, and if D^- is a totally geodesic subspace in $\mathbb{H}_{\mathbb{R}}^n$ of dimension $k^- \in \{1, \dots, n-1\}$, such that $\text{Vol}(\Gamma_{D^-} \backslash D^-)$ is finite, then as $t \rightarrow +\infty$,

$$\frac{m^- m^+ \text{Vol}(\Gamma_{D^-} \backslash D^-)}{c(D^-, D^+)} e^{-(n-1)t} \sum_{x \in D^-} m_t(x) \Delta_x \xrightarrow{*} \text{Vol}_{D^-}. \quad \square \quad (4)$$

Again, for smooth functions ψ with compact support on ∂D^- , there is an error term in the above two equidistribution claims, of the form $O(s^{n-1-\kappa} \|\psi\|_{\ell})$ where $\kappa > 0$ and $\|\psi\|_{\ell}$ is the Sobolev norm of ψ for some $\ell \in \mathbb{N}$.

We end this section by recalling some terminology concerning the isometries of $\mathbb{H}_{\mathbb{R}}^n$. The *translation length* of an isometry γ of $\mathbb{H}_{\mathbb{R}}^n$ is

$$\ell(\gamma) = \inf_{x \in \mathbb{H}_{\mathbb{R}}^n} d(x, \gamma x).$$

If $\ell(\gamma) > 0$, then γ is *loxodromic*. Each loxodromic isometry γ stabilises a unique geodesic line in $\mathbb{H}_{\mathbb{R}}^n$, called the *translation axis* of γ and denoted by $\text{Axis } \gamma$, on which it acts as a translation by $\ell(\gamma)$. The points at infinity of $\text{Axis } \gamma$ are the two fixed points of γ in $\partial_{\infty} \mathbb{H}_{\mathbb{R}}^n$, and we denote by γ^- and γ^+ the attracting and repelling fixed points of γ respectively. Any loxodromic element of Γ is contained in a maximal cyclic subgroup.

An element $\gamma \in \Gamma$ is *primitive* if the cyclic subgroup $\gamma^{\mathbb{Z}}$ it generates is a maximal cyclic subgroup of Γ . A loxodromic element $\gamma \in \Gamma$ is Γ -*reciprocal* if there is an element in Γ that switches the two fixed points of γ . If γ is Γ -reciprocal, then let $\iota_{\Gamma}(\gamma) = 2$, otherwise, we set $\iota_{\Gamma}(\gamma) = 1$. If γ is a primitive loxodromic element of Γ , then the stabiliser of $\text{Axis } \gamma$ is generated by γ , an elliptic element that switches the two points at infinity of the axis of γ if γ is reciprocal, and a (possibly trivial) group of finite order $m_{\Gamma}(\gamma)$, which is the pointwise stabiliser of $\text{Axis } \gamma$, so that

$$m_{\Gamma}(\gamma) = \frac{1}{\iota_{\Gamma}(\gamma)} [\text{Stab}_{\Gamma}(\text{Axis } \gamma) : \gamma^{\mathbb{Z}}]. \quad (5)$$

3 Generalised Mertens' formulas

A classical result, known as *Mertens' formula* (see for example [HaW, Thm. 330], a better error term is due to Walfisz [Wal]), describes the asymptotic behaviour of the average order $\Phi : \mathbb{N} - \{0\} \rightarrow \mathbb{N} - \{0\}$ of Euler's function φ , where $\varphi(n) = \text{Card}((\mathbb{Z}/n\mathbb{Z})^\times)$:

$$\Phi(n) = \sum_{k=1}^n \varphi(k) = \frac{3}{\pi^2} n^2 + O(n \ln n).$$

Corollary 9 provides a geometric proof of Mertens' formula with a less explicit error term, as follows. The modular group $\Gamma_{\mathbb{Q}} = \text{PSL}_2(\mathbb{Z})$ isometrically acts on the upper half plane model of the real hyperbolic plane $\mathbb{H}_{\mathbb{R}}^2$ via Möbius transformations. Let \mathcal{H}_{∞} be the horoball in $\mathbb{H}_{\mathbb{R}}^2$ that consists of all points with vertical coordinate at least 1, and let Γ_{∞} be its stabiliser in $\Gamma_{\mathbb{Q}}$. The images of \mathcal{H}_{∞} under $\Gamma_{\mathbb{Q}}$ different from \mathcal{H}_{∞} are the disks in $\mathbb{H}_{\mathbb{R}}^2$ tangent to the real line at the rational points $\frac{p}{q}$ (where $q > 0$ and $(p, q) = 1$) with Euclidean diameter $\frac{1}{q^2}$. The common perpendicular from \mathcal{H}_{∞} to this disc exists if and only if $q > 1$, its length is $\ln q^2$, and its multiplicity is 1. For every $n \in \mathbb{N} - \{0\}$, the cardinality of the set of rational points $\frac{p}{q} \in]0, 1]$ with $q \leq n$ is $\Phi(n)$. Thus, $\Phi(n) = \mathcal{N}_{\mathcal{H}_{\infty}, \mathcal{H}_{\infty}}(\ln n^2) + 1$. Now $\text{Vol}(\Gamma_{\infty} \backslash \mathcal{H}_{\infty}) = 1$ and $\text{Vol}(\Gamma_{\mathbb{Q}} \backslash \mathbb{H}_{\mathbb{R}}^2) = \frac{\pi}{3}$, and we can apply Corollary 9 to conclude that for some $\kappa > 0$, as $n \rightarrow +\infty$,

$$\Phi(n) = \frac{3}{\pi^2} n^2 + O(n^{2-\kappa}). \quad (6)$$

Let us reformulate Mertens' formula in a way that is easily extendable in more general contexts (see also [PP9]). Let the additive group \mathbb{Z} act on $\mathbb{Z} \times \mathbb{Z}$ by horizontal shears (transvections): $k \cdot (u, v) = (u + kv, v)$, and define

$$\psi(s) = \text{Card } \mathbb{Z} \backslash \{(u, v) \in \mathbb{Z} \times \mathbb{Z} : (u, v) = 1, |v| \leq s\}.$$

We easily have $\psi(n) = 2\Phi(n) + 2$ for every $n \in \mathbb{N} - \{0\}$, so that Mertens' formula (6) is equivalent with $\psi(n) = \frac{6}{\pi^2} s + O(s^{1-\kappa})$. Furthermore, a straightforward application of Corollary 9 shows that as $s \rightarrow +\infty$, we have

$$\frac{\pi^2}{3s} \sum_{(u,v)=1, 1 \leq v \leq s} \Delta_{(\frac{u}{v}, 1)} \xrightarrow{*} \text{Vol}_{\partial \mathcal{H}_{\infty}}.$$

Observing that the pushforward of the measures on $\partial \mathcal{H}_{\infty}$ by the map $(x, 1) \mapsto x$ from $\partial \mathcal{H}_{\infty}$ to \mathbb{R} is linear, is continuous for the weak-star topology, maps the unit Dirac mass at a point p to the unit Dirac mass at $f(p)$ and the volume measure of $\partial \mathcal{H}_{\infty}$ to the Lebesgue measure, we get the well-known result on the equidistribution of the Farey fractions: as $s \rightarrow +\infty$, we have

$$\frac{\pi^2}{6s} \sum_{(u,v)=1, |v| \leq s} \Delta_{\frac{u}{v}} \xrightarrow{*} \text{Leb}_{\mathbb{R}}.$$

In subsections 3.1 and 3.2, we generalise the above results to quadratic imaginary number fields and definite quaternion algebras over \mathbb{Q} .

3.1 A Mertens' formula for the rings of integers of imaginary quadratic number fields

Let K be an imaginary quadratic number field, with ring of integers \mathcal{O}_K , discriminant D_K , zeta function ζ_K and norm \mathbf{n} . Let $\omega_K = \text{Card}(\mathcal{O}_K^\times)$ (with $\omega_K = 2$ if $D_K \neq -3, -4$ for future use). Let \mathfrak{m} be a (nonzero) fractional ideal of \mathcal{O}_K , with norm $\mathbf{n}(\mathfrak{m})$. Note that the action of the additive group \mathcal{O}_K on $\mathbb{C} \times \mathbb{C}$ by the horizontal shears $k \cdot (u, v) = (u + kv, v)$ preserves $\mathfrak{m} \times \mathfrak{m}$.

We consider the counting function $\psi_{\mathfrak{m}} : [0, +\infty[\rightarrow \mathbb{N}$ defined by

$$\psi_{\mathfrak{m}}(s) = \text{Card } \mathcal{O}_K \setminus \{(u, v) \in \mathfrak{m} \times \mathfrak{m} : \mathbf{n}(\mathfrak{m})^{-1} \mathbf{n}(v) \leq s, \mathcal{O}_K u + \mathcal{O}_K v = \mathfrak{m}\}.$$

Note that $\psi_{\mathfrak{m}}$ depends only on the ideal class of \mathfrak{m} and thus we can assume in the computations that \mathfrak{m} is integral. The *Euler function* φ_K of K is defined on the set of (nonzero) integral ideals \mathfrak{a} of \mathcal{O}_K by $\varphi_K(\mathfrak{a}) = \text{Card}((\mathcal{O}_K/\mathfrak{a})^\times)$, see for example [Coh1, §4A]. Thus,

$$\psi_{\mathcal{O}_K}(s) = \sum_{v \in \mathcal{O}_K, 0 < \mathbf{n}(v) \leq s} \varphi_K(v \mathcal{O}_K),$$

and the first claim of the following result, in the special case $\mathfrak{m} = \mathcal{O}_K$, is an analog of Mertens' formula, due to [Gro, Satz 2] (with a better error term), see also [Cos, §4.3]. The equidistribution part of Theorem 10 (stated as Theorem 1 in the introduction) generalises [Cos, Th. 4] which covers the case $\mathfrak{m} = \mathcal{O}_K$ without an explicit proportionality constant but with an explicit estimate on the speed of equidistribution. Note that the ideal class group of \mathcal{O}_K is in general nontrivial, hence the extension to general \mathfrak{m} is interesting.

Theorem 10 *There exists $\kappa > 0$ such that, as $s \rightarrow +\infty$,*

$$\psi_{\mathfrak{m}}(s) = \frac{\pi}{\zeta_K(2) \sqrt{|D_K|}} s^2 + O(s^{2-\kappa}).$$

Furthermore, as $s \rightarrow +\infty$,

$$\frac{|D_K| \zeta_K(2)}{2\pi s^2} \sum_{\substack{(u,v) \in \mathfrak{m} \times \mathfrak{m} \\ \mathbf{n}(\mathfrak{m})^{-1} \mathbf{n}(v) \leq s, \mathcal{O}_K u + \mathcal{O}_K v = \mathfrak{m}}} \Delta_{\frac{u}{v}} \xrightarrow{*} \text{Leb}_{\mathbb{C}},$$

with error term $O(s^{2-\kappa} \|\psi\|_\ell)$ when evaluated on \mathcal{C}^ℓ -smooth functions ψ with compact support on \mathbb{C} , for ℓ big enough.

Proof. The projective action of $\text{PSL}_2(\mathbb{C})$ on the Riemann sphere $\mathbb{P}^1(\mathbb{C}) = \mathbb{C} \cup \{\infty\}$ identifies by the Poincaré extension with the group of orientation preserving isometries of the upper halfspace model of the real hyperbolic space $\mathbb{H}_{\mathbb{R}}^3$. We denote the image in $\text{PSL}_2(\mathbb{C})$ of any subgroup G of $\text{SL}_2(\mathbb{C})$ by \overline{G} and of any element g of $\text{SL}_2(\mathbb{C})$ again by g . The *Bianchi group* $\Gamma_K = \text{SL}_2(\mathcal{O}_K)$ is a (nonuniform) arithmetic lattice in $\text{SL}_2(\mathbb{C})$, whose covolume is given by Humbert's formula (see [EGM, §8.8 and 9.6])

$$\text{Vol}(\overline{\Gamma_K} \backslash \mathbb{H}_{\mathbb{R}}^3) = \frac{|D_K|^{\frac{3}{2}} \zeta_K(2)}{4\pi^2}. \quad (7)$$

Let $\Gamma_{x,y}$ be the stabiliser of $(x, y) \in \mathcal{O}_K \times \mathcal{O}_K$ under the linear action of Γ_K on $\mathcal{O}_K \times \mathcal{O}_K$. In particular, $\Gamma_{1,0}$ is the upper triangular unipotent subgroup of Γ_K , acting on $\mathcal{O}_K \times \mathcal{O}_K$ via

horizontal shears. Furthermore, $x\mathcal{O}_K + y\mathcal{O}_K = u\mathcal{O}_K + v\mathcal{O}_K$ if and only if $(u, v) \in \Gamma_K(x, y)$, and the ideal class group of K corresponds bijectively to the set $\overline{\Gamma_K} \backslash \mathbb{P}^1(K)$ of cusps of the orbifold $\overline{\Gamma_K} \backslash \mathbb{H}_{\mathbb{R}}^3$ by the map induced by $x\mathcal{O}_K + y\mathcal{O}_K \mapsto \rho = \frac{x}{y} \in K \cup \{\infty\}$, see for example Section 7.2 of [EGM] for details.

To prove the first claim (it follows by integration from the second one, but the essential ingredients of the proof of either statement are the same), note that by the above, if $\mathfrak{m} = x\mathcal{O}_K + y\mathcal{O}_K$, then

$$\psi_{\mathfrak{m}}(s) = \text{Card } \Gamma_{1,0} \backslash \{(u, v) \in \Gamma_K(x, y) : |v|^2 \leq \mathfrak{n}(\mathfrak{m})s\}.$$

If $y = 0$, let $\gamma_\rho = \text{id}$ and we may assume that $x = 1$ since $\psi_{\mathfrak{m}}$ depends only on the ideal class of \mathfrak{m} . If $y \neq 0$, let

$$\gamma_\rho = \begin{pmatrix} \rho & -1 \\ 1 & 0 \end{pmatrix} \in \text{SL}_2(\mathbb{C}).$$

Let $\tau \in]0, 1]$. Let \mathcal{H}_∞ be the horoball in $\mathbb{H}_{\mathbb{R}}^3$ that consists of all points with Euclidean height at least $1/\tau$ and let $\mathcal{H}_\rho = \gamma_\rho \mathcal{H}_\infty$. For any $\rho' \in K \cup \{\infty\}$, let $\overline{\Gamma_{\mathcal{H}_{\rho'}}}$ be the stabiliser in $\overline{\Gamma_K}$ of the horoball $\mathcal{H}_{\rho'}$. As in [PP3] after its Equation (4), we fix τ small enough such that \mathcal{H}_∞ and \mathcal{H}_ρ are precisely invariant under $\overline{\Gamma_K}$ (that is, it meets one of its image by an element of $\overline{\Gamma_K}$ only if it coincides with it).

For any $g \in \text{SL}_2(\mathbb{C})$ such that $g\mathcal{H}_\infty$ and \mathcal{H}_∞ are disjoint, it is easy to check using the explicit expression of the Poincaré extension (see for example [Bea, Eq. 4.1.4.]) that the length of the common perpendicular of \mathcal{H}_∞ and $g\mathcal{H}_\infty$ is $|\ln(\tau^{-2}|c|^2)|$, where c is the $(2, 1)$ -entry of g . If $y \neq 0$, then $(u, v) = g(x, y)$ if and only if $(\frac{u}{y}, \frac{v}{y}) = g\gamma_\rho(1, 0)$, and the $(2, 1)$ -entry of $g\gamma_\rho$ is $\frac{v}{y}$. Thus, the length $\ell(\delta_g)$ of the common perpendicular δ_g between \mathcal{H}_∞ and $g\mathcal{H}_\rho = g\gamma_\rho \mathcal{H}_\infty$, if these two horoballs are disjoint, is $|\ln(\frac{|v|^2}{\tau^2|y|^2})|$ if $y \neq 0$ and $|\ln(\tau^{-2}|v|^2)|$ otherwise.

For the rest of the proof, we concentrate on the case $y \neq 0$. The case $y = 0$ is treated similarly. By discreteness, there are only finitely many double classes $[g] \in \Gamma_{1,0} \backslash \Gamma_K / \Gamma_{x,y}$ such that \mathcal{H}_∞ and $g\mathcal{H}_\rho$ are not disjoint or such that $|v| \leq |y|$ or such that the multiplicity of δ_g is different from 1. Since the stabilisers $\Gamma_{1,0}$ and $\Gamma_{x,y}$ do not contain $-\text{id}$, we have

$$\begin{aligned} \psi_{\mathfrak{m}}(s) &= \text{Card } \{[g] \in \Gamma_{1,0} \backslash \Gamma_K / \Gamma_{x,y} : \ell(\delta_g) \leq \ln \frac{\mathfrak{n}(\mathfrak{m})s}{\tau^2|y|^2}\} + O(1) \\ &= 2 \text{Card } \{[g] \in \overline{\Gamma_{1,0}} \backslash \overline{\Gamma_K} / \overline{\Gamma_{x,y}} : \ell(\delta_g) \leq \ln \frac{\mathfrak{n}(\mathfrak{m})s}{\tau^2|y|^2}\} + O(1). \end{aligned}$$

We use [PP3, Lem. 7] with $C = \overline{\Gamma_K}$, $A = \overline{\Gamma_{1,0}}$, $A' = \overline{\Gamma_{\mathcal{H}_\infty}}$, $B = \overline{\Gamma_{x,y}}$, $B' = \overline{\Gamma_{\mathcal{H}_\rho}}$, since there are only finitely many $[g] \in A' \backslash C / B'$ such that $g^{-1}A'g \cap B' \neq \{1\}$. Since $[\overline{\Gamma_{\mathcal{H}_\infty}} : \overline{\Gamma_{1,0}}] = [\overline{\Gamma_{\mathcal{H}_\rho}} : \overline{\Gamma_{x,y}}] = \frac{\omega_K}{2}$, we then have

$$\psi_{\mathfrak{m}}(s) = \frac{\omega_K^2}{2} \text{Card} \{[g] \in \overline{\Gamma_{\mathcal{H}_\infty}} \backslash \overline{\Gamma_K} / \overline{\Gamma_{\mathcal{H}_\rho}} : \ell(\delta_g) \leq \ln \frac{\mathfrak{n}(\mathfrak{m})s}{\tau^2|y|^2}\} + O(1).$$

By [PP3, Lem. 6] and the last equality of the proof of Theorem 4 on page 1055 of loc. cit., we have $\text{Vol}(\overline{\Gamma_{\mathcal{H}_\infty}} \backslash \mathcal{H}_\infty) = \frac{\tau^2 \sqrt{|D_K|}}{2\omega_K}$ and $\text{Vol}(\overline{\Gamma_{\mathcal{H}_\rho}} \backslash \mathcal{H}_\rho) = \frac{\tau^2 \sqrt{|D_K|}}{2\omega_K} \frac{|y|^4}{\mathfrak{n}(\mathfrak{m})^2}$. Thus,

Corollary 9, applied to $n = 3$, $\Gamma = \overline{\Gamma_K}$, $D^- = \mathcal{H}_\infty$ and $D^+ = \mathcal{H}_\rho$, gives, since the pointwise stabilisers of \mathcal{H}_∞ and \mathcal{H}_ρ are trivial,

$$\psi_{\mathfrak{m}}(s) = \frac{\omega_K^2}{2} \frac{2^2 2 \tau^4 |D_K| |y|^4 4\pi^2}{4 \omega_K^2 \mathfrak{n}(\mathfrak{m})^2 4\pi |D_K|^{\frac{3}{2}} \zeta_K(2) \tau^4 |y|^4} \frac{\mathfrak{n}(\mathfrak{m})^2}{s^2} (1 + O(s^{-\kappa})),$$

which, after simplification, proves the first claim.

To prove the second claim when $y \neq 0$, note that Equation (3) in Corollary 9 implies that as $t \rightarrow +\infty$, with the appropriate error term,

$$\frac{|D_K| \zeta_K(2) \omega_K}{2\pi} \frac{\mathfrak{n}(\mathfrak{m})^2}{\tau^2 |y|^4} e^{-2t} \sum_{z \in \partial \mathcal{H}_\infty} m_t(z) \Delta_z \xrightarrow{*} \text{Vol}_{\partial \mathcal{H}_\infty}.$$

Using the change of variable $t = \ln \frac{\mathfrak{n}(\mathfrak{m})s}{\tau^2 |y|^2}$, the fact that the geodesic lines containing the common perpendiculars from \mathcal{H}_∞ to the images of \mathcal{H}_ρ by the elements of $\overline{\Gamma_K}$ are exactly the geodesic lines from ∞ to an element of $\overline{\Gamma_K} \cdot \frac{x}{y}$, and the above facts on disjointness, lengths and multiplicities, this gives, since the map $(u, v) \mapsto u/v$ is ω_K -to-1,

$$\frac{|D_K| \zeta_K(2) \omega_K \tau^2}{2\pi s^2} \frac{1}{\omega_K} \sum_{\substack{(u,v) \in \mathfrak{m} \times \mathfrak{m} \\ \mathfrak{n}(\mathfrak{m})^{-1} |v|^2 \leq s, \ \mathcal{O}_K u + \mathcal{O}_K v = \mathfrak{m}}} \Delta_{(\frac{u}{v}, \frac{1}{\tau})} \xrightarrow{*} \text{Vol}_{\partial \mathcal{H}_\infty},$$

as $s \rightarrow +\infty$. The claim now follows from the observation that the pushforward of $\text{Vol}_{\partial \mathcal{H}_\infty}$ by the endpoint map $(x, \frac{1}{\tau}) \mapsto x$ is $\tau^2 \text{Leb}_{\mathbb{C}}$. The proof of the case $y = 0$ is similar. \square

3.2 A Mertens' formula for maximal orders of rational quaternion algebras

Let \mathbb{H} be Hamilton's quaternion algebra over \mathbb{R} , with $x \mapsto \bar{x}$ its conjugation, $\mathbf{N} : x \mapsto x\bar{x}$ its reduced norm and $\mathbf{Tr} : x \mapsto x + \bar{x}$ its reduced trace. We endow \mathbb{H} with its standard Euclidean structure (making its standard basis orthonormal). Let A be a quaternion algebra over \mathbb{Q} , which is definite ($A \otimes_{\mathbb{Q}} \mathbb{R} = \mathbb{H}$), with reduced discriminant D_A . Let \mathcal{O} be a maximal order in A , and let \mathfrak{m} be a (nonzero) left ideal of \mathcal{O} , with reduced norm $\mathbf{N}(\mathfrak{m})$ (see [Vig] for definitions).

The additive group \mathcal{O} acts on the left on $\mathbb{H} \times \mathbb{H}$ by the horizontal shears (transvections) $o \cdot (u, v) = (u + ov, v)$. Let us consider the following counting function of the generating pairs of elements of \mathfrak{m} , defined for every $s > 0$, by

$$\psi_{\mathfrak{m}}(s) = \text{Card } \mathcal{O} \setminus \{(u, v) \in \mathfrak{m} \times \mathfrak{m} : \mathbf{N}(\mathfrak{m})^{-1} \mathbf{N}(v) \leq s, \ \mathcal{O}u + \mathcal{O}v = \mathfrak{m}\},$$

Theorem 11 *There exists $\kappa > 0$ such that, as s tends to $+\infty$,*

$$\psi_{\mathfrak{m}}(s) = \frac{90 D_A^2}{\pi^2 \zeta(3) \prod_{p|D_A} (p^3 - 1)} s^4 + O(s^{4-\kappa}),$$

with p ranging over positive rational primes. Furthermore,

$$\frac{\pi^2 \zeta(3) \prod_{p|D_A} (p^3 - 1)}{360 D_A s^4} \sum_{\substack{(u,v) \in \mathfrak{m} \times \mathfrak{m} \\ \mathbf{N}(\mathfrak{m})^{-1} \mathbf{N}(v) \leq s, \ \mathcal{O}u + \mathcal{O}v = \mathfrak{m}}} \Delta_{uv^{-1}} \xrightarrow{*} \text{Leb}_{\mathbb{H}}$$

as $s \rightarrow +\infty$, with error term $O(s^{4-\kappa}\|\psi\|_\ell)$ when evaluated on \mathcal{C}^ℓ -smooth functions ψ with compact support on \mathbb{H} , for ℓ big enough.

Proof. We denote by $[\mathfrak{m}]$ the ideal class of a left fractional ideal \mathfrak{m} of \mathcal{O} , and by $\mathcal{O}\mathcal{I}$ the set of left ideal classes of \mathcal{O} , and by $\mathcal{O}_r(I) = \{x \in A : Ix \subset I\}$ the right order of a \mathbb{Z} -lattice I in A (see [Vig] for definitions). Note that $\psi_{\mathfrak{m}}$ depends only on $[\mathfrak{m}]$. For every (u, v) in $A \times A - \{(0, 0)\}$, consider the two left fractional ideals of \mathcal{O}

$$I_{u,v} = \mathcal{O}u + \mathcal{O}v, \quad K_{u,v} = \begin{cases} \mathcal{O}u \cap \mathcal{O}v & \text{if } uv \neq 0, \\ \mathcal{O} & \text{otherwise.} \end{cases}$$

Given $\mathfrak{m}, \mathfrak{m}'$ two left fractional ideals of \mathcal{O} and $s > 0$, let

$$\psi_{\mathfrak{m}, \mathfrak{m}'}(s) = \text{Card } \mathcal{O} \setminus \{(u, v) \in \mathfrak{m} \times \mathfrak{m}' : N(v) \leq N(\mathfrak{m})s, \quad I_{u,v} = \mathfrak{m}, [K_{u,v}] = [\mathfrak{m}']\},$$

so that

$$\psi_{\mathfrak{m}} = \sum_{[\mathfrak{m}'] \in \mathcal{O}\mathcal{I}} \psi_{\mathfrak{m}, \mathfrak{m}'} . \quad (8)$$

We will give in Equation (12) an asymptotic to $\psi_{\mathfrak{m}, \mathfrak{m}'}(s)$ as $s \rightarrow +\infty$, and in Equation (14) the related equidistribution result, for each $[\mathfrak{m}'] \in \mathcal{O}\mathcal{I}$ (interesting in themselves), and then easily infer Theorem 11.

We refer for instance to [Kel], [PP5, §3] for the following properties of quaternionic homographies. Let $\text{SL}_2(\mathbb{H})$ be the group of 2×2 matrices with coefficients in \mathbb{H} and Dieudonné determinant 1. Recall that the Dieudonné determinant of $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ is

$$N(ad) + N(bc) - \text{Tr}(a\bar{c}d\bar{b}). \quad (9)$$

The group $\text{SL}_2(\mathbb{H})$ acts linearly on the left on the right \mathbb{H} -module $\mathbb{H} \times \mathbb{H}$, hence projectively on the left on the right projective line $\mathbb{P}_r^1(\mathbb{H}) = (\mathbb{H} \times \mathbb{H} - \{0\})/\mathbb{H}^\times$. The group $\text{PSL}_2(\mathbb{H}) = \text{SL}_2(\mathbb{H})/\{\pm \text{Id}\}$ identifies by the Poincaré extension procedure with the group of orientation preserving isometries of the upper halfspace model $\mathbb{H} \times]0, +\infty[$ of the real hyperbolic space $\mathbb{H}_{\mathbb{R}}^5$ of dimension 5, with Riemannian metric

$$ds^2(x) = \frac{ds_{\mathbb{H}}^2(z) + dr^2}{r^2}$$

at the point $x = (z, r)$. For any subgroup G of $\text{SL}_2(\mathbb{H})$, we denote by \overline{G} its image in $\text{PSL}_2(\mathbb{H})$.

Let $\Gamma_{\mathcal{O}} = \text{SL}_2(\mathcal{O}) = \text{SL}_2(\mathbb{H}) \cap \mathcal{M}_2(\mathcal{O})$ be the *Hamilton-Bianchi group* of \mathcal{O} , which is a (nonuniform) arithmetic lattice in the connected real Lie group $\text{SL}_2(\mathbb{H})$ (see for instance [PP1, page 1104]). Given $(x, y) \in \mathcal{O} \times \mathcal{O}$, let $\Gamma_{x,y}$ be the stabiliser of (x, y) for the left linear action of $\Gamma_{\mathcal{O}}$. By [PP5, Rem. 7], the map from the set $\overline{\Gamma_{\mathcal{O}}} \setminus \mathbb{P}_r^1(\mathcal{O})$ of cusps of $\overline{\Gamma_{\mathcal{O}}} \setminus \mathbb{H}_{\mathbb{R}}^5$ into $\mathcal{O}\mathcal{I} \times \mathcal{O}\mathcal{I}$ which associates, to the orbit of $[u : v]$ in $\mathbb{P}_r^1(\mathcal{O})$ under $\overline{\Gamma_{\mathcal{O}}}$, the pair of ideal classes $([I_{u,v}], [K_{u,v}])$ is a bijection. We hence fix a (nonzero) element $(x, y) \in \mathcal{O} \times \mathcal{O}$ such that $[I_{x,y}] = [\mathfrak{m}]$ and $[K_{x,y}] = [\mathfrak{m}']$, assuming that $x = 1$ if $y = 0$.

Let us prove the first claim. Since the reduced norm of an invertible element of \mathcal{O} is 1, the index in $\Gamma_{1,0}$ of its unipotent upper triangular subgroup is $|\mathcal{O}^\times|$, and we have

$$\psi_{\mathfrak{m}, \mathfrak{m}'}(s) = |\mathcal{O}^\times| \text{Card } \Gamma_{1,0} \setminus \{(u, v) \in \Gamma_{\mathcal{O}}(x, y) : N(v) \leq N(\mathfrak{m})s\}.$$

Let $\tau \in]0, 1]$. Let \mathcal{H}_∞ be the horoball in $\mathbb{H}_{\mathbb{R}}^5$ consisting of the points of Euclidean height at least $1/\tau$. By [PP1, Lem. 6.7], if c is the $(2, 1)$ -entry of a matrix $g \in \mathrm{SL}_2(\mathbb{H})$ such that \mathcal{H}_∞ and $g\mathcal{H}_\infty$ are disjoint, then the hyperbolic distance between \mathcal{H}_∞ and $g\mathcal{H}_\infty$ is $|\ln(\tau^{-2} \mathbf{N}(c))|$.

Let $\rho = xy^{-1} \in A \cup \{\infty\}$. If $y = 0$, let $\gamma_\rho = \mathrm{id}$, otherwise let

$$\gamma_\rho = \begin{pmatrix} \rho & -1 \\ 1 & 0 \end{pmatrix} \in \mathrm{SL}_2(\mathbb{H}).$$

Let $\mathcal{H}_\rho = \gamma_\rho \mathcal{H}_\infty$, which is a horoball centered at ρ . For every $g \in \Gamma_\theta$ such that \mathcal{H}_∞ and $g\mathcal{H}_\rho$ are disjoint, let δ_g be the common perpendicular from \mathcal{H}_∞ to $g\mathcal{H}_\rho = g\gamma_\rho \mathcal{H}_\infty$, with length $\ell(\delta_g) > 0$. Hence if \mathcal{H}_∞ and $g\mathcal{H}_\rho$ are disjoint, since $(u, v) = g(x, y)$ if and only if $(uy^{-1}, vy^{-1}) = g\gamma_\rho(1, 0)$ when $y \neq 0$, we have $\ell(\delta_g) = |\ln \frac{\mathbf{N}(v)}{\tau^2 \mathbf{N}(y)}|$ if $y \neq 0$ and $\ell(\delta_g) = |\ln(\tau^{-2} \mathbf{N}(v))|$ otherwise. By discreteness, there are only finitely many double classes $[g] \in \Gamma_{1,0} \backslash \Gamma_\theta / \Gamma_{x,y}$ such that \mathcal{H}_∞ and $g\mathcal{H}_\rho$ are not disjoint or such that $\mathbf{N}(v) \leq \mathbf{N}(y)$ or such that the multiplicity of δ_g is different from 1. Assume that $y \neq 0$ (the case $y = 0$ is treated similarly). Then, as $s \rightarrow +\infty$,

$$\begin{aligned} \psi_{\mathbf{m}, \mathbf{m}'}(s) &= |\mathcal{O}^\times| \mathrm{Card}\{[g] \in \Gamma_{1,0} \backslash \Gamma_\theta / \Gamma_{x,y} : \ell(\delta_g) \leq \ln \frac{\mathbf{N}(\mathbf{m})s}{\tau^2 \mathbf{N}(y)}\} + O(1). \\ &= 2|\mathcal{O}^\times| \mathrm{Card}\{[g] \in \overline{\Gamma_{1,0}} \backslash \overline{\Gamma_\theta} / \overline{\Gamma_{x,y}} : \ell(\delta_g) \leq \ln \frac{\mathbf{N}(\mathbf{m})s}{\tau^2 \mathbf{N}(y)}\} + O(1). \end{aligned}$$

Let $\overline{\Gamma_{\mathcal{H}_\rho}}$ be the stabiliser in $\overline{\Gamma_\theta}$ of the horoball \mathcal{H}_ρ . By [PP3, Lem. 7], the number $\psi_{\mathbf{m}, \mathbf{m}'}(s)$ is equal to

$$2|\mathcal{O}^\times| [\overline{\Gamma_{\mathcal{H}_\infty}} : \overline{\Gamma_{1,0}}] [\overline{\Gamma_{\mathcal{H}_\rho}} : \overline{\Gamma_{x,y}}] \mathrm{Card}\{[g] \in \overline{\Gamma_{\mathcal{H}_\infty}} \backslash \overline{\Gamma_\theta} / \overline{\Gamma_{\mathcal{H}_\rho}} : \ell(\delta_g) \leq \ln \frac{\mathbf{N}(\mathbf{m})s}{\tau^2 \mathbf{N}(y)}\} + O(1).$$

Note that $[\overline{\Gamma_{\mathcal{H}_\infty}} : \overline{\Gamma_{1,0}}] = \frac{|\mathcal{O}^\times|}{2}$. If τ is small enough, then \mathcal{H}_∞ and \mathcal{H}_ρ are precisely invariant under $\overline{\Gamma_\theta}$. Hence, using Corollary 9, applied to $n = 5$, $\Gamma = \overline{\Gamma_\theta}$, $D^- = \mathcal{H}_\infty$ and $D^+ = \mathcal{H}_\rho$, we have, since the pointwise stabilisers of \mathcal{H}_∞ and \mathcal{H}_ρ are trivial,

$$\begin{aligned} \psi_{\mathbf{m}, \mathbf{m}'}(s) &= |\mathcal{O}^\times|^2 [\overline{\Gamma_{\mathcal{H}_\rho}} : \overline{\Gamma_{x,y}}] \mathcal{N}_{D^-, D^+} \left(\ln \frac{\mathbf{N}(\mathbf{m})s}{\tau^2 \mathbf{N}(y)} \right) + O(1) \\ &= |\mathcal{O}^\times|^2 [\overline{\Gamma_{\mathcal{H}_\rho}} : \overline{\Gamma_{x,y}}] c(D^-, D^+) \left(\frac{\mathbf{N}(\mathbf{m})s}{\tau^2 \mathbf{N}(y)} \right)^4 (1 + O(e^{-\kappa t})). \end{aligned}$$

By the Remark before Lemma 15 of [PP5], we have $\mathrm{Vol}(\overline{\Gamma_{\mathcal{H}_\infty}} \backslash \mathcal{H}_\infty) = \frac{D_A \tau^4}{8|\mathcal{O}^\times|^2}$, and by Lemma 15, Equation (33) and the centred equation three lines before Remark 19 in [PP5],

$$\mathrm{Vol}(\overline{\Gamma_{\mathcal{H}_\rho}} \backslash \mathcal{H}_\rho) = \frac{D_A \mathbf{N}(y)^4 \tau^4}{16 |\mathcal{O}_r(K_{x,y})^\times| [\overline{\Gamma_{\mathcal{H}_\rho}} : \overline{\Gamma_{x,y}}] \mathbf{N}(\mathbf{m})^4}. \quad (10)$$

By Theorem 3 of [PP5] (due to Emery, see the appendix of [PP5]), we have

$$\mathrm{Vol}(M) = \frac{\zeta(3) \prod_{p|D_A} (p^3 - 1)(p - 1)}{11520}. \quad (11)$$

Recall that $\text{Vol}(\mathbb{S}^4) = \frac{8\pi^2}{3}$. Thus, $\psi_{\mathbf{m}, \mathbf{m}'}(s)$ equals

$$\begin{aligned} & \frac{|\mathcal{O}^\times|^2 [\overline{\Gamma_{\mathcal{H}_\rho}} : \overline{\Gamma_{x,y}}] 2^4 \cdot 4 \cdot 3 \cdot 11520 D_A^2 \tau^8 \mathbf{N}(y)^4 (\mathbf{N}(\mathbf{m})s)^4 (1 + \mathcal{O}(s^{-\kappa}))}{8\pi^2 \zeta(3) \prod_{p|D_A} (p^3 - 1)(p - 1) 8 |\mathcal{O}^\times|^2 16 |\mathcal{O}_r(K_{x,y})^\times| [\overline{\Gamma_{\mathcal{H}_\rho}} : \overline{\Gamma_{x,y}}] \mathbf{N}(\mathbf{m})^4 \mathbf{N}(y)^4 \tau^8} \\ &= \frac{2160 D_A^2}{\pi^2 \zeta(3) |\mathcal{O}_r(K_{x,y})^\times| \prod_{p|D_A} (p^3 - 1)(p - 1)} s^4 (1 + \mathcal{O}(s^{-\kappa})). \end{aligned} \quad (12)$$

By page 134 of [Deu] (see Equation (8) of [PP5]), we get

$$\sum_{[\mathbf{m}'] \in \mathcal{O}^\mathcal{J}} \frac{1}{|\mathcal{O}_r(\mathbf{m}')^\times|} = \frac{1}{24} \prod_{p|D_A} (p - 1). \quad (13)$$

Thus, Equation (8) and the above computations give the first claim of Theorem 11.

To prove the second claim (when $y \neq 0$, the case $y = 0$ being similar), Corollary 9 implies that

$$\frac{\pi^2 \zeta(3) \prod_{p|D_A} (p^3 - 1)(p - 1) |\mathcal{O}_r(K_{x,y})^\times| [\overline{\Gamma_{\mathcal{H}_\rho}} : \overline{\Gamma_{x,y}}] \mathbf{N}(\mathbf{m})^4}{4320 D_A \mathbf{N}(y)^4 \tau^4} e^{-4t} \sum_{z \in \partial \mathcal{H}_\infty} m_t(z) \Delta_z$$

weak-star converges to $\text{Vol}_{\partial \mathcal{H}_\infty}$ as $t \rightarrow +\infty$, with the appropriate error term. Using the change of variable $t = \ln \frac{\mathbf{N}(\mathbf{m})s}{\tau^2 \mathbf{N}(y)}$, the fact that the geodesic lines containing the common perpendicular from ∞ to an image of \mathcal{H}_ρ by an element of $\overline{\Gamma_\rho}$ are exactly the geodesic lines from ∞ to an element of $\overline{\Gamma_\rho} \cdot xy^{-1}$, and the above facts on disjointness, lengths and multiplicities, this gives, since the map $(u, v) \mapsto uv^{-1}$ is $|\mathcal{O}^\times|$ -to-1,

$$\frac{\pi^2 \zeta(3) \prod_{p|D_A} (p^3 - 1)(p - 1) |\mathcal{O}_r(K_{x,y})^\times| [\overline{\Gamma_{\mathcal{H}_\rho}} : \overline{\Gamma_{x,y}}]}{4320 D_A |\mathcal{O}^\times| s^4 \tau^{-4}} \sum_{\substack{(u,v) \in \mathbf{m} \times \mathbf{m}, \mathbf{N}(\mathbf{m})^{-1} \mathbf{N}(v) \leq s \\ I_{u,v} = \mathbf{m}, [K_{u,v}] = [\mathbf{m}']}} \Delta_{(uv^{-1}, \frac{1}{\tau})}$$

weak-star converges to $\text{Vol}_{\partial \mathcal{H}_\infty}$ as $s \rightarrow +\infty$. Since $[\overline{\Gamma_{\mathcal{H}_\rho}} : \overline{\Gamma_{x,y}}] = \frac{|\mathcal{O}^\times|}{2}$ and the pushforward of $\text{Vol}_{\partial \mathcal{H}_\infty}$ by the endpoint map $(x, \frac{1}{\tau}) \mapsto x$ is $\tau^4 \text{Leb}_{\mathbb{H}}$, we have

$$\frac{\pi^2 \zeta(3) \prod_{p|D_A} (p^3 - 1)(p - 1) |\mathcal{O}_r(K_{x,y})^\times|}{8640 D_A s^4} \sum_{\substack{(u,v) \in \mathbf{m} \times \mathbf{m}, \mathbf{N}(\mathbf{m})^{-1} \mathbf{N}(v) \leq s \\ I_{u,v} = \mathbf{m}, [K_{u,v}] = [\mathbf{m}']}} \Delta_{uv^{-1}} \stackrel{*}{\rightarrow} \text{Leb}_{\mathbb{H}}, \quad (14)$$

The second claim of Theorem 11 now follows by dividing both sides by $|\mathcal{O}_r(K_{x,y})^\times|$ and summing over $[\mathbf{m}']$ as above. \square

4 Counting and equidistribution of quadratic irrationals and crossratios

Let K be either \mathbb{Q} or an imaginary quadratic number field, and, respectively, $\widehat{K} = \mathbb{R}$ or $\widehat{K} = \mathbb{C}$. Let \mathcal{O}_K be the ring of integers, D_K the discriminant, and ζ_K the zeta function of K . In this section, we denote by \cdot the action by homographies of the group $\Gamma_K = \text{PSL}_2(\mathcal{O}_K)$

on $\mathbb{P}^1(\widehat{K}) = \widehat{K} \cup \{\infty\}$. For every finite index subgroup G of Γ_K , and every $x \in \widehat{K} \cup \{\infty\}$, let G_x be the stabiliser of x in G , with $\Gamma_x = (\Gamma_K)_x$ to simplify the notation.

For every $\alpha \in \widehat{K}$ which is quadratic irrational over K ,

- denote by α^σ its Galois conjugate over K , by $\text{tr } \alpha = \alpha + \alpha^\sigma$ and $\mathbf{n}(\alpha) = \alpha\alpha^\sigma$ its relative trace and relative norm, and by

$$Q_\alpha(X) = X^2 - \text{tr } \alpha X + \mathbf{n}(\alpha)$$

the standard monic quadratic polynomial with roots α and α^σ ;

- denote by $h(\alpha) = \frac{2}{|\alpha - \alpha^\sigma|}$ the natural complexity of α in an orbit of Γ_K , modulo translations by \mathcal{O}_K , introduced in [PP2] and motivated in [PP4, §4.1] (when α is integral, $\frac{4}{h(\alpha)^2} = \text{Disc } Q_\alpha$ is the discriminant of the minimal polynomial of α);

- if $K = \mathbb{Q}$, let q_α be the least common multiple of the denominators of the rationals $\text{tr } \alpha$ and $\mathbf{n}(\alpha)$, let $D_\alpha = q_\alpha^2((\text{tr } \alpha)^2 - 4\mathbf{n}(\alpha))$ (when α is integral, we have $q_\alpha = 1$ and D_α is the discriminant of the order $\mathbb{Z} + \alpha\mathbb{Z}$), let (t_α, u_α) be the fundamental solution of the Pell-Fermat equation $t^2 - D_\alpha u^2 = 4$, and let $R_\alpha = \text{argcosh } \frac{t_\alpha}{2}$ (which is the regulator of $\mathbb{Z} + \alpha\mathbb{Z}$ when α is integral);

- for every finite index subgroup G of Γ_K , we define the *G-reciprocity index* $\iota_G(\alpha)$ of α as follows: we set $\iota_G(\alpha) = 2$ if α is *G-reciprocal*, that is, if some element of G maps α to α^σ , see [Sar] and [PP4, Prop. 4.3] for characterisations (when $K = \mathbb{Q}$, $G = \Gamma_K$ and α is integral, this is equivalent to saying that the order $\mathbb{Z} + \alpha\mathbb{Z}$ contains a unit of norm -1), and we set $\iota_G(\alpha) = 1$ otherwise.

To conclude with a geometric remark, for every α as above, there exists a unique primitive loxodromic element $\widehat{\alpha} \in \Gamma_K$ such that the repelling fixed point $\widehat{\alpha}^-$ of $\widehat{\alpha}$ is equal to α (see for instance [PP2, Lem. 6.2]). Its attractive fixed point $\widehat{\alpha}^+$ is then α^σ . With the notation at the end of Section 2, we have $\iota_G(\alpha) = \iota_G(\widehat{\alpha})$ for every finite index subgroup G of Γ_K . Finally, for all $\gamma \in \Gamma_K$, we have $\widehat{\gamma\alpha} = \gamma\widehat{\alpha}\gamma^{-1}$.

4.1 Equidistribution and error terms in counting functions of quadratic irrationals

We give in this subsection an error term to the counting asymptotics of [PP4] of the number of quadratic irrationals $\alpha \in \widehat{K}$ over K with complexity $h(\alpha)$ at most s , in an orbit of a finite index subgroup of Γ_K , and we prove an equidistribution result for the set of traces $\text{tr } \alpha$ of these elements, as $s \rightarrow +\infty$.

Theorem 12 *Let α_0 be a real quadratic irrational over \mathbb{Q} and let G be a finite index subgroup of $\Gamma_{\mathbb{Q}}$. There exists $\kappa > 0$ such that, as $s \rightarrow +\infty$,*

$$\begin{aligned} \text{Card}\{\alpha \in G \cdot \alpha_0 + [\Gamma_\infty : G_\infty]\mathbb{Z} : h(\alpha) \leq s\} \\ = \frac{6 [\Gamma_\infty : G_\infty] [\Gamma_{\alpha_0} : G_{\alpha_0}] R_{\alpha_0}}{\pi^2 [\Gamma_{\mathbb{Q}} : G]} s + O(s^{1-\kappa}). \end{aligned}$$

Furthermore, for the weak-star convergence of measures, we have

$$\lim_{s \rightarrow +\infty} \frac{\pi^2 [\Gamma_{\mathbb{Q}} : G]}{3 [\Gamma_{\alpha_0} : G_{\alpha_0}] R_{\alpha_0} s} \sum_{\alpha \in G \cdot \alpha_0 : h(\alpha) \leq s} \Delta_{\text{tr } \alpha} = \text{Leb}_{\mathbb{R}}.$$

Note that the indices (except maybe $[\Gamma_{\alpha_0} : G_{\alpha_0}]$) appearing in the above statement are for instance known when G is the principal congruence subgroup modulo a prime p , or the Hecke congruence subgroup modulo p . When $G = \Gamma_{\mathbb{Q}}$, the first claim is known (with a better error term, see for instance [Coh2, page 164]), but it is new in particular when G is not a congruence subgroup of $\Gamma_{\mathbb{Q}}$. For smooth functions ψ with compact support on \mathbb{R} , there is an error term in the equidistribution claim, of the form $O(s^{1-\kappa}\|\psi\|_{\ell})$ where $\kappa > 0$ and $\|\psi\|_{\ell}$ is the Sobolev norm of ψ for some $\ell \in \mathbb{N}$, see the comment following Theorem 8. Theorem 3 in the Introduction follows from this result.

Proof. The proof of the counting result with an error term is similar to that of [PP4, Théo. 4.4] upon replacing [PP4, Coro. 3.10] by the above Corollary 9 (the factor $\frac{2}{\iota_G(\alpha_0)}$ which appears in [PP4, Théo. 4.4] comes from the fact that we are counting $\alpha \in G \cdot \{\alpha_0, \alpha_0^{\sigma}\}$ therein). Noting that $\text{tr}(\alpha + m) = \text{tr} \alpha + 2m$ for all $m \in \mathbb{Z}$, it also follows by applying the equidistribution result with error term on the quotient of \mathbb{R} by $2[\Gamma_{\infty} : G_{\infty}]\mathbb{Z}$ to the constant function 1.

To prove the second claim, we apply Equation (3) with $n = 2$, $\Gamma = G$, D^- the horoball centred at ∞ consisting of the points with vertical coordinates at least 1 in the upper halfplane model of $\mathbb{H}_{\mathbb{R}}^2$, and D^+ the geodesic line with points at infinity α_0 and α_0^{σ} , whose stabiliser in G we denote by G_{D^+} . The image of D^+ by an element of G is the geodesic line $]\alpha, \alpha^{\sigma}[$ with points at infinity α, α^{σ} for some $\alpha \in G \cdot \alpha_0$. Such an α is uniquely determined if α_0 is not G -reciprocal, otherwise there are exactly two choices, α and α^{σ} . The origin of the common perpendicular from D^- to $]\alpha, \alpha^{\sigma}[$ (which exists except for finitely many G_{∞} -orbits of $\alpha \in G \cdot \alpha_0$) is the point $x_{\alpha} = (\frac{\text{tr} \alpha}{2}, 1) \in \partial D^-$, its length is $\ln h(\alpha)$, and its multiplicity is 1, since $\text{PSL}_2(\mathbb{Z})$ acts freely on $T^1\mathbb{H}_{\mathbb{R}}^2$. The induced Riemannian measure on ∂D^- is the Lebesgue measure. Therefore, by Equation (3) and the value of the constant $c(D^-, D^+)$ given above Corollary 9, we have

$$\lim_{s \rightarrow +\infty} \frac{\text{Vol}(\mathbb{S}^1) \text{Vol}(G \setminus \mathbb{H}_{\mathbb{R}}^2)}{\iota_G(\alpha_0) \text{Vol}(\mathbb{S}^0) \text{Vol}(G_{D^+} \setminus D^+) s} \sum_{\alpha \in G \cdot \alpha_0 : h(\alpha) \leq s} \Delta_{x_{\alpha}} = \text{Leb}_{\partial D^-} . \quad (15)$$

We have

$$\text{Vol}(G \setminus \mathbb{H}_{\mathbb{R}}^2) = [\Gamma_{\mathbb{Q}} : G] \text{Vol}(\Gamma_{\mathbb{Q}} \setminus \mathbb{H}_{\mathbb{R}}^2) = [\Gamma_{\mathbb{Q}} : G] \frac{\pi}{3} . \quad (16)$$

Recall that $\widehat{\alpha}_0 \in \Gamma_{\mathbb{Q}}$ is a primitive loxodromic element with fixed points α_0 and α_0^{σ} . By for instance [Bea, p. 173], the translation length $\ell(\widehat{\alpha}_0)$ of γ_0 satisfies $\cosh \frac{\ell(\widehat{\alpha}_0)}{2} = \frac{|\text{tr} \widehat{\alpha}_0|}{2}$. By for instance the proof of [PP4, Prop. 4.1], we have $|\text{tr} \widehat{\alpha}_0| = t_{\alpha_0}$. Hence

$$\ell(\widehat{\alpha}_0) = 2 R_{\alpha_0} , \quad (17)$$

so that

$$\text{Vol}(G_{D^+} \setminus D^+) = \frac{[\Gamma_{\alpha_0} : G_{\alpha_0}]}{\iota_G(\alpha_0)} \text{Vol}(\Gamma_{\alpha_0} \setminus D^+) = \frac{[\Gamma_{\alpha_0} : G_{\alpha_0}] \ell(\widehat{\alpha}_0)}{\iota_G(\alpha_0)} = \frac{2 [\Gamma_{\alpha_0} : G_{\alpha_0}] R_{\alpha_0}}{\iota_G(\alpha_0)} , \quad (18)$$

The result now follows from Equations (15), (16) and (18) by applying the pushforwards of measures by the map $f : (x, 1) \mapsto 2x$ from ∂D^- to \mathbb{R}^{n-1} (which sends the Lebesgue measure to $\frac{1}{2^{n-1}}$ times the Lebesgue measure). \square

Theorem 13 *Let K be an imaginary quadratic number field, let $\alpha_0 \in \mathbb{C}$ be a quadratic irrational over K , let G be a finite index subgroup of Γ_K , and let Λ be the lattice of $\lambda \in \mathcal{O}_K$ such that $\pm \begin{pmatrix} 1 & \lambda \\ 0 & 1 \end{pmatrix} \in G$. There exists $\kappa > 0$ such that, as $s \rightarrow +\infty$,*

$$\text{Card}\{\alpha \in G \cdot \alpha_0 + \Lambda : h(\alpha) \leq s\} = \frac{\pi^2 [\mathcal{O}_K : \Lambda] [\Gamma_{\alpha_0} : G_{\alpha_0}] \left| \ln \left| \frac{\text{tr } \widehat{\alpha_0} + \sqrt{(\text{tr } \widehat{\alpha_0})^2 - 4}}{2} \right| \right|}{m_{\Gamma_K}(\alpha_0) [\Gamma_K : G] |D_K| \zeta_K(2)} s^2 + O(s^{2-\kappa}),$$

where $\widehat{\alpha_0}$ is a primitive element of Γ_K fixing α_0 with absolute values of eigenvalues different from 1, and $m_G(\alpha_0)$ the number of elements in G fixing α_0 with absolute values of eigenvalues equal to 1. Furthermore, for the weak-star convergence of measures, we have

$$\lim_{s \rightarrow +\infty} \frac{2 |D_K|^{\frac{3}{2}} \zeta_K(2) [\Gamma_K : G] m_{\Gamma_K}(\alpha_0)}{\pi^2 [\Gamma_{\alpha_0} : G_{\alpha_0}] \left| \ln \left| \frac{\text{tr } \widehat{\alpha_0} + \sqrt{(\text{tr } \widehat{\alpha_0})^2 - 4}}{2} \right| \right|} s^2 \sum_{\alpha \in G \cdot \alpha_0 : h(\alpha) \leq s} \Delta_{\text{tr } \alpha} = \text{Leb}_{\mathbb{C}}.$$

For smooth functions ψ with compact support on \mathbb{C} , there is an error term in the above equidistribution claim, of the form $O(s^{2-\kappa} \|\psi\|_{\ell})$ where $\kappa > 0$ and $\|\psi\|_{\ell}$ is the Sobolev norm of ψ for some $\ell \in \mathbb{N}$.

Proof. The proof of the counting claim with an error term is similar to that of [PP4, Théo. 4.6] upon replacing [PP4, Coro. 3.10] by the above Corollary 9, and using the simplification

$$[\Gamma_{\infty} : \Lambda] = [\Gamma_{\infty} : \mathcal{O}_K] [\mathcal{O}_K : \Lambda] = \frac{|\mathcal{O}_K^{\times}|}{2} [\mathcal{O}_K : \Lambda].$$

To prove the second claim, we apply Equation (3) with $n = 3$, $\Gamma = G$, D^- the horoball centred at ∞ consisting of the points with vertical coordinates at least 1 in the upper halfspace model of $\mathbb{H}_{\mathbb{R}}^3$, and D^+ the geodesic line with points at infinity α_0 and α_0^{σ} , whose stabiliser in G we denote by G_{D^+} . The cardinality of the pointwise stabiliser in G of D^+ is $m_G(\alpha_0)$ as defined in the statement of Theorem 13 (we have $m_G(\alpha_0) = m_G(\widehat{\alpha_0})$ with the notation at the end of Section 2).

The set of points in ∂D^- fixed by a nontrivial element of the stabiliser of ∞ in Γ_K is discrete since, for instance, the subgroup of translations in a discrete group of isometries of \mathbb{R}^{n-1} has finite index by Bieberbach's theorem. Hence the multiplicity of the common perpendicular from D^- to an image of D^+ by an element of G is different from 1 only when its origin belongs to a discrete subset S of ∂D^- . Furthermore, since $\Gamma_{D^+} \backslash D^+$ is compact, there exists $\epsilon > 0$ (depending only on Γ_K and α_0) such that a geodesic arc, leaving perpendicularly from an image of D^+ and arriving perpendicularly at another image of D^+ , has length at least ϵ . Hence the number of common perpendiculars from a given point in S to an image of D^+ grows at most linearly in the length. Therefore these common perpendiculars do not contribute asymptotically to the equidistributing sum, and as in the previous proof, we have

$$\lim_{s \rightarrow +\infty} \frac{2 m_G(\alpha_0) \text{Vol}(\mathbb{S}^2) \text{Vol}(G \backslash \mathbb{H}_{\mathbb{R}}^3)}{\iota_G(\alpha_0) \text{Vol}(\mathbb{S}^1) \text{Vol}(G_{D^+} \backslash D^+) s^2} \sum_{\alpha \in G \cdot \alpha_0 : h(\alpha) \leq s} \Delta_{(\frac{\text{tr } \alpha}{2}, 1)} = \text{Leb}_{\partial D^-}. \quad (19)$$

Recall that the translation length $\ell(\gamma_0)$ of an element γ_0 of $\text{PSL}_2(\mathbb{C})$ is

$$\ell(\gamma_0) = 2 \left| \ln \left| \frac{\text{tr } \gamma_0 + \sqrt{(\text{tr } \gamma_0)^2 - 4}}{2} \right| \right| \quad (20)$$

(which is independent of the choice of the square root of the complex number $\text{tr } \gamma_0^2 - 4$ and of the choice of the lift of γ_0 in $\text{SL}_2(\mathbb{C})$ modulo $\{\pm \text{id}\}$). With r the smallest positive integer such that $\widehat{\alpha_0}^r \in G$, we have $[\Gamma_{\alpha_0} : G_{\alpha_0}] = \frac{m_{\Gamma_K}(\alpha_0)}{m_G(\alpha_0)} r$, so that

$$\text{Vol}(G_{D^+} \backslash D^+) = \frac{r}{\iota_G(\alpha_0)} \ell(\widehat{\alpha_0}) = \frac{m_G(\alpha_0) [\Gamma_{\alpha_0} : G_{\alpha_0}]}{m_{\Gamma_K}(\alpha_0) \iota_G(\alpha_0)} \ell(\widehat{\alpha_0}). \quad (21)$$

As in the previous proof, the result now follows from Equations (19), (20) and (7) by applying the pushforwards of measures by the map $(x, 1) \mapsto 2x$ from ∂D^- to \mathbb{C} . \square

Corollary 14 *Let $\phi = \frac{1+\sqrt{5}}{2}$ be the Golden Ratio, let K be an imaginary quadratic number field with $D_K \neq -4$, let \mathfrak{c} be a nonzero ideal in \mathcal{O}_K , and let $\Gamma_0(\mathfrak{c})$ be the Hecke congruence subgroup $\left\{ \pm \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_K : c \in \mathfrak{c} \right\}$. Then there exists $\kappa > 0$ such that, as $s \rightarrow +\infty$,*

$$\text{Card}\{\alpha \in \Gamma_0(\mathfrak{c}) \cdot \phi + \mathcal{O}_K : h(\alpha) \leq s\} = \frac{2\pi^2 k_{\mathfrak{c}} \ln \phi}{|D_K| \zeta_K(2) \mathfrak{n}(\mathfrak{c}) \prod_{\mathfrak{p}|\mathfrak{c}} (1 + \frac{1}{\mathfrak{n}(\mathfrak{p})})} s^2 + O(s^{2-\kappa}),$$

where $k_{\mathfrak{c}}$ is the smallest $k \in \mathbb{N} - \{0\}$ such that the $2k$ -th term of the standard Fibonacci sequence belongs to \mathfrak{c} , and the product is over the prime ideals in \mathcal{O}_K dividing \mathfrak{c} .

Proof. The proof of this corollary, as well as the one of Corollary 4 in the introduction, are similar to the one of [PP4, Coro. 4.7], using the fact proven therein that $m_{\Gamma_0(\mathfrak{c})}(\phi) = 1$, so that (the element $\widehat{\phi}^{-1}$ is denoted by $\gamma_1 = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}$ in loc. cit., and its translation length is $2 \ln \phi$)

$$\frac{[\Gamma_{\phi} : (\Gamma_0(\mathfrak{c}))_{\phi}]}{m_{\Gamma_K}(\phi)} = \frac{[\widehat{\phi}^{\mathbb{Z}} : \widehat{\phi}^{\mathbb{Z}} \cap \Gamma_0(\mathfrak{c})]}{m_{\Gamma_0(\mathfrak{c})}(\phi)} = k_{\mathfrak{c}}. \quad \square$$

Let \mathbb{H} be Hamilton's quaternion algebra over \mathbb{R} , let A be a definite quaternion algebra over \mathbb{Q} , and let \mathcal{O} be a maximal order in A (we refer to Subsection 3.2 for background, as well as for the notation \neg , \mathbb{N} , Tr , D_A , $\text{PSL}_2(\mathbb{H})$, $\text{PSL}_2(\mathcal{O})$, \cdot). For every $x \in \mathbb{H} \cup \{\infty\}$, we denote by G_x the stabiliser of x for the action by homographies of a subgroup G of $\text{PSL}_2(\mathbb{H})$. For every $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{H})$, we denote by X_{γ} the largest real root of

$$2x^3 - c_1 x^2 + 2(c_2 c_3 - 1)x + (c_1 - c_2^2 - c_3^2),$$

where $c_1 = \mathbb{N}(a + d) + \text{Tr}(ad - bc)$, $c_2 = \frac{1}{2} \text{Tr}(a + d)$, $c_3 = \frac{1}{2} \text{Tr}((ad - bc)\bar{a} + (da - cb)\bar{d})$. We will say that an element $\alpha \in \mathbb{H}$ is a *loxodromic quadratic irrational* over \mathcal{O} if there exists $\gamma \in \text{SL}_2(\mathcal{O})$ such that $|X_{\gamma}| \neq 1$ fixing α , in which case we denote by α^{σ} its other fixed point (which exists, is unique, and is independent of such a γ , see below for proofs and a justification of the terminology). Note that by the noncommutativity of \mathbb{H} , there are several types of quadratic equations over \mathcal{O} .

The following result is a counting and equidistribution result in \mathbb{H} of loxodromic quadratic irrationals over \mathcal{O} in a given homographic orbit under $\text{PSL}_2(\mathcal{O})$.

Theorem 15 *Let \mathcal{O} be a maximal order in a definite quaternion algebra over \mathbb{Q} , let $\alpha_0 \in \mathbb{H}$ be a loxodromic quadratic irrational over \mathcal{O} , let G be a finite index subgroup of $\mathrm{PSL}_2(\mathcal{O})$, and let Λ be the lattice of $\lambda \in \mathcal{O}$ such that $\pm \begin{pmatrix} 1 & \lambda \\ 0 & 1 \end{pmatrix} \in G$. There exists $\kappa > 0$ such that, as $\epsilon \rightarrow 0$,*

$$\mathrm{Card}\{\alpha \in G \cdot \alpha_0 + \Lambda : \mathrm{N}(\alpha - \alpha^\sigma) \geq \epsilon\} = \frac{8640 D_A [\mathcal{O} : \Lambda] [\mathrm{PSL}_2(\mathcal{O})_{\alpha_0} : G_{\alpha_0}] |\ln |X_{\gamma_0} + (X_{\gamma_0}^2 - 1)^{1/2}| |}{\zeta(3) m_{\mathrm{PSL}_2(\mathcal{O})}(\alpha_0) [\mathrm{PSL}_2(\mathcal{O}) : G] \prod_{p|D_A} (p^3 - 1)(p - 1)} \epsilon^{-2} + O(\epsilon^{\kappa-2}),$$

where γ_0 is a primitive element of $\mathrm{SL}_2(\mathcal{O})$ fixing α_0 with $|X_{\gamma_0}| \neq 1$, $m_G(\alpha_0)$ is the number of elements γ in G fixing α_0 with $|X_\gamma| = 1$, and the product is over the primes p dividing D_A . Furthermore, the Lebesgue measure $\mathrm{Leb}_{\mathbb{H}}$ on \mathbb{H} is the weak-star limit, as $\epsilon \rightarrow 0$, of the measures

$$\frac{\zeta(3) m_{\Gamma_K}(\alpha_0) [\mathrm{PSL}_2(\mathcal{O}) : G] \prod_{p|D_A} (p^3 - 1)(p - 1) \epsilon^2}{2160 [\mathrm{PSL}_2(\mathcal{O})_{\alpha_0} : G_{\alpha_0}] |\ln |X_{\gamma_0} + (X_{\gamma_0}^2 - 1)^{1/2}| |} \sum_{\alpha \in G \cdot \alpha_0 : \mathrm{N}(\alpha - \alpha^\sigma) \geq \epsilon} \Delta_{\alpha + \alpha^\sigma}.$$

For smooth functions ψ with compact support on \mathbb{H} , there is an error term in the equidistribution claim, of the form $O(\epsilon^{\kappa-2} \|\psi\|_\ell)$ where $\kappa > 0$ and $\|\psi\|_\ell$ is the Sobolev norm of ψ for some $\ell \in \mathbb{N}$.

Proof. First recall (see [PaS, §3]) that for every $\gamma \in \mathrm{SL}_2(\mathbb{H})$, we have $|X_\gamma| \geq 1$, and that γ is loxodromic for its isometric action on the upper halfspace model of $\mathbb{H}_{\mathbb{R}}^5$ by Poincaré's extension (see for instance Equation (14) in [PP5]) if and only if $|X_\gamma| \neq 1$. If this holds, then (see [PaS, §3]) γ is conjugated to $\begin{pmatrix} t_+ & 0 \\ 0 & t_- \end{pmatrix}$ with $\mathrm{N}(t_\pm) = X_\gamma \pm (X_\gamma^2 - 1)^{1/2}$. Since this diagonal matrix acts on the vertical axis $\{0\} \times]0, +\infty[$ in $\mathbb{H}_{\mathbb{R}}^5$ by $(0, r) \mapsto (0, \mathrm{N}(t_+)r)$, the translation length $\ell(\gamma)$ of γ is therefore

$$\ell(\gamma) = |\ln |X_\gamma + (X_\gamma^2 - 1)^{1/2}||. \quad (22)$$

As a side remark, if $\gamma \in \mathrm{SL}_2(\mathcal{O})$ satisfies $|X_\gamma| > 1$, in particular, it has exactly two fixed points, which are the only two solutions of some quadratic equation $ax + b = xcx + xd$ where $a, b, c, d \in \mathcal{O}$ and $\mathrm{N}(ad) + \mathrm{N}(bc) - \mathrm{Tr}(a\bar{c}d\bar{b}) = 1$.

To prove the equidistribution claim of Theorem 15, we apply Equation (3) with $n = 5$, $\Gamma = G$, D^- the horoball centred at ∞ consisting of the points with vertical coordinate at least 1 in $\mathbb{H}_{\mathbb{R}}^5$, and D^+ the geodesic line with points at infinity α_0 and α_0^σ , whose stabiliser in G we denote by G_{D^+} . We define $\iota_G(\alpha_0) = 2$ if there exists an element in G sending α_0 to α_0^σ and $\iota_G(\alpha_0) = 1$ otherwise. Except for finitely many G_∞ -orbits of $\alpha \in G \cdot \alpha_0$, the common perpendicular from D^- to the geodesic line with points at infinity α, α^σ exists and has hyperbolic length $\ln \frac{2}{\mathrm{N}(\alpha - \alpha^\sigma)^{1/2}}$, and its origin is $(\frac{\alpha + \alpha^\sigma}{2}, 1) \in \partial D^-$. As in the proof of Theorem 13, we have

$$\lim_{t \rightarrow +\infty} \frac{4 m_G(\alpha_0) \mathrm{Vol}(\mathbb{S}^4) \mathrm{Vol}(G \backslash \mathbb{H}_{\mathbb{R}}^5)}{\iota_G(\alpha_0) \mathrm{Vol}(\mathbb{S}^3) \mathrm{Vol}(G_{D^+} \backslash D^+) e^{4t}} \sum_{\alpha \in G \cdot \alpha_0 : \mathrm{N}(\alpha - \alpha^\sigma) \geq 4 e^{-2t}} \Delta_{(\frac{\alpha + \alpha^\sigma}{2}, 1)} = \mathrm{Leb}_{\partial D^-}. \quad (23)$$

We have $\text{Vol}(\mathbb{S}^4) = \frac{8\pi^2}{3}$, $\text{Vol}(\mathbb{S}^3) = 2\pi^2$ and, as in the proof of Theorem 13,

$$\text{Vol}(G_{D^+} \backslash D^+) = \frac{[\text{PSL}_2(\mathcal{O})_{\alpha_0} : G_{\alpha_0}] m_G(\alpha_0)}{\iota_G(\alpha_0) m_{\text{PSL}_2(\mathcal{O})}(\alpha_0)} \ell(\gamma_0).$$

Taking $\epsilon = 4e^{-2t}$, the result follows, as in the proof of Theorem 13, from Equations (23), (22) and (11) by applying the pushforwards of measures by the map $(x, 1) \mapsto 2x$ from ∂D^- to \mathbb{H} .

To prove the first claim of Theorem 15, note that by [KO, Prop. 5.5] for instance, we have

$$\text{Vol}((2\Lambda) \backslash \mathbb{H}) = 2^4 [\mathcal{O} : \Lambda] \text{Vol}(\mathcal{O} \backslash \mathbb{H}) = 2^4 [\mathcal{O} : \Lambda] \frac{D_A}{4}.$$

The result then follows by considering the measures induced on the compact quotient $(2\Lambda) \backslash \mathbb{H}$, and by applying the equidistribution result with the error term to the constant function 1. \square

4.2 Relative complexity of loxodromic elements

This subsection is a geometric one, paving the way to the next one, which is arithmetic. It explains in particular why the crossratios play an important geometric role in this paper.

Let $n \geq 2$ and Γ be as in the beginning of Section 2, and let $\gamma_0 \in \Gamma$ be a fixed primitive loxodromic element.

We define the *relative height* of a primitive loxodromic element $\gamma \in \Gamma$ with respect to γ_0 to be the length of the common perpendicular of the translation axes of γ_0 and γ if they are disjoint and 0 otherwise. Note that

$$h_{\gamma_0^\epsilon}(\gamma^{\epsilon'}) = h_{\gamma_0}(\gamma) = h_{g\gamma_0g^{-1}}(g\gamma g^{-1}) = h_{\gamma_0}(g'\gamma g'^{-1})$$

for all $\epsilon, \epsilon' \in \{\pm 1\}$, $g \in \Gamma$ and $g' \in \text{Stab}_\Gamma(\text{Axis } \gamma_0)$. Furthermore, it follows from the properness of the action of Γ as in [PP2, Lemma 3.1] that the set

$$\{[\gamma] \in \gamma_0^\mathbb{Z} \backslash \Gamma / \gamma_1^\mathbb{Z} : h_{\gamma_0}(\gamma\gamma_1\gamma^{-1}) \leq t\}$$

is finite for all primitive loxodromic elements $\gamma_0, \gamma_1 \in \Gamma$ and $t \geq 0$. This shows that the relative height with respect to a given primitive loxodromic element is a reasonable complexity within a given conjugacy class of primitive loxodromic elements of Γ . Theorem 8 easily implies the following counting result of primitive loxodromic elements using their relative heights.

Corollary 16 *If the Bowen-Margulis measure m_{BM} of $\Gamma \backslash \mathbb{H}_{\mathbb{R}}^n$ is finite, then for all primitive loxodromic elements $\gamma_0, \gamma_1 \in \Gamma$, as $t \rightarrow +\infty$,*

$$\sum_{[\gamma] \in \gamma_0^\mathbb{Z} \backslash \Gamma / \gamma_1^\mathbb{Z}, h_{\gamma_0}(\gamma\gamma_1\gamma^{-1}) \leq t} m_{e, \gamma} \sim C_{\gamma_0, \gamma_1} e^{\delta_\Gamma t},$$

where $C_{\gamma_0, \gamma_1} = \iota_\Gamma(\gamma_0) \iota_\Gamma(\gamma_1) m_\Gamma(\gamma_0) m_\Gamma(\gamma_1) \|\sigma_{\text{Axis}(\gamma_0)}\| \|\sigma_{\text{Axis}(\gamma_1)}\| \delta_\Gamma^{-1} \|m_{\text{BM}}\|^{-1}$, with an error term $O(e^{(n-1)t})$ if furthermore $\Gamma \backslash \mathbb{H}_{\mathbb{R}}^n$ is compact or arithmetic. If $\Gamma \backslash \mathbb{H}_{\mathbb{R}}^n$ has finite volume, then

$$C_{\gamma_0, \gamma_1} = \frac{\text{Vol}(\mathbb{S}^{n-2})^2 \ell(\gamma_0) \ell(\gamma_1)}{2^{n-1} (n-1) \text{Vol}(\mathbb{S}^{n-1}) \text{Vol}(\Gamma \backslash \mathbb{H}_{\mathbb{R}}^n)}.$$

In particular, when Γ is torsion free,

$$\text{Card}\{[\gamma] \in \gamma_0^{\mathbb{Z}} \backslash \Gamma / \gamma_1^{\mathbb{Z}} : h_{\gamma_0}(\gamma \gamma_1 \gamma^{-1}) \leq t\} \sim C_{\gamma_0, \gamma_1} e^{\delta_{\Gamma} t}$$

as $t \rightarrow +\infty$.

Proof. By [PP3, Lem. 7], except above finitely many double classes, the canonical map from $\gamma_0^{\mathbb{Z}} \backslash \Gamma / \gamma_1^{\mathbb{Z}}$ to $\text{Stab}_{\Gamma}(\text{Axis}(\gamma_0)) \backslash \Gamma / \text{Stab}_{\Gamma}(\text{Axis}(\gamma_1))$ is a k -to-1 map, where

$$k = [\text{Stab}_{\Gamma}(\text{Axis}(\gamma_0)) : \gamma_0^{\mathbb{Z}}] [\text{Stab}_{\Gamma}(\text{Axis}(\gamma_1)) : \gamma_1^{\mathbb{Z}}] = \iota_{\Gamma}(\gamma_0) \iota_{\Gamma}(\gamma_1) m_{\Gamma}(\gamma_0) m_{\Gamma}(\gamma_1)$$

(using Equation (5)). When $[\gamma]$ varies in a fiber of this map, the quantities $m_{e, \gamma}$ (defined in Section 2) and $h_{\gamma_0}(\gamma \gamma_1 \gamma^{-1})$ are constant. As seen in the proof of Theorem 13, if $\Gamma \backslash \mathbb{H}_{\mathbb{R}}^n$ has finite volume, then $\text{Vol}(\text{Stab}_{\Gamma}(\text{Axis}(\gamma_i)) \backslash \text{Axis}(\gamma_i)) = \frac{\ell(\gamma_i)}{\iota_G(\gamma_i)}$ for $i = 0, 1$. The result then immediately follows from Theorem 8 and Corollary 9. \square

In the end of this subsection, we give an asymptotic formula relating the relative height of two primitive loxodromic elements with the crossratio of its fixed points.

Recall that the *crossratio* of four pairwise distinct points a, b, c, d in $\mathbb{P}_1(\mathbb{R}) = \mathbb{R} \cup \{\infty\}$ or in $\mathbb{P}_1(\mathbb{C}) = \mathbb{C} \cup \{\infty\}$ is

$$[a, b, c, d] = \frac{(c - a)(d - b)}{(c - b)(d - a)},$$

with the standard conventions when one of the points is ∞ . Using Ahlfors's terminology, the *absolute crossratio* of four pairwise distinct points a, b, c, d in the one-point compactification $\mathbb{R}^{n-1} \cup \{\infty\}$ of the standard $n - 1$ -dimensional Euclidean space is

$$\llbracket a, b, c, d \rrbracket = \frac{\|c - a\| \|d - b\|}{\|c - b\| \|d - a\|},$$

where $\|\cdot\|$ denotes the standard Euclidean norm and with conventions analogous to the definition of crossratio when one of the points is ∞ .

We will denote by $]a_-, a_+[$ the (oriented) geodesic line in $\mathbb{H}_{\mathbb{R}}^n$ whose pair of endpoints is a given pair (a_-, a_+) of distinct points in $\mathbb{R}^{n-1} \cup \{\infty\}$. In particular, for every loxodromic element $\gamma \in \Gamma$, we have $\text{Axis}(\gamma) =]\gamma^-, \gamma^+[$.

Lemma 17 (1) *Two geodesic lines $]a_-, a_+[$ and $]b_-, b_+[$ in $\mathbb{H}_{\mathbb{R}}^2$ or $\mathbb{H}_{\mathbb{R}}^3$ are orthogonal if and only if $[a_-, a_+, b_-, b_+] = -1$.*

(2) *For all primitive loxodromic elements $\gamma_0, \gamma \in \Gamma$, as $h_{\gamma_0}(\gamma) \rightarrow +\infty$, we have*

$$e^{h_{\gamma_0}(\gamma)} = \frac{4}{\llbracket \gamma_0^-, \gamma^-, \gamma_0^+, \gamma^+ \rrbracket} + O(1).$$

Proof. (1) This is a classical fact, see for example pages 15 and 31 of [Fen].

(2) This is a corollary of [PP2, Lem. 2.2]. Note that the crossratio in [PP2] is the logarithm of the absolute crossratio in this paper. \square

4.3 Relative complexity of quadratic irrationals

Let K (and its related terminology) be as in the beginning of Section 4. Let α and β be quadratic irrationals in \widehat{K} over K . We define the *relative height* of β with respect to α as

$$h_\alpha(\beta) = \min \left\{ \frac{|\beta - \alpha| |\beta^\sigma - \alpha^\sigma|}{|\beta - \beta^\sigma|}, \frac{|\beta - \alpha^\sigma| |\beta^\sigma - \alpha|}{|\beta - \beta^\sigma|} \right\}. \quad (24)$$

Another expression of this complexity using the absolute crossratios is, if $\beta \notin \{\alpha, \alpha^\sigma\}$,

$$h_\alpha(\beta) = \frac{|\alpha - \alpha^\sigma|}{\max\{\llbracket \alpha, \beta, \alpha^\sigma, \beta^\sigma \rrbracket, \llbracket \alpha, \beta^\sigma, \alpha^\sigma, \beta \rrbracket\}}. \quad (25)$$

Here are some elementary properties of the relative heights.

Lemma 18 *Let α and β in \widehat{K} be quadratic irrationals over K . Then*

- (1) $h_{\alpha^\rho}(\beta^\tau) = h_\alpha(\beta)$ for all $\rho, \tau \in \{\text{id}, \sigma\}$.
- (2) $h_\alpha(\beta) = 0$ if and only if $\beta \in \{\alpha, \alpha^\sigma\}$.
- (3) $h_\alpha(\gamma\beta) = h_\alpha(\beta)$ for every $\gamma \in \text{Stab}_{\Gamma_K}(\{\alpha, \alpha^\sigma\})$.

Proof. We will prove (3), the first two claims being immediate consequences of the definition. We denote by $\dot{}$ the derivative. Assume first that $\gamma \in \Gamma_K$ fixes α (and consequently also α^σ). Then, using a well-known identity for linear fractional transformations (see for example [Ahl, p. 19]) and the well-known fact that $\dot{\gamma}(\alpha) \dot{\gamma}(\alpha^\sigma) = 1$, we have

$$\begin{aligned} \frac{|\gamma\beta - \alpha| |\gamma\beta^\sigma - \alpha^\sigma|}{|\gamma\beta - \gamma\beta^\sigma|} &= \frac{|\gamma\beta - \gamma\alpha| |\gamma\beta^\sigma - \gamma\alpha^\sigma|}{|\gamma\beta - \gamma\beta^\sigma|} \\ &= \sqrt{\frac{|\dot{\gamma}\beta| |\dot{\gamma}\alpha| |\dot{\gamma}\beta^\sigma| |\dot{\gamma}\alpha^\sigma|}{|\dot{\gamma}\beta| |\dot{\gamma}\beta^\sigma|}} \frac{|\beta - \alpha| |\beta^\sigma - \alpha^\sigma|}{|\beta - \beta^\sigma|} = \frac{|\beta - \alpha| |\beta^\sigma - \alpha^\sigma|}{|\beta - \beta^\sigma|}. \end{aligned}$$

The invariance by γ of the second term (seen as a function of β) in the definition of the relative height is checked by a similar computation, replacing α by α^σ . If now $\gamma \in \Gamma_K$ exchanges α and α^σ , then a similar computation gives that γ exchanges the two terms of the definition of the relative height. The result follows. \square

Lemma 19 *For all α, β in \widehat{K} which are quadratic irrationals over K and $s \geq 0$, the set*

$$\{\beta' \in \Gamma_\alpha \backslash \Gamma_K \cdot \beta : h_\alpha(\beta') \leq s\} \quad (26)$$

is finite.

Proof. Note that this set is well defined by Lemma 18 (3). By Equation (25), by Lemma 17 (2), since $h_{\widehat{\alpha}}(\widehat{\beta}) = h_{\widehat{\alpha}}(\widehat{\beta}^{-1})$ and since the fixed points of $\widehat{\alpha}$ and $\widehat{\beta}$ are α, α^σ and β, β^σ respectively, we have

$$h_\alpha(\beta) = \frac{|\alpha - \alpha^\sigma|}{4} e^{h_{\widehat{\alpha}}(\widehat{\beta})} + O(|\alpha - \alpha^\sigma|). \quad (27)$$

Since the set $\{[\gamma] \in \widehat{\alpha}^{\mathbb{Z}} \backslash \widehat{\Gamma} / \widehat{\beta}^{\mathbb{Z}} : h_{\widehat{\alpha}}(\gamma \widehat{\beta} \gamma^{-1}) \leq t\}$ is finite for all $t \geq 0$ (see Subsection 4.2), since $\widehat{\alpha}^{\mathbb{Z}}$ and $\widehat{\beta}^{\mathbb{Z}}$ have finite index in Γ_α and Γ_β respectively, and since $\gamma \widehat{\beta} \gamma^{-1} = \widehat{\gamma} \widehat{\beta}$ for all $\gamma \in \Gamma_K$, this proves the result. \square

This shows that the relative height with respect to a given quadratic irrational α_0 is, indeed, a reasonable complexity for quadratic irrationals in a given orbit of the modular group Γ_K (modulo the stabiliser of α_0).

Recall that for all quadratic irrationals α and β in \widehat{K} over K , we have given the expression of two points $x_\alpha^\pm(\beta)$ in \widehat{K} in Equation (1) of the introduction. We will see in the proof (Lemma 22) of Theorems 20 and 21 that these points are well defined when β varies in a given orbit of Γ_K , except for finitely many Γ_α -classes. We are now going to prove the equidistribution of these points, when counted according to their relative heights. We state separately the cases $K = \mathbb{Q}$ and K a quadratic imaginary number field, but it is natural to give a common proof for both statements.

Theorem 20 *Let $K = \mathbb{Q}$, let G be a subgroup of finite index in $\Gamma_K = \mathrm{PSL}_2(\mathbb{Z})$ and let α_0, β_0 be real quadratic irrationals over \mathbb{Q} . Then there exists $\kappa > 0$ such that, as $s \rightarrow +\infty$,*

$$\begin{aligned} \mathrm{Card}\{\beta \in G_{\alpha_0} \backslash G \cdot \beta_0 : h_{\alpha_0}(\beta) \leq s\} \\ = \frac{24 [\Gamma_{\alpha_0} : G_{\alpha_0}] [\Gamma_{\beta_0} : G_{\beta_0}] R_{\alpha_0} R_{\beta_0} h(\alpha_0)}{\pi^2 [\Gamma_{\mathbb{Q}} : G]} s (1 + O(s^{-\kappa})). \end{aligned}$$

Furthermore, as $s \rightarrow +\infty$, we have the following convergence of measures on $\mathbb{R} - \{\alpha_0, \alpha_0^\sigma\}$:

$$\frac{\pi^2 [\Gamma_{\mathbb{Q}} : G]}{24 [\Gamma_{\beta_0} : G_{\beta_0}] R_{\beta_0} s} \sum_{\beta \in G \cdot \beta_0, h_{\alpha_0}(\beta) \leq s} \Delta_{x_{\alpha_0}^-(\beta)} + \Delta_{x_{\alpha_0}^+(\beta)} \xrightarrow{*} \frac{d\mathrm{Leb}_{\mathbb{R}}(t)}{|Q_{\alpha_0}(t)|}. \quad (28)$$

Theorem 21 *Let K be a quadratic imaginary number field, let G be a subgroup of finite index in $\Gamma_K = \mathrm{PSL}_2(\mathcal{O}_K)$ and let $\alpha_0, \beta_0 \in \mathbb{C}$ be quadratic irrationals over K . Then there exists $\kappa > 0$ such that, as $s \rightarrow +\infty$,*

$$\begin{aligned} \mathrm{Card}\{\beta \in G_{\alpha_0} \backslash G \cdot \beta_0 : h_{\alpha_0}(\beta) \leq s\} \\ = \frac{8\pi^3 [\Gamma_{\alpha_0} : G_{\alpha_0}] [\Gamma_{\beta_0} : G_{\beta_0}] \left| \ln \left| \frac{\mathrm{tr} \widehat{\alpha_0} + \sqrt{\mathrm{tr} \widehat{\alpha_0}^2 - 4}}{2} \right| \right| \left| \ln \left| \frac{\mathrm{tr} \widehat{\beta_0} + \sqrt{\mathrm{tr} \widehat{\beta_0}^2 - 4}}{2} \right| \right|}{m_{\Gamma_K}(\alpha_0) m_{\Gamma_K}(\beta_0) h(\alpha_0)^{-2} [\Gamma_K : G] |D_K|^{\frac{3}{2}} \zeta_K(2)} s^2 (1 + O(s^{-\kappa})). \end{aligned}$$

Furthermore, as $s \rightarrow +\infty$, we have the following convergence of measures on $\mathbb{C} - \{\alpha_0, \alpha_0^\sigma\}$:

$$\frac{m_{\Gamma_K}(\beta_0) [\Gamma_K : G] |D_K|^{\frac{3}{2}} \zeta_K(2)}{16 \pi^2 [\Gamma_{\beta_0} : G_{\beta_0}] \left| \ln \left| \frac{\mathrm{tr} \widehat{\beta_0} + \sqrt{\mathrm{tr} \widehat{\beta_0}^2 - 4}}{2} \right| \right| s^2} \sum_{\beta \in G \cdot \beta_0, h_{\alpha_0}(\beta) \leq s} \Delta_{x_{\alpha_0}^-(\beta)} + \Delta_{x_{\alpha_0}^+(\beta)} \xrightarrow{*} \frac{d\mathrm{Leb}_{\mathbb{C}}(z)}{|Q_{\alpha_0}(z)|^2}. \quad (29)$$

Proof. Let r and r' be the smallest positive integers such that $\widehat{\alpha_0}^r \in G$ and $\widehat{\beta_0}^{r'} \in G$ respectively, so that, as seen at the end of the proof of Theorem 13 (for K as in Theorem 21, but this is clear for $K = \mathbb{Q}$, since then $m_{\Gamma_K}(\alpha_0) = m_G(\alpha_0) = 1$),

$$[\Gamma_{\alpha_0} : G_{\alpha_0}] = \frac{m_{\Gamma_K}(\alpha_0)}{m_G(\alpha_0)} r, \quad (30)$$

and similarly for β_0 .

To prove the counting claims in the above two theorems, we apply Corollary 16 with $n = 2$ or $n = 3$, $\Gamma = G$ (which is arithmetic), $\gamma_0 = \widehat{\alpha_0}^r$ and $\gamma_1 = \widehat{\beta_0}^{r'}$.

Using in the series of equalities below respectively

- the bijection $\beta \mapsto [\beta]$ from $G_{\alpha_0} \backslash G \cdot \beta_0$ to $G_{\alpha_0} \backslash G / G_{\beta_0}$ (with inverse $[\gamma] \mapsto \gamma\beta_0$), Equation (27) and the definition of the complexity $h(\alpha_0)$,
 - the fact that $[G_{\alpha_0} : \widehat{\alpha_0}^{r\mathbb{Z}}] = m_G(\alpha_0)$ and similarly for β_0 , as in the beginning of the proof of Corollary 16,
 - Corollary 16, noting that since $\mathrm{PSL}_2(\mathbb{Z})$ acts freely on $T^1\mathbb{H}_{\mathbb{R}}^2$, all multiplicities $m_{e,\gamma}$ are equal to 1, if $K = \mathbb{Q}$, and otherwise, as seen in the proof of Theorem 13, the multiplicities $m_{e,\gamma}$ different from 1 contribute only in a negligible way to the sums,
- we have, as $s \rightarrow +\infty$,

$$\begin{aligned}
& \mathrm{Card}\{\beta \in G_{\alpha_0} \backslash G \cdot \beta_0 : h_{\alpha_0}(\beta) \leq s\} \\
&= \mathrm{Card}\{[\gamma] \in G_{\alpha_0} \backslash G / G_{\beta_0} : h_{\widehat{\alpha_0}}(\gamma\widehat{\beta_0}\gamma^{-1}) \leq \ln(2h(\alpha_0)s + O(1))\} \\
&= \frac{1 + o(1)}{m_G(\alpha_0)m_G(\beta_0)} \mathrm{Card}\{[\gamma] \in \widehat{\alpha_0}^{r\mathbb{Z}} \backslash G / \widehat{\beta_0}^{r'\mathbb{Z}} : h_{\widehat{\alpha_0}}(\gamma\widehat{\beta_0}\gamma^{-1}) \leq \ln(2h(\alpha_0)s + O(1))\} \\
&= \frac{\mathrm{Vol}(\mathbb{S}^{n-2})^2 r \ell(\widehat{\alpha_0}) r' \ell(\widehat{\beta_0})}{m_G(\alpha_0)m_G(\beta_0) 2^{n-1} (n-1) \mathrm{Vol}(\mathbb{S}^{n-1}) \mathrm{Vol}(G \backslash \mathbb{H}_{\mathbb{R}}^n)} (2h(\alpha_0)s)^{n-1} (1 + O(s^{-\kappa})).
\end{aligned}$$

When $K = \mathbb{Q}$ and $n = 2$, this proves the counting claim in Theorem 20, by using Equations (16), (17) and (30), and noting that $m_{\Gamma_K}(\alpha_0) = m_{\Gamma_K}(\beta_0) = 1$.

When K is quadratic imaginary and $n = 3$, this proves the counting claim in Theorem 21, by using Equations (7), (20) and (30).

To prove the equidistribution claims in the above two theorems, we need to use a stronger geometric equidistribution theorem (also proven in [PP8]) than the one stated in Section 2.

For any $\beta \in G \cdot \beta_0$, let $v_{\alpha_0}(\beta)$ be the initial tangent vector of the common perpendicular from the geodesic line $D^- = \mathrm{Axis}(\widehat{\alpha_0})$ with endpoints α_0 and α_0^σ to that with endpoints β and β^σ , if it exists (and the summations below are on $\beta \in G \cdot \beta_0$ such that it does exist). Let $D^+ = \mathrm{Axis}(\widehat{\beta_0})$ and G_{D^+} its stabiliser in G , so that as already seen $\mathrm{Vol}(G_{D^+} \backslash D^+) = \frac{r' \ell(\widehat{\beta_0})}{\iota_G(\beta_0)}$. By replacing the equidistribution of initial points by the one of the initial tangent vectors (see [PP8, Coro. 20]), and by arguments similar to the ones leading to Equations (15) and (19), we have, by the expression of the skinning measure and Bowen-Margulis measure recalled in Section 2, with $n = 2$ if $K = \mathbb{Q}$ and $n = 3$ otherwise, as $t \rightarrow +\infty$,

$$\frac{2^{n-1} (n-1) m_G(\beta_0) \mathrm{Vol}(\mathbb{S}^{n-1}) \mathrm{Vol}(G \backslash \mathbb{H}_{\mathbb{R}}^n)}{\iota_G(\beta_0) \mathrm{Vol}(\mathbb{S}^{n-2}) \mathrm{Vol}(G_{D^+} \backslash D^+) e^{(n+1)t}} \sum_{\beta \in G \cdot \beta_0, h_{\widehat{\alpha_0}}(\widehat{\beta}) \leq t} \Delta_{v_{\alpha_0}(\beta)} \xrightarrow{*} \mathrm{Vol}_{\partial_+^1 D^-}.$$

Hence, by Equation (27), as $s \rightarrow +\infty$,

$$\frac{(n-1) m_G(\beta_0) \mathrm{Vol}(\mathbb{S}^{n-1}) \mathrm{Vol}(G \backslash \mathbb{H}_{\mathbb{R}}^n)}{\mathrm{Vol}(\mathbb{S}^{n-2}) r' \ell(\widehat{\beta_0}) h(\alpha_0)^{n-1} s^{n-1}} \sum_{\beta \in G \cdot \beta_0, h_{\alpha_0}(\beta) \leq s} \Delta_{v_{\alpha_0}(\beta)} \xrightarrow{*} \mathrm{Vol}_{\partial_+^1 D^-}. \quad (31)$$

We give in the following two lemmas the main computations used to deduce from this the equidistribution claims in the above two theorems.

Lemma 22 *If the geodesic lines $]\alpha_0, \alpha_0^\sigma[$ and $]\beta, \beta^\sigma[$ in $\mathbb{H}_{\mathbb{R}}^n$ are disjoint, then the endpoints of the geodesic line containing their common perpendicular are the points $x_{\alpha_0}^\pm(\beta)$ given by Equation (1), which are hence well defined.*

Proof. Let $]x^-, x^+[$ be this geodesic line. By Lemma 17 (1), and since two nonintersecting geodesic lines have one and only one geodesic line orthogonal to both of them, the points x^- and x^+ are the two solutions of the pair of equations $[x^-, x^+, \alpha_0, \alpha_0^\sigma] = -1 = [x^-, x^+, \beta, \beta^\sigma]$. An easy computation shows that this system is equivalent to

$$\begin{cases} x^- + x^+ = 2 \frac{\mathbf{n}(\alpha_0) - \mathbf{n}(\beta)}{\mathbf{tr} \alpha_0 - \mathbf{tr} \beta} \\ x^- x^+ = \frac{\mathbf{tr} \beta \mathbf{n}(\alpha_0) - \mathbf{tr} \alpha_0 \mathbf{n}(\beta)}{\mathbf{tr} \alpha_0 - \mathbf{tr} \beta}. \end{cases}$$

Solving a quadratic equation gives the result. \square

Lemma 23 *For $n = 2, 3, 5$, for all distinct x, y in $\partial_\infty \mathbb{H}_{\mathbb{R}}^n - \{\infty\}$, the pushforward of the Riemannian measure of the unit normal bundle of the geodesic line $L =]x, y[$ in $\mathbb{H}_{\mathbb{R}}^n$ by the positive endpoint map $v \mapsto z = v_+$ from $\partial_+^1 L$ to $\partial_\infty \mathbb{H}_{\mathbb{R}}^n - \{x, y\}$ is*

$$\frac{\|x - y\|^{n-1}}{\|(z - x)(z - y)\|^{n-1}} d\text{Leb}_{\mathbb{R}^{n-1}}(z).$$

By symmetry, the same result holds for the pushforward by the negative endpoint map $v \mapsto z = v_-$. When $n = 2, 3$, $x = \alpha_0$ and $y = \alpha_0^\sigma$, this measure is $\frac{2^{n-1} d\text{Leb}_{\widehat{K}}(z)}{h(\alpha_0)^{n-1} |Q_{\alpha_0}(z)|^{n-1}}$, with the previous notation.

Proof. Let $\widehat{K} = \mathbb{R}$ if $n = 2$, $\widehat{K} = \mathbb{C}$ if $n = 3$ and $\widehat{K} = \mathbb{H}$ if $n = 5$ (we denote $|z| = \mathbf{N}(z)^{1/2}$ if $z \in \mathbb{H}$), so that $\partial_\infty \mathbb{H}_{\mathbb{R}}^n - \{\infty\} = \widehat{K}$.

If L_∞ is the geodesic line with points at infinity $0 \in \widehat{K}$ and ∞ , then for all $v \in \partial_+^1 L_\infty$, we have $d\text{Vol}_{\partial_+^1 L_\infty}(v) = \frac{dt}{t} d\text{Vol}_{\mathbb{S}^{n-2}}(\sigma)$, where $t > 0$ is the last coordinate of $\pi(v)$ and $\sigma \in \mathbb{S}^{n-2}$ is the parameter of the Euclidean horizontal vector v (note that $d\text{Vol}_{\mathbb{S}^0}$ is the counting measure on $\{-1, 1\}$ if $n = 2$). If $z = v_+ \in \widehat{K} - \{0\}$ is the positive point at infinity of v , then $z = t\sigma$ and

$$d\text{Vol}_{\partial_+^1 L_\infty}(v) = \frac{d\text{Leb}_{\widehat{K}}(z)}{|z|^{n-1}}.$$

We may assume that $x > y$ if $n = 2$. The homography $A : z \mapsto (z - x)(z - y)^{-1}$ maps x to 0 and y to ∞ . Since it may be written $z \mapsto (\lambda z - \lambda x)(\lambda z - \lambda y)^{-1}$ with λ in (the center of) \widehat{K} such that $\lambda^2 = (x - y)^{-1}$ if $\widehat{K} = \mathbb{R}, \mathbb{C}$ and $\lambda = \frac{1}{\mathbf{N}(y - x)}$ if $\widehat{K} = \mathbb{H}$ (see Equation (9) to check the Dieudonné determinant is 1), it is the extension at infinity of an isometry of $\mathbb{H}_{\mathbb{R}}^n$ (by Poincaré's extension), sending $\text{Vol}_{\partial_+^1 L}$ to $\text{Vol}_{\partial_+^1 L_\infty}$. Since $|\dot{A}(z)| = \frac{|x - y|}{|z - y|^2}$, we have

$$(A^{-1})_* \left(\frac{d\text{Leb}_{\widehat{K}}(z)}{|z|^{n-1}} \right) = \frac{|x - y|^{n-1}}{|(z - x)(z - y)|^{n-1}} d\text{Leb}_{\widehat{K}}(z).$$

The result follows. \square

By the continuity of the pushforward maps of measures, by summing the pushforward of Equation (31) both by the positive and by the negative endpoint map, by the comments following Lemma 23, and since $\{x_{\alpha_0}^-(\beta), x_{\alpha_0}^+(\beta)\} = \{v_{\alpha_0}(\beta)_-, v_{\alpha_0}(\beta)_+\}$ by Lemma 22, we have, as $s \rightarrow +\infty$,

$$\frac{(n-1) m_G(\beta_0) \text{Vol}(\mathbb{S}^{n-1}) \text{Vol}(G \backslash \mathbb{H}_{\mathbb{R}}^n)}{2^{n-1} \text{Vol}(\mathbb{S}^{n-2}) r' \ell(\widehat{\beta}_0) s^{n-1}} \sum_{\beta \in G \cdot \beta_0, h_{\alpha_0}(\beta) \leq s} \Delta_{x_{\alpha_0}^-(\beta)} + \Delta_{x_{\alpha_0}^+(\beta)} \xrightarrow{*} \frac{2 d\text{Leb}_{\widehat{K}}(z)}{|Q_{\alpha_0}(z)|^{n-1}}.$$

The equidistribution claims in Theorems 20 and 21 follow, as in the end of the proof of their counting claims. \square

Remark. If $\widehat{K} = \mathbb{H}$, the crossratio of a quadruple of pairwise distinct points (a, b, c, d) in \widehat{K} may be defined as

$$[a, b, c, d] = (c - b)^{-1}(c - a)(d - a)^{-1}(d - b).$$

Though this crossratio does not extend continuously to the quadruples of pairwise distinct points in the one-point compactification $\mathbb{H} \cup \{0\}$, and it is only invariant under $\mathrm{SL}_2(\mathbb{H})$ up to conjugation, it also characterises the orthogonality of the geodesic lines in $\mathbb{H}_{\mathbb{R}}^5$ by the same formula (two geodesic lines $]a_-, a_+[$ and $]b_-, b_+[$ in $\mathbb{H}_{\mathbb{R}}^5$ are orthogonal if and only if $[a_-, a_+, b_-, b_+] = -1$). Furthermore, its absolute crossratio $|c - b|^{-1}|c - a||d - a|^{-1}|d - b|$ (we denote $|z| = \mathrm{N}(z)^{1/2}$ if $z \in \mathbb{H}$) does extend continuously (as already mentioned) and is invariant under $\mathrm{SL}_2(\mathbb{H})$ (see for instance [DeR] for all this).

Let A be a definite quaternion algebra over \mathbb{Q} , let \mathcal{O} be a maximal order in A , let G be a finite index subgroup of $\mathrm{PSL}_2(\mathcal{O})$, and let $\alpha_0, \beta_0 \in \mathbb{H}$ be two loxodromic quadratic irrationals over \mathcal{O} (see the definitions above Theorem 15). We may also define the *relative height* of an element β of $\mathrm{PSL}_2(\mathcal{O}) \cdot \beta_0$ with respect to α_0 by the same formula (24). It is still related to absolute crossratios by Equation (25), and still satisfies the properties of Lemma 18. We do have a counting asymptotic of the elements of $\beta \in G_{\alpha_0} \backslash G \cdot \beta_0$ with relative height at most s with respect to α_0 , as $s \rightarrow +\infty$, analogous to Theorem 20 and 21, and a related equidistribution result, but now, due to the higher difficulty of solving quadratic equations in the noncommutative \mathbb{H} , the points $x_{\alpha_0}^{\pm}(\beta)$ are much less explicit: they can only be defined as the unique two solutions x^-, x^+ of the pair of equations $[x^-, x^+, \alpha_0, \alpha_0^{\sigma}] = -1 = [x^-, x^+, \beta, \beta^{\sigma}]$. We leave the precise statement to the interested reader.

4.4 Counting crossratios and application to generalised Schottky-Klein functions

The aim of this subsection is to generalise the work of Pollicott [Pol], using the asymptotic of crossratios to prove the convergence of Schottky-Klein functions, to a much larger class of Kleinian groups.

Let \widetilde{M} be a complete simply connected Riemannian manifold with (dimension at least 2 and) pinched negative sectional curvature $-b^2 \leq K \leq -1$. Let Γ be a torsion free (only to simplify the statements) nonelementary (that is, not virtually nilpotent) discrete group of isometries of \widetilde{M} , with critical exponent δ_{Γ} and limit set $\Lambda\Gamma$ (defined as in Section 2). Let D^- and D^+ be two geodesic lines in \widetilde{M} with points at infinity in the *domain of discontinuity* $\Omega\Gamma = \partial_{\infty}\widetilde{M} - \Lambda\Gamma$ of Γ .

The Bowen-Margulis measure m_{BM} and skinning measures $\sigma_{D^{\pm}}$ are defined as in Section 2 where $\widetilde{M} = \mathbb{H}_{\mathbb{R}}^n$. We refer to [DaOP] for finiteness criteria of m_{BM} (for instance satisfied if Γ is convex-cocompact, or when Γ is geometrically finite in constant curvature (in which case δ_{Γ} is the Hausdorff dimension of $\Lambda\Gamma$), but there are many more examples), and [Bab] for mixing criteria of m_{BM} under the geodesic flow (for instance satisfied, when m_{BM} is finite, if \widetilde{M} has constant curvature, but there are many more examples).

The following result generalises [Pol, Theo. 1.4] (whose proof used symbolic dynamics and transfer operator techniques), which was only stated for $\widetilde{M} = \mathbb{H}_{\mathbb{R}}^3$, Γ a Schottky group and $D^- = D^+$, with a nonexplicit multiplicative constant.

Theorem 24 *If m_{BM} is finite and mixing, then, as $s \rightarrow +\infty$,*

$$\text{Card}\{\gamma \in \Gamma : d(D^-, \gamma D^+) \leq s\} \sim \frac{\|\sigma_{D^-}\| \|\sigma_{D^+}\|}{\delta_\Gamma \|m_{\text{BM}}\|} e^{\delta_\Gamma s}.$$

Proof. Since the endpoints of D^\pm are in the domain of discontinuity of γ , the support of the measure $\tilde{\sigma}_{D^\pm}$ is compact, hence the skinning measure σ_{D^\pm} is finite. The result then follows from [PP8, Coro. 20], since the stabiliser of D^\pm is trivial. \square

Until the end of Subsection 4.4, we assume that $\widetilde{M} = \mathbb{H}_{\mathbb{R}}^3$. The next result on asymptotic properties of crossratios is deduced from Theorem 24 as Theorem 1.3 is deduced from Theorem 1.4 in [Pol].

Corollary 25 *If m_{BM} is finite, then for all z, ξ in $\Omega\Gamma$, as $\epsilon \rightarrow 0$,*

$$\text{Card}\{\gamma \in \Gamma : |[z, \xi, \gamma z, \gamma \xi] - 1| \geq \epsilon\} \sim \frac{4^{\delta_\Gamma} \|\sigma_{D^-}\| \|\sigma_{D^+}\|}{\delta_\Gamma \|m_{\text{BM}}\|} \epsilon^{-\delta_\Gamma}.$$

The next result is proven as [Pol, Coro. 1.2], and defines a much larger class of holomorphic maps in two variables than the Schottky-Klein functions (see loc. cit. for motivations and references). We denote by Γ_0 a choice of a representative modulo inverse of the non-trivial elements of Γ .

Corollary 26 *If m_{BM} is finite and $\delta_\Gamma < 1$, then the function*

$$(z, \xi) \mapsto (z - \xi) \prod_{g \in \Gamma_0} \frac{(gz - z)(g\xi - z)}{(g\xi - \xi)(g\xi - z)},$$

where the terms in the product are ordered by the distance of their absolute value to 1, converges uniformly on compact subsets of $\Omega\Gamma \times \Omega\Gamma$.

5 Counting and equidistribution around binary quadratic, Hermitian and Hamiltonian forms

Let $Q(x, y) = ax^2 + bxy + cy^2$ be a binary quadratic form, which is primitive integral (its coefficients a, b, c are relatively prime elements in \mathbb{Z}). Let $\text{Disc}_Q = b^2 - 4ac$ be the discriminant of Q . The group $\text{SL}_2(\mathbb{Z})$ acts on the set of integral binary quadratic forms by precomposition, and we denote by $\text{SO}_Q(\mathbb{Z})$ the stabiliser of Q . When Q is positive definite, the following asymptotic is originally due to Gauss with a weaker error bound:

$$\text{Card}\{(u, v) \in \mathbb{Z} \times \mathbb{Z} : (u, v) = 1, Q(u, v) \leq s\} = \frac{12}{\pi \sqrt{-\text{Disc}_Q}} s + O(s^{131/416}),$$

the current error bound is proved in [Hux]. This asymptotic, with a less explicit error bound, can be given a geometric proof using $\text{PSL}_2(\mathbb{Z})$ -orbits of horoballs and points in $\mathbb{H}_{\mathbb{R}}^2$ in the same way as Theorem 34 in Subsection 5.2 below.

When Q is indefinite, the group of automorphs $\text{SO}_Q(\mathbb{Z})$ is infinite, and it is appropriate to count $\text{SO}_Q(\mathbb{Z})$ -orbits of relatively prime representations of integers by Q . Let

$$R_Q = \ln \frac{t_Q + u_Q \sqrt{\text{Disc}_Q}}{2}$$

be the regulator of Q , where (t_Q, u_Q) is the fundamental solution of the Pell-Fermat equation $t^2 - \text{Disc}_Q u^2 = 4$. The asymptotic behavior of this counting function is (see for example [Coh2, p. 164])

$$\text{Card}(\text{SO}_Q(\mathbb{Z}) \setminus \{(u, v) \in \mathbb{Z} \times \mathbb{Z} : (u, v) = 1, |Q(u, v)| \leq s\}) = \frac{12 R_Q}{\pi^2 \sqrt{\text{Disc}_Q}} s + O(\sqrt{s}). \quad (32)$$

A geometric proof of the asymptotic (32) with a less explicit error bound can be obtained from the proof of [PP4, Théo. 4.5], using Corollary 9 instead of [PP4, Cor. 3.9].

The results in Subsection 5.1, generalising the asymptotic (32) to binary Hermitian and Hamiltonian forms, are the versions with error terms and with related equidistribution results of respectively Theorem 4, Theorem 1, Corollary 3 in [PP3], and Theorem 13, Corollary 18, Theorem 1, Corollary 2 in [PP5]. Similarly, the error term in Corollary 6 of [PP3] is $O(s^{2-\kappa})$ for some $\kappa > 0$.

Given an arithmetic group Γ of isometries of a totally geodesic subspace X_Γ of $\mathbb{H}_{\mathbb{R}}^n$, we will denote by $\text{Covol}(\Gamma)$ the volume of the Riemannian orbifold $\Gamma \backslash X_\Gamma$.

5.1 Indefinite binary Hermitian and Hamiltonian forms

Let K and \mathfrak{m} be as in the beginning of Subsection 3.1. Fix a binary Hermitian form $f : \mathbb{C}^2 \rightarrow \mathbb{R}$ with

$$f(u, v) = a|u|^2 + 2\text{Re}(b\bar{u}v) + c|v|^2 \quad (33)$$

which is integral over \mathcal{O}_K (its coefficients $a = a(f), b = b(f), c = c(f)$ satisfy $a, c \in \mathbb{Z}$ and $b \in \mathcal{O}_K$). The group $\text{SL}_2(\mathbb{C})$ acts on the right on the binary Hermitian forms by precomposition, and the action of the Bianchi subgroup $\Gamma_K = \text{SL}_2(\mathcal{O}_K)$ preserves the integral ones. The group $\text{SU}_f(\mathcal{O}_K)$ of automorphs of f consists of the elements $g \in \Gamma_K$ stabilising f , that is, such that $f \circ g = f$.

In this subsection, we assume that f is indefinite (its discriminant $\text{Disc}(f) = |b|^2 - ac$ is positive), a feature which is preserved by the action of $\text{SL}_2(\mathbb{C})$.

Let G be a finite index subgroup of Γ_K . Let $\Gamma_{K,x,y}$ and $G_{x,y}$ be the stabilisers of $(x, y) \in K \times K$ in Γ_K and G respectively. Let $\iota_G = 1$ if $-\text{id} \in G$ and $\iota_G = 2$ otherwise. For every x, y in \mathcal{O}_K not both zero, and for every $s > 0$, let

$$\psi_{f,G,x,y}(s) = \text{Card} \text{SU}_f(\mathcal{O}_K) \cap G \setminus \{(u, v) \in G(x, y) : \mathfrak{n}(\mathcal{O}_K x + \mathcal{O}_K y)^{-1} |f(u, v)| \leq s\}.$$

Theorem 27 *There exists $\kappa > 0$ such that, as s tends to $+\infty$,*

$$\psi_{f,G,x,y}(s) \sim \frac{\pi \iota_G [\Gamma_{K,x,y} : G_{x,y}] \text{Covol}(\text{SU}_f(\mathcal{O}_K) \cap G)}{2 |D_K| \zeta_K(2) \text{Disc}(f) [\Gamma_K : G]} s^2 (1 + O(s^{-\kappa})).$$

If $y \neq 0$, then, as $s \rightarrow +\infty$,

$$\frac{[\Gamma_K : G] |D_K|^{\frac{3}{2}} \zeta_K(2) \text{Disc}(f) \mathfrak{n}(y)}{\pi^2 \iota_G \text{Covol}(\text{SU}_f(\mathcal{O}_K) \cap G) s^2} \sum_{f' \in G \cdot f, 0 < |f'(x, y)| \leq s} \frac{\Delta_{\frac{a(f')x + b(f')y}{f'(x, y)}}}{f'(x, y)} \xrightarrow{*} \text{Leb}_{\mathbb{C}}.$$

If Λ is the lattice of $\lambda \in \mathcal{O}_K$ such that $\pm \begin{pmatrix} 1 & \lambda \\ 0 & 1 \end{pmatrix} \in G$, identifying \mathcal{O}_K with the upper triangular unipotent subgroup of Γ_K , we have, as $s \rightarrow +\infty$,

$$\begin{aligned} \text{Card}\{f' \in \Lambda \backslash G \cdot f : 0 < |a(f')| \leq s\} \\ = \frac{\pi^2 \iota_G[\mathcal{O}_K : \Lambda] \text{Covol}(\text{SU}_f(\mathcal{O}_K) \cap G)}{2 [\Gamma_K : G] |D_K|^{\frac{1}{2}} \zeta_K(2) \text{Disc}(f)} s^2 (1 + O(s^{-\kappa})). \end{aligned}$$

For smooth functions ψ with compact support on \mathbb{C} , there is an error term in the above equidistribution claim, of the form $O(s^{2-\kappa} \|\psi\|_\ell)$ where $\kappa > 0$ and $\|\psi\|_\ell$ is the Sobolev norm of ψ for some $\ell \in \mathbb{N}$.

Proof. The first claim is proven in the same way as [PP3, Thm. 4], using Corollary 9 instead of [PP4, Cor. 4.9].

Let us prove the second claim. Let $n = 3$ and $\hat{K} = \mathbb{C}$. We denote by \overline{H} the image in $\text{PSL}_2(\hat{K})$ of any subgroup H of $\text{SL}_2(\hat{K})$. As in the proof of Theorem 10, we consider the isometric action of $\text{PSL}_2(\hat{K})$ on (the upper halfspace model of) $\mathbb{H}_{\mathbb{R}}^n$. As explained in [PP3, §2] (where the convention for binary Hermitian form was to replace b by \bar{b}), when $a(f) \neq 0$, the totally geodesic hyperplane

$$\mathcal{C}(f) = \{(z, t) \in \mathbb{H}_{\mathbb{R}}^n : f(z, 1) + |a(f)| t^2 = 0\}$$

is preserved by $\text{SU}_f(\mathcal{O}_K)$, its boundary at infinity is the $(n-2)$ -sphere of center $-\frac{b(f)}{a(f)}$ and radius $\frac{\sqrt{\text{Disc}(f)}}{|a(f)|}$, the hyperbolic orbifold $\text{SU}_f(\mathcal{O}_K) \backslash \mathcal{C}(f)$ has finite volume (and so does $(\text{SU}_f(\mathcal{O}_K) \cap G) \backslash \mathcal{C}(f)$), and, for every $g \in \text{SL}_2(\hat{K})$,

$$\mathcal{C}(f \circ g) = g^{-1} \mathcal{C}(f).$$

Let $\rho = xy^{-1} \in K \cup \{\infty\}$, $\gamma_\rho = \begin{pmatrix} \rho & -1 \\ 1 & 0 \end{pmatrix} \in \text{SL}_2(K)$ if $y \neq 0$ and $\gamma_\rho = \text{id}$ otherwise.

We define $R_G(f) = 2$ if there exists an element g in G such that $f \circ g = -f$, and $R_G(f) = 1$ otherwise.

To prove the equidistribution claim of Theorem 27, we apply Equation (3) with $n = 3$, Γ the arithmetic group $\gamma_\rho^{-1} \overline{G} \gamma_\rho$, D^- the horoball $\mathcal{H}_\infty = \{(z, t) \in \mathbb{H}_{\mathbb{R}}^n : t \geq 1\}$ (noting that ∞ is indeed a parabolic fixed point of Γ , since $\rho = \gamma_\rho \infty$ is a parabolic fixed point of \overline{G}) and D^+ the totally geodesic hyperplane $\mathcal{C}(f \circ \gamma_\rho) = \gamma_\rho^{-1} \mathcal{C}(f)$. The stabiliser Γ_{D^+} of D^+ in Γ indeed has finite covolume in D^+ , since (by conjugation for the first equality and by [PP3, Eq. (8)] for the second one)

$$\text{Vol}(\Gamma_{D^+} \backslash D^+) = \text{Vol}(\text{Stab}_{\overline{G}}(\mathcal{C}(f)) \backslash \mathcal{C}(f)) = \frac{1}{R_G(f)} \text{Covol}(\text{SU}_f(\mathcal{O}_K) \cap G). \quad (34)$$

For every $\gamma \in \Gamma$, the origin of the common perpendicular from D^- to $\gamma D^+ = \mathcal{C}(f \circ \gamma_\rho \circ \gamma)$ (when it exists, that is when $a(f \circ \gamma_\rho \circ \gamma) \neq 0$ and the radius of the circle at infinity of $\mathcal{C}(f \circ \gamma_\rho \circ \gamma)$ is strictly less than 1), is $(-\frac{b(f \circ \gamma_\rho \circ \gamma)}{a(f \circ \gamma_\rho \circ \gamma)}, 1)$, and its hyperbolic length is $|\ln(\frac{\sqrt{\text{Disc}(f)}}{|a(f \circ \gamma_\rho \circ \gamma)|})| = \ln(\frac{|a(f \circ \gamma_\rho \circ \gamma)|}{\sqrt{\text{Disc}(f)}})$ (by invariance of the discriminant under $\text{SL}_2(\hat{K})$). Since D^+ has codimension 1 and $\text{PSL}_2(\hat{K})$ preserves the orientation of $\mathbb{H}_{\mathbb{R}}^n$, the pointwise

stabiliser of D^+ in Γ is trivial. Using a similar comment about the multiplicities as in the proof of Theorem 13, we hence have by taking $t = \ln \frac{s}{\sqrt{\text{Disc}(f)}}$ in Equation (3), as $s \rightarrow +\infty$,

$$\frac{(n-1) \text{Vol}(\mathbb{S}^{n-1}) \text{Vol}(\Gamma \backslash \mathbb{H}_{\mathbb{R}}^n)}{\text{Vol}(\mathbb{S}^{n-2}) \text{Vol}(\Gamma_{D^+} \backslash D^+) \left(\frac{s}{\sqrt{\text{Disc}(f)}}\right)^{n-1}} \sum_{\gamma \in \Gamma/\Gamma_{D^+}, 0 < |a(f \circ \gamma_\rho \circ \gamma)| \leq s} \Delta\left(-\frac{b(f \circ \gamma_\rho \circ \gamma)}{a(f \circ \gamma_\rho \circ \gamma)}, 1\right) \xrightarrow{*} \text{Vol}_{\partial D^-}.$$

Since the map from $G \cdot f$ to Γ/Γ_{D^+} defined by $f \circ \gamma \mapsto \gamma_\rho^{-1} \gamma \gamma_\rho \Gamma_{D^+}$ is a $R_G(f)$ -to-1 map, and since the pushforward of measures by the continuous map $(z, 1) \mapsto -z$ from ∂D^- to \widehat{K} sends $\text{Vol}_{\partial D^-}$ to $\text{Leb}_{\widehat{K}}$, we have

$$\frac{(n-1) \text{Vol}(\mathbb{S}^{n-1}) \text{Vol}(\Gamma \backslash \mathbb{H}_{\mathbb{R}}^n) \text{Disc}(f)^{n-1}}{\text{Vol}(\mathbb{S}^{n-2}) \text{Vol}(\Gamma_{D^+} \backslash D^+) R_G(f) s^{n-1}} \sum_{f' \in G \cdot f, 0 < |a(f' \circ \gamma_\rho)| \leq s} \frac{\Delta\left(\frac{b(f' \circ \gamma_\rho)}{a(f' \circ \gamma_\rho)}, 1\right)}{\Delta\left(\frac{b(f' \circ \gamma_\rho)}{a(f' \circ \gamma_\rho)}, 1\right)} \xrightarrow{*} \text{Leb}_{\widehat{K}}. \quad (35)$$

Using Humbert's formula (7), we have

$$\text{Vol}(\Gamma \backslash \mathbb{H}_{\mathbb{R}}^3) = \text{Vol}(\overline{G} \backslash \mathbb{H}_{\mathbb{R}}^3) = [\overline{\Gamma}_K : \overline{G}] \text{Vol}(\overline{\Gamma}_K \backslash \mathbb{H}_{\mathbb{R}}^3) = \frac{[\Gamma_K : G]}{\iota_G} \frac{|D_K|^{\frac{3}{2}} \zeta_K(2)}{4\pi^2}. \quad (36)$$

By Equations (35), (34) and (36), we have

$$\frac{[\Gamma_K : G] |D_K|^{\frac{3}{2}} \zeta_K(2) \text{Disc}(f)}{\iota_G \pi^2 \text{Covol}(\text{SU}_f(\mathcal{O}_K) \cap G) s^2} \sum_{f' \in G \cdot f, 0 < |a(f' \circ \gamma_\rho)| \leq s} \frac{\Delta\left(\frac{b(f' \circ \gamma_\rho)}{a(f' \circ \gamma_\rho)}, 1\right)}{\Delta\left(\frac{b(f' \circ \gamma_\rho)}{a(f' \circ \gamma_\rho)}, 1\right)} \xrightarrow{*} \text{Leb}_{\mathbb{C}}.$$

When $y = 0$, this proves Theorem 7 in the Introduction. By considering the measures induced on the compact quotient $\Lambda \backslash \mathbb{C}$, whose volume is $\frac{|D_K|}{2} [\mathcal{O}_K : \Lambda]$, and by applying Theorem 7 with error term to the constant function 1, we get the last claim of Theorem 27.

When $y \neq 0$, replacing s by $\frac{s}{|y|^2}$, using the pushforward of measures on \mathbb{C} by $z \mapsto -\overline{y^{-1}z}$, which sends $\text{Leb}_{\mathbb{C}}$ to $|y|^2 \text{Leb}_{\mathbb{C}}$, and since $a(f' \circ \gamma_\rho) = f' \circ \gamma_\rho(1, 0) = (y \overline{y})^{-1} f'(x, y)$ and $b(f' \circ \gamma_\rho) = -(a(f') \overline{p} + \overline{b(f')})$ by an easy computation, this gives the equidistribution claim in Theorem 27. \square

For every $s > 0$, we consider the integer (depending only on the ideal class of \mathfrak{m})

$$\psi_{f, \mathfrak{m}}(s) = \text{Card} \left(\text{SU}_f(\mathcal{O}_K) \backslash \left\{ (u, v) \in \mathfrak{m} \times \mathfrak{m} : \mathfrak{n}(\mathfrak{m})^{-1} |f(u, v)| \leq s, \quad u \mathcal{O}_K + v \mathcal{O}_K = \mathfrak{m} \right\} \right),$$

which is the number of nonequivalent \mathfrak{m} -primitive representations by f of rational integers with absolute value at most $s \mathfrak{n}(\mathfrak{m})$.

By taking $G = \Gamma_K$ and $x, y \in K$ such that $\mathfrak{m} = \mathcal{O}_K x + \mathcal{O}_K y$ (which exists, see for instance [EGM, §7]) in the first claim of Theorem 27, we have the following result.

Theorem 28 *There exists $\kappa > 0$ such that, as s tends to $+\infty$,*

$$\psi_{f, \mathfrak{m}}(s) \sim \frac{\pi \text{Covol}(\text{SU}_f(\mathcal{O}_K))}{2 |D_K| \zeta_K(2) \text{Disc}(f)} s^2 (1 + O(s^{-\kappa})). \quad \square$$

Using the results of [MR] (or other ways of obtaining formulas for $\text{Covol}(\text{SU}_f(\mathcal{O}_K))$), one gets very explicit versions of Theorem 28 in special cases. A constant $\iota(f) \in \{1, 2, 3, 6\}$ is defined as follows. If $\text{Disc}(f) \equiv 0 \pmod{4}$, let $\iota(f) = 2$. If the coefficients a and c of the form f as in Equation (33) are both even, let $\iota(f) = 3$ if $\text{Disc}(f) \equiv 1 \pmod{4}$, and let $\iota(f)$ be the remainder modulo 8 of $\text{Disc}(f)$ if $\text{Disc}(f) \equiv 2 \pmod{4}$. In all other cases, let $\iota(f) = 1$. The following result on integral binary Hermitian forms f over $\mathbb{Q}(i)$ follows from the first claim of Theorem 28 using $K = \mathbb{Q}(i)$ as in the proof of [PP3, Coro. 3].

Corollary 29 *There exists $\kappa > 0$ such that, as s tends to $+\infty$,*

$$\begin{aligned} \text{Card } \text{SU}_f(\mathbb{Z}[i]) \setminus \{(u, v) \in \mathbb{Z}[i]^2 : u\mathbb{Z}[i] + v\mathbb{Z}[i] \in \mathbb{Z}[i], |f(u, v)| \leq s\} \\ = \frac{\pi^2}{8 \iota(f) \zeta_{\mathbb{Q}(i)}(2)} \prod_{p \mid \text{Disc}(f)} \left(1 + \left(\frac{-1}{p}\right) p^{-1}\right) s^2 (1 + O(s^{-\kappa})), \end{aligned}$$

where p ranges over the odd positive rational primes and $\left(\frac{-1}{p}\right)$ is a Legendre symbol. \square

Let $\mathbb{H}, A, \mathcal{O}$ and the associated notation, be as in the beginning of Section 3.2. Let $f : \mathbb{H} \times \mathbb{H} \rightarrow \mathbb{R}$ be a binary Hamiltonian form, with

$$f(u, v) = a \mathbf{N}(u) + \text{Tr}(\bar{u} b v) + c \mathbf{N}(v), \quad (37)$$

which is integral over \mathcal{O} (its coefficients $a = a(f), b = b(f), c = c(f)$ satisfy $a, c \in \mathbb{Z}$ and $b \in \mathcal{O}$). In this subsection, we assume that f is indefinite (its discriminant $\text{Disc}(f) = \mathbf{n}(b) - ac$ is positive). We denote by $\Gamma_{\mathcal{O}} = \text{SL}_2(\mathcal{O})$ the Hamilton-Bianchi group of invertible 2×2 matrices with coefficients in \mathcal{O} (see Subsection 3.2 or [PP5, §3] for definitions). The group $\text{SL}_2(\mathbb{H})$ acts on the right on the binary Hermitian forms by precomposition and the action of $\Gamma_{\mathcal{O}}$ preserves the integral ones. The group $\text{SU}_f(\mathcal{O})$ of automorphs of f consists of the elements $g \in \Gamma_{\mathcal{O}}$ stabilising f , that is, such that $f \circ g = f$.

Let G be a finite index subgroup of $\Gamma_{\mathcal{O}}$. For all x, y in A not both zero, and for every $s > 0$, let

$$\psi_{f, G, x, y}(s) = \text{Card } \text{SU}_f(\mathcal{O}) \cap G \setminus \{(u, v) \in G(x, y) : \mathbf{n}(\mathcal{O}x + \mathcal{O}y)^{-1} |f(u, v)| \leq s\}.$$

Let $\Gamma_{\mathcal{O}, x, y}$ and $G_{x, y}$ be the stabilisers of $(x, y) \in A \times A$ for the left linear actions of $\Gamma_{\mathcal{O}}$ and G respectively. Let $K_{x, y}$ be the left fractional ideal \mathcal{O} if $xy = 0$ and $\mathcal{O}x \cap \mathcal{O}y$ otherwise, and $\mathcal{O}_r(K_{x, y})$ its right order. Let $\iota_G = 1$ if $-\text{id} \in G$, and $\iota_G = 2$ otherwise.

Theorem 30 *There exists $\kappa > 0$ such that, as s tends to $+\infty$,*

$$\begin{aligned} \psi_{f, G, x, y}(s) = \\ \frac{540 \iota_G [\Gamma_{\mathcal{O}, x, y} : G_{x, y}] \text{Covol}(\text{SU}_f(\mathcal{O}) \cap G)}{\pi^2 \zeta(3) |\mathcal{O}_r(K_{x, y})^\times| \text{Disc}(f)^2 [\Gamma_{\mathcal{O}} : G] \prod_{p \mid D_A} (p^3 - 1)(1 - p^{-1})} s^4 (1 + O(s^{-\kappa})), \end{aligned}$$

with p ranging over positive rational primes dividing D_A . As $s \rightarrow +\infty$, we have

$$\frac{\zeta(3) [\Gamma_{\mathcal{O}} : G] \text{Disc}(f)^4 \prod_{p \mid D_A} (p^3 - 1)(p - 1)}{2160 \iota_G \text{Covol}(\text{SU}_f(\mathcal{O}) \cap G) s^4} \sum_{f' \in G \cdot f : 0 < |a(f')| \leq s} \frac{\Delta_{\frac{b(f')}{a(f')}}}{a(f')} \stackrel{*}{\sim} \text{Leb}_{\mathbb{H}}$$

and if $y \neq 0$,

$$\frac{\zeta(3) [\Gamma_{\mathcal{O}} : G] \text{Disc}(f)^4 \prod_{p|D_A} (p^3 - 1)(p - 1)}{2160 \iota_G \text{Covol}(\text{SU}_f(\mathcal{O}) \cap G) s^4} \sum_{f' \in G \cdot f, 0 < |f'(x, y)| \leq s} \Delta_{\frac{a(f')x + b(f')y}{f'(x, y)}} \xrightarrow{*} \text{Leb}_{\mathbb{H}}.$$

If Λ is the lattice of $\lambda \in \mathcal{O}$ such that $\pm \begin{pmatrix} 1 & \lambda \\ 0 & 1 \end{pmatrix} \in G$, identifying \mathcal{O} with the upper triangular unipotent subgroup of $\Gamma_{\mathcal{O}}$, we have, as $s \rightarrow +\infty$,

$$\begin{aligned} & \text{Card}\{f' \in \Lambda \backslash G \cdot f : 0 < |a(f')| \leq s\} \\ &= \frac{540 \iota_G D_A \text{Covol}(\text{SU}_f(\mathcal{O}) \cap G)}{\zeta(3) [\Gamma_{\mathcal{O}} : G] \text{Disc}(f)^4 \prod_{p|D_A} (p^3 - 1)(p - 1)} s^4 (1 + O(s^{-\kappa})). \end{aligned}$$

For smooth functions ψ with compact support on \mathbb{H} , there is an error term in the above equidistribution claim, of the form $O(s^{4-\kappa} \|\psi\|_{\ell})$ where $\kappa > 0$ and $\|\psi\|_{\ell}$ is the Sobolev norm of ψ for some $\ell \in \mathbb{N}$.

Proof. The proof is completely analogous to that of Theorem 28, mostly replacing $n = 3$ by $n = 5$, $\widehat{K} = \mathbb{C}$ by $\widehat{K} = \mathbb{H}$, K by A , \mathcal{O}_K by \mathcal{O} , \mathfrak{n} by \mathbb{N} , and references to [PP3] to references to [PP5], so that Equation (35) is still valid, and then one replaces Humbert's formula (7) by Emery's formula (11), and the formula $\text{Vol}(\mathcal{O}_K \backslash \mathbb{C}) = \frac{\sqrt{|D_K|}}{4}$ by $\text{Vol}(\mathcal{O} \backslash \mathbb{H}) = \frac{D_A}{4}$. \square

Given two left fractional ideals $\mathfrak{m}, \mathfrak{m}'$ of \mathcal{O} and $s \geq 0$, let

$$\begin{aligned} & \psi_{f, \mathfrak{m}, \mathfrak{m}'}(s) \\ &= \text{Card} \text{SU}_f(\mathcal{O}) \backslash \left\{ (u, v) \in \mathfrak{m} \times \mathfrak{m} : \frac{|f(u, v)|}{\mathfrak{n}(\mathfrak{m})} \leq s, \mathcal{O}u + \mathcal{O}v = \mathfrak{m}, [K_{u, v}] = [\mathfrak{m}'] \right\}. \end{aligned}$$

The following result follows from the first claim of Theorem 30 as in the proof of [PP5, Coro. 18].

Corollary 31 *There exists $\kappa > 0$ such that, as s tends to $+\infty$,*

$$\psi_{f, \mathfrak{m}, \mathfrak{m}'}(s) \sim \frac{540 \text{Covol}(\text{SU}_f(\mathcal{O}))}{\pi^2 \zeta(3) |\mathcal{O}_r(\mathfrak{m}')^{\times}| \text{Disc}(f)^2 \prod_{p|D_A} (p^3 - 1)(1 - p^{-1})} s^4 (1 + O(s^{-\kappa})). \quad \square$$

For every $s \geq 0$, we consider the integer

$$\psi_{f, \mathfrak{m}}(s) = \text{Card} \text{SU}_f(\mathcal{O}) \backslash \left\{ (u, v) \in \mathfrak{m} \times \mathfrak{m} : \mathbb{N}(\mathfrak{m})^{-1} |f(u, v)| \leq s, \mathcal{O}u + \mathcal{O}v = \mathfrak{m} \right\},$$

which is the number of nonequivalent \mathfrak{m} -primitive representations by f of rational integers with absolute value at most $s \mathbb{N}(\mathfrak{m})$. The next result then follows from Corollary 31 as in the proof given in Corollary 19 of Theorem 1 in [PP5].

Corollary 32 *There exists $\kappa > 0$ such that, as s tends to $+\infty$,*

$$\psi_{f, \mathfrak{m}}(s) = \frac{45 D_A \text{Covol}(\text{SU}_f(\mathcal{O}))}{2 \pi^2 \zeta(3) \text{Disc}(f)^2 \prod_{p|D_A} (p^3 - 1)} s^4 (1 + O(s^{-\kappa})). \quad \square$$

The group of automorphs of the binary Hamiltonian form $(u, v) \mapsto \text{Tr}(\bar{u}v)$ (which is the standard real scalar product on \mathbb{H}) is

$$\text{Sp}_1(\mathcal{O}) = \left\{ g \in \text{SL}_2(\mathcal{O}) : {}^t\bar{g} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} g = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right\},$$

which is an arithmetic lattice in the symplectic group $\text{Sp}_1(\mathbb{H})$ over Hamilton's quaternion algebra. The following result follows from Corollary 32 as in the proof of [PP5, Coro. 2].

Corollary 33 *There exists $\kappa > 0$ such that, as s tends to $+\infty$,*

$$\begin{aligned} \text{Card } \text{Sp}_1(\mathcal{O}) \backslash \{ (u, v) \in \mathcal{O} \times \mathcal{O} : |\text{Tr}(\bar{u}v)| \leq s, \mathcal{O}u + \mathcal{O}v = \mathcal{O} \} = \\ \frac{D_A}{48 \zeta(3)} \prod_{p|D_A} \frac{p^2 + 1}{p^2 + p + 1} s^4 (1 + O(s^{-\kappa})) . \quad \square \end{aligned}$$

5.2 Positive definite binary Hermitian and Hamiltonian forms

Let K and \mathfrak{m} be as in the beginning of Subsection 3.1. Let f be an integral binary Hermitian form over \mathcal{O}_K as in the beginning of Subsection 5.1. In this subsection, we assume that f is positive definite (that is $\text{Disc}(f) < 0$ and $a(f) > 0$), a feature which is preserved by the action of the Bianchi group $\Gamma_K = \text{SL}_2(\mathcal{O}_K)$. Note that the group $\text{SU}_f(\mathcal{O}_K)$ of automorphs of f is then finite.

For every $s > 0$, we consider the following integer (its finiteness is part of the following proof), depending only on the ideal class of \mathfrak{m} ,

$$\psi_{f, \mathfrak{m}}^+(s) = \text{Card } \{ (u, v) \in \mathfrak{m} \times \mathfrak{m} : \mathfrak{n}(\mathfrak{m})^{-1} f(u, v) \leq s, \mathcal{O}_K u + \mathcal{O}_K v = \mathfrak{m} \} ,$$

which is the number of nonequivalent \mathfrak{m} -primitive representations by f of the rational integers at most $s \mathfrak{n}(\mathfrak{m})$.

Theorem 34 *There exists $\kappa > 0$ such that, as s tends to $+\infty$, we have*

$$\psi_{f, \mathfrak{m}}^+(s) \sim \frac{\pi^2}{|D_K| \zeta_K(2) |\text{Disc}(f)|} s^2 (1 + O(s^{-\kappa})) .$$

If G is a finite index subgroup of Γ_K , with $\iota_G = 1$ if $-\text{id} \in G$ and $\iota_G = 2$ otherwise, then, as $s \rightarrow +\infty$,

$$\frac{|\text{SU}_f(\mathcal{O}_K) \cap G| |\Gamma_K : G| |D_K|^{\frac{3}{2}} \zeta_K(2) |\text{Disc}(f)|}{2 \pi^2 \iota_G s^2} \sum_{f' \in G \cdot f, a(f') \leq s} \Delta_{\frac{b(f')}{a(f')}} \stackrel{*}{\sim} \text{Leb}_{\mathbb{C}} .$$

If Λ is the lattice of $\lambda \in \mathcal{O}_K$ such that $\pm \begin{pmatrix} 1 & \lambda \\ 0 & 1 \end{pmatrix} \in G$, identifying \mathcal{O}_K with the upper triangular unipotent subgroup of Γ_K , we have, as $s \rightarrow +\infty$,

$$\begin{aligned} \text{Card} \{ f' \in \Lambda \backslash G \cdot f : a(f') \leq s \} \\ = \frac{\pi^2 \iota_G [\mathcal{O}_K : \Lambda]}{|\text{SU}_f(\mathcal{O}_K) \cap G| |\Gamma_K : G| |D_K| \zeta_K(2) |\text{Disc}(f)|} s^2 (1 + O(s^{-\kappa})) . \end{aligned}$$

For smooth functions ψ with compact support on \mathbb{C} , there is an error term in the above equidistribution claim, of the form $O(s^{2-\kappa}\|\psi\|_\ell)$ where $\kappa > 0$ and $\|\psi\|_\ell$ is the Sobolev norm of ψ for some $\ell \in \mathbb{N}$.

Proof. Let $n = 3$ and $\widehat{K} = \mathbb{C}$.

We start the proof by describing the space of positive definite binary Hermitian forms in hyperbolic geometry terms. Let \mathcal{Q}^+ be the cone of positive definite binary Hermitian forms. The multiplicative group $\mathbb{R}_+ =]0, +\infty[$ acts on \mathcal{Q}^+ by multiplication and we define $\overline{\mathcal{Q}}^+ = \mathcal{Q}^+/\mathbb{R}_+$. The map $\Phi : \overline{\mathcal{Q}}^+ \rightarrow \mathbb{H}_{\mathbb{R}}^n = \widehat{K} \times \mathbb{R}_+$ induced by $f \mapsto (-\frac{b(f)}{a(f)}, \frac{\sqrt{-\text{Disc}(f)}}{a(f)})$ is a homeomorphism, which is (anti-)equivariant in the sense that $\Phi(f \circ g) = g^{-1}\Phi(f)$ for all $g \in \text{SL}_2(\widehat{K})$ and $f \in \mathcal{Q}^+$ (see for example [PP5, Prop. 22] for a proof for positive definite binary Hamiltonian forms, the above claims being obtained by embedding as usual \mathbb{C} in Hamilton's quaternion algebra \mathbb{H}).

Now, let x, y be not both zero in \mathcal{O}_K , with $x = 1$ if $y = 0$, and let $I_{x,y} = \mathcal{O}_K x + \mathcal{O}_K y$. We will more generally study the following counting function of the representations, in a given orbit of G , of integers by f , defined, for all $s \geq 0$, by

$$\psi_{f,G,x,y}^+(s) = \text{Card} \{ (u,v) \in G(x,y) : \mathbf{n}(I_{x,y})^{-1} f(u,v) \leq s \}.$$

Let ρ, γ_ρ be as in the proof of Theorem 27. We will apply Corollary 9 with Γ the arithmetic group $\gamma_\rho^{-1} \overline{G} \gamma_\rho$, D^- the horoball $\mathcal{H}_\infty = \{(z,t) \in \mathbb{H}_{\mathbb{R}}^n : t \geq 1\}$ (which is centered at a parabolic fixed point of Γ , and whose pointwise stabiliser is trivial), and D^+ the singleton consisting of $\Phi(f \circ \gamma_\rho)$.

For any $g \in \Gamma_K$, the distance of $\Phi(f \circ g)$ from \mathcal{H}_∞ is $\ln \frac{a(f \circ g)}{\sqrt{-\text{Disc}(f)}}$, when this number is positive, which is the case except for finitely many right classes of g under the stabiliser of \mathcal{H}_∞ . Note that $\gamma \in \Gamma$ belongs to the stabiliser Γ_{D^+} of D^+ in Γ if and only if γ^{-1} stabilises the positive homothety class of $f \circ \gamma_\rho$, that is, since $\text{SL}_2(\widehat{K})$ preserves the discriminant, if and only if γ^{-1} fixes $f \circ \gamma_\rho$. Thus $\Gamma_{D^+} = \gamma_\rho^{-1}(\text{SU}_f(\mathcal{O}_K) \cap G) \gamma_\rho$, which is also the pointwise stabiliser of D^+ , whose order is hence $|\text{SU}_f(\mathcal{O}_K) \cap G|$.

Let $G_{\mathcal{H}_\rho}$ and $G_{x,y}$ be the stabilisers in G of the horoball $\mathcal{H}_\rho = \gamma_\rho \mathcal{H}_\infty$ in $\mathbb{H}_{\mathbb{R}}^n$ and of the pair (x,y) in $\widehat{K} \times \widehat{K}$. As previously, we denote by \overline{H} the image in $\text{PSL}_2(\widehat{K})$ of any subgroup H of $\text{SL}_2(\widehat{K})$. We use the (surprising !) convention that $\mathbf{n}(y) = 1$ if $y = 0$ to avoid considering cases. The counting result with error term of Corollary 9, and the value of $C(D^-, D^+)$ in case (4) just above it, using cosets and multiplicities arguments already

seen, shows that

$$\begin{aligned}
\psi_{f,G,x,y}^+(s) &= \text{Card} \{ [\gamma] \in G/G_{x,y} : \mathbf{n}(I_{x,y})^{-1} f(\gamma(x,y)) \leq s \} \\
&= \frac{2}{\iota_G} \text{Card} \{ [\gamma] \in \overline{G}/\overline{G_{x,y}} : f \circ \gamma \circ \gamma_\rho(1,0) \leq \frac{s \mathbf{n}(I_{x,y})}{\mathbf{n}(y)} \} \\
&= \frac{2}{\iota_G} [\overline{G_{\mathcal{H}_\rho}} : \overline{G_{x,y}}] \text{Card} \{ [\gamma] \in \overline{G}/\overline{G_{\mathcal{H}_\rho}} : a(f \circ \gamma \circ \gamma_\rho) \leq \frac{s \mathbf{n}(I_{x,y})}{\mathbf{n}(y)} \} + O(1) \\
&= \frac{2}{\iota_G} [\overline{G_{\mathcal{H}_\rho}} : \overline{G_{x,y}}] |\text{SU}_f(\mathcal{O}_K) \cap G| \\
&\quad \text{Card} \{ [\gamma] \in \Gamma_{D^+} \backslash \Gamma / \Gamma_{D^-} : a(f \circ \gamma_\rho \circ \gamma) \leq \frac{s \mathbf{n}(I_{x,y})}{\mathbf{n}(y)} \} + O(1) \\
&= \frac{2}{\iota_G} [\overline{G_{\mathcal{H}_\rho}} : \overline{G_{x,y}}] |\text{SU}_f(\mathcal{O}_K) \cap G| \mathcal{N}_{D^+, D^-} \left(\ln \frac{s \mathbf{n}(I_{x,y})}{\mathbf{n}(y) \sqrt{-\text{Disc}(f)}} \right) + O(1) \\
&= \frac{2 [\overline{G_{\mathcal{H}_\rho}} : \overline{G_{x,y}}] \text{Vol}(\Gamma_{D^-} \backslash D^-)}{\iota_G \text{Vol}(\Gamma \backslash \mathbb{H}_{\mathbb{R}}^n)} \left(\frac{s \mathbf{n}(I_{x,y})}{\mathbf{n}(y) \sqrt{-\text{Disc}(f)}} \right)^{n-1} (1 + O(s^{-\kappa})). \quad (38)
\end{aligned}$$

As seen in the proof of Theorem 10, we have

$$\begin{aligned}
\text{Vol}(\Gamma_{D^-} \backslash D^-) &= \text{Vol}(\overline{G_{\mathcal{H}_\rho}} \backslash \mathcal{H}_\rho) = [\overline{(\Gamma_K)_{\mathcal{H}_\rho}} : \overline{G_{\mathcal{H}_\rho}}] \text{Vol}(\overline{(\Gamma_K)_{\mathcal{H}_\rho}} \backslash \mathcal{H}_\rho) \\
&= [\overline{(\Gamma_K)_{\mathcal{H}_\rho}} : \overline{G_{\mathcal{H}_\rho}}] \frac{\sqrt{|D_K|} \mathbf{n}(y)^2}{2 \omega_K \mathbf{n}(I_{x,y})^2}. \quad (39)
\end{aligned}$$

We have

$$\begin{aligned}
[\overline{(\Gamma_K)_{\mathcal{H}_\rho}} : \overline{G_{\mathcal{H}_\rho}}] [\overline{G_{\mathcal{H}_\rho}} : \overline{G_{x,y}}] &= [\overline{(\Gamma_K)_{\mathcal{H}_\rho}} : \overline{\Gamma_{K,x,y}}] [\overline{\Gamma_{K,x,y}} : \overline{G_{x,y}}] \\
&= \frac{\omega_K}{2} [\Gamma_{K,x,y} : G_{x,y}]. \quad (40)
\end{aligned}$$

Using Equation (36), Equation (38) then gives the following formula, interesting in itself,

$$\psi_{f,G,x,y}^+(s) = \frac{2 \pi^2 [\Gamma_{K,x,y} : G_{x,y}]}{|D_K| \zeta_K(2) |\text{Disc}(f)| [\Gamma_K : G]} s^2 (1 + O(s^{-\kappa})).$$

Since any nonzero ideal \mathfrak{m} in \mathcal{O}_K may be written $\mathfrak{m} = \mathcal{O}_K x_{\mathfrak{m}} + \mathcal{O}_K y_{\mathfrak{m}}$ for some $x_{\mathfrak{m}}, y_{\mathfrak{m}}$ not both zero in \mathcal{O}_K and $\psi_{f,\mathfrak{m}}^+ = \psi_{f,\Gamma_K, x_{\mathfrak{m}}, y_{\mathfrak{m}}}^+$, the first claim of Theorem 34 follows.

Let us now prove the equidistribution result in Theorem 34. The main additional remark is that for all $f' \in G \cdot f$, the initial point of the common perpendicular from \mathcal{H}_∞ to $\Phi(f' \circ \gamma_\rho)$ is $(-\frac{b(f' \circ \gamma_\rho)}{a(f' \circ \gamma_\rho)}, 1)$. Taking $t = \ln \frac{s}{\sqrt{-\text{Disc}(f)}}$ in Equation (3) in Corollary 9, gives, as $s \rightarrow +\infty$,

$$\frac{|\text{SU}_f(\mathcal{O}_K) \cap G| (n-1) \text{Vol}(\Gamma \backslash \mathbb{H}_{\mathbb{R}}^n)}{\left(\frac{s}{\sqrt{-\text{Disc}(f)}} \right)^{n-1}} \sum_{\gamma \in \Gamma / \Gamma_{D^+}, a(f \circ \gamma_\rho \circ \gamma) \leq s} \Delta_{\left(-\frac{b(f \circ \gamma_\rho \circ \gamma)}{a(f \circ \gamma_\rho \circ \gamma)}, 1 \right)} \xrightarrow{*} \text{Vol}_{\partial D^-}.$$

Hence, since the map from $G \cdot f$ to Γ / Γ_{D^+} defined by $f \circ \gamma \mapsto \gamma_\rho^{-1} \gamma \gamma_\rho \Gamma_{D^+}$ is a bijection, using the pushforward of measures by the map $(z, 1) \mapsto -z$ from ∂D^- to \widehat{K} , we have

$$\begin{aligned}
|\text{SU}_f(\mathcal{O}_K) \cap G| (n-1) \text{Vol}(\Gamma \backslash \mathbb{H}_{\mathbb{R}}^n) |\text{Disc}(f)|^{\frac{n-1}{2}} s^{-(n-1)} \sum_{f' \in G \cdot f, a(f' \circ \gamma_\rho) \leq s} \Delta_{\frac{b(f' \circ \gamma_\rho)}{a(f' \circ \gamma_\rho)}} \\
\xrightarrow{*} \text{Leb}_{\widehat{K}}. \quad (41)
\end{aligned}$$

Taking $y = 0$ (then $\gamma_\rho = \text{id}$) and using Equation (36), this gives the equidistribution claim in Theorem 34. The last claim of Theorem 27 follows from it as in the proof of Theorem 27.

The case $y \neq 0$, using Equation (36), gives the following equidistribution result, interesting in itself, as in the end of the proof of Theorem 27:

$$\frac{|\text{SU}_f(\mathcal{O}_K) \cap G| [\Gamma_K : G] |D_K|^{\frac{3}{2}} \zeta_K(2) |\text{Disc}(f)| \mathbf{n}(y)}{2 \pi^2 \iota_G s^2} \sum_{f' \in G \cdot f, f'(x, y) \leq s} \Delta_{\frac{a(f')x + b(f')y}{f'(x, y)}} \xrightarrow{*} \text{Leb}_{\mathbb{C}} . \quad \square$$

Let $\mathbb{H}, A, \mathcal{O}, \mathbf{m}$ and the associated notation be as in the beginning of Section 3.2. Let f be an integral binary Hamiltonian form over \mathcal{O} , as in Subsection 5.1. In this subsection, we assume that f is positive definite (that is $\text{Disc}(f) < 0$ and $a(f) > 0$), a feature which is preserved by the action (on the right) of the Hamilton-Bianchi group $\Gamma_{\mathcal{O}} = \text{SL}_2(\mathcal{O})$ (by precomposition). Note that the group $\text{SU}_f(\mathcal{O})$ of automorphs of f is then finite.

For every $s > 0$, we consider the integer

$$\psi_{f, \mathbf{m}}^+(s) = \text{Card} \left\{ (u, v) \in \mathbf{m} \times \mathbf{m} : \mathbf{N}(\mathbf{m})^{-1} f(u, v) \leq s, \quad \mathcal{O}u + \mathcal{O}v = \mathbf{m} \right\} ,$$

which is the number of nonequivalent \mathbf{m} -primitive representations by f of the rational integers at most $s \mathbf{N}(\mathbf{m})$.

Theorem 35 *There exists $\kappa > 0$ such that, as s tends to $+\infty$,*

$$\psi_{f, \mathbf{m}}^+(s) \sim \frac{60 D_A}{\zeta(3) |\text{Disc}(f)|^2 \prod_{p|D_A} (p^3 - 1)} s^4 (1 + O(s^{-\kappa})) .$$

If G is a finite index subgroup of $\Gamma_{\mathcal{O}}$, with $\iota_G = 1$ if $-\text{id} \in G$ and $\iota_G = 2$ otherwise, then, as $s \rightarrow +\infty$,

$$\frac{|\text{SU}_f(\mathcal{O}) \cap G| [\Gamma_{\mathcal{O}} : G] \zeta(3) |\text{Disc}(f)|^2 \prod_{p|D_A} (p^3 - 1)(p - 1)}{2880 \iota_G s^4} \sum_{f' \in G \cdot f, a(f') \leq s} \Delta_{\frac{b(f')}{a(f')}} \xrightarrow{*} \text{Leb}_{\mathbb{H}} .$$

If Λ is the lattice of $\lambda \in \mathcal{O}$ such that $\pm \begin{pmatrix} 1 & \lambda \\ 0 & 1 \end{pmatrix} \in G$, identifying \mathcal{O} with the upper triangular unipotent subgroup of $\Gamma_{\mathcal{O}}$, we have, as $s \rightarrow +\infty$,

$$\begin{aligned} & \text{Card}\{f' \in \Lambda \backslash G \cdot f : a(f') \leq s\} \\ &= \frac{720 \iota_G [\mathcal{O} : \Lambda] D_A}{|\text{SU}_f(\mathcal{O}) \cap G| [\Gamma_{\mathcal{O}} : G] \zeta(3) |\text{Disc}(f)|^2 \prod_{p|D_A} (p^3 - 1)(p - 1)} s^4 (1 + O(s^{-\kappa})) . \end{aligned}$$

Proof. The proof is completely analogous to that of Theorem 34, mostly replacing $n = 3$ by $n = 5$, $\widehat{K} = \mathbb{C}$ by $\widehat{K} = \mathbb{H}$, K by A , \mathcal{O}_K by \mathcal{O} , \mathbf{n} by \mathbf{N} . Given x, y not both zero in \mathcal{O} , with $x = 1$ if $y = 0$, let $I_{x, y}$ and $K_{x, y}$ be as in the beginning of the proof of Theorem 11.

We also introduce the following counting function of the representations of integers by f , in a given orbit of G :

$$\psi_{f,G,x,y}^+(s) = \text{Card} \left\{ (u,v) \in G(x,y) : \mathbf{N}(I_{x,y})^{-1} f(u,v) \leq s \right\}.$$

Keeping the relevant notation of the proof Theorem 34, Equation (38) is still valid. By Equation (10) where $\tau = 1$, we have

$$\begin{aligned} \text{Vol}(\Gamma_{D^-} \backslash D^-) &= \text{Vol}(\overline{G_{\mathcal{H}_\rho}} \backslash \mathcal{H}_\rho) = [\overline{(\Gamma_\mathcal{O})_{\mathcal{H}_\rho}} : \overline{G_{\mathcal{H}_\rho}}] \text{Vol}(\overline{(\Gamma_\mathcal{O})_{\mathcal{H}_\rho}} \backslash \mathcal{H}_\rho) \\ &= [\overline{(\Gamma_\mathcal{O})_{\mathcal{H}_\rho}} : \overline{G_{\mathcal{H}_\rho}}] \frac{D_A \mathbf{N}(y)^4}{16 |\mathcal{O}_r(K_{x,y})^\times| [\overline{(\Gamma_\mathcal{O})_{\mathcal{H}_\rho}} : \overline{(\Gamma_\mathcal{O})_{x,y}}] \mathbf{N}(I_{x,y})^4}. \end{aligned}$$

By Emery's formula (11), we have

$$\begin{aligned} \text{Vol}(\Gamma \backslash \mathbb{H}_{\mathbb{R}}^n) &= \text{Vol}(\overline{G} \backslash \mathbb{H}_{\mathbb{R}}^5) = [\overline{\Gamma_\mathcal{O}} : \overline{G}] \text{Vol}(\overline{\Gamma_\mathcal{O}} \backslash \mathbb{H}_{\mathbb{R}}^5) \\ &= \frac{[\Gamma_\mathcal{O} : G]}{\iota_G} \frac{\zeta(3) \prod_{p|D_A} (p^3 - 1)(p - 1)}{11520}. \end{aligned}$$

Since $\frac{[\overline{G_{\mathcal{H}_\rho}} : \overline{G_{x,y}}] [\overline{(\Gamma_\mathcal{O})_{\mathcal{H}_\rho}} : \overline{G_{\mathcal{H}_\rho}}]}{[(\Gamma_\mathcal{O})_{\mathcal{H}_\rho} : (\Gamma_\mathcal{O})_{x,y}]} = [\overline{(\Gamma_\mathcal{O})_{x,y}} : \overline{G_{x,y}}] = [(\Gamma_\mathcal{O})_{x,y} : G_{x,y}]$, we hence have the following counting asymptotic, interesting in itself:

$$\psi_{f,G,x,y}^+(s) = \frac{1440 D_A \iota_G [(\Gamma_\mathcal{O})_{x,y} : G_{x,y}]}{\zeta(3) [\Gamma_\mathcal{O} : G] |\mathcal{O}_r(K_{x,y})^\times| |\text{Disc}(f)|^2 \prod_{p|D_A} (p^3 - 1)(p - 1)} s^4 (1 + O(s^{-\kappa})). \quad (42)$$

Given two left fractional ideals $\mathfrak{m}, \mathfrak{m}'$ of \mathcal{O} and $s \geq 0$, let

$$\psi_{f,\mathfrak{m},\mathfrak{m}'}^+(s) = \text{Card} \left\{ (u,v) \in \mathfrak{m} \times \mathfrak{m} : \mathbf{n}(\mathfrak{m})^{-1} f(u,v) \leq s, I_{u,v} = \mathfrak{m}, [K_{u,v}] = [\mathfrak{m}'] \right\}.$$

As explained in the proof of Theorem 11, taking $(x,y) \in \mathcal{O} \times \mathcal{O}$ such that $[I_{x,y}] = [\mathfrak{m}]$ and $[K_{x,y}] = [\mathfrak{m}']$, we have $\psi_{f,\mathfrak{m},\mathfrak{m}'}^+ = \psi_{f,\Gamma_\mathcal{O},x,y}^+$. Hence

$$\psi_{f,\mathfrak{m},\mathfrak{m}'}^+(s) = \frac{1440 D_A}{\zeta(3) |\mathcal{O}_r(\mathfrak{m}')^\times| |\text{Disc}(f)|^2 \prod_{p|D_A} (p^3 - 1)(p - 1)} s^4 (1 + O(s^{-\kappa})).$$

Now, the first claim of Theorem 35 follows from Equation (13), since

$$\psi_{f,\mathfrak{m}}^+ = \sum_{[\mathfrak{m}'] \in \mathcal{O}^\times} \psi_{f,\mathfrak{m},\mathfrak{m}'}^+,$$

The last two assertions of Theorem 35 are proven in the same way as the last two assertions of Theorem 34, since Equation (41) is still valid, and if $y \neq 0$, we furthermore have

$$\begin{aligned} &\frac{|\text{SU}_f(\mathcal{O}) \cap G| [\Gamma_\mathcal{O} : G] \zeta(3) |\text{Disc}(f)|^2 \mathbf{n}(y)^2 \prod_{p|D_A} (p^3 - 1)(p - 1)}{2880 \iota_G s^4} \times \\ &\quad \sum_{f' \in G \cdot f, f'(x,y) \leq s} \Delta_{\frac{a(f')x + b(f')y}{f'(x,y)}} \stackrel{*}{\rightarrow} \text{Leb}_{\mathbb{H}}. \quad \square \end{aligned}$$

5.3 Representation of algebraic integers by integral binary quadratic forms

In this final subsection, we study counting and equidistribution problems of the representations of algebraic integers by quadratic norm forms (for related results, see for instance [Nag], [Odo, Thm. R], [Lan, Chap. VI], as well as [GP, §3.1] for an ergodic approach).

Let K and its associated notation be as in the beginning of Section 4. Let $\alpha_0 \in \widehat{K}$ be a fixed quadratic irrational over \mathcal{O}_K . We also denote by \mathbf{n} the relative norm in $K(\alpha_0)$ over K (which is consistent with the notation of the beginning of Section 4). We consider the (relative) norm form, seen as a map from $K \times K$ to K , defined by

$$N_{\alpha_0} : (u, v) \mapsto \mathbf{n}(u - \alpha_0 v) = u^2 - \mathrm{tr} \alpha_0 uv + \mathbf{n}(\alpha_0) v^2,$$

which is the homogeneous form of the minimal polynomial of α_0 over K . Its values on $\mathcal{O}_K \times \mathcal{O}_K$ belong to \mathcal{O}_K if α_0 is an algebraic integer, and to $a\mathcal{O}_K$ for some $a \in K$ in general. We will study the representations of elements of K by this norm form N_{α_0} , in orbits of (finite index subgroups of) the modular groups $\Gamma_K = \mathrm{SL}_2(\mathcal{O}_K)$ (a minor change of notation from the beginning of Section 4).

We denote by \overline{H} the image in $\mathrm{PSL}_2(\widehat{K})$ of any subgroup H of $\mathrm{SL}_2(\widehat{K})$. We denote by H_P the stabiliser, for any group action of a group H on a set X , of a point P or a subset P in X . The stabiliser $\mathrm{Stab}_{\Gamma_K} N_{\alpha_0}$ of the norm form N_{α_0} , for the action (on the right) of Γ_K by precomposition by the linear action, is exactly $(\Gamma_K)_{\{\alpha_0, \alpha_0^\sigma\}}$.

As a warm-up, when $K = \mathbb{Q}$, we give an equidistribution result of the fractions of the representations, in an aforementioned orbit, of usual rationals by this norm form (which, in this case, is an indefinite quadratic form, hence the classical counting results of the beginning of Section 5 apply).

Theorem 36 *For every finite index subgroup G of $\Gamma_{\mathbb{Q}} = \mathrm{SL}_2(\mathbb{Z})$, with $\iota_G = 1$ if $-\mathrm{id} \in G$ and $\iota_G = 2$ otherwise, we have the following convergence of measures on $\mathbb{R} - \{\alpha_0, \alpha_0^\sigma\}$ as $s \rightarrow +\infty$:*

$$\frac{\pi^2 \iota_G [\Gamma_{\mathbb{Q}} : G]}{12 [(\Gamma_{\mathbb{Q}})_{\infty} : G_{\infty}] s} \sum_{(u,v) \in G(0,1), |\mathbf{n}(u - \alpha_0 v)| \leq s} \Delta_{\frac{u}{v}} \xrightarrow{*} \frac{d\mathrm{Leb}_{\mathbb{R}}(t)}{|Q_{\alpha_0}(t)|}.$$

For smooth functions ψ with compact support on $\mathbb{R} - \{\alpha_0, \alpha_0^\sigma\}$, there is an error term in the above equidistribution claims, of the form $O(s^{1-\kappa} \|\psi\|_{\ell})$ where $\kappa > 0$ and $\|\psi\|_{\ell}$ is the Sobolev norm of ψ for some $\ell \in \mathbb{N}$.

For instance, taking $G = \mathrm{SL}_2(\mathbb{Z})$ gives Theorem 6 in the introduction, and taking for G the Hecke congruence subgroup modulo $k \in \mathbb{N} - \{0\}$, we have, by the index computation of for instance [Shi, p. 24],

$$\frac{\pi^2 k \prod_{p|k} (1 + \frac{1}{p})}{12 s} \sum_{(u,v) \in \mathbb{Z}^2, (u,v)=1, v \equiv 0 \pmod{k}, |\mathbf{n}(u - \alpha_0 v)| \leq s} \Delta_{\frac{u}{v}} \xrightarrow{*} \frac{d\mathrm{Leb}_{\mathbb{R}}(t)}{|Q_{\alpha_0}(t)|}.$$

Proof. We take $n = 2$ and $\widehat{K} = \mathbb{R}$ in this proof. We will apply Equation (4) with Γ the arithmetic group \overline{G} , D^- the geodesic line with points at infinity $\alpha_0, \alpha_0^\sigma$ (on which the group $\widehat{\alpha_0}^{\mathbb{Z}} \cap \overline{G}$ acts with compact quotient) and D^+ the horoball $\mathcal{H}_{\infty} = \{(z, t) \in \mathbb{H}_{\mathbb{R}}^n = \widehat{K} \times \mathbb{R}_+ : t \geq 1\}$ (which is centered at a parabolic fixed point of \overline{G}).

For every $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ in $\mathrm{SL}_2(\mathcal{O}_K)$, the distance ℓ_γ between D^- and γD^+ , when they are disjoint (which is the case except for finitely many double classes of γ in $\Gamma_{D^-} \backslash \Gamma / \Gamma_{D^+}$), is equal to the distance between $D^+ = \mathcal{H}_\infty$ and the geodesic line $\gamma^{-1} D^-$ with points at infinity $\gamma^{-1} \alpha_0, \gamma^{-1} \alpha_0^\sigma$. Hence by [PP4, Lem. 4.2]

$$\ell_\gamma = \left| \ln \frac{|\gamma^{-1} \alpha_0 - \gamma^{-1} \alpha_0^\sigma|}{2} \right| = \ln h(\gamma^{-1} \alpha_0) = \ln (h(\alpha_0) |\mathbf{n}(u - v \alpha_0)|) .$$

Let v_γ be the initial tangent vector of the common perpendicular from D^- to γD^+ , when they are disjoint, and note that its positive point at infinity $(v_\gamma)_+$ is the point at infinity of the horoball γD^+ , which is $\gamma \infty = \frac{a}{c}$.

By coset and multiplicities argument already seen, by the version for initial tangent vectors of Equation (4) seen in Subsection 4.3, we have

$$\frac{\mathrm{Vol}(\mathbb{S}^{n-1}) \mathrm{Vol}(\Gamma \backslash \mathbb{H}_{\mathbb{R}}^n)}{\mathrm{Vol}(\mathbb{S}^{n-2}) \mathrm{Vol}(\Gamma_{D^+} \backslash D^+)} e^t \sum_{\gamma \in \Gamma / \Gamma_{D^+}} \Delta_{v_\gamma} \xrightarrow{*} \mathrm{Vol}_{\partial_+^1 D^-} . \quad (43)$$

We have

$$\mathrm{Vol}(\Gamma \backslash \mathbb{H}_{\mathbb{R}}^n) = [\overline{\Gamma_{\mathbb{Q}}} : \overline{G}] \mathrm{Vol}(\overline{\Gamma_{\mathbb{Q}}} \backslash \mathbb{H}_{\mathbb{R}}^2) = \frac{[\Gamma_{\mathbb{Q}} : G] \pi}{\iota_G} \frac{1}{3} ,$$

and

$$\mathrm{Vol}(\Gamma_{D^+} \backslash D^+) = [\overline{(\Gamma_{\mathbb{Q}})_\infty} : \overline{G_\infty}] \mathrm{Vol}(\overline{(\Gamma_{\mathbb{Q}})_\infty} \backslash D^+) = \frac{[(\Gamma_{\mathbb{Q}})_\infty : G_\infty]}{\iota_G} .$$

Taking $t = \ln(h(\alpha_0)s)$ in Equation (43), since the image of $\mathrm{Vol}_{\partial_+^1 D^-}$ by the pushforward of measures by the positive endpoint map is $\frac{2^{n-1} d \mathrm{Leb}_{\widehat{K}}(z)}{h(\alpha_0)^{n-1} |Q_{\alpha_0}(z)|^{n-1}}$ by (the comment following) Lemma 23, and since the canonical map $G/G_{1,0} \rightarrow \overline{G}/\overline{G}_\infty = \Gamma/\Gamma_{D^+}$ is $\frac{2}{\iota_G}$ -to-1, the result follows. \square

We now assume that K is an imaginary quadratic number field. Let \mathfrak{m} be a (nonzero) fractional ideal of \mathcal{O}_K . Let $\alpha_0 \in \widehat{K}$ be a quadratic irrational over K . We consider the counting function $\overline{\psi}_{\mathfrak{m}}$ of nonequivalent \mathfrak{m} -primitive representations, by the norm form N_{α_0} , of elements of K (necessary integers in K if α_0 is an algebraic integer), defined on $[0, +\infty[$ by

$$s \mapsto \mathrm{Card} \ \mathrm{Stab}_{\Gamma_K N_{\alpha_0}} \backslash \{ (u, v) \in \mathfrak{m} \times \mathfrak{m} : \mathbf{n}(\mathfrak{m})^{-1} |\mathbf{n}(u - v \alpha_0)| \leq s, \ \mathcal{O}_K u + \mathcal{O}_K v = \mathfrak{m} \} .$$

Since for any $b \in \mathcal{O}_K$, we have $\mathbf{n}(b\mathfrak{m}) = |b|^2 \mathbf{n}(\mathfrak{m})$, the counting function $\overline{\psi}_{\mathfrak{m}}$ depends only on the ideal class of \mathfrak{m} .

Theorem 37 *There exists $\kappa > 0$ such that, as s tends to $+\infty$,*

$$\overline{\psi}_{\mathfrak{m}}(s) = \frac{2 \pi^2 h(\alpha_0)^2 \left| \ln \left| \frac{\mathrm{tr} \widehat{\alpha_0} + \sqrt{(\mathrm{tr} \widehat{\alpha_0})^2 - 4}}{2} \right| \right|}{m_{\Gamma_K}(\alpha_0) \iota_{\Gamma_K}(\alpha_0) |D_K| \zeta_K(2)} s^2 + O(s^{2-\kappa}) .$$

Furthermore, for every finite index subgroup G of $\Gamma_K = \mathrm{SL}_2(\mathcal{O}_K)$, with $\iota_G = 1$ if $-\mathrm{id} \in G$ and $\iota_G = 2$ otherwise, we have the following convergence of measures on $\mathbb{C} - \{\alpha_0, \alpha_0^\sigma\}$ as $s \rightarrow +\infty$:

$$\frac{[\Gamma_K : G] |D_K| \zeta_K(2) \iota_G \omega_K}{8 \pi^2 [(\Gamma_K)_\infty : G_\infty] s^2} \sum_{(u,v) \in G(0,1), |\mathbf{n}(u - \alpha_0 v)| \leq s} \Delta_{\frac{u}{v}} \xrightarrow{*} \frac{d \mathrm{Leb}_{\mathbb{C}}(z)}{|Q_{\alpha_0}(z)|^2} .$$

For smooth functions ψ with compact support on \mathbb{C} , there is an error term in the above equidistribution claim, of the form $O(s^{2-\kappa}\|\psi\|_\ell)$ where $\kappa > 0$ and $\|\psi\|_\ell$ is the Sobolev norm of ψ for some $\ell \in \mathbb{N}$.

We leave to the reader a version of this equidistribution claim where $(0, 1)$ is replaced by any fixed pair (x, y) of elements of \mathcal{O}_K which are not both zero (replacing $D^+ = \mathcal{H}_\infty$ by $D^+ = \gamma_\rho \mathcal{H}_\infty$ in its proof, with γ_ρ as in the proof of Theorem 27).

For instance, taking $G = \Gamma_K$, we have

$$\frac{|D_K| \zeta_K(2) \omega_K}{8 \pi^2 s^2} \sum_{(u,v) \in \mathcal{O}_K \times \mathcal{O}_K, \theta_K u + \theta_K v = \theta_K, |\mathbf{n}(u - \alpha_0 v)| \leq s} \Delta_{\frac{u}{v}} \xrightarrow{*} \frac{d \text{Leb}_{\mathbb{C}}(z)}{|Q_{\alpha_0}(z)|^2}.$$

Proof. We will prove a stronger counting claim. Let x, y in \mathcal{O}_K be elements which are not both zero, with $x = 1$ if $y = 0$ and the same strange convention that $\mathbf{n}(y) = 1$ if $y = 0$. Let $I_{x,y} = x\mathcal{O}_K + y\mathcal{O}_K$, and for every $s > 0$, let

$$\overline{\psi}_{G,x,y}(s) = \text{Card } \text{Stab}_G N_{\alpha_0} \setminus \{(u, v) \in G(x, y) : \mathbf{n}(I_{x,y})^{-1} |\mathbf{n}(u - v\alpha_0)| \leq s\}.$$

Let $\rho = xy^{-1} \in K \cup \{\infty\}$, $\gamma_\rho = \begin{pmatrix} \rho & -1 \\ 1 & 0 \end{pmatrix} \in \text{SL}_2(K)$ if $y \neq 0$ and $\gamma_\rho = \text{id}$ otherwise. Let $\mathcal{H}_\infty = \{(z, t) \in \mathbb{H}_{\mathbb{R}}^3 : t \geq 1\}$ and $\mathcal{H}_\rho = \gamma_\rho \mathcal{H}_\infty$. We will apply the counting claim of Corollary 9 with $n = 3$, Γ the arithmetic group \overline{G} , D^- the geodesic line with points at infinity $\alpha_0, \alpha_0^\sigma$ (whose stabiliser is exactly $\text{Stab}_{\Gamma_K} N_{\alpha_0} \cap G$, and acts with compact quotient on D^-) and D^+ the horoball \mathcal{H}_ρ (which is centered at a parabolic fixed point of \overline{G}).

For every $\gamma \in \text{SL}_2(\mathcal{O}_K)$, the distance ℓ_γ between D^- and $\gamma D^+ = \gamma \gamma_\rho \mathcal{H}_\infty$, when they are disjoint, is equal to the distance between \mathcal{H}_∞ and the geodesic line with points at infinity $\gamma_\rho^{-1} \gamma^{-1} \alpha_0, \gamma_\rho^{-1} \gamma^{-1} \alpha_0^\sigma$. Hence again by [PP4, Lem. 4.2], if $\gamma \gamma_\rho(1, 0) = (u', v')$, then $\ell_\gamma = \ln(h(\alpha_0) |\mathbf{n}(u' - v'\alpha_0)|)$. If $y \neq 0$, we have $\gamma \gamma_\rho(1, 0) = \frac{1}{y} \gamma(x, y)$, hence

$$\ell_\gamma = \ln\left(\frac{h(\alpha_0)}{\mathbf{n}(y)} |\mathbf{n}(u - v\alpha_0)|\right)$$

if $\gamma(x, y) = (u, v)$. By the above convention, this is also true if $y = 0$.

As in the proof of Theorem 10, we then have

$$\begin{aligned} \overline{\psi}_{G,x,y}(s) &= \text{Card } \{[\gamma] \in G_{\{\alpha_0, \alpha_0^\sigma\}} \setminus G/G(x, y) : \ell_\gamma \leq \ln \frac{h(\alpha_0) \mathbf{n}(I_{x,y}) s}{\mathbf{n}(y)}\} + O(1) \\ &= \frac{2}{\iota_G} \text{Card } \{[\gamma] \in \overline{G}_{\{\alpha_0, \alpha_0^\sigma\}} \setminus \overline{G}/\overline{G}(x, y) : \ell_\gamma \leq \ln \frac{h(\alpha_0) \mathbf{n}(I_{x,y}) s}{\mathbf{n}(y)}\} + O(1) \\ &= \frac{2}{\iota_G} [\overline{G}_{\mathcal{H}_\rho} : \overline{G}(x, y)] \mathcal{N}_{D^-, D^+} \left(\ln \frac{h(\alpha_0) \mathbf{n}(I_{x,y}) s}{\mathbf{n}(y)} \right) + O(1) \\ &= \frac{2 [\overline{G}_{\mathcal{H}_\rho} : \overline{G}(x, y)]}{\iota_G} \frac{\text{Vol}(\mathbb{S}^1) \text{Vol}(\Gamma_{D^-} \setminus D^-) \text{Vol}(\Gamma_{D^+} \setminus D^+)}{m_G(\alpha_0) \text{Vol}(\mathbb{S}^2) \text{Vol}(\Gamma \setminus \mathbb{H}_{\mathbb{R}}^3)} \times \\ &\quad \left(\frac{h(\alpha_0) \mathbf{n}(I_{x,y}) s}{\mathbf{n}(y)} \right)^2 (1 + O(s^{-\kappa})). \end{aligned} \tag{44}$$

By Equation (21), we have $\text{Vol}(\Gamma_{D^-} \setminus D^-) = \frac{m_G(\alpha_0) [(\overline{\Gamma_K})_{\alpha_0} : \overline{G}_{\alpha_0}]}{m_{\Gamma_K}(\alpha_0) \iota_G(\alpha_0)} \ell(\widehat{\alpha_0})$. By Equation (39), we have $\text{Vol}(\Gamma_{D^+} \setminus D^+) = [\overline{(\Gamma_K)}_{\mathcal{H}_\rho} : \overline{G}_{\mathcal{H}_\rho}] \frac{\sqrt{|D_K|} \mathbf{n}(y)^2}{2 \omega_K \mathbf{n}(I_{x,y})^2}$. By Equation (36), we have

$\text{Vol}(\Gamma \backslash \mathbb{H}_{\mathbb{R}}^3) = \frac{[\Gamma_K : G] |D_K|^{\frac{3}{2}} \zeta_K(2)}{4\pi^2 \iota_G}$. We have $[(\overline{\Gamma_K})_{\alpha_0} : \overline{G_{\alpha_0}}] = \frac{[(\Gamma_K)_{\alpha_0} : G_{\alpha_0}]}{\iota_G}$. By Equation (40), we have $[(\overline{\Gamma_K})_{\mathcal{H}_\rho} : \overline{G_{\mathcal{H}_\rho}}][\overline{G_{\mathcal{H}_\rho}} : \overline{G_{(x,y)}}] = \frac{\omega_K}{2} [(\Gamma_K)_{(x,y)} : G_{(x,y)}]$. Therefore Equation (44) gives the following result, interesting in itself,

$$\overline{\psi}_{G,(x,y)}(s) = \frac{2\pi^2 [(\Gamma_K)_{(x,y)} : G_{(x,y)}] [(\Gamma_K)_{\alpha_0} : G_{\alpha_0}] h(\alpha_0)^2 \left| \ln \left| \frac{\text{tr } \widehat{\alpha_0} + \sqrt{(\text{tr } \widehat{\alpha_0})^2 - 4}}{2} \right| \right|}{m_{\Gamma_K}(\alpha_0) \iota_G(\alpha_0) [\Gamma_K : G] |D_K| \zeta_K(2)} \times s^2(1 + O(s^{-\kappa})).$$

Since any nonzero ideal \mathfrak{m} in \mathcal{O}_K may be written $\mathfrak{m} = x_{\mathfrak{m}}\mathcal{O}_K + y_{\mathfrak{m}}\mathcal{O}_K$ for some $x_{\mathfrak{m}}, y_{\mathfrak{m}}$ not both zero in \mathcal{O}_K and $\overline{\psi}_{\mathfrak{m}} = \overline{\psi}_{\Gamma_K, x_{\mathfrak{m}}, y_{\mathfrak{m}}}$, the first claim of Theorem 37 follows.

The proof of the equidistribution claim is similar to the one in Theorem 36 (which was written with greater care than necessary for this purpose). \square

References

- [Ahl] L. V. Ahlfors. *Möbius transformations in several dimensions*. Ordway Prof. Lect. Math., Univ. Minnesota School Math., 1981.
- [Bab] M. Babillot. *On the mixing property for hyperbolic systems*. Israel J. Math. **129** (2002) 61–76.
- [Bea] A. F. Beardon. *The geometry of discrete groups*. Grad. Texts Math. **91**, Springer-Verlag, 1983.
- [BQ] Y. Benoist and J.-F. Quint. *Random walks on finite volume homogeneous spaces*. Invent. Math. **187** (2012) 37–59.
- [Coh1] H. Cohn. *A classical invitation to algebraic numbers and class fields*. Springer Verlag, 1978.
- [Coh2] H. Cohn. *A second course in number theory*. Wiley, 1962, reprinted as *Advanced number theory*, Dover, 1980.
- [Cos] S. Cosentino. *Equidistribution of parabolic fixed points in the limit set of Kleinian groups*. Erg. Theo. Dyn. Syst. **19** (1999) 1437–1484.
- [DaOP] F. Dal’Bo, J.-P. Otal, and M. Peigné. *Séries de Poincaré des groupes géométriquement finis*. Israel J. Math. **118** (2000) 109–124.
- [DeR] F. Delgove and N. Retailleau. *Sur la classification des hexagones hyperboliques à angles droit en dimension 5*. Mémoire de Licence, Univ. Paris-Sud (Orsay), 2013.
- [Deu] M. Deuring. *Algebren*. 2nd. ed., Erg. Math. Grenz. **41**, Springer Verlag, 1968.
- [EGM] J. Elstrodt, F. Grunewald, and J. Mennicke. *Groups acting on hyperbolic space: Harmonic analysis and number theory*. Springer Mono. Math., Springer Verlag, 1998.
- [Fen] W. Fenchel. *Elementary geometry in hyperbolic space*. de Gruyter Stud. Math. **11**, de Gruyter, 1989.
- [GP] A. Gorodnik and F. Paulin. *Counting orbits of integral points in families of affine homogeneous varieties and diagonal flows*. Prépublication [arXiv:1110.5773].
- [Gro] W. Grotz. *Mittelwert der Eulerschen φ -Funktion und des Quadrates der Dirichletschen Teilerfunktion in algebraischen Zahlkörpern*. Monatsh. Math. **88** (1979) 219–228.

- [Har] G. Harcos. Equidistribution on the modular surface and L -functions. In "Homogeneous flows, moduli spaces and arithmetic", M. Einsiedler et al eds., Clay Math. Proc. **10**, Amer. Math. Soc. 2010, 377–387.
- [HaW] G. H. Hardy and E. M. Wright. *An introduction to the theory of numbers*. Oxford Univ. Press, sixth ed., 2008.
- [Hux] M. N. Huxley. *Exponential sums and lattice points III*. Proc. London Math. Soc. **87** (2003) 591–609.
- [Kel] R. Kellerhals. *Quaternions and some global properties of hyperbolic 5-manifolds*. Canad. J. Math. **55** (2003) 1080–1099.
- [Kim] I. Kim. *Counting, mixing and equidistribution of horospheres in geometrically finite rank one locally symmetric manifolds*. Preprint [arXiv:1103.5003].
- [KO] V. Krafft and D. Osenberg. *Eisensteinreihen für einige arithmetisch definierte Untergruppen von $SL_2(\mathbb{H})$* . Math. Z. **204** (1990) 425–449.
- [Lan] S. Lang. *Algebraic number theory*. Grad Texts Math. 2nd ed., Springer Verlag, 1994.
- [MR] C. Maclachlan and A. Reid. *Parametrizing Fuchsian subgroups of the Bianchi groups*. Canad. J. Math. **43** (1991) 158–181.
- [Nag] T. Nagell. *On the number of representations of an A -number in an algebraic field*. Ark. Mat. **4** (1962) 467–478.
- [Odo] R. W. K. Odoni. *Representations of algebraic integers by binary quadratic forms and norm forms from full modules of extension fields*. J. Numb. Theo. **10** (1978) 324–333.
- [Oh] H. Oh. *Orbital counting via mixing and unipotent flows*. In "Homogeneous flows, moduli spaces and arithmetic", M. Einsiedler et al eds., Clay Math. Proc. **10**, Amer. Math. Soc. 2010, 339–375.
- [OhS1] H. Oh and N. Shah. *Equidistribution and counting for orbits of geometrically finite hyperbolic groups*. J. Amer. Math. Soc. **26** (2013) 511–562.
- [OhS2] H. Oh and N. Shah. *Counting visible circles on the sphere and Kleinian groups*. Preprint [arXiv:1004.2129], to appear in "Geometry, Topology and Dynamics in Negative Curvature" (ICM 2010 satellite conference, Bangalore), C. S. Aravinda, T. Farrell, J.-F. Lafont eds, London Math. Soc. Lect. Notes.
- [OtP] J.-P. Otal and M. Peigné. *Principe variationnel et groupes kleinien*. Duke Math. J. **125** (2004) 15–44.
- [PaS] J. Parker and I. Short. *Conjugacy classification of quaternionic Möbius transformations*. Comput. Meth. Funct. Theo. **9** (2009) 13–25.
- [PP1] J. Parkkonen and F. Paulin. *Prescribing the behaviour of geodesics in negative curvature*. Geom. & Topo. **14** (2010) 277–392.
- [PP2] J. Parkkonen and F. Paulin. *Spiraling spectra of geodesic lines in negatively curved manifolds*. Math. Z. **268** (2011) 101–142.
- [PP3] J. Parkkonen and F. Paulin. *On the representations of integers by indefinite binary Hermitian forms*. Bull. London Math. Soc. **43** (2011) 1048–1058.
- [PP4] J. Parkkonen and F. Paulin. *Équidistribution, comptage et approximation par irrationnels quadratiques*. J. Mod. Dyn. **6** (2012) 1–40.
- [PP5] J. Parkkonen and F. Paulin. *On the arithmetic and geometry of binary Hamiltonian forms*. Appendix by Vincent Emery. Algebra & Number Theory **7** (2013) 75–115.

- [PP6] J. Parkkonen and F. Paulin. *Skinning measure in negative curvature and equidistribution of equidistant submanifolds*. Preprint [arXiv:1202.6398], to appear in Ergodic Theory and Dynamical Systems.
- [PP7] J. Parkkonen and F. Paulin. *Counting arcs in negative curvature*. Preprint [arXiv:1203.0175], to appear in "Geometry, Topology and Dynamics in Negative Curvature" (ICM 2010 satellite conference, Bangalore), C. S. Aravinda, T. Farrell, J.-F. Lafont eds, London Math. Soc. Lect. Notes.
- [PP8] J. Parkkonen and F. Paulin. *Counting common perpendicular arcs in negative curvature*. Preprint [arXiv:1305.1332].
- [PP9] J. Parkkonen and F. Paulin. *Counting and equidistribution in the Heisenberg group*. In preparation.
- [Pol] M. Pollicott. *The Schottky-Klein prime function and counting functions for Fenchel double crosses*. Preprint 2011.
- [Rob] T. Roblin. *Ergodicité et équidistribution en courbure négative*. Mémoire Soc. Math. France, **95** (2003).
- [Sar] P. Sarnak. *Reciprocal geodesics*. Clay Math. Proc. **7** (2007) 217–237.
- [Ser] J.-P. Serre. *Lectures on $N_X(p)$* . Chapman & Hall/CRC Research Notes Math. **11**, CRC Press, 2012.
- [Shi] G. Shimura. *Introduction to the arithmetic theory of automorphic functions*. Princeton Univ. Press, 1971.
- [Vig] M. F. Vignéras. *Arithmétique des algèbres de quaternions*. Lect. Notes in Math. **800**, Springer Verlag, 1980.
- [Wal] A. Walfisz. *Weylsche Exponentialsummen in der neueren Zahlentheorie*. VEB Deutscher Verlag der Wissenschaften, 1963.

Department of Mathematics and Statistics, P.O. Box 35
 40014 University of Jyväskylä, FINLAND.
e-mail: jouni.t.parkkonen@jyu.fi

Département de mathématique, UMR 8628 CNRS, Bât. 425
 Université Paris-Sud, 91405 ORSAY Cedex, FRANCE
e-mail: frederic.paulin@math.u-psud.fr