# HIGHER HAMMING WEIGHTS FOR LOCALLY RECOVERABLE CODES ON ALGEBRAIC CURVES

#### EDOARDO BALLICO AND CHIARA MARCOLLA

ABSTRACT. We study the locally recoverable codes on algebraic curves. In the first part of this article, we provide a bound of generalized Hamming weight of these codes. Whereas in the second part, we propose a new family of algebraic geometric LRC codes, that are LRC codes from Norm-Trace curve. Finally, using some properties of Hermitian codes, we improve the bounds of distance proposed in [1] for some Hermitian LRC codes.

#### 1. Introduction

The v-th generalized Hamming weight  $d_v(C)$  of a linear code C is the minimum support size of v-dimensional subcodes of C. The sequence  $d_1(C), \ldots, d_k(C)$  of generalized Hamming weights was introduced by Wei [37] to characterize the performance of a linear code on the wire-tap channel of type II. Later, the GHWs of linear codes have been used in many other applications regarding the communications, as for bounding the covering radius of linear codes [15], in network coding [26], in the context of list decoding [7, 9], and finally for secure secret sharing [18]. Moreover, in [2] the authors show in which way an arbitrary linear code gives rise to a secret sharing scheme, in [16, 17] the connection between the trellis or state complexity of a code and its GHWs is found and in [4] the author proves the equivalence to the dimension/length profile of a code and its generalized Hamming weight. For these reasons, the GHWs (and their extended version, the relative generalized Hamming weights [21, 19]) play a central role in coding theory. In particular, generalized and relative generalized Hamming weights are studied for Reed-Muller codes [10, 23] and for codes constructed by using an algebraic curve [6] as Goppa codes [24, 38], Hermitian codes [12, 25] and Castle codes [27].

In this paper, we provide a bound on the generalized Hamming weight of locally recoverable codes on the algebraic curves proposed in [1]. Moreover, we introduce a new family of algebraic geometric LRC codes and improve the bounds on the distance for some Hermitian LRC codes.

Locally recoverable codes were introduced in [8] and they have been significantly studied because of their applications in distributed and cloud storage systems [3, 13, 32, 34, 35]. We recall that a code  $C \in (\mathbb{F}_q)^n$  has locality r if every symbol of a codeword c can be recovered from a subset of r other symbols of c.

In other words, we consider a finite field  $K = \mathbb{F}_q$ , where q is a power of a prime, and an [n,k] code C over the field K, where  $k = log_q(|C|)$ . For each  $i \in \{1,\ldots,n\}$  and each  $a \in K$  set  $C(i,a) = \{c \in C \mid c_i = a\}$ . For each  $I \subseteq \{1,\ldots,n\}$  and each  $S \subseteq C$  let  $S_I$  be the restriction of S to the coordinates in I.

**Definition 1.1.** Let C be an [n, k] code over the field K, where  $k = log_q(|C|)$ . Then C is said to have **all-symbol locality r** if for each  $a \in \mathbb{F}_q$  and each  $i \in \{1, \ldots, n\}$  there is  $I_i \subset \{1, \ldots, n\} \setminus \{i\}$  with  $|I_i| \leq r$ , such that for  $C_{I_i}(i, a) \cap C_{I_i}(i, a') = \emptyset$  for all  $a \neq a'$ . We use the notation (n, k, r) to refer to the parameters of this code.

Note that if we receive a codeword c correct except for an erasure at i, we can recover the codeword by looking at its coordinates in  $I_i$ . For this reason,  $I_i$  is called a recovering set for the symbol  $c_i$ .

Let C be an (n, k, r) code, then the distance of this code has to verify the bound proved in [28, 8] that is  $d \leq n - k - \lceil k/r \rceil + 2$ . The codes that achieve this bound with equality are called *optimal* LRC codes [32, 34, 35]. Note that when r = k, we obtain the Singleton bound, therefore optimal LRC codes with r = k are MDS codes. Layout of the paper. This paper is divided as follows. In Section 2 we recall the notions of algebraic geometric codes and the definition of algebraic geometric locally recoverable codes introduced in [1]. In Section 3 we provide a bound on the generalized Hamming weights of the latter codes. In Section 4 we propose a new family of algebraic geometric LRC codes, which are LRC codes from the Norm-Trace curve. Finally, in Section 5 we improve the bounds on the distance proposed in [1] for some Hermitian LRC codes, using some properties of the Hermitian codes.

### 2. Preliminary notions

2.1. Algebraic geometric codes. Let  $K = \mathbb{F}_q$  be a finite field, where q is a power of a prime. Let  $\mathcal{X}$  be a smooth projective absolutely irreducible nonsingular curve over K. We denote by  $K(\mathcal{X})$  the rational functions field on  $\mathcal{X}$ . Let D be a divisor on the curve  $\mathcal{X}$ . We recall that the *Riemann-Roch space* associated to D is a vector space  $\mathcal{L}(D)$  over K defined as

$$\mathcal{L}(D) = \{ f \in K(\mathcal{X}) \mid (f) + D \ge 0 \} \cup \{ 0 \}.$$

where we denote by (f) the divisor of f.

Assume that  $P_1, \ldots, P_n$  are rational points on  $\mathcal{X}$  and D is a divisor such that  $D = P_1 + \ldots + P_n$ . Let G be some other divisor such that  $supp(D) \cap supp(G) = \emptyset$ . Then we can define the algebraic geometric code as follows:

**Definition 2.1.** The algebraic geometric code (or AG code) C(D, G) associated with the divisors D and G is defined as

$$C(D,G) = \{ (f(P_1), \dots, f(P_n)) \mid f \in \mathcal{L}(G) \} \subset K^n.$$

The dual  $C^{\perp}(D,G)$  of C(D,G) is an algebraic geometric code.

In other words an algebraic geometric code is the image of the evaluation map  $\operatorname{Im}(ev_D) = C(D, G)$ , where the evaluation map  $ev_D : \mathcal{L}(G) \to K^n$  is given by

$$ev_D(f) = (f(P_1), \dots, f(P_n)) \in K^n.$$

Note that if  $D = P_1 + \ldots + P_n$  and we denote by  $\mathcal{P} = \{P_1, \ldots, P_n\}$  we can also indicate  $ev_D$  as  $ev_{\mathcal{P}}$ .

2.2. Algebraic geometric locally recoverable codes. In this section we consider the construction of algebraic geometric locally recoverable codes of [1].

Let  $\mathcal{X}$  and  $\mathcal{Y}$  be smooth projective absolutely irreducible curves over K. Let  $g: \mathcal{X} \to \mathcal{Y}$  be a rational separable map of curves of degree r+1. Since g is separable, then there exists a function  $x \in K(\mathcal{X})$  such that  $K(\mathcal{X}) = K(\mathcal{Y})(x)$  and that x satisfies the equation  $x^{r+1} + b_r x^r + \ldots + b_0 = 0$ , where  $b_i \in K(\mathcal{Y})$ . The function x can be considered as a map  $x: \mathcal{X} \to \mathbb{P}_K$ . Let  $h = \deg(x)$  be the degree of x. We consider a subset  $S = \{P_1, \ldots, P_s\} \subset \mathcal{Y}(K)$  of  $\mathbb{F}_q$ -rational points of  $\mathcal{Y}$ , a divisor  $Q_{\infty}$  such that  $supp(Q_{\infty}) \cap supp(S) = \emptyset$  and a positive divisor  $D = tQ_{\infty}$ . We denote by

$$\mathcal{A} = g^{-1}(S) = \{P_{ij}, \text{ where } i = 0, \dots, r, j = 1, \dots, s\} \subset \mathcal{X}(K),$$

where  $g(P_{ij}) = P_i$  for all i, j and assume that  $b_i$  are functions in  $\mathcal{L}(n_i Q_{\infty})$  for some natural numbers  $n_i$  with  $i = 1, \ldots, r$ .

Let  $\{f_1, \ldots, f_m\}$  be a basis of the Riemann-Roch space  $\mathcal{L}(D)$ . By the Riemann-Roch Theorem we have that  $m \geq \deg(D) + 1 - g_{\mathcal{Y}}$ , where  $g_{\mathcal{Y}}$  is the genus of  $\mathcal{Y}$ .

From now on, we assume that  $m = \deg(D) + 1 - g_{\mathcal{V}}$ , where  $\deg(D) = t\ell$ , and we consider the K-subspace V of  $K(\mathcal{X})$  of dimension rm generated by

$$\mathcal{B} = \{ f_j x^i, \ i = 0, \dots, r - 1, \ j = 1, \dots, m \}.$$

We consider the evaluation map  $ev_{\mathcal{A}}: V \to K^{(r+1)s}$ . Then we have the following theorem.

**Theorem 2.2.** The linear space  $C(D,g) = \operatorname{Span}_{K^{(r+1)s}} \langle ev_{\mathcal{A}}(\mathcal{B}) \rangle$  is an (n,k,r) algebraic geometric LRC code with parameters

$$n = (r+1)s$$
  
 $k = rm \ge r(t\ell + 1 - g_{\mathcal{Y}})$   
 $d \ge n - t\ell(r+1) - (r-1)h$ .

*Proof.* See Theorem 3.1 of [1].

The AG LRC codes have an additional property. They are LRC codes (n, k, r) with  $(r+1) \mid n$  and  $r \mid k$ . The set  $\{1, \ldots, n\}$  can be divided into n/(r+1) disjoint subsets  $U_j$  for  $1 \leq j \leq s$  with the same cardinality r+1. For each i the set  $I_i \subseteq \{1, \ldots, n\} \setminus \{i\}$  is the complement of i in the element of the partition  $U_j$  containing j, i.e. for all  $i, j \in \{1, \ldots, n\}$  either  $I_i = I_j$  or  $I_i \cap I_j = \emptyset$ .

Moreover, they have also the following nice property. Fix  $w \in (K)^n$  and denote by  $w_{U_j} = \{w_i, \text{ for any } i \in U_j\}$ . Suppose we receive all the symbols in  $U_j$ . There is a simple linear parity test on the r+1 symbols of  $U_j$  such that if this parity check fails we know that at least one of the symbols in  $U_j$  is wrong. If we are guaranteed (or we assume) that at most one of the symbols in  $U_j$  is wrong and the parity check is OK, then all the symbols in  $U_j$  are correct. Moreover we can recover an erased symbol  $w_i$ , with  $i \in U_j$  using a polynomial interpolation through the points of the recovering set  $w_{U_j}$ .

#### 3. Generalized Hamming weights of AG LRC codes

Let K be a field and let  $\mathcal{X}$  be a smooth and geometrically connected curve of genus  $g \geq 2$  defined over the field K. We also assume  $\mathcal{X}(K) \neq \emptyset$ . We recall the following definitions:

**Definition 3.1** ([29], [30]). The K-gonality  $\gamma_K(\mathcal{X})$  of  $\mathcal{X}$  over a field K is the smallest possible degree of a dominant rational map  $\mathcal{X} \to \mathbb{P}^1_K$ . For any field extension L of K, we define also the L-gonality  $\gamma_L(\mathcal{X})$  of  $\mathcal{X}$  as the gonality of the base extension  $\mathcal{X}_L = \mathcal{X} \times_K L$ . It is an invariant of the function field  $L(\mathcal{X})$  of  $\mathcal{X}_L$ .

Moreover, for each integer i > 0, the *i-th gonality*  $\gamma_{i,L}(\mathcal{X})$  of  $\mathcal{X}$  is the minimal degree z such that there is  $R \in \operatorname{Pic}^z(\mathcal{X})(L)$  with  $h^0(R) \geq i + 1$ . The sequence  $\gamma_{i,\overline{K}}(\mathcal{X})$  is the usual gonality sequence [20]. Moreover, the integer  $\gamma_{1,K}(\mathcal{X}) = \gamma_K(\mathcal{X})$  is the K-gonality of  $\mathcal{X}$ .

Let  $K = \mathbb{F}_q$  a finite field with q elements. Let  $C \subset K^n$  be a linear [n, k] code over K. We recall that the *support* of C is defined as follows

$$supp(C) = \{i \mid c_i \neq 0 \text{ for some } c \in C\}.$$

So  $\sharp supp(C)$  is the number of nonzero columns in a generator matrix for C. Moreover, for any  $1 \leq v \leq k$ , the v-th generalized Hamming weight of C [14, §7.10], [36, §1.1] is defined by

$$d_v(C) = \min\{\sharp supp(\mathcal{D}) \mid \mathcal{D} \text{ is a linear subcode of } C \text{ with } dim(\mathcal{D}) = v\}.$$

In other words, for any integer  $1 \leq v \leq k$ ,  $d_v(C)$  is the v-th minimum support weights, i.e. the minimal integer t such that there are an [n, v] subcode  $\mathcal{D}$  of C and a subset  $S \subset \{1, \ldots, n\}$  such that  $\sharp(S) = t$  and each codeword of  $\mathcal{D}$  has zero coordinates outside S. The sequence  $d_1(C), \ldots, d_k(C)$  of generalized Hamming weights (also called weight hierarchy of C) is strictly increasing (see Theorem 7.10.1 of [14]). Note that  $d_1(C)$  is the minimum distance of the code C.

Let us consider  $\mathcal{X}$  and  $\mathcal{Y}$  smooth projective absolutely irreducible curves over K and let  $g: \mathcal{X} \to \mathcal{Y}$  be a rational separable map of curves of degree r+1. Moreover we take  $r, t, Q_{\infty}, f_1, \ldots, f_m$  and  $\mathcal{A} = g^{-1}(S)$  defined as Section 2.2. So we can construct an (n, k, r) algebraic geometric LRC code C as in Theorem 2.2. For this code we have the following:

**Theorem 3.2.** Let C be an (n, k, r) algebraic geometric LRC code as in Theorem 2.2. For every integer  $v \geq 2$  we have that

$$d_v(C) \ge n - t\ell(r+1) - (r-1)h + \gamma_{v-1,K}(\mathcal{X}).$$

*Proof.* Take a v-dimensional linear subspace  $\mathcal{D}$  of C and call

$$E \subseteq \{P_{ij} \mid i = 0, \dots, r, j = 1, \dots, s\},\$$

the set of common zeros of all elements of  $\mathcal{D}$ . Since  $n - d_v(C) = \sharp(E)$ , we have to prove that  $t\ell(r+1) + (r-1)h - \sharp(E) \geq \gamma_{v-1,K}(X)$ . Fix  $u \in \mathcal{D} \setminus \{0\}$  and let  $F_u$  denote the zeros of u. Note that  $F_u$  is contained in the set  $\{P_{ij} \mid i = 0, \dots, j = 1, \dots, s\}$  by the definition of the code C. We have  $F_u \supseteq E$ . By the definition of the integers  $t, \ell$  and  $h := \deg(x)$ , we have  $\sharp(F_u) \leq t\ell(r+1) + (r-1)h$ . The divisors  $F_u - E$ ,  $u \in \mathcal{D} \setminus \{0\}$  form a family of linearly equivalent non-negative divisors, each of them defined over K. Since  $\dim(\mathcal{D}) = v$ , the definition of  $\gamma_{v-1,\overline{K}}(\mathcal{X})$  gives  $\sharp(F_u) - \sharp(E) \geq \gamma_{v-1,K}(\mathcal{X})$ . This inequality for a single  $u \in \mathcal{D} \setminus \{0\}$  proves the theorem.

See Remark 4.4 for an application of Theorem 3.2.

### 4. LRC codes from Norm-Trace Curve

In this section we propose a new family of Algebraic Geometric LRC codes, that is, a LRC codes from the Norm-Trace curve. Moreover, we compute the  $\mathbb{F}_{q^u}$ -gonality of the Norm-Trace curve.

Let  $K = \mathbb{F}_{q^u}$  be a finite field, where q is a power of a prime. We consider the *norm*  $N_{\mathbb{F}_q}^{\mathbb{F}_{q^u}}$  and the *trace*  $\mathrm{Tr}_{\mathbb{F}_q}^{\mathbb{F}_{q^u}}$ , two functions from  $\mathbb{F}_{q^u}$  to  $\mathbb{F}_q$  defined as

$$N_{\mathbb{F}_q}^{\mathbb{F}_{q^u}}(x) = x^{1+q+\dots+q^{u-1}} \text{ and } Tr_{\mathbb{F}_q}^{\mathbb{F}_{q^u}}(x) = x + x^q + \dots + x^{q^{u-1}}.$$

The Norm-Trace curve  $\chi$  is the curve defined over K by the following affine equation

$$N_{\mathbb{F}_q}^{\mathbb{F}_{q^u}}(x) = Tr_{\mathbb{F}_q}^{\mathbb{F}_{q^u}}(y),$$

that is,

(4.1) 
$$x^{(q^{u}-1)/(q-1)} = y^{q^{u-1}} + y^{q^{u-2}} + \dots + y \text{ where } x, y \in K$$

The Norm-Trace curve  $\chi$  has exactly  $n=q^{2u-1}$  K-rational affine points (see Appendix A of [5]), that we denote by  $\mathcal{P}_{\chi}=\{P_1,\ldots,P_n\}$ . The genus of  $\chi$  is  $g=\frac{1}{2}(q^{u-1}-1)(\frac{q^u-1}{q-1}-1)$ . Note that if we consider u=2, we obtain the Hermitian curve.

Starting from the Norm-Trace curve, we have two different ways to construct Norm-Trace LRC codes.

**Projection on x**. We have to construct a  $q^u$ -ary (n, k, r) LRC codes. We consider the natural projection g(x, y) = x. Then the degree of g is  $q^{u-1} = r + 1$  and the degree of y is  $h = 1 + q + \cdots + q^{u-1}$ .

To construct the codes we consider  $S = \mathbb{F}_{q^u}$  and  $D = tQ_{\infty}$  for some  $t \geq 1$ . Then, using a construction of Theorem 2.2 we find the parameters for these Norm-Trace LRC codes.

**Proposition 4.1.** A family of Norm-Trace LRC codes has the following parameters:

$$n = q^{2u-1}, \quad k = mr = (t+1)(q^{u-1}-1)$$

and

$$d \ge n - tq^{u-1} - (q^{u-1} - 1)(1 + q + \dots + q^{u-1}).$$

**Projection on y.** We have to construct a  $q^u$ -ary (n, k, r) LRC codes. We consider the other natural projection g'(x, y) = y. Then  $\deg(g') = 1 + q + \cdots + q^{u-1} = r + 1$ . In this case we take  $S = \mathbb{F}_{q^u} \setminus M$ , where

$$M = \{ a \in \mathbb{F}_{q^u} \mid a^{q^{u-1}} + a^{q^{u-2}} + \dots + a = 0 \},$$

so  $r = q + \cdots + q^{u-1}$  and  $h = \deg(x) = q^{u-1}$ . Then, using Theorem 2.2 we have the following

**Proposition 4.2.** A family of Norm-Trace LRC codes has the following parameters:

$$n = q^{2u-1} - q^{u-1}, \quad k = mr = (t+1)(q + \dots + q^{u-1})$$

and

$$d \ge n - tq^{u-1} - (q + \dots + q^{u-1}) - q^{u-1}(q^{u-1} + \dots + q - 1).$$

For the Norm-Trace curve  $\chi$  we are able to find the K-gonality of  $\chi$ .

**Lemma 4.3.** Let  $\chi$  be a Norm-Trace curve defined over  $\mathbb{F}_{q^u}$ , where  $u \geq 2$ . We have  $\gamma_{1,\mathbb{F}_{q^u}}(\chi) = q^{u-1}$ .

Proof. The linear projection onto the x axis has degree  $q^{u-1}$  and it is defined over  $\mathbb{F}_q$  and hence over  $\mathbb{F}_{q^u}$ . Thus  $\gamma_{1,\mathbb{F}_{q^u}}(\chi) \leq q^{u-1}$ . Denote by  $z = \gamma_{1,\mathbb{F}_{q^u}}(\chi)$  and assume that  $z \leq q^{u-1} - 1$ . By the definition of K-gonality, there is a non-constant morphism  $w: \chi \to \mathbb{P}^1$  with  $\deg(w) = z$  and defined over  $\mathbb{F}_{q^u}$ . Since  $w(\chi(\mathbb{F}_{q^u})) \subseteq \mathbb{P}^1(\mathbb{F}_{q^u})$ , we get  $\sharp(\chi(\mathbb{F}_{q^u})) \leq z(q^u+1) \leq (q^{u-1}-1)(q^u+1)$ , that is a contradiction.

**Remark 4.4.** By Lemma 4.3, we can apply Theorem 3.2 to the Norm–Trace curve. In fact, we can consider the gonality sequence over K of  $\chi$  to get a lower bound on the second generalized Hamming weight of the two families of Norm–Trace LRC codes:

• Let  $t \ge 1$  and let C be a  $(q^{2u-1}, (t+1)(q^{u-1}-1), q^{u-1}-1)$  Norm-Trace LRC code. Then we have

$$d_2(C) \ge q^{2u-1} + q^{u-1} - tq^{u-1} - (q^{u-1} - 1)(1 + q + \dots + q^{u-1}).$$

• Let  $t \ge 1$  and let C be a Norm-Trace LRC code with parameters  $(q^{2u-1}-q^{u-1}, (t+1)(q+\cdots+q^{u-1}), q+\cdots+q^{u-1})$ . Then we have

$$d_2(C) \ge q^{2u-1} - (t-1)q^{u-1} - (1+q^{u-1})(q+\dots+q^{u-1}).$$

## 5. HERMITIAN LRC CODES

In this section we improve the bound on the distance of Hermitian LRC codes proposed in [1] using some properties of *Hermitian codes* which are a special case of algebraic geometric codes.

5.1. **Hermitian codes.** Let us consider  $K = \mathbb{F}_{q^2}$  a finite field with  $q^2$  elements. The Hermitian curve  $\mathcal{H}$  is defined over K by the affine equation

(5.1) 
$$x^{q+1} = y^q + y \text{ where } x, y \in K.$$

This curve has genus  $g = \frac{q(q-1)}{2}$  and has  $q^3 + 1$  points of degree one, namely a pole  $Q_{\infty}$  and  $n = q^3$  rational affine points, denoted by  $\mathcal{P}_{\mathcal{H}} = \{P_1, \dots, P_n\}$  [31].

**Definition 5.1.** Let  $m \in \mathbb{N}$  such that  $0 \le m \le q^3 + q^2 - q - 2$ . Then the **Hermitian** code C(m,q) is the code  $C(D,mQ_{\infty})$  where

$$D = \sum_{\alpha^{q+1} = \beta^q + \beta} P_{\alpha,\beta}$$

is the sum of all places of degree one (except  $Q_{\infty}$ , that is a point at infinity) of the Hermitian function field  $K(\mathcal{H})$ .

By Lemma 6.4.4. of [33] we have that

$$\mathcal{B}_{m,q} = \{x^i y^j \mid qi + (q+1)j \le m, \ 0 \le i \le q^2 - 1, \ 0 \le j \le q - 1\},\$$

forms a basis of  $\mathcal{L}(mQ_{\infty})$ . For this reason, the Hermitian code C(m,q) could be seen as  $\operatorname{Span}_{\mathbb{F}_{q^2}}\langle ev_{\mathcal{P}_{\mathcal{H}}}(\mathcal{B}_{m,q})\rangle$ . Moreover, the dual of C(m,q) denoted by  $C(m_{\perp},q)=C^{\perp}(m,q)$  is again an Hermitian code and it is well known (Proposition 8.3.2 of [33]) that the degree m of the divisor has the following relation with respect to  $m_{\perp}$ :

$$(5.2) m_{\perp} = n + 2g - 2 - m.$$

The Hermitian codes can be divided in four phases [11], any of them having specific explicit formulas linking their dimension and their distance [22]. In particular we are interested in the first and the last phase of Hermitian codes, which are:

**I Phase:**  $0 \le m_{\perp} \le q^2 - 2$ : Then we have  $m_{\perp} = aq + b$  where  $0 \le b \le a \le q - 1$  and  $b \ne q - 1$ . In this case, the distance is

(5.3) 
$$\begin{cases} d = a+1 & if \quad a > b \\ d = a+2 & if \quad a = b. \end{cases}$$

**IV Phase:**  $n-1 \le m_{\perp} \le n+2g-2$ : In this case  $m_{\perp} = n+2g-2-aq-b$  where a,b are integers such that  $0 \le b \le a \le q-2$  and the distance is

$$(5.4) d = n - aq - b.$$

5.2. Bound on distance of Hermitian LRC codes. Let  $K = \mathbb{F}_{q^2}$  be a finite field, where q is a power of a prime. Let  $\mathcal{X} = \mathcal{H}$  be the Hermitian curve with affine equation as in (5.1). We recall that this curve has  $q^3 \mathbb{F}_{q^2}$ -rational affine points plus one at infinity, that we denoted by  $Q_{\infty}$ .

We consider two of the three constructions of Hermitian LRC codes proposed in [1] and we improve the bound on distance of Hermitian LRC codes using properties of Hermitian codes. In particular, if we find an Hermitian code  $C(m,q) = C_{Her}$  such that  $C_{LRC} \subset C_{Her}$ , then we have  $d_{LRC} \geq d_{Her}$ .

**Projection on x**. By Proposition 4 of [1], we have a family of (n, k, r) Hermitian LRC codes with r = q - 1, length  $n = q^3$ , dimension k = (t - 1)(q - 1) and distance  $d \ge n - tq - (q - 2)(q + 1)$ . Moreover, for these codes, S = K,  $D = tQ_{\infty}$  for some  $1 \le t \le q^2 - 1$  and the basis for the vector space V is

(5.5) 
$$\mathcal{B} = \{x^j y^i \mid j = 0, \dots, t, \ i = 0, \dots, q - 2\}.$$

Using the Hermitian codes, we improve the bound on the distance for any integer t, such that  $q^2 - q + 1 \le t \le q^2 - 1$ .

To find an Hermitian code  $C(m,q) = C_{Her}$  such that  $C_{LRC} \subset C_{Her}$ , we have to compute the set  $\mathcal{B}_{m,q}$ , that is, we have to find m. After that, to compute the distance of C(m,q) we use (5.3) and (5.4).

We consider the first Hermitian phase:  $0 \le m_{\perp} \le q^2 - 2$ , that is,  $q^2 - q + 1 \le t \le q^2 - 1$ . For this phase  $m_{\perp} = aq + b$ , where  $0 \le b \le a \le q - 1$  and the distance of the Hermitian code is either d = a + 1 if a > b or d = a + 2 if a = b. By (5.5), m must be equal to m = qt + (q+1)(q-2) and by (5.2) we have that  $m_{\perp} = n + 2g - 2 - m = q(q^2 - t)$ . So b = 0 and  $a = q^2 - t$  and the distance of the Hermitian code is  $d_{Her} = a + 1 = q^2 - t + 1$ , since a > b. This implies that

(5.6) 
$$d_{LRC} \ge q^2 - t + 1$$
, for any  $t \ge q^2 - q + 1$ .

Note that (5.6) improves the bound on the distance proposed in Proposition 4 of [1] since

$$q^2 - t + 1 > q^3 - tq - (q - 2)(q + 1) \iff t(q - 1) > q(q - 1)^2 + 1 \iff t > q^2 - q$$

We just proved the following:

**Proposition 5.2.** Let  $q^2 - q + 1 \le t \le q^2 - 1$ . It is possible to construct a family of (n, k, r) Hermitian LRC codes  $\{C_t\}_{q^2-q+1 \le t \le q^2-1}$  with the following parameters:

$$n = q^3$$
,  $k = (t - 1)(q - 1)$ ,  $r = q - 1$  and  $d \ge q^2 - t + 1$ .

**Two recovering sets**. In [1] the authors propose an Hermitian code with two recovering sets of size  $r_1 = q - 1$  and  $r_2 = q$ , denoted by LRC(2). They consider

$$L = Span\{x^iy^j, i = 0, \dots, q-2, j = 0, \dots, q-1\}$$

and a linear code C obtained by evaluating the functions in L at the points of  $B = g^{-1}(\mathbb{F}_{q^2}\backslash M)$ , where g(x,y) = x and  $M = \{a \in \mathbb{F}_q \mid a^q + a = 0\}$ . So  $|B| = q^3 - q$ . By Proposition 4.3 of [1], the LRC(2) code has length  $n = (q^2 - 1)q$ , dimension k = (q - 1)q and distance

(5.7) 
$$d \ge (q+1)(q^2 - 3q + 3) = q^3 - 2q^2 + 3.$$

As before, we improve the bound on the distance using Hermitian codes that contains the LRC(2) code. To do this we have to find  $m_{\perp}$ . By L, we have that m=q(q-1)+(q+1)(q-2) so we are in the fourth phase of Hermitian codes because  $m_{\perp}=n+2g-2-m=q^3-q^2+q$ . In this case  $d_{Her}=m_{\perp}-2g+2=q^3+2q+2$ . Since  $|B|=q^3-q$ , we have that

$$(5.8) d_{LRC} \ge d_{Her} - q = q^3 + q + 2.$$

Note that this bound improves bound (5.7). We just proved the following proposition:

**Proposition 5.3.** Let C be a linear code obtained by evaluating the functions in L at the points of B. Then C has the following parameters:

$$n = (q^2 - 1)q$$
,  $k = (q - 1)q$ ,  $r_1 = q - 1$ ,  $r_2 = q$  and  $d \ge q^3 + q + 2$ .

#### ACKNOWLEDGEMENTS

The first author is partially supported by MIUR and GNSAGA of INDAM (Italy).

The authors would like to thank the anonymous referees for their comments.

## References

- [1] A. Barg, I. Tamo, and S. Vlădut. Locally recoverable codes on algebraic curves. arXiv preprint arXiv:1501.04904, 2015.
- [2] H. Chen, R. Cramer, S. Goldwasser, R. De Haan, and V. Vaikuntanathan. Secure computation from random error correcting codes. In *Advances in Cryptology-EUROCRYPT 2007*, pages 291–310. Springer, 2007.
- [3] M. Forbes and S. Yekhanin. On the locality of codeword symbols in non-linear codes. *Discrete Mathematics*, 324:78–84, 2014.
- [4] G. D. Forney. Dimension/length profiles and trellis complexity of linear block codes. *Information Theory, IEEE Transactions on*, 40(6):1741–1752, 1994.
- [5] O. Geil. On codes from Norm-Trace curves. Finite Fields Appl., 9:351–371, 2003.
- [6] O. Geil, S. Martin, R. Matsumoto, D. Ruano, and Y. Luo. Relative generalized Hamming weights of one-point algebraic geometric codes. *Information Theory, IEEE Transactions on*, 60(10):5938-5949, 2014.
- [7] P. Gopalan, V. Guruswami, and P. Raghavendra. List decoding tensor products and interleaved codes. SIAM Journal on Computing, 40(5):1432–1462, 2011.
- [8] P. Gopalan, C. Huang, H. Simitci, and S. Yekhanin. On the locality of codeword symbols. *Information Theory, IEEE Transactions on*, 58(11):6925–6934, 2012.
- [9] V. Guruswami. List decoding from erasures: Bounds and code constructions. *Information Theory, IEEE Transactions on*, 49(11):2826–2833, 2003.
- [10] P. Heijnen and R. Pellikaan. Generalized Hamming weights of q-ary Reed-Muller codes. In *IEEE Trans. Inform. Theory.* Citeseer, 1998.
- [11] T. Høholdt, J. H. van Lint, and R. Pellikaan. Algebraic geometry of codes. In V. S. Pless and W. Huffman, editors, *Handbook of coding theory*, Vol. I, II, pages 871–961. North-Holland, 1998.
- [12] M. Homma and S. J. Kim. The second generalized hamming weight for two-point codes on a Hermitian curve. *Designs, Codes and Cryptography*, 50(1):1–40, 2009.
- [13] C. Huang, M. Chen, and J. Li. Pyramid codes: Flexible schemes to trade space for access efficiency in reliable data storage systems. *ACM Transactions on Storage (TOS)*, 9(1):3, 2013.
- [14] W. C. Huffman and V. Pless. Fundamentals of error-correcting codes. Cambridge university press, 2003.
- [15] H. Janwa and A. K. Lal. On generalized Hamming weights and the covering radius of linear codes. In Applied algebra, algebraic algorithms and error-correcting codes, pages 347–356. Springer, 2007.
- [16] T. Kasami, T. Takata, T. Fujiwara, and S. Lin. On complexity of trellis structure of linear block codes. *Information Theory, IEEE Transactions on*, 39(3):1057–1064, 1993.
- [17] T. Kasami, T. Takata, T. Fujiwara, and S. Lin. On the optimum bit orders with respect to the state complexity of trellis diagrams for binary linear codes. *Information Theory*, *IEEE Transactions on*, 39(1):242–245, 1993.

- [18] J. Kurihara and T. Uyematsu. Strongly-secure secret sharing based on linear codes can be characterized by generalized Hamming weight. In *Communication, Control, and Computing (Allerton), 2011 49th Annual Allerton Conference on*, pages 951–957. IEEE, 2011.
- [19] J. Kurihara, T. Uyematsu, and R. Matsumoto. Secret sharing schemes based on linear codes can be precisely characterized by the relative generalized hamming weight. *IEICE Transactions* on Fundamentals of Electronics, Comications and Computer Sciences, 95(11):2067–2075, 2012.
- [20] H. Lange and G. Martens. On the gonality sequence of an algebraic curve. Manuscripta mathematica, 137(3-4):457–473, 2012.
- [21] Y. Luo, C. Mitrpant, A. J. H. Vinck, and K. Chen. Some new characters on the wire-tap channel of type II. *Information Theory, IEEE Transactions on*, 51(3):1222–1229, 2005.
- [22] C. Marcolla. On structure and decoding of Hermitian codes. PhD thesis, University of Trento, 2013.
- [23] S. Martin and O. Geil. Relative generalized Hamming weights of q-ary Reed-Muller codes. arXiv preprint arXiv:1407.6185, 2014.
- [24] C. Munuera. On the generalized Hamming weights of geometric Goppa codes. IEEE transactions on information theory, 40(6):2092–2099, 1994.
- [25] C. Munuera and D. Ramirez. The second and third generalized Hamming weights of Hermitian codes. Information Theory, IEEE Transactions on, 45(2):709–712, 1999.
- [26] C.-K. Ngai, R. W. Yeung, and Z. Zhang. Network generalized hamming weight. Information Theory, IEEE Transactions on, 57(2):1136–1143, 2011.
- [27] W. Olaya-León and C. Granados-Pinzón. The second generalized hamming weight of certain Castle codes. *Designs, Codes and Cryptography*, 76(1):81–87, 2015.
- [28] D. S. Papailiopoulos and A. G. Dimakis. Locally repairable codes. *Information Theory, IEEE Transactions on*, 60(10):5843–5855, 2014.
- [29] R. Pellikaan. On the gonality of curves, abundant codes and decoding. In *Coding theory and algebraic geometry*, pages 132–144. Springer, 1992.
- [30] B. Poonen. Gonality of modular curves in characteristic p. Mathematical Research Letters, 14(4):691–701, 2007.
- [31] H. G. Ruck and H. Stichtenoth. A characterization of Hermitian function fields over finite fields. *Journal fur die Reine und Angewandte Mathematik*, 457:185–188, 1994.
- [32] N. Silberstein, A. S. Rawat, O. O. Koyluoglu, and S. Vishwanath. Optimal locally repairable codes via rank-metric codes. In *Information Theory Proceedings (ISIT)*, 2013 IEEE International Symposium on, pages 1819–1823. IEEE, 2013.
- [33] H. Stichtenoth. Algebraic function fields and codes. Universitext. Springer-Verlag, Berlin, 1993.
- [34] I. Tamo and A. Barg. A family of optimal locally recoverable codes. *Information Theory*, *IEEE Transactions on*, 60(8):4661–4676, 2014.
- [35] I. Tamo, D. S. Papailiopoulos, and A. G. Dimakis. Optimal locally repairable codes and connections to matroid theory. In *Information Theory Proceedings (ISIT)*, 2013 IEEE International Symposium on, pages 1814–1818. IEEE, 2013.
- [36] M. Tsfasman, S. Vlădut, and D. Nogin. Algebraic geometric codes: basic notions, volume 139. American Mathematical Soc., 1990.
- [37] V. K. Wei. Generalized Hamming weights for linear codes. Information Theory, IEEE Transactions on, 37(5):1412–1418, 1991.
- [38] K. Yang, P. V. Kumar, and H. Stichtenoth. On the weight hierarchy of geometric Goppa codes. *Information Theory, IEEE Transactions on*, 40(3):913–920, 1994.

Edoardo Ballico. Dipartimento di Matematica dell'Università di Trento, Via Sommarive 14, 38123 Povo (TN), Italy

E-mail address: edoardo.ballico@unitn.it

Chiara Marcolla. Dipartimento di Matematica dell'Università di Torino, Via Carlo Alberto 10, 10123 Torino, Italy

E-mail address: chiara.marcolla@unito.it