INVARIANT SUBSETS OF SCATTERED TREES AND THE TREE ALTERNATIVE PROPERTY OF BONATO AND TARDIF

CLAUDE LAFLAMME*, MAURICE POUZET**, AND NORBERT SAUER***

ABSTRACT. A tree is *scattered* if it does not contain a subdivision of the complete binary tree as a subtree. We show that every scattered tree contains a vertex, an edge, or a set of at most two ends preserved by every embedding of T. This extends results of Halin, Polat and Sabidussi.

Calling two trees equimorphic if each embeds in the other, we then prove that either every tree that is equimorphic to a scattered tree T is isomorphic to T, or there are infinitely many pairwise non-isomorphic trees which are equimorphic to T. This proves the tree alternative conjecture of Bonato and Tardif for scattered trees, and a conjecture of Tyomkyn for locally finite scattered trees.

1. Introduction

In this paper we deal with trees, alias connected graphs with no cycle. We follow Diestel [3] for most of the standard graph theory notions. A *ray* is a one-way infinite path.

We first generalizes theorems dealing with automorphisms of trees obtained by Polat and Sabidussi [16]. They proved that every rayless tree has either a vertex or an edge preserved by every automorphism, and that a scattered tree having a set of at least three ends of maximal order contains a rayless tree preserved by every automorphism. (Scattered trees are defined in the abstract and at the end of Section 2. Ends are equivalence classes of rays and defined in Section 5, just before Subsection 5.1. The order of an end, which we usually call rank, is defined and discussed in Subsection 5.4.)

Our first result pertains to scattered trees for which there is no end of maximal order. We prove that if T is such a tree then it contains a rayless tree preserved by every embedding (injective endomorphism) of T. Together with a result of Halin [4], who showed that every rayless tree contains a vertex or an edge preserved by every embedding, we obtain the first main contribution of this paper:

Date: Calgary May-June 2012 – Version June 13, 2016.

²⁰⁰⁰ Mathematics Subject Classification. Partially ordered sets and lattices (06A, 06B).

Key words and phrases. graphs, trees, equimorphy, isomorphy.

^{*}Supported by NSERC of Canada Grant # 10007490.

^{**}Research started while the author visited the Mathematics and Statistics Department of the University of Calgary in June 2012; the support provided is gratefully acknowledged.

^{***}The third author was supported by NSERC of Canada Grant # 10007490.

Theorem 1.1. If a tree T is scattered, then either there is one vertex, one edge, or a set of at most two ends preserved by every embedding of T.

This is proved in Section 7 as a consequence of Lemmas 7.4 and 7.5. Of course the fact that every end in a scattered tree has an order (or rank), which has been proven by Jung in [10], is needed. However, for our proofs we need a more structural development, with a finer differentiation of the types of preserved elements. In this context we will also reprove Jung's theorem (cf. Section 6.1).

The above results pertain to scattered trees with no end of maximal order or at least 3 ends of maximal order. If there are only two ends of maximal order, they must be preserved. However, the situation of two ends preserved by every embedding is quite easy to characterize:

Proposition 1.2. A tree T has a set of two distinct ends preserved by every embedding if and only if it contains a two-way infinite path preserved by every embedding.

The case of an end preserved by every embedding is more subtle. Indeed, this does not imply that the end contains a ray preserved by every embedding.

We provide examples in Subsection 5.8 of scattered trees with such an end but which do not contain an infinite path nor a vertex nor an edge preserved by every embedding.

In order to present our next results, we introduce ¹ the following notions.

Let T be a tree and e be an end. We say that e is preserved forward, resp. backward, by an embedding f of T if there is some ray $C \in e$ such that $f[C] \subseteq C$, resp. $C \subseteq f[C]$. We say that e is almost rigid if it is preserved backward and forward by every embedding, that is every embedding fixes pointwise a cofinite subset of every ray belonging to e.

We recall that two trees T and T' are equimorphic, or twins, if T is isomorphic to an induced subtree of T' and T' is isomorphic to an induced subtree of T. As we will see (cf. Lemma 4.1), if $C := \{x_0, \ldots x_n, \ldots\}$ is a ray, then T decomposes as a sum of rooted trees indexed by C, that is $T = \bigoplus_{x_i \in C} T_{x_i}$ where T_{x_i} is the tree, rooted at x_i , whose vertex set is the connected component of x_i in $T \setminus \{x_{i-1}, x_{i+1}\}$ if $i \ge 1$ and in $T \setminus \{x_{i+1}\}$ if i = 0. If the number of pairwise non-equimorphic rooted trees T_{x_i} is finite, we say that C is regular. We say that an end is regular if it contains some regular ray (in which case all other rays that it contains are regular).

Theorem 1.3. Let T be a tree. If T is scattered and contains exactly one end e of maximal order, then e is preserved forward by every embedding. If T contains a regular end e preserved forward by every embedding, then e contains some ray preserved by every embedding provided that T is scattered or e is not almost rigid.

¹We will use an alternative presentation in Section 9 based on the notion of level function (see Subsection 5.1).

Theorem 1.3 will be proved in Subsection 5.7; it follows from Corollary 5.24, Proposition 5.8 and Proposition 5.12.

Corollary 1.4. Let T be a scattered tree. If T contains a regular almost rigid end e, then e contains a ray fixed pointwise by every embedding. In particular, the set of embeddings of T has a common fixed point.

Examples of trees containing an end preserved backward by every embedding and no end preserved forward by every embedding were obtained independently by Hamann [7] (see Example 5 in Subsection 5.8) and Lehner [14] (see Example 6 in Subsection 5.8). They are reproduced with their permission.

Definition 1.5. A tree T is called stable if either:

- (1) There is a vertex or an edge or a two-way infinite path or a one-way infinite path preserved by every embedding or an almost rigid end;
- (2) Or there exists a non-regular end which is preserved forward by every embedding.

Summarizing Theorems 1.1 and 1.3 we have.

Theorem 1.6. Every scattered tree is stable.

We will apply these results to conjectures of Bonato-Tardif and Tyomkyn.

Definition 1.7. The tree alternative property holds for a tree T if either every tree equimorphic to T is isomorphic to T, or else there are infinitely many pairwise non-isomorphic trees which are equimorphic to T.

Conjecture 1.8 (Bonato-Tardif [1]). The tree alternative property holds for every tree.

Bonato and Tardif [1] proved that their conjecture holds for rayless trees, and their result was extended to rayless graphs by Bonato et al, see [2].

Now let twin(T) denote the set of twins of T up to isomorphism. Note that if T is a tree for which every embedding is an automorphism, that is surjective, then |twin(T)| = 1; in particular this is the case for any locally finite rooted tree. On the other hand a star R with infinitely many vertices is an example of a tree having embeddings which are not automorphisms and with |twin(R)| = 1. Another such example is the tree consisting of two disjoint stars with infinitely many vertices whose roots are adjacent.

As this will be developed in Subsection 5.1, if e is an end in a tree T, there is an orientation of the edges of T which is the (oriented) covering relation of an ordering \leq_e on the set V(T) of vertices of T. Indeed, for every vertex $x \in V(T)$ there is unique ray starting at x and belonging to e; denoting this ray by e(x) we may set $x \leq_e y$ if $y \in e(x)$. If x is a vertex, let $T(\to x)$ be the induced rooted tree rooted at x with $V(T(\to x)) = \{y \in V(T) : y \leq_e x\}$.

We are now ready to state the tree alternative result for stable trees.

Theorem 1.9. (i) The tree alternative conjecture holds for stable trees. In particular:

- (ii) If T is a stable tree which does not contain a vertex or an edge preserved by every embedding or an almost rigid end and which has a non-surjective embedding, then $|twin(T)| = \infty$ unless T is the one-way infinite path.
- (iii) If T has a vertex or an edge or an almost rigid end preserved by every embedding and T is locally finite, then every embedding of T is an automorphism of T.
- (iv) If T has an almost rigid end, then |twin(T)| = 1 if and only if $|twin(T(\to x))| = 1$ for every vertex x. Otherwise $|twin(T)| = \infty$.
- (v) T has a non-regular and not almost rigid end preserved forward by every embedding, then $|twin(T)| \ge 2^{\aleph_0}$.

Theorem 1.9 is proved in Sections 9 with the help of results of Section 8. Note that Theorems 1.9 and 1.6 together with Definition 1.5 imply:

Corollary 1.10. Let T be a scattered tree with $|twin(T)| < 2^{\aleph_0}$. Then there exists a vertex or an edge or a two-way infinite path or a one-way infinite path or an almost rigid end preserved by every embedding of T.

Tyomkyn [18] proved that the tree alternative property holds for rooted trees and conjectured that twin(T) is infinite at the exception of the one-way infinite path, for every locally finite tree T which has a non-surjective embedding.

We obtain from Theorem 1.6 and from Theorem 1.9 we now obtain the second main contribution of the paper:

Theorem 1.11. The tree alternative conjecture holds for scattered trees, and Tyomkyn's conjecture holds for locally finite scattered trees.

Theorem 1.9 clearly holds for finite trees. That Theorem 1.6 holds for finite trees can either be seen directly by successively removing endpoints until an edge or a single vertex remains, or by just applying the Polat, Sabidussi, Halin results mentioned above. Hence, in the process of proving Theorems 1.6 and 1.9, we will mostly assume that our trees T are infinite.

The results of this paper have been presented in part at the workshop on Homogeneous Structures, Banff, Nov. 8-13, 2015 and to the seminar on Discrete Mathematics, Hamburg, Feb. 19. 2016. The paper benefited of remarks from the audiences and we are pleased to thank them; in particular we thank M. Hamann and F. Lehner for their examples of trees.

We further thank M. Hamann for the information he provided on his recent result generalizing Theorem 1.1. This generalization [9] has two parts, and the first one reads as follows:

Theorem 1.12. Let G be a monoid of embeddings of a tree T. Then either:

- (1) There is a vertex, an edge or a set of at most two ends preserved by each member of G;
- (2) Or G contains a submonoid freely generated by two embeddings.

Note that if G is a group of automorphisms, the conclusion is similar: in the second case G contains a subgroup freely generated by two automorphisms

(see Theorem 2.8 and Theorem 3.1 of [8]). For anterior versions of this result, see Tits [17], Pays and Valette [15] and Woess [19].

In his second part, M. Hamann proves that if the first case of the alternative above does not hold, then T contains a subdivision of the binary tree. For doing so, he shows that there are infinitely many embeddings with different directions (the *direction* of an embedding f being an end preserved forward by f with a positive period (see Section 5 for the definition of period)). Then, he observes that there are two embeddings f and g such that none fixes the direction of the other. From that, he builds a subdivision of the binary tree.

We wish to warmly thank the referees for their truly valuable and appreciated contributions.

2. Basic definitions

Let T be a tree. For $x, y \in V(T)$ we write $x \sim y$ if x is adjacent to y. An embedding of T into a tree T' is an injection f of V(T) to V(T') for which $x \sim y$ if and only if $f(x) \sim f(y)$ for all $x, y \in V(T)$. An embedding of T is an embedding of T into T. We write $H \subseteq T$ if H is a subtree of T. That is if H is connected and $V(H) \subseteq V(T)$. If $H \subseteq T$ and f is an embedding of T then f[H]is the subtree of T induced by the set $\{f(h): h \in V(H)\}\$ of vertices. Hence an embedding f is an isomorphism of H to f[H]. An embedding f preserves a subtree H of T if $f[H] \subseteq H$. Note that if H is finite or a two-way infinite path and f preserves H then f[H] = H, in which case f restricted to H is an automorphism of H. The embedding f fixes a vertex x if it preserves it, that is if f(x) = x. Two trees T and R are isomorphic, resp. equimorphic or twins, and we set $T \simeq R$, resp. $T \equiv R$, if there is an isomorphism of T onto R, resp. an embedding of T to R and an embedding of R to T. A rooted tree is a pair (T,r) consisting of a tree T and a vertex $r \in V(T)$, the root. If (T,x)is a rooted tree with root x and (R, y) is a rooted tree with root y, then an embedding of (T,x) to (R,y) is an embedding of T to R which maps x to y. The definitions above extend to rooted trees.

Let P be an ordered set (poset), that is a set equipped with an order relation, denoted by \leq . We say that P is an *ordered forest* if for each element $x \in P$, the set $\downarrow x := \{y \in P : y \leq x\}$ is totally ordered. If furthermore, two arbitrary elements have a common lower bound, then P is an *ordered tree*.

Rooted trees can be viewed as particular types of ordered trees. Indeed, let (T,r) be a rooted tree; if we order T by setting $x \le y$ if x is on the shortest path joining r to y we get an ordered tree with least element r, such that $\downarrow x$ is finite for every $x \in T$. Conversely suppose that P is an ordered tree, with a least element 0, such that $\downarrow x$ is finite for every $x \in P$. Then a) P is a meet-semilattice, that is every pair of elements x and y has a meet (a largest lower bound) that we denote by $x \land y$ and b) the unoriented graph of the covering relation of P is a tree (we recall that an element $y \in P$ covers x, and we note $x \lessdot y$, if $x \lessdot y$ and there is no element z such that $x \lessdot z \lessdot y$).

In the sequel, we will rather consider the dual of the above order on a rooted tree, we will denote it \leq_r and we will denote by P(T,r) the resulting poset.

Let $2^{<\mathbb{N}}$ be the set of finite sequences with entries 0 and 1. For $s \in 2^{<\mathbb{N}}$ we denote by s 0 and s 1 the sequences obtained by adding 0 and 1 on the right of s; we also denote by \square the empty sequence. There are an order and a graph structure on $2^{<\mathbb{N}}$. Ordering $2^{<\mathbb{N}}$ via the initial subsequence ordering, the least element being the empty sequence, we obtain an ordered tree, the binary ordered tree. Considering the (undirected) covering relation we get a tree, the binary tree also known as dyadic tree, which we denote by T_2 . A tree T is scattered (See Figure 1) if no subdivision of the binary tree T_2 is embeddable into T.




FIGURE 1. A tree is *scattered* if it does not embed a subdivision of the binary tree.

3. Automorphisms and embeddings

We recall that an automorphism f of a tree T is called a *rotation* if it fixes some vertex x; it is called an *inversion* if it reverses an edge, and it is called a *translation* if it leaves invariant a two-way infinite path. We recall the following basic result due to Tits:

Theorem 3.1. [17] Every automorphism of a tree is either a rotation, an inversion or a translation.

The short and beautiful proof is worth recalling.

Proof. For two elements $x, y \in V(T)$, let $d_T(x, y)$ be the length of the shortest path joining x to y. Let f be an automorphism of T. Let $x \in V(T)$ with $m := d_T(x, f(x))$ minimum. If f(x) = x, then f is a rotation. If $f(x) \neq x$ and f(f(x)) = x, then m = 1 and f is an inversion. If $f(f(x)) \neq x$, then $C := \bigcup_{n \in \mathbb{Z}} f^{(n)}[C[x, f(x)]]$, where C[x, f(x)] is the the shortest path joining x and f(x), is a two-way infinite path left invariant by f.

The case of embeddings is similar. For this Halin proved the following result:

Theorem 3.2. [5] Let f be an embedding of a tree T into itself. Then either there is:

(1) A fixed point; or

- (2) An edge reversed by f; or
- (3) A ray C with $f[C] \subset C$.

Furthermore, each case excludes the others.

Case (3) of the theorem above can be refined into two parts as follows:

Proposition 3.3. Let f be an embedding of a tree T into itself. Then either there is:

- (1) A fixed point; or
- (2) An edge reversed by f; or
- (3) A two-way infinite path preserved by f on which f fixes no vertex and reverses no edge; or
- (4) A ray preserved by f with a vertex not in the range of f.

Furthermore, each case excludes the others and the edge and path in Cases (2) and Case (3) are unique, whereas in Case (4), the ray has a unique maximal extension to a ray preserved by f.

4. Decomposition of trees

The basic idea for the finite case has been to study the effect of reducing the tree by removing the endpoints and study the effect of this operation on the embeddings, of course in this case the automorphisms, of the tree. If the tree T is infinite, we will have to decompose the tree into larger pieces and then determine properties of the tree in relation to properties of the parts and the way those parts are put together to reconstitute the tree. Of course some notation, which will then be used throughout the paper, has to be introduced to describe those actions. In order to reduce the length of chains of symbols we will often, if we think that no confusion is possible, use the symbol for a tree to also stand for the set of its vertices and the set of vertices of an induced subtree of a tree to stand for the subtree. So for example if C is a path in a tree, then we might write $x \in C$ and $P \subseteq C$ to indicate that x is a vertex of C and C is a subpath of C and so on.

Let T be a tree and X be a subset of T. We denote by $T \setminus X$ the subgraph of T induced by $V(T) \setminus X$. If K is a subtree of T and x a vertex of K, we denote by (T,K)(x) the connected component of x in $T \setminus (K \setminus \{x\})$. Note that $(T,\{x\})(x) = T$ and that since T is a tree, (T,K)(x) is the set of vertices on the paths of T meeting K at x. Note that $(T,K)(x) = \{x\}$ whenever the neighbourhood of x is included in K. We denote by ((T,K)(x),x) the tree (T,K)(x) rooted at x. The tree T then consists of the tree K with the rooted trees ((T,K)(x),x) attached at x for every $x \in V(K)$. In general the trees ((T,K)(x),x) may be given as a set of rooted trees of the form (T_x,r_x) . That is with a function associating the rooted trees to the vertices of K and then we identify the root r_x with x and let the trees (T_x,r_x) disjointly stick out of K.

Here is the formal definition: Let T be a tree, X be a subset of V(T) and $(T_x, r_x)_{x \in X}$ be a family of rooted trees. The *sum* of this family is the tree

denoted by $\bigoplus_{x \in X,T} (T_x, r_x)$, or also simply $\bigoplus_{x \in X,T} (T_x, x)$, which is obtained by first taking isomorphic copies T'_x of those trees which are pairwise disjoint and which contain x instead of r_x as a root. With the understanding that if the trees T_x are already pairwise disjoint and contain x as a root, then we will not take different copies of them. The set of vertices of the tree $\bigoplus_{x \in X,T} (T_x,x)$ is the set $V(T) \cup \bigcup_{x \in X} V(T'_x)$ and the edge set is $E(T) \cup \bigcup_{x \in X} E(T_x)$. If X = V(T) we simply denote this tree by $\bigoplus_{x \in T} (T_x, r_x)$. In some special cases we will simplify even further. For example if $\{(T_n, r_n) : n \in \mathbb{N}\}$ is a set of rooted trees, then $\bigoplus_{\mathbb{N}} T_n := \bigoplus_{n \in \mathbb{N}} T_n$ with the understanding that \mathbb{N} also denotes the ray indexed naturally by \mathbb{N} . As to be expected we then have:

Lemma 4.1. If T is a tree and K a subtree, then $T = \bigoplus_{x \in K} ((T, K)(x), x)$.

Proof. First, the vertex-sets of the trees (T,K)(x) and (T,K)(y) for $x \neq y$ are disjoint and if there is an edge adjacent to a vertex in (T,K)(x) and also adjacent to a vertex in (T,K)(y), then it is the edge (x,y). Indeed, by construction, $(T,K)(x) \cap K = \{x\}$ for each $x \in K$. If (T,K)(x) and (T,K)(y) have a non-empty intersection or there is an edge not equal to (x,y) from (T,K)(x) to (T,K)(y), then the path from x to y would complete a cycle of T.

Let T be a tree and $\emptyset \neq X \subseteq T$. For every vertex $x \in V(T)$, the distance from x to X, denoted by $d_T(x,X)$ is the least integer n such that there is some path of length n from x to some vertex $y \in X$. Hence $d_T(x,X) = 0$ iff $x \in X$. If X induces a subtree, say K, this vertex y is unique; we denote it by $p_{(T,K)}(x)$.

Lemma 4.2. If K is a subtree of T, then the map $p_{(T,K)}$ from T to K is a retraction of reflexive graphs and $p_{(T,K)}^{-1}(y) = (T,K)(y)$ for every $y \in V(K)$.

Proof. Clearly, $p_{(T,K)}(x) = x$ if and only if $x \in V(K)$. Hence $p_{(T,K)} \circ p_{(T,K)} = p_{(T,K)}$ and K is the range of $p_{(T,K)}$. This amounts to say that $p_{(T,K)}$ is a set-retraction of V(T) onto V(K). To conclude that it is a reflexive-graph-retraction, we need to prove that it transforms an edge into an edge or identifies its end vertices. For that, let $x, x' \in V(T)$, $y := p_{(T,K)}(x)$ and $y' := p_{(T,K)}(x')$. Then, as it is easy to see, $d_T(x,x') = d_T(x,y) + d_T(y,y') + d_T(y',x')$. Hence, if $u := \{x,x'\} \in E(T)$ and $\{y,y'\}$ is not an edge, then y = y' as required. The fact that the vertices on the shortest path from a vertex x to K belong to $y := p_{(T,K)}(x)$ ensures that $K_y := ((T,K)(y),y)$ is connected. It follows that T is the sum $\bigoplus_{y \in K} K_y$.

Lemma 4.3. Let T be a tree, C and C' be two infinite paths of T and f be an embedding of C into C'. Then ((T,C)(x),x) is embeddable into ((T,C')(f(x)),f(x)) for every $x \in V(C)$ if and only if f has an extension g to an embedding of T into itself such that $C = g^{-1}(C')$.

Proof. For the direct implication, choose for each $x \in V(C)$ an embedding f_x of ((T,C)(x),x) into ((T,C')(f(x)),f(x)). Observe that $g := \bigcup_{x \in V(C)} f_x$ is an embedding of $\bigoplus_{x \in C} ((T,C)(x),x)$ into $\bigoplus_{y \in C'} ((T,C')(y),y)$ and $C = g^{-1}(C')$. According to Lemma 4.1 $\bigoplus_{x \in C} ((T,C)(x),x) = T$, hence the implication holds.

For the converse, note that if g is an extension of f, then g induces an embedding of ((T,C)(x),x) into ((T,C')(f(x)),f(x)) for every vertex $x \in V(C)$ of degree 2. Thus the converse holds if C is a two-way infinite path. If C is a one-way infinite path, i.e. a ray, let x_0 be its vertex of degree 1. Due to the condition on the map g, it induces an embedding of $((T,C)(x_0),x_0)$ into $((T,C')(f(x_0)),f(x_0))$.

Corollary 4.4. Let f be an embedding of a tree T which preserves a two-way-infinite path D. Then $f((T,D)(x),x)) \subseteq ((T,D)(f(x)),f(x))$ for all $x \in V(D)$.

If f is an embedding which preserves a ray C starting at r_0 with the predecessor of $f(r_0)$ in C not in V(f[T]), then

$$f((T,C)(x),x)\subseteq ((T,C)(f(x)),f(x))$$

for all $x \in V(C)$.

5. Ends

Let T be a tree. Two rays C and C' are called *equivalent* if the set $V(C) \cap V(C')$ of vertices induces a ray, equivalently if $C \setminus F = C' \setminus F$ for some finite subset F of V(T). This relation is an equivalence relation, with classes called *ends*; the equivalence class of a ray C is denoted by end(C). The set of ends is denoted by $\Omega(T)$.




FIGURE 2. An *end* of a tree is an equivalence class of rays.

We note that since T is tree, then for each end $e \in \Omega(T)$ and $x \in V(T)$ there is a unique ray originating at x and belonging to e. We denote it by e(x). For $n \in \mathbb{N}$ we denote by $x \boxplus n$ (or $x \boxplus_e n$ if there is a risk of confusion) the vertex of e(x) at distance n of x. For $y \in e(x)$ we denote by T < e(x), y > the tree rooted at y and whose vertex set is the connected component of y in $V(T) \setminus \{y^-, y^+\}$ where y^- and y^+ are the predecessor and successor of y in e(x). If y = x, this rooted tree will be also denoted by $T(\to x)$, its vertex set is the connected component of x in $V(T) \setminus \{x^+\}$. We note that $T = \bigoplus_{y \in e(x)} T < e(x), y >$.

To each end e is attached an order \leq_e and to $\Omega(T)$ a topology. We discuss their properties below.

5.1. Orders on a tree. Let T be a tree and K be a subtree. Let $x, y \in V(T)$; we set $x \leq_K y$ if y is on the shortest path joining x to some vertex of K.

This relation is an order on V(T), V(K) is the set of maximal elements of (T, \leq_K) . For every edge $u := \{x, y\}$ disjoint from K either $x <_K y$ or $y <_K x$. Indeed, since K is connected, no path containing x and y has its end vertices in K, hence either the shortest path from x to K contains y or the shortest path from y to K contains x. For each vertex $x \in V(K)$, $(T, K)(x) = \{y \in V(T) : y \leq_K x\}$. As a poset, V(T) is the dual of a forest. If K reduces to a vertex $x \in V(T)$, we denote by P(T, x) the resulting poset; its dual satisfies properties a) and b) given in Section 2.

Let $e \in \Omega(T)$. Let $x, y \in V(T)$. We set $x \leq_e y$ if $y \in e(x)$. This relation is an order. The unoriented covering relation of this order is the adjacency relation on T. As a poset, V(T) equipped with this order is the dual of an ordered tree; this dual is a join-semilattice. We denote by $x \vee_e y$ the join of two vertices $x, y \in V(T)$.

Lemma 5.1. Let f be an embedding of a tree T into itself. Then an end e, as an equivalence class, is sent by f into an equivalence class e' of T and furthermore:

f preserves the covering relation, (that is $x \leq_e y$ iff $f(x) \leq_{e'} f(y)$ for all $x, y \in V(T)$).

In particular f is a join-semilattice embedding from (T, \leq_e) into $(T, \leq_{e'})$.

Proof. Since $f(e) \subseteq e'$ we have f(e(x)) = e'(f(x)) for every $x \in V(T)$. Let $x, y \in V(T)$. We have respectively $x \lessdot_e y$ iff $x \sim y$ and $y \in e(x)$ and $f(x) \lessdot_{e'} f(y)$ iff $f(x) \sim f(y)$ and $f(y) \in e'(f(x))$. Since $f(e) \subseteq e'$, $x \lessdot_e y$ implies trivially $f(x) \lessdot_{e'} f(y)$. Conversely, suppose that $f(x) \lessdot_{e'} f(y)$. Let $C \in e$. By deleting some elements of C we may suppose that $x, y \notin C$. Let $z \coloneqq p_{(T,C)}(x)$, $C' \coloneqq f(C)$ and $z' \coloneqq f(z)$. Since embeddings are isometric, we have $z' = p_{(T,C')}(x)$. Since $f(x) \leq_{e'} f(y)$ and $C' \in e'$, f(y) is on the shortest path joining f(x) to C'. This path ending at z' it is the image of the shortest path joining x to z, hence y belongs to this path, proving that $x \leq_e y$; since $f(x) \sim f(y)$, it follows that $x \sim y$ hence $x \lessdot_e y$. From this, we have $f(x \vee_e y) = f(x) \vee_{e'} f(y)$ for all $x, y \in V(T)$. Indeed, let $x, y \in V(T)$. Let $z \coloneqq f(x) \vee_{e'} f(y)$. Since f is order preserving, $z \leq_{e'} f(x \vee_e y)$, hence z is both on the shortest path joining f(x) and $f(x \vee_e y)$. Since f is a one-to one isometry, $z = f(x \vee_e y)$.

Corollary 5.2. An embedding f of T preserves e, that is $f[e] \subseteq e$, iff f is a join-semilattice embedding of (T, \leq_e) .

5.2. Valuation, level function and period. Let T be a tree and e be an end of T. To each embedding f preserving e we attach an integer $d \in \mathbb{Z}$, the period of f. In order to do so we introduce the notion of valuation.

A e-valuation is any map $v:V(T)\to\mathbb{Z}$ such that

(1)
$$x \leqslant_e y \Rightarrow v(y) = v(x) + 1$$

for every $x, y \in V(T)$.

Lemma 5.3. Let e be an end of T. For every $x \in V(T)$, $n \in \mathbb{Z}$ there is a unique e-valuation v such that v(x) = n. In particular two e-valuations differ by some constant.

Proof. We order V(T) by \leq_e . Let $y \in V(T)$. Set $v(y) \coloneqq n + d_T(x, z) - d_T(y, z)$ where $z \coloneqq x \vee_e y$. Let $y' \lessdot_e y$. Observe that $y' \vee x = y \vee x = z$. If follows that v(y) = v(y') + 1. Hence v is an e-valuation. If v' is any valuation, we must have $v'(z) - v'(x) = d_T(x, z)$ and $v'(z) - v'(y) = d_T(y, z)$. Hence, $v'(y) = v'(x) + d_T(x, z) - d_T(y, z)$. Thus v' = v iff v'(x) = v(x). If v is a e-valuation and $k \in \mathbb{Z}$, then v + k is a e-valuation. Hence, if v' is an other e-valuation, set $k \coloneqq v(x) - v'(x)$. Then v' + k is an e-valuation which coincides with v on x. Hence v = v' + k.

From this lemma, it follows that if we choose a vertex $o \in V(T)$, there is a unique e-valuation v such that v(o) = 0. Such a vertex is called the *origin* of the e-valuation.

Lemma 5.4. Let e be an end of T, v be an e-valuation on T and f be an embedding. Then f preserves e iff there is some $d \in \mathbb{Z}$ so that v(f(x)) = v(x) + d for all vertices x of T. Moreover, d is non-negative iff f preserves e forward; d is negative iff f preserves e backward and not forward.

Proof. Set v'(x) := v(f(x)). If f preserves e, then v' is an e-valuation. Hence from Lemma 5.3, v' = v + d for some $d \in \mathbb{Z}$. Conversely, suppose that v(f(x)) = v(x) + d for all vertices x of T. Let $x, y \in V(T)$. We have v(f(y)) - v(f(x)) = v(y) - v(x). In particular, v(f(y)) - v(f(x)) = 1 iff v(y) - v(x) = 1, that is $f(x) \leq_e f(y)$ iff $x \leq_e y$. Since f preserves the covering relation \leq_e , it preserves e (Corollary 5.2). Suppose that f preserves e forward, then there is some $C \in e$ such that $f(C) \subseteq C$. This implies $x \leq_e f(x)$ for $x \in C$, hence $d \geq 0$. Similarly, if f preserves e backward there is some $C \in e$ such that $C \subseteq f(C)$, hence $d \leq 0$. If $C \neq f(C)$, then $d \neq 0$. We prove that if f preserves e, then it preserves e forward or backward according to the sign of d.

Claim 5.5. Let $x \in V(T)$, $z := x \vee_e f(x)$ and k := v(z) - v(x) - d. Then $k \ge 0$ and $f(x \boxplus k) = z$.

Proof of Claim 5.5. Since v(f(x)) = v(x) + d we have k = v(z) - v(f(x)); since $f(x) \leq_e z$ we have $k \geq 0$, hence $z' := x \boxplus k$ is well defined. We have v(z') - v(x) = v(f(z')) - v(f(x)) = v(z) - v(f(x)) = k. Since $f(x) \leq_e f(z')$ and $f(x) \leq_e z$, f(z') and f(z) are comparable. Since v(f(z')) - v(f(x)) = v(z) - v(f(x)) = k, f(z') = z.

Pick any $x \in V(T)$. Let $z := x \vee_e f(x)$ and $z' := x \boxplus k$. According to Claim 5.5, f(z') = z. If $d \ge 0$, then $z' \le_e z$ and hence $f[e(z')] \subseteq e(z')$, proving that e is preserved forward. If d = 0, then z' = z, and e(z') is fixed pointwise by f. If d < 0, then $z <_e z'$ then $e(z') \subseteq f[e(z')]$ and e is preserved downward. \square

The number d is the *period* of f w.r.t. v. According to Lemma 5.3 if v' is an other e-valuation, then f has the same period.

An embedding f preserves a valuation $v:V(T)\to\mathbb{Z}$ if v(f(x))=v(x) for all $x\in V(T)$. In this case, f preserves every other valuation attached to the same end. A map $v:V(T)\to\mathbb{Z}$ is a level function if this is the valuation associated with an end preserved forward by every embedding.

Lemma 5.6. Let v be a valuation associated with an end e. Then v is a level function iff every embedding of T has a non-negative period. In particular, v is a level function provided that e is almost rigid.

Proof. The first assertion follows from Lemma 5.4. The second assertion is obvious. \Box

Lemma 5.7. Let T be a tree, e be an end, f be an embedding with non-negative period d and $S_f := \{x \in V(T) : f(x) \in e(x)\}$. Then

$$(2) x \vee_e f(x) \in S_f$$

and

(3)
$$f(x \vee_e f(x)) = (x \vee_e f(x)) \boxplus d$$

for every $x \in V(T)$.

If d = 0, then S_f is the set of fixed points of f, whereas if d > 0, then S_f is either a ray belonging to e or a two-way infinite path preserved by f.

Proof. Equations (2) and (3) are equivalent. The second implies trivially the first; since $v(f(x \vee_e f(x)) = v(x \vee_e f(x)) + d$ the second equation follows from the first. To prove Equation (2) apply Claim 5.5: there is some $z' \leq_e z$ such that $f(z') = z := x \vee_e f(x)$. This implies $z' \in S_f$ hence $z \in S_f$.

Since $S_f = \{x \in V(T) : x \leq_e f(x)\}$ and f preserves \leq_e , it follows that f preserves S_f that is $f(S_f) \subseteq S_f$. Since f preserves \leq_e , f preserves some ray, say C, belonging to e, hence $C \subseteq S_f$ and thus S_f is non-empty. Suppose d > 0. We claim that S_f is totally ordered. From this, it follows that this is a path (a ray or a two-way infinite path). If this is not the case, S_f contains at least two incomparable elements, say x, y. Let $z := x \vee_e y$. Then let $m := d_T(x, y) = d_T(x, z) + d_T(y, z)$. Since f is an embedding f(x) and f(y) are incomparable. This implies that $f(x) \vee_e f(y) = z$. Since $f(x) = x \boxplus d$ and $f(y) = y \boxplus d$, we have $d_T(f(x), f(y)) = m - 2d$ contradicting the fact that f is an isometry.

5.3. Rays preserved by every embedding. We present two results on regular ends, namely Proposition 5.8 and Proposition 5.12. The proofs are based on the same idea, but the proof of the first one is much simpler.

Proposition 5.8. Le T be a scattered tree and e be an almost rigid regular end. Then e contains a ray preserved by every embedding of T.

Proof. Our aim is to find $x \in V(T)$ such that f(x) = x for every embedding f. Indeed, since e is almost rigid, all embeddings have period zero (Lemma 5.4) hence f(y) = y for every $y \in e(x)$ and every embedding f. In fact, we prove that under the weaker assumption that e is preserved forward by every embedding, then there is some x such that $f(x) \ge x$ for every embedding, from which follows that the ray e(x) is preserved by every embedding.

Pick $u \in V(T)$. Let $n \in \mathbb{N}$, set $x_n := u \boxplus n$ in e(u) and $T_n := T < e(u), x_n > \infty$.

Claim 5.9. There is $k \in \mathbb{N}$ such that for each $n \ge k$ the interval [0, k[contains two integers $k'_n < k''_n$ such that $T_{k'_n}, T_{k''_n}$ and T_n are equimorphic.

Proof of Claim 5.9. The existence of k is immediate: say that two integers n and m are equivalent if T_n and T_m are equimorphic (as rooted trees). The fact that e is regular means that this equivalence relation has only finitely many blocks. Pick two elements in each equivalence class whenever possible and otherwise one element; since the number of these elements is finite, some integer dominates the elements chosen. This integer has the required property.

Let k be given by Claim 5.9. We claim that $x_k \leq_e f(x_k)$ for every embedding f of T. Indeed, suppose not. Let $z \coloneqq x_k \vee_e f(x_k)$ in the join-semilattice (T, \leq_e) . Since $x_k \leq_e z$, $z \in e(u)$ hence there is some $n \in \mathbb{N}$ with $n \geq k$ such that $z = x_n$. Let $k'_n, k''_n < k$ such that $T_{k'_n}, T_{k''_n}$ and T_n are equimorphic as rooted trees. Since $x_n = x_k \vee_e f(x_k)$, the rays $e(x_k)$ and $e(f(x_k))$ meet at x_n (and not before), hence f embeds $T(\to x_k)$ into $T(\to f(x_k))$. In particular, the trees $T_{k'_n}$ and $T_{k''_n}$ embed into T_n . Set $r \coloneqq x_n$, $(R, r) \coloneqq T_n$, $y \coloneqq f(u \boxplus k'_n)$, $x = f(u \boxplus k''_n)$, $C_y \coloneqq f[\{u \boxplus m : k'_n \leq m \leq n\}$. According to Claim 5.10, T_n is non-scattered.

Claim 5.10. Let (R,r) be a rooted tree ordered by $u \leq_r v$ if v is on the unique path C_u joining u to r. Let $x, y \in V(R)$ such that $y <_r x <_r r$, let $R(C_y, x)$ be the tree rooted at x whose vertex set is the connected component of x in $R \setminus \{x^-, x^+\}$ (where x^- and x^+ are the neighbours of x in C_y) and $R(\to y)$ be the tree rooted at y whose vertex set is the connected component of y in $R \setminus \{y+\}$. If there exist embeddings of the rooted tree (R,r) into the rooted trees $R(\to y)$ and $R(C_y, x)$, then R is not scattered.

Proof of Claim 5.10. Let f be an embedding of (R, r) into $R(C_y, x)$ and g be an embedding of (R, r) into $R(\to y)$. Then f maps the path from y to r to a path from z := f(y) to x and g maps the path from y to r to the path from to z' = g(y) to y. Implying that there is an oriented path from z' = g(y) to x. It follows that every copy (R', r) of (R, r) contains three distinct vertices x, z and z' with an oriented path from z to x excluding z' and an oriented path from z to x excluding z' and embeddings of (R, r) into $R'(\to z)$ and into $R'(\to z')$. Implying that (R, r) contains a subdivision of the binary tree.

This concludes the proof of Proposition 5.8.

When we have embeddings with positive period, we do not need that T is scattered, but the proof is more complex.

Let T be a tree and e be an end. Let Emb(T) be the set of embeddings of T, $Emb_e(T)$ be the subset of those preserving e, $Emb_e^+(T)$ be the subset of those with positive period and \mathbf{d} be the greatest common divisor of the periods of members of $Emb_e^+(T)$. Let $u \in V(T)$ and $p \in \mathbb{N}$; we say that p is a period of e(u) if T < e(u), $y > \equiv T < e(u)$, $y \equiv p >$ for every y such that $u <_e y$. If e(u) has a positive period, then there is one which divides all the others, we will call it the period of e(u).

If $f \in Emb_e^+(T)$ and d is the period of f we set $\check{S}_f := \{x \in S_f : d \text{ is a period of } e(x)\}$. Furthermore, we set $\check{S} := \bigcup_{f \in Emb_e^+(T)} \check{S}_f$.

As we will see below, under the existence of embeddings with positive period, the regularity of an end amounts to the fact that it contains some periodic ray.

Lemma 5.11. Let T be a tree and e be a regular end. Then,

- (1) for every embedding $f \in Emb_e^+(T)$, the set \check{S}_f contains some u such that d is a period of e(u);
- (2) all rays e(u) for $u \in \check{S}$ have the same period; in particular this period divides \mathbf{d} ;
- (3) If every embedding of T has a non negative period, then \check{S} contains no two-way infinite path; in particular each \check{S}_f is a ray.

Proof. Item (1). Let $f \in Emb_e^+(T)$ and $u \in S_f$. Set $x_n := u \boxplus n$ and $T_n := T < e(u), x_n > \text{for } n \in \mathbb{N}$. Say that two non-negative integers n, m are equivalent if $T_n \equiv T_m$. Since e is a regular end, e(x) is a regular ray and this means that the equivalence relation above has only finitely many classes. Let d be the period of f. Then $f(x_n) = x_n \boxplus d = x_{n+d}$ for every non-negative integer n. Since f is an embedding of T, T_n is embeddable into T_{n+d} for every n > 0. Hence, the rooted trees $T_{n+d,k}$, for $k \geq 0$, form an increasing sequence w.r.t embeddability. Since the number of equivalence classes is finite, there is some k_n such that all n + d.k, for $k \geq k_n$ are equivalent (pick k_n such that T_{n+d,k_n} is maximal with respect to equimorphy). This means that on the set $\{m \in \mathbb{N} : m \geq n + d.k_n\}$, the congruence class of n modulo d is included into the equivalence class of $n + d.k_n$. By considering an upper bound of $k_1, \ldots k_d$, we get an integer ℓ such that on the set $\{m \in \mathbb{N} : m \geq \ell\}$ each congruence class is included into some equivalence class of our relation above. This means d is a period of $e(x_\ell)$ hence $x_\ell \in \check{S}_f$ and thus $e(x_\ell) \subseteq \check{S}_f$.

Item (2) Indeed, let $u, v \in \check{S}$, k and l be periods of e(u) and e(v) respectively. Let d be the greatest common divisor of k and l. The rays e(u) and e(v) intersect on the ray $e(x \vee_e y)$, hence k and l, and thus d, is a period of that ray. But then d is a period of e(u) and e(v). Taking for k and l the periods of e(u) and e(v), we get k = l, proving our assertion. This common period must divide the period of each $f \in Emb_e^+(T)$, hence it divides \mathbf{d} , proving that Item (2) holds.

Item (3) Suppose that \check{S} contains a two-way infinite path, say D. Let $y \in D$; pick $u \in D$ with $u <_e y$. Since $D \subseteq \check{S}$ there is some $f \in Emb_e^+(T)$ such that $u \in \check{S}_f$. According to Item (2) we have $T < D, y >= T < e(u), y >\equiv T < e(u), y \boxplus \mathbf{d} >= T < D, y \boxplus \mathbf{d} >$. Since $T = \bigoplus_{y \in D} T < D, y >$, every translation of D with period \mathbf{d} extend to an embedding of T. In particular, the translation $t_{-\mathbf{d}}$ on D defined by $t_{-\mathbf{d}}(x) \boxplus \mathbf{d} = x$ for $x \in D$ extends to an embedding of T, hence there are embeddings with negative period which preserve e.

We will denote by \check{E} the set of end points of \check{S} . To avoid confusion, we will use $\check{S}(T)$ and $\check{E}(T)$ when needed.

Proposition 5.12. Le T be a tree admitting a regular and not almost rigid end e preserved forward by every embedding. Then the intersection C_o of all rays e(u) for $u \in \check{E}(T)$ is a ray preserved by every embedding and the distance from the origin o of this ray to each $u \in \check{E}(T)$ is at most \mathbf{d} , where \mathbf{d} is the largest common divisor of the positive periods of the embeddings of T.

Proof. We prove that \check{E} has a join, say o, and that the distance from o to every $u \in \check{E}$ is at most \mathbf{d} . This is a consequence of the following claim.

Claim 5.13. Let $u \in \check{E}$ and $x := u \oplus d$. Then $x \leq_e f(x)$ for every $f \in Emb_e^+(T)$.

From this claim, we have $v \leq x$ for every $v \in \check{E}$, from which it follows that \check{E} has a join, and this join o is majorized by x. Indeed, let $v \in \check{E}$, let $g \in Emb_e^+(T)$ such that v is the origin of \check{S}_g . From the claim, we have $x \leq_e g(x)$ hence $x \in S_g$. The ray e(x) is periodic and its period divides \mathbf{d} , hence it divides the period of g thus $x \in \check{S}_g$ hence $u \leq_e x$. Since $u \leq o \leq x = u + \mathbf{d}$ the distance from u to o is at most \mathbf{d} .

We prove the claim by contradiction. We suppose $x \not \in f(x)$ for some $f \in Emb_e^+(T)$. We build a tree T' equimorphic to T which is of the form $T' := \bigoplus_{n \in \mathbb{Z}} T'_n$ and on which act all translations of \mathbb{Z} of period $k.\mathbf{d}$ for $k \in \mathbb{Z}$. From this, we obtain that there are embeddings of T with negative period, contradicting our hypothesis.

Let C := e(u). According to (2) of Lemma 5.11, C is periodic and its period divides \mathbf{d} . As in Lemma 5.11 set $x_n := u \boxplus n$ and $T_n := T < e(u), x_n > \text{for } n \in \mathbb{N}$. For $n \in \mathbb{Z}$, let T'_n be an isomorphic copy rooted at n of the rooted tree T_m where $m := 1 + (n \mod \mathbf{d})$ and $n \mod \mathbf{d}$ is the residue of $n \mod \mathbf{d}$. On the tree $T' := \bigoplus_{n \in \mathbb{Z}} T'_n$ all translations of \mathbb{Z} of period $k.\mathbf{d}$ for $k \in \mathbb{Z}$ extend to embeddings of T'.

Claim 5.14. T is embeddable into T'.

Proof of Claim 5.14. The ray e(u) is preserved by f. It extends to a maximal path D preserved by f. This path is either a ray or a two-way infinite path. We may embed it on the path on \mathbb{Z} by the map ϕ sending u on -1 and $u \boxplus 1$ on 0. We claim that T < D, x > embeds into $T'_{\varphi(x)}$ for each $x \in D$. From this, it follows that φ extends to an embedding of T into T' as claimed. If $x := u \boxplus (n+1)$, with $n \ge 0$, then $\varphi(x) = n$, $T'_n \equiv T_m$ where $n := 1 + (n \mod \mathbf{d})$. Since \mathbf{d} is a period of e(u), then $T_m \equiv T_{n+1} = T < D, x >$, hence $T < D, x > \equiv T_{\varphi(x)}$. Suppose $x \le_e$. Then some iterate of f say g send g onto some g(x) with g(x) and g(x) and g(x) into g(x) and g(x) but then,

 $T < D, g(x) > \equiv T_{\varphi(g(x))}$. Since d is a multiple of \mathbf{d} , $\varphi(x)$ and $\varphi(g(x))$ are congruent modulo \mathbf{d} , hence $T_{\varphi(x)} \equiv T_{\varphi}(g(x))$. This yields $T < D, x > \leq T_{\varphi(x)}$. \square

Claim 5.15. T' is embeddable into T.

Proof of Claim 5.15.

Subclaim 5.16. There are $z, z', z'' \in e(u) \setminus \{u\}$ such that $z'' <_e z' <_e z$, $z'' \boxplus \mathbf{d} = z'$, $z' \boxplus d = z$, $z'' \vee_e f(z'') = z$, f(z') = z, $T < e(u), z >_{\equiv} T < e(u), z' >_{\equiv} T < e(u), z'' >$.

Proof of Subclaim 5.16. Set $z := x \vee f(x)$. Set $z' := x \boxplus k$ where k := v(z) - v(f(x)) and v is any e-valuation. According to Claim 5.5, we have f(z') = z. Since $z' \boxplus d = z$ and \mathbf{d} divides d, we have $T < e(u), z > \equiv T < e(u), z' >$. Since $z := x \vee f(x)$, we have $x <_e z'$, hence $z' = u \boxplus k'$ with $k' > \mathbf{d}$. Set $k'' := k' - \mathbf{d}$ and $z'' := u \boxplus k''$. Hence $z' = z'' \boxplus \mathbf{d}$. It follows that $T < e(u), z' > \equiv T < e(u), z'' >$. By construction, we have $z'' <_e z' <_e z$; since $u <_e z'' <_e z'$ we have $f(u) <_e f(z'') <_e f(z') = z$. From $z = x \vee_e f(x)$ we have $z = u \vee_e f(u)$, hence $z = z'' \vee_e f(z'')$. This proves our subclaim.

With this subclaim the proof of Claim 5.15 goes as follows.

Let h be an embedding of T < e(u), z > into f[T < e(u), z'' >]. Let H_0 be the path joining z and f(z''), set $H_{n+1} := h[H_n]$ and $H := \bigcup_{n \in \mathbb{N}} H_n$. Then H is an infinite path ending at z and $D := H \cup e(z)$ a two-way infinite path. We define an embedding ψ of the path on \mathbb{Z} onto the path D by setting $s := \mathbf{d} + k - 1$, and for $m \ge 0$, $\psi(s+m) = z \boxplus m$ and $\psi(s-m) = z_m$ where z_m is the unique member of H such that $z_m \boxplus m = z$. We check that T'_n is embeddable into $T < D, \psi(n) >$ for every $n \in \mathbb{Z}$. Let $m \ge 0$. Then $\varphi \circ \psi(s+m) = s+m$. Indeed, $\psi(s+m) = z \boxplus m = u \boxplus (s+m+1)$ and $\varphi(u \boxplus (s+m+1)) = s+m$. For m = 0, we have $T'_s \equiv T_{s+1} = T < e(u), z > \equiv T < e(u), z' >$. The map f embeds T < e(u), z' > into $T < D, z > \equiv T < D, \psi(s) >$. Hence T'_s embeds into $T < D, \psi(s) >$. For m > 0 we have $T'_{s+m} \equiv T_{s+m+1} = T < D, \psi(s+m) >$. Let $m' := m \mod d$. We have $T'_{s-m} \equiv T'_{s-m'} \equiv T_{s+1-m'} \equiv T_{s+1-m'-d} \le T_{s+1} < H, z_{m'} > \le T_{s+1} < H, z_m >$ and $T_{s+1} < H, z_m > \le T < D, \psi(s-m) >$.

The origin of C_o is an isomorphism invariant of T, we will call it the *origin* of T.

- 5.4. The space of ends. Let r be a vertex of T and let $\Omega_r(T)$ be the set of rays starting at r. The map ϕ_r which associate with each end e the unique ray e(r) belonging to e and starting at r is a bijective map of $\Omega(T)$ onto $\Omega_r(T)$. Using this, $\Omega(T)$ can be topologized as a subset of $V(T)^{\mathbb{N}}$, equipped with the product topology, with \mathbb{N} equipped with the discrete topology, as well as a subset of the Cantor space $\mathcal{P}(V(T))$. The major features of $\Omega(T)$ with respect to the problem we consider are the following:
- a) The embeddings of T acts on $\Omega(T)$; more specifically, if f is an embedding of T, then f defined by f(e) := end(f[C]) for some $C \in e$ is a continuous embedding of $\Omega(T)$.

- b) $\Omega(T)$ is topologically scattered (that is every subset contains an isolated point) if and only T is scattered in the sense that no subdivision of the binary tree is embeddable in T (See Theorem 6.10 and [10]).
- c) If T is locally finite, then $\Omega(T)$ is compact, in fact a Stone space (totally disconnected compact space) and $\Omega(T)$ is countable if and only if T is scattered (see [3]).

In somewhat more concrete terms, see also [3]: Let (T,r) be the tree T rooted at T. For every $s \in V(T)$ let $\Omega_{r,s}(T)$ be the set of rays in $\Omega_r(T)$ which contain s. The sets $\Omega_{r,s}(T)$ form a basis of a topology. This topology on $\Omega_r(T)$ is easily seen and of course well known to satisfy Item c). Let $r' \in V(T)$. Let P be the set of vertices on the path between r and r'. For $C \in \Omega_r(T)$ let C'' be the ray induced by the set $V(C) \setminus P$ of vertices. Let C' be the unique extension of C'' to a ray in $\Omega_{r'}(T)$ starting a r'. The map which associates with $C \in \Omega_r(T)$ the ray $C' \in \Omega_{r'}(T)$ is a homeomorphism. Both C and C' are elements of the same end of T. It follows that the set of ends of T inherits via ϕ_r^{-1} a topology homeomorphic to $\Omega_r(T)$ which indeed is independent of the particular vertex $r \in V(T)$.

Let f be an embedding mapping r to r'. Then f is an isomorphism of T to f[T]. Let C be a ray in $\Omega_r(T)$ contained in the end e. Then f[C] is a ray in $\Omega_{r'}(f[T])$ and also a ray in $\Omega_{r'}(T)$. The end of f[T] containing f[C] is a subset of the end, denoted f(e), of T containing f[C]. Hence, f is a homeomorphism of $\Omega_r(T)$ to $\Omega_{r'}(f[T])$. From this, Item a) follows.

This applies to the Cantor-Bendixson derivatives of $\Omega_r(T)$. Let us recall that if X is a topological space, we may denote by Isol(X) the set of isolated points of X and define for each ordinal α , the α -th-derivative $X^{(\alpha)}$ setting $X^{(0)} = X$, $X^{(\alpha)} = X^{(\alpha-1)} \setminus \text{Isol}(X^{(\alpha-1)})$ if α is a successor ordinal and $X^{(\alpha)} = \bigcap_{\beta < \alpha} X^{(\beta)}$ if α is a limit ordinal. The least α such that $X^{(\alpha)} = X^{(\alpha+1)}$ is the Cantor-Bendixson rank of X which we denote by rank(X). The rank or order of an element $x \in X$, denoted by rank(x,X) is the least ordinal α such that $x \notin X^{\alpha+1}$. If X is non-empty, rank(X) is a successor ordinal iff there is a last non-empty derivative, that we denote by $X^{(last)}$. As it is well-known, X is scattered iff $X^{(rank(X))} = \emptyset$; if furthermore, X is compact, then rank(X) is a successor ordinal and $X^{(last)}$, is finite. If $X = \Omega_r(T)$ we denote by $\Omega_r^{(\alpha)}(T)$ the α -th derivative of $\Omega(T)$ and if $rank(\Omega_r(T))$ is a successor ordinal we denote by $\Omega_r^{(last)}(T)$ the last non-empty derivative. For an example, if T is rayless, then $\Omega(T) = \emptyset$ hence then $rank(\Omega(T)) = 0$. If T is an infinite one way path or consists of a set of one-way infinite path originating from some vertex r but otherwise disjoint, then $\Omega_r^{(1)}(T)$ is empty and hence $rank(\Omega_r(T))$ is equal to one. Implying that $\Omega^{(1)}(T)$ is empty and the rank of $\Omega(T)$ is one.

Let f be an embedding of T. Let $r \in V(T)$, and r' := f(r). Since f is a homeomorphism of $\Omega_r(T)$ onto $\Omega_{r'}(f[T])$, a ray C is isolated in $\Omega_r(T)$ iff every f[C] is isolated in $\Omega_{r'}(f[T])$. Hence, the α -derivative of $\Omega_r(T)$ is mapped by f onto the α -derivative of $\Omega_{r'}(f[T])$ and hence the Cantor-Bendixson rank of $\Omega_r(T)$ is equal to the Cantor-Bendixson rank of $\Omega_r(f[T])$. Since f is a

continuous maps of $\Omega_r(T)$ into $\Omega_{r'}(T)$ it maps each $\Omega_r^{(\alpha)}(T)$ into $\Omega_{r'}^{(\alpha)}(T)$. Hence, with the identification of $\Omega(T)$ with $\Omega(T)$ and $\Omega_{r'}(T)$, f preserves each $\Omega^{(\alpha)}(T)$.

We give an illustration of the notion of rank in Subsection 5.6, particularly with a proof of the first part of Theorem 1.3. To do so we need to compute the rank for some scattered trees.

5.5. Operations on trees and computation of the rank. Let us define the following operations on rooted trees that we call respectively successor, sup and sum. If T is a rooted tree, 1+T is the rooted tree obtaining by adding a new vertex, say a, joined to the root of T, and by choosing a as a root of this new tree. If $(T_i)_{i\in I}$ is a family of rooted trees, then their supremum $\bigvee_{i\in I} T_i$ is the tree obtained by identifying all the roots of the T_i 's to a single one. If $(T_n)_{n\in\mathbb{N}}$ is a family of rooted trees, each T_n rooted at n, then $\bigoplus_{n\in\mathbb{N}} T_n$ is the tree rooted at 0 of the sum of the T_n 's over the infinite path on \mathbb{N} . If for some number n_0 the trees T_n with $n > n_0$ are all equal to the empty tree \square , then $\bigoplus_{n\in\mathbb{N}} T_n$ is the tree rooted at 0 of the sum of the T_n 's over the finite path on $\{0,1,\ldots,n_0\}$.

We relate the the rank with the operations sup and sum. For this we will need the following notion:

A sequence $(T_n)_{n\in\mathbb{N}}$ of trees has property (*) if:

- (*) $\inf\{\sup\{rank(\Omega(T_n)): m \leq n \in \mathbb{N}\}: m \in \mathbb{N}\} < \sup\{rank(\Omega(T_n)): n \in \mathbb{N}\}\}$. Note that if the sequence of trees T_n has property (*), then $\max\{rank(\Omega(T_n)): n \in \mathbb{N}\}$ exists and there is a largest number $n_0 \in \mathbb{N}$ with $rank(\Omega(T_{n_0})) = \max\{rank(\Omega(T_n)): n \in \mathbb{N}\}$ and with $rank(\Omega(T_{n_0})) > \sup\{rank(\Omega(T_n)): n_0 < n \in \mathbb{N}\}$. Note that (*) does not hold if and only if the following property (not (*)) holds:
- (not (*)) for every ordinal $\alpha < \sup\{rank(\Omega(T_n)) : n \in \mathbb{N}\}$ and every $n \in \mathbb{N}$ there is an $n < m \in \mathbb{N}$ with $\alpha < rank(\Omega(T_n)) \le \sup\{rank(\Omega(T_n)) : n \in \mathbb{N}\}.$

Lemma 5.17. Let $(T_i)_{i\in I}$ be a family of pairwise disjoint rooted trees. If each $\Omega(T_i)$ is topologically scattered, then $\Omega(\bigvee_{i\in I} T_i)$ is scattered and:

(4)
$$rank(\Omega(\bigvee_{i \in I} T_i)) = sup\{rank(\Omega(T_i)) : i \in I\}.$$

Let $(T_n)_{n\in\mathbb{N}}$ be a family of pairwise disjoint rooted trees, each T_n rooted at n If each $\Omega(T_n)$ is topologically scattered, then $\Omega(\bigoplus_{n\in\mathbb{N}} T_n)$ is topologically scattered and:

(5)
$$rank(\Omega(\bigoplus_{n\in\mathbb{N}} T_n)) = \begin{cases} \max\{rank(\Omega(T_n)) : n \in \mathbb{N}\}, & if (*) \ holds; \\ \sup\{rank(\Omega(T_n)) : n \in \mathbb{N}\} + 1, & if (*) \ does \ not \ hold. \end{cases}$$

Proof. Let $(T_i)_{i \in I}$ be a family of pairwise disjoint rooted trees and let T'_i be the tree arising from T_i by identifying its root with the other trees to obtain the root, say r, of $\Omega(\bigvee_{i \in I} T_i)$. Then T'_i is isomorphic to T_i . Let S be a non-empty

subset of $\Omega(\bigvee_{i\in I} T_i)$. Then there is an $i\in I$ for which $S\cap \Omega_r(T_i')$ is not empty and hence, if $\Omega_r(T_i')$ is scattered contains an isolated chain C of $\Omega_r(T_i')$. This chain C is isolated in $\Omega(\bigvee_{i\in I} T_i)$.

Equation (4): Let $T := \bigvee_{i \in I} T_i$. Observe that $\Omega_r(T)$ is the union of the sets $\Omega_r(T_i)'$ of chains and that these sets are pairwise disjoint. Each $\Omega_r(T_i')$ is a clopen subset of $\Omega(T)$. The result follows.

Let $T := \bigoplus_{n \in \mathbb{N}} T_n$ and $\gamma := \sup\{rank(\Omega_n(T_n)) : n \in \mathbb{N}\}$. For $n \in \mathbb{N}$ denote by T'_n the tree obtained by adding to T_n the path from 0 to n rooted at 0. Let S be a non-empty subset of $\Omega(T)$. Clearly, $\Omega_0(T)$ is the union of the $\Omega_0(T'_n)$ plus the path on \mathbb{N} . Each $\Omega_0(T'_n)$ is a clopen set homeomorphic to $\Omega_n(T_n)$. Hence $\gamma := \sup\{rank(\Omega(T_n)) : n \in \mathbb{N}\} = \sup\{rank(\Omega(T'_n)) : n \in \mathbb{N}\}$ and

(6)
$$\gamma \leq rank(\Omega_0(T)) \leq \gamma + 1$$

and there is an $n \in \mathbb{N}$ with $S \cap \Omega_0(T'_n) \neq \emptyset$ or $S = {\mathbb{N}}$. If $\Omega(T_n)$ is scattered there exists a chain C isolated in $\Omega_0(T'_0)$ which then is also isolated in T.

Property (*) holds: Then $rank(\Omega(T_{n_0})) > \sup\{rank(\Omega(T_n)) : n_0 < n \in \mathbb{N}\} := \delta$ and $\gamma = rank(\Omega(T_{n_0}))$ and every path in $\Omega_0^{(\delta)}(T)$ is a path in $\Omega_0^{(\delta)}(T'_{n_0})$, which is not empty because $\delta < rank(\Omega(T_{n_0}))$. Thus $rank(\Omega_0(T)) = rank(\Omega_0(T_{n_0})) = \max\{rank(\Omega(T_n)) : n \in \mathbb{N}\}.$

Property (*) does not hold: Then γ is not attained or it is attained infinitely many times. In either case the path on \mathbb{N} belongs to $\Omega_0^{(\gamma)}(T)$. With inequality (6) this yields $rank(\Omega_0(T)) = \gamma + 1$, thus the result.

5.6. A set of ends preserved.

Lemma 5.18. If a tree T is infinite, locally finite and scattered, then there is a non-empty finite subset C of $\Omega(T)$ which is preserved by every embedding of T.

Proof. The proof will follow from the following claims.

The space $\Omega(T)$ is non-empty by Kőnig's lemma [11]. Since T is scattered, $\Omega(T)$ is scattered; since it is compact its rank is a successor ordinal and $\Omega(T)^{(last)}$ is finite. Set $\mathcal{C} := \Omega^{(last)}(T)$.

If T is not locally finite, the rank can be a limit ordinal and even if it is a successor ordinal the set $\Omega^{(\infty)}(T)$ is not necessarily finite. If T is scattered but not necessarily locally finite, Theorem 1.1 gives an extension of Lemma 5.18 whose proof is presented in Section 7.1.

Proposition 5.19. Let f be an embedding of a tree T. Suppose that there is a two-way infinite path D preserved by f on which f fixes no vertex and reverses no edge, or if not, a one-way infinite path C preserved by f with a vertex not in the range of f.

In the first case, set $\mathcal{J} := \{end(D^-), end(D^+)\}$ where D^- and D^+ are two paths whose union is D and such that $f[D^+] \subseteq D^+$ and set $\mathcal{J} := \{end(C)\}$ in the second case. If T is scattered, then $rank(\Omega(T))$ is a successor ordinal and

in the first case $end(D^+) \in \Omega^{(last)}(T) \subseteq \mathcal{J}$, whereas $\Omega^{(last)}(T) = \mathcal{J}$ in the second case.

Proof.

Claim 5.20. Every end $e \notin \mathcal{J}$ has an infinite orbit under f. In particular the members of \mathcal{J} are the only ends preserved by f.

Proof of Claim 5.20. In the first case take E = D and in the second case take E equal to a maximal one way path (in fact a maximum one) preserved by f. For $x \in E$, set $T_x := ((T, E)(x), x)$. We have $T = \bigoplus_{x \in E} T_x$ and the map f induces an embedding of T_x into $T_{f(x)}$ for every $x \in V(E)$. Apply Corollary 4.4 and let $e \notin \mathcal{J}$ and $R \in e$. Then R has all but finitely many of its vertices in one of the rooted trees T_x . Then f[R] has all but finitely many of its vertices in $T_{f(x)}$. Because $x \notin f(x)$ the rooted trees T_x and $T_{f(x)}$ have no vertex in common, implying that the end f(e) containing f[R] is different from e. Similarly $f^{(n)}(e)$ is different from $f^{(m)}(e)$ for $n \notin m$.

Claim 5.21. Let $\alpha := \sup\{rank(\Omega_x(T_x)) : x \in E\}$. Then, in the first case $e(D^+) \in \Omega^{(\alpha)}(T) \subseteq \mathcal{J}$, whereas $\Omega^{(\alpha)}(T) = \mathcal{J}$ in the second case. In particular, $rank(\Omega(T)) = \alpha + 1$.

Proof of Claim 5.21. Let $\alpha := \sup\{rank(\Omega_x(T_x)) : x \in E\}$. Since f embeds each T_x into $T_{f(x)}$ for all $x \in E$, $rank(\Omega_x(T_x)) \le rank(\Omega_{f(x)}(T_{f(x)}))$. Let $x, y \in E$. Set $x \le_E y$ if the ray originating at x and belonging to $end(D^+)$ contains y. Since $x \le f(x)$, we have $\sup\{rank(\Omega_y(T_y)) : x \le_E x\} = \alpha$. Let $T_{\ge x} := \bigoplus_{y \in E_x} T_y$, where $E_x := \{y \in E : x \le y\}$.

We claim that $rank(\Omega_x(T_{\geq x})) = \sup\{rank(\Omega_y(T_y)) : y \geq x\} + 1 = \alpha + 1$, a fact which follows from Lemma 5.17. Indeed, label the vertices of E_x as $x_0, \ldots x_n, \ldots$ in an increasing order, set $T'_n := T_x$ and observe that $(T'_n)_{n \in \mathbb{N}}$ does not have property (*). According to (2) of Lemma 5.17, $rank(\Omega(T_{\geq x})) = \sup\{rank(\Omega(T_y)) : x \leq_E y\} + 1 = \alpha + 1$; proving our claim.

Let $D_{\geq x} := D \cap \{y : x \leq_E y\}$. Since $D_{\geq x}$ meets no T_y into infinitely often, it follows from our claim that $\Omega^{(\alpha)}(T_{\geq x}) = \{end(D_{\geq x})\}$. If E is a ray, then for the least element x of E we have $T = T_{\geq x}$, hence $\Omega(T)^{\alpha}) = \mathcal{J}$ as claimed. If E is a double ray, then, since each $\Omega(T_{\geq x})$ is closed into $\Omega(T)$, $rank(\Omega(T_{\geq x})) \leq rank(\Omega(T))$, we infer $\alpha + 1 \leq rank(\Omega(T))$. Furthermore, $\Omega(T) = \bigcup_{x \in E} \Omega(T_{\geq x}) \cup \{end(D^-)\}$, hence we have $end(D^+) \in \Omega^{(\alpha)}(T) \subseteq \mathcal{J}$. In particular, $rank(\Omega^{(\alpha)}(T)) = \alpha + 1$. This proves Claim 5.21.

With this Claim, the proof of the Proposition 5.19 is complete

Remark 5.22. In the first case of Proposition 5.19, we have $rank(end(C^-), \Omega(T)) \leq rank(end(C^+), \Omega(T))$.

This inequality can be proved directly. Let $e := end(C^-)$, $e' := end(C^+)$ and $\alpha := rank(e, \Omega(T))$. Then every neighbourhood of e contains an element e_{β} of $\Omega^{(\beta)}(T)$ for every $\beta < \alpha$. Let U' be a neighbourhood of e' in $\Omega(T)$. Some iterate of f will map e_{β} into some element $e'_{\beta} \in U' \cap \Omega^{(\beta)}(T)$. This implies that $e' \in \Omega^{(\beta)}(T)$. The inequality follows.

Corollary 5.23. If a tree T is scattered, then $rank(\Omega(T))$ is a successor ordinal and $\Omega^{(last)}(T)$ has at most two elements provided that some embedding f does not fix a vertex or an edge.

Proof. By Proposition 3.3, such an f satisfies the hypothesis of Proposition 5.19.

Corollary 5.24. Let T be a scattered tree. If $rank(\Omega(T))$ is a successor ordinal, $|\Omega^{(last)}(T)| = 1$ and $e \in \Omega^{(last)}(T)$, then e is preserved forward by every embedding of T.

Proof. Let f be an embedding of T. Then f preserves $\Omega^{(last)}(T)$. Since $\Omega^{(last)}(T) = \{e\}$, f preserves e. Suppose that e is not preserved forward by f. That is there is some ray $C \in e$ such that no infinite subray C' of C is sent into C' by f. In this case, some subray, say C', extends to a two-way infinite path D which is preserved by f. Indeed, pick a subray C' of C such that $C' \subset f(C')$ and $D := \bigcup_{n \in \mathbb{N}} f^{(n)}(C)$. We may write $D := \{x_n : n \in \mathbb{Z}\}$ in such a way that $C' = D^-$ and $f(x_n) = x_{n+k}$ for some positive k. Let $e' := end(D^+)$. According to Proposition 5.19 we have $e' \in \Omega^{(last)}(T)$. Since $e \neq e'$, this is impossible. \square

- 5.7. **Proof of Theorem 1.3.** According to Corollary 5.24, if T contains exactly one end of maximal rank, then this end is preserved forward. Next, if e contains some regular end e preserved forward by every embedding, then e contains some ray preserved forward provided that T is scattered and e almost rigid (Proposition 5.8) or e is not almost rigid (Proposition 5.12).
- 5.8. **Examples of ends preserved.** We present six examples. The first four examples are scattered trees containing exactly one end preserved by every embedding.

In the first three examples there are embeddings with no fixed point. The third example is due to Polat and Sabidussi [16]. In the fourth example the end is "almost fixed" by every embedding. First a notation: let κ be a cardinal, we set $\bigvee_{\kappa} T$ for the tree sup of κ copies of T, that is the tree T is rooted and the copies are disjoint, and we glue the copies together at their roots. For an example, if T is a 1-element tree, then $\bigvee_{\kappa} T = T$.

Example 1. Let $(T_n^1)_{n\in\omega}$ be the sequence of finite trees defined by induction as follows: T_0^1 is the tree consisting of one vertex rooted at 0; $T_{n+1}^1 := 1 + \bigvee_2 T_n^1$ rooted at n+1, (hence, $|T_n^1| = 2^n$). Set $T_\omega^1 := \bigoplus_{n<\omega} T_n^1$. This tree has an interesting reproducing property: if we delete the end vertices of this tree, we get an isomorphic copy.




FIGURE 3. Example 1: T^1_{ω}

Example 2. Let $(T_n^2)_{n\in\omega}$ be the sequence of trees defined by induction as follows: T_0^2 is the tree consisting of one vertex rooted at 0; $T_{n+1}^2 := \bigvee_{\aleph_0} (1 + T_n^2)$ rooted at n+1, $T_\omega^2 := \bigoplus_{n<\omega} T_n^2$.




FIGURE 4. Example 2: T_{ω}^2

Example 3. Let $(T_n^3)_{n\in\omega}$ be the sequence of trees defined by induction as follows: T_n^3 for n=0,1 is the rooted tree consisting of an infinite path rooted at n; for larger n, $T_{n+1}^3 := 1 + \bigvee_2 (1 + T_n^3)$ rooted at n+1, $T_\omega^3 := \bigoplus_{n<\omega} T_n^3$. (See Fig 2 in Polat and Sabidussi [16].) Equivalently, T_ω^3 is obtained from T_ω^1 by replacing each terminal vertex of T_ω^1 by an infinite path rooted at this vertex. We could as well replace each terminal vertex by two infinite paths rooted at this vertex.




FIGURE 5. Example 3: T_{ω}^3

Example 4. Let T^4_{ω} be the tree obtained by replacing each terminal vertex of T^1_{ω} by the rooted tree $\bigvee_3 2$.




FIGURE 6. Example 4: T_{ω}^4

In all these examples, the non-negative integers form a ray. This ray is not regular, hence its end, say e is not regular. This end is preserved by all embeddings. In the first two examples, this is the only end, hence the rank of the space of ends is 1. This end is preserved forward, but no ray containing

it is preserved. In the third example, the rank of the space of ends is 2. In the first three examples there are embeddings which move the path forward. Hence $|twin(T)| \ge 2^{\aleph_0}$ according to Theorem 1.9. This is not the case in the fourth example. Indeed, there the end is almost rigid (any embedding of T^4_{ω} preserves the level function). Because T^4_{ω} is locally finite, every embedding is an automorphism.

The next two examples are trees containing exactly one end preserved by every embedding, this end being preserved backward.

Example 5. To the rooted binary tree add an infinite ray at the root r (See Figure 3). Let T^5 be the resulting tree. Label the vertices from the root by integers as follows: The root has label 0. The label of a vertex x in the binary tree is the distance from x to r. The label of a vertex x in the ray added is -n if the distance from r is n. To each vertex labelled 0 modulo 3, graft in ten leaves and to each vertex labelled 1 modulo 3 graft in five leaves. For degree reasons, if f is an embedding, then either the ray added is fixed or it is moved backward.




FIGURE 7. Hamann's Example 5

Example 6. In the ternary tree, select an infinite ray, say 0, 1, 2, ..., n, ... (See Figure 8). Subdivide each edge on this ray by adding two vertices. Subdivide all other edges $w := \{u, v\}$ (outside that ray) whose distance from w to that ray is even by adding a vertex. Hence edges touching the ray are subdivided, edges a step further are not subdivided, etc. Let T^6 be the resulting tree. In this tree, the infinite ray with the vertices added is either fixed or moved backward.




FIGURE 8. Lehner's Example 6

6. Scattered trees

In this section we present some characterizations and constructions of scattered trees, e.g. Lemma 6.7 and Theorem 6.10, going back to Laver [13] and Jung [10].

6.1. Scattered trees versus non-scattered trees. Let T be a non scattered tree. The non-scattered kernel of T, written Ker(T), is the union of subsets of the form V(T') where T' is a subtree of T, which contains a subdivision of T_2 .

Notation 6.1. Let T be a tree, $x \in V(T)$ and $y \in V(T) \setminus \{x\}$, then $T \setminus \{x\}(y)$ stands for the connected component of y in the graph $T \setminus \{x\}$.

We have the following decomposition result, whose proof is immediate:

Theorem 6.2. Let T be a non scattered tree. The non-scattered kernel of T is a non-scattered tree such that Ker(Ker(T)) = Ker(T) and (T, Ker(T))(x) is scattered for every $x \in Ker(T)$.

Remark 6.3. Ker(T) is the set of vertices $x \in V(T)$ having at least two distinct neighbours y and y' such that the connected components $T \setminus \{x\}(y)$ of y and $T \setminus \{x\}(y')$ of y' in $T \setminus \{x\}$ are non-scattered.

Indeed, if x is such a vertex, then the tree that is the union of a subdivision of the binary tree taken in $T \setminus \{x\}(y)$ and a subdivision of the binary tree taken in $T \setminus \{x\}(y')$, with a path connecting these two subdivision, is a subdivision of T_2 containing x, hence $x \in Ker(T)$. The converse is immediate: If $x \in Ker(T)$ and T' is a subtree of T which is a subdivision of the binary tree containing x, then x has two neighbours y and y' in T' such that the connected components of y and y' in $T' \setminus \{x\}$ are non-scattered. Hence these connected components in T are non-scattered too.

Corollary 6.4. The non-scattered kernel of a non-scattered tree is preserved by every embedding.

Problem 6.5. If there is some embedding f with $Ker(T) \neq f(Ker(T))$, does T have infinitely many twins?

Proposition 6.6. Let T be a non scattered tree and $x \in Ker(T)$ and T_x be the rooted tree ((T, Ker(T))(x), x). If T_x has κ twins, then T has at least κ twins.

Proof. Let K := Ker(T). Let (T', x) be a twin of T_x and let $S := \{z \in K : ((T, K)(z), z) \equiv ((T, K)(x), x)\}$. Let T(T') be the tree obtained by replacing each ((T, K)(z), z) for $z \in S$ by (T', x).

Then T(T') is equimorphic to T and Ker(T(T')) = K. We claim that if (T',x) and (T'',x) are two twins of ((T,Ker(T))(x),x) which are not isomorphic (as rooted trees), then the trees T(T') and T(T'') are not isomorphic and this will prove the proposition. Suppose by contradiction that there is some isomorphism f from T(T') onto T(T''). Clearly f carries the kernel K of T(T') onto the kernel K of T(T'') and we have $((T(T''),K)(f(z)),f(z)) \simeq ((T(T'),K)(z),z)$. Let $z \in S$. We have by construction $((T(T''),K)(z),z) \simeq (T',x)$. Hence $((T(T''),K)(f(z)),f(z)) \simeq (T',x)$. If $f(z) \notin S$, then ((T(T''),K)(f(z)),f(z)) is not equimorphic to (T'',x) which contradicts the fact that (T',x) and (T'',x) are equimorphic. Hence $f(z) \in S$ and hence $((T(T''),K)(f(z)),f(z)) \simeq (T'',x)$, from which follows that (T',x) and (T'',x) are isomorphic.

6.2. Scattered trees as posets. Let T be a tree, $x \in V(T)$ and P(T,x) be the poset associated with the rooted tree (T,x). For an example, if T_2 is the binary tree and the root is the unique vertex of degree 2, we have the dual of the ordered binary tree.

Lemma 6.7. Let T be a tree and $x \in V(T)$. Then the following properties are equivalent:

- (i) T is scattered;
- (ii) The dual of the ordered binary tree does not embed into P(T,x) as a poset.
- **Proof.** $\neg(ii) \Rightarrow \neg(i)$. Suppose that there is an embedding f of the dual of the ordered binary tree into P(T,x). We construct an embedding g of a subdivision of T_2 into T and hence T is not scattered. For that, we transform first f to a map \check{f} from $2^{<\mathbb{N}}$ into P(T,x). We set $\check{f}(\square) := f(0) \vee f(1)$ and more generally $\check{f}(s) := f(s^{\circ}0) \vee f(s^{\circ}1)$. This map satisfies $\check{f}(s \wedge t) = \check{f}(s) \vee \check{f}(t)$. Now, if for each $s \in 2^{<\mathbb{N}}$ we add the shortest path between $\check{f}(s)$ and $\check{f}(s^{\circ}i)$, for $i \in 2$, the resulting graph is a subdivision of T_2 .
- $\neg(i) \Rightarrow \neg(ii)$. If T is not scattered, let K be a subtree of T which is a subdivision of the binary tree. Let x be a vertex of K having degree 2. Then the dual of the ordered binary tree T_2 embeds into the poset P(T, x).

6.3. Construction of scattered trees. We describe the collection of rooted trees such that $rank(\Omega_r(T)) \leq \alpha$. For that, we need the following notation. Let \mathcal{X} be a collection of rooted trees. We set $\bigoplus_{n\in\mathbb{N}} \mathcal{X}$ for the collection of rooted trees $\bigoplus_{n\in\mathbb{N}} T_n$ such that $T_n \in \mathcal{X}$ for every $n \in \mathbb{N}$, with the convention that this class is reduced to the empty tree if \mathcal{X} is empty. Note that if the empty tree $\Box \in \mathcal{X}$, then \bigoplus -sums over finite paths are in \mathcal{X} and hence $\mathcal{X} \subseteq \bigoplus_{n\in\mathbb{N}} \mathcal{X}$. We set $\mathcal{X}^{suc,\sup}$ for the closure of \mathcal{X} by means of the operations successor and sum. Hence $\mathcal{X}^{suc,\sup}$ is the smallest class \mathcal{Y} of rooted trees which contains \mathcal{X} and such that $T \in \mathcal{Y}$ provided that $T' \in \mathcal{Y}$ if T = 1 + T' or all $T_i \in \mathcal{Y}$ if $T = \bigvee_{i \in I} T_i$. We obtain from Equation (4) that if $rank(\Omega(T)) \leq \alpha$ for all $T \in \mathcal{X}$, then $rank(\Omega(T)) \leq \alpha$ for all $T \in \mathcal{X}^{suc,\sup}$.

Lemma 6.8. Let \mathcal{X} be a collection of rooted trees. Then $\mathcal{X}^{suc, sup}$ is the collection $\mathcal{R}(\mathcal{X})$ of rooted trees for which there exists a rayless rooted tree R with members of \mathcal{X} attached at the endpoints of R. For R empty the tree in $\mathcal{R}(\mathcal{X})$ is a member of \mathcal{X} .

Proof. Clearly $\mathcal{X} \in \mathcal{R}(\mathcal{X})$. Applying the operation + to a tree in $\mathcal{R}(\mathcal{X})$ or the operation $\bigvee_{i \in I}$ to a set of trees $(T_i)_{i \in I}$ results in a tree in $\mathcal{R}(\mathcal{X})$. \square We define inductively a collection $\mathcal{D}_r^{(\alpha)}$ of rooted trees for each ordinal α setting first $\mathcal{D}_r^{(0)} := \{\square\}^{suc, \text{sup}}$. We set $\mathcal{D}_r^{(\alpha)} := (\bigcup_{\beta \in \alpha} \mathcal{D}_r^{(\beta)})^{suc, \text{sup}}$ if α is a non-zero limit ordinal. We set $\mathcal{D}_r^{(\alpha)} = (\bigoplus_{n \in \mathbb{N}} \mathcal{D}_r^{(\beta)})^{suc, \text{sup}}$ where β is such that $\alpha = \beta + 1$. We denote by \mathcal{D}_r the union of all $\mathcal{D}_r^{(\alpha)}$ and by \mathcal{D} the collection of trees obtained

by forgetting the roots of members of \mathcal{D}_r . Unless otherwise stated we assume tacitly that r is the name of the root of the elements of \mathcal{D}_r .

Lemma 6.9. For each ordinal α and $T \in \mathcal{D}_r^{(\alpha)}$, the space $\Omega_r(T)$ is topologically scattered and $rank(\Omega_r(T)) \leq \alpha$.

If $T \in \mathcal{D}_r^{(\alpha)}$ and $y \in V(T)$ and \leq_r is the order of P(T,r) with r as largest element, then the subtree of T induced by the set $\{x \in V(T) : x \leq_r y\}$ of vertices rooted at y is an element of $\mathcal{D}_r^{(\alpha)}$.

Proof. The proof goes by induction on α . Suppose $\alpha = 0$. $\mathcal{D}_r^{(0)} := (\square)^{suc, \text{sup}}$ is just the collection of rayless trees. These are the trees such that $\Omega(T) = \emptyset$, hence they are scattered and $rank(\Omega(T)) = 0$. Suppose that the Lemma holds for all $\beta < \alpha$. If α is a limit ordinal, then induction asserts that for each $T \in \bigcup_{\beta < \alpha} \mathcal{D}_r^{\beta}$, $rank(\Omega(T)) < \alpha$. Since $\mathcal{D}_r^{\alpha} = (\bigcup_{\beta < \alpha} \mathcal{D}_r^{\beta})^{suc, \text{sup}}$, the Equation (4) of Lemma 5.17 ensures that $rank(\Omega(T)) \leq \alpha$ for every $T \in \mathcal{D}_r^{\alpha}$. Suppose that α is a successor ordinal and let β such that $\alpha = \beta + 1$. We have $\mathcal{D}_r^{(\alpha)} = (\bigoplus_{n \in \mathbb{N}} \mathcal{D}_r^{(\beta)})^{suc, \text{sup}}$. Induction asserts that $rank(\Omega_r(T)) \leq \beta$ holds for each $T \in \mathcal{D}_r^{(\beta)}$). According to Equation (5) of Lemma 5.17, $rank(\Omega_r(T)) \leq \beta + 1 = \alpha$ for every T of $(\bigoplus_{n \in \mathbb{N}} \mathcal{D}_r^{(\beta)})$. With Equation (4) we obtain that $rank(\Omega_r(T)) \leq \alpha$ holds for each $T \in \mathcal{D}_r^{(\beta)}$. The induction process verifies via Lemma 5.17 the claim that $\Omega(T)$ is topologically scattered for

all trees $T \in \mathcal{D}$. The second statement obviously for trees in $\mathcal{D}_r^{(0)}$ and then is easily seen to hold throughout the induction process.

Theorem 6.10. Let T be a tree. Then the following properties are equivalent:

- (i) T is scattered;
- (ii) $\Omega(T)$ is topologically scattered;
- (iii) $T \in \mathcal{D}$.

Proof. We proceed by showing that $(i) \Rightarrow (iii) \Rightarrow (ii) \Rightarrow (i)$.

 $\neg iii \Rightarrow \neg i$. Let $T \notin \mathcal{D}$. Choose some vertex in V(T) and name it r. The rooted tree (T, r) is not in \mathcal{D}'_r because $T \notin \mathcal{D}$. Let P(T, r) be the corresponding poset with order denoted by \leq_r . Let F be the set of $y \in V(T)$ for which the set $\{x \geq_r y : x \in V(T)\}$ induces a subtree of T which is an element of \mathcal{D}_r with root at y. We claim that there are two incomparable elements y_0 and y_1 in $V(T) \setminus F$. For otherwise, $V(T) \setminus F$ reduces to a path and T is a sum over this path of members of \mathcal{D}_r hence belongs to \mathcal{D}_r . Applying this property repeatedly we get an embedding of the dual of the ordered binary tree and hence, from Lemma 6.7, a subdivision of the binary tree.

 $(iii) \Rightarrow (ii)$. According to Lemma 6.9, for each ordinal α , and each rooted tree $T \in \mathcal{D}_r^{(\alpha)}$, $\Omega_r(T)$ is topologically scattered.

 $\neg i \Rightarrow \neg ii$. Let A be a subtree of T isomorphic to a subdivision of T_2 . Let r be the root of A (a vertex of degree 2 within A.) Then $\Omega(A, r)$ identifies to the Cantor space, hence is not topologically scattered. (No one-way path in $\Omega_r(A)$ for example is isolated in $\Omega_r(A)$ and hence not isolated in $\Omega_r(T)$.) Or, it is easy to see that $\Omega_r(A)$ is a closed subspace of $\Omega_r(T)$, hence this space is not topologically scattered.

Proposition 6.11. For each ordinal α a tree T is an element of $\mathcal{D}_r^{(\alpha)}$ if and only if $\Omega_r(T)$ is topologically scattered and $rank(\Omega_r(T)) \leq \alpha$.

Proof. The direct implication is Lemma 6.9. Suppose that the reverse implication does not hold. Then there is a minimum ordinal α such that for some rooted tree T, the set $\Omega_r(T)$ of one-way paths is topologically scattered and $rank(\Omega_r(T)) \leq \alpha$ but $T \notin \mathcal{D}_r^{(\alpha)}$. In this case $rank(\Omega_r(T)) = \alpha$. Let P(T,r) be the poset associated with T, which has r as largest element and the order denoted by \leq_r . Let F be the set of vertices $y \in V(T)$ such that the subtree of T induced by the set $\{x \leq_r y : x \in V(T)\} := \downarrow x$ of vertices and rooted at y is in $\mathcal{D}_r^{(\alpha)}$. Since $T \notin \mathcal{D}_r^{(\alpha)}$ the vertex $r \notin F$ and hence F is a proper initial segment of P(T,r) according to the second claim of Lemma 6.9. Note that for $x \in V(T)$ the rank of the set of ends of the subtree induced by the set of vertices $\downarrow x$ is less than or equal to α .

We claim that for every $x \in V(T) \setminus F$ there are two incomparable elements y_0 and y_1 in $V(T) \setminus F$ with $y_0, y_1 \leq x$. For otherwise the initial segment $(\downarrow x) \setminus F$ is a chain in the poset P(T, r), hence there is a path $C := x = x_0 \sim \cdots \sim x_n \sim \ldots$ Let Y_n be the set of neighbours y of x_n which are distinct from x_{n+1} . Note that $Y \subseteq F$. Hence the connected component $T \setminus \{x_n\}(y)$

of $T \setminus \{x_n\}$ containing y is an element in $\mathcal{D}_r^{(\alpha)}$ rooted at y. It follows that the tree $T'_n = ((T,C)(x_n),x_n)$ is an element of $\{T \setminus \{x_n\}(y) : y \in Y_n\}^{suc,\sup}$ and hence an element of $\mathcal{D}_r^{(\alpha)}$. (Remember here that $(T,C)(x_n)$ is the connected component of x_n in $T \setminus (C \setminus \{x_n\})$. And then T'_n is this tree rooted at x_n .) Let T' be the tree induced by $\downarrow x$ rooted at x. If C is finite it is easily seen that then the tree T' is an element of $\{T'_n : x_n \in V(C)\}^{suc,\sup}$ and hence an element of $\mathcal{D}_r^{(\alpha)}$, contradicting the stipulation $x \in V(T) \setminus F$.

Hence C is an infinite path and the tree T' can be written as $T' = \bigoplus_{\{x_n:n\in\mathbb{N}\}} T'_n$. Let α_n be the rank of $\Omega_{x_n}(T'_n)$. Via the minimality of α , we have $rank(\Omega_x(T')) = \alpha$. Each T'_n belongs to $\mathcal{D}_r^{(\alpha)}$. We apply Equation (5). The set M of integers n such that $\alpha_n = \alpha$ must be finite for otherwise $rank(\Omega_x(T')) = \alpha + 1$. Let $n_0 = 0$ if there is no n with $\alpha_n = \alpha$, and otherwise let $n_0 = \max M$. Let T'' be the tree induced by $\downarrow x_{n_0+1}$ rooted at x_{n_0+1} . By construction $T'' \notin \mathcal{D}_r^{(\alpha)}$ but it can be written as $T'' = \bigoplus_{\{x_n:n_0< n\in\mathbb{N}\}} T'_n$. If $\sup\{\alpha_n: n_0 < n \in \mathbb{N}\} = \alpha$, then because $\alpha_n < \alpha$ for all $n_0 < n \in \mathbb{N}$, the sequence $(T'_n)_{n_0< n\in\mathbb{N}}$ does not have property (*). This is not possible because then, according to Equation (5), we would have $rank\Omega(T'') = \alpha + 1$; implying $\sup\{\alpha_n: n_0 < n \in \mathbb{N}\} = \beta < \alpha$. Hence $T'' \in \mathcal{D}_r^{(\beta+1)} \subseteq \mathcal{D}_r^{(\alpha)}$. Again a contradiction.

Hence indeed, for every $x \in V(T) \setminus F$ there are two incomparable elements y_0 and y_1 in $V(T) \setminus F$ with $y_0, y_1 \leq x$. But then T embeds a copy of the dual of the ordered binary tree, and hence, from Lemma 6.7, a subdivision of the binary tree, contradicting the fact that $\Omega_r(T)$ is topologically scattered.

Corollary 6.12. For each ordinal α there is a scattered tree whose space of ends has ordinal rank α .

7. An extension of a result of Polat and Sabidussi

We extend Theorem 3.1 of [16].

Lemma 7.1. Let T be a scattered tree such that $\alpha := rank(\Omega(T))$ is a non-zero limit ordinal and $x \in V(T)$. Then the following properties are equivalent:

- (i) Either x has at least two neighbours y, y' such that $rank(\Omega(T \setminus \{x\}(y))) = rank(\Omega(T \setminus \{x\}(y'))) = \alpha$ or $sup\{rank(\Omega(T \setminus \{x\}(y))) : rank(\Omega(T \setminus \{x\}(y))) < \alpha \text{ and } y \sim x\} = \alpha.$
- (ii) There are two disjoint sets X_1 and X_2 of neighbours of x such that $\alpha = \sup\{rank(\Omega(T \setminus \{x\}(y))) : y \in X_1\} = \sup\{rank(\Omega(T \setminus \{x\}(y))) : y \in X_2\}.$

For $rank(\Omega(T))$ a non-zero limit ordinal let Lim(T) be the set of vertices $x \in V(T)$ such that (i) or (ii) holds. We list some simple properties of Lim(T).

Lemma 7.2. For $\alpha = rank(\Omega(T))$ a non-zero limit ordinal and $a \in V(T)$:

(1) $a \in V(T) \setminus Lim(T)$ iff there is a vertex $y \in V(T)$ such that $y \sim a$ and $rank(\Omega(T \setminus \{a\}(y))) = \alpha$ and some ordinal $\beta < \alpha$ such that $rank((\Omega(T \setminus \{a\})(y')) \leq \beta$ for every other $y' \sim a$.

- (2) If $a \in Lim(T)$ and $a \sim y$ and $rank(\Omega(T \setminus \{a\}(y))) = \alpha$, then $y \in Lim(T)$.
- (3) $Lim(T) = \{a\}$ iff $rank(\Omega(T \setminus \{a\}(y))) < \alpha$ for every $y \sim a$.
- (4) Let $a \in Lim(T)$. Then a has degree 1 in Lim(T) iff there is a unique $y \sim a$ such that $rank(\Omega(T \setminus \{a\}(y))) = \alpha$ and $sup\{rank(\Omega(T \setminus \{y\}(y'))) : y' \sim y \text{ and } y' \neq a\} = \alpha$.

Proof. Item (1). Let $a \in V(T)$. For each neighbour y of a, let T'(y, x) be the rooted tree obtained from $T \setminus \{a\}(y)$ by adding x as a new node and a root. Then $T = \bigvee_{y \approx a} T'(y, x)$, hence by Equation 4:

```
\alpha = rank(\Omega(T)) = \sup\{rank(\Omega(T'(y,x)) : y \sim a\} = \sup\{rank(\Omega(T \setminus \{a\}(y))) : y \sim a\}.
```

- Let $S(a) := \{y \sim a : rank(\Omega(T \setminus \{a\}(y))) < \alpha$. If a has either no neighbour outside S(a) or at least two neighbours outside S(a), then $a \in Lim(T)$. If a has just one neighbour outside S(a), then $a \in Lim(T)$ iff $\sup\{rank(\Omega(T \setminus \{a\}(y))) : y \sim a\} = \alpha$.
- Item (2). We apply Item (1). If $y \notin Lim(T)$ there is a vertex $z \sim y$ such that $rank(\Omega(T \setminus \{y\}(z))) = \alpha$ and some ordinal $\beta < \alpha$ such that $rank(\Omega(T \setminus \{a\}(y'))) \leq \beta$ for every other $y' \sim y$. If $z \neq a$, then $rank(\Omega(T \setminus \{y\}(a))) < \alpha$; but then $a \notin Lim(T)$, a contradiction. Hence z = a and $rank(\Omega(T \setminus \{z\}(y))) \leq \beta < \alpha$.
- Item (3) This is clear.
- Item (4) Suppose that a has degree 1 in Lim(T). The only neighbour y of a in Lim(T) has the stated property.

Lemma 7.3. Let T be a scattered tree such that $\alpha := rank(\Omega(T))$ is a non-zero limit ordinal. Then:

- a) Lim(T) is preserved by every embedding of T.
- b) Lim(T) is a non-empty rayless subtree of T.
- **Proof.** a) Let $a \in Lim(T)$ and f an embedding of T. Ther $rank(\Omega_a(T \setminus \{a\}(x)))rank(\Omega_{f(a)}(T \setminus \{f(a)\}(f(x)))$ for all $x \sim a$. Which with (i) of Lemma 7.1, implies that $f(a) \in Lim(T)$.
- b) We prove first that Lim(T) is rayless. Let $C := a_0, \ldots, a_n, \ldots$ be a one-way infinite path. We claim that C is not contained in Lim(T). Indeed, we have $T = \bigoplus_{a_n \in C} (T, C)(a_n)$. Since α is a limit ordinal, Lemma 5.17 asserts that there is some integer n_0 and some ordinal $\beta < \alpha$ such that $rank(\Omega(T(a_n))) \le rank(\Omega(T(a_{n_0}))) = \alpha$ for $n \le n_0$ and $rank(\Omega(T(a_n))) \le \beta$ for $n > n_0$. It follows that $rank(\Omega(T \setminus \{a_{n_0}\}(a_{n_0+1}))) < \alpha$. This implies that $a_{n_0+1} \notin Lim(T)$, proving our claim.

Next we prove that Lim(T) is non-empty.

Let $a \in T \setminus Lim(T)$. We show that there is an integer $n \in \mathbb{N}$ and a sequence $a_0, \ldots, a_i, \ldots a_n$ with $a_0 := a$ and $rank(\Omega(T \setminus \{a_i\}(a_{i+1}))) = \alpha$ for i < n and

with $a_i \in T \setminus Lim(T)$ and $a_n \in Lim(T)$. Indeed, let a_1 be the unique vertex y given by Lemma 7.2. If $a_1 \in Lim(T)$, set n := 1. Otherwise, suppose a_0, \ldots, a_{i+1} be defined, with $rank(\Omega(T \setminus \{a_i\}(a_{i+1}))) = \alpha$. If $a_{i+1} \in Lim(T)$ set n := i+1. Otherwise, observe with Lemma 7.2 that there is a unique neighbour b of a_{i+1} such that $rank(\Omega(T \setminus \{a_{i+1}\}(b))) = \alpha$. Since furthermore, $rank(\Omega(T \setminus \{x\}(y'))) \leq \beta$ for every other $y' \sim a_{i+1}$ and some ordinal $\beta < \alpha$, it follows that $b \neq a_{i+1}$. If the process does not stop, then T is the sum $\bigoplus_{\{a_n:n\in\mathbb{N}\}} T_{a_n}$, where $T_{a_n} := T \setminus \{a_{n+1}\}(a_n)$ for each $n \in \mathbb{N}$. Since $rank(\Omega(T \setminus \{a_n\}(a_{n+1}))) = \alpha$ for each $n \in \mathbb{N}$, Equation 5 of Lemma 5.17 asserts that $rank(\Omega(T)) = \alpha + 1$. This is impossible.

Finally, we prove that Lim(T) is connected.

Suppose not. Let n be the least integer such that there are two vertices a, b in two different connected components of Lim(T) such that $d_T(a, b) = n$. Let $a := a_0, a_1, \ldots, a_n := b$ be the unique path connecting a and b. We have $n \ge 2$, and $a_i \notin Lim(T)$ for every $i, 1 \le i \le n-1$. Since $a_1 \notin Lim(T)$, there is a unique vertex $c \in V(T)$ such that $c \sim a_1$ and $rank(\Omega(T \setminus \{a_1\}(c))) = \alpha$, and some ordinal $\beta < \alpha$ such that $rank(\Omega(T \setminus \{a_1\}(y'))) \le \beta$ for every other $y' \sim a_1$. Suppose $c \ne a_0$. Then $rank(\Omega(T \setminus \{a_1\}(a_0))) \le \beta$. Hence $rank(\Omega(T \setminus \{a_0\}(y'))) \le \beta$ for every $y' \sim a_0, y' \ne a_1$. Thus $a_0 \notin Lim(T)$. Impossible. Since $c = a_0$, $rank(\Omega(T \setminus \{a_0\}(a_1))) < \alpha$, but then $T \setminus \{a_0\}(a_1)$ does not contain a vertex in Lim(T), contradicting that $b \in Lim(T)$.

Lemma 7.4. Let T be a non-empty scattered tree and $\alpha := rank(\Omega(T))$. If α is either a limit ordinal or a successor ordinal $\alpha' + 1$ and $|\Omega^{(\alpha')}(T)| \geq 3$, then some non-empty rayless subtree A of T is preserved by every embedding of T.

Proof. If $\alpha = 0$, then $\Omega(T) = \emptyset$, that is T is rayless and there is nothing to prove. Suppose $\alpha > 0$. We consider two cases:

Case 1. α is a limit ordinal. The assertion follows from Lemma 7.3.

Case 2. α is a successor ordinal $\alpha' + 1$ and $|\Omega^{(\alpha')}(T)| \geq 3$. We may reproduce verbatim the proof of Theorem 3.1 [16], in which the authors prove that there is some rayless tree A which is preserved by every automorphism of T. For every end in $e \in \Omega^{(\alpha')}(T)$ choose a one-way infinite path C_e so that $V(C_e) \cap V(C'_e) = \emptyset$ for $e \neq e' \in \Omega^{(\alpha')}(T)$. This is possible because every end in $\Omega^{(\alpha')}(T)$ is isolated within $\Omega^{(\alpha')}(T)$. Let $W = \{C_e : e \in \Omega^{(\alpha')}(T)\}$. Choose a vertex in each of the sets V(W) and let A be the set of those vertices. Let G be the subtree of T induced by the set of vertices which are on the path from a vertex in A to a vertex in A together with the vertices in $\bigcup_{C \in W} V(C)$. The subtree G contains a vertex v having degree at least three because $|\Omega^{(\alpha')}(T)| \geq 3$. For every $a \in A$ let a_v be the last vertex on the path from v to a of degree larger than or equal to three. Let a' be the first vertex on the path from a_v to a not equal to a_v . Then a' has valence two in G. Let $A' = \{a' : a \in A\}$ and let C_a be the one way infinite path with endpoint a'. Note that for every $a' \in A'$, every one-way infinite path in the end containing C_a has C_a as a subpath. This implies, because every embedding preserves $|\Omega^{(\alpha')}(T)| \geq 3$, that our embedding maps every path in W into a path in W and hence preserves G. Because every embedding maps every vertex to a vertex of the same degree it has to preserve A' and hence the subtree G' of G induced by the set of vertices $V(G) \setminus \bigcup_{C \in W} V(C)$.

It remains to prove that G' does not contain a one-way infinite path, also called a ray. This is proven in [16]. For completeness sake we provide their proof here: Suppose there is a ray $R \in G'$. Every vertex in R is on a path from a vertex in A' to a vertex in A', implying that R does not have a terminal end so that each vertex on it has valence two in G'. If $x \in V(R)$ and the valence of x in G' is at least three, then there is a path from x to some $x' \in A'$. It follows that $x \in A'$ is not an isolated ray in $x \in A'$ in $x \in A'$ and the valence of $x \in A'$ is not an isolated ray in $x \in A'$ is not an isolated ray in $x \in A'$ in $x \in A$

Polat and Sabidussi [16], (Theorem 2.5), proved that every non-empty rayless tree T has a vertex or an edge fixed by every automorphism of T. One of their ideas is to successively change T to the tree $\Phi(T)$ generated by the vertices of infinite degree. Clearly $\Phi(T)$ is preserved by every embedding, not just automorphism of T. It can be easily checked that the arguments in [16] can be extended from automorphisms to embeddings. Another way to verify the extension of the Theorem of Polat and Sabidussi to embeddings is to specialize Theorem 11.5 of Halin, see [6], from graphs to trees. In any case we obtain:

Lemma 7.5. Every non-empty tree T which does not contain a one-way infinite path, has a vertex or an edge preserved by every embedding of T.

7.1. **Proof of Theorem 1.1.** Let T be a scattered tree. According to Theorem 6.10, the space $\Omega(T)$ is scattered. Let $\alpha := rank(\Omega(T))$ be its Cantor-Bendixson rank. There are two cases:

Case 1: α is a limit ordinal or is a successor ordinal $\alpha' + 1$ and $|\Omega^{(\alpha')}(T)| \ge 3$. According to Lemma 7.4, some non-empty rayless subtree A of T is preserved by every embedding of T. According to Lemma 7.5 the subtree A contains a vertex or an edge preserved under every embedding of A, and hence of T.

Case 2: α is a successor ordinal $\alpha' + 1$ and $|\Omega^{(\alpha')}(T)| \le 2$. The set $\Omega^{(\alpha')}(T)$ is preserved by every embedding. We get the required conclusion.

8. Embeddings of labelled trees

Let Q be a quasi ordered set (qoset), that is, a set equipped with a quasi order that we denote by \leq . We set $q \equiv q'$ if $q' \leq q \leq q' \in Q$ and $equiv(q) := \{q' \in Q : q' \equiv q\}$. A Q-labelled graph is a pair (G, ℓ) where ℓ is a map from V(G) into Q. If (G, ℓ) and (G', ℓ') are two Q-labelled graphs, a Q-isomorphism of (G, ℓ) onto (G', ℓ') is an isomorphism f of G onto G' such that $\ell(x) = \ell(f(x))$ for every $x \in V(G)$. A Q-embedding of (G, ℓ) into (G', ℓ') is an embedding f of V(G) into V(G') such that $\ell(x) \leq \ell(f(x))$ for every $x \in V(G)$. We say that (G, ℓ) and (G', ℓ') are isomorphic and we set $(G, \ell) \simeq (G', \ell')$ if there is a Q-isomorphism of (G, ℓ) to (G', ℓ') . We say that they are equimorphic if each of (G, ℓ) and (G', ℓ') are Q-embeddable in the other; in which case we set $(G, \ell) \equiv (G', \ell')$. If (G, ℓ) is a labelled tree, then

as before we set $twin(G, \ell)$ for the set of equimorphic Q-labelled trees up to isomorphism.

If (G, ℓ) is a one-way labelled path we will assume that V(G) has been enumerated with the numbers in \mathbb{N} in a natural way and then, in order to simplify notation, pretend to have identified V(G) with \mathbb{N} along this enumeration. For a two-way infinite path we will use \mathbb{Z} in the place of \mathbb{N} . Let (G, ℓ) be a one-way or two-way infinite path labelled by a poset Q. The labelling ℓ is r-periodic for $1 \leq |r| \in \mathbb{N}$ if $\ell(n) \leq \ell(n+r)$ for all $n \in \mathbb{N}$. The number r for which ℓ is r-periodic with |r| minimal is the period of ℓ . If ℓ is r-periodic with |r| minimal and ℓ is also -r periodic then the period of ℓ is |r|. If ℓ has period p, then $\mathfrak{d}_{\ell} := |\{n \in V(G) : \ell(n) < \ell(n+p)\}|$.

We may note that the tree alternative property fails badly for labelled trees, even with the assumption that the qoset is a poset. In fact:

Let C be the path on the set \mathbb{N} of non-negative integers and ℓ be a labeling by a two element antichain $Q := \{a, b\}$. Then $|twin(C, \ell)| = 1$ if ℓ is constant or ℓ is not periodic and $|twin(C, \ell)| = p$ if p is the period of ℓ . However:

Theorem 8.1. Let (G, ℓ) be a one-way infinite path or a two-way infinite path labelled by a poset Q. If Q has a minimum, say 0, then the tree alternative property holds. If ℓ is not periodic, then $|twin(G, \ell)| = 1$ and if ℓ is periodic and \mathfrak{d}_{ℓ} is infinite then $|twin(G, \ell)| \geq 2^{\aleph_0}$. countable, then $twin(G, \ell)$ is countable.

If ℓ is periodic and G a one-way infinite path, then $twin(G,\ell)$ is infinite unless ℓ is the constant map with image $\{0\}$, in which case $|twin(G,\ell)| = 1$. If ℓ is periodic and G a two-way infinite path, then $twin(G,\ell)$ is infinite unless $\mathfrak{d}_{\ell} = 0$ in which case $|twin(G,\ell)| = 1$.

Question 8.2. Does the tree alternative property hold for labelled trees labelled with a poset having a least element?

The proof of Theorem 8.1 relies on the following lemmas and Theorem 8.8 below.

Lemma 8.3. Let (G, ℓ) and (G', ℓ') be two Q-labelled connected graphs and g be an embedding of (G, ℓ) into (G', ℓ') . Let $\ell'' : V(G') \to Q$ such that:

(7)
$$\ell(g^{-1}(y)) \le \ell''(y) \le \ell'(y)$$

for every $y \in V(G')$, with the stipulation that the first inequality holds if y does not belong to the range of g. Then:

(8)
$$(G,\ell)$$
 embeds into (G',ℓ'') embeds into (G',ℓ') .

As a consequence, if (G, ℓ) and (G', ℓ') are equimorphic, then (G, ℓ) and (G', ℓ'') are equimorphic.

Proof. From the first inequality in (7), the map g is an embedding of (G, ℓ) into (G', ℓ'') , whereas from the second inequality, the identity map is an embedding of (G', ℓ'') into (G', ℓ') .

We need the following result (see Lemma 4, p.39, [12]). For reader convenience, we give a proof.

Lemma 8.4. Let S be an infinite subset of \mathbb{N} . There is a family \mathcal{A} of 2^{\aleph_0} subsets of S such that for every pair A, A' of distinct subsets of \mathcal{A} , no translate of an infinite subset of A is almost included into A'.

Proof. Start with $X := \{x_n \in S, n \in \omega\}$ where $x_0 = \min(S)$ and $x_{n+1} \ge x_n + n$ (one just needs increasing gaps). Now, let A be an almost disjoint family of 2^{\aleph_0} infinite subsets of X. For any $A \in A$, and n > 0, A + n is almost disjoint from X, and thus almost disjoint from any other A'.

Lemma 8.5. Let Q be a poset with a least element 0. Let (G,ℓ) and (G',ℓ') be two equimorphic Q-labelled locally finite graphs. Let g be an embedding of (G,ℓ) into (G',ℓ') and let $D := \{y \in range(g) : \ell(g^{-1}(y)) < \ell'(y)\} \cup \{y \in V(G') \setminus range(g) : \ell'(y) \neq 0\}$. If D is infinite, then $|twin(G,\ell)| \geq 2^{\aleph_0}$.

Proof. Our aim is to find a family \mathcal{B} of 2^{\aleph_0} subsets B of D such that the Q-labelled graphs (G', ℓ'_B) defined by $\ell'_B(y) := \ell(g^{-1}(y))$ are pairwise isomorphic if $y \in B \cap range(g)$ and $\ell'_B(y) = 0$. Indeed, according to Lemma 8.3, these Q-labelled graphs are equimorphic to (G, ℓ) . For this purpose, we will use the following notation.

Let $x \in V(G')$ and $n \in \mathbb{N}$, set $B_{G'}(x,n) \coloneqq \{y \in V(G') : d_{G'}(x,y) = n\}$. Let $X \subseteq V(G')$, set $Spec(X,x) \coloneqq \{n \in \mathbb{N} : X \cap B_{G'}(x,n) \neq \emptyset\}$. Now, fix a vertex of G', say r. Let $S \coloneqq Spec(D,r)$. Since G' is locally finite, S is an infinite subset of \mathbb{N} . Let A be a family of 2^{\aleph_0} subsets A of S satisfying the properties of Lemma 8.4. For each $A \in A$, set $B \coloneqq \{x \in D : d_{G'}(r,x) \in A\}$, hence Spec(B,r) = A. Let $B' \coloneqq \{B : A \in S\}$. This family yields a collection of 2^{\aleph_0} Q-labelled graphs. Divide this collection into isomorphism classes.

We claim that each class is countable. Picking a representative in each class, we will get a family \mathcal{B} as described above. To prove our claim, suppose by contradiction that some isomorphy class, say \mathcal{C} , is uncountable. For each $(G', \ell'_B) \in \mathcal{C}$ pick $x_B \in V(G')$. Since G' is locally finite, V(G') is countable, hence we may find an uncountable subfamily C' such that all labelled graphs $(G', \ell'_B, x_B) \in \mathcal{C}'$ are pairwise isomorphic and, furthermore all x_B are equal to the same element $x \in V(G')$ (indeed, fix a labelled graph $(G', \ell'_B) \in \mathcal{C}$ and observe that if $(G', \ell'_{B'})$ and $(G', \ell'_{B''})$ are isomorphic to (G', ℓ'_B) and the images of $x_{B'}$ and $x_{B''}$ by some isomorphisms are the same, then the labelled graphs $(G', \ell'_{B'}, x_{B'})$ and $(G', \ell'_{B''}, x_{B''})$ are isomorphic).

Claim 8.6. If $(G', \ell'_{B'}, x) \simeq (G', \ell'_{B''}, x)$, then Spec(B', x) = Spec(B'', x).

Proof of Claim 8.6. Assume that $(G', \ell'_{B'}, x) \simeq (G', \ell'_{B''}, x)$. Let $n \in \mathbb{N}$. We prove that if $n \notin Spec(B', x)$, then $n \notin Spec(B'', x)$; that is, $B'' \cap B_{G'}(x, n) = \emptyset$ implies $B'' \cap B_{G'}(x, n) = \emptyset$. Supposing $B'' \cap B_{G'}(x, n) = \emptyset$ we have $\ell_{B''}(y) = \ell(y)$ for every $y \in B_{G'}(x, n)$. Let h be an isomorphism of $(G', \ell'_{B''}, x)$ onto

 $(G', \ell'_{B''}, x)$. For all $y \in B_{G'}(x, n)$ we have:

(9)
$$\ell(y) = \ell_{B''}(y) = \ell_{B'}(h(y) \le \ell(h(y)).$$

Since h is a self map on V(G') we may iterate it. The iterates satisfies the inequalities above. Since G' is locally finite, $B_{G'}(x,n)$ is finite too. Hence, some iterate of h is the identity, and therefore $\ell(y) = \ell(h(y))$ for all $y \in B_{G'}(x,n)$. It follows that $\ell_{B'}(y) = \ell(y)$ for all $y \in B_{G'}(x,n)$. Hence $B' \cap B_{G'}(x,n) = \emptyset$, proving our claim.

Claim 8.7. If Spec(B',x) = Spec(B'',x), then one of the sets A' := Spec(B',r), A'' := Spec(B'',r) contains an infinite translate of the other.

Proof of Claim 8.7. Let D' := Spec(B',x). For each $n \in D'$ pick $x'_n \in B', x'' \in B''$ such that $d_{G'}(x,x'_n) = d_{G'}(x,x''_n) = n$. Since $|d_{G'}(x,x'_n) - d_{G'}(r,x'_n)| \le d_{G'}(x,r)$ and $|d_{G'}(x,x''_n) - d_{G'}(r,x''_n)| \le d_{G'}(x,r)$ we may find a increasing sequence of integers $(\varphi(n))_{n \in \mathbb{N}}$ such that $d_{G'}(x,x''_{\varphi(n)}) - d_{G'}(r,x''_{\varphi(n)}) = c'$ and $d_{G'}(x,x''_{\varphi(n)}) - d_{G'}(r,x''_{\varphi(n)}) = c''$ for some integer constants c' and c''. It follows that $d_{G'}(r,x'_{\varphi(n)}) - d_{G'}(r,x''_{\varphi(n)})$ is an integer constant. The set $Y' := \{d_{G'}(r,x'_{\varphi(n)}) : n \in \mathbb{N}\}$ is an infinite subset of A' := Spec(B',x) and the set Y'' similarly defined is an infinite subset of A'' := Spec(B'',x). One of the sets Y', Y'' is a translate of the other, as claimed.

To complete the proof of Lemma 8.5, observe that this situation is not possible, because $A', A'' \in A$.

We can rephrase Lemma 8.5 and state it as Theorem 8.8:

Theorem 8.8. Let Q be a poset with a least element 0. Let (G, ℓ) be a locally finite graph such that $|twin(G, \ell)| < 2^{\aleph_0}$. Then, for every embedding g of (G, ℓ) into a twin (G', ℓ') , the set of $x \in V(G)$ such that $\ell(x) \neq \ell'(f(x))$, and the set of $y \in V(G') \setminus range(g)$ such that $\ell'(y) \neq 0$, are finite.

Proof of Theorem 8.1 Let (G,ℓ) be a one way infinite path. The function ℓ is r-constant for $1 \le r \in \mathbb{N}$ if $\ell(n+r) = \ell(n)$ for all $n \in \mathbb{N}$. If there exists such a number r, then ℓ is stepwise constant. Note that the smallest such number r is the period of ℓ . If ℓ is stepwise constant when restricted to some terminal interval $[n,\infty)$ of \mathbb{N} , then ℓ is eventually stepwise constant. Let ℓ_n be the restriction of the labelling ℓ to the interval $[n,\infty)$. If ℓ is eventually stepwise constant there exists a smallest number n, the character of ℓ , for which $\mathfrak{d}_{\ell_n} = 0$. The character of ℓ is an isomorphism invariant. Note that if $\mathfrak{d}_{\ell_n} = 0$, then ℓ_n is stepwise constant and if Q is countable, then there are countably many eventually stepwise constant Q-labelled one way paths.

If ℓ is not periodic, then there is no embedding f of (G,ℓ) with 0 < f(0) and hence $|twin(G,\ell)| = 1$. Let ℓ be periodic with period p. If \mathfrak{d}_l is infinite, then because the function $n \to n + p$ is an embedding, it follows from Theorem 8.8 that $|twin(G,\ell)| \ge 2^{\aleph_0}$.

Let (G, ℓ) be a two-way infinite path. The function ℓ is r-constant for $1 \le r \in \mathbb{N}$ if $\ell(n+r) = \ell(n)$ for all $n \in \mathbb{Z}$. If there exists such a number r, then ℓ is stepwise constant. Note that the smallest such number r is the period of ℓ . The function ℓ is eventually stepwise constant if there are numbers m and n in \mathbb{Z} so that ℓ is stepwise constant when restricted to the one-way infinite path $(-\infty, m)$ as well as to the one-way infinite path $[n, \infty)$. In this case the eventual period of ℓ is the least common multiple of the period of ℓ on $[n, \infty)$.

If ℓ is not periodic, there is no embedding f of (G,ℓ) which is a translation of \mathbb{Z} with $f(0) \neq 0$. This implies that every embedding is an automorphism and hence that $|twin(G,\ell)| = 1$. If ℓ is periodic, say with period p and $\mathfrak{d}_{\ell} = 0$, then ℓ is stepwise constant, in which case $|twin(G,\ell)| = 1$. If \mathfrak{d}_{ℓ} is infinite, then because the function $n \to n + p$ is an embedding, it follows from Theorem 8.8 that $|twin(G,\ell)| \geq 2^{\aleph_0}$.

We look at graphs labelled by a qoset.

Lemma 8.9. Let (G, ℓ) be a Q-labelled connected graph. Let $Q_{\ell} := \{equiv(\ell(x)) : x \in V(G)\}$, and κ be the cardinality of the set Φ of maps φ from Q_{ℓ} into Q such that $\varphi(C) \in C$ for each $C \in Q_{\ell}$. Then

(10)
$$\kappa \le |twin(G,\ell)|.$$

In particular

(11)
$$|equiv(\ell(x))| \le |twin(G,\ell)|.$$

for every $x \in V(G)$.

Proof. Let $\varphi \in \Phi$. Set V := V(G) and define $\ell_{\varphi} : V \to Q$ by setting $\ell_{\varphi}(y) = \varphi(equiv(\ell(y)))$. The Q-labelled graph $G < \varphi > := (G, \ell_{\varphi})$ is equimorphic to (G, ℓ) . Indeed, the identity map on V is an embedding of (G, ℓ) into $G < \varphi >$ and of $G < \varphi >$ into (G, ℓ) . If $\varphi \neq \varphi'$, the Q-labelled graphs $G < \varphi >$ and $G < \varphi' >$ cannot be isomorphic. Otherwise, let $x \in V$ such that $q := \varphi(equiv(l(x))) \neq \varphi'(equiv(l(x))) =: q'$. We would have $q \neq \ell_{\varphi'}(f(x)) = \ell_{\varphi}(x) = q$ for every isomorphism f of $G < \varphi >$ onto $G < \varphi' >$. The inequality follows.

9. Bonato-Tardif and Tyomkyn conjectures and a proof of Theorem 1.9

Lemma 9.1. Let T a tree, A be a subtree of T, $x \in A$ and $T_x := ((T, A)(x), x)$. If f(A) = A for every embedding f of T, then $|twin(T_x)| \le |twin(T)|$.

Proof. Let Q be the set of rooted trees (considered up to isomorphy) which are equimorphic to some T(x) for $x \in A$. Let (A, ℓ) where $\ell(x) := T(x)$. We claim that $|twin(A, \ell)| \leq |twin(T)|$. Indeed, notice first that if (A', ℓ') is a Q-labelled tree, and $(A, \ell') \equiv (A, \ell)$, then $\bigoplus_{x \in A} \ell'(x) \equiv \bigoplus_{x \in A} \ell(x)$. Next, let (A', ℓ') , (A'', ℓ'') be two Q-labelled trees equimorphic to (A, ℓ) . Suppose that $\bigoplus_{x \in A'} \ell'(x) \cong \bigoplus_{x \in A''} \ell''(x) \equiv T$. Let f be an isomorphism of $\bigoplus_{x \in A'} \ell'(x)$ onto $\bigoplus_{x \in A''} \ell''(x)$. The map f induces an embedding from T into T hence, by our

hypothesis, f(A) = A. This implies that $(A, \ell') \simeq (A'', \ell)$, proving our claim. According to Lemma 8.9, we have $|twin((T, A)(x), x)| \leq |twin((T, \ell))|$. From Lemma 9.1 and Tyomkyn's Theorem [18] that the rooted tree alternative conjecture holds we obtain:

Corollary 9.2. Let T be a tree and A be a subtree such that f(A) = A for every embedding of T. If $|twin(T)| < \omega$, then |twin((T, A)(x), x)| = 1 for every $x \in A$.

Let T be a tree and A be a finite subtree. If T' is an other tree and A' is a finite subtree of T', we will say that (T,A) is embeddable into (T',A') if there is an embedding of T into T' which maps A into A'. If n := |A| and $(a_i)_{i < n}$ is an enumeration of A and A' is an n-element subtree of a tree T' with an enumeration $(a'_i)_{i < n}$, we will say that $((T,(a_i)_{i < n})$ is embeddable into $(T',(a'_i)_{i < n})$ if there is an embedding of T into T' which sends each a_i onto a'_i . This allows to define twin((T,A)) and $twin((T,(a_i)_{i < n}))$. We define twin((T,A)) as the set of twins of (T,A) considered up to isomorphy, and similarly $twin((T,(a_i)_{i < n}))$.

Lemma 9.3.

(12)
$$\frac{1}{n!} \cdot |twin((T,(a_i)_{i< n}))| \le |twin((T,A))| \le |twin((T,(a_i)_{i< n}))|.$$

Proof. Note that the map φ which transforms $(T, (a_i)_{i < n})$ to (T, A) preserves isomorphy, namely if $(T', (a'_i)_{i < n}) \simeq (T'', (a''_i)_{i < n})$, then $(T', A') \simeq (T'', A'')$. Hence φ sends isomorphic types onto types. Also $(T', (a'_i)_{i < n}) \equiv (T'', (a''_i)_{i < n})$, then $(T', A') \equiv (T'', A'')$, hence φ sends $twin((T, (a_i)_{i < n}))$ into twin((T, A)). This map is surjective, proving that the second inequality holds. Indeed, let $(T', A') \equiv (T, A)$. Let f, g witness this fact, that is f is an embedding of (T', A') into (T, A) and g is an embedding of (T, A) into (T', A'). Since A is finite, there is a nonnegative integer m such that $(g \circ f)_{\uparrow A}^{(m)}$ is the identity on A'. Let $f' := (g \circ f)^{(m)} \circ g$. Let $(a'_i)_{i < n}$ such that $a_i = f(a'_i)$. Then $(T', (a'_i)_{i < n}) \equiv (T, (a_i)_{i < n})$, proving that the map φ is surjective. Let $(T', A') \in twin((T, A))$. The number of non-isomorphic types in $twin((T, (a_i)_{i < n}))$ which can be sent onto (T', A') is at most n!. The first inequality follows.

From this type of trivial argument we cannot expect to prove that $|twin((T,A))| \leq |twin((T,(a_i)_{i< n}))|$. For an example, there are two graphs G and G' and subsets $A := \{a_0, a_1\}, A' := \{a'_0, a'_1\}$ such that $(G,(a_0,a_1)) \equiv (G',(a'_0,a'_1)), (G,(a_0,a_1)) \not\equiv (G',(a'_0,a'_1))$. The graph G is a path on $\mathbb N$ with two extra vertices a_0 and a_1 , with a_0 linked by an edge to every even integer, a_1 linked by an edge to every odd integer. The graph G' is $G_{-0}, \alpha'_i = \alpha_i$.

9.1. A vertex or an edge preserved.

Proposition 9.4. Let T be a tree. If there is a vertex or an edge preserved by every embedding of T, then |twin(T)| is 1 or ∞ , and if T is also locally finite, then every embedding of T is an automorphism of T.

Proof. Case 1: There is a vertex x preserved by every embedding. In this case twin(T) = twin((T,r)). According to Tyomkyn's Theorem, |twin((T,r))| is 1 or ∞ . Every embedding of a locally finite rooted tree is an automorphism, see [18] Lemma 4.

Case 2: There is an edge $u := \{x, y\}$ preserved by every embedding but there is no vertex fixed by every embedding. According to the inequalities in (12) we have:

(13)
$$\frac{1}{2} \cdot |twin((T,(x,y)))| \le twin((T,u))| \le |twin((T,(x,y)))|.$$

According to Tyomkyn's Theorem, |twin((T,x))| is $1 \text{ or } \infty$. Since in our case twin((T,x)) = twin((T,(x,y))), $|twin((T,\{x,y\}))|$ is $1 \text{ or } \infty$. Since $twin(T) = twin((T,\{x,y\}))$, the result follows. Let T be locally finite and f an embedding of T. If f(x) = x, then f(y) = y and hence f is an automorphism on the tree (T,x) and on the tree (T,y). If f(x) = y then f(y) = x and hence f^2 is the identity, and therefore f is an automorphism.

9.2. A two-way infinite path preserved.

Proposition 9.5. If a tree contains a two-way infinite path preserved by every embedding, then the tree alternative property holds.

Proof. Let T be a tree, D be a two-way infinite path preserved by every embedding. Suppose that $|twin(T)| < \omega$. According to Lemma 9.1 and Tyomkyn's Theorem, |twin((T,D)(x),x))| = 1 for every $x \in D$. Write $T = \bigoplus_{x \in D} ((T,D)(x),x)$. Let Q be the set of rooted trees which embed in some ((T,D)(x),x) for some $x \in D$; we quasi order Q by embeddability of rooted trees. Let (D,ℓ) be the labelled two-way infinite path where $\ell:D \to Q$ is defined by $\ell(x) := ((T,D)(x),x)$. Each embedding of a labelled two-way infinite path yields an embedding of T and conversely. Hence $twin(T) = twin(D,\ell)$. The conclusion follows from Theorem 8.1.

9.3. One end preserved. Let \mathfrak{E} be the set of trees having an end preserved forward by every embedding. We will use the notions introduced in Subsection 5.2. In particular, each $T \in \mathfrak{E}$ has a level function, say lev_T .

Case 1: Every embedding of T preserves the level function. This amounts to the fact that the end e is almost rigid. Let \mathfrak{G} be the set of trees T in \mathfrak{E} with the property that every embedding of T preserves lev_T .

Proposition 9.6. If $T \in \mathfrak{G}$, then the set twin(T) is infinite or a singleton. If in addition T is locally finite, then every embedding of T is an automorphism of T, hence |twin(T)| = 1.

Note that if T is locally finite, every embedding which has a fixed point is surjective. Since T has a level function preserved by every embedding of T, every embedding has a fixed point. Note also that if T is scattered and the ray is regular, then by Theorem 1.3, T has a vertex fixed by every embedding. Hence in this case the value of |twin(T)| is given by Proposition 9.4. The general case needs more work.

For n a number in \mathbb{Z} let $\mathcal{L}_n(T) := \{x \in V(T) : lev_T(x) = n\}$ and let $\mathcal{L}_{\geq n}(T) := \{x \in V(T) : lev_T(x) \geq n\}$. Two trees T and T', with the same origin $o \in V(T) \cap V(T')$ for the level functions, are n-equimorphic if $\mathcal{L}_{\geq n}(T)$ is equal to $\mathcal{L}_{\geq n}(T')$ and if for every $x \in \mathcal{L}_n(T)$ the rooted trees $T(\to x)$ and $T'(\to x)$ are equimorphic. An embedding of a tree T into a tree T' is level preserving if $lev_T(x) = lev_{T'}(f(x))$ for all $x \in V(T)$.

Claim 9.7. Let T and T' be two trees which are n-equimorphic, with common origin o for the level function. If $T \in \mathfrak{G}$, then $T' \in \mathfrak{G}$ and every embedding of T' into T is level preserving and every embedding of T into T' is level preserving.

Proof of Claim 9.7. Let f be an embedding of T into T' with f(x) = x for all $x \in \mathcal{L}_{\geq n}$ and f restricted to $T(\to x)$ an embedding of $T(\to x)$ into $T'(\to x)$ for all $x \in \mathcal{L}_n$. Let g be an embedding of T' into T with g(x) = x for all $x \in \mathcal{L}_{\geq n}$ and g restricted to $T'(\to x)$ an embedding of $T'(\to x)$ into $T(\to x)$ for all $x \in \mathcal{L}_n$. Then $lev_T(x) = lev_{T'}(x)$ for all $x \in \mathcal{L}_{\geq n}(T)$. Because f is then an embedding of the rooted tree $T(\to x)$ into the rooted tree $T'(\to x)$ for every $x \in \mathcal{L}_n(T)$ and $lev_T(x) = lev_{T'}(x)$, the embedding f is level preserving on $T(\to x)$. It follows that f is level preserving. Similarly, the embedding g is level preserving.

Let h be an embedding of T'. Then $g \circ h \circ f$ is an embedding of T and hence level preserving. The embeddings f and g and $g \circ h \circ f$ are level preserving and hence h is level preserving. Let h be an embedding of T' into T. Then $h \circ f$ is an embedding of T and hence level preserving. Because f is level preserving, the embedding h is level preserving.

Lemma 9.8. Let $T \in \mathfrak{G}$. Then $|twin(T(\to x))| \le |twin(T)|$ for all $x \in V(T)$ and if the set twin(T) is finite, then $|twin(T(\to x))| = 1$ for every $x \in V(T)$.

Proof. Let $x \in V(T)$ and $n := lev_T(x)$ and $\kappa := |twin(T(\rightarrow x))|$ and let $(R^{\langle \alpha \rangle}; \alpha \in \kappa)$ be an enumeration of $twin(T(\rightarrow x))$. Let $Y := \{y \in \mathcal{L}_n : T(\rightarrow y) \equiv T(\rightarrow x)\}$. For $\alpha \in \kappa$ let $T^{\langle \alpha \rangle}$ be the tree obtained from T by replacing for every $y \in Y$ the tree $T(\rightarrow y)$ with the tree $T(\rightarrow y)$. Then $T^{\langle \alpha \rangle}$ is equimorphic to $T^{\langle \beta \rangle}$ for all α and β in κ . It follows from Claim 9.7 that every embedding of $T^{\langle \alpha \rangle}$ is level preserving for every $\alpha \in \kappa$. Let α and β in κ and let α be an isomorphism of $T^{\langle \alpha \rangle}$ to $T^{\langle \alpha \rangle}$. It follows from Claim 9.7 that α is level preserving. Hence α maps the set α onto the set α and then induces an isomorphism of $T^{\langle \alpha \rangle}(\rightarrow y)$ onto $T^{\langle \alpha \rangle}(\rightarrow y)$ for every $\alpha \in \gamma$. This implies that $\alpha = \beta$.

If $|twin(T(\to x))| > 1$ for some $x \in V(T)$, then $|twin(T(\to x))|$ is infinite, according to Tyomkyn's Theorem that the rooted tree alternative conjecture holds, and hence twin(T) is infinite.

Two twins T and T' are said to be in position if there are embeddings f of T to T' and g of T' to T for which $S_{g \circ f} = V(T) \cap V(T')$ and f(x) = x and g(x) = x for all $x \in S_{g \circ f}$. The embeddings f and g are the position embeddings. Note that if T and T' are in position with position embeddings f and g, then $T(\to x) \equiv T'(\to x)$ for all $x \in S_{g \circ f}$.

Claim 9.9. Let $T \in \mathfrak{G}$ and let T' be a tree equimorphic to T. Then there exists a tree T'' isomorphic to T' so that the trees T and T'' are in position.

Proof of Claim 9.9. Let \bar{T} be an isomorphic copy of T' with $V(T) \cap V(\bar{T}) = \emptyset$. There are then embeddings \bar{f} of T into \bar{T} and \bar{g} of \bar{T} into T. The embedding $\bar{g} \circ \bar{f}$ of T is level preserving and hence has a set of fixed points $S_{\bar{g} \circ \bar{f}}$. Then \bar{f} is an isomorphism of $S_{\bar{g} \circ \bar{f}}$ into \bar{T} with \bar{g} restricted to $f[S_{\bar{g} \circ \bar{f}}]$ the inverse of \bar{f} .

For every point $x \in \tilde{S}_{g \circ f}$ let $T_x = ((T, S_{\bar{g} \circ \bar{f}})(x), x)$. That is, the subtree of T rooted at x which is the connected component after removing the set $S_{\bar{g} \circ \bar{f}} \setminus \{x\}$ of vertices from T. Similarly let $\bar{T}_{\bar{f}(x)} = ((\bar{T}, \bar{f}[S_{\bar{g} \circ \bar{f}}])(\bar{f}(x)), \bar{f}(x))$. Let

$$T'' = \bigoplus_{x \in S_{\bar{q} \circ \bar{f}}} (\bar{T}_{\bar{f}(x)}, \bar{f}(x)).$$

Then T'' is isomorphic to \overline{T} and hence isomorphic to T'.

Let f be the embedding of T into T'' for which f(x) = x for all $x \in S_{\bar{g} \circ \bar{f}}$ and $f(z) = \bar{f}(z)$ for all $z \in V(T) \setminus S_{\bar{g} \circ \bar{f}}$. Let g be the embedding of T'' into T for which g(x) = x for all $x \in S_{\bar{g} \circ \bar{f}}$ and $g(z) = \bar{g}(z)$ for all $z \in V(T'') \setminus S_{\bar{g} \circ \bar{f}}$.

Let \mathfrak{F} be the subclass of \mathfrak{G} made of trees T for which the set twin(T) is finite. We have to prove that |twin(T)| = 1 for every tree in \mathfrak{F} . It follows from Claim 9.9 that it suffices to prove that if $T \in \mathfrak{F}$ and the pair T and T' are twins in position, then T' is isomorphic to T. If T and T' are in position with position embeddings f of T into T' and g of T' into T and if $x \in S_{g \circ f}$, then $T(\to x) \equiv T'(\to x)$ and hence $T(\to x) \simeq T'(\to x)$, because $|twin(T(\to x))| = 1$.

Let $T \in \mathfrak{F}$ and T' be twins in position with position embeddings f of T into T' and g of T' into T. Let $Y \subseteq S_{g \circ f}$ so that $x \notin C_y$ for all x and y in Y with $x \neq y$. Let the tree obtained from T' by replacing $T'(\to Y)$ with $T(\to Y)$ be the tree:

$$repl(T', T(\rightarrow Y)) \coloneqq \bigoplus_{y \in Y, T' \setminus T'(\rightarrow -Y)} T(\rightarrow y).$$

Claim 9.10. Let $T \in \mathfrak{F}$ and T' be twins in position with position embeddings f of T into T' and g of T' into T. Let $Y \subseteq S_{g \circ f}$ so that $x \notin C_y$ for all x and y in Y with $x \neq y$.

Then T and $T'' = repl(T', T(\to Y))$ are twins in position with position embeddings \bar{f} and \bar{g} for which $\bar{f}(z) = f(z)$ for all $z \in T \setminus T(\to -Y)$ and $\bar{f}(z) = z$ for all $z \in T(\to Y)$ and for which $\bar{g}(z) = g(z)$ for all $z \in T' \setminus T'(\to -Y)$ and $\bar{g}(z) = z$ for all $z \in T'(\to Y)$. There exists, for any choice of isomorphism ι_y of $T'(\to y)$ to $T(\to y)$ for $y \in Y$, an isomorphism β of T' to T'' with $\beta(z) = z$ for all z in $T' \setminus T'(\to -Y)$ and $\beta(z) = \iota(z)$ for $z \in T'(\to y)$ and $y \in Y$.

Proof of Claim 9.10. We have $V(T) \cap V(repl(T', T(\to X))) = V(T) \cap V(T') \cup T(\to X)$ and hence both \bar{f} and \bar{g} fix every vertex in $V(T) \cap V(repl(T', T(\to X)))$. Also \bar{f} and \bar{g} are embeddings.

Definition 9.11. Let $T \in \mathfrak{E}$ and let $x \in V(T)$ and $X \subseteq V(T)$.

Let $T(\to -x) = T(\to x) \setminus \{x\}$ and $T(\to X) = \bigcup_{x \in X} T(\to x)$ and $T(\to -X) = T(\to X) \setminus X$ and $N_T(x) = \{y \in (T) : y \to x\}$. For $y^* \in N_T(x)$ let:

$$T\lfloor y^*,x\rfloor = \bigcup_{y\in (N_T(x)\smallsetminus \{y^*\}} V(T(\to y)) \ \ and \ \ T\lceil y^*,x\rceil = T\smallsetminus T\lfloor y^*,x\rfloor.$$

Claim 9.12. Let $T \in \mathfrak{F}$ and T' be twins in position with position embeddings f of T into T' and g of T' into T. Let $x \in S_{g \circ f}$ and $y^* \in N_T(x) \cap S_{g \circ f}$. If $T(\to y) \subseteq S_{g \circ f}$ for all $y \in N_T(x) \cap S_{g \circ f}$ then:

There exists a pair of twins R and R' in position with $\{y^*, x\} \subseteq V(R) \cap V(R')$ and with:

$$T[y^*, x] = R[y^*, x]$$
 and $T'[y^*, x] = R'[y^*, x]$ and $R(\to x) = R'(\to x)$.

There is an isomorphism δ of T to R with $\delta(z) = z$ for all vertices z in $T[y^*, x]$. There is an isomorphism δ' of T' to R' with $\delta'(z) = z$ for all vertices z in $T'[y^*, x]$. The isomorphism δ restricted to $T(\to x)$ is an isomorphism ϵ of $T(\to x)$ to $R(\to x)$ with $\epsilon(z) = z$ for all vertices z in $T(\to y^*)$. The isomorphism δ' restricted to $T'(\to x)$ is an isomorphism ϵ' of $T'(\to x)$ to $R'(\to x)$ with $\epsilon'(z) = z$ for all vertices z in $T'(\to y^*)$.

Proof of Claim 9.12. If y and z are vertices in $N_T(x)$ let $y \cong z$ if the rooted tree $T(\to y)$ is isomorphic to the rooted tree $T(\to z)$. If y and z are vertices in $N_{T'}(x)$ let $y \cong z$ if the rooted tree $T'(\to y)$ is isomorphic to the rooted tree $T'(\to z)$. If $y \in N_T(x)$ and $z \in N_{T'}(x)$ let $y \cong z$ if the rooted tree $T(\to y)$ is isomorphic to the rooted tree $T'(\to z)$. Because $T(\to x)$ and $T'(\to x)$ are isomorphic there exist partitions $\mathcal{P} = (P_k; k \in K)$ of $N_T(x)$ and $\mathcal{P}' = (P'_k; k \in K)$ of $N_{T'}(x)$ into \cong -equivalence classes so that $y \cong y'$ for every $k \in K$ and every $y \in P_k$ and $y' \in P'_k$ and so that $|P_k| = |P'_k|$ for every $k \in K$. For every $k \in K$ let $\bar{P}_k = P_k \setminus (P_k \cap S_{g \circ f})$ and let $\bar{P}'_k = P'_k \setminus (P_k \cap S_{g \circ f})$. Note that $P_k \cap S_{g \circ f} = P'_k \cap S_{g \circ f}$ for every $k \in K$.

Let $k \in K$. It follows from $|P_k| = |P'_k|$ and $P \cap S_{g \circ f} = P'_k \cap S_{g \circ f}$ that if $|\bar{P}_k| \neq |\bar{P}'_k|$, then $|\bar{P}_k| \leq |P_k \cap S|$ and $|\bar{P}'_k| \leq |P'_k \cap S|$ and $|P_k \cap S_{g \circ f}| = P'_k \cap S_{g \circ f}$ is infinite. Let $I = \{k \in K : |\bar{P}_k| \neq |\bar{P}'_k|\}$ and $J = \{k \in K : |\bar{P}_k| = |\bar{P}'_k|\}$. Let $W = T(\to \bigcup_{k \in J} \bar{P}_k)$ and $W' = T'(\to \bigcup_{k \in J} \bar{P}'_k)$. Note that there is an isomorphism, say ι , of W' to W. Let $(W \cup \{x\}, x)$ be the tree rooted at x for which $(W \cup x, x) \setminus x = W$. Let γ be an embedding of T so that for every $k \in I$:

- (1) For every $k \in I$, the restriction of γ to P_k is an injection of P_k onto $(P_k \cap S) \setminus \{y^*\}$.
- (2) For every $k \in I$, the restriction of γ to $T(\to y)$ for $y \in P_k$ is an isomorphism of $T(\to y)$ to $T(\to \gamma(y))$.
- (3) If $z \in W$ or $z \in T(\to y^*)$, then $\gamma(z) = z$.
- (4) If z is a vertex in $T[y^*, x]$, then $\gamma(z) = z$.

Let γ' be an embedding of T' so that for every $k \in I$:

- (1) For every $k \in I$, the restriction of γ to P'_k is an injection of P'_k onto $(P'_k \cap S) \setminus \{y^*\}.$
- (2) For every $k \in I$, the restriction of γ to $T'(\to y)$ for $y \in P'_k$ is an isomorphism of $T'(\to y)$ to $T'(\to \gamma(y))$.
- (3) If $z \in W'$ or $z \in T'(\rightarrow y^*)$, then $\gamma'(z) = z$.
- (4) If z is a vertex in $T'[y^*, x]$, then $\gamma'(z) = z$.

For $R := \gamma[T]$ and $R'' := \gamma'[T']$ and $U = \bigcup_{k \in I} \bar{P}_k$ and $U' = \bigcup_{k \in I} \bar{P}'_k$ we have $R = T \setminus T(\to U)$ and $R'' = T' \setminus T'(\to U')$. Then $T[y^*, x] = R[y^*, x]$ and $T'[y^*, x] = R''[y^*, x]$. Let f be the embedding of R into R'' given by: f(z) = f(z) for all z in R with $z \in T \setminus W$. For $z \in W$ let $f(z) = \iota^{-1}(z)$. Let \bar{g} be the embedding of R" into R given by: $\bar{g}(z) = g(z)$ for all z in R" with $z \in R'' \setminus W'$. For all $z \in W'$ let $\bar{g}(z) = \iota(z)$. It follows that $S_{\bar{g} \circ \bar{f}} = S_{g \circ f}$ and that R and R" are in position with position embeddings f and \bar{g} .

Next we identify the vertices z in R'' with the vertices $\iota(z)$. $R' = (R'' \setminus W') \oplus_x (W \cup \{x\}, x) = repl(R'', T(\rightarrow \bigcup_{k \in J} \bar{P}_k))$ and use Claim 9.10. Then there exists an isomorphism ν of R'' to R' given by $\nu(z)=z$ for all vertices $z \text{ in } R'' \setminus W' \text{ and } \nu(z) = \iota(z) \text{ for } z \text{ in } W'. \text{ Then } T'[y^*, x] = R''[y^*, x] = R'[y^*, x]$ and $R(\to x) = R'(\to x)$.

Let δ be the isomorphism γ of T to R and let δ' be the isomorphism $\nu \circ \gamma$ of T' to R'. Then $\delta(z) = z$ for all $z \in T(\to y^*)$ and because $\delta(x) = x$ and δ is an isomorphism the restriction of δ to $T(\to x)$ is an isomorphism of $T(\to x)$ to $R(\to x)$. Similar of δ' . Let $\dot{f}(z) = \bar{f}(z)$ for those z in R with z in $R \setminus W$ and f(z) = z for z in W. Let $\check{g}(z) = \bar{g}(z)$ for those z in R' with z in R' \times W and $\check{g}(z) = z$ for z in W. Then R and R' are in position with position embeddings f and \check{q} .

Claim 9.13. Let $T \in \mathfrak{F}$ and T' be twins in position. Then there exists a tree R for which both T and T' are isomorphic to R.

Proof of Claim 9.13. Let f of T to T' and g of T' to T be the position embeddings of T and T'. Let the origin o of the level function of T and T' be in $S_{q \circ f}$ and let $x_n := o \boxplus n$ for every $n \in \mathbb{N}$. We will construct an ω -sequence \mathfrak{R} of pairs of twins in position with: $\mathfrak{R} = (T \simeq R^{(0)}, T' \simeq R'^{(0)}), (R^{(1)}, R'^{(1)}), (R^{(2)}, R'^{(2)}), \dots$ and position embeddings f_n of $R^{\langle n \rangle}$ into $R^{\langle n \rangle}$ and g_n of $R^{\langle n \rangle}$ into $R^{\langle n \rangle}$ and with isomorphism γ_{n+1} of $R^{\langle n+1 \rangle}$ to $R^{\langle n \rangle}$ and γ'_{n+1} of $R^{\langle n+1 \rangle}$ to $R^{\langle n \rangle}$ so that for all $n \in \mathbb{N}$:

- (1) The path $C_o \subseteq R^{(n)}$ and $C_o \subseteq R'^{(n)}$.
- (2) $R^{\langle n \rangle}(\to x_n) = R'^{\langle n \rangle}(\to x_n) \subseteq S_{g_n \circ f_n}.$
- $(3) R^{(n)} \setminus (R^{(n)}(\to -x_n)) = T \setminus (T(\to -x_n)).$
- (4) $\gamma_{n+1}(z) = z$ for all z in $R^{\langle n \rangle}[x_n, x_{n+1}]$. (5) $\gamma'_{n+1}(z) = z$ for all z in $R^{\langle n \rangle}[x_n, x_{n+1}]$.

We proceed by recursion on n. Let $R^{(0)} = T$ and $P^{(0)} = T(\to x_0) = Q^{(0)}$ and let α_0 be the identity map on $T(\to x_0)$ and α'_0 an isomorphism of $T'(\to x_0) := Q'^{(0)}$

to $T(\to x_0)$. Let $R'^{(0)} = repl(T', T(\to \{x_0\}))$ and γ_0 the identity map on T and γ'_0 the isomorphism β given by Claim 9.10 with ι for α'_0 .

Let $Y_n = (N_{R^{(n)}}(x_{n+1}) \cap S_{g_n \circ f_n}) \setminus \{x_n\}$ and ι_y an isomorphism of $R'^{(n)}(\to y)$ to $R^{(n)}(\to y)$ for every $y \in Y_n$. Let $R''^{(n)} = repl(R'^{(n)}, R^{(n)}(\to Y_n))$ and β the isomorphism of $R'^{(n)}$ to $R''^{(n)}$ given by Claim 9.10. Note that then $R''^{(n)}(\to y) = R^{(n)}(\to y)$ and hence $R^{(n)}(\to y) \subseteq S_{\bar{g}\circ\bar{f}}$ for all $y \in N_{R^{(n)}}(x_{n+1}) \cap S_{\bar{g}\circ\bar{f}}$. Note also that β restricted to $R'^{(n)}(\to x_{n+1})$ is an isomorphism of $R''^{(n)}(\to x_{n+1})$ which fixes $R'^{(n)}(\to x_n)$.

We apply Claim 9.12 with $R^{(n)}$ for T and $R''^{(n)}$ for T' and x_{n+1} for x and x_n for y^* to obtain the pair $R^{(n+1)}$ and $R'^{(n+1)}$ of twins in position for which $R^{(n+)}(\to x_{n+1}) = R'^{(n+1)}(\to x_{n+1})$ and:

$$R^{\langle n \rangle}[x_n, x_{n+1}] = R^{\langle n+1 \rangle}[x_n, x_{n+1}]$$
 and $R^{\langle n \rangle}[x_n, x_{n+1}] = R^{\langle n \rangle}[x_n, x_{n+1}].$

Let

$$P^{\langle n+1\rangle} = R^{\langle n+1\rangle}(\to x_{n+1}) \setminus R^{\langle n+1\rangle}(\to -x_n)$$

and note that

$$R^{\langle n+1\rangle} = R^{\langle n\rangle}[x_n, x_{n+1}] \oplus_{x_{n+1}} P^{\langle n+1\rangle} \text{ and } R'^{\langle n+1\rangle} = R'^{\langle n\rangle}[x_n, x_{n+1}] \oplus_{x_{n+1}} P^{\langle n+1\rangle}$$

and that

$$T(\to x_{n+1}) \setminus T(\to -x_n) = R^{\langle n \rangle}(\to x_{n+1}) \setminus R^{\langle n \rangle}(\to x_n) := Q^{\langle n+1 \rangle},$$

$$T'(\to x_{n+1}) \setminus T'(\to -x_n) = R'^{\langle n \rangle}(\to x_{n+1}) \setminus R'^{\langle n \rangle}(\to x_n) := Q'^{\langle n+1 \rangle}.$$

The isomorphism ϵ mapping $R^{\langle n \rangle}(\to x_{n+1})$ to $R^{\langle n+1 \rangle}(\to x_{n+1})$ given by Claim 9.12 fixes the rooted tree $R^{\langle n \rangle}(\to x_n)$ and hence induces an isomorphism α_{n+1} of $Q^{\langle n+1 \rangle}$ to $P^{\langle n+1 \rangle}$. Similarly $\epsilon' \circ \beta$ induces an isomorphism of $R'^{\langle n \rangle}(\to x_{n+1})$ to $R'^{\langle n+1 \rangle}(\to x_{n+1})$, which fixes the rooted tree $R'^{\langle n \rangle}(\to x_n)$ and hence induces an isomorphism α'_{n+1} of $Q'^{\langle n+1 \rangle}$ to $P^{\langle n+1 \rangle}$.

The tree T can be written as a sum $T = \bigoplus_{C_o} Q^{\langle n \rangle}$. That is, the tree T consists of the one-way infinite path C_o with a rooted tree $Q^{\langle n \rangle}$ attached at x_n for every $n \in \mathbb{N}$. Similarly $T' = \bigoplus_{C_o} Q'^{\langle n \rangle}$. Let $R = \bigoplus_{C_o} P^{\langle n \rangle}$. Then $\bigcup_{n \in \mathbb{N}} \alpha_n$ is an isomorphism of T to R and $\bigcup_{n \in \mathbb{N}} \alpha'_n$ is an isomorphism of T' to R.

9.3.1. **Proof of Proposition 9.6.** Assume that the set twin(T) is not infinite. It follows from Lemma 9.8 that then $|twin(T(\to x))| = 1$ for all $x \in V(T)$. Because of Claim 9.9 it suffices to show that if T' is a twin of T and T' are in position, then T and T' are isomorphic, which indeed is the case due to Claim 9.13.

Case 2: Let $T \in \mathfrak{E}$. There exists an embedding which does not preserve lev_T , that is the end is not semi-rigid. Such an embedding has a positive period. We have to deal with two subcases.

Subcase 2.1: The end is not regular.

Lemma 9.14. Let $T \in \mathfrak{E}$ and let f be an embedding of T with period k > 0, then $|twin(T)| \ge 2^{\aleph_0}$.

Proof. For $u \in S_f$ and $x \in e(u)$, set T(f,x) := T < e(u), x > Since the end e of T is no regular, for every $u \in S_f$, the ray e(u) contains an infinite subset L such that T(f,x) is not equimorphic to T(f,f(x)) for all $x \in L$.

There exists a set \mathcal{A} of 2^{\aleph_0} infinite pairwise almost disjoint subsets of L. If S_f induces a two-way infinite path enumerate S_f naturally as $\{x_i : i \in \mathbb{Z}\}$. If S_f induces a one-way infinite path enumerate S_f naturally as $\{x_i : i \in \mathbb{N}\}$. Then $f(x_n) = x_{n+k}$ for every $x_n \in S_f$.

Let $A \in \mathcal{A}$. For each $x_n \in A$, let T_A be the tree obtained from T by replacing each of the rooted trees $T(f, x_{n+k})$ by the rooted tree $T(f, x_n)$ and let $T(f, x_{n+k})$ unchanged if $n \notin A$. According to Lemma 4.3, the map f induces an embedding of T into T_A because f maps the rooted tree $T(f, x_n)$ into the rooted tree $T(f, x_{n+k})$, whereas the identity of T_A is an embedding of T_A into T. Hence T_A and T are equimorphic. Say that two sets A and A' are equivalent and set $A \cong A'$ if T_A is isomorphic to $T_{A'}$. If every isomorphism class of sets in A is countable we obtain 2^{\aleph_0} pairwise non isomorphic trees. Hence it suffices to prove that every isomorphism class of sets in A is countable.

Suppose that there is an uncountable family of subsets $(A_{\alpha})_{\alpha<\omega_1}$ of \mathcal{A} which are all in the same isomorphism class. For $\alpha<\omega_1$ let h_{α} be an isomorphism of T_{A_0} onto $T_{A_{\alpha}}$. The map $h_{\alpha}\circ f$ is an embedding of T. For every $\alpha<\omega_1$ choose a vertex $x_{n_{\alpha}}\in S_f\cap S_{h_{\alpha}\circ f}$ which is not an endpoint of S_f nor of $S_{h_{\alpha}\circ f}$. There is an $x_{\overline{n}}\in S_f$ so that $x_{\overline{n}}=x_{n_{\alpha}}$ for uncountably many α . Let $C:=e(x_{\overline{n}})$. Each of those embeddings $h_{\alpha}\circ f$ has an period which is a number in \mathbb{N} and hence there is an uncountable set \mathcal{B} of $\alpha\in\omega_1$ and a number $l\in\mathbb{N}$ so that $x_{\overline{n}}\in S_{h_{\alpha}\circ f}$ and the period of $h_{\alpha}\circ f$ is the same number l for all $\alpha\in\mathcal{B}$. Let α and β in \mathcal{B} . Then $h_{\alpha}\circ f$ and $h_{\beta}\circ f$ have equal restrictions to C.

Let $n > \overline{n} + l - k$ with $n \in A_{\alpha}$ but $n \notin A_{\beta}$. Then the rooted tree $T(C, x_n)$ is isomorphic to the tree $T_{A_{\alpha}}(C, x_{n+k})$ and is embedded by f into the tree $T(C, x_{n+k})$ which in turn is isomorphic to $T_{A_{\beta}}(C, x_{n+k})$. But the tree $T(C_{x_{n+k}})$ can not be embedded into $T(C, x_n)$, for otherwise the trees $T(C_{x_{n+k}})$ and $T(C, x_n)$ would be equimorphic. Hence $T_{A_{\alpha}}(C, x_{n+k})$ can not be embedded into $T_{A_{\beta}}(C, x_{n+k})$ and are therefore not isomorphic. The embedding f maps the tree $T(C, x_{n-l+k})$ into the tree $T_{A_0}(C, x_{n-l+2k})$. Because h_{β} is an isomorphism of T_{A_0} to $T_{A_{\beta}}$, the trees $T_{A_0}(C, x_{n-l+2k})$ and $T_{A_{\beta}}(C, x_{n+k})$ are isomorphic. Because h_{α} is an isomorphic. This implies the contradiction that $T_{A_{\beta}}(C, x_{n+k})$ and $T_{A_{\alpha}}(C, x_{n+k})$ are isomorphic.

Subcase 2.2: The end is regular. We also assume in this Subcase that T is not the one-way infinite path.

Let $v \in \check{E}$ and f be a proper embedding with v the endpoint of \check{S}_f . We are going to construct infinitely many subtrees $Twin_n$ of T, each of them a twin

of T. Let k be the period of f and note that k is a multiple of \mathbf{d} . For $v \in V(T)$ we will denote by C_v , instead of e(v), the ray originating at v.

In the case that S_f is a one-way infinite path, let u be the endpoint of S_f and $v = u \boxplus l$. Because T is not the one-way infinite path there is a number $0 \le i < k$ for which the tree $T(C_v, v \boxplus i) = T(C_u, u \boxplus (l+i))$ contains at least one vertex different from its root $v \boxplus i$. Let $X_n = \{u \boxplus j : 0 \le j \le l + i + 1 + 3nk\}$ with $n \in \mathbb{N}$. The subtree $Twin_n$ for $n \in \mathbb{N}$ of T is obtained from T by removing all trees $T(\to x)$ from T for which $y \to x \in X_n$ and $x \notin C_u$. Then $Twin_n(C_u, x)$ is the singleton x for every vertex $x \in X$. The identity map on $Twin_n$ embeds the tree $Twin_n$ into T and f^m maps T into $Twin_n$ for m > l + i + 1 + 3n. Hence $Twin_n$ is a twin of T for every $1 \le n \in \mathbb{N}$. The embedding f restricted to $Twin_n$ is an embedding of $Twin_n$ and the vertex $v_n := u \oplus (l+i+1+3nk) \in E(Twin_n)$, is the endpoint of the path $\hat{S}_f(Twin_n)$. It follows from Proposition 5.12 that the origin o_n of the tree $Twin_n$ is of the form $v_n \oplus j_n$ with $0 \le j_n \le 2\mathbf{d} \le 2k$. It follows that the level of o_n as a vertex of T is strictly smaller than the level of o_m for n < m and hence that $o_m \in C_{o_n}$ for n < m. Hence if there is an isomorphism, say h, of $Twin_n$ to $Twin_m$ with n < m, then h translates the path C_{o_n} forward onto the path C_{o_m} mapping o_n to o_m according to Proposition 5.12. Hence lev(h(x)) > lev(x) for all vertices in $Twin_n$. Implying that there is no vertex $x \in V(Twin_n)$ with $h(x) = u \in V(Twin_m)$.

In the case that S_f is a two-way infinite path let u be the vertex in S_f with $u \boxplus k = v$ and denote by C the two-way infinite path S_f . Let $X_n = \{v \oplus j : \mathbf{d} \leq j \leq 3nk\}$. For $x = v \oplus j \in X_n$ let $R^{(x)}$ be the tree $T(C, u \oplus i)$ with $0 \leq i < k$ and i congruent to j modulo k. We obtain the tree $Twin_n$ from the tree T by replacing for each $x \in X_n$ the rooted tree T(C, x) by the rooted tree T(C, x) by the rooted tree T(C, x). That is if T(C, x) if T(C, x) = T(C, x) into T(C, x) = T(C, x). (Note that there is no embedding of $Twin_n$ which moves the path from T(C, x) = T(C, x) into a path in T(C, x) = T(C, x) into T(C, x) = T(C, x). The identity maps T(C, x) = T(C, x) into T(C, x) = T(C

Assume that there is an isomorphism h of $Twin_n$ to $Twin_m$ with $1 \le n < m$. Then h maps C_{o_n} onto C_{o_m} and hence $lev(h(x)) - lev(x) =: l \ge k$ for every $x \in V(Twin_n)$. It is not possible that h maps v into C and hence h maps v into one of the rooted trees $Twin_m(C,x)$ with $x \in X$. Let $r \in C$ be the vertex of smallest level with $h(r) \in C$. Then r is a vertex on the oriented path from $v \boxplus 1$ to o_n . Let $x \in C$ with $x \to r$. Then $h(r) = r \boxplus l$ and h(x) is a vertex in $Twin_m(C,h(x))$ and hence every vertex in $Twin_n(\to x)$ is mapped by h into the tree $Twin_m(C,h(x))$. Let y be the vertex in $Twin_n$ with h(y) = x. Then y is not a vertex in $Twin_n(\to x)$ and hence the vertex, say z, of smallest level in C_y and C has to be a vertex in $C_r(Twin_n)$ and hence $y \in Twin_n(C,z)$. But

then $h(y) \in Twin_m(C, z \boxtimes l)$. Implying that $h(y) \neq x$ and hence that h could not have been an isomorphism. This completes Case 2.

We are now ready to complete the proof of Theorem 1.9.

- 9.4. **Proof of Theorem 1.9.** Recall that stable trees are characterized by four types of subgraphs that are preserved by every embedding.
- (i) In this case, T may contain a two way infinite path or a non-regular end preserved by every embedding. In the first case, the conclusion follows from Theorem 8.1. In the second case, T contains a path preserved by every embedding (Theorem 1.3). The conclusion follows by Subcase 2.2.
- (ii) This case follows from the fact that embeddings of a locally finite trees are automorphism provided they fix a vertex.
- (iii) This case follows from Lemma 9.8.
- (iv) This case follows from Lemma 9.14.

References

- [1] A. Bonato, C. Tardif, Mutually embeddable graphs and the tree alternative conjecture. J. Combin. Theory Ser. B 96 (2006), no. 6, 874–880.
- [2] A. Bonato, H. Bruhn, R. Diestel, P. Sprüssel, Twins of rayless graphs. J. Combin. Theory Ser. B 101 (2011), no. 1, 6065.
- [3] R. Diestel, Graph theory. Fourth edition. Graduate Texts in Mathematics, 173. Springer, Heidelberg, 2010. xviii+437 pp.
- [4] R. Halin, Fixed configurations in graphs with small number of disjoint rays, in R.Bodendiek (Ed) Contemporary Methods in Graph Theory, Bibliographisches Inst., Mannheim, 1990, pp. 639-649.
- [5] R. Halin, Automorphisms and endomorphisms of infinite locally finite graphs. Abh. Math. Sem. Univ. Hamburg 39 (1973), 251–283.
- [6] R. Halin, The structure of rayless graphs. Abh. Math. Sem. Univ. Hamburg 68 (1998), 225–253.
- [7] M. Hamann, Personnal communication, Feb 2016.
- [8] M. Hamann, Group action on metric spaces: fixed points and free subgroups, this volume.
- [9] M. Hamann, Self-embeddings of trees. Manuscript, Feb. 2016.
- [10] H. A. Jung, Wurzelbäume und undendliche Wege in Graphen, Math. Nachr. 41 (1969) 1-22.
- [11] D. König, Theorie der Endlichen und Unendlichen Graphen: Kombinatorische Topologie der Streckenkomplexe. 1936 Leipzig: Akad. Verlag.
- [12] C. Laflamme, M. Pouzet, N. Sauer, I. Zaguia, Pairs of orthogonal countable ordinals, Discrete Maths, 335 (2014) 35-44.
- [13] R. Laver, Better-quasi-orderings and a class of trees. Studies in foundations and combinatorics, pp. 31–48, Adv. in Math. Suppl. Stud., 1, Academic Press, New York-London, 1978.
- [14] F. Lehner, Personnal communication, Feb. 2016.
- [15] I. Pays, A. Valette, Sous-groupes libres dans les groupes d'automorphismes d'arbres, L'enseignement Mathématique, 37 (1991) 151–174.
- [16] N. Polat, G. Sabidussi, Fixed elements of infinite trees, in Graphs and combinatorics (Lyon, 1987; Montreal, PQ, 1988), Discrete Math. 130 (1994), no. 1-3, 97–102.
- [17] J. Tits, Sur le groupe des automorphismes d'un arbre, Essays on topology and related topics, pp 188-211, Springer, New-York, 1970.

- [18] M. Tyomkyn, A proof of the rooted tree alternative conjecture, Discrete Math. 309 (2009) 5963-5967.
- [19] W. Woess, Fixed sets and free subgroups of groups acting on metric spaces, Math. Z. 214 (1993) 425-440.

Mathematics & Statistics Department, University of Calgary, Calgary, Alberta, Canada T2N $1\mathrm{N}4$

 $E ext{-}mail\ address: laflamme@ucalgary.ca}$

Univ. Lyon, Université Claude-Bernard Lyon1, CNRS UMR 5208, Institut Camille Jordan, 43 bd. 11 Novembre 1918, 69622 Villeurbanne Cedex, France and Mathematics & Statistics Department, University of Calgary, Calgary, Alberta, Canada T2N 1N4

 $E ext{-}mail\ address: pouzet@univ-lyon1.fr}$

Mathematics & Statistics Department, University of Calgary, Calgary, Alberta, Canada T2N 1N4

 $E\text{-}mail\ address{:}\ \mathtt{nsauer@ucalgary.ca}$