

# A guide to tropical modifications

Nikita Kalinin

May 14, 2024

## Abstract

This paper surveys *tropical modifications*, which have already become a folklore in tropical geometry. Tropical modifications are used in tropical intersection theory, tropical Hodge theory, and in the study of singularities. They admit interpretations in various contexts, such as hyperbolic geometry, Berkovich spaces, and non-standard analysis.

Our main goal is to mention different points of view, to give references, and to demonstrate the abilities of tropical modifications. We assume that the reader has already met “tropical modifications” somewhere and wants to understand them better.

There are novelties here: a new obstruction to the realizability of non-transversal intersections and a tropical version of Weil’s reciprocity law.

AMS2020 codes: 14T10, 14T20, 14T90, 14B10, 05C99

keywords: tropical geometry, Weil reciprocity, tropical modifications, singularity, Hodge theory

**Substance is by nature prior to its modifications.  
... nothing is granted in addition to the understanding,  
except substance and its modifications.**

*Ethics. Benedictus de Spinoza.*

This paper surveys *tropical modifications*, which have already become a folklore in tropical geometry. Tropical modifications are used in tropical intersection theory, tropical Hodge theory, and in the study of singularities. They admit interpretations in various contexts, such as hyperbolic geometry, Berkovich spaces, and non-standard analysis.

One cannot say better than in [12]: “*Tropical modifications ... can be seen as a refinement of the tropicalization process, and allows one to recover some information ... sensitive to higher order terms.*” Tropical modifications were defined by Mikhalkin in [49].

Our main goal is to mention different points of view, to give references, and to demonstrate the abilities of tropical modifications. We assume that

the reader has already met “tropical modifications” somewhere and wants to understand them better.

There are novelties here: a new obstruction (Theorem 2.3) for realizability of non-transversal intersections and a tropical version of Weil’s reciprocity law (Theorem 2.2). A generalization of tropical momentum is given in Section 2.7.

As a preliminary introduction to tropical geometry, see [10], [11] and [52], where tropical modifications are also discussed. We are glad to mention other texts promoting modifications from different perspectives:

**singularities and tangency:** [12] (examples, construction of curves with inflection points), [15] (repairing the  $j$ -invariant of elliptic curves), [16] (tropical genus two curves from several perspectives), [29] (tropical quartics curves), [17] (tropical bitangent lines to a planar quartic), [24, 25] (cuspidal curves counting), [44] (lifting tropical bitangents), [35, 34, 36] (multiplicity of tropical singular points),

**realizability of intersections:** [63] (intersection theory on tropical surfaces), [14] (lifting divisors, constructing curves without maps to an elliptic curve), [45] (studying tropical self-intersections), [54] (realizability of tropical linear subspaces), [71] (tropical  $K3$  surfaces), [60] (tropical constructions of real reducible curve in the Hirzebruch surfaces and other real algebraic varieties), [7] (constructing tropical curves of big genus in  $\mathbb{T}^n$  and other questions about realizability of big Betti numbers), [30] (lifting non-proper tropical intersections),

**maps between tropical curves:** [19] (presenting metric graphs as tropical planar curves), [33, 20] (gonality of tropical curves), [69] (tropical analog of the double ramification locus), [22] (maps to a tree from a modification of a genus  $g$  curve),

**tropical cohomology theory, Hodge theory:** [32] (tropical Dolbeault cohomology), [18] (Chern–Schwartz–MacPherson cycles of matroids), [4] (Homology of tropical fans), [3] (Hodge theory for tropical varieties).

This paper is organized as follows. We define tropical modifications via multivalued operations. Then, we discuss several examples indicating the principal features of the tropical modifications. We prove several structure theorems and discuss applications. Section 4 summarizes intuitive interpretations of the tropical modifications. So, a curious reader can start there for inspiration and then proceed to Section 1.1. In conclusion, we mention several questions that could be solved using tropical modifications.

## Acknowledgement

I want to thank K. Shaw, E. Brugallé, and G. Mikhalkin, who taught me the concept of tropical modification and all others who commented on this text.

# Contents

<b>1</b>	<b>Definitions and examples</b>	<b>4</b>
1.1	Tropical modification via the graphs of multivalued functions	4
1.2	Modification reveals intersections . . . . .	5
1.3	Tropical modifications as limits of amoebas . . . . .	5
1.4	Modification and non-archimedean valuation . . . . .	6
1.5	Examples . . . . .	8
<b>2</b>	<b>Structural theorems about tropical modification</b>	<b>15</b>
2.1	Multiplicity of an intersection . . . . .	15
2.2	Tropical Weil reciprocity law and the tropical momentum map	17
2.3	Application of the tropical momentum to modifications . . . .	19
2.4	Proof of the tropical Weil theorem . . . . .	21
2.5	Stable intersections, realizable intersections . . . . .	22
2.6	Interpretation with chips . . . . .	24
2.7	Digression: a generalization of the tropical momentum . . . .	28
<b>3</b>	<b>Applications of a tropical modification as a method</b>	<b>29</b>
3.1	Inflection points . . . . .	29
3.2	The category of tropical curves . . . . .	29
3.3	Realization of a collection of lines and $(4,d)$ -nets . . . . .	30
3.4	A point of big multiplicity on a planar curve . . . . .	31
<b>4</b>	<b>Intuitions and interpretations</b>	<b>33</b>
4.1	Hyperbolic approach and moduli spaces . . . . .	36
4.2	Non-standard analysis . . . . .	38
<b>5</b>	<b>Questions</b>	<b>39</b>

# 1 Definitions and examples

## 1.1 Tropical modification via the graphs of multivalued functions

Recall that the tropical semi-ring  $\mathbb{T}$  is defined as

$$\mathbb{T} = \mathbb{R} \cup \{-\infty\}, \mathbb{T}^* = \mathbb{T} \setminus \{-\infty\} = \mathbb{R}.$$

We extend addition from  $\mathbb{R}$  to  $\mathbb{T}$  by the rule  $-\infty + A = -\infty$  for all  $A \in \mathbb{T}$ . We extend the order relation from  $\mathbb{R}$  to  $\mathbb{T}$  by the rule  $-\infty < A$  for all  $A \in \mathbb{R}$ . The fastest way to define the tropical modifications is via **multivalued** tropical addition introduced by Viro in [70].

**Definition 1.1.** The set  $\mathbb{T}$  is equipped with tropical multiplication (usual addition on  $\mathbb{R}$ ), and tropical addition, defined as follows:

- the result of tropical multiplication of  $A$  and  $B$  is  $A + B$ ,
- the result of tropical addition of  $A$  and  $B$  is  $\max(A, B)$  if  $A \neq B$ , and
- the result of tropical addition of  $A$  and  $A$  is  $\{x \in \mathbb{T} | x \leq A\}$ .

We may say, equivalently, that the operation  $\max$  is redefined to be multivalued in the case of equal arguments, i.e.,  $\max(A, A) = \{X | X \leq A\}$ .

**Definition 1.2.** A tropical monomial is a function  $f : \mathbb{T}^n \rightarrow \mathbb{T}$  given by

$$f(X_1, X_2, \dots, X_n) = A + i_1 X_1 + i_2 X_2 + \dots + i_n X_n, \quad (1)$$

where  $A \in \mathbb{T}^*$ ,  $(i_1, i_2, \dots, i_n) \in \mathbb{Z}^n$ . It may be thought of as the tropical version of a monomial  $A X_1^{i_1} X_2^{i_2} \dots X_n^{i_n}$ . A tropical polynomial  $f$  is a *tropical* sum of a finite number of tropical monomials,

$$f(X_1, \dots, X_n) = \max_{(i_1, \dots, i_n) \in I} (A_{i_1 \dots i_n} + i_1 X_1 + i_2 X_2 + \dots + i_n X_n), \quad (2)$$

where  $I$  is a finite subset of  $\mathbb{Z}^n$ ,  $A_{i_1 \dots i_n} \in \mathbb{T}^*$ , and the operation  $\max$  is multivalued.

A point  $X' = (X_1, X_2, \dots, X_n)$  belongs to the *zero set* of a tropical polynomial  $f$  if  $-\infty \in f(X')$ , we say that  $f$  *vanishes* at  $X'$ . A tropical hypersurface (as a set) is the zero set of a tropical polynomial on  $\mathbb{T}^n$ .

**Remark 1.1.** Note that the zero set of  $F$  is the set where the maximum in the right-hand side of (2) is attained at least twice; this coincides with the standard definition of a tropical hypersurface [10]. We provide weights to the faces of the tropical hypersurface of maximal dimension. Namely, the face where two tropical monomials with gradients  $(i_1, \dots, i_n), (j_1, \dots, j_n)$  are maximal among others, has the weight  $\gcd(i_1 - j_1, i_2 - j_2, \dots, i_n - j_n)$ . Refer to [49] for details.

**Definition 1.3.** (Tropical modification) Let  $N$  be a tropical hypersurface in  $\mathbb{T}^n$ , i.e., the zero set of a tropical polynomial  $f$  on  $\mathbb{T}^n$ . The *modification* of  $\mathbb{T}^n$  *along*  $N$  is the set

$$m_N(\mathbb{T}^n) = \{(X, Y) \in \mathbb{T}^n \times \mathbb{T} \mid Y \in f(X)\}, \quad (3)$$

i.e., the graph of the multivalued function  $f$ . For a given tropical variety  $K \subset \mathbb{T}^n$ , a tropical subvariety  $K' \subset m_N(\mathbb{T}^n)$  is called a **modification** of  $K$  if the natural projection  $p : \mathbb{T}^n \times \mathbb{T} \rightarrow \mathbb{T}^n$  restricted to  $K'$  is a tropical morphism  $p : K' \rightarrow K$  of degree one. We write  $K' = m_N(K)$  in this case.

**Proposition 1.1** (cf. [49], 1.5 B,C). The set  $m_N(\mathbb{T}^n)$  coincides with the zero set of the tropical polynomial

$$f'(X_1, \dots, X_n, Y) = \max(f(X_1, \dots, X_n), Y) : \mathbb{T}^n \times \mathbb{T} \rightarrow \mathbb{T}.$$

**Definition 1.4.** For an abstract tropical variety  $M$  and its subvariety  $N \subset M$  defined as the zero set of a tropical function  $f : M \rightarrow \mathbb{T}$ , we define the tropical modification  $m_N(M)$  of  $M$  along  $N$  as the graph of  $f$  in  $M \times \mathbb{T}$ . A subvariety  $K' \subset m_N(M)$  is called a modification  $m_N(K)$  of  $K$  along  $N$  if the natural projection  $K' \rightarrow K$  is a tropical morphism of degree one.

Modifications along not-principal divisors, which can be locally presented as zero sets, can be defined locally and then glued to a global abstract tropical variety in a natural way.

## 1.2 Modification reveals intersections

Consider two algebraic curves  $C_1, C_2 \subset (\mathbb{C}^*)^2$  defined by equations

$$F_1(x, y) = 0, F_2(x, y) = 0,$$

respectively. Let us build the map

$$m_{C_2} : (x, y) \rightarrow (x, y, F_2(x, y)) \in (\mathbb{C}^*)^2 \times \mathbb{C}. \quad (4)$$

The set  $m_{C_2}((\mathbb{C}^*)^2)$  is the graph of  $F_2$ , namely  $z = F_2(x, y)$ . The intersection  $C_1 \cap C_2$  can be easily recovered as  $m_{C_2}(C_1) \cap \{(x, y, 0)\}$ . For the complex curves, this does not seem very interesting. Still, during the tropicalization process, the plane  $(x, y, 0)$  goes to the plane  $(X, Y, -\infty) = \{Z = -\infty\}$ , and the intersection of tropical curves will be represented by certain rays going to minus infinity by  $Z$  coordinate, see examples later.

## 1.3 Tropical modifications as limits of amoebas

Look now at what happens in the limiting procedure. Recall that a tropical curve  $C \subset \mathbb{T}^2$  is the tropical limit of a family  $C_t \subset (\mathbb{C}^*)^2, t \in \mathbb{R}$  of plane algebraic curves if in the Gromov-Hausdorff sense, we have

$$C = \lim_{t \rightarrow \infty} \text{Log}_t(C_t)$$

where we apply  $\text{Log}_t : \mathbb{C} \rightarrow \mathbb{T}$  coordinate-wise, i.e. in the two dimensional case

$$\text{Log}_t(C_t) = \{(\log_t |x|, \log_t |y|) | (x, y) \in C_t\}.$$

Let  $F_t$  be the equation of  $C_t$  in  $(\mathbb{C}^*)^2$ .

**Proposition 1.2.** The tropical modification  $m_C \mathbb{T}^2$  of  $\mathbb{T}^2$  along  $C$  is the tropical limit of surfaces

$$S_t = \{(x, y, F_t(x, y)) \in \mathbb{C}^3 | (x, y) \in \mathbb{C}^*\},$$

i.e.  $m_C \mathbb{T}^2 = \lim_{t \rightarrow \infty} \text{Log}_t S_t$ .

Indeed,  $F_t(x, y)$  assumes all the values of a neighborhood of 0 when  $(x, y)$  approaches  $C_t$ . Therefore, after taking the logarithm near  $C$ , the function  $\log_t |F_t(x, y)|$  will assume all the values of a neighborhood of  $-\infty$ .

In other words, for a general point  $(X, Y) \in \mathbb{T}^2$  we have that  $m_C(X, Y) = (X, Y, Z)$  where  $Z$  is computed as  $Z = \lim \log_t |F_t(x_t, y_t)|$  and  $(x_t, y_t)$  are chosen such that  $X = \lim \log_t |x_t|, Y = \lim \log_t |y_t|$ . But if  $(X, Y) \in C$ , then  $(x_t, y_t)$  can be chosen to be close to  $C_t$ , and thus  $Z$  can assume any value from the interval  $F(X, Y) \subset \mathbb{T}$ .

Consider two families  $C_{1,t}, C_{2,t}$  of complex curves with tropical limits  $C_1, C_2$ . Use notation (4).

**Proposition 1.3.** The tropical limit

$$\lim_{t \rightarrow \infty} \text{Log}_t m_{C_{2,t}}(C_{1,t}),$$

of the curves  $m_{C_{2,t}}(C_{1,t}) \subset m_{C_{2,t}}((\mathbb{C}^*)^2) \subset (\mathbb{C}^*)^3$  is a tropical modification  $m_{C_2} C_1$  of  $C_1$ .

Note that  $m_{C_2} \mathbb{T}^2$  depends only on  $C_2$ . Quite the contrary, for given tropical curves  $C_1, C_2$  we **can** construct different families  $C_{1,t}, C_{2,t}$  and the limit  $\lim_{t \rightarrow \infty} \text{Log}_t m_{C_{2,t}}(C_{1,t})$  can be different, see numerous examples below.

## 1.4 Modification and non-archimedean valuation

We always suppose that an algebraic hypersurface comes with a defining equation. Instead of taking the limit of amoebas we can consider non-Archimedean amoebas of the varieties defined over the valuation fields.

**Definition 1.5.** Let  $M' \subset (\mathbb{K}^*)^n$  be a variety over a valuation field  $\mathbb{K}$ . Let  $N' \subset (\mathbb{K}^*)^n$  be an algebraic hypersurfaces defined by an equation

$$f(x) = 0, x \in (\mathbb{K}^*)^n.$$

Consider the graph of  $f$  on  $M$ , i.e.

$$\{(x, f(x)) | x \in M\} \subset (\mathbb{K}^*)^{n+1}.$$

The modification  $m_N M$  of  $M = \text{Trop}(M')$  along  $N = \text{Trop}(N')$  is the non-Archimedean amoeba of the set  $\{(x, f(x)) | x \in M\} \subset (\mathbb{K}^*)^{n+1}$ .

The approach with limits of amoebas gives the same results as the approach with non-Archimedean amoebas.

**Proposition 1.4.** Consider a tropical variety  $M \subset \mathbb{T}^n$  and a tropical hypersurface  $N$  defined by a tropical polynomial  $F$ . Let  $\mathbb{K}$  be the field of power series in  $t$ , converging for  $t$  in a neighborhood of  $0 \in \mathbb{C}$ , and  $\text{val} : \mathbb{K}^* \rightarrow \mathbb{R}$  be its natural valuation (we use convention that  $\text{val}(a+b) \leq \max(\text{val}(a), \text{val}(b))$ , so, for example,  $\text{val}(t^1 + 2t^2) = -1$ ). Suppose that

$$f \in K[x_1, \dots, x_n], f = \sum_{I \in \mathcal{A}} a_I x^I \text{ and } F = \max_{I \in \mathcal{A}} (\text{val}(a_I) + I \cdot X) : \mathbb{T}^n \rightarrow \mathbb{T}$$

where  $I = (i_1, \dots, i_n) \in \mathbb{Z}_{\geq 0}^n$  are multi-indices. Let  $M' \subset \mathbb{K}^n$  be an affine algebraic variety, and its non-Archimedean amoeba  $\text{Val}(M')$  be  $M$ . For a small (by module) complex number  $\varepsilon$ , we can substitute  $t$  as  $\varepsilon$ . Using this substitution we define  $M'_\varepsilon \subset \mathbb{C}^n$  and  $f_\varepsilon \in \mathbb{C}[x_1, \dots, x_n]$ . Then, the three following objects coincide:

- the limit  $\lim_{\varepsilon \rightarrow 0} \text{Log}_\varepsilon(\{x, f_\varepsilon(x) | x \in \mathbb{C}^n\})$ ,
- non-Archimedean amoeba  $\text{Val}(\{(x, f(x)) | x \in (\mathbb{K}^*)^n\}) \subset \mathbb{T}^{n+1}$  of the graph of  $f$ .
- the tropical modification  $m_N(\mathbb{T}^n)$ .

Additionally, two following objects coincide and equal to a tropical modification  $m_N(M)$ :

- the limit  $\lim_{\varepsilon \rightarrow 0} \text{Log}_\varepsilon(\{x, F_\varepsilon(x) | x \in M'_\varepsilon\})$ ,
- non-Archimedean amoeba  $\text{Val}(\{(x, F(x)) | x \in M'\}) \subset \mathbb{T}^{n+1}$  of the graph of  $f$  on  $M'$ .  $\square$

We repeat again that given **only** tropical curves  $C_1, C_2 \subset \mathbb{T}^2$ , in general it is not possible to uniquely “determine” the image of  $C_1$  after the modification along  $C_2$ . That is why a modification of a curve along another curve is multivalued and rather can be used as a *method*. The strategy being usually applied is the following: given two tropical curves, we lift them in a non-Archimedean field (or present them as limits of complex curves, that is the same), then we construct the graph of the function as above and take the non-Archimedean amoeba of the limit of amoebae. Depending on the conditions we imposed on lifted curves (be smooth or singular, be tangent to each other, etc.), we will have a set of possible results (which often can be described using simple combinatorial conditions) for modification of the first curve along the second curve; see examples below.

If  $C_1$  intersects  $C_2$  transversally, then  $m_{C_1}(C_2)$  is uniquely defined. If not, there are the following conditions:

- one equality (via tropical momentum or tropical Menelaus theorem): the sum of the coordinates of all the legs of  $m_{C_2}(C_1)$  going to minus infinity by  $Z$ -coordinate is fixed, see Proposition 2.4;
- one inequality (subordination of divisors): the valuation of the divisor of the intersection of lifted curves is *subordinate* to the stable intersection of  $C_1$  and  $C_2$  (Theorem 2.3).

Both conditions have higher dimensional analogs.

## 1.5 Examples

In this section we study examples of modification, treated as a method. The reader should not be scared by these horrific equations; they are reverse-engineered, starting from the pictures. All the calculations are quite straightforward.

We start by considering the modification of a curve along itself and discuss the ambiguity that appears in this case. Then, we consider how modifications resolve indeterminacy when the intersection of tropical objects is non-transversal. This example promotes the point of view that a tropical modification is the same as adding a new coordinate. So, if one changes coordinates, one can do it via repetitive modifications as in [25].

In the third example, a modification helps recover the position of the inflection point. Also, the usefulness of the tropical momentum and tropical Menelaus Theorem is demonstrated. The tropical Weil theorem that shortens the combinatorial descriptions of possible modification results is proved in Section 2.2.

In the fourth example we study the influence of a singular point on the Newton polygon of a curve. The identical method suits higher dimension and different types of singularities, but more needs to be done due to complicated combinatorics. In the same example, we describe how to find all possible valuations of the intersections of a line with a curve, knowing only their stable tropical intersection – the key tool here is the Vieta theorem. The same arguments may be applied for non-transversal intersections of tropical varieties of any dimension.

**Example 1.1** (Modification along itself). Consider a tropical horizontal line  $L$ , given by  $\max(1, Y)$ . This is the tropicalization of a line of the type  $y = t^{-1} + o(t^{-1})$ . Note that if we make a modification of a line along itself, then all its points go to the minus infinity (Figure 1, left). Indeed, if  $F(x, y)$  is the equation of  $C$ , then the set of points  $\{(x, y, F(x, y)) | (x, y) \in C\}$  belongs to the plane  $z = 0$ , so

$$\text{Val}(\{(x, y, F(x, y)) | (x, y) \in C\}) \subset \{(X, Y, Z) \in \mathbb{T}^3 | Z = -\infty\}.$$

On the other hand, if we consider two different lines  $C_1, C_2$  (with equations  $y = t^{-1}$  and  $y = t^{-1} + t^3$ ) whose tropicalization is  $L$ , then all the points in







Figure 1: Example of a modification of a line along itself. Let  $L_1, L_2$  be defined by  $y = t^{-1}, y = t^{-1} + t^3$  respectively. In each group of pictures, the bottom picture is the initial  $\mathbb{T}^2$ , the middle picture is  $m_{L_1} \mathbb{T}^2$ , the picture at the back is the projection to  $X, Z$ -plane. On the left we see the modification of  $L_1$  along  $L_1$ , on the right we see the modification of  $L_2$  along  $L_1$ . Red line is the result  $m_{L_1} L_1$  (resp.  $m_{L_1} L_2$ ) of the modification.

$m_{C_1} C_2$  have the valuation  $-3$  of  $Z$  coordinate. Again, we see an ambiguity — even if  $L$  is fixed, we can take different lifts of  $L$  and have different results of the modification. On the other hand we can say that the canonical modification along itself is the result similar to Figure 1, left, i.e. we might define  $m_C C$  as the projection of  $C$  to the plane  $Z = -\infty$ . Nevertheless, it is better to always keep this unambiguity in mind instead of giving a precise definition of  $m_C C$ . More about the question of intersections of two curves having the same tropicalization can be found in [45].



(a) Initial picture is below. In the center we see the limit of the graphs of the logarithm of the functions  $F_{2,t}$ . On the picture behind we see the projection of the graph to the plane  $XZ$ . Numbers on the edges are the corresponding weights.

(b) Notation is the same as for the picture on the left. We see the result of the modification in the case when the stable intersection is the actual intersection. The Newton polygon of the curve  $C$  is depicted below.

Figure 2: Example of a modification along a line.

**Example 1.2** (Modification, root of big multiplicity, Figure 2a). In this example we see two tropical curves with non-transverse intersection which hides tangency and genus. Consider the plane curve  $C$ , given by the following equation:  $F(x, y) = 0$ ,

$$F(x, y) = (x - t^{1/3})^3(x - t^{-2}) + t^{-4}xy^2 + (t^{-4} + 2t^{-5})xy + (t^{-5} + t^{-6})x.$$

Its tropicalization<sup>1</sup> is the curve, given by the set of non-smooth points of

$$\text{Trop}(F) = \max(1, 6 + x, 5 + x + y, 4 + x + 2y, 5/3 + 2x, 2 + 3x, 4x).$$

We want to know what is the intersection of  $C$  with the line  $L$  given by the equation  $y + t^{-1} = 0$ . Tropicalizations of  $C$  and  $L$  are drawn on Figure 2a, below, as well as the Newton polygon of  $C$ . The intersection is not transverse, hence we do not know the tropicalization of  $C \cap L$ .

To deal with that, let us consider the map  $m_L : (x, y) \rightarrow (x, y, y + t^{-1})$ . On Figure 2a, in the middle, we see the tropicalization of the set  $\{(x, y, y +$

<sup>1</sup>One can think that we have a family of curves  $C_t$  (given by  $F_t = 0$ ) with parameter  $t$  and its tropicalization is the limit of amoebas  $\lim_{t \rightarrow 0} \text{Log}_t(\{(x, y) | F_t(x, y) = 0\})$ , or that we have a curve  $C$  over Puiseux series  $\mathbb{C}\{\{t\}\} = \mathbb{K}$  given by  $\sum a_{ij}x^i y^j = 0, a_{ij} \in \mathbb{K}$ . Its non-Archimedean amoeba is given by the set of non-smooth points of the function  $\max_{ij}(\text{val}(a_{ij}) + ix + jy)$ . Both ways lead to the same result.

$t^{-1})\}$  and the tropicalization of the image of  $C$  under the map  $m_L$ . Let  $G(x, z)$  be the equation of the projection of  $m_L(C)$  on the  $xz$ -plane. So,  $F(x, y) = 0$  implies that for the new coordinate  $z = y + t^{-1}$  we have

$$G(x, z) = 0, G(x, z) = (x - t^{1/3})^3(x - t^{-2}) + t^{-4}xz + t^{-4}xz^2. \quad (5)$$

Therefore the curve  $C' = pr_{xz}m_L(C)$  is given by the set of non-smooth points of

$$\max(1, 4 + X + Y, 4 + X + 2Y, 2 + 3X, 4X),$$

we see  $C'$  on the projection onto the plane  $XZ$  on the left part of Figure 2a. Notice that in order to recover the transversal intersection of non-Archimedean amoebas we did nothing else as a change of coordinates.

**Remark 1.2.** Consider the restriction of  $\text{Trop}(F)$  on the line  $Y = 1$ . We obtain  $\max(1, 7 + X, 5/3 + 2X, 2 + 3X, 4X) = \max(1, 7 + X, 4X)$ , whose locus of non-linearity corresponds to the stable intersection of our tropical curves. On the other hand, if we restrict  $F$  on the line  $y + t^{-1} = 0$  and only then take the valuation, we obtain  $\max(1, 3X + 2, 4X)$  because  $F(x, -t^{-1}) = (x - t^{1/3})^3(x - t^{-2})$ , and we see that this agrees with the picture of the modifications.

**Definition 1.6.** As we see in this example, a tropical curve in  $\mathbb{T}^n$  typically contains infinite edges. We call them *legs* of a tropical curve. For each leg we have a canonical parametrization  $(a_0 + p_0s, a_1 + p_1s, a_2 + p_2s)$  where  $a_i \in \mathbb{R}, p_i \in \mathbb{Z}, s \in \mathbb{R}, s \geq 0$ , where the vector  $(p_0, p_1, p_2)$ , the *direction* of the leg, is primitive.

Now, on the tropicalization of  $C'$  we see a vertical leg of weight 3,  $Z$ -coordinate goes to  $-\infty$ . That happens because we have the tangency of order 3 between  $C$  and  $L$ , and  $z$  as a function of  $x$  has a root of order 3.

Note that this leg cannot mean that the point is a singular point of  $C$ , because the curve  $C$  (according to criteria of [46] or, more generally [35]) has no singular points, even though the tropicalization of  $C$  has an edge of multiplicity 3.

Thus, this new tropicalization restores the multiplicity of the intersection. We see that the modification of the plane (i.e. amoeba of the set  $\{(x, y, y + t^{-1})\}$ ) is defined, but in codimension one this procedure shows multiplicities of roots and more unapparent structures such as hidden genus squashed initially onto an edge. One can think that this cycle was close to intersection, but after a change of coordinates it becomes visible on the picture of the amoeba of  $C'$ . More examples of this kind can be found in [15, 16].

**Remark 1.3.** Nevertheless, for a general choice of representatives in Puiseux series for these two tropical curves  $\text{Trop}(C), \text{Trop}(L)$ , after modification we will have Figure 2b, which represents stable intersection of the curves.

**Example 1.3** (Modification, inflection point, momentum map). We consider a curve and its tangent line at an inflection point. Suppose, that the intersection of their tropicalizations is not transverse. How can we recover the presence of the inflection point?

We consider a curve  $C$  with the equation  $F(x, y) = 0$  where

$$F(x, y) = y + t^{-3}xy + (t^{-1} + 4 + 6t + 4t^2 + t^3)x^2 + (-t^{-3} - 3 - t - t^2)xy^2 + (t^{-2} - t^{-1} - 2 + t^2 + t^3)x^2y + x^2y^2,$$

and a line  $L$  with the equation  $y = 1 + tx$ . The equation of the curve is chosen just in such a way that the restriction of  $F$  on the line  $L$  is  $t^2(x-1)^3(x-t^{-1})$ , i.e. the point  $(1, 1+t)$  is the inflection point of the curve and  $L$  is tangent to  $C$  at this point.

Tropicalization of the curve is given by the following equation:

$$\text{Trop}(F) = \max(y, x + y + 3, 2x + 1, 2x + y + 2, x + 2y + 3, 2x + 2y). \quad (6)$$

On Figure 3a we see the non-Archimedean amoeba of the image of the curve under the map  $(x, y) \rightarrow (x, y, y - 1 - tx)$ .

In order to find  $X$ -coordinates of the possible legs we can apply the tropical momentum: see Figure 2.3.


**Definition 1.7.** The momentum of a leg  $(A_0 + P_0s, A_1 + P_1s, A_2 + P_2s)$  with respect to a point  $(B_0, B_1, B_2)$  is the vector product  $(A_0 - B_0, A_1 - B_1, A_2 - B_2) \times (P_0, P_1, P_2)$ .

We will prove a (simple) theorem that *the sum of the moments of the legs, counted with their weights, is zero*. Note, that in our case, all the legs we do not know are of the form  $(X_0, Y_0, Z_0 - s)$ , because they are vertical. Refer to Figure 3b. So, we take the vertex  $O$  of the tropical plane, and sum up the vector products  $OX_i \times X_iY_i$  where  $X_iY_i$  are black legs (that we already know) and red legs (which are all vertical). Computation gives us


$$\begin{aligned} & (-4, 0, 0) \times (-1, 1, 1) + (-4, 0, 0) \times (0, -1, 0) + (0, -1, 0) \times (-1, -1, 0) \\ & + (0, -1, 0) \times (1, 0, 1) + (2, 2, 2) \times (1, 0, 1) + (2, 2, 2) \times (0, 1, 1) \\ & + (X, 0, 0) \times (0, 0, -1) + (0, Y, 0) \times (0, 0, -1) + (Z + 1, Z, 0) \times (0, 0, -1) = 0, \end{aligned}$$

i.e.  $(1, -2, 0) + (Y + Z + 1, X + Z, 0) = 0$ , where  $X$  stands for the sum of the  $X$ -coordinates of the vertical legs situated under the line  $(1 - s, 0, 0)$ ,  $Y$  stands for the sum of the  $Y$ -coordinates of the vertical legs under the line  $(1, -s, 0)$ ,  $Z$  stands for the sum of the  $Y$ -coordinates of the vertical legs under the line  $(1, s, s)$ .

On the left picture we see where the red legs are situated. But, since modification of a tropical curve  $C$  along a tropical curve  $C'$  is not canonically



(a) In the center we see a modification of the picture below, its  $XZ$ -projection is on the right, on the left we see it from a different perspective. Doubled red lines graphically represents a line with weight three.



(b) Application of the generalized tropical Menelaus Theorem: we know the direction of the infinite black rays emanating from the tropical curve (in the center on the left), therefore an application of this theorem gives the sum of  $X$ - and  $Y$ - coordinates of red legs, going vertically to the bottom (these legs present exactly the intersection of two considered curves.)

Figure 3: Example of modification in the case of inflection point. The point  $(0,0)$  on the bottom picture is the tropicalization of the inflection point. We modified the black curve along the blue curve, red parts are the parts becoming visible after the modification.

defined<sup>2</sup>, then, for example, a modification of  $C$  could differ from  $C$  just by adding vertical legs at four vertices of the  $C$ : this would correspond to the stable intersection (which is always realizable in the sense that there exists a curve in Puiseux series, such that the valuation of their intersection is the stable intersection)

**Example 1.4.** Singular point, its unique position, and possible liftings of intersection. Consider a curve  $C'$  defined by the equation  $G(x, y) = 0$ , where

$$\begin{aligned} G(x, y) = & t^{-3}xy^3 - (3t^{-3} + t^{-2})xy^2 + (3t^{-3} + 2t^{-2} - 2t^{-1})xy \\ & - (t^{-3} + t^{-2} - 2t^{-1} - 3t^2)x + t^{-2}x^2y^2 - (2t^{-2} - t^{-1})x^2y \\ & + (t^{-2} - t^{-1} - 3t^2)x^2 + t^{-1}y - (t^{-1} + t^2) + t^2x^3. \end{aligned}$$




Figure 4: The extended Newton polyhedron  $\tilde{\mathcal{A}}$  of the curve  $C'$  is drawn in (A). The projection of its faces gives us the subdivision of the Newton polygon of  $C'$ ; see (B). The tropical curve  $\text{Trop}(C')$  is drawn in (C). The vertices  $A_1, A_2, A_3$  have coordinates  $(-2, 0), (1, 0), (4, 0)$ . The edge  $A_1A_2$  has weight 3, while the edge  $A_2A_3$  has weight 2. The point  $P$  is  $(0, 0) = \text{Val}((1, 1))$ .

Let us make the modification along the line  $y = 1$ . For that we draw the graph of the function  $z(x, y) = y - 1$ .

Note that we can easily find the number (with multiplicities) of the vertical legs. Indeed, each edge from  $A_1, A_2, A_3$  going up in direction  $(i, j)$  becomes after the modification a ray going in the direction  $(i, j, j)$ . Therefore, the total momentum of the vertical legs is the sum of  $Y$ -parts of momenta of the edges going up from  $A_1, A_2, A_3$ , that is, 3. Then, if we know that after the modification our curve has a leg of weight 3, then its unique position can be found from the generalized tropical Menelaus theorem. So, in this

<sup>2</sup>If the intersection  $C \cap C'$  is transverse, then the modification is uniquely defined.

case (the points  $(1, 1)$  is of multiplicity 3 for the curve) the pictures after the modifications is as on Figure 5, left. If  $\text{Val}(C') = C$ , but we do not have the other restricting condition, then the picture after the modification can be as in Figure 5, right top, or right bottom, both cases can be realized.




Figure 5: Refer to Example 1.4. Left bottom picture represents a curve  $C$ . On top of it, a modification of it is depicted, with the projection of the latter on the  $XZ$ -plane. On the right side we see two other possible modification of  $C$ .

## 2 Structural theorems about tropical modification

### 2.1 Multiplicity of an intersection

We will recall here certain facts about intersections of tropical curves.

The *multiplicity*  $m(P)$  of the point  $P$  of the transverse intersection of two lines in directions  $(u_1, u_2), (v_1, v_2) \in \mathbb{Z}^2, \gcd(u_1, u_2) = \gcd(v_1, v_2) = 1$  is  $|u_1 v_2 - u_2 v_1|$ .

Given two tropical curves  $A, B \subset \mathbb{T}^2$  we define their *stable intersection* as follows. Let us choose a generic vector  $v$ . Then we consider the curves  $T_{tv}A$  where  $t \in \mathbb{R}, t \rightarrow 0$  and  $T_{tv}$  is translation by the vector  $tv$ . For a generic small positive  $t$ , the intersection  $T_{tv}A \cap B$  is transversal and consists of points  $P_i^t, i = 1, \dots, k$  with multiplicities  $m(P_i^t)$ .

**Proposition 2.1.** Suppose that a horizontal edge  $E$  of a tropical curve  $C$  contains a point  $P$ . Suppose that on the dual subdivision of the Newton polygon for  $C$  the vertical edge  $d(E)$  is dual to  $E$ . Let the endpoints of  $E$  be  $A_1, A_2$  and two faces  $d(A_1), d(A_2)$  adjacent to  $d(E)$  have no other vertical edges. Let the sum of widths in the horizontal direction of the faces  $d(A_1), d(A_2)$  be equal to  $m$ . Then the stable intersection of  $E$  with a horizontal line through  $E$  is  $m$ .

*Proof.* Refer to Example 1.4 and Figure 2b. Let  $L$  be a tropical line containing  $E$  and let the vertex of  $L$  does not coincide with the endpoints of  $E$ . Making the modification along the line  $L$  we see that the sum  $S$  of vertical components of edges going upward from  $A_1, A_2$  equals the sum  $m$  of the  $y$ -components of them.

Then, the sum of vertical components of edges, going downwards, equals  $S$  by the balancing condition for tropical curves. Sum of  $y$ -components of edges in the vertex  $v$  is exactly the width in the  $(1, 0)$  direction of the dual to  $v$  face  $d(v)$  in the Newton polygon.  $\square$

**Definition 2.1** (cf. [61]). For each connected component  $X$  of  $A \cap B$ , we define the *local stable intersection of  $A$  and  $B$  along  $X$*  as  $A \cdot_X B = \sum_i m(P_i^t)$  for  $t$  close to zero, where the sum runs over  $\{i \mid \lim_{t \rightarrow 0} P_i^t \in X\}$ . For a point  $Q \in A$ , we define  $A \cdot_Q B$  as  $A \cdot_X B$ , where  $X$  is the connected component of  $Q$  in the intersection  $A \cap B$ .

**Proposition 2.2** ([12] Proposition 3.11, see also [58] Corollary 12.12). For two curves  $C_1, C_2 \in \mathbb{K}^2$  we consider a *compact* connected component  $X$  of the intersection  $\text{Trop}(C_1) \cap \text{Trop}(C_2)$ . Then,

$$\sum_{x \in C_1 \cap C_2, \text{Val}(x) \in X} m(x) = \text{Trop}(C_1) \cdot_X \text{Trop}(C_2)$$

where  $m(x)$  is the multiplicity of the point  $x$  in the intersection  $C_1 \cap C_2$ .

*Proof.* Consider the equation  $F(x, y) = 0$  of  $C_2$ . We construct the non-Archimedean amoeba  $m_{C_2} C_1$  of  $\{(x, y, F(x, y)) \mid (x, y) \in C_1\}$ . Then

$$\text{Trop}(C_1) \cdot_X \text{Trop}(C_2)$$

is the sum of the weights of the vertical legs of  $m_{C_2} C_1$  under  $X$ . The latter is equal to  $\sum_{x \in C_1 \cap C_2, \text{Val}(x) \in X} m(x)$ .  $\square$

**Remark 2.1.** For non-compact connected components of the intersection we only have an inequality  $\sum_{x \in C_1 \cap C_2, \text{Val}(x) \in X} m(x) \leq \text{Trop}(C_1) \cdot_X \text{Trop}(C_2)$ . It can be upgraded to equality by considering intersections of  $C_1, C_2$  “at infinity”, in the appropriate compactification of the torus, see [64].

For further discussion about the multiplicity of singular points in the tropical world, see [35].



## 2.2 Tropical Weil reciprocity law and the tropical momentum map

The aim of this section is to establish another fact in tropical geometry, obtained as a word-by-word repetition of a fact in classical algebraic geometry.

Weil reciprocity law can be formulated as

**Theorem 2.1.** Let  $C$  be a complex curve and  $f, g$  two meromorphic functions on  $C$  with disjoint divisors. Then

$$\prod_{x \in C} f(x)^{\text{ord}_g x} = \prod_{x \in C} g(x)^{\text{ord}_f x},$$

where  $\text{ord}_f x$  is the minimal degree in the Taylor expansion (in local coordinates) of the function  $f$  at a point  $x$ :

$$f(z) = a_0(z - x)^{\text{ord}_f x} + a_1(z - x)^{\text{ord}_f x + 1} + \dots, a_0 \neq 0.$$

The products in this theorem are finite because  $\text{ord}_g x, \text{ord}_f x$  equal to zero everywhere except finite number of points.

**Definition 2.2.** Define the term  $[f, g]_x = \frac{f(x)^{\text{ord}_g x}}{g(x)^{\text{ord}_f x}} = \frac{a_n^m}{b_m^n} \cdot (-1)^{nm}$  at a point  $x$ , where  $f(z) = a_n(z - x)^n + \dots, g(z) = b_m(z - x)^m + \dots$  are the Taylor expansions of  $f, g$  at the point  $x$ .

If  $f$  and  $g$  share some points in their zeros and poles sets, then we can restate Theorem 2.1 as  $\prod_{x \in C} [f, g]_x = 1$ .

**Example 2.1.** If  $C = \mathbb{CP}^1$  and  $f, g$  are polynomials

$$f(x) = A \prod_{i=1}^n (x - a_i), g(x) = B \prod_{j=1}^m (x - b_j) \quad (7)$$

with  $a_i \neq b_j$ , then

$$\prod_{x \in C} g(x)^{\text{ord}_f x} = B^{nm} \prod_{i=1, j=1}^{n, m} (a_i - b_j), \prod_{x \in C} f(x)^{\text{ord}_g x} = A^{nm} \prod_{i=1, j=1}^{n, m} (b_j - a_i), \quad (8)$$

and their ratio  $(A/B)^{nm}$  is corrected by the term  $[f, g]_\infty$ , because  $f, g$  have a common pole at infinity.

Khovanskii studied various generalizations of the Weil reciprocity law and reformulated them in terms of logarithmic differentials [39, 40, 41]. The final formulation is for toric surfaces and seems like a tropical balancing condition, what is, indeed, the case. The symbol  $[f, g]_x$  is related with Hilbert

character and link coefficient, and is generalized by Parshin residues. Mazin [47] treated them in geometric context of resolutions of singularities.

In order to study what happens after a modification we consider a tropical version of Weil theorem. We need to define tropical meromorphic function and  $\text{ord}_f x$ , see also [52].

**Definition 2.3** ([49]). A tropical meromorphic function  $f$  on a tropical curve  $C$  is a piece-wise linear function with integer slopes. The points, where the balancing condition is not satisfied, are poles and zeroes, and  $\text{ord}_f x$  is the defect in the balancing condition by definition.

**Example 2.2.** The function  $f(x) = \max(0, 2x)$  on  $\mathbb{TP}^1 = \{-\infty\} \cup \mathbb{R} \cup \{+\infty\}$  has a zero of multiplicity 2 at 0, i.e.  $\text{ord}_f(0) = 2$ , and a pole of multiplicity 2 at  $+\infty$ , i.e.  $\text{ord}_f(+\infty) = -2$ .

**Theorem 2.2.** [A proof is in Section 2.4] Let  $C$  be a compact tropical curve and  $f, g$  be two meromorphic tropical functions on  $C$ . Then

$$\sum_{x \in C} f(x) \cdot \text{ord}_g x = \sum_{x \in C} g(x) \cdot \text{ord}_f x.$$

Word-by-word repetition of the reasoning in Example 2.1 proves this theorem in the case  $C = \mathbb{TP}^1$ , because a tropical polynomial  $f : \mathbb{T} \rightarrow \mathbb{T}$  can be presented as

$$f(X) = \sum_{i=1}^n \max(A_i, X),$$

where  $A_i$  are the tropical roots of  $f$ .

For the general statement there are many proofs (and one can proceed by studying piece-wise linear functions on a graph), we give here the shortest one (and also using tropical modifications), via so-called *tropical momentum*.

Suppose that  $C$  is a planar tropical curve. We list all the edges  $E_1, \dots, E_k$  of  $C$ , suppose that their directions are given by primitive (i.e. non-multiple of another integer vector) integer vectors  $v_1, \dots, v_n$ . Suppose that each edge  $E_i$  has weight  $m_i$  and if  $E_i$  is infinite, then the direction of  $v_i$  is chosen to be “to infinity” (there are two choices and for us the orientation of  $v_i$  will be important). Let  $A$  be a point on the plane. Let us choose a point  $B_i \in E_i$  for each  $i = 1, 2, \dots, k$ .

**Definition 2.4** ([72]). Tropical momentum of an edge  $E_i$  of  $C$  with respect to the point  $A$  is given by  $\rho_A(E_i) = m_i \cdot \det(v_i, AB_i)$ .

**Definition 2.5.** For a point  $A \in \mathbb{R}^2$  define  $\rho_A(C)$  as  $\sum_E \rho_A(E)$  where  $E$  runs over all infinite edges of  $C$ .

**Lemma 2.1** ([72]). If a tropical curve  $C$  has only one vertex, then  $\rho_A(C) = \sum_{i=1}^k \rho_A(E_i) = 0$  for any point  $A$  on the plane.

*Proof.* First of all,  $\sum_{i=1}^k \rho_A(E_i)$  does not depend on the point  $A$ , because if we translate  $A$  by some vector  $u$ , then each summand in  $\rho_A(C)$  will change by  $\det(v_i, u) \cdot w_i$  and the sum of changes is zero because of the balancing condition. Therefore,  $\rho_A(C) = 0$ , because we can place  $A$  in the vertex of this curve.  $\square$

**Lemma 2.2** (Moment condition in [72], also it appeared in [51] under the name Tropical Menelaus Theorem). For an arbitrary plane tropical curve  $C \subset \mathbb{R}^2$  and any point  $A \in \mathbb{R}^2$  the equality  $\rho_A(C) = 0$  holds.

*Proof.* Note that the total momentum for a curve is the sum of momenta for all vertices, because a summand corresponding to an edge between two vertices will appear two times with different signs. So, this lemma follows from the previous one.  $\square$

**Definition 2.6.** We consider a tropical curve  $C \subset \mathbb{T}^3$ . Let  $E_1, E_2, \dots, E_n$  be its infinite edges. We define the momentum of  $C$  with respect to  $A$  as  $\rho(A) = \sum_{i=1}^n (v_i \times AB_i) \cdot m_i$  where  $\times$  stands for the vector product,  $v_i$  is the primitive vectors (in the direction “to infinity”) of an edge  $E_i$ ,  $m_i$  is the weight of  $E_i$ , and  $B_i$  is a point on  $E_i$ .

**Proposition 2.3** (Generalized Tropical Menelaus theorem). For a tropical curve  $C \subset \mathbb{T}^3$  and any point  $A$ , the total tropical momentum  $\rho_A(C)$  of  $C$  with respect to  $A$  is zero.

*Proof.* We proceed as in the planar case. We show that  $\rho_A(C)$  does not depend on  $A$  because of the balancing condition. Indeed, if  $C$  has only one vertex, then the claim is trivial. In general case we sum up the tropical momentum by all the edges, and the terms for internal edges appear two times with different signs, which concludes the proof.  $\square$

An application of this theorem can be found in Example 1.3.

## 2.3 Application of the tropical momentum to modifications

**Example 2.3.** Consider the graph of a tropical polynomial

$$f(X) = \max(A_0, A_1 + X, \dots, A_n + nX).$$

Suppose that we know only  $A_0$  and  $A_n$ . Definitely, the positions of the tropical roots of  $f$  may vary, being dependent on the coefficients of  $f$ . Nevertheless, we can apply the tropical Menelaus theorem for the graph of  $f$ . We will calculate the momentum with respect to  $(0, 0)$ . This graph has one infinite horizontal edge with momentum  $A_0$  and one edge of direction  $(1, n)$  with the momentum  $-A_n$ . Also, for each root  $P_i \in \mathbb{T}$  of  $f$  we have an infinite vertical edge with the momentum  $-P_i$ . Application of the tropical moment theorem gives us  $\sum P_i = A_0 - A_n$ , which is simply a tropical manifestation

of Vieta's theorem — the product of the roots  $p_i$  of a polynomial  $\sum_{i=1}^n a_i x^i$  is  $a_0/a_n$ .

**Example 2.4.** Let  $C$  be a planar tropical curve, such that all its infinite edges are horizontal or vertical. Consider first and second coordinates  $X, Y$  on  $C$  as two tropical functions. Denote these functions  $f = X, g = Y$ . Then, Theorem 2.2 says that  $\rho_{(0,0)}C = 0$ , because a tropical root of  $f$  is represented by a horizontal leg of  $C$ , and the value of  $g$  at this root is exactly the  $Y$ -coordinate of this leg.

On Figure 2b, 3a, *a priori* we know only the sum of the directions of the edges with endpoints on the modified curve. We know that there is no horizontal infinite edges (in these examples). In general, it is possible, if the intersection of our two tropical curves is non-compact. Therefore by Weil theorem (or tropical Menelaus Theorem, it is the same) we know the sum of  $X$ -coordinates of the vertical infinite edges. Thus the sum of the weights for red vertical edges equals the sum of the vertical components of the black edges in the Figure 3b.

**Lemma 2.3.** If we make a modification along a horizontal line, then the total vertical slope of the infinite vertical edges under this horizontal line is the total horizontal slope of the region in the subdivision of the Newton polygon, which is dual to the connected component of the intersection of this line with the curve.

*Proof.* The same as for Proposition 2.1. □

**Lemma 2.4.** If the stable intersection of  $\text{Trop}(C)$  with a horizontal line  $L$  is equal to  $m$ ,  $\text{Trop}(C) \cap L$  is compact, and there exists a point  $q \in C$  with  $\mu_q(C) \geq m$  and  $\text{Val}(q) \in \text{Trop}(C) \cap L$ , then we can uniquely recover the position of  $\text{Val}(q)$ .

*Proof.* Indeed, consider a lift  $l$  of  $L$  which passes through  $q$ . If we make the modification along  $l$ , we obtain a leg of  $m_L(\text{Trop}(C))$  under  $\text{Trop}(C) \cap L$  of the weight at least  $m$ . Since the stable intersection  $\text{Trop}(C) \cap L$  is equal to  $m$ , there is only one leg under  $\text{Trop}(C) \cap L$ . Therefore, the tropical momentum theorem gives us the unique position of this leg (of course, it is evident via balancing — we know all the infinite edges of a tropical curve except one, therefore the coordinates of this last edge can be found via the balancing condition). □

**Proposition 2.4** (see [12], Proposition 4.5). For each compact connected component  $C$  of  $C_1 \cap C_2$  the sum of  $X$  coordinates (and the sum of  $Y$ -coordinates) of the valuations of the intersection points of  $C_1, C_2$  with valuations in  $C$  can be calculated just by looking on behavior of  $C_1$  and  $C_2$  near  $C$ .

Indeed, we use tropical Menelaus theorem, this gives us sum of the momenta of all the legs of  $m_{C_2}C_1$  going to  $-\infty$  by  $Z$ -coordinate.

## 2.4 Proof of the tropical Weil theorem

We carry on with a proof of the tropical Weil theorem. Given two tropical meromorphic functions  $f, g$  on a tropical curve  $C$  we want to define the map  $C \rightarrow \mathbb{T}^2, x \rightarrow (f(x), g(x))$  and then use tropical Menelaus theorem (cf. Example 2.4). Here we have to use tropical modification, because *a priori*, the image of tropical curve under the map  $(f, g) : C \rightarrow \mathbb{T}^2$  with  $f, g$  tropical meromorphic functions, is **not** a plane tropical curve: balancing condition is not satisfied near zeroes and poles of  $f$  and  $g$ , we need to add legs there. Formally, we have to consider a modification  $C'$  of  $C$ , and then extend  $f, g$  on it. Then, if the roots and poles of  $f, g$  will be only at 1-valent vertices, then the image of the map  $C' \rightarrow \mathbb{T}^2$  will be a planar tropical curve.

**Definition 2.7.** We call a triple  $(C, f, g)$  of a tropical curve  $C$  and two meromorphic function  $f, g : C \rightarrow \mathbb{TP}^1$  on it *admissible* if all the zeroes and poles of  $f, g$  are located at different one-valent vertices of  $C$ .

**Lemma 2.5.** Given a triple  $(C, f, g)$  of a tropical curve  $C$  and two meromorphic function  $f, g : C \rightarrow \mathbb{TP}^1$  on it, we always can extend the function  $f, g$  on the modification  $D = m_{\text{div}(g)} m_{\text{div}(f)} C$  of  $C$ , such that the obtained triple  $(D, f', g')$  is admissible and

$$\sum_{x \in C} f(x) \cdot \text{ord}_g x - \sum_{x \in C} g(x) \cdot \text{ord}_f x = \sum_{x \in D} f'(x) \cdot \text{ord}_{g'} x - \sum_{x \in D} g'(x) \cdot \text{ord}_{f'} x. \quad (9)$$

*Proof.* We perform tropical modifications of  $C$  in order to have all zeros and poles of  $f, g$  at the vertices of valency one. Namely, for a point  $p$  such that  $p$  is in the corner locus of  $f$  we add to  $C$  an infinite edge  $l$  emanating from  $p$ . We define  $f$  on  $l$  as the linear function with integer slope such that the sum of slopes of  $f$  over the edges from  $p$  is zero, i.e.  $f(x) = f(p) - \text{ord}_f p \cdot x$  where  $x$  is the coordinate on  $l$  such that  $x = 0$  at  $p$  and then  $x$  grows. We define  $g$  on this edge as the constant  $g(p)$ . We perform this operation for all roots and poles of  $f$ . Then, we do the same procedure along the divisor of  $g$ . If a point in  $C$  belongs to both the divisors of  $f$  and  $g$ , we add only one infinite leg.  $\square$

*Proof of the tropical Weil theorem.* By the lemma above we may suppose that the triple  $(C, f, g)$  is admissible. Now  $f, g$  define a map  $C \rightarrow \mathbb{T}^2$  and the image is a tropical curve  $D = \{(f(x), g(x)) | x \in C\}$ : indeed, at every vertex of the image the balancing condition is satisfied; all one-valent vertices go to infinity by one of the coordinates. Now it is easy to verify that  $g(x) \cdot \text{ord}_f(x)$  is a term in the definition of the momentum of  $D$  with respect to  $(0, 0)$ : if  $\text{ord}_f(x) \neq 0$ , then  $D$  has a horizontal infinite edge, and its  $Y$ -coordinate is  $g(x)$ . Finally,

$$\sum_{x \in D} f(x) \cdot \text{ord}_g x - \sum_{x \in D} g(x) \cdot \text{ord}_f x = \rho((0, 0)) = 0. \quad (10) \quad \square$$

**Remark 2.2.** If  $f, g$  come as limits of complex functions  $f_i, g_i$ , having  $\text{ord}_{f_i}(p_i) = k, \text{ord}_{g_i}(p_i) = m, \lim p_i = p$ , then the tropical limit of the family  $\{(f_i(x), g_i(x)) | x \in C_i\}$  will not have vertical (with multiplicity  $k$ ) and horizontal (with multiplicity  $m$ ) leg from a common divisor point  $p$  of  $f$  and  $g$ , but will have one leg of direction  $(k, m)$ . Nevertheless, because of the tropical Menelaus theorem or the balancing condition, it has no influence on Eq. (10).

## 2.5 Stable intersections, realizable intersections

One may ask if the only obstruction for a modification is the generalized tropical Menelaus theorem. As we will see in this section, not at all.

Let us start with a variety  $M' \subset \mathbb{K}^n$  and a hypersurface  $N' \subset \mathbb{K}^n$  and their non-Archimedean amoebas  $M, N \subset \mathbb{T}^n$ . We suppose that the intersection of  $M$  with a tropical hypersurface  $N$  is not transverse. We ask: how does the non-Archimedean amoeba of intersections of  $M' \cap N'$  look like?

First of all, as a divisor on  $M$  (or  $N$ ) it should be rationally equivalent to the divisor of the stable intersection of  $M$  and  $N$ , as it has been shown for the case of curves in [53]. In the general case it follows from the results of this section.

It is easy to find some additional necessary conditions. Let us restrict the equation  $F$  of  $N'$  on  $M'$ , and take the valuations of all these objects  $M', N', F$ . We get some function  $f = \text{Trop}(F)$  whose behavior on a neighborhood of  $N \cap M$  is fixed but its behavior on  $M$  is under the question.

**Definition 2.8.** Let  $M$  be an abstract tropical variety and an embedding  $\iota : M \rightarrow \mathbb{T}^n$  be its realization as a tropical subvariety of  $\mathbb{T}^n$ . Let  $f$  is a tropical function on  $\mathbb{T}^n$ . We define the pull-back of  $\iota^*(f)$  to  $M$  as  $f \circ \iota$ . We call  $\iota^*(f)$  *frozen* at a point  $p \in M$  if  $f$  is smooth at  $\iota(p)$ .

Note that in general the slopes of  $f$  on  $\iota(M)$  does not coincide with slopes of  $\iota^*(f)$  on  $M$  (Example 2.5). From now on we consider tropical functions which have frozen points, the motivation is explained in the following definition.

**Definition 2.9.** A principal divisor  $P$  on an abstract tropical variety  $M$  is called *subordinate* to a principal divisor  $Q$  (we write  $P \prec Q$ ), which is defined by a tropical meromorphic function  $f$  with frozen points, if  $P$  can be defined by a tropical meromorphic function  $h$ , which satisfies  $h \leq f$  everywhere and  $h = f$  at the points where  $f$  is frozen.

**Remark 2.3.** It is straightforward to verify that the fact of being subordinate depends only on  $P, Q$ , and does not depend on particular choice of  $f, h$  as long as the sets of frozen points in  $M$  is fixed.




Figure 6: Refer to Example 2.5. On the left figure we see the vertical part of the modification of the curve given by  $F(x, y) = (-t^{-1} + t^{5/3} + t^{-1}y) + x(t^{-3}y - (t^{-3} + t^{-5/6}) + x^2(t^{-2}y - t^{-2} + t^{-3/2}) + x^3t^2)$  along the line  $y = 1$ . On the right figure, we see the tropicalization of the restriction of  $F$  on  $y = 1$ , i.e., the function  $\max(3X - 2, 2X + 1.5, X + 5/6, -5/3)$ .

**Example 2.5.** Refer to Example 1.4. Let us start from the tropical curve  $M$  given by

$$\max(3 + X + 3Y, 3 + X + 2Y, 3 + X + Y, 3 + X, 2 + 2X + 2Y, 2 + 2X + Y, 2 + 2X, 1 + Y, 1, 3X - 2)$$

and a horizontal line  $N$  given by  $\max(Y, 0)$ . We want to understand the valuations of possible intersections of  $M' \cap N'$  where  $\text{Trop}(M') = M$ ,  $\text{Trop}(N') = N$ .

We can choose the equation for  $M'$  in the form

$$F(x, y) = (t^{-1} + \alpha_0 + t^{-1}y) + x(t^{-3} + \alpha_1 + t^{-3}y^3) + x^2(t^{-2} + \alpha_2 + t^{-2}y) + x^3(t^2 + \alpha_3),$$

where  $\text{val}(\alpha_0) < 1$ ,  $\text{val}(\alpha_1) < 3$ ,  $\text{val}(\alpha_2) < 2$ ,  $\text{val}(\alpha_3) < -2$ . It is clear, that for any  $A \leq 1$ ,  $B \leq 3$ ,  $C \leq 2$  by choosing  $y$  of the form  $1 + \alpha$ ,  $\text{val}(\alpha) < 0$  and then with careful choice for  $\alpha_1, \alpha_2, \alpha_3$  we can obtain (see Figure 6)

$$f(X) = \text{Val}(F(x, 1 + \alpha)) = \max(A, B + X, C + 2X, -2 + 3X).$$

In this example the set  $X \geq 4$  on  $N$  is frozen for  $\text{Trop}(F)$ , that is why we have a choice for the constant term  $A$ . If the intersection is a compact set (as in Example 1.2), then the constant term is also fixed. Note that for the stable intersection our tropical function is  $\text{Trop}(F)(X, 0) = \max(1, 3 + X, 2 + 2X, -2 + 3X)$  and  $f(X) \leq \text{Trop}(F)(X, 0)$  at every point.

Now we prove the following theorem, whose proof consists only of a reformulation in the language of tropical modifications and staring to the pictures; see Remark 1.2 as an illustration.

Fix an abstract tropical variety  $M$ , its tropical embedding  $\iota : M \rightarrow \mathbb{T}^n$ , and a tropical hypersurface  $N \subset \mathbb{T}^n$ , given by a tropical polynomial  $f$ . As we know, the pullback of the divisor of the stable intersection of  $\iota(M)$  with  $N$  is given by  $\iota^*(f)$ . Note that we supply the function  $\iota^*(f)$  with frozen points, according to Definition 2.8.

**Theorem 2.3.** In the above hypothesis, if  $N'$  and  $M'$  are such that  $\text{Trop}(N') = N$ , and  $N' \subset (\mathbb{K}^*)^n$  is given by an equation  $F = 0$ , and  $\text{Trop}(M') = M$ ,  $M' \subset (\mathbb{K}^*)^n$ , then the pullback of  $\text{Val}(N' \cap M')$  to  $M$  is subordinate (Definition 2.9) to the divisor of  $\iota^*(f)$  (Definition 2.8).

*Proof.* Recall that  $f = \text{Trop}(F)$ ,  $f : \mathbb{T}^n \rightarrow \mathbb{T}$ . Let us make the modification of  $\mathbb{T}^n$  along  $N$ . Look at the image  $m_f(M)$  of  $M$  under this map. Clearly, the valuation of the set  $\{(x, F(x)) | x \in M'\}$  belongs to  $m_f(M)$ , therefore the graph of the function  $\text{Trop}(F|_{M'})$  on  $M$  belongs to  $m_f(M)$ . Also,  $\text{Trop}(F|_{M'})$  coincides with  $f$  at the points where  $f$  is smooth. Therefore the pullback of  $\iota^*(\text{Trop}(F|_{M'}))$  is at most  $\iota^*(f)$  everywhere, and  $\iota^*(\text{Trop}(F|_{M'})) = \iota^*(f)$  at the points where  $\iota^*(f)$  is frozen. So, the divisor of  $\iota^*(\text{Trop}(F|_{M'}))$  on  $M$  is subordinate to the pullback of the stable intersection by definition.  $\square$

The graph of  $\text{Trop}(F|_{M'})$  can be lower than the graph of  $\text{Trop}(F)|_M$  because when we substitute the points on  $M'$  to  $F$ , some cancellation can occur, which are invisible when we consider  $\text{Trop}(F)$  as a function on  $\mathbb{T}^n$ . Recall that if the image of the valuation map  $\text{val}$  is  $\mathbb{T}$ , then we know that  $\text{Trop}(F)(X)$  is the maximum of  $\text{val}(F(x))$  with  $\text{Val}(x) = X$ . On the other hand,  $\text{Trop}(F|_{M'})(X)$  for  $X \in M$  is the maximum of  $\text{val}(F(x))$  with  $\text{Val}(x) = X$  and  $x \in M'$ . Clearly, the latter maximum is at most the former maximum.

**Example 2.6.** Refer to Figure 7. We have the stable intersection  $A + B + C + D$  of the curves given by  $\max(0, Y)$  and  $\max(0, X, 2X - 1, 3X - 3, 4X - 6, X + Y, 2X + Y - 1, 3X + Y - 3)$ . The divisor  $A + B' + C' + D$  is rationally equivalent to  $A + B + C + D$ , but the tropical polynomial, which makes rational equivalence, is bigger than the polynomial  $\iota^*(f) = \max(0, X, 2X - 1, 3X - 3, 4X - 6)$  coming from the restriction of the second equation to the line. Therefore,  $A + B' + C' + D$  can not be the valuation of lifts of these curves. This example shows that our conditions are stronger than in [53].

Note that to realize a divisor subordinate to the stable intersection as a tropicalization of intersection one needs to tune the coefficients of  $F$  to have all necessary cancellations. For the case of a planar curve this can be done along trees, [66]. If a curve contains cycles then new conditions appear.

## 2.6 Interpretation with chips

In the case of curves we can represent a divisor on a curve as a collection of chips. In the last subsection we proved Theorem 2.3 which says that






Figure 7: The divisor  $A + B' + C' + D$  is not-realizable as the valuation of intersection of the lifts of the curves defined by  $\max(0, Y)$  and  $\max(0, X, 2X - 1, 3X - 3, 4X - 6, X + Y, 2X + Y - 1, 3X + Y - 3)$ . On the right we see the function which carries this rational equivalence out, it is bigger than the function for the stable intersection and so violates the Theorem 2.3.

any realizable intersection is subordinate to the stable intersection. So, one might ask for a method to produce **all** the subordinate divisors to a given divisor (though, it is possible that not all of them are realizable as the valuation of an intersection).

Let us start with the stable intersection of two tropical curves, this intersection is a divisor (collection of chips) on the first curve. Then we allow the following movement: pushing continuously together two neighbor chips on an edge, with equal speed. We do not allow the opposite operation — when we slide continuously two points apart from each other (so, the operation in Figure 7 does not provide a subordinate to  $A + B + C + D$  divisor).

This corresponds to the following: we look at the modification of the first curve along the second curve, given by a tropical polynomial  $\text{Trop}(F)$ . By decreasing the coefficients of the monomials in  $\iota^*(f)$  on  $C$ , one by one, we can obtain any function less than  $\iota^*(f)$ .

This reasoning can be applied to the intersection of any two tropical varieties if one of them is a complete intersection. We restrict the equations of the second variety on the first, which gives us a stable intersection, then we have a situation similar to Definition 2.9, and, as above, by decreasing the coefficients of pullbacks of tropical polynomials we can obtain all the subordinate to the stable intersection divisors.

**Example 2.7.** Consider the function  $\max(0, X - 1, 2X - 3)$ . This function defines the divisor on  $\mathbb{T}^1$  with two chips, one at  $X = 1$  and the second at  $X = 2$ . When we decrease the coefficient in the monomial  $X - 1$ , these chips are moving closer. For example, the function  $\max(0, X - 1.3, 2X - 3)$  defines the divisor with chips at the points with the coordinates 1.3 and 1.7.

**Remark 2.4.** Note that if the stable intersection is not compact, then we need to add a chip at infinity (or to treat infinity as a point with one chip). Now let  $A, B$  be two chips,  $A$  is at infinity and  $B$  is on the leg of  $V$  going to  $A$ . Then, the operation “pushing together  $A, B$ ” moves only  $B$  towards infinity (and  $A$  remains unchanged at infinity). This corresponds to decreasing the constant term in Example 2.5.

**Example 2.8.** Big order tangency with only two degrees of freedom. ([12], Lemma 3.15). We consider a line  $y - \alpha x - \beta = 0$ ,  $\text{val}(\alpha) = 0$ ,  $\text{val}(\beta) = 0$  and a curve  $a_0 + a_1 y + a_2 x y^l = 0$  with  $\text{val}(a_0) = 0$ ,  $\text{val}(a_1) = 0$ ,  $\text{val}(a_2) = 0$ .

Clearly, we have non-transversal intersection, we can perform substitution  $y = \alpha x + \beta$ , that gives  $a_0 + a_1(\alpha x + \beta) + a_2 x(\alpha x + \beta)^l = (a_0 + a_1 \beta) + x(a_1 \alpha + a_2 \beta^l) + \sum_{i=2}^{l+1} a_2 \beta^{l+1-i} \alpha^{i-1} x^i$ . The contraction may only appear at two coefficients: the coefficient before  $x$  and the constant term. So we have only two degrees of freedom. Let us present the intersection points as chips. By changing the coefficients  $\alpha, \beta, a_i$  we change the intersection, so we can look at how the chips move. So, when  $\text{val}(a_0 + a_1 \beta) < \text{val}(a_0)$ , this corresponds to the movement in Remark 2.4, one chip moves towards infinity

while the others do not move at all. Also we can push two chips together by decreasing the valuation of  $a_1\alpha + a_2\beta^l$ . Note that  $l - 2$  chips at the point  $(0, 0)$  are unmovable.

Here we have only two degrees of freedom because we have only two degrees of freedom in the equation  $a_0 + a_1y + a_2xy^l = 0$ .

**Question 2.1.** Motivated by the above example, we give the following suggestions, which seem to be reasonable in the question of the realizability of intersections. Suppose that we have a tropical line and a tropical curve defined by a tropical polynomial  $f$ . While defining  $\iota^*(f)$  we keep track of all the monomials  $m_i$  of  $f$  and then in Definition 2.9 we allow  $g$  to contain only monomials of the type  $\iota^*(m_i)$ . I.e. if  $f = \max(a_{ij} + iX + jY)$ , then we only allow  $g$  of the type  $\max(c_{ij} + \iota^*(x^i y^j))$  with  $c_{ij} \leq a_{ij}$  which coincides with  $f$  on the frozen set of  $f$ . We explain why we concentrate on the case when one of the curves is a line. Normally, we can perturb the coefficients of the equations of both curves. If one of the curves is a line, we can always suppose that its equation is fixed. For the general case, one should expect that apart from  $\iota^*(f)$  on  $M$  we can find another thin structure, which is responsible for the deformation of the equation of  $M$  being immersed to  $\mathbb{T}^n$ , something like “a pull-back of the normal bundle”, coming from the map  $\iota$ . As we mentioned before, this program can be carried for the case of planar curves [66].

**Example 2.9.** Difference between a leg of big weight and a root. Take the curve  $C$  given by  $F = 0$  where  $F(x, y) = 1 + (t^{-1} + t)x + (2t^{-1} + t^2 + t^4)x^2 + (t^3 + 2t^4)x^3 + t^{-1}xy + 2t^{-1}x^2y$  and intersect it with the line given by  $t^5x + y + 1 = 0$ .

Performing the tropical modification along the line we see that the resulting curve has a leg of weight three going to  $-\infty$ . But it is not a root of multiplicity three! If we substitute  $y = -1 - t^5x$  to the equation, we will see that the obtained polynomial  $1 + tx + t^2x^2 + t^3x^3$  has three roots with the valuation 1, but they do not coincide. But if we consider the curve  $C'$  given by the equation  $F = 0$ ,  $F(x, y) = 1 + (t^{-1} + 3t)x + (2t^{-1} + 3t^2 + t^4)x^2 + (t^3 + 2t^4)x^3 + t^{-1}xy + 2t^{-1}x^2y$ , we see that  $\text{Trop}(C) = \text{Trop}(C')$  and  $C'$  has a tangency of order three with the line.

The same example can be constructed for a similar Newton polygon

$$\text{ConvHull}\{(0, 0) - (1, 1) - (n, 1) - (n + 1, 0)\},$$

where we also can obtain the tangency of the order  $n + 1$ .

**Question 2.2.** Suppose that the intersection of a tropical line with a tropical curve is a segment. Is it always possible to make a modification in order to have a leg of the weight equal to the local stable intersection (Definition 2.1)? If yes, is it always possible to find the coefficients for the equations in order to have a tangency of the order equal to the stable intersection? Also, we can ask this question for any two curves with non-transverse intersections.

Due to combinatorial restrictions in tropical terms, sometimes we can see that it is impossible to have a singular point with high multiplicity on a curve. Note that even in this case, we can have a leg of big multiplicity after the modification; see Example 2.9.

## 2.7 Digression: a generalization of the tropical momentum

A natural generalization of the vector product (or cross product) in  $\mathbb{R}^3$

$$(x_1, y_1, z_1) \times (x_2, y_2, z_2) = (y_1 z_2 - y_2 z_1, x_2 z_1 - x_1 z_2, y_1 z_2 - y_2 z_1) \quad (11)$$

is the following. Given  $k$  vectors  $v_1, v_2, \dots, v_k \in \mathbb{R}^n, k \leq n$  we consider the vector consisting of all the minors  $k \times k$  of the matrix  $k \times n$  constructed as the matrix with the vectors  $v_1, \dots, v_k$  as rows. We call this vector of minors *generalized cross product of  $v_1, v_2, \dots, v_k$* .

Consider a tropical variety  $V^k \in \mathbb{T}^n, k < n$ . Let us choose a basis in each face of  $V$  of codimension one and zero, i.e. for a face  $F$  we choose a basis in the lattice associated with the integer affine structure of this face. For each face  $G$  of codimension one in  $V$  and the faces  $F_1, F_2, \dots, F_l$  of codimension zero, containing  $G$ , we choose vectors  $v_G(F_i)$  which participate in the balancing condition along  $G$ . Now we can define the sign  $s_G(F) \in \{+1, -1\}$  to be  $+1$  if the basis in  $G$  with added vector  $v_G(F)$  at the last place gives the same orientation in  $F$  as the previously chosen basis in  $F$ , and  $-1$  otherwise.

**Definition 2.10.** Let  $\mathfrak{G}(V)$  be the abelian group generated formally by all the faces  $G$  of  $V$  of codimension one. Now we will describe relations in it. For a face  $F \subset V$  of maximal dimension define  $m(F) \in \mathfrak{G}(V)$  to be the sum  $\sum_{G \subset F} s_G(F) \cdot G$ . For each bounded face  $F \subset V$  of maximal dimension we add the relation  $m(F)$  to  $\mathfrak{G}(V)$ .

**Example 2.10.** Compare Definition 2.11 with the proof of Lemma 2.2. For the case of planar tropical curve  $C$  the group  $\mathfrak{G}(V)$  is generated by all the vertices of  $C$ . Then, each internal edge of  $C$  gives the relation that its ends are equal. Therefore, in that case, the group  $\mathfrak{G}(V)$  is  $\mathbb{Z}$  with generator  $\mathbf{1}$ , and for each unbounded  $F$ , we have  $m(F) = \mathbf{1}$ .

**Definition 2.11.** Let  $A$  be a point in  $\mathbb{T}^n$ . Pick a face  $F$  of  $V$  of codimension zero and let  $B$  be a point in  $F$ . Then, define  $r_A(F)$  as the generalized cross product of the vector  $AB$  and the vectors in the basis in  $F$ . Note that  $r_A(F)$  does not depend on  $B$ . Finally, define

$$\rho_A(F) = r_A(F) \otimes_{\mathbb{Q}} m(F) \in \mathbb{R}^{\binom{n}{k+1}} \otimes_{\mathbb{Q}} \mathfrak{G}(V). \quad (12)$$

**Proposition 2.5.** For any point  $A \in \mathbb{T}^n$  we have  $\sum_F \rho_A(F) = 0$ , where  $F$  runs over all the unbounded faces of  $V$  of the maximal dimension.

*Proof.* The structure of the proof is the same as in Lemma 2.1. Let us only show that  $\sum \rho_A(F)$  does not depend on the point  $A$ . Indeed, for each face  $G \subset V$  of the codimension **one** we consider the terms in  $\sum \rho_A(F)$  which contain  $G$ . It is easy to see, that thanks to the balancing condition along  $G$  and our choice of signs, the sum of these terms is zero.  $\square$

**Question 2.3.** It seems that in a general situation, if  $V$  is a tropical curve, then, again,  $\mathfrak{G}(V)$  is  $\mathbb{Z}$ . On the other hand, it seems that if the dimension of  $V$  is at least two, then  $\mathfrak{G}(V)$  is freely generated by the unbounded faces of  $V$  of codimension 1. Also, it would be nice to state an analog of the tropical Weil theorem in this new context and find its classical algebraic counterpart.

### 3 Applications of a tropical modification as a method

One must say that the name “modification” is used in two different senses: the modification as a well-defined *operation* (defined, e.g. in [49]); and a modification along  $N$  as a *method* that reveals a behavior of other varieties in an infinitesimal neighborhood of  $N$ . Namely, performing the modification of  $M$  along  $N \subset M$ , we will see how  $M$  changes, but the objects of codimension one in  $M$  may behave differently, depending on their behavior near  $N$ . We will clarify this distinction with examples.

#### 3.1 Inflection points

An inflection point of a curve is either its singular point, or a point where the tangent line has order of tangency at least 3. It was known before that the number of real inflection points on a curve of degree  $d$  is at most  $d(d-2)$  and the maximum is attainable. The question attacked in [12] is *which topological types of planar real algebraic curves admits the maximal number of real inflection points?* Using classical way to construct algebraic curves – Viro’s patchworking method – the authors construct examples, for what they study possible local pictures of tropicalizations of inflection points. The property to be verified is tangency, but intersection of tropical curve with a tangent line at some point in most cases is not transversal and it is not visible what is the actual order of tangency. To see that, the authors do tropical modifications.

#### 3.2 The category of tropical curves

For the treatment of this question with tropical harmonic maps see [1, 2]. G. Mikhalkin (lectures, 2011) defines the morphisms in the category of tropical curves as all the maps, satisfying the balancing and Riemann-Hurwitz conditions (see, for example [8]) and subject to the *modifiability* condition:

**Definition 3.1.** A morphism  $f : A \rightarrow B$  of tropical curves  $A, B$  is said to be *modifiable* if for any modification  $B'$  of  $B$  there exists a modification  $A'$  of  $A$  and a lift  $f'$  of  $f$  which makes the obtained diagram commutative.

**Proposition 3.1.** The modifiability condition ensures that a morphism came as a degeneration of maps between complex curves (see Section 4.1).

*Sketch of a proof.* After a number of modifications we may have the map  $f'$  contracting no cycles. Then we construct a family of complex curves  $B_i$  such that  $\lim B_i = B'$  in the hyperbolic sense (see section 4.1). Finally, since  $f'$  should come as a tropicalization of a covering, the complex curves  $A_i$  with  $\lim A_i = A'$  are constructed as coverings  $f_i : A_i \rightarrow B_i$  over  $B_i$  where the combinatorics (ramification profiles, local degrees at points of tori contracting to tropical edges) of  $f_i$  is prescribed by  $f'$ . Balancing and Riemann-Hurwitz conditions follow.  $\square$

### 3.3 Realization of a collection of lines and $(4, d)$ -nets

Which configuration of lines and points in  $\mathbb{P}^2$  with given incidence relation are possible? That is a classical question and even for seemingly easy data the answer is often not clear.

**Definition 3.2.** A  $(4, d)$ -net in  $\mathbb{P}^2$  is four collections by  $d$  lines each of them, such that exactly four lines pass through any point of intersection of two lines from different collections, all these four lines are from different collections.

It is not clear whether a  $(4, d)$ -net exists for  $d \geq 5$ . In [28] the authors proved, using tropical geometry, that there exists no  $(4, 4)$ -net.

One of the key ingredients is the following: if some net exists in the classical world, then it exists in the tropical world. The problem appears: if we have more than three tropical lines through a point on a plane, then the intersection of two of them will be non-transversal. However, thanks to modifications we always can have transversal intersections, but probably in the space of bigger dimensions. For that, we just do modifications along lines that have non-transversal intersection. After these modifications, all intersections become transversal and the modified lines go to infinity. Then, let us think about the following theorem, announced by the authors of [28], from the point of view of modifications:

**Question 3.1.** If some combinatorial data (required dimensions of intersections of linear spaces) can be realized in  $\mathbb{P}^k$  by a collection of linear spaces, does there exists a collection of tropical linear spaces that realize the same combinatorial data in  $T\mathbb{P}^{k'}$  with  $k' \geq k$ ?

Indeed, consider this realization in  $\mathbb{P}^k$ . By passing to the tropical limit we obtain a tropical configuration, but the intersection dimensions may increase. Then, by doing the modifications, we want to repair the correct dimensions. Is it always possible to achieve?

### 3.4 A point of big multiplicity on a planar curve

In its most general form, this question could be formulated as follows: given a cohomological class  $a$  of subvariety  $S$  in a bigger variety, how many singularities  $S$  may have? For example, is it possible for a surface of degree 4 in  $\mathbb{CP}^4$  to have four double points and three two-fold lines?

There are several reasons why tropical geometry may provide tools for such questions. We will demonstrate these tools in the case of curves, where this deed has been already done. Combinatorics of a planar tropical curve is encoded in the subdivision of its Newton polygon. A singular point of multiplicity  $m$  influences a part of the subdivision of area of order  $m^2$  ([35]), what is in accordance with the order of the number of linear conditions  $\binom{m(m+1)}{2}$  that a point of multiplicity  $m$  imposes on the coefficients of the curve's equation. For a general treatment of the tropical singularities, see [35], [36] and Chapters 1,2 in [34].

In this section, we will only demonstrate how to apply modification techniques in this problem, though we will obtain a weaker estimation – but still of order  $m^2$ .

The idea is the following: if a curve  $C$  has a point  $p$  of multiplicity  $m$ , then for each curve  $D$ , passing through  $p$ , the local intersection of  $C$  and  $D$  at  $p$  is at least  $m$ . The multiplicity of a local intersection of  $C$  and  $D$  can be estimated from above by studying the connected component, containing  $\text{Val}(p)$ , of the stable intersection  $\text{Trop}(C) \cap \text{Trop}(D)$  for the non-Archimedean amoebas of  $C$  and  $D$ , see Theorem 2.2.

Here is method: we take the polynomial  $F$  defining  $D$ , and use the fact that the image of  $C$  under the map  $m_D : (x, y) \rightarrow (x, y, F(x, y))$  intersects the plane  $z = 0$  with multiplicity at least  $m$ . That implies **existence** of a modification of  $\text{Trop}(C)$  along  $\text{Trop}(D)$ , which has a leg of weight  $m$  going in the direction  $(0, 0, -1)$ , exactly under the point  $\text{Val}(p)$ . The latter modification is obtained just by taking the non-Archimedean amoeba of  $m_D(C) \subset m_D(\mathbb{P}^2)$ .

Now we reduce the problem to its combinatorial counterpart: is it possible for two given tropical curves, that after the modification along the second, the first curve will have a leg of weight  $m$ , which projects exactly on the given point  $\text{Val}(p)$ ? After some work with intrinsically tropical objects, we will get an estimate of this point's influence on the Newton polygon of the curve.

We are not going to consider this problem in the full generality, so we will have a close look at the simplest interesting example.

**Proposition 3.2.** Suppose that a horizontal edge  $E$  of a tropical curve  $C$  contains a point  $\text{Val}(p)$  where  $p$  is of multiplicity  $m$  for a curve  $C'$  such that  $\text{trop}(C') = C$ . Denote by  $d(E)$  the vertical edge in the dual subdivision of the Newton polygon which is dual to  $E$ . Let the endpoints of  $E$  be  $A_1, A_2$  and two faces  $d(A_1), d(A_2)$  adjacent to  $d(E)$  have no other vertical edges.

Then the sum of widths of the faces  $d(A_1), d(A_2)$  is at least  $m$ , so their total area is at least  $m^2/2$ .

*Proof.* Suppose that  $p$  is of multiplicity  $m$  for  $C'$ . Let us take a line  $D$  through  $p$ , such that  $\text{Trop}(D)$  contains inside its vertical edge the point  $\text{Val}(p)$ . Clearly the local intersection  $\text{Trop}(C') \cap \text{Trop}(D)$  is one point, and the multiplicity of this point should be at least  $m$ . That immediately implies that the weight of  $E$  is at least  $m$ . Hence the lattice length of  $d(E)$  is at least  $m$ .

Let us look at the dual picture in the Newton polygon. Two faces  $d(A_1), d(A_2)$  adjacent to the vertical edge have the sum of width in the  $(1, 0)$  direction at least  $m$  (by Proposition 2.1),  $d(E)$  has length  $m$ , so the sum of the areas of  $d(A_1), d(A_2)$  is at least  $m^2/2$ .  $\square$

**Remark 3.1.** Note that if the stable intersection of  $\text{Trop}(C)$  with the horizontal line is  $m$ , then we can uniquely determine the position of the valuation of the singular point, see Lemma 2.4.

What to do if there is a usual horizontal line  $L$ , a part of  $C$ , through  $\text{Val}(p)$ ? We perform the modification along this horizontal line  $L$ . If a part of the curve goes to the minus infinity, that means that we can divide the equation  $F$  of  $C'$  by an equation of  $D$ . That means that the Newton polygon of  $C$  has two parallel vertical sides. The components of the modification which do not go to the minus infinity do not contribute to the singularity.

However, it is possible that  $d(A_1), d(A_2)$  have other vertical sides besides  $d(E)$ . Let  $\mathfrak{E}$  be the stable intersection of  $\text{Trop}(C)$  and the horizontal line; clearly  $E \subset \mathfrak{E}$ . Now, let us compute the sum of the areas of the faces  $d(V)$  corresponding to vertices  $V$  of  $\text{Trop}(C)$  on  $\mathfrak{E}$ . It is possible that more than two faces correspond to one singular point, if the edge with the singular point has an extension, see again Example 1.4.

Suppose that a tropical curve has edges  $A_1A_2, A_2A_3, \dots, A_{k-1}A_k$  and  $A_1, A_2, \dots, A_k$  are situated on a horizontal interval  $A_1A_k = \mathfrak{E}$ . Suppose that  $p$ , point of multiplicity  $m$ , is on the edge  $A_sA_{s+1}$ . Making a modification along a line containing  $A_1A_k$  in its horizontal ray we estimate only the common width of faces corresponding to  $A_1, A_2, \dots, A_k$ , which gives no good estimate for the sum of areas of  $d(A_i)$ .

But we can make a modification along a quadric.

**Lemma 3.1.** In the above hypothesis, the sum of areas of all faces

$$d(A_1), d(A_2), \dots, d(A_k)$$

is at least  $m/2 + m^2/4$ .

*Proof.* Let  $a_i = \omega_{(1,0)}(d(A_{s+i}))$ ,  $i \geq 1$  be the width of  $i$ -th face (i.e.  $d(A_{s+i})$ ) on the right,  $b_i = \omega_{(1,0)}(d(A_{s-i}))$ ,  $i \geq 0$  be the width of  $i$ -th face (i.e.  $d(A_{s-i})$ )



on the left. Let  $c_i$  be the lattice length of  $i$ -th vertical edge on the right (i.e.  $c_i = \omega_{(0,1)} d(A_{s+i} A_{s+i+1}), i \geq 1$ ),  $d_i$  be the length of the  $i$ -th vertical edge on the left (i.e.  $d_i = \omega_{(0,1)} d(A_{s-i} A_{s-i+1}), i \geq 1$ ). Then, let  $\sum_{i=1}^k a_i = A_k, \sum_{i=1}^k b_i = B_k$ . With the same calculations as above, making the modification along a piece of a quadric with vertices on  $A_{s-j} A_{s+1-j}$  and  $A_{s+i} A_{s+1+i}$  we get  $A_i + c_i + B_j + d_j \geq m$  for all pairs  $i, j$ . Denote  $\min_i (c_i + A_i) = A, \min_i (d_j + B_j) = B$ , so  $A + B \geq m$ .

Then,  $c_i \geq A - A_i, d_j \geq B - B_j$ . Sum  $S$  of areas can be estimated as

$$2S \geq (m + c_1)A_1 + \sum (A_{i+1} - A_i)(c_i + c_{i+1}) + (m + d_1)B_1 + \sum (B_{i+1} - B_i)(d_i + d_{i+1})$$

$$\begin{aligned} 2S &\geq (m + A - A_1)A_1 + \sum (A_{i+1} - A_i)(A - A_i + A - A_{i+1}) + \\ &\quad (m + B - B_1)B_1 + \sum (B_{i+1} - B_i)(B - B_i + B - B_{i+1}) \geq \\ &\quad A_1(m - A) + A^2 + B_1(m - B) + B^2 \geq m + m^2/2. \end{aligned}$$

So,  $S \geq m/2 + m^2/4$ .

□

## 4 Intuitions and interpretations

*La science toujours progresse et jamais ne faillit,  
toujours se hausse et jamais ne dégénère,  
toujours dévoile et jamais n'occulte.*

*Anonyme.*

This section explains why a tropical modification is a natural notion and gives several interpretations of a modification in different contexts. The reader, interested in definitions, examples, and theorems, should directly proceed to the previous sections, and return here only for inspiration or references.

Tropical modifications were introduced in the seminal paper [48] as the main ingredient in the tropical equivalence relation. Namely, two tropical varieties are *equivalent* (tropical counterpart of birational isomorphism) if they are related by a chain of tropical modifications and reverse operations. For the full definition of an abstract tropical variety, refer to [52] and [50].

The underlying idea is as follows. Recall, that a tropical variety  $V$  can be decomposed into a disjoint union of a compact part  $V_c$  and a non-compact part  $V_\infty$ , and  $V = V_c \cup V_\infty$ . Moreover,  $V$  retracts on  $V_c$ . Then, the set  $V_\infty$  consists of “tree-like” unions of hyperplanes’ parts. We call these parts *legs* in the one-dimensional case and *leaves* in general situation. For tropical curves,  $V_\infty$  is a union of half-lines. For example, for a tropical elliptic curve (see Figure 8, left side) the set  $V_c$  is the ellipse, and  $V_\infty$  is the set of trees growing on the ellipse.




Figure 8: On the left side we see a tropical elliptic curve  $V$  which is a part of the analytification of an elliptic curve. The ellipse is  $V_c$  and the union of tree-like pieces is  $V_\infty$ . On the right side we see a tropical rational curve  $V$ , which is equal to  $V_\infty$ . Each point  $x$  of  $V$  can serve as  $V_c$ , because  $V$  contracts onto any of its point  $x \in V$ .

**Remark 4.1.** On a tropical rational<sup>3</sup> variety  $V$ , each point may be chosen as  $V_c$ , see Figure 8 right side.

Consider the tropical limit  $V$  of algebraic varieties  $W_i \subset (\mathbb{C}^*)^n$ , i.e.  $V = \lim_{i \rightarrow \infty} \text{Log}_{t_i}(W_i)$ , where we apply the map  $\text{Log}_{t_i} : \mathbb{C}^* \rightarrow \mathbb{R}, x \rightarrow \log_{t_i} |x|$  coordinate-wise and  $\{t_i\}_{i=1}^\infty$  is a sequence of positive numbers, tending to  $+\infty$ . In this case the set  $V_\infty$  encodes the topological way of how  $W_i$  approach some compactification of  $(\mathbb{C}^*)^n$ . For the moment, the particular choice of the compactification does not matter<sup>4</sup>.

Besides, for  $i$  big enough, the Bergman fan  $B(W_i) := \lim_{t \rightarrow \infty} \text{Log}_t(W_i)$  of  $W_i$  is equal to  $\lim_{t \rightarrow \infty} \frac{1}{t}V$ . The latter limit is obtained by contracting the compact part  $V_c$  of  $V$ , so the Bergman fan can be restored by  $V_\infty$ . Note, that  $V$  came here with a particular immersion to  $\mathbb{R}^n$ .

**Example 4.1.** If curves  $W_i, i = 1, 2, \dots$  in  $(\mathbb{C}^*)^2$  all have branches with asymptotic  $(s^k, s^l)$  with a local parameter  $s \rightarrow \infty$ , then the tropical limit  $V$  of this family lies in  $\mathbb{R}^2$ , and  $V$  has the infinite leg (half-line) in the lattice direction  $(k, l)$ .

Let us suppose that we have an algebraic map  $f : (\mathbb{C}^*)^n \rightarrow (\mathbb{C}^*)^m$ , and  $f$  is in general position with respect to the family  $\{W_i\}$ , i.e. for each  $i$  big enough, the image  $f(W_i)$  is birationally equivalent to  $W_i$ . Let  $V'$  be the tropical limit of the family  $\{f(W_i)\}$ . One can prove that  $V'_\infty$  differs from  $V_\infty$  by adding new half-planes and contracting other half-planes. These half-planes grow along the tropicalization of zeros and poles of  $f$  on  $W_i$  (exactly as

<sup>3</sup>Rational tropical varieties are the contractible ones, as a topological space. They are not well studied even in small dimensions. For example, there exist algebraic three-dimensional cubic hypersurfaces which are not rational. It is not known whether we can see this tropically, because all tropical cubic surfaces are contractible.

<sup>4</sup>For a fixed compactification, see the notion of sedentarity in [65] and [10], p. 44.

in Definition 1.3). This consideration suggests the ideas of *modification* and, subsequently, *tropical equivalence*. The name “modification” was borrowed from complex analysis, and tropical modification is sometimes called “tropical blow-up”.

In Section 3.2 we see how the notion of modifications allows us to define the *category of tropical curves*. This category keeps track of birational isomorphism in the category of complex algebraic curves. See also §2.2, where making modifications for curves simplifies a proof to some extent.

Alternatively, tropical geometry can be thought as studying of skeletons of analytifications of algebraic varieties, see Figure 8, the analytification of an elliptic curve on the left, the analytification of  $\mathbb{P}^1$  on the right. The analytification  $X^{an}$  of a variety  $X$  is the set of all seminorms on functions on  $X$ . Each point  $x \in X$  defines such a seminorm by measuring the order of vanishing of a function at  $x$ , on Figure 8 these points are represented by the ends of leafs (also these valuations represent the norms with “zero” radius). The analytification of an elliptic curve is the injective limit of all modifications of its tropicalization, i.e. we add a leg at every points of a circle, then we add a leg at every points of this new space, etc.

For the sake of shortness, we refer the reader to a nice introduction in Berkovich spaces, with a bit of pictures [5],[68] and to [6] to see how it has been applied to tropical geometry (also, see on the page 7 in [6], using of log reminds hyperbolic approach).

We can obtain a tropical variety  $V$  as the non-Archimedean amoeba of an algebraic variety  $W$  over a non-Archimedean field. This approach (see section §4.2) finally suggests the same idea of equivalence up to modification, because the analytification  $W^{an}$  is the injective limit of all “affine” tropical modifications (i.e. along only principal divisors) of  $V$  (see [57]). Berkovich proved that  $W^{an}$  retracts on a finite polyhedral complex, so  $V_c$  is a deformation retract of  $W^{an}$ . Even better, the metric on  $W^{an}$  agrees with the metric on  $V$  for the case of curves<sup>5</sup> ([6]). For elliptic curves  $V_c$  will be a circle in both tropical and analytical cases, and its length is prescribed by the  $j$ -invariant of the considered curve ([15]).

This connection between tropical geometry and analytic geometry leads to the questions of *lifting* or *realizability*, i.e. what could be the intersection of two varieties  $X, Y$  if we know the intersection of their tropicalizations? If their tropicalizations  $\text{Trop}(X), \text{Trop}(Y)$  intersect transversally, the answer is relatively simple, see [55]. If the intersection of  $\text{Trop}(X), \text{Trop}(Y)$  is non-transverse, then we can lift *the stable intersection* of these tropical varieties, see [56],[58].

This raised the following question: to what extent the only condition for a divisor on a curve to be realizable as an intersection is to be rationally

---

<sup>5</sup>That should be true for varieties of any dimension, modulo integer affine transformations, but no proof has appeared yet. For the skeletons in higher dimensions see [26, 27].

equivalent to the stable intersection (cf. [53], Conjecture 3.4)?

Tropical modification (as a *method*) helps dealing with such questions. It is known that being rationally equivalent to the stable intersection is not enough. We consider other existing obstructions (in fact, equivalent to Vieta theorem) for what can happen in non-transverse tropical intersections, and prove, for that occasion, the tropical Weil reciprocity law by using the tropical momentum Lemma 2.2.

Consequently, modifications are used in tropical intersection theory ([63, 64]), to define the intersection product. Nevertheless, one must use modifications along non-Cartier divisors (Examples 1.1.37, 3.4.18 in [64], for moduli space of five points on rational curve) and even along non-realizable subvarieties – for a proof that they are non-realizable as tropical limits.

As we stated before, one should think that a tropical modification along  $X$  reveals asymptotical behavior of objects near  $X$ . We can find an analogy in non-standard analysis: the tropical line is the hyperreal line, the modification at a point is an approaching this point with *an infinitesimal telescope*, see Figure 10 and Section 4.2. In order to define tropical Hopf manifolds one should also use the modifications to study certain germs [62].

Given a surface with hyperbolic structure, we can make a puncture at  $x$ . This changes the hyperbolic structure and  $x$  goes, in a sense, to “infinity”. A tropical curve can be obtained as a degeneration of hyperbolic structures, and making a puncture at  $x$  results as the modification at the limit of  $x$ , see Section §4.1.

A modification can be described as a graph of a function, if we use the convention about multivalued addition, brought in tropical geometry by Oleg Viro ([70]), see Section 1.1.

The other applications of tropical modification as a *method* are following. Passing to tropical limit squashes a variety, and some local features become invisible. In order to reveal them back we can do a modification (whence also this metaphor “look in an infinitesimal microscope”). For example, modifications allow us to restore transversality between lines if we have lost it during tropicalization (§3.3), then it allows us to see  $(-1)$ -curves on del Pezzo surfaces ([59]). Methods of lifting non-transverse intersections leads us to use modifications in questions about singularities: inflection points – [12], singular points – [46]. As an example (Section 3.4), we use modification in the study of singular points of order  $m$  (but obtain weaker results than in [35]).

## 4.1 Hyperbolic approach and moduli spaces

Consider a tropical curve  $C$  given as the tropical limit of complex curves  $C_i$ . From the point of view of hyperbolic geometry, a modification of  $C$  at a point  $x \in C$  means just making a puncture  $x_i$  in  $C_i$ , with condition that  $x_i \rightarrow x$ . To explain this we need to know how to directly construct tropical curves via limits of Riemann surfaces with hyperbolic structure on them, without

any immersions<sup>6</sup>.

So, for details how tropical geometry can be built on on the ground of hyperbolic geometry, see [43]. Here we briefly sketch the construction.




Figure 9: We draw the tropical limits of Riemann surfaces, and a surface close to the limit. Modification adds a puncture to each curve in the family and a leg to the tropical curve.

The approach, proposed by L. Lang, uses the collar lemma ([13]). This lemma simply says that any closed geodesic of length  $l$  has a collar of width  $\log(\coth(l/4))$  and what is more important, for different closed geodesics their collars do not intersect, see Figure 9. That is also important that smaller geodesics have bigger collars (and, intuitively, a puncture has the collar of infinite width).

Thus, given a family of curves  $C_i$  (of the same genus), we consider a fixed pair-of-pants decomposition by geodesics  $L_i$ . The tropical curve is constructed as follows: its vertices are in one-to-one correspondence with the pair-of-pants, each shared boundary component between two pairs-of-pants correspond to an edge of the tropical curve, and the collar lemma furnishes us with the length of the edges of the tropical curve as the logarithms (with base  $t$ , and  $t \rightarrow \infty$  as the hyperbolic structure degenerates) of widths of the collars of  $L_i$ 's. Compare this approach with [9].

What will happen if we make a puncture? A puncture is the limit of small geodesic circles. Cutting out a disk with radius  $t^{-n}$  adds a leaf of finite length  $n$ , as it is seen from the above description. Therefore, cutting

<sup>6</sup>Usually people consider curves  $C_i$  in toric variety  $X$  and then they consider degeneration of complex structures on  $X$ .

out a point results in adding an infinite edge, i.e. a modification.

That explains why a permanent using of graphs for moduli space problems is actually useful ([42], cf. [38]). Tropical curves describe the part of boundary of a moduli space, and modification corresponds to marking a point (read [21] to see the hyperbolic view on moduli space problems), which are punctures from the hyperbolic point of view (see applications to moduli space of points [50]). Tropical differential forms are also defined in this manner while taking a limit of hyperbolic structure [52].

## 4.2 Non-standard analysis

Non-standard analysis appeared as an attempt to formalize the notion of “infinitesimally small” variables (see §4 of [67] for a nice and short exposition).




Figure 10: Similarity in the pictures while using an infinitesimal microscope (left) and the tropical modification at points 1 and  $1 + \varepsilon$  (right).

There is a way to understand tropical geometry via nonstandard analysis (cf. §1.4 [31]). Figure 10 shows that tropical modifications are similar to “infinitesimal microscope” for the hyperreal line in the terminology of [37], and this interpretation in computational sense is the same as for Berkovich spaces: doing modification at the point  $x = 1$  on a curve is adding a leg to the tropical curve, which ranges points according to their asymptotical distance to  $x = 1$ , i.e.  $\text{val}(x - 1)$ , these pictures are also similar to the hyperbolic ones (Figure 9). Dotted lines represent directions to the end points of the analytifications, we have similar type of branching at all points in Figure 8.

It is worth to note that there are still no applications of this point of view, neither in tropical geometry, nor in non-standard analysis. However, Berkovich spaces can be understood as a modern version of non-standard analysis, and tropical modification has applications there.

We should say that

## 5 Questions

An important feature of tropical geometry is that it erects a bridge from a very geometric things (hyperbolic geometry) to very discrete things as  $p$ -adic valuations and non-Archimedean analysis. As tropical modifications dwell in both realms, we expect their fruitful use in future.

**Question 5.1.** Explore tropical modifications over the phase-tropical numbers [70].

For example, in [23] tropical geometry was used to construct a system of polynomial equations with only 5 monomials and 6 positive solutions, and the authors used non-transverse intersections. Looking at questions of such type with a modification tool over phase tropical numbers looks promising.

**Question 5.2.** Study the tangency between tropical curves and tropical surfaces in  $\mathbb{T}^3$ .

For example, one can ask how many surfaces of a given degree are tangent to a given collection of lines (or twisted cubics) in  $\mathbb{T}^3$ .

## References

- [1] O. Amini, M. Baker, E. Brugallé, and J. Rabinoff. Lifting harmonic morphisms I: metrized complexes and Berkovich skeleta. *Res. Math. Sci.*, 2:2:7, 2015. 29
- [2] O. Amini, M. Baker, E. Brugallé, and J. Rabinoff. Lifting harmonic morphisms II: Tropical curves and metrized complexes. *Algebra Number Theory*, 9(2):267–315, 2015. 29
- [3] O. Amini and M. Piquerez. Hodge theory for tropical varieties. *arXiv preprint arXiv:2007.07826*, 2020. 2
- [4] O. Amini and M. Piquerez. Homology of tropical fans. *arXiv preprint arXiv:2105.01504*, 2021. 2
- [5] M. Baker. An introduction to Berkovich analytic spaces and non-archimedean potential theory on curves. *p-adic Geometry (Lectures from the 2007 Arizona Winter School)*, 2008. 35
- [6] M. Baker, S. Payne, and J. Rabinoff. Nonarchimedean geometry, tropicalization, and metrics on curves. *Algebr. Geom*, 3(1):63–105, 2016. 35
- [7] B. Bertrand, E. Brugallé, and L. L. De Medrano. Planar tropical cubic curves of any genus, and higher dimensional generalisations. *L'Enseignement Mathématique*, 64(3):415–457, 2019. 2

- [8] B. Bertrand, E. Brugallé, and G. Mikhalkin. Tropical open Hurwitz numbers. *Rend. Semin. Mat. Univ. Padova*, 125:157–171, 2011. 29
- [9] B. H. Bowditch and D. B. A. Epstein. Natural triangulations associated to a surface. *Topology*, 27(1):91–117, 1988. 37
- [10] E. Brugallé, I. Itenberg, G. Mikhalkin, and K. Shaw. Brief introduction to tropical geometry. In *Proceedings of the Gökova Geometry-Topology Conference 2014*, pages 1–75. Gökova Geometry/Topology Conference (GGT), Gökova, 2015. 2, 4, 34
- [11] E. Brugallé and K. Shaw. A bit of tropical geometry. *Amer. Math. Monthly*, 121(7):563–589, 2014. 2
- [12] E. A. Brugallé and L. M. López de Medrano. Inflection points of real and tropical plane curves. *J. Singul.*, 4:74–103, 2012. 1, 2, 16, 20, 26, 29, 36
- [13] P. Buser. The collar theorem and examples. *Manuscripta Math.*, 25(4):349–357, 1978. 37
- [14] M. Coppens. A metric graph satisfying  $w_4^1 = 1$  that cannot be lifted to a curve satisfying  $\dim(w_4^1) = 1$ . *Open Mathematics*, 14(1):1–12, 2016. 2
- [15] M. A. Cueto and H. Markwig. How to repair tropicalizations of plane curves using modifications. *Experimental Mathematics*, 25(2):130–164, 2016. 2, 11, 35
- [16] M. A. Cueto and H. Markwig. Tropical geometry of genus two curves. *Journal of Algebra*, 517:457–512, 2019. 2, 11
- [17] M. A. Cueto and H. Markwig. Combinatorics and real lifts of bitangents to tropical quartic curves. *Discrete & Computational Geometry*, 69(3):597–658, 2023. 2
- [18] L. L. de Medrano, F. Rincón, and K. Shaw. Chern–schwartz–macpherson cycles of matroids. *Proceedings of the London Mathematical Society*, 120(1):1–27, 2020. 2
- [19] A. M. M. del Campo and S. Carpentier. Tropical embeddings of metric graphs. *arXiv preprint arXiv:1604.06176*, 2016. 2
- [20] H. Dörvodeli. Construction of tropical morphisms from tropical modifications of nonhyperelliptic genus 3 metric graphs with tree gonality 3 to metric trees. *arXiv preprint arXiv:2301.01989*, 2023. 2
- [21] N. Do. *Intersection theory on moduli spaces of curves via hyperbolic geometry*. PhD thesis, Ph. D. Thesis, University of Melbourne, 2008. 38



- [22] J. Draisma and A. Vargas. Catalan-many tropical morphisms to trees; part i: Constructions. *Journal of symbolic computation*, 104:580–629, 2021. 2
- [23] B. El Hilany. Constructing polynomial systems with many positive solutions using tropical geometry. *Revista Matemática Complutense*, 31:525–544, 2018. 39
- [24] Y. Ganor. *Enumerating Cuspidal Curves on Toric Surfaces*. Tel Aviv University, 2013. 2
- [25] Y. Ganor and E. Shustin. Enumeration of unicuspidal curves of any degree and genus on toric surfaces. *International Mathematics Research Notices*, 2022(21):16464–16523, 2022. 2, 8
- [26] W. Gubler, J. Rabinoff, and A. Werner. Skeletons and tropicalizations. *Adv. Math.*, 294:150–215, 2016. 35
- [27] W. Gubler, J. Rabinoff, and A. Werner. Tropical skeletons. In *Annales de l’Institut Fourier*, volume 67, pages 1905–1961, 2017. 35
- [28] M. H. Gunturkun and A. U. O. Kisisel. Using tropical degenerations for proving the nonexistence of certain nets. *arXiv preprint arXiv:1107.5530*, 2011. 30
- [29] M. A. Hahn, H. Markwig, Y. Ren, and I. Tyomkin. Tropicalized quartics and canonical embeddings for tropical curves of genus 3. *arXiv preprint arXiv:1802.02440*, 2018. 2
- [30] X. He. A generalization of lifting non-proper tropical intersections. *Journal of Pure and Applied Algebra*, 223(2):794–817, 2019. 2
- [31] I. Itenberg, G. Mikhalkin, and E. Shustin. *Tropical algebraic geometry*, volume 35 of *Oberwolfach Seminars*. Birkhäuser Verlag, Basel, second edition, 2009. 38
- [32] P. Jell and V. Wanner. Poincaré duality for the tropical dolbeault cohomology of non-archimedean mumford curves. *Journal of Number Theory*, 2017. 2
- [33] Y. Kageyama. Divisorial condition for the stable gonality of tropical curves. *arXiv preprint arXiv:1801.07405*, 2018. 2
- [34] N. Kalinin. Tropical geometry for Nagata’s conjecture and Legendrian curves. *Thesis, available online at <https://archive-ouverte.unige.ch/unige:80308>*, 2015. 2, 31

- [35] N. Kalinin. The Newton polygon of a planar singular curve and its subdivision. *Journal of Combinatorial Theory, Series A*, 137:226 – 256, 2016. 2, 11, 16, 31, 36
- [36] N. Kalinin. Tropical approach to Nagata’s conjecture in positive characteristic. *Discrete & Computational Geometry*, 58:158–179, 2017. 2, 31
- [37] H. J. Keisler. *Foundations of infinitesimal calculus*, volume 20. Prindle, Weber & Schmidt Boston, 1976. 38
- [38] M. Kerber and H. Markwig. Intersecting Psi-classes on tropical  $M_{0,n}$ . *Int. Math. Res. Not. IMRN*, (2):221–240, 2009. 38
- [39] A. Khovanskii. Logarithmic functional and the Weil reciprocity laws. In *Proceedings of the Waterloo Workshop on Computer Algebra*, pages 85–108. World Scientific, 2006. 17
- [40] A. Khovanskii. *Logarithmic functional and reciprocity laws*. Amer Math Soc, 2008. 17
- [41] A. G. Khovanskii. Newton polygons, curves on torus surfaces, and the converse Weil theorem. *Russian Mathematical Surveys*, 52(6):1251, 1997. 17
- [42] M. Kontsevich. Intersection theory on the moduli space of curves and the matrix airy function. *Communications in Mathematical Physics*, 147(1):1–23, 1992. 38
- [43] L. Lang. Geometry of tropical curves and application to real algebraic geometry, Thesis, available online at. <http://archive-ouverte.unige.ch/unige:43180>, 2014. 37
- [44] Y. Len and H. Markwig. Lifting tropical bitangents. *Journal of Symbolic Computation*, 96:122–152, 2020. 2
- [45] Y. Len and M. Satriano. Lifting tropical self intersections. *Journal of Combinatorial Theory, Series A*, 170:105138, 2020. 2, 9
- [46] H. Markwig, T. Markwig, and E. Shustin. Tropical curves with a singularity in a fixed point. *Manuscripta Math.*, 137(3-4):383–418, 2012. 11, 36
- [47] M. Mazin. Geometric theory of Parshin residues. *C. R. Math. Acad. Sci. Soc. R. Can.*, 32(3):81–96, 2010. 18
- [48] G. Mikhalkin. Enumerative tropical algebraic geometry in  $\mathbb{R}^2$ . *J. Amer. Math. Soc.*, 18(2):313–377, 2005. 33

- [49] G. Mikhalkin. Tropical geometry and its applications. In *International Congress of Mathematicians. Vol. II*, pages 827–852. Eur. Math. Soc., Zürich, 2006. 1, 4, 5, 18, 29
- [50] G. Mikhalkin. Moduli spaces of rational tropical curves. In *Proceedings of Gökova Geometry-Topology Conference 2006*, pages 39–51. Gökova Geometry/Topology Conference (GGT), Gökova, 2007. 33, 38
- [51] G. Mikhalkin. Quantum indices and refined enumeration of real plane curves. *Acta Math*, 219:135–180, 2017. 19
- [52] G. Mikhalkin and I. Zharkov. Tropical curves, their Jacobians and theta functions. In *Curves and abelian varieties*, volume 465 of *Contemp. Math.*, pages 203–230. Amer. Math. Soc., Providence, RI, 2008. 2, 18, 33, 38
- [53] R. Morrison. Tropical images of intersection points. *Collectanea Mathematica*, pages 1–11, 2014. 22, 24, 36
- [54] J. Mundinger. The image of a tropical linear space. *arXiv preprint arXiv:1808.02150*, 2018. 2
- [55] B. Osserman and S. Payne. Lifting tropical intersections. *Doc. Math.*, 18:121–175, 2013. 35
- [56] B. Osserman and J. Rabinoff. Lifting non-proper tropical intersections. *Tropical and Non-Archimedean Geometry*, 605:15–44, 2011. 35
- [57] S. Payne. Analytification is the limit of all tropicalizations. *Math. Res. Lett.*, 16(2-3):543–556, 2009. 35
- [58] J. Rabinoff. Tropical analytic geometry, Newton polygons, and tropical intersections. *Adv. Math.*, 229(6):3192–3255, 2012. 16, 35
- [59] Q. Ren, K. Shaw, and B. Sturmfels. Tropicalization of Del Pezzo Surfaces. *arXiv:1402.5651*, Feb. 2014. 36
- [60] A. Renaudineau. *Constructions de surfaces algébriques réelles*. PhD thesis, Paris 6, 2015. 2
- [61] J. Richter-Gebert, B. Sturmfels, and T. Theobald. First steps in tropical geometry. In *Idempotent mathematics and mathematical physics*, volume 377 of *Contemp. Math.*, pages 289–317. Amer. Math. Soc., Providence, RI, 2005. 16
- [62] M. Ruggiero and K. Shaw. Tropical Hopf manifolds and contracting germs. *Manuscripta Mathematica*, pages 1–60, 2015. 36
- [63] K. Shaw. Tropical surfaces. *arXiv:1506.07407*, 2015. 2, 36

- [64] K. M. Shaw. Tropical Intersection Theory and Surfaces, Thesis, available online at. <http://archive-ouverte.unige.ch/unige:22758>, 2011. 16, 36
- [65] K. M. Shaw. A tropical intersection product in matroidal fans. *SIAM Journal on Discrete Mathematics*, 27(1):459–491, 2013. 34
- [66] M. Sukenaga. Tropical lifting problem for the intersection of plane curves. *Beiträge zur Algebra und Geometrie/Contributions to Algebra and Geometry*, pages 1–35, 2023. 24, 27
- [67] T. Tao. *Compactness and contradiction*. American Mathematical Society, Providence, RI, 2013. 38
- [68] M. Temkin. Introduction to Berkovich analytic spaces. In *Berkovich spaces and applications*, volume 2119 of *Lecture Notes in Math.*, pages 3–66. Springer, Cham, 2015. 35
- [69] M. Ulirsch and D. Zakharov. Tropical double ramification loci. *arXiv preprint arXiv:1910.01499*, 2019. 2
- [70] O. Viro. On basic concepts of tropical geometry. *Proceedings of the Steklov Institute of Mathematics*, 273(1):252–282, 2011. 4, 36, 39
- [71] Y. Yamamoto. Periods of tropical K3 hypersurfaces. *arXiv preprint arXiv:1806.04239*, 2018. 2
- [72] S. Yoshitomi. Jacobian varieties of reduced tropical curves. *arXiv preprint math/0612810*, 2006. 18, 19