## ALMOST ALL PRIMES HAVE A MULTIPLE OF SMALL HAMMING WEIGHT

CHRISTIAN ELSHOLTZ

ABSTRACT. Recent results of Bourgain and Shparlinski imply that for almost all primes p there is a multiple mp that can be written in binary as

$$mp = 1 + 2^{m_1} + \dots + 2^{m_k}, \quad 1 \le m_1 < \dots < m_k,$$

with k = 66 or k = 16, respectively. We show that k = 6 (corresponding to Hamming weight 7) suffices.

We also prove there are infinitely many primes p with a multiplicative subgroup  $A = \langle g \rangle \subset \mathbb{F}_p^*$ , for some  $g \in \{2, 3, 5\}$ , of size  $|A| \gg p/(\log p)^3$ , where the sum-product set  $A \cdot A + A \cdot A$  does not cover  $\mathbb{F}_p$  completely.

## 1. INTRODUCTION

Recently Shparlinski [42] initiated the study of prime divisors of "sparse integers", i.e. of integers which only use a few non-zero digits in a g-ary representation. Let  $g \ge 2$  and  $k \ge 1$  be integers, and let  $\mathcal{D} = (d_i)_{i=0}^k$  be a sequence of k+1 nonzero integers. Let  $\mathcal{S}_{g,k}(\mathcal{D})$  be the set of integers n of the form

(1) 
$$n = d_0 + d_1 g^{m_1} + \dots + d_k g^{m_k},$$

where  $1 \leq m_1 < \cdots < m_k$ . Here the  $d_i$  can be thought of as (but are not necessarily) the *g*-ary digits  $d_i \in \{1, \ldots, g-1\}$  of *n*. Using exponential sums Shparlinski [42] proved: for any fixed  $\delta \in (0, 1/2)$  and every  $k > \max\{15, \frac{1}{\delta} - 1\}$ , for sufficiently large *X*, for almost all primes  $p \leq X$ , (i.e. for all but  $o(\pi(X))$  primes, where  $\pi(X)$  denotes the number of primes  $p \leq X$ ), there exists  $n \in S_{g,k}(\mathcal{D})$  with  $\log n \ll X^{1/2+\delta}$ , such that  $p \mid n$  holds.

Using new bounds of very short exponential sums Bourgain [6, 4] managed to prove the existence of much smaller n: Let  $\delta > 0$ , and  $k > k_0(\delta)$ . Let X be sufficiently large, then for all but  $o(\pi(X))$  primes  $p \leq X$  there exists  $n \in S_{g,k}(\mathcal{D})$ with log  $n \ll X^{\delta}$ , such that  $p \mid n$  holds.

Shparlinski [43] writes, working out  $k_0$  from Bourgain's approach would take "significant efforts", and with an alternative approach using incomplete exponential sums "on average over primes" he worked out an explicit admissible value: Let  $0 < \delta < 1/2$ , then one can take

$$k_0(\delta) = \frac{16}{\delta^2} + 1$$

One may wonder, how much this can be improved. It is clear that  $k \geq \frac{1}{\delta}$  must hold: The number of sparse integers of the type (1) is of order of magnitude

Institute of Analysis and Number Theory, Graz University of Technology Kopernikusgasse 24, A-8010 Graz, Austria. *elsholtz@math.tugraz.at.* 

 $C_{k,g}(\log n)^k \ll X^{\delta k}$ , and this expression must be at least of the same order of magnitude as the number of primes,  $\frac{X}{\log X}$ .

It seems there is no result in the literature that tries to minimize k, even when allowing a larger value n. It follows from the results above that k = 16, respectively k = 66 are admissable. Here we will show that actually k = 6 suffices. As it turns out the proof combines essentially two types of tools from the literature:

a) recent results on sumsets modulo a fixed prime p from additive combinatorics,

b) an old result of Erdős stating that, for most primes, the powers of 2 generate at least  $p^{1/2+o(1)}$  distinct residue classes, and variants, e.g. due to Pappalardi [37], Erdős and Ram Murty [17].

Using these tools the proof is actually quite short. However it seems the new type of view and the method of proof lead to considerable progress on the problem mentioned.

## 2. Discussion of the case g = 2.

The Hamming weight of an integer counts the number of ones in binary representation. During the last years there have been a number of related investigations on different number theoretic questions, such as primes p with all positive values of  $p - 2^i$  composite (see Tao [46]) or quadratic nonresidues with small Hamming weight, (see Dietmann, Elsholtz and Shparlinski [13]).

In this article we show that almost all primes have a nonzero multiple with a bounded Hamming weight of at most 7 (corresponding to k = 6 above).

Let  $T_k$  denote the set of those primes p which divide some integer of the form  $2^{a_1} + \cdots + 2^{a_k} + 1$ , where  $a_1, \ldots, a_k \in \mathbb{N}$ , and let  $T_k(X)$  denote the number of such primes  $p \leq X$ .

A result of Hasse [26] says that the Dirichlet density of primes  $p \leq x$  dividing a number of the form  $2^a + 1$ , where  $a \in \mathbb{N}$ , is  $\frac{17}{24}$ . This was later conjectured for natural density by Krishnamurty, and eventually proved by Odoni [36]. In other words,  $T_1(X) \sim \frac{17}{24}\pi(X) \sim \frac{17}{24}\frac{X}{\log X}$ .

There are extensions of this result to composite moduli, or to more general sequences such as  $a^k + b^k$  due to Ballot [3], Moree [34], Odoni [36] and Wiertelak [49].

Skałba [44] conjectured, (see also Moree [35]):

# Conjecture 1 (Skałba). $T_2(X) \sim \frac{X}{\log X}$ .

Skałba proved a partial result towards this conjecture. To formulate it we introduce some background first. Let  $\operatorname{ord}_p(a)$  denote the multiplicative order of a in  $\mathbb{F}_p^*$ . Erdős [16] conjectured that for each 0 < c < 1 for almost all primes  $\operatorname{ord}_p(2) > p^c$ holds, and proved this for  $c \leq \frac{1}{2} - o(1)$ . There have been several attempts to address Erdős's conjecture. Indeed, on the Generalized Riemann Hypothesis it is known that for almost all primes p:  $\operatorname{ord}_p(2) > \frac{p}{f(p)}$ , where f tends (arbitrarily slowly) to infinity, see Pappalardi [37]. Unconditionally, refining work of Pappalardi [37], Erdős and Murty [17] proved that for almost all primes  $p \leq x$  and for any function  $\varepsilon(x)$  tending to 0 as x tends to infinity,

$$\operatorname{ord}_p(2) \ge p^{\frac{1}{2} + \varepsilon(p)}$$

holds.

Skałba gave a conditional proof of his conjecture:

**Theorem** (Skałba [44]). If  $\operatorname{ord}_p(2) > p^{0.8}$ , then there exist integers a, b such that

$$2^a + 2^b + 1 \equiv 0 \bmod p$$

holds. In particular, if Erdős's conjecture holds with c = 0.8, then Conjecture 1 holds.

The proof made use of Weil bounds [48] on the number of solutions on congruences.

As progress on Erdős's conjecture has been very slow it seems fair to say that Erdős's conjecture with c = 0.8 is currently very far from a solution. The currently known exponent  $c = \frac{1}{2}$ , combined with Skałba's method based on Weil estimates seems insufficient to establish  $T_k(X) \sim \frac{X}{\log X}$ , for any fixed k.

As mentioned in the introduction, the results by Shparlinski give k = 66 and k = 16, respectively. We take a different approach to the problem. Combining methods from additive and multiplicative number theory we prove a result towards Skalba's conjecture, where  $2^a + 2^b + 1$  is replaced by  $2^{a_1} + \cdots + 2^{a_6} + 1$ . In other words, we prove that  $T_6(x) \sim \frac{x}{\log x}$ . This means that almost all primes p have a nonzero multiple mp with Hamming weight at most 7. Observe that the number of integers  $n \leq N$  with Hamming weight at most k is about  $\sum_{i=0}^{k} {t \choose i}$ , where  $t \sim \frac{\log N}{\log 2}$ . Hence there are about  $O((\log N)^6)$  odd integers  $n \leq N$  of Hamming weight at most r. For a given prime p the corresponding multiple mp of small Hamming weight can of course be much larger than the prime. As the proof shows, the exponents  $a_i$  are bounded above by p - 1, and hence  $mp \leq k2^{p-1}p$ . It seems possible to reduce the exponent by a small factor, following the techniques in Cilleruelo and Zumalacárregui [10] and Garaev and Kueh [19], but we do not make an attempt to do this.

One may wonder if this result, that almost all primes have a very sparse binary multiple, has any practical application. Of course, addition and multiplication with sparse integers should be faster than with arbitrary integers, but it seems one would need to find this sparse multiple in the first place and there could be several more issues to overcome.

We also prove similar results, where  $2^a + 2^b + 1$  is replaced e.g. by  $2^{a_1}3^{b_1}5^{c_1} + 2^{a_2}3^{b_2}5^{c_2} + 1$ . A more general formulation of these results is in the next section.

In the opposite direction Skałba proved:

**Theorem** (Skałba [44]). Let  $\Omega(w)$  denote the number of prime factors of w, with multiplicity. If  $\Omega(2^n - 1) < \frac{\log n}{\log 3}$ , then there exists a prime divisor p of  $2^n - 1$  such that for no pair of integers  $(a, b) : 2^a + 2^b + 1 \equiv 0 \mod p$ .

We will extend this to a similar estimate with  $2^a + 2^b$  replaced by a k-fold sum. One may conjecture that for arbitrarily large k there are infinitely many primes m such that no multiple of m can be written as  $\sum_{i=1}^{k} 2^{s_i}$  for nonnegative integers  $s_i$ . This would of course follow if there exist infinitely many Mersenne primes, but the condition  $\Omega(2^n - 1) < \frac{\log n}{\log k}$  in Theorem 5 is much more modest.

In the last section, we study a restriction to sum product estimates.

### 3. Results

3.1. Representation of residue classes, and approximations to Skałba's conjecture.

**Theorem 1.** Let  $r \ge 2$  and  $m \ne 0$  be fixed integers. For almost all primes  $p \le x$  there is a solution of

$$r^{a_1} + r^{a_2} + \dots + r^{a_6} \equiv m \mod p,$$

with integers  $0 \le a_1, \ldots, a_6 . More precisely, with <math>\varepsilon > 0$ , the number of primes  $p \le X$  with no solution of

$$r^{a_1} + r^{a_2} + \dots + r^{a_6} \equiv m \mod p$$

is  $O_r(X^{22/23+\varepsilon})$ .

With r = 2, m = -1 we get the following approximation to Skałba's conjecture.

**Corollary 1.** The following holds:  $T_6(X) \sim \frac{X}{\log X}$ .

Remark 1. In the special case that there exists some i with  $2^i \equiv -1 \mod p$  one can reduce the "6" in Theorem 1 above (with a different size of the exceptional set) to "5", in view of recent work of Shkredov (Theorem 2 of [41]).

**Theorem 2.** Let  $r \ge 2$ . For almost all primes  $p \le X$  there are at least  $p^{4/5}$  classes m which can be represented by

$$r^{a_1} + r^{a_2} \equiv m \bmod p,$$

with integers  $0 \le a_1, a_2 .$ 

Other approximations to Skałba's conjecture are as follows:

**Theorem 3.** Let  $r, s \ge 2$  be coprime integers and let  $m \ne 0$  be a fixed integer. For almost all primes  $p \le X$  there is a solution of

$$r^{a_1}s^{a_2} + r^{a_3}s^{a_4} + r^{a_5}s^{a_6} \equiv m \mod p,$$

with integers  $0 \le a_1, ..., a_6 .$ 

**Theorem 4.** Let  $r, s, t \ge 2$  be mutually coprime integers and let  $m \ne 0$  be a fixed integer. For almost all primes  $p \le X$  there is a solution of

$$r^{a_1}s^{a_2}t^{a_3} + r^{a_4}s^{a_5}t^{a_6} \equiv m \mod p$$

with integers  $0 \le a_1, ..., a_6 .$ 

For completeness we also state the following, which is a direct consequence of a result of Schoen and Shkredov [39].

**Corollary 2.** Let  $r \ge 2$ . If  $\operatorname{ord}_p(r) > p^{3/4}$ , then for some integers  $a_1, a_2$  $r^{a_1} + r^{a_2} + 1 \equiv 0 \mod p$ .

In particular, if Erdős's conjecture holds with c = 0.75, then Conjecture 1 holds. This last result replaces  $\frac{4}{5}$  in Skałba's Theorem (mentioned above) by  $\frac{3}{4}$ .

Remark 2. Another approach to this problem could be the following one: a) Baker and Harman [2] proved that for a positive proportion of primes, the largest prime factor P of p-1 is large:  $P(p-1) \gg p^{0.677}$ . This result is ineffective due to the use of Siegel zeros.

b) Harman [23] also gave an effective result (i.e. not depending on a Siegel zero), namely that for some positive (computable) density of the primes this holds with exponent 0.6105.

From case a) one can deduce that for a positive (ineffective) proportion of primes  $\operatorname{ord}_p(2) \gg p^{0.677}$  holds. In view of the large prime factor we first consider the case in which 2 generates only  $O(p^{1/3})$  residue classes: here we know from the Erdős conjecture with  $c = \frac{1}{2}$ , see also Lemma 1, that this case rarely happens. In the other case 2 generates a subgroup of size at least  $cp^{0.677}$  residue classes. See also [31, Lemma 20].

This would imply (e.g. by Lemma 5) that for these primes  $2^a + 2^b + 2^c + 1 \equiv 0 \mod p$  has a solution. In its current form this consequence would be much weaker than Hasse's result, namely that for a density of 17/24 of the primes even  $2^a + 1 \equiv 0 \mod p$  has a solution. Therefore we do not pursue this path further.

3.2. Primes without multiples of small Hamming weight. As an extension of Skałba's second theorem with regard to multiple sums, and thus a limitation or partial converse to the type of results above, we prove the following:

## Theorem 5. If

(2) 
$$\Omega(2^n - 1) < \frac{\log n}{\log k},$$

then there is some prime factor  $q \mid 2^n - 1$  such that

$$2^{a_1} + \dots + 2^{a_{k-1}} + 1 \not\equiv 0 \mod q$$

for all integer choices of  $a_i$ . Hence all multiples of q have Hamming weight at least k+1.

3.3. A restriction on sum product estimates. Suppose a k-fold sum  $r^{a_1} + \cdots + r^{a_k}$  covers all residue classes, in particular the class 0; here without loss of generality  $a := a_k \leq \cdots \leq a_1$ . Dividing by  $r^a$  one obtains that  $r^{a_1-a} + \cdots + r^{a_{k-1}-a} + 1 \equiv 0 \mod p$ .

Also, the related question has been studied, about the minimal number d such that the mixed d-fold sum-product set  $dAB := \{a_1b_1 + \cdots + a_db_d : a_i \in A, b_i \in B\}$  covers all residue classes.

For random sets  $A \subset \mathbb{Z}/p\mathbb{Z}$  with  $|A| \geq C_{\varepsilon}p^{\frac{1}{2}+\varepsilon}$ , one may expect that  $A + A = \mathbb{Z}/p\mathbb{Z}$ . See for example some discussion (with too optimistic conjectures) in Hart and Iosevich [25], Rudnev [38] and Chapman, Erdoğan, Hart, Iosevich and Koh [9]. The last authors observed that some kind of restriction must occur, they write about subsets  $A \subset \mathbb{F}_q$ : "Due to the misbehavior of the zero element it is not possible for  $A \cdot A + A \cdot A = \mathbb{F}_q$  unless A is a positive proportion of the elements of  $\mathbb{F}_q$ ."

It may be worth recalling that for a prime  $p \equiv 3 \mod 4$  the set A of (nonzero) quadratic residues has positive density but  $0 \notin AA + AA$ . The set of squares are of course an explicit example of a large multiplicative subgroup of  $\mathbb{F}_p^*$  with restricted sumsets in  $\mathbb{F}_p$ . On the other hand this example of size  $|A| = \frac{p-1}{2}$  is extremal, as for any set A with  $|A| \geq \frac{p+1}{2}$  the Cauchy-Davenport theorem guarantees that  $A + A = \mathbb{F}_p$ .

The set of squares is generated by the square  $g^2$  of a primitive root g. But, as  $g^2$  is not a fixed element, following the view point of this paper one can ask if there exists such examples generated by a fixed element, e.g. g = 2 and  $A = \{2^i \mod p : 1 \le i \le \operatorname{ord}_p(2)\}$ . In short, the answer is: "probably yes, but unconditionally we can only prove a slightly weaker result".

#### CHRISTIAN ELSHOLTZ

Let us first describe an explicit and "large" example, where however we cannot prove this type of example will occur infinitely often. We then describe a slightly weaker example, which occurs quite frequently.

Let *a* be fixed, and *p* be a prime with  $\operatorname{ord}_p(a) = \frac{p-1}{2}$ , where  $\frac{p-1}{2}$  is odd. Let  $A = B = \{a^i : 1 \le i \le \frac{p-1}{2}\}$ . Then  $|A||B| \sim \frac{p^2}{4}$  and  $2AB = A^2 + A^2 = A + A$ . But  $0 \notin A + A$ , as otherwise  $a^i \equiv -a^j \mod p$ . But -1 is not in the multiplicative subgroup generated by *a*, as *a* has odd order. The size of  $|A| = \frac{p-1}{2}$  is of course much larger than  $p^{\frac{1}{2}+\varepsilon}$ . For a = 2 such primes are  $p = 7, 23, 47, 71, 79, 103, 167, 191, 199, \ldots$  Alternatively, one can take primes  $p \equiv 3 \mod 4$ , with 2 as primitive root, and  $A = \{4^i : 1 \le i \le \frac{p-1}{2}\}$ . Such primes are  $3, 11, 19, 59, 67, 83, 107, 131, 139, 163, 179, \ldots$  Artin's conjecture on primitive roots (known only on GRH, see Hooley [29]) predicts that such sets of primes have a positive density in the set of primes.

Unconditionally, we are able to prove the existence of an infinite number of primes with a restricted sumset for the following slightly weaker variant:

**Theorem 6.** For a given prime p let s = s(p) and w = w(p) be defined by  $p = 2^s w + 1$ , where w is odd. For almost all positive square-free numbers a > 1, with at most three exceptions, (or almost all primes, with at most 2 exceptions), there exist positive constants c and  $c_1$ , and there exist at least  $c_1 \frac{x}{(\log x)^2}$  many primes  $p \le x$  such that the multiplicative subgroup

$$A = \{ \left( a^{2^s} \right)^i : 1 \le i \le \operatorname{ord}_p(a^{2^s}) \} \subset \mathbb{F}_p^*$$

has the following properties:

- (1)  $|A| > cp/(\log p)^3$  and
- (2)  $A \cdot A + A \cdot A = A + A \neq \mathbb{F}_p$ .

Remark: in this situation  $A = A \cdots A$ , but as  $|A \cdot A|$  is typically much larger than |A| we state it in the form above.

Let us mention a related result by Alon and Bourgain [1]: there is an absolute constant c > 0 so that there are infinitely many primes p and a multiplicative subgroup  $A \subset \mathbb{F}_p^*$  with  $|A| \ge cp^{1/3}$  such that: there are no  $x, y, z \in A$  with x+y=z.

## 4. Proofs

We observe that Erdős's conjecture only makes a statement about the number of residue classes of the form  $r^i \mod p$ , (for fixed r and p), but not about its algebraic structure. Observing that these classes are a multiplicative subgroup of  $\mathbb{F}_p^*$  we now study sumsets of multiplicative subgroups in  $\mathbb{F}_p^*$ . In recent years there has been a considerable number of papers concerned with sumsets modulo primes. Of the many results available we choose those results that seem best suited for our application.

4.1. **Lemmas.** For the proof we essentially use an ingredient (the first two lemmas below) coming from multiplicative number theory and several very recent quantitative versions of results of additive combinatorics.

**Lemma 1** (Erdős-Murty [17, Theorem 5], Pappalardi [37]). Let  $\Gamma \subseteq \mathbb{Q}^*$  be a multiplicative subgroup of rank d. Suppose that  $\Gamma$  is generated by the mutually coprime numbers  $b_1, \ldots, b_d$ . For all primes p not dividing the denominators of

 $b_1, \ldots, b_d$  we define  $f_{\Gamma}(p)$  to be the order of  $\Gamma \mod p$ . Let  $\varepsilon(x)$  be any function tending to zero as  $x \to \infty$ . For all but  $o(\frac{x}{\log x})$  primes  $p \leq x$ :

$$f_{\Gamma}(p) \ge p^{\frac{d}{d+1} + \varepsilon(p)}$$

The number of exceptional primes can be bounded from above in a more precise way; see for example Indlekofer and Timofeev [30] and Ford [18]. However for our situation, a method by Matthews, worked out in detail by Pappalardi, is suitable.

Here  $b_1, \ldots, b_d$ , are multiplicatively independent integers, not being squares, and not being  $\pm 1$ .

**Lemma 2** (Pappalardi [37, Lemma 1.2], Matthews [33]). Suppose that d is a function of t such that  $dt^{-d}$  is bounded. Then

$$\#\{p: f_{\Gamma}(p) \le t\} \ll \frac{t^{1+1/d}}{\log t} 2^{d} d \sum_{i=1}^{d} \log b_{i}$$

uniformly with respect to t, d and  $\{b_1, \ldots, b_d\}$ .

Our application will only need the special case d = 1.

A slightly weaker version of Lemma 1 goes back to Erdős [16], and one can give the following very simple proof:

The sequence  $(2^i \mod p), i = 1, \ldots, n$  is periodic. The order  $\operatorname{ord}_p(2)$  is by definition the length of the period, and one has  $2^{\operatorname{ord}_p(2)} \equiv 1 \mod p$ . This implies that

$$\prod_{p < y} p^{\lfloor \frac{n}{\operatorname{ord}_p(2)} \rfloor} \le \prod_{i=1}^n (2^i - 1) \le 2^{n(n+1)/2},$$

where the product on the left hand side runs over the primes, up to some level y specified below. Taking logarithms one sees that

(3) 
$$\sum_{p < y} \log p \lfloor \frac{n}{\operatorname{ord}_p(2)} \rfloor \le Cn^2$$

for some positive constant C. Assuming that  $\operatorname{ord}_p(2) \leq y^{\frac{1}{2}-\varepsilon}$  for at least  $\delta \frac{y}{\log y}$  of the primes  $p \leq y$ , (for some  $\delta > 0$ ), then

$$\sum_{p < y} \log p\left(\frac{n}{\operatorname{ord}_p(2)} - 1\right) \ge \frac{\delta y}{\log y} \left(\frac{n}{y^{\frac{1}{2} - \varepsilon}} - 1\right) \ge \delta n \frac{y^{\frac{1}{2} + \varepsilon}}{\log y} - \frac{y}{\log y}$$

For  $y = n^2$  this contradicts equation (3), proving that for most primes  $p < n^2$  the powers  $1, 2, 4, \ldots, 2^n$  occupy more than  $y^{\frac{1}{2}-\varepsilon} > p^{\frac{1}{2}-\varepsilon}$  distinct residue classes modulo p. For more general discussions on this see Elsholtz [14].

We now come to the additive ingredients, which are more modern, and much deeper.

**Lemma 3** (Hart [24]). Let  $R \subseteq \mathbb{F}_p^*$  be a multiplicative subgroup such that  $|R| \ge p^{\kappa}$ , where  $\kappa > \frac{11}{23}$ . Then for all sufficiently large p we have  $6R := R + R + R + R + R + R \ge \mathbb{F}_p^*$ .

An earlier version with exponent 41/83 appeared in Schoen and Shkredov [39, Theorem 4.1], and 55/112 in Shkredov [40, Corollary 32].

See also [11, 12, 15] for a number of related results on Waring's problem modulo p.

**Lemma 4** (Hart [24], Theorem 10). Let R be a multiplicative subgroup of  $\mathbb{F}_p^*$  with  $|R| \ll p^{5/9-\varepsilon}$ , then  $|R+R| \gg |R|^{8/5} (\log |R|)^{-3/10}$ .

**Lemma 5** (Schoen and Shkredov [39, Theorem 2.6, l = 3]). Let R be a multiplicative subgroup of  $\mathbb{F}_p^*$ , with  $|R| > p^{2/3}$ . Then  $3R := R + R + R \supseteq \mathbb{F}_p^*$ .

**Lemma 6** (Schoen and Shkredov [39, Theorem 2.6, l = 2]). Let R be a multiplicative subgroup of  $\mathbb{F}_p^*$ , with  $|R| > p^{3/4}$ . Then  $2R := R + R \supseteq \mathbb{F}_p^*$ .

For the case of a twofold sum there was important earlier work by Heath-Brown and Konyagin [28]: Let R be a multiplicative subgroup of  $\mathbb{F}_p^*$ , with  $|R| \gg p^{2/3}$ . Then  $|R + R| \gg p$ . Also, several related results of this type of theorem are due to Bourgain [5], Glibichuk [20], [21], Cochrane and Pinner [11], Hart and Iosevich [25].

4.2. **Proofs of Theorems.** After this preparation the Theorems are straightforward consequences:

The sets  $R = \{r^i : 1 \leq i \leq \operatorname{ord}_p(r)\}, R = \{r^i s^j : 1 \leq i \leq \operatorname{ord}_p(r), 1 \leq j \leq \operatorname{ord}_p(s)\}$ etc. are multiplicative subgroups of  $\mathbb{F}_p^*$  so that the corresponding lemmas apply as follows:

Theorem 1 follows from Lemma 1, with d = 1,  $b_1 = r$  and Lemma 3. (Observe that  $\frac{11}{23} < \frac{1}{2}$ .) Further observe that it suffices to estimate how often  $\operatorname{ord}_p(r) \leq p^{11/23+\varepsilon}$  holds. It follows from Lemma 2 with d = 1,  $b_1 = r$ ,  $t = x^{11/23+\varepsilon/2}$  that

$$\#\{p: \operatorname{ord}_p(r) \le p^{11/23 + \varepsilon/2}\} \le \#\{p: \operatorname{ord}_p(r) \le x^{11/23 + \varepsilon/2}\} \ll_r \frac{x^{22/23 + \varepsilon}}{\log x}$$

Theorem 2 follows from Lemma 1 with d = 1,  $b_1 = r$  and Lemma 4 with  $|R| \ge p^{1/2+\varepsilon}$ , and observing that  $\frac{p^{8\varepsilon/5}}{(\log p)^{3/10}} \gg p^{\varepsilon}$ .

Theorem 3 follows from Lemma 1, with d = 2, and Lemma 5. Theorem 4 follows from Lemma 1, with d = 3, and Lemma 6.

Corollary 2 follows from Lemma 6.

4.3. **Proof of Theorem 5.** Every multiple of an integer of the form  $M = 2^n - 1$  has Hamming weight at least n, see Stolarsky [45, Theorem 2.1] or Wagstaff [47]. Moreover, if

(4) 
$$R := \Omega(2^n - 1) < \frac{\log n}{\log k},$$

then some prime factor  $q \mid 2^n - 1$  also has this property. To see this we adapt Skałba's argument. Let  $M = 2^n - 1 = \prod_{i=1}^r q_i^{t_i}$ . Suppose all  $q_i$  have some multiple  $w_i q_i$  which is a sum of at most k powers of two, that is

$$w_i q_i = \varepsilon_{1,i} 2^{s_1(q_i)} + \dots + \varepsilon_{k,i} 2^{s_k(q_i)},$$

with  $\varepsilon_{j,i} \in \{0,1\}$ , (not all of them being 0), then

$$P := \prod_{i=1}^{r} (w_i q_i)^{t_i} = \prod_{i=1}^{r} \left( \varepsilon_{1,i} 2^{s_1(q_i)} + \dots + \varepsilon_{k,i} 2^{s_k(q_i)} \right)^{t_i}$$

is on the one hand side a multiple of M, and thus has Hamming weight at least n. On the other side, P is a sum of at most  $k \sum_{i=1}^{r} t_i = k^{\Omega(2^n-1)} = k^R < n$  powers of 2, a contradiction. Hence there is some prime divisor  $q \mid 2^n - 1$  for which

$$\varepsilon_i 2^{s_1(q)} + \dots + \varepsilon_k 2^{s_k(q)} \not\equiv 0 \mod q,$$

and hence all nonzero multiples of q have Hamming weight at least k + 1.

#### 4.4. Proof of theorem 6.

*Proof.* In the proof below the  $c_i$  are suitably chosen positive constants.

By Heath-Brown's work [27, Corollary 3] on Artin's primitive root conjecture, for any fixed bound x and any four distinct positive square-free numbers  $a_1, a_2, a_3, a_4$ (not being 1) (or for any three distinct primes) one of these, say a, is a primitive root (i.e.  $\operatorname{ord}_p(a) = p-1$ ) for at least  $c_1 \frac{x}{(\log x)^2}$  such primes  $p \leq x$ . Hence for suitable x there are at least  $c_2 \frac{x}{(\log x)^2}$  such primes  $\frac{x}{2} . Write <math>p-1=2^{s}w$ , where w is odd, and  $y = \frac{c_2}{4} \frac{p}{(\log p)^3}$ . As there are at most  $y \frac{\log x}{\log 2} < \frac{c_2}{2} \frac{x}{(\log x)^2}$  such primes with  $w \leq y$ , there are at least  $\frac{c_2}{2} \frac{x}{(\log x)^2}$  many primes p with  $\operatorname{ord}_p(a^{2^s}) = w \gg \frac{p}{(\log p)^3}$ . Observe that all elements of the form  $a^{2^s i}$  have odd order and that therefore -1is not of this type. As  $a^{2^s i} + 1 \neq 0 \mod p$  it follows: the set  $A = \{a^{2^s i} : 1 \leq i \leq \operatorname{ord}_p(a^{2^s})\}$  is of size  $|A| \gg \frac{p}{(\log p)^3}$  but  $0 \notin A + A = A^2 + A^2$ .

## 5. Open problems

Finally, we state as open questions:

- (1) For primes  $p \notin T_2$ : does  $\operatorname{ord}_p(2) \leq c_{\varepsilon} p^{1/2+\varepsilon}$  hold, for all  $\varepsilon > 0$ , and some constant  $c_{\varepsilon}$ ?
- (2) For  $h \to \infty$ , what can one say about primes  $p \notin T_h$ , but not being Mersenne numbers?
- (3) How can one algorithmically find a "sparse" representation of a multiple of p? Can this be used for other complexity questions?

Finally, we note that there are 231 primes  $p \leq 4 \cdot 10^6$  which are not in  $T_2$ , whereas there are 283 146 primes below 4 million. On the other hand, as conjectured by Skałba [44], proving that there are infinitely many such primes remains an open problem.

#### 6. Acknowledgements

I would like to thank Rainer Dietmann, Igor Shparlinski and the referees for useful comments.

#### References

- Alon, N.; Bourgain, J: Additive patterns in multiplicative subgroups. Geom. Funct. Anal. 24 (2014), no. 3, 721–739.
- Baker, R. C.; Harman, G.: Shifted primes without large prime factors. Acta Arith. 83 (1998), no. 4, 331–361.
- [3] Ballot, C.: Density of prime divisors of linear recurrences. Mem. Amer. Math. Soc. 115 (1995), no. 551.
- Bourgain, J.: New bounds on exponential sums related to the Diffie-Hellman distributions. C. R. Math. Acad. Sci. Paris 338 (2004), no. 11, 825–830.
- [5] Bourgain, J.: Mordell's exponential sum estimate revisited, J. Amer. Math. Soc. 18 (2005), 477–499.
- [6] Bourgain, J.: Estimates on exponential sums related to the Diffie-Hellman distributions. Geom. Funct. Anal. 15 (2005), no. 1, 1–34.
- [7] Bourgain, J.; Katz, N.; Tao, T.: A sum-product estimate in finite fields, and applications, Geometric And Functional Analysis 14, 1, (2004), 27–57,

#### CHRISTIAN ELSHOLTZ

- [8] Bundschuh, P.: Solution of problem 618. Elemente der Mathematik 26 (1971), 43–44.
- Chapman, J., Erdoğan, M., Hart, D., Iosevich, A., Koh, D.: Pinned distance sets, k-simplices, Wolff's exponent in finite fields and sum-product estimates Mathematische Zeitschrift 271(1), (2012) 63–93.
- [10] Cilleruelo, J.; Zumalacárregui, A.: An additive problem in finite fields of powers of elements of large multiplicative order, Revista Matemática Complutense 27, no.2 (2014) 501–508.
- [11] Cochrane, T.; Pinner, C.: Sum-product estimates applied to Waring's problem mod p, Integers 8 (2008), A46, 18pp.
- [12] Cochrane, T., Hart, D., Pinner, C.; Spencer, C.: Waring's number for large subgroups of  $\mathbb{Z}_p^*$ , Acta Arith. 163 no. 4, (2014) 309–325.
- [13] Dietmann, R.; Elsholtz; C.; Shparlinski, I.: On Gaps Between Primitive Roots in the Hamming Metric, Quarterly J. Math. 64 (no 4) (2013), 1043–1055.
- [14] Elsholtz, C.: The distribution of sequences in residue classes. Proc. Amer. Math. Soc. 130 (2002), no. 8, 2247–2250.
- [15] Elsholtz, C.: The number  $\Gamma(k)$  in Waring's problem, Acta Arith. 131 (2008), 43–49.
- [16] Erdős, P.: Bemerkungen zu einer Aufgabe in den Elementen Arch. Math. (Basel) 27 (1976) no. 2, 159–163.
- [17] Erdős, P.; Murty, M. Ram: On the order of a (mod p). Number theory (Ottawa, 1996), 87–97, CRM Proc. Lecture Notes, 19.
- [18] Ford, K.: The distribution of integers with a divisor in a given interval, Annals of Math. (2) 168 (2008), 367–433.
- [19] Garaev, M.Z., Kueh, K.L.: Distribution of special sequences modulo a large prime, Int. J. of Math. and Math. Sci. 50 (2003), 3189–3194.
- [20] Glibichuk, A.A.: Additive properties of product sets in an arbitrary finite field. Arxiv: 0801.2021, preprint 2008
- [21] Glibichuk, A.A.: Sums of powers of subsets of an arbitrary finite field. (Russian) Izv. Ross. Akad. Nauk Ser. Mat. 75 (2011), no. 2, 35–68; translation in Izv. Math. 75 (2011), no. 2, 253–285.
- [22] Glibichuk, A.A.; Rudnev, M.: On additive properties of product sets in an arbitrary finite field. J. Anal. Math. 108 (2009), 159–170.
- [23] Harman, G.: On the greatest prime factor of p-1 with effective constants. Math. Comp. 74 (2005), no. 252, 2035–2041.
- [24] Hart, D.: A note on sumsets of subgroups in  $\mathbb{Z}_p^*$ , Acta Arith. 161 (2013), 387–395.
- [25] Hart, D.; Iosevich, A.: Sums and products in finite fields: An integral geometric viewpoint, Contemp. Math. 464, Amer. Math. Soc., Providence, RI, 2008, pp. 129–135.
- [26] Hasse, H.: Über die Dichte der Primzahlen p, für die eine vorgegebene ganzrationale Zahl  $a \neq 0$  von gerader bzw. ungerader Ordnung mod p ist. Math. Ann. 166 (1966) 19–23.
- [27] Heath-Brown, D. R.: Artin's conjecture for primitive roots. Quart. J. Math. Oxford Ser. 37 (1) (1986), 27–38.
- [28] Heath-Brown, D.R., Konyagin, S.: New bounds for Gauss sums derived from kth powers. and for Heilbronn's exponential sum Q. J. Mathematics, 51 (2) (2000), 221–235.
- [29] Hooley, C.: On Artin's conjecture. J. Reine Angew. Math. 225 (1967), 209–220.
- [30] Indlekofer, K.-H., Timofeev, N.M.: Divisors of Shifted Primes, Publ. Math. Debrecen 60 (2002), 307-345.
- [31] Kurlberg, P., Pomerance, C.: On the periods of the linear congruential and power generators. Acta Arith. 119 (2005), no. 2, 149–169.
- [32] Luca, F., Stanica, P.: Prime divisors of Lucas sequences and a conjecture of Skalba, International Journal of Number Theory 1 (2005), no. 1, 583-591.
- [33] Matthews, C.R.: Counting points modulo p for some finitely generated subgroups of algebraic groups. Bull. London Math. Soc. 14 (1982), 149–154.
- [34] Moree, P.: On the divisors of  $a^k + b^k$ . Acta Arith. 80 (1997), no. 3, 197–212.
- [35] Moree, P.: Artin's primitive root conjecture -a survey -, Integers Volume 12A (2012), #A13.
- [36] Odoni, R. W. K.: A conjecture of Krishnamurthy on decimal periods and some allied problems. J. Number Theory 13 (1981), no. 3, 303–319.
- [37] Pappalardi, F.: On the Order of Finitely Generated Subgroups of  $Q^* \pmod{p}$  and Divisors of p 1, J. Number Theory 57 (2), 1996, 207–222.
- [38] Rudnev, M.: An improved estimate on sums of product sets, arXiv:0805.2696v1, 2008.

11

- [39] Schoen, T.; and Shkredov, I.D.: Additive properties of multiplicative subgroups of  $\mathbb{F}_p$ , Q. J. Oxford, 63 (3) 2012, 713–722.
- [40] Shkredov, I.D.: Some new inequalities in additive combinatorics, Mosc. J. Comb. Number Theory 3:3–4 (2013), 189–239.
- [41] Shkredov, I.D.: On exponential sums over multiplicative subgroups of medium size, Finite Fields Appl. 30 (2014), 72–87.
- [42] Shparlinski, I.E.: Prime divisors of sparse integers. Period. Math. Hungar. 46 (2003), no. 2, 215–222.
- [43] Shparlinski, I.E.: Exponential sums and prime divisors of sparse integers. Period. Math. Hungar. 57 (2008), no. 1, 93–99.
- [44] Skałba, M.: Two conjectures on primes dividing  $2^a + 2^b + 1$ . Elem. Math. 59 (2004), no. 4, 171–173.
- [45] Stolarsky, K.B.: Integers whose multiples have anomalous digital frequencies. Acta Arith. 38 (1980),  $117{-}128$
- [46] Tao, T.: A remark on primality testing and decimal expansions. J. Austral. Math. Soc. 91(03) (2011), 405–413.
- [47] Wagstaff, S., Jr.: Prime Numbers with a Fixed Number of One Bits or Zero Bits in Their Binary Representation, Experiment. Math. Volume 10, Issue 2 (2001), 267–274.
- [48] Weil, A.: Numbers of solutions of equations in finite fields. Bull. Amer. Math. Soc. 55 (1949), 497–508.
- [49] Wiertelak, K.: On the density of some sets of primes. IV. Acta Arith. 43 (1984), no. 2, 177–190.

11A41 Primes

- 11B13 Additive bases, including sumsets
- 11B50 Sequences (mod m)
- 11N25 Distribution of integers with specified multiplicative constraints
- 11P05 Waring's problem and variants