Feedback Vertex Set Inspired Kernel for Chordal Vertex Deletion*

Akanksha Agrawal[†] Daniel Lokshtanov[‡] Pranabendu Misra[§] Saket Saurabh[¶]

Meirav Zehavi[∥]

July 18, 2017

Abstract

Given a graph G and a parameter k, the CHORDAL VERTEX DELETION (CVD) problem asks whether there exists a subset $U \subseteq V(G)$ of size at most k that hits all induced cycles of size at least 4. The existence of a polynomial kernel for CVD was a well-known open problem in the field of Parameterized Complexity. Recently, Jansen and Pilipczuk resolved this question affirmatively by designing a polynomial kernel for CVD of size $\mathcal{O}(k^{161} \log^{58} k)$, and asked whether one can design a kernel of size $\mathcal{O}(k^{10})$. While we do not completely resolve this question, we design a significantly smaller kernel of size $\mathcal{O}(k^{12} \log^{10} k)$, inspired by the $\mathcal{O}(k^2)$ -size kernel for FEEDBACK VERTEX SET. Furthermore, we introduce the notion of the independence degree of a vertex, which is our main conceptual contribution.

 $^{^*}$ A preliminary version of this paper appeared in the proceedings of 28th ACM-SIAM Symposium on Discrete Algorithms (SODA 2017). The research leading to these results received funding from the European Research Council under the European Unions Seventh Framework Programme (FP/2007-2013) / ERC Grant Agreement no. 306992.

[†]University of Bergen, Bergen, Norway. akanksha.agrawal@uib.no.

[‡]University of Bergen, Bergen, Norway. daniello@uib.no.

[§]The Institute of Mathematical Sciences, HBNI, Chennai, India. pranabendu@imsc.res.in.

[¶]University of Bergen, Bergen, Norway. The Institute of Mathematical Sciences, HBNI, Chennai, India. saket@imsc.res.in.

^{||}University of Bergen, Bergen, Norway. meirav.zehavi@uib.no.

1 Introduction

Data reduction techniques are widely applied to deal with computationally hard problems in real world applications. It has been a long-standing challenge to formally express the efficiency and accuracy of these "pre-processing" procedures. The framework of parameterized complexity turns out to be particularly suitable for a mathematical analysis of pre-processing heuristics. Formally, in parameterized complexity each problem instance is accompanied by a parameter k, and we say that a parameterized problem is *fixed-parameter tractable (FPT)* if there is an algorithm that solves the problem in time $f(k) \cdot |I|^{O(1)}$, where |I| is the size of the input and f is a computable function of the parameter k alone. *Kernelization* is the subarea of parameterized complexity that deals with the mathematical analysis of pre-processing heuristics. A parameterized problem is said to admit a *polynomial kernel* if there is a polynomial-time algorithm, that reduces the input instance down to an instance whose size is bounded by a polynomial p(k) in k, while preserving the answer. This reduced instance is called a p(k)-kernel for the problem. Observe that if a problem has a kernelization algorithm, then it is also has an FPT algorithm.

Kernelization appears to be an interesting computational approach not only from a theoretical perspective, but also from a practical perspective. There are many real-world applications where even very simple preprocessing can be surprisingly effective, leading to significant size-reduction of the input. Kernelization is a natural tool for measuring the quality of existing preprocessing rules proposed for specific problems as well as for designing new powerful such rules. The most fundamental question in the field of kernelization is:

Let Π be parameterized problem that admits an FPT algorithm. Then, does Π admit a polynomial kernel?

In recent times, the study of kernelization, centred on the above question, has been one of the main areas of research in parameterized complexity, yielding many new important contributions to theory. These include general results showing that certain classes of parameterized problems have polynomial kernels, and as well as other results that utilize advanced techniques from algebra, matroid theory and topology for data reduction [4, 6, 18, 17, 22, 25, 28, 29, 32, 31, 39, 40]. The development of a framework for ruling out polynomial kernels under certain complexity-theoretic assumptions [5, 7, 14, 20] has added a new dimension to the area, and strengthened its connections to classical complexity theory. We refer to the following surveys [30, 34] and the corresponding chapters in the books [12, 15, 16, 38], for a detailed introduction to the field of kernelization.

An important class of problems, that has led to the development of many upper bound tools and techniques in kernelization, is the class of *parameterized graph deletion problems*. A typical problem of this class is associated with a family of graphs, \mathcal{F} , such as edgeless graphs, forests, cluster graphs, chordal graphs, interval graphs, bipartite graphs, split graphs or planar graphs. The deletion problem corresponding to \mathcal{F} is formally stated as follows.

\mathcal{F} -Vertex (Edge) Deletion	Parameter: k
Input: An undirected graph G and a non-negative integer k .	
Question: Does there exist $S \subseteq V(G)$ (or $S \subseteq E(G)$) such that $ S \leq k$ a	and $G \setminus S$ is in \mathcal{F} ?

Graph deletion problems are also among the most basic problems in graph theory and graph algorithms. Most of these problems are NP-complete [41, 33], and thus they were subject to intense study in various algorithmic paradigms to cope with their intractability [21, 35, 17, 37]. These include, considering a restricted class of inputs, approximation algorithms, parameterized complexity and kernelization.

Some of the most well known results in kernelization are polynomial kernels for graph deletion problems such as FEEDBACK VERTEX SET [40], ODD CYCLE TRANSVERSAL [32, 31], VERTEX COVER [2, 11], PLANAR- \mathcal{F} -DELETION [17], and TREEDEPTH- η -DELETION [22]. A common thread among all these problems, with the exception of ODD CYCLE TRANSVERSAL, is that the corresponding family \mathcal{F} can be characterized by a finite set of forbidden minors that include at least one connected planar graph. It is known that, if \mathcal{F} can be characterized by a finite set of forbidden induced subgraphs, then the corresponding \mathcal{F} -VERTEX DELETION problem immediately admits an FPT algorithm as well as polynomial sized kernel because of its connection to d-HITTING SET [1]. However, if \mathcal{F} is characterized by an *infinite* set of forbidden induced subgraphs, which is the case when \mathcal{F} is the class of chordal graphs, chordal bipartite graphs, interval graphs, proper interval graphs and permutation graphs, our understanding of these problems in the realm of parameterized complexity and kernelization, is still at a nascent stage. While CHORDAL VERTEX DELETION (CVD) was known to be FPT for some time [36, 10], the parameterized complexity of INTERVAL VERTEX DELETION was settled only recently [9, 8]. The parameterized complexity of PERMUTATION VERTEX DELETION and CHORDAL BIPARTITE VERTEX DELETION is still unknown. Coming to the question of polynomial kernels for these problems, the situation is even more grim. Until recently, the only known result was a polynomial kernel for PROPER INTERVAL VERTEX DELETION: Fomin et al. [19] obtained a $\mathcal{O}(k^{53})$ sized polynomial kernel for PROPER INTERVAL VERTEX DELETION, which has recently been improved to $\mathcal{O}(k^4)$ [27]. A dearth of further results in this area has led to the questions of kernelization complexity of CHORDAL VERTEX DELETION and INTERVAL VERTEX DELETION becoming prominent open problems [10, 13, 19, 24, 36].

Jansen and Pilipzuck [26] recently resolved one of these open questions. They showed that CVD admits a polynomial kernel of size $\mathcal{O}(k^{161}\log^{58} k)$, and further posed an open question:

Does CVD admit a kernel of size $\mathcal{O}(k^{10})$?

While we do not completely resolve this question, we design a significantly smaller kernel. In particular, we obtain the following result.

Theorem 1. CVD admits a polynomial kernel of size $\mathcal{O}(k^{12}\log^{10} k)$.

Our Methods. Our result is inspired by the $\mathcal{O}(k^2)$ -size kernel for FEEDBACK VERTEX SET (FVS) (checking whether there exists a k sized vertex subset that intersects all cycles), designed by Thomassé [40]. The kernel for FVS consists of the two following steps.

- 1. Reduce the maximum degree of the graph by using matching based tools (in particular expansion lemma). That is, upper bound the maximum degree Δ of the graph by $\mathcal{O}(k)$.
- 2. When the graph has maximum degree Δ , one can show that if a graph has minimum degree at least 3 (which can be easily achieved for FVS) then any minimum feedback vertex set has size $\mathcal{O}(n/\Delta)$. This together with an upper bound on Δ implies $\mathcal{O}(k^2)$ -size kernel.

Let us now look at the CVD problem. Here, our objective is to check whether there exists a k-sized vertex subset S such that $G \setminus S$ is a chordal graph. However, what is a chordal graph? A graph is chordal if it does not contain any induced cycle of length at least 4. That is, every cycle of length at least 4 has a chord. Thus, FVS is about intersecting all cycles and CVD is about intersecting all chordless cycles. Unfortunately, this apparent similarity stops here! Nevertheless, we are still able to exploit ideas used in the $\mathcal{O}(k^2)$ -size kernel for FVS. Towards this, we define the notion of *independence degree* of vertices and graphs. Roughly speaking, the independence degree of a vertex v is the size of a maximum independent set in its neighborhood (G[N(v)]). The study of this notion is our main conceptual contribution.

As a first step we bound the independence degree of every vertex by $k^{\mathcal{O}(1)}$ – this is similar to the first step of the kernel for FVS. Once we have bounded the independence degree of a graph, we obtain an approximate solution M (also called modulator) and analyze the graph $G \setminus M$. The bound on the independence degree immediately implies that the number of leaves, vertices of degree at least three and the number of maximal degree two paths in the clique forest of $G \setminus M$ is bounded by $k^{\mathcal{O}(1)}$. Then, using ideas similar to those used by Marx [36] to bound the size of a maximal clique while designing the first algorithm for CVD, we reduce the size of each maximal clique in $G \setminus M$ to $k^{\mathcal{O}(1)}$. Finally, we use structural analysis to bound the size of each maximal degree two path, which includes the design of a reduction rule that computes a family of minimum cuts, and thus we obtain our final kernel. We believe that the notion of independent degree is likely to be useful for designing algorithms for other graph modification problems. Not only does it lead to a significant improvement, once it is bounded, it greatly simplifies the analysis of the resulting instance.

Finally, after we obtain a kernel, we show that by reruning our entire kernelization procedure once, we can actually reduce the size of the kernel. Since we rely on an $\mathcal{O}(\log^2 n)$ -approximation algorithm, when we call our kernelization procedure for the first time, we work with an approximate solution of size $\mathcal{O}(k^3 \log^2 k)$; indeed, it can be assumed that $\log n < k \log k$, else the $2^{\mathcal{O}(k \log k)} \cdot n^{\mathcal{O}(1)}$ -time algorithm for CVD by Cao and Marx [10] solves the input instance in polynomial time. However, once we have our first kernel, it holds that $n = \mathcal{O}(k^{22} \log^{12} k)$. At this point, if we reuse the approximation algorithm, we obtain an approximate solution of size $\mathcal{O}(k \log^2 k)$. The size of our kernel depends on the size of the approximate solution; in particular, having an approximate solution of size $\mathcal{O}(k \log^2 k)$ allows us to obtain a kernel of size $\mathcal{O}(k^{12} \log^{10} k)$.

2 Preliminaries

For a positive integer k, we use [k] as a shorthand for $\{1, 2, \ldots, k\}$. Given a function $f : A \to B$ and a subset $A' \subseteq A$, we let $f|_{A'}$ denote the function f restricted to the domain A'.

Parameterized Complexity. In Parameterized Complexity each problem instance is accompanied by a parameter k. A central notion in this field is the one of *fixed-parameter tractability (FPT)*. This means, for a given instance (I, k), solvability in time $f(k)|I|^{\mathcal{O}(1)}$ where f is some computable function of k. A parameterized problem is said to admit a *polynomial kernel* if there is a polynomial-time algorithm (the degree of polynomial is independent of the parameter k), called a *kernelization algorithm*, that reduces the input instance down to an equivalent instance whose size is bounded by a polynomial p(k) in k. Here, two instances are equivalent if one of them is a yes-instance if and only if the other one is a yes-instance. The reduced instance is called a p(k)-kernel for the problem. For a detailed introduction to the field of kernelization, we refer to the following surveys [30, 34] and the corresponding chapters in the books [12, 15, 16, 38].

Kernelization algorithms often rely on the design of reduction rules. The rules are numbered, and each rule consists of a condition and an action. We always apply the first rule whose condition is true. Given a problem instance (I, k), the rule computes (in polynomial time) an instance (I', k') of the same problem where $k' \leq k$. Typically, |I'| < |I|, where if this is not the case, it should be argued why the rule can be applied only polynomially many times. We say that the rule safe if the instances (I, k) and (I', k') are equivalent.

Graphs. Given a graph G, we let V(G) and E(G) denote its vertex-set and edge-set, respectively. In this paper, we only consider undirected graphs. We let n = |V(G)| denote the number of vertices in the graph G, where G will be clear from context. The open neighborhood, or simply the neighborhood, of a vertex $v \in V(G)$ is defined as $N_G(v) = \{w \mid \{v, w\} \in E(G)\}$. The closed neighborhood of v is defined as $N_G[v] = N_G(v) \cup \{v\}$. The degree of v is defined as $d_G(v) = |N_G(v)|$. We can extend the definition of neighborhood of a vertex to a set of vertices as follows. Given a subset $U \subseteq V(G)$, $N_G(U) = \bigcup_{u \in U} N_G(u)$ and $N_G[U] = \bigcup_{u \in U} N_G[u]$. The induced subgraph G[U]is the graph with vertex-set U and edge-set $\{\{u, u'\} \mid u, u' \in U, \text{ and } \{u, u'\} \in E(G)\}$. Moreover, we define $G \setminus U$ as the induced subgraph $G[V(G) \setminus U]$. We omit subscripts when the graph Gis clear from context. An independent set in G is a set of vertices such that there is no edge between any pair of vertices in this set. The independence number of G, denoted by $\alpha(G)$, is defined as the cardinality of the largest independent set in G. A clique in G is a set of vertices such that there is an edge between every pair of vertices in this set.

A path P in G is a subgraph of G where $V(P) = \{x_1, x_2, \ldots, x_\ell\} \subseteq V(G)$ and $E(P) = \{\{x_1, x_2\}, \{x_2, x_3\}, \ldots, \{x_{\ell-1}, x_\ell\}\} \subseteq E(G)$ for some $\ell \in [n]$. The vertices x_1 and x_ℓ are called endpoints of the path P and the remaining vertices in V(P) are called internal vertices of P. We also say that P is a path between x_1 and x_ℓ . A cycle C in G is a subgraph of G where $V(C) = \{x_1, x_2, \ldots, x_\ell\} \subseteq V(G)$ and $E(C) = \{\{x_1, x_2\}, \{x_2, x_3\}, \ldots, \{x_{\ell-1}, x_\ell\}, \{x_\ell, x_1\}\} \subseteq E(G)$, i.e., it is a path with an additional edge between x_1 and x_ℓ . Let P be a path in the graph G on at least three vertices. We say that $\{u, v\} \in E(G)$ is a chord of P if $u, v \in V(P)$ but $\{u, v\} \notin E(P)$. Similarly, for a cycle C on at least four vertices, $\{u, v\} \in E(G)$ is a chord of C if $u, v \in V(C)$ but $\{u, v\} \notin E(C)$. A path P or cycle C is chordless if it has no chords.

The graph G is *connected* if there is a path between every pair of vertices, otherwise G is *disconnected*. A connected graph without any cycles is a *tree*, and a collection of trees is a *forest*. A maximal connected subgraph of G is called a *connected component* of G.

Forest Decompositions. A forest decomposition of a graph G is a pair (F,β) where F is forest, and $\beta: V(T) \to 2^{V(G)}$ is a function that satisfies the following,

- (i) $\bigcup_{v \in V(F)} \beta(v) = V(G),$
- (ii) for any edge $\{v, u\} \in E(G)$ there is a node $w \in V(F)$ such that $v, u \in \beta(w)$,

(iii) and for any $v \in V(G)$, the collection of nodes $T_v = \{u \in V(F) \mid v \in \beta(u)\}$ is a subtree of F.

For $v \in V(F)$, we call $\beta(v)$ the bag of v, and for the sake of clarity of presentation, we sometimes use v and $\beta(v)$ interchangeably. We refer to the vertices in V(F) as nodes. A tree decomposition is a forest decomposition where F is a tree.

Chordal Graphs. A graph G is a *chordal graph* if it has no chordless cycle as an induced subgraph, i.e., every cycle of length at least four has a chord. A *clique forest* of G is a forest decomposition of G where every bag is a maximal clique. The following lemma shows that the class of chordal graphs is exactly the class of graphs which have a clique forest.

Lemma 1 (Theorem 4.8, [23]). A graph G is a chordal graph if and only if G has a clique forest.

Given a subset $U \subseteq V(G)$, we say that U hits a chordless cycle C in G if $U \cap V(C) \neq \emptyset$. Observe that if U hits every chordless cycle of G, then $G \setminus U$ is a chordal graph. Given a $v \in V(G)$, we say that a vertex-set $B \subseteq V(G) \setminus \{v\}$ is a v-blocker if B hits every chordless cycle in G. Observe that the set B must not contain the vertex v.

The CHORDAL VERTEX DELETION (CVD) problem is defined as follows.

CHORDAL VERTEX DELETION (CVD)	Parameter: k
Input: An undirected graph G and a non-negative integer k .	
Question: Does there exist a subset $S \subseteq V(G)$ such that $ S $	$\leq k$ and $G \setminus S$ is a chordal
graph?	

For purposes of approximation, we also formulate this problem as an optimization problem.

CHORDAL VERTEX DELETION (CVD) **Input:** An undirected graph G. **Question:** What is the minimum cardinality of a subset $S \subseteq V(G)$ such that $G \setminus S$ is a chordal graph?

The Expansion Lemma. Let c be a positive integer. A *c-star* is a graph on c + 1 vertices where one vertex, called the center, has degree c, and all other vertices are adjacent to the center and have degree one. A *bipartite graph* is a graph whose vertex-set can be partitioned into two independent sets. Such a partition the vertex-set is called a *bipartition* of the graph. Let G be a bipartite graph with bipartition (A, B). A subset of edges $M \subseteq E(G)$ is called *c-expansion of* A*into* B if

- (i) every vertex of A is incident to exactly c edges of M,
- (ii) and M saturates exactly c|A| vertices in B.

Note that a *c*-expansion saturates all vertices of A, and for each $u \in A$ the set of edges in M incident on u form a *c*-star. The following lemma allows us to compute a *c*-expansion in a bipartite graph. It captures a certain property of neighborhood sets which is very useful for designing kernelization algorithms.

Lemma 2 ([40, 12]). Let G be a bipartite graph with bipartition (A, B) such that there are no isolated vertices in B. Let c be a positive integer such that $|B| \ge c|A|$. Then, there are non-empty subsets $X \subseteq A$ and $Y \subseteq B$ such that

- there is a c-expansion from X into Y,
- and there is no vertex in Y that has a neighbor in $A \setminus X$, i.e. $N_G(Y) = X$.

Further, the sets X and Y can be computed in polynomial time.

3 Kernelization

In this section we prove Theorem 1. First, in Section 3.1, we briefly state results relating to approximate solutions for CVD, which will be relevant to following subsections. In Section 3.2 we address annotations that will be added to the input instance. Next, in Section 3.3, we introduce the notion of the *independent degree* of a vertex, which lies at the heart of the design of our kernelization algorithm. We carefully examine the independent degrees of vertices in our graphs, and show how these degrees can be bounded by a small polynomial in k. In Section 3.4, we consider the clique forest of the graph obtained by removing (from the input graph) the vertices of an approximate solution. In particular, we demonstrate the usefulness of our notion of an independent degree of a vertex – having bounded the independent degree of each vertex, we show that the number of leaves in the clique forest can be bounded in a simple and elegant manner. We also efficiently bound the size of a maximal clique. Then, in Section 3.5, we turn to bound the length of degree-2 paths in the clique forest. This subsection is quite technical, and its proofs are based on insights into the structure of chordal graphs and their clique forests. In particular, we use a reduction rule which computes a collection of minimum cuts rather than one minimum cut which overall allows us to capture the complexity of a degree-2 path using only few vertices. In Section 3.6, we remove annotations introduced in preceding sections. Next, in Section 3.7, we bound the size of our kernel. Finally, in Section 3.8, we show that an alternating application of approximation and kernelization can improve the performance of our kernelization algorithm.

3.1 Approximation

Observe that it can be assumed that $\log n < k \log k$, else the $2^{\mathcal{O}(k \log k)} \cdot n^{\mathcal{O}(1)}$ -time algorithm for CVD by Cao and Marx [10] solves the input instance in polynomial time. Thus, since CVD admits an $\mathcal{O}(\log^2 n)$ -factor approximation algorithm [3], we obtain the following result.

Lemma 3. CVD admits an $\mathcal{O}(k^2 \log^2 k)$ -factor approximation algorithm.

Throughout Section 3, we let APPROX denote a polynomial-time algorithm for CVD that returns approximate solutions of size $f(\mathsf{opt})$ for some function f. Initially, it will denote the algorithm given by Lemma 3. Given an instance of CVD, we say that an approximate solution Dis *redundant* if for every vertex $v \in D$, $D \setminus \{v\}$ is also an approximate solution, that is, $G \setminus (D \setminus \{v\})$ is a chordal graph. Jansen and Pilipczuk [26] showed that given an approximate solution of size g(k) for some function g, one can find (in polynomial time) either a vertex contained in every solution of size at most k or a redundant approximate solution of size $\mathcal{O}(k \cdot g(k))$. Thus, we have the following result.

Corollary 4 ([26]). Given an instance of CVD, one can find (in polynomial time) either a vertex contained in every solution of size at most k or a redundant approximate solution of size $\mathcal{O}(k \cdot f(k))$.

Next, we fix an instance (G, k) of CVD. By relying on APPROX and Corollary 4, we may assume that we have a vertex-set $\widetilde{D} \subseteq V(G)$ that is an approximate solution of size f(k) and a vertex-set $D \subseteq V(G)$ that is a redundant approximate solution of size $c \cdot k \cdot f(k)$ for some constant c independent of the input.

In the following subsection, we will also need to strengthen approximate solutions to be v-blockers for some vertices $v \in V(G)$. To this end, we will rely on the following result.

Lemma 5. Given a vertex $v \in V(G)$, one can find (in polynomial time) either a vertex contained in every solution of size at most k or an approximate solution of size f(k) that is a v-blocker.

Proof. Fix a vertex $v = v_0 \in V(G)$. We define the graph G' by setting $V(G') = V(G) \cup \{v_1, v_2, \ldots, v_{f(k)}\}$, where $v_1, v_2, \ldots, v_{f(k)}$ are new vertices, and $E(G') = E(G) \cup \{\{u, v_i\} : \{u, v\} \in E(G), i \in [f(k)]\} \cup \{(v_i, v_j) : i \neq j \in \{0\} \cup [f(k)]\}$. In other words, G' is the graph G to which we add f(k) copies of the vertex v, which form a clique amongst themselves and v.

We call the algorithm APPROX to obtain an approximate solution S. Since $G'[\{v_0, v_1, \ldots, v_{f(k)}\}]$ is a clique, any chordless cycle in G' contains at most one vertex from $\{v_0, v_1, \ldots, v_{f(k)}\}$. In particular, for any chordless cycle C in G', by replacing v_i by v_0 (in case v_i belongs to C), we obtain a chordless cycle in G. Therefore, if the instance (G, k) admits a solution of size at most k that does not contain v, then this solution is also a solution for the instance (G', k), which implies that the size of S should be at most f(k). Thus, we can next assume that $|S| \leq f(k)$, else we conclude that the vertex v is contained in every solution of size at most k. Since the vertices $v_0, v_1, v_2, \ldots, v_{f(k)}$ have the same neighbor-set, if $\{v_0, v_1, v_2, \ldots, v_{f(k)}\} \cap S \neq \emptyset$, we can assume that $\{v_0, v_1, v_2, \ldots, v_{f(k)}\} \subseteq S$, since otherwise we can take $S \setminus \{v_0, v_1, v_2, \ldots, v_{f(k)}\} \cap S = \emptyset$. In particular, we have that $v \notin S$ and therefore S is a v-blocker.

In light of Lemma 5, we may next assume that for every vertex $v \in V(G)$, we have a vertex-set B_v that is both a v-blocker and an approximate solution of size f(k).

3.2 Irrelevant and Mandatory Edges

During the execution of our kernelization algorithm, we mark some edges in E(G) as *irrelevant* edges. At the beginning of its execution, all of the edges in E(G) are assumed to be relevant

edges. When we mark an edge as an irrelevant edge, we prove that any solution that hits all of the chordless cycles in G that contain only relevant edges also hits all of the chordless cycles in G that contain the irrelevant edge. In other words, we prove that we can safely ignore chordless cycles that contain at least one irrelevant edge. Observe that we cannot simply remove irrelevant edges from E(G) since this operation may introduce new chordless cycles in G. Instead we maintain a set E_I , which contains the edges marked as irrelevant.

We also mark some edges in E(G) as mandatory edges. We will ensure that at least one endpoint of a mandatory edge is present in any solution of size at most k. We let E_M denote the set of mandatory edges.

In some situations, we identify a pair of non-adjacent vertices such that any solution of size at most k must contain at least one of them. Then, we add an edge between the vertices of the pair, and mark this edge as a mandatory edge. The correctness of this operation follows from the observation that any chord correct edge is marked by the addition of this edge contains both vertices of the pair, and since the edge is marked as a mandatory edge, we ensure this chord correct edge will be hit although it may no longer exist. Formally, we have the following reduction rule.

Reduction Rule 1. Given two non-adjacent vertices in G, v and u, such that at least one of them belongs to any solution of size at most k, insert the edge $\{v, u\}$ into both E(G) and E_M .

Hence from now onwards our instance is of the form (G, k, E_I, E_M) , and during the execution of our kernelization algorithm, we will update the sets E_I and E_M . In Section 3.6, we show that we can unmark the edges in $E_I \cup E_M$, obtaining an ordinary instance of CVD. For the sake of simplicity, when E_I and E_M are clear from context, we omit them.

The Number of Mandatory Edges. If a vertex v is incident to at least k + 1 mandatory edges, it must belong any solution of size at most k. Therefore, we may safely apply the following reduction rule.

Reduction Rule 2. If there exists a vertex v incident to at least k+1 mandatory edges, remove v from G and decrement k by 1.

After exhaustively applying the above reduction rule we have the following lemma.

Lemma 6. If $|E_M| > k^2$ then the input instance is a no-instance.

Proof. After the exhaustive application of Reduction Rule 2, any vertex incident to a mandatory edge is incident to at most k such edges. Therefore, any set of at most k vertices from V(G) may intersect at most k^2 mandatory edge. Since every solution of size at most k must intersect every mandatory edge, we deduce that if $|E_M| > k^2$, the input instance is a no-instance.

Thus, we will next assume that $|E_M| \leq k^2$ (else Reduction Rule 2 applies). Moreover, we let D' denote the set $D \cup D$ to which we add every vertex that is an endpoint of a vertex in E_M (recall that D is our approximate solution of size at most f(k) and D is our redundant approximate solution of size $\mathcal{O}(k \cdot f(k))$). Observe that by adding vertices to a redundant approximate solution, it remains a redundant approximate solution, and therefore D' or any other superset of D is such a solution.

3.3 Independent Degree

Given a vertex $v \in V(G)$, we use the notation $N_G^R(v)$ to refer to the set of each vertex $u \in N_G(v)$ such that $\{v, u\}$ does not belong to $E_I \cup E_M$. We remark that in this subsection we identify and mark some edges as irrelevant edges.

Independent Degrees. We start by introducing the notion of the independent degree of vertices and graphs.

Definition 7. Given a vertex $v \in V(G)$, the independent degree of v, denoted by $d_G^I(v)$, is the size of a maximum independent set in the graph $G[N_G^R(v)]$. The independent degree of G, denoted by Δ_G^I , is the maximum independent degree of a vertex in V(G).

Fix $\Delta = (k+3)f(k)$. The objective of this subsection is to investigate the notion of an independent degree, ultimately proving the following result.

Lemma 8. One can construct (in polynomial time) an instance (G', k', E'_I, E'_R) of CVD that is equivalent to the input instance (G, k, E_I, E_R) and such that both $k' \leq k$ and $\Delta^I_{G'} \leq \Delta$.

To this end, we may assume that we are given a vertex $v \in V(G)$ such that $d_G^I(v) > \Delta$. We say that an instance (G', k', E'_I, E'_M) of CVD is *better* than the input instance (G, k, E_I, E_M) if $k' \leq k, V(G') = V(G), E_I \subseteq E'_I, E_M \subseteq E'_M d_{G'}^I(v) \leq \Delta$ and for all $u \in V(G'), d_{G'}^I(u) \leq d_G^I(u)$. To prove the correctness of Lemma 8, it is sufficient to prove the correctness of the following lemma.

Lemma 9. We can construct (in polynomial time) an instance (G', k', E'_I, E'_M) of CVD that is better than the input instance (G, k, E_I, E_M) .

Indeed, to prove Lemma 8, one can repeatedly apply the operation given by Lemma 9 in the context of every vertex $u \in V(G)$ such that $d_G^I(u) > \Delta$. We start with a simple result concerning independent degrees.

Lemma 10. Let $u \in V(G)$ be a vertex such that $d_G^I(u) \ge |B_u|$. Then, one can find (in polynomial time) an independent set in $G[N_G^R(u) \setminus B_u]$ of size at least $d_G^I(u) - |B_u|$.

Proof. Since B_u is a solution, $G \setminus B_u$ is a chordal graph. In particular, $G[N_G^R(u) \setminus B_u]$ is a chordal graph. Since chordal graphs are perfect, MAXIMUM INDEPENDENT SET in chrodal graphs is solvable in polynomial time [23]. This means that we can find a maximum independent set in $G[N_G^R(u) \setminus B_u]$ in polynomial time. Since the size of a maximum independent set in $G[N_G^R(u) \setminus B_u]$ is at least $d_G^I(u)$, the size of a maximum independent set in $G[N_G^R(u) \setminus B_u]$ is at least $d_G^I(u) - |B_u|$, which concludes the correctness of the lemma.

Recall that for any vertex $u \in V(G)$, $|B_u| \leq f(k)$. Thus, we have the following corollary.

Corollary 11. One can find (in polynomial time) an independent set in $G[N_G^R(v) \setminus B_v]$ of size at least $\Delta - f(k)$.

We let I denote the independent set given by Corollary 11.

Independent Components. Let $X = N_G(v) \setminus (B_v \cup I)$ denote the neighbor-set of v from which we remove the vertices of the v-blocker B_v and of the independent set I. We also let $H = G \setminus (\{v\} \cup B_v \cup X)$ denote the graph obtained by removing (from G) the vertex v, the v-blocker B_v and any neighbor of v that does not belong to the independent set I. We define the set independent components of v as the set of each connected component of H that contains at least one vertex from I, and denote this set by \mathcal{A} .

For the set \mathcal{A} , we prove the following lemmata.

Lemma 12. Each connected component $A \in \mathcal{A}$ contains exactly one vertex from I and no other vertex from $N_G(v)$.

Proof. The graph H does not contain any vertex from $N_G(v) \setminus I$, and therefore we only need to prove the first part of the statement of the lemma. Fix a connected component $A \in \mathcal{A}$. By the definition of \mathcal{A} , it is only necessary to prove that A cannot contain (at least) two vertices from I. Suppose, by way of contradiction, that it contains two such vertices, u and w. Let P denote



Figure 1: The relations between v, B_v, X and \mathcal{A} .

the shortest path in A that connects u and w. Since I is an independent set, this path contains at least two edges. Therefore, together with the vertex v, the path P forms a chordless cycle. However, this chordless cycle contain no vertex from B_v , which contradicts the fact that B_v is an approximate solution.

By Lemma 12, for each connected component $A \in \mathcal{A}$ we can let $z(A) \in I$ denote the unique neighbor of v in A. In fact, Corollary 11 and Lemma 12 imply that $\Delta - f(k) \leq |\mathcal{A}|$.

Lemma 13. Every vertex $x \in X$ that is adjacent (in G) to some vertex $y \in V(A)$, where $A \in A$, is also adjacent (in G) to the vertex z(A).

Proof. Let $x \in X$ be a vertex that is adjacent (in G) to some vertex $y \in V(A)$. Suppose, by way of contradiction, that $\{x, z(A)\} \notin E(G)$. Let P denote the shortest path in A that connects y and z(A). This path contains at least two edges. Therefore, together with the vertex v, the path P forms a chordless cycle. However, this chordless cycle contain no vertex from B_v , which contradicts the fact that B_v is an approximate solution.

An illustration of the relations between $v, B_v X$ and \mathcal{A} is given in Fig. 1.

The Bipartite Graph \widehat{H} . To decrease $d_G^I(u)$, we will consider the bipartite graph \widehat{H} , which is defined as follows. We define vertex-set of \widehat{H} by $V(\widehat{H}) = \mathcal{A} \cup B_v$. In this context, we mean that each connected component $A \in \mathcal{A}$ is represented by a vertex in $V(\widehat{H})$, and for the sake of simplicity, we use the symbol A also to refer to this vertex. We partition B_v into two sets, B_c and B_f , where B_c contains the vertices in B_v that are adjacent (in G) to v, while B_f contains the remaining vertices in B_v . Here, the letters c and f stand for "close" and "far". Having this partition, we define the edge-set of \widehat{H} as follows. For every vertex $b \in B_c$ and connected component $A \in \mathcal{A}$ such that $b \in N_G(V(A)) \setminus N_G(z(A))$ (i.e., b is a neighbor of some vertex in Abut not of the vertex z(A)), insert the edge $\{b, A\}$ into $E(\widehat{H})$. Moreover, for every vertex $b \in B_f$ and connected component $A \in \mathcal{A}$ such that $b \in N_G(V(A))$, insert the edge $\{b, A\}$ into $E(\widehat{H})$. An illustration of the bipartite graph \widehat{H} is given in Fig. 2. The motivation behind its definition lies at the following lemma.

Lemma 14. The bipartite graph \hat{H} satisfies the following properties.

1. Suppose that we are given an edge $\{b, A\} \in E(\widehat{H})$ such that $b \in B_c$ and $A \in \mathcal{A}$. Then, the graph G has a chordless cycle defined by the edges $\{v, b\}$ and $\{v, z(A)\}$ and a path in A.



Figure 2: The bipartite graph \hat{H} .

2. Suppose that we are given edges $\{b, A\}, \{b, A'\} \in E(\widehat{H})$ such that $b \in B_f$ and $A, A' \in \mathcal{A}$. Then, the graph G has a chordless cycle defined by the edges $\{v, z(A)\}, \{v, z(A')\},$ the vertex b and paths in A and A'.

Proof. To prove the first item, let y be a vertex in $N_G(b) \cap V(A)$, whose existence is guaranteed by the assumption that $\{b, A\} \in E(\widehat{H})$. The desired chordless cycle can be defined by the edges $\{v, b\}, \{v, z(A)\}, \{b, y\}$ and the edge-set of a shortest path between z(A) and y in A – in particular, by the definition of \mathcal{A} and Lemma 12, v is not adjacent to any vertex on this path, excluding z(A), and since $\{b, A\} \in E(\widehat{H})$, b is not adjacent to z(A).

For the second item, choose $y \in N_G(b) \cap V(A)$ and $y' \in N_G(b) \cap V(A')$, whose existence is guaranteed by the assumption that $\{b, A\}, \{b, A'\} \in E(\widehat{H})$, such that the length of a shortest path between z(A) (z(A')) and y (resp. y') in A (resp. A') is minimum. Observe that the cycle defined $\{v, z(A)\}, \{v, z(A')\}, \{y, b\}, \{y', b\}$ and the shortest paths between z(A) and y in A and z(A') and y' in A', respectively, is a chordless cycle – in particular, by the definition of \mathcal{A} , Lemma 12 and since $b, b' \in B_f$, the cycle cannot have a chord containing v.

Isolated Vertices in \hat{H} . We start investigating the bipartite graph \hat{H} by examining the isolated vertices in \mathcal{A} that it contains. In this context, we need the two following lemmata.

Lemma 15. Let $A \in \mathcal{A}$ be an isolated vertex in \widehat{H} , and denote z = z(A). Then, $N_G(V(A)) = N_G(z) \setminus V(A)$.

Proof. Since $z \in A$, it is clear that $N_G(z) \setminus V(A) \subseteq N_G(V(A))$. Next, we show that $N_G(V(A)) \subseteq N_G(z) \setminus V(A)$. To this end, consider some vertices $y \in V(A)$ and $u \in N_G(y) \setminus V(A)$, and suppose, by way of contradiction, that $u \notin N_G(z)$. Because A is a connected component of H, it holds that $u \in \{v\} \cup B_v \cup X$. Moreover, because A is isolated in \widehat{H} and yet $u \in N_G(V(A)) \setminus N_G(z)$, it further holds that $u \in X$. However, this results in a contradiction to the statement of Lemma 13.

Lemma 16. Let $A \in \mathcal{A}$ be an isolated vertex in \widehat{H} , and denote z = z(A). Then, G does not have a chordless cycle that contains the edge $\{v, z\}$.

Proof. Suppose, by way of contradiction, that G has a chordless cycle C that contains the edge $\{v, z\}$. The cycle C must contain an edge $\{y, u\}$ such that $y \in A$ and $u \notin A$. By Lemma 15, $N_G(V(A)) = N_G(z) \setminus A$, and therefore $\{z, u\} \in E(G)$. Hence, since the cycle C is chordless, it holds that z = y, which, in turn, implies that $u \notin N_G(v)$. We deduce that u must belong to B_f .

However, this implies that $\{A, u\} \in \widehat{H}$, contradicting the assumption that A is an isolated vertex in \widehat{H} .

Lemma 16 leads us to the design of the following reduction rule.

Reduction Rule 3. If the graph \hat{H} contains an isolated vertex $A \in A$, mark the edge $\{v, z\}$ as irrelevant.

After an exhaustive application of this rule, we can assume that \widehat{H} does not contain an isolated vertex $A \in \mathcal{A}$.

Applying the Expansion Lemma. Next, we would like to apply Lemma 2 in the context of the bipartite graph \widehat{H} . Since $|\mathcal{A}| \ge \Delta - f(k) \ge (k+2) \cdot |B_v|$ and we have already ensured that \widehat{H} does not contain any isolated vertex $A \in \mathcal{A}$, this lemma implies that we can find (in polynomial time) subsets $\mathcal{A}^* \subseteq \mathcal{A}$ and $B^* \subseteq B_v$ such that there exists a (k+2)-expansion from B^* into \mathcal{A}^* .

The usefulness of B^* is stated in the following lemma.

Lemma 17. Any solution of size at most k to the input instance that does not contain v contains all of the vertices in B^* .

Proof. Suppose that the input instance is a yes-instance, and suppose it has a solution S of size at most k that does not contain v. Consider some vertex $b \in B^*$. Let $A_1, A_2, \ldots, A_{k+2}$ be the neighbors of b in \widehat{H} that correspond to our (k+2)-expansion. For any choice of A_i and A_j , Lemma 14 implies that if there is no chordless cycle defined by v, b and a path in A_i , then there is a chordless cycle defined by v, b, a path in A_i and a path in A_j . Therefore, if S contains neither v nor b, it must contain at least one vertex from each connected component A_i excluding at most one such component. However, there are k + 2 such components, and since S does not contain v, we deduce that it contains b. The choice of $b \in B^*$ was arbitrary, and therefore we conclude that S contains all of the vertices in B^* .

Decreasing the Independent Degree of v**.** Armed with Lemma 17, we can apply the following reduction rule.

Reduction Rule 4. For each vertex $b \in B^*$, insert the edge $\{b, v\}$ into E(G) (if it is not already present), and mark $\{b, v\}$ as a mandatory edge. Moreover, mark each edge $\{v, z(A)\}$ such that $A \in \mathcal{A}^*$ as an irrelevant edge.

Lemma 18. Reduction Rule 4 is safe.

Proof. First we claim that adding a mandatory edge between v and some $b \in B^*$ is safe. Let G' denote the graph obtained after applying this operation. By Lemma 17, at least one vertex among v and b must be present in any solution of size at most k. Therefore, we may mark the newly added edge $\{v, b\}$ as a mandatory edge. Since this edge is a mandatory edge, any solution of size at most k to (G', k) will hit any new chordless cycle created by adding this edge as well as any chordless cycle C that exists in G even if in G'[V(C)] is not a chordless cycle (since if G'[V(C)] is not a chordless cycle, $\{v, b\} \subseteq V(C)$). By adding the edges one-by-one, arguing that each operation is safe given that the preceding operation was safe, we derive the safeness of the insertion of all of the edges in $\{\{b, v\} : b \in B^*\}$ as well as the marking of these edges as mandatory.

Now, let G' denote the graph obtained after the application of Reduction Rule 4. Let us argue that it is indeed safe to ignore (in G') any chordless cycle containing an edge $\{v, z(A)\}$ such that $A \in \mathcal{A}^*$. To this end, consider such a chordless cycle C and a solution S of size at most k to (G', k). If S contains v, it clearly hits C. Otherwise, we have that S contains all of the vertices in B^* . Recall that by Lemma 12, z(A) is the only neighbor of v in A, and by Lemma 13, $N_G(z(A)) \cap X = N_G(V(A)) \cap X$. Therefore, since the cycle C is chordless, it must contain a vertex from B_v , but then, by Lemma 2, we further deduce that it must contain a vertex from B^* . We thus get that C is hit by S.

Reduction Rule 4 decreases $|N_G^R(v)|$. Moreover, as long as $d_G^R(v) > \Delta$, we can apply this rule. Thus, after an exhaustive application of this rule, it should hold that $d_G^R(v) \leq \Delta$. Furthermore, this rule neither inserts vertices into V(G) nor unmarks edges, and therefore we conclude that Lemma 9 is correct. Denote $\Delta' = \Delta + k$. Thus, by Reduction Rule 2, the size of a maximum independent set in the neighborhood of each vertex in the graph G from which we remove irrelevant edges is bounded by Δ' .

3.4 The Clique Forest

Let F denote the clique forest associated with the chordal graphs $G \setminus D'$. Towards bounding the number of leaves in F, we need the following lemma.

Lemma 19. Let I be an independent set in the graph $G \setminus D'$. Then, there are most $|D'| \cdot \Delta'$ relevant edges between vertices in D' and vertices in I.

Proof. The claim follows from the observation that since $\Delta_G^I \leq \Delta$, $|E_M| \leq k^2$ and I is an independent set, for every vertex $v \in D'$, there are at most Δ' relevant edges between v and vertices in I.

We will also need the following reduction rule.

Reduction Rule 5. If there exists a vertex v in $G \setminus D'$ such that the vertices in $N_G(v)$ which are connected to v via relevant edges form a clique, remove the vertex v from G.

Lemma 20. Reduction Rule 5 is safe.

Proof. The special choice of the vertex v implies that every chordless cycle containing v must contain at least one irrelevant edge, and can therefore be ignored. We thus conclude that it is safe to remove the vertex v from G.

The bound on the number of leaves will follow from a bound on the number of bags containing *private vertices*, which are defined as follows.

Definition 21. A vertex v in $G \setminus D'$ is a private vertex if there exists only one bag in F that contains it.

Lemma 22. The number of bags in F containing private vertices is bounded by $|D'| \cdot \Delta'$.

Proof. Let ℓ denote the number of bags in F containing private vertices. By the definition of a clique forest, if we take exactly one private vertex from each bag in F, we obtain an independent set. Therefore, the graph G contain an independent set of size at least ℓ . Let I denote this independent set. Observe that since each vertex in I is a private vertex, its neighborhood in $G \setminus D'$ forms a clique. Therefore, after an exhaustive application of Reduction Rule 5, each vertex in I must be connected by at least one relevant edge to a vertex in D'. By Lemma 19, we conclude that $\ell \leq |D'| \cdot \Delta'$.

We are now ready to bound the number of leaves and nodes of degree at least 3 in the clique forest F.

Lemma 23. Both the number of leaves in F and the number of nodes of degree at least 3 in F are bounded by $|D'| \cdot \Delta'$.

Proof. Observe that since every leaf in F corresponds to a maximal clique in $G \setminus D'$, it contains a private vertex. Thus, by Lemma 22, the number of leaves is bounded by $|D'| \cdot \Delta'$. Since in a forest, the number of nodes of degree at least 3 is bounded by the number of leaves, we conclude that the lemma is correct.

Next, we turn to bound the size of a bag of F, to which end we prove the correctness of the following lemma.

Lemma 24. In polynomial time we can produce an instance (G', k') equivalent to (G, k) such that $k' \leq k$ and the size of any maximal clique in G' is bounded by $\kappa = c \cdot (|\widetilde{D}|^3 \cdot k + |\widetilde{D}| \cdot \Delta' \cdot (k+2)^3)$. (Here c is some constant independent of the input.)

The proof of this lemma closely follows the approach using which Marx [36] bounds the size of maximal cliques, but requires a few adaptations since the purpose of the paper [36] is to give a parameterized algorithm for CVD, which runs in exponential time, while we need to reduce instances in polynomial time. However, due to the similarity between our proof and this approach, details are deferred to Appendix A.

Observe that the size of each bag of F is bounded by the size of a maximal clique of $G \setminus D'$. Furthermore, since $G \setminus D'$ is a subgraph of G, the size of a maximal clique of $G \setminus D'$ is bounded by the size of a maximal clique of G. Thus, having applied the procedure given by Lemma 24, we have the following result.

Lemma 25. The size of any bag of F is upper bounded by κ .

3.5 The Length of Degree-2 Paths

The Family of Paths \mathcal{P} . Let V_F denote the set of each node of degree at least 3 in the forest F as well as each node whose bag has at least one private vertex. Let \mathcal{P} denote the set of paths whose endpoints belong to V_F and such that all of their internal nodes do not belong to V_F . Clearly, it holds that $|\mathcal{P}| \leq |V_F|$. By Lemmata 22 and 23, we have the following observation.

Observation 26. $|\mathcal{P}| \leq 2|D'| \cdot \Delta'$.

Thus, in light of Lemma 25, by bounding the maximum number of nodes on each path in \mathcal{P} , we can bound the total number of vertices in the graph. To this end, we fix some path $P \in \mathcal{P}$. Moreover, we orient the path from left to right, where the choice of the leftmost and rightmost nodes is arbitrary.

Partitioning the Path P**.** Next, we will partition P into more "manageable paths". To this end, we need the following definition.

Definition 27. We say that a subpath Q of P complies with a vertex $d \in D'$ if at least one of the following conditions holds.

- 1. For every two bags B and B' on Q, both $B \subseteq N_G(d)$ and $B' \subseteq N_G(d)$.
- 2. Let B_1, B_2, \ldots, B_t denote the bags of Q ordered from left to right. Then, at least one of the following condition holds.
 - (a) $B_1 \cap N_G(d) \subseteq B_2 \cap N_G(d) \subseteq \ldots \subseteq B_t \cap N_G(d).$
 - (b) $B_t \cap N_G(d) \subseteq B_{t-1} \cap N_G(d) \subseteq \ldots \subseteq B_1 \cap N_G(d).$

In particular, $N_G(d)$ is a subset of at least one of the two bags B_1 and B_t .

We would like to find a set \mathcal{B} of at most $\mathcal{O}(|D'|)$ bags on the path P such that after their removal from P, the following lemma will be true.

Lemma 28. Each subpath resulting from the removal of the bags in \mathcal{B} from P complies with every vertex in D'.

The rest of this subsubsection concerns the proof of this lemma. To prove it, it is sufficient to show that for each vertex $d \in D'$, we can find $\mathcal{O}(1)$ bags such that after their removal from P, each of the resulting subpaths complies with d. To this end, fix some vertex $d \in D'$.

First, we need the following lemma.

Lemma 29. Let $u, v \in N_G(d) \setminus D'$ be non-adjacent vertices, B_u be a bag containing u such that no bag to its right (on P) contains u, and B_v be a bag containing v such that not bag to its left (on P) contains v. Then, d is adjacent to every vertex in every bag that lies strictly between B_u and B_v .

Proof. Let z be a vertex in a bag that lies strictly between B_u and B_v . Since this bag is not an endpoint of the path P, z belongs to at least two bags that lie between B_u and B_v . Denote the path between B_u and B_v by $B_u = B_1 - B_2 - B_3 - \ldots - B_t = B_v$. Let B_i be the leftmost bag containing z, and let B_j be the rightmost bag containing z. Suppose, by way of contradiction, that $z \notin N_G(d)$.

We claim that there exists a path, P_1 , from u to z such that none of its internal vertices belongs to $B_{i+1} \cup B_{i+2} \ldots \cup B_v$. The proof by induction on the number i. When i = 1, we have that $B_i = B_u$. Then, since the vertices in B_u form a clique, the claim is correct. Now, suppose that the claim holds for $i - 1 \ge 1$, and let us prove it for i.

Since the vertices of the bag B_i form a maximal clique in the graph $G \setminus D'$, and the bag B_i does not contain private vertices, we have that $B_i \subseteq B_{i-1} \cup B_{i+1}$ and $B_i \setminus B_{i+1} \subset B_{i-1}$ is non-empty. Consider a vertex $w \in B_i \setminus B_{i+1}$ and observe that z and w are adjacent in G (since both belong to B_i), and further the vertex w is not present in $B_{i+1} \cup B_{i+2} \ldots \cup B_v$. By induction, there is a path P from u to w whose internal vertices do not belong to $B_i \cup B_{i+1} \ldots \cup B_v$. Appending the edge (w, z) to P gives us the path P_1 .

Similarly, there is a path P_2 from z to v such that none of its internal vertices belongs to $B_u \cup \ldots \cup B_{i-2} \cup B_{i-1}$. Since i < j, the paths P_1 and P_2 have no common vertex except z, and there is no edge between an internal vertex of P_1 and an internal vertex of P_2 .

Let P'_1 be the subpath of P_1 from u' to z, where u' is the last vertex in P_1 adjacent to d. Similarly, let P'_2 be the subpath of P_2 from z to v', where v' is the first vertex in P_2 adjacent to d. We may assume that P'_1 and P'_2 do not contain chords, else we can replace P'_1 and P'_2 by a chordless subpath of P'_1 and chordless subpath of P'_2 , respectively, which will still contain u', z and v'. The cycle $d - P'_1 - z - P'_2 - d$ is a chordless cycle in G, contradicting the fact that $G \setminus (D' \setminus \{d\})$ is a chordal graph. \Box

We also need the following notation. Let B_{ℓ} be the leftmost bag on P that contains a neighbor v_{ℓ} of d such that v_{ℓ} does not belong to any bag to the right of B_{ℓ} . Similarly, let B_r be the rightmost bag on P that contains a neighbor v_r of d such that v_r does not belong to any bag to the left of B_{ℓ} .

Lemma 30. Let B and B' be two bags on P that do not lie on the right of B_{ℓ} and such that B lies on the left of B'. Then, it holds that $B \cap N_G(d) \subseteq B' \cap N_G(d)$.

Proof. Suppose, by way of contradiction, that $B \cap N_G(d) \not\subseteq B' \cap N_G(d)$. However, this implies that B contains a neighbor v of d such that v does not belong to any bag to the right of B, which contradicts the choice of B_{ℓ} .

Lemma 31. Let B and B' be two bags on P that do not lie on the left of B_r and such that B lies to the right of B'. Then, it holds that $B \cap N_G(d) \subseteq B' \cap N_G(d)$.

Proof. The proof of this lemma is symmetric to the proof of Lemma 30.

By Lemmas 29–31, each of the subpaths resulting from the removal of B_{ℓ} and B_r from P complies with d. Thus, we conclude that Lemma 28 is correct.

Handling a Manageable Path. We now examine a subpath of P, denoted by Q, which complies with every vertex $d \in D'$. We will devise reduction rules such that after applying them exhaustively, the number of vertices in the union of the bags of the path Q will be bounded by $\mathcal{O}(\kappa)$.

Let B_1, B_2, \ldots, B_t denote the bags of Q ordered from left to right. Moreover, denote $V(Q) = \bigcup_{i=1}^t B_i$ and $A = \bigcap_{i=1}^t B_i$. We partition D' into two sets D_a and D_p , where $D_a = \{d \in D' : V(Q) \subseteq N_G(d)\}$ and $D_p = D' \setminus D_a$. Here the letters a and p stand for "all" and "partial", respectively. Definition 27 directly implies that the following observation is correct.

Observation 32. For every vertex $d \in D_p$, either (i) $B_1 \cap N_G(d) \subseteq B_2 \cap N_G(d) \subseteq \ldots \subseteq B_t \cap N_G(d)$ or (ii) $B_t \cap N_G(d) \subseteq B_{t-1} \cap N_G(d) \subseteq \ldots \subseteq B_1 \cap N_G(d)$. In particular, either (i) $N_G(d) \cap V(Q) \subseteq B_1$ or (ii) $N_G(d) \cap V(Q) \subseteq B_t$.

We also denote $U = V(Q) \setminus (B_1 \cup B_t)$. It is sufficient to ensure that $|U| = \mathcal{O}(\kappa)$ since then, by Lemma 25, $|V(Q)| = \mathcal{O}(\kappa)$. Thus, we can next suppose that $|U| > \delta$ where $\delta = 2(k+1) + 6\kappa$.

For each pair of non-adjacent vertices in D_a , we apply Reduction Rule 1. The correctness of this operation is given by the following lemma.

Lemma 33. Let u and v be two distinct non-adjacent vertices in D_a . Then, every solution of size at most k contains at least one of the vertices u and v.

Proof. Since $|U| > \delta$ and the size of each bag is bounded by κ , standard arguments on weighted paths imply that there exists $i \in [t]$ such that $|\bigcup_{j=1}^{i} B_j| > \delta/2 - \kappa$ and $|\bigcup_{j=i+1}^{t} B_j| > \delta/2 - \kappa$; indeed, we may iteratively increase i from 1 to t until we reach the first time where it holds that $|\bigcup_{j=1}^{i} B_j| > \delta/2 - \kappa$, in which case it will also hold that $|\bigcup_{j=i+1}^{t} B_j| > \delta/2 - \kappa$. Denote $U^1 = U \cap ((\bigcup_{j=1}^{i-1} B_j) \setminus (B_1 \cup B_i))$ and $U^2 = U \cap ((\bigcup_{j=i+2}^{t} B_j) \setminus (B_{i+1} \cup B_t))$. Then, again since the size of each bag is bounded by κ , it holds that $|U^1|, |U^2| > \delta/2 - 3\kappa \ge k + 1$. Moreover, by the definition of a clique forest, $U^1 \cap U^2 = \emptyset$ and there is no vertex in U^1 that is adjacent to a vertex in U^2 . Thus, for any pair of vertices $x \in U^1$ and $y \in U^2$, the subgraph induced by $\{u, v, x, y\}$ is a chordless cycle. However, any solution of size at most k can only contain at most k vertices from $U^1 \cup U^2$, and thus, to hit all of these chordless cycles, it must contain at least one vertex among u and v.

Thus, from now on we can assume that $G[D_a]$ is a clique. However, by the definition of A, for every vertex in A and every vertex in V(Q), there exists a bag B_i , $i \in [t]$, which contains both of them, and therefore they are adjacent. We thus deduce that the following observation is correct.

Observation 34. Any two distinct vertices $v \in D_a \cup A$ and $u \in D_a \cup V(Q)$ are adjacent.

Let us now examine chordless cycles that contain vertices from U.

Lemma 35. Let C be a chordless cycle in G that contains some vertex $u \in U$. Then, no vertex on V(C) belongs to $D_a \cup A$, and both neighbors of u in C do not belong to $D' \cup A$.

Proof. First, by Observation 32, we have that u does not have neighbors (in G) in D_p , and therefore both neighbors of u in C do not belong to D_p . Thus, it remains to show that no vertex in V(C) belongs to $D_a \cup A$. By Observation 34, if at least one of vertex $v \in V(C)$ belongs to $D_a \cup A$, it is adjacent to u in G and it is either a neighbor of u in C adjacent in G to the other

neighbor of u in C or it is adjacent in G to both neighbors of u in C. In any case, if at least one of vertex $v \in V(C)$ belonged to $D_a \cup A$, the cycle C would have contained a chord, contradicting the supposition that C is a chordless cycle. We thus conclude that the lemma is correct. \Box

Lemma 36. Let C be a chordless cycle in G that contains some vertex $u \in U$. Then, C contains a path between a vertex in $B_1 \setminus A$ and a vertex in $B_t \setminus A$ whose internal vertices belong to U and one of them is u.

Proof. Since D' is an approximate solution, the cycle C must contain at least one vertex that does not belong to U. Thus, by Lemma 35, we deduce that the cycle C contains two subpaths, each between u and a vertex in $(B_1 \cup B_t) \setminus A$, whose only common vertex is u and whose internal vertices belong to U. Moreover, one of these paths must contain an endpoint from $B_1 \setminus A$ and the other from $B_t \setminus A$, else the cycle C contains a chord corresponding to the edge between these two endpoints. Therefore, by concatenating the two paths, we obtain the desired subpath of C.

We continue our examination of chordless cycles that contain vertices from U in the context of separators.

Lemma 37. Let S be a minimal solution that contains at least one vertex from U. Then, there exists $i \in [t-1]$ such that (i) $(B_i \cap B_{i+1}) \setminus A \subseteq S$, and (ii) $S \cap U \subseteq B_i \cap B_{i+1}$.

Proof. **Property** (i). Since S is a minimal solution, G contains a chordless cycle C with a vertex $u \in S \cap U$ and no other vertex from S. By Lemma 36, C contains a path between a vertex in $B_1 \setminus A$ and a vertex in $B_t \setminus A$ whose set of internal vertices includes u and is a subset of U. In particular, since each bag induces a clique while C is a chordless cycle, C contains a path P between a vertex $x \in B_1 \setminus A$ and a vertex in $y \in B_t \setminus A$ whose set of internal vertices is a non-empty subset of $V(Q) \setminus A$, and such that $(V(C) \setminus V(P)) \cap V(Q) = \emptyset$. Thus, x and y are not adjacent. Moreover, by Lemma 35, $V(C) \cap (D_a \cup A) = \emptyset$. Thus, since D' is a redundant approximate solution, C contains a vertex $d \in D_p$.

Suppose, by way of contradiction, that there does not exist $i \in [t-1]$ such that $(B_i \cap B_{i+1}) \setminus A \subseteq$ S. Then, G has a path P' between x and y whose internal vertices belong to $V(Q) \setminus (S \cup A)$. Since $(V(C) \setminus V(P)) \cap V(Q) = \emptyset$, it holds that $(V(C) \setminus V(P)) \cap V(P') = \emptyset$, which implies that C contains a path between x and y whose set of internal vertices is disjoint from the one of V(P'). Observe that if a graph H contains a vertex a with two non-adjacent neighbors, b and c, such that $H \setminus \{a\}$ has a chord less path between b and c with at least one vertex that is not a neighbor of a, then H has a chordless cycle. On the one hand, by the definition of a clique forest, any vertex in $V(G) \setminus (V(Q) \cup D')$ cannot be adjacent to both a vertex in $B_1 \setminus A$ and a vertex in $B_t \setminus A$, and it is adjacent to no vertex in U. On the other hand, Observation 32 implies that any vertex in D_p also satisfies this property. Thus, $V(P') \cap U = \emptyset$, since any vertex in this set fits the above description of the vertex a where $H = G[(V(C) \setminus V(P)) \cup V(P')]$, which is a subgraph of the chordal graph $G \setminus S$. Without loss of generality, we have that P' contains a subpath p - q - r where $p, q \in B_1 \setminus A$ and $r \in B_t \setminus A$. Let P'' denote a shortest path between pand r in $G[(V(C) \setminus V(P)) \cup (V(P') \setminus \{q\})]$, which contains at least two internal vertices (since it must contain a vertex from $V(C) \setminus V(P)$, else P' would have had a chord, and such a vertex cannot be adjacent to both p and r). Every vertex on P'' should be adjacent to q, else q fits the above description of the vertex a. Since P' is a chordless path and $(V(C) \setminus V(P)) \cap V(Q) = \emptyset$, P'' does not contain any vertex from B_1 that is not p, and the only vertex from B_t that is not r and which P'' can contain is the neighbor of r. Overall, this implies that P'' contains a vertex from $V(C) \setminus V(P)$ adjacent to both q and a vertex in $B_t \setminus A$, which is a contradiction.

Property (ii). It remains to show that $S \cap U \subseteq B_i \cap B_{i+1}$. To this end, we consider some arbitrary vertex $u \in S \cap U$ and show that it belongs to $B_i \cap B_{i+1}$. Since S is a minimal solution,

G contains a chordless cycle *C'* such that $V(C') \cap S = \{u\}$. By Lemma 36, *C'* contains a subpath between a vertex in $B_1 \setminus A$ and a vertex in $B_t \setminus A$ whose internal vertices belong to *U*. Thus, by the definition of a clique forest, *C'* must contain a vertex from $(B_i \cap B_{i+1}) \setminus A$. However, we have already shown that $(B_i \cap B_{i+1}) \setminus A \subseteq S$, and therefore it must hold that $u \in B_i \cap B_{i+1}$ (since otherwise we reach a contradiction to the fact that $V(C') \cap S = \{u\}$).

Let \mathcal{W} be the family of each subset $W \subseteq (B_1 \cup B_t) \setminus A$ of size at most k for which there exists an index $i \in [t-1]$ such that $W = (B_i \cap B_{i+1}) \setminus (A \cup U)$ and $|(B_i \cap B_{i+1}) \cap U| \leq k$. We can easily bound the size of the family \mathcal{W} as follows.

Lemma 38. $|W| \le 2k + 1$.

Proof. Let *i* be the smallest index in [t-1] for which there exists $W^i \in \mathcal{W}$ such that $W^i = (B_i \cap B_{i+1}) \setminus (A \cup U)$, and let *j* be the largest or which there exists $W^j \in \mathcal{W}$ such that $W^j = (B_i \cap B_{i+1}) \setminus (A \cup U)$. Then, by the definition of a clique forest, for every set $W \in \mathcal{W}$ it holds that $W \subseteq W^i \cup W^j$. Furthermore, the sets in \mathcal{W} can be sorted by $W_1, W_2, \ldots, W_{|\mathcal{W}|}$ such that for all $r \in [|\mathcal{W}| - 1]$, $W_{r+1} \cap B_1 \subseteq W_r \cap B_1$ and $W_r \cap B_t \subset W_{r+1} \cap B_t$. We thus conclude that $|\mathcal{W}| \leq 2k + 1$.

We proceed by associating a separator with each set $W \in \mathcal{W}$ as follows. First, let \mathcal{I}_W denote the set of all indices $i \in [t-1]$ such that $W = (B_i \cap B_{i+1}) \setminus (A \cup U)$. Now, let i_W denote an index in \mathcal{I}_W that minimizes $|(B_i \cap B_{i+1}) \cap U|$ (if there are several choices, choose one arbitrarily). We further denote $M = \bigcup_{W \in \mathcal{W}} ((B_{i_W} \cap B_{i_W+1}) \cap U)$. Observe that by Lemmata 25 and 38, $|M| = \mathcal{O}(k^2)$. Thus, it is sufficient to argue that there exists a vertex in $U \setminus M$ that can be removed from G (since as long as $|U| > \delta$, we will be able to find such a vertex). To this end, we will need the following lemma.

Lemma 39. Let $u \in U \setminus M$. If (G, k) is a yes-instance, then it has a solution S of size at most k that does not contain the vertex u.

Proof. Suppose that (G, k) is a yes-instance, and let S be a solution of minimum size. Assume that $u \in S$, else we are done. By Lemma 37, there exists $i \in [t-1]$ such that $(B_i \cap B_{i+1}) \setminus A \subseteq S$ and $S \cap U \subseteq B_i \cap B_{i+1}$. Denote $W = (B_i \cap B_{i+1}) \setminus (A \cup U)$. Since $|S| \leq k$ and $W \subseteq S$, we have that $W \in \mathcal{W}$. Denote $R = (B_{iW} \cap B_{iW+1}) \cap U$ and $T = (B_i \cap B_{i+1}) \cap U$. Since $u \in U \setminus M$, it holds that $u \notin R$. Moreover, since $S \cap U \subseteq B_i \cap B_{i+1}$, it holds that $u \in T$. By the definition of i_W , we have that $|R| \leq |T|$. Thus, to show that the lemma is correct, it is sufficient to show that $S' = (S \setminus T) \cup R$ is a solution.

Suppose, by way of contradiction, that S' is not a solution. Then, since S is a solution of minimum size, there exist a chordless cycle C and a vertex $v \in T$ such that $v \in V(C)$ and $V(C) \cap S' = \emptyset$. Since $T \subseteq U$, by Lemma 36, C contains a path between a vertex in $(B_1 \cap B_2) \setminus A$ and a vertex in $(B_{t-1} \cap B_t) \setminus A$ whose internal vertices belong to U. In particular, by the definition of a clique forest, $V(C) \cap ((B_{i_W} \cap B_{i_W+1}) \setminus A) \neq \emptyset$. However, we have that $(B_{i_W} \cap B_{i_W+1}) \setminus A = R \cup W \subseteq S'$, which contradicts the fact that $V(C) \cap S' = \emptyset$.

We are now ready to present our reduction rule.

Reduction Rule 6. Let $u \in U \setminus M$. Remove the vertex u from the graph G and add an edge between any two non-adjacent vertices in $N_G(u)$.

Lemma 40. Reduction Rule 6 is safe.

Proof. Let G' be the graph resulting from the application of this rule. For the forward direction, suppose that (G, k) is a yes-instance, and let S be a solution of minimum size. By Lemma 39, we can assume that $u \notin S$, and therefore $S \subseteq V(G')$. Thus, to show that (G', k) is a yes-instance,

we need to prove that S hits every chordless cycle in G'. Let C be a chordless cycle in G'. Suppose that this cycle does not exist in G, else it is clear that S hits it. Then, C contains an edge between two vertices $v, w \in N_G(u)$ that are non-adjacent in G. Observe that since C is a chordless cycle and $G'[N_G(u)]$ is a clique, $V(C) \cap N_G(u) = \{v, w\}$. Thus, by replacing $\{v, w\}$ by $\{v, u\}$ and $\{u, w\}$, we obtain a chordless cycle in G. Since S hits this cycle and $u \notin S$, it holds that S also hits C.

For the backward direction, suppose that (G', k) is a yes-instance, and let S be a solution of minimum size. To show that (G, k) is a yes-instance, it is sufficient to show that S is also a solution to (G, k). Since $V(G') \subseteq V(G)$, it is clear that $S \subseteq V(G)$. We need to prove that Shits every chordless cycle in G. Let C be a chordless cycle in G. Suppose that this cycle does not exist in G', else it is clear that S hits it. Furthermore, suppose that $G'[V(C) \setminus \{u\}]$ does not contain a chordless cycle, else again it is clear that S hits C. We get that C contains two vertices $v, w \in N_G(u)$ that are not adjacent in G. Since $u \in U$, Observation 32 and the definition of a clique forest imply that $N_G(u) \subseteq V(Q) \cup D_a$. By Observation 34, we deduce that $v, w \notin D_a \cup A$, and that C contains at most one vertex from $D_a \cup A$ (since any two vertices from $D_a \cup A$ are adjacent to each other and to both v and w). Since D' is a redundant approximate solution, C must contain a vertex $p \in D_p$. Since $G'[V(C) \setminus \{u\}]$ does not contain a chordless cycle, the neighbors of p on C belong to $N_G(u)$, and we can assume w.l.o.g that these neighbors are v and w. However, since v and w are not adjacent there cannot be a bag that contains both of them, which results in a contradiction to Observation 32.

3.6 Unmarking Irrelevant and Mandatory Edges

Recall that our instance includes a set E_I of irrelevant edges and a set E_M of mandatory edges. It is clear that we can unmark each irrelevant edge (these edges were marked only for the sake of clarity of the analysis of our kernel). However, to unmark mandatory edge, we need the following operation.

Reduction Rule 7. For every mandatory edge $\{x, y\}$ introduce k + 1 pairs of new vertices, $\{x_1, y_1\}, \{x_2, y_2\}, \ldots, \{x_{k+1}, y_{k+1}\}, and$ for each pair $\{x_i, y_i\}$ add the edges $\{x, x_i\}, \{x_i, y_i\}$ and $\{y_i, y\}$. Moreover, unmark the edge $\{x, y\}$.

Lemma 41. Reduction Rule 7 is safe.

Proof. Let $(G', k, E_I = \emptyset, E'_M)$ be the instance resulting from the application of this rule. Each edge $\{x, y\}$ is contained (in G') in k + 1 cycles of size 4 which do not share vertices other than x and y. Therefore, and solution of size at most k to (G', k) must contain at least one of the vertices x and y. Thus, since any chordless cycle in G is also present in G' and we have only unmarked the edge $\{x, y\}$, we conclude that if (G', k, E_I, E'_M) is a yes-instance, so is (G, k, E_I, E_M) . On the other hand, any solution to (G, k, E_I, E_M) hits all of the chordless cycles in G' since each of these chordless cycles is either present in G or contain both of the vertices x and y. Therefore, if (G, k, E_I, E_M) is a yes-instance, so is (G', k, E_I, E_M) .

Recall that $|E_M| \leq k^2$. Hence, the total number of newly added vertices does not exceed $\mathcal{O}(k^3)$.

3.7 The Number of Vertices in the Kernel

In this section, we obtained an approximate solution D of size f(k) and a redundant approximate solution D of size $\mathcal{O}(k \cdot f(k))$. Then, we examined the clique forest F associated with the chordal graph $G \setminus D'$ where $|D'| = \mathcal{O}(|D| + k^2)$. To this end, we considered a set \mathcal{P} of degree-2 paths that together cover all of the nodes of the forest, and showed that $|\mathcal{P}| = \mathcal{O}(|D'| \cdot \Delta')$. Recall that $\Delta' = \mathcal{O}(k \cdot f(k))$. We removed $\mathcal{O}(|D'|)$ bags, each of size $\kappa = \mathcal{O}(|\tilde{D}|^3 \cdot k + |\tilde{D}| \cdot \Delta' \cdot k^3)$, from each path $P \in \mathcal{P}$, and considered each of the resulting subpaths Q. We showed the number of vertices in the union of the bags of the path Q will be bounded by $\mathcal{O}(\kappa)$. Finally, we added $\mathcal{O}(k^3)$ new vertices to unmark mandatory edges. Thus, we conclude that the number of vertices in our kernel is bounded by

 $\begin{array}{ll} \mathcal{O}(|\mathcal{P}| \cdot |D'| \cdot \kappa) \\ = & \mathcal{O}(|D'|^2 \cdot \Delta' \cdot (|\widetilde{D}|^3 \cdot k + |\widetilde{D}| \cdot \Delta' \cdot k^3)) \\ = & \mathcal{O}(f(k)^3 k^3 \cdot (f(k)^3 k + f(k)^2 k^4)) \\ = & \mathcal{O}(f(k)^5 k^4 \cdot (f(k) + k^3)) \end{array}$

Recall that by Lemma 3, we can assume that $f(k) = \mathcal{O}(k^3 \log^2 k)$. Thus, at this point, we obtain a kernel of size $\mathcal{O}(k^{22} \log^{12} k)$.

3.8 A Better Kernelization Algorithm

Finally, we present a bootstrapping trick that will exploit the nature of our approximation algorithm to obtain a kernel of size $\mathcal{O}(k^{12}\log^{10}k)$. Recall that at this point, where we have already run our kernelization algorithm once, it holds that $n = \mathcal{O}(k^{22}\log^{12}k)$. Now, we again recall that CVD admits an $\mathcal{O}(\log^2 n)$ -factor approximation algorithm [3]. Currently, it holds that $f(k) = \mathcal{O}(k\log^2 k)$ rather than $f(k) = \mathcal{O}(k^3\log^2 k)$. Thus, if we rerun our kernelization procedure, obtaining a kernel of size $\mathcal{O}(f(k)^5k^4 \cdot (f(k) + k^3))$ (see Section 3.7), it now holds that this size is upper bounded by $\mathcal{O}(k^{12}\log^{10}k)$. This concludes the proof of correctness of Theorem 1.

4 Conclusion

In this paper we obtained a polynomial kernel for CVD of size $\mathcal{O}(k^{12} \log^{10} k)$. The new kernel significantly improves over the previously known $\mathcal{O}(k^{161} \log^{58} k)$ sized kernel. We believe that the notion of independence degree and the bootstrapping trick used in our kernelization procedure could be useful in designing polynomial kernels for other \mathcal{F} -VERTEX (EDGE) DELETION problems, where \mathcal{F} is characterized by an infinite set of forbidden induced graphs. We conclude the paper with the following open problems.

- Design a polynomial kernel for CVD of size $\mathcal{O}(k^c)$ for some fixed constant $c \leq 5$.
- Design a constant-factor approximation algorithm for CVD.
- Does there exist an FPT algorithm for CVD with running time $c^k n^{\mathcal{O}(1)}$, for some fixed constant c?

References

- F. N. ABU-KHZAM, A kernelization algorithm for d-Hitting Set, J. Comput. Syst. Sci., 76 (2010), pp. 524–531.
- [2] F. N. ABU-KHZAM, R. L. COLLINS, M. R. FELLOWS, M. A. LANGSTON, W. H. SUTERS, AND C. T. SYMONS, *Kernelization algorithms for the vertex cover problem: Theory and experiments*, in Proceedings of the Sixth Workshop on Algorithm Engineering and Experiments and the First Workshop on Analytic Algorithmics and Combinatorics, New Orleans, LA, USA, January 10, 2004, 2004, pp. 62–69.

- [3] A. AGRAWAL, D. LOKSHTANOV, P. MISRA, S. SAURABH, AND M. ZEHAVI, Polylogarithmic approximation algorithms for weighted-{-deletion problems., Manuscript available upon request, (2017).
- [4] N. ALON, G. GUTIN, E. J. KIM, S. SZEIDER, AND A. YEO, Solving MAX-r-SAT above a tight lower bound, Algorithmica, 61 (2011), pp. 638–655.
- [5] H. L. BODLAENDER, R. G. DOWNEY, M. R. FELLOWS, AND D. HERMELIN, On problems without polynomial kernels, J. Comput. Syst. Sci., 75 (2009), pp. 423–434.
- [6] H. L. BODLAENDER, F. V. FOMIN, D. LOKSHTANOV, E. PENNINKX, S. SAURABH, AND D. M. THILIKOS, (meta) kernelization, J. ACM, 63 (2016), pp. 44:1–44:69.
- [7] H. L. BODLAENDER, B. M. P. JANSEN, AND S. KRATSCH, Preprocessing for treewidth: A combinatorial analysis through kernelization, SIAM J. Discrete Math., 27 (2013), pp. 2108– 2142.
- [8] Y. CAO, Linear recognition of almost interval graphs, in Proceedings of the Twenty-Seventh Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2016, Arlington, VA, USA, January 10-12, 2016, 2016, pp. 1096–1115.
- [9] Y. CAO AND D. MARX, Interval deletion is fixed-parameter tractable, ACM Transactions on Algorithms, 11 (2015), pp. 21:1–21:35.
- [10] —, Chordal editing is fixed-parameter tractable, Algorithmica, 75 (2016), pp. 118–137.
- [11] J. CHEN, I. A. KANJ, AND W. JIA, Vertex cover: further observations and further improvements, Journal of Algorithms, 41 (2001), pp. 280–301.
- [12] M. CYGAN, F. V. FOMIN, Ł. KOWALIK, D. LOKSHTANOV, D. MARX, M. PILIPCZUK, M. PILIPCZUK, AND S. SAURABH, *Parameterized Algorithms*, Springer-Verlag, 2015.
- [13] M. CYGAN, L. KOWALIK, AND M. PILIPCZUK, Open problems from workshop on kernels, URL: http://worker2013.mimuw.edu.pl/slides/worker-opl.pdf, (2013).
- [14] H. DELL AND D. VAN MELKEBEEK, Satisfiability allows no nontrivial sparsification unless the polynomial-time hierarchy collapses, J. ACM, 61 (2014), p. 23.
- [15] R. G. DOWNEY AND M. R. FELLOWS, *Fundamentals of Parameterized Complexity*, Texts in Computer Science, Springer, 2013.
- [16] J. FLUM AND M. GROHE, Parameterized Complexity Theory, Texts in Theoretical Computer Science. An EATCS Series, Springer-Verlag, Berlin, 2006.
- [17] F. V. FOMIN, D. LOKSHTANOV, N. MISRA, AND S. SAURABH, Planar f-deletion: Approximation, kernelization and optimal FPT algorithms, in 53rd Annual IEEE Symposium on Foundations of Computer Science, FOCS 2012, New Brunswick, NJ, USA, October 20-23, 2012, 2012, pp. 470–479.
- [18] F. V. FOMIN, D. LOKSHTANOV, S. SAURABH, AND D. M. THILIKOS, *Bidimensionality and kernels*, in Proceedings of the Twenty-First Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2010, Austin, Texas, USA, January 17-19, 2010, 2010, pp. 503–510.
- [19] F. V. FOMIN, S. SAURABH, AND Y. VILLANGER, A polynomial kernel for proper interval vertex deletion, SIAM J. Discrete Math., 27 (2013), pp. 1964–1976.

- [20] L. FORTNOW AND R. SANTHANAM, Infeasibility of instance compression and succinct PCPs for NP, J. Comput. Syst. Sci., 77 (2011), pp. 91–106.
- [21] T. FUJITO, A unified approximation algorithm for node-deletion problems, Discrete Applied Mathematics, 86 (1998), pp. 213–231.
- [22] A. C. GIANNOPOULOU, B. M. P. JANSEN, D. LOKSHTANOV, AND S. SAURABH, Uniform kernelization complexity of hitting forbidden minors, in Automata, Languages, and Programming - 42nd International Colloquium, ICALP 2015, Kyoto, Japan, July 6-10, 2015, Proceedings, Part I, 2015, pp. 629–641.
- [23] M. C. GOLUMBIC, Algorithmic Graph Theory and Perfect Graphs, Academic Press, New York, 1980.
- [24] P. HEGGERNES, P. VAN 'T HOF, B. M. P. JANSEN, S. KRATSCH, AND Y. VILLANGER, Parameterized complexity of vertex deletion into perfect graph classes, Theor. Comput. Sci., 511 (2013), pp. 172–180.
- [25] B. M. P. JANSEN, Turing kernelization for finding long paths and cycles in restricted graph classes, in Proc. 22th ESA, A. S. Schulz and D. Wagner, eds., vol. 8737 of Lecture Notes in Computer Science, Springer, 2014, pp. 579–591.
- [26] B. M. P. JANSEN AND M. PILIPCZUK, Approximation and kernelization for chordal vertex deletion, in Proceedings of the Twenty-Eight Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2017, 2017, p. to appear.
- [27] Y. KE, Y. CAO, X. OUYANG, AND J. WANG, Unit interval vertex deletion: Fewer vertices are relevant, arXiv preprint arXiv:1607.01162, (2016).
- [28] E. J. KIM, A. LANGER, C. PAUL, F. REIDL, P. ROSSMANITH, I. SAU, AND S. SIKDAR, Linear kernels and single-exponential algorithms via protrusion decompositions, ACM Trans. Algorithms, 12 (2016), p. 21.
- [29] S. KRATSCH, Polynomial kernelizations for MIN $F^+\Pi_1$ and MAX NP, Algorithmica, 63 (2012), pp. 532–550.
- [30] —, Recent developments in kernelization: A survey, Bulletin of the EATCS, 113 (2014).
- [31] S. KRATSCH AND M. WAHLSTRÖM, Representative sets and irrelevant vertices: New tools for kernelization, in 53rd Annual IEEE Symposium on Foundations of Computer Science, FOCS 2012, New Brunswick, NJ, USA, October 20-23, 2012, 2012, pp. 450–459.
- [32] —, Compression via matroids: A randomized polynomial kernel for odd cycle transversal, ACM Trans. Algorithms, 10 (2014), pp. 20:1–20:15.
- [33] J. M. LEWIS AND M. YANNAKAKIS, The node-deletion problem for hereditary properties is np-complete, J. Comput. Syst. Sci., 20 (1980), pp. 219–230.
- [34] D. LOKSHTANOV, N. MISRA, AND S. SAURABH, Kernelization preprocessing with a guarantee, in The Multivariate Algorithmic Revolution and Beyond - Essays Dedicated to Michael R. Fellows on the Occasion of His 60th Birthday, 2012, pp. 129–161.
- [35] C. LUND AND M. YANNAKAKIS, On the hardness of approximating minimization problems, J. ACM, 41 (1994), pp. 960–981.

- [36] D. MARX, Chordal deletion is fixed-parameter tractable, Algorithmica, 57 (2010), pp. 747– 768.
- [37] D. MARX, B. O'SULLIVAN, AND I. RAZGON, Finding small separators in linear time via treewidth reduction, ACM Transactions on Algorithms, 9 (2013), p. 30.
- [38] R. NIEDERMEIER, *Invitation to Fixed-Parameter Algorithms*, vol. 31 of Oxford Lecture Series in Mathematics and its Applications, Oxford University Press, Oxford, 2006.
- [39] M. PILIPCZUK, M. PILIPCZUK, P. SANKOWSKI, AND E. J. VAN LEEUWEN, *Network* sparsification for steiner problems on planar and bounded-genus graphs, in Proc. 55th FOCS, IEEE Computer Society, 2014, pp. 276–285.
- [40] S. THOMASSÉ, A 4k² kernel for feedback vertex set, ACM Trans. Algorithms, 6 (2010), pp. 32:1–32:8.
- [41] M. YANNAKAKIS, Node- and edge-deletion np-complete problems, in Proceedings of the 10th Annual ACM Symposium on Theory of Computing, May 1-3, 1978, San Diego, California, USA, 1978, pp. 253–264.

A Bounding the Size of a Maximal Clique

Let (G, k) be an instance of CVD whose independent degree is bounded Δ , and let D be some (possibly approximate) solution to this instance. In this appendix, we will bound the size of a maximal clique in the graph G. We will show that if the graph admits a chordal deletion set of size at most k, then for any maximal clique K in $G \setminus D$, it is safe to remove all but a bounded number vertices of K. Formally, we will show the following lemma.

Lemma 42. Let (G, k) be an instance of CVD whose independent degree is bounded Δ , and let D be a solution to this instance. Then in polynomial time we can produce an equivalent instance (G', k') such that $k' \leq k$, and the size of any maximal clique in G' is bounded by $c \cdot (|D|^3 \cdot k + |D| \cdot \Delta \cdot (k+2)^3)$. (Here c is some constant independent of the input.)

Our proof of this lemma is an adaptation of the work of Marx [36] with a few modifications. Specifically, the lemmas in [36] construct a so called "necessary set" of vertices with the property that one of the vertices in this set must be part of any solution of size k. We modify these lemmas to ensure that a necessary set output by them always has at most two vertices. Observe that such a necessary set is either a vertex that must be part of any solution of size at most k, or a mandatory edge. We note that Jansen and Pilipczuk [26] also give similar result, inspired by the results of [36]. Given a redundant approximate solution \hat{D} , the size of a maximal clique in $G \setminus \hat{D}$ can be bounded by $\mathcal{O}(|\hat{D}|^3k)$. However the present method for computing a redundant approximate solution implies that $|\hat{D}| \geq |D| \cdot k$, and hence Lemma 42 gives a better upper-bound.

Lemma 43 ([26]). The size of each bag in the clique forest F is bounded by $c \cdot |D'|^3 k$, where c is some constant independent of the input.

For the rest of this subsection, we fix a maximal clique K of $G \setminus D$, which contains more than $c \cdot (|D|^3 \cdot k + |D| \cdot \Delta \cdot (k+2)^3)$ vertices. We will show that we can mark a bounded number of vertices of K so that the following holds. Let X be any set of at most k vertices such that $G \setminus X$ has a chordless cycle H that contains a vertex u of K. If u is an unmarked vertex then there is another chordless cycle H' in $G \setminus X$ that avoids u and contains strictly fewer unmarked vertex in K. This condition implies that we can safely ignore any chordless cycle that includes an unmarked vertex, which further implies that it is safe to delete these vertices from the graph. We shall closely follow the notations and proofs of [36], but in light of the bound on (relevant) independent-degree of vertices in D, we shall modify them appropriately. For a vertex $v \in V(G) \setminus D$, we say that the vertex has the *label* $t \in D$ if v is a neighbor of t. Note that the edge (t, v) could be relevant or irrelevant, and a vertex may have several labels depending on its set of neighbors in D.

A.1 Dangerous vertices and their witnesses in K

Let us begin with the notion of dangerous vertices for the clique K in the graph.

Definition 44. Let $v \in V(G) \setminus (D \cup K)$ be a relevant neighbor of $t \in D$ such that there is a path P from v to $u \in K$ whose internal vertices do not have the label t.

- (i) The vertex v is a t-dangerous vertex for K if the vertex u does not have the label t.
- (ii) The vertex v is a t^{*}-dangerous vertex for K if the vertex u is also a relevant neighbor of t and u, v are not neighbors in G.

The vertex $u \in K$ is called a t-witness (t*-witness) of v, and the path P is called a t-witness (t*-witness) path of v.

In [36], it is shown that there are many vertices in the clique K whose deletion does not affect any of the dangerous vertices in the following sense. We mark a bounded number of vertices in Ksuch that, for any subset X of vertices of size at most k, if there is chordless cycle in $G \setminus X$ that passes through t, a dangerous vertex v and through some unmarked vertex u in K, then there is another chordless cycle in $G \setminus X$ which contains a marked vertex u' and avoids u. This implies that we may ignore any chordless cycles in G that includes an unmarked vertex of K. Note that the definition above differs from [36] in the requirement that v (and u) must be a relevant neighbor of t, since if the edge (t, v) (or the edge (t, u)) were irrelevant then any chordless cycles that contain both t and v (or t and u) can be safely ignored. We have the following bounds on the size of an independent set of dangerous vertices in G.

Lemma 45 (Lemma 11, [36]). If I is any collection of independent t-dangerous vertices, then either $|I| \leq 6k^2$ or we can find a new mandatory edge in the graph, or we can find a vertex that must be part of any solution of size at most k.

The following lemma improves upon Lemma 12 of [36] by using the bound on the independent degree of vertices in D.

Lemma 46 (Lemma 12, [36]). If I is any collection of independent t^* -dangerous vertices, then $|I| \leq \Delta$.

The following lemma shows that if Q is a clique of t-dangerous vertices we require only k + 1 vertices as witnesses for all the vertices of Q.

Lemma 47 (Lemma 13 [36]). Let Q be a clique of t-dangerous vertices. Then we can mark k+1 vertices in K such that for any set X of k vertices, if $v \in Q$ has an unmarked t-witness in $K \setminus X$ then it has a marked t-witness in $K \setminus X$.

In the context of the following lemma, we remark that there is a similar lemma for cliques of t^* -dangerous vertices in [36], but we give a version of it that is more suitable for our purposes.

Lemma 48 (Lemma 14, [36]). Let Q be a clique of t^* dangerous vertices. Then either we can find a vertex that must be part of any solution of size at most k, or we can find a new mandatory edge, or else we can mark $(k + 2)^3$ vertices of K such that for any set X of k vertices, if $v \in Q$ has a unmarked t^* witness in $K \setminus X$ then it also has a marked witness $K \setminus X$. Proof. Our proof is a minor modification of the proof of Lemma 14, [36]. Let T be a clique-tree of $G \setminus D$. Following [36], we say that a vertex v covers a bag x of T if v is contained in the bag x, and then T_v denotes the sub-clique-tree of all the bags which are covered by v. Now, since Qand K are cliques in $G \setminus D$, there are two bags x and y which contain Q and K, respectively. Consider the unique path connecting x and y in T, and suppose that the bags on this path are numbered as $x = 1, 2, \ldots s = y$. Let u_1, u_2, \ldots be vertices of K with label t, and let a_i denote the smallest numbered bag on this path that occurs in T_{u_i} , i.e. the smallest numbered bag containing u_i . Similarly, let v_1, v_2, \ldots be the vertices of Q and let b_i denote the largest numbered bag which is contained in T_{v_i} . It follows that T_{v_i} and T_{u_j} intersect if and only if $a_j \leq b_i$, in which case there is an edge (v_i, u_j) in the graph. We also assume that the collection of a_i and b_i are distinct, which is easily achieved by adding additional bags along this path in T. Further we assume that the vertices u_1, u_2, \ldots and v_1, v_2, \ldots are ordered so that the sequences of a_i and b_j are strictly increasing.

We define a subsequence of b_i and a_j as follows. Let $\beta_1 = 1$, and for ever $j \ge 1$ let α_j be the smallest value such that $a_{\alpha_j} > b_{\beta_j}$. For every $i \ge 2$, let β_i be the smallest value such that $b_{\beta_i} > a_{\alpha_{i-1}}$. Observe that we obtain a strictly increasing sequence, $b_{\beta_1} < a_{\alpha_1} < b_{\beta_2} < a_{\alpha_2} \dots$ Let β_ℓ be the last element of the above sequence which corresponds to a vertex in Q.

Now, let u_s be a witness of a t^* -dangerous vertex v_{β_j} . We will show that a_{α_j} is also a witness for v_{β_j} . Clearly, $a_s > b_{\beta_j}$, which implies $a_s \ge a_{\alpha_j}$. Hence, the t^* -witness path from v_{β_j} to u_s passes through the bag a_{α_j} which contains u_{α_j} (see the proof of Lemma 14, [36]). Further, $a_{\alpha_j} > b_{\beta_j}$ implies that v_{β_j} and a_{α_j} are not neighbors in G. Therefore, a_{α_j} is also a witness for v_{β_j} .

Now, suppose that $\ell \leq (k+2)^2$. Then for each $i = 1, 2, \ldots, \ell$ we mark the k+2 vertices $u_{\alpha_i}, u_{\alpha_i+1}, \ldots, u_{\alpha_i+k+1}$ (if they exist), and we will show that this set of marked vertices contains a sufficient number of witnesses for every vertex in Q. Note that we have marked at most $(k+2)^3$ vertices of K. Consider any set X of k vertices such that a vertex $v_x \in Q$ has a witness path in $G \setminus X$ to some unmarked $u_y \in K$. In other words there is a chordless cycle in $G \setminus X$ which includes the vertex t, a t^* -dangerous vertex v_x and its t^* -witness $u_y \in K$. Since v_x and u_y are non-neighbors, we have that $b_x < a_y$, which implies that there is some j for which $b_x < a_{\alpha_j} \leq a_y$. If a_y is not marked then $y > \alpha_j + k + 1$, and hence $G \setminus X$ contains some $b_x < a_{\alpha_j+r} < a_y$. Now the witness path from v_x to a_y must contain a neighbor of a_{α_j+r} , which implies that a_{α_j+r} is also a witness for v_x , in $G \setminus X$.

Finally, suppose that $\ell > (k+2)^2$. We will show that either the vertex t must be part of any solution of size k, or we can find a new mandatory edge in the graph. Let P_i be a witness path from v_{β_i} to u_{α_i} for every $1 \le i \le \ell$. Consider the collection of chordless cycles $H_{(k+2)\cdot j}$ for $1 \le j \le k+1$, where $H_{(k+2)\cdot j} = (t v_{\beta_{(k+2)\cdot j}} P_{(k+2)\cdot j} u_{\alpha_{(k+2)\cdot j}} t)$. If all these cycles are pairwise vertex disjoint except for the vertex t, then we have obtained a collection of k+1 chordless cycles with t as the only common vertex. Hence t must be part of any solution of size at most k.

Otherwise, let $H_{(k+2)\cdot j}$ and $H_{(k+2)\cdot j'}$ have a common vertex w, for some j < j'. Then observe that w can only be an internal vertex of the paths $P_{(k+2)\cdot j}$ and $P_{(k+2)\cdot j'}$, and therefore it is not a neighbor of t. Since T is a tree-decomposition, this implies that w occurs in every bag numbered between $a_{\alpha_{(k+2)\cdot j}}$ and $b_{\beta_{(k+2)\cdot j'}}$. Since $a_{\alpha_{(k+2)\cdot j}} < b_{\beta_{(k+2)\cdot j+1}} < a_{\alpha_{(k+2)\cdot j+1}} < \dots < b_{\beta_{(k+2)\cdot j+k+1}} < a_{\alpha_{(k+2)\cdot j+k+1}} < b_{\beta_{(k+2)\cdot j'}}$, we have that w occurs in every one of these bags. Observe that we obtain a collection of k + 1 chordless cycles of length 4, namely $(t v_{\beta_{(k+2)\cdot j+i}} w u_{\alpha_{(k+2)\cdot j+i}} t)$ for $i = 1, 2, \dots, k+1$, such that the vertices t and w are the only common vertices. Since t and ware non-adjacent in G, we obtain a new mandatory edge (t, w) in G.

Since $G \setminus D$ is a chordal graph which contains all the dangerous vertices for K, it follows that the graph induced by the set of all *t*-dangerous (t^* -dangerous) vertices forms a chordal graph as well. Since a chordal graph is perfect, it has a clique cover of size α , where α is the cardinality of a maximum independent set in the graph (see [23]). This gives us the following lemma, using the bound on the size of an independent set of dangerous vertices.

- **Lemma 49** (Lemmas 15 & 16, [36]). (i) Either we can find a vertex that must be part of any solution of size at most k, or we can find a new mandatory edge, or we can mark $6k^2(k+1)$ vertices in K such that for any set X of k vertices, if a t-dangerous vertex v has a unmarked witness in $K \setminus X$, then it also has a marked witness in $K \setminus X$.
 - (ii) Either we can find a vertex that must be part of any solution of size at most k, or we can find a new mandatory edge, or we can mark $\Delta \cdot (k+2)^3$ vertices in K such that for any set X of k vertices, if a t*-dangerous vertex has a unmarked witness in $K \setminus X$, then it also has a marked witness in $K \setminus X$.

From the above lemma we conclude that, for any set X of at most k vertices, to test if X intersects all those chordless cycles that pass through t, K and some dangerous vertex v, it is sufficient to consider only the marked vertices of K as witnesses. However the above lemma considers only a single vertex $t \in D$, and we must mark additional witness vertices for each $t \in D$.

- **Lemma 50** (Lemmas 15 & 16, [36]). (i) Either we can find a vertex that must be part of any solution of size at most k, or we can find a new mandatory edge, or we can mark $|D| \cdot 6k^2(k+1)$ vertices in K such that for any set X of k vertices, if a t-dangerous vertex v has a unmarked witness in $K \setminus X$, then it also has a marked witness in $K \setminus X$.
- (ii) Either we can find a vertex that must be part of any solution of size at most k, or we can find a new mandatory edge, or we can mark $|D| \cdot \Delta \cdot (k+2)^3$ vertices in K such that for any set X of k vertices, if a t^{*}-dangerous vertex has a unmarked witness in $K \setminus X$, then it also has a marked witness in $K \setminus X$.

A.2 Fragments of Chordless cycles intersecting K

Now we shall mark vertices for chordless cycles in G that intersect K. If H is a chordless cycle in G, then consider F, P_1, P_2, \ldots, P_s where $F = H \cap D$ and $P_1, P_2 \ldots, P_s$ are the paths in $H \setminus D$. It follows that each P_i has exactly two labels from F on its endpoints, and the internal vertices have no labels from F, and further these paths are pairwise independent (i.e. there is no edge between two vertices that are in two different paths). We call F, P_1, \ldots, P_s the fragments of the chordless cycle H. Since K is a clique, at most one of these paths intersects K, which we assume to be the path P_1 , and further it contains at most two vertices from K. We will show that we can mark a bounded number of vertices in K such that for any chordless cycle H that includes an unmarked vertex (that lies in P_1), there is another chordless cycle H' that avoids this unmarked vertex. Let us first consider the case when P_1 just a single vertex (in K). The following lemma is a close variant of Lemma 18, [36] for this case.

Lemma 51 (Lemma 18, [36]). Let F, P_1, \ldots, P_s be the fragments of H where P_1 is just a single vertex that lies in K. Then either we can find a new mandatory edge, or we can mark $|D|^3 \cdot (k+2)$ vertices in K such that the following holds. For any set X of k vertices such that $G \setminus X$ has a chordless cycle which intersects K in an unmarked vertex, there is another chordless cycle in $G \setminus X$ which does not use any unmarked vertices of K.

Proof. Our proof of this lemma is obtained by modifying proof of Lemma 18, [36]. And note that, as P_1 is a single vertex, F must have at least two vertices. For every $l_1, l_2, l_3 \in D$, mark k + 1 vertices of K which have labels l_1, l_2 and not l_3 . Hence, in total we have marked $|D|^3 \cdot (k+1)$ vertices of K. Marx [36] shows that this is sufficient for the case when F has 3 or more vertices.

When F contains only two vertices l_1, l_2 , we need to mark some additional vertices. Let us recall that P_1 is only a single vertex of K and has the labels l_1, l_2 , and so it follows that l_1, l_2 are non-neighbors. Let x be the bag in the clique-tree of $G \setminus D$, which corresponds to the maximal clique K, and further assume that x is the root of this tree-decomposition. For any chordless cycle H^i such that $H^i \setminus D = P_1^i, P_2^i$ where P_1^i is a single vertex, let w_i be the bag closest to x in the tree decomposition that contains a vertex of P_2^i . Note that $w_i \neq x$, as vertices of P_1^i and P_2^i have no edges between them, and it follows that the vertices of P_2^i are contained in the sub-clique-tree rooted at w_i , and furthermore none of them are present in any bag outside this sub-clique-tree. We can generate a list of these bags w_1, w_2, \ldots , by considering every choice of P_1^i , l_1 and l_2 and selecting those bags of the tree decomposition that have a path from l_1 to l_2 with at least one internal vertex in the non-neighborhood of the choice of P_1^i . Among these bags, select $w_{i_1}, w_{i_2}, \ldots, w_{i_q}$ such that none of its descendants in the clique-tree are in the collection w_1, w_2, \ldots computed above. Note that, by definition no bag selected above is an ancestor or a descendant of another selected bag. Let $H_{i_1}, H_{i_2}, \ldots, H_{i_q}$ be a collection of chordless cycles such that $P_2^{i_j}$ is contained in the sub-clique-tree of w_{i_j} , for $j = 1, 2, \ldots, q$. Note that the collection of $P_2^{i_j}$ are pairwise vertex disjoint, and further there are no edges between the vertices of $P_2^{i_j}$ and $P_2^{i_{j'}}$ for $j \neq j'$. This follows from the fact that no bag outside the sub-clique-tree rooted at w_{i_j} contains any vertex of $P_2^{i_j}$, and that no bag selected in the collection of w_{i_j} is an ancestor or a descendant of another.

Consider the case when $q \leq k + 1$. We define the distance of a vertex v from a bag w in the clique-tree as the minimum of the distance between w and a bag x that contains v. Then for each w_{i_j} , sort the vertices of of K, which have labels l_1, l_2 , according to the distance from w_{i_j} , and mark k + 1 vertices from w_{i_j} . It follows that we mark $|D|^2 \cdot (k+1)^2$ vertices of K. Let us argue that these marked vertices satisfy the requirements. For any set X of at most k vertices, suppose H is a chordless cycle in $G \setminus X$ with fragments F, P_1, P_2 where P_1 is just a single vertex $u \in K$ with labels $l_1, l_2 \in F$. Consider some bag w_i whose sub-clique-tree contains the path P_2 , and note that w_i , or some descendent w_{i_j} was selected in the above collection. Now if u was not marked for w_{i_j}, l_1, l_2 , then any of the vertices that were marked for this tuple has a greater distance from the bags w_{i_j} (and w_i) than u. Since we marked k + 1 vertices for this tuple, at least one of them is present in $K \setminus X$ and let u' be that vertex. It follows that replacing u with u' in H gives us a chordless cycle in $G \setminus X$.

Now, consider the case when $q \ge k+2$. The collection of paths P^{i_j} , along with l_1 and l_2 , defines a collection of chordless cycles such that any solution of size k must pick at least one of l_1 or l_2 . Since l_1, l_2 are non-neighbors, we obtain a new mandatory edge (l_1, l_2) .

Further note that the total number of marked vertices is $|D|^3 \cdot (k+1) + |D|^2 \cdot (k+1)^2$. Since $|D| \ge k$, it follows that the total number of marked vertices is upper bounded by $|D|^3 \cdot (k+2)$. \Box

Now, the only remaining case is when the path P_1 has two or more vertices. For this case, we have the following lemma, which follows from Lemma 19 of [36] where we rely on Lemma 50.

Lemma 52 (Lemma 19, [36]). Let F, P_1, \ldots, P_s be the fragments of a chordless cycle H where P_1 contains at least two vertices, and further at least one of those vertices are in K. Then we can mark at most $|D| \cdot 6k^2(k+1) + |D| \cdot \Delta \cdot (k+2)^3$ vertices in K such that, for any set X of at most k vertices, if $G \setminus X$ contains a chordless cycle that includes an unmarked vertex $u \in K$, then $G \setminus X$ also contains a chordless cycle that avoids u, and it has strictly fewer unmarked vertices of K.

Combining all of the above lemmas we obtain the following.

Lemma 53. Let (G, k) be an instance of CVD whose independent-degree is bounded by Δ . Let D be solution to this instance and let K be a maximal clique in $G \setminus D$. Then in polynomial time,

either we can find a vertex which must be part of any solution of size at most k, or we can find a new mandatory edge, or we can safely remove all but $c \cdot (|D|^3 \cdot k + |D| \cdot \Delta \cdot (k+2)^3)$ vertices of K. (Here c is some constant independent of the input.)

Proof. We apply the Lemmas 50, 51 and 52 to the given instance. If any of them return a vertex that must be part of any solution of size at most k, or a new mandatory edge then we output that vertex or edge. Otherwise, together they mark a maximum of $c' \cdot (|D|^3 \cdot k + |D| \cdot \Delta \cdot (k+2)^3)$ vertices in K, where c' is some constant independent of the input. In addition, let us also mark any unmarked vertex in K which is an endpoint of a mandatory edge in the instance. Since the total number of mandatory edges is always bounded by k^2 , the total number of marked vertices in K does not exceed $c \cdot (|D|^3 \cdot k + |D| \cdot \Delta \cdot (k+2)^3)$, where c is again a constant independent of the input.

Let u be any unmarked vertex in K. We will show that the instances G and $G \setminus u$ are equivalent. Let X be any set of at most k vertices such that $(G \setminus u) \setminus X$ is a chordal graph. If $G \setminus X$ is not chordal, then there is some chordless cycle H in $G \setminus X$. If H does not contain the vertex u, then it is also present in $(G \setminus u) \setminus X$ which is a contradiction. On the other hand if H contains u, then by the above lemmas, we can argue that there is some chordless cycle H' in $G \setminus X$ which avoids u (see Lemma 19, [36]). Therefore H' is present in $(G \setminus u) \setminus X$, which is again a contradiction. Now, by induction, we can show that it is safe to remove all the unmarked vertices in K from the graph.

Now we are ready to prove Lemma 42. Consider any maximal clique in the graph $G \setminus D$ with more than $c \cdot (|D|^3 \cdot k + |D| \cdot \Delta \cdot (k+2)^3)$ vertices, and apply Lemma 53 to it. If it returns a vertex v that must be part of any solution of size k, we remove it from the graph and decrease k by one. Else if it returns a mandatory edge, we add this edge to the graph and mark it as mandatory. We can argue, as before, that both these operations are safe. Otherwise Lemma 53 ensures that it is safe remove all but the $c \cdot (|D|^3 \cdot k + |D| \cdot \Delta \cdot (k+2)^3)$ marked vertices of K. Therefore, we remove these vertices from the graph. Observe that, each application of Lemma 53 either reduces k, or adds a new mandatory edge, or reduces the number of vertices in the graph. Further, we may add at most $k^2 + 1$ new mandatory edges before finding a new vertex that must be part of any solution of size k or concluding that the given instance is a No instance. Hence, in polynomial time, either we bound the size of every maximal clique in the graph or conclude that the given instance is a No instance.