Flexible Bayesian Dynamic Modeling of Correlation and Covariance Matrices

Shiwei Lan*¹, Andrew Holbrook², Gabriel A. Elias³, Norbert J. Fortin⁴, Hernando Ombao⁵, and Babak Shahbaba⁶

 $^{1}\mathrm{Department}$ of Statistics, University of Illinois Urbana-Champaign, Champaign, IL 61820

^{2,6}Department of Statistics, University of California-Irvine, Irvine, CA 92697
 ^{3,4}Center for the Neurobiology of Learning and Memory, Department of Neurobiology and Behavior, University of California-Irvine, Irvine, CA 92697
 ⁵Statistics Program, King Abdullah University of Science and Technology, Thuwal 23955, Saudi Arabia

Abstract

Modeling correlation (and covariance) matrices can be challenging due to the positive-definiteness constraint and potential high-dimensionality. Our approach is to decompose the covariance matrix into the correlation and variance matrices and propose a novel Bayesian framework based on modeling the correlations as products of unit vectors. By specifying a wide range of distributions on a sphere (e.g. the squared-Dirichlet distribution), the proposed approach induces flexible prior distributions for covariance matrices (that go beyond the commonly used inverse-Wishart prior). For modeling real-life spatio-temporal processes with complex dependence structures, we extend our method to dynamic cases and introduce unit-vector Gaussian process priors in order to capture the evolution of correlation among components of a multivariate time series. To handle the intractability of the resulting posterior, we introduce the adaptive Δ -Spherical Hamiltonian Monte Carlo. We demonstrate the validity and flexibility of our proposed framework in a simulation study of periodic processes and an analysis of rat's local field potential activity in a complex sequence memory task.

Keywords: Dynamic covariance modeling; Spatio-temporal models; Geometric methods; Posterior contraction; Δ -Spherical Hamiltonian Monte Carlo

^{*}shiwei@illinois.edu

1 Introduction

Modeling covariance matrices—or more broadly, positive definite (PD) matrices—is one of the most fundamental problems in statistics. In general, the task is difficult because the number of parameters grows quadratically with the dimension of the matrices. The complexity of the challenge increases substantially if we allow dependencies to vary over time (or space) in order to account for the dynamic (non-stationary) nature of the underlying probability model. In this paper, we propose a novel solution to the problem by developing a flexible and yet computationally efficient Bayesian inferential framework for both static and dynamic covariance matrices.

This work is motivated by modeling the dynamic brain connectivity (i.e., associations between brain activity at different regions). In light of recent technical advances that allow the collection of large, multidimensional neural activity datasets, brain connectivity analyses are emerging as critical tools in neuroscience research. Specifically, the development of such analytical tools will help elucidate fundamental mechanisms underlying cognitive processes such as learning and memory, and identify potential biomarkers for early detection of neurological disorders. There are a number of new methods that have been developed (Cribben et al., 2012; Fiecas and Ombao, 2016; Lindquist et al., 2014; Ting et al., 2015; Prado, 2013) but the main limitation of these methods (especially the ones that have a frequentist approach) is a lack of natural framework for inference. Moreover, parametric approaches (e.g. vector auto-regressive models) need to be tested for adequacy for modeling complex brain processes and often have high dimensional parameter spaces (especially with a large number of channels and high lag order). This work provides both a nonparametric Bayesian model and an efficient inferential method for modeling the complex dynamic dependence among multiple stochastic processes that is common in the study of brain connectivity.

Within the Bayesian framework, it is common to use an inverse-Wishart prior on the covariance matrix for computational convenience (Mardia et al., 1980; Anderson, 2003). This choice of prior however is very restrictive (e.g. common degrees of freedom for all components of variance) (Barnard et al., 2000; Tokuda et al., 2011). Daniels (1999); Daniels and Kass (2001) propose uniform shrinkage priors. Daniels and Kass (1999) discuss three hierarchical priors to generalize the inverse-Wishart prior. Alternatively, one may use decomposition strategies for more flexible modeling choices (see Barnard et al. (2000) for more details). For instance, Banfield and Raftery (1993), Yang and Berger (1994), Celeux and Govaert (1995), Leonard and Hsu (1992), Chiu et al. (1996), and Bensmail et al. (1997) propose methods based on the spectral decomposition of the covariance matrix. Another strategy is to use the Cholesky decomposition of the covariance matrix or its inverse, e.g., Pourahmadi (1999, 2000); Liu (1993); Pinheiro and Bates (1996). There are other approaches directly related to correlation, including the constrained model based on truncated distributions (Liechty, 2004), the Cholesky decomposition of correlation matrix using an angular parametrization (Pourahmadi and Wang, 2015), and methods based on

partial autocorrelation and parameterizations using angles (Rapisarda et al., 2007). In general, these methods fail to yield full flexibility and generality; and often sacrifice statistical interpretability.

While our proposed method in this paper is also based on the separation strategy (Barnard et al., 2000) and the Cholesky decomposition, the main distinction from the existing methods is that it represents each entry of the correlation matrix as a product of unit vectors. This in turn provides a flexible framework for modeling covariance matrices without sacrificing interpretability. Additionally, this framework can be easily extended to dynamic settings in order to model real-life spatio-temporal processes with complex dependence structures that evolve over the course of the experiment.

To address the constraint for correlation processes (positive definite matrix at each time having unit diagonals and off-diagonal entries with magnitudes no greater than 1), we introduce unit-vector Gaussian process priors. There are other related works, e.g. generalized Wishart process (Wilson and Ghahramani, 2011), and latent factor process (Fox and Dunson, 2015), that explore the product of vector Gaussian processes. In general they do not grant full flexibility in simultaneously modeling the mean, variance and correlation processes. For example, latent factor based models link the mean and covariance processes through a loading matrix, which is restrictive and undesirable if the linear link is not appropriate, and thus are outperformed by our proposed flexible framework (See more details in Section 4.2.2). Other approaches to model non-stationary processes use a representation in terms of a basis such as wavelets (Nason et al., 2000; Park et al., 2014; Cho and Fryzlewicz, 2015) and the SLEX (Ombao et al., 2005), which are actually inspired by the Fourier representations in the Dahlhaus locally stationary processes Dahlhaus (2000); Priestley (1965). These approaches are frequentist and do not easily provide a framework for inference (e.g., obtaining confidence intervals). The class of time-domain parametric models allows for the ARMA parameters to evolve over time (see, e.g. Rao, 1970) or via parametric latent signals (West et al., 1999; Prado et al., 2001). A restriction for this class of parametric models is that some processes might not be adequately modeled by them.

This main contributions of this paper are: (a.) a sphere-product representation of correlation/covariance matrix is introduced to induce flexible priors for correlation/covariance matrices and processes; (b.) a general and flexible framework is proposed for modeling mean, variance, and correlation processes separately; (c.) an efficient algorithm is introduced to infer correlation matrices and processes; (d.) the posterior contraction of modeling covariance (correlation) functions with Gaussian process prior is studied for the first time.

The rest of the paper is organized as follows. In the next section, we present a geometric view of covariance matrices and extend this view to allow covariance matrices to change over time. In Section 3, we use this geometrical perspective to develop an effective and computationally efficient inferential method for modeling static and dynamic covariance matrices. Using simulated data, we will evaluate our method in Section 4. In Section 5, we apply our proposed method to local field potential (LFP) activity data recorded from the hippocampus of rats performing a complex sequence memory task (Allen et al., 2014,

2016; Ng et al., 2017). In the final section, we conclude with discussions on the limitations of the current work and future extensions.

2 Structured Bayesian Modeling of the Covariance (Correlation) Matrices

To derive flexible models for covariance and correlation matrices, we start with the Cholesky decomposition, form a sphere-product representation, and finally obtain the separation decomposition in Barnard et al. (2000) with correlations represented as products of unit vectors. The sphere-product representation is amenable for the inferential algorithm to handle the resulting intractability, and hence lays the foundation for full flexibility in choosing priors.

Any covariance matrix $\Sigma = [\sigma_{ij}] > 0$ is symmetric positive definite, and hence has a unique Cholesky decomposition $\Sigma = \mathbf{L}\mathbf{L}^{\mathsf{T}}$ where the Cholesky factor $\mathbf{L} = [l_{ij}]$ is a lower triangular matrix such that $\sigma_{ij} = \sum_{k=1}^{\min\{i,j\}} l_{ik} l_{jk}$. We denote the variance vector as $\boldsymbol{\sigma}^2 := [\sigma_1^2, \cdots, \sigma_D^2]^{\mathsf{T}}$, then each variance component, $\sigma_i^2 := \sigma_{ii}$, can be written in terms of the corresponding row \mathbf{l}_i of \mathbf{L} as follows:

$$\sigma_i^2 = \sum_{k=1}^i l_{ik}^2 = ||\mathbf{l}_i||^2, \quad \mathbf{l}_i := [l_{i1}, l_{i2}, \cdots, l_{ii}]$$
 (1)

For Σ to be positive definite, it is equivalent to require all the leading principal minors $\{M_i\}$ to be positive,

$$M_i = \prod_{k=1}^{i} l_{kk}^2 > 0, \ i = 1, \dots, D \iff l_{ii} \neq 0, \ i = 1, \dots, D$$
 (2)

Based on (1) and (2), for $i \in \{1, \dots, D\}$, \mathbf{l}_i can be viewed as a point on a sphere with radius σ_i excluding the equator, denoted as $S_0^{i-1}(\sigma_i) := \{\mathbf{l} \in \mathbb{R}^i | ||\mathbf{l}||_2 = \sigma_i, l_{ii} \neq 0\}$. Therefore the space of the Cholesky factor in terms of its rows can be written as a product of spheres and we require

$$(\mathbf{l}_1, \mathbf{l}_2, \cdots, \mathbf{l}_D) \in \mathcal{S}_0^0(\sigma_1) \times \mathcal{S}_0^1(\sigma_2) \cdots \times \mathcal{S}_0^{D-1}(\sigma_D)$$
(3)

Note that (3) is the sufficient and necessary condition for the matrix $\Sigma = \mathbf{L}\mathbf{L}^{\mathsf{T}}$ to be a covariance matrix.

We present probabilistic models involving covariance matrices in the following generic form:

$$\mathbf{y}|\boldsymbol{\Sigma}(\boldsymbol{\sigma}, \mathbf{L}) \sim \ell(\mathbf{y}; \boldsymbol{\Sigma}(\boldsymbol{\sigma}, \mathbf{L})), \quad \boldsymbol{\Sigma}(\boldsymbol{\sigma}, \mathbf{L}) = \mathbf{L}\mathbf{L}^{\mathsf{T}}$$

$$\boldsymbol{\sigma} \sim p(\boldsymbol{\sigma})$$

$$\mathbf{L}|\boldsymbol{\sigma} \sim p(\mathbf{L}; \boldsymbol{\sigma}), \quad \operatorname{vech}^{\mathsf{T}}(\mathbf{L}) \in \prod_{i=1}^{D} \mathcal{S}_{0}^{i-1}(\sigma_{i})$$

$$(4)$$

where $\boldsymbol{\sigma} := [\sigma_1, \dots, \sigma_D]^\mathsf{T}$, and the half-vectorization in row order, vech^{T}, transforms the lower triangular matrix \mathbf{L} into a vector $(\mathbf{l}_1, \mathbf{l}_2, \dots, \mathbf{l}_D)$. The total dimension of $(\boldsymbol{\sigma}, \mathbf{L})$ is $\frac{D(D+1)}{2}$.

Alternatively, if we separate variances from covariance, then we have a unique Cholesky decomposition for the correlation matrix $\mathbf{P} = [\rho_{ij}] = \mathbf{L}^*(\mathbf{L}^*)^\mathsf{T}$, where the Cholesky factor $\mathbf{L}^* = \mathrm{diag}(\boldsymbol{\sigma}^{-1})\mathbf{L}$ can be obtained by normalizing each row of \mathbf{L} . The magnitude requirements for correlations are immediately satisfied by the Cauchy-Schwarz inequality: $|\rho_{ij}| = \frac{|\sigma_{ij}|}{\sigma_i \sigma_j} = \frac{|\langle \mathbf{l}_i, \mathbf{l}_j \rangle|}{\|\mathbf{l}_i\|_2 \|\mathbf{l}_j\|_2} \le 1$. Thus we require

$$(\mathbf{l}_1^*, \mathbf{l}_2^*, \cdots, \mathbf{l}_D^*) \in \mathcal{S}_0^0 \times \mathcal{S}_0^1 \cdots \times \mathcal{S}_0^{D-1}$$

$$\tag{5}$$

where $S_0^{i-1} := S_0^{i-1}(1)$. Similarly, (5) is the sufficient and necessary condition for $\mathbf{P} = \mathbf{L}^*(\mathbf{L}^*)^\mathsf{T}$ to be a correlation matrix. Then we have the following alternatively structured model for covariance Σ that involves correlation \mathbf{P} explicitly

$$\mathbf{y}|\boldsymbol{\Sigma}(\boldsymbol{\sigma}, \mathbf{L}^*) \sim \ell(\mathbf{y}; \boldsymbol{\Sigma}(\boldsymbol{\sigma}, \mathbf{L}^*)), \quad \boldsymbol{\Sigma}(\boldsymbol{\sigma}, \mathbf{L}^*) = \operatorname{diag}(\boldsymbol{\sigma}) \mathbf{P} \operatorname{diag}(\boldsymbol{\sigma}), \quad \mathbf{P} = \mathbf{L}^* (\mathbf{L}^*)^\mathsf{T}$$

$$\boldsymbol{\sigma} \sim p(\boldsymbol{\sigma})$$

$$\mathbf{L}^* \sim p(\mathbf{L}^*), \quad \operatorname{vech}^\mathsf{T}(\mathbf{L}^*) \in \prod_{i=1}^D \mathcal{S}_0^{i-1}$$
(6)

Note, this direct decomposition $\Sigma = \operatorname{diag}(\sigma) \mathbf{P} \operatorname{diag}(\sigma)$ as a separation strategy is motivated by statistical thinking in terms of standard deviations and correlations (Barnard et al., 2000). This setting is especially relevant if the statistical quantity of interest is correlation matrix \mathbf{P} itself, and we can then skip inference of the standard deviation σ by fixing it to a data-derived point estimate.

In what follows, we will show that the above framework includes the inverse-Wishart prior as a special case, but it can be easily generalized to a broader range of priors for additional flexibility. Such flexibility enables us to better express prior knowledge, control the model complexity and speed up computation in modeling real-life phenomena. This is crucial in modeling spatio-temporal processes with complex structures.

2.1 Connection to the Inverse-Wishart Prior

There are some interesting connections between the spherical product representations (3) (5) and the early development of the Wishart distribution (Wishart, 1928). The original Wishart distribution was derived by orthogonalizing multivariate Gaussian random variables leading to a lower triangular matrix whose elements $\{t_{ij}^*|i\geq j\}$ (analogous to l_{ij} or l_{ij}^*) were called rectangular coordinates. This way, the probability density has a geometric

¹For each $i \in \{1, \dots, D\}$, given σ_i , there are only (i-1) free parameters on $\mathcal{S}_0^{i-1}(\sigma_i)$, so there are totally $\frac{D(D-1)}{2} + D$ free parameters.

interpretation as a product of volumes and approximate densities on a series of spherical shells with radius $\{t_{ii}^*\}$ (See more details in Sverdrup, 1947; Anderson, 2003). Now we demonstrate that the proposed schemes (4) (6) include the commonly used inverse-Wishart prior as a special case in modeling covariances.

Suppose Σ is a random sample from the inverse-Wishart distribution $\mathcal{W}_D^{-1}(\Psi, \nu)$ with the scale matrix $\Psi > 0$ and the degree of freedom $\nu \geq D$. Therefore, $\Sigma^{-1} \sim \mathcal{W}_D(\Psi^{-1}, \nu)$. Denote \mathbf{C} as the Cholesky factor of Ψ^{-1} , i.e. $\Psi^{-1} = \mathbf{C}\mathbf{C}^{\mathsf{T}}$. Then Σ^{-1} has the following Bartlett decomposition (Anderson, 2003; Smith and Hocking, 1972)

$$\Sigma^{-1} = \mathbf{T} \mathbf{T}^{\mathsf{T}}, \quad \mathbf{T} := \mathbf{C} \mathbf{T}^*, \quad t_{ij}^* \sim \begin{cases} \chi_{D-i+1}, & i = j \\ \mathcal{N}(0,1), & i > j \\ \delta_0, & i < j \end{cases}$$
(7)

where the lower triangular matrix \mathbf{T} , named *Bartlett factor*, has the following density (Theorem 7.2.1 of Anderson, 2003)

$$p(\mathbf{T}) = \frac{|\mathbf{\Psi}|^{\nu/2}}{2^{D(\nu-2)/2} \Gamma_D(\nu/2)} \prod_{i=1}^{D} |t_{ii}|^{\nu-i} \exp\left(-\frac{1}{2} \operatorname{tr}(\mathbf{\Psi} \mathbf{T} \mathbf{T}^{\mathsf{T}})\right)$$

with multivariate gamma function defined as $\Gamma_D(x) := \pi^{D(D-1)/4} \prod_{i=1}^D \Gamma[x + (1-i)/2]$.

Now taking the inverse of the first equation in (7) yields the following reversed Cholesky $decomposition^2$

$$\Sigma = \mathbf{U}\mathbf{U}^\mathsf{T}, \qquad \sigma_{ij} = \sum_{k=\max\{i,j\}}^D u_{ik}u_{jk}, \quad \mathrm{vech}(\mathbf{U}^\mathsf{T}) \in \prod_{i=1}^D \mathcal{S}_0^{D-i}(\sigma_i)$$

where $\mathbf{U} := \mathbf{T}^{-\mathsf{T}}$ is an upper triangular matrix. The following proposition describes the density of the reversed Cholesky factor \mathbf{U} of Σ , which enables us to treat the inverse-Wishart distribution as a special instance of strategy (4) or (6).

Proposition 2.1. Assume $\Sigma \sim W_D^{-1}(\Psi, \nu)$. Then its reversed Cholesky factor U has the following density

$$p(\mathbf{U}) = \frac{|\mathbf{\Psi}|^{\nu/2}}{2^{D(\nu-2)/2} \Gamma_D(\nu/2)} |\mathbf{U}|^{-(\nu+D+1)} \prod_{i=1}^D u_{ii}^i \exp\left(-\frac{1}{2} \text{tr}(\mathbf{\Psi} \mathbf{U}^{-\mathsf{T}} \mathbf{U}^{-1})\right)$$

Proof. See Section A in the supplementary file.

²This can be achieved through the *exchange matrix* (a.k.a. reversal matrix, backward identity, or standard involutory permutation) \mathbf{E} with 1's on the anti-diagonal and 0's elsewhere. Note that \mathbf{E} is both involutory and orthogonal, i.e. $\mathbf{E} = \mathbf{E}^{-1} = \mathbf{E}^{\mathsf{T}}$. Let $\mathbf{E}\boldsymbol{\Sigma}\mathbf{E} = \mathbf{L}\mathbf{L}^{\mathsf{T}}$ be the usual Cholesky decomposition. Then $\boldsymbol{\Sigma} = (\mathbf{E}\mathbf{L}\mathbf{E})(\mathbf{E}\mathbf{L}\mathbf{E})^{\mathsf{T}} = \mathbf{U}\mathbf{U}^{\mathsf{T}}$ and define $\mathbf{U} := \mathbf{E}\mathbf{L}\mathbf{E}^{\mathsf{T}}$.

If we normalize each row of U and write

$$\mathbf{U} = \operatorname{diag}(\boldsymbol{\sigma})\mathbf{U}^*, \quad \sigma_i = \sqrt{\sigma_{ii}} = \|\mathbf{u}_i\|, \quad u_{ij}^* = u_{ij}/\sigma_i,$$

then the following joint prior of (σ, \mathbf{U}^*) is inseparable in general:

$$p(\boldsymbol{\sigma}, \mathbf{U}^*) \propto \prod_{i=1}^{D} |\sigma_i u_{ii}^*|^{i - (\nu + D + 1)} \exp \left\{ -\frac{1}{2} \operatorname{tr}(\boldsymbol{\Psi} \operatorname{diag}(\boldsymbol{\sigma}^{-1}) (\mathbf{U}^*)^{-\mathsf{T}} (\mathbf{U}^*)^{-1} \operatorname{diag}(\boldsymbol{\sigma}^{-1})) \right\}$$
(8)

With this result, we can conditionally model variance and correlation factor as $p(\boldsymbol{\sigma}|\mathbf{U}^*)$ and $p(\mathbf{U}^*|\boldsymbol{\sigma})$ respectively, similarly as in our proposed scheme (4) or (6). It is also used to verify the validity of our proposed method (6) (see more details in Section 4.1.1). A similar result exists for the Wishart prior distribution regarding the Cholesky factor. This representation facilitates the construction of a broader class of more flexible prior distributions for covariance matrix detailed below.

2.2 More Flexible Priors

Within the above framework, the only constraint on \mathbf{U} or \mathbf{L} is that it resides on the product of spheres with increasing dimensions. Using this fact, we can develop a broader class of priors on covariance matrices and thus be able to model processes with more complicated dependence in covariance structures. Since $\boldsymbol{\sigma}$ and \mathbf{L}^* have independent priors in (6), in what follows we focus on the scheme (6), and for simplicity, we denote the normalized Cholesky factor as \mathbf{L} . Also, following Barnard et al. (2000), we assume a log-Normal prior on $\boldsymbol{\sigma}$:

$$\log(oldsymbol{\sigma}) \sim \mathcal{N}(oldsymbol{\xi}, oldsymbol{\Lambda})$$

We now discuss priors on **L** that properly reflect the prior knowledge regarding the covariance structure among variables. If two variables, y_i and y_j (assuming i < j) are known to be uncorrelated a priori, i.e. $0 = \rho_{ij} = \langle \mathbf{l}_i, \mathbf{l}_j \rangle$, then we can choose a prior that encourages $\mathbf{l}_i \perp \mathbf{l}_j$, e.g. $l_{jk} \approx 0$ for $k \leq i$. In contrast, if we believe a priori that there is a strong correlation between the two variables, we can specify that \mathbf{l}_i and \mathbf{l}_j be linearly dependent, e.g., by setting $[l_{jk}]_{k \leq i} \approx \pm \mathbf{l}_i$. When there is no prior information, we might assume that components are uncorrelated and specify priors for \mathbf{l}_i that concentrate on the (two) poles of \mathcal{S}_0^{i-1} ,

$$p(\mathbf{l}_i) \propto |l_{ii}|, \quad i = 2, \cdots, D$$
 (9)

Putting more prior probability on the diagonal elements of **L** renders fewer non-zero off-diagonal elements, which in turn leads to a larger number of perpendicular variables; that is, such a prior favors zeros in the correlation matrix **P**. More generally, one can map a probability distribution defined on the simplex onto the sphere and consider the following squared-Dirichlet distribution.




Figure 1: Symmetric squared-Dirichlet distributions $\operatorname{Dir}^2(\boldsymbol{\alpha})$ defined on the 2-sphere with different settings for concentration parameter $\boldsymbol{\alpha} = \alpha \mathbf{1}$. The uniform distribution on the simplex, $\operatorname{Dir}(\mathbf{1})$, becomes non-uniform on the sphere due to the stretch of geometry (left); the symmetric Dirichlet distribution $\operatorname{Dir}(\frac{1}{2}\mathbf{1})$ becomes uniform on the sphere (middle); with α closer to 0, the induced distribution becomes more concentrated on the polar points (right).

Definition 1 (Squared-Dirichlet distribution). A random vector $\mathbf{l}_i \in \mathcal{S}^{i-1}$ is said to have a squared-Dirichlet distribution with parameter $\boldsymbol{\alpha}_i := (\alpha_{i1}, \alpha_{i2}, \cdots, \alpha_{ii})$ if

$$\mathbf{l}_i^2 := (l_{i1}^2, l_{i2}^2, \cdots, l_{ii}^2) \sim \operatorname{Dir}(\boldsymbol{\alpha}_i)$$

Denote $\mathbf{l}_i \sim \mathrm{Dir}^2(\boldsymbol{\alpha}_i)$. Then \mathbf{l}_i has the following density

$$p(\mathbf{l}_i) = p(\mathbf{l}_i^2)|2\mathbf{l}_i| \propto (\mathbf{l}_i^2)^{\alpha_i - 1}|\mathbf{l}_i| = |\mathbf{l}_i|^{2\alpha_i - 1} := \prod_{k=1}^i |l_{ik}|^{2\alpha_{ik} - 1}$$
(10)

Remark 1. This definition includes a large class of flexible prior distributions on the unit sphere that specify different concentrations of probability density through the parameter α_i . For example, the above prior (9) corresponds to $\alpha_i = (\frac{1}{2}, \dots, \frac{1}{2}, 1)$.

To induce a prior distribution for the correlation matrix $\mathbf{P} = \mathbf{L}\mathbf{L}^{\mathsf{T}}$, one can specify priors on row vectors of \mathbf{L} , $\mathbf{l}_i \sim \mathrm{Dir}^2(\boldsymbol{\alpha}_i)$ for $i=2,\cdots,D$. To encourage small correlation, we choose the concentration parameter $\boldsymbol{\alpha}_i$ so that the probability density concentrates around the (two) poles of \mathcal{S}_0^{i-1} , e.g. $0 < \alpha_{ik} \ll \alpha_{ii}$ for k < i. Figure 1 illustrates the density heat maps of some symmetric squared-Dirichlet distributions $\mathrm{Dir}^2(\alpha \mathbf{1})$ on \mathcal{S}^2 . It is interesting that the squared-Dirichlet distribution induces two important uniform prior distributions over correlation matrices from Barnard et al. (2000) in an effort to provide flexible priors for covariance matrices, as stated in the following theorem.

Theorem 2.1 (Uniform distributions). Let $\mathbf{P} = \mathbf{L}\mathbf{L}^{\mathsf{T}}$. Suppose $\mathbf{l}_i \sim \mathrm{Dir}^2(\boldsymbol{\alpha}_i)$, for $i = 2, \dots, D$, are independent, where \mathbf{l}_i is the *i*-th row of \mathbf{L} . We have the following

- 1. If $\alpha_i = (\frac{1}{2}\mathbf{1}_{i-1}^\mathsf{T}, \alpha_{ii}), \ \alpha_{ii} = \frac{(i-2)D-1}{2}$, then **P** follows a marginally uniform distribution, that is, $\rho_{ij} \sim \mathrm{Unif}(-1,1), \quad i \neq j$.
- 2. If $\alpha_i = (\frac{1}{2}\mathbf{1}_{i-1}^\mathsf{T}, \alpha_{ii}), \ \alpha_{ii} = \frac{D-i}{2} + 1$, then **P** follows a jointly uniform distribution, that is, $p(\mathbf{P}) \propto 1$.

Proof. See Section A in the supplementary file.

Another natural spherical prior can be obtained by constraining a multivariate Gaussian random vector to have unit norm. This is later generalized to a vector Gaussian process constrained to a sphere that serves as a suitable prior for modeling correlation processes. Now we consider the following *unit-vector Gaussian* distribution:

Definition 2 (Unit-vector Gaussian distribution). A random vector $\mathbf{l}_i \in \mathcal{S}^{i-1}$ is said to have a unit-vector Gaussian distribution with mean $\boldsymbol{\mu}$ and covariance $\boldsymbol{\Sigma}$ if

$$\mathbf{l}_i \sim \mathcal{N}_i(\boldsymbol{\mu}, \boldsymbol{\Sigma}), \quad with \|\mathbf{l}_i\|_2 = 1$$

Then we denote $l_i \sim \mathcal{N}_i^{\mathcal{S}}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ and l_i has the following (conditional) density

$$p(\mathbf{l}_i| \|\mathbf{l}_i\|_2 = 1) = \frac{1}{(2\pi)^{\frac{i}{2}} |\boldsymbol{\Sigma}|^{\frac{1}{2}}} \exp\left\{-\frac{1}{2}(\mathbf{l}_i - \boldsymbol{\mu})^{\mathsf{T}} \boldsymbol{\Sigma}^{-1} (\mathbf{l}_i - \boldsymbol{\mu})\right\}, \quad \|\mathbf{l}_i\|_2 = 1$$

Remark 2. This conditional density essentially defines the following Fisher-Bingham distribution (a.k.a. generalized Kent distribution, Kent, 1982; Mardia and Jupp, 2009). If $\Sigma = \mathbf{I}$, then the above distribution reduces to the von Mises-Fisher distribution (Fisher, 1953; Mardia and Jupp, 2009) as a special case. If in addition $\mu = \mathbf{0}$, then the above density becomes a constant; that is, the corresponding distribution is uniform on the sphere S_0^{i-1} . See more details in Section E.1 of the supplementary file.

2.3 Dynamically Modeling the Covariance Matrices

We can generalize the proposed framework for modeling covariance/correlation matrices to the dynamic setting by adding subscript t to variables in the model (4) and the model (6), thus called dynamic covariance and dynamic correlation models respectively. We focus the latter in this section. One can model the components of σ_t as independent dynamic processes using, e.g. ARMA, GARCH, or log-Gaussian process. For \mathbf{L}_t , we use vector processes. Since each row of \mathbf{L}_t has to be on a sphere of certain dimension, we require the unit norm constraint for the dynamic process over time. We refer to any multivariate process $\mathbf{l}_i(x)$ satisfying $\|\mathbf{l}_i(x)\| \equiv 1$, $\forall x \in \mathcal{X}$ as unit-vector process (uvP). A unit-vector process can be obtained by constraining an existing multivariate process, e.g. the vector Gaussian process (vGP), as defined below.

Definition 3 (Vector Gaussian process). A *D*-dimensional vector Gaussian process $\mathbf{Z}(x) := (Z_1(x), \dots, Z_D(x))$, with vector mean function $\boldsymbol{\mu}(x) = (\mu_1(x), \dots, \mu_D(x))$, covariance function \mathcal{C} and (*D*-dimensional) cross covariance $\mathbf{V}_{D\times D}$,

$$\mathbf{Z}(x) \sim \mathcal{GP}_D(\boldsymbol{\mu}, \mathcal{C}, \mathbf{V}_{D \times D})$$

is a collection of D-dimensional random vectors, indexed by $x \in \mathcal{X}$, such that for any finite set of indices $\{x_1, \dots, x_N\}$, the random matrix $\widetilde{\mathbf{Z}}_{N \times D} := (\mathbf{Z}(x_1), \dots, \mathbf{Z}(x_N))^\mathsf{T}$ has the following matrix normal distribution

$$\widetilde{\mathbf{Z}}_{N imes D} \sim \mathcal{MN}_{N imes D}(\mathbf{M}_{N imes D}, \mathbf{K}_{N imes N}, \mathbf{V}_{D imes D})$$

where $\mathbf{M}_{N\times D} := (\mathbf{m}_1, \cdots, \mathbf{m}_D)$, and $\mathbf{m}_k = (\mu_k(x_1), \cdots, \mu_k(x_N))^\mathsf{T}$, and \mathbf{K} is the kernel matrix with elements $K_{ij} = \mathcal{C}(x_i, x_j)$.

Remark 3. Note for each $k = 1, \dots D$, we have the following marginal GP

$$Z_k(x) \sim \mathcal{GP}(\boldsymbol{\mu}_k, \mathcal{C})$$

In the above definition, we require a common kernel C for all the marginal GPs, whose dependence is characterized by the cross covariance $\mathbf{V}_{D\times D}$. On the other hand, for any fixed $x^* \in \mathcal{X}$, we have

$$\mathbf{Z}(x^*) \sim \mathcal{N}_D(\boldsymbol{\mu}(x^*), \mathbf{V}_{D \times D})$$

For simplicity, we often consider $\mu \equiv 0$ and $\mathbf{V}_{D \times D} = \mathbf{I}_D$. That is, $Z_k(x) \stackrel{iid}{\sim} \mathcal{GP}(0, \mathcal{C})$ for $k = 1, \dots, D$.

Restricting vGP $\mathbf{Z}(\cdot)$ to sphere yields a unit-vector Gaussian process (uvGP) $\mathbf{Z}^*(\cdot) := \mathbf{Z}(\cdot) | \{ \| \mathbf{Z}(\cdot) \|_2 \equiv 1 \}$, denoted as $\mathbf{Z}^*(\cdot) \sim \mathcal{GP}_D^{\mathcal{S}}(\boldsymbol{\mu}, \mathcal{C}, \mathbf{V})$. Note for any fixed $x^* \in \mathcal{X}$, $\mathbf{Z}^*(x^*) \sim \mathcal{N}_D^{\mathcal{S}}(\boldsymbol{\mu}, \mathbf{V})$. Setting $\mu \equiv 0$, $\mathbf{V} = \mathbf{I}$, and conditioned on the length ℓ_n of each row of $\widetilde{\mathbf{Z}}$, we have

$$p(\widetilde{\mathbf{Z}}^* | \{ \|z_{n \cdot}\| = \ell_n \}) = \frac{\prod_{n=1}^N \ell_n^D}{(2\pi)^{\frac{ND}{2}} |\mathbf{K}|^{\frac{D}{2}}} \exp \left\{ -\frac{1}{2} \operatorname{tr} \left[(\widetilde{\mathbf{Z}}^*)^\mathsf{T} \operatorname{diag}(\{\ell_n\}) \mathbf{K}^{-1} \operatorname{diag}(\{\ell_n\}) \widetilde{\mathbf{Z}}^* \right] \right\}$$

This conditional density is preserved by the inference algorithm in Section 3 and used for defining priors for correlations with all $\ell_n = 1$. For each marginal GP, we select the following exponential function as the common kernel

$$C(x, x') = \gamma \exp(-0.5||x - x'||^s/\rho^s)$$

where s controls the smoothness, the scale parameter γ is given an inverse-Gamma prior, and the correlation length parameter ρ is given a log-normal prior. Figure 2 shows a realization of vector GP \mathbf{Z}_t , unit-vector GP (forming rows of) \mathbf{L}_t and the induced correlation process \mathbf{P}_t respectively.




Figure 2: A realization of vector GP \mathbf{Z}_t (left), unit-vector GP (forming rows of) \mathbf{L}_t (middle) and the induced correlation process \mathbf{P}_t (right).

In what follows, we focus on multivariate time series; therefore, we use the one dimensional time index $t \in \mathcal{X} = \mathbb{R}^+$. The overall dynamic correlation model can be summarized as follows:

$$\mathbf{y}_{t} \sim \mathcal{N}(\boldsymbol{\mu}_{t}, \boldsymbol{\Sigma}_{t}), \quad \boldsymbol{\Sigma}_{t} = \operatorname{diag}(\boldsymbol{\sigma}_{t}) \mathbf{L}_{t} \mathbf{L}_{t}^{\mathsf{T}} \operatorname{diag}(\boldsymbol{\sigma}_{t})$$

$$\boldsymbol{\mu}_{t} \sim \mathcal{GP}_{D}(0, \mathcal{C}_{\mu}, \mathbf{I}), \quad \mathcal{C}_{\mu}(t, t') = \gamma_{\mu} \exp(-0.5 \|t - t'\|^{s} / \rho_{\mu}^{s})$$

$$\operatorname{log} \boldsymbol{\sigma}_{t} \sim \mathcal{GP}_{D}(0, \mathcal{C}_{\sigma}, \mathbf{I}), \quad \mathcal{C}_{\sigma}(t, t') = \gamma_{\sigma} \exp(-0.5 \|t - t'\|^{s} / \rho_{\sigma}^{s})$$

$$\mathbf{l}_{i}(t) \sim \mathcal{GP}_{i}^{\mathcal{S}}(\mathbf{n}_{i}, \mathcal{C}_{L}, \mathbf{I}), \quad \mathcal{C}_{L}(t, t') = \gamma_{L} \exp(-0.5 \|t - t'\|^{s} / \rho_{L}^{s})$$

$$\gamma_{*} \sim \Gamma^{-1}(a_{*}, b_{*}), \quad \operatorname{log} \rho_{*} \sim \mathcal{N}(m_{*}, V_{*}), \quad * = \mu, \sigma, \text{ or } L$$

$$(11)$$

where a constant mean function $\mathbf{n}_i = (0, \dots, 0, 1)$ is used in the uvGP prior for $\mathbf{l}_i(t)$, with mean matrix $\mathbf{M} = \mathbf{1}_N \otimes \mathbf{n}_i^\mathsf{T}$ for the realization $\tilde{\mathbf{l}}_i$. This model (11) captures the spatial dependence in the matrix $\boldsymbol{\Sigma}_t$, which evolves along the time; while the temporal correlation is characterized by various GPs. The induced covariance process $\boldsymbol{\Sigma}_t$ is not a generalized Wishart process (Wilson and Ghahramani, 2011), which only models Cholesky factor of covariance using GP. Though with GP, dynamic covariance model may work similarly as the dynamic correlation model (11), yet the latter provides extra flexibility in modeling the evolution of variances and correlations separately. In general such flexibility could be useful in handling constraints for processes, e.g. modeling the dynamic probability for binary time series.

With this structured model (11), one can naturally model the evolution of variances and correlations separately in order to obtain more flexibility. If the focus is on modeling the correlation among multiple time series, then one can substitute σ_t with a point estimate $\hat{\sigma}$ from one trial and assume a steady variance vector. Alternatively, if sufficient trials are present, one can obtain an empirical estimate, $\hat{\sigma}_t$, from multiple trials at each time point. In the following, we study the posterior contraction of GP modeling in this setting.

2.4 Posterior Contraction Theorem

We now provide a theorem on the posterior contraction of the dynamic covariance model before we conclude this section. Because the posterior contraction for mean regression using Gaussian process has been vastly investigated in the literature (van der Vaart and van Zanten, 2008a, 2009, 2011; Yang and Dunson, 2016), we only investigate the posterior contraction for the covariance regression and set $\mu_t \equiv 0$. We leave the posterior contraction of the dynamic correlation model (11) for future work. Note, the Cholesky decomposition of covariance matrix $\Sigma = \mathbf{L}\mathbf{L}^{\mathsf{T}}$ is unique if all the diagonal entries of \mathbf{L} are positive. Therefore in the remaining of this section, we identify Cholesky factors up to a column-wise sign, i.e. $\mathbf{L} \sim \mathbf{L} \operatorname{diag}(-\sum_{j \in J} e_j)$ for $J \subset \{1, \dots, D\}$ where e_j is the j-th column of identity matrix \mathbf{I}_D .

In most cases, Gaussian process \mathbf{L}_t can be viewed as a tight Borel measurable map in a Banach space, e.g. a space of continuous functions or L_p space. It is well known that the support of a centered GP is equal to the closure of the reproducible kernel Hilbert space (RKHS) \mathbb{H} associated to this process (Lemma 5.1 of van der Vaart and van Zanten, 2008b). Because the posterior distribution necessarily puts all its mass on the support of the prior, the posterior consistency requires the true parameter \mathbf{L}_0 governing the distribution of the data to fall in this support (van der Vaart and van Zanten, 2008a). Following van der Vaart and van Zanten (2008a,b, 2011), we express the rate of the posterior contraction in terms of the concentration function

$$\phi_{\mathbf{L}_0}(\varepsilon) = \inf_{h \in \mathbb{H}: \|h - \mathbf{L}_0\| < \varepsilon} \|h\|_{\mathbb{H}}^2 - \log \Pi(\mathbf{L}: \|\mathbf{L}\| < \varepsilon)$$
(12)

where $\|\cdot\|$ is the norm of the Banach space where the GP **L** takes value, Π is the GP prior and \mathbb{H} is the associated RKHS with norm $\|\cdot\|_{\mathbb{H}}$. Under certain regularity conditions, the posterior contracts with increasing data expressed in n at the rate $\varepsilon_n \to 0$ satisfying

$$\phi_{\mathbf{L}_0}(\varepsilon_n) \le n\varepsilon_n^2 \tag{13}$$

Define $\|\mathbf{L}(t)\|_{\infty} := \max_{1 \leq i,j \leq D} \sup_{t \in \mathcal{X}} |l_{ij}(t)|$. Consider the separable Banach space $L^{\infty}(\mathcal{X})^{D(D+1)/2} := \{\mathbf{L}(t) : \|\mathbf{L}(t)\|_{\infty} < +\infty\}$. Let p be a (centered) Gaussian model, which is uniquely determined by the covariance matrix $\boldsymbol{\Sigma} = \mathbf{L}\mathbf{L}^{\mathsf{T}}$. Therefore the model density is parametrized by \mathbf{L} , hence denoted as $p_{\mathbf{L}}$. Denote $P_{\mathbf{L}}^{(n)} := \bigotimes_{i=1}^{n} P_{\mathbf{L},i}$ as the product measure on $\bigotimes_{i=1}^{n} (\mathcal{X}_{i}, \mathcal{B}_{i}, \mu_{i})$. Each $P_{\mathbf{L},i}$ has a density $p_{\mathbf{L}_{i}}$ with respect to the σ -finite measure μ_{i} . Define the average Hellinger distance as $d_{n}^{2}(\mathbf{L}, \mathbf{L}') = \frac{1}{n} \sum_{i=1}^{n} \int (\sqrt{p_{\mathbf{L},i}} - \sqrt{p_{\mathbf{L}',i}})^{2} d\mu_{i}$. Denote the observations $Y^{(n)} = \{Y_{i}\}_{i=1}^{n}$ with $Y_{i} = \mathbf{y}(t_{i})$. Note they are independent but not identically distributed (inid). Now we state the main theorem of posterior contraction.

Theorem 2.2 (Posterior contraction). Let $\mathbf{L} - \mathbf{I}$ be a Borel measurable, zero-mean tight Gaussian random element in $L^{\infty}(\mathcal{X})^{D(D+1)/2}$ and $P_{\mathbf{L}}^{(n)} = \bigotimes_{i=1}^{n} P_{\mathbf{L},i}$ be the product measure of $Y^{(n)}$ parametrized by \mathbf{L} . Let $\phi_{\mathbf{L}_0}$ be the function in (12) with the uniform norm $\|\cdot\|_{\infty}$.

If \mathbf{L}_0 is contained in the support of \mathbf{L} and $\phi_{\mathbf{L}_0}$ satisfies (13) with $\varepsilon_n \geq n^{-\frac{1}{2}}$, then $\Pi_n(\mathbf{L}: d_n(\mathbf{L}, \mathbf{L}_0) > M_n \varepsilon_n | Y^{(n)}) \to 0$ in $P_{\mathbf{L}_0}^{(n)}$ -probability for every $M \to \infty$.

Proof. See Section B in the supplementary file.

Remark 4. In principle, the smoothness of GP should match the regularity of the true parameter to achieve the optimal rate of contraction (van der Vaart and van Zanten, 2008a, 2011). One can scale GP, e.g. using an inverse-Gamma bandwidth, to get optimal contraction rate for every regularity level so that the resulting estimator is rate adaptive (van der Vaart and van Zanten, 2009, 2011). One can refer to Section 3.2 of (van der Vaart and van Zanten, 2011) for posterior contraction rates using squared exponential kernel for GP. We leave further investigation on contraction rates in the setting of covariance regression to future work.

Remark 5. Here the GP prior \mathbf{L} defines a (mostly finite) probability measure on the space of bounded functions. The true parameter function \mathbf{L}_0 is required to be contained in the support of the prior, the RKHS of \mathbf{L} . The contraction rate depends on the position of \mathbf{L}_0 relative to the RKHS and the small-ball probability $\Pi(\|\mathbf{L}\| < \varepsilon)$.

3 Posterior Inference

Now we obtain the posterior probability of mean μ_t , variance σ_t , Cholesky factor of correlation \mathbf{L}_t , hyper-parameters $\boldsymbol{\gamma} := (\gamma_{\mu}, \gamma_{\sigma}, \gamma_L)$ and $\boldsymbol{\rho} := (\rho_{\mu}, \rho_{\sigma}, \rho_L)$ in the model (11). Denote the realization of processes $\boldsymbol{\mu}_t, \boldsymbol{\sigma}_t, \mathbf{L}_t$ at discrete time points $\{t_n\}_{n=1}^N$ as $\tilde{\boldsymbol{\mu}}_{N\times D}, \tilde{\boldsymbol{\sigma}}_{N\times D}, \tilde{\mathbf{L}}_{N\times D\times D}$ respectively. Transform the parameters $\tilde{\boldsymbol{\tau}} := \log(\tilde{\boldsymbol{\sigma}}), \, \boldsymbol{\eta} := \log(\boldsymbol{\rho})$ for the convenience of calculation. Denote $\tilde{\mathbf{Y}}_{M\times N\times D} := \{\mathbf{Y}_1, \cdots, \mathbf{Y}_M\}$ for M trials, $(\mathbf{Y}_m)_{N\times D} := [\mathbf{y}_{m1}, \cdots, \mathbf{y}_{mN}]^\mathsf{T}$ and $\mathbf{y}_{mn}^* := (\mathbf{y}_{mn} - \boldsymbol{\mu}_n) \circ e^{-\tau_n}$ where \circ is the Hadamard product (a.k.a. Schur product), i.e. the entry-wise product. Let $\mathbf{K}_*(\gamma_*, \eta_*) = \gamma_* \mathbf{K}_{0*}(\eta_*)$ and $\tilde{\mathbf{I}}_i^* := \tilde{\mathbf{I}}_i - \mathbf{1}_N \otimes \mathbf{n}_i^\mathsf{T}$.

3.1 Metropolis-within-Gibbs

We use a Metropolis-within-Gibbs algorithm and alternate updating the model parameters $\widetilde{\mu}, \widetilde{\tau}, \widetilde{\mathbf{L}}, \gamma, \eta$. We now list the parameters and their respective updates one by one.

 (γ) . Note the prior for γ is conditionally conjugate given $* = \mu, \tau$, or L,

$$\gamma_*|\tilde{*}, \eta_* \sim \Gamma^{-1}(a'_*, b'_*), \quad a'_* = a_* + \frac{ND}{2}(\frac{D+1}{2} - \frac{1}{D})^{[*=L]}, \quad b'_* = b_* + \frac{1}{2}\operatorname{tr}(\tilde{*}^\mathsf{T}\mathbf{K}_{0*}(\eta_*)^{-1}\tilde{*})$$

where [condition] is 1 with the condition satisfied and 0 otherwise.

(η). Given * = μ , τ , or L, we could sample η_* using the slice sampler (Neal, 2003), which only requires log-posterior density and works well for scalar parameters,

$$\log p(\eta_*|\tilde{*}, \gamma_*) = -\frac{D(\frac{D+1}{2} - \frac{1}{D})^{[*=L]}}{2} \log |\mathbf{K}_{0*}(\eta_*)| - \frac{\operatorname{tr}(\tilde{*}^\mathsf{T}\mathbf{K}_{0*}(\eta_*)^{-1}\tilde{*})}{2\gamma_*} - \frac{(\eta_* - m_*)^2}{2V_*}$$

By the definition of vGP, we have $\widetilde{\boldsymbol{\mu}}|\gamma_{\mu}, \eta_{\mu} \sim \mathcal{MN}_{N \times D}(\mathbf{0}, \mathbf{K}_{\mu}, \mathbf{I}_{D})$; therefore, $\operatorname{vec}(\widetilde{\boldsymbol{\mu}})|\gamma_{\mu},\eta_{\mu}\sim\mathcal{N}_{ND}(\mathbf{0},\mathbf{I}_{D}\otimes\mathbf{K}_{\mu}).$ On the other hand, one can write

$$\sum_{m=1}^{M} \sum_{n=1}^{N} \mathbf{y}_{mn}^{*} \mathbf{P}_{n}^{-1} \mathbf{y}_{mn}^{*} = \sum_{m=1}^{M} \operatorname{vec}((\mathbf{Y}_{m} - \widetilde{\boldsymbol{\mu}})^{\mathsf{T}})^{\mathsf{T}} \operatorname{diag}(\{\widetilde{\boldsymbol{\Sigma}}_{n}^{-1}\}) \operatorname{vec}((\mathbf{Y}_{m} - \widetilde{\boldsymbol{\mu}})^{\mathsf{T}})$$

$$= \sum_{m=1}^{M} (\operatorname{vec}(\mathbf{Y}_{m}) - \operatorname{vec}(\widetilde{\boldsymbol{\mu}}))^{\mathsf{T}} \widetilde{\boldsymbol{\Sigma}}_{K}^{-1} (\operatorname{vec}(\mathbf{Y}_{m}) - \operatorname{vec}(\widetilde{\boldsymbol{\mu}}))$$

where $\widetilde{\boldsymbol{\Sigma}}_{K}^{-1} := K_{(D,N)} \operatorname{diag}(\{\widetilde{\boldsymbol{\Sigma}}_{n}\})^{-1} K_{(N,D)}$, and $K_{(N,D)}$ is the commutation matrix of size $ND \times ND$ such that for any $N \times D$ matrix \mathbf{A} , $K_{(N,D)} \text{vec}(\mathbf{A}) = \text{vec}(\mathbf{A}^{\mathsf{T}})$ (Tracy and Dwyer, 1969; Magnus and Neudecker, 1979). Therefore, the prior on $\text{vec}(\widetilde{\mu})$ is conditionally conjugate, and we have

$$\operatorname{vec}(\widetilde{\boldsymbol{\mu}})|\widetilde{\mathbf{Y}}, \widetilde{\boldsymbol{\Sigma}}, \gamma_{\mu}, \eta_{\mu} \sim \mathcal{N}_{ND}(\boldsymbol{\mu}', \boldsymbol{\Sigma}')$$

$$\boldsymbol{\mu}' = \boldsymbol{\Sigma}' \widetilde{\boldsymbol{\Sigma}}_{K}^{-1} \sum_{m=1}^{M} \operatorname{vec}(\mathbf{Y}_{m}), \quad \boldsymbol{\Sigma}' = \left(\mathbf{I}_{D} \otimes \mathbf{K}_{\mu}^{-1} + M \widetilde{\boldsymbol{\Sigma}}_{K}^{-1}\right)^{-1}$$

Using a similar argument by matrix Normal prior for $\tilde{\tau}$, we have $\text{vec}(\tilde{\tau})|\gamma_{\tau},\eta_{\tau}\sim$ $\mathcal{N}_{ND}(\mathbf{0}, \mathbf{I}_D \otimes \mathbf{K}_{\tau})$. Therefore, we could use the elliptic slice sampler (ESS, Murray et al., 2010), which only requires the log-likelihood

$$\log p(\widetilde{\boldsymbol{\tau}}; \widetilde{\mathbf{Y}}, \widetilde{\boldsymbol{\mu}}) = -M \mathbf{1}_{ND}^{\mathsf{T}} \text{vec}(\widetilde{\boldsymbol{\tau}}) - \sum_{m=1}^{M} \frac{1}{2} \text{vec}(\mathbf{Y}_{m}^{*})^{\mathsf{T}} \widetilde{\mathbf{P}}_{K}^{-1} \text{vec}(\mathbf{Y}_{m}^{*})$$

where $\widetilde{\mathbf{P}}_K^{-1} := K_{(D,N)} \operatorname{diag}(\{\widetilde{\mathbf{P}}_n\})^{-1} K_{(N,D)}$ and $\mathbf{Y}_m^* := (\mathbf{Y}_m - \widetilde{\boldsymbol{\mu}}) \circ \exp(-\widetilde{\boldsymbol{\tau}})$. $(\widetilde{\mathbf{L}})$. For each $n \in \{1, \dots N\}$, we have $\operatorname{vech}^\mathsf{T}(\mathbf{L}_n) \in \prod_{i=1}^D \mathcal{S}_0^{i-1}$. We could sample from its posterior distribution using the Δ -Spherical Hamiltonian Monte Carlo (Δ -SphHMC) described below. The log-posterior density of L is

$$\log p(\widetilde{\mathbf{L}}|\widetilde{\mathbf{Y}}, \widetilde{\boldsymbol{\mu}}, \widetilde{\boldsymbol{\tau}}, \gamma_L, \eta_L) = -\sum_{n=1}^N \left[M \log |\mathbf{L}_n| + \sum_{m=1}^M \frac{1}{2} \mathbf{y}_{mn}^* ^\mathsf{T} \mathbf{P}_n^{-1} \mathbf{y}_{mn}^* \right] - \frac{1}{2} \sum_{i=2}^D \operatorname{tr}(\widetilde{\mathbf{l}}_i^{*\mathsf{T}} \mathbf{K}_L^{-1} \widetilde{\mathbf{l}}_i^*)$$

The derivative of log-likelihood with respect to \mathbf{L}_n and the derivative of log-prior with respect to \mathbf{l}_i can be calculated as

$$\frac{\partial \log p(\widetilde{\mathbf{L}}; \widetilde{\mathbf{Y}}, \widetilde{\boldsymbol{\mu}}, \widetilde{\boldsymbol{\tau}})}{\partial \mathbf{L}_n} = -M \frac{\mathbf{I}_D}{\mathbf{L}_n} + \sum_{m=1}^{M} \mathrm{tril}(\mathbf{P}_n^{-1} \mathbf{y}_{mn}^* \mathbf{y}_{mn}^* {}^\mathsf{T} \mathbf{L}_n^{-\mathsf{T}}), \ \frac{\partial \log p(\widetilde{\mathbf{L}}|\gamma_L, \eta_L)}{\partial \widetilde{\mathbf{l}}_i} = -\mathbf{K}_L^{-1} \widetilde{\mathbf{l}}_i^*$$

3.2 Spherical HMC

We need an efficient algorithm to handle the intractability in the posterior distribution of $\widetilde{\mathbf{L}}$ introduced by various flexible priors. Spherical Hamiltonian Monte Carlo (SphHMC, Lan et al., 2014; Lan and Shahbaba, 2016) is a Hamiltonian Monte Carlo (HMC, Duane et al., 1987; Neal, 2011) algorithm on spheres that can be viewed as a special case of geodesic Monte Carlo (Byrne and Girolami, 2013), or manifold Monte Carlo methods (Girolami and Calderhead, 2011; Lan et al., 2015). The algorithm was originally proposed to handle norm constraints in sampling so it is natural to use it to sample each row of the Cholesky factor of a correlation matrix with unit 2-norm constraint. The general notation \mathbf{q} is instantiated as \mathbf{l}_i in this section.

Assume a probability distribution with density function $f(\mathbf{q})$ is defined on a (D-1) dimensional sphere with radius r, $\mathcal{S}^{D-1}(r)$. Due to the norm constraint, there are (D-1) free parameters $\mathbf{q}_{-D} := (q_1, \cdots, q_{D-1})$, which can be viewed as the Cartesian coordinates for the manifold $\mathcal{S}^{D-1}_+(r)$. To induce Hamiltonian dynamics on the sphere, we define the potential energy for position \mathbf{q} as $U(\mathbf{q}) := -\log f(\mathbf{q})$. Endowing the canonical spherical metric $\mathbf{G}(\mathbf{q}_{-D}) = \mathbf{I}_{D-1} + \frac{\mathbf{q}_{-D}\mathbf{q}_{-D}^{\mathsf{T}}}{q_D^2}$ on the Riemannian manifold $\mathcal{S}^{D-1}(r)$, we introduce the auxiliary velocity vector $\mathbf{v}|\mathbf{q} \sim \mathcal{N}(\mathbf{0}, \mathbf{G}(\mathbf{q})^{-1})$ and define the associated kinetic energy as $K(\mathbf{v};\mathbf{q}) := -\log f_{\mathcal{N}}(\mathbf{v}|\mathbf{q}) = -\frac{1}{2}\log|\mathbf{G}(\mathbf{q}_{-D})| + \frac{1}{2}\mathbf{v}_{-D}^{\mathsf{T}}\mathbf{G}(\mathbf{q}_{-D})\mathbf{v}_{-D}$ (Girolami and Calderhead, 2011). Therefore the total energy is defined as

$$E(\mathbf{q}, \mathbf{v}) := U(\mathbf{q}) + K(\mathbf{v}; \mathbf{q}) = \tilde{U}(\mathbf{q}) + K_0(\mathbf{v}; \mathbf{q})$$
(14)

where we denote $\tilde{U}(\mathbf{q}) := U(\mathbf{q}) - \frac{1}{2} \log |\mathbf{G}(\mathbf{q}_{-D})| = -\log f(\mathbf{q}) + \log |q_D|$, and $K_0(\mathbf{v}; \mathbf{q}) := \frac{1}{2} \mathbf{v}_{-D}^\mathsf{T} \mathbf{G}(\mathbf{q}_{-D}) \mathbf{v}_{-D} = \frac{1}{2} \mathbf{v}^\mathsf{T} \mathbf{v}$ (Lan and Shahbaba, 2016). Therefore the Lagrangian dynamics with above total energy (14) is (Lan et al., 2015)

$$\dot{\mathbf{q}}_{-D} = \mathbf{v}_{-D}
\dot{\mathbf{v}}_{-D} = -\mathbf{v}_{-D}^{\mathsf{T}} \mathbf{\Gamma}(\mathbf{q}_{-D}) \mathbf{v}_{-D} - \mathbf{G}(\mathbf{q}_{-D})^{-1} \nabla_{\mathbf{q}_{-D}} \tilde{U}(\mathbf{q})$$
(15)

where $\Gamma(\mathbf{q}_{-D}) = r^{-2}\mathbf{G}(\mathbf{q}_{-D}) \otimes \mathbf{q}_{-D}$ is the Christoffel symbols of second kind (see details in Lan and Shahbaba, 2016, for r = 1). A splitting technique is applied to yield the following geometric integrator (Lan et al., 2014; Lan and Shahbaba, 2016), which also includes the last coordinates q_D, v_D :

$$\mathbf{v}^{-} = \mathbf{v} - \frac{h}{2} \mathcal{P}(\mathbf{q}) \mathbf{g}(\mathbf{q})$$

$$\begin{bmatrix} \mathbf{q}' \\ \mathbf{v}^{+} \end{bmatrix} = \begin{bmatrix} r & 0 \\ 0 & \|\mathbf{v}^{-}\|_{2} \end{bmatrix} \begin{bmatrix} \cos(\|\mathbf{v}^{-}\|_{2}r^{-1}h) + \sin(\|\mathbf{v}^{-}\|_{2}r^{-1}h) \\ -\sin(\|\mathbf{v}^{-}\|_{2}r^{-1}h) + \cos(\|\mathbf{v}^{-}\|_{2}r^{-1}h) \end{bmatrix} \begin{bmatrix} r^{-1} & 0 \\ 0 & \|\mathbf{v}^{-}\|_{2}^{-1} \end{bmatrix} \begin{bmatrix} \mathbf{q} \\ \mathbf{v}^{-} \end{bmatrix}$$

$$\mathbf{v}' = \mathbf{v}^{+} - \frac{h}{2} \mathcal{P}(\mathbf{q}') \mathbf{g}(\mathbf{q}')$$

$$(16)$$

where $\mathbf{g}(\mathbf{q}) := \nabla_{\mathbf{q}} \tilde{U}(\mathbf{q}), \ \mathcal{P}(\mathbf{q}) := \mathbf{I}_D - r^{-2} \mathbf{q} \mathbf{q}^{\mathsf{T}}.$ (16) defines a mapping $\mathcal{T}_h : (\mathbf{q}, \mathbf{v}) \mapsto (\mathbf{q}', \mathbf{v}')$. Denote $\|\mathbf{u}\|_{\mathcal{P}(\mathbf{q})}^2 := \mathbf{u}^{\mathsf{T}} \mathcal{P}(\mathbf{q}) \mathbf{u}$. After applying such integrator T times, a proposal $(\mathbf{q}_T, \mathbf{v}_T) = \mathcal{T}_h^T(\mathbf{q}_0, \mathbf{v}_0)$ is accepted with the following probability

$$a_{sphHMC} = 1 \wedge \exp(-\Delta E)$$

$$\Delta E = \tilde{U}(\mathbf{q}_T) - \tilde{U}(\mathbf{q}_0) - \frac{h^2}{8} \left[\|\mathbf{g}(\mathbf{q}_T)\|_{\mathcal{P}(\mathbf{q})}^2 - \|\mathbf{g}(\mathbf{q}_0)\|_{\mathcal{P}(\mathbf{q})}^2 \right]$$

$$- \frac{h}{2} \left[\langle \mathbf{v}_0, \mathbf{g}(\mathbf{q}_0) \rangle + \langle \mathbf{v}_T, \mathbf{g}(\mathbf{q}_T) \rangle \right] - h \sum_{\tau=1}^{T-1} \langle \mathbf{v}_\tau, \mathbf{g}(\mathbf{q}_\tau) \rangle$$
(17)

We can prove the following limiting result (Beskos et al., 2011).

Theorem 3.1. Let $h \to 0$ we have the following energy conservation

$$E(\mathbf{q}(T), \mathbf{v}(T)) - E(\mathbf{q}(0), \mathbf{v}(0)) = \tilde{U}(\mathbf{q}(T)) - \tilde{U}(\mathbf{q}(0)) - \int_0^T \langle \mathbf{v}(t), \mathbf{g}(\mathbf{q}(t)) \rangle dt = 0$$

Proof. See Section C in the supplementary file.

3.3 Adaptive Spherical HMC

There are two tuning parameters in HMC and its variants: the step size h and the number of integration (leapfrog) steps T. Hand tuning heavily relies on domain expertise and could be inefficient. Here, we adopt the 'No-U-Turn' idea from Hoffman and Gelman (2014) and introduce a novel adaptive algorithm that obviates manual tuning of these parameters.

First, for any given step size h, we adopt a rule for setting the number of leapfrog steps based on the same philosophy as 'No-U-Turn' (Hoffman and Gelman, 2014). The idea is to avoid waste of computation occurred (e.g. when the sampler backtracks on its trajectory) without breaking the detailed balance condition for the MCMC transition kernel. $S^{D-1}(r)$ is a compact manifold where any two points $\mathbf{q}(0), \mathbf{q}(t) \in S^{D-1}(r)$ have bounded geodesic distance πr . We adopt the stopping rule for the leapfrog when the sampler exits the orthant of the initial state, that is, the trajectory measured in geodesic distance is at least $\frac{\pi}{2}r$, which is equivalent to $\langle \mathbf{q}(0), \mathbf{q}(t) \rangle < 0$. On the other hand, this condition may not be satisfied within reasonable number of iterations because the geometric integrator (16) does not follow a geodesic (great circle) in general (only the middle part does), therefore we set some threshold T_{max} for the number of tests, and adopt the following 'Two-Orthants' (as the starting and end points occupy two orthants) rule for the number of leapfrogs:

$$T_{2orth} = \min_{\tau \in \{0, \dots, T_{\text{max}}\}} \{ \tau : \langle \mathbf{q}_0, \mathbf{q}_\tau \rangle < 0 \}$$
 (18)

Alternatively, one can stop the leapfrog steps in a *stochastic* way based on the geodesic distance travelled:

$$T_{stoch} = \min_{\tau} \{ \tau : Z_{\tau} = 0 \}, \quad Z_{\tau} \sim \text{Bern}(p_{\tau}), \quad p_{\tau} = \frac{r^{-2} \langle \mathbf{q}_0, \mathbf{q}_{\tau} \rangle + 1}{2}$$
 (19)

These stopping criteria are already time reversible, so the recursive binary tree as in 'No-U-Turn' algorithm (Hoffman and Gelman, 2014) is no longer needed.

Lastly, we adopt the *dual averaging* scheme (Nesterov, 2009) for the adaptation of step size h. See Hoffman and Gelman (2014) for more details. We summarize our *Adaptive Spherical Hamiltonian Monte Carlo (adp-SphHMC)* in the supplementary file.

To sample \mathbf{L} (or \mathbf{L}_t), we could update each row vector $\mathbf{l}_i \in \mathcal{S}_0^{i-1}$ according to (16) (in parallel), and accept/reject vech^T(\mathbf{L}) (or vech^T(\mathbf{L}_t)) simultaneously based on (17) in terms of the sum of total energy of all components. We refer to the resulting algorithm as Δ -Spherical HMC (Δ -SphHMC).

The computational complexity involving GP prior is $\mathcal{O}(N^3)$, and that of the likelihood evaluation is $\mathcal{O}(MD^2)$. MCMC updates of $\tilde{\mu}_{N\times D}$, $\tilde{\sigma}_{N\times D}$, $\tilde{\mathbf{L}}_{N\times D\times D}$ have complexity $\mathcal{O}(ND)$, $\mathcal{O}(ND)$ and $\mathcal{O}(ND^2)$ respectively. To scale up applications to larger dimension D, one could preliminarily classify data into groups, and arrange the corresponding blocks of their covariance/correlation matrix in some 'band' along the main diagonal assuming no correlation among groups. More specifically, we can assume \mathbf{L}_t is w-band lower triangular matrix for each time t, i.e. $l_{ij} = 0$ for i < j or $i - j \ge w$, then the resulting covariance/correlation matrix will be (2w-1)-banded. In this way the complexity of likelihood evaluation and updating $\tilde{\mathbf{L}}$ will be reduced to $\mathcal{O}(MwD)$ and $\mathcal{O}(NwD)$ resepctively. Therefore the total computational cost would scale linearly with the dimension D. This technique will be investigated in Section 4.2.3.

4 Simulation Studies

In this section, we use simulated examples to illustrate the advantage of our structured models for covariance. First, we consider the normal-inverse-Wishart problem. Since there is conjugacy and we know the true posterior, we use this to verify our method and investigate flexible priors in Section 2.2. Then we test our dynamical modeling method in Section 2.3 on a periodic process model. Our model manifests full flexibility compared to a state-of-the-art nonparametric covariance regression model based on latent factor process (Fox and Dunson, 2015).

4.1 Normal-inverse-Wishart Problem

Consider the following example involving inverse-Wishart prior

$$\mathbf{y}_n | \boldsymbol{\Sigma} \sim \mathcal{N}(\boldsymbol{\mu}_0, \boldsymbol{\Sigma}), \quad n = 1, \cdots, N$$

$$\boldsymbol{\Sigma} \sim \mathcal{W}_D^{-1}(\boldsymbol{\Psi}, \nu)$$
(20)

It is known that the posterior of $\Sigma|Y$ is still inverse-Wishart distribution:

$$\Sigma | \mathbf{Y} \sim \mathcal{W}_D^{-1} (\mathbf{\Psi} + (\mathbf{Y} - \boldsymbol{\mu}_0)(\mathbf{Y} - \boldsymbol{\mu}_0)^\mathsf{T}, \nu + N), \qquad \mathbf{Y} = [\mathbf{y}_1, \cdots, \mathbf{y}_N]^\mathsf{T}$$
 (21)




Figure 3: Marginal posterior densities of σ_{ij} in the normal-inverse-Wishart problem. Solid blue lines are estimates by Δ -SphHMC and dashed red lines are estimates by direct sampling. All densities are estimated with 10^6 samples.

We consider dimension D=3 and generate data \mathbf{Y} with $\boldsymbol{\mu}_0=\mathbf{0}, \quad \boldsymbol{\Sigma}=\boldsymbol{\Sigma}_0=\frac{1}{11}(\mathbf{I}+\mathbf{1}\mathbf{1}^\mathsf{T})$ for N=20 data points so that the prior is not overwhelmed by data.

4.1.1 Verification of Validity

Specifying conditional priors based on (8) in the structured model (6), we want to check the validity of our proposed method by comparing the posterior estimates using Δ -SphHMC agains the truth (21).

We sample $\tau := \log(\sigma)$ using standard HMC and \mathbf{U}^* using Δ -SphHMC. They are updated in Metropolis-Within-Gibbs scheme. 10^6 samples are collected after burning the first 10% and subsampling every 1 of 10. For each sample of τ and $\operatorname{vech}(\mathbf{U}^*)$, we calculate $\Sigma = \operatorname{diag}(e^{\tau})\mathbf{U}^*(\mathbf{U}^*)^{\mathsf{T}}\operatorname{diag}(e^{\tau})$. Marginal densities of entries in Σ are estimated with these samples and plotted against the results by direct sampling in Figure 3. Despite of sampling variance, these estimates closely match the results by direct sampling, indicating the validity of our proposed method.

4.1.2 Examining Flexibility of Priors

We have studied several spherical priors for the Cholesky factor of correlation matrix proposed in Section 2.2. Now we examine the flexibility of these priors in providing prior information for correlation with various parameter settings.

With the same data generated according to (20), we now consider the squared-Dirichlet




Figure 4: Marginal posterior, prior (induced from squared-Dirichlet distribution) densities of correlations and MLEs with different settings for concentration parameter α , estimated with 10^6 samples.

prior (10) for L in the structured model (6) with the following setting

$$\tau_{i} = \log(\sigma_{i}) \sim \mathcal{N}(0, 0.1^{2}), \quad i = 1, \cdots, D$$

$$\mathbf{l}_{i} \sim \operatorname{Dir}^{2}(\boldsymbol{\alpha}_{i}), \quad \boldsymbol{\alpha}_{i} = (\alpha \mathbf{1}_{i-1}, \alpha_{0}), \quad i = 2, \cdots, D$$
(22)

where we consider three cases i) $\alpha = 1$, $\alpha_0 = 1$; ii) $\alpha = 0.1$, $\alpha_0 = 1$; iii) $\alpha = 0.1$, $\alpha_0 = 10$. We generate 10^6 prior samples (according to (22)) and posterior samples (by Δ -SphHMC) for **L** respectively and covert them to $\mathbf{P} = \mathbf{L}\mathbf{L}^{\mathsf{T}}$. For each entry of ρ_{ij} , we estimate the marginal posterior (prior) density based on these posterior (prior) samples. The posteriors, priors and maximal likelihood estimates (MLEs) of correlations ρ_{ij} are plotted in Figure 4 for different α 's respectively. In general, the posteriors are compromise between priors and the likelihoods (MLEs). With more and more weight (through α) put around the poles (last component) of each factor sphere, the priors become increasingly dominant that the posteriors (red dash lines) almost fall on priors (blue solid lines) when $\alpha = (0.1, 0.1, 10)$. In this extreme case, the squared-Dirichlet distributions induce priors in favor of trivial (zero) correlations. We have similar conclusion on Bingham prior and von Mises-Fisher prior but results are reported in Section E.1 of the supplementary file.




Figure 5: Estimation of the underlying mean functions μ_t (left in each of 4 subpannels) and covariance functions Σ_t (right in each of 4 subpannels) of 2-dimensional periodic processes. M is the number of trials, and N is the number of discretization points. Dashed lines are true values, solid lines are estimates and shaded regions are 95% credible bands.

4.2 Simulated Periodic Processes

In this section, we investigate the performance of our dynamic model (11) on the following periodic process example

$$y(t) \sim \mathcal{N}_D(\mu(t), \Sigma(t)), \quad \Sigma(t) = L(t)L(t)^{\mathsf{T}} \circ S, \quad t \in [0, 2]$$

 $\mu_i(t) = \sin(it\pi/D), \quad L_{ij}(t) = (-1)^i \sin(it\pi/D)(-1)^j \cos(jt\pi/D), \quad j \le i = 1, \dots, D,$
 $S_{ij} = (|i-j|+1)^{-1}, \quad i, j = 1, \dots, D$

$$(23)$$

Based on the model (23), we generate M trials (process realizations) of data y at N evenly spaced points for t in [0,2], and therefore the whole data set $\{y(t)\}$ is an $M \times N \times D$ array. We first consider D=2 to investigate the posterior contraction phenomena and the model flexibility; then we consider D=100 over a shorter period [0,1] to show the scalability using the 'w-band' structure.

4.2.1 Posterior Contraction

Posterior contraction describes the phenomenon that the posterior concentrates on smaller and smaller neighborhood of the true parameter (function) given more and more data (van der Vaart and van Zanten, 2008a). We investigate such phenomena in both mean




Figure 6: Estimation of the underlying mean functions μ_t (left column), variance functions σ_t (middle column) and correlation function ρ_t (right column) of 2-dimensional periodic processes, using latent factor process model (upper row) and our flexible model (lower row), based on M=10 trials of data over N=200 evenly spaced points. Dashed lines are true values, solid lines are estimates and shaded regions are 95% credible bands.

functions and covariance functions in our model (11) using the following settings i)M = 10, N = 20; ii)M = 100, N = 20; iii)M = 10, N = 200; iv)M = 100, N = 200.

To fit the data using the model (11), we set s = 2, a = (1, 1, 1), $b = (0.1, 10^{-3}, 0.2)$, m = (0, 0, 0) for all settings, V = (1, 0.5, 1) for N = 20 and V = (1, 1, 0.3) for N = 200. We also add an additional nugget of $10^{-5}I_n$ to all the covariance kernel of GPs to ensure non-degeneracy. Following the procedure in Section 3.1, we run MCMC for 1.5×10^5 iterations, burn in the first 5×10^4 and subsample 1 for every 10. Based on the resulting 10^4 posterior samples, we estimate the underlying mean functions and covariance functions and plot the estimates in Figure 5.

Note in Figure 5, both M and N have effect on the amount of data information thereafter on the posterior contraction but the contraction rate may depend on them differently. Both mean and covariance functions have narrower credible bands for more discretization points N (comparing N=20 in the first row with N=200 for the second row). On the other hand, both posteriors contract further with more trials M (comparing M=10 in the first column agains M=100 for the second column). In general the posterior of mean function contracts to the truth faster than the posterior of covariance function. With M=100 trials and N=200 discretization points, both mean and covariance functions are almost recovered by the model (11).




Figure 7: Posterior estimation of the underlying correlation functions P_t (left) and its 2-norm distance to the truth (right) of 100-dimensional periodic processes with 2-band structure, based on M=100 trials of data over N=100 discretization points. Dashed lines are true values, solid lines are estimates and shaded regions are 95% credible bands.

4.2.2 Full Flexibility

Our method (11) grants full flexibility because it models mean, variance and correlation processes separately. This is particularly useful if they behave differently. It contrasts with latent factor based models that the mean and covariance processes together. One of the state-of-the-art models of this type is Bayesian nonparametric covariance regression (Fox and Dunson, 2015):

$$y(x) \sim \mathcal{N}_D(\mu(x), \Sigma(x)), \qquad \mu(x) = \Lambda(x)\psi(x), \quad \Sigma(x) = \Lambda(x)\Lambda(x)^\mathsf{T} + \Sigma_0$$
 (24)

We tweak the simulated example (23) for D=2 to let mean and correlation processes have higher frequency than variance processes, as shown in the dashed lines in Figure 6. We generate M=10 trials of data over N=200 evenly spaced points. In this case, the true mean processes $\mu(x)$ and true covariance processes $\Sigma(x)$ behave differently but are modeled with a common loading matrix $\Lambda(x)$ in model (24). This imposes difficulty on (24) to have a latent factor process $\psi(x)$ that could properly accommodate the heterogeneity in mean and covariance processes. Figure 6 shows that due to this reason, latent factor based model (24) (upper row) fails to generate satisfactory fit for all of the mean, variance and correlation processes. Our fully flexible model (11) (bottom row), on the contrary, successfully produces more accurate characterization for all of them. Note that this artificial example is used to demonstrate the flexibility of our dynamic model (11). For cases that are not as extreme, (24) may performance equally well. See more discussion in Section 6 and more details in Section E.2 in the supplementary file.

4.2.3 Scalability

Now we use the same simulation model (23) for D=100 dimensions to test the scalability of our dynamic model (11). However instead of the full covariance, we only consider a diagonal covariance matrix plus 4 non-zero off-diagonal entries $\sigma_{1,2}$ ($\sigma_{2,1}$) and $\sigma_{99,100}$ ($\sigma_{100,99}$). We focus on the correlation process in this example thus set $\mu_t \equiv 0$ and $\sigma_t \equiv 1$, for $t \in [0,1]$. More specifically when generating data $\{y_t\}$ with (23), if $i \notin \{2,100\}$ we set i-th rows $L_i = S_i = e_i$ with e_i being the i-th row of identity matrix.

To apply our dynamical model (11) in this setting, we let \mathbf{L}_t have 'w-band' structure with w=2 at each time t. Setting s=2, a=1, b=0.1, m=0 and $V=10^{-3}$, N=100 and M=100, we repeat the MCMC runs for 7.5×10^4 iterations, burn in the first 2.5×10^4 and subsample 1 for every 10 to obtain 5×10^3 posterior samples in the end. Based on those samples, we estimate the underlying correlation functions and only plot $\rho_{1,2}$, $\rho_{49,50}$ and $\rho_{99,100}$ in Figure 7. With the 'w-band' structure, we have less entries in the covariance matrix and focus on the 'in-group' correlation. Our dynamical model (11) is sensitive enough to discern the informative non-zero components from the non-informative ones in these correlation functions. Unit-vector GP priors provide flexibility for the model to capture the changing pattern of informative correlations. The left panel of Figure 7 shows that the model (11) correctly identify the non-zero components $\rho_{1,2}$ and $\rho_{99,100}$ and characterize their evolution. The right panel shows that the 2-norm distance between the estimated and true correlation matrices, $\|\hat{P}(t) - P(t)\|_2$, is small, indicating that our dynamic model (11) performs well with higher dimension in estimating complex dependence structure among multiple stochastic processes.

5 Analysis of Local Field Potential Activity

Now we use the proposed model (11) to analyze a local field potential (LFP) activity dataset. The goal of this analysis is to elucidate how memory encoding, retrieval and decision-making arise from functional interactions among brain regions, by modeling how their dynamic connectivity varies during performance of complex memory tasks. Here we focus on LFP activity data recorded from 24 electrodes spanning the dorsal CA1 subregion of the hippocampus as rats performed a sequence memory task (Allen et al., 2014, 2016; Ng et al., 2017; Holbrook et al., 2017). The task involves repeated presentations of a sequence of odors (e.g., ABCDE) at a single port and requires rats to correctly determine whether each odor is presented 'in sequence' (InSeq; e.g., ABCDE; by holding their nosepoke response until the signal at 1.2s) or 'out of sequence' (OutSeq; e.g., ABDDE; by withdrawing their nose before the signal). In previous work using the same dataset, Holbrook et al. (2016) used a direct MCMC algorithm to study the spectral density matrix of LFP from 4 selected channels. However, they did not examine how their correlations varied across time and recording site. These limitations are addressed in this paper.

We focus our analyses on the time window from 0ms to 750ms (with 0 corresponding




Figure 8: LFP signals on "in sequence" and "out of sequence" trials. It is difficult to identify differences between the two conditions based on a mere visual inspection of the LPFs.

to when the rat's nose enters the odor port). Critically, this includes a time period during which the behavior of the animal is held constant (0-500ms) so differences in LFP reflect the cognitive processes associated with task performance, and, to serve as a comparison, a time period near 750ms during which the behavioral state of the animal is known to be different (i.e., by 750ms the animal has already withdrawn from the port on the majority of OutSeq trials, but is still in the port on InSeq trials). We also focus our analyses on two sets of adjacent electrodes (electrodes 20 and 22, and electrodes 8 and 9), which allows for comparisons between probes that are near each other (<1mm; i.e., 20:22 and 8:9) or more distant from each other (>2mm; i.e., 20:8, 20:9, 22:8, and 22:9). Figure 8 shows M=20 trials of these LFP signals from D=4 channels under both InSeq and OutSeq conditions. Our main objective is to quantify how correlations among these LFP channels varied across trial types (InSeq vs OutSeq) and over time (within the first 750ms of trials). To do so, we discretize the time window of 0.75 seconds into N = 300 equally-spaced small intervals. Under each experiment condition (InSeq or OutSeq), we treat all the signals as a 4 dimensional time series and fit them using our proposed dynamic correlation model (11) in order to discover the evolution of their relationship. Note that we model the mean, variance, and correlation processes separately but only report findings about the evolution of correlation among those brain signals.

We set s = 2, a = (1,1,1), b = (1,0.1,0.2), m = (0,0,0), V = (1,1.2,2); and the general results are not very sensitive to the choice of these fine-tuning parameters. We also scale the discretized time points into (0,1] and add an additional nugget of $10^{-5}I_n$




Figure 9: Estimated correlation processes of LFPs (beta) under in-sequence condition (top), out-of-sequence condition (middle) and the (Frobenius) distance between two correlation matrices (bottom).

to the covariance kernel of GPs. We follow the same procedure in Section 3.1 to collect 7.5×10^4 samples, burn in the first 2.5×10^4 and subsample 1 for every 10. The resulting 10^4 samples yield estimates of correlation processes as shown in Figure 9 for beta-filtered traces (20-40Hz) but similar patterns were also observed for theta-filtered traces (4-12Hz; see the supplement). The bottom panel of Figure 9 shows the dissimilarity between correlation processes under different conditions measured by the Frobenius norm of their difference.

Our approach revealed many important patterns in the data. First, it showed that electrodes near each other (20:22 and 8:9) displayed remarkably high correlations in their LFP activity on InSeq and OutSeq trials, whereas correlations were considerably lower among more distant electrodes (20:8, 20:9, 22:8, and 22:9). Second, it revealed that the correlations between InSeq and OutSeq matrices evolved during the presentation of individual trials. These results are consistent with other analyses on learning (see, e.g., Fiecas and Ombao, 2016). As expected, InSeq and OutSeq activity was very similar at the beginning of the time window (e.g., before 350ms), which is before the animal has any information about the InSeq or OutSeq status of the presented odor, but maximally different at the end of the time window, which is after it has made its response on OutSeq trials. Most important, however, is the discovery of InSeq vs OutSeq differences before 500ms, which reveal changes in neural activity associated with the complex cognitive process of identifying if events occurred in their expected order. These findings highlight the sensitivity of our novel approach, as such differences have not been detected with traditional analyses. Interested readers can find more results about all the 12 channels in Section E.3 of the supplementary file.

6 Conclusion

In this paper, we propose a novel Bayesian framework that grants full flexibility in modeling covariance and correlation matrices. It extends the separation strategy proposed by Barnard et al. (2000) and uses the Cholesky decomposition to maintain the positive definiteness of the correlation matrix. By defining distributions on spheres, a large class of flexible priors can be induced for covariance matrix that go beyond the commonly used but restrictive inverse-Wishart distribution. Furthermore, the structured models we propose maintain the interpretability of covariance in terms of variance and correlation. Adaptive Δ -Spherical HMC is introduced to handle the intractability of the resulting posterior. Furthermore, we extend this structured scheme to dynamical models to capture complex dependence among multiple stochastic processes, and demonstrate the effectiveness and efficiency in Bayesian modeling covariance and correlation matrices using a normal-inverse-Wishart problem, a simulated periodic process, and an analysis of LFP data. In addition, we provide both theoretic characterization and empirical investigation of posterior contraction for dynamically covariance modeling, which to our best knowledge, is a first attempt.

In this work, we consider the marginal (pairwise) dependence among multiple stochastic processes. The priors for correlation matrix specified through the sphere-product representation are in general dependent among component variables. For example, the method we use to induce uncorrelated prior between y_i and y_j (i < j) by setting $l_{jk} \approx 0$ for $k \le i$ has a direct consequence that $Cor(y_{i'}, y_j) \approx 0$ for $i' \le i$. In another word, more informative priors (part of the components are correlated) may require careful ordering in $\{y_i\}$. To avoid this issue, one might consider the inverse of covariance (precision) matrices instead. This leads to modeling the conditional dependence, or Markov network (Dempster, 1972; Friedman et al., 2008). Our proposed methodology applies directly to (dynamic) precision matrices/processes, which will be our future direction.

To further scale our method to problems of greater dimensionality in future, one could explore the low-rank structure of covariance and correlation matrices, e.g. by adopting the similar factorization as in (Fox and Dunson, 2015) and assuming vech^T(\mathbf{L}_t) $\in (\mathcal{S}^k)^D$ for some $k \ll D$, or impose some sparse structure on the precision matrices.

We have proved that the posterior of covariance function contracts at a rate given by the general form of concentration function (van der Vaart and van Zanten, 2008a). Empirical evidence (Section 4.2.1) shows that the posterior of covariance contracts slower than that of mean. More theoretical works is needed to compare their contraction rates. Also, future research could involve investigating posterior contraction in covariance regression with respect to the optimal rates under different GP priors.

While our research has generated interesting new findings regarding brain signals during memory tasks, one limitation of our current analysis on LFP data is that it is conducted on a single rat. The proposed model can be generalized to account for variation among rats. In the future, we will apply this sensitive approach to other datasets, including simultaneous LFP recordings from multiple brain regions in rats as well as BOLD fMRI data collected

from human subjects performing the same task.

Acknowledgement

SL is supported by ONR N00014-17-1-2079. AH is supported by NIH grant T32 AG000096. GAE is supported by NIDCD T32 DC010775. NJF is supported by NSF awards IOS-1150292 and BCS-1439267 and Whitehall Foundation award 2010-05-84. HO is supported by NSF DMS 1509023 and NSF SES 1461534. BS is supported by NSF DMS 1622490 and NIH R01 AI107034. We thank Yulong Lu at Duke for discussions on posterior contraction of covariance regression with Gaussian processes.

References

- Allen, T. A., Morris, A. M., Mattfeld, A. T., Stark, C. E., and Fortin, N. J. (2014). A sequence of events model of episodic memory shows parallels in rats and humans. *Hippocampus*, 24(10):1178–1188.
- Allen, T. A., Salz, D. M., McKenzie, S., and Fortin, N. J. (2016). Nonspatial sequence coding in cal neurons. *Journal of Neuroscience*, 36(5):1547–1563.
- Anderson, T. W. (2003). An Introduction to Multivariate Statistical Analysis. Wiley Series in Probability and Statistics. Wiley Interscience, Hoboken, N. J.
- Banfield, J. D. and Raftery, A. E. (1993). Model-based gaussian and non-gaussian clustering. *Biometrics*, pages 803–821.
- Barnard, J., McCulloch, R., and Meng, X.-L. (2000). Modeling covariance matrices in terms of standard deviations and correlations, with application to shrinkage. *Statistica Sinica*, pages 1281–1311.
- Bensmail, H., Celeux, G., Raftery, A. E., and Robert, C. P. (1997). Inference in model-based cluster analysis. *statistics and Computing*, 7(1):1–10.
- Beskos, A., Pinski, F. J., Sanz-Serna, J. M., and Stuart, A. M. (2011). Hybrid monte carlo on hilbert spaces. *Stochastic Processes and their Applications*, 121(10):2201–2230.
- Bingham, C. (1974). An antipodally symmetric distribution on the sphere. *The Annals of Statistics*, pages 1201–1225.
- Byrne, S. and Girolami, M. (2013). Geodesic monte carlo on embedded manifolds. *Scandinavian Journal of Statistics*, 40(4):825–845.
- Celeux, G. and Govaert, G. (1995). Gaussian parsimonious clustering models. *Pattern recognition*, 28(5):781–793.
- Chiu, T. Y., Leonard, T., and Tsui, K.-W. (1996). The matrix-logarithmic covariance model. *Journal of the American Statistical Association*, 91(433):198–210.
- Cho, H. and Fryzlewicz, P. (2015). Multiple-change-point detection for high dimensional time series via sparsified binary segmentation. *Journal of the Royal Statistical Society:* Series B (Statistical Methodology), 77(2):475–507.
- Clausen, T. (1832). Über die Function $\sin \phi + (1/2^2) \sin 2\phi + (1/3^2) \sin 3\phi +$ etc. Journal für die reine und angewandte Mathematik, 8:298–300.

- Cribben, I., Haraldsdottir, R., Atlas, L. Y., Wager, T. D., and Lindquist, M. A. (2012). Dynamic connectivity regression: Determining state-related changes in brain connectivity. NeuroImage, 61(4):907 – 920.
- Dahlhaus, R. (2000). A likelihood approximation for locally stationary processes. *Ann. Statist.*, 28(6):1762–1794.
- Daniels, M. J. (1999). A prior for the variance in hierarchical models. *Canadian Journal of Statistics*, 27(3):567–578.
- Daniels, M. J. and Kass, R. E. (1999). Nonconjugate bayesian estimation of covariance matrices and its use in hierarchical models. *Journal of the American Statistical Association*, 94(448):1254–1263.
- Daniels, M. J. and Kass, R. E. (2001). Shrinkage estimators for covariance matrices. *Biometrics*, 57(4):1173–1184.
- Dempster, A. P. (1972). Covariance selection. *Biometrics*, 28(1):157–175.
- Duane, S., Kennedy, A. D., Pendleton, B. J., and Roweth, D. (1987). Hybrid Monte Carlo. *Physics Letters B*, 195(2):216 222.
- Fiecas, M. and Ombao, H. (2016). Modeling the evolution of dynamic brain processes during an associative learning experiment. *Journal of the American Statistical Association*, 111(516):1440–1453.
- Fisher, N. I., Lewis, T., and Embleton, B. J. (1987). Statistical analysis of spherical data. Cambridge university press.
- Fisher, R. (1953). Dispersion on a sphere. In *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, volume 217, pages 295–305. The Royal Society.
- Fox, E. B. and Dunson, D. B. (2015). Bayesian nonparametric covariance regression. Journal of Machine Learning Research, 16:2501–2542.
- Friedman, J., Hastie, T., and Tibshirani, R. (2008). Sparse inverse covariance estimation with the graphical lasso. *Biostatistics*, 9(3):432–441.
- Ghosal, S. and van der Vaart, A. (2007). Convergence rates of posterior distributions for noniid observations. *Ann. Statist.*, 35(1):192–223.
- Girolami, M. and Calderhead, B. (2011). Riemann manifold Langevin and Hamiltonian Monte Carlo methods. *Journal of the Royal Statistical Society, Series B*, (with discussion) 73(2):123–214.

- Hoffman, M. D. and Gelman, A. (2014). The no-u-turn sampler: Adaptively setting path lengths in hamiltonian monte carlo. *The Journal of Machine Learning Research*, 15(1):1593–1623.
- Holbrook, A., Lan, S., Vandenberg-Rodes, A., and Shahbaba, B. (2016). Geodesic lagrangian monte carlo over the space of positive definite matrices: with application to bayesian spectral density estimation. arXiv preprint arXiv:1612.08224.
- Holbrook, A., Vandenberg-Rodes, A., Fortin, N., and Shahbaba, B. (2017). A bayesian supervised dual-dimensionality reduction model for simultaneous decoding of lfp and spike train signals. *Stat*, 6(1):53–67. sta4.137.
- Kent, J. T. (1982). The fisher-bingham distribution on the sphere. *Journal of the Royal Statistical Society. Series B (Methodological)*, pages 71–80.
- Lan, S. and Shahbaba, B. (2016). Algorithmic Advances in Riemannian Geometry and Applications, chapter 2, pages 25–71. Advances in Computer Vision and Pattern Recognition. Springer International Publishing, 1 edition.
- Lan, S., Stathopoulos, V., Shahbaba, B., and Girolami, M. (2015). Markov chain monte carlo from lagrangian dynamics. *Journal of Computational and Graphical Statistics*, 24(2):357–378.
- Lan, S., Zhou, B., and Shahbaba, B. (2014). Spherical hamiltonian monte carlo for constrained target distributions. volume 32, pages 629–637, Beijing. The 31st International Conference on Machine Learning.
- Leonard, T. and Hsu, J. S. (1992). Bayesian inference for a covariance matrix. *The Annals of Statistics*, pages 1669–1696.
- Liechty, J. C. (2004). Bayesian correlation estimation. Biometrika, 91(1):1–14.
- Lindquist, M. A., Xu, Y., Nebel, M. B., and Caffo, B. S. (2014). Evaluating dynamic bivariate correlations in resting-state fmri: A comparison study and a new approach. *NeuroImage*, 101(Supplement C):531 546.
- Liu, C. (1993). Bartlett's decomposition of the posterior distribution of the covariance for normal monotone ignorable missing data. *Journal of Multivariate Analysis*, 46(2):198–206.
- Magnus, J. R. and Neudecker, H. (1979). The commutation matrix: some properties and applications. *The Annals of Statistics*, pages 381–394.
- Magnus, J. R. and Neudecker, H. (1980). The elimination matrix: some lemmas and applications. SIAM Journal on Algebraic Discrete Methods, 1(4):422–449.

- Mardia, K. V. and Jupp, P. E. (2009). *Directional statistics*, volume 494. John Wiley & Sons.
- Mardia, K. V., Kent, J. T., and Bibby, J. M. (1980). Multivariate analysis (probability and mathematical statistics).
- Minka, T. (1997; revised 12/00). Old and New Matrix Algebra Useful for Statistics. MIT Media Lab note.
- Murray, I., Adams, R. P., and MacKay, D. J. (2010). Elliptical slice sampling. *JMLR:* W&CP, 9:541–548.
- Nason, G. P., Von Sachs, R., and Kroisandt, G. (2000). Wavelet processes and adaptive estimation of the evolutionary wavelet spectrum. *Journal of the Royal Statistical Society:* Series B (Statistical Methodology), 62(2):271–292.
- Neal, R. M. (2003). Slice sampling. Annals of Statistics, 31(3):705–767.
- Neal, R. M. (2011). MCMC using Hamiltonian dynamics. In Brooks, S., Gelman, A., Jones, G., and Meng, X. L., editors, *Handbook of Markov Chain Monte Carlo*, pages 113–162. Chapman and Hall/CRC.
- Nesterov, Y. (2009). Primal-dual subgradient methods for convex problems. *Mathematical programming*, 120(1):221–259.
- Ng, C.-W., Elias, G. A., Asem, J. S., Allen, T. A., and Fortin, N. J. (2017). Nonspatial sequence coding varies along the cal transverse axis. *Behavioural Brain Research*.
- Ombao, H., von Sachs, R., and Guo, W. (2005). Slex analysis of multivariate nonstationary time series. *Journal of the American Statistical Association*, 100(470):519–531.
- Onstott, T. C. (1980). Application of the bingham distribution function in paleomagnetic studies. *Journal of Geophysical Research: Solid Earth*, 85(B3):1500–1510.
- Park, T., Eckley, I. A., and Ombao, H. C. (2014). Estimating time-evolving partial coherence between signals via multivariate locally stationary wavelet processes. *IEEE Transactions on Signal Processing*, 62(20):5240–5250.
- Pinheiro, J. C. and Bates, D. M. (1996). Unconstrained parametrizations for variance-covariance matrices. *Statistics and Computing*, 6(3):289–296.
- Pourahmadi, M. (1999). Joint mean-covariance models with applications to longitudinal data: Unconstrained parameterisation. *Biometrika*, pages 677–690.
- Pourahmadi, M. (2000). Maximum likelihood estimation of generalised linear models for multivariate normal covariance matrix. *Biometrika*, pages 425–435.

- Pourahmadi, M. and Wang, X. (2015). Distribution of random correlation matrices: Hyperspherical parameterization of the cholesky factor. Statistics & Probability Letters, 106:5-12.
- Prado, R. (2013). Sequential estimation of mixtures of structured autoregressive models. Computational Statistics & Data Analysis, 58(Supplement C):58 – 70. The Third Special Issue on Statistical Signal Extraction and Filtering.
- Prado, R., West, M., and Krystal, A. D. (2001). Multichannel electroencephalographic analyses via dynamic regression models with time-varying lag-lead structure. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, 50(1):95–109.
- Priestley, M. (1965). Evolutionary and non-stationary processes. J. Roy. Statist. Soc. Ser. B, 27:204–237.
- Rao, T. S. (1970). The fitting of non-stationary time-series models with time-dependent parameters. *Journal of the Royal Statistical Society. Series B (Methodological)*, 32(2):312–322.
- Rapisarda, F., Brigo, D., and Mercurio, F. (2007). Parameterizing correlations: a geometric interpretation. *IMA Journal of Management Mathematics*, 18(1):55–73.
- Smith, W. and Hocking, R. (1972). Algorithm as 53: Wishart variate generator. *Journal of the Royal Statistical Society. Series C (Applied Statistics)*, 21(3):341–345.
- Sverdrup, E. (1947). Derivation of the wishart distribution of the second order sample moments by straightforward integration of a multiple integral. *Scandinavian Actuarial Journal*, 1947(1):151–166.
- Ting, C. M., Seghouane, A. K., Salleh, S. H., and Noor, A. M. (2015). Estimating effective connectivity from fmri data using factor-based subspace autoregressive models. *IEEE Signal Processing Letters*, 22(6):757–761.
- Tokuda, T., Goodrich, B., Van Mechelen, I., Gelman, A., and Tuerlinckx, F. (2011). Visualizing distributions of covariance matrices. *Columbia Univ.*, New York, USA, Tech. Rep, pages 18–18.
- Tracy, D. S. and Dwyer, P. S. (1969). Multivariate maxima and minima with matrix derivatives. *Journal of the American Statistical Association*, 64(328):1576–1594.
- van der Vaart, A. and van Zanten, H. (2011). Information rates of nonparametric gaussian process methods. J. Mach. Learn. Res., 12:2095–2119.
- van der Vaart, A. W. and van Zanten, J. H. (2008a). Rates of contraction of posterior distributions based on gaussian process priors. *The Annals of Statistics*, 36(3):1435–1463.

- van der Vaart, A. W. and van Zanten, J. H. (2008b). Reproducing kernel Hilbert spaces of Gaussian priors, volume Volume 3 of Collections, pages 200–222. Institute of Mathematical Statistics, Beachwood, Ohio, USA.
- van der Vaart, A. W. and van Zanten, J. H. (2009). Adaptive bayesian estimation using a gaussian random field with inverse gamma bandwidth. *Ann. Statist.*, 37(5B):2655–2675.
- West, M., Prado, R., and Krystal, A. D. (1999). Evaluation and comparison of eeg traces: Latent structure in nonstationary time series. *Journal of the American Statistical Association*, 94(446):375–387.
- Wilson, A. G. and Ghahramani, Z. (2011). Generalised wishart processes. In *Proceedings* of the 27th Conference on Uncertainty in Artificial Intelligence.
- Wishart, J. (1928). The generalised product moment distribution in samples from a normal multivariate population. *Biometrika*, pages 32–52.
- Yang, R. and Berger, J. O. (1994). Estimation of a covariance matrix using the reference prior. *The Annals of Statistics*, pages 1195–1211.
- Yang, Y. and Dunson, D. B. (2016). Bayesian manifold regression. Ann. Statist., 44(2):876–905.

SUPPLEMENTARY MATERIAL

A Connection to Known Priors

The following lemma is essential in proving that our proposed methods (4) (6) generalize existing methods in specifying priors, including the inverse-Wishart distribution, and two uniform distributions (Barnard et al., 2000) as well.

Lemma A.1. Let $\Sigma = UU^T$ be the reversed Cholesky decomposition of Σ . The Jocobian of the transformation $\Sigma \mapsto U$ is

$$\left| \frac{d_h \Sigma}{d_h \mathbf{U}^\mathsf{T}} \right| := \left| \frac{\partial \mathrm{vech} \Sigma}{\partial \mathrm{vech} \mathbf{U}^\mathsf{T}} \right| = 2^D \prod_{i=1}^D |u_{ii}|^i$$

Let $\mathbf{P} = \mathbf{L}\mathbf{L}^\mathsf{T}$ be the Cholesky decomposition of \mathbf{P} . The Jacobian of the transformation $\mathbf{L} \mapsto \mathbf{P}$ is

$$\left| \frac{d_h \mathbf{L}}{d_h \mathbf{P}} \right| := \left| \frac{\partial \text{vech} \mathbf{L}}{\partial \text{vech} \mathbf{P}} \right| = 2^{-D} \prod_{i=1}^{D} |l_{ii}|^{i - (D+1)}$$

Proof. Note we have

$$d\mathbf{\Sigma} = d\mathbf{U}\mathbf{U}^\mathsf{T} + \mathbf{U}d\mathbf{U}^\mathsf{T}$$

Taking vec on both sides and applying its property

$$d\text{vec}\boldsymbol{\Sigma} = (\mathbf{U} \otimes \mathbf{I})d\text{vec}\mathbf{U} + (\mathbf{I} \otimes \mathbf{U})d\text{vec}\mathbf{U}^\mathsf{T}$$

Applying the elimination \mathbf{L}_D on both sides

$$d\text{vech}\boldsymbol{\Sigma} = \mathbf{L}_D[(\mathbf{U} \otimes \mathbf{I})\mathbf{K}_D d\text{vec}\mathbf{U}^\mathsf{T} + (\mathbf{I} \otimes \mathbf{U}) d\text{vec}\mathbf{U}^\mathsf{T}] = \mathbf{L}_D(\mathbf{K}_D + \mathbf{I})(\mathbf{I} \otimes \mathbf{U}) d\text{vec}\mathbf{U}^\mathsf{T}$$
$$= 2\mathbf{L}_D\mathbf{N}_D(\mathbf{I} \otimes \mathbf{U})\mathbf{L}_D^\mathsf{T} d\text{vech}\mathbf{U}^\mathsf{T} = 2\mathbf{L}_D\mathbf{N}_D\mathbf{L}_D^\mathsf{T}\mathbf{D}_D^\mathsf{T}(\mathbf{I} \otimes \mathbf{U})\mathbf{L}_D^\mathsf{T} d\text{vech}\mathbf{U}^\mathsf{T}$$

where \mathbf{K}_D is the *commutation matrix* such that $\mathbf{K}_D \text{vec} \mathbf{A} = \text{vec} \mathbf{A}^\mathsf{T}$ for matrix $\mathbf{A}_{D \times D}$, $\mathbf{N}_D := (\mathbf{K}_D + \mathbf{I})/2$, and \mathbf{D}_D is the *duplication matrix* which is regarded as the inverse of the elimination matrix \mathbf{L}_D . The last equation is by $\mathbf{D}_D \mathbf{L}_D \mathbf{N}_D = \mathbf{N}_D = \mathbf{N}_D^\mathsf{T}$ (Lemma2.1 and Lemma3.5 in Magnus and Neudecker, 1980). Thus according to (Lemma3.4 and Lemma4.1 in Magnus and Neudecker, 1980) we have

$$\left| \frac{d_h \mathbf{\Sigma}}{d_h \mathbf{U}^\mathsf{T}} \right| = \left| \frac{\partial \text{vech} \mathbf{\Sigma}}{\partial \text{vech} \mathbf{U}^\mathsf{T}} \right| = |2 \mathbf{L}_D \mathbf{N}_D \mathbf{L}_D^\mathsf{T} \mathbf{D}_D^\mathsf{T} (\mathbf{I} \otimes \mathbf{U}) \mathbf{L}_D^\mathsf{T}|$$
$$= 2^{D(D+1)/2} |\mathbf{L}_D \mathbf{N}_D \mathbf{L}_D^\mathsf{T}| |\mathbf{L}_D (\mathbf{I} \otimes \mathbf{U}^\mathsf{T}) \mathbf{D}_D| = 2^D \prod_{i=1}^D |u_{ii}^i|$$

By similar argument, we have

$$\begin{aligned} \left| \frac{d_h \mathbf{P}}{d_h \mathbf{L}} \right| &= \left| \frac{\partial \text{vech} \mathbf{P}}{\partial \text{vech} \mathbf{L}} \right| = |2 \mathbf{L}_D \mathbf{N}_D \mathbf{L}_D^\mathsf{T} \mathbf{D}_D^\mathsf{T} (\mathbf{L} \otimes \mathbf{I}) \mathbf{L}_D^\mathsf{T} | \\ &= 2^{D(D+1)/2} |\mathbf{L}_D \mathbf{N}_D \mathbf{L}_D^\mathsf{T} | |\mathbf{L}_D (\mathbf{L}^\mathsf{T} \otimes \mathbf{I}) \mathbf{D}_D| = 2^D \prod_{i=1}^D |l_{ii}|^{D+1-i} . \end{aligned}$$

Thus it completes the proof.

Proof of Proposition 2.1. We know that the density of $\Sigma \sim W_D^{-1}(\Psi, \nu)$ is

$$p_{\mathcal{W}^{-1}}(\boldsymbol{\varSigma};\boldsymbol{\Psi},\nu) = \frac{|\boldsymbol{\Psi}|^{\nu/2}}{2^{D\nu/2}\Gamma_D(\nu/2)} |\boldsymbol{\varSigma}|^{-(\nu+D+1)/2} \exp\left(-\frac{1}{2}\mathrm{tr}(\boldsymbol{\Psi}\boldsymbol{\varSigma}^{-1})\right)$$

By Lemma A.1 we have

$$\begin{split} p(\mathbf{U}) &= p(\boldsymbol{\Sigma}) \left| \frac{d_h \boldsymbol{\Sigma}}{d_h \mathbf{U}^\mathsf{T}} \right| = 2^D p_{\mathcal{W}^{-1}} (\mathbf{U} \mathbf{U}^\mathsf{T}; \boldsymbol{\Psi}, \nu) \prod_{i=1}^D |u_{ii}^i| \\ &= \frac{|\boldsymbol{\Psi}|^{\nu/2}}{2^{D(\nu-2)/2} \Gamma_D(\nu/2)} |\mathbf{U}|^{-(\nu+D+1)} \prod_{i=1}^D u_{ii}^i \exp\left(-\frac{1}{2} \mathrm{tr}(\boldsymbol{\Psi} \mathbf{U}^{-\mathsf{T}} \mathbf{U}^{-1})\right) \,. \end{split}$$

Then the proof is completed.

Proof of Theorem 2.1. To prove the first result, we use Lemma A.1

$$p(\mathbf{P}) = p(\mathbf{L}) \left| \frac{d_h \mathbf{L}}{d_h \mathbf{P}} \right| \propto \prod_{i=2}^{D} |\mathbf{l}_i|^{2\alpha_i - 1} \prod_{i=1}^{D} |l_{ii}|^{i - (D+1)} = \prod_{i=1}^{D} |l_{ii}|^{(i-3)(D+1)}$$

On the other hand, from Equation (8) in Barnard et al. (2000), we have the density of marginally uniform distribution:

$$p(\mathbf{P}) \propto |\mathbf{P}|^{\frac{D(D-1)}{2}-1} (\prod_{i} \mathbf{P}_{ii})^{-\frac{D+1}{2}} = (\prod_{j=1}^{D} l_{jj}^{2})^{\frac{D(D-1)}{2}-1} (\prod_{j=1}^{D} \prod_{i=1}^{j} l_{ii}^{2})^{-\frac{D+1}{2}} = \prod_{j=1}^{D} |l_{jj}|^{(j-3)(D+1)}$$

where \mathbf{P}_{ii} is the *i*-th principal minor of \mathbf{P} . Similarly by Lemma A.1 we can prove the second result

$$p(\mathbf{P}) = p(\mathbf{L}) \left| \frac{d_h \mathbf{L}}{d_h \mathbf{P}} \right| \propto \prod_{i=2}^{D} |\mathbf{l}_i|^{2\alpha_i - 1} \prod_{i=1}^{D} |l_{ii}|^{i - (D+1)} \propto 1.$$

Therefore we have finished the proof.

B Posterior Contraction

For the Gaussian likelihood models $p_i \sim \mathcal{N}(\boldsymbol{\mu}_i(t), \boldsymbol{\Sigma}_i(t))$ for i = 0, 1, we first bound the Hellinger distance, Kullback-Leibler distance and variance distance $V(p_0, p_1) = \mathrm{E}_0(\log(p_0/p_1))^2$ with the uniform norm in the following lemma. Notation \lesssim means "smaller than or equal to a universal constant times".

Lemma B.1. For any bounded measurable functions $\Sigma_i : \mathcal{X} \to \mathbb{R}^{D^2}$ with Cholesky decompositions $\Sigma_i = \mathbf{L}_i \mathbf{L}_i^\mathsf{T}$, assume $\min_{1 \le j \le D} \inf_{t \in \mathcal{X}} |l_{i,jj}(t)| \ge c_0 > 0$, i = 0, 1. Then we have

- $h(p_0, p_1) \lesssim \|\mathbf{L}_0 \mathbf{L}_1\|_{\infty}^{\frac{1}{2}}$
- $K(p_0, p_1) \lesssim ||\mathbf{L}_0 \mathbf{L}_1||_{\infty}$
- $V(p_0, p_1) \lesssim \|\mathbf{L}_0 \mathbf{L}_1\|_{\infty}^2$

Proof. First we calculate

$$\log p_{0} - \log p_{1} = \frac{1}{2} \left\{ \log \frac{|\boldsymbol{\Sigma}_{1}|}{|\boldsymbol{\Sigma}_{0}|} + (\mathbf{y} - \boldsymbol{\mu}_{*})^{\mathsf{T}} \boldsymbol{\Sigma}_{*}^{-1} (\mathbf{y} - \boldsymbol{\mu}_{*}) + ** \right\}$$

$$\boldsymbol{\Sigma}_{*}^{-1} = \boldsymbol{\Sigma}_{1}^{-1} - \boldsymbol{\Sigma}_{0}^{-1}, \quad \boldsymbol{\mu}_{*} = \boldsymbol{\Sigma}_{*} (\boldsymbol{\Sigma}_{1}^{-1} \boldsymbol{\mu}_{1} - \boldsymbol{\Sigma}_{0}^{-1} \boldsymbol{\mu}_{0})$$

$$** = -(\boldsymbol{\mu}_{1} - \boldsymbol{\mu}_{0})^{\mathsf{T}} \boldsymbol{\Sigma}_{1}^{-1} \boldsymbol{\Sigma}_{*} \boldsymbol{\Sigma}_{0}^{-1} (\boldsymbol{\mu}_{1} - \boldsymbol{\mu}_{0})$$
(B.1)

Taking expectation of (B.1) with respect to p_0 yields the following Kullback-Leibler divergence

$$K(p_0, p_1) = \frac{1}{2} \left\{ \operatorname{tr}(\boldsymbol{\Sigma}_1^{-1} \boldsymbol{\Sigma}_0) + (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_0)^\mathsf{T} \boldsymbol{\Sigma}_1^{-1} (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_0) - D + \log \frac{|\boldsymbol{\Sigma}_1|}{|\boldsymbol{\Sigma}_0|} \right\}$$

Consider $\mu_i \equiv 0$. By the non-negativity of K-L divergence we have for general $\Sigma_i > 0$,

$$\log \frac{|\boldsymbol{\Sigma}_0|}{|\boldsymbol{\Sigma}_1|} \le \operatorname{tr}(\boldsymbol{\Sigma}_1^{-1}\boldsymbol{\Sigma}_0 - \mathbf{I})$$
(B.2)

Therefore we can bound K-L divergence

$$K(p_0, p_1) \leq \frac{1}{2} \{ \operatorname{tr}(\boldsymbol{\Sigma}_1^{-1} \boldsymbol{\Sigma}_0 - \mathbf{I}) + \operatorname{tr}(\boldsymbol{\Sigma}_0^{-1} \boldsymbol{\Sigma}_1 - \mathbf{I}) \} \leq C(D, c_0) \|\boldsymbol{\Sigma}_0 - \boldsymbol{\Sigma}_1\|_{\infty} \lesssim \|\mathbf{L}_0 - \mathbf{L}_1\|_{\infty}$$

where we bound each term involving trace

$$\operatorname{tr}(\boldsymbol{\Sigma}_{1}^{-1}\boldsymbol{\Sigma}_{0} - \mathbf{I}) = \operatorname{tr}(\boldsymbol{\Sigma}_{1}^{-1}(\boldsymbol{\Sigma}_{0} - \boldsymbol{\Sigma}_{1})) \leq \|\boldsymbol{\Sigma}_{1}^{-1}\|_{F} \|\boldsymbol{\Sigma}_{0} - \boldsymbol{\Sigma}_{1}\|_{F}$$

$$\leq D^{3/2} \|\boldsymbol{\Sigma}_{1}^{-1}\|_{2} \|\boldsymbol{\Sigma}_{0} - \boldsymbol{\Sigma}_{1}\|_{\infty} \lesssim c_{0}^{-2D} (\|\mathbf{L}_{0}\|_{\infty} + \|\mathbf{L}_{1}\|_{\infty}) \|\mathbf{L}_{0} - \mathbf{L}_{1}\|_{\infty}$$
(B.3)

Note the last inequality holds because

$$c_0^D \leq \Pi_{j=1}^D |l_{jj}| = \left[\Pi_{j=1}^D \lambda_j(\boldsymbol{\Sigma}) \right]^{\frac{1}{2}} \leq \lambda_{\min}^{\frac{1}{2}} \left[\frac{\operatorname{tr}(\boldsymbol{\Sigma})^{D-1}}{D-1} \right]^{\frac{1}{2}} \leq \lambda_{\min}^{\frac{1}{2}} \left[\frac{\|\mathbf{L}\|_F^{2(D-1)}}{D-1} \right]^{\frac{1}{2}}$$
$$\leq \lambda_{\min}^{\frac{1}{2}} \left[\frac{D^{2(D-1)}}{D-1} \right]^{\frac{1}{2}} \|\mathbf{L}\|_{\infty}^{D-1} = \|\boldsymbol{\Sigma}_1^{-1}\|_2^{-\frac{1}{2}} \left[\frac{D^{2(D-1)}}{D-1} \right]^{\frac{1}{2}} \|\mathbf{L}\|_{\infty}^{D-1}$$

and we can write

$$\boldsymbol{\Sigma}_0 - \boldsymbol{\Sigma}_1 = \frac{1}{2} \left[(\mathbf{L}_0 - \mathbf{L}_1)(\mathbf{L}_0 + \mathbf{L}_1)^\mathsf{T} + (\mathbf{L}_0 + \mathbf{L}_1)(\mathbf{L}_0 - \mathbf{L}_1)^\mathsf{T} \right]$$

Now take expectation of squared (B.1) with respect with p_0 to get the following variance distance

$$V(p_0, p_1) = \frac{1}{2} \text{tr}((\boldsymbol{\Sigma}_1^{-1} \boldsymbol{\Sigma}_0 - \mathbf{I})^2) + (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_0)^{\mathsf{T}} \boldsymbol{\Sigma}_1^{-1} \boldsymbol{\Sigma}_0 \boldsymbol{\Sigma}_1^{-1} (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_0) + K^2(p_0, p_1)$$

Consider $\mu_i \equiv 0$ and we can bound the variance distance by similar argument as (B.3)

$$V(p_0, p_1) \le \frac{1}{2} \|\boldsymbol{\Sigma}_1^{-1} (\boldsymbol{\Sigma}_0 - \boldsymbol{\Sigma}_1)\|_F^2 + K^2(p_0, p_1) \le C \|\boldsymbol{\Sigma}_1^{-1}\|_F^2 \|\boldsymbol{\Sigma}_0 - \boldsymbol{\Sigma}_1\|_F^2 + K^2(p_0, p_1) \lesssim \|\mathbf{L}_0 - \mathbf{L}_1\|_{\infty}^2$$

where we use the fact $\|\mathbf{A}\mathbf{B}\|_F \leq \|\mathbf{A}\|_F \|\mathbf{B}\|_F$.

Lastly, the squared Hellinger distance for multivariate Gaussians can be calculated

$$h^{2}(p_{0}, p_{1}) = 1 - \frac{|\boldsymbol{\Sigma}_{0}\boldsymbol{\Sigma}_{1}|^{1/4}}{\left|\frac{\boldsymbol{\Sigma}_{0} + \boldsymbol{\Sigma}_{1}}{2}\right|^{1/2}} \exp \left\{-\frac{1}{8}(\boldsymbol{\mu}_{0} - \boldsymbol{\mu}_{1})^{\mathsf{T}} \left(\frac{\boldsymbol{\Sigma}_{0} + \boldsymbol{\Sigma}_{1}}{2}\right)^{-1} (\boldsymbol{\mu}_{0} - \boldsymbol{\mu}_{1})\right\}$$

Consider $\mu_i \equiv 0$. Notice that $1 - x \leq -\log x$, and by (B.2) we can bound the squared Hellinger distance using similar argument in (B.3)

$$h^{2}(p_{0}, p_{1}) \leq \log \frac{\left|\frac{\boldsymbol{\Sigma}_{0} + \boldsymbol{\Sigma}_{1}}{2}\right|^{1/2}}{|\boldsymbol{\Sigma}_{0} \boldsymbol{\Sigma}_{1}|^{1/4}} \leq \frac{1}{2} \operatorname{tr}(\mathbf{L}_{0}^{-\mathsf{T}} \mathbf{L}_{1}^{-1} (\boldsymbol{\Sigma}_{0} + \boldsymbol{\Sigma}_{1})/2 - \mathbf{I})$$
$$= \frac{1}{4} \left\{ \operatorname{tr}(\mathbf{L}_{1}^{-1} \mathbf{L}_{0} - \mathbf{I}) + \operatorname{tr}(\mathbf{L}_{0}^{-\mathsf{T}} \mathbf{L}_{1}^{\mathsf{T}} - \mathbf{I}) \right\} \lesssim c_{0}^{-D} \|\mathbf{L}_{0} - \mathbf{L}_{1}\|_{\infty}$$

where we use $\|\mathbf{L}^{-1}\|_2 = \lambda_{\min}^{-1}(\mathbf{L}) \le c_0^{-D} \left[\frac{D\|\mathbf{L}\|_{\infty}}{D-1} \right]^{D-1}$.

Define the following coordinate concentration function as in (12)

$$\phi_{l_{0,ij}}(\varepsilon) = \inf_{h_{ij} \in \mathbb{H}_{ij} : \|h_{ij} - l_{0,ij}\|_{ij} < \varepsilon} \|h\|_{\mathbb{H}_{ij}}^2 - \log \Pi(l_{ij} : \|l_{ij}\|_{ij} < \varepsilon)$$
(B.4)

It is easy to see that $\phi_{l_{0,ij}}(\varepsilon) \leq \phi_{\mathbf{L}_0}(\varepsilon)$ for $\forall 1 \leq j \leq i \leq D$. Let $\|\cdot\| := \max_{i,j} \|\cdot\|_{ij}$. For $\varepsilon > 0$, let $N(\varepsilon, B, d)$ denote the minimum number of balls of radius ε that a cover B in a metric space with metric d, which is named ε -covering number for B. Now we are ready to prove the theorem of posterior concentration.

Proof of Theorem 2.2. We use Theorem 4 of Ghosal and van der Vaart (2007) and it suffices to verify three conditions (the entropy condition 3.2, the complementary asertion 3.3 and the prior mass condition 3.4) as follows:

$$\sup_{\varepsilon > \varepsilon_n} \log N(\varepsilon/36, \{ \mathbf{L} \in \Theta_n : d_n(\mathbf{L}, \mathbf{L}_0) < \varepsilon \}, d_n) \le n\varepsilon_n^2$$
(B.5)

$$\frac{\Pi_n(\Theta \backslash \Theta_n)}{\Pi_n(\overline{B}_n(\mathbf{L}_0, \varepsilon_n))} = o(e^{-2n\varepsilon_n^2})$$
 (B.6)

$$\frac{\Pi_n(\mathbf{L} \in \Theta_n : \kappa \varepsilon_n < d_{n,H}(\mathbf{L}, \mathbf{L}_0) < 2\kappa \varepsilon_n)}{\Pi_n(\overline{B}_n(\mathbf{L}_0, \varepsilon_n))} \le e^{n\varepsilon_n^2 \kappa^2/4}, \text{ for large } \kappa$$
 (B.7)

where $\overline{B}_n(\mathbf{L}_0, \varepsilon) := \{ \mathbf{L} \in \Theta : \frac{1}{n} \sum_{i=1}^n K_i(\mathbf{L}_0, \mathbf{L}) \le \varepsilon^2, \frac{1}{n} \sum_{i=1}^n V_i(\mathbf{L}_0, \mathbf{L}) \le \varepsilon^2 \}, \text{ with } \Theta = L^{\infty}(\mathcal{X})^{D(D+1)/2}, K_i(\mathbf{L}_0, \mathbf{L}) = K(P_{\mathbf{L}_0,i}, P_{\mathbf{L},i}) \text{ and } V_i(\mathbf{L}_0, \mathbf{L}) = V(P_{\mathbf{L}_0,i}, P_{\mathbf{L},i}).$

Applying Theorem 2.1 of van der Vaart and van Zanten (2008a) to each Gaussian random element l_{ij} in $\mathbb{B}_{ij} = L^{\infty}(\mathcal{X})$ for $1 \leq j \leq i \leq D$, with $l_{0,ij} \in \overline{\mathbb{H}}_{ij}$, we have C > 2 and the measurable set $B_{n,ij} \subset \mathbb{B}_{ij}$ such that

$$\log N(3\varepsilon_n, B_{n,ij}, \|\cdot\|_{\infty}) \le 6Cn\varepsilon_n^2 \tag{B.8}$$

$$\Pi(l_{ij} \notin B_{n,ij}) \le e^{-Cn\varepsilon_n^2} \tag{B.9}$$

$$\Pi(\|l_{ij} - l_{0,ij}\|_{\infty} < 2\varepsilon_n) \ge e^{-n\varepsilon_n^2}$$
(B.10)

Now set $\Theta_n = \{ \mathbf{L} : l_{ij} \in B_{n,ij} \}$, and $N(\varepsilon_n, \Theta_n, d_n) = \max_{1 \leq j \leq i \leq D} N(3\varepsilon_{n,ij}, B_{n,ij}, \| \cdot \|_{\infty})$. By Lemma B.1 and (B.8), we have the following global entropy bound because $d_n^2(\mathbf{L}, \mathbf{L}') \leq \|\mathbf{L} - \mathbf{L}'\|_{\infty} \leq \varepsilon_n^2$ for $\forall \mathbf{L}, \mathbf{L}' \in \Theta_n$.

$$\log N(\varepsilon_n, \Theta_n, d_n) \le 6Cn(\varepsilon_n^2)^2 \le C'n\varepsilon_n^4 \le n\varepsilon^2$$

which is stronger than the local entropy condition (B.5). Now by Lemma B.1 and (B.10) we have

$$\Pi_n(\overline{B}_n(\mathbf{L}_0, \varepsilon_n)) \ge \Pi_n(\|\mathbf{L}_0 - \mathbf{L}\|_{\infty} \le \varepsilon_n^2, \|\mathbf{L}_0 - \mathbf{L}\|_{\infty}^2 \le \varepsilon_n^2) = \Pi_n(\|\mathbf{L}_0 - \mathbf{L}\|_{\infty} \le \varepsilon_n^2)
= \Pi(\|l_{\operatorname{argmax}} - l_{0,\operatorname{argmax}}\|_{\infty} < \varepsilon_n^2) \ge e^{-n(\varepsilon_n^2/2)^2} = e^{-n\varepsilon_n^4/4} \ge e^{-n\varepsilon_n^2\kappa^2/4}$$

Then (B.7) is immediately satisfied because the numerator is bounded by 1. Finally, by (B.9) we have

$$\Pi_n(\Theta \backslash \Theta_n) \le \sum_{i=1}^D \sum_{j=1}^i \Pi(l_{ij} \notin B_{n,ij}) \le \frac{D(D+1)}{2} e^{-Cn\varepsilon_n^2} = o(e^{-2n\varepsilon_n^2})$$

Then (B.6) holds because the denominator is bounded below by a term $(e^{-n\varepsilon_n^4/4})$ of smaller order. Therefore the proof is completed.

C Spherical Hamiltonian Monte Carlo

C.1 Derivation of the geometric integrator for SphHMC

The Lagrangian dynamics (15) on the sphere $S^{D-1}(r)$ with the first (D-1) coordinates can be split into the following two smaller dynamics:

$$\begin{cases} \dot{\mathbf{q}}_{-D} = \mathbf{v}_{-D} \\ \dot{\mathbf{v}}_{-D} = -\mathbf{v}_{-D}^{\mathsf{T}} \mathbf{\Gamma}(\mathbf{q}_{-D}) \mathbf{v}_{-D} \end{cases}$$

$$\begin{cases} \dot{\mathbf{q}}_{-D} = \mathbf{0} \\ \dot{\mathbf{v}}_{-D} = -\nabla_{\mathbf{q}_{-D}}^{-1} \tilde{U}(\mathbf{q}_{-D}) \end{cases}$$
(C.1b)

where (C.1a) is the equation of geodesic on manifold S^D which has analytical solution; and (C.1b) has analytical solution. Both define volume preserving maps.

The mapping $\mathcal{I}: \mathbf{q}_{-D} \mapsto \mathbf{q} = (\mathbf{q}_{-D}, q_D)$ can be viewed as an imbedding of \mathcal{S}_+^{D-1} into

 \mathbb{R}^D . Denote its Jacobian as $d\mathcal{I}(\mathbf{q}) := \begin{bmatrix} \mathbf{I}_{D-1} \\ -\frac{\mathbf{q}^{\mathsf{T}}_{D}}{q_D} \end{bmatrix}$. Then we have

$$d\mathcal{I}(\mathbf{q})^{\mathsf{T}}d\mathcal{I}(\mathbf{q}) = \mathbf{G}(\mathbf{q}_{-D}), \qquad d\mathcal{I}(\mathbf{q})\mathbf{G}(\mathbf{q}_{-D})^{-1}d\mathcal{I}(\mathbf{q})^{\mathsf{T}} = \mathcal{P}(\mathbf{q}) = \mathbf{I} - r^{-2}\mathbf{q}\mathbf{q}^{\mathsf{T}}$$
$$\nabla_{\mathbf{q}_{-D}}\tilde{U}(\mathbf{q}) = d\mathcal{I}(\mathbf{q})^{\mathsf{T}}\nabla_{\mathbf{q}}\tilde{U}(\mathbf{q}), \qquad \mathbf{v} = d\mathcal{I}(\mathbf{q})\mathbf{v}_{-D}, \qquad \mathbf{v}^{\mathsf{T}}\mathbf{v} = \mathbf{v}_{-D}^{\mathsf{T}}\mathbf{G}(\mathbf{q}_{-D})\mathbf{v}_{-D}$$

Then Equation (C.1a) has the following solution with full coordinates

$$\begin{bmatrix} \mathbf{q}(t) \\ \mathbf{v}(t) \end{bmatrix} = \begin{bmatrix} \mathbf{I} & \mathbf{0} \\ \mathbf{0}^{\mathsf{T}} & r^{-1} \| \mathbf{v}(0) \|_{2} \end{bmatrix} \begin{bmatrix} \cos(r^{-1} \| \mathbf{v}(0) \|_{2}t) & \sin(r^{-1} \| \mathbf{v}(0) \|_{2}t) \\ -\sin(r^{-1} \| \mathbf{v}(0) \|_{2}t) & \cos(r^{-1} \| \mathbf{v}(0) \|_{2}t) \end{bmatrix} \begin{bmatrix} \mathbf{I} & \mathbf{0} \\ \mathbf{0}^{\mathsf{T}} & r \| \mathbf{v}(0) \|_{2}^{-1} \end{bmatrix} \begin{bmatrix} \mathbf{q}(0) \\ \mathbf{v}(0) \end{bmatrix} \\
= \begin{bmatrix} \mathbf{q}(0) \cos(r^{-1} \| \mathbf{v}(0) \|_{2}t) + r\mathbf{v}(0) \| \mathbf{v}(0) \|_{2}^{-1} \sin(r^{-1} \| \mathbf{v}(0) \|_{2}t) \\ -r^{-1}\mathbf{q}(0) \| \mathbf{v}(0) \|_{2} \sin(r^{-1} \| \mathbf{v}(0) \|_{2}t) + \mathbf{v}(0) \cos(r^{-1} \| \mathbf{v}(0) \|_{2}t) \end{bmatrix} \tag{C.2}$$

and Equation (C.1b) has the following solution in full coordinates

$$\mathbf{q}(t) = \mathbf{q}(0)$$

$$\mathbf{v}(t) = \mathbf{v}(0) - \frac{t}{2}d\mathcal{I}(\mathbf{q}(0))\nabla_{\mathbf{q}_{-D}}^{-1}\tilde{U}(\mathbf{q}(0)) = \mathbf{v}(0) - \frac{t}{2}\mathcal{P}(\mathbf{q})\nabla_{\mathbf{q}}\tilde{U}(\mathbf{q}(0))$$
(C.3)

So numerically updating (C.3) for h/2, updating (C.2) for h and updating (C.3) for another h/2 yield the integrator (16).

C.2 Reformulating Acceptance

At the end of the numerical simulation, a proposal $(\mathbf{q}_T, \mathbf{v}_T)$ is accepted according to the following probability

$$a_{sphHMC} = 1 \wedge \exp(-\Delta E), \qquad \Delta E = E(\mathbf{q}_T, \mathbf{v}_T) - E(\mathbf{q}_0, \mathbf{v}_0)$$
 (C.4)

Such classic definition of acceptance probability can be reformulated by replacing ΔE in (C.4) with

$$\Delta E = \sum_{\tau=1}^{T} \Delta E_{\tau}$$
 $\Delta E_{\tau} = E(\mathbf{q}_{\tau}, \mathbf{v}_{\tau}) - E(\mathbf{q}_{\tau-1}, \mathbf{v}_{\tau-1})$

With (16) we can write

$$\begin{split} &\Delta E' = E(\mathbf{q}', \mathbf{v}') - E(\mathbf{q}, \mathbf{v}) \\ &= \tilde{U}(\mathbf{q}') - \tilde{U}(\mathbf{q}) + \frac{1}{2} \mathbf{v}'^{\mathsf{T}}_{-D} \mathbf{G}(\mathbf{q}'_{-D}) \mathbf{v}'_{-D} - \frac{1}{2} \mathbf{v}^{\mathsf{T}}_{-D} \mathbf{G}(\mathbf{q}_{-D}) \mathbf{v}_{-D} \\ &= \Delta \tilde{U} - \frac{1}{2} \|\mathbf{v}\|_2^2 + \frac{1}{2} \|\mathbf{v}^+ - \frac{h}{2} \mathcal{P}(\mathbf{q}') \nabla_{\mathbf{q}} \tilde{U}(\mathbf{q}') \|_2^2 \\ &= \Delta \tilde{U} - \frac{1}{2} \|\mathbf{v}\|_2^2 + \frac{1}{2} \mathbf{v}^{+\mathsf{T}} \mathbf{v}^+ - \frac{h}{2} \mathbf{v}^{+\mathsf{T}} \mathcal{P}(\mathbf{q}') \nabla_{\mathbf{q}} \tilde{U}(\mathbf{q}') + \frac{h^2}{8} \nabla_{\mathbf{q}} \tilde{U}(\mathbf{q}')^{\mathsf{T}} \mathcal{P}(\mathbf{q}') \nabla_{\mathbf{q}} \tilde{U}(\mathbf{q}') \\ &= \Delta \tilde{U} - \frac{1}{2} \|\mathbf{v}\|_2^2 + \frac{1}{2} \|\mathbf{v}^-\|_2^2 - \frac{h}{2} \mathbf{v}^{+\mathsf{T}} \nabla_{\mathbf{q}} \tilde{U}(\mathbf{q}') + \frac{h^2}{8} \|\nabla_{\mathbf{q}} \tilde{U}(\mathbf{q}')\|_{\mathcal{P}(\mathbf{q}')}^2 \\ &= \Delta \tilde{U} - \frac{1}{2} \|\mathbf{v}\|_2^2 - \frac{h}{2} \mathbf{v}^{+\mathsf{T}} \nabla_{\mathbf{q}} \tilde{U}(\mathbf{q}') + \frac{h^2}{8} \|\nabla_{\mathbf{q}} \tilde{U}(\mathbf{q}')\|_{\mathcal{P}(\mathbf{q}')}^2 + \frac{1}{2} \|\mathbf{v}\|_2^2 - \frac{h}{2} \mathbf{v}^{\mathsf{T}} \nabla_{\mathbf{q}} \tilde{U}(\mathbf{q}) + \frac{h^2}{8} \|\nabla_{\mathbf{q}} \tilde{U}(\mathbf{q}')\|_{\mathcal{P}(\mathbf{q}')}^2 - \frac{h}{2} \mathbf{v}^{\mathsf{T}} \nabla_{\mathbf{q}} \tilde{U}(\mathbf{q}) + \frac{h^2}{8} \|\nabla_{\mathbf{q}} \tilde{U}(\mathbf{q}')\|_{\mathcal{P}(\mathbf{q}')}^2 - \frac{h}{2} \mathbf{v}^{\mathsf{T}} \nabla_{\mathbf{q}} \tilde{U}(\mathbf{q}) \|_{\mathcal{P}(\mathbf{q})}^2 \\ &= \Delta \tilde{U} - \frac{h}{2} \left[\mathbf{v}'^{\mathsf{T}} \nabla_{\mathbf{q}} \tilde{U}(\mathbf{q}') + \mathbf{v}^{\mathsf{T}} \nabla_{\mathbf{q}} \tilde{U}(\mathbf{q}) \right] - \frac{h^2}{8} \left[\|\nabla_{\mathbf{q}} \tilde{U}(\mathbf{q}')\|_{\mathcal{P}(\mathbf{q}')}^2 - \|\nabla_{\mathbf{q}} \tilde{U}(\mathbf{q})\|_{\mathcal{P}(\mathbf{q})}^2 \right] \end{split}$$

where $\mathcal{P}(\mathbf{q}')\mathbf{v}^+ = \mathbf{v}^+$, $\mathcal{P}(\mathbf{q})\mathbf{v}^- = \mathbf{v}^-$, and $\|\mathbf{v}^+\|_2^2 = \|\mathbf{v}^-\|_2^2$. Accumulating the above terms over $\tau = 1, \dots, T$ yields the reformulated acceptance probability (17).

We now prove the energy conservation theorem 3.1 (Beskos et al., 2011).

Proof of Theorem 3.1. With the second equation of Lagrangian dynamics (15) we have

$$-\langle \mathbf{v}(t), \mathbf{g}(\mathbf{q}(t)) \rangle = \mathbf{v}(t)^{\mathsf{T}} \nabla_{\mathbf{q}} \tilde{U}(\mathbf{q}(t)) = \mathbf{v}_{-D}(t)^{\mathsf{T}} d\mathcal{I}(\mathbf{q})^{\mathsf{T}} \nabla_{\mathbf{q}} \tilde{U}(\mathbf{q}(t)) = \mathbf{v}_{-D}(t)^{\mathsf{T}} \nabla_{\mathbf{q}_{-D}} \tilde{U}(\mathbf{q}(t))$$

$$= \mathbf{v}_{-D}(t)^{\mathsf{T}} \mathbf{G}(\mathbf{q}_{-D}(t)) \left[\dot{\mathbf{v}}_{-D}(t) + \mathbf{v}_{-D}^{\mathsf{T}}(t) \mathbf{\Gamma}(\mathbf{q}_{-D}(t)) \mathbf{v}_{-D}(t) \right]$$

$$= \mathbf{v}_{-D}(t)^{\mathsf{T}} \mathbf{G}(\mathbf{q}_{-D}(t)) \dot{\mathbf{v}}_{-D}(t) + \frac{1}{2} \mathbf{v}_{-D}(t)^{\mathsf{T}} d\mathbf{G}(\mathbf{q}_{-D}(t)) \mathbf{v}_{-D}(t)$$

$$= \frac{d}{dt} \frac{1}{2} \mathbf{v}_{-D}(t)^{\mathsf{T}} \mathbf{G}(\mathbf{q}_{-D}(t)) \mathbf{v}_{-D}(t) = \frac{d}{dt} \frac{1}{2} \|\mathbf{v}(t)\|_{2}^{2}$$

Then we have the first equality hold because

$$-\int_{0}^{T} \langle \mathbf{v}(t), \mathbf{g}(\mathbf{q}(t)) \rangle dt = \frac{1}{2} \|\mathbf{v}(T)\|_{2}^{2} - \frac{1}{2} \|\mathbf{v}(0)\|_{2}^{2}$$

Lastly, from the first equation of Lagrangian dynamics (15)

$$\tilde{U}(\mathbf{q}(T)) - \tilde{U}(\mathbf{q}(0)) = \int_0^T \dot{\tilde{U}}(\mathbf{q}(t)) = \int_0^T \langle \dot{\mathbf{q}}(t), \nabla_{\mathbf{q}} \tilde{U}(\mathbf{q}(t)) \rangle dt = \int_0^T \langle \mathbf{v}(t), \mathbf{g}(\mathbf{q}(t)) \rangle dt$$

Therefore the second equality is proved.

Algorithm 1 Adaptive Spherical HMC (adp-SphHMC)

```
Given \mathbf{q}_0, a_0, N, N^{\mathrm{adapt}}.
Set h_0 = 1 or using Algorithm 4 of Hoffman and Gelman (2014), \mu = \log(10h_0), \overline{h}_0 = 1, \overline{A}_0 = 1
0, \gamma = 0.05, n_0 = 10, \kappa = 0.75.
for n = 1 to N do
    Sample a new velocity \mathbf{v}_{n-1} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_D), and set \mathbf{v}_{n-1} = \mathcal{P}(\mathbf{q}_{n-1})\mathbf{v}_{n-1}.
    Set \mathbf{q}^{(0)} = \mathbf{q}_{n-1}, \, \mathbf{v}^{(0)} = \mathbf{v}_{n-1}.
    for \tau = 0 to T - 1 (T = T_{2orth} \text{ or } T_{stoch}) do
        Run leapfrog step (16) to update (\mathbf{q}^{(\tau+1)}, \mathbf{v}^{(\tau+1)}) \leftarrow \mathcal{T}_{h_{n-1}}(\mathbf{q}^{(\tau)}, \mathbf{v}^{(\tau)}).
        if Stopping criterion (18) (or (19)) is satisfied then
        end if
    end for
    Accept the proposal (\mathbf{q}^{(T)}, \mathbf{v}^{(T)}) with probability a_n^{sphHMC} in (17) and set \mathbf{q}_n = \mathbf{q}^{(T)}; otherwise
set \mathbf{q}_n = \mathbf{q}_{n-1}.
if n \leq N^{\mathrm{ad}apt} then
        Set \overline{A}_n = \left(1 - \frac{1}{n+n_0}\right) \overline{A}_{n-1} + \frac{1}{n+n_0} (a_0 - a_n).
        Set \log h_n = \mu - \frac{\sqrt{n}}{\gamma} \overline{A}_n, and \log \overline{h}_n = n^{-\kappa} \log h_n + (1 - n^{-\kappa}) \log \overline{h}_{n-1}.
        Set h_n = \overline{h}_{N^{adpat}}.
    end if
end for
```

D Gradient Calculation in Normal-inverse-Wishart Problem

We use the representation (6) and derive log-posterior (log-likelihood and log-prior) and the corresponding gradients for (20) using matrix calculus.

D.1 Gradients of log-likelihood

Denote $\mathbf{y}_n^* := (\mathbf{y}_n - \boldsymbol{\mu}_0)/\boldsymbol{\sigma}$. Then the log-likelihood becomes

$$\ell(\mathbf{y}^*; \boldsymbol{\sigma}, \mathbf{P}) = -N\mathbf{1}_D^\mathsf{T} \log \boldsymbol{\sigma} - \frac{N}{2} \log |\mathbf{P}| - \frac{1}{2} \sum_{n=1}^N \mathbf{y}_n^{*\mathsf{T}} \mathbf{P}^{-1} \mathbf{y}_n^*$$

 $\left[\frac{\partial \ell}{\partial \tau}\right]$. We calculate the gradient of log-likelihood with respect to σ

$$\frac{\partial \ell}{\partial \sigma_k} = -N\sigma_k^{-1} + \sum_{n=1}^N \sum_i \frac{y_{ni}^*}{\sigma_i} \delta_{ik}(\mathbf{P}^{-1}\mathbf{y}_n^*)_i, \quad i.e. \frac{\partial \ell}{\partial \boldsymbol{\sigma}} = -N\boldsymbol{\sigma}^{-1} + \sum_{n=1}^N \operatorname{diag}(\mathbf{y}_n^*/\boldsymbol{\sigma})(\mathbf{P}^{-1}\mathbf{y}_n^*)$$

And with the transformation $\tau = \log(\sigma)$ it becomes

$$\frac{\partial \ell}{\partial \boldsymbol{\tau}} = \frac{d\boldsymbol{\sigma}^{\mathsf{T}}}{d\boldsymbol{\tau}} \frac{\partial \ell}{\partial \boldsymbol{\sigma}} = \operatorname{diag}(\boldsymbol{\sigma}) \left[-\frac{N}{\boldsymbol{\sigma}} + \sum_{n=1}^{N} \operatorname{diag}(\mathbf{y}_{n}^{*}/\boldsymbol{\sigma})(\mathbf{P}^{-1}\mathbf{y}_{n}^{*}) \right] = -N\mathbf{1}_{D} + \sum_{n=1}^{N} \operatorname{diag}(\mathbf{y}_{n}^{*})(\mathbf{P}^{-1}\mathbf{y}_{n}^{*})$$

 $\left[\frac{\partial \ell}{\partial \mathbf{U}^*} \left(\frac{\partial \ell}{\partial \mathbf{L}}\right)\right]$. When $\mathbf{P} = \mathbf{U}^*(\mathbf{U}^*)^\mathsf{T}$, $\frac{1}{2}\log|\mathbf{P}| = \log|\mathbf{U}^*| = \mathbf{1}_D^\mathsf{T}\log|\operatorname{diag}(\mathbf{U}^*)|$ and thus we have

$$\frac{\partial \ell}{\partial \mathbf{U}^*} = -\frac{N\mathbf{I}_D}{\mathbf{U}^*} + \sum_{n=1}^{N} \frac{dg_n(\widetilde{\mathbf{U}})}{d\mathbf{U}^*}$$

where $\frac{\mathbf{I}_D}{\mathbf{U}^*} = \operatorname{diag}(\{(u_{ii}^*)^{-1}\})$ is a diagonal matrix formed by element-wise division, $\widetilde{\mathbf{U}} := (\mathbf{U}^*)^{-1}$ and $g_n(\widetilde{\mathbf{U}}) := -\frac{1}{2}\mathbf{y}_n^*\mathsf{T}\widetilde{\mathbf{U}}\mathsf{T}\widetilde{\mathbf{U}}\mathbf{y}_n^*$.

Taking differential directly on $g_n(\mathbf{U}^*) := -\frac{1}{2}\mathbf{y}_n^{*\mathsf{T}}(\mathbf{U}^*)^{-\mathsf{T}}(\mathbf{U}^*)^{-1}\mathbf{y}_n^*$, and noting that differential and trace operators are exchangeable, we have

$$dg_n(\mathbf{U}^*) = -\frac{1}{2} \operatorname{tr}(\mathbf{y}_n^{*\mathsf{T}} d(\mathbf{U}^*)^{-\mathsf{T}} (\mathbf{U}^*)^{-1} \mathbf{y}_n^* + \mathbf{y}_n^{*\mathsf{T}} (\mathbf{U}^*)^{-\mathsf{T}} d(\mathbf{U}^*)^{-1} \mathbf{y}_n^*)$$

$$= \frac{1}{2} \left[\operatorname{tr}(\mathbf{y}_n^{*\mathsf{T}} (\mathbf{U}^*)^{-\mathsf{T}} d(\mathbf{U}^*)^{\mathsf{T}} \mathbf{P}^{-1} \mathbf{y}_n^*) + \operatorname{tr}(\mathbf{y}_n^{*\mathsf{T}} \mathbf{P}^{-1} d\mathbf{U}^* (\mathbf{U}^*)^{-1} \mathbf{y}_n^*) \right]$$

$$= \operatorname{tr}(\mathbf{y}_n^{*\mathsf{T}} \mathbf{P}^{-1} d\mathbf{U}^* (\mathbf{U}^*)^{-1} \mathbf{y}_n^*) = \operatorname{tr}((\mathbf{U}^*)^{-1} \mathbf{y}_n^* \mathbf{y}_n^{*\mathsf{T}} \mathbf{P}^{-1} d\mathbf{U}^*)$$

Conversion from differential to normal derivative form in the numerator layout (Minka, 1200) yields

$$\frac{\partial g_n(\mathbf{U}^*)}{\partial (\mathbf{U}^*)^\mathsf{T}} = \mathrm{tril}((\mathbf{U}^*)^{-1}\mathbf{y}_n^*\mathbf{y}_n^{\mathsf{T}}\mathbf{P}^{-1}), \quad i.e., \quad \frac{\partial g_n(\mathbf{U}^*)}{\partial \mathbf{U}^*} = \mathrm{triu}(\mathbf{P}^{-1}\mathbf{y}_n^*\mathbf{y}_n^{\mathsf{T}}(\mathbf{U}^*)^{-\mathsf{T}})$$

Finally, we have

$$\frac{\partial \ell}{\partial \mathbf{U}^*} = -\frac{N\mathbf{I}_D}{\mathbf{U}^*} + \operatorname{triu}(\mathbf{P}^{-1} \sum_{n=1}^{N} \mathbf{y}_n^* \mathbf{y}_n^{*\mathsf{T}} (\mathbf{U}^*)^{-\mathsf{T}})$$

When $\mathbf{P} = \mathbf{L}\mathbf{L}^{\mathsf{T}}$, by similar argument as above, we have

$$\frac{\partial \ell}{\partial \mathbf{L}} = -\frac{N\mathbf{I}_D}{\mathbf{L}} + \mathrm{tril}(\mathbf{P}^{-1} \sum_{n=1}^{N} \mathbf{y}_n^* \mathbf{y}_n^{*\mathsf{T}} \mathbf{L}^{-\mathsf{T}})$$

D.2 Gradients of log-priors

The logarithm of conditional prior $p(\boldsymbol{\sigma}|\mathbf{U}^*)$ after transformation $\boldsymbol{\tau} = \log(\boldsymbol{\sigma})$ becomes

$$\log p(\boldsymbol{\tau}|\mathbf{U}^*) = \log p(\boldsymbol{\sigma}|\mathbf{U}^*) + \log \left| \frac{d\boldsymbol{\sigma}}{d\boldsymbol{\tau}} \right| = \sum_{i=1}^{D} (i - (\nu + D))\tau_i - \frac{1}{2} \operatorname{tr}(\boldsymbol{\Psi}\operatorname{diag}(e^{-\boldsymbol{\tau}})\mathbf{P}^{-1}\operatorname{diag}(e^{-\boldsymbol{\tau}}))$$

 $\left[\frac{d}{d\tau}\log p(\tau|\mathbf{U}^*)\right]$. We calculate the derivative of $\log p(\tau|\mathbf{U}^*)$ with respect to τ

$$\frac{d}{d\tau}\log p(\tau|\mathbf{U}^*) = \mathbf{i} - (\nu + D) + \frac{dg(\tau)}{d\tau}$$

where $\mathbf{i} = [1, \dots, D]^{\mathsf{T}}$, and $g(\boldsymbol{\tau}) = -\frac{1}{2} \mathrm{tr}(\boldsymbol{\Psi} \operatorname{diag}(e^{-\boldsymbol{\tau}}) \mathbf{P}^{-1} \operatorname{diag}(e^{-\boldsymbol{\tau}}))$. Noting that differential and trace operators are exchangeable, we have

$$dg(\boldsymbol{\tau}) = -\frac{1}{2} \operatorname{tr}(\boldsymbol{\Psi} d \operatorname{diag}(e^{-\boldsymbol{\tau}}) \mathbf{P}^{-1} \operatorname{diag}(e^{-\boldsymbol{\tau}}) + \boldsymbol{\Psi} \operatorname{diag}(e^{-\boldsymbol{\tau}}) \mathbf{P}^{-1} d \operatorname{diag}(e^{-\boldsymbol{\tau}}))$$

$$= \frac{1}{2} \left[\operatorname{tr}(\mathbf{P}^{-1} \operatorname{diag}(e^{-\boldsymbol{\tau}}) \boldsymbol{\Psi} \operatorname{diag}(e^{-\boldsymbol{\tau}}) \operatorname{diag}(d\boldsymbol{\tau})) + \operatorname{tr}(\boldsymbol{\Psi} \operatorname{diag}(e^{-\boldsymbol{\tau}}) \mathbf{P}^{-1} \operatorname{diag}(e^{-\boldsymbol{\tau}}) \operatorname{diag}(d\boldsymbol{\tau})) \right]$$

$$= \sum_{i=1}^{D} d\tau_{i} \sum_{j=1}^{D} \psi_{ij} e^{-\tau_{j}} \rho^{ji} e^{-\tau_{i}}$$

Thus

$$\frac{dg(\tau)}{d\tau} = \operatorname{diag}(\boldsymbol{\Psi}\operatorname{diag}(e^{-\tau})\mathbf{P}^{-1})\operatorname{diag}(e^{-\tau}) = \operatorname{diag}(\boldsymbol{\Psi}\operatorname{diag}(e^{-\tau})\mathbf{P}^{-1}) \circ e^{-\tau}$$

where diag acting on a vector forms a diagonal matrix while the action a matrix means extracting the diagonal vector. \circ is the Hadamard product (a.k.a. Schur product), i.e. the entrywise product.

 $\left[\frac{d}{d\mathbf{U}^*}\log p(\mathbf{U}^*|\boldsymbol{\tau})\right]$. Now consider the derivative of $\log p(\mathbf{U}^*|\boldsymbol{\tau})$ with respect to the matrix \mathbf{U}^* . We have

$$\frac{d}{d\mathbf{U}^*} \log p(\mathbf{U}^* | \boldsymbol{\tau}) = \frac{\operatorname{diag}(\mathbf{i} - (\nu + D + 1))}{\mathbf{U}^*} + \frac{dg(\mathbf{U}^*)}{d\mathbf{U}^*}$$

where $g(\mathbf{U}^*) = -\frac{1}{2} \text{tr}(\mathbf{\Psi} \operatorname{diag}(e^{-\tau})(\mathbf{U}^*)^{-\mathsf{T}}(\mathbf{U}^*)^{-1} \operatorname{diag}(e^{-\tau}))$, and $\frac{\operatorname{diag}(\mathbf{i})}{\mathbf{U}^*}$ is a diagonal matrix formed by element-wise division.

Again by the exchangeability between differential and trace, we have

$$\begin{split} & dg(\mathbf{U}^*) \\ &= -\frac{1}{2} \mathrm{tr}(\boldsymbol{\Psi} \operatorname{diag}(e^{-\boldsymbol{\tau}}) d(\mathbf{U}^*)^{-\mathsf{T}} (\mathbf{U}^*)^{-1} \operatorname{diag}(e^{-\boldsymbol{\tau}}) + \boldsymbol{\Psi} \operatorname{diag}(e^{-\boldsymbol{\tau}}) (\mathbf{U}^*)^{-\mathsf{T}} d(\mathbf{U}^*)^{-1} \operatorname{diag}(e^{-\boldsymbol{\tau}})) \\ &= \frac{1}{2} \left[\mathrm{tr}(\boldsymbol{\Psi} \operatorname{diag}(e^{-\boldsymbol{\tau}}) (\mathbf{U}^*)^{-\mathsf{T}} d(\mathbf{U}^*)^{\mathsf{T}} \mathbf{P}^{-1} \operatorname{diag}(e^{-\boldsymbol{\tau}})) + \mathrm{tr}(\boldsymbol{\Psi} \operatorname{diag}(e^{-\boldsymbol{\tau}}) \mathbf{P}^{-1} d\mathbf{U}^* (\mathbf{U}^*)^{-1} \operatorname{diag}(e^{-\boldsymbol{\tau}})) \right] \\ &= \frac{1}{2} \left[\mathrm{tr}(\operatorname{diag}(e^{-\boldsymbol{\tau}}) \mathbf{P}^{-1} d\mathbf{U}^* (\mathbf{U}^*)^{-1} \operatorname{diag}(e^{-\boldsymbol{\tau}}) \boldsymbol{\Psi}) + \mathrm{tr}(\boldsymbol{\Psi} \operatorname{diag}(e^{-\boldsymbol{\tau}}) \mathbf{P}^{-1} d\mathbf{U}^* (\mathbf{U}^*)^{-1} \operatorname{diag}(e^{-\boldsymbol{\tau}})) \right] \\ &= \mathrm{tr}((\mathbf{U}^*)^{-1} \operatorname{diag}(e^{-\boldsymbol{\tau}}) \boldsymbol{\Psi} \operatorname{diag}(e^{-\boldsymbol{\tau}}) \mathbf{P}^{-1} d\mathbf{U}^*) \end{split}$$

Therefore we have

$$\frac{dg(\mathbf{U}^*)}{d(\mathbf{U}^*)^{\mathsf{T}}} = \operatorname{tril}((\mathbf{U}^*)^{-1}\operatorname{diag}(e^{-\boldsymbol{\tau}})\boldsymbol{\Psi}\operatorname{diag}(e^{-\boldsymbol{\tau}})\mathbf{P}^{-1}),$$

that is,

$$\frac{dg(\mathbf{U}^*)}{d\mathbf{U}^*} = \operatorname{triu}(\mathbf{P}^{-1}\operatorname{diag}(e^{-\tau})\boldsymbol{\Psi}\operatorname{diag}(e^{-\tau})(\mathbf{U}^*)^{-\mathsf{T}}),$$

 $\left[\frac{d}{d\tau}\log p(\tau), \frac{d}{d\mathbf{L}}\log p(\mathbf{L})\right]$. Lastly, the log-priors for (22) and their gradients after transformation $\boldsymbol{\tau} := \log(\boldsymbol{\sigma})$ are calculated

$$\log p(\boldsymbol{\tau}) = -\frac{1}{2}\boldsymbol{\tau}^{\mathsf{T}}\boldsymbol{\tau}, \qquad \qquad \frac{d}{d\boldsymbol{\tau}}\log p(\boldsymbol{\tau}) = -\boldsymbol{\tau}$$
$$\log p(\mathbf{l}_i) = \log p(\mathbf{l}_i^2) + \mathbf{1}_i^{\mathsf{T}}\log|2\mathbf{l}_i| = (2(\boldsymbol{\alpha}_i - 1) + \mathbf{1}_i)^{\mathsf{T}}\log|\mathbf{l}_i|, \quad \frac{d}{d\mathbf{l}_i}\log p(\mathbf{l}_i) = \frac{2(\boldsymbol{\alpha}_i - 1) + \mathbf{1}_i}{\mathbf{l}_i}$$

The bottom row can be written as

$$\log p(\mathbf{L}) = \sum_{i=1}^{D} (2\alpha_i - 1)^{\mathsf{T}} \log |\mathbf{l}_i|, \quad \frac{d}{d\mathbf{L}} \log p(\mathbf{L}) = \frac{2\alpha - 1}{\mathbf{L}}$$

where $\frac{1}{\mathbf{L}}$ denotes a lower-triangular matrix with l_{ij}^{-1} being its (i,j) entry $(i \geq j)$.

E More Numerical Results

E.1 Flexibility of von Mises-Fisher Prior and Bingham Prior

Definition E.1 (Fisher-Bingham / Kent distribution). The probability density function of the Kent distribution for the random vector $\mathbf{l}_i \in \mathcal{S}^{i-1}$ is given by

$$p(\mathbf{l}_i) \propto \exp\left\{\kappa {\boldsymbol{\gamma}_1}^\mathsf{T} \mathbf{l}_i + \sum_{k=2}^i \beta_k ({\boldsymbol{\gamma}_k}^\mathsf{T} \mathbf{l}_i)^2\right\}$$

where $\sum_{k=2}^{i} \beta_k = 0$ and $0 \le 2|\beta_k| < \kappa$ and the vectors $\{\gamma_k\}_{k=1}^i$ are orthonormal.

Remark E.1. The parameters κ and γ_1 are called the concentration and the mean direction parameter, respectively. The greater the value of κ , the higher the concentration of the distribution around the mean direction γ_1 . The choice of γ_1 could impact our priors when modeling correlations. Parameters $\{\beta_k\}_{k=2}^i$ determine the ellipticity of the contours of equal probability. The vectors $\{\gamma_k\}_{k=2}^i$ determine the orientation of the equal probability contours on the sphere.

Remark E.2. If $\beta_k = 0$ for $k = 2, \dots, i$, then this distribution reduces to von Mises-Fisher distribution (Fisher, 1953; Mardia and Jupp, 2009), denoted as vMF (κ, γ_1) . If $\kappa = 0$, then it defines an antipodally symmetric distribution, named Bingham distribution (Bingham, 1974), denoted as Bing(**A**), with $\mathbf{l}_i^\mathsf{T} \mathbf{A} \mathbf{l}_i = \sum_{k=2}^i \beta_k (\gamma_k^\mathsf{T} \mathbf{l}_i)^2$.




Figure E.1: Marginal posterior, prior (induced from von Mises-Fisher distribution) densities of correlations and MLEs with different settings for concentration parameter κ , estimated with 10^6 samples.

As before, to induce smaller correlations, one can put higher prior probabilities for \mathbf{l}_i on the poles of \mathcal{S}^{i-1} . For example, we might consider $\mathbf{l}_i \sim \text{vMF}(\kappa, \mathbf{n}_i)$, or $\mathbf{l}_i \sim \text{Bing}(\zeta \operatorname{diag}(\mathbf{n}_i))$, where $\mathbf{n}_i := (0, \dots, 0, 1)^\mathsf{T}$ is denoted as the north pole. Now let's consider the following von Mises-Fisher prior (Fisher et al., 1987; Fisher, 1953; Mardia and Jupp, 2009) for \mathbf{l}_i , the *i*-th row of the Cholesky factor \mathbf{L} of correlation matrix \mathbf{P} in the structured model (6).

Definition E.2 (Von Mises-Fisher distribution). The probability density function of the von Mises-Fisher distribution for the random vector $\mathbf{l}_i \in \mathcal{S}^{i-1}$ is given by

$$p(\mathbf{l}_i) = C_i(\kappa) \exp(\kappa \boldsymbol{\mu}^\mathsf{T} \mathbf{l}_i)$$

where $\kappa \geq 0$, $\|\boldsymbol{\mu}\| = 1$ and the normalization constant $C_i(\kappa)$ is equal to

$$C_i(\kappa) = \frac{\kappa^{i/2-1}}{(2\pi)^{i/2} I_{i/2-1}(\kappa)}$$

where I_v denotes the modified Bessel function of the first kind at order v. Denote $\mathbf{l}_i \sim \text{vMF}(\kappa, \boldsymbol{\mu})$.

Since we have no prior knowledge about the mean direction μ , we choose $\mu = \mathbf{n}_i = (\mathbf{0}_{i-1}, 1)^\mathsf{T}$ that favors the polar direction, i.e.

$$\mathbf{l}_i \sim \text{vMF}(\kappa, \mathbf{n}_i), \qquad p(\mathbf{l}_i) \propto \exp(\kappa l_{ii}), \quad i = 2, \cdots, D$$




Figure E.2: Marginal posterior, prior (induced from Bingham distribution) densities of correlations and MLEs with different settings for concentration parameter ζ , estimated with 10^6 samples.

where we consider i) $\kappa = 1$; ii) $\kappa = 10$; iii) $\kappa = 100$. With the von Mises-Fisher prior, we have

$$\log p(\mathbf{L}) = \sum_{i=1}^{D} \kappa l_{ii} = \kappa \operatorname{tr}(\mathbf{L}), \quad \frac{d}{d\mathbf{L}} \log p(\mathbf{L}) = \kappa \mathbf{I}$$

We repeat the experiment in Section 4.1.2 with the von Mises-Fisher prior for l_i . The posteriors, priors and maximal likelihood estimates (MLE) of correlations with different κ 's are plotted in Figure E.1 respectively. With larger concentration parameter κ , the posterior is pulled more towards 0.

Finally, we consider the following Bingham prior (Bingham, 1974; Onstott, 1980) for l_i in the structured model (6).

Definition E.3 (Bingham distribution). The probability density function of the Bingham distribution for the random vector $\mathbf{l}_i \in \mathcal{S}^{i-1}$ is given by

$$p(\mathbf{l}_i) = {}_1F_1(\frac{1}{2}; \frac{n}{2}; \mathbf{Z})^{-1} \exp(\mathbf{l}_i^\mathsf{T} \mathbf{M} \mathbf{Z} \mathbf{M}^\mathsf{T} \mathbf{l}_i)$$

where **M** is an orthogonal orientation matrix, **Z** is a diagonal concentration matrix, and ${}_{1}F_{1}(\cdot;\cdot;\cdot)$ is a confluent hypergeometric function of matrix argument. Denote $\mathbf{l}_{i} \sim \operatorname{Bing}(\mathbf{M},\mathbf{Z})$.

Note, according to Bingham (1974), this distribution is defined for **Z** up to an arbitrary scalar matrix $\zeta_0 \mathbf{I}$. Therefore, we consider $\mathbf{M} = \mathbf{I}$ and $\mathbf{Z} = \zeta \operatorname{diag}(\mathbf{n}_i)$ that favors the polar




Figure E.3: Simulated data y over the underlying mean functions μ_t (left), the variance functions Σ_t , and the correlation functions P_t (right) of 10-dimension periodic processes.

direction, i.e.

$$\mathbf{l}_i \sim \operatorname{Bing}(\mathbf{I}, \zeta \operatorname{diag}(\mathbf{n}_i)), \qquad p(\mathbf{l}_i) \propto \exp(\zeta l_{ii}^2), \quad i = 2, \cdots, D$$

where we consider i) $\zeta = 1$; ii) $\zeta = 10$; iii) $\zeta = 100$. The log-prior and its gradient are calculated as follows

$$\log p(\mathbf{L}) = \sum_{i=1}^{D} \zeta l_{ii}^2 = \zeta \|\operatorname{diag}(\mathbf{L})\|^2, \quad \frac{d}{d\mathbf{L}} \log p(\mathbf{L}) = 2\zeta \operatorname{diag}(\mathbf{L})$$

We repeat the above experiment with the Bingham prior for \mathbf{l}_i . The posteriors, priors and maximal likelihood estimates (MLE) of correlations with different ζ 's are plotted in Figure E.2 respectively. With larger concentration parameter ζ , the posteriors are pulled more towards the induced priors and concentrate on 0.

E.2 More Comparison to Latent Factor Process Model

The example of simulated periodic process in Section 4.2.2 is consider for D=2 for simplicity and convenience of visualization. Here we consider higher dimension D=10. The purpose here is not to show the scalability, but rather to investigate the robustness of our dynamic model (11) in terms of full flexibility.

We generate M=20 trials of data over N=100 evenly spaced points over [0,1]. The true mean, variance and correlation functions are modified from the example (23) using the Clausen functions (Clausen, 1832). Seen from Figure E.3, they behave more intricately with higher heterogeneity among those processes. This could impose further challenge for latent factor based models like (24) compared to D=2. We repeat the experiments in Section 4.2.2 and compare our dynamic model (11) with the latent factor process model (24) by Fox and Dunson (2015). To aid the visualization, we subtract the estimated process




Figure E.4: Estimated error functions of the underlying mean μ_t (left column), variance σ_t (middle column) and correlation ρ_t (right column) of 10-dimensional periodic processes, using latent factor process model (upper row) and our flexible model (lower row), based on M=20 trials of data over N=100 evenly spaced points. Solid lines are estimated errors and shaded regions are 95% credible bands.

from their true values and plot the error functions in Figure E.4. Even if we have tried our best to tune the parameters, e.g. L, the number of basis functions, and k, the size of latent factors, the latent factor process model (Fox and Dunson, 2015) is outperformed by our flexible dynamic model (11) in reducing estimation errors.

E.3 More Results on the Analysis of LFP data

In Section 5, we studied the LFP data collected from the hippocampus of rats performing a complex sequence memory task. Figure E.5 shows 12 locations from CA1 subregion of the hippocampus of the rat where LFP signals are recorded. Figure E.6 shows the theta-filtered traces (4-12Hz; left panel) and the estimated correlation processes under different experiment conditions (InSeq vs OutSeq; right panel). Here we observe a similar dynamic pattern of correlation matrices under two conditions that diverge after 500ms, indicating the neural activity associated with the cognitive process of identifying whether events occurred in their expected order.

We also did a study of the correlation evolution on the full 12 channels and revealed the block structure of those channels and the same changing pattern under different experimental conditions discovered with the chosen 4 channels in Section 5. This video demonstrates the result of 12 channels https://www.youtube.com/watch?v=NMUUicOIDsM.




Figure E.5: Locations of recorded LFP signals in CA1 subregion of the rat's hippocampus.




Figure E.6: Results of LFP theta signals: data (left), estimation of correlations (right).