Input-to-State Stability of Periodic Orbits of Systems with Impulse Effects via Poincaré Analysis

Sushant Veer, Rakesh, and Ioannis Poulakakis

Abstract—In this paper we investigate the relation between robustness of periodic orbits exhibited by systems with impulse effects and robustness of their corresponding Poincaré maps. In particular, we prove that input-to-state stability (ISS) of a periodic orbit under external excitation in both continuous and discrete time is equivalent to ISS of the corresponding 0-input fixed point of the associated forced Poincaré map. This result extends the classical Poincaré analysis for asymptotic stability of periodic solutions to establish orbital input-to-state stability of such solutions under external excitation. In our proof, we define the forced Poincaré map, and use it to construct ISS estimates for the periodic orbit in terms of ISS estimates of this map under mild assumptions on the input signals. As a consequence of the availability of these estimates, the equivalence between exponential stability (ES) of the fixed point of the 0input (unforced) Poincaré map and ES of the corresponding orbit is recovered. The results can be applied naturally to study the robustness of periodic orbits of continuous-time systems as well. Although our motivation for extending classical Poincaré analysis to address ISS stems from the need to design robust controllers for limit-cycle walking and running robots, the results are applicable to a much broader class of systems that exhibit periodic solutions.

Index Terms—Poincaré map, systems with impulse effects, limit cycles, input-to-state stability, robustness.

I. INTRODUCTION

SYSTEMS with impulse effects (SIEs) are characterized by a set of ordinary differential equations (ODEs) and a discrete map that reinitializes the ODEs when the corresponding solution reaches a switching surface, possibly resulting in discontinuous evolution. These systems arise in a broad range of fields; a non-exhaustive list of examples includes impact mechanics [1], modeling of population dynamics [2], communication [3], and legged robotics [4]; a collection of methods for analyzing SIEs can be found in [5].

In this paper, we study the stability properties of limit cycles exhibited by SIEs under external excitation. Our interest in this specific class of systems arises from dynamically-stable legged robots, where periodic walking gaits are modeled as limit cycles of SIEs. This approach has been successful in generating asymptotically stable periodic gaits for bipedal robots through a variety of methods, including hybrid zero dynamics [6], [7], geometric control [8], virtual holonomic constraints [9], to name a few. Recent extensions of these methods resulted in generating continuums of limit-cycle gaits for bipedal walkers [10], [11], and switching among them [12]–[14], to enlarge the behavioral repertoire of these robots in order to accomplish tasks that require adaptability to typical human-centric environments [15], and human (or robot) collaborators [16]. Practical use of these robots demands robustness to external disturbances, which has led many researchers—including the authors of the present paper—to analyze [17]–[19] and design [14], [20], [21] controllers that enhance the robustness of limit-cycle walking gaits. With this being our motivation, we develop in this paper a framework for rigorously analyzing the robustness of limit cycles, by relating orbital input-to-state stability (ISS) for hybrid limit cycles of SIEs with the corresponding Poincaré map.

1

The notion of ISS has been widely used to study robustness of equilibrium points in continuous [22], discrete [23], and hybrid [24] systems. Intuitively, the solutions emanating in a neighborhood of an ISS equilibrium point remain bounded when the external inputs are bounded. In addition, when the inputs vanish, these solutions converge back to the equilibrium. Beyond equilibrium points, ISS can be naturally applied to study robustness of zero-invariant sets by considering the point-to-set distance [22], [25]. Establishing ISS in this context poses a considerable challenge, which, in the case of SIEs, is exacerbated by the hybrid nature of the system. However, for periodic orbits-such as those of interest in this paper-we show here that the problem can be reduced to studying ISS of an unforced (0-input) fixed point of a discrete dynamical system, thus avoiding direct analysis of hybrid solutions. This discrete system arises through the Poincaré map construction suitably extended to incorporate external inputs, thereby resulting in the definition of a forced Poincaré map.

Numerous results exist that analyze forced Poincaré maps of systems evolving under the influence of external inputs; in the context of SIEs, examples include [18], [19], [26], in which the input signals are not necessarily periodic. However, the exact relation of conclusions deduced on the basis of the Poincaré map to properties of the underlying periodic orbit has not been explicitly discussed in the relevant literature. Indeed, rigorous results that relate the stability properties of the Poincaré map with those of the corresponding periodic orbit are restricted to systems without inputs; e.g., [27, Theorem 6.4] addresses local asymptotic stability (LAS) of periodic orbits in continuous systems, while [28, Theorem 1], [5, Theorem 13.1] address LAS and [29, Theorem 1] local exponential stability (LES) of such orbits in SIEs. The relation between the behavior of periodic orbits of SIEs under external inputs and the corresponding *forced* Poincaré map is at the core of this paper.

Specifically, the main contribution of this work (Theorem 1) is that ISS of a limit cycle exhibited by a SIE is equivalent

S. Veer and I. Poulakakis are with the Department of Mechanical Engineering and Rakesh is with the Department of Mathematical Sciences, University of Delaware, Newark, DE, 19716 USA e-mail: {veer, rakesh, poulakas}@udel.edu.

This work is supported in part by NSF CAREER Award IIS-1350721 and by NRI-1327614.

to ISS of a 0-input fixed point of the corresponding forced Poincaré map. This result significantly simplifies analysis, as it replaces the problem of establishing ISS of a hybrid limit cycle with the simpler problem of checking asymptotic stability of a 0-input fixed point of a discrete dynamical system defined by the corresponding Poincaré map (Theorem 2). To ensure the level of generality required by practical applications, we consider inputs affecting both the continuous and the discrete dynamics of the system. The continuous-time inputs belong in the (Banach) space of continuous bounded functions under the supremum norm. The resulting forced Poincaré map is a nonlinear functional defined over an infinite-dimensional function space, thus significantly extending prior work that considers finite dimensional disturbances; see [18], [19], [30] for example. Finally, the proof of the main result provides an explicit connection between ISS estimates of the forced Poincaré map and those of the hybrid orbit.

The results presented in this paper generalize previous contributions such as [29, Theorem 1], which is widely used to establish exponential stability (ES) of a hybrid limit cycle when the fixed point of the corresponding Poincaré map is ES. Indeed, [29, Theorem 1] can be obtained as a consequence of Theorem 1 of Section III below. Furthermore, Proposition 1 of Section III and Lemma 9 of Section V complete crucial arguments that were omitted in the proof of [29, Theorem 1]. Moreover, our results can offer useful tools for the design of robust controllers for limit cycles of SIEs. For example, the methods in [31] that are based on Poincaré map analysis can be supported using Theorem 1. As a final note, the results of this paper can naturally be applied to study ISS of limit cycles of continuous-time nonlinear systems under external excitation. Hence, their relevance extends to other bioinspired robotsincluding aerial robots with flapping wings [32] and robot snakes [33]-which, like legged robots, realize locomotion through periodic forceful interactions with their environment.

II. BACKGROUND

This section introduces the class of systems with impulse effects pertinent to this paper, and develops a forced Poincaré map suitable for studying periodic orbits of such systems under the influence of continuous and discrete exogenous inputs; such inputs could be command or disturbance signals. We begin with a few notes on the notation used in the paper.

A. Notation

Let \mathbb{R} and \mathbb{Z} denote the sets of real and integer numbers, and \mathbb{R}_+ and \mathbb{Z}_+ the corresponding subsets that include the non-negative reals and integers, respectively. For any $x \in \mathbb{R}^n$, the Euclidean norm is represented as ||x||. An open ball of radius $\delta > 0$ centered at x is denoted by $B_{\delta}(x)$. The pointto-set distance of x from $\mathcal{A} \subseteq \mathbb{R}^n$ is defined as $\operatorname{dist}(x, \mathcal{A}) :=$ $\inf_{y \in \mathcal{A}} ||x - y||$. We use $\mathcal{P}(\mathcal{A})$ to represent the power set of \mathcal{A} , and \mathcal{A}^c to denote the complement of \mathcal{A} with respect to \mathbb{R}^n .

For any interval $E \subseteq \mathbb{R}$ let $u : E \to \mathbb{R}^p$ be a function that represents the continuous-time inputs. The norm of u is defined as $||u||_{\infty} := \sup_{t \in E} ||u(t)||$. The set of continuoustime inputs we work with belongs to $\mathcal{U} := \{u : E \to \mathbb{R}^p\}$ $\mathbb{R}^p \mid u$ is continuous, $\|u\|_{\infty} \in \mathbb{R}_+$ }. Discrete-time inputs $\bar{v} : \mathbb{Z}_+ \to \mathbb{R}^q$ correspond to sequences $\bar{v} = \{v_k\}_{k=0}^{\infty}$ with $v_k \in \mathbb{R}^q$ for $k \in \mathbb{Z}_+$. The norm of \bar{v} is defined as $\|\bar{v}\|_{\infty} := \sup_{k \in \mathbb{Z}_+} \|v_k\|$. The discrete inputs belong to $\mathcal{V} := \{\bar{v} : \mathbb{Z}_+ \to \mathbb{R}^q \mid \|\bar{v}\|_{\infty} \in \mathbb{R}_+\}$. With an abuse of notation we use $\|\cdot\|_{\infty}$ to denote the norm for both \mathcal{U} and \mathcal{V} . No ambiguity arises because the meaning of $\|\cdot\|_{\infty}$ depends on whether the argument is continuous or discrete.

A function $\alpha : \mathbb{R}_+ \to \mathbb{R}_+$ belongs to class \mathcal{K} if it is continuous, strictly increasing, and $\alpha(0) = 0$. A function $\beta : \mathbb{R}_+ \times \mathbb{R}_+ \to \mathbb{R}_+$ belongs to class \mathcal{KL} if it is continuous, $\beta(\cdot, t)$ belongs to \mathcal{K} for any fixed $t \ge 0$, $\beta(s, \cdot)$ is strictly decreasing, and $\lim_{t\to\infty} \beta(s, t) = 0$, for any fixed $s \ge 0$.

B. Forced Systems With Impulse Effects

We are interested in studying the stability of periodic orbits exhibited by systems with impulse effects under externally applied inputs. These systems are characterized by alternating continuous and discrete phases. The evolution of the state $x \in \mathbb{R}^n$ during the continuous phase is governed by an ODE

$$\dot{x}(t) = f(x(t), u(t))$$
, (1)

where the input $u : \mathbb{R}_+ \to \mathbb{R}^p$ is an element of \mathcal{U} defined in Section II-A and $u(t) \in \mathbb{R}^p$ is its value. The vector field f in the right-hand side of (1) satisfies the following assumption: A.1) $f : \mathbb{R}^n \times \mathbb{R}^p \to \mathbb{R}^n$ is twice continuously differentiable¹.

Local existence and uniqueness of solutions of (1) for a fixed u follows from [34, Theorem 3.1] based on assumption A.1 and the continuity of u as a function of t. We denote the flow of (1) starting from the initial state x(0) and evolving under the influence of the input u by $\varphi(t, x(0), u)$.

The continuous phase terminates when the flow of (1) reaches a set $S \subset \mathbb{R}^n$ defined as

$$\mathcal{S} := \{ x \in \mathbb{R}^n \mid H(x) = 0 \} \quad , \tag{2}$$

where it is assumed that

A.2) $S \neq \emptyset$, $H : \mathbb{R}^n \to \mathbb{R}$ is twice continuously differentiable, and for all $\hat{x} \in S$, $\frac{\partial H}{\partial x}|_{\hat{x}} \neq 0$, i.e., S is a co-dimension 1 embedded submanifold in \mathbb{R}^n ; see [35, p. 431].

For future use we define the sets $S^+ := \{x \in \mathbb{R}^n \mid H(x) > 0\}$ and $S^- := \{x \in \mathbb{R}^n \mid H(x) < 0\}.$

The intersection of the flow of (1) with S initiates the discrete phase, which is governed by the mapping

$$x^{+} = \Delta(x^{-}, v) \quad \text{for } x^{-} \in \mathcal{S} \quad , \tag{3}$$

where x^- , x^+ are the states right before and after impacting S, respectively, and $v \in \mathbb{R}^q$ is a member of the discrete input \bar{v} that belongs in \mathcal{V} defined in Section II-A. It is assumed that A.3) $\Delta : \mathbb{R}^n \times \mathbb{R}^q \to \mathbb{R}^n$ is continuously differentiable.

Putting together the continuous and discrete phases (1) and (3), the *forced system with impulse effects* takes the form

$$\Sigma : \begin{cases} \dot{x}(t) = f(x(t), u(t)) & \text{if } x(t) \notin \mathcal{S} \\ x^+(t) = \Delta(x^-(t), v) & \text{if } x^-(t) \in \mathcal{S} \end{cases}$$

$$\tag{4}$$

¹Whenever we state that a function is k-times continuously differentiable, it applies to all its arguments.

where $x^-(t) := \lim_{\tau \nearrow t} x(\tau)$ and $x^+(t) := \lim_{\tau \searrow t} x(\tau)$.

At any time instant for which it exists, the solution of (4) evolves according to either (1) or (3). This allows us to represent the hybrid flow of (4) as the solution of (1) which, on approaching S, is interrupted by the discrete map (3). Let $\psi(t, x(0), u, \bar{v})$ denote the flow of (4) for some initial state x(0), continuous input u, and discrete input \bar{v} . Adapting the definition in [28, Section III-A], $\psi(t, x(0), u, \bar{v})$ as a function of time $t \in [0, t_f), t_f \in \mathbb{R}_+ \cup \{\infty\}$, satisfies the following: (i) it is right continuous² on $[0, t_f)$; (ii) left limits exist at each point in $(0, t_f)$; and (iii) there exists a discrete subset $\mathcal{T} \subset [0, t_{\rm f})$ such that (a) for every $t \notin \mathcal{T}$, $\psi(t, x(0), u, \bar{v})$ satisfies (1) for the input u considered, and (b) for $t \in \mathcal{T}$, $x^{-}(t) = \lim_{\tau \nearrow t} \psi(\tau, x(0), u, \bar{v}) \in \mathcal{S}$ and $x^+(t) = \lim_{\tau \searrow t} \psi(\tau, x(0), u, \bar{v}) = \Delta(x^-(t), v)$ where v is a member of the sequence \bar{v} . Note that right continuity implies that at time $t \in \mathcal{T}$ the solution attains the value $x^{+}(t)$ and not $x^{-}(t)$; that is, $x(t) = \psi(t, x(0), u, \bar{v}) = x^{+}(t)$. Moreover, Proposition 2 in Section III below ensures that in the neighborhood of distinct locally input-to-state stable periodic orbits-such as those of interest in this work-the solutions of (4) exist for arbitrary $t_{\rm f} > 0$, they do not possess consecutive discrete jumps (beating) and do not exhibit Zeno behavior; see [5], [36] for definitions.

Let $x^* \in S$ and $T^* \in (0, \infty)$ such that the following hold: A.4) dist $(\Delta(x^*, 0), S) > 0$, and $\Delta(x^*, 0) \in S^+$. The choice $\Delta(x^*, 0) \in S^+$ does not result in loss of generality; if $\Delta(x^*, 0) \in S^-$ re-define S with $\hat{H}(x) := -H(x)$.

A.5) $\varphi(t, \Delta(x^*, 0), 0)$ exists for all $t \in [0, T^*]$ and $\varphi(T^*, \Delta(x^*, 0), 0) = x^*$.

Using assumptions A.4-A.5 define

$$\mathcal{O} := \{\varphi(t, \Delta(x^*, 0), 0) \mid t \in [0, T^*)\} , \qquad (5)$$

and suppose further that \mathcal{O} satisfies the following assumptions: A.6) Let $\overline{\mathcal{O}}$ be the closure of \mathcal{O} , then $S \cap \overline{\mathcal{O}} = \{x^*\}$.

A.7) \mathcal{O} is transversal to \mathcal{S} at x^* , i.e., $L_f H(x^*, 0) := \frac{\partial H}{\partial x}\Big|_{x^*} f(x^*, 0) < 0.$

It follows from our assumptions that \mathcal{O} is a bounded unforced $(u \equiv 0, \bar{v} \equiv 0)$ hybrid periodic orbit of Σ that exhibits only one impact with S at x^* and has period T^* . Moreover, \mathcal{O} is not a closed curve; see Fig. 1 for a geometric illustration.

C. Forced Poincaré Map

The Poincaré map is a common tool used for analyzing systems with periodic orbits. Given a Poincaré section—which is an embedded submanifold transversal to the orbit—the Poincaré map returns consecutive intersections of the system's flow with the Poincaré section. Here, we study the map which returns the intersection of the solution of (4) with S under the influence of the external inputs u and \bar{v} . Consequently, it is natural to call this map the *forced* Poincaré map.



Fig. 1. Geometric illustration of O. The switching surface S is in grey.

As was mentioned in Section II-B, when the input uaffecting (1) is a *fixed* signal from \mathcal{U} , existence and uniqueness of the solution emanating from $x(0) \in \mathbb{R}^n$ can be established over an interval $J \subseteq \mathbb{R}_+$ with $0 \in J$, by [34, Theorem 3.1] applied on the time-varying vector field $\hat{f}(t, x) := f(x, u(t))$. To develop the forced Poincaré map, however, we need to compare solutions with different initial conditions and different inputs. To do this, it is important to be able to consider the forced solution $\varphi(t, x(0), u)$ of (1) as a mapping from $J \times \mathbb{R}^n \times \mathcal{U}$ to \mathbb{R}^n , interpreting u as an infinite-dimensional "parameter" residing in the Banach space $(\mathcal{U}, \|\cdot\|_{\infty})$. We can then analyze variations of $\varphi(t, x(0), u)$ with respect to its arguments, including u. The following lemma shows that, over its maximal interval of existence, the solution $\varphi(t, x(0), u)$ of (1) is continuously differentiable in its arguments, with differentiability understood in the Fréchet sense [37, p. 333].

Lemma 1. Let $f : \mathbb{R}^n \times \mathbb{R}^p \to \mathbb{R}^n$ in (1) be continuously differentiable, and let $u \in \mathcal{U}$ with \mathcal{U} as in Section II-A. Then, the solution $\varphi : J \times \mathbb{R}^n \times \mathcal{U} \to \mathbb{R}^n$ is continuously differentiable in its arguments in the Fréchet sense.

The proof makes use of Banach calculus [37] and is presented in Appendix A. We only note here that, for notational convenience, we use the same symbol u to denote both the finite-dimensional values of the input function at given instants and the infinite-dimensional input signal as a function in the Banach space $(\mathcal{U}, \|\cdot\|_{\infty})$; the distinction will always be clear through the domain of definition of the corresponding map.

Let $T_{\mathrm{I}}: \mathcal{S} \times \mathcal{U} \times \mathbb{R}^q \to \mathbb{R}_+ \cup \{\infty\}$ be the time-to-impact map defined as

$$T_{\mathbf{I}}(x, u, v) := \begin{cases} \inf\{t \ge 0 \mid \varphi(t, \Delta(x, v), u) \in \mathcal{S}\}, \\ & \text{if } \exists t : \varphi(t, \Delta(x, v), u) \in \mathcal{S} \\ \infty, & \text{otherwise} \end{cases}$$
(6)

Lemma 2 below establishes that the time-to-impact function $T_{\rm I}$ is well defined and continuously differentiable in x, u and v. Note that the dependence of $T_{\rm I}$ on u is to be understood with u interpreted as a function in $(\mathcal{U}, \|\cdot\|_{\infty})$. The proof is based on the implicit mapping theorem [37, Chapter XIV, Theorem 2.1] and on Lemma 1 above, and is presented in Appendix A.

Lemma 2. Consider (6). Suppose that (4) satisfies assumptions A.1-A.7. Then, there exists a $\delta > 0$ such that $T_{\rm I}$ is continuously differentiable for any $x \in B_{\delta}(x^*) \cap S$, $u \in U$ with $||u||_{\infty} < \delta$, and $v \in B_{\delta}(0)$.

³Without loss of generality, we use t = 0 as the initial time to avoid confusion with t_0 , which, in our notation, is the instant of the first impact.

²To avoid the state having to take two values at impact, a choice is to be made as to whether the state just before or just after impact—i.e., x^- or x^+ , respectively—is included in the solution. The former corresponds to left continuity and the latter to right continuity of ψ as a function of time. We assume here that ψ is right continuous with respect to t; note however that the results that follow hold regardless of this choice [5].

We are now ready to define the forced Poincaré map $P: \mathcal{S}\times\mathcal{U}\times\mathbb{R}^q\to\mathcal{S}$ as

$$P(x, u, v) := \varphi(T_{\mathbf{I}}(x, u, v), \Delta(x, v), u)$$

From Lemmas 1 and 2 it follows that P is well defined and continuously differentiable for any $x \in B_{\delta}(x^*) \cap S$, $u \in \mathcal{U}$ with $||u||_{\infty} < \delta$, and $v \in B_{\delta}(0)$.

Let $\psi(t, x(0), u, \bar{v})$ be a solution of (4) and $x(t) = \psi(t, x(0), u, \bar{v})$ be the value of the state at time t. For $k \in \mathbb{Z}_+$, let t_k be the instant at which the (k + 1)-th "intersection" of x(t) with S occurs. Define $u_k(t) := u(t)$ for $t_k \leq t < t_{k+1}$, and let v_k be the k-th element of the sequence \bar{v} . Then, the forced Poincaré map gives rise to the forced discrete system

$$x_{k+1} = P(x_k, u_k, v_k) , (7)$$

where $x_k := \lim_{t \neq t_k} x(t)$. The discrete system (7) captures the evolution of the system from *just before* an impact with Sto *just before* the next impact, assuming that the next impact occurs. It should be emphasized that the state x(t) does *not* attain⁴ the value x_k at t_k because ψ has been assumed right continuous in t; in fact, $x(t_k) = \Delta(x_k, v_k) = \lim_{t \geq t_k} x(t) \neq$ $\lim_{t \neq t_k} x(t) = x_k$. Let x^* be as in assumption A.6, then x^* is the 0-input fixed point of (7), i.e., $x^* = P(x^*, 0, 0)$.

For future use, we also define $\hat{T}_{I} : S^{+} \times \mathcal{U} \to \mathbb{R}_{+} \cup \{\infty\}$ as the time-to-impact function for solutions of (1) starting from states in S^{+} as

$$\hat{T}_{I}(x,u) :=
\begin{cases}
\inf\{t \ge 0 \mid \varphi(t,x,u) \in \mathcal{S}\}, & \text{if } \exists t : \varphi(t,x,u) \in \mathcal{S} \\
\infty, & \text{otherwise}
\end{cases}$$
(8)

It is noted that for any point $w \in \mathcal{O}$ there exists a $\delta > 0$ such that \hat{T}_{I} is continuously differentiable for any $x \in B_{\delta}(w)$ and any $u \in \mathcal{U}$ with $||u||_{\infty} < \delta$. The proof similar to that of Lemma 2, and it is not presented for brevity.

Finally, Remark 1 clarifies the relation between ψ , φ and the sequence $\{x_k\}_{k=0}^{\infty}$ and Remark 2 indicates that the results are valid even when the input u is a piecewise continuous signal, as long as the points of discontinuity are at t_k .

Remark 1. Let $\psi(t, x(0), u, \bar{v})$ be a solution of (4) that exists for all $t \ge 0$ and $x(t) = \psi(t, x(0), u, \bar{v})$ be the value of the state at time t. Then, if $t_0 := \hat{T}_I(x(0), u)$ is the time instant of the first crossing of S we have

$$x(t) = \varphi(t, x(0), u_{[0,t_0)}), \text{ for } 0 \le t < t_0$$

Moreover, if the sequence $\{x_k\}_{k=0}^{\infty}$ is the solution of (7) for the initial state $x_0 := \lim_{t \neq t_0} x(t)$ and the sequence of input functions $\{u_k\}_{k=0}^{\infty}$ are as defined above, then for all $k \in \mathbb{Z}_+$

$$x(t) = \varphi(t, \Delta(x_k, v_k), u_k), \text{ for } t_k \le t < t_{k+1} ,$$

where v_k is the k-th element of \bar{v} and

$$t_{k+1} = t_k + T_{\mathrm{I}}(x_k, u_k, v_k) = t_0 + \sum_{j=0}^k T_{\mathrm{I}}(x_j, u_j, v_j)$$
.

⁴Right continuity of ψ in t implies that, in general, there is no t for which $x(t) \in S$; see related comments in Section II-B and in [4, Section 4.1.2].

Remark 2. The results of the paper hold when u is discontinuous, as long as each u_k is a continuous function in U that agrees with u over the interval $[t_k, t_{k+1})$.

D. Pertinent Stability Definitions

Notions of orbital stability that will be studied in this paper are introduced here. We begin with local input-to-state stability (LISS) of the periodic orbit.

Definition 1. The periodic orbit \mathcal{O} of (4) is orbitally LISS if there exists $a \delta > 0$, $\alpha_1, \alpha_2 \in \mathcal{K}$, and $\beta \in \mathcal{KL}$ such that $x(t) = \psi(t, x(0), u, \overline{v})$ satisfies for all $t \in [0, t_f)$, $t_f \in \mathbb{R}_+ \cup \{\infty\}$,

$$\operatorname{dist}(x(t), \mathcal{O}) \leq \beta(\operatorname{dist}(x(0), \mathcal{O}), t) + \alpha_1(\|u\|_{\infty}) + \alpha_2(\|\bar{v}\|_{\infty}) , \qquad (9)$$

for any $x(0) \in S^+$ with $\operatorname{dist}(x(0), \mathcal{O}) < \delta$, $u \in \mathcal{U}$ with $||u||_{\infty} < \delta$, and $\bar{v} \in \mathcal{V}$ with $||\bar{v}||_{\infty} < \delta$.

Proposition 2 in Section III below asserts that shrinking $\delta > 0$ in Definition 1 guarantees that all ensuing hybrid solutions exist for all time; that is, $t_f > 0$ in Definition 1 can be chosen arbitrarily large. Since we focus on local properties of distinct periodic orbits \mathcal{O} of (4), we work with solutions in a small enough neighborhood of \mathcal{O} that satisfy (9) for all $t \geq 0$.

Besides LISS, we will briefly consider local exponential stability (LES) of \mathcal{O} . As above, Proposition 2 of Section III below ensures that in a small enough neighborhood of \mathcal{O} solutions of (4) exist over arbitrarily long intervals. Hence, the definition below assumes existence of solutions for all $t \ge 0$.

Definition 2. The periodic orbit \mathcal{O} of (4) is LES if there exists $a \ \delta > 0$, N > 0, and $\omega > 0$ such that $x(t) = \psi(t, x(0), 0, 0)$ satisfies for all $t \ge 0$,

$$\operatorname{dist}(x(t), \mathcal{O}) \le N \mathrm{e}^{-\omega t} \operatorname{dist}(x(0), \mathcal{O}) ,$$

for any $x(0) \in S^+$ with $dist(x(0), \mathcal{O}) < \delta$.

In addition to orbital stability, we also present notions of stability for the discrete system (7).

Definition 3. The system (7) is LISS if there exists a $\delta > 0$, $\alpha_1, \alpha_2 \in \mathcal{K}$, and $\beta \in \mathcal{KL}$, such that for all $k \in \mathbb{Z}_+$,

$$||x_k - x^*|| \le \beta(||x_0 - x^*||, k) + \alpha_1(||u||_{\infty}) + \alpha_2(||\bar{v}||_{\infty}) ,$$
(10)

is satisfied for any $x_0 \in S$ with $||x_0 - x^*|| < \delta$, $u \in U$ with $||u||_{\infty} < \delta$, and $\bar{v} \in V$ with $||\bar{v}||_{\infty} < \delta$.

Finally, the 0-input fixed point x^* of (7) satisfies $x^* = P(x^*, 0, 0)$, and it is locally asymptotically stable (LAS) or LES if it satisfies the following definition.

Definition 4. The 0-input fixed point x^* of (7) is LAS if there exists a $\delta > 0$ and $\beta \in \mathcal{KL}$ such that for all $k \in \mathbb{Z}^+$,

$$||x_k - x^*|| \le \beta(||x_0 - x^*||, k)$$

is satisfied for any $x_0 \in S$ with $||x_0 - x^*|| < \delta$. Furthermore, if there exists a N > 0, and $0 < \rho < 1$ such that

$$\beta(\|x_0 - x^*\|, k) \le N\rho^k \|x_0 - x^*\|$$

then x^* is a LES 0-input fixed point of (7).

III. MAIN RESULTS

In this section we present the main results of this paper. First, we introduce an important proposition on the geometric relation between \mathcal{O} and \mathcal{S} . This proposition allows us to express bounds on the orbital distance of any $x \in \mathcal{S}$ from \mathcal{O} equivalently based on the Euclidean distance of x from x^* , and vice-versa. The importance of this proposition becomes clear by observing the distance metrics used in Definition 1 and Definition 3. Hence, it serves as an important bridge between the orbital notions of stability and the Poincaré map's stability.

Proposition 1. Let S be defined as in (2) and satisfy assumption A.2. Let O be defined as in (5) and satisfy assumptions A.4-A.7. Then, there exists a $0 < \lambda < 1$ such that

$$\lambda \|x - x^*\| \le \operatorname{dist}(x, \mathcal{O}) \le \|x - x^*\| \quad (11)$$

for all $x \in S$.

The proof of Proposition 1 is detailed in Section IV below.

Proposition 1 can be used to show that solutions in a small enough neighborhood of a LISS periodic orbit O and for sufficiently small continuous and discrete input signals do not exhibit beating and Zeno behavior, and exist indefinitely. This statement is made precise by the following proposition, a proof of which can be found in Appendix A.

Proposition 2. Consider the system (4) which satisfies assumptions A.1-A.7. Suppose that the solutions of (4), denoted by $x(t) = \psi(t, x(0), u, \bar{v})$ and defined in Section II-B, satisfy Definition 1. Then, there exists a $\delta > 0$ such that for all $x(0) \in S^+$ with $\operatorname{dist}(x(0), \mathcal{O}) < \delta$, $u \in \mathcal{U}$ with $||u||_{\infty} < \delta$, and $\bar{v} \in \mathcal{V}$ with $||\bar{v}||_{\infty} < \delta$ the following holds:

- (i) x(t) has no consecutive discrete jumps,
- (ii) x(t) does not exhibit Zeno behavior, and
- (iii) x(t) exists for all $t \ge 0$.

Now we are ready to present the main result of the paper.

Theorem 1. Consider the system (4) which satisfies assumptions A.1-A.3 and possesses a periodic orbit O that is defined as in (5) and satisfies assumptions A.4-A.7. Then, the following are equivalent.

- (i) \mathcal{O} is an LISS orbit of (4);
- (ii) x^* is an LISS fixed point of (7).

It is straightforward to note that in the absence of inputs $(u \equiv 0, \bar{v} \equiv 0)$, Theorem 1 reduces to the Poincaré result for asymptotic stability of periodic orbits of systems with impulse effects, providing an alternative proof to [28, Theorem 1]. Note though that the proof detailed in the following sections explicitly constructs the class- \mathcal{KL} functions involved in the definitions, thereby providing useful insight on the rates of convergence. The following result can be stated as an immediate corollary of Theorem 1.

Corollary 1. Under the assumptions of Theorem 1, the following are equivalent

- (i) \mathcal{O} is an LES 0-input orbit of (4);
- (ii) x^* is an LES 0-input fixed point of (7).

The following remarks are in order.

Remark 3. The equivalence between ES of a periodic orbit and ES of the corresponding fixed point of the associated Poincaré map has been discussed in [29, Theorem 1], which has been subsequently used in a number of relevant publications, e.g., [7], [38]–[40], and many more. However, [29, Theorem 1] is proved only for initial states in the Poincaré section, as noted above [29, Equation (6)], rather than for initial states in a neighborhood of the entire orbit, as Definition 2 requires. Furthermore, Proposition 1, which is crucial for commuting between Definition 2 and Definition 4 is omitted in the proof of [29, Theorem 1], resulting in the estimate in [29, Equation (6)] being incomplete; the final estimate should have been expressed in terms of dist(x, O), which requires the use of Proposition 1.

Remark 4. It should be emphasized that the results of this paper can be used to study limit-cycle solutions of continuoustime forced systems like (1) by replacing the discrete update map Δ with the identity map for the x component and the zero map for the v component.

Theorem 1 can be used to establish LISS of a periodic orbit of (4) on the basis of LISS of a fixed point of the associated Poincaré map (7). However, in many applications-see Section VI for an example-the lack of analytical expressions for the forced Poincaré map makes it challenging to establish LISS for a fixed point of it. To alleviate this issue, the following theorem provides a tool for establishing that a 0-input fixed point x^* of the forced Poincaré map (7) is LISS by showing that x^* is a LAS fixed point of the unforced Poincaré map. Hence, one can simply linearize the unforced Poincaré map and compute the eigenvalues of the associated linearization. If all the eigenvalues lie within the unit disc, the corresponding fixed point is a LAS fixed point of the unforced Poincaré map. Then, Theorem 2 ensures that x^* is a LISS fixed point of the forced Poincaré map and Theorem 1 establishes LISS of the associated periodic orbit. A result similar to Theorem 2 can be found in [22] for continuous-time systems; however, to the best of our knowledge, we have not seen such a result for discrete systems and we provide it below.

Theorem 2. Consider the discrete dynamical system (7). Let $\delta > 0$ such that P is continuously differentiable in the Fréchet sense for $x \in B_{\delta}(x^*) \cap S$, $u \in U$ with $||u||_{\infty} < \delta$, and $v \in B_{\delta}(0)$. Then, the following are equivalent.

- (i) x^* is an LISS fixed point of (7);
- (ii) x^* is an LAS 0-input fixed point of (7).

A proof for Theorem 2 is presented in Section V.

IV. PROOF OF PROPOSITION 1

The proof of Proposition 1 is organized in a sequence of lemmas. We begin with a lemma which establishes that for \mathcal{O} , the point-to-set distance is equal to the minimum Euclidean distance over the closure of the orbit. As the minimum will be attained by some point in $\overline{\mathcal{O}}$, Lemma 3 allows us to work with the Euclidean distance from that point instead of dealing with $\inf_{y \in \mathcal{O}} ||x - y||$.

Lemma 3. Let \mathcal{O} be defined as in (5) and satisfy assumptions A.4-A.7, then for all $x \in \mathbb{R}^n$, we have

$$\operatorname{dist}(x, \mathcal{O}) := \inf_{y \in \mathcal{O}} \|x - y\| = \min_{y \in \overline{\mathcal{O}}} \|x - y\|$$

Proof. Let $x \in \mathbb{R}^n$. The fact that $\overline{\mathcal{O}} = \mathcal{O} \cup \{x^*\}$ implies

$$\min_{y \in \overline{O}} \|x - y\| = \min\{\|x - x^*\|, \inf_{y \in O} \|x - y\|\} .$$
(12)

On the other hand,

$$\inf_{y \in \mathcal{O}} \|x - y\| \le \inf_{y \in \mathcal{O}} (\|x - x^*\| + \|x^* - y\|) = \|x - x^*\| \quad (13)$$

because $\inf_{y \in \mathcal{O}} ||x^* - y|| = 0$ due to $x^* \in \overline{\mathcal{O}}$. The result follows from (12) in view of (13).

To simplify notation in the proofs that follow, the closure of the orbit $\overline{\mathcal{O}}$ is parameterized as a function $y(\tau)$ in "time" like coordinates τ taking values in a closed interval [0,T]. In more detail, let $\varphi^{-}(t, x^*, 0)$ be the solution of the 0-input continuous system (1) backwards in time from the initial state x^* . The flow is chosen to be backwards so that x^* is at $\tau = 0$. This is primarily for convenience of notation; the flow can be chosen forwards in time starting from $\Delta(x^*, 0)$ as well. Then, let $\tau := t/s$, where s > 0 is a scaling constant, and define the function $y : [0,T] \to \overline{\mathcal{O}}$ by

$$y(\tau) := \varphi^{-}(s\tau, x^*, 0) \tag{14}$$

where

$$T := T^*/s \quad . \tag{15}$$

The scaling is performed to ensure that in the Taylor expansion of $y(\tau)$ about $\tau = 0$, the first derivative is a vector of unit magnitude. This is done only to simplify notation in the future. Note that $y(\tau)$ should be viewed as a parameterization of the set $\overline{\mathcal{O}}$ and *not* as a solution of the system Σ in (4). In fact, this section only deals with geometric properties of \mathcal{O} and S and does not study the dynamical system as such. The following lemma provides some useful properties of $y(\tau)$.

Lemma 4. The map $\tau \mapsto y(\tau)$ is bijective and three-times continuously differentiable in τ .

The proof of Lemma 4 can be found in Appendix B.

Let⁵ $\tau_{\rm m}: \mathbb{R}^n \to \mathcal{P}(\overline{\mathcal{O}})$ be a set-valued map defined as

$$\tau_{\rm m}(x) := \arg\min_{\tau \in [0,T]} \|x - y(\tau)\| ,$$
(16)

where $x \in \mathbb{R}^n$ and $y(\tau)$, T as defined in (14) and (15), respectively. Intuitively, the map $x \mapsto \tau_m(x)$ returns the *set* $\tau_m(x)$ of "times" τ that "realize" the points $y(\tau)$ on \overline{O} that are nearest to x. Hence, for any $\tau_{\min} \in \tau_m(x)$, we have

$$||x - y(\tau_{\min})|| \le ||x - y(\tau)||$$
 for all $\tau \in [0, T]$. (17)

The next lemma shows that by selecting x sufficiently close to x^* , the points on $\overline{\mathcal{O}}$ nearest to x also remain close to x^* .

Lemma 5. Let $\tau_{\rm m}$ be defined as in (16). Then, for every $\epsilon > 0$ there exists $\delta > 0$ such that $||x - x^*|| < \delta$ implies $\tau_{\rm min} < \epsilon$ for all $\tau_{\rm min} \in \tau_{\rm m}(x)$.

⁵Recall from Section II-A that $\mathcal{P}(\cdot)$ is the power set of its argument.

6

Proof. Let $0 < \epsilon < T$. The map $y(\tau)$ is continuous, so $y([\epsilon, T])$ is compact (and thus closed) in \mathbb{R}^n . Hence, its complement $y([\epsilon, T])^c$ in \mathbb{R}^n is open and contains $y(0) = x^*$ using the injectivity of $y(\tau)$ from Lemma 4. As a result, there exists a $\delta > 0$ such that $B_{2\delta}(x^*) \subset y([\epsilon, T])^c$.

It can be seen that for any $x \in B_{\delta}(x^*)$, the points on $\overline{\mathcal{O}}$ closest to x will be within $B_{2\delta}(x^*)$. This follows by a contradiction argument. Indeed, take any $x \in B_{\delta}(x^*)$ and let τ_{\min} be any element of the set⁶ $\tau_{\mathrm{m}}(x)$ defined in (16). Assume $y(\tau_{\min})$ is outside $B_{2\delta}(x^*)$ so that $||x - y(\tau_{\min})|| \ge \delta$ by the reverse triangle inequality. But, since $y(0) = x^* \in \overline{\mathcal{O}}$, by (17) we have $||x - y(\tau_{\min})|| \le ||x - x^*|| < \delta$, thus resulting in a contradiction. It follows that for any $\tau_{\min} \in \tau_{\mathrm{m}}(x)$, $y(\tau_{\min}) \in B_{2\delta}(x^*) \subset y([\epsilon, T])^c$, and thus $\tau_{\min} < \epsilon$ as a consequence of the injectivity of $y(\tau)$ by Lemma 4. As this holds for any $0 < \epsilon < T$, the result trivially holds $\forall \epsilon > 0$. \Box

Next, we present a lemma which shows that the lower bound of Proposition 1 holds locally around x^* .

Lemma 6. Let S be defined as in (2) and satisfy assumption A.2. Let O be as in (5) and satisfy assumptions A.4-A.7. Then, there exists $\delta > 0$ and $0 < \overline{\lambda} < 1$ such that

$$\operatorname{dist}(x, \mathcal{O}) \ge \lambda \|x - x^*\| \quad :$$

for all $x \in B_{\delta}(x^*) \cap S$.

Proof. This proof is structured as follows. We begin by establishing the desired inequality for states restricted to the vector spaces $T_{x^*}\mathcal{O}$ (tangent line to \mathcal{O} at x^*) and $T_{x^*}\mathcal{S}$ (tangent plane to \mathcal{S} at x^*) and subsequently introduce non-linearities one-by-one. First, we extend the result to \mathcal{O} and $T_{x^*}\mathcal{S}$, and finally, we extend the result to \mathcal{O} and \mathcal{S} . A geometric illustration of the setup can be seen in Fig. 2.

Performing Taylor's expansion [37, p. 349] of $y(\tau)$ about $\tau = 0$, we get

$$y(\tau) = x^* + \tau\nu + \tau^2 r(\tau) \tag{18}$$

where ν is a unit vector and $\tau^2 r(\tau)$ is the remainder. The scaling factor s in the definition of τ above (14) is chosen to ensure that ν has unit length. Let $y'(\tau)$, $r'(\tau)$ be shorthand for $dy/d\tau$ and $dr/d\tau$, respectively. As $y(\tau)$ is three-times continuously differentiable by Lemma 4, $r(\tau)$ and $r'(\tau)$ are continuous on $\tau \in [0, T]$, hence there exists $M_r \ge 0$ such that $\|r(\tau)\| \le M_r$ and $\|r'(\tau)\| \le M_r$ for all $\tau \in [0, T]$. (i) $T_{x^*}\mathcal{O}$ and $T_{x^*}\mathcal{S}$

The tangent line of \mathcal{O} at x^* is $T_{x^*}\mathcal{O} := \{x^* + \tau\nu \mid \tau \in \mathbb{R}\}$ as illustrated in Fig. 2. Given any $z \in T_{x^*}\mathcal{S}$, we have $\operatorname{dist}(z, T_{x^*}\mathcal{O}) = \inf_{\tau \in \mathbb{R}} ||z - (x^* + \tau\nu)||$ and the point on $T_{x^*}\mathcal{O}$ closest to z can be obtained by projecting the vector $z - x^*$ along the unit vector ν . Specifically, the point on $T_{x^*}\mathcal{O}$ closest to z is given by $x^* + \hat{\tau}_{\min}(z)\nu$ with

$$\hat{\tau}_{\min}(z) = \langle z - x^*, \nu \rangle \quad , \tag{19}$$

where $\langle \cdot, \cdot \rangle$ represents the inner product in \mathbb{R}^n . Consider now the right triangle with vertices at x^* , z, and $x^* + \hat{\tau}_{\min}(z)\nu$. Then, $||z - (x^* + \hat{\tau}_{\min}(z)\nu)|| = ||z - x^*|| \cdot |\sin(\theta(z))|$, where

⁶It is straightforward to note that $\tau_{\rm m}(x) \neq \emptyset$ since $\overline{\mathcal{O}} \neq \emptyset$ and compact.



Fig. 2. The switching surface S is in grey, the tangent plane $T_{x^*}S$ is in light grey, the curved line is the orbit O, and the dashed line is $T_{x^*}O$.

 $\theta(z)$ is the angle between $z - x^*$ and ν . By transversality of \mathcal{O} and \mathcal{S} at x^* given by assumption A.7, $\theta(z)$ will never be 0 or π , so $\min_{\|z-x^*\|=1} |\sin(\theta(z))| =: \mu$ satisfies $0 < \mu \leq 1$. Thus, for all $z \in T_{x^*}\mathcal{S}$, we have

$$dist(z, T_{x^*}\mathcal{O}) = \inf_{\tau \in \mathbb{R}} ||z - (x^* + \tau\nu)|| = ||z - (x^* + \hat{\tau}_{\min}(z)\nu)|$$
$$= ||z - x^*|| \cdot |\sin(\theta(z))| \ge \mu ||z - x^*|| \quad (20)$$

(ii) \mathcal{O} and $T_{x^*}\mathcal{S}$

Now we extend the result to \mathcal{O} and $T_{x^*}\mathcal{S}$. Choose $\delta > 0$ such that $z \in B_{\delta}(x^*) \cap T_{x^*}\mathcal{S}$ implies $\tau_{\min} < T$ with T as in (15) for all $\tau_{\min} \in \tau_{\mathrm{m}}(z)$; Lemma 5 guarantees that such a δ exists. Next, split the set $B_{\delta}(x^*) \cap T_{x^*}\mathcal{S}$ into two subsets:

(a) $E_1 := \{z \in B_{\delta}(x^*) \cap T_{x^*}\mathcal{S} \mid \exists \tau_{\min} \in \tau_{\mathrm{m}}(z) : \tau_{\min} = 0\}$ (b) $E_2 := \{z \in B_{\delta}(x^*) \cap T_{x^*}\mathcal{S} \mid \forall \tau_{\min} \in \tau_{\mathrm{m}}(z), \tau_{\min} \neq 0\}.$ Clearly, E_1 and E_2 are disjoint and $E_1 \cup E_2 = B_{\delta}(x^*) \cap T_{x^*}\mathcal{S}.$ If $z \in E_1$, then $0 \in \tau_{\mathrm{m}}(z)$ and thus

dist
$$(z, \mathcal{O}) = \inf_{y \in \mathcal{O}} ||z - y|| = ||z - y(\tau_{\min})|| = ||z - x^*||$$
. (21)

For convenience, from here on when we use τ_{\min} , it is understood that τ_{\min} can be any element of $\tau_{m}(z)$. Also, we will drop the functional dependence of $\hat{\tau}_{\min}$ on z; see (19).

If $z \in E_2$ then $\tau_{\min} > 0$ and the vector from z to the nearest point on \mathcal{O} must be orthogonal to the orbit. Hence, we have $\langle z - y(\tau_{\min}), y'(\tau_{\min}) \rangle = 0$, which, on using (18), gives

$$\langle z - x^* - \tau_{\min}\nu - \tau_{\min}^2 r(\tau_{\min}),$$

$$\nu + 2\tau_{\min}r(\tau_{\min}) + \tau_{\min}^2 r'(\tau_{\min}) \rangle = 0 \quad . \tag{22}$$

Next, we use (22) to derive the following important estimate,

$$\tau_{\min} = \langle z - x^*, \nu \rangle + 2\tau_{\min} \langle r(\tau_{\min}), z - x^* \rangle + \tau_{\min}^2 a(\tau_{\min}, z) ,$$
(23)
where $a(\tau, z) := \langle r'(\tau), z - x^* \rangle - \langle 3\nu, r(\tau) \rangle - \tau(\langle \nu, r'(\tau) \rangle + \langle 2r(\tau), r(\tau) \rangle) - \tau^2 \langle r(\tau), r'(\tau) \rangle.$ Since $r(\tau)$ and $r'(\tau)$ are bounded as discussed below (18), $z \in B_{\delta}(x^*) \cap T_{x^*}\mathcal{S}$, and $\tau_{\min} < T$, we have, for some constant $c > 0$, that

$$\tau_{\min} \le \|z - x^*\| + c\tau_{\min}(\|z - x^*\| + \tau_{\min}).$$
(24)

Using $\epsilon = 1/(4c)$ in Lemma 5 there exists a $\delta < 1/(4c)$ (shrink δ if necessary) such that for $||z - x^*|| < \delta$, we have $\tau_{\min} < 1/(4c)$. Then from (24) we have

$$\tau_{\min} \le 2 \|z - x^*\|.$$
 (25)

Next, noting $\hat{\tau}_{\min} = \langle z - x^*, \nu \rangle$ by (19), from (23) we obtain

$$\begin{aligned} |\hat{\tau}_{\min} - \tau_{\min}| &\leq |2\tau_{\min}\langle r(\tau_{\min}), z - x^*\rangle + \tau_{\min}^2 a(\tau_{\min}, z)| \\ &\leq c(\tau_{\min}||z - x^*|| + \tau_{\min}^2) \end{aligned}$$

where the last inequality follows by using bounds similar to those that led to (24). Using (25) in the above inequality and updating the constant⁷ c > 0 accordingly, we have

$$|\hat{\tau}_{\min} - \tau_{\min}| \le c ||z - x^*||^2$$
, (26)

provided $||z - x^*|| < \delta$.

Turning our attention to dist(z, O) and using Lemma 3,

$$\inf_{y \in \mathcal{O}} \|z - y\| = \|z - y(\tau_{\min})\| \\
= \|z - x^* - \tau_{\min}\nu - \tau_{\min}^2 r(\tau_{\min})\|$$
(27)

$$= \|z - x^* - \hat{\tau}_{\min}\nu + (\hat{\tau}_{\min} - \tau_{\min})\nu - \tau_{\min}^2 r(\tau_{\min})\| \quad (28)$$

$$\geq \|z - x^* - \hat{\tau}_{\min}\nu\| - |\hat{\tau}_{\min} - \tau_{\min}| - \tau_{\min}^2 \|r(\tau_{\min})\|$$
(29)
$$\geq \mu \|z - x^*\| - c\|z - x^*\|^2 ,$$
(30)

where (27) is obtained by using (18); (28) is obtained by adding and subtracting $\hat{\tau}_{\min}\nu$; (29) is obtained by using the reverse triangle inequality; and (30) is obtained by the boundedness of $r(\tau)$, (20), (26), and (25). Again we update the constant c > 0 accordingly. Further we can write (30) as

$$\inf_{y \in \mathcal{O}} \|z - y\| \ge (\mu/2) \|z - x^*\| + \|z - x^*\| (\mu/2 - c\|z - x^*\|)$$

Choosing $||z - x^*|| \le \mu/(2c)$ (shrink $\delta > 0$ if necessary) gives

$$dist(z, \mathcal{O}) := \inf_{y \in \mathcal{O}} \|z - y\| \ge (\mu/2) \|z - x^*\| \quad , \qquad (31)$$

for all $z \in E_2$.

Putting together (21) and (31) for the sets E_1 and E_2 , respectively, and noting that $\min\{1, \mu/2\} = \mu/2$ as $\mu/2 \le 1/2 < 1$ (see below (19) to recall the meaning of μ) gives

$$dist(z, \mathcal{O}) := \inf_{y \in \mathcal{O}} \|z - y\| \ge (\mu/2) \|z - x^*\| \quad , \qquad (32)$$

for all $z \in B_{\delta}(x^*) \cap T_{x^*}S$. (*iii*) \mathcal{O} and S

Here, we extend the result to $x \in B_{\delta}(x^*) \cap S$. First, note that if $x \in S$ is a point in the neighborhood of x^* and $z \in T_{x^*}S$ is the projection of x on $T_{x^*}S$, then Appendix C shows that there exists a constant c > 0 such that

$$||x - z|| \le c ||x - x^*||^2 .$$
(33)

Since $||z - x^*|| = ||z - x + x - x^*|| \le ||x - x^*|| + c||x - x^*||^2$ by the triangle inequality and (33), choosing $x \in S$ so that $||x - x^*|| < \delta$ for a sufficiently small $\delta > 0$ ensures that the corresponding z satisfies (32). Then,

$$\inf_{y \in \mathcal{O}} \|x - y\| = \inf_{y \in \mathcal{O}} \|x - z + z - y\|$$

$$\geq \inf_{y \in \mathcal{O}} \|z - y\| - \|x - z\|$$

$$\geq (\mu/2) \|z - x^*\| - \|x - z\| \qquad (34)$$

$$\geq (\mu/2) \|x - x^*\| - (1 + \mu/2) \|x - z\| \qquad (35)$$

$$\geq (\mu/2) \|x - x^*\| - c\|x - x^*\|^2 \quad . \tag{36}$$

where (34) follows from (32); (35) from the reverse triangle inequality on $||z - x^*|| = ||z - x + x - x^*||$; and (36) follows from (33) with c > 0 updated accordingly. Write (36) as

$$\inf_{y \in \mathcal{O}} \|x - y\| \ge (\mu/4) \|x - x^*\| + \|x - x^*\| (\mu/4 - c\|x - x^*\|),$$

⁷Intermediate constants of no particular importance are used as c while updating the meaning of c as we proceed with the proof.

and choose $||x - x^*|| \le \mu/(4c)$, shrinking δ if necessary. The result follows by letting $\overline{\lambda} = \mu/4 < 1$.

Now we present the proof of Proposition 1, which essentially extends Lemma 6 to the entire S.

Proof of Proposition 1. The upper bound on $dist(x, \mathcal{O})$ in Proposition 1 follows, for all $x \in S$, directly from (13).

For the lower bound, we begin by applying Lemma 6 to establish the existence of $\delta > 0$ and $0 < \overline{\lambda} < 1$ such that

$$\operatorname{dist}(x, \mathcal{O}) \ge \bar{\lambda} \|x - x^*\| \quad , \tag{37}$$

for all $x \in S$ with $||x - x^*|| < \delta$. To obtain a lower bound that holds for all $x \in S$, we first consider the case where S is unbounded; then, the case where S is bounded follows easily.

Let S be unbounded and distinguish the following regions. (i) $\mathcal{R}_{I} := \{x \in S \mid ||x - x^{*}|| > \delta'\}$ for $\delta' > \delta$

We will show that a $\delta' > \delta$ exists so that for all $x \in \mathcal{R}_{I}$ a lower bound for dist (x, \mathcal{O}) similar to (37) can be found. First note that, by the definition (2), the surface S is closed. Furthermore, by assumption A.6 we have $\overline{\mathcal{O}} \cap S = \{x^*\}$, and thus the only limit point that \mathcal{O} and S share is x^* . Hence, dist $(x, \mathcal{O}) > 0$ for all $x \in S \setminus \{x^*\}$, as these points are in the complement of the closure of \mathcal{O} in \mathbb{R}^n , and dist $(x, \mathcal{O})/||x - x^*|| > 0$ is well defined for all $x \in S \setminus \{x^*\}$. We claim that

$$\lim_{\|x\|\to\infty, x\in\mathcal{S}\setminus\{x^*\}} \operatorname{dist}(x,\mathcal{O})/\|x-x^*\| = 1 \quad , \qquad (38)$$

from which it follows easily that there exists $\delta' > 0$ (expand δ' if necessary to ensure $\delta' > \delta$) such that

$$dist(x, \mathcal{O}) \ge (1/2) ||x - x^*||$$
, (39)

for all $x \in \mathcal{R}_{\mathrm{I}}$. To show the claim (38), take any $x \in \mathcal{S} \setminus \{x^*\}$, let $\tau_{\min} \in \tau_{\mathrm{m}}(x)$ and define $M_{\mathcal{O}} := \max_{y_1, y_2 \in \overline{\mathcal{O}}} \|y_1 - y_2\|$ so that $\|x^* - y(\tau_{\min})\| \leq M_{\mathcal{O}}$. Then, $\operatorname{dist}(x, \mathcal{O}) := \|x - y(\tau_{\min})\| \geq \|x - x^*\| - \|x^* - y(\tau_{\min})\|$ implies

$$\frac{\operatorname{dist}(x,\mathcal{O})}{\|x-x^*\|} \ge 1 - \frac{\|x^* - y(\tau_{\min})\|}{\|x-x^*\|} \ge 1 - \frac{M_{\mathcal{O}}}{\|x-x^*\|} , \quad (40)$$

and dist $(x, \mathcal{O}) = ||x - y(\tau_{\min})|| \le ||x - x^*|| + ||x^* - y(\tau_{\min})||$ implies

$$\frac{\operatorname{dist}(x,\mathcal{O})}{\|x-x^*\|} \le 1 + \frac{\|x^* - y(\tau_{\min})\|}{\|x-x^*\|} \le 1 + \frac{M_{\mathcal{O}}}{\|x-x^*\|} \quad .$$
(41)

As a result, for any sequence of points $x_n \in S \setminus \{x^*\}$ such that $||x_n - x^*|| \to \infty$, it follows from (40) and (41) that

$$1 \le \liminf_{n \to \infty} \frac{\operatorname{dist}(x_n, \mathcal{O})}{\|x_n - x^*\|} \le \limsup_{n \to \infty} \frac{\operatorname{dist}(x_n, \mathcal{O})}{\|x_n - x^*\|} \le 1 ,$$

implying $\lim_{n\to\infty} \operatorname{dist}(x_n, \mathcal{O})/||x_n - x^*|| = 1$, which by [41, Theorem 4.2] proves the claim (38).

(ii) $\mathcal{R}_{\mathrm{II}} := \{ x \in \mathcal{S} \mid \delta \le ||x - x^*|| \le \delta' \}$

With $\delta > 0$ provided by Lemma 6 and $\delta' > \delta$ selected as in case (i), let $\hat{\lambda} := \min_{x \in \mathcal{R}_{\mathrm{II}}} \left(\operatorname{dist}(x, \mathcal{O}) / ||x - x^*|| \right) > 0$, which is well defined since $\operatorname{dist}(x, \mathcal{O}) / ||x - x^*|| > 0$ is continuous over the compact set $\mathcal{R}_{\mathrm{II}}$. Hence,

$$\operatorname{dist}(x, \mathcal{O}) \ge \lambda \|x - x^*\| \quad , \tag{42}$$

for all $x \in \mathcal{R}_{II}$.

Finally, combining (37), (39), and (42) by choosing $\lambda = \min{\{\bar{\lambda}, 1/2, \hat{\lambda}\}}$ gives

$$\operatorname{dist}(x, \mathcal{O}) := \inf_{y \in \mathcal{O}} \|x - y\| \ge \lambda \|x - x^*\|,$$

for all $x \in S$, completing the proof when S is unbounded.

For the case where S is bounded, we can choose $\delta' > 0$ sufficiently large to ensure that $S \subset B_{\delta'}(x^*)$. Then, an argument analogous to that used in *(ii)* above establishes the desired lower bound for this case, thereby completing the proof of Proposition 1.

V. PROOF OF THEOREM 1 AND THEOREM 2

In the proof of Theorem 1, we will compare different solutions based on their initial states and inputs. With this in mind, we present the following lemma, which is a straightforward adaptation of [34, Theorem 3.5] and its proof will be omitted.

Lemma 7. Suppose f in (1) satisfies assumption A.1. Let $u_1 \in U$ and $x_1(t)$ be the solution of

$$\dot{x}(t) = f(x(t), u_1(t)), \quad x_1(0) = a_1$$

which exists for all $t \in [0, \hat{T}]$. Then, there exists L > 0 and $\delta > 0$ such that if $||a_2 - a_1|| < \delta$ and $||u_2 - u_1||_{\infty} < \delta$,

 $\dot{x}(t) = f(x(t), u_2(t)), \quad x_2(0) = a_2$

has a unique solution $x_2(t)$ for all $t \in [0, \hat{T}]$. Further,

$$||x_1(t) - x_2(t)|| \le e^{LT} ||a_1 - a_2|| + (e^{LT} - 1)||u_1 - u_2||_{\infty},$$
(43)

for all $t \in [0, \hat{T}]$.

The following remark extends the unperturbed (zero-input) solution $\varphi(t, \Delta(x^*, 0), 0)$ of (1) that starts from $\Delta(x^*, 0)$.

Remark 5. By assumption A.5, the solution $\varphi(t, \Delta(x^*, 0), 0)$ of (1) exists for all $t \in [0, T^*]$, where $T^* = T_I(x^*, 0, 0)$ is the period of \mathcal{O} . Hence, by [34, Theorem 3.1], there exists $\overline{T} \in (T^*, \infty)$ so that $\varphi(t, \Delta(x^*, 0), 0)$ can be extended over the interval $[T^*, \overline{T}]$. As $L_f H(\varphi(T^*, \Delta(x^*, 0), 0), 0) < 0$ from assumption A.7 while $L_f H(\varphi(t, \Delta(x^*, 0), 0), 0)$ is continuous in time, it follows that for a sufficiently small $\overline{T} > T^*$, we can ensure $L_f H(\varphi(t, \Delta(x^*, 0), 0), 0) < 0$ for all $T^* \leq t \leq \overline{T}$, so $\varphi(t, \Delta(x^*, 0), 0) \in S^-$ over the interval $(T^*, \overline{T}]$.

Lemma 8 below shows that the perturbed solution $\varphi(t, \Delta(x, v), u)$ of (1) is locally well defined over the interval $[0, \overline{T}]$ where the unperturbed solution $\varphi(t, \Delta(x^*, 0), 0)$ can be extended. Moreover, the lemma provides a *linear* upper bound on the distance of $\varphi(t, \Delta(x, v), u)$ from \mathcal{O} that will be used in proving Theorem 1. The proof of Lemma 8 is in Appendix D.

Lemma 8. Under the assumptions of Theorem 1, there exist $\delta > 0$, and $\underline{T} > 0$, $\overline{T} > T^*$ with $T^* = T_I(x^*, 0, 0)$ being the period of \mathcal{O} , such that for all $x \in B_{\delta}(x^*) \cap \mathcal{S}$, $u \in \mathcal{U}$ with $\|u\|_{\infty} < \delta$, and $\overline{v} \in \mathcal{V}$ with $\|\overline{v}\|_{\infty} < \delta$ the following hold

- (i) The perturbed solution $\varphi(t, \Delta(x, v), u)$ of (1), where v is an element of \overline{v} , exists and is unique for $t \in [0, \overline{T}]$.
- (ii) The solution $\varphi(t, \Delta(x, v), u)$ crosses S in finite time $T_{\rm I}(x, u, v)$ given by (6), and $0 < \underline{T} < T_{\rm I}(x, u, v) < \overline{T}$,

where $\underline{T} < T^* < \overline{T}$. Moreover, it does so transversally to S with $L_f H(\varphi(T_I(x, u, v), \Delta(x, v), u), u) < 0$. (iii) There exists a c > 0 such that

$$\sup_{0 \le t < T_{\mathrm{I}}} \operatorname{dist}(\varphi(t, \Delta(x, v), u), \mathcal{O}) \le c \|x - x^*\| + c \|u\|_{\infty} + c \|\bar{v}\|_{\infty}.$$
(44)

In proving Theorem 1, we will also need Lemma 9 which is an "orbital" analogue of Lemma 8. Lemma 9 does not require x to be confined on S so that any $x \in S^+$ can be used, as long as it is sufficiently close to \mathcal{O} .

Lemma 9. Under the assumptions of Theorem 1, there exist $\delta > 0$ and $\overline{T} > T^*$ with $T^* = T_I(x^*, 0, 0)$ being the period of \mathcal{O} , such that for all $x \in S^+$ with $\operatorname{dist}(x, \mathcal{O}) < \delta$ and $u \in \mathcal{U}$ with $||u||_{\infty} < \delta$, the following hold

- (i) The perturbed solution $\varphi(t, x, u)$ of (1) exists and is unique for $t \in [0, \overline{T} \xi]$ where $\xi \in [0, T^*]$.
- (ii) The solution $\varphi(t, x, u)$ crosses S in finite time $\hat{T}_{I}(x, u)$ given by (8), and $0 < \hat{T}_{I}(x, u) < \overline{T}$. Moreover, it does so transversally to S with $L_{f}H(\varphi(\hat{T}_{I}(x, u), x, u), u) < 0$.
- (iii) There exists a c > 0 such that

$$\sup_{0 \le t < \hat{T}_{\mathbf{I}}} \operatorname{dist}(\varphi(t, x, u), \mathcal{O}) \le c \operatorname{dist}(x, \mathcal{O}) + c \|u\|_{\infty}.$$
(45)

The proof of Lemma 9 is similar to the proof of Lemma 8, albeit more technical as it requires the construction of a suitable open cover for $\overline{\mathcal{O}}$; thus, it is relegated to Appendix D.

Now we are ready to provide the proof of Theorem 1.

Proof of Theorem 1. We first show (i) \implies (ii) and then (ii) \implies (i) by explicitly constructing class- \mathcal{KL} and class- \mathcal{K} functions to satisfy Definitions 1 and 3. To avoid ambiguity in notation, we use $x(0) \in \mathbb{R}^n$ as the initial condition for the system with impulse effects (4) and $x_0 \in S$ as the initial condition for the discrete system (7).

 $(i) \Longrightarrow (ii)$

Assume that \mathcal{O} is a LISS orbit of (4), and let $x(t) = \psi(t, x(0), u, \bar{v})$ be a solution of (4) that satisfies Definition 1 for some $\delta_{\Sigma} > 0$ and for suitable functions $\alpha_1, \alpha_2 \in \mathcal{K}$, $\beta \in \mathcal{KL}$. Choose δ_{Σ} sufficiently small to further ensure that Lemma 9 is satisfied. Then, $t_0 := \hat{T}_1(x(0), u)$ is finite, and $x_0 := \lim_{t \neq t_0} x(t)$ is well-defined. On arriving at \mathcal{S} , the solution jumps to $x(t_0) = \Delta(x_0, v_0)$ where v_0 is the first element of the sequence \bar{v} . However, to establish the estimate (10) in Definition 3 we need x_0 to appear in the RHS instead of $x(t_0)$. To do this we will use the fact that, by assumption A.3, the map Δ is continuously differentiable and thus locally Lipschitz. Hence, there exists a $\delta_{\Delta} > 0$ for which the Lipschitz condition holds uniformly for all $x \in B_{\delta_{\Delta}}(x^*) \cap \mathcal{S}$ and $\|\bar{v}\|_{\infty} < \delta_{\Delta}$ for some constant $L_{\Delta} > 0$, so that

$$dist(\Delta(x_0, v_0), \mathcal{O}) = dist(\Delta(x_0, v_0), \mathcal{O}) - dist(\Delta(x^*, 0), \mathcal{O})$$

$$\leq \|\Delta(x_0, v_0) - \Delta(x^*, 0)\|$$

$$\leq L_{\Delta}(\|x_0 - x^*\| + \|\bar{v}\|_{\infty}) , \qquad (46)$$

where we used $dist(\Delta(x^*, 0), \mathcal{O}) = 0$ and the fact that $dist(\cdot, \mathcal{O})$ is Lipschitz continuous with constant equal to 1. Finally, to guarantee that the time to impact is well defined for subsequent intersections of x(t) with S, choose $\delta_T > 0$ so that Lemma 8 is satisfied for T > 0 and $\overline{T} > T^*$.

Now we refine the selection of δ so that the aforementioned properties hold uniformly along the entire orbit. Pick $0 < \delta < \min\{\delta_{\Sigma}, \delta_T, \delta_{\Delta}\}$ sufficiently small to also ensure that $\beta(\delta, 0) + \alpha_1(\delta) + \alpha_2(\delta) < \min\{\delta_{\Sigma}, \lambda\delta_T, \lambda\delta_{\Delta}\}$; here, by Proposition 1, $\lambda \in (0, 1)$ is a constant such that (11) holds for all $x \in S$. Then, for dist $(x(0), \mathcal{O}) < \delta$, $||u||_{\infty} < \delta$, and $||\bar{v}||_{\infty} < \delta$,

$$dist(x(t), \mathcal{O}) \leq \beta(dist(x(0), \mathcal{O}), t) + \alpha_1(||u||_{\infty}) + \alpha_2(||\bar{v}||_{\infty})$$
$$\leq \beta(\delta, 0) + \alpha_1(\delta) + \alpha_2(\delta)$$
$$< \min\{\delta_{\Sigma}, \lambda\delta_T, \lambda\delta_{\Delta}\}$$

for all $t \geq 0$. Now, by continuity of the distance function, $\lim_{t \nearrow t_k} \operatorname{dist}(x(t), \mathcal{O}) = \operatorname{dist}(x_k, \mathcal{O})$, and this choice of $\delta > 0$ ensures that $\operatorname{dist}(x_k, \mathcal{O}) < \lambda \delta_{\Delta}$. Hence, by Proposition 1, we get $||x_k - x^*|| < \delta_{\Delta}$ for all $k \in \mathbb{Z}_+$ and, since this δ also guarantees $||\bar{v}||_{\infty} < \delta_{\Delta}$, the bound (46) holds for all $k \in \mathbb{Z}_+$. Similarly, this choice of δ ensures $||x_k - x^*|| < \delta_T$ for all $k \in \mathbb{Z}_+$, ensuring that $\underline{T} < T_{\mathrm{I}}(x_k, u_k, v_k) < \overline{T}$ uniformly for all $k \in \mathbb{Z}_+$. Finally, this δ guarantees that $\operatorname{dist}(x_0, \mathcal{O}) < \delta_{\Sigma}$, allowing the use of (9) with the same β , α_1 , α_2 as above. Putting all these together, for all $k \in \mathbb{Z}_+$ we can write

$$\begin{aligned} &|x_{k} - x^{*}\| \\ &\leq \lambda^{-1} \operatorname{dist}(x_{k}, \mathcal{O}) \\ &\leq \lambda^{-1}(\beta(\operatorname{dist}(\Delta(x_{0}, v_{0}), \mathcal{O}), t_{k} - t_{0}) + \alpha_{1}(\|u\|_{\infty}) \\ &+ \alpha_{2}(\|\bar{v}\|_{\infty})) \\ &\leq \lambda^{-1}(\beta(L_{\Delta}(\|x_{0} - x^{*}\| + \|\bar{v}\|_{\infty}), t_{k} - t_{0}) + \alpha_{1}(\|u\|_{\infty}) \\ &+ \alpha_{2}(\|\bar{v}\|_{\infty})) \\ &\leq \lambda^{-1}(\beta(2L_{\Delta}\|x_{0} - x^{*}\|, t_{k} - t_{0}) + \alpha_{1}(\|u\|_{\infty}) \\ &+ \alpha_{2}(\|\bar{v}\|_{\infty})) \\ &\leq \lambda^{-1}(\beta(2L_{\Delta}\|x_{0} - x^{*}\|, k\underline{T}) + \alpha_{1}(\|u\|_{\infty}) + \alpha_{2}(\|\bar{v}\|_{\infty})) \end{aligned}$$

where the first inequality follows from Proposition 1; the second from (9) with the solution starting from $\Delta(x_0, v_0)$; the third from (46); the fourth follows by [42, Lemma 14] on $\beta(L_{\Delta}(||x_0 - x^*|| + ||\bar{v}||_{\infty}), t_k - t_0)$ followed by absorbing the $||\bar{v}||_{\infty}$ term in α_2 ; and the last inequality follows from the fact that β is monotonically decreasing in time and $t_{k+1} - t_k = T_1(x_k, u_k, v_k) \geq \underline{T}$ for all $k \in \mathbb{Z}_+$. Noting that $(1/\lambda)\beta \in \mathcal{KL}$, $(1/\lambda)\alpha_1 \in \mathcal{K}$ and $(1/\lambda)\alpha_2 \in \mathcal{K}$, and comparing the last inequality with (10), we get that the solution of (7) satisfies Definition 3, thus completing the proof. (ii) \Longrightarrow (i)

Assume that x^* is a LISS fixed point of (7), and let $\{x_k\}_{k\in\mathbb{Z}_+}$ with $x_k \in S$ for all $k \in \mathbb{Z}_+$ be a solution of (7) that satisfies Definition 3 for some $\delta_1 > 0$. Then, for $x_0 \in B_{\delta_1}(x^*) \cap S$, $\|u\|_{\infty} < \delta_1$, and $\|\bar{v}\|_{\infty} < \delta_1$, there exist suitable functions $\alpha_1, \alpha_2 \in \mathcal{K}, \beta \in \mathcal{KL}$ such that (10) is satisfied. This implies $\|x_k - x^*\| \leq \beta(\delta_1, 0) + \alpha_1(\delta_1) + \alpha_2(\delta_1)$ for all $k \in \mathbb{Z}_+$, and thus δ_1 can be chosen (shrinking it if necessary) so that Lemma 8 is satisfied for $\varphi(t, \Delta(x_k, v_k), u_k)$ for all integers $k \geq 0$. Then, there exist $\overline{T}_1 > T^*$ (obtained as in Remark 5) so that $T_1(x_k, u_k, v_k) < \overline{T}_1$ for all $k \in \mathbb{Z}_+$.

The setting above provides a uniform (over k) upper bound \overline{T}_1 to the impact times $T_I(x_k, u_k, v_k)$ defining the intervals

 $[t_k, t_{k+1})$, where $t_{k+1} = t_k + T_I(x_k, u_k, v_k)$ for $k \in \mathbb{Z}_+$. However, to establish a relation between the discrete-time solution of (7) and the continuous-time solution (4), we also need to address the interval $[0, t_0)$. By Lemma 9, there exists $\delta > 0$ such that, if $x(0) \in S^+$ satisfies $dist(x(0), \mathcal{O}) < \delta$ and $u \in \mathcal{U}$ satisfies $||u||_{\infty} < \delta$, the solution crosses S in finite time $t_0 := \hat{T}_I(x(0), u)$, and there exists a bound $\overline{T}_2 > T^*$ so that $t_0 < \overline{T}_2$. We need to make sure that δ can be selected in a way that x_0 satisfies $||x_0 - x^*|| < \delta_1$ so that Lemma 8 continues to hold. As before, let $x(t) = \psi(t, x(0), u, \overline{v})$; see Remark 1 for the form of x(t) over the intervals $[t_k, t_{k+1})$. From Lemma 9(*iii*) we have

$$\sup_{0 \le t < t_0} \operatorname{dist}(x(t), \mathcal{O}) \le c_1 \operatorname{dist}(x(0), \mathcal{O}) + c_1 \|u\|_{\infty} , \quad (47)$$

for some $c_1 > 0$, which since $x_0 := \lim_{t \nearrow t_0} x(t)$ and the function $\operatorname{dist}(x, \mathcal{O})$ is continuous in x, implies that

$$\operatorname{dist}(x_0, \mathcal{O}) \le c_1 \operatorname{dist}(x(0), \mathcal{O}) + c_1 \|u\|_{\infty} \quad .$$
(48)

Since $x_0 \in S$, Proposition 1 implies $\lambda ||x_0 - x^*|| \leq \operatorname{dist}(x_0, \mathcal{O})$ for $\lambda \in (0, 1)$, and by (48) we have $\lambda ||x_0 - x^*|| \leq c_1 \operatorname{dist}(x(0), \mathcal{O}) + c_1 ||u||_{\infty} < 2c_1 \delta$. Hence, by the inequality above, choosing $\delta < \min\{\delta_1, \lambda \delta_1/(2c_1)\}$ in Lemma 9 ensures that Lemma 8 continues to hold. Furthermore, in what follows, we define $\overline{T} := \max\{\overline{T}_1, \overline{T}_2\}$.

The analysis above shows that, under the assumption of x^* being a LISS fixed point of (7), there exist a $\delta > 0$ such that, for $x(0) \in S^+$ with $dist(x(0), \mathcal{O}) < \delta$ and for $u \in \mathcal{U}$ with $||u||_{\infty} < \delta$, Lemma 9 implies $t_0 < \overline{T}$ where $t_0 := \hat{T}_I(x(0), u)$, and that the solution satisfies the bound (47). If, in addition, $\overline{v} \in \mathcal{V}$ with $||\overline{v}||_{\infty} < \delta$, Lemma 8 implies $t_{k+1} - t_k < \overline{T}$ with $t_{k+1} = t_k + T_I(x_k, u_k, v_k)$ for all $k \in \mathbb{Z}_+$. In this case, the solution satisfies the bound

$$\sup_{t_k \le t < t_{k+1}} \operatorname{dist}(x(t), \mathcal{O}) \le c_2 \|x_k - x^*\| + c_2 \|u\|_{\infty} + c_2 \|\bar{v}\|_{\infty},$$
(49)

for some $c_2 > 0$ and for all $k \in \mathbb{Z}_+$. Substituting (10) in (49), and (for compactness of notation) defining $\alpha(u, \bar{v}) :=$ $\hat{\alpha}_1(||u||_{\infty}) + \hat{\alpha}_2(||\bar{v}||_{\infty})$ with $\hat{\alpha}_1(||u||_{\infty}) := c_2\alpha_1(||u||_{\infty}) +$ $c_2||u||_{\infty}$ and $\hat{\alpha}_2(||\bar{v}||_{\infty}) := c_2\alpha_2(||\bar{v}||_{\infty}) + c_2||\bar{v}||_{\infty}$, results in

$$dist(x(t), \mathcal{O}) \le c_2 \beta(\|x_0 - x^*\|, k) + \alpha(u, \bar{v})$$

$$\le c_2 \beta(dist(x_0, \mathcal{O})/\lambda, k) + \alpha(u, \bar{v})$$
(50)

for all $t_k \leq t < t_{k+1}$ and $k \in \mathbb{Z}_+$. Note that (50) was obtained by using Proposition 1 for $\lambda \in (0, 1)$.

With this information, we now construct suitable class- \mathcal{KL} and class- \mathcal{KL} functions to prove that the orbit \mathcal{O} is LISS in the system (4). We distinguish the following cases.

Case (a): $t \in [t_k, t_{k+1})$ for any $k \ge 1$.

By Remark 1 we have that $t_{k+1} \leq (k+2)\overline{T}$. Pick $c_3 > 0$ such that for all $k \geq 1$, $c_3(k+2)\overline{T} \leq k$. This can be ensured by selecting $c_3 \leq 1/(3\overline{T})$. Using this in (50) implies

$$dist(x(t), \mathcal{O}) \leq c_2 \beta (dist(x_0, \mathcal{O})/\lambda, c_3(k+2)T) + \alpha(u, \bar{v}) \\ \leq c_2 \beta (dist(x_0, \mathcal{O})/\lambda, c_3t) + \alpha(u, \bar{v}) , \quad (51)$$

which is obtained by using the fact $t < t_{k+1} \leq (k+2)\overline{T}$. The estimate (51) holds for all $k \in \mathbb{Z}_+$ with $k \geq 1$, and thus it holds for all $t \geq t_1$.

Case (b): $t \in [t_k, t_{k+1})$ for k = 0.

Choose $\gamma := \ln(2)/(2\overline{T})$ so that $c_2\beta(\operatorname{dist}(x_0, \mathcal{O})/\lambda, 0) \leq 2c_2\beta(\operatorname{dist}(x_0, \mathcal{O})/\lambda, 0)e^{-\gamma t}$ over the interval $[t_0, t_0 + \overline{T}) \subset [0, 2\overline{T}]$. Then, since $t_1 < t_0 + \overline{T}$, the estimate (50) implies that, for all $t \in [t_0, t_1)$,

$$dist(x(t), \mathcal{O}) \le c_2 \beta(dist(x_0, \mathcal{O})/\lambda, 0) + \alpha(u, \bar{v}) \le 2c_2 \beta(dist(x_0, \mathcal{O})/\lambda, 0) e^{-\gamma t} + \alpha(u, \bar{v}) .$$
(52)

Now we can combine the estimates (51) and (52) to obtain a bound for all $t \ge t_0$ of the distance from \mathcal{O} of a solution starting from x_0 at time t_0 . Note that the function

$$\hat{\beta}(\operatorname{dist}(x_0, \mathcal{O}), t) := c_2 \max\{2\beta(\operatorname{dist}(x_0, \mathcal{O})/\lambda, 0)e^{-\gamma t}, \\ \beta(\operatorname{dist}(x_0, \mathcal{O})/\lambda, c_3 t)\},\$$

is continuous, monotonically increasing in dist (x_0, \mathcal{O}) , and monotonically decreasing in t because the individual functions in the max have the same properties; hence, $\hat{\beta} \in \mathcal{KL}$. Upper bounding (51) and (52) with $\hat{\beta}$ and remembering that $\alpha(u, \bar{v}) := \hat{\alpha}_1(\|u\|_{\infty}) + \hat{\alpha}_2(\|\bar{v}\|_{\infty})$ results in

$$\operatorname{dist}(x(t),\mathcal{O}) \leq \hat{\beta}(\operatorname{dist}(x_0,\mathcal{O}),t) + \hat{\alpha}_1(\|u\|_{\infty}) + \hat{\alpha}_2(\|\bar{v}\|_{\infty}) ,$$
(53)

which holds for all $t \ge t_0$.

To complete the proof of this case, we need an estimate in which the class- \mathcal{KL} function in the RHS of (53) depends on dist $(x(0), \mathcal{O})$ and not dist (x_0, \mathcal{O}) ; recall that x(0) is the initial state of the solution of (4), i.e., $\psi(t, x(0), u, \bar{v})$, while x_0 is the first intersection of ψ with S. To remedy this, use (48) noting that $\hat{\beta}(\text{dist}(x_0, \mathcal{O}), t)$ in (53) is a class- \mathcal{K} function for any fixed t, followed by [42, Lemma 14] to get

$$\begin{split} \hat{\beta}(\operatorname{dist}(x_0,\mathcal{O}),t) &\leq \hat{\beta}(c_1\operatorname{dist}(x(0),\mathcal{O}) + c_1 \|u\|_{\infty},t) \\ &\leq \hat{\beta}(2c_1\operatorname{dist}(x(0),\mathcal{O}),t) + \hat{\beta}(2c_1 \|u\|_{\infty},t) \\ &\leq \hat{\beta}(2c_1\operatorname{dist}(x(0),\mathcal{O}),t) + \hat{\beta}(2c_1 \|u\|_{\infty},0). \end{split}$$

Use this inequality in (53) and with an abuse of notation absorb the second term of the above inequality in $\hat{\alpha}_1(||u||_{\infty})$ to obtain

$$\operatorname{dist}(x(t), \mathcal{O}) \leq \hat{\beta}(2c_1 \operatorname{dist}(x(0), \mathcal{O}), t) + \hat{\alpha}_1(\|u\|_{\infty}) + \hat{\alpha}_2(\|\bar{v}\|_{\infty}) .$$
(54)

However, (54) merely holds for $t \ge t_0$ and not for all $t \ge 0$. To address this issue we consider the following case.

Case (c): $t \in [0, t_0)$.

We use the bound (47), which is a consequence of Lemma 9. Employing a trick similar to the one used for constructing the class \mathcal{KL} function in *Case* (b), let⁸ $\gamma := \ln(2)/(2\overline{T})$. Then, $c_1 \operatorname{dist}(x(0), \mathcal{O}) \leq 2c_1 \operatorname{dist}(x(0), \mathcal{O}) e^{-\gamma t}$ over the interval $[0, \overline{T}]$. Then, since $t_0 \leq \overline{T}$, (47) gives for $t \in [0, t_0)$,

$$\operatorname{dist}(x(t), \mathcal{O}) \le 2c_1 \operatorname{dist}(x(0), \mathcal{O}) e^{-\gamma t} + c_1 \|u\|_{\infty} \quad (55)$$

⁸In fact, for this case, $\gamma = \ln(2)/\overline{T}$ would suffice, but we use the same γ as in *Case (b)* to avoid introducing additional constants.

We now combine the bound (54) for $t \ge t_0$ with the bound (55) for $t \in [0, t_0)$ to construct class- \mathcal{KL} and class- \mathcal{K} functions that satisfy Definition 1. Indeed, defining

$$\tilde{\beta}(\operatorname{dist}(x(0), \mathcal{O}), t) := \max\{2c_1 \operatorname{dist}(x(0), \mathcal{O}) e^{-\gamma t}, \\ \hat{\beta}(2c_1 \operatorname{dist}(x(0), \mathcal{O}), t)\},\$$

and $\tilde{\alpha}_1(||u||_{\infty}) := \hat{\alpha}_1(||u||_{\infty}) + c_1||u||_{\infty}$ and $\tilde{\alpha}_2(||\bar{v}||_{\infty}) := \hat{\alpha}_2(||\bar{v}||_{\infty})$ implies that the solution x(t) satisfies Definition 1 for all $t \ge 0$, thereby completing the proof of Theorem 1. \Box

Next, we present a proof of Corollary 1.

Proof of Corollary 1. The proof is identical to that of Theorem 1 with $u \equiv 0$, $\bar{v} \equiv 0$. Ony note that in proving (ii) \Longrightarrow (i) we choose $\omega > 0$ such that $\rho = e^{-\omega T}$ as in Definition 4. \Box

Finally, we present the proof of Theorem 2.

Proof of Theorem 2. The proof of $(i) \Longrightarrow (ii)$ trivially follows by substituting u = 0 and v = 0 in Definition 3 to recover Definition 4 that establishes LAS. For $(ii) \Longrightarrow (i)$, we assume that x^* is a LAS fixed point of the 0-input system. By [43, Theorem 1(1)], there exists a smooth (discrete) Lyapunov function $V : \mathbb{R}^n \to \mathbb{R}_+$ so that for all $x \in B_{\delta}(x^*) \cap S$,

$$V(P(x,0,0)) - V(x) \le -\alpha(||x - x^*||)$$
, (56)

with $\alpha \in \mathcal{K}$. Since V is smooth and P is continuously differentiable, $V \circ P : \mathcal{S} \times \mathcal{U} \times \mathbb{R}^q \to \mathbb{R}_+$ is locally Lipschitz, and $\delta > 0$ can be chosen (shrink if necessary) so that the Lipschitz condition holds uniformly for some $L_{PV} > 0$ for all $x \in B_{\delta}(x^*) \cap \mathcal{S}, u \in \mathcal{U}, ||u||_{\infty} < \delta$, and $v \in B_{\delta}(0)$. Then,

$$V(P(x, u, v)) - V(x) = V(P(x, 0, 0)) - V(x) + V(P(x, u, v)) - V(P(x, 0, 0)) \leq -\alpha(||x - x^*||) + L_{PV} ||u||_{\infty} + L_{PV} ||\bar{v}||_{\infty} ,$$
 (57)

where the inequality follows from (56) and the Lipschitz continuity of $V \circ P$. Hence, V is a LISS Lyapunov function as required by [23, Definition 3.2] and from [43, Lemma 3.5] it follows that the system is LISS.

VI. EXAMPLE: LISS OF A BIPEDAL WALKER

This section presents an example of applying Theorems 1 and 2 to establish ISS for a periodic walking gait for the biped of Fig. 3. The model is underactuated, having five degrees of freedom (DOF) and four actuators; two actuators are placed at the knees and two at the hip. Ground contact is modeled as a passive pivot. More details about the model along with the mechanical properties used here can be found in [4, Table 6.3].

As shown in Fig. 3, we choose $q := (q_1, q_2, q_3, q_4, q_5)$ as coordinates for the configuration space Q. Let Γ be the actuator inputs and $F_e \in \mathcal{F} := \{F_e : \mathbb{R}_+ \to \mathbb{R}^2 \mid ||F_e||_{\infty} < \infty, F_e \text{ is continuous}\}$ be an external force acting at the torso as shown in Fig. 3. Further, let $x \in \mathcal{T}Q := \{(q, \dot{q}) \mid q \in Q, \dot{q} \in \mathbb{R}^5\}$ be the state. The swing phase dynamics is

$$\dot{x} = f(x) + g(x)\Gamma + g_{\rm e}(x)F_{\rm e} \quad . \tag{58}$$

This phase terminates when the swing foot makes contact with the ground. This occurs when $x \in S := \{(q, \dot{q}) \in$



Fig. 3. Robot model with a choice of generalized coordinates.

 $\mathcal{TQ} \mid p_v(q) = 0$ }, where $p_v(q)$ is the height of the swing foot from the ground. The ensuing impact map Δ takes the states x^- just before to the states x^+ just after impact, under the influence of an impulsive disturbance $F_{\rm I} \in \mathbb{R}^2$ applied at the same point as $F_{\rm e}$. The resulting system takes the form

$$\Sigma: \begin{cases} \dot{x} = f(x) + g(x)\Gamma + g_{e}F_{e}(t) & \text{if } x \notin \mathcal{S} \\ x^{+} = \Delta(x^{-}, F_{I}) & \text{if } x^{-} \in \mathcal{S} \end{cases},$$
(59)

in which Δ is smooth. Note that $F_{\rm e}$ and $F_{\rm I}$ are viewed as continuous and discrete disturbances, respectively. There is a variety of methods available for designing control laws $\Gamma(x)$ that result in asymptotically stable limit-cycle gaits in the absence of the disturbances; here, we use the method in [17].

Let $\overline{F}_{I} := \{F_{I,k}\}_{k=0}^{\infty}$ be the sequence of impulsive disturbances and $\{F_{e,k}\}_{k=0}^{\infty}$ be the sequence of continuous inputs as in Section II-C. Then, (59) in closed loop with $\Gamma(x)$, gives rise to the forced Poincaré map $P : S \times \mathcal{F} \times \mathbb{R}^{2} \to S$

$$x_{k+1} = P(x_k, F_{e,k}, F_{I,k}) , \qquad (60)$$

which captures the dynamics of (59) as it goes through S. A simple calculation shows that all the eigenvalues of the linearization of the Poincaré map about a 0-input fixed point x^* are within the unit disk so that x^* is LAS. Then, Theorem 2 implies that x^* is a LISS fixed point of the forced Poincaré map (60) and Theorem 1 ensures that the corresponding periodic orbit \mathcal{O} is LISS in the presence of disturbances.

Figure 4 shows the behavior of the model when F_e is a horizontal sinusoidal force and \bar{F}_I consists of horizontal impulses $\bar{F}_{I,k}$ uniformly sampled from the interval $[-\|\bar{F}_I\|_{\infty}, \|\bar{F}_I\|_{\infty}]$ for a predetermined $\|\bar{F}_I\|_{\infty} < \infty$. It can be seen that the solution $\{x_k\}_{k=0}^{\infty}$ of (60) in Fig. 4(a) as well as the solution $x(t) = \psi(t, x(0), F_e, \bar{F}_I)$ of (59) in Fig. 4(b), asymptotically converge to 0 in the absence of disturbances (blue); are ultimately bounded when $\|F_e\|_{\infty} = 5$ N, $\|\bar{F}_I\|_{\infty} = 0.1$ N.s (gray) and $\|F_e\|_{\infty} = 10$ N, $\|\bar{F}_I\|_{\infty} = 0.2$ N.s (red), with the ultimate bound for the former being smaller than that of the latter, indicating LISS behavior.

VII. CONCLUSION

This paper presents a method for analyzing robustness of limit cycles exhibited by systems with impulse effects. It is



Fig. 4. Response of the biped from an initial condition away from the orbit with $F_e = 0$ N, $\|\bar{F}_I\|_{\infty} = 0$ N.s in blue; $F_e = [5\sin(4t) \ 0]^T$ N, $\|\bar{F}_I\|_{\infty} = 0.1$ N.s in gray; and $F_e = [10\sin(4t) \ 0]^T$ N, $\|\bar{F}_I\|_{\infty} = 0.2$ N.s in red. (a) Evolution of $\|x_k - x^*\|$ over step number k where $\{x_k\}_{k=0}^{\infty}$ is the solution of (60). (b) Supremum deviation of $x(t) = \psi(t, x(0), F_e, \bar{F}_I)$, the solution of (59), from the orbit \mathcal{O} over each step.

shown that ISS of the limit cycle is equivalent to that of the forced Poincaré map. This result allows us to analyze the robustness of *hybrid* limit cycles by merely analyzing a discrete dynamical system. The proof of this result, provides ISS estimates that could be used to quantify the robustness. Furthermore, exploiting the availability of these estimates, we establish an equivalence between ES of the limit cycle and the 0-input Poincaré map. The overarching goal of this work is to develop a framework within which the robustness of periodic orbits can be rigorously analyzed.

APPENDIX A

This appendix provides proofs to Lemmas 1 and 2 and to Proposition 2, clarifying properties of the forced solution $\varphi(t, x(0), u)$ of (1) and $\psi(t, x(0), u, \bar{v})$ of (4).

Proof of Lemma 1. The [37, proof relies on Theorem 5.2, p. 377]. To apply this result, define $F: \mathbb{R}_+ \times \mathbb{R}^n \times \mathcal{U} \to \mathbb{R}^n$ as F(t, x, u) := f(x, G(t, u)), where $G : \mathbb{R}_+ \times \mathcal{U} \rightarrow \mathbb{R}^p$ is G(t, u) = u(t). Then, [37, Theorem 5.2, p. 377] states that φ is continuously differentiable in x and u if F is so. Note that, in view of the procedure in [37, p. 369] for treating time-dependent vector fields, the statement of [37, Theorem 5.2, p. 377] would require F to be continuously differentiable in t. However, the proof of [37, Theorem 5.2, p. 377] only uses continuity of F in t. Hence, below we show that F is continuously differentiable in x and u, but only continuous in t.

We begin by showing that G is continuous in t and continuously differentiable in u. Continuity of G in t is clear. Continuity of G in $u \in U$ follows from the fact that $\hat{G} : U \to \mathbb{R}^p$ defined by $\hat{G}(u) := G(t, u) = u(t)$ is bounded and linear for each *fixed* time $t \in \mathbb{R}_+$; by [44, p. 257, Theorem 1] this implies that \hat{G} is continuous in u, thus G is also continuous in u. Indeed, linearity is immediate by the definition of \hat{G} , while boundedness follows from $\|\hat{G}(u)\| = \|u(t)\| \le \|u\|_{\infty}$, implying that the operator norm is upper bounded by 1 for any $t \in \mathbb{R}_+$. As a result, G(t, u) is continuous in both arguments. Furthermore, G is linear with respect to u, and using [37, p. 339, Theorem 3.1], we have that the Fréchet (partial) derivative of G with respect to u is continuous, for it is G itself. Thus, G(t, u) is continuous in t and continuously differentiable in u. Using this fact with the assumption that f is continuously differentiable, it follows that F is continuous in t and continuously differentiable in (x, u). The result then follows from [37, Theorem 5.2, p. 377].

Proof of Lemma 2. Before proceeding with the proof, note that even though the domain of $T_{\rm I}$ is restricted to $S \times \mathcal{U} \times \mathbb{R}^q$, $T_{\rm I}$ is well-defined on $\mathbb{R}^n \times \mathcal{U} \times \mathbb{R}^q$ since Δ and φ are well-defined maps for any $x \in \mathbb{R}^n$. Hence, we will consider this extended domain of $T_{\rm I}$ in the proof, which follows from the implicit mapping theorem [37, Chapter XIV, Theorem 2.1].

Let $\tilde{H}(t, x, u, v) := H \circ \varphi(t, \Delta(x, v), u)$ where $x \in \mathbb{R}^n$, $u \in \mathcal{U}$, and $v \in \mathbb{R}^q$. From Lemma 1 in Appendix A, we have that the solution φ is continuously differentiable in all its arguments in the Fréchet sense. Using this with assumption A.2 we have that \tilde{H} is continuously differentiable. From assumption A.5 it follows that $\tilde{H}(T^*, x^*, 0, 0) = 0$. Further, from assumption A.7 we have $\partial \tilde{H}/\partial t|_{(T^*, x^*, 0, 0)} \neq 0$. Next, noting that $(\mathbb{R}^n \times \mathbb{R}^q, \|\cdot\|)$ and $(\mathcal{U}, \|\cdot\|_{\infty})$ are Banach spaces, we can use [37, Chapter XIV, Theorem 2.1] to establish the existence of a unique map $T_1(x, u, v)$ which satisfies $\tilde{H}(T_1(x, u, v), x, u, v) = 0$ for a sufficiently small $\delta > 0$ such that $x \in B_{\delta}(x^*)$, $u \in \mathcal{U}$ with $\|u\|_{\infty} < \delta$, and $v \in B_{\delta}(0)$. Additionally, since $\tilde{H}(t, x, u, v)$ is continuously differentiable with respect to its arguments, so is $T_1(x, u, v)$. As this holds for any $x \in B_{\delta}(x^*)$, it also holds for any $x \in B_{\delta}(x^*) \cap S$. \Box

Proof of Proposition 2. We restrict attention to initial conditions x(0) that result in solutions that hit S; otherwise the solution is continuous and it can be extended indefinitely, trivially excluding the occurrence of Zeno and beating phenomena.

We first show parts (i) and (ii) simultaneously. Let $x(t) = \psi(t, x(0), u, \bar{v})$ be a solution of (4) that is defined over some interval $[0, t_f)$ and satisfies Definition 1. As in Remark 1, let t_k and t_{k+1} denote two subsequent impact times so that $t_{k+1} - t_k = T_1(x_k, u_k, v_k)$, where $x_k := \lim_{t \neq t_k} x(t) \in S$, $u_k(t) = u(t)$ for $t \in [t_k, t_{k+1})$, and v_k is the k-th element of the sequence \bar{v} . We will show that there exists $\underline{T} > 0$ such that $t_{k+1} - t_k > \underline{T}$ for all $k \in \mathbb{Z}_+$ with $[t_k, t_{k+1}) \subset [0, t_f)$. By Definition 1, there exists $\delta > 0$ so that $dist(x(0), \mathcal{O}) < \delta$, $u \in \mathcal{U}$ with $||u||_{\infty} < \delta$, and $\bar{v} \in \mathcal{V}$ with $||\bar{v}||_{\infty} < \delta$ imply

$$\operatorname{dist}(x(t), \mathcal{O}) \le \beta(\delta, 0) + \alpha_1(\delta) + \alpha_2(\delta) \quad , \qquad (61)$$

for all $t \in [0, t_f)$. Furthermore, by continuity of the distance function, we have $\lim_{t \nearrow t_k} \operatorname{dist}(x(t), \mathcal{O}) = \operatorname{dist}(x_k, \mathcal{O})$, and (61) in view of Proposition 1 implies that, for some $\lambda \in (0, 1)$,

$$\|x_k - x^*\| \le \frac{1}{\lambda} \left(\beta(\delta, 0) + \alpha_1(\delta) + \alpha_2(\delta)\right) \quad , \qquad (62)$$

for all $k \in \mathbb{Z}_+$ with $[t_k, t_{k+1}) \subset [0, t_f)$. The result now follows from continuity of the time-to-impact function by Lemma 2. Indeed, as in the proof of part (ii) of Lemma 5, continuity of $T_{\rm I}$ implies that, for some $\underline{T} > 0$, there exists a $\delta_T >$ 0 such that $x \in B_{\delta_T}(x^*) \cap S$, $u \in \mathcal{U}$ with $||u||_{\infty} < \delta_T$, and $v \in B_{\delta_T}(0)$ imply $\underline{T} < T_{\mathrm{I}}(x, u, v)$. Choosing δ in (62) so that $\frac{1}{\lambda} \left(\beta(\delta, 0) + \alpha_1(\delta) + \alpha_2(\delta) \right) < \delta_T$ ensures that $x_k \in$ $B_{\delta_T}(x^*) \cap \mathcal{S}$ for all $k \in \mathbb{Z}_+$ with $[t_k, t_{k+1}) \subset [0, t_f)$. Such choice of δ is always possible since the upper bound in (62) is a class- \mathcal{K} function of δ . Shrinking δ further (if necessary) to ensure that $\delta < \delta_T$ guarantees that for $x_k \in B_{\delta}(x^*) \cap S$, $u \in \mathcal{U}$ with $||u||_{\infty} < \delta$, and $v \in B_{\delta}(0)$ we have that $T_{I}(x_{k}, u_{k}, v_{k}) >$ <u>*T*</u> for all discrete events k of the solution. As a result, t_{k+1} – $t_k > \underline{T}$ for all $k \in \mathbb{Z}_+$ with $[t_k, t_{k+1}) \subset [0, t_f)$. This ensures that any two discrete events are punctuated by a time gap of \underline{T} , thereby precluding solutions that are purely discrete or eventually discrete, or exhibit Zeno behavior, completing the proof of (i) and (ii).

To prove part (*iii*), from (61) it is clear that x(t) is trapped in a compact set. Neglecting – as was mentioned at the beginning of the proof – the trivial case where the continuous solution never approaches S, using arguments similar to the proof of [34, Theorem 3.3] the solution can be extended until it reaches S. At this point, a well-defined discrete jump occurs that ensures the post-discrete-event state is still trapped within the same compact set and lies outside S because of part (*i*); hence, the solution must flow again according to the continuous dynamics until it reaches S. Additionally (*ii*) ensures the absence of Zeno behavior. Hence, we can propagate this argument forward for all time to obtain (*iii*).

APPENDIX B

Proof of Lemma 4. By assumption A.1, f(x,0) is twice continuously differentiable; thus, its backward flow $\varphi^{-}(s\tau, x^*, 0) = y(\tau)$ is three-times continuously differentiable. Surjectivity of $y(\tau)$ is obvious from the definition of $y(\tau)$ in (14). Injectivity follows from a contradiction argument. Assume $y(\tau)$ is not injective, then there exist $\tau_1 < \tau_2$ in [0,T] with $y(\tau_1) = y(\tau_2)$. Let $f^{-}(x,0) := -f(x,0)$ be the vector field for the backwards flow. If $f^{-}(y(\tau_1),0) \neq f^{-}(y(\tau_2),0)$ then f would not be well defined. If $f^{-}(y(\tau_1),0) = f^{-}(y(\tau_2),0)$, we have: *Case* (i): $0 < \tau_1 < \tau_2$

Since $y(\tau_1) = y(\tau_2)$, we return to the same state after an interval $\tau_2 - \tau_1 > 0$. Hence, $E := \{y(\tau) \mid \tau_1 \le \tau \le \tau_2\} \subset \overline{\mathcal{O}}$ is a periodic orbit of the backwards-flow continuous system and we have the following sub-cases:

- (a) $y(\tau) \in E$ for all $\tau \in [\tau_1, T]$: The orbit E would also exist in the forward flow. As the periodic orbit is an invariant set under the 0-input continuous dynamics (1), the forward flow starting from $y(T) = \Delta(x^*, 0)$ will be trapped in E and never reach S, contradicting assumption A.5, according to which the solution must reach S in finite time T^* .
- (b) There exists $\hat{\tau} \in [\tau_1, T]$ such that $y(\hat{\tau}) \notin E$: This contradicts uniqueness of the backwards solution, as starting from $y(\tau_1) \in E$, one solution flows to $y(\hat{\tau}) \notin E$ while the other gets trapped in E.

Case (ii): $0 = \tau_1 < \tau_2$

Note that $H(y(\tau))$ and $L_{f^-}H(y(\tau,0))$ are continuous in τ . Additionally, from assumption A.6-A.7, $H(y(0)) = H(x^*) = 0$ and $L_{f^-}H(y(0), 0) > 0$ (flipped sign from assumption A.7 due to the flow being backwards in time), thus there exists a $\delta > 0$ such that $L_{f^-}H(y(\tau), 0) > 0$ for all $\tau \in [0, \delta)$. Hence for the interval $(0, \delta)$ we have $H(y(\tau)) > 0$, i.e., $\{y(\tau) \mid \tau \in (0, \delta)\} \subset S^+$. Again using continuity of $H(y(\tau))$ and $L_{f^-}H(y(\tau), 0)$ at τ_2 we have that $H(y(\tau))$ is strictly increasing in the interval $(\tau_2 - \delta, \tau_2 + \delta)$ (shrink $\delta > 0$ if necessary to ensure $\delta < \tau_2 - \delta$) but $H(y(\tau_2)) = 0$, hence, $H(y(\tau)) < 0$ for all $\tau \in (\tau_2 - \delta, \tau_2)$, i.e., $\{y(\tau) \mid \tau \in (\tau_2 - \delta, \tau_2)\} \subset S^-$. Thus, for some $\bar{\tau}$ such that $\delta < \bar{\tau} < \tau_2 - \delta$, the solution must cross over from S^+ to S^- at a point other than x^* , resulting in a contradiction to assumption A.6.

APPENDIX C

First note that S is a twice continuously differentiable embedded submanifold in \mathbb{R}^n defined by H(x) = 0. Clearly, $H(x^*) = 0$, and, without loss of generality, assume that for the *n*-th coordinate $\frac{\partial H}{\partial x_n}\Big|_{x^*} \neq 0$, which follows from assumption A.2. Hence, using the implicit function theorem we can write $x_n = h(\bar{x})$ where $\bar{x} := (x_1, ..., x_{n-1})$ for $\bar{x} \in B_{\delta}(\bar{x}^*)$ where $x^* = (\bar{x}^*, h(\bar{x}^*))$. As a result, the local coordinates of states $x \in S$ in a neighborhood of x^* are $x = (\bar{x}, h(\bar{x}))$. The Taylor expansion of $h(\bar{x})$ at \bar{x}^* gives

$$h(\bar{x}) = h(\bar{x}^*) + A(\bar{x} - \bar{x}^*) + O(\|\bar{x} - \bar{x}^*\|^2) , \qquad (63)$$

where $A = \frac{\partial h}{\partial \bar{x}}|_{\bar{x}^*}$. Let $z \in T_{x^*}S$ be the projection of x on $T_{x^*}S$ along $(x_1, x_2, ..., x_{n-1})$, then its coordinates are $z = (\bar{x}, h(\bar{x}^*) + A(\bar{x} - \bar{x}^*))$. Hence, for $x \in B_{\delta}(x^*) \cap S$ and $z \in T_{x^*}S$, there exists a c > 0 such that,

$$\begin{aligned} \|x - z\| &= \|(\bar{x}, h(\bar{x})) - (\bar{x}, h(\bar{x}^*) + A(\bar{x} - \bar{x}^*))\| \\ &= |O(\|\bar{x} - \bar{x}^*\|^2)| \le c \|\bar{x} - \bar{x}^*\|^2 \le c \|x - x^*\|^2 . \end{aligned}$$

APPENDIX D

Proof of Lemma 8. We begin with part (i). By Remark 5, $\varphi(t, \Delta(x^*, 0), 0)$ exists and is unique over $[0, \overline{T}]$ with $\overline{T} > T^*$. Lemma 7 establishes the existence of $\delta_1 > 0$ for which, when $x \in S$ and $v \in \mathbb{R}^q$ are such that $||\Delta(x, v) - \Delta(x^*, 0)|| < \delta_1$, and when $||u||_{\infty} < \delta_1$, the perturbed solution $\varphi(t, \Delta(x, v), u)$ exists and is unique over the same interval $[0, \overline{T}]$. By the continuity of Δ following from assumption A.3, there exists a $\delta_2 > 0$ for which $||x - x^*|| < \delta_2$ and $||v|| < \delta_2$ guarantee $||\Delta(x, v) - \Delta(x^*, 0)|| < \delta_1$. Hence, choosing $\delta = \min{\{\delta_1, \delta_2\}}$ we have that, for $x \in B_{\delta}(x^*) \cap S$, $u \in \mathcal{U}$ with $||u||_{\infty} < \delta$ and $\overline{v} \in \mathcal{V}$ with $||\overline{v}||_{\infty} < \delta$, the perturbed solution $\varphi(t, \Delta(x, v), u)$ of (1) exists and is unique over $[0, \overline{T}]$, thus proving part (i).

To prove part (ii), let $\epsilon_T := \overline{T} - T^* > 0$. By continuity of T_1 there exists a $\delta_T > 0$ such that for $x \in B_{\delta_T}(x^*) \cap S$, $u \in \mathcal{U}$ with $||u||_{\infty} < \delta_T$, and $v \in B_{\delta_T}(0)$, we have that $|T_1(x, u, v) - T_1(x^*, 0, 0)| < \epsilon_T$, which implies that $\underline{T} < T_1(x, u, v) < \overline{T}$ where $\underline{T} := T^* - \epsilon_T = 2T^* - \overline{T} > 0$ (shrink \overline{T} if necessary to ensure that $\overline{T} < 2T^*$). To show transversality, define $\ell(x, u, v) := \frac{\partial H}{\partial x}|_{\varphi(T_1(x, u, v), \Delta(x, v), u)} f(\varphi(T_1(x, u, v), \Delta(x, v), u), u)$ so that $\ell^* := \ell(x^*, 0, 0) < 0$ by assumption A.7, and let $\epsilon_\ell \in (0, -\ell^*)$. By continuity of ℓ , there is a $\delta_\ell > 0$ such that for $x \in B_{\delta_\ell}(x^*) \cap S$, $u \in \mathcal{U}$ with $||u||_{\infty} < \delta_\ell$, and $v \in B_{\delta_\ell}(0)$, we have that $|\ell(x, u, v) - \ell(x^*, 0, 0)| < \epsilon_\ell$, implying $\ell(x, u, v) < \epsilon_\ell + \ell^* < 0$. Choosing $\delta = \min\{\delta_1, \delta_2, \delta_T, \delta_\ell\}$ with $\delta_1, \delta_2 > 0$ as in the proof of part (i) completes the proof of part (ii).

Finally, for part (*iii*), we begin by setting up the region within which we work. Let $w(t) = \varphi(t, \Delta(x^*, 0), 0)$ be the unperturbed (zero-input) solution of (1), which by Remark 5 is well-defined over the interval $[0,\overline{T}]$ with $\overline{T} > T^*$. Let $\delta > 0$ be as in the proof of part (*ii*) so that, for $x \in B_{\delta}(x^*) \cap$ S, $u \in \mathcal{U}$ with $||u||_{\infty} < \delta$ and $v \in B_{\delta}(0)$, the perturbed solution $\varphi(t, \Delta(x, v), u)$ exists and is unique over $[0,\overline{T}]$ and $T_{\mathrm{I}}(x, u, v) < \overline{T}$. Furthermore, for this choice of δ , Lemma 7 guarantees the existence of L > 0 such that

$$\begin{aligned} \|\varphi(t,\Delta(x,v),u) - w(t)\| &\leq \mathrm{e}^{L\overline{T}} \|\Delta(x,v) - \Delta(x^*,0)\| \\ &+ (\mathrm{e}^{L\overline{T}} - 1) \|u\|_{\infty}, \end{aligned}$$

for all $t \in [0, \overline{T}]$. Since Δ is continuously differentiable from assumption A.3, it is locally Lipschitz. Hence, shrinking δ (if necessary) ensures that the Lipschitz condition can be satisfied uniformly over $B_{\delta}(x^*) \cap S$ and $\|\overline{v}\|_{\infty} < \delta$ for some constant $L_{\Delta} > 0$, thereby resulting in the following estimate

$$\sup_{0 \le t \le \overline{T}} \|\varphi(t, \Delta(x, v), u) - w(t)\|$$

$$\leq e^{L\overline{T}} L_{\Delta}(\|x - x^*\| + \|\overline{v}\|_{\infty}) + (e^{L\overline{T}} - 1)\|u\|_{\infty}.$$
(64)

In addition, since by Lemma 2 the map $T_{\rm I}$ is continuously differentiable and thus locally Lipschitz. Shrinking δ further (if necessary) ensures that there exists a $L_T > 0$ such that

$$|T_{\mathrm{I}}(x, u, v) - T_{\mathrm{I}}(x^{*}, 0, 0)| \le L_{T}(||x - x^{*}|| + ||u||_{\infty} + ||\bar{v}||_{\infty})$$
(65)

for all $x \in B_{\delta}(x^*) \cap S$, $u \in \mathcal{U}$ with $||u||_{\infty} < \delta$ and $\bar{v} \in \mathcal{V}$ with $||\bar{v}||_{\infty} < \delta$. In what follows, we work in such region so that the perturbed solution $\varphi(t, \Delta(x, v), u)$ and the corresponding time to impact $T_{\mathrm{I}}(x, u, v)$ satisfy (64) and (65), respectively.

Now, let us consider the distance of the perturbed solution $\varphi(t, \Delta(x, v), u)$ from \mathcal{O} . For any $t \in [0, \overline{T}]$ we have

$$dist(\varphi(t, \Delta(x, v), u), \mathcal{O})$$

$$:= \inf_{y \in \mathcal{O}} \|\varphi(t, \Delta(x, v), u) - w(t) + w(t) - y\|$$

$$\leq \|\varphi(t, \Delta(x, v), u) - w(t)\| + dist(w(t), \mathcal{O}) , \quad (66)$$

where (66) is obtained by the triangle inequality. Regarding the term $dist(w(t), \mathcal{O})$ in (66), note that when $t \in [0, T^*)$ we have $dist(w(t), \mathcal{O}) = 0$, while for $t \in [T^*, \overline{T}]$ we have

$$dist(w(t), \mathcal{O}) := \inf_{y \in \mathcal{O}} ||w(t) - x^* + x^* - y|| \leq ||w(t) - x^*|| + \inf_{y \in \mathcal{O}} ||x^* - y|| = ||w(t) - x^*|| ,$$
(67)

since $\inf_{y \in \mathcal{O}} ||x^* - y|| = 0$. We distinguish the following cases. *Case (a):* Assume that $T_{\mathrm{I}}(x, v, u) < T^*$. Then, for all $t \in [0, T_{\mathrm{I}}(x, v, u)]$, we have $\operatorname{dist}(w(t), \mathcal{O}) = 0$ and application of $\sup_{0 < t < T_{\mathrm{I}}}$ on (66) results in the bound

$$\sup_{0 \le t < T_{\mathrm{I}}} \operatorname{dist}(\varphi(t, \Delta(x, v), u), \mathcal{O})$$

$$\leq \sup_{0 \le t < T_{\mathrm{I}}} \|\varphi(t, \Delta(x, v), u) - w(t)\| \quad . \tag{68}$$

Since $T^* < \overline{T}$, for this case we have $[0, T_{\mathrm{I}}(x, u, v)) \subset [0, \overline{T}]$, and the bound (64) implies

$$\sup_{0 \le t < T_{\mathrm{I}}} \operatorname{dist}(\varphi(t, \Delta(x, v), u), \mathcal{O})$$
$$\le \mathrm{e}^{L\overline{T}} L_{\Delta}(\|x - x^*\| + \|\overline{v}\|_{\infty}) + (\mathrm{e}^{L\overline{T}} - 1) \|u\|_{\infty}.$$

Case (b): Assume that $T_{I}(x, v, u) \ge T^{*}$. Then, (67) implies

$$\sup_{0 \le t < T_1} \operatorname{dist}(w(t), \mathcal{O}) \le \sup_{T^* \le t < T_1} \|w(t) - x^*\| \quad . \tag{69}$$

Applying $\sup_{0 \le t \le T_1}$ on (66) followed by (69) results in

$$\sup_{0 \le t < T_{\mathrm{I}}} \operatorname{dist}(\varphi(t, \Delta(x, v), u), \mathcal{O})$$

$$\leq \sup_{0 \le t < T_{\mathrm{I}}} \|\varphi(t, \Delta(x, v), u) - w(t)\|$$

$$+ \sup_{T^* \le t < T_{\mathrm{I}}} \|w(t) - x^*\| .$$
(70)

Regarding the first term in the RHS of (70), the upper bound (64) can be used since $[0, T_{\rm I}) \subset [0, \overline{T}]$. Next, we look at the second term. Let $K := \max_{T^* \leq t \leq \overline{T}} ||f(w(t), 0)||$. As mentioned in Remark 5, we have $L_f \overline{H}(w(t), 0) < 0$ for all $T^* \leq t \leq \overline{T}$ which implies that $f(w(t), 0) \neq 0$ for all $t \in [T^*, \overline{T}]$, hence K > 0. Use K in the following,

$$\sup_{T^* \le t < T_{\mathrm{I}}} \|w(t) - x^*\| = \sup_{T^* \le t < T_{\mathrm{I}}} \left\| \int_{T^*}^{t} f(w(s), 0) \, ds \right\|$$

$$\le K |T_{\mathrm{I}}(x, u, v) - T_{\mathrm{I}}(x^*, 0, 0)|$$

$$\le K L_{T}(\|x - x^*\| + \|u\|_{\infty} + \|\bar{v}\|_{\infty}) , \qquad (71)$$

where (65) was used. The desired estimate in Case (b) is then found by combining (64) and (71) according to (70).

Finally, Case (a) and (b) can be combined for an appropriate c > 0 to obtain (44), thereby completing the proof.

Proof of Lemma 9. We first prove Lemma 9(*i*), (*ii*) simultaneously. We begin by using Lemma 7 to show that, for any point of \mathcal{O} , there exists an open ball of initial conditions and inputs for which a unique (forced) solution exists over a time interval that is sufficiently long to cross S transversally in finite time. Then, compactness of $\overline{\mathcal{O}}$ is used to extract a finite open subcover of $\overline{\mathcal{O}}$, based on which an upper bound on the time to cross S is established uniformly in a tube around \mathcal{O} .

Let $w(t) = \varphi(t, \Delta(x^*, 0), 0)$ be the unperturbed (zeroinput) solution of (1) starting from $\Delta(x^*, 0)$. By Remark 5, w(t) is well-defined over an interval $[0,\overline{T}]$ with $\overline{T} > T^*$. Then define $w_e := w(T)$ which is in the open set S^- . Hence, there exists a $\delta_{\rm e} > 0$ such that $B_{\delta_{\rm e}}(w_{\rm e}) \subset S^-$. Given an arbitrary time $\xi \in [0,T^*]$ and the corresponding point $w(\xi) \in \mathcal{O}$, Lemma 7 ensures that there exists a $\delta_{\xi} > 0$ such that, for any $x \in B_{\delta_{\xi}}(w(\xi))$ and $u \in \mathcal{U}$ with $||u||_{\infty} < \delta_{\xi}$, the (perturbed) solution $\varphi(t, x, u)$ exists and is unique over the interval $t \in [0, T - \xi]$. Furthermore, for some L > 0, (43) of Lemma 7 gives $\|\varphi(t,x,u) - w(\xi+t)\| \leq (2e^{L(\overline{T}-\xi)}-1)\delta_{\xi}$ for all $t \in [0, \overline{T} - \xi]$, which implies that δ_{ξ} can be chosen sufficiently small to ensure that $\sup_{0 < t < \overline{T} - \xi} \|\varphi(t, x, u) - w(t + \xi)\| < \delta_{e}$. As a result, $\varphi(\overline{T} - \xi, x, u) \in B_{\delta_e}(w_e)$, from which it follows that the perturbed solution has crossed S during the interval $[0, \overline{T}_x]$ with $\overline{T}_x := \overline{T} - \xi$; clearly, \overline{T}_x depends on the choice of $w(\xi)$. Finally, the fact that the solution crosses S transversally can be shown similarly to the proof of part (ii) of Lemma 8.

Now, to extend these results over the entire orbit, construct an open cover of $\overline{\mathcal{O}}$ by using the open balls $B_{\delta_{\xi}}(w(\xi))$ for each $\xi \in [0, T^*]$. As $\overline{\mathcal{O}}$ is compact, there exists $\xi_1, \xi_2, ..., \xi_N \in$ $[0,T^*]$, and $\delta_i := \delta_{\xi_i}, w_i := w(\xi_i)$ such that $\bigcup_{i=1}^N B_{\delta_i}(w_i) \supset$ $\overline{\mathcal{O}}$ is a finite sub-cover. Choose $0 < \delta < \min_{1 \le i \le N} \delta_i$ such that $\{x \in \mathbb{R}^n \mid \operatorname{dist}(x, \mathcal{O}) < \delta\} \subset \bigcup_{i=1}^N B_{\delta_i}(w_i)$. Choosing such $\delta > 0$ is always possible. Indeed, since $\bigcup_{i=1}^{N} B_{\delta_i}(w_i)$ is bounded, its boundary $\mathcal{B} := \partial(\bigcup_{i=1}^N B_{\delta_i}(w_i))$ is compact. Let $0 < \delta < \min_{x \in \mathcal{B}} \operatorname{dist}(x, \mathcal{O})/2$ (shrink if necessary to ensure $\delta < \min_{1 \le i \le N} \delta_i$) where the min is well-defined because $\operatorname{dist}(x, \mathcal{O})$ is continuous. Note that $\{x \in \mathbb{R}^n \mid \operatorname{dist}(x, \mathcal{O}) < \delta\}$ is connected because O is connected; hence, this choice of δ ensures that $\{x \in \mathbb{R}^n \mid \operatorname{dist}(x, \mathcal{O}) < \delta\} \subset \bigcup_{i=1}^N B_{\delta_i}(w_i)$. With this choice of δ , any solution $\varphi(t, x, u)$ such that $x \in S^+$ with $\operatorname{dist}(x, \mathcal{O}) < \delta$ and $u \in \mathcal{U}$ with $||u||_{\infty} < \delta$ exists and is unique for $t \in [0, T - \xi_i]$, where ξ_i corresponds to the ball $B_{\delta_i}(w_i)$ that includes x (note that x may belong to multiple such balls; picking any one would suffice). Moreover, the solution will cross S transversally and in time within $\max_{1 \le i \le N} (\overline{T} - \xi_i) \le 1$ \overline{T} . This completes the proof of Lemma 9(*i*), (*ii*).

The proof of Lemma 9(*iii*) closely follows the proof of Lemma 8(*iii*), with the difference that, instead of comparing the solution $\varphi(t, x, u)$ with $\varphi(t, \Delta(x^*, 0), 0)$ as was the case in Lemma 8(*iii*), now we compare $\varphi(t, x, u)$ with $\varphi(t, w(\xi_{\min}), 0) = w(t + \xi_{\min})$, where $\xi_{\min} \in [0, T^*]$ is such that⁹ dist $(x, \mathcal{O}) = ||x - w(\xi_{\min})||$. To do this comparison, we use $T^* - \xi_{\min}$ and $\overline{T} - \xi_{\min}$ instead of T^* and \overline{T} , and we use Claim 1 and Claim 2 below to generalize estimates related to \hat{T}_{I} and Lemma 7, respectively, from an open-ball around a point in \mathbb{R}^n to an entire open neighborhood of \mathcal{O} in S^+ . In the end, we replace $||x - w(\xi_{\min})||$ with dist (x, \mathcal{O})

Claim 1: Let \hat{T}_{I} be as in (8). There exist $L_{\hat{T}} > 0$ and $\delta > 0$ such that for $x \in S^{+}$ with $\operatorname{dist}(x, \mathcal{O}) < \delta$ and $u \in \mathcal{U}$ with $\|u\|_{\infty} < \delta$, the following is satisfied

$$|\hat{T}_{\mathrm{I}}(x,u) - \hat{T}_{\mathrm{I}}(w(\xi_{\min}),0)| \le L_{\hat{T}}(||x - w(\xi_{\min})|| + ||u||_{\infty}).$$

⁹Such ξ_{\min} exists by Lemma 3 and depends on x since $\xi_{\min} \in \arg\min_{\xi \in [0,T]} ||x - w(\xi)||$; we do not display this dependence here.

Claim 2: There exist L > 0 and $\delta > 0$ such that for any $x \in S^+$ with $\operatorname{dist}(x, \mathcal{O}) < \delta$ and $u \in \mathcal{U}$ with $||u||_{\infty} < \delta$, the following is satisfied for $0 \le t \le \overline{T} - \xi_{\min}$,

$$\|\varphi(t,x,u) - w(t + \xi_{\min})\| \leq e^{L\overline{T}} \|x - w(\xi_{\min})\| + (e^{L\overline{T}} - 1) \|u\|_{\infty}.$$

Finally, we present proofs of Claim 1 and 2.

Proof of Claim 1. For any $\xi \in [0, T^*]$, there exists a $\delta_{\xi} > 0$ such that \hat{T}_{I} is continuously differentiable, and thus locally Lipschitz for $x \in B_{\delta_{\xi}}(w(\xi))$ and $u \in \mathcal{U}$ with $||u||_{\infty} < \delta_{\xi}$. Shrink (if necessary) δ_{ξ} to ensure that the Lipschitz condition is satisfied uniformly for some $L_{\hat{T},\xi} > 0$. Construct an opencover for $\overline{\mathcal{O}}$ and extract a finite subcover $\bigcup_{i=1}^{N} B_{\delta_i/2}(w_i)$ as above, but now use $\delta_i/2$. Let $x \in B_{\delta_i/2}(w_i)$ for some i, then $w(\xi_{\min}) \in B_{\delta_i}(w_i)$ for the same *i*. This holds because $||x - w(\xi_{\min})|| \le ||x - w_i|| < \delta_i/2$, hence using the triangle inequality $||w(\xi_{\min}) - w_i|| \le ||x - w(\xi_{\min})|| + ||w_i - x|| < \delta_i$. Therefore, for any $x \in \bigcup_{i=1}^N B_{\delta_i/2}(w_i)$, we can write the Lipschitz condition with the constant $L_{\hat{T},\xi_i}$ replaced by $L_{\hat{T}} =$ $\max_{1 \leq i \leq N} L_{\hat{T},\xi_i}$. Choosing $0 < \delta < \min_{1 \leq i \leq N} \delta_i/2$ such that $\{x \in \mathcal{S}^+ \mid \operatorname{dist}(x, \mathcal{O}) < \delta\} \subset \bigcup_{i=1}^N B_{\delta_i/2}(w_i)$ and $u \in \mathcal{U}$ with $||u||_{\infty} < \delta$ completes the proof.

Proof of Claim 2. The proof is similar to [34, Theorem 3.5]; hence, we only provide a sketch. Note that L is the Lipschitz constant of f on the compact set $\{(x, \hat{u}) \in \mathbb{R}^n \times \mathbb{R}^p \mid ||x - w(t)|| \le \epsilon, \forall t \in [0, T^*], \text{ and } ||\hat{u}|| \le \epsilon\}$ where \hat{u} denotes the value of $u \in \mathcal{U}$ at a certain time instant and should not be confused with the function itself. The proof follows by the Gronwall-Bellman inequality [34, Lemma A.1], which gives

$$\begin{aligned} \|\varphi(t, x, u) - w(t + \xi_{\min})\| & (72) \\ &\leq e^{L(\overline{T} - \xi_{\min})} \|x - w(\xi_{\min})\| + (e^{L(\overline{T} - \xi_{\min})} - 1)\|u\|_{\infty}, \\ &\leq e^{L\overline{T}} \|x - w(\xi_{\min})\| + (e^{L\overline{T}} - 1)\|u\|_{\infty}. \end{aligned}$$

for $t \in [0, \overline{T} - \xi_{\min}]$. Choosing $\delta = \epsilon/(2e^{L\overline{T}})$ such that $\operatorname{dist}(x, \mathcal{O}) = ||x - w(\xi_{\min})|| < \delta$ and $||u||_{\infty} < \delta$ ensures the solution remains trapped in the compact ϵ -neighborhood of $\overline{\mathcal{O}}$ for $t \in [0, T^* - \xi_{\min}]$ and hence the inequality holds. \Box

REFERENCES

- [1] D. Baĭnov and P. S. Simeonov, *Systems with impulse effect: stability, theory, and applications.* Chichester, U.K.: Ellis Horwood, 1989.
- [2] G. Ballinger and X. Liu, "Permanence of population growth models with impulsive effects," *Mathematical and Computer Modelling*, vol. 26, no. 12, pp. 59–72, 1997.
- [3] T. Yang and L. O. Chua, "Impulsive stabilization for control and synchronization of chaotic systems: theory and application to secure communication," *IEEE Tr. on Circuits and Systems I: Fundamental Theory and Applications*, vol. 44, no. 10, pp. 976–988, 1997.
- [4] E. R. Westervelt, J. W. Grizzle, C. Chevallereau, J. H. Choi, and B. Morris, *Feedback Control of Dynamic Bipedal Robot Locomotion*. CRC Press, 2007.
- [5] W. M. Haddad, V. Chellaboina, and S. G. Nersesov, *Impulsive and hybrid dynamical systems*. Princeton University Press, 2006.
- [6] E. R. Westervelt, J. W. Grizzle, and D. E. Koditschek, "Hybrid zero dynamics of planar biped walkers," *IEEE Tr. on Automatic Control*, vol. 48, no. 1, pp. 42–56, 2003.
- [7] A. Ames, K. Galloway, J. Grizzle, and K. Sreenath, "Rapidly exponentially stabilizing control Lyapunov functions and hybrid zero dynamics," *IEEE Tr. on Automatic Control*, vol. 59, no. 4, pp. 876–891, 2014.

- [8] R. Gregg and M. Spong, "Reduction-based control of three-dimensional bipedal walking robots," *Int. J. of Robotics Research*, vol. 29, no. 6, pp. 680–702, 2010.
- [9] L. B. Freidovich, U. Mettin, A. S. Shiriaev, and M. W. Spong, "A passive 2-DOF walker: Hunting for gaits using virtual holonomic constraints," *IEEE Tr. on Robotics*, vol. 25, no. 5, pp. 1202–1208, 2009.
- [10] H. Razavi, A. M. Bloch, C. Chevallereau, and J. W. Grizzle, "Symmetry in legged locomotion: a new method for designing stable periodic gaits," *Autonomous Robots*, pp. 1–24, 2016.
- [11] S. Veer, M. S. Motahar, and I. Poulakakis, "Generation of and switching among limit-cycle bipedal walking gaits," in *Proc. of IEEE Conf. on Decision and Control*, 2017, pp. 5827–5832.
- [12] C. O. Saglam and K. Byl, "Switching policies for metastable walking," in Proc. of IEEE Conf. on Decision and Control, 2013, pp. 977–983.
- [13] M. S. Motahar, S. Veer, and I. Poulakakis, "Composing limit cycles for motion planning of 3D bipedal walkers," in *Proc. of IEEE Conf. on Decision and Control*, 2016, pp. 6368–6374.
- [14] S. Veer, M. S. Motahar, and I. Poulakakis, "Adaptation of limitcycle walkers for collaborative tasks: A supervisory switching control approach," in *Proc. of IEEE/RSJ Int. Conf. on Intelligent Robots and Systems*, 2017, pp. 5840–5845.
- [15] —, "Almost driftless navigation of 3D limit-cycle walking bipeds," in Proc. of IEEE/RSJ Int. Conf. on Intelligent Robots and Systems, 2017, pp. 5025–5030.
- [16] M. S. Motahar, S. Veer, and I. Poulakakis, "Steering a 3d limit-cycle walker for collaboration with a leader," in *Proc. of IEEE/RSJ Int. Conf.* on Intelligent Robots and Systems, 2017, pp. 5251–5256.
- [17] S. Veer, M. S. Motahar, and I. Poulakakis, "On the adaptation of dynamic walking to persistent external forcing using hybrid zero dynamics control," in *Proc. of IEEE/RSJ Int. Conf. on Intelligent Robots and Systems*, 2015, pp. 997–1003.
- [18] —, "Local input-to-state stability of dynamic walking under persistent external excitation using hybrid zero dynamics," in *Proc. of American Control Conf.*, 2016, pp. 4801–4806.
- [19] S. Kolathaya and A. D. Ames, "Parameter to state stability of Control Lyapunov Functions for hybrid system models of robots," *Nonlinear Analysis: Hybrid Systems*, 2016.
- [20] Q. Nguyen and K. Sreenath, "Optimal robust control for bipedal robots through control lyapunov function based quadratic programs," in *Proc.* of Robotics: Science and Systems, 2015.
- [21] K. A. Hamed, B. G. Buss, and J. W. Grizzle, "Exponentially stabilizing continuous-time controllers for periodic orbits of hybrid systems: Application to bipedal locomotion with ground height variations," *Int. J. of Robotics Research*, vol. 35, no. 8, pp. 977–999, 2016.
- [22] E. D. Sontag and Y. Wang, "New characterizations of input-to-state stability," *IEEE Tr. on Automatic Control*, vol. 41, no. 9, pp. 1283–1294, 1996.
- [23] Z.-P. Jiang and Y. Wang, "Input-to-state stability for discrete-time nonlinear systems," *Automatica*, vol. 37, no. 6, pp. 857–869, 2001.
- [24] C. Cai and A. R. Teel, "Characterizations of input-to-state stability for hybrid systems," *Systems & Control Letters*, vol. 58, no. 1, pp. 47–53, January 2009.
- [25] E. D. Sontag, "The ISS philosophy as a unifying framework for stabilitylike behavior," in *Nonlinear control in the year 2000 volume 2*. Springer, 2001, pp. 443–467.
- [26] A. E. Martin and R. D. Gregg, "Stable, robust hybrid zero dynamics control of powered lower-limb prostheses," *IEEE Tr. on Automatic Control*, vol. 62, no. 8, pp. 3930–3942, 2017.
- [27] H. K. Khalil, Nonlinear Systems. New York: Macmillan Publishing Company, 1992.
- [28] J. W. Grizzle, G. Abba, and F. Plestan, "Asymptotically stable walking for biped robots: Analysis via systems with impulse effects," *IEEE Tr.* on Automatic Control, vol. 46, no. 1, pp. 51–64, 2001.
- [29] B. Morris and J. Grizzle, "A restricted Poincaré map for determining exponentially stable periodic orbits in systems with impulse effects: Application to bipedal robots," in *Proc. of IEEE Conf. on Decision and Control and European Control Conference*, 2005, pp. 4199–4206.
- [30] A. L. Fradkov and A. Y. Pogromsky, Introduction to control of oscillations and chaos. World Scientific, 1998, vol. 35.
- [31] K. A. Hamed and R. D. Gregg, "Decentralized feedback controllers for robust stabilization of periodic orbits of hybrid systems: Application to bipedal walking," *IEEE Tr. on Control Systems Technology*, vol. 25, no. 4, pp. 1153–1167, July 2017.
- [32] A. Ramezani, S. U. Ahmed, J. Hoff, S.-J. Chung, and S. Hutchinson, "Describing robotic bat flight with stable periodic orbits," in *Biomimetic and Biohybrid Systems. Living Machines 2017*, M. Mangan,

M. Cutkosky, A. Mura, P. Verschure, T. Prescott, and N. Lepora, Eds. Springer, 2017, pp. 394–405.

- [33] P. Liljeback, K. Y. Pettersen, Ø. Stavdahl, and J. T. Gravdahl, "Controllability and stability analysis of planar snake robot locomotion," *IEEE Tr. on Automatic Control*, vol. 56, no. 6, pp. 1365–1380, 2011.
- [34] H. K. Khalil, *Nonlinear Systems*, 3rd ed. Upper Saddle River, New Jersey: Prentice hall, 2002.
- [35] J. Guckenheimer and P. Holmes, Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields. Springer-Verlag, 1996.
 [36] R. Goebel, R. G. Sanfelice, and A. R. Teel, Hybrid Dynamical Systems:
- *modeling, stability, and robustness.* Princeton University Press, 2012.
- [37] S. Lang, *Real and functional analysis*, 3rd ed. Springer-Verlag, 1993.[38] I. Poulakakis and J. W. Grizzle, "The spring loaded inverted pendulum
- as the hybrid zero dynamics of an asymmetric hopper," *IEEE Tr. on Automatic Control*, vol. 54, no. 8, pp. 1779–1793, 2009.
- [39] K. Sreenath, H.-W. Park, I. Poulakakis, and J. W. Grizzle, "Embedding active force control within the compliant hybrid zero dynamics to achieve stable, fast running on MABEL," *The Int. J. of Robotics Research*, vol. 32, no. 3, pp. 324–345, 2013.
- [40] A. D. Ames, R. D. Gregg, and M. W. Spong, "A geometric approach to three-dimensional hipped bipedal robotic walking," in *Proc. of IEEE Conf. on Decision and Control*, 2007, pp. 5123–5130.
- [41] W. Rudin, *Principles of mathematical analysis*. McGraw-Hill New York, 1964, vol. 3.
- [42] E. Sarkans and H. Logemann, "Input-to-state stability of discrete-time Lur'e systems," *SIAM Journal on Control and Optimization*, vol. 54, no. 3, pp. 1739–1768, 2016.
- [43] Z.-P. Jiang and Y. Wang, "A converse Lyapunov theorem for discretetime systems with disturbances," *Systems & Control Letters*, vol. 45, no. 1, pp. 49–58, 2002.
- [44] H. Royden and P. Fitzpatrick, *Real Analysis*, 4th ed. Prentice Hall, 2010.



Sushant Veer is a PhD Candidate in the Department of Mechanical Engineering at the University of Delaware. He received his B.Tech in Mechanical Engineering from the Indian Institute of Technology Madras (IIT-M) in 2013. His research interests lie in the control of complex dynamical systems with application to dynamically stable robots.



Rakesh received an MA in Mathematics from Delhi University in 1978 and a PhD in Mathematics from Cornell University in 1986. He is currently a Professor in the Department of Mathematical Sciences at University of Delaware. His main area of interest is Inverse Problems for Hyperbolic PDEs.



Ioannis Poulakakis received his Ph.D. in Electrical Engineering (Systems) from the University of Michigan, MI, in 2009. From 2009 to 2010 he was a post-doctoral researcher with the Department of Mechanical and Aerospace Engineering at Princeton University, NJ. Since September 2010 he has been with the Department of Mechanical Engineering at the University of Delaware, where he is currently an Associate Professor. His research interests lie in the area of dynamics and control with applications to

robotic systems, particularly dynamically dexterous legged robots. Dr. Poulakakis received the National Science Foundation CAREER Award in 2014.