ON THE DISCONTINUITY OF THE π_1 -ACTION

JEREMY BRAZAS

ABSTRACT. We show the classical π_1 -action on the *n*-th homotopy group can fail to be continuous for any *n* when the homotopy groups are equipped with the natural quotient topology. In particular, we prove the action $\pi_1(X) \times \pi_n(X) \to \pi_n(X)$ fails to be continuous for a one-point union $X = A \vee \mathbb{H}_n$ where *A* is an aspherical space such that $\pi_1(A)$ is a topological group and \mathbb{H}_n is the (n - 1)-connected, n-dimensional Hawaiian earring space \mathbb{H}_n for which $\pi_n(\mathbb{H}_n)$ is a topological abelian group.

1. INTRODUCTION

For a based topological space *X*, let $\Omega^n(X)$ denote the space of based maps $S^n \to X$ with the compact-open topology. For $n \ge 1$, the n-th homotopy group $\pi_n(X)$ may be endowed with the natural quotient topology inherited from $\Omega^n(X)$ so that the map $q : \Omega^n(X) \to \pi_n(X)$, q(f) = [f] identifying based homotopy classes is a topological quotient map.

If *X* has the homotopy type of a CW-complex, then $\pi_n(X)$ is a discrete group for all $n \ge 1$ [9, 15]. More generally, $\pi_n(X)$ can fail to be a topological group in any dimension. For instance, if \mathbb{H} is the Hawaiian earring, then $\pi_1(\mathbb{H})$ fails to be a topological group [13]. Other counterexamples for n = 1 related to free topological groups appear in [3]. The higher homotopy groups are slightly better behaved since if \mathbb{H}_n is the n-dimensional Hawaiian earring, $\pi_n(\mathbb{H}_n)$, $n \ge 2$ is a topological group isomorphic to the direct product $\prod_{n=1}^{\infty} \mathbb{Z}$ of discrete groups [16]. The improved behavior for $n \ge 2$ is essentially because, as suggested in [2], these groups are commutative in an infinitary sense. Despite this case, examples in [14] show that multiplication in $\pi_n(X)$ can still fail to be continuous.

Although multiplication in homotopy groups can fail to be continuous, π_n : **Top**_{*} \rightarrow **qTopGrp** is a homotopy invariant functor from the category of based spaces to the category of quasitopological groups¹ (abelian when $n \ge 2$) and continuous group homomorphisms. The topological structure of $\pi_1(X)$ is studied in depth in [3] and [7].

In the current paper, we address the continuity of the natural action of $\pi_1(X)$ on $\pi_n(X)$. For each $n \ge 1$, we construct a compact metric space $X \subseteq \mathbb{R}^{n+1}$ such that the action $\pi_1(X) \times \pi_n(X) \to \pi_n(X)$ is not continuous. Our example is constructed as a one-point union $X = A \vee \mathbb{H}_n$ where:

- $\pi_1(A)$ is a topological group and A is aspherical, i.e. $\pi_n(A) = 0$ for $n \ge 2$,
- \mathbb{H}_n is (n-1)-connected and $\pi_n(\mathbb{H}_n)$ is a topological abelian group.

Key words and phrases. quasitopological homotopy group, quasitopological fundamental group, continuous group action.

¹A *quasitopological group* G is a group equipped with a topology such that inversion is continuous and multiplication is continuous in each variable. See [1] for a general theory of quasitopological groups.

A fruitful approach to giving the homotopy groups a topology that makes them genuine topological groups is to apply a reflection functor τ to $\pi_n(X)$, which deletes the smallest set of open sets from the quotient topology so that one obtains continuity of multiplication. See [5] for more on this construction where the resulting topological groups are denoted $\pi_n^{\tau}(X)$. For n = 1, this topology has helped to prove new results in topological group theory [6]. An analogous construction has been carried out for the universal path space in [19]. Since writing the paper [5], the author has been unable to confirm the continuity of the π_1 -action for these "topological groups." Since these groups are topological groups, the case n = 1 is trivial since the π_1 -action is just the action by conjugation and continuity follows from the continuity of the group operations.

Problem 1. For any n > 1 and space *X*, must the natural action $\pi_1^{\tau}(X) \times \pi_n^{\tau}(X) \rightarrow \pi_n^{\tau}(X)$ be continuous?

Acknowledgements. The author is grateful to Paul Fabel for helpful conversations related to this topic.

2. Separate continuity of the π_1 -action

Given based topological spaces (X, x_0) and (Y, y_0) , let Y^X denote the space of basepoint-preserving maps $f : (X, x_0) \to (Y, y_0)$ with the compact-open topology generated by subbasic sets $\langle K, U \rangle = \{f \in Y^X | f(K) \subseteq U\}$ for $K \subseteq X$ compact and $U \subseteq Y$ open. If $x_0 \in A \subseteq X$, let $Y^{(X,A)} \subseteq Y^X$ be the subspace of maps f for which $f(A) = y_0$. Let [X, Y] be the set of basepoint-preserving homotopy classes of maps equipped with the quotient topology with respect to the map $Y^X \to [X, Y], f \mapsto [f]$ that identifies homotopy classes. In particular, $Y^{S^n} = \Omega^n(Y)$ and $\pi_n(Y) = [S^n, Y]$. For $\alpha_1, \alpha_2, ..., \alpha_k \in \Omega^n(Y), \alpha_1 \cdot \alpha_2 \cdots \alpha_k$ denotes the map $S^n \to Y$ which is the standard k-fold concatenation representing the product $[\alpha_1][\alpha_2] \cdots [\alpha_n]$ in $\pi_n(Y)$.

We recall the definition of the π_1 -action as given in [20]. Suppose the space X with basepoint $x_0 \in X$ is well-pointed, i.e. $\{x_0\} \to X$ is a cofibration. Set $Z = I \times \{x_0\} \cup X \times \{0\}$ where I = [0, 1] is the unit interval. The condition that (X, x_0) is well-pointed is equivalent to the existence of a retraction $r : X \times I \to Z$. Fixing such a retraction, let $i_1 : X \to X \times \{1\}$, $i_1(x) = (x, 1)$ be the inclusion.

For the moment, we identify $\Omega(Y)$ with the mapping space $Y^{(I,[0,1])}$. Two maps $\alpha \in \Omega(Y)$ and $\gamma \in Y^X$ uniquely determine a map $f : Z \to Y$ by $f(t, x_0) = \alpha^-(t)$ and $f(x, 0) = \gamma(x)$. Let $\alpha * \gamma = f \circ r \circ i : X \to Y$ and note this is a based map. This construction is well-defined on homotopy classes in the sense that $\pi_1(Y) \times [X, Y] \to [X, Y]$, $([\alpha], [\gamma]) \mapsto [\alpha * \gamma]$ is a group action on the set [X, Y].

Proposition 2. The map $\Omega(Y) \times Y^X \to Y^X$, $(\alpha, \gamma) \mapsto \alpha * \gamma$ is continuous and the group action $\pi_1(Y) \times [X, Y] \to [X, Y]$ is continuous in each variable.

Proof. Let $A = \{(x_0, 0), (x_0, 1)\} \subseteq X \times I$. Since there is a canonical homeomorphisms $Y^{S^1} \times Y^X \cong Y^{S^1 \vee X} \cong Y^{(Z,A)}, (\alpha, \gamma) \mapsto f$, it suffices check the map $Y^{(Z,A)} \to Y^X$, $f \mapsto f \circ r \circ i$ is continuous. It is a fact from elementary topology that if $g : X \to X'$ is a map, then the induced function $Y^{X'} \to Y^X$, $f \mapsto f \circ g$ is continuous. We apply this fact to the following maps:

(1) the map $r : X \times I \to Z$ induces a map $Y^Z \to Y^{X \times I}$ which restricts to a well-defined (since *r* is a retraction) continuous map $Y^{(Z,A)} \to Y^{(X \times I,A)}$ on subspaces,

(2) the map $i_1 : X \to X \times I$, induces a map $Y^{X \times I} \to Y^X$ which restricts to a map $Y^{(X \times I, A)} \to Y^X$.

The composition of these continuous maps agrees with the function $(\alpha, \gamma) \mapsto \alpha * \gamma$. Therefore, it is continuous.

For the group action, consider the commuting diagram



where the top map is the continuous action of loops and where q_1 and q_2 are the respective quotient maps. Although $q_1 \times q_2$ is not necessarily a quotient map, if we fix $\alpha \in \Omega(Y)$ and replace $\Omega(Y)$ and $\pi_1(Y)$ with the one-point subspaces $\{\alpha\}$ and $\{[\alpha]\}$ respectively, then the corresponding restriction of $q_1 \times q_2$ is a quotient map (since it is just q_2 times a trivial map). Since the top composition is continuous, it follows from the universal property of quotient spaces that the bottom map $\{[\alpha]\} \times [X, Y] \rightarrow [X, Y]$, which is the restriction of the group action, is continuous. Fixing γ in a similar fashion shows that the restriction $\pi_1(Y) \times \{[\gamma]\} \rightarrow [X, Y]$ is continuous. Thus the action is continuous in each variable.

Corollary 3. If $\pi_1(X)$ or $\pi_n(X)$ is discrete, then the π_1 -action $\pi_1(X) \times \pi_n(X) \to \pi_n(X)$ is continuous.

Corollary 4. Suppose X is a metric space and $\pi_1(X)$ or $\pi_n(X)$ are first countable, then the π_1 -action $\pi_1(X) \times \pi_n(X) \to \pi_n(X)$ is continuous.

Proof. Since *X* is metric, $\Omega(X) \times \Omega^n(X)$ is first countable. If $\pi_1(X) \times \pi_n(X)$ is first countable, then according to the general theory of *k*-spaces [8], the product of quotient maps $q_1 \times q_2 : \Omega(X) \times \Omega^n(X) \to \pi_1(X) \times \pi_n(X)$ is a quotient map. Applying the argument used in the second half of Proposition 2, the continuity of the action follows from the continuity of $* : \Omega(X) \times \Omega^n(X) \to \Omega^n(X)$ and the universal property of quotient maps.

3. The space A

Let $D \subseteq \mathbb{R}^2$ be the closed disk having $[0, 1] \times \{0\}$ as a diameter. For each $m \ge 1$, let $C_m \subseteq \mathbb{R}^2$ be the circle having $[0, \frac{m+1}{m}] \times \{0\}$ as a diameter. We write C_{∞} for the boundary of D and note $C_m \to C_{\infty}$ in the Hausdorff metric. Let $e_m = (\frac{m+1}{m}, 0) \in C_m$ and $e_{\infty} = (1, 0) \in C_{\infty}$.

Define $A = D \cup \bigcup_{m \ge 1} C_m$. We denote the origin (0, 0) by **0** and take this point to be the basepoint of A. Observe that A is compact but not locally path-connected at any point of $C_{\infty} \setminus \mathbf{0}$. For $m \in \{1, 2, ..., \infty\}$, let ℓ_m be a simple closed curve traversing C_m once in the counterclockwise direction such that $\ell_m \to \ell_{\infty}$ uniformly. Set $x_m = [\ell_m] \in \pi_1(A)$. Since $\ell_m \to \ell_{\infty}$, ℓ_{∞} is null-homotopic, and $\Omega(A) \to \pi_1(A)$ is continuous, x_m converges to the trivial element $1 \in \pi_1(A)$.

4



FIGURE 1. The space $A \subseteq \mathbb{R}^2$.

Remark 5. The space *A* is aspherical, i.e. $\pi_n(A) = 0$ for $n \ge 2$. To see this, one could apply general results about planar sets such as those in [10], however, it is simpler to utilize the locally path-connected coreflection lpc(A) which has the same underlying set as *A* and topology generated by the path components of open sets in *A*. In general, *X* and lpc(X) have naturally isomorphic homotopy and homology groups. In particular, since lpc(A) is homotopy equivalent to a wedge of circles, lpc(A) is aspherical and it follows that *A* is aspherical.

Lemma 6. Suppose $z_m \to z$ is a convergent sequence in a Hausdorff space Z such that there exists an open neighborhood U of z so that none of the points z_m lie in the path component of U containing z. If K is compact and locally path-connected and $f : K \to Z$ is a map, then $z_m \in f(K)$ for at most finitely many m.

Proof. Suppose $f(k_n) = z_{m_n}$ for $m_1 < m_2 < \dots$ Since *K* is compact, we may replace k_n with a convergent subsequence. Hence $k_n \rightarrow k$ in *K* where we must have f(k) = z. Consider the neighborhood *U* of *z* as in the statement of the lemma. Find a path-connected neighborhood *V* of *k* in *K* such that $f(V) \subseteq U$. Now infinitely many k_n lie in *V* and so infinitely many z_{m_n} must lie in the same path component of *U* as *z*; a contradiction.

Certainly, *A* enjoys the hypothesis of the previous lemma as applied to the convergent sequence $e_m \rightarrow e_{\infty}$. We now show that similar results exist for convergent sequences of maps.

Lemma 7. Suppose $\alpha_n : [0,1] \to A$ is a sequence of maps such that $|\{e_1, e_2, ...\} \cap \alpha_n([0,1])| \to \infty$. Then there is no map $\alpha : [0,1] \to A$ for which $\alpha_n \to \alpha$ uniformly.

Proof. Suppose α_n satisfies the hypothesis of the lemma. Without loss of generality, we may assume $|\{e_1, e_2, ...\} \cap \alpha_n([0, 1])| \ge n$. The open set $U = A \setminus \mathbf{0}$ has open path components $D \setminus \mathbf{0}$ and $C_m \setminus \mathbf{0}$, $m \ge 1$. For the moment, fix $n \ge 1$. The open set $\alpha_n^{-1}(U)$ is a countable disjoint union of open intervals. By assumption, we may find components $(a_1, b_1), ..., (a_n, b_n)$ of $\alpha_n^{-1}(U)$, naturally ordered within [0, 1], and points $t_{2k} \in (a_k, b_k)$ such that $\alpha(t_{2k}) = e_{m_k}$ for *n*-distinct integers $m_1, m_2, ..., m_n$. Let $t_{2k-1} = a_k$ for k = 1, ..., n and $t_{2n+1} = b_n$. Now we have $t_0 < t_1 < t_2 < ... < t_{2n+1}$ where $f(t_{2k-1}) = \mathbf{0}$ and $\alpha(t_{2k}) = e_{m_k}$. In particular, $||\alpha_n(t_j) - \alpha_n(t_{j+1})|| > 1$ for all $n \ge 1$, j = 0, 1, ..., 2n.

Suppose, to obtain a contradiction, that $\alpha_n \to \alpha$ uniformly for some $\alpha : [0, 1] \to X$. Since the set $\{\alpha_n | n \ge 1\}$ is uniformly equicontinuous, there is a $\delta > 0$ such that $\|\alpha_n(s) - \alpha_n(t)\| < 1$ for all $n \ge 1$ and $s, t \in [0, 1]$ with $|s - t| < \delta$. Find n such that $\frac{1}{n} < \delta$ and recall the previous paragraph applied to α_n . Since the intervals (a_k, b_k) , k = 1, ..., n are disjoint, at least one of them, call it (a_{k_0}, b_{k_0}) , must have diameter less than $\frac{1}{n}$. Since $t_{2k_0} \in (a_{k_0}, b_{k_0}) = (t_{2k_0-1}, b_{k_0})$, we have $|t_{2k_0-1} - t_{2k_0}| < \frac{1}{n} < \delta$ and thus $||\alpha_n(t_{2k_0-1}) - \alpha_n(t_{2k_0})|| < 1$; a contradiction of the last sentence in the previous paragraph.

Lemma 8. Suppose K is a Peano continuum and $f_n : K \to A$ is a sequence of maps such that $|\{e_1, e_2, ...\} \cap f_n(K)| \to \infty$. Then there is no map $f : K \to A$ for which $f_n \to f$ uniformly.

Proof. Suppose f_n satisfies the hypothesis of the lemma. Since K is a Peano continuum, there is an onto map $\beta : [0, 1] \to K$. Let $\alpha_n = f_n \circ \beta$. If $f_n \to f$ uniformly, then $\alpha_n \to \alpha = f \circ \beta$ uniformly and $|\{e_1, e_2, ...\} \cap \alpha_n([0, 1])| \to \infty$; a contradiction of Lemma 7.

Proposition 9. $\pi_1(A)$ is freely generated by the set $\{x_1, x_2, ...\}$ and is a Hausdorff topological group.

Proof. Note that $A_k = D \cup \bigcup_{1 \le m \le k} C_m$ is a retract of A by maps $\rho_k : A \to A_k$ collapsing $C_{k+1}, C_{k+2}, ...$ homeomorphically onto C_{∞} . The space A_k is homotopy equivalent to the wedge of circles $\bigcup_{1 \le m \le k} C_m$ so $\pi_1(A_k)$ is freely generated by the set $\{x_1, ..., x_k\}$. The inclusions $A_1 \to A_2 \to ...$ induce a directed system of discrete free groups $\pi_1(A_k) = F(x_1, ..., x_k)$ such that $\varinjlim_k \pi_1(A_k) = \varinjlim_k F(x_1, ..., x_k) = F(x_1, x_2, ...)$. Moreover, the inclusions $A_n \to A$ induce a canonical homomorphism $\phi : \varinjlim_k \pi_1(A_k) \to \pi_1(A)$. We check that ϕ is a group isomorphism. If $\alpha : S^1 \to A$ is a loop, then by Lemma 6, α has image in $A \setminus \{e_{M+1}, e_{M+2}, ...\}$ for some M. Since $A \setminus \{e_{M+1}, e_{M+2}, ...\}$ deformation retracts onto A_M , α is homotopic to a loop in A_M . This verifies the surjectivity of ϕ . Suppose $\alpha : S^1 \to A_k$ is a loop which contracts by a null-homotopy $h : S^1 \times [0, 1] \to A$. Once again applying Lemma 6, h must have image in $A \setminus \{e_{M+1}, e_{M+2}, ...\}$ for some M > k. Composing h with a deformation retract of $A \setminus \{e_{M+1}, e_{M+2}, ...\}$ onto A_M , we obtain a null-homotopy of α within A_M . Thus α represents the trivial element in $\varinjlim_k \pi_1(A_k)$, proving injectivity.

A proof from first principles that $\pi_1(A)$ is a Hausdorff topological group is much more technical; we discuss the ingredients of the proof here, referring to more general work. Since *A* is a planar set, $\pi_1(A)$ is Hausdorff according to [7, Corollary 20]. To see that $\pi_1(A)$ is a topological group, we call upon results in [3]. The space $C_{\infty} \cup \bigcup_n C_n$ is precisely the space in [3, Example 4.24] whose fundamental group is shown to be the free Markov topological group generated by the subspace $\{x_1, x_2, ..., x_{\infty}\} \subset \pi_1(A)$. It is known that if a 2-cell e^2 is attached to a path-connected space *Y*, then the inclusion $Y \to Y \cup e^2$ induces a homomorphism $\pi_1(Y) \to \pi_1(Y \cup e^2)$ which is a topological quotient map [4, Lemma 4.4]. Since *A* is obtained from $C_{\infty} \cup \bigcup_m C_m$ by attaching a 2-cell with boundary C_{∞} , $\pi_1(A)$ is the topological quotient group of the topological group $\pi_1(C_{\infty} \cup \bigcup_m C_m)$ by factoring the normal closure of $\{[\ell_{\infty}]\}$. Since the quotient of a topological group equipped with the quotient topology is a topological group, $\pi_1(A)$ is a topological group. \Box

Remark 10. One can conclude from the previous proof that $\pi_1(A)$ is isomorphic to the free Graev topological group generated by the space $\{x_1, x_2, ..., x_\infty\}$ with basepoint x_∞ .

4. The space \mathbb{H}_n

Fix $n \ge 2$ and let $S_m \subseteq \mathbb{R}^{n+1}$ be the n-dimensional sphere centered at (1/m, 0, ..., 0) with radius 1/m. Now $\mathbb{H}_n = \bigcup_{m\ge 1} S_m$ is the *n*-dimensional Hawaiian earring space with basepoint $\mathbf{0} = (0, 0, ..., 0)$. Let $y_m \in \pi_n(\mathbb{H}_n)$ be the homotopy class of the canonical homeomorphism $\gamma_m : S^n \to S_m \subset \mathbb{H}_n$ that generates $\pi_n(S_m) \cong \mathbb{Z}$.



FIGURE 2. The 2-dimensional Hawaiian earring $\mathbb{H}_2 \subseteq \mathbb{R}^3$.

We note some technical achievements regarding \mathbb{H}_n :

- (1) [12] \mathbb{H}_n is (n 1)-connected.
- (2) [12] If $r_M : \mathbb{H}_n \to S_M$ is the natural retraction collapsing $\bigcup_{m \neq M} S_m$ to **0**, then the induced homomorphism $\pi_n(\mathbb{H}_n) \to \prod_{m \geq 1} \pi_n(S_m), [\alpha] \mapsto ((r_1)_{\#}([\alpha]), (r_2)_{\#}([\alpha]), ...)$ is an isomorphism. In particular, if $\gamma : S^n \to \mathbb{H}_n$ is a map such that $r_m \circ \gamma$ is null-homotopic for all *m*, then γ is null-homotopic. Thus $\pi_n(\mathbb{H}_n) \cong \prod_{m \geq 1} \mathbb{Z}$.
- (3) [16] The group isomorphism π_n(ℍ_n) → Π_{m≥1} Z is an isomorphism of topological groups when Π_{m≥1} Z is viewed as the direct product of discrete groups.

Although not explicitly stated in [12], it follows that $\pi_n(\mathbb{H}_n)$ is isomorphic to the n-th singular homology group $H_n(\mathbb{H}_n)$. In particular, since \mathbb{H}_n is (n - 1)connected, the Hurewicz Theorem implies that the Hurewicz homomorphism $h : \pi_n(\mathbb{H}_n) \to H_n(\mathbb{H}_n)$ is an isomorphism. One might initially be concerned about the applicability of the Hurewicz Theorem to spaces that are not CW-complexes, however, the theory of CW-approximations and weak homotopy equivalences (see p.352-357 of [18]) permits the application.

Remark 11. If $Y = \bigvee_{i=1}^{m} S^1 \vee \bigvee_{j=1}^{k} S^n$ is a finite wedge of 1-spheres and *n*-spheres, then the universal covering space \widetilde{Y} may be viewed as the Caley graph *T* with vertex set $F(x_1, ..., x_m)$ with a copy of $\bigvee_{i=1}^{k} S^n$ attached at each vertex. By collapsing the tree

to a point, one observes that $\pi_n(Y) \cong \pi_n(\widetilde{Y}) \cong \bigoplus_{w \in F(x_1,...,x_m)} \pi_n(\bigvee_{j=1}^k S^n) \cong \bigoplus_{\mathbb{N}} \mathbb{Z}$. In the following lemma, we seek to obtain an analogous result when we replace $\bigvee_{j=1}^k S^n$ with the infinite shrinking wedge \mathbb{H}_n , however, care must be taken since collapsing edges will not necessarily preserve homotopy type.

Lemma 12. Let *T* be a finite tree. For each vertex *v* of *T*, attach a copy of \mathbb{H}_n to *T* by the identification $v \sim \mathbf{0}$. Denote the resulting space *Z*. Then *Z* is (n - 1)-connected and the quotient map $\eta : Z \to Z/T$ induces a continuous group isomorphism $\eta_{\#} : \pi_n(Z) \to \pi_n(Z/T)$. In particular, $\pi_n(Z) \cong \prod_v \prod_{n \ge 1} \mathbb{Z} \cong \prod_N \mathbb{Z}$.

Proof. The continuity of $\eta_{\#}$ follows from functorality of π_n . Our proof of the other claims is carried out by induction on the number of edges of *T*. The lemma follows from the results on \mathbb{H}_n above when *T* has a single vertex.

Suppose the lemma holds for all trees with *n* vertices. Suppose *T* has n + 1 vertices. Let $T' \subset T$ be a subtree with *n* edges and *E* be the remaining edge of *T* so that $T' \cup E = T$. For convenience, identify *E* with [0, 1] so that $\{1\} = T' \cap E$ and the vertex 1 is an end. Let $K_1, ..., K_p$ be the copies of \mathbb{H}_n attached to the vertices of *T'* so that $Z' = T' \cup K_1 \cup ... \cup K_p$. Let K_0 denote the copy of \mathbb{H}_n attached at 0. By the induction hypothesis *Z'* is (n-1)-connected, the quotient map $\eta' : Z' \to Z'/T'$ induces an isomorphism $(\eta')_{\#} : \pi_n(Z') \to \pi_n(Z'/T')$. By choosing a homeomorphism $Z'/T' \cong \mathbb{H}_n$, we may identify $\pi_n(Z'/T')$ and $H_n(Z'/T')$ with $\prod_{m\geq 1} \mathbb{Z}$.

Since *Z* is the one-point union of (n - 1)-connected spaces $K_0 \cup [0, 1/2]$ and $[1/2, 1] \cup Z'$, which are both locally contractible at 1/2, *Z* is also (n - 1)-connected. Since both *Z* and Z'/T' are (n - 1)-connected, it now suffices (by the Hurewicz Theorem) to show that $\eta_* : H_n(Z) \to H_n(Z/T)$ is an isomorphism.

Set $U = K_0 \cup [0, 2/3)$ and $V = (1/3, 1] \cup Z'$. These open sets deformation retract onto K_0 and Z' respectively. Since $U \cap V$ is contractible, the Mayer-Vietoris sequence for the triad (Z, U, V) implies that the inclusions $i : K_0 \to Z$ and $j : Z' \to Z$ induce an isomorphism $\phi = i_* - j_* : H_n(K_0) \oplus H_n(Z') \to H_n(Z)$.

The inclusions $k : K_0 \to Z/T$, $l : Z'/T' \to Z/T$ induce a homomorphism $k_* - l_* : H_n(K_0) \oplus H_n(Z'/T') \to H_n(Z/T)$, which makes the bottom square in following diagram commute:



To ensure the top map is an isomorphism it suffices to show that $k_* - l_*$ is an isomorphism. However, by identifying $H_n(K_0) \oplus H_n(Z'/T')$ with $\prod_{m\geq 1} \mathbb{Z} \oplus \prod_{m\geq 1} \mathbb{Z}$ and $H_n(Z/T)$ with $\prod_{m\geq 1} \mathbb{Z}$, we see that $k_* - l_* : \prod_{m\geq 1} \mathbb{Z} \oplus \prod_{m\geq 1} \mathbb{Z} \to \prod_{m\geq 1} \mathbb{Z}$ is a coordinate shuffling isomorphism determined by a homeomorphism $Z/T \cong \mathbb{H}_n$.

Corollary 13. Let T be a tree with vertex set V and basepoint $v_0 \in V$ and Z be the space T with a copy K_v of \mathbb{H}_n attached at each $v \in V$. If A_v is the arc in T from v_0 to v, then the inclusion maps $A_v \cup K_v \to Z$ induce a group isomorphism $\rho : \bigoplus_{v \in V} \pi_n(A_v \cup K_v) \to \pi_n(Z)$. In particular, $\pi_n(Z) \cong \bigoplus_{v \in V} \prod_N \mathbb{Z}$ as a group.

Proof. Lemma 12 is precisely the case when *T* is finite. When *T* is infinite, ρ is defined on elements as $\rho(\sum_{v} n_v[\delta_v]) = \left[\prod_{n_v \neq 0} \delta_v^{n_v}\right]$ where all but finitely many $n_v \in \mathbb{Z}$ are non-zero. Every map $S^n \to Z$ and null-homotopy of a map $S^n \to Z$ must have image in a subspace $Z_F \subset Z$ consisting of a finite subtree $F \subset T$ and all K_v with $v \in F$. Hence the finite case directly implies the infinite case.

Remark 14. It is well-known that the homomorphism $\eta_{\#}$ from Lemma 12 fails to be surjective if we allow n = 1.

5. The space
$$X = A \vee \mathbb{H}_n$$

We now fix $n \ge 1$ and consider the one-point union $X = A \lor \mathbb{H}_n$, which may be embedded as a compact subspace of \mathbb{R}^{n+1} . The main difficulty in what follows is ensuring that $\pi_1(X)$ and $\pi_n(X)$ are T_1 .



FIGURE 3. The space $X = A \vee \mathbb{H}_n$ for n = 2.

Proposition 15. The inclusion $A \to X$ induces an isomorphism $\pi_1(A) \to \pi_1(X)$ of topological groups.

Proof. Since the space \mathbb{H}_n is first countable and locally simply connected, it follows from a theorem of Griffiths [17] that the inclusions $A \to X$ and $\mathbb{H}_n \to X$ induce an isomorphism $\pi_1(A) * \pi_1(\mathbb{H}_n) \to \pi_1(X)$ from the free product. Since \mathbb{H}_n is simply connected, $\pi_1(A) \to \pi_1(X)$ is a group isomorphism. This isomorphism is continuous by the functorality of π_1 . Lastly, the retraction $X \to A$ collapsing \mathbb{H}_n to a point induces a retraction $\pi_1(X) \to \pi_1(A)$ of quasitopological groups for which $\pi_1(A) \to \pi_1(X)$ is a section. Thus $\pi_1(A) \to \pi_1(X)$ is a topological embedding and is therefore an isomorphism of quasitopological groups. Since $\pi_1(A)$ is a topological group (Prop. 9), so is $\pi_1(X)$.

Recall that $A_m = D \cup C_1 \cup ... \cup C_m$. Let $X_m = A_m \vee \mathbb{H}_n$ and $Y_m = C_1 \cup ... \cup C_m \vee \mathbb{H}_n$ so that $X_m = Y_m \vee D$. Note that the inclusion $Y_m \to X_m$ is a homotopy equivalence which induces an isomorphism $\pi_r(Y_m) \cong \pi_r(X_m)$ of quasitoplogical groups for all $r \ge 1$. Hence, we may replace X_m with Y_m when necessary.

Consider the finite CW-complex subspaces: $X_{m,k} = A_m \cup \bigcup_{1 \le j \le k} S_j$ and $Y_{m,k} = C_1 \cup ... \cup C_m \cup \bigcup_{1 \le j \le k} S_j$ so that $X_{m,k} = Y_{m,k} \vee D$. We may view $X_m = \varprojlim_k X_{m,k}$ and

 $Y_m = \lim_{k \to \infty} Y_{m,k}$ using the inverse sequences



where the retractions $f_{k+1,k}$ and $g_{k+1,k}$ collapse S_{k+1} to a point. Moreover, the natural retractions $f_k : X_m \to X_{m,k}$ and $g_k : Y_m \to Y_{m,k}$ induce continuous homomorphisms $\psi : \pi_n(X_m) \to \varprojlim_k \pi_n(X_{m,k})$ and $\phi : \pi_n(Y_m) \to \varprojlim_k \pi_n(Y_{m,k})$ respectively which make the following diagram commute:

$$\pi_{n}(X_{m}) \xrightarrow{\psi} \varprojlim_{k} \pi_{n}(X_{m,k})$$

$$\cong \uparrow \qquad \uparrow \qquad \uparrow \cong$$

$$\pi_{n}(Y_{m}) \xrightarrow{\phi} \varprojlim_{k} \pi_{n}(Y_{m,k})$$

Here, $\pi_n(X_{m,k})$ and $\pi_n(Y_{m,k})$ are discrete groups since $X_{m,k}$ and $Y_{m,k}$ are locally contractible [15] and the inverse limit groups have the natural inverse limit topology (as a subspace of the direct product). The vertical morphisms are homeomorphisms and group isomorphisms.

Lemma 16. $\psi : \pi_n(X_m) \to \varprojlim_k \pi_n(X_{m,k})$ and $\phi : \pi_n(Y_m) \to \varprojlim_k \pi_n(Y_{m,k})$ are continuous group isomorphisms.

Proof. As noted above, it suffices to prove the lemma for either ψ or ϕ ; we apply previous results to prove it for ϕ .

For n = 1, Y_m is one-dimensional so the injectivity of ϕ follows from the π_1 -shape injectivity of one-dimensional spaces [11].

Suppose $n \ge 2$. Although Y_m is not locally contractible at its basepoint, it is locally simply connected and therefore admits a universal covering space. Let $p: \widetilde{Y_m} \to Y_m$ and $p_k: \widetilde{Y_{m,j}} \to Y_{m,k}$ be respective the universal covering maps. We view $\widetilde{Y_{m,k}}$ as the Caley graph on the free group $F(x_1, ..., x_m)$ with a copy of $S_1 \lor ... \lor S_k$ attached at each vertex. Similarly, we view $\widetilde{Y_m}$ as the Caley graph on $F(x_1, ..., x_m)$ with a copy of \mathbb{H}_n attached at each vertex.

Since $\widetilde{Y_{m,j+1}}$ is simply connected, the map $g_{j+1,j} \circ p_{j+1}$ induces a retraction $\widetilde{g}_{j+1,j}$: $\widetilde{Y_{m,j+1}} \rightarrow \widetilde{Y_{m,j}}$ such that $p_j \circ \widetilde{g}_{j+1,j} = g_{j+1,j} \circ p_{j+1}$. In particular, $\widetilde{g}_{j+1,j}$ collapses each copy of S_{j+1} to the vertex at which it is adjoined. We now have an inverse sequence of the maps p_j such that $p = \lim_{j \to \infty} p_j$:



The canonical retractions $\widetilde{g}_j : \widetilde{Y}_m \to \widetilde{Y}_{m,j}$ collapse the copies of $S_{j+1}, S_{j+2}, ...$ to the vertex at which they are adjoined and together induce a continuous homomorphism $\widetilde{\phi} : \pi_n(\widetilde{Y}_m) \to \varprojlim_j \pi_n(\widetilde{Y}_{m,j})$. Since covering maps induce isomorphisms on $\pi_n, n \ge 2$ and an inverse limit of isomorphisms is an isomorphism, we see that the vertical morphisms in the lower square of the following diagram are isomorphisms.



Hence, to show that ϕ is a group isomorphism it suffices to show that ϕ is a group isomorphism.

From Remark 11, we may identify

$$\pi_n(\widetilde{Y_{m,k}}) \cong \bigoplus_{w \in F(x_1, \dots, x_m)} \prod_{j=1}^{\kappa} \mathbb{Z}$$

with elements represented as sums $\sum_{w} n_w(s_1^w, ..., s_k^w)$ where $s_j^k \in \mathbb{Z}$ and all but finitely many $n_w \in \mathbb{Z}$ are zero. Under this identification, the projection map $(\widetilde{g}_{k+1,k})_{\#} : \pi_n(\widetilde{Y}_{m,k+1}) \to \pi_n(\widetilde{Y}_{m,k})$ becomes a countable summand

$$\bigoplus_{w \in F(x_1, \dots, x_m)} \prod_{j=1}^{k+1} \mathbb{Z} \to \bigoplus_{w \in F(x_1, \dots, x_m)} \prod_{j=1}^k \mathbb{Z}, \qquad \sum_w n_w(s_1^w, \dots, s_k^w, s_{k+1}^w) \mapsto \sum_w n_w(s_1^w, \dots, s_k^w)$$

of the projection map $\prod_{j=1}^{k+1} \mathbb{Z} \to \prod_{j=1}^{k} \mathbb{Z}$. By Corollary 13, there is a natural isomorphism

$$\pi_n(\widetilde{X_m}) \cong \bigoplus_{w \in F(x_1, \dots, x_m)} \prod_{k \ge 1} \mathbb{Z}.$$

With this identification, ϕ becomes identified with the isomorphism

$$\bigoplus_{x \in F(x_1, \dots, x_m)} \prod_{k \ge 1} \mathbb{Z} \cong \lim_{k} \bigoplus_{w \in F(x_1, \dots, x_m)} \prod_{j=1}^{k} \mathbb{Z}$$

defined on elements as $\sum_{w} n_w(s_1^w, s_2^w, ...) \mapsto (\sum_{w} n_w(s_1^w, ..., s_k^w))_k$.

Corollary 17. $\pi_n(X_m) \cong \pi_n(Y_m) \cong \bigoplus_{\mathbb{N}} \prod_{\mathbb{N}} \mathbb{Z}$ as groups.

Corollary 18. As a topological space, $\pi_n(X_m)$ is Hausdorff.

Proof. Any space which continuously injects into a Hausdorff space is Hausdorff. Since an inverse limit of discrete groups is Hausdorff and $\psi : \pi_n(X_m) \rightarrow \lim_{j \to \infty} \pi_n(X_{m,j})$ is a continuous injection by the previous lemma, $\pi_n(X_m)$ is Hausdorff.

Now for each *m*, we consider the retractions $R_{m+1,m} : X_{m+1} \to X_m$ which send C_{m+1} homeomorphically to C_{∞} . This forms an inverse system for which $X = \underset{k \to m}{\lim} X_m$. The retractions $R_m : X \to X_m$ that collapse each circle of $\bigcup_{k>m} C_k$ to C_{∞} so that $R_{m+1} \circ R_{m+1,m} = R_m$ induce a canonical continuous homomorphism $\Psi : \pi_n(X) \to \underset{m}{\lim} \pi_n(X_m)$.

Lemma 19. $\Psi : \pi_n(X) \to \varprojlim_m \pi_n(X_m)$ is a continuous injection.

Proof. Continuity of Ψ is immediate from the construction. Suppose $\gamma : S^n \to X$ is a map such that $\gamma_m = R_m \circ \gamma : S^n \to X_m$ is null-homotopic for each m. By Lemma 6, γ has image in $X \setminus \{e_{m_0+1}, e_{m_0+2}, ...\}$ for some $m_0 \ge 1$. Since $X \setminus \{e_{m_0+1}, e_{m_0+2}, ...\}$ deformation retracts onto X_{m_0} , γ is homotopic (by a homotopy in X) to a loop γ' with image in X_{m_0} . Since $\gamma' = R_{m_0} \circ \gamma' \simeq R_{m_0} \circ \gamma$ in X_{m_0} , γ' is null homotopic in X_{m_0} . Thus γ is null-homotopic in X.

Corollary 20. $\pi_n(X)$ is Hausdorff.

Proof. By Corollary 18, $\pi_n(X_m)$ is Hausdorff. Since the inverse limit of Hausdorff spaces is Hausdorff, $\lim_{m} \pi_n(X_m)$ is a Hausdorff group. By Lemma 19, $\pi_n(X)$ continuously injects into a Hausdorff group and is therefore Hausdorff.

Remark 21. To explicitly identify the isomorphism class of $\pi_n(X)$ as a group, replace *X* with its locally path-connected coreflection and contract the disk *D* to obtain the weakly homotopy equivalent space $X' = \bigcup_{m \ge 1} C_m \vee \mathbb{H}_n$. By applying Corollary 13 to the universal covering space of *X'*, we have group isomorphisms $\pi_n(X) \cong \pi_n(X') \cong \bigoplus_{w \in F(x_1, x_2, ...)} \prod_{k \ge 1} \mathbb{Z} \cong \bigoplus_{\mathbb{N}} \prod_{\mathbb{N}} \mathbb{Z}$.

6. Discontinuity of the π_1 -action

Finally, we show that for the *n*-dimensional space $X \subseteq \mathbb{R}^{n+1}$ considered in the previous sections, the π_1 -action, denoted here as the function $a : \pi_1(X) \times \pi_n(X) \rightarrow \pi_n(X), a(x, y) = x * y$, is not continuous.

Theorem 22. For any $n \ge 1$, the natural group action $a : \pi_1(X) \times \pi_n(X) \to \pi_n(X)$ is not continuous.

Note that the case n = 1 indicates that if $X = A \vee \mathbb{H}$ is the one-point union of A and the Hawaiian earring, then the action by conjugation in $\pi_1(X)$ is discontinuous.

Recall that $\pi_1(X)$ is freely generated by the set $\{x_1, x_2, ...\}$ and $y_j = [\gamma_j]$ is a generator of $\pi_n(S_j)$. Let $C = \{(x_i x_{i+1} ... x_{i+j}) * y_j^i | i, j \ge 1\}$. We show that *C* is closed in $\pi_n(X)$ but that $a^{-1}(C)$ is not closed in $\pi_1(X) \times \pi_n(X)$. Our choice of *C* is inspired Fabel's argument in [14].

Proposition 23. If $\gamma : S^n \to X$ represents $(x_i x_{i+1} \dots x_{i+j}) * y_{j'}^i$ then $\bigcup_{k=i}^{i+j} C_k \subset \gamma(S^n)$. In particular, $\{e_i, e_{i+1}, \dots, e_{i+j}\} \subseteq \gamma(S^n)$.

Proof. As we have argued previously, γ is homotopic to a loop δ with image in $X_{m_0} \cap Im(\gamma)$ for some $m_0 \ge i + j$. Let \widetilde{X}_{m_0} be the universal covering space of X_{m_0} . Again, we identify \widetilde{X}_{m_0} with the Caley graph of the free group $F(x_1, ..., x_{m_0})$ (with vertices identified with elements of $F(x_1, ..., x_{m_0})$ and basepoint the identity element e) with a copy of the simply connected space $D \lor \mathbb{H}_n$ attached at each vertex. Since S^n is simply connected, there is a unique lift $\delta : (S^n, (1, 0, ..., 0)) \to (\widetilde{X}_m, e)$. Since the

Caley graph is uniquely arc-wise connected, in order for δ to represent $(x_i x_{i+1} \dots x_{i+j}) * y_j^i$, the image of δ must contain the unique minimal edge path from the vertex e to the vertex $(x_i x_{i+1} \dots x_{i+j})$ along the edges $[e, x_i], [x_i, x_i x_{i+1}], \dots, [x_i x_{i+1} \dots x_{i+j-1}, x_i x_{i+1} \dots x_{i+j}]$. These edges are mapped respectively by the covering map onto $C_i, C_{i+1}, \dots, C_{i+j}$, so we have $\bigcup_{k=i}^{i+j} C_k \subset \delta(S^n) \subset \gamma(S^n)$.

Proposition 24. $a^{-1}(C)$ is not closed in $\pi_1(X) \times \pi_n(X)$.

Proof. Since *C* does not contain the identity of $\pi_n(X)$, the identity (1, 1) of $\pi_1(X) \times \pi_n(X)$ is not in $a^{-1}(C)$. We show that (1, 1) is a limit point of $a^{-1}(C)$. Let $V \times U$ be a basic open neighborhood of (1, 1) in $\pi_1(X) \times \pi_n(X)$. Let $q_i : \Omega^i(X) \to \pi_i(X)$ be the natural quotient map. Find an open neighborhood *W* of x_0 such that $\langle S^n, W \rangle \subseteq q_n^{-1}(U)$ in $\Omega^n(X)$. Since all but finitely many of the spheres $S_j \subset \mathbb{H}_n$ lie in *W* all but finitely many of the generators $\gamma_j : S^n \to X$ have image in $\langle S^n, W \rangle$. It follows that there exists a *J* such that if γ_j^i is the map γ_j concatenated with itself *i*-times, then $\gamma_j^i \in \langle S^n, W \rangle$ for all $j \ge J$ and all $i \ge 1$. Thus for homotopy classes, we have $y_i^i \in U$ for all $j \ge J$ and $i \ge 1$.

Now that *J* is fixed, notice that the sequence $\ell_i \cdot \ell_{i+1} \cdot \ldots \cdot \ell_{i+J}$, $i \ge 1$ of loops in $\Omega(X)$ converges uniformly to the null-homotopic loop $\ell_{\infty} \cdot \ell_{\infty} \cdots \ell_{\infty}$ (with *J* + 1-factors). The continuity of $q_1 : \Omega(X) \to \pi_1(X)$ now gives $x_i x_{i+1} \cdots x_{i+J} \to 1$ as a sequence in $\pi_1(X)$. Thus there is an *I* such that $x_I x_{I+1} \cdots x_{I+J} \in V$. Since we also have $y_J^I \in U$, it follows that $((x_I x_{I+1} \ldots x_{I+J}), y_I^I) \in (V \times U) \cap a^{-1}(C)$.

Proposition 25. *C* is closed in $\pi_n(X)$.

Proof. Since $q_n : \Omega^n(X) \to \pi_n(X)$, $q_n(\gamma) = [\gamma]$ is quotient, it suffices to show $q_n^{-1}(C)$ is closed in the metric space $\Omega^n(X)$ of based maps $S^n \to X$. Suppose $\delta_k \to \delta$ is a convergent sequence in $\Omega^n(X)$ and $q_n(\delta_k) = [\delta_k] \in C$ for all k. In particular, suppose $[\delta_k] = (x_{i_k}x_{i_k+1}...x_{i_k+j_k}) * y_{j_k}^{i_k}$ for sequences of integers i_k , $j_k \in \mathbb{N}$. It suffices to show $[\delta] \in C$. To start, we will show that both sequences i_k and j_k must be eventually constant.

Suppose j_k is unbounded. If necessary, replace j_k with a subsequence to ensure that $j_1 < j_2 < ...$ By Proposition 23, we have $\{e_{i_k}, e_{i_k+1}, ..., e_{i_k+j_k}\} \subset \delta_k(S^n)$ for each k. Let $R : X \to A$ be the retraction collapsing \mathbb{H}_n to the basepoint and set $f_k = R \circ \delta_k$ and $f = R \circ \delta$. We have $f_k \to f$ uniformly and $\{e_{i_k}, e_{i_k+1}, ..., e_{i_k+j_k}\} \subset f_k(S^n)$, which is a contradiction of Lemma 8. Thus j_k must be bounded.

Since $J = \{j_k | k \ge 1\}$ is a finite set, $\bigvee_{j \in J} S_j \subset \mathbb{H}_n$ is a finite wedge of n-spheres. In particular, $\pi_n (\bigvee_{j \in J} S_j) = \prod_{j \in J} \mathbb{Z}$ is a discrete abelian group freely generated by $\{y_j | j \in J\}$. Since $\delta_k \to \delta$ and q_n is continuous, we have $[\delta_k] \to [\delta]$ in $\pi_n(X)$. Consider the retraction $r : X \to \bigvee_{j \in J} S_j$ which collapses A and all n-spheres S_j , $j \notin J$ to the basepoint. By functorality, the homomorphism $r_{\#} : \pi_n(X) \to \pi_n (\bigvee_{j \in J} S_j)$ is continuous and thus $r_{\#}([\delta_k]) = y_{j_k}^{i_k} \to r_{\#}([\delta])$. However, the only sequences that converge in a discrete group are eventually constant. Thus i_k must be eventually constant and, in particular, bounded.

Since both i_k and j_k are bounded, the set $\{[\delta_k]|k \ge 1\} = \{(x_{i_k}x_{i_k+1}...x_{i_k+j_k})*y_{j_k}^{i_k}|k \ge 1\}$ is a finite subset of $\pi_n(X)$. By Corollary 20, $\pi_n(X)$ is Hausdorff. Since finite subsets of

Hausdorff spaces are closed, $\{[\delta_k]|k \ge 1\}$ is closed in $\pi_n(X)$. Finally, since $[\delta_k] \to [\delta]$, we must have $[\delta] \in \{[\delta_k]|k \ge 1\} \subset C$.

References

- A. Arhangel'skii, M. Tkachenko, Topological Groups and Related Structures, Series in Pure and Applied Mathematics, Atlantis Studies in Mathematics, 2008.
- [2] M.G. Barratt, J. Milnor, An example of anomalous singular theory, Proc. Amer. Math. Soc. 13 (1962) 293–297.
- [3] J. Brazas, *The topological fundamental group and free topological groups*, Topology Appl. 158 (2011) 779–802.
- [4] J. Brazas, Homotopy Mapping Spaces, Ph.D. Dissertation, University of New Hampshire, 2011.
- [5] J. Brazas, The fundamental group as a topological group, Topology Appl. 160 (2013) 170-188.
- [6] J. Brazas, Open subgroups of free topological groups, Fundamenta Mathematicae 226 (2014) 17-40.
- [7] J. Brazas, P. Fabel, On fundamental groups with the quotient topology, J. Homotopy and Related Structures 10 (2015) 71–91.
- [8] R. Brown, Topology and Groupoids, Booksurge PLC, 2006.
- [9] J.S. Calcut, J.D. McCarthy, Discreteness and homogeneity of the topological fundamental group, Topology Proceedings 34 (2009), 339–349.
- [10] J. Cannon, G. Conner, A. Zastrow, One-dimensional sets and planar sets are aspherical, Topology Appl. 120 (2002) 23–45.
- [11] K. Eda, K. Kawamura, The fundamental groups of one-dimensional spaces, Topology Appl. 87 (1998) 163–172.
- [12] K. Eda, K. Kawamura, Homotopy and Homology Groups of the n-dimensional Hawaiian earring, Fund. Math. 165 (2000) 17–28.
- [13] P. Fabel, Multiplication is discontinuous in the Hawaiian earring group, Bull. Polish Acad. Sci. Math. 59 (2011) 77–83
- [14] P. Fabel, Compactly generated quasitopological homotopy groups with discontinuous multiplication, Topology Proc. 40 (2012) 303–309.
- [15] H. Ghane, Z. Hamed, B. Mashayekhy, H. Mirebrahimi, *Topological homotopy groups*, Bull. Belg. Math. Soc. Simon Stevin, 15:3 (2008), 455–464.
- [16] H. Ghane, Z. Hamed, B. Mashayekhy, H. Mirebrahimi, On topological homotopy groups of n-Hawaiian like spaces, Topology Proc, 36 (2010), 255-266.
- [17] H.B. Griffiths, The fundamental group of two spaces with a point in common, Quart. J. Math. Oxford, 5 (1954). 175–190.
- [18] A. Hatcher, Algebraic Topology, Cambridge Univ. Press, 2002.
- [19] Z. Virk, A. Zastrow, A new topology on the universal path space, Topology Appl. 231 (2017), 186–196.
- [20] G.W. Whitehead, *Elements of Homotopy Theory*, Springer Verlag, Graduate Texts in Mathematics. 1978.

Department of Mathematics; West Chester University; 25 University Ave.; West Chester, PA, 19383