Efficiently Realizing Interval Sequences

Amotz Bar-Noy, Keerti Choudhary, David Peleg, Dror Rawitz[§]

Abstract

We consider the problem of realizable interval-sequences. An interval sequence comprises of n integer intervals $[a_i, b_i]$ such that $0 \le a_i \le b_i \le n-1$, and is said to be graphic/realizable if there exists a graph with degree sequence, say, $D = (d_1, \ldots, d_n)$ satisfying the condition $a_i \le d_i \le b_i$, for each $i \in [1, n]$. There is a characterisation (also implying an O(n) verifying algorithm) known for realizability of interval-sequences, which is a generalization of the Erdös-Gallai characterisation for graphic sequences. However, given any realizable interval-sequence, there is no known algorithm for computing a corresponding graphic certificate in $o(n^2)$ time.

In this paper, we provide an $O(n \log n)$ time algorithm for computing a graphic sequence for any realizable interval sequence. In addition, when the interval sequence is non-realizable, we show how to find a graphic sequence having minimum deviation with respect to the given interval sequence, in the same time. Finally, we consider variants of the problem such as computing the most regular graphic sequence, and computing a minimum extension of a length p non-graphic sequence to a graphic one.

1 Introduction

The *Graph Realization problem* for a property P deals with the following existential question: Does there exist a graph that satisfies the property P? Its fundamental importance is apparent, ranging from better theoretical understanding, to network design questions (such as constructing networks with certain desirable connectivity properties). Some very basic, yet challenging, properties that have been considered in past are degree sequences [9, 18, 20], eccentricites [6, 24], connectivity and flow [16, 12, 10, 11].

One of the earliest classical problems studied in this domain is that of graphic sequences. A sequence of n positive integers, $D = (d_1, \ldots, d_n)$, is said to be graphic if there exists an n vertex graph G such that D is identical to the sequence of vertex degrees of G. The problem of realizing graphic sequences and counting the number of non-isomorphic realizations of a given graphic-sequence, is particularly of interest due to many practical applications, see [27] and reference therein. In 1960, Erdös and Gallai [9] gave a characterization (also implying an O(n) verifying algorithm) for graphic sequences. Havel and Hakimi [18, 20] gave a recursive algorithm that given a sequence D of integers computes a realizing graph, or proves that the sequence is non-graphic, in optimal time $O(\sum_i d_i)$. Recently, Tripathi et al. [28] provided a constructive proof of Erdös and Gallai's [9] characterization.

We consider a generalization of the graphic sequence problem where instead of specifying precise degrees, we are given a *range* (or interval) of possible degree values for each vertex. Formally, an *interval-sequence* is a sequence of n intervals $S = ([a_1, b_1], \ldots, [a_n, b_n])$, also represented as S = (A, B), where $A = (a_1, \ldots, a_n)$ and $B = (b_1, \ldots, b_n)$, and $0 \le a_i \le b_i \le n - 1$ for every *i*. It is said to be *realizable*

^{*}City University of New York (CUNY), USA. E-mail: amotz@sci.brooklyn.cuny.edu

[†]Tel Aviv University, Israel. E-mail: keerti.choudhary@cs.tau.ac.il

[‡]Weizmann Institute of Science, Israel. E-mail: david.peleg@weizmann.ac.il

[§]Bar Ilan University, Israel. E-mail: dror.rawitz@biu.ac.il

if there exists a sequence $D = (d_1, \ldots, d_n)$ that is graphic and satisfies the condition $a_i \leq d_i \leq b_i$, for $1 \leq i \leq n$. Two questions that are natural to ask here are:

Question 1 (Verification). *Find an efficient algorithm for verifying the realizability of any given intervalsequence* S?

Question 2 (Graphic Certificate). *Given a realizable interval-sequence S, compute a certificate (that is, a graphic sequence D) realizing it.*

Cai et al. [7] extended Erdös and Gallai's work by providing an easy to verify characterization for realizable interval-sequences, thereby resolving Question 1. Their result crucially uses the (g, f)-Factor Theorem of Lovász [25]. Garg et al. [15] provided a constructive proof of the characterisation of Cai et al. [7] for realizable interval sequences. In [22], Hell and Kirkpatrick provided an algorithm based on Havel and Hakimi's work for computing a graph that realizes an interval sequence (if exists). For non-realizable interval sequences S, their algorithm computes a graph whose deviation $\delta(D, S)$ (see Section 2 for definition) with respect to L1-norm is minimum. The time complexity of their algorithm is $O(\sum_{i=1}^{n} b_i)$ (which can be as high as $\Theta(n^2)$).

Our Contributions. In this paper we introduce a new approach for representing and analyzing the interval sequence realization problem. Our algorithms are based on a novel divide and conquer methodology, wherein we show that partitioning a realizable interval sequence along any levelled sequence (a new class of sequences introduced herein) guarantees that at least one of the new child interval sequences is also realizable. This enables us to present an $O(n \log n)$ time algorithm for computing a graphic certificate (if exists) for any given interval sequence. While the problem was well studied, to the best of our knowledge there was no known $o(n^2)$ time algorithm for computing graphic certificate. In addition, given an interval sequence S, our algorithm can obtain in the same time a degree-certificate corresponding to graphs with minimum (resp. maximum) possible edges. Specifically, we obtain the following result.

Theorem 1. There exists an algorithm that for any integer $n \ge 1$ and any length n interval sequence S, computes a graphic sequence D realizing S, if exists, in $O(n \log n)$ time.

Moreover, our algorithm can also output in the same time graphic certificates corresponding to a sparsest as well as densest possible graph (i.e. graphs with minimum and maximum possible edges), realizing S.

We also investigate the problem of efficiently computing graphic sequences having the least possible L_1 -deviation in the scenario when the input interval sequence is non-realizable. We must point out here that till now there was also no sub-quadratic time algorithm known for computing even the deviation $\delta(D, S)$. Our result for deviation minimizing certificate can be formalized as follows.

Theorem 2. There exists an algorithm that for any integer $n \ge 1$ and any length n non-realizable interval sequence S, outputs in $O(n \log n)$ time a graphic sequence D minimizing the deviation $\delta(D, S)$.

Our new approach enables us to tackle also an optimization version of the problem in which it is required to compute the "most regular" sequence realizing the given interval sequence S, using the natural measure of the minimum sum of pairwise degree differences, $\sum_{i,j} |d_i - d_j|$, as our regularity measure. To the best of our knowledge, this problem was not studied before and is not dealt with directly by the existing approaches to the interval sequence problem. Specifically, we obtain the following.

Theorem 3. There exists an algorithm that for any integer $n \ge 1$ and any length n realizable interval sequence S, computes the most regular graphic sequence realizing interval sequence S (i.e., the one minimizing the sum of pairwise degree difference), in time $O(n^2)$.

The tools developed in this paper allows us to study other interesting applications, such as computing a minimum extension of non-graphic sequences to graphic ones (see Section 6).

Related work. Kleitman and Wang [23], and Fulkerson-Chen-Anstee [2, 8, 14] solved the problem of degree realization for directed graphs, wherein, for each vertex both the in-degree and out-degree is specified. In [19], Nichterlein and Hartung proved the NP-completeness of the problem when the additional constraint of acyclicity is imposed. Over the years, various extensions of the degree realization problems were studied as well, cf. [1, 30]. The *Subgraph Realization problem* considers the restriction that the realizing graph must be a subgraph (*factor*) of some fixed input graph. For an interesting line of work on graph factors, refer to [29, 3, 21, 17]. The subgraph realization problems are generally harder. For instance, it is very easy to compute an *n*-vertex connected graph whose degree sequence consists of all values 2, however, the same problem for subgraph-realization is NP-hard (since it reduces to Hamiltonian-cycle problem).

Lesniak [24] provided a characterization for the sequence of eccentricities of an *n*-vertex graph. Behzad et al. [6] studied the problem of characterizing the set comprising of vertex-eccentricity values of general graphs (the sequence problem remains open). Fujishige et al. [13] considered the problem of realizing graphs and hypergraphs with given cut specifications. Realization problems related to various criteria of relative satisfaction are considered in [5]. Several other realization problems are surveyed in [4].

Organization of the Paper. In Section 2, we present the notation and definitions. In Section 3, we discuss the main ideas and tools that help us to construct graph certificates for interval sequence problem. Section 4 presents our $O(n \log n)$ time algorithm for computing graphic certificate with minimum deviation. Section 5 provides a quadratic-time algorithm for computing the most regular certificate. We discuss the applications in Section 6. The appendix includes linear time verification algorithms implied by the work of Erdös and Gallai [9], and Cai et al. [7].

2 Preliminaries

A sequence is defined to be an *n*-element vector whose entries are non-negative integers. For any sequence $D = (d_1, \ldots, d_n)$, define $\min(D) = \min_{i=1}^n \{d_i\}$, $\max(D) = \max_{i=1}^n \{d_i\}$, $\sup(D) = \sum_{i=1}^n d_i$, and parity $D = \operatorname{sum}(D) \mod 2$. Given any two sequences $X = (x_1, \ldots, x_n)$ and $Y = (y_1, \ldots, y_n)$, we say that $X \leq Y$ if $x_i \leq y_i$ for $1 \leq i \leq n$. Any two sequences X and Y are said to be similar if they are identical up to permutation of the elements (i.e., their sorted versions are identical). A sequence D is said to *lie* in an interval-sequence (A, B), denoted by $D \in (A, B)$, if $A \leq D \leq B$. We define $\min(X, Y) = (\min\{x_1, y_1\}, \ldots, \min\{x_n, y_n\})$, and $\max(X, Y) = (\max\{x_1, y_1\}, \ldots, \max\{x_n, y_n\})$. The L₁-distance of the pair (X, Y) is defined as L₁ $(X, Y) = \sum_{i=1}^n |y_i - x_i|$.

Denote by \top and \bot the *n*-length sequences all whose entries are respectively n - 1 and 0. Given a sequence $D = (d_1, \ldots, d_n)$ and an integer $k \in [1, n]$, define the vectors X(D) and Y(D) by setting for $1 \le k \le n$:

$$X_k(D) \triangleq \sum_{i=1}^k d_i$$
, and $Y_k(D) \triangleq k(k-1) + \sum_{i=k+1}^n \min(d_i, k)$

For any sequence $D = (d_1, \ldots, d_n)$, the *spread* of D is defined as $\phi(D) = \sum_{1 \le r < s \le n} |d_r - d_s|$, and it always lies in the range $[0, n^3]$. A sequence D is said to be more *regular* than another sequence D' if $\phi(D) < \phi(D')$. For any two integers $x \le y$, $[x, y] = \{x, x + 1, \ldots, y\}$. For any $I \subseteq [1, n]$, define D[I] to be the subsequence of D consisting of elements d_i , for $i \in I$; and define E_I to be the *characteristic vector* of I, namely, the sequence (e_1, e_2, \ldots, e_n) such that $e_i = 1$ if $i \in I$, and $e_i = 0$ otherwise. For any sequence $D = (d_1, \ldots, d_n)$ and an interval-sequence $S = ([a_1, b_1], \ldots, [a_n, b_n])$, the upper and lower deviation of D, is respectively defined as

$$\delta_U(D, S) = \sum_{i=1}^n \max\{0, (d_i - b_i)\}, \text{ and } \delta_L(D, S) = \sum_{i=1}^n \max\{0, (a_i - d_i)\}.$$

The deviation of D is defined as $\delta(D, S) = \delta_U(D, S) + \delta_L(D, S)$. For any vertex x in an undirected simple graph H, define $\deg_H(x)$ to be the degree of x in H, and define $N_H(x) = \{y \mid (x, y) \in E(H)\}$ to be the neighbourhood of x in H.

We next state the Erdös and Gallai [9] characterisation for realizable (graphic) sequences, and Cai et al. [7] characterisation for realizable interval sequences. An O(n)-time implementation of the both theorems is provided in the Appendix A.

Theorem 4 (Erdös and Gallai [9]). A non-increasing sequence $D = (d_1, \ldots, d_n)$ is graphic if and only if (i) $X_n(D)$ is even, and (ii) $X(D) \leq Y(D)$.

Theorem 5 (Cai et al. [7]). Let $S = ([a_1, b_1], \dots, [a_n, b_n]) = (A, B)$ be an interval-sequence such that A is non-increasing and for any index $1 \le i < n$, $b_{i+1} \le b_i$ whenever $a_i = a_{i+1}$. For each $k \in [1, n]$, define $W_k(S) = \{i \in [k+1, n] \mid b_i \ge k+1\}$. Then S is realizable if and only if $X(A) \le Y(B) - \varepsilon(S)$, where, $\varepsilon(S)$ is defined by setting

$$\varepsilon_k(\mathcal{S}) = \begin{cases} 1 & \text{if } a_i = b_i \text{ for } i \in W_k(\mathcal{S}) \text{ and } \sum_{i \in W_k(\mathcal{S})} (b_i + k|W_k(\mathcal{S})|) \text{ is odd,} \\ 0 & \text{otherwise.} \end{cases}$$

3 Main Tools

In this section, we develop some crucial tools that help us in efficient computation of certificate for a realizable interval-sequence. These tools will help us to search a graphic sequence in $O(n \log n)$ time using a clever divide and conquor methodology. Also they aid in searching for the maximally regular sequence in just quadratic time.

Levelling operation. Given a sequence $D = (d_1, \ldots, d_n)$ and a pair of indices $\alpha \neq \beta$ satisfying $d_{\alpha} > d_{\beta}$, we define $\pi(D, \alpha, \beta) = D^* = (d_1^*, \ldots, d_n^*)$ to be a sequence obtained from D by decrementing d_{α} by 1 and incrementing d_{β} by 1 (i.e., $d_{\alpha}^* = d_{\alpha} - 1$, $d_{\beta}^* = d_{\beta} + 1$, and $d_k^* = d_k$ for $k \neq \alpha, \beta$). This operation is called the *levelling operation* on D for the indices α and β . The operation essentially "levels" (or "flattens") the sequence D, making it more uniform.

We now discuss some properties of levelling operations.

Lemma 1. Any levelling operation on a sequence D that results in a non-similar sequence, reduces its spread $\phi(D)$ by a value at least two.

Proof. Let $D = (d_1, d_2, ..., d_n)$ and $Z = (z_1, ..., z_n) = \pi(D, \alpha, \beta)$, be a sequence obtained from D by performing a levelling operation on a pair of indices α, β such that $d_\alpha > d_\beta$. If $d_\alpha = d_\beta + 1$, then it is easy

to verify that D and Z are similar. If $d_{\alpha} \geq d_{\beta} + 2$, then

$$\begin{split} \phi(Z) &= |z_{\alpha} - z_{\beta}| + \sum_{s \neq \alpha, \beta} (|z_{\alpha} - z_{s}| + |z_{\beta} - z_{s}|) + \sum_{\substack{1 \le r < s \le n, \\ r, s \notin \{\alpha, \beta\}}} |z_{r} - z_{s}| \\ &= |d_{\alpha} - d_{\beta}| - 2 + \sum_{\substack{s \neq \alpha, \beta \text{ s.t.} \\ d_{s} \notin (d_{\beta}, d_{\alpha})}} (|d_{\alpha} - d_{s}| + |d_{\beta} - d_{s}|) \\ &+ \sum_{\substack{s \neq \alpha, \\ \beta \text{ s.t.} d_{s} \in (d_{\beta}, d_{\alpha})}} (|d_{\alpha} - d_{s}| + |d_{\beta} - d_{s}| - 2) + \sum_{\substack{1 \le r < s \le n, \\ r, s \notin \{\alpha, \beta\}}} |d_{r} - d_{s}| \\ &\leq \left(\sum_{1 \le r < s \le n} |d_{r} - d_{s}|\right) - 2 = \phi(D) - 2 \,. \end{split}$$

Thus, the claim follows.

Lemma 2 (Corollary 3.1.4, [26]). *The levelling operations preserves graphicity, that is, if we perform a levelling operation on a graphic sequence, then the resulting sequence is also graphic.*

Proof. Let $D = (d_1, \ldots, d_n)$ be a graphic sequence, and $\pi(D, \alpha, \beta) = D^* = (d_1^*, \ldots, d_n^*)$ for some indices α, β satisfying $d_\alpha > d_\beta$. If $d_\alpha = 1 + d_\beta$, then D^* is similar to D, and thus also graphic. So for the rest the proof let us focus on the case $d_\alpha \ge 2 + d_\beta$. Let G = (V, E) be a graph realising the sequence D, and let x_α and x_β be two vertices in G having degrees respectively d_α and d_β . Since $|N_G(x_\alpha)| \ge 2 + |N_G(x_\beta)|$, there must exists at least one neighbour, say w, of vertex x_α that does not lie in set $\{x_\beta\} \cup N_G(x_\beta)$. Let $G^* = (V, E^*)$ be a graph obtained from G by deleting the edge (w, x_α) , and adding a new edge (w, x_β) . Observe that the degree of all vertices other than x_α and x_β are identical in graphs G and G^* , also $\deg_{G^*}(x_\alpha) = \deg_G(x_\alpha) - 1$, and $\deg_{G^*}(x_\beta) = \deg_G(x_\beta) + 1$. Therefore G^* is a graph realising the profile D^* , and thus the claim follows.

Levelled sequences. A sequence D is said to be *levelled with respect to the integer-sequence* S = (A, B) if (i) $A \leq D \leq B$, and (ii) the spread of D cannot be decreased by a levelling operation, i.e., for any two indices $\alpha \neq \beta$ satisfying $d_{\alpha} > d_{\beta}$ and $A \leq \pi(D, \alpha, \beta) \leq B$, we have $\phi(\pi(D, \alpha, \beta)) = \phi(D)$. See Figure 1.

The volume of a sequence D lying between A and B with respect to $\mathcal{S} = (A, B)$ is defined as

$$\operatorname{VOL}(D, \mathcal{S}) \triangleq \mathbf{L}_1(D, A)$$
,

and is invariant of levelling operations applied to D. In other words, applying a levelling operation to a sequence D may reduce its spread but preserves its volume. Note that the volume lies in the range $[0, \mathbf{L}_1(A, B)]$.

Lemma 3. For any S = (A, B), a sequence D satisfying $A \le D \le B$ can be transformed into a levelled sequence D^* having the same volume VOL(D, S) by a repeated application of (at most $O(n^3)$) levelling operations.¹

¹We remark that the algorithms presented later on generate a desired levelled sequence using more efficient methods than the one implicit in the proof, and are therefore faster.

Proof. By Lemma 1, every levelling operation that results in a new (non-similar) sequence decreases the spread by at least two. Since the spread of any sequence D is always non-negative and finite (specifically, $O(n^3)$), it is possible to perform $(O(n^3))$ levelling operations on D so that the resultant sequence D^* is levelled. Since the levelling operation preserves the volume, $VOL(D^*, S)$ must be same as VOL(D, S). \Box

Any graphic sequence D realizing the interval sequence S = (A, B) by Lemma 3 can be altered by $O(n^3)$ levelling operations to obtain a levelled sequence lying between A and B. The resultant sequence by Lemma 2 remains graphic, thus the following theorem is immediate.

Theorem 6. For any realizable interval sequence S = (A, B) there exists a graphic sequence realizing S which is a levelled sequence.

Characterizing and Computing Levelled sequences. Given any interval sequence S = (A, B) and a real number $\ell \in [\min(A), \max(B)]$, let²

$$F(\ell, \mathcal{S}) \triangleq \sum_{i \in [1,n]} (\min\{\ell, b_i\} - \min\{\ell, a_i\}) .$$

Observe that $F(\cdot, S)$ is a non-decreasing function in the range $(\min(A), \max(B))$. Hence we may define the corresponding inverse function as $F^{-1}(L, S) = \min\{\ell \mid F(\ell, S) = L\}$.

Given any interval sequence S = (A, B), we define $I(\ell, S) \triangleq \{i \in [1, n] \mid a_i < \ell < b_i\}$.

We conclude this section by providing the following theorems for characterising and computing levelled sequences.

Theorem 7. Consider an interval sequence S = (A, B). Let L be an integer in $[0, \mathbf{L}_1(A, B)]$ and $\ell \ge 0$ be such that $\ell = F^{-1}(L, S)$. Then the collection of levelled sequences that have volume L with respect to S is equal to the collection of sequences $D = (d_1, \ldots, d_n)$ satisfying the following three conditions: (a) $d_i = b_i$ for any i satisfying $b_i \le \ell$;

- (a) $a_i = b_i$ for any i satisfying $b_i \leq c$,
- (b) $d_i = a_i$ for any i satisfying $a_i \ge \ell$; and
- (c) Among all indices lying in set $I(\ell, S)$, exactly $F(\ell, S) F(\lfloor \ell \rfloor, S)$ indices *i* satisfy $d_i = \lceil \ell \rceil$, and the remaining indices *i* satisfy $d_i = \lfloor \ell \rfloor$.

Proof. We first make some observations about levelled sequences. Let $D = (d_1, \ldots, d_n)$ be a levelled sequence with respect to S = (A, B) having volume L, then

- (i) For any *i* with $d_i > \lceil \ell \rceil$, $d_i = a_i$.
- (ii) For any *i* with $d_i < \lfloor \ell \rfloor$, $d_i = b_i$.

To prove claim (i), suppose, to the contrary, that there exists an index i_0 such that $d_{i_0} > \max\{\lceil \ell \rceil, a_{i_0}\}$. If there exists some $j_0 \neq i_0$ such that $d_{j_0} < \min\{\lceil \ell \rceil, b_{j_0}\}$. Then one can perform a levelling operation on the pair (i_0, j_0) , to obtain a new sequence D^* that lies between A and B, and has spread ϕ strictly less than that of sequence D. This contradicts the fact that D was levelled. So let us assume $d_j \ge \min\{\lceil \ell \rceil, b_j\}$ for every $j \neq i_0$. Then

$$d_j - a_j \geq \min\{\lceil \ell \rceil, b_j\} - a_j \text{ for each } j \neq i_0$$
.

For i_0 we have

$$d_{i_0} - a_{i_0} > \max\{\lceil \ell \rceil, a_{i_0}\} - a_{i_0} \ge \lceil \ell \rceil - a_{i_0} \ge \min\{\lceil \ell \rceil, b_{i_0}\} - a_{i_0}$$

²One can think of S as representing a collection of n connected vessels, each in the shape of a unit column closed at both ends, then $F(\ell, S)$ is the amount of fluid that will fill this connected vessel system to level ℓ .



Figure 1: Illustration of a levelled sequence D (in red) satisfying L = VOL(D, S) = 33. For $\ell = 7.5$, $F(\ell = 7.5, S) = 33$, $F(\lfloor \ell \rfloor = 7, S) = 30$, and $F(\lceil \ell \rceil = 8, S) = 36$. The segments contributing to $F(\ell = 7.5, S)$, i.e., the parts of the connected vessel system filled with fluid, are shown in blue. The values in D at all indices in set $I(\ell, S)$ differ by at most one as they lie in the set $\{\lfloor \ell \rfloor, \lceil \ell \rceil\}$.

Combining above two inequalities, we get

$$F(\ell, \mathcal{S}) = L = \sum_{1 \le i \le n} (d_i - a_i) > \sum_{\substack{1 \le i \le n, \\ a_i \le \lceil \ell \rceil}} (\min\{\lceil \ell \rceil, b_i\} - a_i) = F(\lceil \ell \rceil, \mathcal{S}),$$

where the inequality follows from the fact that $d_{i_0} - a_{i_0} > 0$. (This is because if $a_{i_0} < \lceil \ell \rceil$, then index i_0 contributes a positive value of $d_{i_0} - a_{i_0}$ to the first sum and zero to the second.) This violates the fact that F is non-decreasing. The proof of claim (ii) follows in similar manner as that of claim (i), and is thus omitted.

We now show that each levelled sequence D w.r.t. S satisfies conditions (a), (b) and (c). We show that for any *i* satisfying $b_i \leq \ell$ (or equivalently, $b_i \leq \lfloor \ell \rfloor$), $d_i = b_i$. To prove this, let us assume to the contrary that $d_i < b_i$. Then $d_i < \lfloor \ell \rfloor$, which implies $d_i = b_i$, violating our assumption $d_i < b_i$. Similarly, it follows that for any *i* satisfying $a_i \geq \ell$ (or equivalently, $a_i \geq \lceil \ell \rceil$), $d_i = a_i$.

Next for any $i \in I(\ell, S)$, we have $d_i \in \{\lfloor \ell \rfloor, \lceil \ell \rceil\}$. Indeed, if $d_i < \lfloor \ell \rfloor$, then $b_i = d_i < \ell$, violating the fact $\ell < b_i$. Similarly, if $d_i > \lceil \ell \rceil$, then $a_i = d_i > \ell$, violating the fact $a_i < \ell$. Since the degrees are integral, the indices in $I(\ell, S)$ are partitioned into $L - F(\lfloor \ell \rfloor, S)$ indices with degree $d_i = \lceil \ell \rceil$, and $|I(\ell, S)| - (L - F(\lfloor \ell \rfloor, S))$ indices i with degree $d_i = \lfloor \ell \rfloor$.

To prove the converse, note that all sequences satisfying conditions (a), (b) and (c) in the theorem are similar and have volume L.

Theorem 8. Given an interval sequence S = (A, B) consisting of *n*-pairs, and an integer $L \in [0, \mathbf{L}_1(A, B)]$, a levelled sequence D having volume L with respect to S can be computed in O(n) time.

Algorithm 1: LEVELLED-SEQUENCE(A, B, L)

1 if $L \notin [0, \mathbf{L}_1(A, B)]$ then Return Null; 2 $A', B' \leftarrow 0;$ 3 for i = 1 to n do $A'[a_i] \leftarrow A'[a_i] + 1;$ 4 $B'[b_i] \leftarrow B'[b_i] + 1;$ 5 6 for k = 1 to n do $Q[k] \leftarrow Q[k-1] + A'[k-1] - B'[k-1];$ 7 $k \leftarrow 1, L' \leftarrow L;$ 8 while $(Q[k] \leq L')$ do $L' \leftarrow L' - Q[k]$ and $k \leftarrow k + 1$; 9 $\ell \leftarrow k - 1 + L'/Q[k];$ 10 for i = 1 to n do if $(a_i \ge \ell)$ then $d_i = a_i$; 11 if $(b_i \leq \ell)$ then $d_i = b_i$; 12 if $(a_i < \ell < b_i)$ then 13 if L' > 0 then $d_i = \lceil \ell \rceil$ and $L' \leftarrow L' - 1$; 14 else $d_i = |\ell|;$ 15 16 Return D;

Proof. For any $k \in [1, n]$, let Q[k] denote the size of the set $\{i \mid a_i < k \leq b_i\}$. In order to efficiently compute Q, we define two *n*-sequences A' and B', where A'[k] and B'[k] count the number of intervals i such that $a_i = k$ and $b_i = k$, resp. That is, $A'[k] = |\{i : a_i = k\}|$ and $B'[k] = |\{i : b_i = k\}|$. Observe that Q[k] - Q[k-1] = A'[k-1] - B'[k-1], for every $k \in [1, n]$.

Notice for any integer ℓ , $F(\ell, S) = \sum_{k \leq \ell} Q[k]$. Let $\ell' = \max\{k \in [0, n] \mid \sum_{i \leq k} Q[i] \leq L\}$, so that, ℓ' is the maximum integer for which $F(\ell', S) \leq L$. Also let

$$\ell = \ell' + \frac{L - \sum_{i \le \ell'} Q[i]}{Q[\ell' + 1]} \,.$$

It is not hard to verify that $\ell = F^{-1}(L, S)$. Thus a levelled sequence of volume L can be computed in O(n) time using Theorem 7. In Algorithm 1 we present the pseudocode of our implementation that can be seen as filling the vessels with fluid until reaching the desired level ℓ .

The correctness of Algorithm 1 follows from above description, also it is easy to verify that the total run-time of the algorithm is O(n).

4 An $O(n \log n)$ time algorithm for Graphic Certificate

In this section, we present an algorithm for computing a certificate for interval sequence that takes just $O(n \log n)$ time. If the input interval S = (A, B) is realizable, our algorithm computes a graphic sequence $D \in S$, otherwise it computes a sequence minimizing the deviation value $\delta(D, S)$. We begin by considering the case where the sequence S is realizable (since it is simpler to understand given Theorems 7 and 8), and then we move to the case where S is non-realizable. Then characterization of [7] implies an O(n) time verification algorithm for realizability of interval sequence. (For details refer to the Appendix).

4.1 Realizable Interval Sequences

First we show that any two levelled sequences after an appropriate reordering of their elements are coordinatewise comparable.

Lemma 4. For any interval sequence S = (A, B), and any two levelled sequences $C, D \in S$ satisfying $VOL(D, S) \leq VOL(C, S)$, the following holds.

- 1. $D' \leq C$, for some sequence $D' \in S$ similar to D.
- 2. $D \leq C''$, for some sequence $C'' \in S$ similar to C.

Proof. We show how to transform $D = (d_1, \dots, d_n)$ into sequence $D' = (d'_1, \dots, d'_n) \in S$ such that $D' \leq C$. Let $\ell_D = F^{-1}(\text{VOL}(D, S), S)$ and $\ell_C = F^{-1}(\text{VOL}(C, S), S)$. Since $F(\cdot, S)$ is a non-decreasing function, we have that $\ell_D \leq \ell_C$.

Let us first consider the case where ℓ_C and ℓ_D are both non-integral, and $\lfloor \ell_C \rfloor = \lfloor \ell_D \rfloor = (\operatorname{say} \ell_1)$ and $\lceil \ell_C \rceil = \lceil \ell_D \rceil = (\operatorname{say} \ell_2)$. By Theorem 7, for any index $i \in [1, n]$, (i) $a_i \geq \ell_D$ (or $a_i \geq \ell_C$) implies $d_i = a_i = c_i$; (ii) $b_i \leq \ell_D$ (or $b_i \leq \ell_C$) implies $d_i = b_i = c_i$. Also, among indices in set $I_0 = I(\ell_D, S) =$ $I(\ell_C, S)$, (i) exactly $L_D - F(\lfloor \ell_D \rfloor, S)$ indices i satisfy $d_i = \ell_2$ (let I_D denote the set of these indices) and the remaining indices i satisfy $d_i = \ell_1$; (ii) exactly $L_C - F(\lfloor \ell_C \rfloor, S)$ indices i satisfy $c_i = \ell_2$ (let I_C denote the set of these indices) and the remaining indices i satisfy $c_i = \ell_1$. Since $L_D \leq L_C$, it follows that $|I_D| \leq |I_C|$, however, observe that I_D need not be a subset of I_C . We set D' to be the sequence that satisfy the condition that (i) $d'_i = d_i$, for each $i \notin I_0$, and (ii) for indices in I_0 , at any arbitrary $|I_D|$ indices lying in I_C, d'_i take the value ℓ_2 , and at remaining $|I_0| - |I_D|$ indices d'_i take the value ℓ_1 . It is easy to verify that Dand D' are similar, and $D' \leq C$.

The remaining case is when $\lceil \ell_D \rceil \leq \lfloor \ell_C \rfloor$. For any index $i \in [1, n]$, $d_i \leq \lceil \ell_D \rceil$ and $c_i \geq \lfloor \ell_C \rfloor$, implies $d_i \leq c_i$. Observe that by Theorem 7, (i) for an index i, $d_i > \lceil \ell_D \rceil$ implies $d_i = a_i (\leq c_i)$; and (ii) for an index i, $c_i < \lfloor \ell_C \rfloor$ implies $c_i = b_i (\geq d_i)$. Therefore, for each index i, $d_i \leq c_i$. So in this case, we set D' to be D. The construction of sequence C'' follows similarly.

Next lemma shows significance of partitioning an interval-sequence using a levelled sequence.

Lemma 5. Let C and D be any two levelled sequences lying in an interval sequence S = (A, B), and having volume L_C and L_D , respectively. Also assume D is a graphic sequence. Then, (a) $L_D \leq L_C$ implies (A, C) is a realizable interval sequence. (b) $L_D \geq L_C$ implies (C, B) is a realizable interval sequence.

Proof. We provide proof of the case $L_D \leq L_C$ (the proof of part (b) will follow in a similar fashion). By Lemma 4, we can transform $D = (d_1, \dots, d_n)$ into another levelled sequence $D' = (d'_1, \dots, d'_n) \in S$ such that D' is similar to D and $D' \leq C$. Since $D' \leq C$, and D' is a graphic sequence, it follows that (A, C) is realizable interval sequence.

From Lemma 5, and the fact that each realizable interval-sequence contains a levelled graphic sequence (see Theorem 6), we obtain following.

Theorem 9. For any realizable interval sequence S = (A, B), and any levelled sequence $C \in S$, at least one of the interval-sequences (A, C) and (C, B) is realizable.

The above theorem provides a divide-and-conquer strategy to search for a levelled graphic sequence for realizable interval-sequences as shown in Algorithm 2. Let (A_0, B_0) be initialized to (A, B). We compute a levelled sequence C_0 having volume $\lfloor \mathbf{L}_1(A_0, B_0)/2 \rfloor$ using Theorem 7. It follows from Theorem 9, either (A_0, C_0) or (C_0, B_0) must be a realizable interval-sequence. If (A_0, C_0) is realizable then we replace

 B_0 by C_0 ; otherwise (C_0, B_0) must be realizable, so we replace A_0 by C_0 . We continue this process (of replacements) until $\mathbf{L}_1(A_0, B_0)$ decreases to a value smaller than 2. In the end, the interval sequence (A_0, B_0) contains at most two sequences, namely A_0 and B_0 . If A_0 is graphic then we return A_0 , otherwise we return B_0 . The correctness of the algorithm is immediate from the description. It is also easy to verify that the algorithms outputs a graphic-certificate with the least possible number of edges.

Algorithm 2: CERTIFICATE-REALIZABLE(*A*, *B*)

1 Initialize interval sequence (A_0, B_0) to (A, B);

2 while $L_1(A_0, B_0) \ge 2$ do

- 3 | $C_0 \leftarrow$ a levelled sequence of volume $\lfloor \mathbf{L}_1(A_0, B_0)/2 \rfloor$;
- 4 **if** (Interval-sequence (A_0, C_0) is realizable) then $B_0 \leftarrow C_0$;
- 5 else $A_0 \leftarrow C_0$;
- 6 if A_0 is graphic then Return A_0 ;
- 7 else Return B_0 ;

To analyze the running time, observe that the L_1 -distance between A_0 and B_0 decreases by (roughly) a factor of 2 in each call of the while loop, so it follows that number of iterations is $O(\log n)$. Verifying if an interval sequence is realizable, or a sequence D is graphic can be performed in O(n) time, using Theorem 11. Also in O(n) time we can generate a levelled sequence of any given volume L by Theorem 8. Thus, the total time complexity of the algorithm is $O(n \log n)$.

We obtain the following result:

Theorem 10. There exists an algorithm that for any integer $n \ge 1$ and any n-length interval sequence S = (A, B), computes a graphic sequence $D \in (A, B)$, if it exists, in $O(n \log n)$ time.

*Moreover, our algorithm can also output graphic certificates corresponding to a sparsest and densest*³ *possible graph (i.e. having minimum and maximum possible edges), realizing S, in the same time.*

4.2 Non-Realizable Sequences

In this subsection we consider the scenario where S is non-realizable, our goal is to compute a graphic sequence D minimizing the deviation $\delta(D, S)$ with respect to the given interval sequence S.

As a first step, we show that in order to search a sequence D minimizing $\delta(D, S)$, it suffices to search a sequence $D \ge A$ that minimizes the value $\delta_U(D, S)$.

Lemma 6. $\min\{\delta(D, S) \mid D \text{ is graphic }\} = \min\{\delta_U(D, S) \mid D \text{ is graphic, } D \geq A\}$, for any interval sequence S = (A, B).

Proof. Let $D = (d_1, \ldots, d_n)$ be a graphic sequence minimizing the value $\delta(D, S)$, and in case of ties take that D for which $\delta_L(D, S)$ is the lowest. Let us suppose there exists an index $i \in [1, n]$ such that $d_i < a_i$. Consider the graph G realizing the sequence D, and let v_i denote the *i*th vertex of G, so that, $\deg(v_i) = d_i$. Observe that $|N_G(v_i)| \neq n - 1$, since $d_i < a_i \leq n - 1$. For any vertex $v_j \notin N_G(v_i)$, $d_j = \deg(v_j)$ must be at least b_j , because otherwise adding (v_i, v_j) to G reduces $\delta(D, S)$. Thus for any vertex $v_j \notin N_G(v_i)$, adding (v_i, v_j) to G, decreases $\delta_L(D, S)$ and increases $\delta_U(D, S)$ by a value exactly 1. However, by our choice D was a sequence minimizing $\delta_L(D, S)$, thus $\delta_L(D, S)$ must be zero. The claim follows from the fact that $D \geq A$ and $\delta(D, S) = \delta_U(D, S)$.

³Note that the sparsest (resp. densest) possible graph realizing an interval sequence S need not be unique.

By the previous lemma, our goal is to find a graphic sequence D in the interval sequence (A, \top) minimizing $\delta(D, S)$. Notice that if D is graphic, then the interval sequence (A, R), where $R = \max(D, B)$, is realizable. Also, $\delta(D, S) = \sup(R - B)$. Hence, in order to compute a graphic sequence with minimum deviation, we define \mathcal{R} to be the set of all sequence $R \in [B, \top]$ such that (i) the interval sequence (A, R) is realizable, and (ii) $\sup(R - B)$ is minimized.

The following lemma shows the significance of the set \mathcal{R} in computing a certificate with minimum deviation.

Lemma 7. For any $R \in \mathcal{R}$, and any graphic sequence D_0 lying in the interval sequence (A, R), we have $\delta(D_0, S) = \min{\{\delta(D, S) \mid D \text{ is graphic }\}} = \sup(R - B).$

Proof. Let D^* be a graphic sequence minimizing the value $\delta(D, S)$. By Lemma 6, we may assume that D^* belongs to (A, \top) . Observe that $\delta(D^*, S) = \operatorname{sum}(R^* - B)$, where $R^* = \max\{B, D^*\}$. By the choice of D^* we have that $\operatorname{sum}(R^* - B) = \delta(D^*, S) \leq \delta(D_0, S) = \delta_U(D_0, S) \leq \operatorname{sum}(R - B)$, where the last inequality follows from the fact that $D_0 \in (A, R)$. By definition of \mathcal{R} , we have that $\operatorname{sum}(R^* - B) \geq \operatorname{sum}(R - B)$, and therefore $\delta(D^*, S) = \delta(D_0, S) = \operatorname{sum}(R - B) = \operatorname{sum}(R^* - B)$. Thus R^* also lies in the set \mathcal{R} . The lemma follows from the fact that $\delta(D^*, S) = \min\{\delta(D, S) \mid D \text{ is graphic }\}$.

Next, let \mathcal{R}_L be the set of all levelled sequences in \mathcal{R} with respect to interval sequence (B, \top) .

Lemma 8. $\mathcal{R}_L \neq \emptyset$.

Proof. Clearly, $\mathcal{R} \neq \emptyset$. Consider any sequence $R = (r_1, \ldots, r_n) \in \mathcal{R}$. Suppose there exists $\alpha, \beta \in [1, n]$ such that $r_\alpha - r_\beta \ge 1$ and $R' = \pi(R, \alpha, \beta) \in [B, \top]$. Observe that $\operatorname{sum}(R' - B) = \operatorname{sum}(R - B)$. It remains to show that (A, R') is realizable. Indeed, if $D \in (A, R)$ is a graphic sequence, then either (i) $D = (d_1, \ldots, d_n)$ lies in (A, R'), or (ii) $d_\alpha - d_\beta \ge 1$ and $D' = \pi(D, \alpha, \beta)$ lies in (A, R'). Since levelling operation preserves graphicity, D' is graphic. Thus $R' \in \mathcal{R}$, which shows that \mathcal{R} is closed under the levelling operation, and hence \mathcal{R}_L is non-empty.

Algorithm 3: CERTIFICATE-NON-REALIZABLE(*A*, *B*)

 $\begin{array}{ll} \mathbf{1} & (M_1, M_2) \leftarrow (B, \top); \\ \mathbf{2} & \mathbf{while} \ \mathbf{L_1}(M_1, M_2) \geq 2 \ \mathbf{do} \\ \mathbf{3} & \begin{bmatrix} M_0 \leftarrow a \ \text{levelled sequence of volume} \ \lfloor \mathbf{L_1}(M_1, M_2)/2 \rfloor; \\ \mathbf{4} & \begin{bmatrix} M_0 \leftarrow a \ \text{levelled sequence} \ (A, M_0) \ \text{is realizable} \ \mathbf{then} \ M_2 \leftarrow M_0; \\ \mathbf{5} & \begin{bmatrix} e \ \text{lse} \ M_1 \leftarrow M_0; \\ \mathbf{6} \ \text{if} \ (A, M_1) \ \text{is realizable then} \ R \leftarrow M_1; \\ \mathbf{7} \ \text{else} \ R \leftarrow M_2; \\ \mathbf{8} \ \text{Return CERTIFICATE-REALIZABLE}(A, R) \\ \end{array}$

We now describe the algorithm for computing a graphic sequence with minimum deviation (refer to Algorithm 3 for a pseudocode). Recall that we assume that (A, B) is a non-realizable interval sequence. The first step is to compute a levelled sequence $R \in \mathcal{R}_L$, and the second is to use Algorithm 2 to find a graphic sequence in (A, R).

We initialize two sequences M_1 and M_2 , resp., to B and \top , and these sequences serve as lower and upper boundaries for sequence R. The pair (M_1, M_2) is updated as long as $sum(M_2 - M_1) \ge 2$ as follows. We compute a levelled sequence M_0 having volume $\lfloor \mathbf{L}_1(M_1, M_2)/2 \rfloor$ with respect to the interval sequence (M_1, M_2) using Theorem 8. There are two cases: **Case 1.** (A, M_0) is realizable.

Consider any sequence $R \in (M_1, M_2)$ that lies in \mathcal{R}_L . Since (A, M_0) is realizable, from the definition of R it follows that sum $(R - B) \leq \text{sum}(M_0 - B)$. As R and M_0 both belong to (M_1, M_2) , by Lemma 4, there exists a sequence R_0 similar to R lying in interval $(M_1, M_2) \subseteq (B, \top)$ such that $R_0 \leq M_0$. It is easy to check that $R_0 \in \mathcal{R}_L$, thus the search range of R which was (M_1, M_2) can be narrowed down to (M_1, M_0) , so we reset M_2 to M_0 .

Case 2. (A, M_0) is not realizable.

Consider any $R \in \mathcal{R}_L$, we first show that $sum(R - B) > sum(M_0 - B)$. Let us assume on the contrary, $sum(R - B) \le sum(M_0 - B)$. In such a case, by Lemma 4, there exists a sequence R' similar to R lying in interval $(M_1, M_2) \subseteq (B, \top)$ such that $R' \le M_0$. Also $R' \in \mathcal{R}_L$. Since, by definition of \mathcal{R}_L , (A, R') is realizable, it violates the fact that (A, M_0) is not realizable. Now as R, M_0 both belong to (M_1, M_2) , by Lemma 4, there exists a sequence R_0 similar to R lying in interval (M_1, M_2) , by Lemma 4, there exists a sequence R_0 similar to R lying in interval $(M_1, M_2) \subseteq (B, \top)$ such that $R_0 \ge M_0$. Also $R_0 \in \mathcal{R}_L$, thus the search range of R can be narrowed down to (M_0, M_2) , so we reset M_1 to M_0 .

We continue the process of shrinking the range (M_1, M_2) until $\mathbf{L}_1(M_1, M_2)$ decreases to a value smaller than 2. Finally there exists in range (M_1, M_2) at most two sequences, namely M_1 and M_2 . If (A, M_1) is graphic then we set R to M_1 , otherwise we set R to M_2 .

The running time analysis is similar to the one for Algorithm 2. Since the L_1 -distance between M_1 and M_2 decreases by a factor of 2 in each successive call of the while loop of the algorithm, it follows that number of times the while loops run is $O(\log n)$. Verifying if an interval sequence is realizable, or a sequence D is graphic can be performed in O(n) time, using Theorem 11. Also it takes O(n) time to generate a levelled sequence of any given volume L by Theorem 8. Finally, the running time of Algorithm 2 is $O(n \log n)$. Thus, the total time complexity of algorithm is $O(n \log n)$.

This completes the proof of Theorem 1.

5 Most Regular Certificate in $O(n^2)$ time

In this section, we present an $O(n^2)$ -time algorithm for computing a most-regular certificate with respect to a given interval sequence S = (A, B). We assume that S is realizable. Our algorithm involves a subroutine that given an integer $z \in [\min(A), \max(B) - 1]$, computes a most-regular graphic-sequence, say D, satisfying the condition $z \leq \ell = F^{-1}(\operatorname{VOL}(D, S), S) \leq z + 1$. The following lemma is immediate from Theorem 7.

Lemma 9. Any levelled sequence $\overline{D} = (\overline{d}_1, \ldots, \overline{d}_n)$ of volume L with respect to interval sequence S = (A, B), satisfies $z \leq \ell = F^{-1}(L, S) \leq z + 1$ if and only if $\overline{d}_i = a_i$ for $a_i \geq z + 1$, $\overline{d}_i = b_i$ for $b_i \leq z$, and $\overline{d}_i \in \{z, z + 1\}$ for remaining indices i.

We partition the set [1, n] into three sets I_1, I_2, I_3 such that $I_1 = \{i \in [1, n] \mid a_i \ge z + 1\}, I_2 = \{i \in [1, n] \mid a_i \le z \text{ and } z + 1 \le b_i\}$, and $I_3 = \{i \in [1, n] \mid b_i \le z\}$. Also, using integer sort in linear time, we rearrange the pairs in (A, B) along with the corresponding sets I_1, I_2, I_3 so that (i) for any $i \in I_1, j \in I_2, k \in I_3$, we have i < j < k, and (ii) the sub-sequences $A[I_1]$ and $B[I_3]$ are sorted in the non-increasing order.

We initialize $D_z = (d_{z,1}, d_{z,2}, \dots, d_{z,n})$ by setting $d_{z,i}$ to : a_i if $i \in I_1$, z if $i \in I_2$, and b_i if $i \in I_3$. The sequence D_z is sorted in non-increasing order, since the sub-sequences $A[I_1]$ and $B[I_3]$ are sorted in non-increasing order. Let $\alpha = |I_1|$ and $\beta = |I_1| + |I_2|$, so that $I_2 = [\alpha + 1, \alpha + 2, \dots, \beta]$. We would search all those indices $i \in [\alpha, \beta]$ such that on incrementing $d_{\alpha+1}, \ldots, d_i$ to value z + 1, the resulting sequence is graphic; or equivalently, the sequence $D_z + E_{[\alpha+1,i]}$ is graphic. Note that for any index $i \in [\alpha, \beta]$, (i) the sequence $D_z + E_{[\alpha+1,i]}$ is non-increasing, and (ii) $A \leq D_z + E_{[\alpha+1,i]} \leq B$. The next lemma, which follows from the definition of ϕ , will be used to compute $\phi(D_z + E_{[\alpha+1,i]})$ from $\phi(D_z)$.

Lemma 10. For any index $i \in [\alpha + 1, \beta]$, $\phi(D_z + E_{[\alpha+1,i]}) = \phi(D_z) + (i - \alpha)(n - i - \alpha)$.

Proof. Let $D_z^i = D_z + E_{[\alpha+1,i]}$, for $i \in [\alpha+1,\beta]$.

$$\begin{split} \phi(D_z^i) &= \sum_{j=1}^{n-1} \sum_{k=j+1}^n \left| d_{z,j}^i - d_{z,k}^i \right| \\ &= \sum_{j \leq \alpha} \left(\sum_{k>j, k \not\in [\alpha+1,i]} |d_{z,j} - d_{z,k}| + \sum_{k \in [\alpha+1,i]} |d_{z,j} - d_{z,k} - 1| \right) + \\ &\sum_{j \in [\alpha+1,i]} \left(\sum_{k \in [j+1,i]} |d_{z,j} - d_{z,k}| + \sum_{k>i} |d_{z,j} + 1 - d_{z,k}| + \right) + \\ &\sum_{j>i} \sum_{k>j} |d_{z,j} - d_{z,k}| \\ &= \sum_{j=1}^{n-1} \sum_{k=j+1}^n |d_{z,j} - d_{z,k}| - \alpha \cdot (i - \alpha) + (i - \alpha) \cdot (n - i) \\ &= \phi(D_z) + (i - \alpha) \cdot (n - i - \alpha) \;, \end{split}$$

and the lemma follows.

For each z we compute the vectors $X(D_z)$ and $Y(D_z)$ using Theorem 11 in the Appendix. For each integer $k \in [1, n]$, let

AVOID
$$(k) = \left\{ i \in [\alpha, \beta] \mid X_k \left(D_z + E_{[\alpha+1,i]} \right) > Y_k \left(D_z + E_{[\alpha+1,i]} \right) \right\}$$
, and
AVOID $= \bigcup_{k=1}^n \operatorname{AVOID}(k)$.

Observe that by Theorem 4, for any $i \in [\alpha, \beta]$, the sequence $D_z + E_{[\alpha+1,i]}$ is graphic if and only if *i* does not lie in the set AVOID, and $\text{parity}(D_z + E_{[\alpha+1,i]}) = 0$. The following lemma shows that the set AVOID(k), for any index k, is computable in O(1) time.

Lemma 11. For each $k \in [1, n]$, AVOID(k) is a contiguous sub-interval of [1, n], and is computable in O(1) time.

Proof. For each $k \in [1, n]$, let

$$\begin{aligned} &\text{AVOID}_{1}(k) = [\alpha, \beta] \cap [\alpha + 1 + Y_{k}(D_{z}) - X_{k}(D_{z}), k] \\ &\text{AVOID}_{2}(k) = [\alpha, \beta] \cap \begin{cases} [k, n] & k \leq z \text{ and } \max\{k - \alpha, 0\} + X_{k}(D_{z}) > Y_{k}(D_{z}); \\ [k, \max\{k, \alpha\} + X_{k}(D_{z}) - Y_{k}(D_{z}) - 1] & k \geq z + 1; \\ \emptyset & \max\{k - \alpha, 0\} + X_{k}(D_{z}) \leq Y_{k}(D_{z}). \end{aligned}$$

We fix an index $k \in [1, n]$ for the rest of the proof. Our goal will be to show that the set AVOID(k) is the union of the interval AVOID₁(k) and AVOID₂(k), and is thus computable in O(1) time. Consider any $i \in [\alpha, \beta] \cap [1, k]$. Since $\alpha \le i \le k$, we have that

$$X_k(D_z + E_{[\alpha+1,i]}) = \sum_{j=1}^k d_{z,j} + (i-\alpha) = X_k(D_z) + (i-\alpha) ,$$

$$Y_k(D_z + E_{[\alpha+1,i]}) = k(k-1) + \sum_{j=k+1}^n \min(d_{z,j},k) = Y_k(D_z) .$$

Observe $i \in AVOID(k)$ if and only if $X_k(D_z + E_{[\alpha+1,i]}) \ge Y_k(D_z + E_{[\alpha+1,i]}) + 1$, which is equivalent to

$$\alpha + 1 + Y_k(D_z) - X_k(D_z) \le i \,.$$

It follows that

AVOID
$$(k) \cap [1, k] = [\alpha, \beta] \cap [\alpha + 1 + Y_k(D_z) - X_k(D_z), k] = AVOID_1(k)$$
.

Next consider any $i \in [\alpha, \beta] \cap [k, n]$. As $k \leq i \leq \beta$, we have that

$$\begin{aligned} X_k(D_z + E_{[\alpha+1,i]}) &= \sum_{j=1}^k d_{z,j} + \max\left\{0, k - \alpha\right\} = X_k(D_z) + \max\left\{0, k - \alpha\right\} ,\\ Y_k(D_z + E_{[\alpha+1,i]}) &= \begin{cases} Y_k(D_z) & k \le z, \\ Y_k(D_z) + (i - \alpha) & k \ge z + 1. \end{cases} \end{aligned}$$

We have the following three different cases:

Case 1: $k \leq z$ and $\max\{0, k - \alpha\} + X_k(D_z) > Y_k(D_z)$: In this case $X_k(D_z + E_{[\alpha+1,i]}) = X_k(D_z) + \max\{0, k - \alpha\}$ and $Y_k(D_z + E_{[\alpha+1,i]}) = Y_k(D_z)$, hence $X_k(D_z + E_{[\alpha+1,i]}) > Y_k(D_z + E_{[\alpha+1,i]})$, implying that $i \in \text{AVOID}_2(k)$. Therefore, in this case $\text{AVOID}_2(k) = [\alpha, \beta] \cap [k, n]$.

- Case 2: $\max\{0, k \alpha\} + X_k(D_z) \le Y_k(D_z)$: In this case $X_k(D_z + E_{[\alpha+1,i]}) = X_k(D_z) + \max\{0, k - \alpha\} \le Y_k(D_z) \le Y_k(D_z + E_{[\alpha+1,i]})$, thus $\operatorname{Avoid}_2(k) = \emptyset$.
- Case 3: $k \ge z + 1$:

In this case $Y_k(D_z + E_{[\alpha+1,i]}) = Y_k(D_z) + (i - \alpha)$. Observe that, $i \in AVOID(k)$ if and only if $X_k(D_z + E_{[\alpha+1,i]}) \ge Y_k(D_z + E_{[\alpha+1,i]}) + 1$. The latter is satisfied if and only if $X_k(D_z) + \max\{0, k - \alpha\} \ge Y_k(D_z) + (i - \alpha) + 1$, or if

$$i \le \max\left\{\alpha, k\right\} + X_k(D_z) - Y_k(D_z) - 1$$

Hence, $\operatorname{AVOID}_2(k) = [\alpha, \beta] \cap [k, \max\{k, \alpha\} + X_k(D_z) - Y_k(D_z) - 1].$

The lemma follows.

Algorithm 4 presents the procedure for computing the most-regular certificate. For each $k \in [1, n]$, AVOID(k) is a contiguous sub-interval of [1, n], therefore, the union AVOID = $\bigcup_{k=1}^{n} \text{AVOID}(k)$ can be computed in linear time using simple stack based data-structure, once the intervals are sorted in order of their endpoints⁴ using integer sort. Let \mathcal{I}_z denote the set obtained by removing from $[\alpha, \beta] \setminus \text{AVOID}$ each index *i*

⁴We say $[r, s] \leq [r', s']$ if either (i) r < r', or (ii) r = r' and $s \leq s'$.

Algorithm 4: MOST-REGULAR-CERTIFICATE(A, B)

1 OPT $\leftarrow \infty$; 2 foreach $z \in [\min(A), \max(B) - 1]$ do $I_1 \leftarrow \{i \in [1, n] \mid a_i \ge z + 1\};$ 3 $I_2 \leftarrow \{i \in [1,n] \mid a_i \le z, z+1 \le b_i\};$ 4 $I_3 \leftarrow \{i \in [1, n] \mid b_i \leq z\};$ 5 Rearrange the pairs in (A, B) along with the corresponding sets I_1, I_2, I_3 so that (i) for any 6 triplet (i, j, k) satisfying $i \in I_1, j \in I_2, k \in I_3$, we have i < j < k, and (ii) the sub-sequences $A[I_1]$ and $B[I_3]$ are sorted in the non-increasing order; Initialize $D_z = (d_1, d_2, \dots, d_n)$, where for $i \in I_1$, $d_i = a_i$; for $i \in I_2$, $d_i = z$; and for $i \in I_3$, 7 $d_i = b_i;$ if $z = \min(A)$ then set $\phi(D_z) = \sum_{1 \le r < s \le n} |a_r - a_s|$; 8 Compute $X(D_z), Y(D_z)$ using Theorem 11; 9 Let $\alpha = |I_1|$ and $\beta = |I_1| + |I_2|$; 10 for k = 1 to n do 11 AVOID₁(k) = $[\alpha, \beta] \cap [\alpha + 1 + Y_k(D_z) - X_k(D_z), k];$ 12 if $\max\{k - \alpha, 0\} + X_k(D_z) > Y_k(D_z)$ and $k \le z$ then $\operatorname{AvOID}_2(k) = [\alpha, \beta] \cap [k, n];$ 13 else if $\max\{k - \alpha, 0\} + X_k(D_z) \le Y_k(D_z)$ then $\operatorname{Avoid}_2(k) = \emptyset$; 14 else AVOID₂(k) = $[\alpha, \beta] \cap [k, \max\{k, \alpha\} + X_k(D_z) - Y_k(D_z) - 1];$ 15 $AVOID(k) = AVOID_1(k) \cup AVOID_2(k);$ 16 Compute AVOID = $\bigcup_{k=1}^{n} \text{AVOID}(k)$; 17 foreach $i \in [\alpha, \beta] \setminus \text{AVOID}$ do 18 if $(\text{parity}(D_z) = (i - \alpha) \mod 2)$ then 19 Compute $\phi(D_z + E_{[\alpha+1,i]}) = \phi(D_z) + (i - \alpha)(n - i - \alpha);$ 20 $OPT = \min\{OPT, \phi(D_z + E_{[\alpha+1,i]})\};\$ 21 Set $\phi(D_{z+1}) = \phi(D_z) + (\beta - \alpha)(n - \beta - \alpha);$ 22 23 Return OPT and the corresponding graphic sequence;

for which parity $(D_z + E_{[\alpha+1,i]}) = \text{parity}(\text{sum}(D_z) + (i-\alpha))$ is non-zero. Since $\text{sum}(D_z)$ (or parity (D_z)) is computable in O(n), the set \mathcal{I}_z can be computed in O(n) time as well. Note that $D_z + E_{[\alpha+1,i]}$ is graphic if and only if $i \in \mathcal{I}_z$. By Lemma 10, for any index $i \in \mathcal{I}_z$, the value $\phi(D_z + E_{[\alpha+1,i]})$ is computable in O(1) time, once we know $\phi(D_z)$. This shows that in just O(n) time, we can compute the spread of all the levelled sequences D satisfying $z \leq F^{-1}(\text{VOL}(D), \mathcal{S}) < z + 1$, and also find a sequence having the minimum spread. All that remains is to efficiently computing $\phi(D_z)$ for each $z \in [\min(A), \max(B)]$. Observe that $D_{\min(A)} = A$, and so $\phi(D_{\min(A)}) = \sum_{1 \leq r < s \leq n} |a_r - a_s|$ is computable in $O(n^2)$ time. Next by Lemma 10, for any $z \in [\min(A), \max(B) - 1]$, $\phi(D_{z+1}) = \phi(D_z) + (\beta - \alpha)(n - \beta - \alpha)$ is computable in O(1) time. Since z can take $\max(B) - \min(A) - 1$ values, our algorithm in total takes $O(n^2 + n(\max(B) - \min(A) - 1)) = O(n^2)$ time.

This completes the proof of Theorem 3.

6 Applications and Extensions

In this section, we discuss some related problems whose solutions follow as immediate application of our interval sequence work.

Problem 1 (Minimum Graphic extensions). Given a sequence $A = (a_1, \ldots, a_p)$ find the minimum integer $n \geq p$ such that a super sequence $D = (a_1, \ldots, a_p, d_{p+1}, d_{p+2}, \ldots, d_n)$ of sequence A is realizable.

Solution: Let M denote the value $\max(A) = \max_{i \in [1,p]} a_i$. For any $n \ge p$, let $S_n = ([a_1, a_1], \ldots, [a_p, a_p], [1, n], \ldots, [1, n])$ denote the sequence obtained by appending n - p copies of interval [1, n] to interval sequence (A, A). Let n_0 the denote the length of a minimum graphic extension of A. Observe that $n_0 \in [\max\{p, M\}, p + M]$. The lower limit is due to the fact that the length of minimum graphic extension of A must be at least $\max\{p, M\}$; the upper limit holds since one can have a bipartite graph with partitions $X = \{x_1, \ldots, x_p\}$ and $Y = \{y_1, \ldots, Y_M\}$ of length p and M, and for $i \in [1, p]$, connect the vertex x_i to vertices y_1, \ldots, y_{a_i} . It turns out that we need to find the smallest integer $n \in [\max\{p, M\}, p + M]$ such that S_n is graphic. The minimum n can be obtained by a binary search over the range $[\max\{p, M\}, p + d]$ and using Theorem 5; this takes $O(\max\{p, M\} \log \max\{p, M\})$ time. Once n_0 is known, the optimal graphic extension can be computed using Theorem 10 for searching graphic certificate in $O(\max\{p, M\} \log \max\{p, M\})$ time.

Problem 2. Given $A = (a_1, ..., a_n)$, find a graphic sequence $D = (d_1, ..., d_n)$ whose chebyshev distance $(L_{\infty} \text{ distance})$ from A is minimum.

Solution: The above problem can be reduced to interval sequence problem, as we need to find smallest non-negative integer $c \in [1, n]$ such that $S_c = ([a_1 - c, a_1 + c], \dots, [a_n - c, a_n + c])$ is realizable. To find the minimum c, we do a binary search with help of Theorem 5 for verification; this takes $O(n \log n)$ time. Once optimal c is known, the sequence D can be computed using Theorem 10 to search graphic certificate in S_c , thus the time complexity for computing sequence D is $O(n \log n)$.

Problem 3. Given $A = (a_1, \ldots, a_n)$, find minimum fraction ϵ and a graphic sequence $D = (d_1, \ldots, d_n)$ satisfying $a_i(1 - \epsilon) \le d_i \le a_i(1 + \epsilon)$.

Solution: Again we need to find smallest non-negative fraction ϵ such that the interval sequence $S_{\epsilon} = ([a_1(1-\epsilon), a_1(1+\epsilon)], \dots, [a_n(1-\epsilon), a_n(1+\epsilon)])$ is realizable. To find the minimum ϵ , we do a binary search with help of Theorem 5; this takes $O(n \log n)$ time. Once ϵ is known, using Theorem 10, sequence D can be computed in $O(n \log n)$ time.

References

- [1] Martin Aigner and Eberhard Triesch. Realizability and uniqueness in graphs. *Discrete Mathematics*, 136:3–20, 1994.
- [2] Richard Anstee. Properties of a class of (0,1)-matrices covering a given matrix. *Can. J. Math.*, pages 438–453, 1982.
- [3] Richard Anstee. An algorithmic proof of tutte's f-factor theorem. J. Algorithms, 6(1):112–131, 1985.
- [4] Amotz Bar-Noy, Keerti Choudhary, David Peleg, and Dror Rawitz. Realizability of graph specifications: Characterizations and algorithms. In 25th SIROCCO, volume 11085 of LNCS, pages 3–13, 2018.

- [5] Amotz Bar-Noy, Keerti Choudhary, David Peleg, and Dror Rawitz. Graph profile realizations and applications to social networks. In 13th WALCOM, volume 11355 of LNCS, pages 1–12, 2019.
- [6] M. Behzad and James E. Simpson. Eccentric sequences and eccentric sets in graphs. Discrete Mathematics, 16(3):187 – 193, 1976.
- [7] Mao-cheng Cai, Xiaotie Deng, and Wenan Zang. Solution to a problem on degree sequences of graphs. *Discrete Mathematics*, 219(1-3):253–257, 2000.
- [8] Wai-Kai Chen. On the realization of a (p,s)-digraph with prescribed degrees. *Journal of the Franklin Institute*, 281(5):406 422, 1966.
- [9] Paul Erdös and Tibor Gallai. Graphs with prescribed degrees of vertices [hungarian]. *Matematikai Lapok*, 11:264–274, 1960.
- [10] A. Frank. Augmenting graphs to meet edge-connectivity requirements. SIAM J. Discrete Math., 5:25– 43, 1992.
- [11] A. Frank. Connectivity augmentation problems in network design. In *Mathematical Programming: State of the Art*, pages 34–63. Univ. Michigan, 1994.
- [12] H. Frank and W. Chou. Connectivity considerations in the design of survivable networks. *IEEE Trans. Circuit Theory*, CT-17:486–490, 1970.
- [13] Satoru Fujishige and Sachin B. Patkar. Realization of set functions as cut functions of graphs and hypergraphs. *Discrete Mathematics*, 226(1-3):199–210, 2001.
- [14] D.R. Fulkerson. Zero-one matrices with zero trace. *Pacific J. Math.*, 12:831 836, 1960.
- [15] Ankit Garg, Arpit Goel, and Amitabha Tripathi. Constructive extensions of two results on graphic sequences. *Discrete Applied Mathematics*, 159(17):2170–2174, 2011.
- [16] R.E. Gomory and T.C. Hu. Multi-terminal network flows. J. Soc. Industrial & Applied Math., 9, 1961.
- [17] Jiyun Guo and Jianhua Yin. A variant of Niessen's problem on degree sequences of graphs. Discrete Mathematics and Theoretical Computer Science, Vol. 16 no. 1 (in progress)(1):287–292, May 2014. Graph Theory. URL: https://hal.inria.fr/hal-01179211.
- [18] S. Louis Hakimi. On realizability of a set of integers as degrees of the vertices of a linear graph –I. SIAM J. Appl. Math., 10(3):496–506, 1962.
- [19] Sepp Hartung and André Nichterlein. Np-hardness and fixed-parameter tractability of realizing degree sequences with directed acyclic graphs. SIAM J. Discrete Math., 29(4):1931–1960, 2015.
- [20] V. Havel. A remark on the existence of finite graphs [in Czech]. Casopis Pest. Mat., 80:477–480, 1955.
- [21] Katherine Heinrich, Pavol Hell, David G. Kirkpatrick, and Guizhen Liu. A simple existence criterion for g ; ffactors, with applications to [a, b]factors. *Discrete Mathematics*, 85:313–317, 1990.
- [22] Pavol Hell and David G. Kirkpatrick. Linear-time certifying algorithms for near-graphical sequences. *Discrete Mathematics*, 309(18):5703–5713, 2009.

- [23] Daniel J. Kleitman and D. L. Wang. Algorithms for constructing graphs and digraphs with given valences and factors. *Discrete Mathematics*, 6(1):79–88, 1973.
- [24] Linda Lesniak. Eccentric sequences in graphs. *Periodica Mathematica Hungarica*, 6(4):287–293, 1975.
- [25] László Lovász. Subgraphs with prescribed valencies. J. Comb. Theory, 8:391-416, 1970.
- [26] N.V.R. Mahadev and U.N. Peled. *Threshold Graphs and Related Topics*. Annals of Discrete Mathematics. Elsevier Science, 1995.
- [27] Akutsu Tatsuya and Hiroshi Nagamochi. Comparison and enumeration of chemical graphs. *Computational and structural biotechnology*, 5, 2013.
- [28] Amitabha Tripathi, Sushmita Venugopalan, and Douglas B. West. A short constructive proof of the erdos-gallai characterization of graphic lists. *Discrete Mathematics*, 310(4):843–844, 2010.
- [29] W. T. Tutte. Graph factors. *Combinatorica*, 1(1):79–97, Mar 1981.
- [30] D.L. Wang and D.J. Kleitman. On the existence of *n*-connected graphs with prescribed degrees (n > 2). *Networks*, 3:225–239, 1973.

Appendix

A An O(n) time implementation of Theorem 4 and Theorem 5

Erdös and Gallai [9] presented a characterisation for graphic sequences; and later Cai et al. [7] presented a similar characterisation for realizable interval-sequences (Theorem 4 and Theorem 5). For completeness, we present simple O(n) algorithm for efficiently computing vectors X(D), Y(D), and $\varepsilon(S)$ for any given sequence $D = (d_1, d_2, \ldots, d_n)$ and interval-sequence $S = (A, B) = ([a_1, b_1], \ldots, [a_n, b_n])$. Recall

$$\begin{aligned} X_k(D) &= \sum_{i=1}^k d_i; \ Y_k(D) = k(k-1) + \sum_{i=k+1}^n \min(d_i, k); \ W_k(\mathcal{S}) = \{i \in [k+1, n] \mid b_i \ge k+1\}; \\ \varepsilon_k(\mathcal{S}) &= \begin{cases} 1 & \text{if } a_i = b_i \text{ for } i \in W_k(\mathcal{S}) \text{ and } \sum_{i \in W_k(\mathcal{S})} (b_i + k|W_k(\mathcal{S})|) \text{ is odd.} \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

For any $k \in [1, n]$, j(k) is defined to be 0 if $k > d_1$, else $j(k) \in [1, n]$ is the largest index such that $d_1, d_2, \ldots, d_{j(k)} \ge k$. Then

$$Y_k(D) = k(k-1) + \sum_{i=k+1}^{j(k)} k + \sum_{i=j(k)+1}^n d_i = k(j(k)-1) + \sum_{i=j(k)+1}^n d_i.$$

Observe that j(k) is a non-increasing sequence.

Algorithm 5: Computation of vectors

1 Let $D = (d_1, d_2, ..., d_n)$ be the input sequence; 2 Set $X_1(D) = d_1, j(1) = n$, and $W_1(D) = 0$; 3 for k = 2 to n do 4 $X_k(D) = X_{k-1}(D) + d_k$; 5 while $(d_{j(k)} \not\geq k)$ do 6 j(k) = j(k) - 1; 7 $W_k(D) = W_{k-1}(D) + d_{j(k)}$; 8 $Y_k(D) = k(j(k) - 1) + W_k(D)$; 9 return (X(D), Y(D)).

It is easy to verify that the Algorithm 5 takes linear time, and correctly computes the vectors X(D) and Y(D). Also similarly $\varepsilon(S)$ is computable in O(n) time. So the following theorem is immediate.

Theorem 11. Given any sequence D and interval-sequence S = (A, B) of length n, the vectors X(D), Y(D), and $\varepsilon(S)$ are all computable in O(n) time.