

Relational Width of First-Order Expansions of Homogeneous Graphs with Bounded Strict Width

Michał Wrona 

Theoretical Computer Science Department, Faculty of Mathematics and Computer Science,
Jagiellonian University, Poland
wrona@tcs.uj.edu.pl

Abstract

Solving the algebraic dichotomy conjecture for constraint satisfaction problems over structures first-order definable in countably infinite finitely bounded homogeneous structures requires understanding the applicability of local-consistency methods in this setting. We study the amount of consistency (measured by relational width) needed to solve $\text{CSP}(\mathbb{A})$ for first-order expansions \mathbb{A} of countably infinite homogeneous graphs $\mathcal{H} := (A; E)$, which happen all to be finitely bounded. We study our problem for structures \mathbb{A} that additionally have bounded strict width, i.e., for which establishing local consistency of an instance of $\text{CSP}(\mathbb{A})$ not only decides if there is a solution but also ensures that every solution may be obtained from a locally consistent instance by greedily assigning values to variables, without backtracking.

Our main result is that the structures \mathbb{A} under consideration have relational width exactly $(2, \mathbb{L}_{\mathcal{H}})$ where $\mathbb{L}_{\mathcal{H}}$ is the maximal size of a forbidden subgraph of \mathcal{H} , but not smaller than 3. It beats the upper bound: $(2m, 3m)$ where $m = \max(\text{arity}(\mathbb{A}) + 1, \mathbb{L}, 3)$ and $\text{arity}(\mathbb{A})$ is the largest arity of a relation in \mathbb{A} , which follows from a sufficient condition implying bounded relational width given in [10]. Since $\mathbb{L}_{\mathcal{H}}$ may be arbitrarily large, our result contrasts the collapse of the relational bounded width hierarchy for finite structures \mathbb{A} , whose relational width, if finite, is always at most $(2, 3)$.

2012 ACM Subject Classification Theory of computation \rightarrow Complexity classes; Theory of computation \rightarrow Problems, reductions and completeness

Keywords and phrases Constraint Satisfaction, Homogeneous Graphs, Bounded Width, Strict Width, Relational Width, Computational Complexity

Digital Object Identifier 10.4230/LIPIcs.STACS.2020.35

Funding The research was partially supported by NCN grant number 2014/14/A/ST6/00138.

1 Introduction

The constraint satisfaction problem (CSP) is one of the most important problems in theoretical and applied computer science and at the same time it is a general framework in which many other computational problems may be formalized. Given a number of constraints imposed on variables one asks if there is a global solution, i.e., a function assigning values to variables so that all the constraints are simultaneously satisfied. Boolean satisfiability and graph colouring are among the most prominent examples of NP-hard problems that can be formalized as CSPs and hence the CSP is NP-hard in general. Thus, one considers the problem $\text{CSP}(\mathbb{A})$ parametrized by a relational structure (called also a constraint language, a language or a template) \mathbb{A} . (In this paper, \mathbb{A} is always over a finite signature). A longstanding open problem in this area was to verify the Feder-Vardi [20] conjecture which states that for every finite \mathbb{A} the problem $\text{CSP}(\mathbb{A})$ is either in P or it is NP-complete. After over thirty years of work and a number of important partial results this so-called Dichotomy Conjecture was confirmed independently in [26] and [16]. In both cases the proof was carried out in the so-called universal-algebraic approach to the complexity of CSPs [22, 17]. The approach not only provided appropriate tools but also suggested the delineation. This so-called algebraic



© Michał Wrona;

licensed under Creative Commons License CC-BY

37th International Symposium on Theoretical Aspects of Computer Science (STACS 2020).

Editors: Christophe Paul and Markus Bläser; Article No. 35; pp. 35:1–35:32

Leibniz International Proceedings in Informatics



LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

dichotomy conjecture [17] saying that $\text{CSP}(\mathbb{A})$ is hard under the condition that the algebra corresponding to \mathbb{A} lacks interesting operations also has been confirmed in both proofs.

The universal-algebraic approach to finite-domain constraint satisfaction problems has been generalized to capture the computational complexity in many other similar settings. In particular, the complexity of $\text{CSP}(\mathbb{A})$ depends on the algebra corresponding to \mathbb{A} when \mathbb{A} is ω -categorical [11], i.e., all countable models of the first-order theory of \mathbb{A} are isomorphic. In particular, all structures first-order definable in (reducts of) (countably infinite) homogeneous structures over finite signatures are ω -categorical structure. (A structure is homogeneous if every isomorphism between its finite substructures may be extended to an automorphism of a structure.) Considering these infinite structures significantly broadens the class of problems that may be captured within the CSP framework. In particular, the order over rational numbers $(\mathbb{Q}, <)$, which is homogeneous, gives rise to $\text{CSP}(\mathbb{Q}; <)$ that can be seen as the digraph acyclicity problem. The latter cannot be expressed as the CSP over a finite template. Furthermore a number of problems of interest in qualitative reasoning may be captured by $\text{CSP}(\mathbb{A})$ where \mathbb{A} is a reduct of a homogeneous structure \mathbb{B} . It concerns constraint satisfaction problems in formalisms such as Allen's interval algebra or RCC-5, see [6] for a survey. Many of the homogeneous structures \mathbb{B} of interest are finitely bounded, i.e., there exists a finite unique minimal set $\mathcal{F}_{\mathbb{B}}$ of finite structures over the signature of \mathbb{B} such that a finite structure Δ embeds into \mathbb{B} if and only if none of the structures in $\mathcal{F}_{\mathbb{B}}$ embeds into Δ . A dichotomy for algebras corresponding to reducts of countably infinite finitely bounded homogeneous structures was proved in [3]. As in the finite case, it suggests the delineation between polynomial-time solvable and NP-hard CSPs. Although the complexity dichotomy is still far from being obtained, the algebraic dichotomy conjecture for reducts of finitely bounded homogeneous structures is known to hold in the number of cases including the reducts of $(\mathbb{N}, =)$ [7], $(\mathbb{Q}, <)$ [8], the random partial order [23] or a countably infinite homogeneous graph [13, 14, 9].

Theoretical research on CSPs is focused not only on providing classifications of computational complexity but also on settling the limits of applicability of widely known algorithms or algorithmic techniques such as establishing local consistency. This method is used not only for finite CSP but is also considered to be the most important (if not the only) algorithmic technique for qualitative CSPs [25]. The algebraic characterization of finite structures \mathbb{A} with bounded width [2], i.e., for which $\text{CSP}(\mathbb{A})$ can be solved by establishing local consistency, is considered to be an important step towards solving the Feder-Vardi conjecture. Thus, in order to understand the complexity of CSPs for reducts \mathbb{A} of finitely bounded homogeneous structures, we need to characterize \mathbb{A} with bounded width and to understand how different notions of consistency relate to each other for templates under consideration. The focus of this paper is on the latter.

The amount of consistency needed to solve $\text{CSP}(\mathbb{A})$ for \mathbb{A} with bounded width is measured here [1] and here [15] by relational width. The relational width of \mathbb{A} is a pair of numbers (k, l) with $k \leq l$ (for the exact definition we refer the reader to Section 4). The following question was of interest for finite structures.

► **Question 1.** *What is the exact relational width of \mathbb{A} with bounded width?*

Question 1 for finite \mathbb{A} was completely answered in [1] where it was proved that \mathbb{A} with bounded width has always either relational width $(1, 1)$ or $(2, 3)$, see [15] for another proof. Both proofs rely, however, on the algebraic characterization of structures \mathbb{A} with bounded width. Although the notion of bounded width has been generalized to ω -categorical structures [5], according to our knowledge, no algebraic characterization of bounded width for such structures is within sight. Nevertheless the algebraic characterization of strict bounded

width has been quite easily lifted from finite [20] to infinite domains [5]. (Again, for a detailed definition we refer the reader to Section 4.) A reduct of a finitely bounded homogeneous structure has bounded strict width if and only if it is preserved by so-called oligopotent quasi near-unanimity operation. This algebraic characterization gives us a hope to answer the following question analogous to Question 1.

► **Question 2.** *What is the exact relational width of reducts \mathbb{A} of finitely bounded homogeneous structures with bounded strict width?*

In this paper we answer Question 2 for first-order expansions of countably infinite homogeneous graphs. We believe that our method may be used to provide the general answer in the near future. We note that the answer to Question 2 would be not only a nice theoretical result but should be also of particular interest for structures that give rise to constraint satisfaction problems in qualitative reasoning. In this context strict width is called *local-to-global consistency* and has been widely studied see, e.g., [19].

1.1 Our results

In contrast to all homogeneous structures, all countably infinite homogeneous graphs are well understood and have been classified in [24]. It happens that every such graph \mathcal{H} is also finitely bounded, i.e., in each case there exists a finite unique minimal set of finite graphs $\mathcal{F}_{\mathcal{H}}$ such that a finite graph G embeds into \mathcal{H} if and only if none of the graphs in $\mathcal{F}_{\mathcal{H}}$ embeds into G . We will write $\mathbb{L}_{\mathcal{H}}$ for the maximum of the number 3 and the size of the largest finite structure in $\mathbb{L}_{\mathcal{H}}$. Perhaps the best known example of a homogeneous graph is the random graph that is determined up to isomorphism by the two properties of being homogeneous and universal (i.e., it contains all countable graphs as induced subgraphs). Equivalently, the random graph is a unique countably infinite graph which has this extension property: for all disjoint finite subsets U, U' of the domain there exists an element v such that v is adjacent to all members of U and to none in U' . In this case the finite set of bounds consists of a single directed edge and a loop, and hence \mathbb{L}_G for the random graph G is 3. Furthermore, the family of homogeneous graphs contains universal countable k -clique free graphs H_k with $k \geq 3$, called also Henson graphs, in which case \mathcal{F}_{H_k} contains also a k -clique, and hence \mathbb{L}_{H_k} is k or the graphs C_n^s that are disjoint sums of n cliques of size s where $1 \leq n, s \leq \omega$ and either n or s equals ω . Observe that $\mathcal{F}_{C_n^s}$ contains a graph on three vertices with two edges and one non-edge as well as a null graph over $n+1$ vertices in case n is finite or a $(s+1)$ -clique in the case where s is finite. Thus, $\mathbb{L}_{C_n^s}$ is either 3, $n+1$ or $s+1$. All remaining homogeneous graphs are the complements of graphs H_k or C_n^s . In this paper we prove the following.

Main result. *Let \mathbb{A} be a first-order expansion of a countably infinite homogeneous graph \mathcal{H} such that \mathbb{A} has bounded strict width. Then \mathbb{A} has relational width $(2, \mathbb{L}_{\mathcal{H}})$.*

In fact, we obtain a more general result. Some sufficient conditions implying that a first-order expansion of a homogeneous graph \mathcal{H} has relational width $(2, \mathbb{L}_{\mathcal{H}})$ are given in Section 5. In particular, the conditions cover all languages under consideration preserved by binary canonical operations considered in [13, 14, 9] where an analysis of algebras corresponding to reducts of homogeneous graphs and the computational dichotomy is provided. Our result: relational width $(2, \mathbb{L}_{\mathcal{H}})$ beats the upper bound $(2m, 3m)$, where $m = \max(\text{arity}(\mathbb{A}) + 1, \mathbb{L}, 3)$ and $\text{arity}(\mathbb{A})$ is the largest arity of a relation in \mathbb{A} , that can be easily obtained from the proof of Theorem 4.10 in [10].

We believe that measuring relational width of structures with bounded width is interesting in its own rights. Nevertheless, our research has complexity consequences. As in the finite case, it was proved in [5] that $\text{CSP}(\mathbb{A})$ for an ω -categorical \mathbb{A} with strict width k may be solved by establishing $(k, k+1)$ -consistency and hence in time $O(n^{k+1})$ where n is the number of variables in an instance. Our main result implies that such $\text{CSP}(\mathbb{A})$ for a first-order expansion \mathbb{A} of a homogeneous graph \mathcal{H} may be solved by establishing $(2, \mathbb{L}_{\mathcal{H}})$ -minimality, and hence in time $O(n^m)$ where $m = \max(\mathbb{L}_{\mathcal{H}}, \text{arity}(\mathbb{A}))$.

1.2 Outline of the paper

We start with general preliminaries in Section 2. Then we review canonical operations providing tractability for reducts of homogeneous graphs, Section 3. Bounded (relational) width, strict width and other notions related to local consistency are provided in Section 4. There we also give a number of examples explaining the applicability of our main result. The proof of the main result is divided into Section 5 and Section 6. In the former one, we give a number of sufficient conditions implying relational width $(2, \mathbb{L}_{\mathcal{H}})$, while in the latter one we show that the sufficient conditions are satisfied whenever a first-order expansion of \mathcal{H} has bounded strict width. In Section 5 we additionally show that the sufficient condition are also satisfied by first-order expansions of homogeneous graphs preserved by the studied binary canonical operations. As a consequence, we obtain that all tractable (whose CSP is solvable in polynomial time) reducts of \mathcal{H} where \mathcal{H} is $C_1^\omega, C_\omega^1, C_\omega^\omega$ or H_k with $k \geq 3$ have bounded relational width $(2, \mathbb{L}_{\mathcal{H}})$ and hence can be solved by establishing $(2, \mathbb{L}_{\mathcal{H}})$ -minimality.

2 Preliminaries

We write $t = (t[1], \dots, t[n])$ for a tuple of elements and $[n]$ to denote the set $\{1, \dots, n\}$.

2.1 Relations, languages and formulas

In this paper we consider first-order expansions $\mathbb{A} := (A; E, R_1, \dots, R_k)$ over a finite signature τ of homogeneous graphs, called also (constraint) languages or templates, where all R_1, \dots, R_k have a first-order definition in $(A; E)$. We assume that \mathbb{A} contains $=$ and N whenever N is pp-definable in \mathbb{A} . Relations E and N refer always to a homogeneous graph \mathcal{H} known from the context. For the sake of presentation we usually do not distinguish between a relation symbol R in the signature of \mathbb{A} and the relation $R^{\mathbb{A}}$ and use the former symbol for both. We often write O, O_1, O_2, \dots for elements of $\{E, N, =\}$ and $\underline{E}, \underline{N}, \underline{O}, \underline{O}_1, \underline{O}_2$ to denote relations $(E \cup =), (N \cup =), (O \cup =), (O_1 \cup =), (O_2 \cup =)$, respectively.

For a structure \mathbb{A} over domain A and a tuple $t \in A^k$, the orbit of t in \mathbb{A} is the relation $\{(\alpha(t[1]), \dots, \alpha(t[k])) \mid \alpha \in \text{Aut}(\mathbb{A})\}$ where $\text{Aut}(\mathbb{A})$ is the set of automorphisms of \mathbb{A} . In particular, E, N and $=$ are orbits of pairs, called also orbitals. We would like to note that all structures considered in this paper are ω -categorical. By a theorem proved independently by Ryll-Nardzewski, Engeler and Svenonius, a structure \mathbb{A} is ω -categorical if and only if its automorphism group is oligomorphic, i.e., for every n the number of orbits of n -tuples is finite. See [21] for a textbook on model theory.

A primitive-positive (pp-)formula is a first-order formula built exclusively out of existential quantifiers \exists , conjunction \wedge and atomic formulas $R(x_1, \dots, x_k)$ where R is a k -ary relation symbol and x_1, \dots, x_k are variables, not necessarily pairwise different.

2.2 The universal-algebraic approach

We say that an operation $f : A^n \rightarrow A$ is a polymorphism of an m -ary relation R iff for any m -tuples $t_1, \dots, t_n \in R$, it holds that the tuple $(f(t_1[1], \dots, t_n[1]), \dots, f(t_1[m], \dots, t_n[m]))$ is also in R . We write $f(t_1, \dots, t_n)$ as a shorthand for $(f(t_1[1], \dots, t_n[1]), \dots, f(t_1[m], \dots, t_n[m]))$. An operation f is a polymorphism of \mathbb{A} if it is a polymorphism of every relation in \mathbb{A} . If $f : A^n \rightarrow A$ is a polymorphism of \mathbb{A} , R , we say that f preserves \mathbb{A} , R , otherwise that f violates \mathbb{A} , R . A set of polymorphisms of an ω -categorical structure \mathbb{A} forms an algebraic object called an oligomorphic locally closed clone [4], which in particular contains an oligomorphic permutation group [18].

► **Theorem 1.** ([11]) *Let \mathbb{A} be a countable ω -categorical structure. Then R is preserved by the polymorphisms of \mathbb{A} if and only if it has a primitive-positive definition in \mathbb{A} , i.e., a definition via a primitive-positive formula.*

We say that a set of operations F generates a set of operations G if every $g \in G$ is in the smallest locally-closed clone containing F . We write $\overline{\text{Aut}(\mathbb{A})}$ to denote the clone generated by the automorphisms of the structure \mathbb{A} . An operation f of an oligomorphic clone F is called oligopotent if $\{g\}$ where $g(x) := f(x, \dots, x)$ is generated by the permutations in F . We say that a k -ary operation f is a weak near-unanimity operation if $f(y, x, \dots, x) = f(x, y, x, \dots, x) = \dots = f(x, \dots, x, y)$ for all $x, y \in A$ and that f is a quasi near-unanimity operation (short, qnu-operation) if it is a weak near-unanimity and it additionally satisfies $f(x, \dots, x) = f(x, \dots, x, y)$ for all $x, y \in A$. We say that a k -ary operation f is a weak near-unanimity operation modulo $\overline{\text{Aut}(\mathbb{A})}$ if there exist $e_1, \dots, e_k \in \overline{\text{Aut}(\mathbb{A})}$ such that: $e_1(f(y, x, \dots, x)) = e_2(f(x, y, x, \dots, x)) = \dots = e_k(f(x, \dots, x, y))$ for all $x, y \in A$.

2.3 The constraint satisfaction problem

We define the CSP to be a computational problem whose instance \mathcal{I} is a triple $(\mathcal{V}, \mathcal{C}, A)$ where $\mathcal{V} = \{v_1, \dots, v_n\}$ is a set of variables, \mathcal{C} is a set of constraints each of which is of the form $((v_{i_1}, \dots, v_{i_k}), R)$ where $\{v_{i_1}, \dots, v_{i_k}\} \subseteq \mathcal{V}$ is the scope of the constraint and $R \subseteq A^k$. The question is whether there is a solution $\mathbf{s} : \mathcal{V} \rightarrow A$ to \mathcal{I} satisfying $(\mathbf{s}(v_{i_1}), \dots, \mathbf{s}(v_{i_k})) \in R$ for all $((v_{i_1}, \dots, v_{i_k}), R) \in \mathcal{C}$. Further, we define $\text{CSP}(\mathbb{A})$ for a constraint language \mathbb{A} to be the CSP restricted to instances where all relations come from \mathbb{A} .¹

We define the projection of $((v_{i_1}, \dots, v_{i_k}), R)$ to the set $\{w_1, \dots, w_l\} \subseteq \{v_{i_1}, \dots, v_{i_k}\}$ to be the constraint $(\{w_1, \dots, w_l\}, R')$ where the relation R' is given by $(R'(w_1, \dots, w_l) \equiv \exists x_1 \dots \exists x_m R(v_{i_1}, \dots, v_{i_k}))$ and $\{x_1, \dots, x_m\} = \{v_{i_1}, \dots, v_{i_k}\} \setminus \{w_1, \dots, w_m\}$. Let $W \subseteq \mathcal{V}$. An assignment $\mathbf{a} : W \rightarrow A$ is a partial solution to \mathcal{I} if \mathbf{a} satisfies all projections of constraints in \mathcal{I} to variables in W .

It is very well known that adding pp-definable relations to the template does not change the complexity of the problem.

► **Proposition 2.** *Let $\mathbb{A} = (A; R_1, \dots, R_l)$ be a relational structure, and let R be a relation that has a primitive-positive definition in \mathbb{A} . Then $\text{CSP}(\mathbb{A})$ and $\text{CSP}(A, R, R_1, \dots, R_l)$ are log-space equivalent.*

¹ Equivalently, one defines an instance of $\text{CSP}(\mathbb{A})$ as a conjunction φ of atomic formulae over the signature of \mathbb{A} . Then the question is whether φ is satisfiable in \mathbb{A} .

B_1	$=$	E	N
$=$	$=$	E	N
E	E	E	E
N	N	E	N

B_2	$=$	E	N
$=$	$=$	E	E
E	E	E	E
N	E	E	E

B_3	$=$	E	N
$=$	$=$	N	N
E	N	E	N
N	N	N	N

■ **Figure 1** Three exemplary binary behaviours: B_1 , B_2 , and B_3 .

2.4 Efficient entailment

We say that a formula φ_1 entails a formula φ_2 both over free variables x_1, \dots, x_n if $(\forall x_1 \dots \forall x_n (\varphi_1(x_1, \dots, x_n) \implies \varphi_2(x_1, \dots, x_n)))$ is a valid sentence. Furthermore, we say that an n -ary relation R entails φ over free variables x_1, \dots, x_n if $R(x_1, \dots, x_n)$ entails φ . These definitions are quite standard but for the purposes of this paper we need a stronger notion of entailment.

► **Definition 3.** We say that a quaternary relation R efficiently entails $\psi := (S_1(x_1, x_2) \implies S_2(x_3, x_4))$ where S_1, S_2 are binary relations if R entails ψ and R contains

1. a tuple t_1 such that $(t_1[1], t_1[2]) \in S_1$ and $(t_1[3], t_1[4]) \in S_2$, and
2. a tuple t_2 such that $(t_2[1], t_2[2]) \notin S_1$ and $(t_2[3], t_2[4]) \notin S_2$.

We say that a quaternary relation R is a $[(S_1(x_1, x_2) \implies S_2(x_3, x_4)), (\varphi)]$ -relation if R efficiently entails $(S_1(x_1, x_2) \implies S_2(x_3, x_4))$ and entails φ or that a quaternary relation R is a $[(S_1(x_1, x_2) \implies S_2(x_3, x_4))]$ -relation if R efficiently entails $(S_1(x_1, x_2) \implies S_2(x_3, x_4))$.

3 Canonical Operations over Reducts of Homogeneous Graphs

The polymorphisms that appear in the complexity classifications of CSPs of reducts of homogeneous graphs display some regularities in the sense defined below.

Let $f : A^k \rightarrow A$, and let G be a permutation group on A . We say that f is canonical with respect to G if for all $m \in N, \alpha_1, \dots, \alpha_k \in G$ and m -tuples a_1, \dots, a_k , there exists $\beta \in G$ such that $\beta f(\alpha_1(a_1), \dots, \alpha_k(a_k)) = f(a_1, \dots, a_k)$. Equivalently, this means that f induces an operation $\xi^{\text{typ}}(f)$, called a k -ary behaviour, on orbits of m -tuples under G , by defining $\xi^{\text{typ}}(f)(O_1, \dots, O_k)$ as the orbit of $f(a_1, \dots, a_k)$ where a_i is any m -tuple in O_i . In what follows we are mainly interested in operations that are canonical with respect to $\text{Aut}(\mathcal{H})$ where \mathcal{H} is a homogeneous graph. Therefore we usually say simply canonical. See [12] for a survey on canonical operations. Three simple binary behaviors are presented in Figure 1. According to Definition 6, a binary injection f such that $\xi^{\text{typ}}(f)$ is

- B_1 is said to be of behavior max and balanced,
- B_2 is said to be E -constant,
- B_3 is said to be of type min and N -dominated.

We introduce the following notation. Let $R_1, \dots, R_k \subseteq A^2$ be binary relations. We write $R_1 \cdots R_k$ for the binary relation on A^k defined so that: $R_1 \cdots R_k(a_1, a_2)$ holds for k -tuples $a_1, a_2 \in A^k$ if and only if $R_i(a_1[i], a_2[i])$ holds for all $i \in [k]$. Here, we can find the list of all binary behaviours of interest.

► **Definition 4.** Let (A, E) be a countably infinite homogeneous graph. We say that a binary injective operation $f : A^2 \rightarrow A$ is

- balanced if for all $a, b \in A^2$ we have that $E=(a, b)$ and $=E(a, b)$ implies $E(f(a), f(b))$ as well as $N=(a, b)$ and $=N(a, b)$ implies $N(f(a), f(b))$,

- *E-dominated (N-dominated) if for all $a, b \in A^2$ with $\neq(a, b)$ or $=\neq(a, b)$ we have that $E(f(a), f(b))$ ($N(f(a), f(b))$);*
- *of behaviour min if for all $a, b \in A^2$ with $\neq(a, b)$ we have $E(f(a), f(b))$ iff $EE(a, b)$;*
- *of behaviour max if for all $a, b \in A^2$ with $\neq(a, b)$ we have $N(f(a), f(b))$ iff $NN(a, b)$;*
- *of behaviour projection if there exists $i \in [2]$ such that for all $a, b \in A^2$ with $\neq(a, b)$ we have $E(f(a), f(b))$ iff $E(a[i], b[i])$;*
- *of behaviour xor if for all $a, b \in A^2$ with $\neq(a, b)$ the relation $E(f(a), f(b))$ holds iff $EN(a, b)$ or $NE(a, b)$ holds;*
- *of behaviour xnor if for all $a, b \in A^2$ with $\neq(a, b)$ the relation $E(f(a), f(b))$ holds iff $EE(a, b)$ or $NN(a, b)$ holds;*
- *E-constant if the image of f is a clique,*
- *N-constant if the image of f is an independent set.*

We now turn to ternary behaviours of interest.

► **Definition 5.** Let $(A; E)$ be a countably infinite homogeneous graph. We say that a ternary injective operation $f : A^3 \rightarrow A$ is of behaviour

- *majority if for all $a, b \in D^3$ satisfying $\neq\neq(a, b)$ we have that $E(f(a), f(b))$ if and only if $EEE(a, b), EEN(a, b), ENE(a, b)$, or $NEE(a, b)$,*
- *minority if for all $a, b \in D^3$ satisfying $\neq\neq(a, b)$ we have that $N(f(a), f(b))$ if and only if $NNN(a, b), EEN(a, b), ENE(a, b)$, or $NEE(a, b)$.*

Furthermore, let B be a binary behavior. A ternary function is hyperplanely of behaviour B if the binary functions $(x, y) \rightarrow f(x, y, c)$, $(x, z) \rightarrow f(x, c, z)$, and $(y, z) \rightarrow f(c, y, z)$ have behavior B for all $c \in D$.

4 Consistency and Minimality

This section is devoted to the formal introduction of consistency and width notions. The main algorithm we are interested in is based on establishing minimality.

► **Definition 6.** Let $l \geq k > 0$ be natural numbers. An instance $\mathcal{I} = (\mathcal{V}, \mathcal{C}, A)$ of the CSP is (k, l) -minimal if:

1. Every at most l -element set of variables is contained in the scope of some constraint in \mathcal{I} .
2. For every set W with $|W| \leq k$ and every pair of constraints C_1 and C_2 in \mathcal{C} whose scopes contain W , the projections of the constraints C_1 and C_2 to W are the same.

We say that \mathcal{I} is trivial if it contains a constraint with an empty relation. Otherwise, we say that \mathcal{I} is non-trivial.

As in the finite case, one may transform an instance \mathcal{I} into an equivalent instance, i.e. with the same set of solutions by simply introducing at most $O(|\mathcal{V}|^l)$ new constraints so that the first condition in Definition 6 was satisfied and then by repeatedly removing orbits of tuples from constraints until the second condition is satisfied. Similarly to the finite CSP we have the following.

► **Proposition 7.** Let \mathbb{A} be an ω -categorical relational structure. Then for every instance \mathcal{I} of $\text{CSP}(\mathbb{A})$ and $l \geq k > 0$ there exists an instance \mathcal{I}' of the CSP with the same sets of solutions as \mathcal{I} such that \mathcal{I}' is (k, l) -minimal.

For fixed (k, l) and \mathbb{A} , the process of establishing (k, l) -minimality, i.e., transforming \mathcal{I} into \mathcal{I}' takes time $O(|\mathcal{V}|^m)$ where $m = \max(l, \text{arity}(\mathbb{A}))$ is the maximum of l and the greatest arity of a relation in \mathbb{A} . If \mathcal{I}' is trivial, then both \mathcal{I} and \mathcal{I}' have no solutions.

We are now ready to define the relational width.

► **Definition 8.** *We say that \mathbb{A} has relational width (k, l) if and only if \mathcal{I} has a solution provided any (k, l) -minimal instance of the CSP equivalent to \mathcal{I} is non-trivial. We say that \mathbb{A} has (relational) bounded width ² if there exist (k, l) such that \mathbb{A} has relational width (k, l) .*

Finite structures with bounded relational width admit an algebraic characterization [2]. It is known that a finite structure \mathbb{A} has bounded (relational) width if and only if it has a four-ary polymorphism f and a ternary polymorphism g that are weak near-unanimity operations and such that $f(y, x, x, x) = g(y, x, x)$ for all $x, y \in A$. We have a similar sufficient condition for reducts of finitely bounded homogeneous structures.

► **Theorem 9.** ([10]) *Let \mathbb{A} be a finite-signature reduct of a finitely bounded homogeneous structure \mathbb{B} . Suppose that \mathbb{A} has a four-ary polymorphism f and a ternary polymorphism g that are canonical with respect to $\text{Aut}(\mathbb{B})$ and are weak near-unanimity operations modulo $\overline{\text{Aut}(\mathbb{B})}$, and such that there are operations $e_1, e_2 \in \overline{\text{Aut}(\mathbb{B})}$ with $e_1(f(y, x, x, x)) = e_2(g(y, x, x))$ for all $x, y \in A$. Then $\text{CSP}(\mathbb{A})$ has bounded relational width.*

A slight change in the proof of the above theorem gives us the upper bound for relational width of infinite structures under consideration.

► **Corollary 10.** *Let \mathbb{A} be a finite-signature reduct of a finitely bounded homogeneous structure \mathbb{B} . Suppose that \mathbb{A} has a four-ary polymorphism f and a ternary polymorphism g that are canonical with respect to $\text{Aut}(\mathbb{B})$, that are weak near-unanimity operations modulo $\overline{\text{Aut}(\mathbb{A})}$, and such that there are operations $e_1, e_2 \in \overline{\text{Aut}(\mathbb{B})}$ with $e_1(f(y, x, x, x)) = e_2(g(y, x, x))$ for all $x, y \in A$. Then \mathbb{A} has relational width $(2m, 3m)$ where $m = \max(\text{arity}(\mathbb{A}) + 1, \text{arity}(\mathbb{B}) + 1, \mathbb{L}_{\mathbb{B}}, 3)$.*

We now use Corollary 10 to provide the upper bound of the relational width for reducts of homogeneous graphs preserved by binary canonical operations considered in [13, 14, 9].

► **Proposition 11.** *Let \mathbb{A} be a reduct of a countably infinite homogeneous graph \mathcal{H} preserved by a binary injection:*

1. *of behaviour \max which is either balanced or E -dominated, or*
2. *of behaviour \min which is either balanced or N -dominated, or*
3. *which is E -constant, or*
4. *which is N -constant.*

Then it has relational width $(2m, 3m)$ where $m = \max(\text{arity}(\mathbb{A}) + 1, \mathbb{L}_{\mathcal{H}}, 3)$.

In Section 6.1, we use our approach to show that the exact relational width of structures under consideration in Proposition 11 is $(2, \mathbb{L}_{\mathcal{H}})$. The same is proved for first-order expansions of homogeneous graphs with bounded strict width.

Strict width is defined as follows. A (k, l) -minimal instance \mathcal{I} of the CSP is called globally consistent, if every partial solution of \mathcal{I} can be extended to a total solution of \mathcal{I} .

► **Definition 12.** *We say that \mathbb{A} has strict width k if for some $l \geq k \geq 2$ all instances of $\text{CSP}(\mathbb{A})$ that are (k, l) -minimal are globally consistent. We say that \mathbb{A} has bounded strict width if it has strict width k for some k .³*

² We note that the definition of bounded width provided in [5] is equivalent to ours.

³ Our definition of strict width slightly varies from a definition in [5] but again both definitions are equivalent.

We have the following algebraic characterization of ω -categorical structures with bounded strict width.

► **Theorem 13.** [5, 4] *Let \mathbb{A} be an ω -categorical language. Then the following are equivalent.*

1. \mathbb{A} has strict width k .
2. \mathbb{A} has an oligopotent $(k+1)$ -ary quasi near-unanimity operation as a polymorphism.

For a $(2, k)$ -minimal instance over variables $\mathcal{V} = \{v_1, \dots, v_n\}$ we write $\mathcal{I}_{i,j}$ with $i, j \in [n]$ to denote a subset of $\{E, N, =\}$ such that the projection of all constraints having v_i, v_j in its scope to $\{v_i, v_j\}$ equals $\bigcup_{O \in \mathcal{I}_{i,j}} O$. We will say that an instance is *simple* if $|\mathcal{I}_{i,j}| = 1$ for all $i, j \in [n]$.

We will now show that a simple non-trivial $(2, \mathbb{L}_{\mathcal{H}})$ -instance of $\text{CSP}(\mathbb{A})$ for a first-order expansion \mathbb{A} of a homogeneous graph \mathcal{H} always has a solution and that this amount of consistency is necessary.

► **Observation 14.** *Let \mathcal{I} be a simple non-trivial $(2, \mathbb{L}_{\mathcal{H}})$ -minimal instance of the CSP equivalent to an instance of $\text{CSP}(\mathcal{H}')$ where \mathcal{H}' is the expansion of \mathcal{H} containing all orbitals pp-definable in \mathcal{H} . Then \mathcal{I} has a solution.*

On the other hand, for every homogeneous graph \mathcal{H} there exists a simple non-trivial $(1, \mathbb{L}_{\mathcal{H}})$ -minimal instance \mathcal{I}_1 equivalent to an instance of $\text{CSP}(\mathcal{H}')$ and a simple non-trivial $(2, \mathbb{L}_{\mathcal{H}} - 1)$ -minimal instance \mathcal{I}_2 equivalent to an instance of $\text{CSP}(\mathcal{H}')$ that have no solutions.

Proof. We start from proving the first part of the observation. Define Δ to be a finite structure over the domain consisting of variables $\{v_1, \dots, v_n\}$ in \mathcal{I} and the signature $\tau \subseteq \{E, N, =\}$ such that $(v_i, v_j) \in R^\Delta$ for $i, j \in [n]$ and $R \in \tau$ if φ_I contains a constraint $((v_i, v_j), R)$. Since \mathcal{I} is $(2, 3)$ -minimal, we have the following.

► **Observation 15.** *The binary relation $\sim := \{(v_i, v_j) \mid \mathcal{I}_{i,j} \subseteq \{=\}\} \cup \bigcup_{i \in [n]} \{(v_i, v_i)\}$ is an equivalence relation.*

We claim that there is an embedding from Δ / \sim to \mathcal{H}' . Assume the contrary. Since \mathcal{H} is finitely bounded, there exists G over variables $\{w_1, \dots, w_l\}$ in $\mathcal{F}_{\mathcal{H}'}$ such that G embeds into Δ / \sim and $l \leq \mathbb{L}_{\mathcal{H}}$. Since \mathcal{I} is $(2, \mathbb{L}_{\mathcal{H}})$ -minimal, there is a constraint C in \mathcal{I} whose scope contains $\{w_1, \dots, w_l\}$ and the corresponding relation is empty. It contradicts with the assumption that \mathcal{I} is non-trivial. Thus, Δ / \sim embeds into \mathcal{H} , and in consequence \mathcal{I} has a solution. It completes the proof of the first part of the observation.

For the second part of the observation, we select \mathcal{I}_1 to be $\{((v_1, v_2), E), ((v_1, v_2), N)\}$. Indeed, every subset of variables of \mathcal{I}_1 is in the scope of some constraint. The projection of each constraint to $\{v_1\}$ or $\{v_2\}$ is the set of all vertices in \mathcal{H} . It follows that \mathcal{I}_1 is $(1, \mathbb{L}_{\mathcal{H}})$ -consistent. Clearly \mathcal{I}_1 has no solutions.

We now turn to \mathcal{I}_2 . If $\mathbb{L}_{\mathcal{H}} > 3$ and $G = ([n], E)$ is a forbidden subgraph of size $n = \mathbb{L}_{\mathcal{H}}$ consider an instance \mathcal{I}'_2 over variables $\{v_1, \dots, v_n\}$ containing a constraint $((v_i, v_j), E)$ if $(i, j) \in E^G$ and a constraint $((v_i, v_j), N)$ if $(i, j) \notin E^G$. Let \mathcal{I}_2 be a $(2, \mathbb{L}_{\mathcal{H}} - 1)$ -minimal instance of the CSP equivalent to \mathcal{I}'_2 . By the minimality of $\mathcal{F}_{\mathcal{H}}$, we have that no induced subgraph of G is in $\mathcal{F}_{\mathcal{H}}$. It follows that \mathcal{I}_2 is non-trivial but, clearly, \mathcal{I}_2 has no solutions. If $\mathbb{L}_{\mathcal{H}} = 3$, then we select \mathcal{I}_2 to be an instance such that $\mathcal{C} = \{((v_1, v_2), =), ((v_2, v_3), =), ((v_1, v_3), E)\}$. It is again straightforward to check that \mathcal{I}_2 is $(2, 2)$ -minimal. Yet, it has no solutions. It completes the proof of the observation. ◀

We complete this section by giving some examples of first-order expansions of homogeneous graphs with bounded strict width.

► **Proposition 16.** *Let \mathbb{A} be a first-order expansion of the random graph $\mathcal{H} = (A; E)$ such that every relation in \mathbb{A} is pp-definable as a conjunction of clauses of the form:*

$$(x_1 \neq y_1 \vee \dots \vee x_k \neq y_1 \vee R(y_1, y_2) \vee y_2 \neq z_1 \vee \dots \vee y_2 \neq z_l),$$

where $R \in \{E, N\}$. Then \mathbb{A} has bounded strict-width.

And here comes another example.

► **Proposition 17.** *The constraint language $\mathbb{A} = (A; E, N, R)$ where $(A; E)$ is C_2^ω and $R(x_1, x_2, x_3) \equiv ((E(x_1, x_2) \wedge N(x_2, x_3)) \vee (N(x_1, x_2) \wedge E(x_2, x_3)))$ has bounded strict width.*

5 Conditions Sufficient for Low Relational Width

In order to show that a non-trivial $(2, \mathbb{L}_{\mathcal{H}})$ -minimal instance \mathcal{I} of $\text{CSP}(\mathbb{A})$ for first-order expansions \mathbb{A} of a homogeneous graph \mathcal{H} has a solution, we always use one scheme. We take advantage of the fact that certain quaternary and ternary relations are not pp-definable in \mathbb{A} and we carefully narrow down $\mathcal{I}_{i,j}$ for $i, j \in [n]$ so that we end up with a simple non-trivial instance \mathcal{I}' which is a 'subinstance' of \mathcal{I} in the following sense: for every $C = ((x_1, \dots, x_r), R)$ in \mathcal{I} we have $C' = ((x_1, \dots, x_r), R') \in \mathcal{I}'$ such that $R' \subseteq R$. Since \mathcal{I}' is simple, by Observation 14, it has a solution. This solution is clearly a solution to the original instance \mathcal{I} .

We shrink an instance of $\text{CSP}(\mathbb{A})$ using one of three different sets of relations presented in the three lemmas below.

► **Lemma 18.** *Let $\{O_1, O_2\}$ be $\{E, N\}$ and \mathbb{A} be a first-order expansion of a homogeneous graph \mathcal{H} such that none of the following types of relations is pp-definable in \mathbb{A} :*

1. $[(O_1(x_1, x_2) \implies \underline{O_2}(x_3, x_4))]$ -relations,
2. $[(O_1(x_1, x_2) \implies x_3 = x_4)]$ -relations, and
3. $[(O_2(x_1, x_2) \implies x_3 = x_4), (\underline{O_2}(x_1, x_2) \wedge \underline{O_2}(x_3, x_4))]$ -relations.

Then \mathbb{A} has relational width $(2, \mathbb{L}_{\mathcal{H}})$.

Before we discuss the 'shrinking' strategy that stands behind Lemma 18, consider a non-trivial instance \mathcal{I} of some \mathbb{A} under consideration in the lemma and a constraint $((x_1, \dots, x_r), R)$ for which there are i_1, j_1, i_2, j_2 such that $v_{i_1}, v_{j_1}, v_{i_2}, v_{j_2} \in \{x_1, \dots, x_r\}$ and $O_1 \in \mathcal{I}_{i_1, j_1}$, $O_1 \in \mathcal{I}_{i_2, j_2}$. Since \mathcal{I} is non-trivial and $(2, 3)$ -minimal the relation $(R'(x_1, \dots, x_r) \equiv (R(x_1, \dots, x_r) \wedge O_1(v_{i_1}, v_{j_1})))$ is non-empty. But also $(R''(x_1, \dots, x_r) \equiv (R(x_1, \dots, x_r) \wedge O_1(v_{i_1}, v_{j_1}) \wedge O_1(v_{i_2}, v_{j_2})))$ is non-empty. Indeed, otherwise since $O_1 \cup \underline{O_2} = A^2$, the structure \mathbb{A} would define a relation from Item 1 or Item 2. Generalizing the argument, one can easily transform \mathcal{I} to a non-trivial \mathcal{I}' where every $\mathcal{I}'_{i,j} = \{O_1\}$ whenever $\mathcal{I}_{i,j}$ contains O_1 . Using a similar reasoning and Item 3, and taking care of some details, we have to skip here, one can then transform \mathcal{I}' to \mathcal{I}'' so that $\mathcal{I}''_{i,j} = O_2$ whenever $\mathcal{I}'_{i,j}$ contains O_2 . Since \mathcal{I}'' is simple and non-trivial, we can use Observation 14 to argue that both \mathcal{I}'' and \mathcal{I} has a solution.

The next lemma considers a specific situation where \mathcal{H} is a disjoint sum of ω edges and languages under consideration are preserved by oligopotent qnu-operations.

► **Lemma 19.** *Let \mathbb{A} be a first-order expansion of C_ω^2 preserved by an oligopotent qnu-operation and such that none of the following types of relations is pp-definable in \mathbb{A} :*

1. $[(N(x_1, x_2) \implies \underline{E}(x_3, x_4))]$ -relations,
2. $[(N(x_1, x_2) \implies \underline{E}(x_3, x_4)), (\underline{E}(x_3, x_4))]$ -relations,
3. $[(N(x_1, x_2) \implies x_3 = x_4)]$ -relations,

4. $[(O_1(x_1, x_2) \implies O_2(x_3, x_4)), (\underline{E}(x_1, x_2) \wedge N(x_2, x_3) \wedge \underline{E}(x_3, x_4))]$ -relations where the set $\{O_1, O_2\}$ equals $\{E, =\}$.
Then \mathbb{A} has relational width $(2, 3)$.

The shrinking strategy for Lemma 19 is as follows. We start with a non-trivial $(2, 3)$ -minimal instance \mathcal{I} and use Items 1–3 to transform it into a non-trivial $(2, 3)$ -minimal \mathcal{I}' such that $\mathcal{I}'_{i,j} = \{N\}$ whenever $\mathcal{I}_{i,j}$ contains N . Since \underline{E} fo-definable in C_ω^2 is transitive and \mathcal{I}' is $(2, 3)$ -minimal, it is easy to show that the graph over variables $\{v_1, \dots, v_n\}$ and edges $\mathcal{I}'_{i,j}$ with $i, j \in [n]$ is a disjoint union of components K_1, \dots, K_κ such that for all $k \in [\kappa]$ and all $v_i, v_j \in K_k$ it holds that $\mathcal{I}'_{i,j} \subseteq \{E, =\}$ and whenever v_i, v_j are in different components, then $\mathcal{I}'_{i,j} = \{N\}$. Now, any \mathcal{I}'_{K_i} — the instance \mathcal{I}' restricted to variables in K_i , which is in fact an instance of $\text{CSP}(\Delta)$ for Δ over two-elements (some edge in C_ω^2 , different for every $i \in [\kappa]$) preserved by a near-unanimity operation, is shown to have a solution \mathbf{s}_i . It follows by the characterization of relational width for finite structure. In order to prove that solution $\mathbf{s} := \bigcup_{i \in [\kappa]} \mathbf{s}_i$ is the solution to \mathcal{I}' and hence to \mathcal{I} we use the fact that relations from Item 4 are not pp-definable in \mathbb{A} .

Finally, we turn to the case where \mathcal{H} is a disjoint sum of two infinite cliques and the structures \mathbb{A} have oligopotent qnu-operations as polymorphisms.

► **Lemma 20.** *Let \mathbb{A} be a first-order expansion of C_ω^2 preserved by an oligopotent qnu-operation and such that \mathbb{A} pp-defines neither \underline{N} nor $[(O(x_1, x_2) \rightarrow x_3 = x_4)]$ for any $O \in \{E, N\}$. Then \mathbb{A} has relational width $(2, 3)$.*

Clearly any tuple over C_ω^2 takes some of its values from one equivalence class in C_ω^2 and the remaining values from the other class. In order to prove Lemma 20, we consider a non-trivial $(2, 3)$ -minimal instance \mathcal{I} of $\text{CSP}(\mathbb{A})$ but this time we also assume without loss of generality that there are no i, j with $\mathcal{I}_{i,j} = \{=\}$. Since \mathbb{A} does not define \underline{N} we have that for all $i, j \in [n]$ the set $\mathcal{I}_{i,j}$ contains E . Then we transform \mathcal{I} to \mathcal{I}_B of $\text{CSP}(\Delta)$ where Δ is over the domain $\{0, 1\}$ by replacing any tuple t in any relation in any constraint in \mathcal{I} by a tuple over $\{0, 1\}$ so that all values in one equivalence class are replaced by 0 and all values in the other equivalence class are replaced by 1. Since Δ is preserved by a near-unanimity operation, and hence has bounded relational width we have that the $(2, 3)$ -minimal \mathcal{I}_B has a solution $\mathbf{s}_B : \{v_1, \dots, v_m\} \rightarrow \{0, 1\}$. We use \mathbf{s}_B to transform \mathcal{I} to \mathcal{I}' so that we set $\mathcal{I}'_{i,j}$ to $\{N\}$ whenever $\mathbf{s}_B(v_i) \neq \mathbf{s}_B(v_j)$. No $[(N(x_1, x_2) \rightarrow x_3 = x_4)]$ -relations are pp-definable in \mathbb{A} , and hence we have that $\mathcal{I}'_{i,j}$ for any $i, j \in [n]$ contains E . Since \mathbb{B} pp-defines no $[(E(x_1, x_2) \rightarrow x_3 = x_4)]$ -relations we may transform \mathcal{I}' into \mathcal{I}'' so that $\mathcal{I}''_{i,j}$ is E whenever $\mathcal{I}'_{i,j} \neq \{N\}$. Thus, \mathcal{I}'' is a simple non-trivial $(2, 3)$ -minimal instance. It follows by Observation 14 that both \mathcal{I}'' and \mathcal{I} have a solution. Again, we skipped many details but our goal was rather to convey some intuitions that stand behind the proofs of the lemmas in this section.

6 Constraint Languages with Low Relational Width

In this section we employ lemmas from Section 5 to provide the exact characterization of relational width of first-order expansions of homogeneous graphs with bounded strict width and first-order expansions of homogeneous graphs preserved by binary canonical operations from Proposition 11. In order to prove the former, we also show which quaternary relations of interest are violated by ternary injections used in the complexity classification (see Subsection 6.2). To rule out some other relations, we have to use oligopotent qnu-operations directly (see Subsection 6.3).

6.1 Binary Injections and Low Relational Width

We start with first-order expansions \mathbb{A} of homogeneous graphs \mathcal{H} whose tractability has been shown in Proposition 8.22 in [13], Proposition 6.2 in [14] as well as Proposition 37 and Theorem 62 in [9].

► **Lemma 21.** *Let \mathbb{A} be a first-order expansion of a countably infinite homogeneous graph \mathcal{H} preserved by a binary injection:*

1. *of behaviour max which is either balanced or E -dominated, or*
2. *of behaviour min which is either balanced or N -dominated, or*
3. *which is E -constant, or*
4. *which is N -constant.*

Then \mathbb{A} has relational width $(2, \mathbb{L}_{\mathcal{H}})$.

The above lemma gives an opportunity to reformulate the dichotomy results for reducts \mathbb{A} of $C_{\omega}^1, C_1^{\omega}, C_{\omega}^{\omega}$ and H_k for any $k \geq 3$.

► **Corollary 22.** *Let \mathbb{A} be a reduct of a homomorphism graph \mathcal{H} which is $C_{\omega}^1, C_1^{\omega}, C_{\omega}^{\omega}$ or H_k for any $k \geq 3$. Then either $\text{CSP}(\mathbb{A})$ is NP-complete or \mathbb{A} has relational width $(2, \mathbb{L}_{\mathcal{H}})$.*

Proof. We have that any tractable first-order expansion of $(\mathbb{N}; =, \neq)$ is preserved by a binary injection [7]. It follows that every tractable reduct of C_1^{ω} is either preserved by a constant operation or is a first-order expansion of C_1^{ω} and preserved by a binary injection which is of behaviour max and E -dominated. A similar reasoning holds for reducts of C_{ω}^1 with a difference that we replace E with N . Further, every reduct of C_{ω}^{ω} is either homomorphically equivalent to a reduct of $(\mathbb{N}; =)$ or pp-defines both E and N , see Theorem 4.5 [14]. In the former case we are done, while in the latter a tractable \mathbb{A} is preserved by a binary injection of behaviour min and balanced, Corollary 7.5 in [14]. The corollary follows by Lemma 21. By Proposition 15 and Lemma 17 in [9], a tractable reduct of H_k with $k \geq 3$ is either homomorphically equivalent to a reduct of $(\mathbb{N}; =)$ or pp-defines both E and N . In the former case we are done while in the latter, we have that $\text{CSP}(\mathbb{A})$ is in P when it is preserved by a binary injection of behaviour min and N -dominated (see Theorem 38 in [9]). Again, the corollary follows by Lemma 21. ◀

6.2 Types of Relations violated by Ternary Canonical Operations

Here we look at quaternary relations of interest violated by canonical ternary operations. We start with ternary injections of behaviour majority.

► **Lemma 23.** *Let \mathbb{A} be a reduct of a countably infinite homogeneous graph preserved by a ternary injection of behaviour majority which additionally is:*

- *hyperplanely balanced and of behaviour projection, or*
- *hyperplanely E -constant, or hyperplanely N -constant, or*
- *hyperplanely of behaviour max and E -dominated, or*
- *hyperplanely of behaviour min and N -dominated.*

Then \mathbb{A} pp-defines no $[(O(x_1, x_2) \implies x_3 = x_4)]$ -relations with $O \in \{E, N\}$.

We continue with ternary injections of behaviour minority.

► **Lemma 24.** *Let \mathbb{A} be a reduct of a countably infinite homogeneous graph preserved by a ternary injection of behaviour minority which additionally is:*

- *hyperplanely balanced and of behaviour projection,*

- hyperplanely of behaviour projection and E -dominated, or
- hyperplanely of behaviour projection and N -dominated, or
- hyperplanely balanced of behaviour xnor , or
- hyperplanely balanced of behaviour xor .

Then \mathbb{A} does not pp-define $[(O(x_1, x_2) \implies x_3 = x_4)]$ -relations with $O \in \{E, N\}$.

We are already in the position to prove that another large family of $\text{CSP}(\mathbb{A})$ under consideration may be solved by establishing minimality.

► **Corollary 25.** *Let \mathbb{A} be a first-order expansion of C_2^ω preserved by a canonical ternary injection of behaviour minority which is hyperplanely balanced of behaviour xnor and an oligopotent qnu-operation. Then \mathbb{A} has relational width $(2, 3)$.*

Proof. By Lemma 24, the structure \mathbb{A} does not pp-define $[(O(x_1, x_2) \implies x_3 = x_4)]$ with $O \in \{E, N\}$. Since a canonical ternary injection of behaviour minority which is hyperplanely balanced of behaviour xnor does not preserve $\underline{\underline{N}}$, the result follows by appealing to Lemma 20. ◀

The third lemma of this subsection takes care of the third kind of ternary operations that occurs in complexity classifications of CSPs for reducts of homogeneous graphs.

► **Lemma 26.** *Let \mathbb{A} be a reduct of a countably infinite homogeneous graph preserved by a ternary canonical operation h with $h(N, \cdot, \cdot) = h(\cdot, N, \cdot) = h(\cdot, \cdot, N) = N$ and which behaves like a minority on $\{E, =\}$, i.e., h satisfies the behaviour B such that $B(E, E, E) = B(E, =, =) = B(=, E, =) = B(=, =, E) = E$ and $B(=, =, =) = B(=, E, E) = B(E, =, E) = B(E, E, =)$ equals $=$. Then \mathbb{A} pp-defines none of the following types of relations:*

- $[(N(x_1, x_2) \implies \underline{E}(x_3, x_4))]$ -relations,
- $[(N(x_1, x_2) \implies x_3 = x_4)]$ -relations,
- $[(N(x_1, x_2) \implies E(x_3, x_4)), (\underline{E}(x_3, x_4))]$ -relations.

6.3 Types of Relations violated by Oligopotent QNUs

Here we provide a list of quaternary relations of interest violated by ternary canonical operations and oligopotent qnu-operations. We start with the case where the considered homogeneous graph is the random graph.

► **Lemma 27.** *Let \mathbb{A} be a first-order expansion of the random graph preserved by a ternary injection of behaviour majority which additionally satisfies one of the conditions in Lemma 23 or of behaviour minority which additionally satisfies one of the conditions in Lemma 24, and an oligopotent qnu-operation. Then \mathbb{A} pp-defines at most one of the following:*

1. either a $[(E(x_1, x_2) \implies \underline{\underline{N}}(x_3, x_4))]$ -relation or
2. a $[(N(x_1, x_2) \implies \underline{E}(x_3, x_4))]$ -relation.

Here comes the corollary.

► **Corollary 28.** *Let \mathbb{A} be a first-order expansion of the random graph preserved by a ternary injection of behaviour majority which additionally satisfies one of the conditions in Lemma 23 or of behaviour minority which additionally satisfies one of the conditions in Lemma 24 and an oligopotent qnu-operation. Then \mathbb{A} has relational width $(2, 3)$.*

Proof. By appeal to Lemma 27, it follows that there are $\{O_1, O_2\} = \{E, N\}$ such that \mathbb{A} does not pp-define a $[(O_1(x_1, x_2) \implies \underline{O_2}(x_3, x_4))]$ -relation. By Lemmas 23 and 24, \mathbb{A} pp-defines neither $[(O_1(x_1, x_2) \implies x_3 = x_4)]$ -relations nor $[(O_2(x_1, x_2) \implies x_3 = x_4)]$ -relations. Since $\mathbb{L}_{\mathcal{H}}$ in the case where \mathcal{H} is the random graph equals 3, the result follows by Lemma 18. ◀

We now turn to the case where the considered homogeneous graph is the disjoint union of ω edges.

► **Lemma 29.** *Let \mathbb{A} be a first-order expansion of C_ω^2 preserved by a ternary canonical operation h with $h(N, \cdot, \cdot) = h(\cdot, N, \cdot) = h(\cdot, \cdot, N) = N$ and which behaves like a minority on $\{E, =\}$ and an oligopotent qnu-operation. Then \mathbb{A} pp-defines neither*

- *a $[(E(x_1, x_2) \implies (x_3 = x_4)), (\underline{E}(x_1, x_2) \wedge N(x_2, x_3) \wedge \underline{E}(x_3, x_4))]$ -relation nor*
- *a $[(x_1 = x_2) \implies E(x_3, x_4)], (\underline{E}(x_1, x_2) \wedge N(x_2, x_3) \wedge \underline{E}(x_3, x_4))]$ -relation.*

Then we provide another similar lemma.

► **Lemma 30.** *Let \mathbb{A} be a first-order expansion of C_ω^2 preserved by a ternary canonical operation h with $h(N, \cdot, \cdot) = h(\cdot, N, \cdot) = h(\cdot, \cdot, N) = N$ and which behaves like a minority on $\{E, =\}$ and an oligopotent qnu-operation. Then \mathbb{A} pp-defines neither*

- *a $[(E(x_1, x_2) \implies E(x_3, x_4)), (\underline{E}(x_1, x_2) \wedge N(x_2, x_3) \wedge \underline{E}(x_3, x_4))]$ -relation nor*
- *a $[(x_1 = x_2) \implies (x_3 = x_4)], (\underline{E}(x_1, x_2) \wedge N(x_2, x_3) \wedge \underline{E}(x_3, x_4))]$ -relation.*

Then comes the corollary.

► **Corollary 31.** *Let \mathbb{A} be a first-order expansion of C_ω^2 preserved by a ternary canonical operation h with $h(N, \cdot, \cdot) = h(\cdot, N, \cdot) = h(\cdot, \cdot, N) = N$ and which behaves like a minority on $\{E, =\}$ and an oligopotent qnu-operation. Then \mathbb{A} has relational width $(2, 3)$.*

Proof. By Lemmas 26, 29 and 30 none of the types of relations mentioned in Lemma 19 is pp-definable in \mathbb{A} . Appealing to Lemma 19 completes the proof of the corollary. ◀

6.4 The Main Result

Here we prove our main result.

► **Theorem 32.** *Let \mathbb{A} be a first-order expansion of a countably infinite homogeneous graph \mathcal{H} which has bounded strict width. Then \mathbb{A} has relational width $(2, \mathbb{L}_{\mathcal{H}})$.*

Proof. By the classification of Lachlan and Woodrow [24], we have that \mathcal{H} is either the random graph, a Henson graph H_k with a forbidden k -clique where $k \geq 3$, a disjoint set of n cliques of size s denoted by C_n^s or a complement of either C_n^s or H_k . The case where \mathcal{H} is C_1^ω , C_ω^1 , C_ω^ω or H_k with $k \geq 3$ follows by Corollary 22. If C_n^s is such that $3 \leq n < \omega$ or $3 \leq s < \omega$, then by Theorem 60 in [9], a first-order expansion \mathbb{A} of C_n^s is either homomorphically equivalent to a reduct of $(\mathbb{N}; =)$ or is not preserved by an oligopotent qnu-operation and we are done. If \mathbb{A} is a first-order expansion of C_2^ω , then by Theorem 61 in [9] either it is homomorphically equivalent to a reduct of $(\mathbb{N}; =)$ or is not preserved by an oligopotent qnu-operation or pp-defines both E and N and is preserved by a canonical ternary injection of behaviour minority which is hyperplanely balanced of behaviour xnor and then \mathbb{A} has relational width $(2, 3)$ by Corollary 25. If \mathbb{A} is a first-order expansion of C_ω^2 , then by Theorem 62 in [9], we have that either \mathbb{A} is homomorphically equivalent to a reduct of $(\mathbb{N}; =)$, or it is not preserved by an oligopotent qnu-operation or it pp-defines both E and N and is preserved by a canonical binary injection of behaviour min that is N -dominated or a ternary canonical operation h with $h(N, \cdot, \cdot) = h(\cdot, N, \cdot) = h(\cdot, \cdot, N) = N$ and which behaves like a minority on $\{E, =\}$. In the former case the language \mathbb{A} has relational width $(2, 3)$ by Lemma 21, in the latter by Corollary 31.

The remaining case is where \mathbb{A} is a first-order expansion of the random graph G preserved by an oligopotent qnu-operation. By Theorem 6.1 in [13] we have that a first-order expansion of G is either homomorphically equivalent to a reduct of $(\mathbb{N}; =)$ and then we are done or pp-defines both E and N in which case, by Theorem 9.3 in [13], we have that:

- \mathbb{A} is preserved by a binary injection of behaviour max which is either balanced or E-dominated, by a binary injection of behaviour min which is either balanced or N -dominated, by a binary injection which is E-constant, or a binary injection which is N -constant, and then the theorem follows by Lemma 21, or
- \mathbb{A} is preserved by a ternary injection of behaviour majority which additionally satisfies one of the conditions in Lemma 23 or of behaviour minority which additionally satisfies one of the conditions in Lemma 24, and then the theorem holds by Corollary 28.

It completes the proof of the theorem. ◀

7 Summary and Future Work

In this paper we proved in particular that:

1. every first-order expansion of a homogeneous graph \mathcal{H} preserved by a canonical binary operation considered in [13, 14, 9] and
2. every first-order expansion of a homogeneous graph \mathcal{H} with bounded strict width has relational width exactly $(2, \mathbb{L}_{\mathcal{H}})$.

A nice consequence of the former result is that all tractable reducts of $C_{\omega}^1, C_1^{\omega}, C_{\omega}^{\omega}$ and H_k with $k \geq 3$ have relational width exactly $(2, \mathbb{L}_{\mathcal{H}})$, and thus all tractable $\text{CSP}(\mathbb{A})$ may be solved by establishing $(2, \mathbb{L}_{\mathcal{H}})$ -minimality. Nevertheless, we find the latter result to be the main result of this paper. It is for the following reason.

Our general strategy is that we show that constraint languages \mathbb{A} under consideration do not express ‘too many implications’, i.e., quaternary relations that efficiently entail formulas of the form $(R_1(x_1, x_2) \implies R_2(x_3, x_4))$, see definitions in Section 2.4 and lemmas in Section 5 and then use these facts in order to find a strategy of how to shrink a non-trivial $(2, \mathbb{L}_{\mathcal{H}})$ -minimal instance of the CSP so that it became a simple instance. In this paper, in order to show that certain relations are not pp-definable in \mathbb{A} we employ in particular some binary and ternary canonical operations. We believe that it is not in fact necessary and theorems analogous to Theorem 32 may be obtained for large families of constraint languages using only the fact that structures \mathbb{A} under consideration are preserved by oligopotent qnu-operations. Thereby we believe that Question 2 may be answered in full generality.

References

- 1 Libor Barto. The collapse of the bounded width hierarchy. *J. Log. Comput.*, 26(3):923–943, 2016.
- 2 Libor Barto and Marcin Kozik. Constraint satisfaction problems solvable by local consistency methods. *J. ACM*, 61(1):3:1–3:19, 2014. An extended abstract appeared in the Proceedings of the Symposium on Foundations of Computer Science (FOCS’09).
- 3 Libor Barto and Michael Pinsker. The algebraic dichotomy conjecture for infinite domain constraint satisfaction problems. In *Proceedings of the 31st Annual ACM/IEEE Symposium on Logic in Computer Science, (LICS)*, pages 615–622, 2016.
- 4 Manuel Bodirsky and Hubert Chen. Oligomorphic clones. *Algebra Universalis*, 57(1):109–125, 2007.
- 5 Manuel Bodirsky and Víctor Dalmau. Datalog and constraint satisfaction with infinite templates. *J. Comput. Syst. Sci.*, 79(1):79–100, 2013.
- 6 Manuel Bodirsky and Peter Jonsson. A model-theoretic view on qualitative constraint reasoning. *J. Artif. Intell. Res. (JAIR)*, 58:339–385, 2017.
- 7 Manuel Bodirsky and Jan Kára. The complexity of equality constraint languages. *Theory Comput. Syst.*, 43(2):136–158, 2008.

- 8 Manuel Bodirsky and Jan Kára. The complexity of temporal constraint satisfaction problems. *Journal of the ACM*, 57(2):1–41, 2009. An extended abstract appeared in the Proceedings of the Symposium on Theory of Computing (STOC’08).
- 9 Manuel Bodirsky, Barnaby Martin, Michael Pinsker, and András Pongrácz. Constraint satisfaction problems for reducts of homogeneous graphs. *CoRR*, abs/1602.05819, 2016. An extended abstract appeared in the Proceedings of the International Colloquium on Automata, Languages, and Programming (ICALP’16).
- 10 Manuel Bodirsky and Antoine Mottet. A dichotomy for first-order reducts of unary structures. *Logical Methods in Computer Science*, 14(2), 2018. A preliminary version of the paper appeared in the proceedings of LICS’16.
- 11 Manuel Bodirsky and Jaroslav Nešetřil. Constraint satisfaction with countable homogeneous templates. *J. Log. Comput.*, 16(3):359–373, 2006.
- 12 Manuel Bodirsky and Michael Pinsker. Reducts of Ramsey structures. *AMS Contemporary Mathematics*, vol. 558 (Model Theoretic Methods in Finite Combinatorics), pages 489–519, 2011.
- 13 Manuel Bodirsky and Michael Pinsker. Schaefer’s theorem for graphs. *J. ACM*, 62(3):19:1–19:52, 2015. An extended abstract appeared in the Proceedings of the Symposium on Theory of Computing (STOC’11).
- 14 Manuel Bodirsky and Michał Wrona. Equivalence constraint satisfaction problems. In *Proceedings of Computer Science Logic*, volume 16 of *LIPICS*, pages 122–136. Dagstuhl Publishing, September 2012.
- 15 Andrei A. Bulatov. Bounded relational width. manuscript, 2009.
- 16 Andrei A. Bulatov. A dichotomy theorem for nonuniform csps. In *Proceedings of the Symposium on Foundations of Computer Science (FOCS)*, pages 319–330. IEEE Computer Society, 2017.
- 17 Andrei A. Bulatov, Andrei A. Krokhin, and Peter G. Jeavons. Classifying the complexity of constraints using finite algebras. *SIAM Journal on Computing*, 34:720–742, 2005. A preliminary version of the paper appeared in the proceedings of the Symposium on Logic in Computer Science (ICALP’00).
- 18 Peter J. Cameron. *Oligomorphic Permutation Groups*. Cambridge University Press, Cambridge, 1990.
- 19 Rina Dechter. From local to global consistency. *Artificial Intelligence*, 55(1):87–108, 1992.
- 20 Tomás Feder and Moshe Y. Vardi. The computational structure of monotone monadic SNP and constraint satisfaction: a study through Datalog and group theory. *SIAM Journal on Computing*, 28:57–104, 1999. A preliminary version appeared in the proceedings of the Symposium on Theory of Computing (STOC’93).
- 21 Wilfrid Hodges. *A shorter model theory*. Cambridge University Press, Cambridge, 1997.
- 22 Peter Jeavons, David Cohen, and Marc Gyssens. Closure properties of constraints. *Journal of the ACM*, 44(4):527–548, 1997.
- 23 Michael Kompatscher and Van Trung Pham. A complexity dichotomy for poset constraint satisfaction. In *34th Symposium on Theoretical Aspects of Computer Science, STACS’17*, pages 47:1–47:12, 2017.
- 24 A. H. Lachlan and Robert E. Woodrow. Countable ultrahomogeneous undirected graphs. *Trans. Amer. Math. Soc.*, 262:51–94, 1980.
- 25 Jochen Renz. Implicit constraints for qualitative spatial and temporal reasoning. In *Principles of Knowledge Representation and Reasoning: Proceedings of the Thirteenth International Conference, KR’12*, 2012.
- 26 Dmitriy Zhuk. A proof of CSP dichotomy conjecture. In *Proceedings of the Symposium on Foundations of Computer Science (FOCS)*, pages 331–342. IEEE Computer Society, 2017.

A Proof of Proposition 7

The algorithm of establishing (k, l) -minimality is quite straightforward and adopted from the world of finite CSP. First for every l -element subset $\{v_{i_1}, \dots, v_{i_l}\}$ of the variables $\mathcal{V} = \{v_1, \dots, v_n\}$ in φ_I that does not occur in any constraint in φ_I we introduce a constraint $((v_{i_1}, \dots, v_{i_l}), A^l)$. Then until the process stabilizes, for every $W \subseteq \mathcal{V}$ of size at most k and every two constraints $((v_{i_1}, \dots, v_{i_l}), R_1), ((v_{j_1}, \dots, v_{j_k}), R_2)$ such that W is a subset of both $\{x_1, \dots, x_a\}$ and $\{y_1, \dots, y_b\}$ we replace R_1 with $R'_1(v_{i_1}, \dots, v_{i_l}) \equiv \exists z_1 \dots \exists z_c R_1(v_{i_1}, \dots, v_{i_l}) \wedge R_2(v_{j_1}, \dots, v_{j_k})$ where $\{z_1, \dots, z_c\} = \{v_{i_1}, \dots, v_{i_l}\} \setminus \{v_{j_1}, \dots, v_{j_k}\}$; and we replace R_2 with $R'_2(v_{j_1}, \dots, v_{j_k}) \equiv \exists z_1 \dots \exists z_c R_1(v_{i_1}, \dots, v_{i_l}) \wedge R_2(v_{j_1}, \dots, v_{j_k})$ where $\{z_1, \dots, z_c\} = \{v_{i_1}, \dots, v_{i_l}\} \setminus \{v_{j_1}, \dots, v_{j_k}\}$.

We introduce $O(|\mathcal{V}|^l)$ new constraints and then we do the replacement at most the number of constraints multiplied by the number of pairwise different orbits of tuples of the length of the longest constraint. Indeed, in every step of the algorithm at least one orbit from at least one constraint is removed. Since there are at most $O(|\mathcal{V}|^l)$ constraints and the length of every constraint is bounded by a constant: $\max(l, m)$ where m is the largest arity in the signature of \mathbb{A} . It follows that for fixed (k, l) the process of establishing (k, l) -minimality may be performed in time $O(|\mathcal{V}|^l)$. \square

B Proof of Corollary 10

Here we use notions and results from [10]. Let (b_1, \dots, b_m) be a tuple of elements in \mathbb{B} . A quantifier-free (qf-) type of (b_1, \dots, b_m) , also called an m -type in \mathbb{B} , is the set of all quantifier-free formulas $\phi(z_1, \dots, z_m)$ such that $\mathbb{B} \models \phi(b_1, \dots, b_m)$. The structures \mathbb{B} under considerations have finite relational signatures, and hence there are only finitely many m -types in \mathbb{B} . The set of all m -types in \mathbb{B} will be denoted by $T_{\mathbb{B}, m}$.

Define now $T_{\mathbb{B}, m}(\mathbb{A})$ to be a finite structure whose elements are the m -types $T_{\mathbb{B}, m}$ of \mathbb{B} and relations are as follows.

- We have a unary constraint for each relation R of \mathbb{A} of arity r with a definition $\chi(z_1, \dots, z_r)$ in \mathbb{B} . For $i : [r] \rightarrow [m]$ we write $\langle \chi(z_{i(1)}, \dots, z_{i(r)}) \rangle$ to denote the unary relation that consists of all the types that contain $\chi(z_{i(1)}, \dots, z_{i(r)})$, and add all such relations to $T_{\mathbb{B}, m}(\mathbb{A})$ ⁴.
- We also have a binary relation $\text{Comp}_{i,j}$ for all $r \in [m]$ and $i, j : [r] \rightarrow [m]$. Define $\text{Comp}_{i,j}$ to be the binary relation that contains the pairs (p, q) of m -types such that for every quantifier-free formula $\chi(z_1, \dots, z_s)$ of \mathbb{B} and $t : [s] \rightarrow [r]$, the formula $\chi(z_{it(1)}, \dots, z_{it(s)})$ is in p iff $\chi(z_{jt(1)}, \dots, z_{jt(s)})$ is in q .

In the proof of Theorem 3.1 in [10], an instance \mathcal{I}_A of $\text{CSP}(\mathbb{A})$ over variables $\mathcal{V} = \{v_1, \dots, v_n\}$ is transformed into an instance \mathcal{I}_T of $\text{CSP}(T_{\mathbb{B}, m}(\mathbb{A}))$ over variables that are increasing functions I from $[m]$ to \mathcal{V} . The intention is that $h : \mathcal{V} \rightarrow A$ is a solution to \mathcal{I}_A if and only if $g : I \rightarrow T_{\mathbb{B}, m}$ such that $g(v)$ is the type of $(h(v(1)), \dots, h(v(m)))$ is a solution to \mathcal{I}_T .

- The instance \mathcal{I}_T has a unary constraint for every constraint $((j(1), \dots, j(r)), R)$ in \mathcal{I}_A where R is a relation of \mathbb{A} with a qf-definition $\chi(z_1, \dots, z_r)$ over \mathbb{B} and $j : [r] \rightarrow V$. Let $v \in I$ be such that $\text{Im}(j) \subseteq \text{Im}(v)$ and U be the relation symbol of $T_{\mathbb{B}, m}(\mathbb{A})$ that denotes the unary relation $\langle \chi(z_{v^{-1}j(1)}, \dots, z_{v^{-1}j(r)}) \rangle$. We then add $((v), U)$ to \mathcal{I}_T .

⁴ As in [10], we use functions to index tuples.

- Furthermore, \mathcal{I}_T has a binary constraint $((v, s), \text{Comp}_{v^{-1}k, s^{-1}k})$ for all $v, s \in I$ and bijections $k : [r] \rightarrow \text{Im}(v) \cap \text{Im}(s)$.

By the proof of Theorem 3.1 in [10] we have the following.

► **Fact 33.** *The assignment $h : \mathcal{V} \rightarrow A$ is a solution to \mathcal{I}_A if and only if $g : I \rightarrow T_{\mathbb{B},m}$ such that $g(v)$ is the type of $(h(v(1), \dots, h(v(m)))$ is a solution to \mathcal{I}_T .*

By Lemma 4.7 in [10], the structure $T_{\mathbb{B},m}(\mathbb{A})$ has a ternary and a four-ary polymorphism required for bounded width. It follows that $T_{\mathbb{B},m}(\mathbb{A})$ has bounded width $(2, 3)$, and hence $\text{CSP}(T_{\mathbb{B},m}(\mathbb{A}))$ is solvable by establishing $(2, 3)$ -minimality. For the proof of the corollary we will show that the algorithm of establishing minimality transforms an instance \mathcal{I}_A of $\text{CSP}(\mathbb{A})$ to a non-trivial $(2m, 3m)$ -minimal instance iff the algorithm of establishing minimality transforms an instance \mathcal{I}_T of $\text{CSP}(T_{\mathbb{B},m}(\mathbb{A}))$ to a $(2, 3)$ -minimal instance.

The right-to-left implication is easy to observe. Indeed, if an instance \mathcal{I}_T of $\text{CSP}(T_{\mathbb{B},m}(\mathbb{A}))$ has a $(2, 3)$ -minimal non-trivial equivalent instance of the CSP, then \mathcal{I}_T has a solution. By Fact 33, we have that \mathcal{I}_A has also a solution and hence there is a $(2m, 3m)$ -minimal non-trivial instance of the CSP equivalent to \mathcal{I}_A .

For the left-to-right implication consider a non-trivial $(2m, 3m)$ -minimal instance \mathcal{I}'_A of the CSP equivalent to \mathcal{I}_A that is obtained from \mathcal{I}_A by first adding an extra constraint $((j(1), \dots, j(r)), A^r)$ for all $m \leq r \leq 3m$ and all $j : [r] \rightarrow [m]$ and then establishing $(2m, 3m)$ -minimality. In consequence we obtain a $(2m, 3m)$ -minimal instance with the following constraints:

- $((j(1), \dots, j(r)), R')$ for every constraint of the form $((j(1), \dots, j(r)), R)$ in \mathcal{I}_A where $R' \subseteq R$, and
- a constraint $((j(1), \dots, j(r)), S')$ for all $m \leq r \leq 3m$ and $j : [r] \rightarrow \mathcal{V}$ originating from a constraint $((j(1), \dots, j(r)), A^r)$.

The next step is to construct \mathcal{I}'_T from \mathcal{I}'_A and \mathcal{I}_T . To this end, for $u_1, \dots, u_k \in I$ and a constraint $C = (\bar{x}, R)$ whose scope contain $\text{Im}(u_1), \dots, \text{Im}(u_k)$ we define $\text{Types}(C, u_1, \dots, u_k)$ to be a set of k -tuples of m -types which for every tuple $t \in R$ contains $(\text{type}_1, \dots, \text{type}_k)$ where type_i is the type of the projection of t to $\text{Im}(u_i)$. We are now ready to define \mathcal{I}'_T .

1. For a constraint $C = ((j(1), \dots, j(m)), S')$ in \mathcal{I}'_A we have $(j, \text{Types}(C, j))$ in \mathcal{I}'_T .
2. For a constraint $C = ((j(1), \dots, j(r)), S')$ with $m < r \leq 2m$ and all $u, v \in I$ with $\text{Im}(u), \text{Im}(v) \subseteq \text{Im}(j)$ we have a binary constraint $((u, v), \text{Types}(C, u, v))$ in \mathcal{I}'_T .
3. For a constraint $C = ((j(1), \dots, j(r)), S')$ with $m + 2 < r \leq 3m$ and all $u, v, w \in I$ with $\text{Im}(u), \text{Im}(v), \text{Im}(w) \subseteq \text{Im}(j)$ we have a ternary constraint $((u, v, w), (\text{Types}(C, u, v, w)))$ in \mathcal{I}'_T .
4. Let $((v), U)$ be a unary constraint in \mathcal{I}_T and $(j, \text{Types}(C, j))$ a constraint from Item 1 above. Then \mathcal{I}'_T contains $((v), U \cap \text{Types}(C, j))$.
5. Let $((v, s), \text{Comp}_{v^{-1}k, s^{-1}k})$ be a binary constraint in \mathcal{I}_T and $((v, s), \text{Types}(C, v, s))$ a constraint from Item 2. Then \mathcal{I}'_T contains $((v, s), \text{Comp}_{v^{-1}k, s^{-1}k} \cap \text{Types}(C, v, s))$.

In order to complete the proof of the corollary, we need to show that \mathcal{I}'_T is non-trivial, $(2, 3)$ -minimal and equivalent to \mathcal{I}_T .

► **Fact 34.** *The instance \mathcal{I}'_T is equivalent to \mathcal{I}_T .*

Proof. By Fact 33 we have that $h : \mathcal{V} \rightarrow A$ is a solution to \mathcal{I}_A if and only if $g : I \rightarrow T_{\mathbb{B},m}$ such that $g(v)$ is the type of $(h(v(1), \dots, h(v(m)))$ is a solution to \mathcal{I}_T . We need to show that every such g is also a solution to \mathcal{I}'_T . Since h satisfies all new constraints in \mathcal{I}_A , i.e., constraints

of the form $((j(1), \dots, j(r)), S')$ for all $m \leq r \leq 3m$ and $j : [r] \rightarrow \mathcal{V}$ originating from a constraint $((j(1), \dots, j(r)), A^r)$, it is straightforward to check that g satisfies all constraints in \mathcal{I}'_T from Items 1–3. Since g satisfies these new constraints and all constraints in \mathcal{I}_T , it also satisfies constraints in \mathcal{I}'_T defined in Items 4 and 5. ◀

► **Fact 35.** *The instance \mathcal{I}'_T is non-trivial.*

Proof. We need to show that all relations associated to constraints in \mathcal{I}'_T are non-empty. Since \mathcal{I}'_A is non-trivial, this statement clearly holds for all constraints from Items 1–3. Since all constraints in \mathcal{I}'_T from Item 1 are non-empty, we have that for every $v \in I$ there is an m -type which satisfies all unary constraints in \mathcal{I}_T of the form $((v), U)$. It follows that relations in constraints in \mathcal{I}'_T in Item 4 are non-empty. Finally observe that for every constraint $((v, s), R)$ in \mathcal{I}'_T from Item 2, the binary relation R is contained in all relations of the form $\text{Comp}_{v^{-1}k, s^{-1}k}$. It follows that relations in constraints defined in Item 5 are also non-empty. It completes the proof of the fact. ◀

► **Fact 36.** *The instance \mathcal{I}'_T is $(2, 3)$ -minimal.*

Proof. Clearly all $s, u, v \in I$ occur together in the scope of some constraint in \mathcal{I}'_T from Item 3. For Condition 2 in Definition 6, observe both of the following:

- for all $v \in I$, the constraint (v, U) in \mathcal{I}'_T in Item 4 and all constraints (v, U') in Item 1, we have that $U = U'$, and
- for all $v, u \in I$, the constraint $((v, u), R)$ in \mathcal{I}'_T in Item 5 and all constraints $((v, u), R')$ in Item 2, we have that $R = R'$.

Thus, in order to satisfy Condition 2, it is enough to restrict to the case where C_1, C_2 are constraints from Items 1–3. But Condition 2 in Definition 6 holds for all pairs of constraints in Items 1–3 because \mathcal{I}'_A is $(2m, 3m)$ -minimal. It completes the proof of the fact. ◀

C Proof of Proposition 11

Let h be any binary canonical operation from the formulation of the lemma. We will show that in any case:

- $f(x_1, x_2, x_3, x_4) = h(h(h(x_1, x_2), x_3), x_4)$ and
- $g(x_1, x_2, x_3) = h(h(x_1, x_2), x_3)$

are weak near-unanimity operations modulo $\overline{\text{Aut}(\mathcal{H})}$ where \mathcal{H} is a homogeneous graph and that they satisfy the condition in Corollary 10. Clearly, f and g are canonical. In order to complete the proof of the proposition, we will show that all :

- $f(y, x, x, x), f(x, y, x, x), f(x, x, y, x)$ and $f(x, x, x, y)$,
- $g(y, x, x), g(x, y, x), g(x, x, y)$, and
- $f(y, x, x, x)$ and $g(y, x, x)$ have the same behaviour.

Observe that if a behaviour B of h is min or max, then it is a simillattice operation on the three-element set $\{E, N, =\}$. It is therefore straightforward to check that all operations obtained from f and g mentioned in the items above are of the same behaviour as h . Thus, the proposition follows in this case where h is of behaviour min or max.

If h is E -constant or N -constant, then all operations mentioned in the items above are E -constant or N -constant, respectively. Thus, the proposition follows also in this case. ◻

D Proof of Observation 15

The relation \sim is clearly reflexive and symmetric. Assume on the contrary that it is not transitive. Then there are $i, j, k \in [n]$ such that $\mathcal{I}_{i,j}, \mathcal{I}_{j,k} \subseteq \{=\}$ and $\mathcal{I}_{i,k}$ contains $O \in \{E, N\}$. Since \mathcal{I} is $(2, \mathbb{L})$ -minimal and $\mathbb{L} \geq 3$, it contains a constraint C whose scope contains $\{v_i, v_j, v_k\}$. But C is not satisfiable by $a : \{v_i, v_j, v_k\} \rightarrow D$ such that $a(v_i) = a(v_j)$, $a(v_j) = a(v_k)$ and $(a(v_i), a(v_k)) \in O$. It contradicts $(2, \mathbb{L})$ -minimality of \mathcal{I} and completes the proof of the observation. \square

E Proof of Proposition 16

In order to prove the proposition, we will show that any \mathbb{A} is preserved by an oligopotent qnu-operation, see Theorem 13. We start with a couple of definitions.

A k -tuple t with $k \geq 3$ has the main value a if all but at most one entry in t takes the value a . Note that not all tuples have the main value, but if a tuple has the main value, then it is unique. It is straightforward to see that every qnu-operation sends all tuples with the same main value to the same element of the image. For a natural number $k \geq 3$, define \sim_k to be the equivalence relation on A^k such that $u, v \in A^k$ satisfy $u \sim_k v$ if and only if u and v have the same main value. We first state a simple observation.

► **Observation 37.** *Let $p = 2m + 1$ for some natural number m and $R \in \{E, N\}$. Then there exists an oligopotent quasi near-unanimity operation f of arity p such that all of the following hold:*

1. $f(u) = f(v)$ if and only if $u \sim_p v$;
2. f preserves both E and N ;
3. if u has the main value c and v has the main value d , then $(f(u), f(v)) \in R$ if and only if $(c, d) \in R$;
4. if u has the main value c and v has no main value and there exists $I \subseteq [p]$ with $|I| \geq m + 1$ such that $(c, v[i]) \in R$ for all $i \in I$ then $(f(u), f(v)) \in R$.

Proof. Consider a graph G_f whose vertices are the equivalence classes of \sim_p and $[u]_{\sim_p}$ and $[v]_{\sim_p}$ in A^p / \sim_p are connected by an edge if one of the following holds:

- u has the main value c , v has the main value d and $(c, d) \in E$,
- u has the main value c , v has no main value and $I = \{i \in [2m + 1] \mid (c, v_i) \in E\}$ satisfies $|I| \geq m + 1$,
- neither u nor v has the main value and $(u[i], v[i]) \in E$ for all $i \in [2m + 1]$.

Since G_f is a countable graph, it clearly embeds into the random graph by an embedding f . It is now straightforward to check that f satisfies all 4 items in the formulation of the observation. ◀

For every relation R in \mathbb{A} set φ_R to be some definition of R given by the conjunction of clauses of the form given in the formulation of the proposition and $r = k + l$ be the maximal number of disequalities over all clauses in all φ_R for all R in \mathbb{A} . Let m be a natural number satisfying $m \geq r + 2$. We will prove that \mathbb{A} is preserved by a $(2m + 1)$ -ary oligopotent qnu-operation f from Observation 37. By appeal to Theorem 13, it will imply that \mathbb{A} has bounded strict width.

In fact, by Theorem 1, it is enough to prove that f preserves any relation defined by a single clause of the desired form with the number of disequalities $(k + l)$ less than r . Assume on the contrary that f violates some R defined by a single clause ψ of the considered form over variables $X = \{x_1, \dots, x_k, y_1, y_2, z_1, \dots, z_l\}$ and set $a(v)$ to be the

vector $(a_1(v), \dots, a_{2m+1}(v))$ for all $v \in X$ where $a_1, \dots, a_{2m+1} : X \rightarrow A$ are such that all a_i with $i \in [2m+1]$ satisfy ψ but $h : X \rightarrow A$ defined so that $h(v) = f(a_1(v), \dots, a_{2m+1}(v))$ for all $v \in X$ does not satisfy ψ . In particular, $h(y_1) = h(x_i)$ for all $i \in [k]$, $h(y_2) = h(z_i)$ for all $i \in [l]$ and $(h(y_1), h(y_2)) \notin R$. Assume without loss of generality that all $a(x_1), \dots, a(x_k)$ are different than $a(y_1)$ and that all $a(z_1), \dots, a(z_l)$ are different than $a(y_2)$. Then we have both of the following:

- either $\{x_1, \dots, x_k\}$ is empty or $a(y_1), a(x_1), \dots, a(x_k)$ have all the same main value c , and
- either $\{z_1, \dots, z_l\}$ is empty or $a(y_2), a(z_1), \dots, a(z_l)$ have the same main value d .

If both $\{x_1, \dots, x_k\}$ and $\{z_1, \dots, z_l\}$ are empty, then ψ is $R(y_1, y_2)$ and we have the contradiction. Indeed, by Item 2 in Observation 37, f preserves both E and N . Next, turn to the case where exactly one holds: either $\{x_1, \dots, x_k\}$ or $\{z_1, \dots, z_l\}$ is empty. Assume without loss of generality that $\{z_1, \dots, z_l\}$ is empty. We now consider a couple of cases. Observe that in this case $a(y_1)$ has the main value c . If $a(y_2)$ also has the main value, then for all but at most two $i \in [2m+1]$ we have $(a_i(y_1), a_i(y_2)) \in R$. Since $m \geq 2$, by the definition of f , it follows that $(h(y_1), h(y_2)) \in R$. Thus, we may assume that $a(y_2)$ does not have the main value. Set $I \subseteq [2m+1]$ to be the set of indices such that $i \in I$ if and only if $(a_i(x_j) = a_i(y_1))$ for all $j \in [k]$. Observe that $(a_i(y_1), a_i(y_2)) \in R$ for all $i \in I$. If $(a_1(y_1), \dots, a_{2m+1}(y_1))$ is a constant tuple, then $|I| \geq m+1$, then, by Item 4 in Observation 37, we have $(h(y_1), h(y_2)) \in R$. If $a(y_1)$ is not a constant tuple, then either i such that $a_i(y_1) \neq c$ is in I or not. Observe that in the first case $I = [2m+1]$ and that it contradicts the assumption. In the other case, since $m \geq k+1$, we have that $|I| \geq m+1$ and for all $i \in I$ it holds $a_i(y_1) = c$. Clearly, for all $i \in I$ we have also that $(a_i(y_1), a_i(y_2)) \in R$. This time, by Item 4 in Observation 37, we have that $(h(y_1), h(y_2)) \in R$. It completes the proof of the case where either $\{x_1, \dots, x_k\}$ or $\{z_1, \dots, z_l\}$ is empty.

Finally, we turn to the case where neither $\{x_1, \dots, x_k\}$ nor $\{z_1, \dots, z_l\}$ is empty. Then both $a(y_1)$ and $a(y_2)$ have the main values c and d , respectively. Since $(c, d) \in R$ we have by Item 3 in Observation 37 that $(h(y_1), h(y_2)) \in R$. It completes the proof of the proposition.

F Proof of Proposition 17

In order to prove that $\mathbb{A} = (A; E, N, R)$ where $(A; E)$ is C_2^ω and $R(x_1, x_2, x_3) \equiv ((E(x_1, x_2) \wedge N(x_2, x_3)) \vee (N(x_1, x_2) \wedge E(x_2, x_3)))$ has bounded strict width, we will show that it is preserved by an oligopotent qnu-operation. To this end, we will see elements of A as pairs in $\{0, 1\} \times \mathbb{N}$ so that $((d_1, e_1), (d_2, e_2)) \in E$ if and only if $d_1 = d_2$. Let now $m : \{0, 1\}^3 \rightarrow \{0, 1\}$ be the majority operation, i.e., it satisfies $m(0, 0, 0) = m(1, 0, 0) = m(0, 1, 0) = m(0, 0, 1) = 0$ and $m(1, 1, 1) = m(0, 1, 1) = m(1, 0, 1) = m(0, 1, 1) = 1$ and $i : \mathbb{N}^3 \rightarrow \mathbb{N}$, where \mathbb{N} is the set of natural numbers, be any injection. We claim that $f : A^3 \rightarrow A$ defined so that it satisfies: $f((d_1, e_1), (d_2, e_2), (d_3, e_3)) = (m(d_1, d_2, d_3), i(e_1, e_2, e_3))$ is an oligopotent qnu-operation.

The operation is oligopotent since it clearly preserves E and N . In order to see that R is preserved by f consider elements (e_i^j, d_i^j) in A with $i, j \in [3]$ such that for all $j \in [3]$ we have $((d_1^j, e_1^j), (d_2^j, e_2^j), (d_3^j, e_3^j)) \in R$. We have to show that $(f((d_1^1, e_1^1), (d_1^2, e_1^2), (d_1^3, e_1^3)), f((d_2^1, e_1^1), (d_2^2, e_1^2), (d_2^3, e_1^2)), f((d_3^1, e_1^1), (d_3^2, e_1^2), (d_3^3, e_1^2)))$ is in R . Observe that two out of $\{d_1^2, d_2^2, d_3^2\}$ need to have the same value. Assume without loss of generality that $d_1^2 = d_2^2 = 0$. Further, if $d_1^1 = d_2^1 = 0$ and $d_1^3 = d_2^3 = 1$ or $d_1^1 = d_2^1 = 1$ and $d_1^3 = d_2^3 = 0$, then we are clearly done. Hence, without loss of generality we may assume that $d_1^1 = 0, d_2^1 = 1$ and $d_1^3 = 1, d_2^3 = 0$. But in this case exactly one out of $\{d_1^3, d_3^3\}$ equals 0 and still f preserves R . \square

G

 Proof of Lemma 18

Let \mathcal{I} be a non-trivial $(2, \mathbb{L})$ -minimal instance of the CSP equivalent to an instance of $\text{CSP}(\mathbb{A})$ and let φ_I be a conjunction of atomic formulas containing $R(v_{i_1}, \dots, v_{i_r})$ for every constraint of the form $((v_{i_1}, \dots, v_{i_r}), R)$ in \mathcal{I} ⁵. Then consider

$$\varphi_S := \varphi_S^{O_1} \wedge \varphi_S^{O_2} \wedge \varphi_S^=$$

where

- $\varphi_S^{O_1} := \bigwedge_{\mathcal{I}_{i,j} \ni O_1} O_1(v_i, v_j)$,
- $\varphi_S^{O_2} := \bigwedge_{\mathcal{I}_{i,j} = \{O_2, =\}} O_2(v_i, v_j) \wedge \bigwedge_{\mathcal{I}_{i,j} = \{O_2\}} O_2(v_i, v_j)$, and
- $\varphi_S^= := \bigwedge_{\mathcal{I}_{i,j} = \{=\}} v_i = v_j$.

In order to prove that \mathcal{I} has a solution, we will show that $\varphi_I \wedge \varphi_S$ is satisfiable. Let $\varphi_I^{O_1}$ be φ_I where every atomic formula ψ is replaced by ψ^{O_1} equivalent to $\psi \wedge \eta_S^{O_1}$ where $\eta_S^{O_1}$ is $\varphi_S^{O_1}$ restricted to conjuncts involving variables in ψ . We first prove that every $\psi \wedge \eta_S^{O_1}$ defines a non-empty relation. Assume the contrary. Then there exists ψ , a minimal non-empty subset \mathbb{I}^{O_1} of $\{(i, j) \mid \mathcal{I}_{i,j} \ni O_1\}$ and (i_3, i_4) with $P_{i_3, i_4} \ni O_1$ not in \mathbb{I}^{O_1} such that $(\psi \wedge \bigwedge_{(i,j) \in \mathbb{I}^{O_1}} O_1(v_i, v_j))$ entails $\underline{O_2}(v_{i_3}, v_{i_4})$. Let $(i_1, i_2) \in \mathbb{I}^{O_1}$ and $\{y_1, \dots, y_l\}$ be all variables occurring in ψ . Set

$$R(x_1, x_2, x_3, x_4) \equiv (\exists y_1 \dots \exists y_l (\psi \wedge \bigwedge_{(i,j) \in \mathbb{I}^{O_1} \setminus \{(i_1, i_2)\}} O_1(v_i, v_j) \wedge \bigwedge_{j \in [4]} x_j = v_{i_j})). \quad (1)$$

Since \mathbb{A} is a first-order expansion of \mathcal{H} and in consequence contains both E and N , we have that \mathbb{A} pp-defines R . Furthermore, observe that R efficiently entails $(O_1(x_1, x_2) \implies \underline{O_2}(x_3, x_4))$. The relation R clearly entails $(O_1(x_1, x_2) \implies \underline{O_2}(x_3, x_4))$. Since \mathbb{I}^{O_1} is minimal, we have that R contains a tuple t_1 such that $(t_1[1], t_1[2]) \in O_1$ and $(t_1[3], t_1[4]) \in \underline{O_2}$. On the other hand, the minimality of \mathbb{I}^{O_1} implies that R contains a tuple t_2 satisfying $(t_2[3], t_2[4]) \in O_1$. Since R entails $(O_1(x_1, x_2) \implies \underline{O_2}(x_3, x_4))$ we have that $(t_2[1], t_2[2]) \notin O_1$. It follows that R is a $[(O_1(x_1, x_2) \implies \underline{O_2}(x_3, x_4))]$ -relation and contradicts the assumption that \mathbb{A} does not pp-define such relations. Thus, every ψ^{O_1} is satisfiable.

It follows that for every atomic formula ψ^{O_1} in $\varphi_I^{O_1}$ the projection of ψ^{O_1} to any $\{v_i, v_j\}$ with $\mathcal{I}_{i,j} \ni O_1$ equals O_1 . Observe that the following is also true: the projection of every ψ^{O_1} to $\{v_i, v_j\}$ with $O_2 \subseteq \mathcal{I}_{i,j} \subseteq \underline{O_2}$ contains O_2 . Indeed, otherwise there exists a minimal \mathbb{I}^{O_1} of $\{(i, j) \mid \mathcal{I}_{i,j} \ni O_1\}$ and (i_3, i_4) with \mathcal{I}_{i_3, i_4} satisfying $O_2 \subseteq P_{i_3, i_4} \subseteq \underline{O_2}$ such that $(\psi \wedge \bigwedge_{(i,j) \in \mathbb{I}^{O_1}} O_1(v_i, v_j))$ entails $(v_{i_3} = v_{i_4})$. Let $(i_1, i_2) \in \mathbb{I}^{O_1}$ and $\{y_1, \dots, y_l\}$ be all variables occurring in ψ . In the same way as above, we argue that $R(x_1, x_2, x_3, x_4)$ defined in the same way as the relation in (1) contains a tuple t_1 such that $(t_1[1], t_1[2]) \in O_1$ and $(t_1[3] = t_1[4])$ as well as a tuple t_2 satisfying $(t_2[1], t_2[2]) \in \underline{O_2}$ and $(t_1[3], t_1[4]) \in O_2$. It implies that R efficiently entails $(O_1(x_1, x_2) \rightarrow x_3 = x_4)$, which contradicts the assumption that R does not pp-define $[(O_1(x_1, x_2) \rightarrow x_3 = x_4)]$ -relations.

In the second step of the transformation of the instance, we obtain $\varphi_I^{O_2}$ from $\varphi_I^{O_1}$ by replacing every atomic formula ψ^{O_1} with ψ^{O_2} equivalent to $(\psi^{O_1} \wedge \eta_S^{O_2})$ where $\eta_S^{O_2}$ is $\varphi_S^{O_2}$ restricted to variables in ψ . We claim that a relation defined by any $(\psi^{O_1} \wedge \eta_S^{O_2})$ is non-empty. Assume on the contrary that it is not the case for some $(\psi^{O_1} \wedge \eta_S^{O_2})$. Define \mathbb{I}^{O_2} to be the set of all $(i, j) \in [n]^2$ where $\{v_i, v_j\}$ are in the scope of ψ^{O_2} and the projection

⁵ It is well-known that all such relations R are pp-definable in \mathbb{A}

of ψ^{O_1} to $\{v_i, v_j\}$ contains $\underline{O_2}$. Since ψ^{O_1} is satisfiable and ψ^{O_2} is not and since the projection of ψ^{O_1} to $\{v_i, v_j\}$ with $\mathcal{I}_{i,j}$ satisfying $O_2 \subseteq \mathcal{I}_{i,j} \subseteq \underline{O_2}$ contains O_2 , there exists a minimal non-empty $\mathbb{J} \subseteq \mathbb{I}^{O_2}$ and $(i_3, i_4) \in \mathbb{I}^{O_2} \setminus \mathbb{J}$ such that $(\psi^{\overline{O_1}} \wedge \bigwedge_{(i,j) \in \mathbb{J}} O_2(v_i, v_j))$ entails $(v_{i_3} = v_{i_4})$. This time, having in mind that ψ^{O_1} is equivalent to $\psi \wedge \eta_S^{O_1}$ we can pp-define a $[(O_2(x_1, x_2) \implies x_3 = x_4), (O_2(x_1, x_2) \wedge \underline{O_2}(x_3, x_4))]$ -relation, which again contradicts the assumption that such relations are not pp-definable in \mathbb{A} .

Observe that in the obtained $\varphi_I^{O_2}$ a projection of any constraint ψ^{O_2} to any pair of variables $\{v_i, v_j\}$ in the scope of ψ equals E, N or $=$ and the projections agree in different atomic formulas. Thus, in fact \mathcal{I}' obtained from $\varphi_I^{O_2}$ by replacing atomic formulas of the form $R(v_{i_1}, \dots, v_{i_r})$ with constraints $((v_{i_1}, \dots, v_{i_r}), R)$ is a $(2, \mathbb{L})$ -minimal instance equivalent to an instance of $\text{CSP}((D; E, N, =))$ where $(D; E)$ is \mathcal{H} . By Observation 14, it follows that \mathcal{I}' , and hence, by Proposition 7, \mathcal{I} has a solution. \square .

H Proof of Lemma 19

Let \mathcal{I} be a $(2, 3)$ -minimal non-trivial instance of the CSP equivalent to an instance of $\text{CSP}(\mathbb{A})$ and let φ_I be a conjunction of atomic formulas obtained from \mathcal{I} by replacing every constraint of the form $((v_{i_1}, \dots, v_{i_r}), R)$ with an atomic formula $R(v_{i_1}, \dots, v_{i_r})$. We need to prove that φ_I is satisfiable. For an atomic formula ψ in φ_I we define ψ^N to be equivalent to $\psi \wedge \eta^N$ where η^N is the conjunction of $\bigwedge_{\mathcal{I}_{i,j} \supseteq \{N\}} N(v_i, v_j)$ restricted to variables occurring in ψ . We first show that φ_I^N obtained from φ_I by replacing every ψ with ψ^N is satisfiable.

Assume first on the contrary that there exists ψ^N in φ_I^N defining an empty relation. Then there exists a non-empty and minimal set \mathbb{I}^N of pairs of indices (i, j) satisfying $\mathcal{I}_{i,j} \supseteq \{N\}$ and $(i_3, i_4) \notin \mathbb{I}^N$ but with $P_{i_3, i_4} \supseteq \{N\}$ such that $(\psi \wedge \bigwedge_{(i,j) \in \mathbb{I}^N} N(v_i, v_j))$ entails $\underline{E}(v_{i_3}, v_{i_4})$. Let $\{y_1, \dots, y_l\}$ be all variables occurring in ψ and $(i_1, i_2) \in \mathbb{I}^N$. Then:

$$R(x_1, x_2, x_3, x_4) \equiv (\exists y_1 \dots \exists y_l (\psi \wedge \bigwedge_{(i,j) \in \mathbb{I}^N \setminus \{(i_1, i_2)\}} N(v_i, v_j) \wedge \bigwedge_{j \in [4]} x_j = v_{i_j})) \quad (2)$$

entails $(N(x_1, x_2) \implies \underline{E}(x_3, x_4))$. Since \mathbb{I} is minimal and non-empty, the relation R contains a tuple t_1 such that $(t_1[1], t_1[2]) \in N$ and $(t_1[3], t_1[4]) \in \underline{E}$ as well as a tuple t_2 satisfying $(t_2[1], t_2[2]) \in \underline{E}$ and $(t_2[3], t_2[4]) \in N$. It follows that \mathbb{A} pp-defines a $[(N(x_1, x_2) \implies \underline{E}(x_3, x_4))]$ -relation, which contradicts the assumption of the lemma.

We will now show that \mathcal{I}^N obtained from φ_I^N by replacing every atomic formula of the form $R(v_{i_1}, \dots, v_{i_r})$ by $((v_{i_1}, \dots, v_{i_r}), R)$ is $(2, 3)$ -minimal. Since \mathcal{I} is $(2, 3)$ -minimal, clearly every subset of at most 3 variables is in the scope of some constraint. Further, if the projection of any constraint in \mathcal{I}^N to $\{v_i, v_j\}$ with $i, j \in [n]$ contains N , it equals N , and hence the projections of any two constraint in \mathcal{I}^N to $\{v_i, v_j\}$ with $\mathcal{I}_{i,j} \supseteq \{N\}$ and $i, j \in [n]$ agree. In order to complete the proof of the fact that \mathcal{I}^N satisfy Property 2 in Definition 6, we concentrate on $\mathcal{I}_{i,j}$ that do not contain N . For all such $i, j \in [n]$ and all $O \in \mathcal{I}_{i,j}$ we will show that $(\psi^N \wedge O(v_i, v_j))$ is satisfiable. If $|\mathcal{I}_{i,j}| = 1$, we are done by the fact that every ψ^N is satisfiable. Thus, we turn to the case where $\mathcal{I}_{i,j} = \{E, =\}$. Assume on the contrary that there are $\psi^N, \{v_{i_3}, v_{i_4}\}$ in the scope of ψ^N and $O \in \{E, =\}$ such that $(\psi^N \wedge O(v_{i_3}, v_{i_4}))$ is not satisfiable. It follows that there is a non-empty and minimal $\mathbb{I}^N \subseteq \{(i, j) \mid \mathcal{I}_{i,j} \supseteq \{N\}\}$ such that $(\psi \wedge \bigwedge_{(i,j) \in \mathbb{I}^N} N(v_i, v_j))$ entails $O_1(v_{i_3}, v_{i_4})$ where $O_1 \in \{E, =\} \setminus \{O\}$. Hence $R(x_1, x_2, x_3, x_4)$ defined by the formula (2) entails $(N(x_1, x_2) \implies O_1(x_3, x_4))$. Again, using the minimality of \mathbb{I}^N we can show that R contains a tuple t_1 such that $(t_1[1], t_1[2]) \in N$ and $(t_1[3], t_1[4]) \in O_1$ as well as a tuple t_2 satisfying $(t_2[1], t_2[2]) \in \underline{E}$ and $(t_2[3], t_2[4]) \in O$. It follows that R efficiently entails $(N(x_1, x_2) \implies O_1(x_3, x_4))$ and that it entails $\underline{E}(x_3, x_4)$.

Since \mathbb{A} contains N , the relation R is a $[(N(x_1, x_2) \implies O_1(x_3, x_4)), (\underline{E}(x_3, x_4))]$ -relation pp-definable in \mathbb{A} . Now, regardless of whether O_1 is E or $=$, we have a contradiction with the assumptions of the lemma.

From now on we assume that \mathcal{I}^N is $(2, 3)$ -minimal and we set $P_{i,j}^N$ for $i, j \in [n]$ to be the set of orbitals $O \in \{E, N, =\}$ contained in the projection of any constraint in \mathcal{I}^N to $\{v_i, v_j\}$. Actually $P_{i,j}^N = \mathcal{I}_{i,j}$ if $N \notin \mathcal{I}_{i,j}$ and $P_{i,j}^N = \{N\}$ otherwise. If $|P_{i,j}^N| = 1$ for all $i, j \in [n]$, then \mathcal{I}^N is a non-trivial $(2, 3)$ -minimal instance of the CSP equivalent to an instance of $\text{CSP}((D; E, N, =))$ where $(D; E)$ is C_ω^2 . Since $\mathbb{L}_{C_\omega^2}$ is 3, it follows by Observation 14 that \mathcal{I}^N and, by Proposition 7, that \mathcal{I} has a solution. It completes the proof of the lemma in the case where $|P_{i,j}^N| = 1$ for all $i, j \in [n]$.

From now on we assume that there exists $i, j \in [n]$ such that $|P_{i,j}^N| = 2$. In particular, we have that \underline{E} is definable in \mathbb{A} . Since \mathcal{I}^N is $(2, 3)$ -minimal, it is easy to see that $\mathcal{I}_{i,j}^N$ with $i, j \in [n]$ respects the transitivity of \underline{E} in the sense that for all $i, j, k \in [n]$: if $P_{i,j}^N \cap \{E, =\} \neq \emptyset$ and $P_{j,k}^N \cap \{E, =\} \neq \emptyset$, then $P_{i,k}^N \cap \{E, =\} \neq \emptyset$. Set now J_1, \dots, J_m to be the partition of \mathcal{V} such that for all $l \in [m]$ and $i, j \in J_l$ we have that $P_{i,j}^N \cap \{E, =\} \neq \emptyset$. Then construct φ^i for all $i \in [m]$ to be a formula that for every atomic formula ψ^N in φ_I^N contains ψ^i equivalent to $(\exists v_{j_1} \dots \exists v_{j_l} \psi^N)$ where $\{j_1, \dots, j_l\} = [n] \setminus J_i$. Observe that every φ^i is equivalent to an instance \mathcal{I}^i of $\text{CSP}(\mathbb{A}_i)$ such that \mathbb{A}_i has a pp-definition in \mathbb{A} and hence is also preserved by an oligopotent qnu-operation f . Furthermore, the core \mathbb{A}_i^c of \mathbb{A}_i for every $i \in [m]$ is finite. Assume without loss of generality that \mathbb{A}_i^c admits a fo-definition over a single edge $\{a_{2i-1}, a_{2i}\}$ in A and that f is idempotent on $\{a_1, \dots, a_{2m}\}$. Let h_i with $i \in [m]$ be a homomorphism from \mathbb{A}_i to \mathbb{A}_i^c that is, without loss of generality, idempotent on $\{a_{2i-1}, a_{2i}\}$. Then, since f preserves \underline{E} , we have that $h_i(f(x_1, \dots, x_k))$ restricted to $\{a_{2i-1}, a_{2i}\}$ is a near-unanimity operation preserving \mathbb{A}_i^c . Since an instance of the CSP equivalent to φ^i is $(2, 3)$ -minimal also when evaluated in \mathbb{A}_i^c , it follows by Theorem 10 that φ^i for all $i \in [l]$ has a solution s_i in $\{a_{2i-1}, a_{2i}\}$.

For all $l \in [m]$ and $i, j \in J_l$ we set $O_{i,j}$ to E if $s(v_i) \neq s(v_j)$ and to $=$ otherwise. In order to complete the proof of the lemma we will show that $(\varphi_I^N \wedge \bigwedge_{l \in [m]} \bigwedge_{i,j \in J_l} O_{i,j}(v_i, v_j))$ is satisfiable. Assume the contrary. Then there exists a ψ^N in φ_I^N and a minimal non-empty subset \mathbb{J} of pairs of indices $(i, j) \in \bigcup_{l \in [m]} J_l \times J_l$ of variables occurring in ψ^N such that $(\psi^N \wedge \bigwedge_{(i,j) \in \mathbb{J}} O_{i,j}(v_i, v_j))$ is not satisfiable. Since s_i is a solution to φ^i for all $i \in [m]$ we have that \mathbb{J} contains at least two pairs: $(i_1, i_2) \in J_a$ and $(i_3, i_4) \in J_b$ with $a \neq b$. Set y_1, \dots, y_l to be all variables occurring in ψ and consider now the relation

$$R(x_1, x_2, x_3, x_4) \equiv (\exists y_1 \dots y_m (\psi^N \wedge \bigwedge_{(i,j) \in \mathbb{J} \setminus \{(i_1, i_2), (i_3, i_4)\}} O_{i,j}(v_i, v_j) \wedge \bigwedge_{k \in [4]} x_k = v_{i_k})).$$

Since $a \neq b$, the relation R entails $N(x_2, x_3)$ and $(O_{i_1, i_2}(x_1, x_2) \implies O'_{i_3, i_4}(x_3, x_4))$ where $O'_{i_3, i_4} \in \{E, =\} \setminus \{O_{i_3, i_4}\}$. By the minimality of \mathbb{J} we have that R contains a tuple t_1 such that $(t_1[1], t_1[2]) \in O_{i_1, i_2}$ and $(t_1[3], t_1[4]) \in O'_{i_3, i_4}$ as well as a tuple t_2 satisfying $(t_2[1], t_2[2]) \in O'_{i_1, i_2}$ where $O'_{i_1, i_2} \in \{E, =\} \setminus \{O_{i_1, i_2}\}$ and $(t_2[3], t_2[4]) \in O_{i_3, i_4}$. Since \mathbb{A} contains E, N , we have that it pp-defines a $[(O_{i_1, i_2}(x_1, x_2) \implies O'_{i_3, i_4}(x_1, x_2)), (\underline{E}(x_1, x_2) \wedge N(x_2, x_3) \wedge \underline{E}(x_3, x_4))]$ -relation where $O_{i_1, i_2}, O'_{i_3, i_4} \in \{E, =\}$. It contradicts the assumptions of the lemma and proves that s satisfies φ_I , and hence it is a solution to \mathcal{I} . It completes the proof of the lemma. \square

I Proof of Lemma 20

Let \mathcal{I} be a non-trivial $(2, 3)$ -minimal instance of the CSP equivalent to an instance of $\text{CSP}(\mathbb{A})$ and let $0, 1 \in A$ be two elements in two different infinite cliques in C_2^ω and $g : A \rightarrow \{0, 1\}$ a

function that sends all elements in the infinite clique containing 0 to 0 and all elements in the infinite clique containing 1 to 1. Define the structure $g(\mathbb{A})$ over the domain $\{0, 1\}$ to be a structure in which every R in \mathbb{A} is replaced by $g(R)$ in which in turn every tuple $t \in R$ is replaced by $g(t)$. Since f preserves E and hence $\underline{E}(x_1, x_2) \equiv (\exists y E(x_1, y) \wedge E(y, x_2))$, we may assume without loss of generality that f sends (a_1, \dots, a_k) for a_1, \dots, a_k in the same infinite clique to an element in the same infinite clique. It follows that $f'(x_1, \dots, x_k) := g(f(x_1, \dots, x_k))$ restricted to $\{0, 1\}$ is a near-unanimity operation preserving $g(\mathbb{A})$. Consider then $g(\mathcal{I})$ which is an instance of $\text{CSP}(g(\mathbb{A}))$ obtained from \mathcal{I} by replacing every constraint $((v_{i_1}, \dots, v_{i_r}), R)$ by $((v_{i_1}, \dots, v_{i_r}), g(R))$. Notice that $g(\mathcal{I})$ is $(2, 3)$ -consistent. Clearly for every set of three variables, there is a constraint containing these variables in its scope. Further, for any two variables v_i, v_j with $i, j \in [n]$, the projection of any constraint

$$((v_{k_1}, \dots, v_{k_r}), g(R))$$

to $\{v_i, v_j\}$ contains $\{(0, 0), (1, 1)\}$ if the projection of $((v_{k_1}, \dots, v_{k_r}), R)$ to $\{v_i, v_j\}$ contains $=$ or E and $\{(0, 1), (1, 0)\}$ if the projection of $((v_{k_1}, \dots, v_{k_r}), R)$ to $\{v_i, v_j\}$ contains N . Hence $g(\mathcal{I})$ is $(2, 3)$ -minimal. By Theorem 10 we have that $g(\mathcal{I})$ has a solution s_g . For every constraint C we let

- \mathbb{I}_C^d to be the set of pairs of indices $(i, j) \in [n]^2$ of variables occurring in C such that $s_g(v_i) \neq s_g(v_j)$ and
- \mathbb{I}_C^s to be the set of pairs of indices $(i, j) \in [n]^2$ of variables occurring in C such that $s_g(v_i) = s_g(v_j)$ and $\mathcal{I}_{i,j} \supseteq \{E\}$.

Consider now \mathcal{I}^N obtained from \mathcal{I} by replacing every constraint $C := ((v_{i_1}, \dots, v_{i_r}), R)$ in \mathcal{I} by $C^N := ((v_{i_1}, \dots, v_{i_r}), R')$ where $R'(v_{i_1}, \dots, v_{i_r}) \equiv (R(v_{i_1}, \dots, v_{i_r}) \wedge \bigwedge_{(i,j) \in \mathbb{I}_C^d} N(v_i, v_j))$. Since s_g is the solution to $g(\mathcal{I})$, we have that every R' is non-empty. Furthermore, we claim that the projection of every C^N to $\{v_i, v_j\}$ with $(i, j) \in \mathbb{I}_C^s$ contains E . Assume on the contrary that there is a constraint C^N and a pair $(i_3, i_4) \in \mathbb{I}_C^s$ such that the projection of C^N to $\{v_i, v_j\}$ does not contain E . It follows that there exists a minimal non-empty subset $\mathbb{J}_C^d \subseteq \mathbb{I}_C^d$ such that $(\eta_\psi := \bigwedge_{(i,j) \in \mathbb{J}_C^d} N(v_i, v_j) \wedge R')$ entails $\underline{N}(v_{i_3}, v_{i_4})$ and does not entail $N(v_{i_3}, v_{i_4})$. Since \mathbb{A} does not pp-define \underline{N} , we have that η_ψ entails $(v_{i_3} = v_{i_4})$. Let $(i_1, i_2) \in \mathbb{J}_C^d$ and $\{y_1, \dots, y_l\}$ be the scope of C . Then

$$R(x_1, x_2, x_3, x_4) := (\exists y_1 \dots \exists y_l \left(\bigwedge_{(i,j) \in \mathbb{J}_C^d \setminus \{(i_1, i_2)\}} N(v_i, v_j) \wedge \psi \wedge \bigwedge_{j \in [4]} v_{i_j} = x_j \right))$$

entails $(N(x_1, x_2) \implies (x_3 = x_4))$. Since s_g is a solution to $g(\mathcal{I})$, the relation R clearly contains a tuple t_1 with $(t_1[1], t_1[2]) \in N$ and $(t_1[3] = t_1[4])$. On the other hand, the set \mathbb{J}_C^d is minimal, and hence R contains a tuple t_2 satisfying $(t_2[3] \neq t_2[4])$ and $(t_2[1], t_2[2]) \notin N$. We obtain that R is a $[(N(x_1, x_2) \rightarrow (x_3 = x_4))]$ -relation. Since \mathbb{A} contains N , it pp-defines R . It contradicts the assumption of the lemma. We have that the projection of every C^N to any $(i, j) \in \mathbb{I}_C^s$ contains E .

In the next step we construct \mathcal{I}^E which is obtained from \mathcal{I}^N by replacing every constraint $C^N := ((v_{k_1}, \dots, v_{k_r}), R)$ by $C^E := ((v_{k_1}, \dots, v_{k_r}), R')$ where $R'(v_{k_1}, \dots, v_{k_r}) \equiv (R(v_{k_1}, \dots, v_{k_r}) \wedge \bigwedge_{(i,j) \in \mathbb{I}_C^s} E(v_i, v_j))$. We claim that every such R' is non-empty. Assume the contrary. Then there exists a minimal and non-empty $\mathbb{J}_C^s \subseteq \mathbb{I}_C^s$ and $(i_3, i_4) \in \mathbb{I}_C^s \setminus \mathbb{J}_C^s$ such that $(\eta_C^E := R'(v_{k_1}, \dots, v_{k_r}) \wedge \bigwedge_{(i,j) \in \mathbb{J}_C^s} E(v_i, v_j))$ entails $\underline{N}(v_{i_3}, v_{i_4})$ and does not entail $N(v_{i_3}, v_{i_4})$. Again, since \mathbb{A} does not pp-define \underline{N} , the formula η_ψ^E entails $(v_{i_3} = v_{i_4})$. Let

$(i_1, i_2) \in \mathbb{J}_C^s$. Then

$$R(x_1, x_2, x_3, x_4) := (\exists v_{k_1} \cdots \exists v_{k_r} (\psi^N \wedge \bigwedge_{(i,j) \in \mathbb{J}_C^s \setminus \{(i_1, i_2)\}} E(v_i, v_j) \wedge \bigwedge_{j \in [4]} v_{i_j} = x_j))$$

entails $(E(v_{i_1}, v_{i_2}) \implies (v_{i_3} = v_{i_4}))$. Since the projection of every C^N to any $(i, j) \in \mathbb{J}_C^s$ contains E , the relation R contains a tuple t_1 with $(t_1[1], t_1[2]) \in E$ and $(t_1[3] = t_1[4])$. On the other hand, since \mathbb{J}_C^s is minimal there is a tuple $t_2 \in R$ with $(t_2[3] \neq t_2[4])$ and $(t_2[1], t_2[2]) \notin E$. Thus, R is a $[(E(x_1, x_2) \rightarrow x_3 = x_4)]$ -relation. Since \mathbb{A} contains E and N , we have that R is pp-definable in \mathbb{A} . It contradicts the assumption of the lemma and completes the proof of the fact that every R' is non-empty.

Since \mathbb{A} does not pp-define \underline{N} , for every $i, j \in [n]$, the projection of any C^E to $\{v_i, v_j\}$ is N if $(i, j) \in \mathbb{I}_C^d$, it is E if $(i, j) \in \mathbb{J}_C^s$ and $=$ otherwise that is when $\mathcal{I}_{i,j} = \{=\}$. Thus, \mathcal{I}^E is a $(2, 3)$ -minimal instance equivalent to an instance of $\text{CSP}((D; E, N, =))$, where $(D; E)$ is C_2^ω . Since $\mathbb{L}_{C_2^\omega}$ is 3, it follows by Observation 14 that \mathcal{I}^E and in consequence, by Proposition 7, \mathcal{I} has a solution. It completes the proof of the lemma. \square

J Proof Of Lemma 21

We will show that in any of the considered cases \mathbb{A} pp-defines none of the relations mentioned in the formulation of Lemma 18 where $O_1 = E, O_2 = N$ in Cases 1 and 3 and $O_1 = N, O_2 = E$ in Cases 2 and 4.

Assume first on the contrary that \mathbb{A} pp-defines a $[(O_1(x_1, x_2) \implies \underline{O_2}(x_3, x_4))]$ -relation. The relation R contains:

- a tuple t_1 such that $(t_1[1], t_1[2]) \in O_1$ and $(t_1[3], t_1[4]) \in \underline{O_2}$, and
- a tuple t_2 such that $(t_2[1], t_2[2]) \in \underline{O_2}$ and $(t_2[3], t_2[4]) \in \underline{O_1}$.

Observe that $t := f(t_1, t_2)$, where f is any operation from the formulation of the lemma, satisfies $(t[1], t[2]) \in O_1$ and $(t[3], t[4]) \in O_1$ in any case. But containment of $t \in R$ contradicts the fact that R efficiently entails $(O_1(x_1, x_2) \implies \underline{O_2}(x_3, x_4))$.

Then we turn to showing that $[(O_1(x_1, x_2) \implies x_3 = x_4)]$ -relations R cannot be pp-definable in \mathbb{A} . Assume on the contrary that the relation R contains

- a tuple t_1 such that $(t_1[1], t_1[2]) \in O_1$ and $t_1[3] = t_1[4]$, and
- a tuple t_2 such that $(t_2[1], t_2[2]) \in \underline{O_2}$ and $(t_2[3] \neq t_2[4])$.

This time, regardless of the case, $t := \overline{f}(t_1, t_2)$ satisfies $(t[1], t[2]) \in O_1$ and $(t[3] \neq t[4])$. It contradicts the assumption that R efficiently entails $(O_1(x_1, x_2) \implies x_3 = x_4)$.

Finally we turn to $[(O_2(x_1, x_2) \implies x_3 = x_4), (\underline{O_2}(x_1, x_2) \wedge \underline{O_2}(x_3, x_4))]$ -relations. Assume on the contrary that R is such a relation and is preserved by one of operations f mentioned in the formulation of the lemma. The relation R contains:

- a tuple t_1 such that $(t_1[1], t_1[2]) \in O_2$ and $t_1[3] = t_1[4]$, and
- a tuple t_2 such that $(t_2[1] = t_2[2])$ and $(t_2[3], t_2[4]) \in O_2$.

If f is of behaviour min or max and O_1 -dominated or O_1 -constant, then clearly $\underline{O_2}$ and in consequence R are not preserved by f . If f is of behaviour min or max and balanced, then $t := f(t_1, t_2)$ satisfies $(t[1], t[2]) \in O_1$ and $(t[3] \neq t[4])$. This contradicts the fact that R efficiently entails $(O_2(x_1, x_2) \implies x_3 = x_4)$ and completes the proof of the lemma. \square

K Proof of Lemma 23

$\mathbb{A} [(O(x_1, x_2) \implies x_3 = x_4)]$ -relation R contains:

- a tuple t_1 such that $(t_1[1], t_1[2]) \in O$ and $(t_1[3] = t_1[4])$, and

- a tuple t_2 such that $(t_2[1], t_2[2]) \notin O$ and $(t_2[3] \neq t_2[4])$,

Let f be a ternary injections of behaviour majority specified in the formulation of the lemma. To reach a contradiction, it is enough to show that f applied, perhaps multiple times, to t_1, t_2 produces a tuple t such that $(t[1], t[2]) \in O$ and $(t[3] \neq t[4])$. Indeed, it implies that R does not entail $(O(x_1, x_2) \implies x_3 = x_4)$ and contradicts the assumption.

Let $t := f(t_1, t_1, t_2)$. Since f is an injection and $t_2[3] \neq t_2[4]$, it always holds that $t[3] \neq t[4]$. If $t_2[1] \neq t_2[2]$, then since f is of behaviour majority, we have that $(t[1], t[2]) \in O$. If $t_2[1] = t_2[2]$, then since $(t_1[1], t_1[2]) \in O$, it is straightforward to check that $(t[1], t[2]) \in O$ for f that is

- hyperplanely balanced and of behaviour projection,
- hyperplanely of behaviour max and E -dominated,
- hyperplanely of behaviour min and N -dominated,
- hyperplanely E -constant if $O = E$,
- hyperplanely N -constant if $O = N$.

For the remaining two cases: the case where f is hyperplanely E -constant and O is N and the case where f is hyperplanely N -constant and O is E , observe that $t' = f(t_1, t_1, t)$ satisfies $(t'[1], t'[2]) \in O, t'[3] \neq t'[4]$. It completes the proof of the lemma. \square

L Proof of Lemma 24

Assume on the contrary that \mathbb{A} pp-defines a $[(O(x_1, x_2) \implies x_3 = x_4)]$ -relation R . The relation R contains

- a tuple t_1 such that $(t_1[1], t_1[2]) \in O$ and $t_1[3] = t_1[4]$, and
- a tuple t_2 such that $(t_2[1], t_2[2]) \notin O$ and $t_1[3] \neq t_1[4]$.

The proof goes along the lines of the proof of Lemma 23. Again, to reach a contradiction, we will produce a tuple t satisfying $(t[1], t[2]) \in O$ and $(t[3] \neq t[4])$. It contradicts that R is a $[(O(x_1, x_2) \implies x_3 = x_4)]$ -relation and will complete the proof of the lemma.

Consider $t = f(t_1, t_2, t_2)$ where f is any operation from the formulation of the lemma. Since $t_2[3] \neq t_2[4]$ and f is an injection, we have that $t[3] \neq t[4]$. Furthermore, since f is of behaviour minority we have $(t[1], t[2]) \in O$ always when $(t_2[1] \neq t_2[2])$. Thus, we turn to the case where $t_2[1] = t_2[2]$. If f is hyperplanely balanced and of behaviour projection, then $(t[1], t[2]) \in O$ and we are done. Further, if f is hyperplanely of behaviour projection and E -dominated, or hyperplanely balanced of behaviour xnor we obtain $(t[1], t[2]) \in E$ and hence we are done in the case where O is E . Similarly, if f is hyperplanely of behaviour projection and N -dominated, or hyperplanely balanced of behaviour xor we obtain $(t[1], t[2]) \in N$ and in this case we are done where O is N . Thus, either we are done or we have t such that $(t[1], t[2]) \in O_1$ with $\{O_1\} = \{E, N\} \setminus \{O\}$ and $(t[3] \neq t[4])$. Now $t' := f(t_1, t, t)$ is the desired tuple satisfying $(t'[1], t'[2]) \in O$ and $(t'[3] \neq t'[4])$. It completes the proof of the lemma. \square

M Proof of Lemma 26

Assume first on the contrary that a $[(N(x_1, x_2) \implies \underline{E}(x_3, x_4))]$ -relation R is pp-definable in \mathbb{A} and let t_1 be a tuple in R with $(t_1[1], t_1[2]) \in N$ and $(t_1[3], t_1[4]) \in \underline{E}$ and let $t_2 \in R$ satisfy $(t_2[1], t_2[2]) \in \underline{E}$ and $(t_2[3], t_2[4]) \in N$. Then for $t = h(t_1, t_1, t_2)$ we have that $(t[1], t[2]), (t[3], t[4]) \in N$. It contradicts the fact that R efficiently entails $(N(x_1, x_2) \implies \underline{E}(x_3, x_4))$ and completes the proof for the first case.

In the second case we again assume the contrary and consider a $[(N(x_1, x_2) \implies x_3 = x_4)]$ -relation R . This time we have $t_1 \in R$ with $(t_1[1], t_1[2]) \in N$ and $(t_1[3] = t_1[4])$ and $t_2 \in R$

with $(t_2[1], t_2[2]) \in \underline{E}$ and $(t_2[3] \neq t_2[4])$. Now, regardless of whether $(t_2[3], t_2[4]) \in N$ or $(t_2[3], t_2[4]) \in E$ the tuple $t = h(t_1, t_1, t_2)$ satisfies $(t[1], t[2]) \in N$ and $(t[3] \neq t_4)$. It contradicts the fact that R is a $[(N(x_1, x_2) \implies x_3 = x_4)]$ -relation.

The proof of the third case goes in the same way. We assume on the contrary that R pp-definable in \mathbb{A} is a $[(N(x_1, x_2) \implies E(x_3, x_4)), (\underline{E}(x_3, x_4))]$ -relation. This time we have t_1 with $(t_1[1], t_1[2]) \in N$ and $(t_1[3], t_1[4]) \in E$ and $(t_2[1], t_2[2]) \notin N$ and $(t_2[3] = t_2[4])$. Observe that $t := f(t_1, t_1, t_2)$ satisfies $(t[1], t[2]) \in N$ and $(t[3] = t[4])$, which contradicts that R is a $[(N(x_1, x_2) \implies E(x_3, x_4)), (\underline{E}(x_3, x_4))]$ -relation. \square

N Proof of Lemma 27

In what follows, a quaternary tuple t is called an OP-tuple with $O, P \in \{E, N, =\}$ if $(t[1], t[2]) \in O$ and $(t[3], t[4]) \in P$. A tuple is called a constant tuple if all its entries have the same value, injective if all its entries are pairwise different.

Assume now on the contrary that \mathbb{A} preserved by a ternary injection of behaviour minority or majority and a k -ary oligopotent qnu-operation f pp-defines both a $[(E(x_1, x_2) \rightarrow \underline{N}(x_3, x_4))]$ -relation R_{EN} and a $[(N(x_1, x_2) \implies \underline{E}(x_3, x_4))]$ -relation R_{NE} . Since by Lemmas 23 and 24, the structure \mathbb{A} pp-defines no $[(O(x_1, x_2) \rightarrow x_3 = x_4)]$ -relations with $O \in \{E, N\}$, the relation R_{EN} contains an EN-tuple. On the other hand R_{EN} contains a tuple t such that $(t[1], t[2]) \in \underline{N}$ and $(t[3], t[4]) \in E$. This time by Lemmas 23 and 24, we have that R_{EN} must have an NE-tuple. In a similar way we show that R_{NE} contains both an EN-tuple and a NE-tuple.

In the remainder of the proof we first show that a relation R_{NE} (resp., R_{EN}) pp-defines a $[(N(x_1, x_2) \rightarrow \underline{E}(x_3, x_4))]$ -relation R'_{NE} (resp. a $[(E(x_1, x_2) \rightarrow \underline{N}(x_3, x_4))]$ -relation R'_{EN}) that contains an EN-tuple and a NE-tuple of some desirable properties. Finally, we will show that both R'_{EN} and R'_{NE} contain either an EE-tuple or a NN-tuple. It follows that either R'_{EN} is not an $[(E(x_1, x_2) \rightarrow \underline{N}(x_3, x_4))]$ -relation or R'_{NE} is not a $[(N(x_1, x_2) \rightarrow \underline{E}(x_3, x_4))]$ -relation. It will complete the proof of the lemma. We continue with an easy observation on EN-tuples and NE-tuples.

► **Observation 38.** *Let t be either an EN-tuple or a NE-tuple. Then there are at most two different $i, j \in [4]$ such that $t[i] = t[j]$.*

Proof. The proof is performed by the inspection of cases. Since t is an EN-tuple or a NE-tuple, we have $(t[1], t[2]) \in O_1$, $(t[3], t[4]) \in O_2$ where $\{O_1, O_2\} = \{E, N\}$. Hence, in particular O_1, O_2 are two different orbitals. In the case where $t[2] = t[3]$, we have that $(t[1], t[3]) \in O_1$ and $(t[2], t[4]) \in O_2$. Since O_1 and O_2 are different, it follows that $t[1] \neq t[4]$. Furthermore, at most one may be true: either $(t[1] = t[3])$ or $(t[2] = t[4])$. Indeed, otherwise $(t[1], t[4])$ would be in both O_1 and O_2 . If either $(t[1] = t[3])$ or $(t[2] = t[4])$, then since $(t[1] \neq t[2])$ and $(t[3] \neq t[4])$, we have that $t[1] \neq t[4]$. The remaining case to be considered is when $t[1] = t[4]$. Then $(t[2], t[4]) \in O_1$ and $(t[1], t[3]) \in O_2$. It implies that $t[2] \neq t[3]$ and completes the proof of the lemma. ◀

We now prove that R_{EN} (resp. R_{NE}) pp-defines R'_{EN} (resp. R'_{NE}) of the desired properties.

► **Observation 39.** *The relation R_{EN} (resp. R_{NE}) pp-defines a $[(E(x_1, x_2) \implies \underline{N}(x_3, x_4))]$ -relation R'_{EN} (resp. a $[(N(x_1, x_2) \implies \underline{E}(x_3, x_4))]$ -relation R'_{NE}) that contains*

1. *either both an EN-tuple t_{EN} and a NE-tuple t_{NE} such that for all $t \in \{t_{EN}, t_{NE}\}$ we have $(t[2] = t[3])$, or*
2. *both an injective EN-tuple t_{EN} and an injective NE-tuple t_{NE} .*

Proof. Notice first that whenever $R \in \{R_{NE}, R_{EN}\}$ contains an OP-tuple t with $\{O, P\} = \{E, N\}$ which satisfies $t[i] \neq t[j]$ for some different $i, j \in [4]$, then R contains also a PO-tuple t' with $t'[i] \neq t'[j]$. Indeed, if R is preserved by a ternary injection g of behaviour majority, then the desired t' is the result of $g(t, t'', t'')$ where t'' is some PO-tuple in R . On the other hand, if R is preserved by a ternary injection g of behaviour minority, the desired t' is the result of $g(t, t, t'')$ where t'' is some PO-tuple in R .

It follows, by Observation 38 that either there is some exactly one $\{i, j\}$ such that all EN-tuples and all NE-tuples t in R satisfy $t[i] = t[j]$ or R contains an injective EN-tuple and an injective NE-tuple. In the latter case we are done since for the desired $R'_{NE}(R'_{EN})$ we simply take $R_{NE}(R_{EN})$ while in the former case $\{i, j\}$ is $\{2, 3\}$, $\{1, 3\}$, $\{2, 4\}$, or $\{1, 4\}$.

- If $\{i, j\} = \{2, 3\}$, then we set $R'(x_1, x_2, x_3, x_4) \equiv R(x_1, x_2, x_3, x_4)$
- If $\{i, j\} = \{1, 3\}$, then we set $R'(x_1, x_2, x_3, x_4) \equiv R(x_2, x_1, x_3, x_4)$.
- If $\{i, j\} = \{2, 4\}$, then we set $R'(x_1, x_2, x_3, x_4) \equiv R(x_1, x_2, x_4, x_3)$.
- If $\{i, j\} = \{1, 4\}$, then we set $R'(x_1, x_2, x_3, x_4) \equiv R(x_2, x_1, x_4, x_3)$.

It is straightforward to check that in each of the four cases R' efficiently entails the formula $(E(x_1, x_2) \implies \underline{N}(x_3, x_4))$ (resp., $(N(x_1, x_2) \implies \underline{E}(x_3, x_4))$) if R efficiently entails $(E(x_1, x_2) \implies \underline{N}(x_3, x_4))$ (resp., $(N(x_1, x_2) \implies \underline{E}(x_3, x_4))$) and contains both an EN-tuple t_{EN} and a NE-tuple t_{NE} such that for all $t \in \{t_{EN}, t_{NE}\}$ we have $(t[2] = t[3])$. It completes the proof of the observation. \blacktriangleleft

In Section 3 we allowed to write $R_1 \cdots R_k(a_1, a_2)$ for k -tuples $a_1, a_2 \in A^k$ if and only if $R_i(a_1[i], a_2[i])$ holds for all $i \in [k]$. With reference to this notation we say that a_1, a_2 are $\{R_1, \dots, R_k\}$ -connected. Clearly, the relations $\{R_1, \dots, R_k\}$ do not have to be pairwise different. We write $n_{R_i(a_1, a_2)}$ to denote the number of coordinates at which R_j with $j \in [k]$ equals R_i . For example, we write $NE = (a, b)$ for $a, b \in A^3$ if $(a[1], b[1]) \in N$, $(a[2], b[2]) \in N$, and $a[3] = b[3]$ and say that a, b are $\{E, N, =\}$ -connected. In this case we have that $n_{E(a, b)} = n_{N(a, b)} = n_{=(a, b)} = 1$.

We say that a pair of tuples u, s is constant-injective if u is a constant tuple, s is an injective tuple and no entry in s equals the main value of u .

► **Observation 40.** *Let $R \in \{R'_{EN}, R'_{NE}\}$ and $m \in [k]$. If for all $\{E, N\}$ -connected constant-injective pairs $u, s \in A^k$ with $n_{E(u, s)} = m$ we have $(f(u), f(s))$ in E (resp. in N), then either*

- *for all constant-injective pairs $u, s \in A^k$ with $n_{E(u, s)} \in \{k - m, k - m + 1\}$ it holds $((f(u), f(s))$ in N (resp. in E), or*
- *R contains either an EE-tuple or a NN-tuple.*

Proof. We only handle the case where $(f(u), f(s)) \in E$ for all $\{E, N\}$ -connected constant-injective $u, s \in A^k$ with $n_{E(u, s)} = m$ and we show that in this case if R does not satisfy the first item, then R contains an EE-tuple. The proof of the other case is analogous. The difference is that we show the existence of a NN-tuple. Assume that there is a $\{E, N\}$ -connected constant-injective $u, s \in A^k$ such that $n_{E(u, s)} \in \{(k - m), (k - m + 1)\}$ and $(f(u), f(s)) \notin N$. Since f preserves a first-order expansion \mathbb{A} of the random graph, it also preserves E, N and \neq . Thus, we may assume that $(f(u), f(s)) \in E$.

Let now $t \in A^k$ be a tuple with the same main value as u such that none of the values in t occurs in s , $n_{E(t, s)} = k - m$ and t, s are $\{E, N\}$ -connected. If already $n_{E(u, s)} = k - m$, then we take t to be u , otherwise $n_{E(u, s)} = k - m + 1$ and then using the extension property we simply change a value of one of the entries in u so that the obtained t satisfy the desired properties. Further, either Case 1 or Case 2 in Observation 39 holds.

For Case 1 select a constant tuple u' with the main value d such that for all $i \in [k]$ the tuple $(t[i], s[i], s[i], u'[i])$ equals either t_{EN} or t_{NE} . By Observation 38 we may choose u' such that d is different from all values in t and s and hence the existence of u' follows by the extension property of the random graph. In particular, we have that $n_{E(s, u')} = k$. By the assumption, it follows that $(f(s), f(u')) \in E$. On the other hand, since t has the same main value as u , it follows that $(f(t), f(s)) \in E$. The tuples t_{EN} and t_{NE} are in R and hence $(t[i], s[i], s[i], u'[i])$ is in R for every $i \in [k]$ which is pp-definable in \mathbb{A} and hence preserved by f . It follows that R contains an EE-tuple. It completes the proof of the claim for Case 1 in Observation 39. In the proof for Case 2 we choose first an injective s' with values not occurring in t, s and then a constant u' with the main value d which again does not occur in none of t, s, s' so that for all $i \in [k]$ the tuple $(t[i], s[i], s'[i], u'[i])$ equals either t_{EN} or t_{NE} . Since in this case both t_{EN} and t_{NE} are injective we can clearly find first such s' and u using the extension property of the random graph. Then using the same reasoning as in Case 1 we show that also in Case 2 we obtain an EE-tuple in R . It completes the proof of the observation. \blacktriangleleft

We continue the proof of the lemma. Since f preserves E , we have that $(f(u), f(s)) \in E$ for all constant injective pairs $u, s \in A^k$ with $n_{E(u, s)} = k$. By one application of Observation 40, it follows that either R contains the desired EE-tuple or $(f(u), f(s)) \in N$ for all constant-injective pairs u, s with $n_{E(u, s)} = 1$. In the latter case, by another application of the claim we have that R contains the desired NN-tuple or $(f(u), f(s)) \in E$ for all constant-injective pairs u, s with $n_{E(u, s)} = n - 1$. By inductively continuing this process we obtain that $(f(u), f(s)) \in E$ for all constant-injective pairs u, s with $n_{E(u, s)} = i$ where $i \in [k]$, and on the other hand that $(f(u), f(s)) \in N$ for all constant-injective pairs u, s with $n_{E(u, s)} = k - i$ where $i \in [k]$ or R contains either an EE-tuple or a NN-tuple. Since $(f(u), f(s))$ cannot be in both E and N , we obtain that either an EE-tuple or a NN-tuple is in R . It follows that both R'_{EN} and R'_{NE} contain either an EE-tuple or a NN-tuple. Hence either R'_{EN} does not efficiently entail $(E(x_1, x_2) \rightarrow \underline{N}(x_3, x_4))$ or R'_{NE} does not efficiently entail $(E(x_1, x_2) \rightarrow \underline{N}(x_3, x_4))$. It implies the contradiction and completes the proof of the lemma. \square

O Proof of Lemma 29

Assume on the contrary that \mathbb{A} pp-defines R which is either a $[(E(x_1, x_2) \implies (x_3 = x_4), (\underline{E}(x_1, x_2) \wedge N(x_2, x_3) \wedge \underline{E}(x_3, x_4)))]$ -relation or a $[(x_1 = x_2) \implies E(x_3, x_4), (\underline{E}(x_1, x_2) \wedge N(x_2, x_3) \wedge \underline{E}(x_3, x_4))]$ -relation. In any case R contains both an E -tuple $t_{E=}$ and an $=E$ -tuple $t_{=E}$ such that for all $t \in \{t_{E=}, t_{=E}\}$ we have $(t[2], t[3]) \in N$. In the remainder of the proof we show that R preserved by h and the qnu-operation f from the formulation of the lemma containing both these tuples contains also an EE-tuples t_{EE} and a $==$ -tuple $t_{==}$. Observe that to this end it is enough to show that at least one of these tuples is in R . Indeed, $h(t_{E=}, t_{=E}, t_{EE})$ produces an $==$ -tuple and $h(t_{E=}, t_{=E}, t_{==})$ produces an EE -tuple. A relation that contains $t_{EE}, t_{E=}, t_{=E}$ and $t_{==}$ efficiently entails neither $(E(x_1, x_2) \implies (x_3 = x_4))$ nor $((x_1 = x_2) \implies E(x_3, x_4))$. It contradicts the assumption that R does efficiently entail one of these relations and will complete the proof of the lemma. We let k to be the arity of the qnu-operation f and continue with a simple observation.

► **Observation 41.** *Let $m \in [k]$. If for all $\{E, =\}$ -connected $u, s \in A^k$ such that u is constant and $n_{E(u, s)} = m$ it holds that $(f(u), f(s)) \in E$ (resp. $(f(u), f(s)) \in N$). Then either*

- *for all $\{E, =\}$ -connected $u, s \in A^k$ such that u is constant and $n_{E(u, s)} = \{k - m, k - m + 1\}$ we have $f(u) = f(s)$ (resp. $(f(u), f(s)) \in E$), or*

■ R contains an EE-tuple (resp. an ==-tuple).

Proof. We only handle the case where $(f(u), f(s)) \in E$ for all $\{E, =\}$ -connected $u, s \in A^k$ such that u is constant and $n_{E(u,s)} = m$. Assume that there exists a constant $u \in A^k$ and $s \in A^k$ such that u, s are $\{E, =\}$ -connected, $n_{E(u,s)} \in \{k-m, k-m+1\}$, and $(f(s) \neq f(u))$. Since f preserves R and therefore \underline{E} , we may assume that $(f(u), f(s)) \in E$. We will show that in this case R contains an EE-tuple. Select t to be a k -tuple with the same main value as u and such that $n_{E(t,s)} = k-m$. Observe that all entries in t, s comes from a single edge in C_ω^2 . Then, clearly, there exists $s', u' \in A^k$ of values originating from any other edge in C_ω^2 such that $n_{E(u',s')} = m$ and such that for all $i \in [k]$ it holds that $(t[i], s[i], s'[i], u'[i])$ is either $t_{E=}$ or $t_{=E}$. By the assumption, $(f(s'), f(u')) \in E$, and hence $(f(t), f(s), f(s'), f(u'))$ is an EE-tuple in R . It completes the proof of the claim. ◀

Since f preserves E , we have that $(f(u), f(s)) \in E$ for all $u, s \in A^k$ with $n_{E(u,s)} = k$. By one application of Observation 41, it follows that either R contains the desired EE-tuple or $(f(u) = f(s))$ for all $\{E, =\}$ -connected $u, s \in A^k$ where u is constant and $n_{E(u,s)} = 1$. If the application did not return the desired tuple, then by another application of the claim we have that $(f(u), f(s)) \in E$ for all $\{E, =\}$ -connected $u, s \in A^k$ such that u is constant and $n_{E(u,s)} = n-1$. By inductively continuing this process we obtain that $(f(u), f(s)) \in E$ for all $\{E, =\}$ -connected $u, s \in A^k$ such that u is constant and $n_{E(u,s)} = i$ where $i \in [k]$, and on the other hand that $(f(u) = f(s))$ for all constant u and all s with $n_{E(u,s)} = k-i$ where $i \in [k]$. Since $(f(u), f(s))$ cannot be in both E and $=$, we obtain a contradiction. Thus, in consequence R contains all: an EE-tuple and an ==-tuple. It implies a contradiction and completes the proof of the lemma. ◻

P Proof of Lemma 30

The proof goes along the lines of the proof of Lemma 29. We assume on the contrary that \mathbb{A} pp-defines R which is either a $[(E(x_1, x_2) \implies (E(x_3, x_4)), (\underline{E}(x_1, x_2) \wedge N(x_2, x_3) \wedge \underline{E}(x_3, x_4)))]$ -relation or a $[(x_1 = x_2) \implies (x_3 = x_4), (\underline{E}(x_1, x_2) \wedge N(x_2, x_3) \wedge \underline{E}(x_3, x_4))]$ -relation and observe that in any case R contains both an EE-tuple t_{EE} and a ==-tuple $t_{==}$ such that for all $t \in \{t_{EE}, t_{==}\}$ we have $(t[2], t[3]) \in N$. We will show that such R preserved by the oligopotent qnu-operation f from the formulation of the lemma and h contains additionally an E=-tuple $t_{E=}$ and an =E-tuple $t_{=E}$. In fact, it is enough to show that R contains one of these tuples. Indeed, $h(t_{EE}, t_{==}, t_{E=})$ produces an =E-tuple and $h(t_{EE}, t_{==}, t_{=E})$ an E=-tuple. A relation containing $\{t_{EE}, t_{==}, t_{E=}, t_{=E}\}$ efficiently entails neither $(E(x_1, x_2) \implies (E(x_3, x_4)))$ nor $((x_1 = x_2) \implies (x_3 = x_4))$. It contradicts the assumption and will complete the proof of the lemma. We let k to be the arity of f and continue with a simple observation.

► **Observation 42.** Let $m \in [k]$. If for all $\{E, =\}$ -connected $u, s \in A^k$ such that u is constant and $n_{E(u,s)} = m$ it holds that $(f(u), f(s)) \in E$. Then either

- for all $\{E, =\}$ -connected $u, s \in A^k$ such that u is constant and $n_{E(u,s)} = m-1$ we have $(f(u), f(s)) \in E$, or
- R contains an E=-tuple.

Proof. Assume on the contrary that there exists a constant tuple $u \in A^k$ and $s \in A^k$ such that u, s are $\{E, =\}$ -connected, $n_{E(u,s)} = m-1$ and $(f(s), f(u)) \notin E$. Since f preserves R and hence \underline{E} , we have $(f(s) = f(u))$. We will show that in this case R contains an E=-tuple. Select t to be a k -tuple with the same main value as u and such that $n_{E(t,s)} = m$. Observe that all entries in t, s come from a single edge in C_ω^2 . Then, clearly, there exists $s', u' \in A^k$

of values originating from any other edge in C_ω^2 such that $n_{E(s',u')} = m$ and such that for all $i \in [k]$ it holds that $(u'[i], s'[i], s[i], t[i])$ is either t_{EE} or $t_{==}$. Since t has the same main-value as u we have $(f(s) = f(t))$. Since $n_{E(s',u')} = m$, by the assumption, $(f(s'), f(u')) \in E$. It follows that $(f(u'), f(s'), f(s), f(t))$ is an E -tuple in R . It completes the proof of the observation. \blacktriangleleft

The operation f preserves E and hence $(f(u), f(s)) \in E$ for all $u, s \in A^k$ with $n_{E(u,s)} = k$. By Observation 42, it follows that either R contains an E -tuple or $(f(u), f(s)) \in E$ for all constant $u \in A^k$ and $s \in A^k$ such that u, s are $\{E, =\}$ -connected and $n_{E(u,s)} = k - 1$. By inductively continuing this argument, we either obtain the desired E -tuple or we have $(f(u), f(s)) \in E$ for all constant $u \in A^k$ and $s \in A^k$ such that u, s are $\{E, =\}$ -connected and $n_{E(u,s)} = k - 1, k - 2, \dots, 1$. The latter contradicts the fact that f is a quasi near-unanimity operation. Indeed, for any constant u and t with the same main-value as u which are $\{E, =\}$ -connected and $n_{E(u,s)} = 1$, we have $(f(u), f(s)) \in E$ and in consequence $(f(u) \neq f(s))$. It completes the proof of the lemma. \square