# UNIFORM LIMITS OF $\mathcal{B}_1^{**}$ -FUNCTIONS

#### PIOTR SWOROWSKI, WALDEMAR SIEG

ABSTRACT. We characterise the class of uniform limits of functions from Pawlak's class  $\mathcal{B}_{1}^{**}$ . The resulting class  $u\mathcal{S}_{1}$ , which contains functions with the oscillation rank one, is discussed in connection with its linear span. We apply a general, topological space, setting in our discussion.

#### 1. Introduction

The so-called Baire-one-star ( $\mathcal{B}_1^*$ ) property of a real-valued function, defined on an interval in the real line, has received certain attention over recent couple of decades, often in connection with generalised differentiation and integration, where it is investigated if some generalised primitives or indefinite integrals necessarily obey to this, stronger than Baire-one, property; see, e.g., [2]. A different kind of application reveals a work by one of the authors [10], where the  $\mathcal{B}_1^*$  property was used as a main tool for a characterisation of integrability analogous to the Lebesgue theorem on the Riemann integral.<sup>1</sup>

Under the Baire-one-star name, and in generalised differentiation connection, the property was introduced by Richard J. O'Malley in [11], but is also known as the Baire half property after Harvey Rosen's paper [13], or as piecewise continuity [3]. (See [6] for more info, various descriptions of  $\mathcal{B}_1^*$  and related properties.) For the present note we can accept the following definition: an  $f: X \to Y, X, Y$  topological spaces, has the  $\mathcal{B}_1^*$  property if for every closed  $D \subset X$ , there is an open  $O \subset X$ ,  $D \cap O \neq \emptyset$ , such that the restriction  $f \upharpoonright (D \cap O)$  is continuous (in its domain).

Certain limit properties of  $\mathcal{B}_1^*$  make this class technically useful in many Baire hierarchy related problems. A general result found in Kuratowski's monograph [7, pp. 294–295] says that, for an arbitary ordinal  $\alpha$ , any Baire class  $\alpha$  function can be obtained as the uniform limit of a sequence of Baire- $\alpha$  functions all whose ranges are discrete sets. For  $\alpha=1$  and in, e.g., the  $\mathbb{R} \to \mathbb{R}$  case, the discrete range property implies  $\mathcal{B}_1^*$  property. Thus one can claim that Baire class one functions are exactly limits of uniformly convergent sequences of Baire-one-star functions. Hence, for approximation problems, the  $\mathcal{B}_1^*$  property may be seen, in a sense, exactly as useful as  $\mathcal{B}_1$ .

The main objective of this note is to provide a simple characterisation for the uniform closure of the class of  $\mathcal{B}_1^{**}$ -functions, a subclass of  $\mathcal{B}_1^*$ , first considered by Ryszard J. Pawlak in [12]. Since the characterisation theorem allows a simple and brief proof, we strove for presenting it in as much as possible general setting.

1

 $Key\ words\ and\ phrases.$  Baire-one-star function, Pawlak's class  $\mathcal{B}_1^{**}$ , Małek's classes  $\mathcal{S}_{\alpha}$ , classes  $u\mathcal{S}_{\alpha}$ , oscillation rank, Katětov–Tong insertion theorem.

<sup>&</sup>lt;sup>1</sup>Quite recently,  $B_1^*$  functions received some attention in connection with dynamical systems, as so-called *k*-continuous functions or right  $B_1$  compositors [1].

### 2. The characterisation

Let  $(X, \mathcal{T})$  be a topological space,  $(Y, \rho)$  a metric space. For a function  $f: X \to Y$  and  $x \in X$  we denote with  $\omega_f(x)$  the oscillation of f at x, that is,  $\omega_f(x) = \inf_O \sup_{\xi \in O} \varrho(f(x), f(\xi))$ , where the inf ranges over all neighbourhoods  $O \ni x$ . This should not be confused with the oscillation of f on a subset  $Z \subset X$ , which is the diameter of f(Z). (Clearly,  $\omega_f(x) = 0$  iff f is continuous at x.) For two functions  $f, g: X \to Y$  we set  $\varrho_\infty(f, g) = \sup_{\xi \in X} \varrho(f(\xi), g(\xi))$ .

By definition, a function  $f: X \to Y$  is  $\mathcal{B}_1^{**}$  if the restriction of f to the set of its discontinuity points,  $D_f$ , if nonempty, is continuous.<sup>2</sup> The notation  $\mathcal{B}_1^{**}(X,Y)$  stands for the class of all  $\mathcal{B}_1^{**}$ -functions defined over X with values in Y. If Y is the real line with the natural metric, we write instead just  $\mathcal{B}_1^{**}(X)$ .

**Theorem 1.** Assume  $f_n \rightrightarrows f$ , where all  $f_n \in \mathcal{B}_1^{**}(X,Y)$ . The following condition holds:

(**\***)

if  $\varepsilon > 0$ , the restriction of f to the set  $D_f^{\varepsilon} = \{ \xi \in X : \omega_f(\xi) \geq \varepsilon \}$  is continuous.

*Proof.* Let  $x \in D_f^{\varepsilon}$ . Suppose to the contrary that, for some  $\eta > 0$ , in every  $O \in \mathcal{T}$ ,  $O \ni x$ , there is a  $\xi \in D_f^{\varepsilon}$  such that  $\varrho(f(x), f(\xi)) \ge \eta$ . Fix an  $n \in \mathbb{N}$  large enough so that  $\varrho_{\infty}(f, f_n) < \min \{\eta/4, \varepsilon/2\}$ . Then, if  $\xi$  is chosen for a given  $O \in \mathcal{T}$ ,  $O \ni x$ , as above, we have

$$\varrho(f_n(x), f_n(\xi)) \ge \varrho(f(x), f(\xi)) - \varrho(f_n(\xi), f(\xi)) - \varrho(f_n(x), f(x)) > \eta - 2 \cdot \frac{\eta}{4} = \frac{\eta}{2},$$
  
so  $x \in D_{f_n}$ . Moreover,

$$\omega_{f_n}(\xi) > \omega_f(\xi) - 2 \cdot \frac{\varepsilon}{2} \ge 0.$$

It means that all  $\xi \in D_{f_n}$  and so  $f_n \upharpoonright D_{f_n}$  is discontinuous at x, which arrives in contradiction with  $f_n \in \mathcal{B}_1^{**}(X,Y)$ .

For the opposite implication we assume the domain space X is  $T_5$  (i.e., hereditarily normal and  $T_1$ ).

**Theorem 2.** Assume  $f: X \to \mathbb{R}$  satisfies  $(\star)$ . Then there is a sequence  $(f_n)_{n=1}^{\infty} \subset \mathcal{B}_1^{**}(X)$  such that  $f_n \rightrightarrows f$ .

*Proof.* Let  $\varepsilon > 0$ . Note that  $D_f^{\varepsilon}$  is a closed subset of X. At every  $x \in U_f^{\varepsilon} = X \setminus D_f^{\varepsilon}$  define  $h_1(x)$  and  $h_2(x)$  to be correspondingly

$$\inf_{O} \sup_{\xi \in O} f(\xi) \qquad \text{and} \qquad \sup_{O} \inf_{\xi \in O} f(\xi),$$

where the outer extrema range over all neighbourhoods  $O \ni x$ . Note that  $h_1$  and  $h_2$ , as functions defined on  $U_f^{\varepsilon}$ , are respectively upper and lower semicontinuous. It is easy to check that

$$f(x) + \varepsilon > f(x) + \omega_f(x) = h_1(x) \ge f(x) \ge h_2(x) = f(x) - \omega_f(x) > f(x) - \varepsilon,$$

whence  $h_2(x) + \varepsilon > f(x) > h_1(x) - \varepsilon$ ,  $x \in U_f^{\varepsilon}$ . As the subspace  $U_f^{\varepsilon}$  of X is  $T_4$ , from the Katětov-Tong insertion theorem [4, 14], there exists a continuous function  $g \colon U_f^{\varepsilon} \to \mathbb{R}$  such that, at every  $x \in U_f^{\varepsilon}$ ,

$$f(x) + \varepsilon \ge h_2(x) + \varepsilon \ge g(x) \ge h_1(x) - \varepsilon \ge f(x) - \varepsilon.$$

<sup>&</sup>lt;sup>2</sup>In such case, notice,  $D_f$  must be nowhere dense in X, so that f has also  $\mathcal{B}_1^*$  property.

So,  $|f(x) - g(x)| \leq \varepsilon$ ,  $x \in U_f^{\varepsilon}$ . Define  $\tilde{f}$  as f over  $D_f^{\varepsilon}$  and as g over  $U_f^{\varepsilon}$ . Obviously,  $\varrho_{\infty}(f, \tilde{f}) \leq \varepsilon$ . Note that  $\tilde{f} \in \mathcal{B}_1^{**}(X)$ . Indeed,  $D_{\tilde{f}} \subset D_f^{\varepsilon}$ , so the restriction  $\tilde{f} \upharpoonright_{D_{\tilde{f}}} = f \upharpoonright_{D_{\tilde{f}}}$  is continuous.  $\square$ 

**Corollary 1.** Let X be a  $T_5$  topological space and  $f: X \to \mathbb{R}$ . The following two statements are equivalent:

- f satisfies  $(\star)$ ;
- there is a sequence  $(f_n)_{n=1}^{\infty} \subset \mathcal{B}_1^{**}(X)$  with  $f_n \rightrightarrows f$  (abbr.  $f \in u\mathcal{B}_1^{**}(X)$ ).

# **Problem 3.** Is $T_5$ essential for Theorem 2?

The problem boils down to finding the function g (cf. the proof). Despite the Katětov-Tong theorem can be applied only if the underlying space X is  $T_5$  (and so  $U_f^{\varepsilon}$  is  $T_4$ ), the existence of g can be granted in some particular topological spaces that are far from  $T_5$ , e.g., in  $\mathbb{R}$  with co-countable topology, which isn't even Hausdorff.

The condition  $(\star)$  in terms of [5] means that the oscillation rank of f,  $\beta(f)$ , equals (at most) one. The iterates of  $D_f^{\varepsilon}$  considered in the next section correspond to larger (still finite) values of  $\beta(f)$ , namely,  $f \in u \mathcal{S}_n$  iff  $\beta(f) \leq n$ . The oscillation rank has been introduced, with finite and countable values, in order to serve for classifying members of the Baire one class. The range of  $\beta(f)$  in the present note is confined to  $\mathbb{N}$ , that is, to f living in the first small Baire class [5].

# 3. Iterates of $D_f^{arepsilon}$ and the linear span of $u\mathcal{B}_1^{**}$

We proceed with a discussion on sums of functions from  $u\mathcal{B}_1^{**}$ . To this end, we need to recall some concepts (although with modified notation) from [8, 9] and then extend them. Let X be a topological space,  $f\colon X\to\mathbb{R}$ , and  $D_f^0$  the closure of set of discontinuity points of f. Moreover, let  $f^0=f\upharpoonright\operatorname{cl} D_f^0$ . By induction, given  $f^{0,\ldots,0}$  (or briefly  $f^{n*0}$ ) and  $D_f^{0,\ldots,0}$  (or  $D_f^{n*0}$ ),  $n\in\mathbb{N}$  is the number of upper indices here, we set  $D_f^{(n+1)*0}$  to be the closure of the set of discontinuity points of  $f^{n*0}\colon D_f^{n*0}\to\mathbb{R}$ . We denote with  $\mathcal{S}_n(X)$  (or  $\mathcal{S}_n$  when the domain is understood) the class of all f on X such that  $D_f^{(n+1)*0}=\emptyset$ . Note that (cf. page 2)  $\mathcal{B}_1^{**}(X)=\mathcal{S}_1(X)$ .

Now let  $\varepsilon>0$ . Recall that  $D_f^\varepsilon$  is the set of points  $x\in X$  such that  $\omega_f(x)\geq \varepsilon$ , and let  $f^\varepsilon$  be  $f\upharpoonright D_f^\varepsilon$ . Inductively, given arbitrary  $\varepsilon_1,\dots,\varepsilon_n,\varepsilon_{n+1}>0,\ n\in\mathbb{N}$ , we set  $D_f^{\varepsilon_1,\dots,\varepsilon_n,\varepsilon_{n+1}}$  to be the set of points  $x\in D_f^{\varepsilon_1,\dots,\varepsilon_n}$  such that the oscillation of  $f^{\varepsilon_1,\dots,\varepsilon_n}$  (defined on  $D_f^{\varepsilon_1,\dots,\varepsilon_n}$ ) at x, written as  $\omega_f^{\varepsilon_1,\dots,\varepsilon_n}(x)$ , is  $\geq \varepsilon_{n+1}$ ;  $f^{\varepsilon_1,\dots,\varepsilon_n,\varepsilon_{n+1}}=f\upharpoonright D_f^{\varepsilon_1,\dots,\varepsilon_n,\varepsilon_{n+1}}$ ,  $U_f^{\varepsilon_1,\dots,\varepsilon_n}=X\setminus D_f^{\varepsilon_1,\dots,\varepsilon_n}$ . We will apply the following abbreviation in case all bounds  $\varepsilon_1,\dots,\varepsilon_n$  are equal:

$$D_f^{\overbrace{\varepsilon,\ldots,\varepsilon}^n}=D_f^{n*\varepsilon},\qquad f^{\overbrace{\varepsilon,\ldots,\varepsilon}^n}=f^{n*\varepsilon},\qquad \omega_f^{\overbrace{\varepsilon,\ldots,\varepsilon}^n}=\omega_f^{n*\varepsilon}.$$

The class  $u\mathcal{S}_n(X)$ , or just  $u\mathcal{S}_n$ , is defined as the class of all functions  $f\colon X\to\mathbb{R}$  such that for arbitrary  $\varepsilon_1,\ldots,\varepsilon_{n+1},\,D_f^{\varepsilon_1,\ldots,\varepsilon_{n+1}}=\emptyset$  (note that it is tantamount to  $D_f^{(n+1)*\varepsilon}=\emptyset$  for every  $\varepsilon>0$ ). By this definition and by Corollary 1, provided X is  $T_5,\,u\mathcal{B}_1^{**}(X)=u\mathcal{S}_1(X)$ .

**Theorem 4.** If a sequence  $(f_m)_{m=1}^{\infty} \subset S_n(X)$  tends uniformly to an f, then  $f \in uS_n(X)$ .

Proof. We will proceed by induction on n. The case n=1 is the content of Theorem 1. Assume the claim holds for some n and take a sequence  $(f_m)_{m=1}^{\infty} \subset \mathbb{S}_{n+1}(X)$ ,  $f_m \rightrightarrows f$ . Take an  $\varepsilon > 0$ . For almost all m,  $||f - f_m||_{\infty} < \varepsilon/2$  and so (for these m)  $D_f^{\varepsilon} \subset D_{f_m}$ ; the latter means  $f_m$  is in  $\mathbb{S}_n$  on  $D_f^{\varepsilon}$ . By the assumption, as  $f_m \rightrightarrows f$  on  $D_f^{\varepsilon}$ ,  $f^{\varepsilon} \in u \mathbb{S}_n(D_f^{\varepsilon})$ , so  $D_f^{(n+1)*\varepsilon} = \emptyset$ .

**Theorem 5.** Let X be  $T_5$ ,  $n \in \mathbb{N}$ . If  $f \in uS_n(X)$ , then there exists a sequence  $(f_m)_{m=1}^{\infty} \subset S_n(X)$  such that  $f_m \rightrightarrows f$ .

Proof. Again induction on n. The result holds for n=1 (Theorem 2). Assume the claim is fulfilled for some n and consider an  $f \in u \mathbb{S}_{n+1}(X)$  and  $\varepsilon > 0$ . By definition,  $f^{\varepsilon} \in u \mathbb{S}_{n}(D_{f}^{\varepsilon})$ . Apply the assumption to  $f^{\varepsilon}$  and pick a function  $g \in \mathbb{S}_{n}(D_{f}^{\varepsilon})$  with  $\|f^{\varepsilon} - g\|_{\infty} < \varepsilon$ . Moreover, as  $\omega_{f}(x) < \varepsilon$  at every  $x \in U_{f}^{\varepsilon}$ , arguing like in the proof of Theorem 2, we can arrive to a continuous function  $h \colon U_{f}^{\varepsilon} \to \mathbb{R}$  such that (over  $U_{f}^{\varepsilon}$ )  $\|f - h\|_{\infty} < \varepsilon$ . So, g and h taken jointly form a function  $\tilde{f}$  on X satisfying  $\|f - \tilde{f}\|_{\infty} < \varepsilon$ . Moreover,  $\tilde{f} \in \mathbb{S}_{n+1}(X)$ . Indeed,  $\operatorname{cl} D_{\tilde{f}} \subset D_{f}^{\varepsilon}$  and so  $\tilde{f} \cap D_{\tilde{f}} \subset D_{\tilde{f}}$ . Thus  $\tilde{f} \cap D_{\tilde{f}} \subset D_{\tilde{f}}$ . The proof is over.

**Theorem 6.** Let X be  $T_5$ ,  $f \in uS_n(X)$ , and  $g \in uS_m(X)$ . Then  $f+g \in uS_{n+m}(X)$ .

*Proof.* By Theorem 5 there are sequences  $(f_s)_{s=1}^{\infty} \subset S_n(X)$ ,  $(g_s)_{s=1}^{\infty} \subset S_m(X)$  such that  $f_s \rightrightarrows f$  and  $g_s \rightrightarrows g$ . By [8],  $f_s + g_s \in S_{n+m}(X)$  for every s. Since  $f_s + g_s \rightrightarrows f + g$ , from Theorem 4,  $f + g \in uS_{n+m}(X)$ .

As it is shown in [8],  $S_n + S_m = S_{n+m}$  over any topological space. In particular, every function  $f \in S_n$  can be written as the sum of at most n functions from  $\mathcal{B}_1^{**} = S_1$ . We were unable to prove the analogous result for  $uS_n$ , however we show that every  $f \in S_n$  can be decomposed into finitely many terms from  $uS_1$  (Theorem 9).

We will say a space X has the property (T) if the following assertion, more general than the Tietze extension theorem is true: Let  $Y \subset X$  be a closed subspace,  $f: Y \to \mathbb{R}$  an arbitrary function. Then there exists a continuous function  $g: X \setminus Y \to \mathbb{R}$  such that if  $h = f \cup g$ ,  $\omega_f(x) = \omega_h(x)$  at every  $x \in Y$ . Clearly, every space with the property (T) must be  $T_4$ . It is easy to see that, e.g.,  $\mathbb{R}$  with the natural topology has this property.

**Problem 7.** Does every  $T_4$  topological space have property (T)?

**Lemma 8.** Let X be a  $T_6$  space. Then every open subset  $O \subset X$  can be written as a union  $\bigcup_{k=1}^{\infty} E_k$  of closed sets  $E_k$  such that  $E_k \subset \text{int } E_{k+1}$  for every  $k \in \mathbb{N}$ .

Proof. By the  $T_6$  property of X we have  $O = \bigcup_{k=1}^{\infty} D_k$ , where all  $D_k$ 's are closed. Set  $E_1 = D_1$ . Assume  $E_k \supset D_k$  is defined and take two open and disjoint sets U, V with  $E_k \cup D_{k+1} \subset U$  and  $X \setminus O \subset V$ . Define  $E_{k+1} = \operatorname{cl} U$ . Clearly,  $D_k \subset E_k \subset U \subset \operatorname{int} \operatorname{cl} U = \operatorname{int} E_{k+1} \subset E_{k+1} \subset X \setminus V \subset O$  and so  $O = \bigcup_{k=1}^{\infty} D_k \subset \bigcup_{k=1}^{\infty} E_k \subset O$ .  $\square$ 

**Theorem 9.** Let  $f \in uS_{n+1}(X)$ ,  $n \in \mathbb{N}$ , X be a  $T_6$  space with property (T). Then there exist functions  $f_1, f_3 \in uS_n(X)$ ,  $f_2 \in uS_1(X)$ , such that  $f = f_1 + f_2 + f_3$ .

*Proof.* Set  $D = D_f^1$ . Then both restrictions  $f \upharpoonright D$  and  $f \upharpoonright (X \setminus D)$  are in  $u \mathcal{S}_n$ .

The term  $f_1$  is defined as follows:  $f_1 = f$  on D, while on  $X \setminus D$  it is extended continuously so that  $\omega_{f_1}(x) = \omega_f^1(x)$  at every  $x \in D$  (property (T)). Clearly,  $f_1 \in$ 

 $u\mathcal{S}_n(X)$ . Indeed, if  $\varepsilon > 0$ , then  $D_{f_1}^{\varepsilon} \subset D$  and so  $D_{f_1}^{\varepsilon} = D_f^{1,\varepsilon}$ ; hence the restriction of  $f_1$  to  $D_{f_1}^{n*\varepsilon} = D_f^{1,n*\varepsilon}$  (if nonempty) is continuous.

Write down  $X \setminus D = \bigcup_{k=1}^{\infty} E_k$ , where the sequence  $(E_k)_{k=1}^{\infty}$  of closed sets is as in Lemma 8. Note that the restriction of  $f - f_1$  (like of f alone) to the set  $X \setminus D$  is in  $uS_n(X \setminus D)$ . Thus,  $(f - f_1) \upharpoonright (D_f^{n*1} \cap E_1) = g_1$  is continuous. Moreover, for the same reason,  $(f - f_1) \upharpoonright (D_f^{n*1} \cap \operatorname{int} E_1 \cup D_f^{n*1/2} \cap (E_2 \setminus \operatorname{int} E_1)) = g_2$  is a continuous function defined on a closed subset of  $D_f^{n*1/2} \cap E_2$ . And so on by induction—note that the restriction of  $f - f_1$  to the (closed) set

$$\bigcup_{s=0}^{k} \left( D_f^{n*1/(s+1)} \cap (E_{s+1} \setminus \operatorname{int} E_s) \right) \subset D_f^{n*1/(k+1)} \cap E_{k+1}, \qquad k \in \mathbb{N},$$

(here  $E_1 \setminus \text{int } E_0$  we accept as int  $E_1$ ), called  $g_{k+1}$ , is continuous. Having in mind the property of  $(E_k)_{k=1}^{\infty}$  described in Lemma 8, one can claim that also the limit  $\lim_{k\to\infty} g_k$ , which is the restriction of  $f-f_1$  to the set

(1) 
$$\bigcup_{s=0}^{\infty} \left( D_f^{n*1/(s+1)} \cap (E_{s+1} \setminus \operatorname{int} E_s) \right),$$

is a continuous function and its domain, (1), is a closed subset of  $X \setminus D$ . Thus, from the normality of  $X \setminus D$ , there is a continuous extension g of  $\lim_{k\to\infty} g_k$  defined on  $X \setminus D$ . Set  $f_2$  as g on  $X \setminus D$  and as 0 on D. Clearly,  $f_2 \in \mathcal{B}_1^{**} \subset u\mathcal{S}_1$ .

It remains to check that  $f_3=f-f_1-f_2$  is a  $u\mathcal{S}_n$  function. Indeed, for arbitrary positive  $\varepsilon<1$  and i (as  $f_1$  and  $f_2$  were continuous in  $X\setminus D$ ),  $D_f^{i*\varepsilon}\setminus D=D_{f_3}^{i*\varepsilon}\setminus D$ . Hence  $D_{f_3}^{n*\varepsilon}\setminus D=\emptyset$  (since f is  $u\mathcal{S}_n$  on  $X\setminus D$ ). Take  $k\in\mathbb{N}$  such that  $\varepsilon>1/k$ . By definition,  $f_3(x)=0$  at all  $x\in D$  and also at every  $x\in D_f^{n*1/k}\setminus E_{k-1}\subset D_f^{n*\varepsilon}\setminus E_{k-1}$ . As  $E_{k-1}$  is a closed set,  $f_3^{n*\varepsilon}$  is continuous at every x (from its domain) that lies in D. Concluding,  $f_3^{n*\varepsilon}$  is a continuous function and so  $f_3\in u\mathcal{S}_n$ .

By induction we obtain

**Corollary 10.** Let f and X be as in the theorem above. Then f can be written as the sum of at most  $2^{n+1} - 1$  many terms from  $uS_1(X)$ .

It may be of some interest to have the bound  $2^{n+1} - 1$  above improved. It might be well even n + 1.

## References

- A. Alikhani-Koopaei, On chain recurrent sets of typical bounded Baire one functions, Topology and its Applications, 257 (2019), 1–10.
- R. A. Gordon, Some comments on an approximately continuous Khintchine integral, Real Analysis Exchange, 20 №2 (1994/95), 831–841.
- [3] J. E. Jayne, C. A. Rogers, First level Borel functions and isomorphisms, Jour. Math. Pures et Appl., 61 (1982), 177–205.
- [4] M. Katětov, On real-valued functions in topological spaces, Fundamenta Math., 38 (1951), 85–91.
- [5] A. S. Kechris, A. Louveau, A classification of Baire class 1 functions, Transactions of the American Mathematical Society, 318 №1 (1990), 209–236.
- [6] B. Kirchheim, Baire one star functions, Real Analysis Exchange, 18 №2 (1992/93), 385–399.
- [7] C. Kuratowski, Topologie, 4<sup>th</sup> French ed., vol. 1, Monografie Matematyczne, no. 20, PWN, Warszawa, 1958.

- [8] A. Łukasiewicz (neé Małek), A classification of Baire one star functions in topological spaces, Atti Semin. Mat. Fis. Univ. Modena Reggio Emilia, 54 № 1-2 (2006), 175–181.
- [9] A. Małek, A classification of Baire one star functions, Real Analysis Exchange, 32 №1 (2006/07), 205–211.
- [10] A. Maliszewski, P. Sworowski, A characterization of  $H_1$ -integrable functions, Real Analysis Exchange, 28 Nº1 (2002/03), 93–104.
- [11] R. J. O'Malley, Baire\*1, Darboux functions, Proceedings of the American Mathematical Society, 60 (1976), 187–192.
- [12] R. J. Pawlak, On some class of functions intermediate between the class B<sub>1</sub><sup>\*</sup> and the family of continuous functions, Tatra Mt. Math. Publ., 19 (2000), 135–144.
- [13] H. Rosen, Darboux Baire-.5 functions, Proceedings of the American Mathematical Society, 110 №1 (1990), 285–286.
- [14] H. Tong, Some characterizations of normal and perfectly normal spaces, Duke Math J., 19 №2 (1952), 289–292.

Kazimierz Wielki University Institute of Mathematics Powstańców Wielkopolskich 2 85-090 Bydgoszcz Poland

E-MAIL: piotrus@ukw.edu.pl E-MAIL: waldeks@ukw.edu.pl