LARGE DEVIATIONS PRINCIPLES FOR SYMPLECTIC DISCRETIZATIONS OF STOCHASTIC LINEAR SCHRÖDINGER EQUATION

CHUCHU CHEN, JIALIN HONG, DIANCONG JIN, AND LIYING SUN

ABSTRACT. In this paper, we consider the large deviations principles (LDPs) for the stochastic linear Schrödinger equation and its symplectic discretizations. These numerical discretizations are the spatial semi-discretization based on spectral Galerkin method, and the further full discretizations with symplectic schemes in temporal direction. First, by means of the abstract Gärtner–Ellis theorem, we prove that the observable $B_T = \frac{u(T)}{T}$, T>0 of the exact solution u is exponentially tight and satisfies an LDP on $L^2(0,\pi;\mathbb{C})$. Then, we present the LDPs for both $\{B_T^M\}_{T>0}$ of the spatial discretization $\{u^M\}_{M\in\mathbb{N}}$ and $\{B_N^M\}_{N\in\mathbb{N}}$ of the full discretization $\{u^M\}_{M,N\in\mathbb{N}}$, where $B_T^M = \frac{u^M(T)}{T}$ and $B_N^M = \frac{u_N^M}{N\tau}$ are the discrete approximations of B_T . Further, we show that both the semi-discretization $\{u^M\}_{M\in\mathbb{N}}$ and the full discretization $\{u^M\}_{M,N\in\mathbb{N}}$ based on temporal symplectic schemes can weakly asymptotically preserve the LDP of $\{B_T\}_{T>0}$. These results show the ability of symplectic discretizations to preserve the LDP of the stochastic linear Schrödinger equation, and first provide an effective approach to approximating the LDP rate function in infinite dimensional space based on the numerical discretizations.

1. Introduction

The stochastic Schrödinger equation, as an important stochastic Hamiltonian partial differential equation, is widely used to model the propagation of dispersive waves in inhomogeneous or random media (see e.g., [12]), and possesses the infinite dimensional stochastic symplectic geometric structure. To numerically inherit the geometric structure of the stochastic Schrödinger equation, [2] proposes the infinite dimensional stochastic symplectic algorithms and considers the semi-discretizations, such as the stochastic symplectic Runge–Kutta methods. Moreover, the full discretizations based on the stochastic symplectic methods in temporal direction are also proposed (see e.g., [2, 4, 5, 10, 11] and references therein). The numerical experiments show that the stochastic symplectic discretizations are more stable in the long-time simulation than the non-symplectic ones. In this paper, we aim to deepen the understanding of the long-time asymptotical behavior and probabilistic characteristics of stochastic symplectic methods from the perspective of LDP. More precisely, we study the LDPs for both the stochastic linear Schrödinger equation and its numerical discretizations, and investigate the ability of symplectic discretizations to asymptotically preserve the LDP of the original system.

The theory of large deviations has been applied to many other branches of sciences, for example statistical physics, finance, engineering information theory ([14, 15]). It is concerned with the exponential decay of probabilities of very rare events, where the decay rate is characterized by the LDP rate function. In some cases, LDP rate functions describe steady rate and fluctuations of physical quantities, such as the entropy or free energy of statistical systems (see e.g., [9]).

Key words and phrases. large deviations principle; symplectic discretizations; stochastic Schrödinger equation; rate function; exponential tightness.

This work is supported by National Natural Science Foundation of China (Nos. 91630312, 11971470, 11871068, 11926417, 11711530017).

In this paper, we consider the following stochastic linear Schrödinger equation

$$du = i\Delta u dt + i\alpha dW(t), \qquad t > 0,$$

$$u(0) = u_0 \in H_0^1(0, \pi),$$
(1.1)

where $\alpha > 0$, Δ is the Laplace operator with the Dirichlet boundary condition, and W is an $L^2(0,\pi;\mathbb{R})$ -valued Q-Wiener process defined on a complete filtered probability space $(\Omega,\mathscr{F},\{\mathscr{F}_t\}_{t\geq 0},\mathbf{P})$ with $\{\mathscr{F}_t\}_{t\geq 0}$ satisfying the usual conditions. The mass $\|u\|_{H^0}^2 = \int_0^{\pi} |u(x)|^2 dx$ of (1.1) is an important physical quantity with $H^0 := L^2(0,\pi;\mathbb{C})$, which is conservative if $\alpha = 0$. However, in the stochastic setting, it grows linearly in the mean sense, i.e., $\mathbf{E} \| u(T) \|_{H^0}^2 = \mathbf{E} \| u_0 \|_{H^0}^2 + \alpha^2 T \operatorname{tr}(Q)$. Markov's inequality yields that the quantity $||B_T||_{H^0}$ tends to zero in probability, where $B_T := \frac{u(T)}{T}$. In order to characterize the speed of convergence or give an exponential tail estimate, we investigate the LDP of $\{B_T\}_{T>0}$ on H^0 . Our idea is to use the abstract Gärtner-Ellis theorem, which involves the existence of the logarithmic moment generating function and exponential tightness. The Gaussian property of the exact solution on H^0 with the real inner product is analyzed to give the logarithmic moment generating function of $\{B_T\}_{T>0}$. A prerequisite of the exponential tightness is to find the compact subset of H^0 , under the non-compactness of the Schrödinger group, such that the probabilities of $\{B_T\}_{T>0}$ escaping from the compact subset is exponentially small. This relies on two skills: One is that the regularity of u on H^1 gives a series of compact sets in H^0 , and the other is that the Fernique theorem yields the estimate of probability that B_T hits these compact sets on an exponential scale. Utilizing the property of reproducing kernel Hilbert space, we obtain the explicit expression of the large deviations rate function I of $\{B_T\}_{T>0}$.

The large deviations rate functions characterize the essential decay rate of the probability of rare events. It is important for a numerical discretization to preserve the rate function in certain sense. Thus, for a numerical discretization of (1.1), it is natural to ask:

- (P1) Does the discrete approximation of $\{B_T\}_{T>0}$, associated with the numerical discretization of (1.1), satisfy the LDP?
- (P2) If so, which kind of numerical discretizations can preserve the LDP of the original system, namely preserve the LDP rate function, exactly or asymptotically?

This paper aims to deal with the above problems. We are faced with two major difficulties in the numerical analysis. One is how to define the preservation for the LDP of an infinite dimensional stochastic differential equation by its numerical discretizations. Unlike the LDP of the original system in infinite dimensional spaces, the space concerning the LDP of a numerical discretization is finite dimensional. Therefore one needs a reasonable definition to link these two spaces. Another difficulty arises from the symplectic discretizations of the stochastic Schrödinger equation, including the general formulation in high dimensional case and the combination with the theory of large deviations.

Concerning these issues, we first apply the spectral Galerkin method to (1.1) and get the spatial semi-discretization (see (4.1))

$$du^{M}(t) = i\Delta_{M}u^{M}(t)dt + i\alpha P_{M}dW(t), \qquad t > 0,$$

$$u^{M}(0) = P_{M}u_{0} \in H_{M}.$$
(1.2)

Here $H_M = \text{span}\{e_1, e_2, \dots, e_M\}$, where $e_k, k = 1, 2, \dots$ are the eigenfunctions of Q and form an orthonormal basis of H^0 . In fact, (1.2) is a symplectic discretization and can be rewritten into a stochastic Hamiltonian system (see (5.1)):

$$dP^{M}(t) = \mathcal{M}Q^{M}(t)dt,$$

$$dQ^{M}(t) = -\mathcal{M}P^{M}(t)dt + \alpha \mathcal{Q}d\beta(t),$$
(1.3)

where $u^M = P^M + iQ^M$. We define by $B_T^M = \frac{u^M(T)}{T}$, T > 0 a discrete approximation of the observable B_T for (1.2). Following the arguments of dealing with the LDP for $\{B_T\}_{T>0}$, we prove that for each $M \in \mathbb{N}$, $\{B_T^M\}_{T>0}$ obeys an LDP on H_M with the good rate function \widetilde{I}^M . Note that \widetilde{I}^M and I have different domains, which brings the difficulty to define and study the preservation of the LDP for $\{B_T\}_{T>0}$ by $\{u^M\}_{M\in\mathbb{N}}$. A possibility is to transfer the LDP of $\{B_T^M\}_{T>0}$ on H_M to H^0 . This can be solved by means of Lemma 3.6 which reveals the relationship between LDPs of a stochastic process on some space and that on subspaces. This is to say, $\{B_T^M\}_{T>0}$ also satisfies the LDP on H^0 with a rate function I^M . However, we also note that the valid domain, on which I^M takes finite values, is a proper subset of the valid domain of I. Hence, we introduce the definition of weakly asymptotical preservation for LDP (see Definition 4.2) in the sense that I is well approximated by I^M for some sufficiently large M. Further, we prove that $\{u^M\}_{M\in\mathbb{N}}$ weakly asymptotically preserves the LDP of $\{B_T\}_{T>0}$ based on the strong continuity of $\{P_M\}_{M\in\mathbb{N}}$.

Next, we attempt to show that the full discretization based on a large class of temporal symplectic discretization can weakly asymptotically preserve the LDP of $\{B_T\}_{T>0}$. In order to give the general formula of symplectic discretizations for the high dimensional system (1.3), an argument of dimensionality reduction is applied. More precisely, we divide (1.3) into M subsystems (see (5.3)). Then we obtain a class of full discretizations $\{u_n^M\}_{M,n\in\mathbb{N}}$ based on the temporal symplectic discretizations of (1.3) by combining the symplectic discretizations in [3] for every 2-dimensional subsystem. For this full discretization, we define a discrete approximation $B_N^M = \frac{u_N^M}{N\tau}$ of B_T , with τ being the temporal stepsize, and give the LDP of $\{B_N^M\}_{N\in\mathbb{N}}$ based on the Gärtner–Ellis theorem and the contraction theorem. Further, we study whether $\{u_n^M\}_{M,n\in\mathbb{N}}$ can weakly asymptotically preserve the LDP (see Definition 5.5) of $\{B_T\}_{T>0}$, which depends on the asymptotical behavior of the modified rate function $I_{mod}^{M,\tau}$ of $\{B_N^M\}_{N\in\mathbb{N}}$. Notice that I^M is a good approximation of I, it suffices to prove that for each $M\in\mathbb{N}$, $\{u_n^M\}_{n\in\mathbb{N}}$ can asymptotically preserve the LDP of $\{B_T^M\}_{T>0}$, i.e., the modified rate function $I_{mod}^{M,\tau}$ converges to I^M pointwise as τ tends to zero. Similar to [3], under certain convergence condition of numerical approximations, we obtain $\lim_{\tau \to 0} I_{mod}^{M,\tau}(\cdot) = I^{M}(\cdot)$. Combining the asymptotical convergence of I^{M} to I, we deduce our main conclusion that the full discretization $\{u_n^M\}_{M,n\in\mathbb{N}}$, based on the spatial spectral Galerkin approximation and temporal symplectic discretizations, can weakly asymptotically preserve the LDP of $\{B_T\}_{T>0}$. That is to say, we obtain a good approximation of the LDP rate function of $\{B_T\}_{T>0}$ based on the symplectic discretizations. To the best of our knowledge, this is the first result of approximating the LDP rate function in infinite dimensional space based on the numerical discretizations. We partially answer the open problem proposed by [3].

The paper is organized as follows. In Section 2, some useful notations and preliminaries are introduced. In Section 3, we give an introduction on the LDP in general topological vector spaces, and prove that $\{B_T\}_{T>0}$ satisfies an LDP on H^0 . The weakly asymptotical preservations of LDP for $\{B_T\}_{T>0}$ by the spectral Galerkin approximation and the further full discretizations based on the temporal symplectic discretizations are given in Sections 4 and 5, respectively. Section 6 generalizes the LDP of $\{B_T\}_{T>0}$ to the case of complex-valued noises. Future work is discussed in Section 7.

2. Preliminaries

We begin with some notations. Throughout this paper, denote by $H^s = H^s(0,\pi)$ and $H^s(0,\pi;\mathbb{R})$, the classical Sobolev space of complex-valued functions and the classical Sobolev space of real-valued functions, respectively. In particular, denote $H^0 = L^2(0,\pi;\mathbb{C})$, $H^1_0(0,\pi) = \{f \in H^1(0,\pi) | f(0) = f(\pi) = 0\}$, $U^0 = L^2(0,\pi;\mathbb{R})$ and $U^1 = H^1(0,\pi;\mathbb{R})$. For a linear operator A from some Hilbert space onto itself, let $\lambda_k(A)$ be the kth eigenvalue of A. For a complex number z, let $\Re z$ and $\Im z$ be its real part

and imaginary part, respectively. And denote by i the imaginary unit. Let $(U, \|\cdot\|_U, \langle\cdot,\cdot\rangle_U)$ and $(H, \|\cdot\|_H, \langle\cdot,\cdot\rangle_H)$ be two separable Hilbert spaces. Then $\mathcal{L}_2(U, H)$ denotes the Banach spaces consisting of all the Hilbert–Schmidt operators from U to H, with the norm $\|A\|_{\mathcal{L}_2(U,H)} = \left(\sum_{k=1}^{\infty} \|Af_k\|_H^2\right)^{\frac{1}{2}}$, where $\{f_k\}_{k\in\mathbb{N}}$ is any orthonormal basis of U. Denote the real inner product by $\langle f,g\rangle_{\mathbb{R}} = \Re\int_0^\pi f(x)\bar{g}(x)dx$, and the complex inner product by $\langle f,g\rangle_{\mathbb{C}} = \int_0^\pi f(x)\bar{g}(x)dx$ for $f,g\in H^0$.

and the complex inner product by $\langle f,g\rangle_{\mathbb{C}}=\int_0^\pi f(x)\bar{g}(x)dx$ for $f,g\in H^0$. For a given $M\in\mathbb{N}$, \mathbb{C}^M denotes the space of M-dimensional complex-valued vectors. Define the inner product on \mathbb{C}^M by $\langle u,v\rangle_{\mathbb{R}}=\sum\limits_{k=1}^M\left(\Re u_k\Re v_k+\Im u_k\Im v_k\right)$, and the norm by $\|u\|=\sqrt{\langle u,u\rangle_{\mathbb{R}}}$ for any $u=(u_1,u_2,\ldots,u_M),\ v=(v_1,v_2,\ldots,v_M)\in\mathbb{C}^M$. $R=\mathcal{O}(h^p)$ stands for $|R|\leq Ch^p$, for all sufficiently small h>0. $f(h)\sim h^p$ means that f(h) and h^p are equivalent infinitesimal. For the random variables X,Y, $\mathbf{Var}(X)$ denotes the covariance operator of X and X.

In order to investigate the stochastic Schrödinger equation (1.1), we introduce the definition and properties of the noise. Let $e_k(x) = \sqrt{\frac{2}{\pi}}\sin(kx)$, then $\{e_k\}_{k\in\mathbb{N}}$ forms an orthonormal basis of both $(H^0, \langle \cdot, \cdot \rangle_{\mathbb{C}})$ and $(U^0, \langle \cdot, \cdot \rangle_{\mathbb{R}})$. Assume that Q is a nonnegative symmetric operator on U^0 with $Qe_k = \eta_k e_k$ for some non-increasing sequence $\{\eta_k\}_{k\in\mathbb{N}}$. Then W has the expansion $W(t) = \sum_{k\geq 1} \sqrt{\eta_k} \beta_k(t) e_k$. Q can be extended to H^0 by defining $Qf = Q(\Re f) + iQ(\Im f)$ for every $f \in H^0$ and the extended operator is still denoted by Q, if no confusion occurs. Noting that $\Delta e_k = -k^2 e_k$, $k = 1, 2 \dots$, we have that $\Delta Q = Q\Delta$.

Let $S(t) = e^{it\Delta}$ be the unitary C_0 -group generated by A. Throughout the paper, we assume that $Q^{\frac{1}{2}} \in \mathcal{L}_2(U^0, U^1)$ and $u_0 \in H_0^1(0, \pi)$, then (1.1) admits a unique mild solution in $H_0^1(0, \pi)$ (see e.g., [1]):

$$u(t) = S(t)u_0 + i\alpha \int_0^t S(t-s)dW(s).$$
(2.1)

Next, we give some results about the property of the distribution of exact solution (2.1). These results are based on the following proposition.

Proposition 2.1. [7, Proposition 4.28] Let W be a U-valued Q-Wiener process and $\mathcal{N}_W^2(0,T;L_0^2)$ denote the set

$$\left\{\Phi: [0,T] \times \Omega \to \mathcal{L}_2(Q^{\frac{1}{2}}(U),H) \middle| \Phi \text{ is predicable and } \mathbf{E} \int_0^T \left\|\Phi(s) \circ Q^{\frac{1}{2}}\right\|_{\mathcal{L}_2(U,H)}^2 ds < \infty\right\},$$

where H is a separable Hilbert space. Assume that $\Phi_1, \Phi_2 \in \mathcal{N}^2_W(0,T;L^2_0)$, then the correlation operators

$$V(t,s) = \mathbf{Cor}(\Phi_1 \cdot W(t), \Phi_2 \cdot W(s)), \quad t,s \in [0,T]$$

are given by the formula

$$V(t,s) = \mathbf{E} \int_0^{t \wedge s} \Phi_2(r) Q(\Phi_1(r))^* dr.$$

Here, the operator V(t,s) is defined by

$$\langle V(t,s)a,b\rangle_H = \mathbf{E} \langle \Phi_1 \cdot W(t),a\rangle_H \langle \Phi_2 \cdot W(s),b\rangle_H, \qquad a,b \in H.$$

It follows from (2.1) that

$$u(t) = S(t)u_0 + i\alpha \int_0^t (\cos((t-s)\Delta)) + i\sin((t-s)\Delta)) dW(s)$$

$$= S(t)u_0 - \alpha \int_0^t \sin((t-s)\Delta)dW(s) + i\alpha \int_0^t \cos((t-s)\Delta)dW(s)$$

=: $S(t)u_0 - \alpha W_{\sin}(t) + i\alpha W_{\cos}(t)$.

Noting that $\langle f, g \rangle_{\mathbb{R}} = \langle \Re f, \Re g \rangle_{\mathbb{R}} + \langle \Im f, \Im g \rangle_{\mathbb{R}}$, we have that for each $h = \Re h + i \Im h \in H^0$,

$$\langle u(t), h \rangle_{\mathbb{R}} = \langle S(t)u_0, h \rangle_{\mathbb{R}} - \alpha \langle W_{\sin}(t), \Re h \rangle_{\mathbb{R}} + \alpha \langle W_{\cos}(t), \Im h \rangle_{\mathbb{R}}.$$
 (2.2)

Hence,

$$\mathbf{E} \langle u(t), h \rangle_{\mathbb{R}} = \langle S(t)u_0, h \rangle_{\mathbb{R}}. \tag{2.3}$$

It follows from Proposition 2.1 that

$$W_{\sin}(t) \sim \mathcal{N}\left(0, \int_0^t \sin^2((t-s)\Delta)Qds\right), \qquad W_{\cos}(t) \sim \mathcal{N}\left(0, \int_0^t \cos^2((t-s)\Delta)Qds\right), \qquad (2.4)$$

$$\mathbf{Cor}\left(W_{\sin}(t), W_{\cos}(t)\right) = \int_0^t \sin((t-s)\Delta)\cos((t-s)\Delta)Qds.$$

Using the above formulas and $\Delta Q = Q\Delta$, one has

$$\mathbf{Var} \langle u(t), h \rangle_{\mathbb{R}} = \alpha^{2} \left\langle \int_{0}^{t} \sin^{2}((t-s)\Delta)Qds\Re h, \Re h \right\rangle_{\mathbb{R}} + \alpha^{2} \left\langle \int_{0}^{t} \cos^{2}((t-s)\Delta)Qds\Im h, \Im h \right\rangle_{\mathbb{R}} - 2\alpha^{2} \left\langle \int_{0}^{t} \sin((t-s)\Delta)\cos((t-s)\Delta)Qds\Re h, \Im h \right\rangle_{\mathbb{R}}.$$
(2.5)

Since Δ is invertible, we have

$$\int_0^t \sin^2((t-s)\Delta)ds = \frac{1}{2} \int_0^t (I - \cos(2(t-s)\Delta)) ds = \frac{tI}{2} - \frac{\Delta^{-1}}{4} \sin(2t\Delta), \tag{2.6}$$

$$\int_0^t \cos^2((t-s)\Delta)ds = \frac{1}{2} \int_0^t (I + \cos(2(t-s)\Delta)) ds = \frac{tI}{2} + \frac{\Delta^{-1}}{4} \sin(2t\Delta), \tag{2.7}$$

$$\int_0^t \sin(2(t-s)\Delta)ds = \frac{\Delta^{-1}}{2} \left[I - \cos(2t\Delta) \right]. \tag{2.8}$$

Combining (2.5), (2.6), (2.7) and (2.8) leads to

$$\mathbf{Var} \langle u(t), h \rangle_{\mathbb{R}} = \frac{t\alpha^{2}}{2} \left(\langle Q\Re h, \Re h \rangle_{\mathbb{R}} + \langle Q\Im h, \Im h \rangle_{\mathbb{R}} \right) - \frac{\alpha^{2}}{4} \left[\left\langle \Delta^{-1} \sin(2t\Delta) Q\Re h, \Re h \right\rangle_{\mathbb{R}} - \left\langle \Delta^{-1} \sin(2t\Delta) Q\Im h, \Im h \right\rangle_{\mathbb{R}} \right] - \frac{\alpha^{2}}{2} \left\langle \Delta^{-1} \left(I - \cos(2t\Delta) \right) Q\Re h, \Im h \right\rangle_{\mathbb{R}}.$$
 (2.9)

3. LDP for B_T of stochastic linear Schrödinger equation

In this section, we study the LDP for $\{B_T\}_{T>0}$ by means of the abstract Gärtner–Ellis theorem. As a corollary, we give the exponential tail estimate of the mass of (1.1). Throughout this section, let \mathcal{X} be a locally convex Hausdorff topological vector space and \mathcal{X}^* be its dual space.

3.1. **Introduction on LDP.** In this part, we recall some concepts upon LDP and useful theorems and lemmas in studying the LDP of a family of probability measures. First we introduce the definitions of rate function and LDP (see e.g., [3]).

Definition 3.1. A real-valued function $I: \mathcal{X} \to [0, \infty]$ is called a rate function, if it is lower semicontinuous, i.e., for each $a \in [0, \infty)$, the level set $I^{-1}([-\infty, a])$ is a closed subset of \mathcal{X} . If all level sets $I^{-1}([-\infty, a])$, $a \in [0, \infty)$, are compact, then I is called a good rate function.

Definition 3.2. Let I be a rate function and $\{\mu_{\epsilon}\}_{{\epsilon}>0}$ be a family of probability measures on \mathcal{X} . We say that $\{\mu_{\epsilon}\}_{{\epsilon}>0}$ satisfies an LDP on \mathcal{X} with the rate function I if

(LDP1)
$$\liminf_{\epsilon \to 0} \epsilon \ln(\mu_{\epsilon}(U)) \ge -\inf I(U) \quad \text{for every open } U \subset \mathcal{X},$$

(LDP2)
$$\limsup_{\epsilon \to 0} \epsilon \ln(\mu_{\epsilon}(C)) \le -\inf I(C) \quad \text{for every closed } C \subset \mathcal{X}.$$

Analogously, we say that a family of random variables $\{Z_{\epsilon}\}_{\epsilon>0}$ valued on \mathcal{X} satisfies an LDP with the rate function I if its distribution satisfies the lower bound LDP (LDP1) and upper bound LDP (LDP2) in Definition 3.2 for the rate function I.

Generally speaking, we need to investigate the logarithmic moment generating function and the exponential tightness of $\{\mu_{\epsilon}\}_{\epsilon>0}$, when we derive the LDP of $\{\mu_{\epsilon}\}_{\epsilon>0}$. Especially, if the state space \mathcal{X} is finite dimensional, the existence of logarithmic moment generating function implies the exponential tightness. However, when \mathcal{X} is infinite dimensional, the exponential tightness of $\{\mu_{\epsilon}\}_{\epsilon>0}$ can not be ignored.

Definition 3.3. [8, Page 8] A family of probability measures $\{\mu_{\epsilon}\}$ on \mathcal{X} is exponentially tight if for every $\alpha < \infty$, there exists a compact set $K_{\alpha} \subset \mathcal{X}$ such that

$$\limsup_{\epsilon \to 0} \epsilon \ln \mu_{\epsilon}(K_{\alpha}^{c}) < -\alpha. \tag{3.1}$$

Theorem 3.4. [8, Corollary 4.6.14] Let $\{\mu_{\epsilon}\}_{\epsilon>0}$ be an exponentially tight family of Borel probability measures on \mathcal{X} . Suppose the logarithmic moment generating function $\Lambda(\cdot) = \lim_{\epsilon \to 0} \epsilon \Lambda_{\mu_{\epsilon}}(\cdot/\epsilon)$ is finite valued and Gateaux differentiable, where $\Lambda_{\mu_{\epsilon}}(\lambda) := \ln \int_{\mathcal{X}} e^{\lambda(x)} \mu_{\epsilon}(dx)$, $\lambda \in \mathcal{X}^*$. Then $\{\mu_{\epsilon}\}_{\epsilon>0}$ satisfies the LDP in \mathcal{X} with the convex, good rate function $\Lambda^*(x) = \sup_{\lambda \in \mathcal{X}^*} \{\lambda(x) - \Lambda(\lambda)\}$.

Theorem 3.4 can be viewed as the abstract Gärtner–Ellis theorem. The following two lemmas are useful to derive new LDPs based on a given LDP. The first lemma is also called the contraction principle, which produces a new LDP on another space based on the known LDP via a continuous mapping. The second one gives the relationship between the LDP of $\{\mu_{\epsilon}\}_{\epsilon>0}$ on \mathcal{X} and that on the subspaces of \mathcal{X} .

Lemma 3.5. [8, Theorem 4.2.1] Let \mathcal{Y} be another Hausdorff topological space, $f: \mathcal{X} \to \mathcal{Y}$ be a continuous function, and $I: \mathcal{X} \to [0, \infty]$ be a good rate function.

(a) For each $y \in \mathcal{Y}$, define

$$\tilde{I}(y) \triangleq \inf \{ I(x) : x \in \mathcal{X}, y = f(x) \}.$$

Then $\tilde{I}(y)$ is a good rate function on \mathcal{Y} , where as usual the infimum over the empty set is taken as ∞ .

(b) If I controls the LDP associated with a family of probability measures $\{\mu_{\epsilon}\}$ on \mathcal{X} , then $\tilde{I}(y)$ controls the LDP associated with the family of probability measures $\{\mu_{\epsilon} \circ f^{-1}\}$ on \mathcal{Y} .

Lemma 3.6. [8, Lemma 4.1.5] Let E be a measurable subset of \mathcal{X} such that $\mu_{\epsilon}(E) = 1$ for all $\epsilon > 0$. Suppose that E is equipped with the topology induced by \mathcal{X} . If E is a closed subset of \mathcal{X} and $\{\mu_{\epsilon}\}_{\epsilon>0}$ satisfies the LDP on E with the rate function I, then $\{\mu_{\epsilon}\}_{\epsilon>0}$ satisfies the LDP on \mathcal{X} with the rate function $\tilde{I}(y)$ such that $\tilde{I}(y) = I$ on E and $\tilde{I}(y) = \infty$ on E^c .

Proposition 3.7. [8, Lemma 1.2.15] Let N be a fixed integer. Then, for every $a_{\epsilon}^{i} \geq 0$,

$$\limsup_{\epsilon \to 0} \epsilon \ln \left(\sum_{i=1}^{N} a_{\epsilon}^{i} \right) = \max_{i=1,\dots,N} \limsup_{\epsilon \to 0} \epsilon \ln a_{\epsilon}^{i}.$$

Proposition 3.7 is an important tool in deriving (LDP1) and (LDP2). Furthermore, we need to make use of the following proposition in stochastic calculus.

Proposition 3.8. [6, Proposition 1.13] Assume that \widetilde{Q} is a nonnegative symmetric operator on a real separable Hilbert space H with finite trace. Let $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n \geq \cdots$ be the eigenvalues of \widetilde{Q} . Define the determinant of $(I - 2\epsilon \widetilde{Q})$ by setting $\det(I - 2\epsilon \widetilde{Q}) := \lim_{n \to \infty} \prod_{k=1}^{n} (1 - 2\epsilon \lambda_k) := \prod_{k=1}^{\infty} (1 - 2\epsilon \lambda_k)$. Let $\mu = \mathcal{N}(0, \widetilde{Q})$ be the symmetric Gaussian measure on H. Then for every $\epsilon \in \mathbb{R}$,

$$\int_{H} e^{\epsilon \|x\|_{H}^{2}} \mu(dx) = \begin{cases} \left[\det(I - 2\epsilon \widetilde{Q}) \right]^{-1/2}, & if \ \epsilon < \frac{1}{2\lambda_{1}}, \\ +\infty, & otherwise. \end{cases}$$
(3.2)

3.2. **LDP** for $\{B_T\}_{T>0}$. In this subsection, we show the LDP for $\{B_T\}_{T>0}$ of (1.1) by using Theorem 3.4, where $B_T := \frac{u(T)}{T}$ with u(T) being the solution of (1.1) at time T. The regime of Gärtner–Ellis theorem is applicable to the real Banach space. Given that the exact solution $\{u(t)\}_{t\geq 0}$ takes values in H^0 , the space of complex-valued functions, we use the real inner product to establish the LDP of $\{B_T\}_{T>0}$ on H^0 .

Theorem 3.9. $\{B_T\}_{T>0}$ satisfies an LDP on H^0 with the good rate function

$$I(x) = \begin{cases} \frac{1}{\alpha^2} \left\| Q^{-\frac{1}{2}} x \right\|_{H^0}^2, & if \ x \in Q^{\frac{1}{2}}(H^0), \\ +\infty, & otherwise, \end{cases}$$

where $Q^{-\frac{1}{2}}$ is the pseudo inverse of $Q^{\frac{1}{2}}$.

Proof. We divide the proof into three steps.

Step 1: The logarithmic moment generating function of $\{B_T\}_{T>0}$

For each $\lambda \in H^0$, define the mapping $\lambda' : H^0 \to \mathbb{R}$ by $\lambda'(x) = \langle x, \lambda \rangle_{\mathbb{R}}$, $x \in H^0$. Then by Riesz representation theorem, $\{\lambda'\}_{\lambda \in H^0}$ forms the set of all real bounded linear functionals of H^0 , i.e., $\{\lambda'\}_{\lambda \in H^0} = (H^0)^*$. Since $\langle u(t), \lambda \rangle_{\mathbb{R}}$ is Gaussian, it follows from (2.3) and (2.9) that

$$\Lambda(\lambda') = \lim_{T \to \infty} \frac{1}{T} \ln \mathbf{E} e^{T\langle B_T, \lambda \rangle_{\mathbb{R}}} = \lim_{T \to \infty} \frac{1}{T} \ln \mathbf{E} e^{\langle u(T), \lambda \rangle_{\mathbb{R}}}
= \lim_{T \to \infty} \frac{1}{T} \left[\mathbf{E} \langle u(T), \lambda \rangle_{\mathbb{R}} + \frac{1}{2} \mathbf{Var} \langle u(T), \lambda \rangle_{\mathbb{R}} \right]
= \frac{\alpha^2}{4} \left(\langle Q \Re \lambda, \Re \lambda \rangle_{\mathbb{R}} + \langle Q \Im \lambda, \Im \lambda \rangle_{\mathbb{R}} \right)
= \frac{\alpha^2}{4} \left\| Q^{\frac{1}{2}} \lambda \right\|_{H^0}^2,$$
(3.3)

where we use the facts $\|\sin(t\Delta)\|_{\mathcal{L}(H^0)} \le 1$, $\|\cos(t\Delta)\|_{\mathcal{L}(H^0)} \le 1$ and $\|\Delta^{-1}\|_{\mathcal{L}(H^0)} = 1$.

Step 2: Exponential tightness of $\{B_T\}_{T>0}$

In order to obtain the exponential tightness of $\{B_T\}_{T>0}$ (see Definition 3.3), it suffices to show that there exists a family of compact sets $\{K_L\}_{L>0}$ such that

$$\lim_{L \to \infty} \limsup_{T \to \infty} \frac{1}{T} \ln \mathbf{P} \left(B_T \in K_L^c \right) = -\infty. \tag{3.4}$$

Define $K_L = \{f \in H^1 \mid ||f||_{H^1} \leq L\}$. Then K_L is the compact set of H^0 . Recall that $u(T) = S(T)u_0 - \alpha W_{\sin}(T) + i\alpha W_{\cos}(T)$. Thus

$$\mathbf{P}\left(B_T \in K_L^c\right) = \mathbf{P}\left(\|u(T)\|_{H^1} > LT\right)$$

$$\leq \mathbf{P}\left(\|S(T)u_0\|_{H^1} > \frac{TL}{3}\right) + \mathbf{P}\left(\alpha\|W_{\sin}(T)\|_{U^1} > \frac{TL}{3}\right) + \mathbf{P}\left(\alpha\|W_{\cos}(T)\|_{U^1} > \frac{TL}{3}\right). \tag{3.5}$$

Since the first term in (3.5) is 0 for sufficiently large T, we only need to estimate the second and third terms in (3.5).

Since $Q^{\frac{1}{2}} \in \mathcal{L}_2(U^0, U^1)$, Q is also the finite trace operator on U^1 . Then we obtain from (2.6) that

$$W_{\sin}(T) \sim \mathcal{N}\left(0, \int_0^t \sin^2((t-s)\Delta)Qds\right) = \mathcal{N}\left(0, \left(\frac{TI}{2} - \frac{\Delta^{-1}\sin(2T\Delta)}{4}\right)Q\right) \text{ on } U^1.$$

Further, it holds that

$$\frac{W_{\sin(T)}}{\sqrt{T}} \sim \mathcal{N}\left(0, \left(\frac{I}{2} - \frac{\Delta^{-1}\sin(2T\Delta)}{4T}\right)Q\right) \text{ on } U^{1}.$$
(3.6)

By Markov's inequality, for each $\varepsilon > 0$,

$$\mathbf{P}\left(\alpha \|W_{\sin}(T)\|_{U^{1}} > \frac{TL}{3}\right) = \mathbf{P}\left(\left\|\frac{W_{\sin}(T)}{\sqrt{T}}\right\|_{U^{1}} > \frac{\sqrt{T}L}{3\alpha}\right)$$

$$= \mathbf{P}\left(\exp\left\{\varepsilon \left\|\frac{W_{\sin}(T)}{\sqrt{T}}\right\|_{U^{1}}^{2}\right\} > \exp\left\{\frac{\varepsilon TL^{2}}{9\alpha^{2}}\right\}\right)$$

$$\leq e^{-\frac{\varepsilon TL^{2}}{9\alpha^{2}}}\mathbf{E}\exp\left\{\varepsilon \left\|\frac{W_{\sin}(T)}{\sqrt{T}}\right\|_{U^{1}}^{2}\right\}.$$
(3.7)

Notice that $\lambda_k \left(\left(\frac{I}{2} - \frac{\Delta^{-1} \sin(2T\Delta)}{4T} \right) Q \right) = \left(\frac{1}{2} - \frac{\sin(2Tk^2)}{4Tk^2} \right) \eta_k = \frac{1}{2} \left(1 - \frac{\sin(2Tk^2)}{2Tk^2} \right) \eta_k < \eta_k \le \eta_1$. It follows from Proposition 3.8 that for each $0 < \varepsilon < \frac{1}{2\eta_1}$,

$$\mathbf{E} \exp \left\{ \varepsilon \left\| \frac{W_{\sin}(T)}{\sqrt{T}} \right\|_{U^{1}}^{2} \right\} = \left[\det \left(I - 2\varepsilon \left(\frac{I}{2} - \frac{\Delta^{-1} \sin(2T\Delta)}{4T} \right) Q \right) \right]^{-\frac{1}{2}}$$

$$< \left[\det (I - 2\varepsilon Q) \right]^{-\frac{1}{2}} = C(\varepsilon, Q), \tag{3.8}$$

where we have used the fact that $[\det(I-2\varepsilon Q)]^{-\frac{1}{2}}=(\prod_{k=1}^{\infty}(1-2\varepsilon\eta_k))^{-\frac{1}{2}}$ is monotonically increasing with respect to η_k for every $k=1,2,\ldots$ Combining (3.8) with (3.7) yields

$$\limsup_{T \to \infty} \frac{1}{T} \ln \mathbf{P} \left(\alpha \| W_{\sin}(T) \|_{U^1} > \frac{TL}{3} \right) \le \limsup_{T \to \infty} \frac{1}{T} \ln \left(e^{-\frac{\varepsilon T L^2}{9\alpha^2}} C(\varepsilon, Q) \right) = -\frac{\varepsilon L^2}{9\alpha^2}. \tag{3.9}$$

In addition, it holds that

$$\frac{W_{\cos(T)}}{\sqrt{T}} \sim \mathcal{N}\left(0, \left(\frac{I}{2} + \frac{\Delta^{-1}\sin(2T\Delta)}{4T}\right)Q\right) \text{ on } U^1.$$

Then $\lambda_k \left(\left(\frac{I}{2} + \frac{\Delta^{-1} \sin(2T\Delta)}{4T} \right) Q \right) = \frac{1}{2} \left(1 + \frac{\sin(2Tk^2)}{2Tk^2} \right) \eta_k < \eta_k \le \eta_1$. Analogous to the proof of (3.9), one has that for $0 < \varepsilon < \frac{1}{2\eta_1}$,

$$\limsup_{T \to \infty} \frac{1}{T} \ln \mathbf{P} \left(\alpha \| W_{\cos}(T) \|_{U^1} > \frac{TL}{3} \right) \le -\frac{\varepsilon L^2}{9\alpha^2}. \tag{3.10}$$

Combining (3.9), (3.10), (3.5) and Proposition 3.7, we obtain

$$\limsup_{T \to \infty} \frac{1}{T} \ln \mathbf{P} \left(B_T \in K_L^c \right) \le \max \left\{ -\frac{\varepsilon L^2}{9\alpha^2}, -\frac{\varepsilon L^2}{9\alpha^2} \right\} = -\frac{\varepsilon L^2}{9\alpha^2}.$$

Accordingly, we have

$$\lim_{L \to \infty} \limsup_{T \to \infty} \frac{1}{T} \ln \mathbf{P} \left(B_T \in K_L^c \right) = -\infty, \tag{3.11}$$

which proves the exponential tightness of $\{B_T\}_{T>0}$.

It is verified that $\Lambda(\lambda')$ is finite valued and Gateaux differentiable. In fact, $\Lambda(\lambda')$ is Fréchet differentiable, and its Fréchet derivative is $\mathcal{D}\Lambda(\lambda')(\cdot) = \frac{\alpha^2}{2}\langle Q\lambda, \cdot \rangle_{\mathbb{R}}$. Due to Theorem 3.4, $\{B_T\}_{T>0}$ satisfies an LDP on H^0 with the good rate function Λ^* . It remains to give the explicit expression of the Fenchel–Legendre transform Λ^* of Λ .

Step 3: The explicit expression of Λ^*

Before giving the expression of Λ^* , we recall the concept of reproducing kernel Hilbert space (RKHS). Let μ be a centered Gaussian measure on a separable Banach space E. An arbitrary $\varphi \in E^*$ can be identified with an element of the Hilbert space $L^2(\mu) := L^2(E, \mathcal{B}(E), \mu; \mathbb{R})$. Denote by $\overline{E^*} = \overline{E^*}^{L^2(\mu)}$ the closure of E^* in $L^2(\mu)$. Define a mapping $J : \overline{E^*} \to E$ by setting

$$J(\varphi) = \int_E x \varphi(x) \mu(dx), \quad \forall \quad \varphi \in \overline{E^*}.$$

Then the image \mathcal{H}_{μ} of J in E, $\mathcal{H}_{\mu} = J(\overline{E^*})$ is the RKHS of μ with the scalar product

$$\langle J(\varphi), J(\psi) \rangle_{\mathcal{H}_{\mu}} = \int_{E} \varphi(x) \psi(x) \mu(dx).$$

Further, if $\mu = \mathcal{N}(0, \widetilde{Q})$ is a Gaussian measure on some Hilbert space H with \widetilde{Q} being a nonnegative symmetric operator with finite trace, then the RKHS \mathcal{H}_{μ} of μ is $\mathcal{H}_{\mu} = \widetilde{Q}^{\frac{1}{2}}(H)$ with the norm $\|x\|_{\mathcal{H}_{\mu}} = \|\widetilde{Q}^{-\frac{1}{2}}x\|_{H}$. We refer to [7, Section 2.2.2] for more details of the RKHS.

In our case, $\mu = \mathcal{N}(0, Q)$. The mapping $J : \overline{(H^0)^*}^{L^2(\mu)} \to H^0$ is

$$J(h) = \int_{H^0} zh(z)\mu(dz).$$

Then $\mathcal{H}_{\mu} = J\left(\overline{(H^0)^*}^{L^2(\mu)}\right) = Q^{\frac{1}{2}}(H^0)$. It follows from the properties of Gaussian measure that

$$\int_{H^0} \langle \lambda, x \rangle_{\mathbb{R}}^2 \, \mu(dx) = \langle Q\lambda, \lambda \rangle_{\mathbb{R}} = \left\| Q^{\frac{1}{2}} \lambda \right\|_{H^0}^2.$$

Thus, $\Lambda(\lambda') = \frac{\alpha^2}{4} \left\| Q^{\frac{1}{2}} \lambda \right\|_{H^0}^2 = \frac{\alpha^2}{4} \left\| \lambda' \right\|_{L^2(\mu)}^2$. Recall that

$$\Lambda^*(x) = \sup_{\lambda' \in (H^0)^*} \left\{ \lambda'(x) - \Lambda(\lambda') \right\}.$$

For a given $x \in H^0$, if $\Lambda^*(x) < +\infty$, then there exists a constant $C(x) < +\infty$ such that $\lambda'(x) \leq \frac{\alpha^2}{4} \|\lambda'\|_{L^2(\mu)}^2 + C(x)$. Define the linear functional x^{**} on $\left((H^0)^*, \|\cdot\|_{L^2(\mu)}\right) \subseteq \overline{(H^0)^*}^{L^2(\mu)}$ by $x^{**}(\lambda') = \lambda'(x)$, for every $\lambda' \in (H^0)^*$. Then we have $\sup_{\lambda' \in (H^0)^*, \|\lambda'\|_{L^2(\mu)} \leq 1} x^{**}(\lambda') \leq \frac{\alpha^2}{4} + C(x)$. It

means that x^{**} is a bounded linear functional on $(H^0)^*$, $\|\cdot\|_{L^2(\mu)}$. By Hahn–Banach theorem and the fact that $(H^0)^*$, $\|\cdot\|_{L^2(\mu)}$ is dense in $\overline{(H^0)^*}^{L^2(\mu)}$, x^{**} can be uniquely extended to $\overline{(H^0)^*}^{L^2(\mu)}$. (In fact, for each $\lambda' \in \overline{(H^0)^*}^{L^2(\mu)}$, take $\lambda'_n \in (H^0)^*$ such that $\lambda'_n \to \lambda'$ in the norm $\|\cdot\|_{L^2(\mu)}$. Then the extended functional is $x^{**}(\lambda') = \lim_{n \to \infty} x^{**}(\lambda'_n)$.) The extended functional is still denoted by

 x^{**} . In this way, for every $x \in H^0$ satisfying $\Lambda^*(x) < +\infty$, we obtain a bounded linear functional on $\overline{(H^0)^*}^{L^2(\mu)}$ such that $x^{**}(\lambda') = \lambda'(x)$ for each $\lambda' \in (H^0)^*$. By Riesz representation theorem, there exists some $h \in \overline{(H^0)^*}^{L^2(\mu)}$ such that $x^{**}(\lambda') = \langle \lambda', h \rangle_{L^2(\mu)}$ for each $\lambda' \in \overline{(H^0)^*}^{L^2(\mu)}$. Hence, $\lambda'(x) = \langle \lambda', h \rangle_{L^2(\mu)}$ for each $\lambda' \in (H^0)^*$. Further, we have that

$$\lambda'(x) = \int_{H^0} h(z)\lambda'(z)\mu(dz) = \lambda'\left(\int_{H^0} zh(z)\mu(dz)\right) = \lambda'(J(h)), \qquad \forall \quad \lambda' \in (H^0)^*.$$

By the arbitrariness of λ' , x = J(h). Hence, $\Lambda^*(x) < +\infty$ implies that $x \in \mathcal{H}_{\mu} = J\left(\overline{(H^0)^*}^{L^2(\mu)}\right) = Im(Q^{\frac{1}{2}})$, where $Im(Q^{\frac{1}{2}})$ is the image of $Q^{\frac{1}{2}}$.

On the other hand, if $H^0 \ni x = J(h)$ for some $h \in \overline{(H^0)^*}^{L^2(\mu)}$, then

$$\Lambda^*(x) = \Lambda^*(J(h)) = \sup_{\lambda' \in (H^0)^*} \left\{ \lambda'(J(h)) - \frac{\alpha^2}{4} \|\lambda'\|_{L^2(\mu)}^2 \right\}$$
$$= \sup_{\lambda' \in (H^0)^*} \left\{ \langle \lambda', h \rangle_{L^2(\mu)} - \frac{\alpha^2}{4} \|\lambda'\|_{L^2(\mu)}^2 \right\}.$$

Noting the continuity of $\langle \lambda', h \rangle_{L^2(\mu)} - \frac{\alpha^2}{4} \|\lambda'\|_{L^2(\mu)}^2$ with respect to λ' in the norm $\|\cdot\|_{L^2(\mu)}$, and that $\left((H^0)^*, \|\cdot\|_{L^2(\mu)}\right)$ is dense in $\overline{(H^0)^*}^{L^2(\mu)}$, we have

$$\Lambda^*(x) = \sup_{g \in \overline{(H^0)^*}^{L^2(\mu)}} \left\{ \langle g, h \rangle_{L^2(\mu)} - \frac{\alpha^2}{4} \|g\|_{L^2(\mu)}^2 \right\}
\leq \sup_{g \in \overline{(H^0)^*}^{L^2(\mu)}} \left\{ \frac{1}{2} \left[\frac{\alpha^2}{2} \|g\|_{L^2(\mu)}^2 + \frac{2}{\alpha^2} \|h\|_{L^2(\mu)}^2 \right] - \frac{\alpha^2}{4} \|g\|_{L^2(\mu)}^2 \right\}
= \frac{1}{\alpha^2} \|h\|_{L^2(\mu)}^2.$$

Taking $g = \frac{2}{\alpha^2}h$ leads to $\Lambda^*(x) \geq \frac{1}{\alpha^2} \|h\|_{L^2(\mu)}^2$. Thus, we obtain

$$\Lambda^*(x) = \frac{1}{\alpha^2} \|h\|_{L^2(\mu)}^2 = \frac{1}{\alpha^2} \|x\|_{\mathcal{H}_{\mu}}^2 = \frac{1}{\alpha^2} \|Q^{-\frac{1}{2}}x\|_{H^0}^2.$$
 (3.12)

Finally we have

$$\Lambda^*(x) = \begin{cases} \frac{1}{\alpha^2} \left\| Q^{-\frac{1}{2}} x \right\|_{H^0}^2, & \text{if } x \in Q^{\frac{1}{2}}(H^0), \\ +\infty, & \text{otherwise,} \end{cases}$$

which completes this proof.

Similar to the proof of [1, Proposition 3.1], we obtain $\mathbf{E} \|u(T)\|_{H^0}^2 = \mathbf{E} \|u_0\|_{H^0}^2 + \alpha^2 T \operatorname{tr}(Q)$, where $\operatorname{tr}(Q) = \sum_{k=1}^{\infty} \eta_k$. Then, by Markov's inequality, one has that for each R > 0 and sufficiently large T

$$\mathbf{P}\left(\|u(T)\|_{H^0}^2 \ge T^2 R^2\right) \le \frac{\mathbf{E}\|u(T)\|_{H^0}^2}{T^2 R^2} \le \frac{C}{T},\tag{3.13}$$

for some constant C independent of T. In what follows, we show that the probability of the tail event of the mass $||u(T)||_{H^0}^2$ in (3.13) can be exponentially small. More precisely, by Lemma 3.5 and Theorem 3.9, we immediately obtain the LDP of $\{||B_T||_{H^0}\}_{T>0}$, which yields the following corollary.

Corollary 3.10. If Q is an injection, then it holds that

(1) $\{\|B_T\|_{H^0}\}_{T>0}$ satisfies an LDP on $\mathbb{R}^+ := [0, +\infty)$ with the good rate function

$$J(y) = \frac{1}{\alpha^2} \inf_{z \in H^0, \|Q^{\frac{1}{2}}z\|_{H^0} = y} \|z\|_{H^0}^2, \qquad y \ge 0.$$

(2) For every R > 0 and $\varepsilon > 0$, there is some T_0 such that

$$\mathbf{P}\left(\left\|u(T)\right\|_{H^{0}}^{2} \ge T^{2}R^{2}\right) \le \exp\left\{-T\left(\inf_{y \ge R}J(y) - \varepsilon\right)\right\}, \qquad \forall \quad T \ge T_{0},$$
and $\inf_{y > R}J(y) \in (0, +\infty).$

Proof. (1) Since the mapping $\|\cdot\|_{H^0}: H^0 \to \mathbb{R}^+$ is continuous, it follows from Lemma 3.5 and Theorem 3.9 that $\{\|B_T\|_{H^0}\}_{T>0}$ satisfies an LDP on \mathbb{R}^+ with the good rate function

$$\begin{split} J(y) &= \inf_{x \in H^0, \ \|x\|_{H^0} = y} I(x) = \inf_{x \in Q^{\frac{1}{2}}(H^0), \ \|x\|_{H^0} = y} I(x) \\ &= \frac{1}{\alpha^2} \inf_{x \in Q^{\frac{1}{2}}(H^0), \ \|x\|_{H^0} = y} \left\| Q^{-\frac{1}{2}} x \right\|_{H^0}^2 \\ &= \frac{1}{\alpha^2} \inf_{z \in H^0, \ \left\| Q^{\frac{1}{2}} z \right\|_{H^0} = y} \left\| z \right\|_{H^0}^2, \end{split}$$

where we have used the assumption that Q is an injection. This proves the first conclusion.

(2) Clearly, the set $\left\{z \in H^0, \ \left\|Q^{\frac{1}{2}}z\right\|_{H^0} = y\right\}$ is nonempty for every $y \geq 0$. Hence, $J(y) < +\infty$ for every $y \geq 0$. Accordingly, $\inf_{y \geq R} J(y) < +\infty$ for each R > 0. In addition, we claim J(y) > 0 for each y > 0. In fact, if for some $y_0 > 0$, $J(y_0) = 0$, then there is a sequence $\{z_n\}_{n \in \mathbb{N}} \subseteq H^0$ such that $\|Q^{\frac{1}{2}}z_n\|_{H^0} = y_0$ and $\lim_{n\to\infty} \|z_n\|_{H^0} = 0$. Noting that $Q^{\frac{1}{2}}$ is a continuous operator, then we have $y_0 = \lim_{n \to \infty} \left\| Q^{\frac{1}{2}} z_n \right\|_{H^0} = 0$, which yields a contradiction. Hence, we prove the claim. Using the fact that a good rate function can achieve its infimum on every nonempty closed set (see e.g., [8, Page 4]), we have that for each R > 0, there is some $y_R \ge R$ such that $\inf_{y \ge R} J(y) = J(y_R) > 0$. It remains to prove (3.14). Since $\{\|B_T\|_{H^0}\}_{T>0}$ satisfies the LDP with the rate function J, we obtain that for each fixed R > 0,

$$\limsup_{T \to \infty} \frac{1}{T} \ln \mathbf{P} \left(\frac{\|u(T)\|_{H^0}}{T} \ge R \right) \le -\inf_{y \ge R} J(y).$$

The above formula implies that for every $\varepsilon > 0$, there is a $T_0 > 0$ such that

$$\frac{1}{T}\ln\mathbf{P}\left(\frac{\|u(T)\|_{H^0}}{T}\geq R\right)\leq -\inf_{y\geq R}J(y)+\varepsilon, \quad \forall \quad T\geq T_0.$$

Hence we have that

$$\mathbf{P}\left(\|u(T)\|_{H^0}^2 \ge T^2 R^2\right) = \mathbf{P}\left(\frac{\|u(T)\|_{H^0}}{T} \ge R\right) \le \exp\left\{-T\left(\inf_{y \ge R} J(y) - \varepsilon\right)\right\}, \quad \forall \quad T \ge T_0.$$

This completes the proof.

Remark 3.11. For sufficiently large L > 0, one can always find R and T such that $T^2R^2 \le L$. Then by (3.11) one has that $\mathbf{P}\left(\|u(T)\|_{H^0}^2 \ge L\right) \le \mathbf{P}\left(\|u(T)\|_{H^0}^2 \ge T^2R^2\right) \le \exp\left\{-T\left(\inf_{y\ge R}J(y)-\varepsilon\right)\right\}$. This indicates that the probability of the tail event of the mass of (1.1) is exponentially small on a sufficiently large time.

4. LDP FOR THE SPATIAL SPECTRAL GALERKIN APPROXIMATION

In the previous section, we derive the LDP of $\{B_T\}_{T>0}$ for the continuous system (1.1). In order to obtain a valid approximation for the rate function I of $\{B_T\}_{T>0}$, we apply the spatial spectral Galerkin method to (1.1), and study the LDP of $\{B_T^M\}_{T>0}$ of spectral Galerkin approximation. Here, B_T^M is a discrete approximation of B_T , which will be specified later.

For $M \in \mathbb{N}$, we define the finite dimensional subspace $H_M := \operatorname{span} \{e_1, e_2, \dots, e_M\}$ of $(H^0, \langle \cdot, \cdot \rangle_{\mathbb{C}})$ and the projection operator $P_M : H^0 \to H_M$ by $P_M x = \sum_{k=1}^M \langle x, e_k \rangle_{\mathbb{C}} e_k$ for each $x \in H^0$. Then P_M is also a projection operator from $(U^0, \langle \cdot, \cdot \rangle_{\mathbb{R}})$ onto U_M such that $P_M x = \sum_{k=1}^M \langle x, e_k \rangle_{\mathbb{R}} e_k$ for each $x \in U^0$. Denote $\Delta_M = \Delta P_M$. Using the above notations, we get the following spectral Galerkin approximation:

$$du^{M}(t) = \mathbf{i}\Delta_{M}u^{M}(t)dt + \mathbf{i}\alpha P_{M}dW(t), \qquad t > 0,$$

$$u^{M}(0) = P_{M}u_{0} \in H_{M}.$$
(4.1)

It is verified that (4.1) admits a unique mild solution on H_M given by

$$u^{M}(t) = S_{M}(t)u^{M}(0) + i\alpha \int_{0}^{t} S_{M}(t-s)P_{M}dW(s), \tag{4.2}$$

where $S_M(t) = e^{it\Delta_M}$ is the unitary C_0 -group generated by $i\Delta_M$.

For the spatial discretization (4.1), we define $B_T^M = \frac{u^M(T)}{T}$ which is viewed as a discrete approximation for B_T . In what follows, we study the LDP of $\{B_T^M\}_{T>0}$ and whether $\{u^M\}_{M\in\mathbb{N}}$ can asymptotically preserve the LDP of B_T .

4.1. **LDP for** $\{B_T^M\}_{T>0}$. Following the ideas of deriving the LDP of $\{B_T\}_{T>0}$, in this part, we give the LDP of $\{B_T^M\}_{T>0}$. For this end, we first consider the logarithmic moment generating function $\Lambda^M(\lambda) = \lim_{T\to\infty} \frac{1}{T} \ln \mathbf{E} \exp \{T\langle \lambda, B_T^M\rangle_{\mathbb{R}}\}$, for each $\lambda \in H_M$. Then, we study the exponential tightness of $\{B_T^M\}_{T>0}$. Finally, by means of Theorem 3.4, we obtain the LDP of $\{B_T^M\}_{T>0}$. Hereafter we use the notation $K(a_1,\ldots,a_m)$ to denote some constant dependent on the parameters a_1,\ldots,a_m but independent of T and N, which may vary from one line to another.

Theorem 4.1. For each fixed $M \in \mathbb{N}$, $\{B_T^M\}_{T>0}$ satisfies an LDP on H^0 with the good rate function $I^M(\cdot)$ given by

$$I^{M}(x) = \begin{cases} \frac{1}{\alpha^{2}} \left\| Q_{M}^{-\frac{1}{2}} x \right\|_{H^{0}}^{2}, & if \ x \in Q_{M}^{\frac{1}{2}}(H^{0}), \\ +\infty, & otherwise, \end{cases}$$
(4.3)

where $Q_M := QP_M$ and $Q_M^{-\frac{1}{2}}$ is the pseudo inverse of $Q_M^{\frac{1}{2}}$ on H_M , i.e., $Q_M^{-\frac{1}{2}}x = \underset{z}{\operatorname{argmin}} \{ \|z\|_{H^0} : z \in H_M, Q_M^{\frac{1}{2}}z = x \}$ for every $x \in H_M$.

Proof. Noting that $S_M(t) = \cos(t\Delta_M) + i\sin(t\Delta_M)$, we have

$$u^{M}(T) = S_{M}(T)u^{M}(0) - \alpha \int_{0}^{T} \sin((T-s)\Delta_{M})P_{M}dW(s) + i\alpha \int_{0}^{T} \cos((T-s)\Delta_{M})P_{M}dW(s)$$
$$=: S_{M}(T)u^{M}(0) - \alpha W_{\sin}^{M}(T) + i\alpha W_{\cos}^{M}(T). \tag{4.4}$$

Notice that for each T > 0, $W_{\sin}^M(T)$ is a Gaussian random variable taking values on $(U_M, \langle \cdot, \cdot \rangle_{\mathbb{R}})$. By Proposition 2.1, the covariance operator $\mathbf{Var}(W_{\sin}^M(T))$ of $W_{\sin}^M(T)$ is

$$\mathbf{Var}(W_{\sin}^{M}(T)) = \int_{0}^{T} \sin^{2}((T-s)\Delta_{M})Q_{M}ds$$

$$= \frac{Q_{M}}{2} \int_{0}^{T} \left[I - \cos(2(T-s)\Delta_{M})\right]ds$$

$$= \frac{TQ_{M}}{2} - \frac{Q_{M}\Delta_{M}^{-1}}{4}\sin(2T\Delta_{M}), \tag{4.5}$$

where $Q_M = QP_M$. Similarly, we have that

$$W_{\text{cos}}^M(T) \sim \mathcal{N}(0, \mathbf{Var}(W_{\text{cos}}^M(T)))$$
 on U_M (4.6)

with $\mathbf{Var}(W_{\cos}^M(T)) = \frac{TQ_M}{2} + \frac{1}{4}Q_M\Delta_M^{-1}\sin(2T\Delta_M)$. And the correlation operator $\mathbf{Cor}\left(W_{\sin}^M(T), W_{\cos}^M(T)\right)$ is

$$\mathbf{Cor}\left(W_{\sin}^{M}(T), W_{\cos}^{M}(T)\right) = \frac{Q_{M}\Delta_{M}^{-1}}{4}\left[I - \cos(2T\Delta_{M})\right]. \tag{4.7}$$

For each $\lambda \in H_M$, we write it as $\lambda = \Re \lambda + i \Im \lambda$ with $\Re \lambda$, $\Im \lambda \in U_M$. Then by (4.4),

$$\langle u^M(T), \lambda \rangle_{\mathbb{R}} = \langle S_M(T)u^M(0), \lambda \rangle_{\mathbb{R}} - \alpha \langle W_{\sin}^M(T), \Re \lambda \rangle_{\mathbb{R}} + \alpha \langle W_{\cos}^M(T), \Im \lambda \rangle_{\mathbb{R}}. \tag{4.8}$$

Hence, we obtain

$$\left| \mathbf{E} \langle u^M(T), \lambda \rangle_{\mathbb{R}} \right| = \left| \langle S_M(T) u^M(0), \lambda \rangle_{\mathbb{R}} \right| \le K(\lambda). \tag{4.9}$$

It follows from (4.5), (4.6), (4.7) and (4.8) that

$$\begin{aligned} \mathbf{Var}\langle u^{M}(T), \lambda \rangle_{\mathbb{R}} &= \alpha^{2} \mathbf{Var} \left\langle W_{\sin}^{M}(T), \Re \lambda \right\rangle_{\mathbb{R}} + \alpha^{2} \mathbf{Var} \left\langle W_{\cos}^{M}(T), \Im \lambda \right\rangle_{\mathbb{R}} \\ &- 2\alpha^{2} \mathbf{Var} \left(\left\langle W_{\sin}^{M}(T), \Re \lambda \right\rangle_{\mathbb{R}}, \left\langle W_{\cos}^{M}(T), \Im \lambda \right\rangle_{\mathbb{R}} \right) \\ &= \alpha^{2} \left\langle \mathbf{Var}(W_{\sin}^{M}(T)) \Re \lambda, \Re \lambda \right\rangle_{\mathbb{R}} + \alpha^{2} \left\langle \mathbf{Var}(W_{\cos}^{M}(T)) \Im \lambda, \Im \lambda \right\rangle_{\mathbb{R}} \\ &- 2\alpha^{2} \left\langle \mathbf{Cor}(W_{\sin}^{M}(T), (W_{\cos}^{M}(T)) \Re \lambda, \Im \lambda \right\rangle_{\mathbb{R}} \\ &= \frac{\alpha^{2} T}{2} \left\langle Q_{M} \Re \lambda, \Re \lambda \right\rangle_{\mathbb{R}} + \frac{\alpha^{2} T}{2} \left\langle Q_{M} \Im \lambda, \Im \lambda \right\rangle_{\mathbb{R}} - \frac{\alpha^{2}}{4} \left\langle Q_{M} \Delta_{M}^{-1} \sin(2T\Delta_{M}) \Re \lambda, \Re \lambda \right\rangle_{\mathbb{R}} \\ &+ \frac{\alpha^{2}}{4} \left\langle Q_{M} \Delta_{M}^{-1} \sin(2T\Delta_{M}) \Im \lambda, \Im \lambda \right\rangle_{\mathbb{R}} - \frac{\alpha^{2}}{2} \left\langle Q_{M} \Delta_{M}^{-1} (I - \cos(2T\Delta_{M})) \Re \lambda, \Im \lambda \right\rangle_{\mathbb{R}} \\ &=: \frac{\alpha^{2} T}{2} \left\langle Q_{M} \Re \lambda, \Re \lambda \right\rangle_{\mathbb{R}} + \frac{\alpha^{2} T}{2} \left\langle Q_{M} \Im \lambda, \Im \lambda \right\rangle_{\mathbb{R}} + R(T) \end{aligned} \tag{4.10}$$

with $|R(T)| \leq K(M, \lambda)$. Using (4.9) and (4.10), we have that, for every $\lambda \in H_M$,

$$\begin{split} \Lambda^{M}(\lambda) &= \lim_{T \to \infty} \frac{1}{T} \ln \mathbf{E} e^{T \left\langle B_{T}^{M}, \lambda \right\rangle_{\mathbb{R}}} = \lim_{T \to \infty} \frac{1}{T} \ln \mathbf{E} e^{\left\langle u^{M}(T), \lambda \right\rangle_{\mathbb{R}}} \\ &= \lim_{T \to \infty} \frac{1}{T} \left(\mathbf{E} \left\langle u^{M}(T), \lambda \right\rangle_{\mathbb{R}} + \frac{1}{2} \mathbf{Var} \left\langle u^{M}(T), \lambda \right\rangle_{\mathbb{R}} \right) \end{split}$$

$$\begin{split} &= \frac{1}{2} \left(\frac{\alpha^2}{2} \langle Q_M \Re \lambda, \Re \lambda \rangle_{\mathbb{R}} + \frac{\alpha^2}{2} \langle Q_M \Im \lambda, \Im \lambda \rangle_{\mathbb{R}} \right) \\ &= \frac{\alpha^2}{4} \left\| Q_M^{\frac{1}{2}} \lambda \right\|_{H^0}^2. \end{split}$$

Analogous to the proof of (3.12), we have

$$(\Lambda^{M})^{*}(x) = \sup_{\lambda \in H_{M}} \left\{ \langle \lambda, x \rangle - \Lambda^{M}(\lambda) \right\} = \begin{cases} \frac{1}{\alpha^{2}} \left\| Q_{M}^{-\frac{1}{2}} x \right\|_{H^{0}}^{2}, & \text{if } x \in Q_{M}^{\frac{1}{2}}(H_{M}), \\ +\infty & \text{otherwise.} \end{cases}$$

Next, we show that $\{B_T^M\}_{T>0}$ is exponentially tight. Define $K_L = \{f \in H_M \mid ||f||_{H^0} \leq L\}$, then K_L is the compact subset of H_M . It follows from (4.4) that

$$\mathbf{P}\left(B_{T}^{M} \in K_{L}^{c}\right) = \mathbf{P}\left(\|u^{M}(T)\|_{H^{0}} > LT\right)$$

$$\leq \mathbf{P}\left(\|S_{M}(T)u_{0}^{M}\|_{H^{0}} > \frac{TL}{3}\right) + \mathbf{P}\left(\alpha\|W_{\sin}^{M}(T)\|_{U^{0}} > \frac{TL}{3}\right) + \mathbf{P}\left(\alpha\|W_{\cos}^{M}(T)\|_{U^{0}} > \frac{TL}{3}\right). \quad (4.11)$$

By (4.5), we have

$$\frac{W_{\sin}^{M}(T)}{\sqrt{T}} \sim \mathcal{N}\left(0, \left(\frac{I}{2} - \frac{\Delta_{M}^{-1}\sin(2T\Delta_{M})}{4T}\right)Q_{M}\right) \quad \text{on } U_{M}.$$
(4.12)

Hence, we obtain

$$\lambda_k \left(\mathbf{Var} \left(\frac{W_{\sin}^M(T)}{\sqrt{T}} \right) \right) = \left(\frac{1}{2} - \frac{\sin(2Tk^2)}{4Tk^2} \right) \eta_k = \frac{1}{2} \left(1 - \frac{\sin(2Tk^2)}{2Tk^2} \right) \eta_k < \eta_k, \qquad k = 1, 2, \dots, M.$$

For every $0 < \varepsilon < \frac{1}{2n_1}$, it follows from Proposition 3.8 that

$$\mathbf{E} \exp \left\{ \varepsilon \left\| \frac{W_{\sin}^M(T)}{\sqrt{T}} \right\|_{U^0}^2 \right\} = \left[\det \left(I - 2\varepsilon \mathbf{Var} \left(\frac{W_{\sin}^M(T)}{\sqrt{T}} \right) \right) \right]^{-\frac{1}{2}} < \left[\det (I - 2\varepsilon Q_M) \right]^{-\frac{1}{2}} = C(\varepsilon, Q_M).$$

The above formula yields

$$\mathbf{P}\left(\alpha \|W_{\sin}^{M}(T)\|_{U^{0}} > \frac{TL}{3}\right) = P\left(\exp\left\{\varepsilon \left\|\frac{W_{\sin}^{M}(T)}{\sqrt{T}}\right\|_{U^{0}}^{2}\right\} > \exp\left\{\frac{\varepsilon TL^{2}}{9\alpha^{2}}\right\}\right)$$

$$\leq e^{-\frac{\varepsilon TL^{2}}{9\alpha^{2}}}\mathbf{E}\exp\left\{\varepsilon \left\|\frac{W_{\sin}^{M}(T)}{\sqrt{T}}\right\|_{U^{0}}^{2}\right\} \leq e^{-\frac{\varepsilon TL^{2}}{9\alpha^{2}}}C(\varepsilon, Q_{M}). \tag{4.13}$$

Similarly, one has

$$\mathbf{P}\left(\alpha \|W_{\cos}^{M}(T)\|_{U^{0}} > \frac{TL}{3}\right) \leq e^{-\frac{\varepsilon TL^{2}}{9\alpha^{2}}} C(\varepsilon, Q_{M}). \tag{4.14}$$

According to Proposition 3.7, (4.13) and (4.14), we have

$$\limsup_{T \to \infty} \frac{1}{T} \ln \mathbf{P} \left(B_T^M \in K_L^c \right) \le -\frac{\varepsilon L^2}{9\alpha^2}, \qquad 0 < \varepsilon < \frac{1}{2\eta_1},$$

where we have used the fact that $\mathbf{P}\left(\|S_M(T)u_0^M\|_{H^0} > \frac{TL}{3}\right) = 0$ for sufficiently large T. Then, we obtain

$$\lim_{L \to \infty} \limsup_{T \to \infty} \frac{1}{T} \ln \mathbf{P} \left(B_T^M \in K_L^c \right) = -\infty,$$

which implies the exponential tightness of $\{B_T^M\}_{T>0}$.

Notice that $\Lambda^M(\cdot)$ is Fréchet differentiable and $\mathcal{D}\Lambda^M(\lambda)(\cdot) = \frac{\alpha^2}{2} \langle Q_M \lambda, \cdot \rangle$ for each $\lambda \in H_M$. Then it follows from Theorem 3.4 that $\{B_T^M\}_{T>0}$ satisfies an LDP on H_M with the good rate function

$$\widetilde{I}^{M}(x) = (\Lambda^{M})^{*}(x) = \begin{cases} \frac{1}{\alpha^{2}} \left\| Q_{M}^{-\frac{1}{2}} x \right\|_{H^{0}}^{2}, & \text{if } x \in Q_{M}^{\frac{1}{2}}(H_{M}), \\ +\infty, & x \in H_{M} \setminus Q_{M}^{\frac{1}{2}}(H_{M}). \end{cases}$$

Clearly, H_M is the closed subspace of H^0 and for each T>0, $\mathbf{P}(B_T^M\in H^M)=1$. Thus, using Lemma 3.6 and the fact $Q_M^{\frac{1}{2}}(H_M)=Q_M^{\frac{1}{2}}(H^0)$, we conclude that $\{B_T^M\}_{T>0}$ satisfies an LDP on H^0 with the good rate function

$$I^{M}(x) = \begin{cases} \frac{1}{\alpha^{2}} \left\| Q_{M}^{-\frac{1}{2}} x \right\|_{H^{0}}^{2}, & \text{if } x \in Q_{M}^{\frac{1}{2}}(H^{0}), \\ +\infty, & \text{otherwise.} \end{cases}$$

4.2. Weakly asymptotical preservation for the LDP of $\{B_T\}_{T>0}$. In the last subsection, we obtain the LDP for $\{B_T^M\}_{T>0}$ of the spectral Galerkin approximation $\{u^M(T)\}_{T>0}$. It is natural to consider whether I^M converges to I pointwise as M tends to infinity. In [3], authors give the definition of asymptotical preservation for the LDP of the original system, i.e., the discrete rate functions of numerical methods converge to that of the original system in the pointwise sense. In our case, since generally $Q_M^{\frac{1}{2}}(H^0) \subsetneq Q^{\frac{1}{2}}(H^0)$, it can not be assured that I^M converges to I pointwise. However, the sequence $\left\{Q_M^{\frac{1}{2}}(H^0)\right\}_{M\in\mathbb{N}}$ of sets converges to $Q^{\frac{1}{2}}(H^0)$ by the fact $\lim_{M\to\infty}Q_M^{\frac{1}{2}}x=Q^{\frac{1}{2}}x$ for each $x\in H^0$. It is hoped that I^M is a good approximation of I when M is large enough. Thus, we give the following definition.

Definition 4.2. For a spatial semi-discretization $\{u^M\}_{M\in\mathbb{N}}$ of (1.1), denote $B_T^M = \frac{u^M(T)}{T}$. Assume that $\{B_T^M\}_{T>0}$ satisfies an LDP on H^0 with the rate function I^M for all sufficiently large M. Then we say that $\{u^M\}_{M\in\mathbb{N}}$ weakly asymptotically preserves the LDP of $\{B_T\}_{T>0}$ if for each $x \in Q^{\frac{1}{2}}(H^0)$ and $\varepsilon > 0$, there exist $x_0 \in H^0$ and $M \in \mathbb{N}$ such that

$$||x - x_0||_{H^0} < \varepsilon, \qquad |I(x) - I^M(x_0)| < \varepsilon,$$
 (4.15)

where I is the rate function of $\{B_T\}_{T>0}$.

Theorem 4.3. For the spectral Galerkin approximation (4.1), $\{u^M\}_{M\in\mathbb{N}}$ weakly asymptotically preserves the LDP of $\{B_T\}_{T>0}$, i.e., (4.15) holds.

Proof. This problem is discussed in the following two cases.

Case 1: There are infinitely many 0 in $\{\eta_k\}_{k\in\mathbb{N}}$, i.e., for some $l\in\mathbb{N}$, $\eta_l>\eta_{l+1}=\eta_{l+2}=\cdots=0$. For this case, Q degenerates to a finite-rank operator. If $M\geq l$, then $Q_M=Q$. Hence, it holds that $I^M(x)=I(x)$ for every $x\in H^0$, which implies (4.15). We say that $\{u^M\}_{M\in\mathbb{N}}$ exactly preserves the LDP of $\{B_T\}_{T>0}$ for this case (see [3, Definition 4.1]).

Case 2: There are finitely many 0 in $\{\eta_k\}_{k\in\mathbb{N}}$.

Notice that for each finite $M \in \mathbb{N}$, $\eta_1 \geq \eta_2 \geq \cdots \geq \eta_M > 0$. We denote $y = Q^{-\frac{1}{2}}x$ and define

 $x_M := Q_M^{\frac{1}{2}} y$. Further, we have

$$\begin{split} Q_{M}^{-\frac{1}{2}}x_{M} = & argmin \left\{ \|z\|_{H^{0}}: \ z \in H_{M}, \ Q_{M}^{\frac{1}{2}}z = x_{M} \right\} \\ = & argmin \left\{ \|z\|_{H^{0}}: \ z \in H_{M}, \ Q^{\frac{1}{2}}z = Q^{\frac{1}{2}}P_{M}y \right\} \\ = & argmin \left\{ \|z\|_{H^{0}}: \ z \in H_{M}, \ \sqrt{\eta_{k}}\langle z, e_{k}\rangle_{\mathbb{C}} = \sqrt{\eta_{k}}\langle y, e_{k}\rangle_{\mathbb{C}}, \ k = 1, 2 \dots, M \right\} \\ = & P_{M}y. \end{split}$$

The above formula yields

$$\lim_{M \to \infty} \left| I^M(x_M) - I(x) \right| = \frac{1}{\alpha^2} \lim_{M \to \infty} \left| \|P_M y\|_{H^0}^2 - \|y\|_{H^0}^2 \right| = 0. \tag{4.16}$$

In addition, it holds that

$$\lim_{M \to \infty} x_M = \lim_{M \to \infty} Q_M^{\frac{1}{2}} y = \lim_{M \to \infty} P_M Q^{\frac{1}{2}} y = Q^{\frac{1}{2}} y = x. \tag{4.17}$$

Thus, it follows from (4.16) and (4.17) that for each $x \in Q^{\frac{1}{2}}(H^0)$ and $\varepsilon > 0$, there exist sufficiently large M and $x_0 = Q_M^{\frac{1}{2}}\left(Q^{-\frac{1}{2}}x\right)$ such that (4.15) holds.

Combining Case 1 and Case 2, we complete the proof.

Remark 4.4. As is seen in the proof of Theorem 4.3, for every $x \in Q^{\frac{1}{2}}(H^0)$ and sufficiently large M, $I^M(Q_M^{\frac{1}{2}}Q^{-\frac{1}{2}}x)$ is a good approximation of I(x).

5. LDP BY SPATIO-TEMPORAL FULL DISCRETIZATION

In this section, we investigate the LDP for the full discretizations, spatially by the spectral Galerkin method and temporally by the symplectic methods or non-symplectic ones. We show that the full discretization weakly asymptotically preserves the LDP of $\{B_T\}_{T>0}$ when using a symplectic method in temporal direction, while it does not share this property for a temporal non-symplectic method. These results indicate that the modified rate function of the full discretization, based on the spatial spectral Galerkin method and a temporal symplectic method, is a good approximation of I.

5.1. **Full discretization.** Since the spectral Galerkin approximation $\{u^M(t)\}_{t\geq 0}$ takes values in H_M , it holds that $u^M(t) = \sum_{k=1}^M \langle u^M(t), e_k \rangle_{\mathbb{C}} e_k$. Denote $U^M(t) = (\langle u^M(t), e_1 \rangle_{\mathbb{C}}, \langle u^M(t), e_2 \rangle_{\mathbb{C}}, \cdots, \langle u^M(t), e_M \rangle_{\mathbb{C}})^{\top}$. Let $U^{M,k}(t)$ be the kth component of $U^M(t)$. It follows from (4.1) that

$$dU^{M,k}(t) = -ik^2 U^{M,k}(t) dt + i\alpha \sqrt{\eta_k} d\beta_k(t), \qquad k = 1, 2, \dots, M.$$

Then, we obtain a \mathbb{C}^M -valued SDE

$$dU^{M}(t) = -i\mathcal{M}U^{M}(t)dt + i\alpha \mathcal{Q}d\beta(t),$$

where $\mathcal{M} = \operatorname{diag}(1, 2^2, \dots, M^2) \in \mathbb{R}^{M \times M}$, $\mathcal{Q} = \operatorname{diag}(\sqrt{\eta_l}, \sqrt{\eta_2}, \dots, \sqrt{\eta_M}) \in \mathbb{R}^{M \times M}$, and $\beta(t) = (\beta_1(t), \beta_2(t), \dots, \beta_M(t))^\top \in \mathbb{R}^M$. Further, using the notation $U^M(t) = P^M(t) + iQ^M(t)$ with $P^M(t) = \Re U^M(t)$ and $Q^M(t) = \Im U^M(t)$, we obtain a 2M-dimensional stochastic Hamiltonian system

$$dP^{M}(t) = \mathcal{M}Q^{M}(t)dt,$$

$$dQ^{M}(t) = -\mathcal{M}P^{M}(t)dt + \alpha \mathcal{Q}d\beta(t),$$
(5.1)

which is equivalent to the system (4.1) with $\langle u^M(t), e_k \rangle_{\mathbb{C}} = P^{M,k}(t) + iQ^{M,k}(t)$, where $P^{M,k}$ and $Q^{M,k}$ are the kth arguments of P^M and Q^M , respectively. Hence, in order to obtain the numerical method for (4.1), we only need to consider discretizing the equivalent system (5.1).

Denote by $\{(p_n^M, q_n^M)\}_{n \in \mathbb{N}}$ the numerical approximation of $\{(P^M(t), Q^M(t))\}_{t \geq 0}$. Let F be the linear function from \mathbb{C}^M to H^0 defined by

$$F(z) = \sum_{k=1}^{M} z_k e_k, \quad \forall \quad z = (z_1, z_2, \dots, z_M) \in \mathbb{C}^M.$$
 (5.2)

Then we obtain the numerical solution $\{u_n^M\}_{n\in\mathbb{N}}$ with $u_n^M:=F(p_n^M+iq_n^M)$. Further, we define $B_N^M=\frac{u_N^M}{N\tau}$ (see [3]), where τ is the temporal stepsize. Then B_N^M is a discrete approximation of B_T . To give the LDP for $\{B_N^M\}_{N\in\mathbb{N}}$, our idea is to first investigate the LDP of $\{A_N^M\}_{N\in\mathbb{N}}$, where $A_N^M=\frac{p_N^M+iq_N^M}{N\tau}$. Then noting that $B_N^M=F(A_N^M)$, combining the LDP of $\{A_N^M\}_{N\in\mathbb{N}}$ on \mathbb{C}^M and the contraction principle (Lemma 3.5), we derive the LDP of $\{B_N^M\}_{N\in\mathbb{N}}$. More precisely, we divide (5.1) into the following M subsystems

$$d\begin{pmatrix} P^{M,k}(t) \\ Q^{M,k}(t) \end{pmatrix} = k^2 \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} P^{M,k}(t) \\ Q^{M,k}(t) \end{pmatrix} dt + \alpha_k \begin{pmatrix} 0 \\ 1 \end{pmatrix} d\beta_k(t), \qquad k = 1, 2, \dots, M,$$
 (5.3)

where $\alpha_k = \alpha \sqrt{\eta_k}$, k = 1, 2, ..., M. For each $k \in \{1, 2, ..., M\}$, we consider the general numerical method in the following form

$$\begin{pmatrix} p_{n+1}^{M,k} \\ q_{n+1}^{M,k} \end{pmatrix} = \begin{pmatrix} a_{11}(k^2\tau) & a_{12}(k^2\tau) \\ a_{21}(k^2\tau) & a_{22}(k^2\tau) \end{pmatrix} \begin{pmatrix} p_n^{M,k} \\ q_n^{M,k} \end{pmatrix} + \alpha_k \begin{pmatrix} b_1(k^2\tau) \\ b_2(k^2\tau) \end{pmatrix} \delta\beta_{k,n}, \tag{5.4}$$

where $\delta\beta_{k,n} = \beta_k(t_{n+1}) - \beta_k(t_n)$ with $t_n = n\tau$, n = 1, 2, ..., and functions a_{ij} , $b_i : (0, \infty) \to \mathbb{R}$, i, j = 1, 2 are determined by a concrete method. In addition, we require $b_1^2(h) + b_2^2(h) \neq 0$ for all sufficiently small h. Hence, we finally obtain the numerical solution $\{(p_n^M, q_n^M)\}_{n \in \mathbb{N}}$ generated by (5.4), with $(p_n^{M,k}, q_n^{M,k})$ being the kth component of (p_n^M, q_n^M) , n = 1, 2, ... By defining functions

$$A(h) := \begin{pmatrix} a_{11}(h) & a_{12}(h) \\ a_{21}(h) & a_{22}(h) \end{pmatrix}, \qquad B(h) := \begin{pmatrix} b_1(h) \\ b_2(h) \end{pmatrix}, \qquad \forall \quad h > 0,$$
 (5.5)

we rewrite (5.4) as

$$\begin{pmatrix} p_{n+1}^{M,k} \\ q_{n+1}^{M,k} \end{pmatrix} = A(k^2 \tau) \begin{pmatrix} p_n^{M,k} \\ q_n^{M,k} \end{pmatrix} + \alpha_k B(k^2 \tau) \delta \beta_{k,n}, \qquad n = 0, 1, 2 \dots$$
 (5.6)

with $p_0^{M,k} + iq_0^{M,k} = \langle u^M(0), e_k \rangle_{\mathbb{C}}$.

Next we introduce some concrete temporal discretizations taking the form (5.6).

Example 5.1 (Midpoint Scheme). Applying midpoint scheme to (4.1) yields

$$u_{n+1}^{M} = u_{n}^{M} + \frac{1}{2} i \tau \Delta_{M} \left(u_{n}^{M} + u_{n+1}^{M} \right) + i \alpha P_{M} \delta W_{n}, \qquad n = 0, 1, 2 \dots,$$

with

$$A^{1}(h) := \frac{1}{4+h^{2}} \begin{pmatrix} 4-h^{2} & 4h \\ -4h & 4-h^{2} \end{pmatrix}, \qquad B^{1}(h) := \frac{2}{4+h^{2}} \begin{pmatrix} h \\ 2 \end{pmatrix}, \qquad \forall \quad h > 0.$$

Here $\delta W_n := W(t_{n+1}) - W(t_n)$.

Example 5.2 (Exponential Euler Method). The exponential Euler method for (4.1) is

$$u_{n+1}^M = S_M(\tau)u_n^M + \boldsymbol{i}\alpha S_M(\tau)P_M\delta W_n, \qquad n = 0, 1, 2\dots,$$

with

$$A^2(h) := \left(\begin{array}{cc} \cos(h) & \sin(h) \\ -\sin(h) & \cos(h) \end{array} \right), \qquad B^2(h) := \left(\begin{array}{c} \sin(h) \\ \cos(h) \end{array} \right), \qquad \forall \quad h > 0.$$

Example 5.3 (Backward Euler–Maruyama Method). The backward Euler–Maruyama method for (4.1) reads

$$u_{n+1}^M = u_n^M + i\tau \Delta_M u_{n+1}^M + i\alpha P_M \delta W_n, \qquad n = 0, 1, 2 \dots,$$

with

$$A^3(h):=\frac{1}{1+h^2}\left(\begin{array}{cc} 1 & h \\ -h & 1 \end{array}\right), \qquad B^3(h):=\frac{1}{1+h^2}\left(\begin{array}{c} h \\ 1 \end{array}\right), \qquad \forall \quad h>0.$$

Next, we give our main assumptions on functions A and B, which will be used to derive the LDP of $\{B_N^M\}_{N\in\mathbb{N}}$.

Assumption 1. There is some $h_1 > 0$ such that

$$4 \det(A(h)) - (\operatorname{tr}(A(h)))^2 > 0, \quad \forall \quad h < h_1,$$

where tr(A) and det(A) denote the trace and the determinant of A, respectively.

We will use **Assumption 1** to give the general expression of the method (5.4), following the idea of [3].

Assumption 2. There is some $h_2 > 0$ such that for all $h < h_2$, det(A(h)) = 1.

One can show that the numerical method generated by (5.4) is symplectic if and only if **Assumption 2** holds. In fact, $\{(p_n^M, q_n^M)\}_{n \in \mathbb{N}}$ generated by (5.4) is symplectic for all sufficiently small $\tau > 0$ if and only if for all sufficiently small $\tau > 0$, $dp_{n+1}^M \wedge dq_{n+1}^M = dp_n^M \wedge dq_n^M$, i.e.,

$$\sum_{k=1}^{M} dp_{n+1}^{M,k} \wedge dq_{n+1}^{M,k} = \sum_{k=1}^{M} dp_n^{M,k} \wedge dq_n^{M,k}, \qquad n = 1, 2, \dots$$

According to (5.4), it holds that $dp_{n+1}^{M,k} \wedge dq_{n+1}^{M,k} = \left(a_{11}(k^2\tau)a_{22}(k^2\tau) - a_{12}(k^2\tau)a_{21}(k^2\tau)\right)dp_n^{M,k} \wedge dq_n^{M,k}$. Hence, the method generated by (5.4) is symplectic for all sufficiently small $\tau > 0$ if and only if for all sufficiently small $\tau > 0$, $k = 1, 2, \ldots, M$,

$$a_{11}(k^2\tau)a_{22}(k^2\tau) - a_{12}(k^2\tau)a_{21}(k^2\tau) = 1,$$

which is equivalent to that there is some $h_0 > 0$ such that

$$a_{11}(h)a_{22}(h) - a_{12}(h)a_{21}(h) = 1, \quad \forall \quad h < h_0,$$

i.e., **Assumption 2** holds.

Assumption 3. There exist some $\eta \in (0,1)$ and some $h_3 > 0$ such that

$$|c(h)| < (1 - \eta)\sqrt{a(h)b(h)}, \quad \forall \quad h < h_3.$$

Here, functions $a, b, c: (0, \infty) \to \mathbb{R}$ are defined by

$$a = (a_{11}b_1 + a_{12}b_2 - b_1)^2 + b_1(a_{11}b_1 + a_{12}b_2)(2 - \operatorname{tr}(A)),$$

$$b = (a_{21}b_1 - a_{11}b_2 + b_2)^2 - b_2(a_{21}b_1 - a_{11}b_2)(2 - \operatorname{tr}(A)),$$

$$c = \frac{1}{2} (a_{21}b_1 - a_{11}b_2) b_1 \operatorname{tr}(A) + b_1 b_2 \left(\frac{1}{2} (\operatorname{tr}(A))^2 - 1 \right) - (a_{11}b_1 + a_{12}b_2) (a_{21}b_1 - a_{11}b_2) - \frac{1}{2} \operatorname{tr}(A) (a_{11}b_1 + a_{12}b_2) b_2.$$

Assumption 3 is used to give the explicit expression of the rate functions of $\{A_N^M\}_{N\in\mathbb{N}}$ and $\{B_N^M\}_{N\in\mathbb{N}}$. In fact, a(h),b(h)>0 for sufficiently small h, whose proof is similar to those of Lemmas 3.2 and 5.1 in [3]. In addition, we have the following property.

Remark 5.4. Under **Assumption 2**, $c = \frac{a_{11} - a_{22}}{2} \left[a_{12}b_2^2 - a_{21}b_1^2 + b_1b_2(a_{11} - a_{22}) \right]$.

This is because under **Assumption 2**, $det(A) = a_{11}a_{22} - a_{12}a_{21} = 1$. Then it follows that

$$c = b_1^2 \left(\frac{1}{2} a_{21} \operatorname{tr}(A) - a_{11} a_{21} \right) + b_2^2 \left(a_{11} a_{12} - \frac{1}{2} a_{12} \operatorname{tr}(A) \right)$$

$$+ b_1 b_2 \left[\frac{1}{2} (\operatorname{tr}(A))^2 - 1 - (a_{12} a_{21} - a_{11}^2) - a_{11} \operatorname{tr}(A) \right]$$

$$= b_1^2 \left(\frac{1}{2} a_{21} \operatorname{tr}(A) - a_{11} a_{21} \right) + b_2^2 \left(a_{11} a_{12} - \frac{1}{2} a_{12} \operatorname{tr}(A) \right)$$

$$+ b_1 b_2 \left[\frac{1}{2} (a_{11} - a_{22})^2 + a_{11} a_{22} - a_{12} a_{21} - 1 \right]$$

$$= \frac{a_{11} - a_{22}}{2} \left[a_{12} b_2^2 - a_{21} b_1^2 + b_1 b_2 \left(a_{11} - a_{22} \right) \right].$$

When we investigate the LDP of $\{B_N^M\}_{N\in\mathbb{N}}$ via temporal non-symplectic methods, we give the following assumption (see [3]).

Assumption 4. There is some $h_4 > 0$ such that for all $h < h_4$, $\det(A(h)) < 1$.

In addition, when investigating the asymptotical preservation of $\{u_n^M\}_{M,n\in\mathbb{N}}$ for the LDP of $\{B_T\}_{T>0}$, we give the following assumption concerning the convergence of the numerical method.

Assumption 5.
$$|a_{11} - 1| + |a_{22} - 1| + |a_{12} - h| + |a_{21} + h| = \mathcal{O}(h^2)$$
, and $|b_1| + |b_2 - 1| = \mathcal{O}(h)$.

One can prove that under **Assumption 5**, $\{(p_n^M, q_n^M)\}_{n \in \mathbb{N}}$ corresponding to (5.4) has at least first order convergence in mean-square sense. For more details, one refers to [3].

It is verified that the methods in Examples 1 and 2 are symplectic satisfying **Assumptions 1-3** and 5. And the method in Example 3 is non-symplectic satisfying **Assumptions 1 and 4**.

To characterize the asymptotical preservation of $\{u_n^M\}_{M,n\in\mathbb{N}}$ for the LDP of $\{B_T\}_{T>0}$, we give the following definition (see [3] for the similar definition).

Definition 5.5. For a spatio-temporal full discretization $\{u_n^M\}_{M,n\in\mathbb{N}}$ of (1.1) with temporal stepsize τ , denote $B_N^M = \frac{u_N^M}{N\tau}$. Assume that for each fixed $M \in \mathbb{N}$, $\{B_N^M\}_{N\in\mathbb{N}}$ satisfies an LDP on H^0 with the rate function $I^{M,\tau}$. We call $I_{mod}^{M,\tau} := \frac{I^{M,\tau}}{\tau}$ the modified rate function. Then $\{u_n^M\}_{M,n\in\mathbb{N}}$ is said to weakly asymptotically preserves the LDP of $\{B_T\}_{T>0}$ if for each $x \in Q^{\frac{1}{2}}(H^0)$ and $\varepsilon > 0$, there exist $x_0 \in H^0$, M > 0 and $\tau > 0$ such that

$$||x - x_0||_{H^0} < \varepsilon, \qquad \left| I(x) - I_{mod}^{M,\tau}(x_0) \right| < \varepsilon.$$
 (5.7)

With the above preparation, we give our main results of this paper. That is, for the full discretization $\{u_n^M\}_{M,n\in\mathbb{N}}$ with $u_n^M=F(p_n^M+iq_n^M)$, where $\{p_n^M,q_n^M\}_{M,n\in\mathbb{N}}$ is the numerical solution

corresponding to (5.4), when the temporal discretization is symplectic, it weakly asymptotically preserves the LDP of $\{B_T\}_{T>0}$, while it does not possess this property for a temporal non-symplectic discretization.

Theorem 5.6. If Assumptions 1, 2 and 5 hold, then

(1) For each fixed $M \in \mathbb{N}$ with $\eta_M > 0$, we have that for all sufficiently small stepsize τ , $\{B_N^M\}_{N\in\mathbb{N}}$ satisfies an LDP on H^0 with the good rate function given by

$$I^{M,\tau}(x)$$

$$= \begin{cases} \sum_{k=1}^{M} \frac{\tau\left(4 - \left(\operatorname{tr}(\mathbf{A}(\mathbf{k}^{2}\tau))\right)^{2}\right)}{4[a(k^{2}\tau)b(k^{2}\tau) - c^{2}(k^{2}\tau)]\alpha_{k}^{2}} \left[b(k^{2}\tau)(\Re\langle x, e_{k}\rangle_{\mathbb{C}})^{2} + a(k^{2}\tau)(\Im\langle x, e_{k}\rangle_{\mathbb{C}})^{2} \\ + 2c(k^{2}\tau)\Re\langle x, e_{k}\rangle_{\mathbb{C}}\Im\langle x, e_{k}\rangle_{\mathbb{C}}\right], & if \ x \in H_{M}, \\ +\infty, & otherwise. \end{cases}$$
(5.8)

(2) For each fixed $M \in \mathbb{N}$ with $\eta_M > 0$, $\{u_N^M\}_{N \in \mathbb{N}}$ asymptotically preserves the LDP of $\{B_T^M\}_{T>0}$, i.e., the modified rate function satisfies

$$\lim_{\tau \to 0} I_{mod}^{M,\tau}(x) = I^{M}(x). \tag{5.9}$$

(3) $\{u_n^M\}_{M,n\in\mathbb{N}}$ weakly asymptotically preserves the LDP for $\{B_T\}_{T>0}$ of (1.1), i.e., (5.7) holds.

Theorem 5.7. If **Assumptions 1 and 4** hold, then for each $M \in \mathbb{N}$, $\{B_N^M\}_{N \in \mathbb{N}}$ satisfies an LDP on H^0 with the good rate function

$$I_{ns}^{M,\tau}(x) = \begin{cases} 0, & \text{if } x = 0, \\ +\infty, & \text{otherwise.} \end{cases}$$

Moreover, $\{u_n^M\}_{M,n\in\mathbb{N}}$ can not weakly asymptotically preserve the LDP for $\{B_T\}_{T>0}$ of (1.1), i.e., (5.7) does not hold.

For the method in Example 1, we have that for each $M \in \mathbb{N}$, there is sufficiently small τ , $\{B_N^M\}_{N\in\mathbb{N}}$ satisfies an LDP on H^0 with the good rate function

$$I_1^{M,\tau}(x) = \begin{cases} \frac{\tau}{\alpha^2} \left\| \left(I + \frac{\tau^2 \Delta_M^2}{4} \right) Q_M^{-\frac{1}{2}} x \right\|, & \text{if } x \in Q_M^{\frac{1}{2}}(H_0), \\ +\infty, & \text{otherwise.} \end{cases}$$

Hence, $\lim_{\tau \to 0} I_{1,mod}^{M,\tau}(x) := \lim_{\tau \to 0} I_1^{M,\tau}(x)/\tau = I^M(x)$ for each $x \in H^0$. In addition, for each h > 0, **Assumptions 1 and 2** hold. These verify that midpoint scheme satisfies both the conditions and the first conclusion of Theorem 5.6. Finally, combining Theorem 4.3, we have that the full discretization spatially, by a spatial Galerkin method and temporally by the midpoint scheme, weakly asymptotically the LDP of $\{B_T\}_{T>0}$.

Next we give the rate functions of $\{B_N^M\}_{N\in\mathbb{N}}$ when using the methods in Examples 1-3.

• Midpoint Scheme The rate function of $\{B_N^M\}_{N\in\mathbb{N}}$ is

$$I_1^{M,\tau}(x) = \begin{cases} \frac{\tau}{\alpha^2} \left\| \left(I + \frac{\tau^2 \Delta_M^2}{4} \right) Q_M^{-\frac{1}{2}} x \right\|, & \text{if } x \in Q_M^{\frac{1}{2}}(H_0), \\ +\infty, & \text{otherwise.} \end{cases}$$

• Exponential Euler Method The rate function of $\{B_N^M\}_{N\in\mathbb{N}}$ is

$$I_2^{M,\tau}(x) = \begin{cases} \frac{\tau}{\alpha^2} \left\| Q_M^{-\frac{1}{2}} x \right\|, & \text{if } x \in Q_M^{\frac{1}{2}}(H_0), \\ +\infty, & \text{otherwise.} \end{cases}$$

In particular, we note that if Q is a finite rank operator, i.e., there is $l \in \mathbb{N}$ such $\eta_{l+1} = \eta_{l+2} = \cdots = 0$, then $I_{2,mod}^{l,\tau} = I$. This indicates when noise takes values in finite dimensional space, this full discretization preserves exactly the LDP of $\{B_T\}_{T>0}$.

• Backward Euler–Maruyama Method The rate function of $\{B_N^M\}_{N\in\mathbb{N}}$ is

$$I_3^{M,\tau}(x) = \begin{cases} 0, & \text{if } x = 0, \\ +\infty, & \text{otherwise.} \end{cases}$$

5.2. **Proof of Theorem 5.6.** In this part, we consider the LDP of $\{B_N^M\}_{N\in\mathbb{N}}$ for the full discretizations of (1.1), spatially by the spectral Galerkin method (4.1) and temporally by symplectic methods. To this end, we let **Assumption 2** hold throughout this part. Firstly, for every fixed $k \in \{1, 2, ..., M\}$, we derive the limit $\Lambda_k(z) := \lim_{N\to\infty} \frac{1}{N} \ln \mathbf{E} \exp\left\{\frac{1}{\tau} \left\langle z, p_N^{M,k} + iq_N^{M,k} \right\rangle_{\mathbb{R}}\right\}$ for $z \in \mathbb{C}$, to give the expression of the logarithmic moment generating function $\Lambda^{M,\tau}(\lambda) = \lim_{N\to\infty} \frac{1}{N} \ln \mathbf{E} \exp\left\{N \left\langle \lambda, A_N^M \right\rangle_{\mathbb{R}}\right\}$ of $\{A_N^M\}_{N\in\mathbb{N}}$. Then using Theorem 3.4, we obtain the LDP of $\{A_N^M\}_{N\in\mathbb{N}}$ for symplectic methods. Further, the contraction principle (Lemma 3.5) leads to the LDP of $\{B_N^M\}_{N\in\mathbb{N}}$ with $B_N^M := F(A_N^M)$. Finally combining the convergence condition (**Assumption 5**), we prove that $\{u_n^M\}_{M,n\in\mathbb{N}}$ weakly asymptotically preserves the LDP of $\{B_T\}_{T>0}$, which completes the proof of Theorem 5.6.

Lemma 5.8. If **Assumptions 1 and 2** hold, then for each fixed $M \in \mathbb{N}$, we have that for all sufficiently small stepsize τ ,

$$\Lambda^{M,\tau}(\boldsymbol{\lambda}) = \sum_{k=1}^{M} \Lambda_k(\lambda_k) = \sum_{k=1}^{M} \frac{\alpha_k^2}{4\tau \sin^2(\theta_k)} \left\{ a(k^2\tau)(\Re \lambda_k)^2 + b(k^2\tau)(\Im \lambda_k)^2 - 2c(k^2\tau)\Re \lambda_k \Im \lambda_k \right\},\tag{5.10}$$

where a, b, c are given in **Assumption 3**. Moreover, $\Lambda^{M,\tau}$ is finite valued and Gateaux differentiable. Proof. For each $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_M) \in \mathbb{C}^M$, we have

$$\left\langle oldsymbol{\lambda}, p_N^M + oldsymbol{i} q_N^M
ight
angle_{\mathbb{R}} = \sum_{k=1}^M \left\langle \lambda_k, p_N^{M,k} + oldsymbol{i} q_N^{M,k}
ight
angle_{\mathbb{R}} = \sum_{k=1}^M \left(\Re \lambda_k p_N^{M,k} + \Im \lambda_k q_N^{M,k}
ight).$$

Thus, the logarithmic moment generating function for $\{A_N^M\}_{N\in\mathbb{N}}$ is

$$\begin{split} \Lambda^{M,\tau}(\boldsymbol{\lambda}) &= \lim_{N \to \infty} \frac{1}{N} \ln \mathbf{E} \exp \left\{ N \left\langle \boldsymbol{\lambda}, A_N^M \right\rangle_{\mathbb{R}} \right\} \\ &= \lim_{N \to \infty} \frac{1}{N} \ln \mathbf{E} \exp \left\{ \frac{1}{\tau} \left\langle \boldsymbol{\lambda}, p_N^M + \boldsymbol{i} q_N^M \right\rangle_{\mathbb{R}} \right\} \\ &= \lim_{N \to \infty} \frac{1}{N} \ln \mathbf{E} \exp \left\{ \sum_{k=1}^M \frac{1}{\tau} \left(\Re \lambda_k p_N^{M,k} + \Im \lambda_k q_N^{M,k} \right) \right\} \end{split}$$

$$= \lim_{N \to \infty} \frac{1}{N} \ln \prod_{k=1}^{M} \mathbf{E} \exp \left\{ \frac{1}{\tau} \left(\Re \lambda_{k} p_{N}^{M,k} + \Im \lambda_{k} q_{N}^{M,k} \right) \right\}$$

$$= \sum_{k=1}^{M} \lim_{N \to \infty} \frac{1}{N} \ln \mathbf{E} \exp \left\{ \frac{1}{\tau} \left(\Re \lambda_{k} p_{N}^{M,k} + \Im \lambda_{k} q_{N}^{M,k} \right) \right\}$$

$$= \sum_{k=1}^{M} \Lambda_{k}(\lambda_{k}), \tag{5.11}$$

where we have used the fact that $\left\{\left(p_n^{M,k},q_n^{M,k}\right)\right\}_{n\in\mathbb{N}},\ k=1,2,\ldots,M,$ are mutually independent stochastic processes as a result of the independence of $\{\beta_k(t)\}_{t\geq 0},\ k=1,2\ldots,M.$

To acquire the expression $\Lambda_k(\cdot)$, we need to give the general formula of $\left\{\left(p_n^{M,k},q_n^{M,k}\right)\right\}_{n\in\mathbb{N}}$ (Hereafter, we always fix some $k\in\{1,2,\ldots,M\}$ without extra statement). It follows from the recurrence formula (5.6) that

$$\begin{pmatrix} p_n^{M,k} \\ q_n^{M,k} \end{pmatrix} = (A(k^2\tau))^n \begin{pmatrix} p_0^{M,k} \\ q_0^{M,k} \end{pmatrix} + \alpha_k \sum_{j=0}^{n-1} (A(k^2\tau))^{n-j-1} B(k^2\tau) \delta \beta_{k,j}, \qquad n = 0, 1, 2 \dots$$

Let $\theta_k \in (0,\pi)$ be the parameter such that

$$\cos(\theta_k) = \frac{\operatorname{tr}(A(k^2\tau))}{2\sqrt{\det(A(k^2\tau))}}, \quad \sin(\theta_k) = \frac{\sqrt{4\det(A(k^2\tau)) - (\operatorname{tr}(A(k^2\tau)))^2}}{2\sqrt{\det(A(k^2\tau))}}.$$
 (5.12)

Then under **Assumption 1**, one has (also see [3, Sect. 3]) that for sufficiently small τ ,

$$(A(k^2\tau))^n = \begin{pmatrix} -\det(A(k^2\tau))\hat{\alpha}_{n-1}^k + a_{11}(k^2\tau)\hat{\alpha}_n^k & a_{12}(k^2\tau)\hat{\alpha}_n^k \\ a_{21}(k^2\tau)\hat{\alpha}_n^k & \hat{\alpha}_{n+1}^k - a_{11}(k^2\tau)\hat{\alpha}_n^k \end{pmatrix},$$

where $\hat{\alpha}_n^k = \left[\det(A(k^2\tau))\right]^{\frac{n-1}{2}} \sin(n\theta_k)/\sin(\theta_k)$. In this way, we obtain the following expression of the general formula of $\left\{\left(p_n^{M,k},q_n^{M,k}\right)\right\}_{n\in\mathbb{N}}$

$$p_n^{M,k} = -\det(A)\hat{\alpha}_{n-1}^k p_0^{M,k} + \hat{\alpha}_n^k \left(a_{11} p_0^{M,k} + a_{12} q_0^{M,k} \right)$$

$$+ \alpha_k \sum_{j=0}^{n-1} \left[-\det(A)\hat{\alpha}_{n-2-j}^k b_1 + (a_{11}b_1 + a_{12}b_2)\hat{\alpha}_{n-1-j}^k \right] \delta \beta_{k,j}$$
(5.13)

and

$$q_n^{M,k} = a_{21} \hat{\alpha}_n^k p_0^{M,k} + \hat{\alpha}_{n+1}^k q_0^{M,k} - a_{11} \hat{\alpha}_n^k q_0^{M,k} + \alpha_k \sum_{j=0}^{n-1} \left[(a_{21}b_1 - a_{11}b_2) \hat{\alpha}_{n-1-j}^k + b_2 \hat{\alpha}_{n-j}^k \right] \delta \beta_{k,j},$$
(5.14)

where det(A), a_{ij} , b_i , i, j = 1, 2, are computed at $k^2\tau$. For convenience, when no confusion occurs, we always omit the argument $k^2\tau$ of det(A), a_{ij} , b_i , i, j = 1, 2.

Since **Assumption 2** holds,

$$\cos(\theta_k) = \frac{\operatorname{tr}(A(k^2\tau))}{2}, \qquad \sin(\theta_k) = \frac{\sqrt{4 - (\operatorname{tr}(A(k^2\tau)))^2}}{2}, \qquad \hat{\alpha}_n^k = \frac{\sin(n\theta_k)}{\sin(\theta_k)}. \tag{5.15}$$

It follows from (5.13), (5.14) and (5.15) that

$$\mathbf{E}p_{N}^{M,k} = -\hat{\alpha}_{N-1}^{k} p_{0}^{M,k} + \hat{\alpha}_{N}^{k} \left(a_{11} p_{0}^{M,k} + a_{12} q_{0}^{M,k} \right)$$

$$= \frac{1}{\sin(\theta_{k})} \left[-\sin((N-1)\theta_{k}) p_{0}^{M,k} + \sin(N\theta_{k}) \left(a_{11} p_{0}^{M,k} + a_{12} q_{0}^{M,k} \right) \right]$$
(5.16)

and

$$\mathbf{E}q_N^{M,k} = \hat{\alpha}_{N+1}^k q_0^{M,k} + \hat{\alpha}_N^k \left(a_{21} p_0^{M,k} - a_{11} q_0^{M,k} \right)$$

$$= \frac{1}{\sin(\theta_k)} \left[\sin((N+1)\theta_k) q_0^{M,k} + \sin(N\theta_k) \left(a_{21} p_0^{M,k} - a_{11} q_0^{M,k} \right) \right]. \tag{5.17}$$

In addition, we obtain

$$\mathbf{Var}(p_N^{M,k}) = \tau \alpha_k^2 \sum_{j=0}^{N-1} \left[-\hat{\alpha}_{N-2-j}^k b_1 + (a_{11}b_1 + a_{12}b_2)\hat{\alpha}_{N-1-j}^k \right]^2$$

$$= \frac{\tau \alpha_k^2}{\sin^2(\theta_k)} \sum_{j=0}^{N-1} \left[b_1^2 \sin^2((j-1)\theta_k) + (a_{11}b_1 + a_{12}b_2)^2 \sin^2(j\theta_k) -2(a_{11}b_1 + a_{12}b_2)b_1 \sin(j\theta_k) \sin((j-1)\theta_k) \right].$$

Using the fact $2\sin(\alpha)\sin(\beta) = \cos(\alpha - \beta) - \cos(\alpha + \beta)$, we have

$$\mathbf{Var}(p_N^{M,k}) = \frac{\tau \alpha_k^2 N}{2\sin^2(\theta_k)} \left[b_1^2 + (a_{11}b_1 + a_{12}b_2)^2 - 2(a_{11}b_1 + a_{12}b_2)b_1\cos(\theta_k) \right] + R_1(k), \tag{5.18}$$

where

$$R_1(k) = \frac{\tau \alpha_k^2}{\sin^2(\theta_k)} \sum_{j=0}^{N-1} \left[-\frac{b_1^2}{2} \cos(2(j-1)\theta_k) - \frac{(a_{11}b_1 + a_{12}b_2)^2}{2} \cos(2j\theta_k) + (a_{11}b_1 + a_{12}b_2)b_1 \cos((2j-1)\theta_k) \right].$$

By the facts $\sum_{n=1}^{N} \cos((2n+1)\theta) = \frac{\sin((2N+2)\theta)-\sin(2\theta)}{2\sin(\theta)}$ and $\sum_{n=1}^{N} \cos((2n)\theta) = \frac{\sin((2N+1)\theta)-\sin(\theta)}{2\sin(\theta)}$, $\left|\sum_{j=0}^{N-1} \cos(2(j-1)\theta_k)\right| + \left|\sum_{j=0}^{N-1} \cos(2j\theta_k)\right| \leq K(\tau, M)$ (Recall that we use the notation $K(\tau, M)$ to denote the constant dependent on τ, M , but independent of N). Hence, we obtain $|R_1| \leq K(\tau, M)$. Similarly, one has

$$\mathbf{Var}(q_N^{M,k}) = \frac{\tau \alpha_k^2 N}{2\sin^2(\theta_k)} \left[b_2^2 + (a_{21}b_1 - a_{11}b_2)^2 + 2(a_{21}b_1 - a_{11}b_2)b_2\cos(\theta_k) \right] + R_2$$
 (5.19)

with $|R_2| \leq K(\tau, M)$, and

$$\mathbf{Cor}(p_N^{M,k}, q_N^{M,k}) = -\frac{\tau \alpha_k^2 N}{2\sin^2(\theta_k)} \left[(a_{21}b_1 - a_{11}b_2)b_1 \cos(\theta_k) + b_1 b_2 \cos(2\theta_k) - (a_{11}b_1 + a_{12}b_2)(a_{21}b_1 - a_{11}b_2) - (a_{11}b_1 + a_{12}b_2)b_2 \cos(\theta_k) \right] + R_3$$
 (5.20)

with $|R_3| \leq K(\tau, M)$. It follows from (5.16) and (5.17) that

$$\left| \mathbf{E} \left\langle \lambda_k, p_N^{M,k} + i q_N^{M,k} \right\rangle_{\mathbb{R}} \right| = \left| \Re \lambda_k \mathbf{E} p_N^{M,k} + \Im \lambda_k \mathbf{E} q_N^{M,k} \right| \le K(\tau, M, \lambda_k).$$
 (5.21)

Further, (5.18), (5.19) and (5.20) give

$$\mathbf{Var}\left\langle \lambda_k, p_N^{M,k} + oldsymbol{i}q_N^{M,k}
ight
angle_{\mathbb{R}}$$

$$= (\Re \lambda_k)^2 \mathbf{Var}(p_N^{M,k}) + (\Im \lambda_k)^2 \mathbf{Var}(q_N^{M,k}) + 2\Re \lambda_k \Im \lambda_k \mathbf{Cor} \left(p_N^{M,k}, q_N^{M,k} \right)$$

$$= \frac{\tau \alpha_k^2 N (\Re \lambda_k)^2}{2 \sin^2(\theta_k)} \left[b_1^2 + (a_{11}b_1 + a_{12}b_2)^2 - 2(a_{11}b_1 + a_{12}b_2)b_1 \cos(\theta_k) \right]$$

$$+ \frac{\tau \alpha_k^2 N (\Im \lambda_k)^2}{2 \sin^2(\theta_k)} \left[b_2^2 + (a_{21}b_1 - a_{11}b_2)^2 + 2(a_{21}b_1 - a_{11}b_2)b_2 \cos(\theta_k) \right]$$

$$- \frac{\Re \lambda_k \Im \lambda_k \tau \alpha_k^2 N}{\sin^2(\theta_k)} \left[(a_{21}b_1 - a_{11}b_2)b_1 \cos(\theta_k) + b_1b_2 \cos(2\theta_k) \right]$$

$$- (a_{11}b_1 + a_{12}b_2)(a_{21}b_1 - a_{11}b_2) - (a_{11}b_1 + a_{12}b_2)b_2 \cos(\theta_k) \right] + R$$

$$(5.22)$$

with $|R| \leq K(\tau, M, \lambda_k)$. Noting $\left\langle \lambda_k, p_N^{M,k} + iq_N^{M,k} \right\rangle_{\mathbb{R}}$ is Gaussian, we have that for each $\lambda_k \in \mathbb{C}$,

$$\Lambda_{k}(\lambda_{k}) = \lim_{N \to \infty} \frac{1}{N} \ln \mathbf{E} \exp \left\{ \frac{1}{\tau} \left\langle \lambda_{k}, p_{N}^{M,k} + iq_{N}^{M,k} \right\rangle_{\mathbb{R}} \right\}
= \lim_{N \to \infty} \frac{1}{N} \left(\frac{1}{\tau} \mathbf{E} \left\langle \lambda_{k}, p_{N}^{M,k} + iq_{N}^{M,k} \right\rangle_{\mathbb{R}} + \frac{1}{2\tau^{2}} \mathbf{Var} \left\langle \lambda_{k}, p_{N}^{M,k} + iq_{N}^{M,k} \right\rangle_{\mathbb{R}} \right)
= \frac{\alpha_{k}^{2} (\Re \lambda_{k})^{2}}{4\tau \sin^{2}(\theta_{k})} \left[b_{1}^{2} + (a_{11}b_{1} + a_{12}b_{2})^{2} - 2(a_{11}b_{1} + a_{12}b_{2})b_{1} \cos(\theta_{k}) \right]
+ \frac{\alpha_{k}^{2} (\Im \lambda_{k})^{2}}{4\tau \sin^{2}(\theta_{k})} \left[b_{2}^{2} + (a_{21}b_{1} - a_{11}b_{2})^{2} + 2(a_{21}b_{1} - a_{11}b_{2})b_{2} \cos(\theta_{k}) \right]
- \frac{\Re \lambda_{k} \Im \lambda_{k} \alpha_{k}^{2}}{2\tau \sin^{2}(\theta_{k})} \left[(a_{21}b_{1} - a_{11}b_{2})b_{1} \cos(\theta_{k}) + b_{1}b_{2}(2\cos(\theta_{k})^{2} - 1)
- (a_{11}b_{1} + a_{12}b_{2})(a_{21}b_{1} - a_{11}b_{2}) - (a_{11}b_{1} + a_{12}b_{2})b_{2} \cos(\theta_{k}) \right].$$
(5.23)

Then, noting that $\operatorname{tr}(A(k^2\tau)) = 2\cos(\theta_k)$, we rewrite (5.23) as

$$\Lambda_k(\lambda_k) = \frac{\alpha_k^2}{4\tau \sin^2(\theta_k)} \left\{ a(k^2\tau)(\Re \lambda_k)^2 + b(k^2\tau)(\Im \lambda_k)^2 - 2c(k^2\tau)\Re \lambda_k \Im \lambda_k \right\}. \tag{5.24}$$

By (5.11), we get the expression (5.10).

In addition, for each $\hat{\lambda}$, $z \in \mathbb{C}^{\hat{M}}$, the Gateaux derivative of $\Lambda^{M,\tau}$ is given by

$$\mathcal{G}\Lambda^{M,\tau}(\boldsymbol{\lambda})(z) = \sum_{k=1}^{M} \frac{\alpha_k^2}{4\tau \sin^2(\theta_k)} \left[2a(k^2\tau)\Re\lambda_k\Re z_k + 2b(k^2\tau)\Im\lambda_k\Im z_k - 2c(k^2\tau)\left(\Re\lambda_k\Im z_k + \Im\lambda_k\Re z_k\right) \right].$$

According to Theorem 3.4, in order to give the LDP of $\{A_N^M\}_{N\in\mathbb{N}}$, it remains to show that $\{A_N^M\}_{N\in\mathbb{N}}$ is exponentially tight. As is mentioned in Section 3, we will use the finiteness of logarithmic moment generating function to derive the exponential tightness. In fact, we have the following lemma

Lemma 5.9. If **Assumptions 1 and 2** hold, then for each fixed $M \in \mathbb{N}$, we have that for all sufficiently small stepsize τ , $\{A_N^M\}_{N\in\mathbb{N}}$ satisfies an LDP with the good rate function $(\Lambda^{M,\tau})^*(z) = \sup_{\boldsymbol{\lambda}\in\mathbb{C}^M} \{\langle \boldsymbol{\lambda},z\rangle_{\mathbb{R}} - \Lambda^{M,\tau}(\boldsymbol{\lambda})\}.$

Proof. It follows from Lemma 5.8 that for each $\lambda \in \mathbb{C}^M$

$$\Lambda^{M,\tau}(\lambda) = \lim_{N \to \infty} \frac{1}{N} \ln \mathbf{E} \exp \left\{ N \left\langle \lambda, A_N^M \right\rangle_{\mathbb{R}} \right\} < \infty.$$
 (5.25)

In particular, we take $\lambda = (0, \dots, 0, 1, 0, \dots, 0)$ in (5.25) with 1 being its kth component. Then we obtain

$$\zeta_{k,1} := \lim_{N \to \infty} \frac{1}{N} \ln \mathbf{E} \exp\left\{ N \Re A_N^{M,k} \right\} < \infty, \tag{5.26}$$

where $A_N^{M,k}$ is the kth argument of A_N^M . Taking $\lambda = (0, \dots, 0, -1, 0, \dots, 0)$ in (5.25) with -1 being its kth component yields

$$\zeta_{k,2} := \lim_{N \to \infty} \frac{1}{N} \ln \mathbf{E} \exp\left\{-N\Re A_N^{M,k}\right\} < \infty.$$
 (5.27)

For each L > 0, using Markov's inequality one has

$$\mathbf{P}\left(\Re A_N^{M,k} > \frac{L}{2M}\right) = \mathbf{P}\left(\exp\left\{N\Re A_N^{M,k}\right\} > \exp\left\{\frac{NL}{2M}\right\}\right) \leq \exp\left\{-\frac{NL}{2M}\right\}\mathbf{E}\exp\left\{N\Re A_N^{M,k}\right\}$$

and

$$\mathbf{P}\left(\Re A_N^{M,k}<-\frac{L}{2M}\right)=\mathbf{P}\left(\exp\left\{-N\Re A_N^{M,k}\right\}>\exp\left\{\frac{NL}{2M}\right\}\right)\leq \exp\left\{-\frac{NL}{2M}\right\}\mathbf{E}\exp\left\{-N\Re A_N^{M,k}\right\}.$$

Hence, (5.26) leads to

$$\limsup_{N \to \infty} \frac{1}{N} \ln \mathbf{P} \left(\Re A_N^{M,k} > \frac{L}{2M} \right) \le -\frac{L}{2M} + \zeta_{k,1},$$

and (5.27) leads to

$$\limsup_{N \to \infty} \frac{1}{N} \ln \mathbf{P} \left(\Re A_N^{M,k} < -\frac{L}{2M} \right) \le -\frac{L}{2M} + \zeta_{k,2}.$$

Combining the above formulas and Proposition 3.7, we have

$$\limsup_{N\to\infty} \frac{1}{N} \ln \mathbf{P} \left(|\Re A_N^{M,k}| > \frac{L}{2M} \right) \le \max \left\{ -\frac{L}{2M} + \zeta_{k,1}, -\frac{L}{2M} + \zeta_{k,2} \right\} = -\frac{L}{2M} + \zeta_k', \qquad (5.28)$$

with $\zeta'_k = \max\{\zeta_{k,1}, \zeta_{k,2}\}$. By taking $\lambda = (0, \dots, 0, i, 0, \dots, 0)$ (resp. $\lambda = (0, \dots, 0, -i, 0, \dots, 0)$) in (5.25) with i (resp. -i) being its kth component, and repeating the above procedure, we have

$$\limsup_{N \to \infty} \frac{1}{N} \ln \mathbf{P} \left(|\Im A_N^{M,k}| > \frac{L}{2M} \right) \le -\frac{L}{2M} + \zeta_k'', \tag{5.29}$$

for some $\zeta_k'' < \infty$.

Further, it holds that for every k = 1, 2, ..., M,

$$\mathbf{P}\left(\left\|A_N^{M,k}\right\| > \frac{L}{M}\right) \leq \mathbf{P}\left(\left|\Re A_N^{M,k}\right| > \frac{L}{2M}\right) + \mathbf{P}\left(\left|\Im A_N^{M,k}\right| > \frac{L}{2M}\right),$$

which together with (5.28), (5.29) and Proposition 3.7 yields

$$\limsup_{N \to \infty} \frac{1}{N} \ln \mathbf{P} \left(\left\| A_N^{M,k} \right\| > \frac{L}{M} \right) \le -\frac{L}{2M} + \zeta_k, \tag{5.30}$$

with $\zeta_k = \max\{\zeta_k', \zeta_k''\}$. For L > 0, define $K_L = \{z \in \mathbb{C}^M : ||z|| \leq L\}$, which is a compact subset of \mathbb{C}^M . Then it holds that

$$\mathbf{P}\left(A_{N}^{M} \in K_{L}^{c}\right) = \mathbf{P}\left(\left\|A_{N}^{M}\right\| > L\right) \leq \mathbf{P}\left(\sum_{k=1}^{M} \left\|A_{N}^{M,k}\right\| > L\right) \leq \mathbf{P}\left(\bigcup_{k=1}^{M} \left\{\left\|A_{N}^{M,k}\right\| > \frac{L}{M}\right\}\right)$$

$$\leq \sum_{k=1}^{M} \mathbf{P}\left(\left\|A_{N}^{M,k}\right\| > \frac{L}{M}\right). \tag{5.31}$$

Substituting (5.30) into (5.31) and using Proposition 3.7, one has

$$\limsup_{N \to \infty} \frac{1}{N} \ln \mathbf{P} \left(A_N^M \in K_L^c \right) \le -\frac{L}{2M} + \max_{k=1,2,\dots,M} \zeta_k.$$

Then, one immediately has

$$\lim_{L \to \infty} \limsup_{N \to \infty} \frac{1}{N} \ln \mathbf{P} \left(A_N^M \in K_L^c \right) = -\infty,$$

which implies the exponential tightness of $\{A_N^M\}_{N\in\mathbb{N}}$. By Lemma 5.8, the exponential tightness of $\{A_N^M\}_{N\in\mathbb{N}}$ and Theorem 3.4, we complete the proof.

Lemma 5.10. Let **Assumptions 1, 2 and 3** hold. For each fixed $M \in \mathbb{N}$ with $\eta_M > 0$, we have that for all sufficiently small stepsize τ ,

$$(\Lambda^{M,\tau})^*(z) = \sum_{k=1}^M \frac{\tau \left(4 - (\operatorname{tr}(A(k^2\tau)))^2\right)}{4 \left[a(k^2\tau)b(k^2\tau) - c^2(k^2\tau)\right] \alpha_k^2} \left[b(k^2\tau)(\Re z_k)^2 + a(k^2\tau)(\Im z_k)^2 + 2c(k^2\tau)\Re z_k\Im z_k\right].$$
(5.32)

Proof. It follows from (5.10) that the Fenchel-Legendre transform of $\Lambda^{M,\tau}$ is

$$(\Lambda^{M,\tau})^*(z) = \sup_{\boldsymbol{\lambda} \in \mathbb{C}^M} \left\{ \langle \boldsymbol{\lambda}, z \rangle_{\mathbb{R}} - \Lambda^{M,\tau}(\boldsymbol{\lambda}) \right\}$$

$$= \sup_{\lambda_1 \in \mathbb{C}} \sup_{\lambda_2 \in \mathbb{C}} \cdots \sup_{\lambda_M \in \mathbb{C}} \left\{ \sum_{k=1}^M \langle \lambda_k, z_k \rangle_{\mathbb{R}} - \Lambda_k(\lambda_k) \right\}$$

$$= \sum_{k=1}^M \sup_{\lambda_k \in \mathbb{C}} \left\{ \langle \lambda_k, z_k \rangle_{\mathbb{R}} - \Lambda_k(\lambda_k) \right\} =: \sum_{k=1}^M \Lambda_k^*(z_k).$$
 (5.33)

According to (5.24),

$$\begin{split} &\Lambda_k^*(z_k) = \sup_{\lambda_k \in \mathbb{C}} \left\{ \langle \lambda_k, z_k \rangle_{\mathbb{R}} - \Lambda_k(\lambda_k) \right\} \\ &= \sup_{(\Re \lambda_k, \Im \lambda_k) \in \mathbb{R}^2} \left\{ \Re \lambda_k \Re z_k + \Im \lambda_k \Im z_k - \frac{\alpha_k^2}{4\tau \sin^2(\theta_k)} \left[a(k^2\tau)(\Re \lambda_k)^2 + b(k^2\tau)(\Im \lambda_k)^2 - 2c(k^2\tau)\Re \lambda_k \Im \lambda_k \right] \right\} \\ &= \sup_{(x,y) \in \mathbb{R}^2} \left\{ (\Re z_k) x + (\Im z_k) y - \frac{\alpha_k^2}{4\tau \sin^2(\theta_k)} \left[a(k^2\tau) x^2 + b(k^2\tau) y^2 - 2c(k^2\tau) xy \right] \right\} \\ &= : \sup_{(x,y) \in \mathbb{R}^2} f_k(x,y). \end{split}$$

Under **Assumption 3**, if τ is sufficiently small, then for each $k = 1, 2, ..., M, x, y \in \mathbb{R}$,

$$\left| \frac{2c(k^2\tau)xy}{a(k^2\tau)x^2 + b(k^2\tau)y^2} \right| < (1-\eta) \frac{2\sqrt{a(k^2\tau)b(k^2\tau)}|xy|}{a(k^2\tau)x^2 + b(k^2\tau)y^2} \le (1-\eta) \frac{a(k^2\tau)x^2 + b(k^2\tau)y^2}{a(k^2\tau)x^2 + b(k^2\tau)y^2} = 1-\eta,$$

which implies $1 - \frac{2c(k^2\tau)xy}{a(k^2\tau)x^2 + b(k^2\tau)y^2} > \eta$ for every $x, y \in \mathbb{R}$. Then, we have

$$\lim_{(x,y)\to\infty} f_k(x,y) = \lim_{(x,y)\to\infty} \left(a(k^2\tau)x^2 + b(k^2\tau)y^2 \right) \left\{ \frac{(\Re z_k)x + (\Im z_k)y}{a(k^2\tau)x^2 + b(k^2\tau)y^2} - \frac{\alpha_k^2}{4\tau \sin^2(\theta_k)} \left[1 - \frac{2c(k^2\tau)xy}{a(k^2\tau)x^2 + b(k^2\tau)y^2} \right] \right\} = -\infty,$$

which along with the continuity of f_k , implies that there exist x_k, y_k satisfying $-\infty < x_k, y_k < +\infty$ such that $\sup_{(x,y)\in\mathbb{R}^2} f_k(x,y) = f_k(x_k,y_k)$. Then, it holds that

$$\frac{\partial f_k(x_k, y_k)}{\partial x} = \Re z_k - \frac{\alpha_k^2}{4\tau \sin^2(\theta_k)} \left[2a(k^2\tau)x_k - 2c(k^2\tau)y_k \right] = 0,$$

$$\frac{\partial f_k(x_k, y_k)}{\partial y} = \Im z_k - \frac{\alpha_k^2}{4\tau \sin^2(\theta_k)} \left[2b(k^2\tau)y_k - 2c(k^2\tau)x_k \right] = 0.$$

For a given $M \in \mathbb{N}$ with $\eta_M > 0$, $\alpha_k = \alpha \sqrt{\eta_k} > 0$, $k = 1, 2, \dots, M$. Then, we obtain

$$x_{k} = \frac{2\tau \sin^{2}(\theta_{k}) \left(\Re z_{k} b(k^{2}\tau) + \Im z_{k} c(k^{2}\tau)\right)}{\left[a(k^{2}\tau)b(k^{2}\tau) - c^{2}(k^{2}\tau)\right] \alpha_{k}^{2}},$$
$$y_{k} = \frac{2\tau \sin^{2}(\theta_{k}) \left(\Im z_{k} a(k^{2}\tau) + \Re z_{k} c(k^{2}\tau)\right)}{\left[a(k^{2}\tau)b(k^{2}\tau) - c^{2}(k^{2}\tau)\right] \alpha_{k}^{2}},$$

which leads to

$$\begin{split} \Lambda_k^*(z_k) = &\Re z_k \frac{2\tau \sin^2(\theta_k) \left(\Re z_k b + \Im z_k c\right)}{(ab - c^2) \,\alpha_k^2} + \Im z_k \frac{2\tau \sin^2(\theta_k) \left(\Im z_k a + \Re z_k c\right)}{(ab - c^2) \,\alpha_k^2} \\ &- \frac{\alpha_k^2}{4\tau \sin^2(\theta_k)} \left[a \frac{4\tau^2 \sin^4(\theta_k) \left(\Re z_k b + \Im z_k c\right)^2}{(ab - c^2)^2 \,\alpha_k^4} + b \frac{4\tau^2 \sin^4(\theta_k) \left(\Im z_k a + \Re z_k c\right)^2}{(ab - c^2)^2 \,\alpha_k^4} \right. \\ &- 2c \frac{4\tau^2 \sin^4(\theta_k) \left(\Re z_k b + \Im z_k c\right) \left(\Im z_k a + \Re z_k c\right)}{(ab - c^2)^2 \,\alpha_k^4} \right] \\ = & \frac{2\tau \sin^2(\theta_k)}{(ab - c^2) \,\alpha_k^2} \left[b(\Re z_k)^2 + 2c\Re z_k \Im z_k + a(\Im z_k)^2 \right] - \frac{\tau \sin^2(\theta_k)}{(ab - c^2)^2 \,\alpha_k^2} \left[a(\Re z_k b + \Im z_k c)^2 + b(\Im z_k a + \Re z_k c)^2 - 2c(\Re z_k b + \Im z_k c) \left(\Im z_k a + \Re z_k c\right) \right]. \end{split}$$

Direct computations give

$$a(\Re z_k b + \Im z_k c)^2 + b(\Im z_k a + \Re z_k c)^2 - 2c(\Re z_k b + \Im z_k c)(\Im z_k a + \Re z_k c)$$

= $(ab - c^2) \left[b(\Re z_k)^2 + 2c\Re z_k \Im z_k + a(\Im z_k)^2 \right].$

In this way, we have

$$\Lambda_k^*(z_k) = \frac{\tau \sin^2(\theta_k)}{[a(k^2\tau)b(k^2\tau) - c^2(k^2\tau)]\alpha_k^2} \left[b(k^2\tau)(\Re z_k)^2 + a(k^2\tau)(\Im z_k)^2 + 2c(k^2\tau)\Re z_k \Im z_k \right]. \tag{5.34}$$

By (5.15), (5.33) and (5.34), we complete the proof.

Now we give the proof of Theorem 5.6.

Proof of Theorem 5.6:

(1) Clearly, F is a continuous mapping from \mathbb{C}^M to H^0 (see (5.2)). By Lemmas 3.5 and 5.9, we deduce that $\{B_N^M\}_{N\in\mathbb{N}}$, with $B_N^M=F(A_N^M)$, satisfies an LDP on H^0 with the good rate function

$$\begin{split} I^{M,\tau}(x) &= (\Lambda^{M,\tau})^*(F^{-1}(x)) \\ &= \begin{cases} \sum_{k=1}^M \frac{\tau \left(4 - (\operatorname{tr}(A(k^2\tau)))^2\right)}{4[a(k^2\tau)b(k^2\tau) - c^2(k^2\tau)]\alpha_k^2} \left[b(k^2\tau)(\Re\left\langle x, e_k\right\rangle_{\mathbb{C}})^2 + a(k^2\tau)(\Im\left\langle x, e_k\right\rangle_{\mathbb{C}})^2 \\ &\qquad \qquad + 2c(k^2\tau)\Re\left\langle x, e_k\right\rangle_{\mathbb{C}}\Im\left\langle x, e_k\right\rangle_{\mathbb{C}}\right], & \text{if } x \in H_M, \\ +\infty, & \text{otherwise.} \end{cases} \end{split}$$

(2) Denote $J_{mod}^{M,\tau}(z) = \frac{(\Lambda^{M,\tau}(z))^*}{\tau}$. Then $I_{mod}^{M,\tau}(x) = J_{mod}^{M,\tau}(F^{-1}(x))$. It follows from **Assumptions 2 and 5** that $a_{12} \sim h$, $a_{21} \sim -h$ and $2 - \operatorname{tr}(A) = 1 + a_{11}a_{22} - a_{12}a_{21} - a_{11} - a_{22} = (a_{11} - 1)(a_{22} - 1) - a_{12}a_{21} \sim h^2$. Hence $4 - (\operatorname{tr}(A))^2 = (2 + \operatorname{tr}(A))(2 - \operatorname{tr}(A)) \sim 4h^2$. In addition, it holds that $a_{11}b_1 + a_{12}b_2 - b_1 = (a_{11} - 1)b_1 + a_{12}b_2 \sim h$. These imply $a \sim h^2$. Further, $a_{21}b_1 - a_{11}b_2 + b_2 = \mathcal{O}(h^2)$, $a_{21}b_1b_2(2 - \operatorname{tr}(A)) = \mathcal{O}(h^4)$, $a_{11}b_2^2(2 - \operatorname{tr}(A)) \sim h^2$, and hence $b \sim h^2$. Similarly, we have $c = \mathcal{O}(h^3)$, which leads to $ab - c^2 \sim h^4$. These implies that under **Assumptions 5**, **Assumptions 3** holds. Accordingly, it follows from (5.32) that for each $z \in \mathbb{C}^M$

$$\lim_{\tau \to 0} J_{mod}^{\tau}(z) = \lim_{\tau \to 0} \sum_{k=1}^{M} \frac{\left(4 - (\operatorname{tr}(A))^{2}\right)}{4 \left[a(k^{2}\tau)b(k^{2}\tau) - c^{2}(k^{2}\tau)\right] \alpha_{k}^{2}} \left[b(k^{2}\tau)(\Re z_{k})^{2} + a(k^{2}\tau)(\Im z_{k})^{2} + 2c(k^{2}\tau)\Re z_{k}\Im z_{k}\right]$$

$$= \sum_{k=1}^{M} \lim_{\tau \to 0} \frac{4(k^{2}\tau)^{4} \left((\Re z_{k})^{2} + (\Im z_{k})^{2}\right) + \mathcal{O}(\tau^{5})}{4(k^{2}\tau)^{4}\alpha_{k}^{2}} = \sum_{k=1}^{M} \frac{\|z_{k}\|^{2}}{\alpha_{k}^{2}}.$$
(5.35)

Hence,

$$\lim_{\tau \to 0} I_{mod}^{M,\tau}(x) = \lim_{\tau \to 0} J_{mod}^{\tau}(F^{-1}(x)) = \begin{cases} \sum_{k=1}^{M} \frac{\|\langle x, e_k \rangle_{\mathbb{C}}\|^2}{\alpha_k^2}, & \text{if } x \in H_M, \\ +\infty, & \text{otherwise.} \end{cases}$$

Note that for each $x \in H_M$,

$$\sum_{k=1}^{M} \frac{\|\langle x, e_{k} \rangle_{\mathbb{C}} \|^{2}}{\alpha_{k}^{2}} = \frac{1}{\alpha^{2}} \sum_{k=1}^{M} \frac{\|\langle x, e_{k} \rangle_{\mathbb{C}} \|^{2}}{\eta_{k}} = \frac{1}{\alpha^{2}} \left\| Q^{-\frac{1}{2}} P_{M} x \right\|_{H^{0}}^{2} = \frac{1}{\alpha^{2}} \left\| Q^{-\frac{1}{2}} x \right\|_{H^{0}}^{2}.$$

In this way, we have

$$\lim_{\tau \to 0} I_{mod}^{M,\tau}(x) = \begin{cases} \frac{1}{\alpha^2} \left\| Q^{-\frac{1}{2}} x \right\|_{H^0}^2, & \text{if } x \in H_M, \\ +\infty, & \text{otherwise.} \end{cases}$$

$$(5.36)$$

Since $\eta_1 \geq \eta_2 \geq \cdots \geq \eta_M > 0$, $Q_M^{\frac{1}{2}}(H^0) = H_M$. Hence I^M becomes

$$I^{M}(x) = \begin{cases} \frac{1}{\alpha^{2}} \left\| Q^{-\frac{1}{2}} x \right\|_{H^{0}}^{2}, & \text{if } x \in H_{M}, \\ +\infty, & \text{otherwise.} \end{cases}$$

By the above formula and (5.36), $\lim_{\tau \to 0} I_{mod}^{M,\tau}(x) = I^M(x)$.

- (3) Case 1: There are finitely many 0 in $\{\eta_k\}_{k\in\mathbb{N}}$. In this case, for each $M \in \mathbb{N}$, $\eta_1 \geq \eta_2 \geq \cdots \geq \eta_M > 0$. Thus, (5.9) and the second case in the proof of Theorem 4.3 yield (5.7).
- Case 2: There are infinitely many 0 in $\{\eta_k\}_{k\in\mathbb{N}}$, i.e., for some $l\in\mathbb{N}$, $\eta_l>\eta_{l+1}=\eta_{l+2}=\cdots=0$. For this case, we take M=l and obtain that $I^M(x)=I(x)$ (see the first case in the proof of Theorem 4.3). Then, it follows from (5.9) that (5.7) holds.
- 5.3. **Proof of Theorem 5.7.** In this part, we consider the LDP of $\{B_N^M\}_{N\in\mathbb{N}}$ for full discretizations of (1.1), based on the spatial spectral Galerkin method (4.1) and temporal non-symplectic methods. Theorem 5.7 indicates that $\{u_n^M\}_{M,n\in\mathbb{N}}$ can not weakly asymptotically preserve the LDP of $\{B_T\}_{T>0}$.

Proof of Theorem 5.7: Recall $\hat{\alpha}_n^k = \left[\det(A(k^2\tau))\right]^{\frac{n-1}{2}} \sin(n\theta_k)/\sin(\theta_k)$. Under **Assumption 4**, for sufficiently small τ , $\left|\hat{\alpha}_n^k\right| \leq R_{k,\tau}^{n-1}/\sin(\theta_k)$ for some constant $R_{k,\tau} < 1$, $k = 1, 2, \ldots, M$. Denote

 $T_{M,\tau} = \max_{k=1,2,...,M} R_{k,\tau}$ and then $T_{M,\tau} < 1$. By (5.13) and (5.14), we have

$$\begin{split} \left| \mathbf{E} p_N^{M,k} \right| &= \left| -\det(A) \hat{\alpha}_{N-1}^k p_0^{M,k} + \hat{\alpha}_N^k \left(a_{11} p_0^{M,k} + a_{12} q_0^{M,k} \right) \right| \\ &\leq \frac{1}{\sin(\theta_k)} \left| p_0^{M,k} \right| \left(T_{M,\tau}^{N-2} + T_{M,\tau}^{N-1} |a_{11}| \right) + \frac{1}{\sin(\theta_k)} \left| q_0^{M,k} \right| |a_{12}| T_{M,\tau}^{N-1} \\ &\leq K(M,\tau). \end{split}$$

Similarly, one has $\left|\mathbf{E}q_N^{M,k}\right| \leq K(M,\tau)$. It follows from (5.13) that

$$\mathbf{Var}(p_N^{M,k}) = \tau \alpha_k^2 \sum_{j=0}^{N-1} \left[-\det(A) \hat{\alpha}_{N-2-j}^k b_1 + (a_{11}b_1 + a_{12}b_2) \hat{\alpha}_{N-1-j}^k \right]^2.$$

Then, Hölder's inequality and the fact $|\hat{\alpha}_n^k| \leq R_{k,\tau}^{n-1}/\sin(\theta_k)$ yield

$$\begin{split} \left| \mathbf{Var}(p_N^{M,k}) \right| &\leq K(M,\tau) \sum_{j=0}^{N-1} \left[\left(\hat{\alpha}_{N-2-j}^k \right)^2 + \left(\hat{\alpha}_{N-1-j}^k \right)^2 \right] \\ &\leq K(M,\tau) \sum_{j=0}^{N-1} \left(T_{M,\tau}^{2(N-2-j)} + T_{M,\tau}^{2(N-1-j)} \right) \\ &= K(M,\tau) \sum_{j=0}^{N-1} \left(T_{M,\tau}^{2j} + T_{M,\tau}^{2(j-1)} \right) \leq K(M,\tau), \end{split}$$

where we use the fact $\sum_{k=0}^{N-1} r^k < \frac{1}{1-r}$ for each $r \in (0,1)$. Analogously, we obtain

$$\left| \mathbf{Var}(p_N^{M,k}) \right| \leq K(M,\tau), \qquad \left| \mathbf{Cor}(p_N^{M,k},q_N^{M,k}) \right| \leq K(M,\tau).$$

Thus, combining the above estimates, we have

$$\left| \mathbf{E} \left\langle \lambda_k, p_N^{M,k} + m{i} q_N^{M,k}
ight
angle_{\mathbb{R}}
ight| + \left| \mathbf{Var} \left\langle \lambda_k, p_N^{M,k} + m{i} q_N^{M,k}
ight
angle_{\mathbb{R}}
ight| < K(M, au,m{\lambda}), \quad k=1,2,\ldots,M.$$

Following the proof of Lemma 5.8, one can show that the logarithmic moment generating function for $\{A_N^M\}_{N\in\mathbb{N}}$ is $\Lambda^{M,\tau}=0$. Then, we conclude that $\{A_N^M\}_{N\in\mathbb{N}}$ satisfies an LDP on \mathbb{C}^M with the good rate function

$$R(z) = \begin{cases} 0, & \text{if } z = 0, \\ +\infty, & \text{otherwise.} \end{cases}$$
 (5.37)

Combining (5.37) and Lemma 3.5, we have that $\{B_N^M\}_{N\in\mathbb{N}}$ satisfies an LDP on H^0 with the good rate function

$$I_{ns}^{M,\tau}(x) = R(F^{-1}(x)) = \begin{cases} 0, & \text{if } x = 0, \\ +\infty, & \text{otherwise.} \end{cases}$$
 (5.38)

It can be verified that (5.7) does not hold.

6. Extension to the case of complex-valued noises

In this part, we study the LDP of $\{B_T\}_{T>0}$ for the stochastic Schrödinger equation (1.1) driven by complex-valued noises. Let W_1 be a U^0 -valued Q_1 -Winner process and W_2 a U^0 -valued Q_2 -Winner process, such that $W_1(t) = \sum_{k=1}^{\infty} Q_1^{\frac{1}{2}} e_k \beta_k^{(1)}(t)$ and $W_2(t) = \sum_{k=1}^{\infty} Q_2^{\frac{1}{2}} e_k \beta_k^{(2)}(t)$. Here Q_1 and Q_2 are two nonnegative symmetric operators on U^0 with finite traces. $\left\{\beta_k^{(1)}(t)\right\}_{t\geq 0}$, $k=1,2,\ldots$ are mutually independent standard Brownian motions, and $\left\{\beta_k^{(2)}(t)\right\}_{t\geq 0}$, $k=1,2,\ldots$ is another family of mutually independent standard Brownian motions. In addition, we assume that $\left\{\beta_k^{(1)}(t)\right\}_{t\geq 0}$ and $\left\{\beta_j^{(2)}(t)\right\}_{t\geq 0}$ mutually independent for all $k,j=1,2,\ldots$ with $k\neq j$. Also assume that for all $k\in\mathbb{N}$, $t>s\geq 0$, $\left(\beta_k^{(1)}(t)-\beta_k^{(1)}(s),\beta_k^{(2)}(t)-\beta_k^{(2)}(s)\right)$ obey the two-dimensional normal distribution with expectation (0,0) and covariance matrix

$$\left(\begin{array}{cc} t-s & \rho(t-s) \\ \rho(t-s) & t-s \end{array}\right),\,$$

for some constant $\rho \in [-1, 1]$. The driving process for stochastic Schrödinger equation (1.1) is $W(t) = W_1(t) + iW_2(t)$.

Let $\mathcal{N}_{W_1}^2(0,T;L_0^2)$ denote the set

$$\left\{\Phi: [0,T]\times\Omega \to \mathcal{L}_2(Q_1^{\frac{1}{2}}(U^0),U^0) \middle| \Phi \text{ is predicable and } \mathbf{E} \int_0^T \left\|\Phi(s)\circ Q_1^{\frac{1}{2}}\right\|_{\mathcal{L}_2(U^0,U^0)}^2 \mathrm{d}s < \infty\right\},$$

and $\mathcal{N}_{W_2}^2(0,T;L_0^2)$ denote the set

$$\left\{\Phi: [0,T]\times\Omega \to \mathcal{L}_2(Q_2^{\frac{1}{2}}(U^0),U^0) \middle| \Phi \text{ is predicable and } \mathbf{E} \int_0^T \left\|\Phi(s)\circ Q_2^{\frac{1}{2}}\right\|_{\mathcal{L}_2(U^0,U^0)}^2 \mathrm{d}s < \infty\right\}.$$

Before giving the LDP of $\{B_T\}_{T>0}$, we first give the following proposition.

Proposition 6.1. Assume that $\Phi_1 \in \mathcal{N}^2_{W_1}(0,T;L_0^2)$, $\Phi_2 \in \mathcal{N}^2_{W_2}(0,T;L_0^2)$. Then the correlation operators

$$V(t,s) = \mathbf{Cor}(\Phi_1 \cdot W_1(t), \Phi_2 \cdot W_2(s)), \qquad t, s \in [0,T]$$

are given by the formula

$$V(t,s) = \rho \mathbf{E} \int_0^{t \wedge s} \Phi_2(r) Q_2^{\frac{1}{2}} Q_1^{\frac{1}{2}} (\Phi_1(r))^* dr.$$

Proof. For simplicity, we take t=s. For each $a,b\in U^0$ and $\sigma>r\geq 0$, it follows from the independence of $\left\{\beta_k^{(1)}(t)\right\}_{t\geq 0}$ and $\left\{\beta_j^{(2)}(t)\right\}_{t\geq 0}$ with $k\neq j$ that

$$\mathbf{E} \langle W_1(\sigma) - W_1(r), a \rangle_{U^0} \langle W_2(\sigma) - W_2(r), b \rangle_{U^0}$$

$$= \mathbf{E} \left(\sum_{k=1}^{\infty} \left(\beta_k^{(1)}(\sigma) - \beta_k^{(1)}(r) \right) \left\langle Q_1^{\frac{1}{2}} e_k, a \right\rangle_{U^0} \right) \left(\sum_{j=1}^{\infty} \left(\beta_j^{(2)}(\sigma) - \beta_j^{(2)}(r) \right) \left\langle Q_2^{\frac{1}{2}} e_j, b \right\rangle_{U^0} \right)$$

$$= \sum_{k=1}^{\infty} \mathbf{E} \left(\beta_k^{(1)}(\sigma) - \beta_k^{(1)}(r) \right) \left(\beta_k^{(2)}(\sigma) - \beta_k^{(2)}(r) \right) \left\langle e_k, Q_1^{\frac{1}{2}} a \right\rangle_{U^0} \left\langle e_k, Q_2^{\frac{1}{2}} b \right\rangle_{U^0}$$

$$= \rho(\sigma - r) \left\langle Q_1^{\frac{1}{2}} a, Q_2^{\frac{1}{2}} b \right\rangle_{U^0} = \rho(\sigma - r) \left\langle Q_2^{\frac{1}{2}} Q_1^{\frac{1}{2}} a, b \right\rangle_{U^0}. \tag{6.1}$$

We first prove that the conclusion hold in the case that both Φ_1 and Φ_2 are elementary processes. For this end, assume that there is a partition $0 = t_0 < t_1 < \cdots < t_N = t$, $N \in \mathbb{N}$, such that

$$\Phi_1(r) = \sum_{n=0}^{N-1} \Phi_1^n \mathbf{1}_{(t_n, t_{n+1}]}(r), \qquad \Phi_2(r) = \sum_{n=0}^{N-1} \Phi_2^n \mathbf{1}_{(t_n, t_{n+1}]}(r),$$

where $\Phi_i^n: \Omega \to \mathcal{L}(U^0, U^0)$ is \mathcal{F}_{t_n} -measurable, and Φ_i^n takes only a finite number of values in $L(U^0, U^0)$, $i = 1, 2, 0 \le n \le N - 1$. Then we have that for each $a, b \in U^0$,

$$\mathbf{E} \left\langle \int_{0}^{t} \Phi_{1}(r) dW_{1}(r), a \right\rangle_{U^{0}} \left\langle \int_{0}^{t} \Phi_{2}(r) dW_{2}(r), b \right\rangle_{U^{0}} \\
= \mathbf{E} \left(\sum_{j=0}^{N-1} \left\langle \Phi_{1}^{j}(W_{1}(t_{j+1}) - W_{1}(t_{j})), a \right\rangle_{U^{0}} \right) \left(\sum_{k=0}^{N-1} \left\langle \Phi_{2}^{k}(W_{2}(t_{k+1}) - W_{2}(t_{k})), b \right\rangle_{U^{0}} \right) \\
= \sum_{j=0}^{N-1} \sum_{k=0}^{N-1} \mathbf{E} \left\langle W_{1}(t_{j+1}) - W_{1}(t_{j}), (\Phi_{1}^{j})^{*} a \right\rangle_{U^{0}} \left\langle W_{2}(t_{k+1}) - W_{2}(t_{k}), (\Phi_{2}^{j})^{*} b \right\rangle_{U^{0}} \\
= : \sum_{j,k=0}^{N-1} \mathbf{E} S_{j,k}. \tag{6.2}$$

If $k \neq j$, we claim $\mathbf{E}S_{j,k} = 0$. For this end, we may assume that k > j without loss of generality. Then $\left\langle W_1(t_{j+1}) - W_1(t_j), (\Phi_1^j)^* a \right\rangle_{U^0}$ and Φ_2^k are \mathcal{F}_{t_k} -measurable. In addition $(W_2(t_{k+1}) - W_2(t_k))$ is \mathcal{F}_{t_k} -independent. It follows from the properties of conditional expectation that

$$\mathbf{E}(S_{j,k}|\mathcal{F}_{t_k})
= \left\langle W_1(t_{j+1}) - W_1(t_j), (\Phi_1^j)^* a \right\rangle_{U^0} \mathbf{E} \left[\left\langle W_2(t_{k+1}) - W_2(t_k), (\Phi_2^j)^* b \right\rangle_{U^0} \middle| \mathcal{F}_{t_k} \right]
= \left\langle W_1(t_{j+1}) - W_1(t_j), (\Phi_1^j)^* a \right\rangle_{U^0} \left(\mathbf{E} \left\langle W_2(t_{k+1}) - W_2(t_k), u \right\rangle_{U^0} \right) \Big|_{u = (\Phi_2^j)^* b}
= 0,$$

which leads to

$$\mathbf{E}S_{i,k} = \mathbf{E}\left(\mathbf{E}(S_{i,k}|\mathcal{F}_{t_k})\right) = 0, \qquad k \neq j. \tag{6.3}$$

Similarly, using (6.1) we obtain

$$\begin{split} &\mathbf{E}(S_{k,k}|\mathcal{F}_{t_k}) \\ &= \left(\mathbf{E} \left\langle W_1(t_{k+1}) - W_1(t_k), u \right\rangle_{U^0} \left\langle W_2(t_{k+1}) - W_2(t_k), v \right\rangle_{U^0} \right) \Big|_{u = (\Phi_1^k)^* a, v = (\Phi_2^j)^* b} \\ &= \rho(t_{k+1} - t_k) \left\langle \left. Q_2^{\frac{1}{2}} Q_1^{\frac{1}{2}} u, v \right\rangle_{U^0} \right|_{u = (\Phi_1^k)^* a, v = (\Phi_2^k)^* b} \\ &= \rho(t_{k+1} - t_k) \left\langle \left. \Phi_2^k Q_2^{\frac{1}{2}} Q_1^{\frac{1}{2}} (\Phi_1^k)^* a, b \right\rangle_{U^0}. \end{split}$$

Hence, it holds that

$$\mathbf{E}S_{k,k} = \rho(t_{k+1} - t_k)\mathbf{E}\left\langle \Phi_2^k Q_2^{\frac{1}{2}} Q_1^{\frac{1}{2}} (\Phi_1^k)^* a, b \right\rangle_{U^0}.$$
 (6.4)

Substituting (6.3) and (6.4) into (6.2) yields

$$\begin{split} \mathbf{E} \left\langle \int_0^t \Phi_1(r) dW_1(r), a \right\rangle_{U^0} \left\langle \int_0^t \Phi_2(r) dW_2(r), b \right\rangle_{U^0} \\ = & \rho \sum_{k=0}^{N-1} (t_{k+1} - t_k) \mathbf{E} \left\langle \Phi_2^k Q_2^{\frac{1}{2}} Q_1^{\frac{1}{2}} (\Phi_1^k)^* a, b \right\rangle_{U^0} \\ = & \rho \mathbf{E} \left\langle \int_0^t \Phi_2(r) Q_2^{\frac{1}{2}} Q_1^{\frac{1}{2}} (\Phi_1(r))^* a, b \right\rangle_{U^0}, \end{split}$$

which proves the conclusion when Φ_i , i = 1, 2, are elementary processes.

If Φ_i , i=1,2, are general processes, one can take elementary process $\Phi_i^{(n)}$ such that

$$\lim_{n \to \infty} \mathbf{E} \int_0^T \left\| \left(\Phi_i^{(n)}(s) - \Phi_i(s) \right) \circ Q_i^{\frac{1}{2}} \right\|_{\mathcal{L}_2(U^0, U^0)}^2 ds = 0, \qquad i = 1, 2.$$

Then by a standard argument of approximation, one can prove that the conclusion holds for any $\Phi_1 \in \mathcal{N}^2_{W_1}(0,T;L^2_0), \ \Phi_2 \in \mathcal{N}^2_{W_2}(0,T;L^2_0)$ (see also the proof of [7, Proposition 4.28]).

Similar to the case of real-valued noises, we assume that $Q_i^{\frac{1}{2}} \in \mathcal{L}_2(U^0, U^1)$, i = 1, 2. Then, we have the following results.

Theorem 6.2. Under the above conditions, $\{B_T\}_{T>0}$ satisfies an LDP on H^0 with the good rate function

$$I(x) = \begin{cases} \frac{1}{\alpha^2} \left\| \widetilde{Q}^{-\frac{1}{2}} x \right\|_{H^0}^2, & if \ x \in \widetilde{Q}^{\frac{1}{2}}(H^0), \\ +\infty, & otherwise, \end{cases}$$

where $\widetilde{Q} = Q_1 + Q_2$.

Proof. This proof is analogous to that of Theorem 3.9. Hence we only give the sketch of proof. The main difference lies in the computation of the variance $\mathbf{Var} \langle u(t), h \rangle_{\mathbb{R}}$. In fact, it holds that

$$u(t) = S(t)u_0 - \alpha \int_0^t \sin((t-s)\Delta)dW_1(s) - \alpha \int_0^t \cos((t-s)\Delta)dW_2(s)$$
$$+ i\alpha \int_0^t \cos((t-s)\Delta)dW_1(s) - i\alpha \int_0^t \sin((t-s)\Delta)dW_2(s).$$

Hence, for each $h \in H^0$.

$$\langle u(t), h \rangle_{\mathbb{R}} = \langle S(t)u_0, h \rangle_{\mathbb{R}} - \alpha \left\langle \int_0^t \sin((t-s)\Delta) dW_1(s), \Re h \right\rangle_{\mathbb{R}} - \alpha \left\langle \int_0^t \cos((t-s)\Delta) dW_2(s), \Re h \right\rangle_{\mathbb{R}} + \alpha \left\langle \int_0^t \cos((t-s)\Delta) dW_1(s), \Im h \right\rangle_{\mathbb{R}} - \alpha \left\langle \int_0^t \sin((t-s)\Delta) dW_2(s), \Im h \right\rangle_{\mathbb{R}}.$$

Using Proposition 6.1, one has

 $\operatorname{Var} \langle u(t), h \rangle_{\mathbb{R}}$

$$\begin{split} &=\alpha^2\left\langle\int_0^t\sin^2((t-s)\Delta)Q_1\mathrm{d}s\Re h,\Re h\right\rangle_{\mathbb{R}} + \alpha^2\left\langle\int_0^t\cos^2((t-s)\Delta)Q_2\mathrm{d}s\Re h,\Re h\right\rangle_{\mathbb{R}} \\ &+\alpha^2\left\langle\int_0^t\cos^2((t-s)\Delta)Q_1\mathrm{d}s\Im h,\Im h\right\rangle_{\mathbb{R}} + \alpha^2\left\langle\int_0^t\sin^2((t-s)\Delta)Q_2\mathrm{d}s\Im h,\Im h\right\rangle_{\mathbb{R}} \\ &+2\alpha^2\rho\left\langle\int_0^t\sin((t-s)\Delta)\cos((t-s)\Delta)Q_2^{\frac{1}{2}}Q_1^{\frac{1}{2}}\mathrm{d}s\Re h,\Re h\right\rangle_{\mathbb{R}} \\ &-2\alpha^2\left\langle\int_0^t\sin((t-s)\Delta)\cos((t-s)\Delta)Q_1\mathrm{d}s\Re h,\Im h\right\rangle_{\mathbb{R}} \\ &+2\alpha^2\rho\left\langle\int_0^t\sin^2((t-s)\Delta)Q_2^{\frac{1}{2}}Q_1^{\frac{1}{2}}\mathrm{d}s\Re h,\Im h\right\rangle_{\mathbb{R}} -2\alpha^2\rho\left\langle\int_0^t\cos^2((t-s)\Delta)Q_2^{\frac{1}{2}}Q_1^{\frac{1}{2}}\mathrm{d}s\Im h,\Re h\right\rangle_{\mathbb{R}} \\ &+2\alpha^2\left\langle\int_0^t\sin((t-s)\Delta)\cos((t-s)\Delta)Q_2\mathrm{d}s\Re h,\Im h\right\rangle_{\mathbb{R}} \\ &+2\alpha^2\left\langle\int_0^t\sin((t-s)\Delta)\cos((t-s)\Delta)Q_2^{\frac{1}{2}}Q_1^{\frac{1}{2}}\mathrm{d}s\Im h,\Im h\right\rangle_{\mathbb{R}} \\ &-2\alpha^2\rho\left\langle\int_0^t\sin((t-s)\Delta)\cos((t-s)\Delta)Q_2^{\frac{1}{2}}Q_1^{\frac{1}{2}}\mathrm{d}s\Im h,\Im h\right\rangle_{\mathbb{R}} \\ &=\frac{t\alpha^2}{2}\left(\left\langle\widetilde{Q}\Re h,\Re h\right\rangle_{\mathbb{R}} + \left\langle\widetilde{Q}\Im h,\Im h\right\rangle_{\mathbb{R}}\right) + \widetilde{R}, \\ &=\frac{t\alpha^2}{2}\left\|\widetilde{Q}_1^{\frac{1}{2}}\lambda\right\|_{H^0}^2 + \widetilde{R}, \end{split}$$

where $|\widetilde{R}| \leq K(Q_1, Q_2, \Delta)$ with $K(Q_1, Q_2, \Delta)$ independent of t. Similar to the proof of Theorem 3.9, we finish the proof by means of the abstract Gärtner–Ellis theorem.

Remark 6.3. In Theorem 6.2, we give the LDP of $\{B_T\}_{T>0}$. Similarly, the LDP for $\{B_N^M\}_{M,N\in\mathbb{N}}$ of numerical method can also be obtained in the case of complex-valued noises.

7. Future work

The calculation of large deviations rate functions is an interesting and important problem. One of the common techniques of approximating the large deviations rate functions is by the Legendre transform of the approximated logarithmic moment generating functions which may be obtained by, e.g., Monte–Carlo methods provided the prior distributions of observables are known ([13]). For a stochastic system, the prior distributions of the considered observables are generally unknown, the approximated logarithmic moment generating functions can be obtained by the combination of numerical discretizations and Monte–Carlo methods. Do all of numerical discretizations work? Theorem 5.6 of this paper shows that the full discretizations $\{u_n^M\}_{M,n\in\mathbb{N}}$, based on the temporal symplectic discretizations and the spatial spectral Galerkin approximation, can weakly asymptotically preserve the LDP of $\{B_T\}_{T>0}$. This result indicates that for an observable associated with a stochastic Hamiltonian partial differential equation, the symplectic discretization is a prior choice. What is the convergence between the rate functions and their numerical approximations? How to combine other techniques, e.g., the adaptive sampling algorithm (see [9]) and multi-level Monte–Carlo methods, to improve the computational efficiency?

REFERENCES

- [1] R. Anton and D. Cohen. Exponential integrators for stochastic Schrödinger equations driven by Itô noise. *J. Comput. Math.*, 36(2):276–309, 2018.
- [2] C. Chen and J. Hong. Symplectic Runge-Kutta semidiscretization for stochastic Schrödinger equation. SIAM J. Numer. Anal., 54(4):2569–2593, 2016.

- [3] C. Chen, J. Hong, D. Jin, and L. Sun. The superiority of stochastic symplectic methods for a linear stochastic oscillator via large deviations principles. arXiv:1906.03451.
- [4] J. Cui, J. Hong, Z. Liu, and W. Zhou. Stochastic symplectic and multi-symplectic methods for nonlinear Schrödinger equation with white noise dispersion. J. Comput. Phys., 342:267–285, 2017.
- [5] J. Cui, J. Hong, Z. Liu, and W. Zhou. Strong convergence rate of splitting schemes for stochastic nonlinear Schrödinger equations. *J. Differential Equations*, 266(9):5625–5663, 2019.
- [6] G. Da Prato. An introduction to infinite-dimensional analysis. Universitext. Springer-Verlag, Berlin, 2006. Revised and extended from the 2001 original by Da Prato.
- [7] G. Da Prato and J. Zabczyk. Stochastic equations in infinite dimensions, volume 152 of Encyclopedia of Mathematics and its Applications. Cambridge University Press, Cambridge, second edition, 2014.
- [8] A. Dembo and O. Zeitouni. Large deviations techniques and applications, volume 38 of Stochastic Modelling and Applied Probability. Springer-Verlag, Berlin, 2010. Corrected reprint of the second (1998) edition.
- [9] G. Ferré and H. Touchette. Adaptive sampling of large deviations. J. Stat. Phys., 172(6):1525–1544, 2018.
- [10] J. Hong and X. Wang. Invariant measures for stochastic nonlinear Schrödinger equations: numerical approximations and symplectic structures, volume 2251 of Lecture Notes in Mathematics. Springer, Singapore, 2019.
- [11] S. Jiang, L. Wang, and J. Hong. Stochastic multi-symplectic integrator for stochastic nonlinear Schrödinger equation. Commun. Comput. Phys., 14(2):393–411, 2013.
- [12] V. Konotop and L. Vázquez. Nonlinear random waves. World Scientific Publishing Co., Inc., River Edge, NJ, 1994.
- [13] C. M. Rohwer, F. Angeletti, and H. Touchette. Convergence of large-deviation estimators. Phys. Rev. E, 92:052104, Nov 2015.
- [14] H. Touchette. The large deviation approach to statistical mechanics. Phys. Rep., 478(1-3):1–69, 2009.
- [15] S. R. S. Varadhan. Large deviations. Ann. Probab., 36(2):397–419, 2008.

ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA; SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 100049, CHINA E-mail address: chenchuchu@lsec.cc.ac.cn

ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA; SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 100049, CHINA *E-mail address*: hjl@lsec.cc.ac.cn

ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA; SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 100049, CHINA *E-mail address*: diancongjin@lsec.cc.ac.cn (Corresponding author)

ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA; SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 100049, CHINA *E-mail address*: liyingsun@lsec.cc.ac.cn