NONZERO-SUM RISK-SENSITIVE CONTINUOUS-TIME STOCHASTIC GAMES WITH ERGODIC COSTS.

MRINAL K. GHOSH, SUBRATA GOLUI, CHANDAN PAL, AND SOMNATH PRADHAN

ABSTRACT. We study nonzero-sum stochastic games for continuous time Markov decision processes on a denumerable state space with risk-sensitive ergodic cost criterion. Transition rates and cost rates are allowed to be unbounded. Under a Lyapunov type stability assumption, we show that the corresponding system of coupled HJB equations admits a solution which leads to the existence of a Nash equilibrium in stationary strategies. We establish this using an approach involving principal eigenvalues associated with the HJB equations. Furthermore, exploiting appropriate stochastic representation of principal eigenfunctions, we completely characterize Nash equilibria in the space of stationary Markov strategies.

Keywords: Nonzero-sum game, risk-sensitive ergodic cost criterion, stationary strategies, coupled HJB equations, Fan's fixed point theorem, Nash equilibrium.

1. INTRODUCTION

We consider a nonzero-sum stochastic game on the infinite time horizon for continuous time Markov decision processes (CTMDPs) on a denumerable state space. The performance evaluation criterion is exponential of integral cost which addresses the decision makers (i.e., players) attitude towards risk. In other words we address the problem of nonzero-sum risk sensitive stochastic games involving continuous time Markov decision processes. In the literature of stochastic games involving continuous time Markov decision processes, one usually studies the integral of the cost [13], [14], [15] which is the so called risk-neutral situation. In the exponential of integral cost, the evaluation criterion is multiplicative as opposed to the additive nature of evaluation criterion in the integral of cost case. This difference makes the risk sensitive case significantly different from its risk neutral counterpart. The study of risk sensitive criterion was first introduced in [3]; see [29] and the references therein. This criterion is studied extensively in the context of MDP both in discrete and continuous times; see, for instance [5], [6], [7], [9], [17], [18], [26], [30], and the references therein. The corresponding results for stochastic (dynamic) games are limited. Notable exceptions are [1], [2], [10]. In discrete time and discrete state space the risk-sensitive zero-sum stochastic games with bounded cost and transition rates have been studied by Basu and Ghosh [2] and nonzero-sum games in [1]. For CTDMPs, zero-sum stochastic games with risk-sensitive costs for bounded cost and bounded transition rates have been studied in [10]. One can

see [12], [28], and the references therein for finite horizon risk-sensitive nonzero-sum games for CTMDPs. Recently risk sensitive continuous time Markov decision processes have been studied in [4], [11], [24], [25]. In this present paper we extend the results of the above four papers to nonzero-sum stochastic games. Using principal eigenvalue approach, under a Lyapunov type stability assumption, we have shown that the corresponding system of coupled HJB equations admits a solution which in turn leads to the existence of Nash equilibrium in stationary strategies. Also, exploiting the stochastic representation of principal eigenfunction we completely characterize all possible Nash equilibria in the space of stationary Markov strategies. The main motivation for studying this kind of games arises from their applications to many interesting problems, such as controlled birth-and-death systems, telecommunication and queueing systems in which the transition and cost rates may be both unbounded.

Our main contribution in this paper is the following. We establish the existence and characterization of Nash equilibria under a blanket Lyapunov type stability assumption. To be more specific, we study ergodic nonzero sum risk-sensitive stochastic games for CTMDPs having the following features: (a) the transition and the cost rates may be unbounded (b) state space is countable (c) at any state of the system the space of admissible actions is compact (d) the strategies may be history dependent. To our knowledge, these results are new in the literature of ergodic non-zero sum risk-sensitive games for CTMDPs. Similar risk-sensitive game problems for discrete time Markov decision processes have been studied under small costs and geometric ergodicity assumption in [2].

The rest of this paper is organized as follows: Section 2 deals with the problem description and preliminaries. The ergodic cost criterion is analyzed in Section 3. Under a Lyapunov type stability assumption(s), we first establish the existence of a solution to the corresponding coupled Hamilton-Jacobi-Bellman (HJB) equations. This in turn leads to the existence of a Nash equilibrium in stationary strategies (see Theorem 3.2). In Section 4, we present an illustrative example.

2. The game model

For the sake of notational simplicity we treat two player game. The N-player game for $N \ge 3$, is analogous. The continuous-time two-person nonzero-sum stochastic game model which consists of the following elements

$$\{S, U_1, U_2, (U_1(i) \subset U_1, U_2(i) \subset U_2, i \in S), \bar{\pi}_{ij}(u_1, u_2), \bar{c}_1(i, u_1, u_2), \bar{c}_2(i, u_1, u_2)\},$$
(2.1)

where each component is described below:

- S, called the state space, is assumed to be the set of all positive integers endowed with the discrete topology, i.e. $S =: \{1, 2, \dots\}$.
- U_1 and U_2 are the action sets for players 1 and 2, respectively. The action spaces U_1 and U_2 are assumed to be Borel spaces with the Borel σ -algebras $\mathcal{B}(U_1)$ and $\mathcal{B}(U_2)$, respectively.
- For each $i \in S$, $U_1(i) \in \mathcal{B}(U_1)$ and $U_2(i) \in \mathcal{B}(U_2)$ denote the sets of admissible actions for players 1 and 2 in state *i*, respectively. Let $K := \{(i, u_1, u_2) | i \in S, u_1 \in U_1(i), u_2 \in U_2(i)\}$, which is a Borel subset of $S \times U_1 \times U_2$. Throughout this paper, we assume that

(A1)(a) For each $i \in S$, the admissible action spaces $U_k(i), k = 1, 2$, are nonempty and compact subsets of U_k .

The transition rates \$\bar{\pi}_{ij}(u_1, u_2)\$, \$(u_1, u_2) \in U_1(i) \times U_2(i)\$, \$i, j \in S\$, satisfy the condition \$\bar{\pi}_{ij}(u_1, u_2)\$ ≥ 0 for all \$i \neq j\$, \$(u_1, u_2) \in U_1(i) \times U_2(i)\$. Also, we assume that:
(A1)(b) The transition rates \$\bar{\pi}_{ij}(u_1, u_2)\$ are conservative, i.e.,

$$\sum_{j \in S} \bar{\pi}_{ij}(u_1, u_2) = 0 \text{ for } i \in S \text{ and } (u_1, u_2) \in U_1(i) \times U_2(i)$$

and

$$\bar{\pi}_i := \sup_{(u_1, u_2) \in U_1(i) \times U_2(i)} [-\bar{\pi}_{ii}(u_1, u_2)] < \infty.$$

• Finally, the measurable function $\bar{c}_k : K \to \mathbb{R}_+$ denotes the cost rate function for player k, k = 1, 2.

We consider a continuous time Markov decision processes (CTMDPs) $\{Y(t)\}_{t\geq 0}$ with state space S and controlled rate matrix $\Pi_{u_1,u_2} = (\bar{\pi}_{ij}(u_1, u_2))$. To construct the underlying CTMDPs Y(t) (as in [[19], [22], [27]) we introduce some notations: let $S_{\Delta} := S \cup \{\Delta\}$ (with some $\Delta \notin S$), $\Omega_0 := (S \times (0, \infty))^{\infty}$, $\Omega_m := (S \times (0, \infty))^m \times S \times (\{\infty\} \times \{\Delta\})^{\infty}$ for $m \geq 1$ and $\Omega := \bigcup_{m=0}^{\infty} \Omega_m$. Let \mathcal{F} be the Borel σ -algebra on Ω . Then we obtain the measurable space (Ω, \mathcal{F}) . For some $m \geq 1$, and sample $\omega := (i_0, \theta_1, i_1, \cdots, \theta_m, i_m, \cdots) \in \Omega$, define

$$T_0(\omega) := 0, \ T_n(\omega) := T_{n-1}(\omega) + \theta_n, \ T_{\infty}(\omega) := \lim_{n \to \infty} T_n(\omega).$$

Using $\{T_m\}$, we define the state process $\{Y(t)\}_{t\geq 0}$ as

$$Y(t) := \sum_{m \ge 0} I_{\{T_m \le t < T_{m+1}\}} i_m + I_{\{t \ge T_\infty\}} \Delta, \text{ for } t \ge 0 \text{ (with } T_0 := 0).$$
(2.2)

Here, I_E denotes the indicator function of a set E, and we use the convention that 0+z =: zand 0z =: 0 for all $z \in S_{\Delta}$. Obviously, Y(t) is right-continuous on $[0, \infty)$. From (2.2), we see that $T_m(\omega)$ $(m \ge 1)$ denotes the *m*-th jump moment of $\{Y(t)\}_{t\ge 0}$ and i_{m-1} is the state of the process on $[T_{m-1}(\omega), T_m(\omega))$, $\theta_m(\omega) = T_m(\omega) - T_{m-1}(\omega)$ plays the role of sojourn time at state i_{m-1} , and the sample path $\{Y(t)(\omega)\}_{t\ge 0}$ has at most denumerable states $i_m(m = 0, 1, \cdots)$. The process after T_{∞} is regarded to be absorbed in the state Δ . Thus, let $q(\cdot | \Delta, u_1^{\Delta}, u_2^{\Delta}) :\equiv 0, U_1^{\Delta} := U_1 \cup \{u_1^{\Delta}\}, U_2^{\Delta} := U_2 \cup \{u_2^{\Delta}\}, U_1(\Delta) := \{u_1^{\Delta}\}, U_2(\Delta) := \{u_2^{\Delta}\}$. Also, assume that $\bar{c}_k(\Delta, u_1, u_2) :\equiv 0$ (\bar{c}_k is the running cost function for kth player) for all $(u_1, u_2) \in U_1^{\Delta} \times U_2^{\Delta}$, where $u_1^{\Delta}, u_2^{\Delta}$ are isolated points. Moreover, let $\mathcal{F}_t := \sigma(\{T_m \leq s, Y(T_m) \in S\} : 0 \leq s \leq t, m \geq 0)$ for all $t \geq 0, \mathcal{F}_{s-} =: \bigvee_{t < s} \mathcal{F}_t$, and $\tilde{\mathcal{F}} := \sigma(A \times \{0\}, B \times (s, \infty) : A \in \mathcal{F}_0, B \in \mathcal{F}_{s-})$ which denotes the σ -algebra of predictable sets on $\Omega \times [0, \infty)$ related to $\{\mathcal{F}_t\}_{t \geq 0}$.

To complete the specification of a risk-sensitive stochastic game problem, we need, of course, to introduce an optimality criterion. This requires to define the class of strategies as below.

Definition 2.1. A admissible strategy for player 1, denoted by $v_1 = \{v_1(t)\}_{t\geq 0}$, is a transition probability $v_1(du_1|\omega, t)$ from $(\Omega \times [0, \infty), \tilde{\mathcal{F}})$ onto $(U_1^{\Delta}, \mathcal{B}(U_1^{\Delta}))$, such that $v_1(U_1(Y(t-)(\omega))|\omega, t) =$ 1. The set of all admissible strategies for player 1 is denoted by \mathcal{A}_1 . A strategy $v_1 \in$ \mathcal{A}_1 , is called a Markov for player 1 if $v_1(t)(\omega) = v_1(t, Y(t-)(w))$, i.e., $v_1(du_1|\omega, t) =$ $v_1(du_1|Y(t-)(w), t)$ for every $w \in \Omega$ and $t \geq 0$, where $Y(t-)(w) := \lim_{s\uparrow t} Y(s)(w)$. We denote by \mathcal{M}_1 the family of all Markov strategies for player 1. If the Markov strategy v_1 for player 1 does not have any explicit time dependency then it is called a stationary Markov strategy. The set of such strategies for player 1 is denoted by \mathcal{S}_1 . The sets of all admissible strategies \mathcal{A}_2 , all Markov strategies \mathcal{M}_2 and all stationary strategies \mathcal{S}_2 for player 2 are defined similarly.

To avoid the explosion of the state process $\{Y(t)\}_{t\geq 0}$, we need the following assumption imposed on the transition rates, which had been widely used in CTMDPs; see, for instance, [[17], [18], [19], [20]] and references therein.

Assumption 2.1. There exists a Lyapunov function $\tilde{W}: S \to [1, \infty)$ such that

- (i) $\sum_{j \in S} \tilde{W}(j) \overline{\pi}_{ij}(u_1, u_2) \leq C_1 \tilde{W}(i) + C_2 \text{ for all } (u_1, u_2) \in U_1(i) \times U_2(i) \text{ and } i \in S \text{ with some constants } C_1 \neq 0, C_2 \geq 0;$
- (ii) $\bar{\pi}_i \leq C_3 \tilde{W}(i)$ for all $i \in S$ with some positive constant C_3 .

For the rest of this article we are going to assume that Assumption 2.1 holds. Note that if $\sup_{i \in S} \bar{\pi}_i < \infty$ then Assumption 2.1 holds. In this case we can choose \tilde{W} to be a suitable constant. Also note that under Assumption 2.1, for any initial state $i \in S$ and any pair of strategies $(v_1, v_2) \in \mathcal{A}_1 \times \mathcal{A}_2$, Theorem 4.27 in [23] yields the existence of a unique probability measure denoted by $P_i^{v_1, v_2}$ on (Ω, \mathcal{F}) . Let $E_i^{v_1, v_2}$ be the expectation operator with respect to $P_i^{v_1, v_2}$. Also, from [[16], pp.13-15], we know that $\{Y(t)\}_{t\geq 0}$ is a Markov process under any $(v_1, v_2) \in \mathcal{M}_1 \times \mathcal{M}_2$ (in fact, strong Markov).

For any compact metric space A, let $\mathcal{P}(A)$ denote the space of probability measures on A with Prohorov topology. Let $V_k = \mathcal{P}(U_k)$ and $V_k(i) = \mathcal{P}(U_k(i))$ for $i \in S$ and k = 1, 2.

For each $i, j \in S$, $k = 1, 2, v_1 \in V_1(i)$ and $v_2 \in V_2(i)$, the associated transition and cost rates are defined, respectively, as follows:

$$\pi_{ij}(v_1, v_2) := \int_{U_1(i)} \int_{U_2(i)} \bar{\pi}_{ij}(u_1, u_2) v_1(du_1) v_2(du_2),$$
$$c_k(v_1, v_2) := \int_{U_1(i)} \int_{U_2(i)} \bar{c}_k(u_1, u_2) v_1(du_1) v_2(du_2).$$

Note that for $k = 1, 2, v_k \in S_k$ can be identified with a map $v_k : S \to V_k$ such that for each $j \in S, v_k(j) \in V_k(j)$ for each $j \in S$. The sets S_1 and S_2 are endowed with product topology.

We list the commonly used notations below.

- For any finite set $\mathcal{D} \subset S$, we define $\mathcal{B}_{\mathcal{D}} = \{f : S \to \mathbb{R} \mid f \text{ is borel measurable function and } f(i) = 0 \quad \forall i \in \mathcal{D}^c\}.$
- Given any real-valued function $\mathcal{V} \geq 1$ on S, we define a Banach space $(L^{\infty}_{\mathcal{V}}, \|\cdot\|^{\infty}_{\mathcal{V}})$ of \mathcal{V} -weighted functions by

$$L_{\mathcal{V}}^{\infty} = \left\{ u: S \to \mathbb{R} \mid \|u\|_{\mathcal{V}}^{\infty} := \sup_{i \in S} \frac{|u(i)|}{\mathcal{V}(i)} < \infty \right\}.$$

• $L_{\mathcal{V}}^{1,\infty}$ denotes the subset of $L_{\mathcal{V}}^{\infty}$ consists of function u such that $\|u\|_{\mathcal{V}}^{\infty} \leq 1$.

For k = 1, 2, let $\bar{c}_k : S \times U_1 \times U_2 \to [0, \infty)$ be the running cost function for the *k*th player, i.e., when state of the system is *i* and the actions (u_1, u_2) are chosen by the players, then the cost incurred by the *k*th player is $\bar{c}_k(i, u_1, u_2)$. By choosing appropriate strategies, each player wants to minimize his/her accumulated cost over infinite time horizon.

For a pair of admissible strategies (v_1, v_2) , the risk-sensitive ergodic cost for player k is given by

$$\rho_k^{v_1, v_2}(i) := \limsup_{T \to \infty} \frac{1}{T} \ln E_i^{v_1, v_2} \left[e^{\int_0^T c_k(Y(t), v_1(t), v_2(t)) dt} \right],$$
(2.3)

where Y(t) is the CTMDP corresponding to $(v_1, v_2) \in \mathcal{A}_1 \times \mathcal{A}_2$ and $E_i^{v_1, v_2}$ denotes the expectation with respect to the law of the process Y(t) with initial condition Y(0) = i.

Since we are allowing our transition and cost rates to be unbounded, to guarantee the finiteness of $\rho_k^{v_1,v_2}$ for k = 1, 2, we need the following Assumption.

Assumption 2.2. We assume that the CTMDP $\{Y(t)\}_{t\geq 0}$ is irreducible under every pair of stationary Markov strategies $(v_1, v_2) \in S_1 \times S_2$. Furthermore, suppose there exist a constant $C_4 > 0$ and a Lyapunov function $W: S \to [1, \infty)$ such that one of the following hold.

(a) When the running cost is bounded: For some positive constant $\gamma > \max\{\|c_1\|_{\infty}, \|c_2\|_{\infty}\}$ and a finite set \mathcal{K} it holds that

$$\sup_{(u_1,u_2)\in U_1(i)\times U_2(i)}\sum_{j\in S} W(j)\overline{\pi}_{ij}(u_1,u_2) \le C_4 I_{\mathcal{K}}(i) - \gamma W(i) \ \forall i\in S.$$

(b) When the running cost is unbounded: For some norm-like function $\ell: S \to \mathbb{R}_+$ and a finite set \mathcal{K} it holds that

$$\sup_{(u_1,u_2)\in U_1(i)\times U_2(i)}\sum_{j\in S} W(j)\overline{\pi}_{ij}(u_1,u_2) \le C_4 I_{\mathcal{K}}(i) - \ell(i)W(i) \ \forall i\in S.$$

Also, the functions $\ell(\cdot) - \max_{(u_1, u_2) \in U_1(\cdot) \times U_2(\cdot)} c_k(\cdot, u_1, u_2)$, k = 1, 2, are norm-like.

Definition 2.2. A pair of strategies $(v_1^*, v_2^*) \in A_1 \times A_2$ is called a Nash equilibrium if

$$\rho_1^{v_1^*, v_2^*}(i) \leq \rho_1^{v_1, v_2^*}(i) \text{ for all } v_1 \in \mathcal{A}_1 \text{ and } i \in S$$

and

$$\rho_2^{v_1^*, v_2^*}(i) \leq \rho_2^{v_1^*, v_2}(i) \text{ for all } v_2 \in \mathcal{A}_2 \text{ and } i \in S.$$

We wish to establish the existence of a Nash equilibrium in stationary strategies. To ensure the existence of a Nash equilibrium, we assume the following:

- Assumption 2.3. (i) For any fixed $i, j \in S, k=1,2, \overline{\pi}_{ij}(u_1, u_2)$ and $\overline{c}_k(i, u_1, u_2)$ are continuous in $(u_1, u_2) \in U_1(i) \times U_2(i)$.
 - (ii) $\sum_{j \in S} W(j) \overline{\pi}_{ij}(u_1, u_2)$ is continuous in $(u_1, u_2) \in U_1(i) \times U_2(i)$ for any given $i \in S$, where W is as Assumption 2.2.
 - (iii) There exists $i_0 \in S$ such that $\overline{\pi}_{i_0j}(u_1, u_2) > 0$ for all $j \neq i_0$ and $(u_1, u_2) \in U_1(j) \times U_2(j)$.

We now proceed to establish the existence of a Nash equilibrium in stationary strategies. To this end we first outline a procedure for establishing the existence of a Nash equilibrium. Suppose player 2 announces that he is going to employ a strategy $v_2 \in S_2$. In such a scenario, player 1 attempts to minimize

$$\rho_1^{v_1,v_2}(i) = \limsup_{T \to \infty} \frac{1}{T} \ln E_i^{v_1,v_2} \left[e^{\int_0^T c_1(Y(t),v_1(t),v_2(Y(t-)))dt} \right],$$

over $v_1 \in \mathcal{A}_1$. Thus for player 1 it is a continuous time Markov decision problem (CTMDP) with risk sensitive ergodic cost. This problem has been studied in [4], [11], [24], [25]. In particular under certain assumptions, it is shown in [4], [24], [25], that the following Hamilton-Jacobi-Bellman (HJB) equation

$$\begin{cases} \rho_1 \ \hat{\psi}_1(i) = \inf_{v_1 \in V_1(i)} \left[\sum_{j \in S} \pi_{ij}(v_1, v_2(i)) \hat{\psi}_1(j) + c_1(i, v_1, v_2(i)) \hat{\psi}_1(i) \right] \\ \hat{\psi}_1(i_0) = 1, \end{cases}$$

has a suitable solution $(\rho_1, \hat{\psi}_1)$, where ρ_1 is a scalar and $\hat{\psi}_1 : S \to \mathbb{R}$ has suitable growth rate; i_0 is a fixed element of S. Furthermore it is shown in [4], [24], [25] that

$$\rho_1 = \inf_{v_1 \in \mathcal{A}_1} \limsup_{T \to \infty} \frac{1}{T} \ln E_i^{v_1, v_2} \left[e^{\int_0^T c_1(Y(t), v_1(t), v_2(Y(t-))) dt} \right],$$

and if $v_1^* \in \mathcal{S}_1$ is such that for $i \in S$

$$\inf_{v_1 \in V_1(i)} \left[\sum_{j \in S} \pi_{ij}(v_1, v_2(i)) \hat{\psi}_1(j) + c_1(i, v_1, v_2(i)) \hat{\psi}_1(i) \right]$$

=
$$\sum_{j \in S} \pi_{ij}(v_1^*(i), v_2(i)) \hat{\psi}_1(j) + c_1(i, v_1^*(i), v_2(i)) \hat{\psi}_1(i),$$

then $v_1^* \in S_1$ is an optimal control for player 1, i.e., for any $i \in S$

$$\rho_1 = \limsup_{T \to \infty} \frac{1}{T} \ln E_i^{v_1^*, v_2} \left[e^{\int_0^T c_1(Y(t), v_1^*(Y(t-)), v_2(Y(t-))) dt} \right].$$

In other words, given that player 2 is using the strategy $v_2 \in S_2$, $v_1^* \in S_1$ is an optimal response for player 1. Clearly v_1^* depends on v_2 and moreover there may be several optimal responses for player 1 in S_1 . Analogous results holds for player 2 if player 1 announces that he is going to use a strategy $v_1 \in S_1$. Hence given a pair of strategies $(v_1, v_2) \in S_1 \times S_2$, we can find a set of pairs of optimal responses $\{(v_1^*, v_2^*) \in S_1 \times S_2\}$ via the appropriate pair of HJB equations described above. This defines a set-valued map. Clearly any fixed point of this set-valued map is a Nash equilibrium.

The above discussion leads to the following procedure for finding a pair of Nash equilibrium strategies. Suppose that there exist a pair of stationary strategies $(v_1^*, v_2^*) \in S_1 \times S_2$, a pair of scalars (ρ_1^*, ρ_2^*) and a pair of functions $(\hat{\psi}_1^*, \hat{\psi}_2^*)$ with appropriate growth conditions, satisfying the following coupled HJB equations:

$$\begin{cases} \rho_1^* \ \hat{\psi}_1^*(i) = \inf_{v_1 \in V_1(i)} \left[\sum_{j \in S} \pi_{ij}(v_1, v_2^*(i)) \hat{\psi}_1^*(j) + c_1(i, v_1, v_2^*(i)) \hat{\psi}_1^*(i) \right] \\ = \sum_{j \in S} \pi_{ij}(v_1^*(i), v_2^*(i)) \hat{\psi}_1^*(j) + c_1(i, v_1^*(i), v_2^*(i)) \hat{\psi}_1^*(i) \\ \hat{\psi}_1^*(i_0) = 1, \\ \rho_2^* \ \hat{\psi}_2^*(i) = \inf_{v_2 \in V_2(i)} \left[\sum_{j \in S} \pi_{ij}(v_1^*(i), v_2) \hat{\psi}_2^*(j) + c_2(i, v_1^*(i), v_2) \hat{\psi}_2^*(i) \right] \\ = \sum_{j \in S} \pi_{ij}(v_1^*(i), v_2^*(i)) \hat{\psi}_2^*(j) + c_2(i, v_1^*(i), v_2^*(i)) \hat{\psi}_2^*(i) \\ \hat{\psi}_2^*(i_0) = 1, \end{cases}$$

where as before $i_0 \in S$ is a fixed point. Then it can be shown that (v_1^*, v_2^*) is a pair of Nash equilibrium and (ρ_1^*, ρ_2^*) is the pair of corresponding Nash values. Thus the main result of our paper is to establish that the above coupled HJB equations has suitable solutions.

Remark 2.1. Note that the similar stochastic optimal control problem has been studied in [11], [25] for bounded cost and bounded transition rates. But in our game model transition and cost rates are unbounded. Analogous MDP problems are treated in [4].

3. COUPLED HJB EQUATIONS AND EXISTENCE OF NASH EQUILIBRIUM

By the definition of weak convergence of probability measures, one can easily get the following result, which will be crucial for the existence of Nash equilibrium; we omit the details.

Lemma 3.1. Under Assumptions 2.1, 2.2, and 2.3, the functions

$$c_k(i, v_1, v_2), \ k = 1, 2 \ and \ \sum_{j \in S} \pi_{ij}(v_1, v_2)\phi(j)$$

are continuous on $V_1(i) \times V_2(i)$ for each fixed $\phi \in L^{\infty}_W$ and $i \in S$.

Let $\mathcal{D}_n \subset S$ be an increasing sequence of finite sets such that $\bigcup_n \mathcal{D}_n = S$ and $i_0 \in \mathcal{D}_n$ for each $n \ge 1$. In the next lemma we show the existence of eigenpairs to certain equations in \mathcal{D}_n for each $n \in \mathbb{N}$.

Lemma 3.2. Grant Assumptions 2.1, 2.2, and 2.3. Then for each $n \in \mathbb{N}$, the following hold.

(1) For $\hat{v}_2 \in S_2$, there exists an eigenpair $(\rho_{1,n}, \psi_{1,n}) \in \mathbb{R} \times \mathcal{B}^+_{\mathcal{D}_n}$, satisfying

$$\begin{cases} \rho_{1,n}\psi_{1,n}(i) = \inf_{v_1 \in V_1(i)} \left[\sum_{j \in S} \psi_{1,n}(j)\pi_{ij}(v_1, \hat{v}_2(i)) + c_1(i, v_1, \hat{v}_2(i))\psi_{1,n}(i) \right] \text{ for } i \in \mathcal{D}_n, \\ \psi_{1,n}(i_0) = 1. \end{cases}$$

$$(3.1)$$

Moreover, we have

$$0 \le \liminf_{n \to \infty} \rho_{1,n} \le \limsup_{n \to \infty} \rho_{1,n} \le \inf_{v_1 \in \mathcal{A}_1} \limsup_{T \to \infty} \frac{1}{T} \ln E_{i_0}^{v_1, \hat{v}_2} \Big[e^{\int_0^T c_1(Y(t), v_1(t), \hat{v}_2(Y(t-))) dt} \Big].$$
(3.2)

(2) Similarly, for $\hat{v}_1 \in S_1$, there exists an eigenpair $(\rho_{2,n}, \psi_{2,n}) \in \mathbb{R} \times \mathcal{B}_{\mathcal{D}_n}^+$, satisfying

$$\begin{cases} \rho_{2,n}\psi_{2,n}(i) = \inf_{v_2 \in V_2(i)} \left[\sum_{j \in S} \psi_{2,n}(j)\pi_{ij}(\hat{v}_1(i), v_2) + c_2(i, \hat{v}_1(i), v_2)\psi_{2,n}(i) \right] \text{ for } i \in \mathcal{D}_n, \\ \psi_{2,n}(i_0) = 1. \end{cases}$$

$$(3.3)$$

Moreover, we have

$$0 \le \liminf_{n \to \infty} \rho_{2,n} \le \limsup_{n \to \infty} \rho_{2,n} \le \inf_{v_2 \in \mathcal{A}_2} \limsup_{T \to \infty} \frac{1}{T} \ln E_{i_0}^{\hat{v}_1, v_2} \Big[e^{\int_0^T c_2(Y(t), \hat{v}_1(Y(t-)), v_2(t)) dt} \Big].$$
(3.4)

Proof. Follows by analogous arguments as in [4, Lemma 3.1, Lemma 3.3]. We omit the details. \Box

Next by taking limit $n \to \infty$ in the equations we show that the limiting equations admit eigenpairs in appropriate spaces. In particular, we have the following theorem.

Theorem 3.1. Grant Assumptions 2.1, 2.2, and 2.3. Then the following hold.

(1) For $\hat{v}_2 \in S_2$, there exists a unique minimal eigenpair $(\rho_1, \psi_1) \in \mathbb{R}_+ \times L_W^{1,\infty}, \psi_1 > 0$, satisfying

$$\begin{cases} \rho_1 \psi_1(i) = \inf_{v_1 \in V_1(i)} \left[\sum_{j \in S} \psi_1(j) \pi_{ij}(v_1, \hat{v}_2(i)) + c_1(i, v_1, \hat{v}_2(i)) \psi_1(i) \right] \text{ for } i \in S, \\ \psi_1(i_0) = 1. \end{cases}$$

$$(3.5)$$

Moreover, we have

$$\rho_1 = \inf_{v_1 \in \mathcal{A}_1} \limsup_{T \to \infty} \frac{1}{T} \ln E_i^{v_1, \hat{v}_2} \Big[e^{\int_0^T c_1(Y(t), v_1(t), \hat{v}_2(Y(t-))) dt} \Big] (:= \rho_1^{\hat{v}_2} = \inf_{v_1 \in \mathcal{A}_1} \rho_1^{v_1, \hat{v}_2}), \quad (3.6)$$

and there exists a finite set $\mathbb{B}_1 \supset \mathbb{K}$, such that

$$\psi_{1}(i) = \inf_{v_{1} \in \mathcal{S}_{1}} E_{i}^{v_{1},\hat{v}_{2}} \left[e^{\int_{0}^{\hat{\tau}(\mathcal{B}_{1})} (c_{1}(Y(t),v_{1}(Y(t-)),\hat{v}_{2}(Y(t-))) - \rho_{1})dt} \psi_{1}(Y(\hat{\tau}(\mathcal{B}_{1}))) \right] (:=\psi_{1}^{\hat{v}_{2}}(i)) \ \forall i \in \mathcal{B}_{1}^{c},$$

$$(3.7)$$

where $\hat{\tau}(\mathcal{B}_1) = \tau(\mathcal{B}_1^c) = \inf\{t : Y(t) \in \mathcal{B}_1\} =: \tilde{\tau}_1.$

(2) Similarly, for $\hat{v}_1 \in S_1$, there exists a unique minimal eigenpair $(\rho_2, \psi_2) \in \mathbb{R}_+ \times L^{1,\infty}_W$, $\psi_2 > 0$ satisfying

$$\begin{cases} \rho_2 \psi_2(i) = \inf_{v_2 \in V_2(i)} \left[\sum_{j \in S} \psi_2(j) \pi_{ij}(\hat{v}_1(i), v_2) + c_2(i, \hat{v}_1(i), v_2) \psi_2(i) \right] \text{ for } i \in S, \\ \psi_2(i_0) = 1. \end{cases}$$

$$(3.8)$$

Moreover, we have

$$\rho_2 = \inf_{v_2 \in \mathcal{A}_2} \limsup_{T \to \infty} \frac{1}{T} \ln E_i^{\hat{v}_1, v_2} \Big[e^{\int_0^T c_2(Y(t), \hat{v}_1(Y(t-)), v_2(t)) dt} \Big] (:= \rho_2^{\hat{v}_1} = \inf_{v_2 \in \mathcal{A}_2} \rho_2^{\hat{v}_1, v_2}), \quad (3.9)$$

and there exists a finite set $\mathbb{B}_2 \supset \mathbb{K}$, such that

$$\psi_{2}(i) = \inf_{v_{2} \in \mathcal{S}_{2}} E_{i}^{\hat{v}_{1},v_{2}} \left[e^{\int_{0}^{\hat{\tau}(\mathcal{B}_{2})} (c_{2}(Y(t),\hat{v}_{1}(Y(t-)),v_{2}(Y(t-))) - \rho_{2})dt} \psi_{2}(Y(\hat{\tau}(\mathcal{B}_{2}))) \right] (:= \psi_{2}^{\hat{v}_{1}}(i)) \ \forall i \in \mathcal{B}_{2}^{c},$$

$$(3.10)$$

where
$$\hat{\tau}(\mathcal{B}_2) = \tau(\mathcal{B}_2^c) = \inf\{t : Y(t) \in \mathcal{B}_2\} =: \tilde{\tau}_2.$$

Proof. Since $c_1 \geq 0$, using Assumption 2.2, we deduce that there exists a finite set \mathcal{B}_1 containing \mathcal{K} such that

• Under Assumption 2.2 (a)

 $(\sup_{(u_1,u_2)\in U_1(i)\times U_2(i)}c_1(i,u_1,u_2)-\rho_{1,n})<\gamma\quad\forall\ i\in \mathcal{B}_1^c\quad\text{and all n large enough}\,.$

• Under Assumption 2.2 (b)

$$\left(\sup_{(u_1,u_2)\in U_1(i)\times U_2(i)}c_1(i,u_1,u_2)-\rho_{1,n}\right)<\ell(i)\quad\forall\ i\in\mathcal{B}_1^c\quad\text{and all n large enough }.$$

Then applying Itô-Dynkin formula, from Assumption 2.2, we have the following estimates:

• Under Assumption 2.2(a):

$$E_i^{v_1,\hat{v}_2} \left[e^{\hat{\tau}(\mathcal{B}_1)\gamma} W(Y(\hat{\tau}(\mathcal{B}_1))) \right] \le W(i) \ \forall i \in \mathcal{B}_1^c.$$
(3.11)

• Under Assumption 2.2(b):

$$E_{i}^{v_{1},\hat{v}_{2}}\left[e^{\int_{0}^{\hat{\tau}(\mathcal{B}_{1})}\ell(Y(t))dt}W(Y(\hat{\tau}(\mathcal{B}_{1})))\right] \leq W(i) \ \forall i \in \mathcal{B}_{1}^{c}.$$
(3.12)

Now as in [4, Lemma 3.4], using the Lyapunov function W we construct a barrier. Then following arguments similar to [4, Lemma 3.4] and letting $n \to \infty$, there exists $(\rho_1, \psi_1) \in \mathbb{R}_+ \times L_W^{1,\infty}$, $\psi_1 > 0$, satisfying (3.5). By truncating the running cost c_1 , one can show that ρ_1 satisfies (3.6) (see, [4, Lemma 3.5]).

Next we prove the stochastic representation (3.7). Applying Itô-Dynkin formula and Fatou's lemma, for any minimizing selector v_1^* of (3.5) we have

$$\psi_{1}(i) \geq E_{i}^{v_{1}^{*},\hat{v}_{2}} \left[e^{\int_{0}^{\hat{\tau}(\mathcal{B}_{1})} (c_{1}(Y(t),v_{1}^{*}(Y(t-)),\hat{v}_{2}(Y(t-))) - \rho_{1})dt} \psi_{1}(Y(\hat{\tau}(\mathcal{B}_{1}))) \right]$$

$$\geq \inf_{v_{1}\in\mathcal{S}_{1}} E_{i}^{v_{1},\hat{v}_{2}} \left[e^{\int_{0}^{\hat{\tau}(\mathcal{B}_{1})} (c_{1}(Y(t),v_{1}(Y(t-)),\hat{v}_{2}(Y(t-))) - \rho_{1})dt} \psi_{1}(Y(\hat{\tau}(\mathcal{B}_{1}))) \right] \forall i \in \mathcal{B}_{1}^{c}. \quad (3.13)$$

Again, by applying Itô-Dynkin formula, from (3.1) for any $v_1 \in S_1$, T > 0 and $i \in \mathcal{D}_n \cap \mathcal{B}_1^c$ it follows that

$$\begin{split} \psi_{1,n}(i) &\leq E_{i}^{v_{1},\hat{v}_{2}} \left[e^{\int_{0}^{\hat{\tau}(\mathcal{B}_{1})\wedge\tau(\mathcal{D}_{n})\wedge T}(c_{1}(Y(t),v_{1}(Y(t-)),\hat{v}_{2}(Y(t-)))-\rho_{1,n})dt}\psi_{1,n}(Y(\hat{\tau}(\mathcal{B}_{1})\wedge\tau(\mathcal{D}_{n})\wedge T)) \right] \\ &\leq E_{i}^{v_{1},\hat{v}_{2}} \left[e^{\int_{0}^{\hat{\tau}(\mathcal{B}_{1})}(c_{1}(Y(t),v_{1}(Y(t-)),\hat{v}_{2}(Y(t-)))-\rho_{1,n})dt}\psi_{1,n}(Y(\hat{\tau}(\mathcal{B}_{1})))I_{\{\hat{\tau}(\mathcal{B}_{1})\leq\tau(\mathcal{D}_{n})\wedge T\}} \right] \\ &+ E_{i}^{v_{1},\hat{v}_{2}} \left[e^{\int_{0}^{T}(c_{1}(Y(t),v_{1}(Y(t-)),\hat{v}_{2}(Y(t-)))-\rho_{1,n})dt}\psi_{1,n}(Y(T))I_{\{T\leq\hat{\tau}(\mathcal{B}_{1})\wedge\tau(\mathcal{D}_{n})\}} \right]. \quad (3.14) \end{split}$$

Using (3.11) and the fact that $\psi_{1,n} \leq W$ (by our construction), we have

$$E_{i}^{v_{1},\hat{v}_{2}}\left[e^{\int_{0}^{T}(c_{1}(Y(t),v_{1}(Y(t-)),\hat{v}_{2}(Y(t-)))-\rho_{1,n})dt}\psi_{1,n}(Y(T))I_{\{T\leq\hat{\tau}(\mathcal{B}_{1})\wedge\tau(\mathcal{D}_{n})\}}\right]$$

$$\leq e^{(\|c_{1}\|_{\infty}-\rho_{1,n}-\gamma)T}E_{i}^{v_{1},\hat{v}_{2}}\left[e^{T\gamma}W(Y(T))I_{\{T\leq\hat{\tau}(\mathcal{B}_{1})\wedge\tau(\mathcal{D}_{n})\}}\right]$$

$$\leq e^{(\|c_{1}\|_{\infty}-\rho_{1,n}-\gamma)T}W(i).$$

Thus, letting $T \to \infty$ from (3.14) we get

$$\psi_{1,n}(i) \le E_i^{v_1,\hat{v}_2} \left[e^{\int_0^{\hat{\tau}(\mathcal{B}_1)} (c_1(Y(t),v_1((Y(t-))),\hat{v}_2(Y(t-))) - \rho_{1,n})dt} \psi_{1,n}(Y(\hat{\tau}(\mathcal{B}_1))) I_{\{\hat{\tau}(\mathcal{B}_1) \le \tau(\mathcal{D}_n)\}} \right]$$

Now, since $\psi_{1,n} \leq W$ using (3.11) by dominated convergence theorem it follows that

$$\psi_1(i) \le E_i^{v_1, \hat{v}_2} \left[e^{\int_0^{\hat{\tau}(\mathcal{B}_1)} (c_1(Y(t), v_1(Y(t-)), \hat{v}_2(Y(t-))) - \rho_1) dt} \psi_1(Y(\hat{\tau}(\mathcal{B}_1))) \right] \, \forall i \in \mathcal{B}_1^c \,. \tag{3.15}$$

10

Since $v_1 \in S_1$ is arbitrary, combining (3.13) and (3.15), we obtain (3.7). Also, it it clear from the proof that for any minimizing selector v_1^* of (3.5) we have

$$\psi_1(i) = E_i^{v_1^*, \hat{v}_2} \left[e^{\int_0^{\hat{\tau}(\mathcal{B}_1)} (c_1(Y(t), v_1^*(Y(t-)), \hat{v}_2(Y(t-))) - \rho_1) dt} \psi_1(Y(\hat{\tau}(\mathcal{B}_1))) \right] \, \forall i \in \mathcal{B}_1^c.$$
(3.16)

Using (3.12) it is easy to check that the same conclusion holds under Assumption 2.2(b).

Now exploiting the stochastic representation (3.7), we show that $(\rho_1, \psi_1) \in \mathbb{R}_+ \times L_W^{1,\infty}$ is the minimal eigenpair. Suppose $(\hat{\rho}_1, \hat{\psi}_1) \in \mathbb{R}_+ \times L_W^{1,\infty}, \hat{\psi}_1 > 0$ is an eigenpair satisfying

$$\begin{cases} \hat{\rho}_1 \hat{\psi}_1(i) = \inf_{v_1 \in V_1(i)} \left[\sum_{j \in S} \hat{\psi}_1(j) \pi_{ij}(v_1, \hat{v}_2(i)) + c_1(i, v_1, \hat{v}_2(i)) \hat{\psi}_1(i) \right] \text{ for } i \in S, \\ \hat{\psi}_1(i_0) = 1. \end{cases}$$

$$(3.17)$$

We want to show that $\rho_1 \leq \hat{\rho}_1$. If not suppose that $\rho_1 > \hat{\rho}_1$. Then, for any minimizing selector \hat{v}_1^* of (3.17), applying Itô-Dynkin formula and Fatou's lemma, we obtain

$$\hat{\psi}_{1}(i) \geq E_{i}^{\hat{v}_{1}^{*},\hat{v}_{2}} \left[e^{\int_{0}^{\hat{\tau}(\mathcal{B}_{1})} (c_{1}(Y(t),\hat{v}_{1}^{*}(Y(t-)),\hat{v}_{2}(Y(t-))) - \hat{\rho}_{1})dt} \hat{\psi}_{1}(Y(\hat{\tau}(\mathcal{B}_{1}))) \right] \,\forall i \in \mathcal{B}_{1}^{c} \,. \tag{3.18}$$

Whereas from (3.7), we have

$$\psi_1(i) \le E_i^{\hat{v}_1^*, \hat{v}_2} \left[e^{\int_0^{\hat{\tau}(\mathcal{B}_1)} (c_1(Y(t), \hat{v}_1^*(Y(t-)), \hat{v}_2(Y(t-))) - \hat{\rho}_1) dt} \psi_1(Y(\hat{\tau}(\mathcal{B}_1))) \right] \, \forall i \in \mathcal{B}_1^c.$$
(3.19)

Let $\hat{\kappa} := \min_{\mathcal{B}_1} \frac{\hat{\psi}_1}{\psi_1}$. Hence, from (3.18) and (3.19) it follows that $(\hat{\psi}_1 - \hat{\kappa}\psi_1) \ge 0$ in S and $(\hat{\psi}_1 - \hat{\kappa}\psi_1)(\tilde{i}_0) = 0$ for some $\tilde{i}_0 \in \mathcal{B}_1$. Now, combining (3.5) and (3.17) we deduce that

$$\left[\sum_{j\neq\tilde{i}_{0}}(\hat{\psi}_{1}-\hat{\kappa}\psi_{1})(j)\pi_{\tilde{i}_{0}j}(\hat{v}_{1}^{*}(\tilde{i}_{0}),\hat{v}_{2}(\tilde{i}_{0}))\right] \equiv 0.$$
(3.20)

Since Y(t) is irreducible under (\hat{v}_1^*, \hat{v}_2) , in view of (3.20) it is clear that $(\hat{\psi}_1 - \hat{\kappa}\psi_1) \equiv 0$. Again, since $\hat{\psi}_1(i_0) = \psi_1(i_0) = 1$, we get $\hat{\psi}_1 \equiv \psi_1$. But this is a contradiction to the fact that $\rho_1 > \hat{\rho}_1$. Thus we deduce that $(\rho_1, \psi_1) \in \mathbb{R}_+ \times L_W^{1,\infty}$ is the minimal eigenpair. Following the above argument one can show that any eigenfunction satisfying (3.7) is unique upto a scalar multiplication. Also, by the similar argument, one can show that there exists a minimal eigenpair $(\rho_2, \psi_2) \in \mathbb{R}_+ \times L_W^{1,\infty}$ satisfying (3.8), (3.9) and (3.10). This completes the proof.

To proceed further we establish some technical results needed later.

Lemma 3.3. Let Assumptions 2.1, 2.2, and 2.3 hold. Then the maps $\hat{v}_1 \to \psi_2^{\hat{v}_1}$ from $S_1 \to L_W^{\infty}$, $\hat{v}_1 \to \rho_2^{\hat{v}_1}$ from $S_1 \to \mathbb{R}_+$, $\hat{v}_2 \to \psi_1^{\hat{v}_2}$ from $S_2 \to L_W^{\infty}$, and $\hat{v}_2 \to \rho_1^{\hat{v}_2}$ from $S_2 \to \mathbb{R}_+$ are continuous.

Proof. Let $\{v_{2,n}\}$ be a sequence in S_2 such that $v_{2,n} \to \tilde{v}_2$ in S_2 , i.e., for each $i \in S$, $v_{2,n}(i) \to \tilde{v}_2(i)$ in $V_2(i)$. Now by Theorem 3.1, there exists $(\rho_1^{v_{2,n}}, \psi_1^{v_{2,n}}) \in \mathbb{R}_+ \times L_W^{1,\infty}, \ \psi_1^{v_{2,n}} > 0$ satisfying

$$\rho_1^{v_{2,n}} \psi_1^{v_{2,n}}(i) = \inf_{v_1 \in V_1(i)} \left[\sum_{j \in S} \psi_1^{v_{2,n}}(j) \pi_{ij}(v_1, v_{2,n}(i)) + c_1(i, v_1, v_{2,n}(i)) \psi_1^{v_{2,n}}(i) \right], \quad (3.21)$$

with $\psi_1^{v_{2,n}}(i_0) = 1$. Now, since $\psi_1^{v_{2,n}} \in L_W^{1,\infty}$, by a standard diagonalization argument, there exists a function $\psi_1^* \in L_W^{1,\infty}$ such that $\psi_1^{v_{2,n}}(i) \to \psi_1^*(i)$ as $n \to \infty$ for all $i \in S$. Also, $\{\rho_1^{v_{2,n}}\}$ is a bounded sequence. Hence, along a suitable subsequence (without loss of generality denoting by the same notation) $\rho_1^{v_{2,n}} \to \rho_1^*$. Now from (3.21), for any $v_1 \in V_1(i)$ we deduce that

$$\rho_1^{v_{2,n}}\psi_1^{v_{2,n}}(i) \leq \left[\sum_{j\in S}\psi_1^{v_{2,n}}(j)\pi_{ij}(v_1,v_{2,n}(i)) + c_1(i,v_1,v_{2,n}(i))\psi_1^{v_{2,n}}(i)\right].$$

This implies that

$$\rho_{1}^{v_{2,n}}\psi_{1}^{v_{2,n}}(i) - \psi_{1}^{v_{2,n}}(i)\pi_{ii}(v_{1}, v_{2,n}(i)) \leq \left[\sum_{j\neq i}\psi_{1}^{v_{2,n}}(j)\pi_{ij}(v_{1}, v_{2,n}(i)) + c_{1}(i, v_{1}, v_{2,n}(i))\psi_{1}^{v_{2,n}}(i)\right]$$

$$(3.22)$$

Note that

$$\sum_{j \neq i} \psi_1^{v_{2,n}}(j) \pi_{ij}(v_1, v_{2,n}(i)) \le \sum_{j \neq i} W(j) \pi_{ij}(v_1, v_{2,n}(i)).$$
(3.23)

Thus, using Lemma 3.1, generalized Fatou's lemma in [21, Lemma 8.3.7] and taking $n \to \infty$ in (3.22), we get

$$\rho_1^* \psi_1^*(i) \le \left[\sum_{j \in S} \psi_1^*(j) \pi_{ij}(v_1, \tilde{v}_2(i)) + c_1(i, v_1, \tilde{v}_2(i)) \psi_1^*(i) \right].$$

Hence,

$$\rho_1^* \psi_1^*(i) \le \inf_{v_1 \in V_1(i)} \left[\sum_{j \in S} \psi_1^*(j) \pi_{ij}(v_1, \tilde{v}_2(i)) + c_1(i, v_1, \tilde{v}_2(i)) \psi_1^*(i) \right].$$
(3.24)

Since $V_1(i)$ is compact, there exist $v_{1,n}^*, v_1^* \in S_1$ such that $v_{1,n}^* \to v_1^*$ satisfying

$$\rho_1^{v_{2,n}}\psi_1^{v_{2,n}}(i) = \left[\sum_{j\in S}\psi_1^{v_{2,n}}(j)\pi_{ij}(v_{1,n}^*(i), v_{2,n}(i)) + c_1(i, v_{1,n}^*(i), v_{2,n}(i))\psi_1^{v_{2,n}}(i)\right].$$
 (3.25)

Now, using Lemma 3.1, the dominated convergent theorem and passing $n \to \infty$ in (3.25), we obtain

$$\rho_1^*\psi_1^*(i) = \left[\sum_{j \in S} \psi_1^*(j)\pi_{ij}(v_1^*(i), \tilde{v}_2(i)) + c_1(i, v_1^*(i), \tilde{v}_2(i))\psi_1^*(i)\right],$$

Therefore

$$\rho_1^* \psi_1^*(i) \ge \inf_{v_1 \in V_1(i)} \left[\sum_{j \in S} \psi_1^*(j) \pi_{ij}(v_1, \tilde{v}_2(i)) + c_1(i, v_1, \tilde{v}_2(i)) \psi_1^*(i) \right].$$
(3.26)

Hence, from (3.24), and (3.26), it follows that

$$\rho_1^* \psi_1^*(i) = \inf_{v_1 \in V_1(i)} \left[\sum_{j \in S} \psi_1^*(j) \pi_{ij}(v_1, \tilde{v}_2(i)) + c_1(i, v_1, \tilde{v}_2(i)) \psi_1^*(i) \right].$$
(3.27)

Since $\rho_1^{\tilde{v}_2}$ is the minimal eigenvalue corresponding to \tilde{v}_2 of (3.27), we have $\rho_1^* \ge \rho_1^{\tilde{v}_2}$. Suppose $\rho_1^* > \rho_1^{\tilde{v}_2}$. Now, from Theorem 3.1, for any minimizing $\hat{v}_1 \in \mathcal{S}_1$ of (3.5), there exists a finite set $\mathcal{B}_1 \supset \mathcal{K}$, such that

$$\psi_1(i) = E_i^{\hat{v}_1, \tilde{v}_2} \left[e^{\int_0^{\hat{\tau}(\mathcal{B}_1)} (c_1(Y(t), \hat{v}_1(Y(t)), \tilde{v}_2(Y(t))) - \rho_1^{\tilde{v}_2}) dt} \psi_1(Y(\hat{\tau}(\mathcal{B}_1))) \right] \,\forall i \in \mathcal{B}_1^c, \tag{3.28}$$

where $\hat{\tau}(\mathcal{B}_1) = \inf\{t : Y(t) \in \mathcal{B}_1\} =: \tilde{\tau}_1$. Since $\rho_1^* > \rho_1^{\tilde{v}_2}$, by similar arguments as in [4, Lemma 3.4] we deduce that

$$\psi_1^*(i) \le E_i^{\hat{v}_1, \tilde{v}_2} \left[e^{\int_0^{\hat{\tau}(\mathcal{B}_1)} (c_1(Y(t), \hat{v}_1(Y(t)), \tilde{v}_2(Y(t))) - \rho_1^{\tilde{v}_2}) dt} \psi_1^*(Y(\hat{\tau}(\mathcal{B}_1))) \right] \,\forall i \in \mathcal{B}_1^c.$$
(3.29)

From (3.28) and (3.29), we obtain

$$(\psi_1 - \psi_1^*)(i) \ge E_i^{\hat{v}_1, \tilde{v}_2} \left[e^{\int_0^{\hat{\tau}(\mathcal{B}_1)} (c_1(Y(t), \hat{v}_1(Y(t)), \tilde{v}_2(Y(t))) - \rho_1^{\tilde{v}_2}) dt} (\psi_1 - \psi_1^*) (Y(\hat{\tau}(\mathcal{B}_1))) \right] \,\forall i \in \mathcal{B}_1^c.$$

$$(3.30)$$

Now choosing an appropriate constant θ (e.g., $\theta = \max_{\mathcal{B}_1} \frac{\psi_1}{\psi_1^*}$), we have $(\psi_1 - \theta \psi_1^*) \ge 0$ in \mathcal{B}_1 and for some $\hat{i}_0 \in \mathcal{B}_1$, $(\psi_1 - \theta \psi_1^*)(\hat{i}_0) = 0$. Thus, in view of (3.30), we get $(\psi_1 - \theta \psi_1^*) \ge 0$ in S. Now combining (3.5) and (3.27), we get

$$\rho_1^{\tilde{v}_2}(\psi_1 - \theta\psi_1^*)(\hat{i}_0) \ge \left[\sum_{j \in S} (\psi_1 - \theta\psi_1^*)(j)\pi_{\hat{i}_0 j}(\hat{v}_1(\hat{i}_0), \tilde{v}_2(\hat{i}_0)) + c_1(\hat{i}_0, \hat{v}_1(\hat{i}_0), \tilde{v}_2(\hat{i}_0))(\psi_1 - \theta\psi_1^*)(\hat{i}_0)\right]$$

This implies that

$$\sum_{j \neq \hat{i}_0} (\psi_1 - \theta \psi_1^*)(j) \pi_{\hat{i}_0 j}(\hat{v}_1(\hat{i}_0), \tilde{v}_2(\hat{i}_0)) = 0.$$
(3.31)

Since, $\{Y(t)\}_{t\geq 0}$ is irreducible under $(\hat{v}_1, \tilde{v}_2) \in S_1 \times S_2$, from (3.31) it follows that $\psi_1 \equiv \theta \psi_1^*$. But this is a contradiction to the fact that $\rho_1^* > \rho_1^{\tilde{v}_2}$. Hence, we deduce that $\rho_1^* = \rho_1^{\tilde{v}_2}$. This proves the continuty of the map $\hat{v}_2 \to \rho_1^{\hat{v}_2}$. Since $\psi_1^{\hat{v}_{2,n}}(i_0) = 1$ for all $n \geq 1$, we have $\psi_1^*(i_0) = 1$. Hence by Theorem 3.1, we have ψ_1^* is the unique solution of (3.5). Thus $\psi_1^* = \psi_1^{\tilde{v}_2}$. This proves the continuity of the map $\hat{v}_2 \to \psi_1^{\hat{v}_2}$. Continuity of other maps follows by the similar argument.

Fix $\hat{v}_2 \in \mathcal{S}_2$. For each $i \in S, v_1 \in V_1(i)$, set

$$\tilde{F}_1(i, v_1, \hat{v}_2(i)) = \left[\sum_{j \in S} \psi_1^{\hat{v}_2}(j) \pi_{ij}(v_1, \hat{v}_2(i)) + c_1(i, v_1, \hat{v}_2(i)) \psi_1^{\hat{v}_2}(i)\right]$$

where $\psi_1^{\hat{v}_2}$ is the solution of (3.5) corresponding to the strategy $\hat{v}_2 \in \mathcal{S}_2$. Let

$$\tilde{H}(\hat{v}_2) = \left\{ \hat{v}_1^* \in \mathcal{S}_1 : \tilde{F}_1(i, \hat{v}_1^*(i), \hat{v}_2(i)) = \inf_{v_1 \in V_1(i)} \tilde{F}_1(i, v_1, \hat{v}_2(i)) \ \forall \ i \in S \right\}.$$

Then by the compactness of each $V_1(i)$, it follows that $\tilde{H}(\hat{v}_2)$ is a non empty subset of S_1 . It is obvious that, $\tilde{H}(\hat{v}_2)$ is convex and closed. Since S_1 is compact, $\tilde{H}(\hat{v}_2)$ is also compact. Similarly, for $i \in S$, $\hat{v}_1 \in S_1$, $v_2 \in V_2(i)$, we set

$$\tilde{F}_2(i, \hat{v}_1(i), v_2) = \left[\sum_{j \in S} \psi_2^{\hat{v}_1}(j) \pi_{ij}(\hat{v}_1(i), v_2) + c_2(i, \hat{v}_1(i), v_2) \psi_2^{\hat{v}_1}(i) \right], \ i \in S,$$

where $\psi_2^{\hat{v}_1}$ is the solution of (3.8) corresponding to the strategy $\hat{v}_1 \in \mathcal{S}_1$. Let

$$\tilde{H}(\hat{v}_1) = \left\{ \hat{v}_2^* \in \mathcal{S}_2 : \tilde{F}_2(i, \hat{v}_1(i), \hat{v}_2^*(i)) = \inf_{v_2 \in V_2(i)} \tilde{F}_2(i, \hat{v}_1(i), v_2) \ \forall \ i \in S \right\}.$$

Then by analogous arguments, $\tilde{H}(\hat{v}_1)$ is nonempty, convex and is a compact subset of \mathcal{S}_2 . Next set

$$\tilde{H}(\hat{v}_1, \hat{v}_2) = \tilde{H}(\hat{v}_2) \times \tilde{H}(\hat{v}_1).$$

From the above argument it is clear that $\tilde{H}(\hat{v}_1, \hat{v}_2)$ is nonempty, convex, and is a compact subset of $S_1 \times S_2$. Therefore we may define a map from $S_1 \times S_2 \to 2^{S_1} \times 2^{S_2}$.

3.1. The existence of Nash equilibria. Next lemma proves upper-semicontinuity of certain set valued map. This result will be useful in establishing existence of a Nash equilibrium in the space of stationary Markov strategies.

Lemma 3.4. Let Assumptions 2.1, 2.2, and 2.3 hold. Then the map $(\hat{v}_1, \hat{v}_2) \rightarrow \tilde{H}(\hat{v}_1, \hat{v}_2)$ from $S_1 \times S_2 \rightarrow 2^{S_1} \times 2^{S_2}$ is upper semicontinuous.

Proof. Let $\{(v_1^m, v_2^m)\} \in S_1 \times S_2$ and $(v_1^m, v_2^m) \to (\hat{v}_1, \hat{v}_2)$ in $S_1 \times S_2$, i.e., for each $i \in S$, $(v_1^m(i), v_2^m(i)) \to (\hat{v}_1(i), \hat{v}_2(i))$ in $V_1(i) \times V_2(i)$. Let $\overline{v}_1^m \in \tilde{H}(v_2^m)$. Then $\{\overline{v}_1^m\} \subset S_1$. Since S_1 is compact, it has a convergent subsequence (denoted by the same sequence by an abuse of notation), such that

$$\overline{v}_1^m \to \overline{v}_1$$
 in \mathcal{S}_1 .

Then $(\overline{v}_1^m, v_2^m) \to (\overline{v}_1, \hat{v}_2)$ in $\mathcal{S}_1 \times \mathcal{S}_2$. Note that

$$\sum_{j \neq i} \pi_{ij}(\overline{v}_1^m, v_2^m(i)) \psi_1^{v_2^m}(j) \le \sum_{j \neq i} \pi_{ij}(\overline{v}_1^m, v_2^m(i)) W(j).$$

Thus from [[21], Lemma 8.3.7], Assumption 2.3 and the (product) topology of S_k , k = 1, 2, it follows that for each $i \in S$,

$$\sum_{j \in S} \pi_{ij}(\overline{v}_1^m, v_2^m(i)) \psi_1^{v_2^m}(j) + c_1(i, \overline{v}_1^m, v_2^m(i)) \psi_1^{v_2^m}(i)$$

converges to

$$\sum_{j\in S} \pi_{ij}(\overline{v}_1, \hat{v}_2(i)) \psi_1^{\hat{v}_2}(j) + c_1(i, \overline{v}_1, \hat{v}_2(i)) \psi_1^{\hat{v}_2}(i).$$

Hence we have

$$\lim_{m \to \infty} \tilde{F}_1(i, \overline{v}_1^m(i), v_2^m(i)) = \tilde{F}_1(i, \overline{v}_1(i), \hat{v}_2(i)).$$
(3.32)

Now fix $\tilde{v}_1 \in S_1$ and consider the sequence (\tilde{v}_1, v_2^m) . Using the analogous arguments as above, we conclude that

$$\lim_{m \to \infty} \tilde{F}_1(i, \tilde{v}_1(i), v_2^m(i)) = \tilde{F}_1(i, \tilde{v}_1(i), \hat{v}_2(i)).$$
(3.33)

Since $\overline{v}_1^m \in H(v_2^m)$, for any m we have

$$\tilde{F}_1(i, \tilde{v}_1(i), v_2^m(i)) \ge \tilde{F}_1(i, \overline{v}_1^m(i), v_2^m(i)).$$

Thus, in view of (3.32) and (3.33), taking $m \to \infty$ in the above equation, for any $\tilde{v}_1 \in S_1$ we get

$$\tilde{F}_1(i,\tilde{v}_1(i),\hat{v}_2(i)) \ge \tilde{F}_1(i,\overline{v}_1(i),\hat{v}_2(i)).$$

Therefore, $\overline{v}_1 \in \tilde{H}(\hat{v}_2)$. Suppose $\overline{v}_2^m \in \tilde{H}(v_1^m)$ and along a subsequence $\overline{v}_2^m \to \overline{v}_2$ in \mathcal{S}_2 . Then, by the similar arguments as above one can show that $\overline{v}_2 \in \tilde{H}(\hat{v}_1)$. This proves that the map $(\hat{v}_1, \hat{v}_2) \to \tilde{H}(\hat{v}_1, \hat{v}_2)$ is upper-semicontinuous.

Theorem 3.2. Grant Assumptions 2.1, 2.2, and 2.3. Then there exists a Nash equilibrium in the space of stationary Markov strategies $S_1 \times S_2$.

Proof. By Lemma 3.4 and Fan's fixed point theorem [8], there exists a fixed point $(\hat{v}_1^*, \hat{v}_2^*) \in S_1 \times S_2$, for the map $(\hat{v}_1, \hat{v}_2) \to \tilde{H}(\hat{v}_1, \hat{v}_2)$ from $S_1 \times S_2 \to 2^{S_1} \times 2^{S_2}$, i.e.,

 $(\hat{v}_1^*, \hat{v}_2^*) \in \tilde{H}(\hat{v}_1^*, \hat{v}_2^*).$

This implies that $(\rho_1^{\hat{v}_2^*}, \psi_1^{\hat{v}_2^*}), (\rho_2^{\hat{v}_1^*}, \psi_2^{\hat{v}_1^*})$ satisfy the following coupled HJB equations:

$$\begin{cases} \rho_1^{\hat{v}_2^*} \psi_1^{\hat{v}_2^*}(i) &= \inf_{v_1 \in V_1(i)} \left[\sum_{j \in S} \pi_{ij}(v_1, \hat{v}_2^*(i)) \psi_1^{\hat{v}_2^*}(j) + c_1(i, v_1, \hat{v}_2^*(i)) \psi_1^{\hat{v}_2^*}(i) \right] \\ &= \left[\sum_{j \in S} \pi_{ij}(\hat{v}_1^*(i), \hat{v}_2^*(i)) \psi_1^{\hat{v}_2^*}(j) + c_1(i, \hat{v}_1^*(i), \hat{v}_2^*(i)) \psi_1^{\hat{v}_2^*}(i) \right], \end{cases}$$
(3.34)
$$\psi_1^{\hat{v}_2^*}(i_0) = 1$$

and

$$\begin{cases} \rho_{2}^{\hat{v}_{1}^{*}}\psi_{2}^{\hat{v}_{1}^{*}}(i) &= \inf_{v_{2} \in V_{2}(i)} \left[\sum_{j \in S} \pi_{ij}(\hat{v}_{1}^{*}(i), v_{2})\psi_{2}^{\hat{v}_{1}^{*}}(j) + c_{2}(i, \hat{v}_{1}^{*}(i), v_{2})\psi_{2}^{\hat{v}_{1}^{*}}(i) \right] \\ &= \left[\sum_{j \in S} \pi_{ij}(\hat{v}_{1}^{*}(i), \hat{v}_{2}^{*}(i))\psi_{2}^{\hat{v}_{1}^{*}}(j) + c_{2}(i, \hat{v}_{1}^{*}(i), \hat{v}_{2}^{*}(i))\psi_{2}^{\hat{v}_{1}^{*}}(i) \right], \qquad (3.35)$$
$$\psi_{2}^{\hat{v}_{2}^{*}}(i_{0}) = 1.$$

Now by Theorem 3.1, from (3.34), it follows that

Similarly, from (3.35), we have

Thus, from equations (3.36) and (3.37), we get

$$\rho_1^{v_1, \hat{v}_2^*} \ge \rho_1^{\hat{v}_1^*, \hat{v}_2^*}, \ \forall \ v_1 \in \mathcal{A}_1,
\rho_2^{\hat{v}_1^*, v_2} \ge \rho_2^{\hat{v}_1^*, \hat{v}_2^*}, \ \forall \ v_2 \in \mathcal{A}_2.$$

Hence $(\hat{v}_1^*, \hat{v}_2^*) \in \mathcal{S}_1 \times \mathcal{S}_2$ is a Nash equilibrium. This completes the proof.

Now we prove a converse of Theorem 3.2.

Theorem 3.3. Let Assumptions 2.1, 2.2, and 2.3 hold. If $(\underline{v}_1^*, \underline{v}_2^*) \in S_1 \times S_2$ is a Nash equilibrium, *i.e.*,

$$\rho_1^{v_1,\underline{v}_2^*} \ge \rho_1^{\underline{v}_1^*,\underline{v}_2^*}, \ \forall \ v_1 \in \mathcal{A}_1,$$
$$\rho_2^{\underline{v}_1^*,v_2} \ge \rho_2^{\underline{v}_1^*,\underline{v}_2^*}, \ \forall \ v_2 \in \mathcal{A}_2.$$

Then $\underline{v}_1^* \in S_1$ is a minimizing selector of (3.5) (corresponding to fixed strategy $\underline{v}_2^* \in S_2$ of player 2) and $\underline{v}_2^* \in S_2$ is a minimizing selector of (3.8) (corresponding to fixed strategy $\underline{v}_1^* \in S_1$ of player 1).

Proof. Applying analogous arguments as in [[4], Lemma 3.4 and Remark 3.1], one can prove that for the given pair $(\underline{v}_1^*, \underline{v}_2^*) \in S_1 \times S_2$, there exists a eigenpair $(\rho_1^{\underline{v}_1^*, \underline{v}_2^*}, \psi_1^{\underline{v}_1^*, \underline{v}_2^*}) \in \mathbb{R} \times L_W^{\infty}$, $\psi_1^{\underline{v}_1^*} > 0$ and $\rho_1^{\underline{v}_1^*, \underline{v}_2^*} \ge 0$ satisfying

$$\begin{cases} \rho_1^{\underline{v}_1^*,\underline{v}_2^*}\psi_1^{\underline{v}_1^*,\underline{v}_2^*}(i) = \sum_{j\in S} \pi_{ij}(\underline{v}_1^*(i),\underline{v}_2^*(i))\psi_1^{\underline{v}_1^*,\underline{v}_2^*}(j) + c_1(i,\underline{v}_1^*(i),\underline{v}_2^*(i))\psi_1^{\underline{v}_1^*,\underline{v}_2^*}(i), \\ \psi_1^{\underline{v}_1^*,\underline{v}_2^*}(i_0) = 1. \end{cases}$$
(3.38)

Also, for given $\underline{v}_2^* \in S_2$, there exists a minimal eigenpair $(\rho_1^{\underline{v}_2^*}, \psi_1^{\underline{v}_2^*}) \in \mathbb{R}_+ \times L_W^{\infty}, \psi_1^{\underline{v}_2^*} > 0$, satisfying

$$\begin{cases} \rho_1^{\underline{v}_2^*}\psi_1^{\underline{v}_2^*}(i) = \inf_{v_1 \in V_1(i)} \left[\sum_{j \in S} \pi_{ij}(v_1, \underline{v}_2^*(i))\psi_1^{\underline{v}_2^*}(j) + c_1(i, v_1, \underline{v}_2^*(i))\psi_1^{\underline{v}_2^*}(i) \right], \\ \psi_1^{\underline{v}_2^*}(i_0) = 1. \end{cases}$$
(3.39)

Since $\rho_1^{\underline{v}_2^*}$ is a minimal eigenvalue of (3.39), corresponding to \underline{v}_2^* , we have

$$\rho_1^{\underline{v}_2^*} = \inf_{v_1 \in \mathcal{A}_1} \rho_1^{v_1, \underline{v}_2^*}.$$
(3.40)

Also, we have

$$\rho_1^{v_1, \underline{v}_2^*} \ge \rho_1^{\underline{v}_1^*, \underline{v}_2^*}, \ \forall \ v_1 \in \mathcal{A}_1$$

Hence,

$$\inf_{v_1 \in \mathcal{A}_1} \rho_1^{v_1, \underline{v}_2^*} \ge \rho_1^{\underline{v}_1^*, \underline{v}_2^*}.$$
(3.41)

So, by (3.40) and (3.41), we obtain

$$\rho_1^{\underline{v}_2^*} \ge \rho_1^{\underline{v}_1^*, \underline{v}_2^*}.$$

Also, from (3.40), we have

$$\rho_1^{\underline{v}_2^*} \le \rho_1^{\underline{v}_1^*, \underline{v}_2^*}.$$

Hence, we deduce that

$$\rho_1^{\underline{v}_2^*} = \rho_1^{\underline{v}_1^*, \underline{v}_2^*}.\tag{3.42}$$

Now, applying Ito-Dynkin formula, from (3.38), it follows that

$$\psi_1^{\underline{v}_1^*,\underline{v}_2^*}(i) = E_i^{\underline{v}_1^*,\underline{v}_2^*} \left[e^{\int_0^{T \wedge \hat{\tau}(\mathcal{B}_1)} (c_1(Y(t),\underline{v}_1^*(Y(t)),\underline{v}_2^*(Y(t))) - \rho_1^{\underline{v}_1^*,\underline{v}_2^*}) dt} \psi_1^{\underline{v}_1^*,\underline{v}_2^*} (Y(T \wedge \hat{\tau}(\mathcal{B}_1))) \right] \,\forall i \in \mathcal{B}_1^c,$$

where \mathcal{B}_1 is as in Theorem 3.1. Now, by Fatou's Lemma, taking $T \to \infty$ in the above equation, we get

$$\psi_{1}^{\underline{v}_{1}^{*},\underline{v}_{2}^{*}}(i) \geq E_{i}^{\underline{v}_{1}^{*},\underline{v}_{2}^{*}} \left[e^{\int_{0}^{\hat{\tau}(\mathcal{B}_{1})}(c_{1}(Y(t),\underline{v}_{1}^{*}(Y(t)),\underline{v}_{2}^{*}(Y(t))) - \rho_{1}^{\underline{v}_{1}^{*},\underline{v}_{2}^{*}}) dt} \psi_{1}^{\underline{v}_{1}^{*},\underline{v}_{2}^{*}}(Y(\hat{\tau}(\mathcal{B}_{1}))) \right] \forall i \in \mathcal{B}_{1}^{c}.$$

$$(3.43)$$

Again, using (3.39), from Theorem 3.1, it follows that

$$\psi_1^{\underline{v}_2^*}(i) \le E_i^{\underline{v}_1^*, \underline{v}_2^*} \left[e^{\int_0^{\hat{\tau}(\mathcal{B}_1)} (c_1(Y(t), \underline{v}_1^*(Y(t)), \underline{v}_2^*(Y(t))) - \rho_1^{\underline{v}_2^*}) dt} \psi_1^{\underline{v}_2^*} (Y(\hat{\tau}(\mathcal{B}_1))) \right] \,\forall i \in \mathcal{B}_1^c.$$
(3.44)

So, by (3.43) and (3.44), we obtain

$$\psi_{1}^{\underline{v}_{1}^{*},\underline{v}_{2}^{*}}(i) - \psi_{1}^{\underline{v}_{2}^{*}}(i) \\ \geq E_{i}^{\underline{v}_{1}^{*},\underline{v}_{2}^{*}} \left[e^{\int_{0}^{\hat{\tau}(\mathcal{B}_{1})}(c_{1}(Y(t),\underline{v}_{1}^{*}(Y(t)),\underline{v}_{2}^{*}(Y(t))) - \rho_{1}^{\underline{v}_{2}^{*}})dt} (\psi_{1}^{\underline{v}_{1}^{*},\underline{v}_{2}^{*}} - \psi_{1}^{\underline{v}_{2}^{*}})(Y(\hat{\tau}(\mathcal{B}_{1}))) \right] \forall i \in \mathcal{B}_{1}^{c}.$$

$$(3.45)$$

Now arguing as in the proof of Lemma 3.3, we obtain $\psi_1^{\underline{v}_1^*,\underline{v}_2^*}(i) \equiv \psi_1^{\underline{v}_2^*}$. Thus, from (3.38) and (3.39) it follows that \underline{v}_1^* is a minimizing selector of (3.5) (for fixed strategy $\underline{v}_2^* \in S_2$ of player 2). Following similar arguments one can show that \underline{v}_2^* is a minimizing selector of (3.8) (for fixed strategy $\underline{v}_1^* \in S_1$ of player 1). This completes the proof. \Box

4. Example

In this section, we present an illustrative example in wherein transition rates are unbounded and cost rates are nonnegative and unbounded.

Example 4.1. Consider a shop which deals with only one type of product for buying and selling. Suppose there are two workers, say, player 1 and player 2 for buying and selling the products, respectively. The number of stocks in the shop is a finite subset of the set of natural numbers \mathbb{N} at each time $t \geq 0$. There are 'natural' buying and selling rates, say $\tilde{\mu}$ and λ , respectively, and buying parameters h_1 controlled by player 1 and selling parameters h_2 controlled by player 2. When the state of the system is $i \in S := \{1, 2, \dots\}$ (i.e., number of items in the shop), player 1 takes an action u_1 from a given set $U_1(i)$, which may increase $(h_1(i, u_1) \geq 0)$ or decrease $(h_1(i, u_1) \leq 0)$ the buying rate. These actions produce a payoff denoted by $r_1(i, u_1)$ per unit time. Similarly, if the state is $i \in S$, player 2 takes an action u_2 from a set $U_2(i)$ to decrease $(h_2(i,b) \leq 0)$ or to increase $(h_2(i,b) \geq 0)$ the selling rate. These actions result in a payoff denoted by $r_2(i, u_2)$ per unit time. We assume that when the stock of items in the shop becomes 1, the first player may buy any number of stocks of that item as much as he/she likes depending upon the availability of cash. In addition, we assume that player k, (k = 1, 2) 'gets' a reward $r_k(i) := p_k i$ or incurs a cost $r_k(i) := p_k i$ for each unit of time during which the system remains in the state $i \in S$, where $p_k > 0$ is a fixed reward fee, and $p_k < 0$, a fixed cost fee, per stock, from the owner.

We next formulate this model as a continuous-time Markov game. The corresponding transition rate $\overline{\pi}_{ij}(u_1, u_2)$ and payoff rate $\overline{c}_k(i, u_1, u_2)$ for player k(k = 1, 2) are given as follows: for $(1, u_1, u_2) \in K$ (K as in the game model (2.1)).

$$\overline{\pi}_{1j}(u_1, u_2) > 0 \ \forall j \ge 2, \ such \ that \ \sum_{j \in S} \overline{\pi}_{1j}(u_1, u_2) = 0, \ and \ \overline{\pi}_{1j}(u_1, u_2) \le e^{-2\theta j} \ \forall \ j \ge 2,$$

$$(4.1)$$

where $\theta > 0$ is a constant.

Also, for $(i, u_1, u_2) \in K$ with $i \geq 2$,

$$\overline{\pi}_{ij}(u_1, u_2) = \begin{cases} \lambda i + h_2(i, u_2), & \text{if } j = i - 1\\ -\tilde{\mu}i - \lambda i - h_1(i, u_1) - h_2(i, u_2), & \text{if } j = i\\ \tilde{\mu}i + h_1(i, u_1), & \text{if } j = i + 1\\ 0, & \text{otherwise.} \end{cases}$$

$$\bar{c}_1(i, u_1, u_2) := ip_1 - r_1(i, u_1), \ \bar{c}_2(i, u_1, u_2) = ip_2 - r_2(i, u_2) \ for \ (i, u_1, u_2) \in K.$$
 (4.2)

We now investigate conditions under which there exists a Nash-equilibrium. To this end we make following assumptions:

(I) For each $i \in S$, $U_1(i) = U_2(i) = [0, L]$, L > 0 is a constant.

- (II) Let $\lambda \ge \tilde{\mu} > 0$, $\tilde{\mu}i + h_1(i, u_1) > 0$, and $\lambda i + h_2(i, u_2) > 0$ for all $(i, u_1, u_2) \in K$ with $i \ge 2$; and assume that $h_1(1, u_1) > 0$ and $h_2(1, u_2) = 0$ for all $(u_1, u_2) \in U_1(i) \times U_2(i)$.
- (III) The functions $h_1(i, u_1)$, $h_2(i, u_2)$, $r_1(i, u_1)$, $r_2(i, u_2)$, and $\overline{\pi}_{11}(u_1, u_2)$ are continuous in (u_1, u_2) for each fixed $i \in S$. Suppose there exists a finite set \mathcal{K} such that $h_k(i, u_k) = \frac{u_k}{e^{\theta i}} I_{\mathcal{K}}(i)$ and $1 \in \mathcal{K}$. Also assume that $\inf_{(u_1, u_2) \in U_1(\cdot) \times U_2(\cdot)} r_k(\cdot, u_k)$ is norm like function for k = 1, 2.
- (IV) Suppose $ip_k r_k(i, u_k) \ge 0 \ \forall i \in S, (u_1, u_2) \in U_1(i) \times U_2(i) \ and \ (1 e^{-\theta})\lambda + (1 e^{\theta})\tilde{\mu} > p_k \ for \ k = 1, 2.$

Proposition 4.1. Under conditions (I)-(IV), the above controlled system satisfies the Assumptions 2.1, 2.2, and 2.3. Hence by Theorem 3.2, there exists a Nash-equilibrium.

Proof. Take a Lyapunov function as $V(i) := e^{\theta i}$ for $i \in S$ for some $\theta > 0$ as described earlier. Then, we have $V(i) \ge 1$ for all $i \in S$. Now for each $i \ge 2$, and $(u_1, u_2) \in U_1(i) \times U_2(i)$, we have

$$\begin{split} \sum_{j \in S} \overline{\pi}_{ij}(u_1, u_2) V(j) &= \overline{\pi}_{i(i-1)}(u_1, u_2) V(i-1) + V(i) \overline{\pi}_{ii}(u_1, u_2) + V(i+1) \overline{\pi}_{i(i+1)}(u_1, u_2) \\ &= e^{\theta i} \left[(\lambda i + h_2(i, u_2)) e^{-\theta} - (i\tilde{\mu} + \lambda i + h_1(i, u_1) + h_2(i, u_2)) + (\tilde{\mu} i + h_1(i, u_1)) e^{\theta} \right] \\ &= e^{\theta i} i \left[\tilde{\mu}(e^{\theta} - 1) + \lambda(e^{-\theta} - 1) + \frac{e^{\theta} h_1(i, u_1) + e^{-\theta} h_2(i, u_2) - h_1(i, u_1) - h_2(i, u_2)}{i} \right] \\ &= i V(i) [\tilde{\mu}(e^{\theta} - 1) + \lambda(e^{-\theta} - 1)] + \left[u_1(e^{\theta} - 1) + u_2(e^{-\theta} - 1) \right] I_{\mathcal{K}}(i) \\ &\leq i V(i) [\tilde{\mu}(e^{\theta} - 1) + \lambda(e^{-\theta} - 1)] + L(e^{\theta} - 1) I_{\mathcal{K}}(i). \end{split}$$
(4.3)

Now for every $\theta > 0$, we know

$$\lambda(e^{-\theta} - 1) + \tilde{\mu}(e^{\theta} - 1) < 0 \Leftrightarrow \tilde{\mu} < \lambda e^{-\theta}.$$

Let $[\tilde{\mu}(e^{\theta}-1)+\lambda(e^{-\theta}-1)] = -\alpha$ for some $\alpha > 0$. Also, let $\ell(i) = i\alpha$ and $C_4 = \max\left\{L(e^{\theta}-1), \frac{e^{-2\theta}}{1-e^{-\theta}}\right\}$ (see (4.5)). Then for $i \ge 2$, $\sup_{(u_1,u_2)\in U_1(i)\times U_2(i)}\sum_{i\in S}V(j)\overline{\pi}_{ij}(u_1,u_2) \le C_4I_{\mathcal{K}}(i) - \ell(i)V(i) \ \forall i \in S.$ (4.4)

Also, we have

$$\sum_{j \in S} \overline{\pi}_{1j}(u_1, u_2) V(j) < \overline{\pi}_{11}(u_1, u_2) e^{\theta} + \sum_{j \ge 2} e^{-2\theta j} e^{\theta j} \le \overline{\pi}_{11}(u_1, u_2) e^{\theta} + \frac{e^{-2\theta}}{1 - e^{-\theta}} < \infty.$$
(4.5)

Since $-\ell(i) < 1$ for all $i \in S$. Hence from (4.4) and (4.5), for $i \ge 1$, we have

$$\sum_{j \in S} \overline{\pi}_{ij}(u_1, u_2) V(j) \le C_1 V(i) + C_2, \text{ where } C_1 = 1 \text{ and } C_2 = C_4.$$
(4.6)

For $i \geq 2$,

$$-\overline{\pi}_{ii}(u_1, u_2) = \tilde{\mu}i + \lambda i + h_1(i, u_1) + h_2(i, u_2)$$

$$\leq i(\tilde{\mu} + \lambda) + 2L$$

$$\leq \frac{1}{\theta}(\tilde{\mu} + \lambda)V(i) + 2LV(i)$$

$$= (2L + \tilde{\mu} + \lambda)\frac{1}{\theta}V(i)$$

$$= C_3V(i). \qquad (4.7)$$

Take $W = \tilde{W} = V$. Now for k = 1, 2

$$\ell(i) - \sup_{(u_1, u_2) \in U_1(i) \times U_2(i)} \bar{c}_k(i, u_1, u_2) = \alpha i - ip_k + \inf_{u_k \in U_k(i)} r_k(i, u_k)$$
$$= i\beta_k + \inf_{u_k \in U_k(i)} r_k(i, u_k).$$
(4.8)

We see that from condition (IV), that $\beta_k = \alpha - p_k \ge 0$. So, $\ell(i) - \sup_{(u_1, u_2) \in U_1(i) \times U_2(i)} \bar{c}_k(i, u_1, u_2)$ is norm-like function for k = 1, 2. Now by (4.6), we say Assumption 2.1 (i) holds. Also by (4.1) and (4.7), Assumption 2.1 (ii) is verified.

Now we verify Assumption 2.2. By (4.4), (4.5) and (4.8), it is easy to see that Assumption 2.2 is satisfied.

Now by condition (III) and (4.2), we say $c_k(i, u_1, u_2)$ and $\overline{\pi}_{ij}(u_1, u_2)$ are continuous in $(u_1, u_2) \in U_1(i) \times U_2(i)$ for each fixed $i, j \in S$ and for k = 1, 2. So, Assumption 2.3 (i) is verified. By (4.3) and (4.5) and condition (III), we say that Assumption 2.3 (ii) is verified. Also, from (4.1) it is easy to see that Assumption 2.3 (iii) is satisfied.

Hence by Theorem 3.2 there exists a Nash-equilibrium for this controlled process. \Box

5. Acknowledgment

The research work of Mrinal K. Ghosh is partially supported by UGC Centre for Advanced Study. The research work of Somnath Pradhan is partially supported by a National Postdoctoral Fellowship PDF/2020/001938.

References

- A. BASU AND M. K. GHOSH, Nonzero-sum risk-sensitive stochastic games on a countable state space, Math. Oper. Res., 43(2) (2018), pp. 516-532.
- [2] A. BASU AND M. K. GHOSH, Zero-sum risk-sensitive stochastic games on a countable state space, Stochastic Process. Appl., 124(1) (2014), pp. 961-983.
- [3] R. BELLMAN, Dynamic Programming, Princeton University Press, Princeton, N. J., (1957).
- [4] A. BISWAS AND S. PRADHAN, Ergodic risk-sensitive control of Markov processes on countable state space revisited, ArXiv e-prints 2104.04825 (2021), available at https://arxiv.org/abs/2104.04825.
- [5] R. CAVAZOS-CADENA AND E. FERNANDEZ-GAUCHERAND, Controlled Markov chains with risksensitive criteria: average cost, optimality equations, and optimal solutions, Math. Methods Oper. Res., 49 (1999), pp. 299-324.

20

- [6] G. B. Di MASI AND L. STETTNER, Risk-sensitive control of discrete-time Markov processes with infinite horizon, SIAM J. Control Optim., 38(1)(1999), pp. 61-78.
- [7] G. B. Di MASI AND L. STETTNER, Infinite horizon risk-sensitive control of discrete time Markov processes under minorization property, SIAM J. Control Optim., 46(1) (2007), pp. 231-252.
- [8] K. FAN, Fixed-point and minimax theorems in locally convex topological linear spaces, Proc. Nat. Acad. Sc., 38 (1952), pp. 121-126.
- [9] W. H. FLEMING, AND D. HERNANDEZ-HERNANDEZ, Risk-sensitive control of finite state machines on an infinite horizon, SIAM J. Control Optim. 35(5) (1997), pp. 1790-1810.
- [10] M. K. GHOSH, K. S. KUMAR, AND C. PAL, Zero-sum risk-sensitive stochastic games for continuoustime Markov chains, Stoch. Anal. Appl., 34 (2016), pp. 835-851.
- [11] M. K. GHOSH AND S. SAHA, Risk-sensitive control of continuous-time Markov chains, Stochastics, 86 (2014), pp. 655-675.
- [12] S. GOLUI AND C. PAL, Continuous-time zero-sum games for Markov chains with risk-sensitive finite-horizon cost criterion, Stoch. Anal. Appl., (2021), available at https://doi.org/10.1080/07362994.2021.1889381.
- [13] X. P. GUO AND O. HERNANDEZ-LERMA, Zero-sum games for continuous-time Markov chains with unbounded transition and average payoff rates, J. Appl. Probab., 40(2) (2003), pp. 327-345.
- [14] X. P. GUO AND O. HERNANDEZ-LERMA, Nonzero-sum games for continuous-time Markov chains with unbounded discounted payoffs, J. Appl. Probab., 42(2) (2005) pp. 303-320.
- [15] X. P. GUO AND O. HERNANDEZ-LERMA, Zero-sum games for continuous-time jump Markov processes in Polish spaces: discounted payoffs, Adv. in Appl. Probab. 39(3) (2007) pp. 645-668.
- [16] X. P. GUO AND O. HERNANDEZ-LERMA, Continuous-Time Markov decision processes: Theory and Applications, Stoch. modelling and Appl. Probab., Springer, Berlin, 62 (2009).
- [17] X. P. GUO AND Z. W. LIAO, Risk-sensitive discounted continuous-time Markov decision processes with unbounded rates, SIAM J. Control Optim., 57 (2019), pp. 3857-3883.
- [18] X. P. GUO, Q. LIU, AND Y. ZHANG, Finite horizon risk-sensitive continuous-time Markov decision processes with unbounded transition and cost rates, 4OR, 17 (2019), pp. 427-442.
- [19] X. P. GUO AND A. PIUNOVSKIY, Discounted continuous-time Markov decision processes with constraints: Unbounded transition and loss rates, Math. Oper. Res., 36 (2011), pp. 105-132.
- [20] X. P. GUO AND X. SONG, Discounted continuous-time constrained Markov decision processes in polish spaces, Ann. Appl. Probab., 21 (2011), pp. 2016-2049.
- [21] O. HERNANDEZ-LERMA, J. LASSERRE, Further topics on discrete-time Markov control processes, Springer, New York, (1999).
- [22] M. Y. KITAEV, Semi-Markov and jump Markov controlled models: Average cost criterion, SIAM Theory Probab. Appl., 30 (1995), pp. 272-288.
- [23] M. Y. KITAEV AND V.V. RYKOV, Controlled Queueing Systems, CRC Press, Boca Raton, (1995).
- [24] K.S. KUMAR AND C. PAL, Risk-sensitive control of jump process on denumerable state space with near monotone cost, Appl. Math. Optim., 68 (2013), pp. 311-331.
- [25] K.S. KUMAR AND C. PAL, Risk-sensitive control of continuous-time Markov processes with denumerable state space, Stoch. Anal. Appl., 33 (2015), pp. 863-881.
- [26] C. PAL AND S. PRADHAN, Risk sensitive control of pure jump processes on a general state space, Stochastics, 91(2) (2019), pp. 155-174.
- [27] A. PIUNOVSKIY AND Y. ZHANG, Discounted continuous-time Markov decision processes with unbounded rates: The convex analytic approach, SIAM J. Control Optim., 49 (2011), pp. 2032-2061.
- [28] Q. WEI, Nonzero-sum risk-sensitive finite-horizon continuous-time stochastic games, Statistics & Probability Letters, 147 (2019), pp. 96-104.
- [29] P. WHITTLE, Risk-Sensitive Optimal Control, Wiley-Inter science Series in Systems and Optimization, John Wiley & Sons Ltd., Chichester, (1990).
- [30] Y. ZHANG, Continuous-time Markov decision processes with exponential utility, SIAM J. Control Optim., 55 (2017), pp. 2636-2660.

Department of Mathematics, Indian Institute of Science, Bangalore-560012, India. $Email \ address: \ mkg@iisc.ac.in$

Department of Mathematics, Indian Institute of Technology Guwahati, Guwahati, Assam, India

Email address: golui@iitg.ac.in

Department of Mathematics, Indian Institute of Technology Guwahati, Guwahati, Assam, India

Email address: cpal@iitg.ac.in

Department of Mathematics, Indian Institute of Science Education and Research, Pune, Maharashtra-411008, India

Email address: somnath@iiserpune.ac.in

22