A General Framework for Learning-Based Distributionally Robust MPC of Markov Jump Systems

Mathijs Schuurmans and Panagiotis Patrinos

Abstract—We present a learning model predictive control (MPC) scheme for chance-constrained Markov jump systems with unknown switching probabilities. Using samples of the underlying Markov chain, ambiguity sets of transition probabilities are estimated which include the true conditional probability distributions with high probability. These sets are updated online and used to formulate a time-varying, risk-averse optimal control problem. We prove recursive feasibility of the resulting MPC scheme and show that the original chance constraints remain satisfied at every time step. Furthermore, we show that under sufficient decrease of the confidence levels, the resulting MPC scheme renders the closed-loop system mean-square stable with respect to the truebut-unknown distributions, while remaining less conservative than a fully robust approach. Finally, we show that the value function of the learning MPC converges from above to its nominal counterpart as the sample size grows to infinity. We illustrate our approach on a numerical example.

I. INTRODUCTION

A. Background, motivation and related work

Due to the ubiquitous nature of stochastic uncertainty in processes arising in virtually all branches of science and engineering, control of dynamical systems perturbed by stochastic processes is a long-standing topic of research. Model predictive control (MPC) – stochastic MPC in particular – has been a popular and successful tool in this endeavor, due to its ability to naturally include probabilistic information directly into the control design via the cost, the dynamics and the constraints [2]–[4]. In classical stochastic MPC, however, it is typically assumed that the distribution of the underlying stochastic process is known, although in practice, this is usually not the case. If the disturbance takes values on a bounded set, the absence of full distributional knowledge can be taken into account by designing the controller under the worst-case realization of the stochastic disturbance. This approach is referred to as robust MPC [2], [4].

An obvious drawback of robust approaches is that the complete disregard of the probabilistic nature of the disturbance can be rather crude, resulting in a tendency for overly conservative decisions. As an alternative approach, one may simply compute an empirical estimate of the disturbance distribution and replace the true value by this estimate in the optimal control problem. Although this is a reasonable approach given a sufficient amount of data, for more moderate sample sizes, there may be a significant misestimation of the underlying distributions—often referred to as *ambiguity*. It is well known that this is likely to cause degradation of the resulting performance when evaluated on new samples from the true distribution. This phenomenon is known as the *optimizer's curse* [5]. To account for

M. Schuurmans and P. Patrinos are with the Department of Electrical Engineering (ESAT-STADIUS), KU Leuven, Kasteelpark Arenberg 10, 3001 Leuven, Belgium. Email: {mathijs.schuurmans, panos.patrinos}@esat.kuleuven.be

This work was supported by: FWO projects: No. G086318N; No. G086518N; Fonds de la Recherche Scientifique – FNRS, the Fonds Wetenschappelijk Onderzoek–Vlaanderen under EOS Project No. 30468160 (SeLMA), Research Council KU Leuven C1 project No. C14/18/068 and the Ford–KU Leuven Research Alliance project No. KUL0023.

A preliminary version of this work has been presented at the 59th IEEE Conference on Decision and Control [1].

this ambiguity, one could, instead of a point estimate, construct a set of all distributions (an *ambiguity set*) that are in some specific sense consistent with the data. By accounting for the worst-case distribution within this set, the decision maker is protected against the limitations of the finite sample size.

This approach, known as distributionally robust (DR) optimization [6], addresses the drawbacks of the above approaches by utilizing available data, but only to the extent that it is statistically meaningful. As more data is gathered online and ambiguity sets get updated accordingly, it is expected that these sets will shrink, so that the optimal decisions gradually become less conservative. This, among other desirable properties, has caused an increasing popularity of DR methods in recent years, initially mostly in stochastic programming and operations research communities [5], [7]–[10] and more recently in (optimal) control [11]-[16] as well. See also [17] for a comprehensive review. Much of the earlier work focuses on the study of particular classes of ambiguity sets, each modelling certain structural assumptions on the underlying distribution. Although most of our analysis does not require a particular family of ambiguity sets, we will, for concreteness, put particular emphasis on ambiguity sets that are written as a divergence ball around an empirical estimate, as this family of sets is a natural choice in the setting at hand. This is described in Section III, where a table containing several choices for the divergence is provided.

As the focus of research in data-driven and learning-based control is gradually shifting towards real-life, safety-critical applications, there has been an increasing concern for safety guarantees of data-driven methods, which are valid in a finite data regime. This has led to a variety of different approaches besides distributionally robust methodologies, each valid under different assumptions on the data-generating process and the controlled systems. For instance, this has led to data-driven variants of tube-based MPC [18], [19], Gaussian-process based estimation with reachability-based safe set constraints [20], Data-enabled predictive control ("DeePC") [21] combining Willems' fundamental lemma with MPC for linear systems, or techniques based on Koopman operators [22]. We refer to [23] for a recent survey.

In this work, we allow for general (possibly nonlinear) dynamics under stochastic disturbances with unknown distribution, and subject to chance constraints. However, we restrict our attention to finitelysupported stochastic disturbances. One of the advantages of this construction is that the predicted evolution of the system can be represented on a scenario tree, which allows us to explicitly (and without approximation) optimize over closed-loop control policies, rather than open-loop sequences. This property helps combat excessive conservatism due to accumulation of uncertainty over the prediction horizon [24]–[26]. Motivated by similar considerations, [27] and [28] utilize scenario trees to approximate the realizations of continuous disturbances. [28] then considers safety separately by projecting the computed control action onto a set of control actions that keep the state within safe robust control invariant (RCI) set, similarly to [20]. This projection requires the additional solution of a mixed-integer quadratic program (MIOP), whenever the used RCI set is polyhedral. In our setting, however, we consider the switching behavior inherent to the system, allowing us to provide

safety guarantees directly through the application of MPC theory on the joint controller-learner system.

We will in particular assume that the underlying disturbance process is a Markov chain, leading to a system class commonly referred to as Markov jump systems. Control of this class of systems has been widely studied and has been used to model systems stemming from a wide range of applications [25], [29], [30]. In the known distribution case, stability analysis of nonlinear stochastic MPC for this system class has been performed from a worst-case perspective [31], in mean-square sense [30] and in the more general risk-square sense [32], [33]. We emphasize here the distinction between risk-averse and DR approaches, where the former optimizes a given coherent risk measure with respect to the true distribution, whereas the latter constructs a data-driven ambiguity set with respect to which the stochastic cost is robustified. By the dual risk representation [34, Thm. 6.4], every ambiguity set induces some coherent risk measure and vice-versa, leading both approaches to solve the same class of optimization problems. However, the statistical interpretation and thus, the corresponding guarantees differ significantly.

Indeed, by the mentioned equivalence, the notion of risk-square stability in [32] guarantees mean-square stability (MSS) with respect to all the distributions within the 'ambiguity set' induced by the used risk measure. In practice, however, this is insufficient to guarantee MSS with respect to the true-but-unknown distribution, as it is impossible to construct a nontrivial ambiguity set that contains the true distribution with certainty. However, we will show that by careful design of a data-driven sequence of ambiguity sets – which only contain the true distributions with high probability – this concept can be extended to show MSS, as well as recursive constraint satisfaction with respect to the true distribution, under some additional assumptions.

Other data-driven methods have been proposed to design controllers for *unknown transition probabilities* [35], [36]. However, these works are restricted to a simpler, *unconstrained* setting involving only linear state-feedback policies. Furthermore, related risk-averse and DR techniques have been proposed for Markov decision processes (MDPs) [37]–[40], although these consider discrete states and actions, allowing one to solve directly the Bellman equation over all admissible policies. Unfortunately, these techniques become intractable in the present setting involving continuous states and actions.

We finally study the convergence of the optimal value function of our learning controller to the nominal counterpart. This property, known as asymptotic consistency, has recently been studied in the stochastic optimization literature for (static) distributionally robust optimization problems under Wasserstein ambiguity [5], [41]. A common assumption in this line of work is Lipschitz continuity of the cost/constraint functions with respect to the random variable. This assumption is not suitable for our purposes, since we consider discrete random variables $w \in W$ for which a suitable norm may not exist. Instead, we will in some cases need to resort to a uniform boundedness assumption, which serves a similar purpose. In the nonconvex case, the authors of [41] base their analysis on [42], in which the ambiguity sets are not assumed to be random. An additional assumption is added that the constraint boundary has probability zero, such that almost everywhere, the constraint is continuous. This assumption helps in dealing with the discontinuity of the stepfunction at 0 which is inherent to chance constraints. Alternatively, the chance constraints can be replaced by risk constraints involving the average value-at-risk [43], which also circumvents this issue. Besides the mentioned differences in assumptions, additional care is required to handle the multistage nature of the stochastic optimization problems considered here. Specifically, both the optimal cost and the

feasible set are defined recursively through the Bellman operator (see Section V), causing more complex characterizations of the optimal value function as well as reduced freedom in selecting the problem parameters to ensure its required properties as compared to a static two-stage stochastic program.

B. Contributions

Summarizing the previous discussion, we highlight the following contributions of our work. (i) We present a general online learning DR-MPC framework for Markov switching systems with unknown transition probabilities. The resulting closed-loop system satisfies the (chance) constraints of the original stochastic problem and allows for online improvement of performance based on observed data. Thus, we extend the recently developed framework of risk-averse MPC [32], [33], [44] to an online learning setting, in which the involved risk measures are selected and calibrated automatically based on their dual (DR) interpretation. To this end, we formalize the procedure for estimating and updating the corresponding ambiguity sets as a dynamical system, which we refer to as the learning system. We present conditions on this learning system to ensure its convergence and to obtain meaningful statistical guarantees on the resulting controllers with respect to the unknown underlying distributions . (ii) We provide sufficient conditions for recursive feasibility and mean-square stability of the DR-MPC law, with respect to the true-but-unknown distribution. To this end, we state the problem in terms of an augmented state vector including the state of the previously mentioned *learning system*. The dynamics of this so-called *learner state* can be easily expressed for common choices for the ambiguity set. This idea, which is closely related to that of information states [45, Ch. 5] allows us to formulate the otherwise time-varying optimal control problem as a dynamic programming recursion, facilitating stability analysis of the original control system and the learning system jointly. (iii) We provide sufficient conditions under which the value of the DR problem converges from above to that of the nominal optimal control problem, extending existing results in stochastic optimization to the constrained, multi-stage, dynamical setting.

C. Notation

Let \mathbb{N} denote the set of natural numbers and $\mathbb{N}_{>0} := \mathbb{N} \setminus \{0\}$. For two naturals $a,b\in\mathbb{N}$ with $a\leq b$, we denote $\mathbb{N}_{[a,b]}:=\{n\in\mathbb{N}\}$ $\mathbb{N} \mid a \leq n \leq b$ and similarly, we introduce the shorthand $w_{[a,b]} := (w_t)_{t=a}^b$ to denote a sequence of variables indexed from a to b. We denote the extended real line by $\overline{\mathbb{R}} := \mathbb{R} \cup \{\pm \infty\}$ and the set of *nonnegative* (extended) real numbers by \mathbb{R}_+ (and $\overline{\mathbb{R}}_+$). The cardinality of a (finite) set W is denoted by |W|. We write $f: X \rightrightarrows Y$ to denote that f is a set-valued mapping from X to Y. A function is lower semicontinuous (lsc) if its epigraph is closed. Given a matrix $P \in \mathbb{R}^{n \times m}$, we denote its (i, j)'th element by P_{ij} and its i'th row as $P_{i:} \in \mathbb{R}^m$. The i'th element of a vector x is denoted x_i . $\mathbf{vec}(M)$ denotes the vertical concatenation of the columns of a matrix M. We denote the vector in \mathbb{R}^k with all elements one as $\mathbf{1}_k := (1)_{i=1}^k$ and the probability simplex of dimension k as $\Delta_k := \{ p \in \mathbb{R}_+^k \mid p^\top \mathbf{1}_k = 1 \}.$ We define the function $\mathbf{1}_{x=y} = 1$ if x = y and 0 otherwise. The indicator function $\delta_X : \mathbb{R}^n \to \overline{\mathbb{R}}$ of a set $X \subseteq \mathbb{R}^n$ is defined by $\delta_X(x) = 0$ if $x \in X$ and ∞ otherwise. The level set of a function $V:\mathbb{R}^n\to\overline{\mathbb{R}}$ is denoted $\mathbf{lev}_{<\varepsilon} V := \{x \in \mathbb{R}^n \mid V(x) \le \varepsilon\}$. The interior of a set X is denoted int X. We denote the positive part of a quantity x as $[x]_{+} := \max\{0, x\}$, where max is taken element-wise. We say that a function $\phi: \mathbb{R}_+ \to \mathbb{R}_+$ belongs to the class of \mathcal{K}_{∞} functions if it is continuous, strictly increasing, unbounded, and zero at zero [4].

Finally, given a nonempty, proper cone \mathcal{K} , the generalized inequality $a \preccurlyeq_{\mathcal{K}} b$ is equivalent to $b-a \in \mathcal{K}$. $\mathcal{K}^* := \{y \mid \langle x,y \rangle \geq 0, \ \forall x \in \mathcal{K}\}$ denotes the dual cone of \mathcal{K} .

II. PROBLEM STATEMENT AND STRUCTURAL ASSUMPTIONS

Let $\mathbf{w}:=(w_t)_{t\in\mathbb{N}}$ denote a discrete-time, time-homogeneous Markov chain defined on some probability space 1 (Ω,F,\mathbb{P}) and taking values on $W:=\mathbb{N}_{[1,d]}.$ The transition kernel governing the Markov chain is denoted by $P=(P_{ij})_{i,j\in W},$ where $P_{ij}=\mathbb{P}[w_t=j\mid w_{t-1}=i].$ We refer to w_t as the mode of the chain at time t. For simplicity, we will assume that the initial mode is known to be i, so $p_0=(1_{w=i})_{w\in W}.$ Therefore, the Markov chain is fully characterized by its transition kernel. Finally, we will assume that the Markov chain is ergodic.

Assumption II.1 (Ergodicity). The Markov chain $(w_t)_{t\in\mathbb{N}}$ is ergodic, i.e., there exists a value $k\in\mathbb{N}_{>0}$, such that $P^k>0$ elementwise.

This assumption, stating that every mode is reachable from any other mode in k steps, ensures that every mode of the chain gets visited infinitely often [46, Ex. 8.7]. This will allow us to guarantee convergence of the proposed learning MPC scheme to its nominal counterpart. (See Section V.)

We will consider discrete-time systems with dynamics of the form

$$x_{t+1} = f(x_t, u_t, w_{t+1}),$$
 (1)

where $x_t \in \mathbb{R}^{n_x}$, $u_t \in \mathbb{R}^{n_u}$ are the state and control action at time t, respectively. We will assume that the state x_t and mode w_t are observable at time t. This is equivalent to the more common notation $x_{t+1} = f(x_t, u_t, w_t)$, assuming w_{t-1} is observable. However, as we will consider w_t to be part of the system state at time t, the notation of (1) will be more convenient.

Since w_t is drawn from a Markov chain, such systems are commonly referred to as Markov jump systems. Whenever $f(\cdot,\cdot,w)$ is a linear function, (1) describes a Markov jump linear system [29]. Since the state x_t and mode w_t are observable at time t, the distribution of x_{t+1} depends solely on the conditional switching distribution P_{w_t} ; for a given control action u_t .

For a given state-mode pair (x, w), we will impose probabilistic constraints of the form

$$\mathsf{AV}@\mathsf{R}^{P_{w:}}_{\alpha}\big[g_i(x,u,w,v)\mid x,w\big]\leq 0,\; i\in\mathbb{N}_{[1,n_g]} \tag{2}$$

where $v \sim P_w$: is randomly drawn from the Markov chain \mathbf{w} in mode $w, g_i : \mathbb{R}^{n_x} \times \mathbb{R}^{n_u} \times W^2 \to \mathbb{R}$ are constraint functions with corresponding constraint violation rates α_i , and AV@R denotes the (conditional) average value-at-risk. The conditional AV@R (at level $\alpha \in [0,1]$ and with reference distribution $p \in \Delta_d$) of the random variable $\xi : W^2 \to \mathbb{R}$ is defined as

$$\mathsf{AV}@\mathsf{R}^p_\alpha[\xi(w,v)\mid w]$$

$$= \begin{cases} \min_{t \in \mathbb{R}} t + 1/\alpha \mathbb{E}_p \left\{ \left[\xi(w, v) - t \right]_+ \mid w \right\}, & \alpha \neq 0 \\ \max_{v \in W} \left\{ \xi(w, v) \right\}, & \alpha = 0, \end{cases}$$
(3)

and it has the useful property that if $p=P_{w:}$, then the following implication holds tightly [34, sec. 6.2.4]

$$\mathsf{AV@R}^p_\alpha[\xi(w,v)\mid w] \le 0 \Rightarrow \mathbb{P}[\xi(w,v) \le 0\mid w] \ge 1 - \alpha. \tag{4}$$

By exploiting the dual risk representation [34, Thm 6.5], the left-hand inequality in (4) can be formulated in terms of only linear

constraints [44]. As such, it can be used as a tractable surrogate for chance constraints which would lead to nonconvex, non-smooth constraints [43]. By appropriate choices of α_i and g_i , constraint (2) can be used to encode robust constraints ($\alpha_i = 0$) or probabilistic constraints ($0 < \alpha_i < 1$) on the state, the control action, or both. Note that it additionally covers chance constraints on the successor state f(x, u, v) under input u, conditioned on the current values x and x and x are notation, we will without loss of generality assume that x and x are notation, we written as

$$\mathcal{U}(x,w) := \left\{ u \in U : \mathsf{AV}@\mathsf{R}^{Pw:}_{\alpha} \big[g(x,u,w,v) \mid x,w \big] \leq 0 \right\}, \quad (5)$$

where $U \subseteq \mathbb{R}^{n_u}$ is a nonempty, closed set.

Ideally, our goal is to synthesize – by means of a stochastic MPC scheme – a stabilizing control law $\kappa_N:\mathbb{R}^{n_x}\times W\to\mathbb{R}^{n_u}$, such that for the closed loop system $x_{t+1}=f(x_t,\kappa_N(x_t,w_t),w_{t+1})$, it holds almost surely that $\kappa_N(x_t,w_t)\in\mathcal{U}(x_t,w_t)$, for all $t\in\mathbb{N}$. Consider a sequence of N control laws $\pi=(\pi_k)_{k=0}^{N-1}$, referred to as a *policy* of length N. Given a stage cost $\ell:\mathbb{R}^{n_x}\times\mathbb{R}^{n_u}\times W\to\mathbb{R}_+$, and a terminal cost $V_f:\mathbb{R}^{n_x}\times W\to\mathbb{R}_+$ and corresponding terminal set $\mathcal{X}_f\colon \overline{V_f}(x,w):=V_f(x,w)+\delta_{\mathcal{X}_f}(x,w)$, we can assign to each such policy π , a cost

$$V_N^{\pi}(x, w) := \mathbb{E}\left[\sum_{k=0}^{N-1} \ell(x_k, u_k, w_k) + \overline{V_f}(x_N, w_N)\right],$$
 (6)

where $x_{k+1} = f(x_k, u_k, w_{k+1})$, $u_k = \pi_k(x_k, w_k)$ and $(x_0, w_0) = (x, w)$, for $k \in \mathbb{N}_{[0, N-1]}$. This defines the following stochastic optimal control problem (OCP).

Definition II.2 (Stochastic OCP). For a given state-mode pair (x, w), the optimal cost of the stochastic OCP is

$$V_N(x,w) = \min_{\pi} V_N^{\pi}(x,w)$$
 (7a)

subject to

$$x_0 = x, w_0 = w, \pi = (\pi_k)_{k=0}^{N-1},$$
 (7b)

$$x_{k+1} = f(x_k, \pi_k(x_k, w_k), w_{k+1}),$$
 (7c)

$$\pi_k(x_k, w_k) \in \mathcal{U}(x_k, w_k), \ \forall k \in \mathbb{N}_{[0, N-1]}. \tag{7d}$$

We denote by $\Pi_N(x, w)$ the corresponding set of minimizers.

To ensure existence of a solution to (7) (and its DR counterpart, defined in Section IV), we will impose the following (standard) regularity conditions [4], [32].

Assumption II.3 (Problem regularity). The following are satisfied for all $w, v \in W$:

- (i) Functions $\ell(\cdot,\cdot,w):\mathbb{R}^{n_x}\times\mathbb{R}^{n_u}\to\mathbb{R}_+$, $V_{\mathbf{f}}(\cdot,w):\mathbb{R}^{n_x}\to\mathbb{R}_+$, $f(\cdot,\cdot,w)$, and $g(\cdot,\cdot,w,v)$, $i\in\mathbb{N}_{[1,n_q]}$ are continuous;
- (ii) U and \mathcal{X}_f are closed;
- (iii) f(0,0,w) = 0, $\ell(0,0,w) = 0$, $0 \in \mathcal{U}(0,w)$, and $\overline{V_f}(0,w) = 0$;
- (iv) One of the following is satisfied:
 - 1) U is compact; or
 - 2) $\ell(x, u, w) \ge c(\|u\|)$ with $c \in \mathcal{K}_{\infty}$, for all $(x, u) \in \mathbb{R}^{n_x} \times U$.

Let $(\pi_k^{\star}(x,w))_{k=0}^{N-1} \in \Pi_N(x,w)$, so that the stochastic MPC control law is given by $\kappa_N(x,w) = \pi_0^{\star}(x,w)$. Sufficient conditions on the terminal cost $\overline{V_{\rm f}}$ and its effective domain ${\bf dom} \, \overline{V_{\rm f}} = \mathcal{X}_{\rm f}$ to ensure mean-square stability of the closed-loop system, have been studied for a similar problem set-up in [30], among others.

Both designing and computing such a stochastic MPC law requires knowledge of the probability distribution governing the state dynamics (1), or equivalently, of the transition kernel P. In the absence of this knowledge, these probabilities are to be estimated from a finitely-sized data set and therefore subject to some level of ambiguity. Our

¹For an explicit construction of (Ω, F, \mathbb{P}) , we refer to [46, Thm. 8.1].

goal is to devise an MPC scheme which uses the available data in a principled manner, while explicitly taking this ambiguity into account.

To this end, we will introduce the notion of a *learner state*, which is very similar in spirit to the concept of an *information state*, commonly used in control of partially observed Markov decision processes [47], where – in contrast to our approach – it is typically adopted in a Bayesian setting. In both cases, however, it can be regarded as an internal state of the controller that stores all the information required to build (a set of) conditional distributions over the next state, given the observed data. We will make this more precise in the next section. Equipped with such a learning system, our aim is to find a data-driven approximation to the stochastic OCP defined by (7), which asymptotically attains the optimal cost while preserving stability and constraint satisfaction during closed-loop operation.

The remainder of this work is organized as follows. Section III formalizes the assumed learning system and presents and several classes of ambiguity sets found in the literature that fit the framework. In Section IV, we use this learning system to construct a distributionally robust counterpart to the optimal control problem in terms of the ingredients introduced above. Section V contains a theoretical analysis of the proposed scheme; and in Section VI, we illustrate the approach on some numerical examples.

III. DATA-DRIVEN AMBIGUITY SETS

A. Abstract learning system

As mentioned in the previous section, we model the procedure that maps the observed data into a set of transition probabilities as a generic Markovian system, which we refer to as the *learning system*. We first state the required structure in a compact, abstract notation and later provide a concrete example, which will suffice in many practical cases.

Assumption III.1 (Learning system). Given a sequence $w_{[0,t]}$ sampled from the Markov chain \mathbf{w} , we can compute

- 1) a statistic $s_t: W^{t+1} \to \mathcal{S} \subseteq \mathbb{R}^{n_s}$, with \mathcal{S} compact, accompanied by a vector of confidence parameters $\boldsymbol{\beta}_t = (\beta_{t,i})_{i=1}^{n_{\beta}} \in \mathcal{I} := [0,1]^{n_{\beta}}$, for which there exist some Markovian dynamics \mathcal{L} and C such that $s_{t+1} = \mathcal{L}(s_t, \boldsymbol{\beta}_t, w_t, w_{t+1})$ and $\boldsymbol{\beta}_{t+1} = C(\boldsymbol{\beta}_t), t \in \mathbb{N}$;
- 2) an ambiguity set $A: S \times W \times [0,1] \Rightarrow \Delta_d: (s,w,\beta) \mapsto A_{\beta}(s,w)$, mapping s_t , w_t and the component $\beta_{t,i}$ to a convex subset of the d-dimensional probability simplex Δ_d , such that for all $t \in \mathbb{N}$, $w \in W$, and for all $i \in \mathbb{N}_{[1,n_{\beta}]}$,

$$\mathbb{P}[P_{w:} \in \mathcal{A}_{\beta_{t,i}}(s_t, w)] \ge 1 - \beta_{t,i}. \tag{8}$$

We will refer to s_t and β_t as the learner state and the confidence vector at time t, respectively.

Remark III.2 (Learner dynamics \mathcal{L} , C). The existence of the dynamics \mathcal{L} and C implies that the system with the augmented state consisting of both the original system state-mode pair (x_t, w_t) and the learner-confidence pair (s_t, β_t) , is Markovian. This assumption aids the theoretical analysis in Section V and is not restrictive in practice, as it essentially only requires that finite memory is needed for the method, which is the case for all implementable methods. For concreteness, typical examples for \mathcal{L} and C, which are valid for many practical use cases, are presented in Example III.8 and Example III.6, respectively.

Remark III.3 (confidence levels). We consider a vector of confidence levels, rather than a single value. This is motivated by the fact that one would often wish to assign separate confidence levels to ambiguity sets corresponding to the cost function, and to those corresponding

to the n_g chance constraints (See Definition IV.3). Accordingly, we will assume that $n_\beta = n_g + 1$.

In order to ensure reasonable behavior of the learning system, we impose the following restrictions on the choice of the learning dynamics and the confidence levels.

Assumption III.4. There exists a stationary learner state s^* such that $s^* = \mathcal{L}(s^*, \boldsymbol{\beta}, w, v)$, for all $(\boldsymbol{\beta}, w, v) \in \mathcal{I} \times W^2$, and that from any initial state s_0 , $\lim_{t\to\infty} s_t = s^*$, a.s.

Assumption III.5. The confidence dynamics $\beta_{t+1} = C(\beta_t)$ is chosen such that (i) $\sum_{t=0}^{\infty} \beta_t < \infty$; and (ii) $\lim_{t \to \infty} \frac{\log \beta_t}{t} = 0$, element-wise.

Assumption III.4 imposes that asymptotically, the learner settles down to some value which is no longer modified by additional data. It is natural to expect that in such a state, the learner has acquired perfect knowledge of the underlying transition kernel and the ambiguity sets $\mathcal{A}_{\beta_{t,i}}(s^\star,w), i\in\mathbb{N}_{[1,n_\beta]}$ have all converged to a singleton. However, this is not necessarily the case. For instance, the trivial case where $\mathcal{S}=\{s^\star\}$ and $\mathcal{A}_\beta(s,w)=\Delta_d, \, \forall (s,w)\in\mathcal{S}\times W$ satisfies Assumption III.4, but under these conditions , no learning occurs and, in fact, a robust MPC scheme is recovered. In Section V-D, we will pose an additional assumption on the learning system, which excludes this case, but allows us to show consistency of the learning controller.

Assumption III.5 states that the probability of obtaining an ambiguity set that contains the true conditional distribution (expressed by (8)) increases sufficiently fast (condition (i)). This assumption will be of crucial importance in showing stability (see Section V-C). In addition, it places a lower bound on the convergence rate of the confidence levels (condition (ii)), which is crucial in establishing asymptotic consistency of the scheme (see Section V-D), since it will allow convergence of the ambiguity sets, as we discuss in Remark III.12. To fix ideas, we keep the following example in mind as a suitable choice for the confidence dynamics throughout the article.

Example III.6 (Confidence dynamics). A suitable family of sequences for the confidence levels satisfying Assumption III.5 (assuming $n_{\beta} = 1$ for simplicity²) is obtained as

$$\boldsymbol{\beta}_t = b(1+t)^{-q}, \ t \in \mathbb{N}, \tag{9}$$

with parameters $0 < b \le 1, \, q > 1$. Indeed, as $\, q > 1$, this sequence is summable, and furthermore

$$\lim_{t\to\infty}\frac{-q\log(b(1+t))}{t}=0.$$

Using (9), a straightforward calculation reveals that β_t can be updated recursively:

$$\begin{split} \boldsymbol{\beta}_{t+1}^{-1/q} &= b^{-1/q} (2+t) = b^{-1/q} + \boldsymbol{\beta}_t^{-1/q} \\ \iff \boldsymbol{\beta}_{t+1} &= (b^{-1/q} + \boldsymbol{\beta}_t^{-1/q})^{-q} \\ &= b \boldsymbol{\beta}_t (b^{1/q} + \boldsymbol{\beta}_t^{1/q})^{-q} =: C(\boldsymbol{\beta}_t), \end{split}$$

Thus, it additionally satisfies the requirements of Assumption III.1. \triangle

The learner state s_t will in most practical cases be composed of a data-driven estimator for the transition kernel and some parameter calibrating the size of the ambiguity set, based on statistical information. Indeed, ambiguity sets are very often defined as the set of distributions that lie within some radius from an empirical estimate using a particular distance metric or divergence. We will

²For $n_{\beta} > 1$, the same construction can be repeated element-wise.

refer to such ambiguity sets as *divergence-based* ambiguity sets. For the current setting concerning finitely supported distributions, two notable examples of such divergences are the total variation (TV) metric [11], [48], [49] and the Kullback-Leibler (KL) divergence [7]. In the following section, we show that these divergences can be used to design a learning system satisfying our assumptions, and illustrate that from these two cases, several other divergence-based ambiguity sets can be constructed.

B. Divergence-based ambiguity sets

Our goal is to obtain for each mode w of the Markov chain, a data-driven subset of the probability simplex, containing the wth row of the transition kernel P with high probability. Given a sequence $\widehat{w}_{[1,t]} \in W^t$ of $t \in \mathbb{N}$ samples drawn from the Markov chain, d individual datasets $\widehat{W}_{t,i} := \{\widehat{w}_{k+1} \mid \widehat{w}_k = i, k \in \mathbb{N}_{[1,t]}\}, i \in W$ can be obtained by partitioning the set of observed transitions by the mode they originated in. As such, each $\widehat{W}_{t,i}$ contains t_i i.i.d. draws from the distribution P_i . Ambiguity sets can now be constructed for each individual row i, using concentration inequalities based on the data in $\widehat{W}_{t,i}$.

With this set-up, we now consider the following instance of a learning system .

Definition III.7 (Empirical learner). Let the learner state be composed as $s_t = (\mathbf{vec} \, \hat{P}_t, \gamma_t) \in \mathcal{S} = \Delta_d^d \times [0, 1]^d$, where \hat{P}_t denotes the empirical transition probability matrix at time t, that is,

$$\widehat{P}_{t,ij} = \begin{cases} \frac{1}{t_i} \sum_{w \in \widehat{W}_{t,i}} \mathbf{1}_{w=j} & \textit{if } t_i > 0 \\ \frac{1}{d} & \textit{Otherwise,} \end{cases}$$

and $\gamma_t = (\frac{1}{t_i+1})_{i \in W}$ is a vector containing the inverse of the mode-specific sample sizes.³

For this instance of a learning system, we can now easily derive an explicit characterization of \mathcal{L} .

Example III.8 (Dynamics of the empirical learner). The learner state s_t is composed of $s_t = (\mathbf{vec} \, \widehat{P}_t, \gamma_t)$. For the update of the empirical distribution \widehat{P}_t , note that if $w_t \neq i$, then trivially, $\widehat{P}_{t+1,i:} = \widehat{P}_{t,i:}$. Otherwise, we may use the following well-known construction. Let $e_w \in \mathbb{R}^d$ denote the w'th standard basis vector in \mathbb{R}^d , then

$$\begin{split} \widehat{P}_{t+1,i:} &= \frac{1}{t_i+1} \sum_{w \in \widehat{W}_{t+1,i}} e_w \\ &= \frac{1}{t_i+1} \left(\sum_{w \in \widehat{W}_{t,i}} e_w + e_{w_{t+1}} \right) \\ &= \frac{1}{t_i+1} \left(t_i \widehat{P}_{t,i:} + e_{w_{t+1}} \right) \\ &= (1 - \gamma_{t,i}) \widehat{P}_{t,i:} + \gamma_{t,i} e_{w_{t+1}}. \end{split}$$

Thus, for all $i \in W$, we define

$$\mathcal{L}_{1,i}(\widehat{P}_t, \gamma_t, w_t) := \begin{cases} (1 - \gamma_{t,i}) \widehat{P}_{t,i:} + \gamma_{t,i} e_{w_{t+1}} & \text{if } w_t = i \\ \widehat{P}_{t,i:} & \text{otherwise.} \end{cases}$$

Similarly, $\gamma_{t+1,i}=\gamma_{t,i}$ if $w_{t+1}\neq i$. Otherwise, it follows from the definition of $\gamma_{t,i}$ that $\gamma_{t+1,i}=\frac{1}{t_i+2}\implies \gamma_{t+1,i}=\frac{\gamma_{t,i}}{1+\gamma_{t,i}}$, resulting in

$$\mathcal{L}_{2,i}(\gamma_t, w_{t+1}) := \begin{cases} \frac{\gamma_{t,i}}{1 + \gamma_{t,i}} & \text{if } w_{t+1} = i\\ \gamma_{t,i} & \text{otherwise.} \end{cases}$$
(11)

³The inversion results in simpler updates and renders S robustly positive invariant, i.e, $s_t \in S \implies s_{t+1} \in S$.

Concatenating (10)–(11), we obtain the Markovian update required by Assumption III.1

$$\mathcal{L}(s_t, w_t, w_{t+1}) = ((\mathcal{L}_{1,i}(\widehat{P}_t, \gamma_t, w_t))_{i \in W}, (\mathcal{L}_{2,i}(\gamma_t, w_{t+1}))_{i \in W}).$$

Furthermore, this system satisfies Assumption III.4. Indeed, given ergodicity of the Markov chain (Assumption II.1), the Borel-Cantelli lemma [46, Thm. 4.3] in conjunction with [50, Lem. 6] guarantees that with probability 1, there exists a finite time T, such that for all t>T and for all $i\in W$, it holds that $t_i\geq ct$, where c>0 is a constant depending on specific properties of the Markov chain, and t_i denotes the number of visits to mode i. That is, all modes are visited infinitely often, and as a result, both $\lim_{t\to\infty} \gamma_t=0$ and $\lim_{t\to\infty} \widehat{P}_t=P$, which are indeed fixed points of (10)–(11). \triangle

We can now associate with the newly defined empirical learner the following wide class of ambiguity sets, which take the form of a ball around the empirical estimate in some given statistical divergence.

Definition III.9 (Divergence-based ambiguity set). Consider the empirical learner with state $s_t = (\operatorname{vec} \widehat{P}_t, \gamma_t)$. We say that an ambiguity set $\mathcal{A}_{\beta_t}(s_t, w)$ is a divergence-based ambiguity set if it can be expressed in the form

$$\mathcal{A}_{\beta_t}(s_t, w) := \{ p \in \Delta_d \mid \mathcal{D}(\widehat{P}_{t, w}, p) \le r(\gamma_{t, w}^{-1} - 1, \beta_t) \}, \forall w \in W$$

where $\mathcal{D}: \Delta_d \times \Delta_d \to \mathbb{R}_+$ is some statistical divergence—and $r: \mathbb{R}_+ \times [0,1] \to \mathbb{R}_+$ is a given function that returns a radius, given a sample size and a confidence level.

Statistically meaningful values for the radius r under different choices of divergences can be obtained using the following standard results.

Proposition III.10 (Concentration inequalities). Let $p \in \Delta_d$ denote a distribution on the probability simplex and $\widehat{p} = \frac{1}{m} \sum_{t=0}^{m-1} (\mathbf{1}_{wt=i})_{i=1}^d$ the empirical distribution based on m i.i.d. draws $w_t \sim p$. Then, $\mathbb{P}\big[(\frac{1}{2} \| p - \widehat{p} \|_1)^2 > r^{\mathrm{TV}}(m, \beta) \big] \leq \beta$, with

$$r^{\text{TV}}(m,\beta) = \frac{d\log 2 - \log \beta}{2m}.$$
 (12)

Similarly, it holds that $\mathbb{P}[\mathcal{D}_{\mathrm{KL}}(\widehat{p},p) > r^{\mathrm{KL}}(m,\beta)] \leq \beta$, with

$$r^{\text{KL}}(m,\beta) = \frac{d \log m - \log \beta}{m},\tag{13}$$

where $\mathcal{D}_{\mathrm{KL}}(p,q) := \sum_{i=1}^{d} p_i \log \frac{p_i}{q_i}$ denotes the KL divergence from q to p.

The bound on the TV distance (12) is known as the Bretagnolle-Huber-Carol inequality [51, Thm. A.6.6].

Remark III.11. Expression (13) for the KL radius is a well-known result from the field of information theory, obtained through the so-called method-of-types [52], [53]. A slight improvement can be obtained by replacing $d\log m$ by $\log \binom{m+d-1}{d-1}$. Moreover, in [54], an even sharper result for (13) is derived. In fact, this improved concentration bound in the KL divergence was used in the same work to improve upon the TV concentration bound (12) for $\frac{m}{d} \ll 1$, using Pinsker's inequality [55], which relates the TV distance between distributions $p,q \in \Delta_d$ to the KL divergence as $\|p-q\|_1^2 \leq 2\mathcal{D}_{\mathrm{KL}}(p,q)$. Of course, these improved bounds can be readily used in practice to replace those in proposition III.10. However, for the theoretical discussion, these modifications are inconsequential. For this reason, we opt to develop the ideas for the simpler, more commonly used forms.

Besides Pinsker's inequality, there exist several other inequalities relating different statistical divergences (see for instance [56] for a comprehensive overview). Based on these relations, one can derive

OVERVIEW OF FIND EITHES OF COMMON BIVETGENCE BASED AMBIGGITT SETS.						
Divergence	$\mathcal{D}(\widehat{p},p)$	radius $r(m, \beta)$	Conic representation			
Total variation (TV) Kullback-Leibler (KL)	$egin{aligned} \ p-\widehat{p}\ _1 \ \mathcal{D}_{ ext{KL}}(\widehat{p},p) \end{aligned}$	$rac{2\sqrt{r^{ ext{TV}}(m,eta)}}{r^{ ext{KL}}(m,eta)}$	Linear Exponential			
Jensen-Shannon (JS)	$1/2\left(\mathcal{D}_{\mathrm{KL}}\left(\widehat{p},rac{p+\widehat{p}}{2} ight)+\mathcal{D}_{\mathrm{KL}}\left(\widehat{p},rac{p+\widehat{p}}{2} ight) ight)$	$rac{1}{2}r^{ ext{KL}}(m,eta)$	Exponential			
(Squared) Hellinger (H)	$\sum_{i}\left(\sqrt{\widehat{p}_{i}}-\sqrt{\widehat{p}_{i}} ight)^{2}$	$r^{ ext{KL}}(m,eta)$	Quadratic			

TABLE I OVERVIEW OF PROPERTIES OF COMMON DIVERGENCE-BASED AMBIGUITY SETS

from proposition III.10 several divergence-based ambiguity sets defined through other statistical divergences; For instance, since the squared Hellinger divergence is upper bounded by the KL divergence, (13) can be used as a radius for Hellinger divergence-based ambiguity sets. A summary of the resulting radii is provided in Table I. The rightmost column in this table refers to the conic representation of the induced ambiguity sets (cf. (23)), which determines the complexity of the resulting optimal control problems (see Appendix A for more details). Other works that have used these divergences (which belong to the class of ϕ -divergences) for distributionally robust optimization are [57]–[59]. In these works, however, the radii are either selected as a tuning parameter or calibrated using asymptotic arguments, leading to approximate ambiguity sets, which satisfy the coverage condition (8) only as the sample size tends to infinity. By contrast, the radii given in Table I are valid for any sample size.

We conclude the section by proposing a useful extension of the learner state in the case of divergence-based ambiguity sets.

Remark III.12 (Radius as part of the learner state). For divergencebased ambiguity sets, it is often convenient to augment the learner state $s_t = (\mathbf{vec}\,\hat{P}_t, \gamma_{t,i})$ with the computed radii $r_{t,i} := r(\gamma_{t,i}^{-1} - \gamma_{t,i}^{-1})$ $(1, \beta_t)_{i \in W}$, for which the recursive update is obtained simply by composition of r with the previously designed \mathcal{L} and C. It can be easily verified using Proposition III.10 that this quantity also converges to the fixed point $\lim_{t\to\infty} r_{t,i} = 0$, $\forall i$. Indeed, Recall from Example III.8 that $\gamma_{t,i}^{-1} - 1 \sim t$ as $t \to \infty$. Using a radius function r based on either (12) or (13), we obtain

$$\lim_{t \to \infty} r_{t,i} = \lim_{t \to \infty} r(t, \beta_t) = \lim_{t \to \infty} \frac{-\log \beta_t}{t} = 0,$$

where the last equality follows from Assumption III.5.

IV. LEARNING MODEL PREDICTIVE CONTROL

Given a learning system satisfying Assumption III.1, we define the augmented state $y_t = (x_t, s_t, \boldsymbol{\beta}_t) \in \mathcal{Y} := \mathbb{R}^{n_x} \times \mathcal{S} \times \mathcal{I}$, which evolves over time according to the dynamics

$$y_{t+1} = \tilde{f}(y_t, w_t, u_t, w_{t+1}) := \begin{bmatrix} f(x_t, u_t, w_{t+1}) \\ \mathcal{L}(s_t, \beta_t, w_t, w_{t+1}) \\ C(\beta_t) \end{bmatrix}, \quad (14)$$

with $w_{t+1} \sim P_{w_t}$; for $t \in \mathbb{N}$. Furthermore, it will be convenient to define the process $z_t = (y_t, w_t) \in \mathcal{Z} := \mathcal{Y} \times W$. Consequently, the objective is now to obtain a feedback law $\kappa: \mathcal{Z} \to \mathbb{R}^{n_u}$. To this end, we will formulate a DR counterpart to the stochastic OCP (7), in which the expectation operator in the cost and the conditional probabilities in the constraint will be replaced by operators that account for ambiguity in the involved distributions.

A. Ambiguity and risk

In order to reformulate the cost function (6), we first introduce an ambiguous conditional expectation operator, leading to a formulation akin to the Markovian risk measures utilized in [32], [60]. Consider a function $\xi: \mathbb{Z} \times W \to \overline{\mathbb{R}}$, defining a stochastic process $(\xi_t)_{t \in \mathbb{N}} =$ $(\xi(z_t, w_{t+1}))_{t \in \mathbb{N}}$ on (Ω, F, \mathbb{P}) , and suppose that the augmented state $z_t = z = (x, s, \boldsymbol{\beta}, w)$ is given. Let $\beta \in [0, 1]$ denote an arbitrary component of β . The *ambiguous* conditional expectation of $\xi(z, v)$, given z is then

$$\rho_{s,w}^{\beta}[\xi(z,v)] := \max_{p \in \mathcal{A}_{\beta}(s,w)} \mathbb{E}_{p}[\xi(z,v)|z]$$

$$= \max_{p \in \mathcal{A}_{\beta}(s,w)} \sum_{v \in W} p_{v}\xi(z,v).$$
(15)

Linear

Trivially, it holds that if the w'th row of the transition matrix lies in the corresponding ambiguity set, i.e., $P_{w:} \in \mathcal{A}_{\beta}(s, w)$, then

$$\rho_{s,w}^{\beta}[\xi(z,v)] \ge \mathbb{E}_{P_w:}[\xi(z,v) \mid z]$$

$$= \sum_{v \in W} P_{wv}\xi(z,v).$$
(16)

Note that the function $\rho_{s,w}^{\beta}$ defines a coherent risk measure [34, Sec. 6.3]. We say that $\rho_{s,w}^{\beta}$ is the risk measure *induced by* the ambiguity set $\mathcal{A}_{\beta}(s, w)$.

A similar construction can be carried out for the chance constraints (5). We robustify the average value-at-risk with respect to the reference distribution, defining

$$\bar{\rho}_{s,w}^{\beta,\widehat{\alpha}}[\xi(z,v)] := \max_{p \in \mathcal{A}_{\beta}(s,w)} \mathsf{AV}@\mathsf{R}_{\widehat{\alpha}}^{p}[\xi(z,v) \mid z] \leq 0. \tag{17}$$

The function $\bar{\rho}_{s,w}^{\beta,\widehat{\alpha}}$ in turn defines a coherent risk measure. Note that we have replaced the AV@R parameter α by $\widehat{\alpha}$. The reason for this is that the ambiguity set only contains the true distribution with high probability. Considering this fact, it is natural to expect that α needs to be tightened to some extent in order to ensure that the original chance constraint remains satisfied. We make this precise in the following result.

Proposition IV.1. Let $\beta, \alpha \in [0,1]$, be given values with $\beta < \alpha$. Consider the random variable $s:\Omega\to\mathcal{S}$, denoting an (a priori unknown) learner state satisfying Assumption III.1, i.e., $\mathbb{P}[P_w] \in$ $\mathcal{A}_{\beta}(s,w)] \geq 1-\beta$. If the parameter $\widehat{\alpha}$ is chosen to satisfy $0 \leq \widehat{\alpha} \leq \beta$ $\frac{\alpha - \beta}{1 - \beta} \le 1$, then, for an arbitrary function $g : \mathcal{Z} \times W \to \mathbb{R}$, the following implication holds:

$$\bar{\rho}_{s,w}^{\beta,\widehat{\alpha}}[g(z,v)] \leq 0, \text{ a.s.} \Rightarrow \mathbb{P}[g(z,v) \leq 0 \mid x,w] \geq 1-\alpha. \tag{18}$$

Proof. If $\bar{\rho}_{s,w}^{\beta,\widehat{\alpha}}[q(z,v)] < 0$, a.s., then (4) and (17) imply that

$$\mathbb{P}[g(z,v) \leq 0 \mid x, w, P_{w:} \in \mathcal{A}_{\beta}(s,w)] \geq 1 - \widehat{\alpha}, \text{a.s.}$$

Wasserstein* (W) $\min_{\Pi \in \mathbf{R}_{+}^{d \times d}} \left\{ \sum_{i,j \in W} \Pi_{ij} K_{ij} \mid \Pi \mathbf{1}_{d} = p, \Pi^{\top} \mathbf{1}_{d} = \widehat{p} \right\} \quad \max_{i,j \in W} K_{ij} \sqrt{r^{\mathrm{TV}}(m,\beta)}$ *Assumes W is a metric space. $K \in \mathbf{R}^{d \times d}$ is a symmetric distance kernel with $K_{ij} = \mathbf{dist}(i,j), \ \forall i,j \in W$.

Therefore,

$$\mathbb{P}[g(z,v) \leq 0 \mid x,w]$$

$$\geq \mathbb{P}[g(z,v) \leq 0 \mid x,w,P_{w:} \in \mathcal{A}_{\beta}(s,w)]\mathbb{P}[P_{w:} \in \mathcal{A}_{\beta}(s,w)]$$

$$\geq (1-\widehat{\alpha})(1-\beta).$$

Requiring that $(1-\widehat{\alpha})(1-\beta) \geq (1-\alpha)$ then immediately yields the sought condition. \Box

Notice that the implication (18) in Proposition IV.1 provides an a priori guarantee, since the learner state is considered to be random. In other words, the statement is made before the data is revealed. Indeed, for a given learner state s and mode w, the ambiguity set $\mathcal{A}_{\beta}(s,w)$ is fixed and therefore, the outcome of the event $E=\{P_w: \in \mathcal{A}_{\beta}(s,w)\}$ is determined. Whether (18) then holds for these fixed values, depends on the outcome of E. This is naturally reflected through the above condition on $\widehat{\alpha}$, which implies that $\widehat{\alpha} \leq \alpha$, and thus tightens the chance constraints that are imposed conditioned on a fixed s. Hence, the possibility that for this particular s, the ambiguity set may not include the conditional distribution, is accounted for. This tightening can be mitigated by decreasing β , at the cost of a larger ambiguity set. A more detailed study of this trade-off is left for future work.

B. Distributionally robust model predictive control

We are now ready to describe the DR counterpart to the OCP (7), which, when solved in receding horizon fashion, yields the proposed learning MPC scheme.

Consider a given augmented state $z=(x,s,\boldsymbol{\beta},w)\in\mathcal{Z}$. Hereafter, we will assume that $\boldsymbol{\beta}=(\beta,\overline{\beta})$, where component β is related to the cost function and $\overline{\beta}$ is reserved for the constraints.

We use (17) to define the DR set of feasible inputs $\widehat{\mathcal{U}}(z)$ in correspondence to (5), as

$$\widehat{\mathcal{U}}(z) = \left\{ u \in U \, \middle| \, \bar{\rho}_{s,w}^{\overline{\beta},\widehat{\alpha}}[g(x,u,w,v)] \le 0 \right\}. \tag{19}$$

Remark IV.2. The parameter $\widehat{\alpha}$ remains to be chosen in relation to the confidence levels $\pmb{\beta}$ and the original violation rates α . In light of Proposition IV.1, $\widehat{\alpha} = \frac{\alpha - \overline{\beta}}{1 - \overline{\beta}}$ yields the least conservative choice. This choice is valid as long as it is ensured that $\overline{\beta} < \alpha$.

Using (15), we express the DR cost of a policy $\pi = (\pi_k)_{k=0}^{N-1}$ as

$$\begin{split} \widehat{V}_{N}^{\pi}(z) &:= \ell(x_{0}, u_{0}, w_{0}) + \rho_{s_{0}, w_{0}}^{\beta_{0}} \left[\ell(x_{1}, u_{1}, w_{1}) \right. \\ &+ \rho_{s_{1}, w_{1}}^{\beta_{1}} \left[\cdots + \rho_{s_{N-2}, w_{N-2}}^{\beta_{N-2}} \left[\ell(x_{N-1}, u_{N-1}, w_{N-1}) \right. \\ &+ \rho_{s_{N-1}, w_{N-1}}^{\beta_{N-1}} \left[\widehat{V}_{\mathbf{f}}(x_{N}, s_{N}, \boldsymbol{\beta}_{N}, w_{N}) \right] \right] \dots \right] \right], \quad (20 \end{split}$$

where $z_0=z, z_{k+1}=\widetilde{f}(z_k,u_k,w_{k+1})$ and $u_k=\pi_k(z_k)$, for all $k\in\mathbb{N}_{[0,N-1]}$. In Section V, conditions on the terminal cost $\widehat{V}_{\mathrm{f}}:\mathcal{Z}\to\overline{\mathbb{R}}_+:(x,s,\pmb{\beta},w)\mapsto V_{\mathrm{f}}(x,w)+\delta_{\widehat{\mathcal{X}}_{\mathrm{f}}}(x,s,\pmb{\beta},w)$ and its domain are provided in order to guarantee recursive feasibility and stability of the MPC scheme defined by the following OCP.

Definition IV.3 (DR-OCP). Given an augmented state $z \in \mathcal{Z}$, the optimal cost of the distributionally robust optimal control problem (DR-OCP) is

$$\widehat{V}_N(z) = \min \widehat{V}_N^{\pi}(z) \tag{21a}$$

subject to

$$(x_0, s_0, \boldsymbol{\beta}_0, w_0) = z, \ \pi = (\pi_k)_{k=0}^{N-1},$$
 (21b)

$$z_{k+1} = (\tilde{f}(z_k, \pi_k(z_k), w_{k+1}), w_{k+1}), \tag{21c}$$

$$\pi_k(z_k) \in \widehat{\mathcal{U}}(z_k), \ \forall w_{[0,k]} \in W^k,$$
 (21d)

for all $k \in \mathbb{N}_{[0,N-1]}$. We denote by $\widehat{\Pi}_N(z)$ the corresponding set of minimizers.

Remark IV.4. Note that the definition of \widehat{V}_f implicitly imposes the terminal constraint $z_N \in \widehat{\mathcal{X}}_f$, a.s.

We now define the learning MPC law analogously to the stochastic case as

$$\widehat{\kappa}_N(z) = \widehat{\pi}_0^*(z), \tag{22}$$

where $(\widehat{\pi}_k^{\star}(z))_{k=0}^{N-1} \in \widehat{\Pi}_N(z)$. At every time t, the learning MPC scheme thus consists of repeatedly (i) solving (21) to obtain a control action $u_t = \widehat{\kappa}_N(z_t)$ and applying it to the system (1); (ii) observing the outcome of $w_{t+1} \in W$ and the corresponding next state $x_{t+1} = f(x_t, u_t, w_{t+1})$; and (iii) updating the learner state $s_{t+1} = \mathcal{L}(s_t, w_t, w_{t+1})$ and the confidence levels $\beta_{t+1} = C(\beta_t)$, gradually decreasing the size of the ambiguity sets.

Note that in its general form, (21) is a non-smooth, infinite-dimensional optimization problem. However, provided that the involved risk measures are *conic risk measures* (as defined by Definition IV.5), problem (21) can be reformulated as a finite-dimensional, smooth nonlinear program.

Definition IV.5 (Conic risk measure [44]). We say that an ambiguity set $A \subseteq \Delta_d$ is conic representable if it can be written in the form

$$\mathcal{A} = \{ p \in \Delta_d \mid \exists \nu : Ep + F\nu \preccurlyeq_{\mathcal{K}} b \}, \tag{23}$$

with matrices E, F and vector b of suitable dimensions, and a proper cone K. The coherent risk measure induced by a conic representable ambiguity set is called a conic risk measure.

Since ambiguity sets inducing coherent risk measures are convex by construction, many classes of ambiguity sets can be represented using conic inequalities. For completeness, we state the conic representations for the ambiguity sets summarized in Table I, as well as the reformulation of (21) in Appendix A and B, respectively.

V. THEORETICAL ANALYSIS

A. Dynamic programming

To facilitate theoretical analysis of the proposed MPC scheme, we follow an approach similar to [32] and represent (21) as a dynamic programming recursion. We define the Bellman operator \mathbf{T} as $\mathbf{T}(\widehat{V})(z) := \min_{u \in \widehat{\mathcal{U}}(z)} \ell(x,u,w) + \rho_{s,w}^{\beta}[\widehat{V}(\widetilde{f}(z,u,v),v)],$ where $z = (x,s,\beta,w) \in \mathcal{Z},$ with $\boldsymbol{\beta} = (\beta,\overline{\beta})$ as before, are fixed quantities and $v \sim P_{w}$. We denote by $\mathbf{S}(\widehat{V})(z)$ the corresponding set of minimizers. The optimal cost \widehat{V}_N of (21) is obtained through the iteration,

$$\widehat{V}_k = \mathbf{T}\,\widehat{V}_{k-1}, \ \widehat{V}_0 = \widehat{V}_f, \ k \in \mathbb{N}_{[1,N]}.$$
(24)

Similarly, $\widehat{\mathcal{Z}}_k := \operatorname{\mathbf{dom}} \widehat{V}_k$ is given recursively by

$$\widehat{\mathcal{Z}}_k = \left\{ z \, \middle| \, \exists u \in \widehat{\mathcal{U}}(z) : (\widetilde{f}(z,u,v),v) \in \widehat{\mathcal{Z}}_{k-1}, \, \forall v \in W \right\}.$$

Now consider the stochastic closed-loop system

$$y_{t+1} = \tilde{f}^{\widehat{\kappa}_N}(z_t, w_{t+1}) := \tilde{f}(z_t, \widehat{\kappa}_N(z_t), w_{t+1}),$$
 (25)

where $\hat{\kappa}_N(z_t) \in \mathbf{S}(\hat{V}_{N-1})(z_t)$ is an optimal control law obtained by solving the DR-OCP of horizon N in receding horizon.

B. Constraint satisfaction and recursive feasibility

In order to show existence of $\widehat{\kappa}_N \in \mathbf{S} \, \widehat{V}_{N-1}$ at every time step, Proposition V.4 will require that $\widehat{\mathcal{X}}_{\mathbf{f}}$ is a robust control invariant set. We define robust control invariance for the augmented control system under consideration as follows.

Definition V.1 (Robust control invariance). A set $\mathcal{R} \subseteq \mathcal{Z}$ is an RCI set for the system (14) if for all $z \in \mathcal{R}$, $\exists u \in \widehat{\mathcal{U}}(z)$ such that $(\widetilde{f}(z,u,v),v) \in \mathcal{R}, \forall v \in W$. Similarly, \mathcal{R} is a robust positive invariant (RPI) set for the closed-loop system (25) if for all $z \in \mathcal{R}$, $(\widetilde{f}^{\widehat{\kappa}_N}(z,v),v) \in \mathcal{R}, \forall v \in W$.

Since $\widehat{\mathcal{U}}$ consists of conditional risk constraints, our definition of robust invariance provides a distributionally robust counterpart to the notion of *stochastic* robust invariance in [61]. This notion is less conservative than the following, more classical notation of robust invariance.

Definition V.2 (Classical robust control invariance). A set $\mathcal{R}_x \subseteq \mathbb{R}^{n_x} \times W$ is RCI for system (1) in the classical sense if for all $x \in \mathcal{R}_x$,

$$\exists u : g(x, u, w, v) \le 0, \ f(x, u, v) \in \mathcal{X}_{f}(v), \ \forall v \in W.$$
 (26)

In fact, for any set \mathcal{R}_x as in Definition V.2, the set $\mathcal{R}_x \times \mathcal{S} \times \mathcal{I} \times W$ is covered by Definition V.1, as illustrated in Example V.3. On the other hand, our notion of robust control invariance is stricter than that of *uniform control invariance* considered in [32], which only requires successor states to remain in the invariant set for modes v in the *cover* of the given mode w, i.e., the set of modes v for which $P_{wv} > 0$. This flexibility is not available in the current setting, as the transition kernel is assumed to be unknown, so the cover of a mode cannot be determined with certainty.

Example V.3 (Classical robust invariant set). Suppose that the terminal constraint set \mathcal{X}_f of the nominal problem is a robust control invariant set in the classical sense and define for convenience $\mathcal{X}_f(w) := \{x \mid (x,w) \in \mathcal{X}_f\}$. Then, if $\widehat{\mathcal{X}}_f$ is chosen such that $\widehat{\mathcal{X}}_f(w) := \{y \mid (y,w) \in \widehat{\mathcal{X}}_f\} = \mathcal{X}_f(w) \times \mathcal{S} \times \mathcal{I}$, $\widehat{\mathcal{X}}_f$ is RCI for the augmented system (14) according to Definition V.1. Indeed, since $\mathsf{AV}@\mathsf{R}^p_\alpha[g(x,u,w,v)] \leq \max_v g(x,u,w,v)$ for all $\alpha \in [0,1]$ and $p \in \Delta_d$, (26) implies that for all $z \in \widehat{\mathcal{X}}_f$, there exists $u \in \widehat{\mathcal{U}}(z)$, such that $\widehat{f}(z,u,v) \in \widehat{\mathcal{X}}_f(v)$.

Proposition V.4 (Recursive feasibility). If $\widehat{\mathcal{X}_f}$ is an RCI set for (14), then (21) is recursively feasible. That is, feasibility of DR-OCP (21) for some $z \in \mathcal{Z}$, implies feasibility for $z^+ = (\widehat{f}^{\widehat{\kappa}_N}(z,v),v)$, for all $v \in W, N \in \mathbb{N}_{>0}$.

Proof. The proof follows from a straightforward inductive argument on the prediction horizon N. We first show that if $\widehat{\mathcal{X}}_f$ is RCI, then so is $\widehat{\mathcal{Z}}_N$. This is done by induction on the horizon N of the OCP. Base case (N=0). Trivial, since $\widehat{\mathcal{Z}}_0=\widehat{\mathcal{X}}_f$.

Induction step $(N\Rightarrow N+1)$. Suppose that for some $N\in\mathbb{N}$, $\widehat{\mathcal{Z}}_N$ is RCI for (14). Then, by definition of $\widehat{\mathcal{Z}}_{N+1}$, there exists for each $z\in\widehat{\mathcal{Z}}_{N+1}$, a nonempty set $\widehat{\mathcal{U}}_N^\star(z)\subseteq\widehat{\mathcal{U}}(z)$ such that for every $u\in\widehat{\mathcal{U}}_N^\star(z)$ and for all $v\in W$, it holds that $z^+\in\widehat{\mathcal{Z}}_N$, where $z^+=\widehat{f}(z,u,v)$. Furthermore, the induction hypothesis $(\widehat{\mathcal{Z}}_N)$ is RCI), implies that there also exists a $u^+\in\widehat{\mathcal{U}}(z^+)$ such that $\widehat{f}(z^+,u^+,v^+)\in\widehat{\mathcal{Z}}_N(v^+), \forall v^+\in W$. Therefore, z^+ satisfies the conditions defining $\widehat{\mathcal{Z}}_{N+1}$. In other words, $\widehat{\mathcal{Z}}_{N+1}$ is RCI.

The claim follows from the fact that for any N>0 and $z\in\widehat{\mathcal{Z}}_N$, $u=\widehat{\kappa}_N(z)\in\mathbf{S}(\widehat{V}_{N-1})(z)\subseteq\widehat{\mathcal{U}}_{N-1}^\star(z)$, as any other choice of u would yield infinite cost in the definition of the Bellman operator. \square

Corollary V.5 (Chance constraint satisfaction). If the conditions for Proposition V.4 hold, then by Proposition IV.1, the stochastic process $(z_t)_{t\in\mathbb{N}} = (x_t, s_t, \boldsymbol{\beta}_t, w_t)_{t\in\mathbb{N}}$ satisfying dynamics (25) satisfies the nominal chance constraints

$$\mathbb{P}[g(x_t, \widehat{\kappa}_N(z_t), w_{t+1}) > 0 \mid x_t, w_t] < \alpha,$$

a.s., for all $t \in \mathbb{N}$.

We conclude this section by emphasizing that although the MPC scheme guarantees closed-loop constraint satisfaction, it does so while being less conservative than a fully robust approach, which is recovered by taking $\mathcal{A}_{\beta}(s,w) = \Delta_d$ for all $(s,w,\beta) \in \mathcal{S} \times W \times [0,1]$. It is apparent from (17) and (19), that for all other choices of the ambiguity set, the set of feasible control actions will be larger (in the sense of set inclusion).

C. Stability

In this section, we will provide sufficient conditions on the control setup under which the origin is mean-square stable (MSS) for (25), i.e., $\lim_{t\to\infty} \mathbb{E}[\|x_t\|^2] = 0$ for all x_0 in some specified compact set containing the origin.

Our main stability result, stated in Theorem V.7, hinges in large on the following lemma, which relates *risk-square stability* [32, Thm. 6] of the origin for the autonomous system (25) (with respect to a statistically determined ambiguity set) to stability in the mean-square sense (with respect to the true distribution).

Lemma V.6 (Distributionally robust MSS condition). Suppose that Assumption III.5 holds and that there exists a nonnegative, proper function $V: \mathcal{Z} \to \overline{\mathbb{R}}_+$, such that (i) $\operatorname{dom} V$ is RPI for (25) and $\operatorname{dom} V(\cdot, s, \beta, w)$ is compact and contains the origin for all $(s, \beta, w): \operatorname{dom} V(\cdot, s, \beta, w) \neq \emptyset$ (ii) $\rho_{s,w}^{\beta}[V(\tilde{f}^{\widehat{\kappa}_N}(z, v), v)] - V(z) \leq -c||x||^2$, for some c > 0, for all $z \in \operatorname{dom} V$; (iii) V is uniformly bounded on its domain. Then, $\lim_{t\to\infty} \mathbb{E}[||x_t||^2] = 0$ for all $z_0 \in \operatorname{dom} V$, where $(z_t)_{t\in\mathbb{N}} = (x_t, s_t, \beta_t, w_t)_{t\in\mathbb{N}}$ is the stochastic process governed by dynamics (25).

Theorem V.7 (MPC stability). Suppose that Assumptions II.3 and III.5 are satisfied and the following statements hold. (i) $\mathbf{T}\,\widehat{V}_{\mathrm{f}} \leq \widehat{V}_{\mathrm{f}}; \ (ii) \quad c\|x\|^2 \leq \ell(x,u,w) \ \text{for some} \ c > 0, \ \text{for all} \ z = (x,s,\beta,w) \in \operatorname{dom}\widehat{V}_N \ \text{and all} \ u \in \widehat{\mathcal{U}}(z); \ (iii) \quad \widehat{V}_N \ \text{is locally bounded on its domain. Then, the origin is MSS for the MPC-controlled system (25), over all RPI sets <math>\overline{\mathcal{Z}} \subseteq \operatorname{dom}\widehat{V}_N \ \text{such that} \ \text{for all} \ (s,\beta,w): (x,s,\beta,w) \in \overline{\mathcal{Z}}, \ \text{the projection} \ \{x \mid (x,s,\beta,w) \in \overline{\mathcal{Z}}\} \ \text{is compact and contains the origin.}$

Proof. The proof is along the lines of that of [32, thm. 6] and shows that \hat{V}_N satisfies the conditions of Lemma V.6. Details are in the Appendix.

The results in this section indicate that after an appropriate choice of the learning system, the thusly defined risk measures can be used to design an MPC controller using existing techniques (e.g., those presented in [32]). Corresponding stability guarantees (assuming known transition probabilities) then translate directly into stability guarantees under an ambiguously estimated transition kernel.

D. Out-of-sample bounds and consistency

We now turn our attention to analyzing the value function of the DR-OCP in relation to the nominal (stochastic) OCP. We will show that under quite general assumptions, the former provides an upper bound to the latter with high probability (Theorem V.9). Furthermore, under appropriate constraint qualifications, we will show that the optimal value of the DR-OCP converges to that of the nominal problem as the sample size increases, see Theorem V.13. In the particular case where the constraints do not depend on the distribution, we can relax the constraint qualification to obtain a similar result. We include this as a separate statement, as it permits a more direct and illustrative proof using dynamic programming.

Given an arbitrary state-mode pair (x, w), initial value of the learning state s_0 and confidence β_0 , the stochastic process defined by the optimal value of the DR-OCP (21), i.e., $\widehat{V}_N^{(t)}(x,w) := \widehat{V}_N(x,s_t,\pmb{\beta}_t,w), \ t \in \mathbb{N}$ serves as a sequential approximation of the optimal value $V_N(x, w)$ of the horizon-N nominal OCP (7). This section will establish sufficient conditions under which $\widehat{V}_N^{(t)}$ bounds V_N from above, and for which it converges to V_N^{\star} almost surely—a property which we refer to as asymptotic consistency. The former guarantee will provide a performance certificate in the sense that the true optimal cost (under full knowledge of the distribution) will be no worse than the cost predicted by solving its DR counterpart. Of course, this guarantee is also provided by a robust (minimax) scheme (obtained by taking $A_{\beta_t} \equiv \Delta_d, \forall t$). However, such an approach is non-adaptive and therefore lacks consistency. On the other hand, a sample-average approximation (in which the ambiguity set is replaced by a singleton containing only the empirical distribution) may under similar conditions be consistent, but it provides no safety guarantees/performance bounds.

Below, we denote $\mathcal{X}_f(w) = \{x \mid (x, w) \in \mathcal{X}_f\}$ and similarly $\widehat{\mathcal{X}_{\mathrm{f}}}(w) = \{y \mid (y, w) \in \widehat{\mathcal{X}_{\mathrm{f}}}\}.$ We will also pose the following assumptions in the remainder of the section.

Assumption V.8. (i) The risk levels $\hat{\alpha}_t$ are chosen according to the upper bound of Proposition IV.1, i.e., $\widehat{\alpha}_t = \frac{\alpha - \overline{\beta}_t}{1 - \overline{\beta}_t}$ and $\overline{\widehat{\beta}_t} < \overline{\beta}_t$

(ii) $\widehat{\mathcal{X}_f}$ is constructed in relation to the original problem such that for all $w \in W$, $\widehat{\mathcal{X}_f}(w) = \mathcal{X}_f(w) \times \mathcal{S} \times \mathcal{I}$, and \mathcal{X}_f is RCI for system (1) in the sense of Definition V.2.

Theorem V.9 (Performance guarantee). Suppose that assumption V.8 holds. Then, for any initial learner state $s_0 = s \in S$ and any initial confidence level $\beta_0 = \beta \in \mathcal{I}$,

(i) the value function of the DR-OCP of horizon $N \geq 0$ asymptotically upper bounds the true value function. That is,

$$\mathbb{P}[\widehat{V}_{N}^{(t)}(x, w) \ge V_{N}(x, w), \forall (x, w) \in \text{dom } V_{N}] \ge 1 - \gamma_{N}^{(t)}, \quad (27)$$

for all $t \in \mathbb{N}$, with $\gamma_N^{(t)} = d \sum_{k=t}^{t+N} \|\beta_k\|_1$. (ii) If, furthermore, \mathcal{A}_β is selected such that

$$\beta' \le \beta \implies \mathcal{A}_{\beta}(s, w) \subseteq \mathcal{A}_{\beta'}(s, w), \quad \forall (s, w) \in \mathcal{S} \times W, \quad (28)$$

then, (27) holds with $\gamma_N^{(t)} = d \sum_{k=t}^{t+N} \|\boldsymbol{\beta}_k\|_{\infty}$.

Theorem V.9 guarantees that with high probability, the DR value function provides an upper bound for the value function under full knowledge of the distribution. The corresponding violation rate $\gamma_N^{(t)}$ can be tuned using the user-specified confidence levels β_t .

Note that the violation rate $\gamma_N^{(t)}$ increases with the prediction horizon. This is to be expected, since we essentially require the ambiguity set to contain the true switching distribution for all predicted time steps, which becomes increasingly difficult as the horizon length increases. However, due to the summability of the confidence levels β_t (cf. Assumption III.5), the violation rate $\gamma_N^{(t)}$ will converge to a finite value as $N \to \infty$. Similarly, as $t \to \infty$ for fixed $N, \gamma_N^{(t)}$ converges to zero at a summable rate. We will use this fact in corollary V.10 to obtain a stronger guarantee asymptotically.

Before stating the asymptotic extension of Theorem V.9, we briefly highlight the sharper bound for $\gamma_N^{(t)}$ stated in Theorem V.9-(ii). This result requires that for a given learner state s, the size of the ambiguity set scales monotonically with the required confidence level. This is satisfied for the described divergence-based ambiguity sets in Table I. Indeed, the center of the divergence balls are given by the empirical distribution and therefore independent of the confidence level β . The radii, by proposition III.10, are monotone decreasing functions of β . Thus, the intersection of a collection of such ambiguity sets is equal to the ambiguity set with the largest value of β (and thus, the smallest radius).

Corollary V.10. Under the same conditions as theorem V.9, we have with probability one that,

$$\widehat{V}_{N}^{(t)}(x,w) \ge V_{N}(x,w)$$
 for all sufficiently large $t,$ (29)

for all $(x, w) \in \operatorname{dom} V_N$.

Proof. For fixed $(x, w) \in \operatorname{dom} V_N$, theorem V.9 guarantees that $\begin{array}{l} \mathbb{P}[\widehat{V}_N^{(t)}(x,w) < V_N(x,w)] \leq \gamma_N^{(t)}, \text{ where due to assumption III.5,} \\ \sum_{t=0}^{\infty} \gamma_N^{(t)} = d \sum_{k=0}^N \sum_{t=0}^{\infty} \|\boldsymbol{\beta}_{t+k}\|_1 < \infty. \text{ The claim then follows from the Borel-Cantelli lemma [46, Thm. 4.3].} \end{array}$

Having established a performance bound on the true cost, will now demonstrate consistency of the method, starting with the special case where the constraints are independent of the learner state (Theorem V.12), before tackling the general case in Theorem V.13. To this end, we make the following assumption on the learner state and the corresponding ambiguity set.

Assumption V.11 (Ambiguity decrease). There exists a sequence $\{\delta_t\}_{t\in\mathbb{N}}$ with $\lim_{t\to\infty} \delta_t = 0$, such that

$$\sup_{p,q \in \mathcal{A}_{\beta_{t,i}}(s_t,w)} \|p-q\|_1 \le \delta_t \quad a.s., \qquad \forall w \in W, \, \forall i \in \mathbb{N}_{[1,n_{\beta}]},$$

Assumption V.11 states that the ambiguity sets "shrink" to a singleton with probability one. Since the ambiguity is expected to decrease as more information is observed, this is a rather natural assumption, which is satisfied by most classes of ambiguity sets, such as the ones discussed in Section III (cf. Remark III.12).

Theorem V.12 (Asymptotic consistency with hard constraints). Suppose that all constraints are hard constraints, i.e., $\alpha = 0$, so that $\widehat{\mathcal{U}}(z) = \mathcal{U}(x,w)$ for all $z = (x,s,\beta,w)$. Then, for any state-mode pair $(x, w) \in \operatorname{dom} V_N$, any initial learner state $s_0 = s \in \mathcal{S}$ and any initial confidence level $\beta_0 = \beta \in \mathcal{I}$, the optimal cost of the DR-OCP of horizon $N \geq 0$ almost surely converges from above to the true optimal cost. That is, with probability one,

$$\lim_{t \to \infty} \hat{V}_{N}^{(t)}(x, w) = V_{N}(x, w), \tag{30}$$

for all $(x, w) \in \operatorname{dom} V_N$.

In the more general case, where aside from the cost, also the constraints are probabilistic and therefore dependent on the learner state, some additional assumptions on the problem ingredients are required.

Theorem V.13 (Asymptotic consistency under chance constraints). Let $s^* \in \mathcal{S}$ denote a stationary learner state (cf. Assumption III.4) and suppose that for a given state-mode pair $(x, w) \in \operatorname{dom} V_N$, the following hold:

- (i) Assumption V.8 holds, and $\mathcal{X}_{f}(w)$ is closed and convex;
- (ii) the costs $\ell(\cdot,\cdot,w)$, $V_f(\cdot,w)$, constraints $q(\cdot,\cdot,w,v)$ and dynamics $f(\cdot, \cdot, w, v)$ are continuously differentiable;
- (iii) the ambiguity set $A_{\beta}(s, w)$ is conic representable with convex cone K and parameters $E_w(s,\beta)$, $F_w(s,\beta)$ and $b_w(s,\beta)$ that depend smoothly on s and β ;
- (iv) Robinson's constraint qualification [62, Def. 2.86] holds for (37), for initial states (x, w) $s^{\iota} = s^{\star}, \beta^{\iota} = 0$.

Then,
$$\lim_{t\to\infty} \widehat{V}_N^{(t)}(x,w) = V_N(x,w)$$
, a.s.

Proof. Let x and w be fixed and consider the scenario tree formulation of the DR-OCP (37), with parameters $\theta := (s^{\iota}, \beta^{\iota})_{\iota \in \mathcal{N}}$, where $\mathcal{N} := \mathbf{nod}([0, N-1])$ denotes the set of non-leaf nodes of the tree⁴. Problem (37) can then be compactly written in the form (as we specify below)

$$\tilde{V}_{x,w}(\theta) := \min_{\zeta} \Psi(\zeta)$$
 subj. to $\Gamma(\zeta,\theta) \in K$. (31)

By Proposition B.2, we have that if the parameter vector $\theta = \theta_t := (s_t^\iota, \beta_t^\iota)_{\iota \in \mathcal{N}}$ is dynamically feasible, i.e., it satisfies (38) with values $(s_t^0, \beta_t^0) = (s_t, \beta_t)$ at the root node, then

$$\tilde{V}_{x,w}(\theta_t) = \hat{V}_N^{(t)}(x,w), \quad \forall t \in \mathbb{N}.$$
 (32)

Our proof now consists of two main steps. First, we observe that in the limit point $\theta^* := \lim_{t \to \infty} \theta_t$ (which exists by Assumptions III.4 and III.5), $\tilde{V}_{x,w}(\theta^*)$ coincides with $V_N(x,w)$. Secondly, we show that the mapping $\tilde{V}_{x,w}(\cdot)$ is continuous at θ^* .

I Assumption III.5 ensures that $\lim_{t\to\infty}\beta_t=\beta^*=0$ and consequently, by Condition (i), $\lim_{t\to\infty}\widehat{\alpha}_t=\alpha$. By Assumption V.11 and the requirement (8), the Borel-Cantelli lemma [46, Thm. 4.3] implies that for every sequence $(p_t\in\mathcal{A}_{\mathcal{B}_t}(s_t,w))_{t\in\mathbb{N}}$, $\lim_{t\to\infty}p_t=P_w$:, a.s. Furthermore, as $s_t\to s^*$, it follows by Condition (iii) that the mapping $(s,\beta)\mapsto\mathcal{A}_{\beta}(s,w)$ is continuous for all $w\in W$ and therefore $\mathcal{A}_0(s^*,w)=\{P_w:\}$, $\forall w\in W$. Thus, by their definitions (15)–(17), $\rho_{s^*,w}^{\beta^*}$ and $\bar{\rho}_{s^*,w}^{\bar{\beta}^*}$ reduce to \mathbb{E}_{P_w} : and $\mathrm{AV@R}_{\alpha}^{P_w}$:, respectively. Finally, by Condition (i), $(x,s,\beta,w)\in\widehat{\mathcal{X}_f}$ \iff $(x,w)\in\mathcal{X}_f$. Therefore, the DR-OCP (21) reduces to the nominal counterpart (7), or equivalently $V_{x,w}(\theta^*)=V_N(x,w)$.

II In order to show that $\tilde{V}_{x,w}$ is continuous at θ^* , we will show that Ψ and Γ are continuously differentiable and K is a closed convex set. Invoking furthermore Condition (iv), continuity of $\tilde{V}_{x,w}$ then follows from [62, Prop. 4.4]. By inspection of (37a) it is clear that Ψ is a linear function, satisfying the requirements. We now proceed to demonstrate that the constraints (37b)–(37g) admit the desired representation as well.

- 1) The constraints (37b)–(37d), and (37g) can be directly combined into the form $\Gamma_1(\zeta,\theta) \in K_1 := \{0\} \times \mathbb{R}^{n_1}_+ \times \widehat{\mathcal{X}}_{\mathrm{f}}$, where Γ_1 is a concatenation of the functions $\ell(\cdot,\cdot,w), V_{\mathrm{f}}(\cdot,w)$, and $f(\cdot,\cdot,w,v)$ and therefore continuously differentiable by Condition (ii). K_1 is convex due to Condition (i).
- 2) Finally, we consider the remaining constraints (37e) and (37f). Using (36), a conic risk epigraph constraint $(\xi, \gamma) \in \operatorname{epi} \tilde{\rho}^{\iota}$ with parameters $\tilde{E}(\theta^{\iota})$, $\tilde{F}(\theta^{\iota})$, $\tilde{b}(\theta^{\iota})$ and cone $\tilde{\mathcal{K}}$ can be written in the desired form

$$\tilde{\Gamma}_2(\xi, \chi, \theta^{\iota}) \in \tilde{K}_2 := \{0\} \times \tilde{\mathcal{K}}^* \times \mathbb{R}_+^{n_2} \tag{33}$$

with χ an auxiliary variable and

$$\tilde{\Gamma}_2(\xi,\gamma,y,\theta^\iota) := \left[\begin{smallmatrix} \tilde{E}(\theta^\iota) & \tilde{F}(\theta^\iota) \end{smallmatrix} I - \tilde{b}(\theta^\iota) \end{smallmatrix} \right]^\top y + \left[\begin{smallmatrix} 0 \\ -1 \end{smallmatrix} \right] \gamma + \left[\begin{smallmatrix} -I \\ 0 \end{smallmatrix} \right] \xi,$$

which is differentiable provided that $\tilde{E}(\theta^{\iota})$, $\tilde{F}(\theta^{\iota})$ and $\tilde{b}(\theta^{\iota})$ are differentiable. This is ensured exactly by Condition (iii), for the cost risk measure $\rho_{s,w}^{\beta}$, and thus (37e) is of the form (33).

Invoking Proposition B.1, $\bar{\rho}_{s,w}^{\bar{\beta},\widehat{\alpha}}$ is conic representable with parameters

$$\overline{E}_{w}(s,\overline{\beta}) = \begin{bmatrix} E_{\widehat{\alpha}} \\ 0 \end{bmatrix}, \ \overline{F}_{w}(s,\overline{\beta}) = \begin{bmatrix} -B & 0 \\ E_{w}(s,\overline{\beta}) \ F_{w}(s,\overline{\beta}) \end{bmatrix},
\overline{b}_{w}(s,\overline{\beta}) = \begin{bmatrix} b' \\ b_{w}(s,\overline{\beta}) \end{bmatrix}, \ \overline{\mathcal{K}} = \mathbb{R}_{+}^{2(d+1)} \times \mathcal{K},$$
(34)

with $E_{\widehat{\alpha}} = \begin{bmatrix} \mathbf{1}_d - \mathbf{1}_d \ \widehat{\alpha} I - I \end{bmatrix}^{\top}$, and $B, \ b'$ constant. Condition (i) requires that $\widehat{\alpha} = \frac{\alpha - \overline{\beta}}{1 - \overline{\beta}}$ is continuously differentiable in $\overline{\beta}$ for all $\overline{\beta} < 1$. The case $\overline{\beta} = 1$ is excluded by design and furthermore inconsequential as $\overline{\beta} \to 0$. As a result, (37f), i.e., constraints $(g(x, u, w, v), 0) \in \operatorname{epi} \overline{\rho}_{s,w}^{\overline{\beta}, \widehat{\alpha}}$ can be written in the form (33), replacing ξ with g(x, u, w, v) — which preserves continuous differentiability, due to Condition (ii) — and replacing the risk parameters $\widetilde{E}(\theta^t)$, $\widetilde{F}(\theta^t)$ and $\widetilde{b}(\theta^t)$ and $\widetilde{b}(\theta^t)$ and $\widetilde{b}(\theta^t)$ and $\widetilde{b}(\theta^t)$ and $\widetilde{b}(\theta^t)$ and $\widetilde{b}(\theta^t)$.

Given the established differentiability of Γ , the final requirement of [62, Prop. 4.4] is equivalent to Condition (iv), and thus, the result applies.

Combining I and II, we conclude that
$$\lim_{t\to\infty} \tilde{V}_{x,w}(\theta_t) = V_N(x,w)$$
, and the claim follows from (32).

We conclude this section with a few brief remarks regarding the conditions of Theorem V.13. First, we note that using the learning system described in Section III (including the ambiguity radius as part of the learner state as suggested in Remark III.2), Condition (iii) is satisfied for all divergence-based ambiguity sets considered in Table I. Indeed, in the conic formulations provided in Appendix A, we find that in all cases, the empirical distribution and the ambiguity radius enter linearly in the final conic form of the constraints. Second, we remark that Robinson's constraints qualification (Condition (iv)) can be regarded as a generalization of the more well-known Mangasarian-Fromowitz constraint qualification [62, eq. 2.191] (see also [63, Prop. 3.3.8] or [64, 4.10]), which is in turn a generalization of the linear independence constraint qualification (LICQ). It is a very common regularity assumption, ensuring several useful properties such as boundedness of Lagrange multipliers. Of main importance for the purpose of showing consistency under probabilistic constraints, however, is that it provides metric regularity of the (now parametric) feasible set, which implies that the distance from the feasible set can upper bounded by a multiple of the constraint violation.

VI. ILLUSTRATIVE EXAMPLES

A. Ambiguity sets

To illustrate the concentration inequalities provided in Proposition III.10 and Table I, we select a sequence of confidence levels $\beta_t = e^{-d}(t+1)^2$ satisfying summability (Assumption III.5) and we plot the radii corresponding to the considered divergences as a function of the sample size t (see Fig. 1). For comparison, we recursively estimate compute the empirical estimate \widehat{p}_t of a fixed probability vector $p \in \Delta_d$ and plot the empirical upper and lower β_t -quantile of $\mathcal{D}(\widehat{p}_t,p)$ over 200 Monte-Carlo runs. For the Wasserstein distance, a quadratic kernel $K_{wv} = (w-v)^2$, $\forall w,v \in \mathbb{N}_{[1,d]}$ was used. For all divergences, the given bounds provide reasonable approximations, but in particular, we note that the total variation bound is almost tight. Furthermore, it only requires linear constraints in its conic representation (23), making it an attractive choice in terms of both statistical and computational complexity.

B. Distributionally robust MPC

We consider a Markov jump linear system⁵ $x_{t+1} = A(w_{t+1})x_t + B(w_{t+1})u_t$, with

$$A(w) = \begin{bmatrix} 1 + \frac{w-1}{d} & 0.01\\ 0.01 & 1 + 2.5 \frac{w-1}{d} \end{bmatrix}, \ B(w) = I, w \in \mathbb{N}_{[1,d]}$$
 (35)

The state $x_t \in \mathbb{R}^2$ of this system, inspired by [66], models the deviation of temperatures from some nominal value of two adjacent

⁴See Appendix B for scenario tree related notation.

⁵For more extensive simulation results, involving nonlinear dynamics and nonconvex constraints, see [65]




Fig. 1. Radii of the ambiguity sets versus sample size t. The shaded area delineate the empirical β_t upper and lower quantiles of $\mathcal{D}(\widehat{p}_t,p)$ for different divergences \mathcal{D} , computed over 200 Monte-Carlo runs. The dashed lines represent the theoretical upper bounds given in Table I. $p_i = 1/d$, $\forall i \in \mathbb{N}_{[1,d]}, d = 5$ and $\beta_t = e^{-d}(t+1)^{-2}$.

servers in a data center. The actuators $u_t \in \mathbb{R}^2$ correspond to the amount of heating $(u_t \geq 0)$ or cooling $(u_t < 0)$ applied to the corresponding machines. The mode w models the load on the servers. If w=1, the system is idle and no heat is generated. If w=d, then the processors are fully occupied and a maximum amount of heat is added to the system. Note that the second server generates more heat under increasing loads. The true-but-unknown transition probabilities are computed as

$$P_{ij} = \frac{e^{-(j-i/2)^2}}{\sum_{w=1}^{d} e^{-(w-i/2)^2}}, \ \forall i, j \in \mathbb{N}_{[1,d]}.$$

As in [66], we will use a mode-independent quadratic cost $\ell(x, u, w) = ||x||_2^2 + 10^3 ||u||_2^2$.

We impose hard constraints $-1.5 \le u \le 1.5$ on the actuation and (nominally) impose robust chance constraints

$$\mathsf{AV} @ \mathsf{R}_{\alpha}^{P_{w_t:}}[H_{i:}x_{t+1} - h_i \mid x_t] \leq \alpha \text{ with } H = \begin{bmatrix} I_{n_x} \\ \mathbf{1}_{n_x}^\top \end{bmatrix}, h = \begin{bmatrix} 1_{n_x} \\ 0.5 \end{bmatrix},$$

for all $t\in\mathbb{N}_{[0,N-1]}$, and $\alpha=0.19$. Hence, in this example, we have $g_i(x,u,w,v)=H_{i:}(A(v)x+B(v)u)-h_i$.

We compute stabilizing terminal ingredients offline using standard techniques from robust control. We compute a robust quadratic Lyapunov function $V_{\mathbf{f}}(x) = x^{\top}Q_{\mathbf{f}}x$ along with a local linear control gain K, such that $V_{\mathbf{f}}((A(w)+B(w)K)x) \leq -\ell(x,Kx), \forall w \in W$ by solving a linear matrix inequality (LMI) as in [67]. The RCI terminal set $\mathcal{X}_{\mathbf{f}}$ is computed as the level set $\mathcal{X}_{\mathbf{f}} = \mathbf{lev}_{\leq \varepsilon} V_{\mathbf{f}}$, where $\varepsilon = \min_i \{h_i/\|Q_{\mathbf{f}}^{-1/2}H_{i:}\|_2^2\}$ is the largest value such that $\mathbf{lev}_{\leq \varepsilon} V_{\mathbf{f}}$ lies inside the polyhedral set $\{x \in \mathbb{R}^{n_x} \mid H(A(w)+B(w)K) \leq h, \forall w \in W\}$.

For the DR controllers below we choose confidence levels $\beta_t = (\beta_t, \bar{\beta}_t)$ with $\beta_t = \bar{\beta}_t = 0.19t^{-2} < \alpha$ for the cost and the constraints, respectively, ensuring that Assumption III.5 is satisfied. For simplicity, we use identical confidence levels $\bar{\beta}_t$ for all the constraints.

We compare the proposed DR-MPC controller with (i) the (nominal) stochastic MPC controller (see (7)), which we call **omniscient** as it has access to the true transition matrix P; and (ii) the **robust** MPC controller, obtained by solving (37), taking the ambiguity set $A_{\beta}(s,w) = A_{\overline{\beta}}(s,w) = \Delta_d$ to be the entire probability simplex, regardless of the mode or learner state. Both the LMIs involved in the offline computation of the terminal ingredients as the online risk-averse optimal control problem (37) are solved using MOSEK [68] through the CVXPY [69] interface.

We fix the number of modes to d=3, and take N=5. All computations were performed on an Intel Core i7-7700K CPU at 4.20GHz.

1) Timings: To obtain an indication of the comparative computational burden of the different divergence-based ambiguity sets under consideration, we solve the described DR-OCP using the considered divergences 10 times each, for random initial states. Table II reports the average and maximum observed solver time. As expected, the TV and Wasserstein divergence result require the least amount of time, as

they introduce only linear constraint. The Hellinger divergence, which introduces second-order cone constraints results in slightly longer run times. The KL and JS divergence both introduce exponential cone constraints, resulting in the most computationally demanding OCPs.

TABLE II SOLVER TIMES [ms] FOR (21) USING DIFFERENT DIVERGENCE-BASED AMBIGUITY SETS

	TV	Wasserstein	KL	JS	Hellinger
avg.	50.14	50.63	225.6	112.00	61.31
max.	52.20	52.02	235.02	118.79	62.03

2) Closed-loop simulation: Motivated by previous experiments, we now select the TV ambiguity set, and perform a more extensive closed-loop simulation. Fixing the initial state at $x = \begin{bmatrix} 0.5 & 0.5 \end{bmatrix}^T$, we perform 50 Monte-Carlo simulations of the described MPC problems for 30 steps. As the simulation time is rather short, we initialize the DR controller with 10 and 100 offline observations of the Markov chain to obtain more interesting comparisons. Hence, the simulation below essentially compares the controller responses after a sudden disturbance after 10 and 100 time steps. All considered controllers are recursively feasible and mean-square stabilizing by construction. By the nature of the problem set-up, the optimal behavior is to just barely stabilize the system with minimal control effort. However, the larger the uncertainty on the state evolution, the more the controller is forced to drive the states further away from the constraint boundary, leading to larger control actions and consequently, larger costs.




Fig. 2. Control effort and second component of the state vector over 50 monte-carlo simulations. Full lines depict the means over the realizations and the shaded areas are delineated by the 0.05 and 0.95 quantiles.

This behavior can be observed in Fig. 2 and 3. Fig. 2 shows the controls and states over time and Fig. 3 presents the distribution of the closed-loop costs (sum of the stage costs over the simulation time). In the first time step, the robust controller takes the largest step, driving the state the furthest from the constraint boundary. As illustrated in Fig. 2 (right), this is particularly pronounced for the second component of the state vector, as it is more sensitive to the mode (cf. (35)). The *omniscient* stochastic MPC, by contrast, has perfect knowledge of the transition probabilities, and by consequence




Fig. 3. Box plot of the closed-loop cost over 50 monte-carlo simulations. The annotated lines show the mean. The whiskers depict the 0.05 and 0.95 quantiles.

is able to more slowly drive the state to the origin, reducing the control effort considerably. The DR controller naturally 'interpolates' between these behaviors. Initially, it performs only marginally better than the robust controller (due to the very limited number of online learning steps). As it gets access to increasing sample sizes, however, it gradually approximates the behavior of the omniscient controller, while guaranteeing satisfaction of the constraints throughout.




Fig. 4. Relative suboptimality versus sample size for the example system (35). The dashed line depicts the relative suboptimality of the robust controller: $(V_r - V^*)/V^*$.

3) Asymptotic consistency: To illustrate the consistency results from Section V-D, we fix the initial state-mode pair $x_0 = [0.25\ 0.25\]^{\mathsf{T}}$, $w_0 = 1$ and recompute the solution to problem (37) to obtain $\widehat{V}^{(t)} := \widehat{V}_N^{(t)}(x_0, w_0)$ for increasing sample sizes t. For comparison, we compute (i) the true value $V^\star := V_N(x_0, w_0)$ by solving the stochastic MPC problem (7), using the true transition probabilities; and (ii) the robust value function V_r , obtained by solving (37), taking the ambiguity set $\mathcal{A}_\beta(s,w) = \Delta_d$ to be the entire probability simplex, regardless of the mode or learner state.

Figure 4 shows the relative difference between the DR value $\widehat{V}^{(t)}$ and the true value V^{\star} for the different statistical divergences . At very low sample sizes, the DR controllers achieve the same cost as the robust controller. However, as more data is gathered and the ambiguity set is updated, $\widehat{V}^{(t)}$ approaches V^{\star} from above.

VII. CONCLUSION

We presented a distributionally robust MPC strategy for Markov jump systems with unknown transition probabilities subject to general probabilistic constraints. We proved closed-loop constraint satisfaction, mean-square stability, and consistency of the resulting controller for a broad range of data-driven ambiguity sets.

REFERENCES

- M. Schuurmans and P. Patrinos, "Learning-Based Distributionally Robust Model Predictive Control of Markovian Switching Systems with Guaranteed Stability and Recursive Feasibility," arXiv:2009.04422, Sept. 2020.
- [2] B. Kouvaritakis and M. Cannon, *Model Predictive Control*. Advanced Textbooks in Control and Signal Processing, Cham: Springer International Publishing, 2016.

[3] A. Mesbah, "Stochastic Model Predictive Control: An Overview and Perspectives for Future Research," *IEEE Control Systems Magazine*, vol. 36, pp. 30–44, Dec. 2016.

- [4] J. B. Rawlings, D. Q. Mayne, and M. M. Diehl, Model Predictive Control: Theory, Computation, and Design. Madison, Wisconsin: Nob Hill Publishing, second ed., 2017.
- [5] P. Mohajerin Esfahani and D. Kuhn, "Data-driven distributionally robust optimization using the Wasserstein metric: Performance guarantees and tractable reformulations," *Mathematical Programming*, vol. 171, pp. 115–166, Sept. 2018.
- [6] J. Dupačová, "The minimax approach to stochastic programming and an illustrative application," *Stochastics*, vol. 20, pp. 73–88, Jan. 1987.
- [7] B. P. G. Van Parys, P. M. Esfahani, and D. Kuhn, "From Data to Decisions: Distributionally Robust Optimization Is Optimal," *Management Science*, Nov. 2020.
- [8] R. Gao and A. J. Kleywegt, "Distributionally Robust Stochastic Optimization with Wasserstein Distance," arXiv:1604.02199, Apr. 2016.
- [9] W. Wiesemann, D. Kuhn, and M. Sim, "Distributionally Robust Convex Optimization," *Operations Research*, vol. 62, pp. 1358–1376, Dec. 2014.
- [10] D. Bertsimas, V. Gupta, and N. Kallus, "Data-driven robust optimization," *Mathematical Programming*, vol. 167, pp. 235–292, Feb. 2018.
- [11] M. Schuurmans, P. Sopasakis, and P. Patrinos, "Safe Learning-Based Control of Stochastic Jump Linear Systems: A Distributionally Robust Approach," in 58th IEEE Conference on Decision and Control (CDC), pp. 6498–6503, Dec. 2019.
- [12] P. Coppens, M. Schuurmans, and P. Patrinos, "Data-driven distributionally robust LQR with multiplicative noise," in *Learning for Dynamics* and Control, pp. 521–530, PMLR, July 2020.
- [13] I. Yang, "Wasserstein Distributionally Robust Stochastic Control: A Data-Driven Approach," arXiv:1812.09808, Dec. 2018.
- [14] A. Hakobyan and I. Yang, "Wasserstein Distributionally Robust Motion Control for Collision Avoidance Using Conditional Value-at-Risk," arXiv:2001.04727, Jan. 2020.
- [15] A. Hakobyan and I. Yang, "Distributionally robust risk map for learningbased motion planning and control: A semidefinite programming approach," arXiv:2105.00657, May 2021.
- [16] J. Coulson, J. Lygeros, and F. Dörfler, "Regularized and Distributionally Robust Data-Enabled Predictive Control," in 2019 IEEE 58th Conference on Decision and Control (CDC), pp. 2696–2701, Dec. 2019.
- [17] H. Rahimian and S. Mehrotra, "Distributionally Robust Optimization: A Review," arXiv:1908.05659, Aug. 2019.
- [18] A. Aswani, H. Gonzalez, S. S. Sastry, and C. Tomlin, "Provably safe and robust learning-based model predictive control," *Automatica*, vol. 49, pp. 1216–1226, May 2013.
- [19] L. Hewing and M. N. Zeilinger, "Scenario-Based Probabilistic Reachable Sets for Recursively Feasible Stochastic Model Predictive Control," *IEEE Control Systems Letters*, vol. 4, pp. 450–455, Apr. 2020.
- [20] J. F. Fisac, A. K. Akametalu, M. N. Zeilinger, S. Kaynama, J. Gillula, and C. J. Tomlin, "A General Safety Framework for Learning-Based Control in Uncertain Robotic Systems," *IEEE Transactions on Automatic Control*, vol. 64, pp. 2737–2752, July 2019.
- [21] J. Coulson, J. Lygeros, and F. Dörfler, "Data-Enabled Predictive Control: In the Shallows of the DeePC," arXiv:1811.05890, Mar. 2019.
- [22] X. Zhang, J. Liu, X. Xu, S. Yu, and H. Chen, "Robust Learning-Based Predictive Control for Discrete-Time Nonlinear Systems With Unknown Dynamics and State Constraints," *IEEE Transactions on Systems, Man, and Cybernetics: Systems*, pp. 1–14, 2022.
- [23] L. Hewing, K. P. Wabersich, M. Menner, and M. N. Zeilinger, "Learning-Based Model Predictive Control: Toward Safe Learning in Control," *Annual Review of Control, Robotics, and Autonomous Systems*, vol. 3, no. 1, 2020.
- [24] D. Bernardini and A. Bemporad, "Stabilizing Model Predictive Control of Stochastic Constrained Linear Systems," *IEEE Transactions on Automatic Control*, vol. 57, pp. 1468–1480, June 2012.
- [25] D. Bernardini and A. Bemporad, "Scenario-based model predictive control of stochastic constrained linear systems," in 48th IEEE Conference on Decision and Control (CDC) Held Jointly with 2009 28th Chinese Control Conference, pp. 6333–6338, IEEE, Dec. 2009.
- [26] S. Lucia, T. Finkler, and S. Engell, "Multi-stage nonlinear model predictive control applied to a semi-batch polymerization reactor under uncertainty," *Journal of Process Control*, vol. 23, pp. 1306–1319, Oct. 2013.
- [27] C. Leidereiter, A. Potschka, and H. G. Bock, "Quadrature-based scenario tree generation for Nonlinear Model Predictive Control," *IFAC Proceedings Volumes*, vol. 47, no. 3, pp. 11087–11092, 2014.

- [28] A. D. Bonzanini, J. A. Paulson, and A. Mesbah, "Safe learning-based model predictive control under state-and input-dependent uncertainty using scenario trees," in *Proceedings of the IEEE Conference on Decision* and Control. Jeju Island, Republic of Korea. Submitted, 2020.
- [29] O. L. d. V. Costa, M. D. Fragoso, and R. P. Marques, *Discrete-time Markov jump linear systems*. Probability and its applications, London: Springer, 2005.
- [30] P. Patrinos, P. Sopasakis, H. Sarimveis, and A. Bemporad, "Stochastic model predictive control for constrained discrete-time Markovian switching systems," *Automatica*, vol. 50, pp. 2504–2514, Oct. 2014.
- [31] S. Lucia, S. Subramanian, D. Limon, and S. Engell, "Stability properties of multi-stage nonlinear model predictive control," *Systems & Control Letters*, vol. 143, p. 104743, Sept. 2020.
- [32] P. Sopasakis, D. Herceg, A. Bemporad, and P. Patrinos, "Risk-averse model predictive control," *Automatica*, vol. 100, pp. 281–288, Feb. 2019.
- [33] S. Singh, Y.-L. Chow, A. Majumdar, and M. Pavone, "A Framework for Time-Consistent, Risk-Sensitive Model Predictive Control: Theory and Algorithms," arXiv:1703.01029, Apr. 2018.
- [34] A. Shapiro, D. Dentcheva, and A. Ruszczyński, *Lectures on stochastic programming: modeling and theory*. SIAM, 2009.
- [35] R. L. Beirigo, M. G. Todorov, and A. M. S. Barreto, "Online TD(λ) for discrete-time Markov jump linear systems," in 57th IEEE Conference on Decision and Control (CDC), pp. 2229–2234, Dec. 2018.
- [36] S. He, M. Zhang, H. Fang, F. Liu, X. Luan, and Z. Ding, "Reinforcement learning and adaptive optimization of a class of Markov jump systems with completely unknown dynamic information," *Neural Computing and Applications*, Apr. 2019.
- [37] E. Derman and S. Mannor, "Distributional Robustness and Regularization in Reinforcement Learning," July 2020.
- [38] H. Xu and S. Mannor, "Distributionally Robust Markov Decision Processes," in *Advances in Neural Information Processing Systems 23* (J. D. Lafferty, C. K. I. Williams, J. Shawe-Taylor, R. S. Zemel, and A. Culotta, eds.), pp. 2505–2513, Curran Associates, Inc., 2010.
- [39] S. Mannor, O. Mebel, and H. Xu, "Robust MDPs with k-Rectangular Uncertainty," *Mathematics of Operations Research*, vol. 41, pp. 1484– 1509, Nov. 2016.
- [40] M. Ahmadi, U. Rosolia, M. D. Ingham, R. M. Murray, and A. D. Ames, "Constrained Risk-Averse Markov Decision Processes," arXiv:2012.02423, Dec. 2020.
- [41] A. Cherukuri and A. R. Hota, "Consistency of Distributionally Robust Risk- and Chance-Constrained Optimization under Wasserstein Ambiguity Sets," arXiv:2012.08850, Dec. 2020.
- [42] S. Guo, H. Xu, and L. Zhang, "Convergence Analysis for Mathematical Programs with Distributionally Robust Chance Constraint," SIAM Journal on Optimization, vol. 27, pp. 784–816, Jan. 2017.
- [43] A. Nemirovski, "On safe tractable approximations of chance constraints," *European Journal of Operational Research*, vol. 219, no. 3, pp. 707–718, 2012.
- [44] P. Sopasakis, M. Schuurmans, and P. Patrinos, "Risk-averse risk-constrained optimal control," in 18th European Control Conference (ECC), pp. 375–380, June 2019.
- [45] D. P. Bertsekas, Dynamic Programming and Optimal Control. Vol. 1. Athena Scientific Optimization and Computation Series, Belmont, Mass: Athena Scientific, third ed., 2005.
- [46] P. Billingsley, Probability and Measure. Wiley Series in Probability and Mathematical Statistics, New York: Wiley, third ed., 1995.
- [47] V. Krishnamurthy, Partially Observed Markov Decision Processes: From Filtering to Controlled Sensing. Cambridge: Cambridge University Press 2016
- [48] R. Jiang and Y. Guan, "Risk-Averse Two-Stage Stochastic Program with Distributional Ambiguity," *Operations Research*, vol. 66, pp. 1390–1405, Oct. 2018.
- [49] H. Rahimian, G. Bayraksan, and T. Homem-de-Mello, "Effective Scenarios in Multistage Distributionally Robust Optimization with a Focus on Total Variation Distance," arXiv:2109.06791, Sept. 2021.
- [50] G. Wolfer and A. Kontorovich, "Minimax Learning of Ergodic Markov Chains," in *Algorithmic Learning Theory*, pp. 903–929, Mar. 2019.
- [51] A. W. van der Vaart and J. A. Wellner, Weak Convergence and Empirical Processes: With Applications to Statistics. New York: Springer, 2000.
- [52] I. Csiszar, "The method of types," *IEEE Transactions on Information Theory*, vol. 44, pp. 2505–2523, Oct. 1998.
- [53] T. M. Cover and J. A. Thomas, Elements of Information Theory. Hoboken, N.J. Wiley-Interscience, 2nd ed., 2006.
- [54] J. Mardia, J. Jiao, E. Tánczos, R. D. Nowak, and T. Weissman, "Concentration inequalities for the empirical distribution of discrete distributions: Beyond the method of types," *Information and Inference: A Journal of the IMA*, Nov. 2019.

- [55] I. Csiszár and J. Körner, Information Theory: Coding Theorems for Discrete Memoryless Systems. Cambridge; New York: Cambridge University Press, 2nd ed ed., 2011.
- [56] A. L. Gibbs and F. E. Su, "On Choosing and Bounding Probability Metrics," *International Statistical Review*, vol. 70, no. 3, pp. 419–435, 2002.
- [57] G. Bayraksan and D. K. Love, "Data-Driven Stochastic Programming Using Phi-Divergences," in *The Operations Research Revolution* (D. Aleman, A. Thiele, J. C. Smith, and H. J. Greenberg, eds.), pp. 1–19, INFORMS, Sept. 2015.
- [58] A. Ben-Tal, D. den Hertog, A. De Waegenaere, B. Melenberg, and G. Rennen, "Robust Solutions of Optimization Problems Affected by Uncertain Probabilities," *Management Science*, vol. 59, pp. 341–357, Nov. 2012.
- [59] İ. Yanıkoğlu and D. den Hertog, "Safe Approximations of Ambiguous Chance Constraints Using Historical Data," INFORMS Journal on Computing. Nov. 2012.
- [60] A. Ruszczyński, "Risk-averse dynamic programming for Markov decision processes," *Mathematical Programming*, vol. 125, pp. 235–261, Oct. 2010.
- [61] M. Korda, R. Gondhalekar, J. Cigler, and F. Oldewurtel, "Strongly feasible stochastic model predictive control," in 50th IEEE Conference on Decision and Control and European Control Conference, pp. 1245– 1251, Dec. 2011
- [62] J. F. Bonnans and A. Shapiro, Perturbation Analysis of Optimization Problems. Springer Series in Operations Research, New York: Springer, 2000
- [63] D. P. Bertsekas, Nonlinear Programming. Belmont, Mass: Athena Scientific, second ed., 1999.
- [64] J. O. Royset and R. J.-B. Wets, An Optimization Primer. Springer Series in Operations Research and Financial Engineering, Cham, Switzerland: Springer, 2021.
- [65] M. Schuurmans, A. Katriniok, C. Meissen, H. E. Tseng, and P. Patrinos, "Safe, Learning-Based MPC for Highway Driving under Lane-Change Uncertainty: A Distributionally Robust Approach," June 2022.
- [66] B. Recht, "A Tour of Reinforcement Learning: The View from Continuous Control," *Annual Review of Control, Robotics, and Autonomous Systems*, vol. 2, no. 1, pp. 253–279, 2019.
- [67] M. V. Kothare, V. Balakrishnan, and M. Morari, "Robust constrained model predictive control using linear matrix inequalities," *Automatica*, vol. 32, no. 10, pp. 1361–1379, 1996.
- [68] MOSEK ApS, The MOSEK optimization toolbox for MATLAB manual. Version 8.1., 2017.
- [69] S. Diamond and S. Boyd, "CVXPY: A Python-embedded modeling language for convex optimization," *Journal of Machine Learning Research*, vol. 17, no. 83, pp. 1–5, 2016.
- [70] A. Shapiro, "On Duality Theory of Conic Linear Problems," in Semi-Infinite Programming (P. Pardalos, M. Á. Goberna, and M. A. López, eds.), vol. 57, pp. 135–165, Boston, MA: Springer US, 2001.
- [71] G. C. Pflug and A. Pichler, Multistage Stochastic Optimization. Springer Series in Operations Research and Financial Engineering, Cham: Springer International Publishing, 2014.
- [72] M. Schuurmans, A. Katriniok, H. E. Tseng, and P. Patrinos, "Learning-Based Risk-Averse Model Predictive Control for Adaptive Cruise Control with Stochastic Driver Models," in *IFAC 2020 World Congress*, (Berlin), pp. 15337–15342, 2020.
- [73] R. T. Rockafellar and R. J. B. Wets, Variational Analysis, vol. 317 of Grundlehren Der Mathematischen Wissenschaften. Berlin, Heidelberg: Springer Berlin Heidelberg, 1998.



Mathijs Schuurmans obtained a Bachelor's degree (BSc) in Electrical and Mechanical Engineering and a Master's (MSc) in Mathematical Engineering from KU Leuven, Leuven, Belgium in 2016 and 2018, respectively. He is currently a PhD candidate at the Department of Electrical Engineering (ESAT) of KU Leuven. His research is focused on data-driven model predictive control of stochastic systems, focusing on distributionally robust approaches for safety-critical applications in autonomous driving.



Panagiotis Patrinos Panagiotis (Panos) Patrinos is associate professor at the Department of Electrical Engineering (ESAT) of KU Leuven, Belgium. In 2014 he was a visiting professor at Stanford University. He received his PhD in Control and Optimization, M.S. in Applied Mathematics and M.Eng. in Chemical Engineering from the National Technical University of Athens in 2010, 2005 and 2003, respectively. After his PhD he held postdoc positions at the University of Trento and IMT Lucca, Italy, where he became

an assistant professor in 2012. His current research interests lie in the intersection of optimization, control and learning. In particular he is interested in the theory and algorithms for structured nonconvex optimization as well as learning-based, model predictive control with a wide range of applications including autonomous vehicles, machine learning and signal processing. He is the co-recipient of the 2020 best paper award in International Journal of Circuit Theory & Applications

APPENDIX

A. Conic representations of divergence-based ambiguity sets

In this section, we briefly present conic representations for the divergences included in Table I. These reformulations, which may not be unique, follow from rather straightforward manipulations, but are included here for completeness. In the following, \widehat{p} represents a given empirical distribution and r represents a given radius. p is the candidate distribution over which the expectation is maximized. Below, we describe the set $\{p \in \Delta_d \mid \mathcal{D}(\widehat{p},p) \leq r\}$ for different divergences \mathcal{D} . Note that in all cases, we have the constraint $p \in \Delta_d$, which is represented by linear equality and inequality constraints.

Total variation:

$$||p - \widehat{p}||_1 \le r \iff \exists \nu \in \mathbb{R}^d : \begin{cases} -\nu_i \le p_i - \widehat{p}_i \le \nu_i, \ \forall i \in \mathbb{N}_{[1,d]} \\ \sum_{i=1}^d \nu_i \le r. \end{cases}$$

Kullback-Leibler:

$$\begin{split} & \sum_{i=1}^{d} \widehat{p}_{i} \log \frac{\widehat{p}_{i}}{p_{i}} \leq r \\ & \iff \exists \nu \in \mathbb{R}^{d} : \begin{cases} \widehat{p}_{i} \log \frac{\widehat{p}_{i}}{p_{i}} \leq \nu_{i} & \forall i \in \mathbb{N}_{[1,d]} \\ \sum_{i=1}^{d} \nu_{i} \leq r, \end{cases} \\ & \iff \exists \nu \in \mathbb{R}^{d} : \begin{cases} (-\nu_{i}, \widehat{p}_{i}, p_{i}) \in \mathcal{K}_{\mathbf{exp}} & \forall i \in \mathbb{N}_{[1,d]} \\ \sum_{i=1}^{d} \nu_{i} \leq r, \end{cases} \end{split}$$

where $\mathcal{K}_{\mathbf{exp}} = \{(x_1, x_2, x_3) : x_1 \leq x_2 \log\left(\frac{x_3}{x_2}\right), x_2 > 0\}$ denotes the exponential cone, the dual cone of which is $\mathcal{K}^*_{\mathbf{exp}} := \{(u, v, w) : u < 0, w > 0, -u \log(-\frac{u}{w}) + u - v \leq 0\}.$ Jensen-Shannon:

$$\mathcal{D}_{J}(\widehat{p}, p) := \frac{1}{2} \left(\mathcal{D}_{KL} \left(p, \frac{1}{2} (\widehat{p} + p) \right) + \mathcal{D}_{KL} \left(\widehat{p}, \frac{1}{2} (\widehat{p} + p) \right) \right) \le r$$

$$\iff \sum_{i=1}^{d} p_{i} \log \left(\frac{2p_{i}}{p_{i} + \widehat{p}_{i}} \right) + \widehat{p}_{i} \log \left(\frac{2\widehat{p}_{i}}{p_{i} + \widehat{p}_{i}} \right) \le 2r$$

$$\iff \exists x, y \in \mathbb{R}^{d} : \begin{cases} \sum_{i} x_{i} + y_{i} \le 2r \\ (-x_{i}, p_{i}, \frac{1}{2} (p_{i} + \widehat{p}_{i})) \in \mathcal{K}_{\mathbf{exp}} \\ (-y_{i}, \widehat{p}_{i}, \frac{1}{2} (p_{i} + \widehat{p}_{i})) \in \mathcal{K}_{\mathbf{exp}}. \end{cases}$$

Hellinger: Using the fact that

$$\begin{split} \mathcal{D}^2_{\mathrm{H}}(\widehat{p},p) &= \textstyle \sum_{i=1}^d (\sqrt{p_i} - \sqrt{\widehat{p}_i})^2 = 2(1 - \textstyle \sum_{i=1}^d \sqrt{p_i \widehat{p}_i}), \\ \text{we have } \mathcal{D}^2_{\mathrm{H}}(\widehat{p},p) \leq r \iff \widehat{r} &:= 1 - \frac{r}{2} \leq \textstyle \sum_{i=1}^d \sqrt{p_i \widehat{p}_i}, \text{ and thus} \\ \mathcal{D}^2_{\mathrm{H}}(\widehat{p},p) \leq r \iff \exists \nu \in \mathbb{R}^d : \begin{cases} \widehat{r} \leq \textstyle \sum_{i=1}^d \sqrt{\widehat{p}_i} \nu_i \\ \nu_i^2 \leq p_i, \end{cases} & \forall i \in \mathbb{N}_{[1,d]}. \end{split}$$

The constraint $\nu_i^2 \leq p_i$ can be reformulated as

$$||(2\nu_i, p_i - 1)||_2 \le p_i + 1 \iff (p_i + 1, 2\nu_i, p_i - 1) \in \mathcal{K}_{\mathbf{SO}},$$

with $\mathcal{K}_{SO} := \{x = (t, y) \in \mathbb{R}^{1+n} \mid t \geq ||y||_2\}$ the (self-dual) second order (or quadratic) cone.

Wasserstein: Assuming that W is a metric space with distance metric **dist**: $W \times W \rightarrow \mathbb{R}_+$, then we define the fixed-support q-Wasserstein distance, for q > 0 as

$$\mathcal{D}_{\mathrm{W},q}(p,\widehat{p}) := \min_{\Pi \in \mathbb{R}_+^{d \times d}} \left\{ \left(\sum_{i,j \in W} \Pi_{ij} K_{ij}^q \right)^{1/q} \middle| \begin{array}{c} \Pi \mathbf{1}_d = p \\ \Pi^\top \mathbf{1}_d = \widehat{p} \end{array} \right\},$$

where $K := (\mathbf{dist}(i,j))_{i,j \in W} \in \mathbb{R}_+^{d \times d}$ is the distance kernel induced by \mathbf{dist} . Since $\mathcal{D}_{\mathrm{W},q}(p,\widehat{p}) \leq r$ is equivalent to $\mathcal{D}_{\mathrm{W},q}(p,\widehat{p})^q \leq \widetilde{r} := r^q$, the q-Wasserstein case can be reduced to the 1-Wasserstein case with distance kernel $\widetilde{K} := K^q$. Thus, we may drop the subscript q without loss of generality. From the definition, we immediately obtain the representation

$$\mathcal{D}_{\mathrm{W},1}(p,\widehat{p}) \le r \iff \exists \Pi \ge 0 : \begin{cases} \sum_{i,j \in W} \Pi_{ij} \widetilde{K}_{ij} \le \widetilde{r}, \\ \Pi \mathbf{1}_d = p, \ \Pi^\top \mathbf{1}_d = \widehat{p}, \end{cases}$$

consisting only of linear constraints

B. Conic reformulation over scenario trees

By definition, a conic risk measure ρ is given as the optimal value of a standard conic program (CP). Under strong duality, which holds if the CP is strictly feasible [70, Prop. 2.1], its epigraph $\operatorname{epi} \rho := \{(G, \gamma) \in \mathbb{R}^{d+1} \mid \gamma \geq \rho[G]\}$ can be characterized as [44]

$$\mathbf{epi}\,\rho = \left\{ (G, \gamma) \in \mathbb{R}^{d+1} \,\middle|\, \begin{array}{l} \exists y : E^\top y = G, F^\top y = 0, \\ y \in \mathcal{K}^*, \gamma \ge b^\top y \end{array} \right\}. \quad (36)$$

Aside from the ambiguity sets described in Appendix A, it is not difficult to show that the worst-case average value-at-risk (17) over a conic representable ambiguity set also defines a conic risk measure:

Proposition B.1. Let $\mathcal{A}=\{p\in\Delta_d\mid \exists \nu: \overline{E}p+\overline{F}\nu\preccurlyeq_{\mathcal{K}}\bar{b}\}$ be a conic-representable ambiguity set. Then, the risk measure $\bar{\rho}=\max_{p\in\mathcal{A}}\mathsf{AV@R}^p_{\alpha}$ is a conic risk measure.

Proof. For any reference distribution $p \in \Delta_d$, the ambiguity set $\mathcal{A}_{\mathsf{AV@R}}$ inducing $\mathsf{AV@R}^p_\alpha$ can be written in the form (23) with $E = \begin{bmatrix} \mathbf{1}_d - \mathbf{1}_d & \alpha I - I \end{bmatrix}^\top$, F = 0, $\mathcal{K} = \mathbb{R}^{2(d+1)}_+$ the nonnegative orthant, and $b = \begin{bmatrix} 1 - 1 & p^\top & 0 \end{bmatrix}^\top$ (which is of the form b = b' + Bp) [44]. Writing out the definition of $\max_{p \in \mathcal{A}} \mathsf{AV@R}^p_\alpha$ and rearranging terms yields $\max_{p \in \mathcal{A}} \mathsf{AV@R}^p_\alpha[z] = \max_\mu \left\{ \mu^\top z \ \middle| \ \exists \nu : \begin{bmatrix} E \\ 0 \end{bmatrix} \mu + \begin{bmatrix} -B & 0 \\ \overline{E} & \overline{F} \end{bmatrix} \nu \preccurlyeq_{\mathbb{R}^2_+(d+1)} \times \mathcal{K}} \begin{bmatrix} b' \\ \overline{b} \end{bmatrix} \right\}$, which is exactly of the form (23).

Thus, if for all $(s,w,\beta)\in\mathcal{S}\times W\times[0,1]$, $\mathcal{A}_{\beta}(s,w)$ is conic representable, then $\rho_{s,w}^{\beta}$ and $\bar{\rho}_{s,w}^{\beta,\alpha}$ are conic risk measures. Therefore, problem (21) can be cast to a finite dimensional optimization problem, as we now describe.

Since W is a finite set, the possible realizations of $w_{[0,N]}$ can be enumerated and represented on a *scenario tree*. A scenario tree with horizon N represents the natural filtration of (Ω, F, \mathbb{P}) induced by $w_{[0,N]}$ [71]. Any adapted stochastic process (z_t) can be represented on such a scenario tree. We denote the value of z_t corresponding to a node ι in the tree as z^t .

The set of nodes in the tree are partitioned into time steps or *stages*. The set of nodes at a stage k is denoted by $\mathbf{nod}(k)$, and similarly, for $k_0 < k_1 \in \mathbb{N}_{[0,N]}$, $\mathbf{nod}([k_0,k_1]) = \bigcup_{k=k_0}^{k_1} \mathbf{nod}(k)$. For a given node $\iota \in \mathbf{nod}(t)$, $t \in \mathbb{N}_{[0,N-1]}$, we call a node $\iota_+ \in \mathbf{nod}(t+1)$ that can be reached from ι in one step a *child* node, denoted $\iota_+ \in \mathbf{ch}(\iota)$. An N-step policy π can thus be identified with a collection of control actions $\mathbf{u} = \{u^\iota \mid \iota \in \mathbf{nod}([0,N-1])\}$. It therefore

suffices to optimize over a finite number of decision variables rather than infinite-dimensional control laws.

Proposition B.2 (Finite-dimensional reformulation). Given an initial state (x, w) and parameters $(s^{\iota}, \beta^{\iota})_{\iota \in \mathbf{nod}([0, N-1])}$, consider an N-stage scenario tree and the corresponding optimal control

$$\underset{\xi \in T}{\text{minimize}} \ \tau^0 + \xi^0 \tag{37a}$$

subj. to
$$x^0 = x, w^0 = w, x^{\iota_+} = f(x^{\iota}, u^{\iota}, w^{\iota_+}),$$
 (37b)

$$\ell(x^{\iota}, u^{\iota}, w^{\iota}) \le \tau^{\iota}, \tag{37c}$$

$$V_{\mathbf{f}}(x^{\iota_N}, w^{\iota_N}) \le \xi^{\iota_N} + \tau^{\iota_N}, \tag{37d}$$

$$(\tau^{\iota_+} + \xi^{\iota_+}, \xi^{\iota}) \in \operatorname{\mathbf{epi}} \rho^{\beta^{\iota}}_{s^{\iota}, w^{\iota}}, \tag{37e}$$

$$(g(x^{\iota}, u^{\iota}, w^{\iota}, w^{\iota+}), 0) \in \operatorname{\mathbf{epi}} \bar{\rho}_{s\iota}^{\overline{\beta}{\iota}, \alpha\iota}^{\overline{\iota}}, \tag{37f}$$

$$(x^{\iota_N}, s^{\iota_N}, \boldsymbol{\beta}^{\iota_N}, w^{\iota_N}) \in \widehat{\mathcal{X}_{\mathbf{f}}}, \tag{37g}$$

for $\iota \in \mathbf{nod}([0, N-1])$, $\iota_+ \in \mathbf{ch}(\iota)$, and $\iota_N \in \mathbf{nod}(N)$, where ${\boldsymbol x}=(x^\iota)_{\iota\in\mathbf{nod}([0,N])}$ and ${\boldsymbol u}$ as defined above. If the parameters $(s^{\iota}, \beta^{\iota})$ satisfy for all $\iota \in \mathbf{nod}([0, N-1])$ that

$$s^{t+} = \mathcal{L}(s^t, \boldsymbol{\beta}^t, w^t, w^{t+}) \text{ and } \boldsymbol{\beta}^{t+} = C(\boldsymbol{\beta}^t), \tag{38}$$

then the optimal cost of (37) is equal to $\widehat{V}_N(z)$.

Proof. The claim is a direct application of the results in [44].

If (i) the costs $\ell(\cdot,\cdot,w)$, $V_{\rm f}(\cdot,w)$, the constraint mappings $g(\cdot,\cdot,w,v)$ and terminal set $\widehat{\mathcal{X}_{\mathrm{f}}}$ are convex; (ii) the ambiguity sets $\mathcal{A}_{\beta^{\iota}}(s^{\iota}, w^{\iota})$ are conic representable; and (iii) the dynamics $f(\cdot, \cdot, w)$ are affine for all $w \in W$, then it follows from Proposition B.1 that both $\rho_{s^{\iota},w^{\iota}}^{\beta^{\iota}}$ and $\bar{\rho}_{s^{\iota},w^{\iota}}^{\bar{\beta}^{\iota},\hat{\alpha}^{\iota}}$ are conic risk measures and thus (37) can be reduced to a convex conic program. See Section VI for a numerical illustration, as well as [72] for a case study in a slightly simplified setting. Note that since the learner and confidence dynamics \mathcal{L} and Care eliminated before solving the optimization problem, they need not be affine for the problem to remain convex. For nonlinear dynamics $f(\cdot, \cdot, v)$, the problem is no longer convex but can in practice still be solved effectively with standard NLP solvers.

C. Technical Lemma

Lemma C.1 (Infimum convergence). Consider a sequence of proper, lsc functions $V^{(t)}: \mathbb{R}^n \to \overline{\mathbb{R}}, t \in \mathbb{N}$ and a proper, lsc, levelbounded function $V: \mathbb{R}^n \to \overline{\mathbb{R}}$. Suppose that

- (i) (Eventual upper bound) there exists a T ∈ N, such that for all t > T, and for all u, V^(t)(u) ≥ V(u);
 (ii) (Pointwise convergence) V^(t) P/→ V. That is, for all u,
- $\lim_{t} V^{(t)}(u) = V(u).$

Then, $\lim_t \inf_u V^{(t)}(u) = \inf_u V(u)$.

Proof. By (i) it follows that for any sequence $u_t \to \bar{u}$,

$$\liminf_{t} V^{(t)}(u_t) = \liminf_{\substack{u \to \overline{u} \\ t \to \infty}} V^{(t)}(u) \geq \liminf_{u \to \overline{u}} V(u) \geq V(\overline{u}),$$

where the first inequality follows from Condition (i), and the second inequality follows from lower semicontinuity of V. Moreover, fixing $(u_t)_{t\in\mathbb{N}}$ to be the constant sequence $u_t=\bar{u}$, it follows from (ii) that $\limsup_{t} V^{(t)}(u_t) = \lim_{t} V^{(t)}(\bar{u}) \leq V(\bar{u})$. Invoking [73, Prop. 7.2], we conclude that $V^{(t)} \stackrel{\text{e}}{\to} V$, i.e., $V^{(t)}$ epi-converges to V. Secondly, from Condition (i) and the level-boundedness of V, it follows that $(V^{(t)})_{t\in\mathbb{N}}$ is eventually level-bounded [73, Ex. 7.32]. The claim then follows from [73, Thm. 7.33].

D. Deferred proofs

Proof of Lemma V.6.

Let $(z_t)_{t\in\mathbb{N}} = (x_t, s_t, \boldsymbol{\beta}_t, w_t)_{t\in\mathbb{N}}$ denote the stochastic process satisfying dynamics (25), for some initial state $z_0 \in \operatorname{\mathbf{dom}} V$. For ease of notation, let us define $V_t := V(z_t), t \in \mathbb{N}$. Due to nonnegativity

$$\mathbb{E}\left[\sum_{t=0}^{k-1} c \|x_t\|^2\right] \le \mathbb{E}\left[V_k + \sum_{t=0}^{k-1} c \|x_t\|^2\right]$$
$$= \mathbb{E}\left[V_k - V_0 + \sum_{t=0}^{k-1} c \|x_t\|^2\right] + V_0,$$

where the second equality follows from the fact that V_0 is deterministic. By linearity of the expectation, we can in turn write

$$\mathbb{E}\left[V_{k}-V_{0}+c\sum_{t=0}^{k-1}\|x_{t}\|^{2}\right] = \mathbb{E}\left[\sum_{t=0}^{k-1}V_{t+1}-V_{t}+c\|x_{t}\|^{2}\right]$$
$$=\sum_{t=0}^{k-1}\mathbb{E}\left[V_{t+1}-V_{t}+c\|x_{t}\|^{2}\right].$$

Therefore.

$$\mathbb{E}\left[c\sum_{t=0}^{k-1} \|x_t\|^2\right] - V_0 \le \sum_{t=0}^{k-1} \mathbb{E}\left[V_{t+1} - V_t\right] + c \,\mathbb{E}\left[\|x_t\|^2\right]. \tag{39}$$

Recall that β_t denotes the coordinate of β_t corresponding to the risk measures in the cost function (20). Defining the event $E_t := \{\omega \in \Omega \mid$ $P_{w_t(\omega)} \in \mathcal{A}_{\beta_t}(s_t(\omega), w_t(\omega))$, and its complement $\neg E_t = \Omega \setminus E_t$, we can use the law of total expectation to write

$$\mathbb{E}\left[V_{t+1} - V_{t}\right] = \mathbb{E}\left[V_{t+1} - V_{t} \mid E_{t}\right] \mathbb{P}[E_{t}] + \mathbb{E}\left[V_{t+1} - V_{t} \mid \neg E_{t}\right] \mathbb{P}[\neg E_{t}].$$

By condition (8), $\mathbb{P}[\neg E_t] < \beta_t$. From Conditions (i) and (iii), it follows that $z_t \in \operatorname{\mathbf{dom}} V$, $\forall t \in \mathbb{N}_{[0,k]}$ and that there exists a $\overline{V} \geq 0$ such that $V(z) \leq \overline{V}$, for all $z \in \operatorname{dom} V$. Therefore, $\mathbb{E}[V_{t+1} V_t \mid \neg \mathbb{E}_t] \leq \overline{V}$. Finally, by Condition (ii), $\mathbb{E}[V_{t+1} - V_t \mid E_t] \leq$ $\mathbb{E}[-c||x_t||^2 \mid E_t]$. Thus,

$$\mathbb{E}\left[V_{t+1} - V_t\right] \le \mathbb{E}\left[-c\|x_t\|^2 \mid E_t\right] \mathbb{P}[E_t] + \overline{V}\beta_t.$$

This allows us to simplify expression (39) as

$$\begin{split} & \mathbb{E}\left[c\sum_{t=0}^{k-1}\|x_t\|^2\right] - V_0 \\ & \leq \sum_{t=0}^{k-1} - c\,\mathbb{E}\left[\|x_t\|^2 \mid E_t\right]\,\mathbb{P}[E_t] + \overline{V}\beta_t + c\,\mathbb{E}\left[\|x_t\|^2\right] \\ & \leq \sum_{t=0}^{k-1} - c\,\mathbb{E}\left[\|x_t\|^2 \mid E_t\right]\,\mathbb{P}[E_t] + \overline{V}\beta_t \\ & + c\,\mathbb{E}\left[\|x_t\|^2 \mid E_t\right]\,\mathbb{P}[E_t] + c\,\mathbb{E}\left[\|x_t\|^2 \mid \neg E_t\right]\,\mathbb{P}[\neg E_t] \\ & = \sum_{t=0}^{k-1}\overline{V}\beta_t + c\,\mathbb{E}\left[\|x_t\|^2 \mid \neg E_t\right]\,\mathbb{P}[\neg E_t] \\ & \leq \sum_{t=0}^{k-1}\beta_t(\overline{V} + c\,\mathbb{E}\left[\|x_t\|^2 \mid \neg E_t\right]). \end{split}$$

Since $\operatorname{dom} V(\cdot, s_t, \boldsymbol{\beta}_t, w_t)$ was assumed to be compact and to contain the origin, there exists an $r \ge 0$ such that $||x_t||^2 \le r$, $\forall t \in$ $\mathbb{N}_{[0,k]}$. Therefore,

$$\mathbb{E}\left[\sum_{t=0}^{k-1} \|x_t\|^2\right] \le \frac{V_0}{c} + \left(\frac{\overline{V}}{c} + r\right) \sum_{t=0}^{k-1} \beta_t,$$

which remains finite as $k \to \infty$, since $(\beta_t)_{t \in \mathbb{N}}$ is summable. Thus, necessarily $\lim_{t\to\infty} \mathbb{E}[\|x_t\|^2] = 0$.

Proof of Theorem V.7.

First, note that using the monotonicity of coherent risk measures [34, Sec. 6.3, (R2)], a straightforward inductive argument allows us to show that under Condition (i),

$$\mathbf{T}\,\widehat{V}_N < \widehat{V}_N, \quad \forall N \in \mathbb{N}.$$
 (40)

Since $\overline{\mathcal{Z}} \subseteq \operatorname{dom} \widehat{V}_N$, recall that by definition (24), we have for any $z = (x, s, \beta, w) \in \overline{\mathcal{Z}}$ that

$$\widehat{V}_N(z) = \ell(x, \widehat{\kappa}_N(z), w) + \rho_{w,s}^{\beta} \left[\widehat{V}_{N-1} \left(\widehat{f}^{\widehat{\kappa}_N}(z, v), v \right) \right],$$

where β denotes the component of β corresponding to the cost. Therefore, we may write

$$\begin{split} & \rho_{w,s}^{\beta} \left[\widehat{V}_{N}(\widehat{f}^{\widehat{\kappa}_{N}}(z,v),v) \right] - \widehat{V}_{N}(z) \\ & = \rho_{w,s}^{\beta} \left[\widehat{V}_{N}(\widehat{f}^{\widehat{\kappa}_{N}}(z,v),v) \right] - \ell(x,\widehat{\kappa}_{N}(z),w) \\ & - \rho_{w,s}^{\beta} \left[\widehat{V}_{N-1}(\widehat{f}^{\widehat{\kappa}_{N}}(z,v),v) \right] \leq -\ell(x,\widehat{\kappa}_{N}(z),w) \leq -c \|x\|^{2}, \end{split}$$

where the first inequality follows by (40) and monotonicity of coherent risk measures. The second inequality follows from Condition (ii). Combined with Condition (iii), this implies that $V: z \mapsto \widehat{V}_N(z) + \delta_{\overline{Z}}(z)$ satisfies the conditions of Lemma V.6 and the assertion follows. \Box

For the following, it will be convenient to define $Q_N^{(t)}$ and Q_N as

$$Q_{N}^{(t)}(x, u, w) := \ell(x, u, w) + \rho_{s_{t}, w}^{\beta_{t}}[\widehat{V}_{N-1}^{(t+1)}(f(x, u, v), v)],$$

$$Q_{N}(x, u, w) := \ell(x, u, w) + \mathbb{E}_{P_{w}}[V_{N-1}(f(x, u, v), v)|x, w],$$
(41)

and let $\widehat{\mathcal{U}}^{(t)}(x,w) := \widehat{\mathcal{U}}(x,s_t,\boldsymbol{\beta}_t,w)$, so we may write

$$\widehat{V}_N^{(t)}(x,w) = \inf_{u \in \widehat{\mathcal{U}}^{(t)}(x,w)} Q_N^{(t)}(x,u,w) \text{ and} \qquad (42\text{a})$$

$$V_N(x, w) = \inf_{u \in \mathcal{U}(x, w)} Q_N(x, u, w). \tag{42b}$$

Proof of Theorem V.9.

We will show (27) by induction on N. For N=0, we have $\widehat{V}_0^{(t)}\equiv V_{\rm f}\equiv V_0$, thus, (27) holds with $\gamma_0^{(t)}=0,\, \forall t\in \mathbb{N}$, and the claim holds trivially. For the induction step, we define the events

$$A^{(t)} := \{ P_w : \in \bigcap_{i=1}^{n_{\beta}} \mathcal{A}_{\beta_{t,i}}(s_t, w), \ \forall w \in W \}, \tag{43a}$$

$$B_N^{(t)} := \{ \hat{V}_N^{(t)}(x, w) \ge V_N(x, w), \ \forall (x, w) \in \mathbb{R}^{n_x} \times W \},$$
 (43b)

for $N \in \mathbb{N}, t \in \mathbb{N}$. The induction hypothesis now reads

$$\mathbb{P}[B_{N-1}^{(t)}] \ge \gamma_{N-1}^{(t)} = d\sum_{k=0}^{N-1} \|\beta_{t+k}\|_1, \quad \forall t \in \mathbb{N},$$
 (44)

and our goal is to show that this implies that $\mathbb{P}[B_N^{(t)}] \geq \gamma_N^{(t)}, \forall t \in \mathbb{N}$. Given the occurrence of event $B_{N-1}^{(t+1)}$, the monotonicity of risk measures [34, Sec. 6.3, (R2)] ensures that $Q_N^{(t)}(x,u,w) \geq \ell(x,u,w) + \rho_{s_t,w}^{\beta_t}[V_{N-1}(f(x,u,v),v)]$, for all $(x,u,w) \in \mathbb{R}^{n_x} \times \mathbb{R}^{n_u} \times W$, and $t \in \mathbb{N}$. Furthermore, conditional on event $A^{(t)}$, (16) implies that $\rho_{s_t,w}^{\beta_t} \geq \mathbb{E}_{P_{w:}}$ and $\bar{\rho}_{s_t,w}^{\bar{\beta}_t,\hat{\alpha}_t} \geq \mathsf{AV@R}_{\alpha}^{P_w:}$, uniformly. Combining this fact with (5) and (19), we obtain the implication

$$B_{N-1}^{(t+1)}, A^{(t)} \implies \begin{cases} Q_N^{(t)}(x, w, u) \geq Q_N(x, w, u), \\ \widehat{\mathcal{U}}^{(t)}(x, w) \subseteq \mathcal{U}(x, w), \end{cases}$$

 $\forall (x, u, w) \in \mathbb{R}^{n_x} \times \mathbb{R}^{n_u} \times W$, and hence,

$$\begin{split} \widehat{V}_N^{(t)}(x,w) &= \min_{u \in \widehat{\mathcal{U}}^{(t)}(x,w)} Q_N^{(t)}(x,u,w) \geq \min_{u \in \widehat{\mathcal{U}}^{(t)}(x,w)} Q_N(x,u,w) \\ &\geq \min_{u \in \mathcal{U}(x,w)} Q_N(x,u,w) = V_N(x,w), \end{split}$$

which describes exactly the event $B_N^{(t)}$. Thus, we have shown that $\mathbb{P}[B_N^{(t)}] \geq \mathbb{P}[A^{(t)}, B_{N-1}^{(t+1)}]$. By the union bound, we now obtain

$$\mathbb{P}[B_N^{(t)}] \ge \mathbb{P}[A^{(t)}, B_{N-1}^{(t+1)}] \ge 1 - (\mathbb{P}[\neg A^{(t)}] + \mathbb{P}[\neg B_{N-1}^{(t+1)}])$$

$$> 1 - (d||\beta_t||_1 + \gamma_{N-1}^{(t+1)}),$$
(45)

where in the final inequality, $\mathbb{P}[\neg B_{N-1}^{(t+1)}]$ was bounded using the induction hypothesis (44) and $\mathbb{P}[\neg A^{(t)}]$ was replaced by another application of the union bound:

$$\mathbb{P}[\neg A^{(t)}] = \mathbb{P}\left[\exists w \in W : P_w : \notin \bigcap_{i=1}^{n_{\beta}} \mathcal{A}_{\beta_{t,i}}(s_t, w)\right]$$

$$\leq \sum_{w \in W} \sum_{i=1}^{n_{\beta}} \mathbb{P}[P_w : \notin \mathcal{A}_{\beta_{t,i}}(s_t, w)]$$

$$\leq d\sum_{i=1}^{n_{\beta}} \beta_{t,i} = d\|\beta_t\|_1. \tag{46}$$

Thus, substituting the expression for $\gamma_{N-1}^{(t+1)}$ from the induction hypothesis (44) into the result (45), we obtain that (27) holds with

$$\gamma_N^{(t)} := d\|\boldsymbol{\beta}_t\|_1 + \gamma_{N-1}^{(t+1)} = d\sum_{k=0}^N \|\boldsymbol{\beta}_{t+k}\|_1,$$

which establishes (i). Under the conditions of (ii), namely that (28) holds, it follows from definition (43a) that

$$\mathbb{P}\left[\neg A^{(t)}\right] = \mathbb{P}\left[P_{w:} \notin \mathcal{A}_{\beta_t^{\star}}(s_t, w)\right] \le d\beta_t^{\star},\tag{47}$$

with $\beta_t^{\star} := \max_{i \in \mathbb{N}_{[1,n_{\beta}]}} \{\beta_{t,i}\} = \|\beta_t\|_{\infty}$. Statement (ii) is then established by the same inductive argument, replacing the expression for $\gamma_{N-1}^{(t)}$ in (44), and replacing (46) with (47).

Proof of Theorem V.12.

By Assumption V.8, we have for N=0 that $\widehat{V}_0^{(t)}\equiv V_0\equiv \overline{V}_{\rm f}$ and there is nothing to prove. The general case, N>0, is proved by induction. Assume that (30) holds for some $N\geq 0$. We will now demonstrate that this implies that it also holds for N+1. To this end, we will show that the sequence $(Q_{N+1}^{(t)}(x,\cdot,w))_{t\in\mathbb{N}}$ and the function $Q_{N+1}(x,\cdot,w)$, satisfy the conditions of Lemma C.1. Under Assumption II.3, and using [73, Thm. 3.31], it follows from [32, Prop. 2] that Q_N and $Q_{N+1}^{(t)}$, are proper, lsc, and level-bounded in u locally uniformly in x, for all $w\in W$.

Let us introduce the shorthand for the worst-case conditional distribution $p_t^*(u) = (p_t^*, u)_{v \in W}$:

$$p_t^{\star}(u) \coloneqq \underset{p \in \mathcal{A}_{\beta_t}(s_t, w)}{\mathbf{argmax}} \sum_{v \in W} p_v \widehat{V}_N^{(t+1)}(f(x, u, v), v),$$

where we have omitted the dependence on the constant x and w. Combining Corollary V.10 with Assumption III.5, the Borel-Cantelli lemma [46, Thm. 4.3] guarantees that w.p. 1, there exists a finite $T_N \in \mathbb{N}$, such that for all $t > T_N$, $P_w \in \mathcal{A}_{\beta_{t,i}}(s_t,w)$, for all $w \in W$ and $i \in \mathbb{N}_{[1,n_\beta]}$, and furthermore, $\widehat{V}_N^{(t)} \geq V_N$, which implies that $Q_{N+1}^{(t)} \geq Q_{N+1}$. Moreover, by the induction hypothesis (i.e., (30) holds for N), there exists for every $\epsilon > 0$, a $T_\epsilon \geq T_N$, such that for all $t > T_\epsilon$,

$$\begin{split} Q_{N+1}^{(t)}(x,u,w) &- Q_{N+1}(x,u,w) \\ &= \sum_{v \in W} p_{t,v}^{\star}(u) \hat{V}_{N}^{(t+1)}(f(x,u,v),v) - P_{wv} V_{N}(f(x,u,v),v) \\ &\leq \sum_{v \in W} p_{t,v}^{\star}(u) (V_{N}(f(x,u,v),v) + \epsilon) - P_{wv} V_{N}(f(x,u,v),v) \\ &= \sum_{v \in W} (p_{t,v}^{\star}(u) - P_{wv}) V_{N}(f(x,u,v),v) + p_{t,v}^{\star}(u) \epsilon \\ &\leq \sum_{v \in W} \delta_{t} V_{N}(f(x,u,v),v) + \epsilon, \end{split} \tag{48}$$

where the final inequality is due to Assumption V.11 and the fact that for all $t>T_N$, $P_w\colon\in \mathcal{A}_{\beta_t}(w,s_t)$. As $\delta_t\to 0$, the first term in (48) can be made arbitrarily small by increasing t, provided that $V_N(f(x,u,v),v)<\infty$, for all $v\in W$, hence establishing pointwise convergence $Q_{N+1}^{(t)}\stackrel{\mathcal{P}}{\to}Q_{N+1}$ whenever $\operatorname{dom}V_N$ is RCI for (1), which in turn holds if \mathcal{X}_f is RCI by Proposition V.4. The sequence $(Q_{N+1}^{(t)}(x,\cdot,w))_{t\in\mathbb{N}}$ and the function $Q_{N+1}(x,\cdot,w)$ thus satisfy the conditions of Lemma C.1, which establishes (30) for N+1. \square