Jakub Rydval ⊠©

Technische Universität Wien, Vienna, Austria

— Abstract

The infinite-domain CSP dichotomy conjecture extends the finite-domain CSP dichotomy theorem to reducts of finitely bounded homogeneous structures. Every finitely bounded homogeneous structure is uniquely described by a universal first-order sentence, and every reduct of such a structure by a sentence of the logic SNP. By Fraïssé's Theorem, testing the existence of a finitely bounded homogeneous structure for a given universal first-order sentence is equivalent to testing the amalgamation property for the class of its finite models. The present paper motivates a complexity-theoretic view on the classification problem for finitely bounded homogeneous structures. We show that this metaproblem is EXPSPACE-hard or PSPACE-hard, depending on whether the input is specified by a universal sentence or a set of forbidden substructures. By relaxing the input to SNP sentences and the question to the existence of a structure with a finitely bounded homogeneous expansion, we obtain a different meta-problem, closely related to the question of homogenizability. We show that this second meta-problem is already undecidable, even if the input SNP sentence comes from the Datalog fragment and uses at most binary relation symbols. As a byproduct of our proof, we also get the undecidability of some other properties for Datalog programs, e.g., whether they can be rewritten in the logic MMSNP, whether they solve some finite-domain CSP, or whether they define a structure with a homogeneous Ramsey expansion in a finite relational signature.

2012 ACM Subject Classification Theory of computation \rightarrow Logic; Theory of computation \rightarrow Computational complexity and cryptography

Keywords and phrases constraint satisfaction problems, finitely bounded, homogeneous, amalgamation property, universal, SNP, homogenizable

Funding Jakub Rydval: This research was funded in whole or in part by the Austrian Science Fund (FWF) [I 5948]. For the purpose of Open Access, the authors have applied a CC BY public copyright licence to any Author Accepted Manuscript (AAM) version arising from this submission.

Acknowledgements The author thanks Manuel Bodirsky and Simon Knäuer for many inspiring discussions on the topic.

1 Introduction

Strict NP (SNP) is an expressive fragment of existential second-order logic and thus, by Fagin's Theorem, of the complexity class NP. If one only considers structures over a finite relational signature, then SNP can be obtained from the universal fragment of first-order logic simply by allowing existential quantification over relation symbols at the beginning of the quantifier prefix. In particular, universal first-order formulas themselves are SNP formulas. Despite the name, SNP already has the full power of NP [31]. In addition, this logic class has many connections to *Constraint Satisfaction Problems* (CSPs), which we use as the primary source of motivation for the present article. The CSP of a relational structure \mathfrak{B} , denoted by $CSP(\mathfrak{B})$, is (the membership problem for) the class of all finite structures which homomorphically map to \mathfrak{B} . Every computational decision problem is polynomial-time equivalent to a CSP [13]. Many practically relevant problems like Boolean satisfiability or graph colorability can even be formulated as a *finite-domain* CSP, i.e., where the *template* \mathfrak{B} can be chosen finite. The basic link from SNP to CSP is that every sentence of the monotone fragment of this logic defines a finite disjoint union of CSPs of (possibly

infinite) relational structures [9]. There are, however, some more nuanced connections, such as the one that led to the formulation of the *Feder-Vardi conjecture*, now known as the finitedomain CSP dichotomy theorem [51]. In their seminal work [31], Feder and Vardi showed that the *Monotone Monadic* fragment of SNP (MMSNP) exhibits a dichotomy between P and NP-completeness if and only if the seemingly less complicated class of all finite-domain CSPs exhibits such a dichotomy,¹ they also conjectured the latter to be true. The logic class MMSNP contains all finite-domain CSPs, and many other interesting combinatorial problems, e.g., the problem of deciding whether the vertices of a given graph can be 2coloured without obtaining any monochromatic triangle [41]. The Feder-Vardi conjecture was confirmed in 2017 independently by Bulatov and Zhuk [24, 50].

There is a yet unconfirmed generalization of the Feder-Vardi conjecture, to CSPs of reducts of finitely bounded homogeneous structures, formulated by Bodirsky and Pinsker in 2011 [20]. Here we refer to it as the Bodirsky-Pinsker conjecture. A structure with a finite relational signature is *finitely bounded* if the class of all finite structures embeddable into it is definable by a universal first-order sentence, and homogeneous if every isomorphism between two of its finite substructures extends to an automorphism. *Reducts* of such structures are obtained simply by removing some of the original relations. A prototypical example of a structure that satisfies both finite boundedness and homogeneity is $(\mathbb{Q}; <)$, the set of rational numbers equipped with the usual countable dense linear order without endpoints. It is a folklore fact that the class of reducts of finitely bounded homogeneous structures is closed under taking expansions of structures by first-order definable relations [9]. Roughly said, the condition imposed on the structures within the scope of the Bodirsky-Pinsker conjecture ensures that the CSP is in NP and that its template enjoys some of the universal-algebraic properties that have played an essential role in the proofs of the Feder-Vardi conjecture [7]. At the same time, it covers CSP-reformulations of many natural problems in qualitative reasoning, as well as all problems definable in MMSNP.

Every finitely bounded homogeneous structure is uniquely described by a universal firstorder sentence, and every reduct of such a structure by a sentence of the logic SNP. The CSPs of both kinds of structures are always definable in the monotone fragment of SNP. By Fraïssé's theorem, asking whether a given universal first-order sentence describes a finitely bounded homogeneous structure is equivalent to asking whether the class of its finite models has the Amalgamation Property (AP). This question has been considered many times in the context of the Lachlan-Cherlin classification programme for homogeneous structures [4, 35, 26], and is known to be decidable in the case of binary signatures [37, 16]. It also appears as an open problem in Bodirsky's book on infinite-domain constraint satisfaction [9]. Whether a given SNP sentence describes a reduct of a finitely bounded homogeneous structure is a different question, closely related to *homogenizability* [2, 5, 29, 33]. To the best of our knowledge, neither of the two questions is known to be decidable in general. Hence, it is unclear which CSPs actually fall within the scope of the Bodirsky-Pinsker conjecture. Besides CSPs, these questions are also relevant to other areas of theoretical computer science such as verification of database-driven systems [22] or description logics with concrete domains [38, 6].

The Bodirsky-Pinsker conjecture. In 2016, Bodirsky and Mottet presented an elegant tool for lifting tractability from finite-domain constraint satisfaction to the infinite [19],

¹ The correspondence between MMSNP and finite-domain CSP was initially only achieved up to randomized reductions, but it was later derandomized by Kun [36].

hereby establishing the first general link between the Feder-Vardi and the Bodirsky-Pinsker conjecture. Since then, their method has been used numerous times to prove new or reprove old complexity classification results for infinite-domain CSPs. One prominent such example is the universal-algebraic proof of the complexity dichotomy for MMSNP [17]. Conveniently enough, every MMSNP sentence defines a finite union of CSPs of structures within the scope of the Bodirky-Pinsker conjecture, so the two meta-questions from the previous paragraph were not relevant in this context. There is a prospect that the methods from [19] will also prove useful in proving a dichotomy for the even more general logic class *Guarded Monotone* SNP (GMSNP) introduced in [8]. Also GMSNP enjoys the above mentioned property of MMSNP [16], and hence avoids the two meta-questions.

However, outside of GMSNP there exists a regime where the methods from [19] definitely fall short, and where the two meta-questions become relevant. Consider for instance the dichotomy for *temporal CSPs*, i.e., for CSPs of structures with domain \mathbb{Q} and whose relations are definable by a Boolean combination of formulas of the form (x = y) or (x < y), obtained by Bodirsky and Kára in 2010 [15]. At the present time, these problems are already very well understood; tractable temporal CSPs can always be solved by an algorithm that repeatedly searches for a set of potential minimal elements among the input variables, where each instance of the search is performed using an oracle for a tractable finite-domain CSP. The latter is generally determined by the shape of the Boolean combinations. E.g., in the case of $CSP(\mathbb{Q}; \{(x, y, z) \in \mathbb{Q}^3 \mid (x = y < z) \lor (y = z < x) \lor (z = x < y)\})$, solving the finite-domain CSP in question amounts to solving linear equations modulo 2 [15, 21]. It is known that the tractability results from [15] cannot be obtained using the reduction from [19].

In 2022, Mottet and Pinsker introduced the machinery of smooth approximations [43], which vastly generalizes the methods in [19]. The last section of their paper is devoted to temporal CSPs, and the authors manage to reprove a significant part of the dichotomy on just a few pages. They achieve this by applying some of their general results to first-order expansions of $(\mathbb{Q}; <)$ and obtaining either NP-hardness for the CSP, or one of the two types of symmetry that played a fundamental role in the original proof from [15]. This symmetry can then be used to prove correctness of the reduction to a finite-domain CSP described above, but only under an explicit usage of the homogeneity of $(\mathbb{Q}; <)$ (see Proposition 3.1 in [21] and the last section of [43]). In contrast to the methods in [19] which only use homogeneity as a blackbox, this approach can be described as *language-dependent*.

A similar situation occurs in the case of phylogeny CSPs [14], which capture decision problems concerning the existence of a binary tree satisfying certain constraints imposed on its leaves. Tractable phylogeny CSPs are strikingly similar to tractable temporal CSPs; they can always be solved by an algorithm that repeatedly searches for a subdivision of the input variables into two parts, representing the two different branches below the root of a binary tree, where each instance of this search is performed using an oracle for a tractable finite-domain CSP. However, for tractable phylogeny CSPs, already the homogeneity of the infinite-domain CSP template is both sufficient and necessary for proving the correctness of the reduction to the finite-domain CSP (Theorem 6.13 and Lemma 6.12 in [14]). We can therefore speak of a case of extreme language-dependency. Temporal and phylogeny CSPs are special cases of CSPs of structures obtainable from the universal homogeneous binary tree [11] by specifying relations using first-order formulas. Achieving a complexity dichotomy in this context will require a non-trivial combination of the methods from [15] and [14].

An optimal way of approaching the conjecture would be to gain a very good understanding of the class of reducts of finitely bounded homogeneous structures, e.g., through some

sort of a classification. However, it is unclear how realistic this prospect is as model-theoretic properties often tend to be undecidable [25]. We remark that homogeneity is a vital part of the Bodirsky-Pinsker conjecture; this assumption can be weakened or strengthened but not dropped entirely, as otherwise we get a class that provably does not have a dichotomy [31, 9].

Contributions. In the present paper, we confirm the intractability of the two meta-questions. More specifically, we show that the first question is PSPACE-hard (Theorem 6) or EXPSPACE-hard (Theorem 5), depending on the encoding of the input, and that the second question is undecidable (Theorem 11). Theorems 6 and 5 are proved in Section 3.1 by taking a proof-theoretic perspective on the AP for classes defined by universal Horn sentences. We show that, for some of these classes, the failures of the AP are in a 1:1 correspondence with the rejecting runs of certain Datalog programs verifying instances of the rectangle tiling problem. Here, by Datalog we mean the monotone Horn fragment of SNP. Theorem 11 is proved in Section 4, by analyzing model-theoretic properties of a very natural encoding of context-free grammars into Datalog sentences. As a byproduct of the proof, we also get the undecidability of some other properties for Datalog programs, e.g., whether they can be rewritten in the logic MMSNP, whether they solve some finite-domain CSP, or whether they define a structure with a homogeneous Ramsey expansion in a finite relational signature.

Our results highlight the need for a fundamentally new language-independent approach to the Bodirsky-Pinsker conjecture. The first steps in this direction were taken in the recent works of Mottet and Pinsker [43] and Bodirsky and Bodor [10]. Yet, the issues mentioned in the previous paragraphs are, as of now, unaddressed. To keep our results as general as possible, we formulate them for some reasonable promise relaxations of the two questions, i.e., where a subclass and a superclass of the positive instances are being separated from each other with the promise that the input never belongs to the complement of the subclass within the superclass. Omitted proofs or their parts can be found in the appendix.

2 Preliminaries

Relational structures. The set $\{1, \ldots, n\}$ is denoted by [n], and we use the bar notation \bar{t} for tuples. A (relational) signature τ is a set of relation symbols, each $R \in \tau$ with an associated natural number called arity. We say that τ is binary if it consists of symbols of arity ≤ 2 . A (relational) τ -structure \mathfrak{A} consists of a set A (the domain) together with the relations $R^{\mathfrak{A}} \subseteq A^k$ for each $R \in \tau$ with arity k. An expansion of \mathfrak{A} is a σ -structure \mathfrak{B} with A = B such that $\tau \subseteq \sigma$, $R^{\mathfrak{B}} = R^{\mathfrak{A}}$ for each relation symbol $R \in \tau$. Conversely, we call \mathfrak{A} a reduct of \mathfrak{B} . The union of two τ -structures \mathfrak{A} and \mathfrak{B} is the τ -structure $\mathfrak{A} \cup \mathfrak{B}$ with domain $A \cup B$ and relations of the form $R^{\mathfrak{A} \cup \mathfrak{B}} := R^{\mathfrak{A}} \cup R^{\mathfrak{B}}$ for every $R \in \tau$.

A homomorphism $h: \mathfrak{A} \to \mathfrak{B}$ for τ -structures $\mathfrak{A}, \mathfrak{B}$ is a mapping $h: A \to B$ that preserves each relation of \mathfrak{A} , i.e., if $\overline{t} \in \mathbb{R}^{\mathfrak{A}}$ for some k-ary relation symbol $\mathbb{R} \in \tau$, then $h(\overline{t}) \in \mathbb{R}^{\mathfrak{B}}$. We write $\mathfrak{A} \to \mathfrak{B}$ if \mathfrak{A} maps homomorphically to \mathfrak{B} . The Constraint Satisfaction Problem (CSP) of \mathfrak{A} , denoted by CSP(\mathfrak{A}), is defined as the class of all finite structures which homomorphically map to \mathfrak{A} . An embedding is a homomorphism $h: \mathfrak{A} \to \mathfrak{B}$ that additionally satisfies the following condition: for every k-ary relation symbol $\mathbb{R} \in \tau$ and $\overline{t} \in A^k$ we have $h(\overline{t}) \in \mathbb{R}^{\mathfrak{B}}$ only if $\overline{t} \in \mathbb{R}^{\mathfrak{A}}$. We write $\mathfrak{A} \hookrightarrow \mathfrak{B}$ if \mathfrak{A} embeds to \mathfrak{B} . The age of \mathfrak{A} , denoted by age(\mathfrak{A}), is the class of all finite structures which embed to \mathfrak{A} . A substructure of \mathfrak{A} is a structure \mathfrak{B} over $\mathbb{B} \subseteq A$ such that the inclusion map $i: \mathbb{B} \to A$ is an embedding. An isomorphism is a surjective embedding. Two structures \mathfrak{A} and \mathfrak{B} are isomorphic if there exists an isomorphism from \mathfrak{A} to \mathfrak{B} . An automorphism is an isomorphism from \mathfrak{A} to \mathfrak{A} . The

orbit of a tuple $\bar{t} \in A^k$ in \mathfrak{A} is the set $\{g(\bar{t}) \mid g \text{ is an automorphism of } \mathfrak{A}\}$. A structure \mathfrak{A} is ω -categorical if, for every $k \geq 1$, there are only finitely many orbits of k-tuples in \mathfrak{A} .

First-order logic. We assume that the reader is familiar with classical *first-order* logic as well as with basic preservation properties of first-order formulas, e.g., that every firstorder formula ϕ is preserved by isomorphisms; by embeddings if ϕ is existential, and by homomorphisms if ϕ is existential positive. For a first-order sentence Φ , we denote the class of all its finite models by $fm(\Phi)$. We say that a first-order formula is k-ary if it has k free variables. For a first-order formula ϕ , we use the notation $\phi(\bar{x})$ to indicate that the free variables of ϕ are among \bar{x} . This does not mean that the truth value of ϕ depends on each entry in \bar{x} . We assume that equality = as well as the nullary predicate symbol \perp for falsity are always available when building first-order formulas. Thus, atomic τ -formulas, or τ -atoms for short, over a relational signature τ are of the form \perp , (x = y), and $R(\bar{x})$ for some $R \in \tau$. We say that a formula is *equality-free* if it does not contain any occurrence of the default equality predicate. If ϕ is a disjunction of possibly negated τ -atoms, then we define the Gaifman graph of ϕ as the undirected graph whose vertex set consists of all free variables of ϕ and where two distinct variables x, y form an edge if and only if they appear jointly in a negative atom of ϕ . Let Φ be a universal τ -sentence in prenex normal form whose quantifier-free part ϕ is in CNF. We call Φ Horn if every clause of ϕ is Horn, i.e., contains at most one positive disjunct. We call Φ complete if the Gaifman graph of each clause of ϕ is complete. It is a folklore fact that, if Φ is complete, then fm(Φ) is preserved by unions.

SNP and its fragments. An $SNP \tau$ -sentence is a second-order sentence Φ of the form $\exists X_1, \ldots, X_n \forall \bar{x}. \phi$ where ϕ is a quantifier-free formula in CNF over $\tau \cup \{X_1, \ldots, X_n\}$. We call Φ monadic if X_i is unary for every $i \in [n]$; monotone if ϕ does not contain any positive τ -atoms (in particular no positive equality atoms); and guarded if, for every positive atom β there exists a negative atom α containing all variables of β . Note that all notions from the previous paragraph easily transfer to SNP sentences viewed as universal sentences in an extended signature. The monadic monotone and the guarded monotone fragments of SNP are denoted by MMSNP and GMSNP, respectively. The monotone Horn fragment of SNP is commonly known as the logic programming language Datalog. When we say that a Datalog program Φ solves the CSP of a structure \mathfrak{B} , we simply mean that fm(Φ) = CSP(\mathfrak{B}).

Homogeneity, homogenizability, and finite boundedness. A countable structure \mathfrak{S} is homogeneous if every isomorphism between two finite substructures of \mathfrak{S} extends to an automorphism of \mathfrak{S} . Clearly, every homogeneous structure in a finite relational signature is ω -categorical. Homogeneous structures arise as limit objects of well-behaved classes of finite structures in the sense of Theorem 1. Let \mathcal{K} be a class of finite structures in a finite relational signature τ closed under isomorphisms and substructures. We say that \mathcal{K} has the amalgamation property (AP) if, for all $\mathfrak{B}_1, \mathfrak{B}_2 \in \mathcal{K}$ there exists $\mathfrak{C} \in \mathcal{K}$ with $f_1 \colon \mathfrak{B}_1 \hookrightarrow \mathfrak{C}$ and $f_2 \colon \mathfrak{B}_2 \hookrightarrow \mathfrak{C}$ such that $f_1|_{B_1 \cap B_2} = f_2|_{B_1 \cap B_2}$. We refer to \mathfrak{C} as an amalgam of \mathfrak{B}_1 and \mathfrak{B}_2 . Note that, for a class closed under isomorphisms and substructures, the AP is implied by the property of being closed under unions (also called *free amalgams*).

▶ Theorem 1 (Fraïssé). For a class \mathcal{K} of finite structures in a finite relational signature τ , the following are equivalent:

 $- \mathcal{K}$ is the age of an up to isomorphism unique countable homogeneous τ -structure;

 \blacksquare \mathcal{K} is closed under isomorphisms, substructures, and has the AP.

As already mentioned in the introduction, the structure $(\mathbb{Q}; <)$ is homogeneous because every local isomorphism can be extended to an automorphism using a piecewise affine transformation. Its age is the class of all finite strict linear orders.

A countable structure \mathfrak{S} is homogenizable if it is a reduct of a homogeneous structure \mathfrak{H} over a finite relational signature such that \mathfrak{S} and \mathfrak{H} have the same sets of automorphisms [29]. Whenever this happens, by the theorem of Ryll-Nardzewski, all relations of \mathfrak{H} are firstorder definable in \mathfrak{S} [32]. One might say that \mathfrak{S} already has all the relations necessary for homogeneity but they perhaps do not all have names. A prototypical example of this phenomenon is the universal "homogeneous" binary tree, which is homogenizable but not homogeneous, see, e.g., Proposition 3.2 in [11]. We call a class \mathcal{K} of finite structures in a finite relational signature τ homogenizable if it forms the age of a homogenizable structure.

For a class \mathcal{N} of finite structures in a finite relational signature τ , the class $\operatorname{Forb}_e(\mathcal{N})$ consists of all finite τ -structures which do not embed any member of \mathcal{N} . Following the terminology in [39], we say that a class \mathcal{K} of finite structures in a finite relational signature is *finitely bounded* if there exists a finite \mathcal{N} such that $\mathcal{K} = \operatorname{Forb}_e(\mathcal{N})$. We refer to \mathcal{N} as a set of *bounds* for \mathcal{K} , and define the *size* of \mathcal{N} as the sum of the cardinalities of the domain and the relations of all structures in \mathcal{N} . A structure \mathfrak{S} is *finitely bounded* if its age is finitely bounded. We say that a class \mathcal{K} is *finitely bounded homogenizable* if it forms the age of a reduct \mathfrak{R} of a finitely bounded homogeneous structure \mathfrak{H} such that \mathfrak{R} and \mathfrak{H} have the same sets of automorphisms. Sufficient conditions for finitely bounded homogenizability were provided by Hubička and Nešetřil [34], generalizing previous work of Cherlin, Shelah, and Shi [27].

3 The first meta-problem

By Theorem 1, every homogeneous structure is uniquely described by its age. Consequently, every finitely bounded homogeneous structure is uniquely described by a finite set of bounds. It is known that the question whether $\operatorname{Forb}_e(\mathcal{N})$ has the AP for a given finite set of bounds \mathcal{N} can be tested algorithmically in the case where the signature is binary [37]. This decidability result is based on the following observation. A *one-point amalgamation diagram* is an input $\mathfrak{B}_1, \mathfrak{B}_2$ to the AP where $|B_1 \setminus B_2| = |B_2 \setminus B_1| = 1$.

▶ Proposition 2 ([37]). A class of finite relational τ -structures that is closed under isomorphisms and substructures has the AP if and only if it has the AP restricted to one-point amalgamation diagrams.

As a consequence of Proposition 2, if τ is binary and $\operatorname{Forb}_e(\mathcal{N})$ does not have the AP, then the size of a smallest counterexample to the AP is polynomial in the size of \mathcal{N} [16]. Such a counterexample can be non-deterministically guessed and verified using a coNP-oracle, which places the problem at the second level of the polynomial hierarchy (Theorem 15 in [6]).

There is a second, arguably more practical, equivalent definition of finite boundedness. Namely, a class \mathcal{K} of finite structures in a finite relational signature is finitely bounded if and only if there exists a universal sentence Φ such that $\mathcal{K} = \operatorname{fm}(\Phi)$. Using this definition, it is easy to see that (\mathbb{Q} ; <) is finitely bounded because its age, the class of all finite strict linear orders, admits a finite universal axiomatization (irreflexivity, transitivity, and totality). From a complexity-theoretical perspective, the two definitions are equivalent only up to a single-exponential blow-up in one direction. Given a finite set of bounds \mathcal{N} , we can obtain a universal sentence Φ of size polynomial in the size of \mathcal{N} satisfying $\operatorname{fm}(\Phi) = \operatorname{Forb}_e(\mathcal{N})$ by describing each structure in \mathcal{N} up to isomorphism using a quantifier-free formula. However,

given a universal sentence Φ , it can be the case that a smallest \mathcal{N} satisfying $\operatorname{fm}(\Phi) = \operatorname{Forb}_e(\mathcal{N})$ is of size single-exponential in the size of Φ . The reason is that obtaining \mathcal{N} from Φ is comparable to rewriting Φ in DNF. Consequently, the algorithm from [16] only gives us a relatively weak upper bound for the case where the inputs are specified by universal sentences.

▶ **Proposition 3.** Let Φ be a universal sentence over a finite binary relational signature τ . If fm(Φ) does not have the AP, then the size of a smallest counterexample to the AP is at most single-exponential in the size of Φ . Consequently, the question whether fm(Φ) has the AP is decidable in coNEXPTIME.

The upper bound provided by Proposition 3 is not unreasonable since a smallest counterexample to the AP might be of size exponential in the size of the input sentence even if the signature is binary. Very little progress has been done on signatures containing symbols of arities larger than 2. In particular, it is not even known whether the AP is decidable for finitely bounded classes in general. The scenario where this is not the case is not unrealistic since the closely related joint embedding property is undecidable already for finitely bounded classes of graphs [23]. If the AP turns out to be undecidable as well (for finitely bounded classes), then the Bodirsky-Pinsker conjecture addresses a class of structures with an undecidable membership problem. However, it seems that, at least to some extent, the decidability issue can be ignored by only using homogeneity as a blackbox. This was demonstrated in the recent work [44] on the complexity of CSPs of homogeneous uniform hypergraphs, whose classification remains an open problem [4]. We remark that the question is already open in the following case, where we can only prove PSPACE-hardness.

▶ **Theorem 4.** Given a universal Horn sentence Φ over a finite relational signature that is binary except for one ternary symbol, the question whether fm(Φ) has the AP is PSPACE-hard.

The next theorem states that testing the AP becomes properly harder than in the binary case if we do not impose any restrictions on the input (unless the exponential hierarchy collapses). The fact that Theorem 5 also holds for the strong version of the AP might be of independent interest to model-theorists. The *strong* version of the AP is when $\mathfrak{C} \in \mathcal{K}$ and $f_i: \mathfrak{B}_i \hookrightarrow \mathfrak{C}$ for $i \in \{1, 2\}$ can always be chosen so that $f_1(B_1) \cap f_2(B_2) = f_1(B_1 \cap B_2)$.

Theorem 5. Given a universal sentence Φ over a finite signature, the question whether $fm(\Phi)$ has the strong AP or not even the AP is EXPSPACE-hard, even when the number of variables in Φ is bounded by a constant.

The next theorem is a variant of Theorem 5 in the setting where the input is specified by a set of bounds instead of a universal sentence. This setting can be compared to the situation where, in Theorem 5, the quantifier-free part of Φ is required to be in DNF. As a consequence, we cannot profit from succintness of general universal sentences, which leads to a weaker PSPACE lower bound on the complexity.

▶ **Theorem 6.** Given a finite set \mathcal{N} of finite structures over a finite signature, the question whether Forb_e(\mathcal{N}) has the strong AP or not even the AP is PSPACE-hard, even when the domain size for the structures in \mathcal{N} is bounded by a constant.

3.1 Proofs of Theorem 5 and Theorem 6

Our proofs of Theorem 5 and Theorem 6 are based on the fact that, if Φ is a universal Horn sentence such that fm(Φ) does not have the AP, then every counterexample to the AP has

the form of a particular Horn clause which can be derived from Φ in a syntactical manner. By padding each Horn clause in Φ with auxiliary negative atoms and hereby increasing its "degree of completeness," we gain some control over the form of the counterexamples to the AP. When this is performed in a careful and systematic way, the counterexamples to the AP can be brought into a 1:1 correspondence with the rejecting runs of certain Datalog programs. In our case, such programs verify the validity of tilings w.r.t. given input parameters to a bounded tiling problem. This is the main technical contribution of the present article.

Let Φ be an equality-free universal Horn sentence over a relational signature τ , and let $\phi(\bar{x}), \psi(\bar{x}, \bar{y})$ be equality-free conjunctions of atomic τ -formulas. We write $\psi(\bar{x}, \bar{y}) \leq_{\Phi} \phi(\bar{x})$ if, for every atomic τ -formula $\chi(\bar{x})$ other than equality, $\Phi \models \forall \bar{x}, \bar{y}(\psi(\bar{x}, \bar{y}) \Rightarrow \chi(\bar{x}))$ implies $\Phi \models \forall \bar{x}(\phi(\bar{x}) \Rightarrow \chi(\bar{x}))$.

Lemma 7. Let Φ be an equality-free universal Horn sentence over a relational signature τ . Then the following are equivalent:

- **1.** fm(Φ) has the strong AP.
- **2.** fm(Φ) has the AP.
- **3.** If $\phi(\bar{x})$, $\phi_1(\bar{x}, y_1)$, and $\phi_2(\bar{x}, y_2)$ are equality-free conjunctions of atomic formulas, where y_1 and y_2 are distinct variables not contained in \bar{x} , such that, for both $i \in \{1, 2\}$, every atom in $\phi_i(\bar{x}, y_i)$ contains the variable y_i and $\phi(\bar{x}) \land \phi_i(\bar{x}, y_i) \leq_{\Phi} \phi(\bar{x})$, then $\phi(\bar{x}) \land \phi_1(\bar{x}, y_1) \land \phi_2(\bar{x}, y_2) \leq_{\Phi} \phi(\bar{x}) \land \phi_1(\bar{x}, y_1)$.

Let Φ be an equality-free universal Horn sentence and ψ a Horn clause over a relational signature τ . An *SLD-derivation of* ψ from Φ is a finite sequence of Horn clauses $\psi_0, \ldots, \psi_s = \psi$ such that ψ_0 is a conjunct in Φ and, for every $i \in [s]$, there exists a Horn clause ϕ_i which is, up to renaming of variables, a conjunct in Φ , and such that ψ_i is obtained from ψ_{i-1} by replacing a negative atom of ψ_{i-1} that appears positively in ϕ_i with all negative atoms of ϕ_i . We say that ψ_i is a *resolvent* of ψ_{i-1} and ϕ_i . We call ψ a *weakening* of a clause ψ' if ψ' can be obtained from ψ by removing any amount of atoms. In particular, ψ is a weakening of itself. There exists an *SLD-deduction* of ψ from Φ , written as $\Phi \vdash \psi$, if ψ is a tautology or a weakening of a Horn clause ψ' that has an SLD-derivation from Φ . The following theorem presents a fundamental property of equality-free universal Horn sentences.

▶ **Theorem 8** (Theorem 7.10 in [45]). Let Φ be an equality-free universal Horn sentence and ψ an equality-free Horn clause, both in a fixed signature τ . Then $\Phi \models \psi$ if and only if $\Phi \vdash \psi$.

The hardness proofs are by polynomial-time reductions from the complements of two well-known bounded versions of the tiling problem. Consider the signature τ consisting of the two binary symbols P_h, P_v , as well as the four unary symbols P_ℓ, P_r, P_t, P_b . For natural numbers $m, n \geq 1$, the τ -structure $\Re_{m,n}$ has the domain $[m] \times [n]$ and relations

$$\begin{split} P_{\ell}^{\mathfrak{R}_{m,n}} &\coloneqq [m] \times \{1\}, \; P_{r}^{\mathfrak{R}_{m,n}} \coloneqq [m] \times \{n\}, \; P_{b}^{\mathfrak{R}_{m,n}} \coloneqq \{1\} \times [n], \; P_{t}^{\mathfrak{R}_{m,n}} \coloneqq \{m\} \times [n], \\ P_{h}^{\mathfrak{R}_{m,n}} &\coloneqq \{((i,j), (i+1,j)) \mid i \in [n-1], j \in [m]\}, \\ P_{v}^{\mathfrak{R}_{m,n}} &\coloneqq \{((i,j), (i,j+1)) \mid i \in [n], j \in [m-1]\}. \end{split}$$

The rectangle tiling problem asks whether, given a natural number n and a finite τ -structure \mathfrak{T} , there exists a natural number m such that $\mathfrak{R}_{m,n} \to \mathfrak{T}$. In contrast to the better-known NP-complete square tiling problem, one dimension of the tiling grid is not part of the input and is existentially quantified instead. As a result, the problem becomes PSPACE-complete [28, 49].² One can further increase the complexity by allowing a succinct encoding

 $^{^{2}}$ In [49], this problem is called the corridor tiling problem.

of the space bound. The input remains the same but now we ask for a rectangle tiling with 2^n columns. Analogously to the natural complete problems based on Turing machines, this yields a decision problem that is complete for the complexity class EXPSPACE [47].

Theorem 5 is proved by polynomial-time reduction from the complement of the exponential rectangle tiling problem. From every input, we construct a universal sentence Φ of polynomial size such that fm(Φ) has the AP if and only if there exists no exponential rectangle tiling satisfying the given parameters. The sentence Φ is almost Horn but, sometimes, disjunctions and equality atoms are used in premises of implications to represent exponentially many Horn clauses in a universal sentence of polynomial size. In the text that follows, we allow ourselves to still call such sentences Horn. Our encoding is very compact; each row, i.e., an ordered sequence of 2^n -many tiles, is represented using a constant amount of variables. This is achieved by storing the information about each individual row in binary using (n + 1)-ary atoms whose entries always contain at most three variables. We refer to the variables representing rows of the tiling as *path* nodes. In order to check the tiling from bottom to top, i.e., parse a chain of path nodes, we require each pair of subsequent path nodes to be verified by a set of 2^n -many *verifier* nodes. This process ensures the vertical consistency of the tiling as well as the presence of 2^n -many tiles in every row. The precise number of verifier nodes is achieved using combinations of n pairs of unary atoms.

To control the occurrence of amalgamation failures, we first introduce a binary symbol E and two unary symbols L, R. Atoms with these symbols serve no other purpose than to ensure that *almost* each conjunct in Φ is complete, i.e., defines a class of structures that is preserved by taking unions and hence has the AP. More concretely, the premise of almost every Horn clause in Φ has a subformula of the form $L(y_1) \wedge R(y_2) \wedge \bigwedge_{i \in [k]} E(y_1, x_i) \wedge E(y_2, x_i) \wedge \bigwedge_{j \in [k]} E(x_i, x_j)$ making the Horn clause almost complete, with the exception of one potentially missing edge in the Gaifman graph between y_1 and y_2 . Our intention is to make this missing edge the only place at which potential faulty one-point amalgamation diagrams can be built (see Figure 1). The sentence Φ is defined as $\Phi_1 \wedge \Phi_2$, where the two parts are described below.

The first part Φ_1 does not yet explain how our reduction works, but ensures that it does not fall apart, e.g., due to ill-behaved identifications of variables. For every $\alpha \in T$, the signature τ contains an (n + 1)-ary symbol T_{α} . The first n arguments in a T_{α} -atom serve as binary counters, and the last argument carries a given path node p. Suppose that the variables 0 and 1 represent the bits 0 and 1, respectively. Then each atomic formula $T_{\alpha}(c_1,\ldots,c_n,p)$ with $c_1,\ldots,c_n \in \{0,1\}$ represents the situation in which a tile α is present in the *p*-th row and in the $(1+\sum_{k\in[n]}(c_k=1)\cdot 2^{n-k})$ -th column. First, we want to ensure the horizontal consistency of the tiling. To this end, for every pair $(\alpha, \beta) \in T^2 \setminus P_h^{\mathfrak{T}}$, we include in Φ_1 a complete Horn sentence without positive atoms ensuring that two horizontally adjacent positions in the p-th row cannot be tiled with α and β . Here, we encode the successor relation w.r.t. binary addition using a combination of equalities: (c_{n+1},\ldots,c_{2n}) is the successor of (c_1, \ldots, c_n) if and only if there exists $j \in [n]$ such that $c_i = c_{i+n} \in \{0, 1\}$ for every $i \in [j-1]$, $c_j = 0$ and $c_{n+j} = 1$, and $c_i = 1$ and $c_{i+n} = 0$ for every $i \in [n] \setminus [j]$. This encoding only makes sense if 0 and 1 truly represent the bits 0 and 1, so we introduce a simple mechanism (in terms of a Horn sentence) for distinguishing between the variables 0 and 1. Next, we want to ensure that every position in the p-th row is occupied by at most one tile. To this end, for every $(\alpha, \beta) \in T^2$ with $\alpha \neq \beta$, we include

$$\forall p, 0, 1, c_1, \dots, c_n \big(T_\alpha(c_1, \dots, c_n, p) \land T_\beta(c_1, \dots, c_n, p) \land \bigwedge_{i \in [n]} (c_i = 0 \lor c_i = 1) \Rightarrow \bot \big)$$

as a conjunct in Φ_1 . Finally, we want to ensure that each verifier node v represents at most

one number from $[2^n]$. For every $i \in [n]$, τ contains two unary symbols \mathbf{O}_i and $\mathbf{1}_i$ which will be used to encode numbers in binary. We include $\forall v (\bigvee_{i \in [n]} \mathbf{O}_i(v) \land \mathbf{1}_i(v) \Rightarrow \bot)$ as the last conjunct in Φ_1 . Now the idea is that the combinations of atomic formulas $\mathbf{O}_i(v)$ and $\mathbf{1}_i(v)$ at a verifier node v will be compared with the combinations of 0 and 1 in atomic formulas $T_{\alpha}(c_1, \ldots, c_n, p)$. The Horn sentences in the second part of Φ will be formulated so that verifying the presence of all 2^n atoms of the form $T_{\alpha}(c_1, \ldots, c_n, p_1)$ at a path node p_1 using verifier nodes v_1, \ldots, v_{2^n} is the only possible way to progress to a next path node p_2 . As a consequence, we do not need to add an explicit requirement for rows, represented by path nodes, to be completely tiled from left to right. For the same reason, we also do not need to add an explicit requirement for verifier nodes to represent at least one number from $[2^n]$.

We now proceed with the sentence Φ_2 , which explains how the parsing of a tiling actually works. The parsing of a tiling starts from a path node p representing a row whose leftmost position contains a tile that can be present in the bottom left corner of a tiling grid. This must be confirmed by a verifier node, in which case a 6-ary Q_b -atom is derived, representing the fact that the leftmost column of the p-th row has been checked. To this end, we include in Φ_2 suitable Horn sentences for every $\alpha \in P_{\ell}^{\mathfrak{T}} \cap P_b^{\mathfrak{T}}$. These sentences form the non-complete part of Φ ; we intentionally leave a missing edge between y_1 and y_2 in the Gaifman graph to enable the formation of potential AP-counterexamples. Using 2^n -many verifier nodes and propagation of Q_b -atoms, the whole bottom row is checked for the presence of tiles. Their horizontal consistency already follows from the conditions imposed on path nodes by Φ_2 and needs not to be checked during this step. To this end, we include in Φ_2 suitable Horn sentences for every $\alpha \in P_{b}^{\mathfrak{T}}$. After the *p*-th row has been checked by a 2^{*n*}-th verifier node, we mark p with a Q-atom indicating that the parsing can progress to a successor path node. To this end, we include in Φ_2 suitable Horn sentences for every $\alpha \in P_b^{\mathfrak{T}} \cap P_r^{\mathfrak{T}}$. The successor relation for path nodes is represented by the binary symbol S, and the certificate of vertical verification for pairs (p_1, p_2) of successive path nodes is represented by the 7-ary symbol Q_v . For every $(\alpha, \beta) \in P_v^{\mathfrak{T}} \cap (P_\ell^{\mathfrak{T}})^2$, we include in Φ_2 a Horn sentence verifying the vertical consistency of the leftmost positions in the rows p_1 and p_2 and deriving the first Q_v -atom containing p_1 and p_2 , but only if there is a Q-atom containing p_1 . Next, for every $(\alpha,\beta) \in P_v^{\mathfrak{T}}$, we include in Φ_2 a Horn sentence verifying the vertical consistency for the intermediate positions by deriving further Q_v -atoms containing p_1 and p_2 . And finally, for every $(\alpha, \beta) \in P_v^{\mathfrak{T}} \cap (P_r^{\mathfrak{T}})^2$, we include in Φ_2 a Horn sentence verifying the vertical consistency of the leftmost positions in the rows p_1 and p_2 and deriving the first Q-atom containing p_2 only. The top row is verified using a 6-ary symbol Q_t similarly as the bottom row; however, the verification of the rightmost position in the top row results in the derivation of \perp .

Proof of Theorem 5. We first argue that Φ is equivalent to a particular Horn sentence, to which we can apply Lemma 7. By construction, the quantifier-free part of each conjunct in Φ has the form of an implication where the premise possibly also contains instances of disjunction, which are not allowed in Horn clauses, but no instances of negation. Therefore, it can be rewritten as a conjunction of Horn clauses by converting the premise into positive DNF and then considering each disjunct as a separate premise. As a result, the size of the sentence increases exponentially, but this does not matter for the purpose of the proof. Subsequently, all equality atoms can be eliminated by replacing each variable c_i with either 0 or 1. We denote the resulting Horn sentence by $\overline{\Phi}$ and the two parts stemming from Φ_1 and Φ_2 by $\overline{\Phi}_1$ and $\overline{\Phi}_2$, respectively.

"⇒" Suppose that there exists a tiling $f: \mathfrak{R}_{m,2^n} \to \mathfrak{T}$. Guided by f, we define a onepoint amalgamation diagram $\mathfrak{B}_1, \mathfrak{B}_2 \in \mathrm{fm}(\overline{\Phi})$ which has no amalgam in $\mathrm{fm}(\overline{\Phi})$ (see Figure 1). The domains are $B_i := \{y_i, p_1, \ldots, p_m, v_1, \ldots, v_{2^n}, 0, 1\}$ for $i \in \{1, 2\}$, and the relations are




Figure 1 An illustration of an AP-counterexample representing a valid tiling of an exponential rectangle with m rows and 2^n columns.

given by the following conjunctions of atomic formulas. We require $T_{\alpha}(c_1, \ldots, c_n, p_j)$ for $c_1, \ldots, c_n \in \{0, 1\}$ if and only if $f(1 + \sum_{k \in [n]} (c_k = 1) \cdot 2^{n-k}, j) = i$. Next, we require all of the *L*, *R*, and *E*-atoms necessary for enabling the Horn clauses in $\mathfrak{B}_1 \cup \mathfrak{B}_2$. Finally, we require $\bigwedge_{i \in [m-1]} S(p_i, p_{i+1})$ to define a successor chain through path nodes, and $O_i(v_j)$ or $\mathbf{1}_i(v_j)$ if and only if $j = 1 + \sum_{k \in [n]} \lambda_k \cdot 2^{n-k}$ for $\lambda_1, \ldots, \lambda_n \in \{0, 1\}$ and $\lambda_i = 0$ or $\lambda_i = 1$, respectively. Clearly, the tiling atoms are placed correctly and the verifier nodes correctly represent values in $[2^n]$. Since $R^{\mathfrak{B}_1} = \emptyset$ and $L^{\mathfrak{B}_2} = \emptyset$, we have $\mathfrak{B}_1, \mathfrak{B}_2 \models \overline{\Phi}_2$. Since f is horizontally consistent, we have $\mathfrak{B}_1, \mathfrak{B}_2 \models \overline{\Phi}_1$, i.e., $\mathfrak{B}_1, \mathfrak{B}_2 \in \mathrm{fm}(\overline{\Phi})$. But since f is also vertically consistent and $\overline{\Phi}_2$ is a universal Horn sentence, we have $\mathfrak{C} \not\models \overline{\Phi}_2$ for every τ -structure \mathfrak{C} with a homomorphism from $\mathfrak{B}_1 \cup \mathfrak{B}_2$. Hence, $\mathrm{fm}(\overline{\Phi})$ does not have the AP.

" \Leftarrow " Suppose that fm($\overline{\Phi}$) does not have the AP. Then there exists a counterexample to item (3) in Lemma 7, i.e., there exists a Horn clause ψ of the form $\phi(\bar{x}) \wedge \phi_1(\bar{x}, y_1) \wedge \phi_2(\bar{x}, y_2) \Rightarrow \chi$, where ϕ, ϕ_1 , and ϕ_2 satisfy the prerequisites of item (3) in Lemma 7 and $\chi(\bar{x}, y_1)$ is an atomic τ -formula other than equality, such that

$$\overline{\Phi} \models \forall \overline{x}, y_1, y_2 (\phi \land \phi_1 \land \phi_2 \Rightarrow \chi), \tag{1}$$

$$\Phi \not\models \forall \bar{x}, y_1 (\phi \land \phi_1 \Rightarrow \chi). \tag{2}$$

We choose ψ minimal with respect to the number of its atomic subformulas. By Theorem 8, ψ has an SLD-deduction from $\overline{\Phi}$. Note that, by (2), $\chi(\bar{x}, y_1)$ cannot be a subformula of $\phi(\bar{x}) \wedge \phi_1(\bar{x}, y_1)$. Also, $\chi(\bar{x}, y_1)$ cannot be a subformula of $\phi_2(\bar{x}, y_2)$ because every atom in $\phi_2(\bar{x}, y_2)$ contains the variable y_2 which does not appear in $\chi(\bar{x}, y_1)$. Hence, $\chi(\bar{x}, y_1)$ is not a subformula of $\phi(\bar{x}) \wedge \phi_1(\bar{x}, y_1) \wedge \phi_2(\bar{x}, y_2)$, i.e., ψ is not a tautology. Consequently, ψ is a weakening of a Horn clause ψ' which has an SLD-derivation $\psi'_0, \ldots, \psi'_s = \psi'$ from $\overline{\Phi}$. Recall that every atom from ψ' appears in ψ .

 \triangleright Claim 9. $\overline{\Phi}_2 \vdash \psi'$.

Proof. We start by showing that ψ'_0 is a conjunct of $\overline{\Phi}_2$. Suppose, on the contrary, that ψ'_0 is a conjunct of $\overline{\Phi}_1$. Then ψ' does not contain any positive atom because, by construction, Horn sentences in $\overline{\Phi}_1$ do not contain any positive atoms. By construction, in $\overline{\Phi}$ there is no Horn clause containing a positive atom that occurs negatively in a Horn clause from $\overline{\Phi}_1$, i.e., it is impossible to take resolvents of ψ'_0 and Horn clauses from $\overline{\Phi}$. It follows that s = 0. Also, every Horn clause from $\overline{\Phi}_1$ is complete. Since there is no edge between y_1 and y_2 in the

Gaifman graph of ψ , and, for $i \in \{1, 2\}$, each atom in ϕ_i contains the variable y_i , either ϕ_1 or ϕ_2 must be empty. Since ψ' does not contain any positive atom and $\phi(\bar{x}) \wedge \phi_i(\bar{x}, y_i) \leq \overline{\Phi} \phi(\bar{x})$ for both $i \in \{1, 2\}$, we get a contradiction to (2). Thus, ψ'_0 must be a conjunct of $\overline{\Phi}_2$. Since no Horn clause in $\overline{\Phi}_1$ contains a positive atom, no Horn clause from $\overline{\Phi}_1$ can be used as a resolvent. We conclude that ψ' has an SLD-derivation from $\overline{\Phi}_2$.

 \triangleright Claim 10. ψ' contains no positive atoms, and no atoms with a symbol from $\{Q_b, Q_v, Q_t, Q\}$.

Proof. By the construction of $\overline{\Phi}_2$, for every $i \in [s]$, if ψ'_{i-1} contains variables z_1, z_2 such that

every atomic subformula with a symbol from $\{Q_b, Q_v, Q_t, Q\}$ contains z_1 in its 1st and z_2 in its 2nd argument, respectively, (3)

then this is also the case for ψ'_i , for the same variables z_1, z_2 up to renaming. Since every possible choice of ψ'_0 from $\overline{\Phi}_2$ initially satisfies (3), it follows via induction that (3) holds for $\psi' = \psi'_s$ for some variables z_1, z_2 . Next, we show that $\{z_1, z_2\} = \{y_1, y_2\}$ holds for the pair z_1, z_2 satisfying (3) for ψ' . Suppose, on the contrary, that both z_1 and z_2 are among \bar{x}, y_1 or \bar{x}, y_2 . By construction, the only Horn clauses in $\overline{\Phi}_2$ that are not complete have the property that the incompleteness is only due to one missing edge in the Gaifman graph between two distinguished variables satisfying (3). Therefore, for every $i \in [s], \psi'_i$ is a resolvent of ψ'_{i-1} and a Horn clause from $\overline{\Phi}_2$ which is almost complete except possibly for one missing edge in the Gaifman graph between a pair of variables which must be substituted for the pair (z_1, z_2) satisfying (3) for ψ'_{i-1} . Since the variables y_1 and y_2 do not appear together in any atom in ψ' and $\{z_1, z_2\} \neq \{y_1, y_2\}$, they also do not appear together in any atom during the SLD-derivation. Then it follows from the fact that ϕ, ϕ_1 , and ϕ_2 satisfy the prerequisites of item (3) in Lemma 7 that we already have $\overline{\Phi}_2 \vdash \forall \overline{x}, y_1 (\phi \land \phi_1 \Rightarrow \chi)$, a contradiction to (2). Since $\{z_1, z_2\} = \{y_1, y_2\}$ holds for the pair z_1, z_2 satisfying (3) for ψ', ψ' cannot contain any negative atoms with a symbol from $\{Q_b, Q_v, Q_t, Q\}$. Suppose that the conclusion of ψ'_0 is not \perp . Then, by construction, ψ' contains a positive atom with a symbol from $\{Q_b, Q_v, Q_t, Q\}$. But then, since z_1, z_2 with $\{z_1, z_2\} = \{y_1, y_2\}$ satisfy (3) for ψ' , the said positive atom in ψ' contains both variables y_1 and y_2 . This leads to a contradiction to (1) where we assume that the positive atom in ψ may may only contain variables from \bar{x}, y_1 . Thus the conclusion of ψ'_0 is \perp , which means that χ equals \perp due to the minimality assumption.

It remains to show that the existence of such ψ' implies the existence of a tiling $f: \mathfrak{R}_{m,2^n} \to$ \mathfrak{T} . By the first and the second claim, the conclusion of ψ'_0 is \perp . Since ψ' does not contain any atoms with a symbol from $\{Q_b, Q_v, Q_t, Q\}$, the last such atom must have been eliminated from ψ'_{s-1} by taking a resolvent with one of the incomplete Horn clauses in $\overline{\Phi}_2$. By the construction of $\overline{\Phi}_2$, to obtain ψ' through an SLD-derivation $\psi'_0, \ldots, \psi'_s = \psi'$ from $\overline{\Phi}_2$, all Horn clauses introduced in the definition of $\overline{\Phi}_2$ must have been used in the intended order. Recall that we have replaced each variable c_i in Φ_2 with either 0 or 1 while rewriting Φ_2 as an Horn sentence. Every Horn clause from $\overline{\Phi}_2$ has the property that every positive atomic subformula has a symbol from $\{Q_b, Q_v, Q_t, Q\}$ and contains all variables that appear in a negative atomic subformula, with the following two exceptions. First, verifier nodes are not carried over in any atoms because their only contribution is the encoding of a unique number. Second, after a pair of successive rows has been checked by deriving a Q_v -atom containing a 2^{n} -th verifier node, the variable representing the lower row is not carried over in any atom because it is no longer needed. Since ϕ, ϕ_1 , and ϕ_2 satisfy the prerequisites of item (3) in Lemma 7, no ill-behaved variable identifications might have occurred during the SLD-derivation above as otherwise, we would have $\overline{\Phi}_1 \models \forall \overline{x}, y_1(\phi \land \phi_1 \Rightarrow \chi)$, a

contradiction to (2). Consequently, the SLD-derivation must have the full intended length

contradiction to (2). Consequently, the SLD-derivation must have the full intended length $(2^n + 1) \cdot m$ for some $m \geq 1$, because every intermediate stage starts and ends with verifier nodes encoding the numbers 2^n and 1, respectively, and one can only progress in steps which decrement the encoded number by one. Clearly, the SLD-derivation witnesses the existence of $f: \mathfrak{R}_{m,2^n} \to \mathfrak{T}$.

We continue with the proof of Thoeorem 6. This time, we reduce from the complement of the basic rectangle tiling problem. As in the proof of Theorem 5, we first include in τ the two auxiliary unary symbols L, R and the binary auxiliary symbol E. However, the encoding of the tiles is slightly different. We include in τ a symbol I of arity $\lceil \log_2 |T| \cdot n \rceil + 1$. The first $\lceil \log_2 |T| \cdot n \rceil$ entries serve as binary counters to represent the pairs $(i, \alpha) \in [n] \times T$ in binary, and the last entry carries a path node representing a row of the tiling grid. Each pair $(i, \alpha) \in [n] \times T$ is to be interpreted as the fact that the *i*-th column in the row represented by a particular path node contains the tile α . The reason for this choice of encoding is that we aim to construct a universal sentence Φ which is equivalent to a set of forbidden substructures \mathcal{N} of size polynomial in $|T| \cdot n$. To achieve this, we use a constant number of symbols whose arity is logarithmic in the size of the input.

Suppose that 0 and 1 are two variables representing the bits 0 and 1, respectively. For each pair $(i, \alpha) \in [n] \times T$, the ternary formula TILE_{*i*, α}(0, 1, *p*) is the *I*-atom whose last entry contains the variable *p* and the first $\lceil \log_2 |T| \cdot n \rceil$ entries contain the variables 0 and 1 in the unique way that represents the number $i \cdot \alpha$ in binary when read from left to right. Note that the number of such formulas is polynomial in the size of the input to the tiling problem. In contrast to the proof of Theorem 5, it is not necessary to introduce any verifier nodes as the number of columns in the tiling grid is polynomial in the size of the input. The sentence Φ is defined similarly as in the proof of Theorem 5, so we only provide a general overview and highlight the main differences. We want each row to be horizontally consistent. For all $i \in [n-1]$ and $(\alpha, \beta) \in T^2 \setminus P_h^{\mathfrak{T}}$, we include the following sentence as a conjunct in Φ_1 :

$$\forall p, 0, 1(\psi(0, 1) \land \text{TILE}_{i,\alpha}(0, 1, p) \land \text{TILE}_{i+1,\beta}(0, 1, p) \Rightarrow \bot)$$

where $\psi(0, 1)$ represents a simple mechanism for distinguishing between 0 and 1. We also want each position in a given row to be occupied by at most one tile. For all $i \in [n], \alpha, \beta \in T$, we include the following sentence as a conjunct in Φ_1 :

$$\forall p, 0, 1 (\text{TILE}_{i,\alpha}(0, 1, p) \land \text{TILE}_{i,\beta}(0, 1, p) \Rightarrow \bot).$$

As in the proof of Theorem 6, we do *not* need to include an explicit condition stating that each row must be completely tiled from left to right. For the purpose of verifying the validity of a tiling, we include in τ a 5-ary symbol Q, a ($\lceil \log_2 |T| \cdot n \rceil + 4$)-ary symbol Q_v , and two ($\lceil \log_2 |T| \cdot n \rceil + 3$)-ary symbols Q_b, Q_t . For each pair $(i, \alpha) \in [n] \times T$, the formulas

BOT<sub>*i*,
$$\alpha$$</sub>(0, 1, *p*, *y*₁, *y*₂), TOP_{*i*, α} (0, 1, *p*, *y*₁, *y*₂), and VERT_{*i*, α} (0, 1, *p*₁, *p*₂, *y*₁, *y*₂)

are defined analogously to $\text{TILE}_{i,\alpha}(0, 1, p)$ but using the Q_b, Q_t and Q_v -atoms instead. We use them to verify the bottom row, top row, and the vertical consistency of a given tiling. In contrast to $\text{TILE}_{i,\alpha}$, the parameter α is not important in $\text{BOT}_{i,\alpha}$, $\text{TOP}_{i,\alpha}$, and $\text{VERT}_{i,\alpha}$.

We now explain how to convert Φ into a set of forbidden substructures. Let \mathcal{N} be the class of all τ -structures with the domain [i] for some $i \in [6]$ that do not satisfy Φ . Since Φ only uses six variables, we have $\operatorname{fm}(\Phi) = \operatorname{Forb}_e(\mathcal{N})$. It remains to show that there exists a polynomial that bounds the size of \mathcal{N} . Since there is a constant number of domains of structures in \mathcal{N} and their sizes are also constant, it is enough to show that there exists a

polynomial that bounds the number of structures in \mathcal{N} . The only non-constant parameters in the construction are the four symbols I, Q_b, Q_t , and Q_v whose arity grows logarithmically with $|T| \cdot n$. Thus, there exists a constant c such that the number of structures in \mathcal{N} is bounded by $c \cdot (2^{\lceil \log_2 |T| \cdot n \rceil})^4 \leq c \cdot (2 \cdot |T| \cdot n)^4$. The rest is analogous to the proof of Theorem 5.

4 The second meta-problem

Every reduct \mathfrak{R} of a finitely bounded homogeneous structure \mathfrak{H} is uniquely described by an SNP sentence, which can be obtained from a universal sentence for $\operatorname{age}(\mathfrak{H})$ by existentially quantifying all the surplus predicates upfront. This is (arguably) the most natural representation for such structures. The second meta-problem asks whether a given SNP τ -sentence Φ is logically equivalent to an SNP τ -sentence $\Psi = \exists Y_1, \ldots, Y_m \forall \overline{y}. \psi$ such that $\operatorname{fm}(\forall \overline{y}. \psi)$ has the AP in the extended signature $\tau \cup \{Y_1, \ldots, Y_m\}$. We are additionally interested in the refinement of the question where we require the homogeneous structure from Theorem 1 associated to $\operatorname{fm}(\forall \overline{y}. \psi)$ to have the same set of automorphisms as its reduct to the original signature τ . This amounts to asking whether $\operatorname{fm}(\Phi)$ is finitely bounded homogenizable.

MMSNP was presented in [31] as a large subclass of SNP which has a dichotomy between P and NP-completeness if and only if the class of all finite-domain CSPs has one. The latter has been confirmed, and the dichotomy for MMSNP has received a new universal-algebraic proof within the programme attacking the Bodirsky-Pinsker conjecture [18]. The new proof relies on the observation that every MMSNP sentence Φ is equivalent to a finite disjunction $\Phi_1 \vee \cdots \vee \Phi_n$ of MMSNP sentences such that, for every $i \in [n]$, there exists a reduct \mathfrak{R}_i of a finitely bounded homogeneous structure \mathfrak{H}_i such that $\mathrm{fm}(\Phi_i) = \mathrm{CSP}(\mathfrak{R}_i)^3$. Moreover, the structure \mathfrak{H} can be chosen so that its age has the *Ramsey property*, which plays an essential role in an argument in [18] showing that the authors correctly identified all of the tractable cases. The exact definition of this property is not essential to the present article and is therefore omitted. GMSNP was first introduced in [40] in its seemingly weaker form MMSNP₂, as a generalization of MMSNP where "monadic" second-order variables may also range over atomic formulas. It was later shown that relaxing the above requirement for monotone SNP to guardedness does not result in a more expressive logic [8]. There is a prospect that GMSNP will also have dichotomy between P and NP-completeness since it enjoys similar model-theoretic properties as MMSNP [16].

Theorem 11 is the most general version of our undecidability result. The second item might give the impression that one cannot effectively distinguish between CSPs of reducts of finitely bounded structures and CSPs of reducts of finitely bounded homogeneous structures. Recall that the former class does not have a dichotomy [31, 9]. However, as indicated by the formulation of the second item, all CSPs of reducts of finitely bounded homogeneous structures in the proof of Theorem 11 are in fact finite-domain CSPs, for which there is a dichotomy. Therefore, Theorem 11 merely shows that SNP sentences are an exceptionally bad choice of an input to the question, albeit one that is sometimes used (see, e.g., [5, 42]).

Theorem 11. Given a Datalog sentence Φ over a finite binary signature, it is undecidable whether:

or Φ is not even logically equivalent to any GMSNP sentence;

^{1.} Φ is logically equivalent to a monadic Datalog sentence,

³ This observation was first made in [12], see the proof of Theorem 7.

- **2.** Φ simultaneously satisfies the following three conditions:
 - = fm(Φ) is the CSP of a finite structure,
 - = fm(Φ) is a finitely bounded homogenizable class,
 - $= fm(\Phi)$ is the age of a reduct of a finitely bounded homogeneous Ramsey structure,
 - or $fm(\Phi)$ is not even the CSP or the age of any ω -categorical structure.

4.1 A proof of Theorem 11

As usual, the *Kleene plus* and the *Kleene star* of a finite set of symbols Σ , denoted by Σ^+ and Σ^* , are the sets of all finite words over Σ of lengths ≥ 1 and ≥ 0 , respectively.

A context-free grammar (CFG) is a 4-tuple $\mathfrak{G} = (N, \Sigma, P, S)$ where N is a finite set of non-terminal symbols, Σ is a finite set of terminal symbols, P is a finite set of production rules of the form $A \to w$ where $A \in N$ and $w \in (N \cup \Sigma)^+$, $S \in N$ is the start symbol. For $u, v \in (N \cup \Sigma)^+$ we write $u \to_G v$ if there are $x, y \in (N \cup \Sigma)^+$ and $(A \to w) \in P$ such that u = xAy and v = xwy. The language of \mathfrak{G} is $L(\mathfrak{G}) := \{w \in \Sigma^+ \mid S \to_{\mathfrak{G}}^* w\}$, where $\to_{\mathfrak{G}}^*$ denotes the transitive closure of $\to_{\mathfrak{G}}$. Note that with this definition the empty word ϵ can never be an element of $L(\mathfrak{G})$; some authors use a modified definition that also allows rules that derive ϵ , but for our purposes the difference is not essential. A context-free grammar is called (left-)regular if its production rules are always of the form $A \to a$ or $A \to Ba$ for non-terminal symbols A, B and a terminal symbol a. For a finite set Σ , we call a set $L \subseteq \Sigma^+$ regular if it is the language of a regular grammar with terminal symbols Σ .

▶ **Example 12.** Consider the CFG \mathscr{G} with a single terminal symbol a, non-terminal symbols S, A, B, C, and production rules $S \to a$, $S \to aa$, $S \to aaa$, $S \to Aa$, $A \to Ba$, $B \to Ca$, $C \to Ca$, and $C \to a$. Clearly, \mathscr{G} is not regular. However, $L(\mathscr{G}) = \{a\}^+$ is regular.

Let $\mathfrak{G} = (N, \Sigma, P, S)$ be a CFG. The signature τ_{Σ} consists of the unary symbols I, Tand the binary symbols R_a for every $a \in \Sigma$, and the signature τ_N consists of a binary symbol R_a for every element $a \in N$. For $a_1 \dots a_n \in \Sigma^+$, we set $\phi_{a_1 \dots a_n}(x_1, \dots, x_{n+1}) \coloneqq$ $\bigwedge_{i \in [n]} R_{a_i}(x_i, x_{i+1})$. Let $\overline{\Phi}_{\mathfrak{G}}$ be the universal Horn sentence over the signature $\tau_{\Sigma} \cup \tau_N$ whose quantifier-free part contains, for every $(A, w) \in P$, the Horn clause $\phi_w(x_1, \dots, x_{|w|+1}) \Rightarrow$ $R_A(x_1, x_{|w|+1})$, and additionally the Horn clause $I(x_1) \wedge R_S(x_1, x_2) \wedge T(x_2) \Rightarrow \bot$. Then $\Phi_{\mathfrak{G}}$ is the Datalog sentence obtained from $\overline{\Phi}_{\mathfrak{G}}$ by existentially quantifying all symbols from τ_N upfront. This encoding of CFGs into Datalog programs is standard (Exercise 12.26 in [1]), and the correspondence provided by the next lemma can be shown via a straightforward induction. For a proof, we refer the reader to [46].

▶ Lemma 13. For a τ_{Σ} -structure \mathfrak{A} , we have $\mathfrak{A} \models \Phi_{\mathfrak{F}}$ if and only if, for every $w \in L(\mathfrak{F})$,

$$\mathfrak{A} \models \forall x_1, \dots, x_{|w|+1} (I(x_1) \land \phi_w(x_1, \dots, x_{|w|+1}) \land T(x_{|w|+1}) \Rightarrow \bot).$$

The following lemma is proved by establishing a connection between the well-known Myhill-Nerode correspondence and ω -categoricity, under the addition of several auxiliary results from [30, 16].

▶ Lemma 14. Let 𝔅 be a context-free grammar. Then the following are equivalent:

- **1.** $L(\mathcal{G})$ is regular.
- **2.** $\Phi_{\mathfrak{F}}$ is equivalent to monadic Datalog sentence.
- **3.** $\Phi_{\mathfrak{G}}$ is equivalent to a GMSNP sentence.
- **4.** fm($\Phi_{\mathfrak{F}}$) is the CSP of a finite structure.

- **5.** fm($\Phi_{\mathfrak{F}}$) is the CSP of an ω -categorical structure.
- **6.** $\operatorname{fm}(\Phi_{\mathfrak{F}})$ is the age of a reduct of a finitely bounded homogeneous Ramsey structure.
- 7. fm($\Phi_{\mathfrak{F}}$) is the age of an ω -categorical structure.
- **8.** fm($\Phi_{\mathfrak{F}}$) is finitely bounded homogenizable.

Proof of Theorem 11. It is well-known that the questions whether $L(\mathcal{G})$ is regular for a given context-free grammar \mathcal{G} is undecidable, see, e.g., Theorem 6.6.6 in [48]. Hence, all eight equivalent conditions in Lemma 14 are undecidable for \mathcal{G} .

— References -

- 1 Serge Abiteboul, Richard Hull, and Victor Vianu. *Foundations of databases*, volume 8. Addison-Wesley Reading, 1995.
- 2 Ove Ahlman. Homogenizable structures and model completeness. Archive for Mathematical Logic, 55(7-8):977–995, 2016.
- 3 Alfred V. Aho, John E. Hopcroft, and Jeffrey D. Ullman. The Design and Analysis of Computer Algorithms. Addison-Wesley, 1974.
- 4 Reza Akhtar and Alistair H. Lachlan. On countable homogeneous 3-hypergraphs. Arch. Math. Log., 34(5):331–344, 1995.
- 5 Albert Atserias and Szymon Toruńczyk. Non-homogenizable classes of finite structures. Computer Science Logic 2016, page 1, 2016.
- 6 Franz Baader and Jakub Rydval. Using model theory to find decidable and tractable description logics with concrete domains. *Journal of Automated Reasoning*, 66(3):357–407, 2022.
- 7 Libor Barto, Michael Kompatscher, Miroslav Olšák, Trung Van Pham, and Michael Pinsker. The equivalence of two dichotomy conjectures for infinite domain constraint satisfaction problems. In Proceedings of the 32nd Annual ACM/IEEE Symposium on Logic in Computer Science – LICS'17, 2017. available at arXiv:1612.07551.
- 8 Meghyn Bienvenu, Balder ten Cate, Carsten Lutz, and Frank Wolter. Ontology-based data access: A study through disjunctive Datalog, CSP, and MMSNP. ACM Trans. Database Syst., 39(4):33, 2014.
- 9 Manuel Bodirsky. *Complexity of Infinite-Domain Constraint Satisfaction*. Cambridge University Press, 2021.
- 10 Manuel Bodirsky and Bertalan Bodor. Canonical polymorphisms of ramsey structures and the unique interpolation property. In 2021 36th Annual ACM/IEEE Symposium on Logic in Computer Science (LICS), pages 1–13. IEEE, 2021.
- 11 Manuel Bodirsky, David Bradley-Williams, Michael Pinsker, and András Pongrácz. The universal homogeneous binary tree. J. Log. Comput., 28(1):133–163, 2018.
- 12 Manuel Bodirsky and Víctor Dalmau. Datalog and constraint satisfaction with infinite templates. *Journal on Computer and System Sciences*, 79:79–100, 2013. A preliminary version appeared in the proceedings of the Symposium on Theoretical Aspects of Computer Science (STACS'05).
- 13 Manuel Bodirsky and Martin Grohe. Non-dichotomies in constraint satisfaction complexity. In Luca Aceto, Ivan Damgard, Leslie Ann Goldberg, Magnús M. Halldórsson, Anna Ingólfsdóttir, and Igor Walukiewicz, editors, *Proceedings of the International Colloquium on Automata, Languages and Programming (ICALP)*, Lecture Notes in Computer Science, pages 184–196. Springer Verlag, July 2008.
- 14 Manuel Bodirsky, Peter Jonsson, and Trung Van Pham. The complexity of phylogeny constraint satisfaction problems. ACM Transactions on Computational Logic (TOCL), 18(3):1– 42, 2017.
- 15 Manuel Bodirsky and Jan Kára. The complexity of temporal constraint satisfaction problems. Journal of the ACM (JACM), 57(2):1–41, 2010.

- 16 Manuel Bodirsky, Simon Knäuer, and Florian Starke. Asnp: A tame fragment of existential second-order logic. In Beyond the Horizon of Computability: 16th Conference on Computability in Europe, CiE 2020, Fisciano, Italy, June 29–July 3, 2020, Proceedings 16, pages 149–162. Springer, 2020.
- 17 Manuel Bodirsky, Florent Madelaine, and Antoine Mottet. A universal-algebraic proof of the complexity dichotomy for Monotone Monadic SNP. In *Proceedings of the Symposium on Logic* in Computer Science – LICS'18, pages 105–114, 2018. available at ArXiv:1802.03255.
- 18 Manuel Bodirsky, Florent Madelaine, and Antoine Mottet. A proof of the algebraic tractability conjecture for monotone monadic snp. *SIAM Journal on Computing*, 50(4):1359–1409, 2021.
- 19 Manuel Bodirsky and Antoine Mottet. Reducts of finitely bounded homogeneous structures, and lifting tractability from finite-domain constraint satisfaction. In *Proceedings of the 31th Annual IEEE Symposium on Logic in Computer Science LICS'16*, pages 623–632, 2016. available at ArXiv:1601.04520.
- 20 Manuel Bodirsky, Michael Pinsker, and András Pongrácz. Projective clone homomorphisms. *The Journal of Symbolic Logic*, 86(1):148–161, 2021.
- 21 Manuel Bodirsky and Jakub Rydval. On the descriptive complexity of temporal constraint satisfaction problems. *Journal of the ACM*, 70(1):1–58, 2022.
- 22 Mikołaj Bojańczyk, Luc Segoufin, and Szymon Toruńczyk. Verification of database-driven systems via amalgamation. In Proceedings of the 32nd ACM SIGMOD-SIGACT-SIGAI symposium on Principles of database systems, pages 63–74, 2013.
- 23 S. Braunfeld. The undecidability of joint embedding and joint homomorphism for hereditary graph classes. Discrete Mathematics & Theoretical Computer Science, 21, 2019. available at arXiv:1903.11932.
- 24 Andrei A Bulatov. A dichotomy theorem for nonuniform csps. In 2017 IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS), pages 319–330. IEEE, 2017.
- 25 Gregory Cherlin. Forbidden substructures and combinatorial dichotomies: Wqo and universality. Discrete Mathematics, 311(15):1543–1584, 2011.
- **26** Gregory Cherlin. *Homogeneous ordered graphs, metrically homogeneous graphs, and beyond,* volume 1. Cambridge University Press, 2022.
- 27 Gregory Cherlin, Saharon Shelah, and Niandong Shi. Universal graphs with forbidden subgraphs and algebraic closure. Advances in Applied Mathematics, 22(4):454–491, 1999.
- 28 Bogdan S. Chlebus. Domino-tiling games. Journal of Computer and System Sciences, 32(3):374–392, 1986.
- 29 Jacinta Covington. Homogenizable relational structures. Illinois Journal of Mathematics, 34(4):731-743, 1990.
- 30 Péter L. Erdös, Claude Tardif, and Gábor Tardos. Caterpillar dualities and regular languages. SIAM Journal on Discrete Mathematics, 27(3):1287–1294, 2013.
- 31 Tomás Feder and Moshe Y Vardi. The computational structure of monotone monadic snp and constraint satisfaction: A study through datalog and group theory. SIAM Journal on Computing, 28(1):57–104, 1998.
- 32 Wilfrid Hodges. A shorter model theory. Cambridge University Press, Cambridge, 1997.
- 33 Jan Hubička and Jaroslav Nešetřil. All those ramsey classes (ramsey classes with closures and forbidden homomorphisms). Advances in Mathematics, 356:106791, 2019.
- 34 Jan Hubička and Jaroslav Nešetřil. Homomorphism and embedding universal structures for restricted classes. *Multiple-Valued Logic and Soft Computing*, 27(2-3):229–253, 2016. available at arXiv:0909.4939.
- 35 Julia F. Knight and Alistair H. Lachlan. Shrinking, stretching, and codes for homogeneous structures. In Classification Theory: Proceedings of the US-Israel Workshop on Model Theory in Mathematical Logic held in Chicago, Dec. 15–19, 1985, pages 192–229. Springer, 2006.
- **36** Gábor Kun. Constraints, MMSNP, and expander relational structures. *Combinatorica*, 33(3):335–347, 2013.

- 37 A.H. Lachlan. Homogeneous structures. In Proceedings of the International Congress of Mathematicians, volume 1, pages 314–321, Berkeley, 1986. AMS.
- 38 Carsten Lutz and Maja Miličić. A tableau algorithm for description logics with concrete domains and general tboxes. Journal of Automated Reasoning, 38(1-3):227–259, 2007.
- 39 Dugald Macpherson. A survey of homogeneous structures. Discrete Mathematics, 311(15):1599–1634, 2011.
- 40 Florent Madelaine. Universal structures and the logic of forbidden patterns. Logical Methods in Computer Science, 5(2), 2009.
- 41 Florent R Madelaine and Iain A Stewart. Some problems not definable using structure homomorphisms. Ars Combinatoria, 67:153–160, 2003.
- 42 Antoine Mottet. Promise and infinite-domain constraint satisfaction. In 32nd EACSL Annual Conference on Computer Science Logic (CSL 2024). Schloss-Dagstuhl-Leibniz Zentrum für Informatik, 2024.
- 43 Antoine Mottet and Michael Pinsker. Smooth approximations and csps over finitely bounded homogeneous structures. In *Proceedings of the 37th Annual ACM/IEEE Symposium on Logic in Computer Science*, pages 1–13, 2022. available at arXiv:2011.03978.
- 44 Antoine Mottet, Michael Pinsker, and Tomáš Nagy. An order out of nowhere: a new algorithm for infinite-domain CSPs. *arXiv preprint*, 2023. arXiv:2301.12977.
- 45 Shan-Hwei Nienhuys-Cheng and Roland Wolf. Foundations of Inductive Logic Programming, volume 1228 of Lecture Notes in Computer Science. Springer, 1997.
- 46 André Schrottenloher, Jakub Rydval, and Manuel Bodirsky. Universal horn sentences and the joint embedding property. *Discrete Mathematics & Theoretical Computer Science*, 23, 2022.
- 47 François Schwarzentruber. The Complexity of Tiling Problems. arXiv preprint, 2019. arXiv:1907.00102.
- 48 Jeffrey Shallit. A second course in formal languages and automata theory. Cambridge University Press, 2008.
- 49 Peter van Emde Boas. The convenience of tilings. In Complexity, Logic, and Recursion Theory, pages 331–363. CRC Press, 2019.
- 50 Dmitriy Zhuk. A proof of CSP dichotomy conjecture. In 58th IEEE Annual Symposium on Foundations of Computer Science, FOCS 2017, Berkeley, CA, USA, October 15-17, 2017, pages 331–342, 2017.
- **51** Dmitriy Zhuk. A proof of the csp dichotomy conjecture. *Journal of the ACM (JACM)*, 67(5):1–78, 2020.

A Omitted proofs

For a tuple \bar{t} indexed by a set I, the value of \bar{t} at the position $i \in I$ is denoted by $\bar{t}[i]$.

The disjoint union of two τ -structure \mathfrak{A} and \mathfrak{B} is obtained by taking the union of two copies of \mathfrak{A} and \mathfrak{B} with disjoint domains. We call Φ connected if the Gaifman graph of each clause of ϕ is complete connected. It is a folklore fact that, if Φ is connected, then fm(Φ) is preserved by disjoint unions. Note that connectivity easily transfers to SNP sentences viewed as universal sentences in an extended signature.

A.1 **Proof of Proposition 3**

▶ **Proposition 3.** Let Φ be a universal sentence over a finite binary relational signature τ . If fm(Φ) does not have the AP, then the size of a smallest counterexample to the AP is at most single-exponential in the size of Φ . Consequently, the question whether fm(Φ) has the AP is decidable in coNEXPTIME.

Proof. Suppose that $fm(\Phi)$ does not have the AP. By Proposition 2, we may assume that a counterexample to the AP for $fm(\Phi)$ is a one-point amalgamation diagram formed by two structures $\mathfrak{B}_1, \mathfrak{B}_2 \in fm(\Phi)$. Let k be the number of variables in Φ . We define \mathcal{N} as the set of all τ -structures \mathfrak{A} such that $\mathfrak{A} \not\models \Phi$, $|A| \leq k$, and A = [|A|]. Clearly, $fm(\Phi) = \operatorname{Forb}_e(\mathcal{N})$ and the size of \mathcal{N} is at most single-exponential in the size of Φ . Therefore, by the estimates on the size of a smallest counterexample from the proof of Theorem 4 in [16], we can assume that the size of \mathfrak{B}_1 and \mathfrak{B}_2 is at most single-exponential in the size of Φ . First, we must verify that $\mathfrak{B}_1, \mathfrak{B}_2 \in fm(\Phi)$. This can be done in time exponential in the size of Φ simply by evaluating the quantifier-free part of Φ on all possible inputs. Subsequently, we must verify that no amalgam $\mathfrak{C} \in fm(\Phi)$ of \mathfrak{B}_1 and \mathfrak{B}_2 can be obtained either by identifying b_1 and b_2 , or by adding (b_1, b_2) or (b_2, b_1) to some relations of $\mathfrak{B}_1 \cup \mathfrak{B}_2$. This can also be done in time exponential in the size of Φ because only single-exponentially many structures \mathfrak{C} need to be checked. In sum, the existence of a counterexample can be tested in NEXPTIME, which is what we had to show.

A.2 Proof of Lemma 7

Lemma 7. Let Φ be an equality-free universal Horn sentence over a relational signature τ . Then the following are equivalent:

- **1.** fm(Φ) has the strong AP.
- **2.** fm(Φ) has the AP.
- **3.** If $\phi(\bar{x})$, $\phi_1(\bar{x}, y_1)$, and $\phi_2(\bar{x}, y_2)$ are equality-free conjunctions of atomic formulas, where y_1 and y_2 are distinct variables not contained in \bar{x} , such that, for both $i \in \{1, 2\}$, every atom in $\phi_i(\bar{x}, y_i)$ contains the variable y_i and $\phi(\bar{x}) \land \phi_i(\bar{x}, y_i) \leq_{\Phi} \phi(\bar{x})$, then $\phi(\bar{x}) \land \phi_1(\bar{x}, y_1) \land \phi_2(\bar{x}, y_2) \leq_{\Phi} \phi(\bar{x}) \land \phi_1(\bar{x}, y_1)$.

Proof. "(1) \Rightarrow (2)" This direction is trivial.

"(2) \Rightarrow (3)" Let $\phi(\bar{x}), \phi_1(\bar{x}, y_1)$, and $\phi_2(\bar{x}, y_2)$ be as in item (3). Let $\Psi(\bar{x}), \Psi_1(\bar{x}, y_1)$, and $\Psi_2(\bar{x}, y_2)$ be the conjunctions of all *R*-atoms for $R \in \tau$ implied by $\Phi \wedge \phi(\bar{x}), \Phi \wedge \phi(\bar{x}) \wedge \phi_1(\bar{x}, y_1)$, and $\Phi \wedge \phi(\bar{x}) \wedge \phi_2(\bar{x}, y_2)$, respectively. If $\Phi \wedge \Psi_1(\bar{x}, y_1)$ or $\Phi \wedge \Psi_2(\bar{x}, y_2)$ is unsatisfiable, then $\Phi \wedge \Psi(\bar{x})$ is unsatisfiable by the assumption and we are done. So suppose that both are formulas are satisfiable. Define \mathfrak{B}_1 and \mathfrak{B}_2 as the structures whose domains consist of the variables $\{y_1, \bar{x}[1], \ldots\}$ and $\{y_2, \bar{x}[1], \ldots\}$, respectively, and where \bar{z} is a tuple of a relation for $R \in \tau$ if the conjunct $R(\bar{z})$ is contained in Ψ_1 or Ψ_2 , respectively. Since we have $\phi(\bar{x}) \wedge \phi_i(\bar{x}, y_i) \leq \Phi \phi(\bar{x})$ for both $i \in \{1, 2\}$, the induced substructures of \mathfrak{B}_1 and \mathfrak{B}_2 on $B_1 \cap B_2$ coincide. Note that, by construction, \mathfrak{B}_1 and \mathfrak{B}_2 satisfy every Horn clause in Φ . Since Φ is universal Horn, this implies that $\mathfrak{B}_1, \mathfrak{B}_2 \in \mathrm{fm}(\Phi)$. Since $\mathrm{fm}(\Phi)$ has the AP, there exists $\mathfrak{C} \in \mathrm{fm}(\Phi)$ together with embeddings $f_i: \mathfrak{B}_i \hookrightarrow \mathfrak{C}$ for both $i \in \{1,2\}$ such that $f_1|_{B_1\cap B_2} = f_2|_{B_1\cap B_2}$. By the construction of \mathfrak{B}_1 and \mathfrak{B}_2 , it follows that $\Phi \not\models$ $\forall \bar{x}, y_1, y_2(\phi(\bar{x}) \land \phi_1(\bar{x}, y_1) \land \phi_2(\bar{x}, y_2) \Rightarrow \bot)$. Let $\chi(\bar{x}, y_1)$ be an atomic τ -formula other than equality such that $\Phi \models \forall \bar{x}, y_1, y_2(\phi(\bar{x}) \land \phi_1(\bar{x}, y_1) \land \phi_2(\bar{x}, y_2) \Rightarrow \chi(\bar{x}, y_1))$. By the construction of \mathfrak{B}_1 and \mathfrak{B}_2 , and because f_1 and f_2 are homomorphisms, there exist a tuple \bar{z} over B_1 such that $\mathfrak{C} \models \chi(f_1(\bar{z}))$. Since f_1 is an embedding, we must also have $\mathfrak{B}_1 \models \chi(\bar{z})$. Thus, by the construction of \mathfrak{B}_1 and \mathfrak{B}_2 , it follows that $\Phi \models \forall \bar{x}, y_1 (\phi(\bar{x}) \land \phi_1(\bar{x}, y_1) \Rightarrow \chi(\bar{x}, y_1))$.

"(3) \Rightarrow (1)" Let $\mathfrak{B}_1, \mathfrak{B}_2 \in \mathrm{fm}(\Phi)$ be an arbitrary one-point amalgamation diagram. Let y_1 and y_2 be the unique elements contained in $B_1 \setminus B_2$ and $B_2 \setminus B_1$, respectively. We construct a structure $\mathfrak{C} \in \mathrm{fm}(\Phi)$ with embeddings $f_i: \mathfrak{B}_i \hookrightarrow \mathfrak{C}$ for both $i \in \{1, 2\}$ as follows. Let \bar{x} be a tuple of variables representing the elements of $B_1 \cap B_2$ in some order, and let $\phi(\bar{x})$ be the conjunction of all R-atoms for $R \in \tau$ which hold in $\mathfrak{B}_1 \cap \mathfrak{B}_2$. Moreover, for

both $i \in \{1,2\}$, let $\phi_i(\bar{x},y_i)$ be the conjunction of all R-atoms for $R \in \tau$ which contain the variable y_i and hold in \mathfrak{B}_i . By construction, for both $i \in \{1,2\}$, we have $\phi(\bar{x}) \wedge \phi(\bar{x})$ $\phi_i(\bar{x}, y_i) \leq_{\Phi} \phi(\bar{x})$ because $\mathfrak{B}_i \models \Phi$. Let $\Psi(\bar{x}, y_1, y_2)$ be the conjunction of all *R*-atoms for $R \in \tau$ implied by $\Phi \wedge \phi(\bar{x}) \wedge \phi_1(\bar{x}, y_1) \wedge \phi_2(\bar{x}, y_2)$. We claim that $\Phi \wedge \Psi$ is satisfiable: otherwise, $\Phi \models \forall \bar{x}, y_1, y_2(\phi(\bar{x}) \land \phi_1(\bar{x}, y_1) \land \phi_2(\bar{x}, y_2) \Rightarrow \bot)$, and then item (3) implies that $\Phi \models \forall \bar{x}, y_1(\phi(\bar{x}) \land \phi_1(\bar{x}, y_1) \Rightarrow \bot)$, which is impossible since $\mathfrak{B}_1 \models \Phi$. Define \mathfrak{C} as the structure with domain $\{y_1, y_2, \bar{x}[1], \dots\}$ and such that $R^{\mathfrak{C}}$ contains a tuple \bar{z} if and only if Ψ contains the conjunct $R(\bar{z})$. For both $i \in \{1, 2\}$, let f_i be the identity map on B_i . By definition, we have $f_1|_{B_1 \cap B_2} = f_2|_{B_1 \cap B_2}$ and $f_1(B_1) \cap f_2(B_2) = f_1(B_1 \cap B_2) = f_2(B_1 \cap B_2)$. We claim that, for both $i \in \{1, 2\}$, f_i is an embedding from \mathfrak{B}_i to \mathfrak{C} . It is clear from the construction of \mathfrak{C} that f_i is a homomorphism. Suppose for contradiction that there exists $R \in \tau$ and a tuple \bar{z} over B_i such that $\bar{z} \notin R^{\mathfrak{B}_i}$ while $f_i(\bar{z}) \in R^{\mathfrak{C}}$. For the sake of notation, we assume that i = 1; the case that i = 2 can be shown analogously. Note that the construction of \mathfrak{C} implies that $\Phi \models \forall \bar{x}, y_1, y_2(\phi(\bar{x}) \land \phi_1(\bar{x}, y_1) \land \phi_2(\bar{x}, y_2) \Rightarrow R(\bar{z}))$. Then item (3) implies that $\Phi \models \forall \bar{x}, y_1(\phi(\bar{x}) \land \phi_1(\bar{x}, y_1) \Rightarrow R(\bar{z}))$, a contradiction to $\mathfrak{B}_1 \in \mathrm{fm}(\Phi)$. Thus, f_i is an embedding from \mathfrak{B}_i to \mathfrak{C} . This concludes the proof of the strong AP restricted to one-point amalgamation diagrams. Now the statement follows similarly as in Proposition 2.

A.3 Proof of Theorem 4

▶ **Theorem 4.** Given a universal Horn sentence Φ over a finite relational signature that is binary except for one ternary symbol, the question whether fm(Φ) has the AP is PSPACE-hard.

Proof. We reduce from the universality problem for left-regular grammars, which is known to be PSPACE-complete [3]. Let $\mathcal{G} = (N, \Sigma, P, S)$ be a left-regular grammar.

First, consider the Horn sentence Φ_1 over the signature $\tau_1 = \{L, I, T, E\} \cup \{R_a \mid a \in \Sigma \cup N\}$ in prenex normal form whose quantifier-free part consists of the Horn clause

$$I(x_1) \wedge R_S(x_1, x_2) \wedge T(x_2) \Rightarrow \bot, \tag{4}$$

as well as

$$L(y_1) \wedge I(y_1) \wedge E(y_1, x_2) \wedge R_B(y_1, x_1) \wedge R_a(x_1, x_2) \Rightarrow R_A(y_1, x_2), \qquad (A \to Ba \in P)$$
$$L(y_1) \wedge I(y_1) \wedge R_a(y_1, x_1) \Rightarrow R_A(y_1, x_1). \qquad (A \to a \in P)$$

The proof of the following claim is straightforward and left to the reader.

 \triangleright Claim 15. For every $w = a_1 \dots a_n \in \Sigma^+$ and $A \in N$, we have $A \to_{\mathscr{G}}^* w$ if and only if

$$\Phi_1 \models \forall x_1, \dots, x_n, y_1 \left(\begin{array}{c} L(y_1) \wedge I(y_1) \wedge R_{a_1}(y_1, x_1) \\ \wedge \bigwedge_{i \in [n-1]} R_{a_i}(x_i, x_{i+1}) \wedge E(y_1, x_{i+1}) \end{array} \right) \Rightarrow R_A(y_1, x_n) \right)$$
(5)

Secondly, consider the Horn sentence Φ_2 over the signature $\tau_2 = \{L, R, I, T, E, Q\} \cup \{R_a \mid a \in \Sigma\}$ defined as the universally quantified conjunction of all Horn clauses of the form

$$L(y_1) \wedge R(y_2) \wedge I(y_1) \wedge R_a(y_1, x_1) \wedge E(y_2, x_1) \Rightarrow Q(y_1, y_2, x_1),$$
(6)

$$Q(y_1, y_2, x_1) \wedge R_a(x_1, x_2) \wedge E(y_1, x_2) \wedge E(y_2, x_2) \Rightarrow Q(y_1, y_2, x_2), \qquad (a \in \Sigma)$$
(7)

$$Q(y_1, y_2, x_1) \wedge T(x_1) \Rightarrow \bot.$$
(8)

The proof of the following claim is also straightforward and left to the reader.




Figure 2 An illustration of the AP-counterexamples from Claim 16.

 \triangleright Claim 16. Let $\phi(\bar{x})$ be a conjunction of $(\tau \setminus \{Q\})$ -atoms. Then $\Phi_2 \models \forall \bar{x} (\phi(\bar{x}) \Rightarrow \bot)$ if and only if there exists $a_1 \dots a_n \in \Sigma^+$ such that ϕ contains the premise of

$$\left. \begin{array}{c} R_{a_1}(y_1, x_1) \wedge \bigwedge_{i \in [n-1]} R_{a_{i+1}}(x_i, x_{i+1}) \wedge E(y_1, x_{i+1}) \\ \wedge I(y_1) \wedge T(x_n) \wedge \bigwedge_{i \in [n]} E(y_2, x_i) \wedge L(y_1) \wedge R(y_2) \end{array} \right\} \Rightarrow \bot$$

$$(9)$$

as a subformula modulo renaming of the variables (see Figure 2).

We set $\Phi := \Phi_1 \wedge \Phi_2$, and show that $\operatorname{fm}(\Phi)$ has the AP if and only if $L(\mathscr{G}) = \Sigma^+$. We remark that the roles of L and I are (coincidentally) identical in our construction, but we keep both symbols for the sake of staying consistent with the previous proofs.

" \Rightarrow " Suppose that fm(Φ) has the AP. Let $a_1 \ldots a_n \in \Sigma^+$ be arbitrary. Consider the formulas $\phi(x_1, \ldots, x_n), \phi_1(x_1, \ldots, x_n, y_1)$, and $\phi_2(x_1, \ldots, x_n, y_2)$ given by

$$\begin{split} \phi &\coloneqq T(x_n) \land \bigwedge_{i \in [n-1]} R_{a_{i+1}}(x_i, x_{i+1}), \\ \phi_1 &\coloneqq L(y_1) \land I(y_1) \land R_{a_1}(y_1, x_1) \land \bigwedge_{i \in [n-1]} E(y_1, x_{i+1}), \\ \text{and} \quad \phi_2 &\coloneqq R(y_2) \land \bigwedge_{i \in [n]} E(y_2, x_i). \end{split}$$

By Claim 16, we have

$$\Phi_2 \models \forall x_1, \dots, x_n, y_1, y_2(\phi \land \phi_1 \land \phi_2 \Rightarrow \bot).$$
⁽¹⁰⁾

We claim that $\Phi \wedge \phi \wedge \phi_2$ is satisfiable and $\phi \wedge \phi_2 \leq \Phi \phi$. Let $\chi(x_1, \ldots, x_n)$ be an atomic τ -formula other than equality such that $\Phi \models \forall x_1, \ldots, x_n, y_2(\phi \wedge \phi_2 \Rightarrow \chi)$. Note that all Horn clauses in Φ contain an $\{I, Q\}$ -atom in their premise while $\phi \wedge \phi_2$ contains no $\{I, Q\}$ -atoms. As a consequence, $\phi \wedge \phi_2 \Rightarrow \chi$ must be a tautology, i.e., χ is a subformula of $\phi \wedge \phi_2$. In particular, χ cannot be of the form \bot . Since every atom in ϕ_2 contains the variable y_2 while χ does not, χ is a subformula of ϕ . It follows that $\Phi \models \forall x_1, \ldots, x_n(\phi \Rightarrow \chi)$. Since χ was chosen arbitrarily, this confirms our claim.

Since fm(Φ) has the AP and we already have $\phi \wedge \phi_2 \leq_{\Phi} \phi$, by Lemma 7, it cannot be the case that $\phi \wedge \phi_1 \leq_{\Phi} \phi$. Otherwise, (10) would lead to a contradiction to the satisfiability of $\Phi \wedge \phi \wedge \phi_2$ via Lemma 7. Thus, there must exist an atomic formula $\chi(x_1, \ldots, x_n)$ other than equality such that

$$\Phi \models \forall x_1, \dots, x_n, y_1(\phi \land \phi_1 \Rightarrow \chi) \quad \text{and} \quad \Phi \not\models \forall x_1, \dots, x_n(\phi \Rightarrow \chi). \tag{11}$$

By Theorem 8, we have $\Phi \vdash \forall x_1, \ldots, x_n, y_1 (\phi \land \phi_1 \Rightarrow \chi)$. Clearly, by (11), $\phi \land \phi_1 \Rightarrow \chi$ is not a tautology because every atom in ϕ_1 contains the variable y_1 while χ does not. Thus,

 $\phi \wedge \phi_1 \Rightarrow \chi$ is a weakening of a Horn clause ψ that has an SLD-derivation $\psi_0, \ldots, \psi_s = \psi$ from Φ modulo renaming variables. Since every Horn clause in Φ_2 contains an $\{Q, R\}$ -atom in its premise while Horn clauses in Φ_1 , as well as $\phi \wedge \phi_1$, do not contain any $\{Q, R\}$ -atoms, Φ_2 cannot contribute anything during the SLD-derivation step. Therefore, the SLD-derivation $\psi_0, \ldots, \psi_s = \psi$ is from Φ_1 . Note that, if a variable appears in an *I*-atom in the premise of a Horn clause from Φ_1 whose conclusion is not \perp , then this variable also appears in an atom from the conclusion. Moreover, no Horn clause from Φ_1 contains an *I*-atom in its conclusion. Since y_1 is the only variable which appears in an *I*-atom in $\phi \wedge \phi_1$ and χ does not contain the variable y_1, ψ_0 must be of the form (4). Thus, we have that

$$\Phi_1 \vdash \forall x_1, \dots, x_n, y_1(\phi \land \phi_1 \Rightarrow \bot).$$
(12)

Let ϕ' be the conjunction of atomic formulas obtained from ϕ by removing the atom $T(x_n)$. Note that x_n is the only variable which appears in a *T*-atom in $\phi \wedge \phi_1$ and no Horn clause from Φ_1 contains an *T*-atom in its conclusion. Since ψ_0 is of the form (4), we must have

$$\Phi_1 \models \forall x_1, \dots, x_n, y_1 \big(\phi' \land \phi_1 \Rightarrow R_S(y_1, x_n) \big),$$

otherwise a counterexample to (12) can be easily constructed. But then it follows from Claim 15 that $a_1 \ldots a_n \in L(\mathcal{G})$ and we are done.

" \Leftarrow " We prove the contrapositive. We assume that fm(Φ) does not have the AP. Then there exists a counterexample to Lemma 7, i.e., there exists a Horn clause ψ of the form $\phi(\bar{x}) \wedge \phi_1(\bar{x}, y_1) \wedge \phi_2(\bar{x}, y_2) \Rightarrow \chi$, where ϕ, ϕ_1 , and ϕ_2 satisfy the prerequisites of Lemma 7 and $\chi(\bar{x}, y_1)$ is an atomic τ -formula other than equality, such that

$$\Phi \models \forall \bar{x}, y_1, y_2 \big(\phi \land \phi_1 \land \phi_2 \Rightarrow \chi \big) \tag{13}$$

and
$$\Phi' \not\models \forall \bar{x}, y_1 (\phi \land \phi_1 \Rightarrow \chi).$$
 (14)

We choose ψ minimal with respect to the number of its atomic subformulas.

Our proof strategy is as follows. First we show that ψ encodes a single word $w \in \Sigma^+$ in the sense of Claim 16. Then we show that the word w may not be contained in $L(\mathcal{G})$, because otherwise a part of the counterexample would encode w in the sense of Claim 15 which would lead to a contradiction.

 \triangleright Claim 17. The formula $\Phi \land \phi(\bar{x}) \land \phi_i(\bar{x}, y_i)$ is satisfiable for both $i \in \{1, 2\}$.

Proof. Suppose for contradiction that $\Phi \models \forall x_1, \ldots, x_n, y_i (\phi \land \phi_i \Rightarrow \bot)$ for some $i \in \{1, 2\}$. Since the conclusion of $\phi \land \phi_i \Rightarrow \bot$ does not contain the variable y_i and $\phi(\bar{x}) \land \phi_i(\bar{x}, y_i) \leq \Phi$ $\phi(\bar{x})$ for both $i \in [2]$, it follows that $\Phi \models \forall \bar{x} (\phi(\bar{x}) \Rightarrow \bot)$. But this yields a contradiction to (14). Thus, the statement holds.

- \triangleright Claim 18. The formula ψ
- \blacksquare is not a tautology,
- \blacksquare has an SLD-deduction from Φ_2 ,
- \blacksquare only contains τ_2 -atoms.

Proof. First, we claim that $\Phi_1 \vdash \psi$ or $\Phi_2 \vdash \psi$. By Theorem 8, we have $\Phi \vdash \psi$. Note that $\chi(\bar{x}, y_1)$ cannot be a subformula of $\phi(\bar{x}) \land \phi_1(\bar{x}, y_1)$, by (14). Also note that $\chi(\bar{x}, y_1)$ cannot be a subformula of $\phi_2(\bar{x}, y_2)$, because every atom in $\phi_2(\bar{x}, y_2)$ contains the variable y_2 . Hence, $\chi(\bar{x}, y_1)$ is not a subformula of $\phi(\bar{x}) \land \phi_1(\bar{x}, y_1) \land \phi_2(\bar{x}, y_2)$, i.e., ψ is not a tautology.

Thus, ψ is a weakening of a Horn clause ψ' which has an SLD-derivation $\psi'_0, \ldots, \psi'_s = \psi'$ from Φ modulo renaming variables. Note that the Horn clauses in Φ have the property that, depending on whether they come from Φ_1 or from Φ_2 , they either contain no Q-atoms or no R_A -atoms for $A \in N$, respectively. This applies in particular to ψ'_0 which is a conjunct of Φ . Since the conclusion of each Horn clause in Φ_1 is an R_A -atom for $A \in N$ and the conclusion of each Horn clause in Φ_2 is a Q-atom, the property of ψ'_0 from above propagates inductively to every ψ'_i for $i \in [k]$. But this means that ψ' has an SLD-derivation from Φ_1 or from Φ_2 . Hence, ψ has an SLD-deduction from Φ_1 or from Φ_2 , which concludes the claim.

Next, we claim that $\Phi_2 \vdash \psi$. Suppose, on the contrary, that $\Phi_1 \vdash \psi$. Let ϕ', ϕ'_1 , and ϕ'_2 be the formulas obtained from ϕ, ϕ_1 , and ϕ_2 , respectively, by removing all *Q*-atoms. Since $\Phi_1 \vdash \psi$, the SLD-derivation sequence ψ'_0, \ldots, ψ'_s from the first paragraph contains no *Q*-atoms. Thus, all *Q*-atoms occurring in ψ come from the weakening step, which means that

$$\Phi_1 \vdash \forall \bar{x}, y_1, y_2 (\phi' \land \phi'_1 \land \phi'_2 \Rightarrow \chi). \tag{15}$$

We show that, for both $i \in [2]$,

$$\phi'(\bar{x}) \wedge \phi'_i(\bar{x}, y_i) \le_{\Phi_1} \phi'(\bar{x}). \tag{16}$$

Let $\eta(\bar{x})$ be an atomic τ_1 -formula other than equality such that $\Phi_1 \models \forall \bar{x}, y_i (\phi' \land \phi'_i \Rightarrow \eta)$ for some $i \in [2]$. Since $\phi(\bar{x}) \land \phi_i(\bar{x}, y_i) \leq \phi(\bar{x})$ for both $i \in [2]$, we have $\Phi \models \forall \bar{x} (\phi \Rightarrow \eta)$. Clearly, for both $i \in [2]$, η cannot be a subformula of ϕ_i because every atom in ϕ_i contains the variable y_i . If η is a subformula of ϕ , then trivially $\Phi_1 \models \forall \bar{x}(\phi' \Rightarrow \eta)$ and the claim holds. So suppose that this is not the case. By Theorem 8, we have $\Phi \vdash \forall \bar{x}(\phi \Rightarrow \eta)$. Since $\phi \Rightarrow \eta$ is not a tautology, it is a weakening of a Horn clause which has an SLD-derivation from Φ modulo renaming variables. Since $\Phi \wedge \phi$ is satisfiable by Claim 17, η neither is of the form \perp nor was it obtained by replacing \perp with a different τ_1 -atom during the weakening step. Consequently, η can only be an R_A -atom for some $A \in N$ because other symbols from τ_1 do not appear in the conclusion of any Horn clause in Φ . But then, since R_A -atoms only appear in the conclusion of Horn clauses from Φ_1 , and no Horn clause from Φ_2 contains an atom in its conclusion that would appear in the premise of a Horn clause from Φ_1 , it must be the case that $\Phi_1 \vdash \forall \bar{x}(\phi \Rightarrow \eta)$. Consequently, $\Phi_1 \vdash \forall \bar{x}(\phi' \Rightarrow \eta)$ because $\{Q, R\}$ -atoms do not appear in any Horn clause from Φ_1 . Since η was chosen arbitrarily, we conclude that (16) indeed holds. Now we come back to the SLD-derivation sequence ψ'_0, \ldots, ψ'_s . Since ψ'_0 is a Horn clause from Φ_1 , it follows from the minimality assumption for ψ that either χ is an R_A -atom for some $A \in N$, or χ equals \perp . In both cases, (16), (15), and (14) witness that $fm(\Phi_1)$ does not have AP through an application of Lemma 7. But this is in contradiction to the fact that $fm(\Phi_1)$ is preserved by unions because Φ_1 is complete. Thus, $\Phi_1 \vdash \psi$ does not hold, and $\Phi_2 \vdash \psi$ holds instead.

Since we have $\Phi_2 \vdash \psi$, the premise $\phi \land \phi_1 \land \phi_2$ of ψ can only contain symbols from τ_2 , otherwise we could remove all $(\tau_1 \setminus \tau_2)$ -atoms and get a contradiction to the minimality of ψ . Since ψ'_0 is a Horn clause from Φ_2 , it also follows from the minimality assumption for ψ that either χ is an Q-atom, or χ is \bot . Thus, ψ only contains symbols from τ_2 .

Let ψ' be a Horn clause such that ψ is a weakening of ψ' and there is an SLD-derivation $\psi'_0, \ldots, \psi'_s = \psi'$ from Φ_2 modulo renaming variables. Recall that, since ψ is a weakening of ψ' , every atom from the premise of ψ' also appears in the premise of ψ .

 \triangleright Claim 19. ψ' does not contain any Q-atoms and its conclusion equals \perp .

Proof. Note that, by the construction of Φ_2 , for every $i \in [k]$, if there exist variables z_1, z_2 in ψ'_{i-1} such that

every Q-atom contains z_1 in its first and z_2 in its second argument, respectively, (17)

then ψ'_i also satisfies (17) for the same variables z_1, z_2 up to renaming. Since every possible choice of ψ'_0 from Φ_2 initially satisfies (17) for some variables, it follows via induction that (17) must hold for $\psi' = \psi'_s$ for some variables. Also note that (6) is the only Horn clause in Φ_2 that is not complete, but the incompleteness is only due to one missing edge in the Gaifman graph between the two distinguished variables satisfying (17) for (6).

We claim that $\{z_1, z_2\} = \{y_1, y_2\}$ holds for the pair z_1, z_2 satisfying (17) for ψ' . Suppose, on the contrary, that both z_1 and z_2 are among \bar{x}, y_1 or \bar{x}, y_2 . For every $i \in [k], \psi'_i$ is a resolvent of ψ'_{i-1} and a Horn clause from Φ_2 which is almost complete except for one missing edge in the Gaifman graph between a pair of variables which must be substituted for the pair (z_1, z_2) satisfying (17) for ψ'_{i-1} . Since the variables y_1 and y_2 do not appear together in any atom in ψ' and $\{z_1, z_2\} \neq \{y_1, y_2\}$, they also do not appear together in any atom during the SLD-derivation. Then it follows from the fact that ϕ, ϕ_1 , and ϕ_2 satisfy the prerequisites of Lemma 7 that we already have $\Phi_2 \vdash \forall \bar{x}, y_1(\phi \land \phi_1 \Rightarrow \chi)$, a contradiction to (14). Therefore, $\{z_1, z_2\} = \{y_1, y_2\}.$

We claim that ψ'_0 is of the form (8). Otherwise, ψ'_0 is of the form (6) or (7), in which case the conclusion of ψ' is a Q-atom. But then, since z_1, z_2 with $\{z_1, z_2\} = \{y_1, y_2\}$ satisfy (17) for ψ' , the conclusion of ψ' would be an atom containing both variables y_1 and y_2 . This leads to a contradiction to (13) where we assume that the conclusion of ψ' may only contain variables from \bar{x}, y_1 . The claim implies that the conclusion of ψ' equals \perp . Hence, ψ' does not contain any Q-atoms at all. \triangleleft

By Claim 16, there exists $a_1, \ldots, a_n \in \Sigma^+$ such that the premise of ψ' contains the premise of (5) as a subformula modulo renaming and indentifying variables. Since ϕ, ϕ_1 , and ϕ_2 satisfy the prerequisites of Lemma 7, it must be the case that

- $= \phi(\bar{x}) \text{ has } T(x_n) \wedge \bigwedge_{i \in [n-1]} R_{a_{i+1}}(x_i, x_{i+1}) \text{ as a subformula,}$ $= \phi_1(\bar{x}, y_1) \text{ has } L(y_1) \wedge I(y_1) \wedge R_{a_1}(y_1, x_1) \wedge \bigwedge_{i \in [n-1]} E(y_1, x_{i+1}) \text{ as a subformula, and}$ $= \phi_2(\bar{x}, y_2)$ has $R(y_2) \wedge \bigwedge_{i \in [n]} E(y_2, x_i)$ as a subformula,

where the variables need not all be distinct and the roles of ϕ_1 and ϕ_2 might be swapped. Otherwise, we get a contradiction to Claim 17. Note that, if $L(\mathcal{G}) = \Sigma^+$, then Claim 15 together with (4) implies that

$$\Phi_1 \models \forall x_1, \dots, x_{n+1} (\phi \land \phi_1 \Rightarrow \bot).$$
⁽¹⁸⁾

If some variables among x_1, \ldots, x_n are identified in $\phi \land \phi_1 \land \phi_2$, then we still have (18) even if we perform the same identification of variables. But then we get a contradiction to Claim 17. Thus, $L(\mathcal{G}) \neq \Sigma^+$.

We have thus found a reduction from the PSPACE-hard universality problem for \mathcal{G} to the decidability problem of the AP for $fm(\Phi)$; note that Φ is universal Horn and can be computed from G in polynomial time.

A.4 Proof of Lemma 14

▶ Lemma 14. Let 𝔅 be a context-free grammar. Then the following are equivalent:

1. $L(\mathcal{G})$ is regular.

- **2.** $\Phi_{\mathcal{G}}$ is equivalent to monadic Datalog sentence.
- **3.** $\Phi_{\mathfrak{G}}$ is equivalent to a GMSNP sentence.
- **4.** fm($\Phi_{\mathfrak{F}}$) is the CSP of a finite structure.
- **5.** fm($\Phi_{\mathfrak{F}}$) is the CSP of an ω -categorical structure.
- **6.** $\operatorname{fm}(\Phi_{\mathfrak{F}})$ is the age of a reduct of a finitely bounded homogeneous Ramsey structure.
- 7. fm($\Phi_{\mathcal{G}}$) is the age of an ω -categorical structure.
- **8.** fm($\Phi_{\mathfrak{F}}$) is finitely bounded homogenizable.

Before we can prove the lemma, we need to introduce some additional terminology and one model-theoretic result from [16].

A deterministic finite automaton (DFA) is a 5-tuple $\mathcal{A} = (Q, \Sigma, \delta, q_0, F)$ where Q is a finite set of states, Σ is a finite set of input symbols, $\delta : Q \times \Sigma \to Q$ is a transition function, $q_0 \in Q$ is a distinguished starting state, and $F \subseteq Q$ is a distinguished set of final states. The language of \mathcal{A} is $L(\mathcal{A}) \coloneqq \{a_1 \dots a_n \in \Sigma^+ \mid \delta(a_n, \dots \delta(a_2, \delta(a_1, q_0)) \dots) \in F\}$. Note that, as in the case of CFGs, the empty word ϵ can never be an element of $L(\mathcal{A})$ according to the present definition.

Let Σ be a finite set of symbols and L a subset of Σ^+ . The *Myhill-Nerode equivalence* relation on Σ^* , denoted by \sim_L , is defined by $w_1 \sim_L w_2$ if there is no $w \in \Sigma^*$ such that $|\{w_1w, w_2w\} \cap L| = 1$. The following correspondence is well-known.

▶ **Theorem 20** (Myhill-Nerode). For every finite set Σ and every $L \subseteq \Sigma^+$, the following are equivalent:

- \blacksquare L is regular;
- \blacksquare L is accepted by a DFA;
- $\sim_L has finitely many classes.$
- ▶ Example 21. $\{a^m b^n \mid m, n \ge 1\}$ is regular while $\{a^n b^n \mid n \ge 1\}$ is not.

Let $\mathcal{A} = (Q, \Sigma, \delta, q_0, F)$ be a DFA. The signature τ_{Σ} is defined as before, and the signature τ_Q consists of the unary symbols R_q for every $q \in Q$ that is *reachable from* q_0 , i.e., there exists a word $a_1 \ldots a_n \in \Sigma^+$ such that $q = \delta(a_n, \ldots \delta(a_2, \delta(a_1, q_0)) \ldots)$. Note that q_0 is not necessarily reachable from itself. Let $\overline{\Phi}_{\mathcal{A}}$ be the universal Horn sentence over the signature $\tau_{\Sigma} \cup \tau_Q$ whose quantifier-free part contains: for every $a \in \Sigma$, the Horn clause

$$I(x_1) \land R_a(x_1, x_2) \Rightarrow R_{\delta(q_0, a)}(x_2)$$

for every $(q, a) \in Q \times \Sigma$ such that q is reachable from q_0 , the Horn clause

$$R_q(x_1) \wedge R_a(x_1, x_2) \Rightarrow R_{\delta(q,a)}(x_2),$$

and, for every $q \in F$ that is reachable from q_0 , the Horn clause

 $R_q(x_1) \wedge T(x_1) \Rightarrow \bot.$

Then $\Phi_{\mathscr{A}}$ is the connected monadic Datalog sentence obtained from $\overline{\Phi}_{\mathscr{A}}$ by existentially quantifying all symbols from τ_Q upfront. The following lemma can be proved similarly as Lemma 13, and its proof is omitted.

▶ Lemma 22. For a τ_{Σ} -structure \mathfrak{A} , we have $\mathfrak{A} \models \Phi_{\mathfrak{A}}$ if and only if, for every $w \in L(\mathfrak{A})$,

$$\mathfrak{A} \models \forall x_1, \dots, x_{|w|+1} \Big(I(x_1) \land \phi_w(x_1, \dots, x_{|w|+1}) \land T(x_{|w|+1}) \Rightarrow \bot \Big).$$

To the following model-theoretic correspondence was essentially proved in [16].

▶ **Theorem 23** ([16]). Every GMSNP sentence is equivalent to a finite disjunction of connected GMSNP sentences. If Φ is a connected GMSNP sentence, then there exists a reduct \mathfrak{R} of a finitely bounded homogeneous Ramsey structure with $\operatorname{fm}(\Phi) = \operatorname{age}(\mathfrak{R}) = \operatorname{CSP}(\mathfrak{R})$.

Proof of Theorem 23, second part only. The fact that every GMSNP sentence is equivalent to a finite disjunction of connected GMSNP sentences was proved in [16] in great detail, so we refer the reader there and start already with a connected GMSNP sentence.

Let $\Phi := \exists X_1, \ldots, X_n \forall \bar{x}. \phi$ be an arbitrary connected GMSNP sentence and τ its signature. Set $\sigma := \{X_1, \ldots, X_n\}$. We define two new signatures τ' and σ' by introducing two new symbols R^+ and R^- for every $R \in \tau \cup \sigma$. Then we obtain a new GMSNP sentence $\Phi' := \exists X_1^+, X_1^-, \ldots, X_n^+, X_n^- \forall \bar{x}. \phi'$ over τ' as follows. In every clause of ϕ , we replace each negative atom $\neg R(\bar{x})$ the negative atom $\neg R^+(\bar{x})$ and each positive atom $R(\bar{x})$ with the negative atom $\neg R^-(\bar{x})$. The new GMSNP sentence is still connected, and additionally has the property that none of its clauses contains positive atoms. It follows that $\operatorname{fm}(\forall \bar{x}. \phi')$ is defined by forbidden homomorphic images from finitely many connected finite $\tau' \cup \sigma'$ -structures. By Theorem 1.3 in [34], there exists a particular finitely bounded class \mathcal{A} over a larger signature ρ which has the AP and whose $\tau' \cup \sigma'$ -reducts are precisely the members of $\operatorname{fm}(\forall \bar{x}. \phi')$. Let < be a fresh binary symbol not contained in ρ . We define \mathcal{R} as the class of all $\{<\} \cup \rho$ -expansions of structures from \mathcal{A} where < interprets as a linear order. Clearly, \mathcal{R} is still finitely bounded. By Theorem 2.11 in [33], \mathcal{R} is a Ramsey class (see also Theorem A.3 in [18]). Let \mathfrak{F} be the homogeneous structure from Theorem 1 associated to \mathcal{R} .

We say that a set $S \subseteq F$ is correctly labeled if, for every k-ary $R \in \tau' \cup \sigma'$ and every k-ary tuple \overline{t} over S, we have $\mathfrak{F} \models R^+(\overline{t}) \Leftrightarrow \neg R^-(\overline{t})$. Now consider the $\{<\} \cup \tau \cup \sigma \cup \rho$ -expansion $\overline{\mathfrak{F}}$ of \mathfrak{F} where each k-ary $R \in \tau \cup \sigma$ interprets as the k-ary relation defined by the quantifier-free first-order formula $R^+(x_1, \ldots, x_k) \land \{x_1, \ldots, x_k\}$ is correctly labeled. Since first-order definable relations are preserved by automorphisms, $\overline{\mathfrak{F}}$ is still homogeneous. We define \mathfrak{B} as the τ -reduct of $\overline{\mathfrak{F}}$.

" \Rightarrow " Suppose that $\mathfrak{A} \in \operatorname{CSP}(\mathfrak{B})$, i.e., there exists a homomorphism $h: \mathfrak{A} \to \mathfrak{B}$. We define $\overline{\mathfrak{A}}$ as the expansion of \mathfrak{A} obtained by pulling back the relations from $\overline{\mathfrak{F}}$ through h. Clearly h is a homomorphism from $\overline{\mathfrak{A}}$ to $\overline{\mathfrak{F}}$. Let $\neg \psi_1 \lor \cdots \lor \neg \psi_k \lor \psi_{k+1} \lor \cdots \lor \psi_\ell$ be a clause in Φ , where each ψ_i is a positive atomic formula. Suppose that $\overline{\mathfrak{A}} \models \psi_1 \land \cdots \land \psi_k(\overline{a})$. Since his a homomorphism, we have $\overline{\mathfrak{F}} \models \psi_1 \land \cdots \land \psi_k(e(\overline{a}))$. For every atom ψ of the form $R(\overline{x})$ for $R \in \tau' \cup \sigma'$, we define ψ^+ as $R^+(\overline{x})$ and ψ^- as $R^-(\overline{x})$. By the definition of \mathfrak{B} , we have $\overline{\mathfrak{F}} \models \psi_1^+ \land \cdots \land \psi_k^+(e(\overline{a}))$. By the definition of $\overline{\mathfrak{F}}$, we have $\overline{\mathfrak{F}} \models \neg \psi_{k+1}^- \lor \cdots \lor \neg \psi_\ell^-(e(\overline{a}))$, i.e., there exists $i \in [\ell] \setminus [k]$ such that $\overline{\mathfrak{F}} \models \neg \psi_i^-(e(\overline{a}))$. Since Φ is guarded, there exists $j \in [k]$ such that all variables \overline{x}_i of ψ_i are among the variables \overline{x}_j of ψ_j . By the assumption that the the set $\{h(\overline{x}_j)[1], \ldots\}$ is correctly labeled, we have $\overline{\mathfrak{F}} \models \psi_i^+(e(\overline{a}))$. Clearly $\{h(\overline{x}_i)[1], \ldots\}$ is also correctly labeled, and hence $\overline{\mathfrak{F}} \models \psi_i(e(\overline{a}))$. Since Φ is monotone, ψ_i is of the form $R(\overline{x})$ for $R \in \sigma'$. Hence, by the definition of $\overline{\mathfrak{A}}$, we have $\overline{\mathfrak{A}} \models \psi_i(e(\overline{a}))$.

"⇐" Suppose that $\mathfrak{A} \models \Phi$, and let \mathfrak{A}_{Φ} be a $\tau \cup \sigma$ -expansion of \mathfrak{A} witnessing this fact. We define \mathfrak{A}' as the $\tau' \cup \sigma'$ structure with domain A and where R^+ interprets as $R^{\mathfrak{A}_{\Phi}}$ and R^- as the complement of $R^{\mathfrak{A}_{\Phi}}$ in \mathfrak{A}_{Φ} . Then clearly $\mathfrak{A}' \models \forall \bar{x}. \phi'$. Consequently, \mathfrak{A}' is a reduct of a substructure $\overline{\mathfrak{A}}$ of $\overline{\mathfrak{F}}$. Clearly, the τ -reduct of $\overline{\mathfrak{A}}$ is isomorphic to \mathfrak{A} . It follows that $\mathfrak{A} \in \operatorname{age}(\mathfrak{B})$.

Proof of Lemma 14. " $(2) \Rightarrow (3)$ " This direction is trivial.

"(4) \Rightarrow (5)" This direction is also trivial.

"(6) \lor (8) \Rightarrow (7)" This direction is well-known and easy to see [32].

"(3) \Rightarrow (6)" Since $\Phi_{\mathscr{G}}$ is connected, fm($\Phi_{\mathscr{G}}$) is preserved by disjoint unions. Since $\Phi_{\mathscr{G}}$ is equivalent to a GMSNP sentence, by the first part of Theorem 23, $\Phi_{\mathscr{G}}$ is equivalent to a disjunction $\Phi_1 \lor \cdots \lor \Phi_k$ of connected GMSNP sentences. We may assume that k is minimal with this property. If $k \geq 2$, then, by the minimality of k, there exist $\mathfrak{A}_1, \mathfrak{A}_2 \in \Phi_{\mathscr{G}}$ such that $\mathfrak{A}_i \models \Phi_i \land \bigwedge_{j \in [k] \setminus \{i\}} \neg \Phi_j$ for $i \in \{1, 2\}$. But then the disjoint union of \mathfrak{A}_1 and \mathfrak{A}_2 does not satisfy $\Phi_{\mathscr{G}}$, a contradiction. Hence, k = 1. Now the statement follows from the second part of Theorem 23.

"(1) \Rightarrow (2) \wedge (4) \wedge (8)" By Theorem 20, there exists a DFA \mathscr{A} such that $L(\mathscr{G}) = L(\mathscr{A})$. By Lemma 13 and Lemma 22, we have fm($\Phi_{\mathscr{G}}$) = fm($\Phi_{\mathscr{A}}$). This implies item (2) because $\Phi_{\mathscr{A}}$ is in connected monadic Datalog. In fact, the sentence falls into an even stricter fragment which was called *caterpillar Datalog* in [30]. By Theorem 4.1 in [30], there exists a finite structure whose CSP equals fm($\Phi_{\mathscr{A}}$). This implies item (4). Finally, we show that fm($\Phi_{\mathscr{G}}$) is finitely bounded homogenizable. Since the universal sentence $\overline{\Phi}_{\mathscr{A}}$ is complete, the class fm($\overline{\Phi}_{\mathscr{A}}$) is closed under unions, and therefore has the AP. Let \mathfrak{H} be the homogeneous structure from Theorem 1 associated with fm($\overline{\Phi}_{\mathscr{A}}$). By the definition of fm($\overline{\Phi}_{\mathscr{A}}$), the signature τ_Q only contains unary symbols R_q for those $q \in Q$ which are reachable from q_0 . Let $Q_0 \subseteq Q$ be the set of states reachable from q_0 and, for every $q \in Q_0$, let $w_q \in \Sigma^+$ be an arbitrary word witnessing that q is reachable from q_0 . Note that, if $q_1, q_2 \in Q_0$ are distinct, then $w_{q_1} \neq w_{q_2}$. For every $w \in \Sigma^+$, consider the unary formula

$$\phi_w^I(x_1) \coloneqq \exists x_2, \dots, x_{|w|+1} \big(I(x_{|w|+1}) \land \phi_w(x_{|w|+1}, \dots, x_1) \big).$$

We show that, for every $q \in Q_0$, the unary formulas $R_q(x)$ and $\phi_{w_q}^I(x)$ define the same relation in \mathfrak{H} . Then, for every $q \in Q_0$, the relation $R_q^{\mathfrak{H}}$ is first-order definable in the τ_{Σ} reduct of \mathfrak{H} and is therefore preserved by all of its automorphisms. This implies item (8). First, suppose that $\mathfrak{H} \models \phi_{w_q}^I(x)$ for some $x \in H$. Since w_q witnesses that q is reachable from q_0 and $\mathfrak{H} \models \overline{\Phi}_{\mathfrak{A}}$, it follows that $\mathfrak{H} \models R_q(x)$. Next, suppose that $\mathfrak{H} \models R_q(x)$ for some $x \in H$. Since \mathfrak{A} is deterministic, by the definition of $\overline{\Phi}_{\mathfrak{A}}$, there exists a structure \mathfrak{A} in the signature of \mathfrak{H} with domain $\{x_{|w_q|+1}, \ldots, x_1\}$ that satisfies $\mathfrak{A} \models \overline{\Phi}_{\mathfrak{A}} \wedge I(x_{|w_q|+1}) \wedge \phi_{w_q}(x_{|w_q|+1}, \ldots, x_1)$ and whose substructure on $\{x_1\}$ is isomorphic to the substructure of \mathfrak{H} on $\{x\}$. Since $\mathfrak{A} \models \overline{\Phi}_{\mathfrak{A}}$, there exists an embedding $e: \mathfrak{A} \hookrightarrow \mathfrak{H}$. Since \mathfrak{H} is homogeneous, there exists an automorphism f of \mathfrak{H} such that $f \circ e(x_1) = x$. Since automorphisms preserve first-order definable relations, it follows that $\mathfrak{H} \models \phi_{w_q}^I(x)$.

" $(5) \vee (7) \Rightarrow (1)$ " Let \mathfrak{D} be an ω -categorical structure such that $\operatorname{fm}(\Phi_{\mathfrak{G}}) = \operatorname{age}(\mathfrak{D})$ or $\operatorname{fm}(\Phi_{\mathfrak{G}}) = \operatorname{CSP}(\mathfrak{D})$. Suppose, on the contrary, that $L(\mathfrak{G})$ is not regular. For every $w \in \Sigma^+$, consider the unary formula

$$\phi_w^T(x_1) \coloneqq \exists x_2, \dots, x_{|w|+1} \big(\phi_w(x_1, \dots, x_{|w|+1}) \wedge T(x_{|w|+1}) \big).$$

Since $L(\mathfrak{G})$ is not regular, by Theorem 20, $\sim_{L(\mathfrak{G})}$ has infinitely many classes. Let w_1, \ldots be the representatives of the classes. For every $i \geq 1$ let $R_i \subseteq D$ be the unary relation defined by $\phi_{w_i}^I(x_1)$ in \mathfrak{D} . For every $i \geq 1$, the formula $\phi_{w_i}^I(x_1)$ is satisfiable in \mathfrak{D} because it does not contain any *T*-atom. Since first-order formulas are preserved under automorphisms, for every $i \geq 1$, R_i is a non-empty union of unary orbits. By the definition of $\sim_{L(\mathfrak{G})}$, for every pair $i \neq j$, there exists $w \in \Sigma^*$ such that $|\{w_i w, w_j w\} \cap L(\mathfrak{G})| = 1$. By Lemma 13, exactly one of the relations R_i and R_j contains an element satisfying $\phi_w^T(x_1)$ in \mathfrak{D} . In other words, there exists an orbit that is contained in one of the two relations but not in the other. This can only be the case for all $i \neq j$ if there are infinitely many unary orbits, a contradiction to ω -categoricity. Therefore, $L(\mathfrak{G})$ is regular.

B Proofs of Theorem 5 and Theorem 6

In the next two subsections, we only provide the exact specifications of the conjuncts in Φ_1 and Φ_2 . The actual proof of the theorems remains almost unchanged, except for the small detail that the structures $\mathfrak{B}_1, \mathfrak{B}_2$ from the first part must additionally satisfy the formula N(0, 1). The presence of this atom will be explained later, but we can already reveal that it has to do with the simple mechanism for distinguishing between the variables 0 and 1.

B.1 Inputs specified by universal sentences

The first part Φ_1 does not explain how our reduction works, but ensures that it does not fall apart, e.g., due to ill-behaved identifications of variables. For every $\alpha \in T$, the signature τ contains an (n + 1)-ary symbol T_{α} . The first n arguments in a T_{α} -atom serve as binary counters, and the last argument carries a given path node p. Suppose that the variables 0 and 1 represent the bits 0 and 1, respectively. Then each atomic formula $T_{\alpha}(c_1, \ldots, c_n, p)$ with $c_1, \ldots, c_n \in \{0, 1\}$ represents the situation in which a tile α is present in the p-th row and in the $(1 + \sum_{k \in [n]} (c_k = 1) \cdot 2^{n-k})$ -th column. First, we want to ensure the horizontal consistency of the tiling. To this end, for every pair $(\alpha, \beta) \in T^2 \setminus P_h^{\mathfrak{T}}$, we include in Φ_1 a complete Horn sentence ensuring that two horizontally adjacent positions in the p-th row cannot be tiled with α and β . Here, we encode the successor relation w.r.t. binary addition using a combination of equalities: $(c_{n+1}, \ldots, c_{2n})$ is the successor of (c_1, \ldots, c_n) if and only if there exists $j \in [n]$ such that

■ $c_i = c_{i+n} \in \{0, 1\}$ for every $i \in [j-1]$, ■ $c_j = 0$ and $c_{n+j} = 1$, and ■ $c_i = 1$ and $c_{i+n} = 0$ for every $i \in [n] \setminus [j]$.

This encoding only makes sense if 0 and 1 truly represent the bits 0 and 1, so we introduce a simple mechanism (in terms of a Horn sentence) that distinguishes between the variables 0 and 1. We include in τ the binary symbol N that will later be used to distinguish between 0 and 1, and the following sentences as conjuncts in Φ_1 :

$$\forall x, y (N(x,y) \land N(y,x) \Rightarrow \bot), \tag{19}$$

and, for every pair $(\alpha, \beta) \in T^2 \setminus P_h^{\mathfrak{T}}$, the sentence

$$\forall p, 0, 1, c_1, \dots, c_{2n}$$

$$\begin{pmatrix} N(0, 1) \land T_{\alpha}(c_1, \dots, c_n, p) \land T_{\beta}(c_{n+1}, \dots, c_{2n}, p) \\ \land \left(\bigvee_{j \in [n]} \bigwedge_{i \in [j-1]} (c_i = 0 \land c_{i+n} = 0 \lor c_i = 1 \land c_{i+n} = 1) \\ \land c_j = 0 \land c_{j+n} = 1 \land \bigwedge_{i \in [n] \setminus [j]} c_i = 1 \land c_{i+n} = 0 \end{pmatrix}$$

$$\downarrow$$

$$\downarrow$$

Next, we want to ensure that every position in the *p*-th row is occupied by at most one tile. To this end, for every $(\alpha, \beta) \in T^2$ with $\alpha \neq \beta$, we include

$$\forall p, 0, 1, c_1, \dots, c_n \big(T_\alpha(c_1, \dots, c_n, p) \land T_\beta(c_1, \dots, c_n, p) \land \bigwedge_{i \in [n]} (c_i = 0 \lor c_i = 1) \Rightarrow \bot \big)$$

as a conjunct in Φ_1 . Finally, we want to ensure that each verifier node v represents at most one number from $[2^n]$. For every $i \in [n]$, τ contains two unary symbols O_i and 1_i which

will be used to encode numbers in binary. We include $\forall v (\bigvee_{i \in [n]} O_i(v) \land \mathbf{1}_i(v) \Rightarrow \bot)$ as the last conjunct in Φ_1 . Now the idea is that the combinations of atomic formulas $O_i(v)$ and $\mathbf{1}_i(v)$ at a verifier node v will be compared with the combinations of 0 and 1 in atomic formulas $T_{\alpha}(c_1, \ldots, c_n, p)$. The Horn sentences in the second part of Φ will be formulated so that verifying the presence of all 2^n atoms of the form $T_{\alpha}(c_1, \ldots, c_n, p_1)$ at a path node p_1 using verifier nodes is the only possible way to progress to a next path node p_2 . As a consequence, we do not need to add an explicit requirement for rows, represented by path nodes, to be completely tiled from left to right. For the same reason, we also do not need to add an explicit requirement for verifier nodes to represent at least one number from $[2^n]$.

We now proceed with the sentence Φ_2 , which explains how the parsing of a tiling actually works. We use $SUCC(v_1, v_2)$ as a shortcut for the formula

$$\bigvee_{j\in[n]} \bigwedge_{i\in[j-1]} \left(\mathbf{0}_i(v_1) \wedge \mathbf{0}_i(v_2) \vee \mathbf{1}_i(v_1) \wedge \mathbf{1}_i(v_2) \right) \\ \wedge \mathbf{0}_j(v_1) \wedge \mathbf{1}_j(v_2) \wedge \bigwedge_{i\in[n]\setminus[j]} \mathbf{1}_i(v_1) \wedge \mathbf{0}_i(v_2).$$

We will need SUCC (v_1, v_2) to represent the successor relation for verifier nodes. We will also need an auxiliary formula for testing whether a sequence of bits represents the same number as a particular verifier node. This is achieved using the formula $EQ(v, 0, 1, c_1, \ldots, c_n)$, which is defined as

$$\bigwedge\nolimits_{j\in[n]} \left(0_j(v) \wedge c_j = 0 \vee 1_j(v) \wedge c_j = 1 \right)$$

Finally, we use $PAD(y_1, y_2, x_1, \dots, x_k)$ as a shortcut for the formula

$$L(y_1) \wedge R(y_2) \wedge \bigwedge_{i \in [k]} E(y_1, x_i) \wedge E(y_2, x_i) \wedge \bigwedge_{j \in [k]} E(x_i, x_j)$$

The parsing of a tiling starts from a path node p representing a row whose leftmost position contains a tile that can be present in the bottom left corner of a tiling grid. This must be confirmed by a verifier node, in which case a 6-ary Q_b -atom is derived, representing the fact that the leftmost column of the p-th row has been checked. To this end, we include in Φ_2 suitable Horn sentences for every $\alpha \in P_{\ell}^{\mathfrak{T}} \cap P_b^{\mathfrak{T}}$:

$$\left(\underbrace{\begin{array}{c} N(0,1) \wedge T_{\alpha}(c_{1}, \dots, c_{n}, p) \wedge \bigwedge_{j \in [n]} c_{j} = 0 \\ \wedge \operatorname{EQ}(v, 0, 1, c_{1}, \dots, c_{n}) \wedge \operatorname{PAD}(y_{1}, y_{2}, p, z, 0, 1) \\ & \downarrow \\ Q_{b}(y_{1}, y_{2}, p, v, 0, 1) \end{array}}_{Q_{b}(y_{1}, y_{2}, p, v, 0, 1)} \right).$$

$$(20)$$

Note that, by (19), the presence of the atom N(0, 1) in (20) creates an asymmetry between 0 and 1. At this point, it might not be clear in which sense 0 and 1 represent 0 and 1, respectively, and not the other way around. However, a detailed inspection of the construction reveals that this will be achieved implicitly through the use of EQ and SUCC. These sentences form the only non-complete part of Φ ; we intentionally leave a missing edge between y_1 and y_2 in the Gaifman graph to enable potential formation of AP failures.

Using 2^n -many verifier nodes and propagation of Q_b -atoms, the whole bottom row is checked for the presence of tiles. Their horizontal consistency already follows from the conditions imposed on path nodes by Φ_2 and needs not to be checked during this step. To

this end, we include in Φ_2 suitable Horn sentences for every $\alpha \in P_b^{\mathfrak{T}}$:

$$\left(\underbrace{\begin{array}{c} T_{\alpha}(c_{1},\ldots,c_{n},p) \land \mathrm{EQ}(v_{1},0,1,c_{1},\ldots,c_{n}) \land \mathrm{SUCC}(v_{1},v_{2}) \\ \land Q_{b}(y_{1},y_{2},p,v_{1},0,1) \land \mathrm{PAD}(y_{1},y_{2},p,v_{1},v_{2},0,1) \\ & \downarrow \\ Q_{b}(y_{1},y_{2},p,v_{2},0,1) \end{array} \right)$$

After the *p*-th row has been checked by a 2^n -th verifier node, we mark the variable *p* with a Q-atom that indicates that the parsing can progress to a successor path node. To this end, we include in Φ_2 suitable Horn sentences for every $\alpha \in P_b^{\mathfrak{T}} \cap P_r^{\mathfrak{T}}$:

$$\forall y_1, y_2, p, v, 0, 1, c_1, \dots, c_n \\ \begin{pmatrix} T_{\alpha}(c_1, \dots, c_n, p) \land \bigwedge_{j \in [n]} c_j = 1 \land \mathrm{EQ}(v, 0, 1, c_1, \dots, c_n) \\ \land Q_b(y_1, y_2, p, v, 0, 1) \land \mathrm{PAD}(y_1, y_2, p, v, 0, 1) \\ & \downarrow \\ Q(y_1, y_2, p, 0, 1) \end{pmatrix}.$$

The successor relation for path nodes is represented by the binary symbol S, and the certificate of vertical verification for pairs (p_1, p_2) of successive path nodes is represented by the 7-ary symbol Q_v . For every $(\alpha, \beta) \in P_v^{\mathfrak{T}} \cap (P_\ell^{\mathfrak{T}})^2$, we include in Φ_2 a Horn sentence verifying the vertical consistency of the leftmost positions in the rows p_1 and p_2 and deriving the first Q_v -atom containing p_1 and p_2 , but only if there is a Q-atom containing p_1 :

$$\forall y_1, y_2, p_1, p_2, v, 0, 1, c_1, \dots, c_{2n} \\ \begin{pmatrix} S(p_1, p_2) \land \mathrm{EQ}(v, 0, 1, c_1, \dots, c_n) \land \bigwedge_{j \in [n]} c_j = 0 \land \mathrm{EQ}(v, 0, 1, c_{n+1}, \dots, c_{2n}) \\ \land \bigwedge_{j \in [n]} c_{j+n} = 0 \land T_{\alpha}(c_1, \dots, c_n, p_1) \land T_{\beta}(c_{n+1}, \dots, c_{2n}, p_2) \\ \land Q(y_1, y_2, p_1, 0, 1) \land \mathrm{PAD}(y_1, y_2, p_1, p_2, v, 0, 1) \\ & \downarrow \\ Q_v(y_1, y_2, p_1, p_2, v, 0, 1) \end{pmatrix} .$$

Next, for every $(\alpha, \beta) \in P_v^{\mathfrak{T}}$, we include in Φ_2 a Horn sentence verifying the vertical consistency for the intermediate positions by deriving further Q_v -atoms containing p_1 and p_2 :

 $\forall y_1, y_2, p_1, p_2, v_1, v_2, 0, 1, c_1, \dots, c_{2n}$

$$\left(\underbrace{\begin{array}{c} \text{SUCC}(v_1, v_2) \land \text{EQ}(v_1, 0, 1, c_1, \dots, c_n) \land \text{EQ}(v_1, 0, 1, c_{n+1}, \dots, c_{2n}) \\ \land T_{\alpha}(c_1, \dots, c_n, p_1) \land T_{\beta}(c_{n+1}, \dots, c_{2n}, p_2) \land Q_v(y_1, y_2, p_1, p_2, v_1, 0, 1) \\ \land \text{PAD}(y_1, y_2, p_1, p_2, v_1, v_2, 0, 1) \\ & \downarrow \\ Q_v(y_1, y_2, p_1, p_2, v_2, 0, 1) \end{array}\right)}.$$

And finally, for every $(\alpha, \beta) \in P_v^{\mathfrak{T}} \cap (P_r^{\mathfrak{T}})^2$, we include in Φ_2 a Horn sentence verifying the vertical consistency of the leftmost positions in the rows p_1 and p_2 and deriving the first

Q-atom containing p_2 only:

 $\forall y_1, y_2, p_1, p_2, v, 0, 1, c_1, \dots, c_{2n}$

$$\begin{pmatrix} EQ(v, 0, 1, c_1, \dots, c_n) \land \bigwedge_{j \in [n]} c_j = 1 \land EQ(v, 0, 1, c_{n+1}, \dots, c_{2n}) \\ \land \bigwedge_{j \in [n]} c_{j+n} = 1 \land T_{\alpha}(c_1, \dots, c_n, p_1) \land T_{\beta}(c_{n+1}, \dots, c_{2n}, p_2) \\ \land Q_v(y_1, y_2, p_1, p_2, v, 0, 1) \land PAD(y_1, y_2, p_1, p_2, v, 0, 1) \\ \downarrow \\ Q(y_1, y_2, p_2, 0, 1) \end{pmatrix}$$

The top row is verified using a 6-ary symbol Q_t similarly as the bottom row; however, the verification of the rightmost position in the top row results in the derivation of \perp . We include the following sentences as conjuncts in Φ_2 : for every $\alpha \in P_{\ell}^{\mathfrak{T}} \cap P_t^{\mathfrak{T}}$, the sentence

$$\forall y_1, y_2, p, v, 0, 1, c_1, \dots, c_n$$

$$\left(\underbrace{\begin{array}{c} \operatorname{EQ}(v,0,1,c_1,\ldots,c_n) \land \bigwedge_{j \in [n]} c_j = 0 \land T_{\alpha}(c_1,\ldots,c_n,p) \\ \land Q(y_1,y_2,p,0,1) \land \operatorname{PAD}(y_1,y_2,p,v,0,1) \\ & \downarrow \\ Q_t(y_1,y_2,p,v,0,1) \end{array}}_{Q_t(y_1,y_2,p,v,0,1)} \right),$$

for every $\alpha \in P_t^{\mathfrak{T}}$, the sentence

$$\forall y_1, y_2, p, v_1, v_2, 0, 1, c_1, \dots, c_n \\ \left(\underbrace{\begin{array}{c} \text{SUCC}(v_1, v_2) \land \text{EQ}(v_1, 0, 1, c_1, \dots, c_n) \land T_{\alpha}(c_1, \dots, c_n, p) \\ \land Q_t(y_1, y_2, p, v_1, 0, 1) \land Q(y_1, y_2, p, 0, 1) \land \text{PAD}(y_1, y_2, p, v_1, v_2, 0, 1) \end{array}}_{\substack{\downarrow \\ Q_t(y_1, y_2, p, v_2, 0, 1)}} \right),$$

and, for every $\alpha \in P_t^{\mathfrak{T}} \cap P_r^{\mathfrak{T}}$, the sentence

$$\forall y_1, y_2, p, v, 0, 1, c_1, \dots, c_n \\ \begin{pmatrix} EQ(v, 0, 1, c_1, \dots, c_n) \land \bigwedge_{j \in [n]} c_j = 1 \land T_\alpha(c_1, \dots, c_n, p) \\ \land Q_t(y_1, y_2, p, v, 0, 1) \land Q(y_1, y_2, p, 0, 1) \land PAD(y_1, y_2, p, v, 0, 1) \\ \downarrow \\ \downarrow \end{pmatrix} \right).$$
(21)

B.2 Inputs specified by sets of bounds

We include in τ a symbol I of arity $\lceil \log_2 |T| \cdot n \rceil + 1$. The first $\lceil \log_2 |T| \cdot n \rceil$ entries serve as binary counters to represent the pairs $(i, \alpha) \in [n] \times T$ in binary, and the last entry carries a path node representing a row of the tiling grid. Each pair $(i, \alpha) \in [n] \times T$ is to be interpreted as the fact that the *i*-th column in the row represented by a particular path node contains the tile α . The reason for this choice of encoding is that we aim to construct a universal sentence Φ which is equivalent to a set of forbidden substructures \mathcal{N} of size polynomial in $|T| \cdot n$. To achieve this, we use a constant number of symbols whose arity is logarithmic in the size of the input.

Suppose that 0 and 1 are two variables representing the bits 0 and 1, respectively. For each pair $(i, \alpha) \in [n] \times T$, the ternary formula TILE_{*i*, α}(0, 1, *p*) is the *I*-atom whose last entry contains the variable *p* and the first $\lceil \log_2 |T| \cdot n \rceil$ entries contain the variables 0 and 1 in the unique way that represents the number $i \cdot \alpha$ in binary when read from left to right. Note that the number of such formulas is polynomial in the size of the input to the tiling problem. In contrast to the proof of Theorem 5, it is not necessary to introduce any verifier nodes as the number of columns in the tiling grid is polynomial in the size of the input. The sentence $\Phi := \Phi_1 \wedge \Phi_2$ is defined similarly as in the proof of Theorem 5, so we only provide a general overview and highlight the main differences.

We start with Φ_1 . As before, we include the following sentence as a disjunct in Φ_1 :

$$\forall x, y \big(N(x, y) \land N(y, x) \Rightarrow \bot \big).$$

We want each row to be horizontally consistent. For all $i \in [n-1]$ and $(\alpha, \beta) \in T^2 \setminus P_h^{\mathfrak{T}}$, we include the following sentence as a conjunct in Φ_1 :

$$\forall p, \mathbf{0}, \mathbf{1} (N(\mathbf{0}, \mathbf{1}) \land \mathrm{TILE}_{i,\alpha}(\mathbf{0}, \mathbf{1}, p) \land \mathrm{TILE}_{i+1,\beta}(\mathbf{0}, \mathbf{1}, p) \Rightarrow \bot).$$

We also want each position in a given row to be occupied by at most one tile. For all $i \in [n], \alpha, \beta \in T$, we include the following sentence as a conjunct in Φ_1 :

$$\forall p, 0, 1 (\text{TILE}_{i,\alpha}(0, 1, p) \land \text{TILE}_{i,\beta}(0, 1, p) \Rightarrow \bot)$$

As in the proof of Theorem 6, we do *not* need to include an explicit condition stating that each row must be completely tiled from left to right.

We continue with Φ_2 . For the purpose of verifying the validity of a tiling, we include in τ a 5-ary symbol Q, a ($\lceil \log_2 |T| \cdot n \rceil + 4$)-ary symbol Q_v , and two ($\lceil \log_2 |T| \cdot n \rceil + 3$)-ary symbols Q_b, Q_t . For each pair $(i, \alpha) \in [n] \times T$, the formulas

BOT<sub>*i*,
$$\alpha$$</sub>(0, 1, *p*, *y*₁, *y*₂), TOP_{*i*, α} (0, 1, *p*, *y*₁, *y*₂), and VERT_{*i*, α} (0, 1, *p*₁, *p*₂, *y*₁, *y*₂)

are defined analogously to $\text{TILE}_{i,\alpha}(0, 1, p)$ but using the Q_b, Q_t and Q_v -atoms instead. We use them to verify the bottom row, top row, and the vertical consistency of a given tiling. In contrast to $\text{TILE}_{i,\alpha}$, the parameter α is not important in $\text{BOT}_{i,\alpha}$, $\text{TOP}_{i,\alpha}$, and $\text{VERT}_{i,\alpha}$. The formula $\text{PAD}(y_1, y_2, x_1, \ldots, x_k)$ is defined as in the previous subsection.

The parsing of a tiling starts from a path node p representing a row whose leftmost position contains a tile that can be present in the bottom left corner of a tiling grid. We include the following sentences as conjuncts in Φ_2 . For all $\alpha \in P_{\ell}^{\mathfrak{T}} \cap P_b^{\mathfrak{T}}$, $\gamma \in T$:

$$\forall y_1, y_2, p, 0, 1$$

for all $i \in [n-1], \alpha, \beta \in P_b^{\mathfrak{T}}, \gamma \in T$:

 $\forall y_1, y_2, p, 0, 1$

$$\left(\begin{array}{c} \operatorname{TILE}_{i,\alpha}(0,1,p) \wedge \operatorname{TILE}_{i+1,\beta}(0,1,p) \\ \wedge \operatorname{BOT}_{i,\gamma}(0,1,p,y_1,y_2) \wedge \operatorname{PAD}(y_1,y_2,0,1,p) \\ & \\ \downarrow \\ & \\ \operatorname{BOT}_{i+1,\gamma}(0,1,p,y_1,y_2) \end{array}\right)$$

and, for all $\alpha \in P_r^{\mathfrak{T}} \cap P_b^{\mathfrak{T}}$, $\gamma \in T$:

$$\forall y_1, y_2, p, 0, 1$$

$$\left(\underbrace{\text{TILE}_{n,\alpha}(0, 1, p) \land \text{BOT}_{n,\gamma}(0, 1, p, y_1, y_2) \land \text{PAD}(y_1, y_2, 0, 1, p)}_{\Downarrow} \right)$$

$$\downarrow$$

$$Q(0, 1, p, y_1, y_2)$$

The successor relation for path nodes is represented by the binary symbol S. We include the following sentences as conjuncts in Φ_2 . For all $(\alpha, \beta) \in P_v^{\mathfrak{T}} \cap (P_{\ell}^{\mathfrak{T}})^2$, $\gamma \in T$:

$$\begin{array}{c} \forall y_1, y_2, p_1, p_2, 0, 1 \\ \\ \left(\underbrace{\begin{array}{c} S(p_1, p_2) \land \text{TILE}_{1,\alpha}(0, 1, p_1) \land \text{TILE}_{1,\beta}(0, 1, p_2) \\ \land Q(0, 1, p_1, y_1, y_2) \land \text{PAD}(y_1, y_2, 0, 1, p_1, p_2) \\ \end{array}}_{\downarrow \downarrow} \\ \\ \text{VERT}_{1,\gamma}(0, 1, p_1, p_2, y_1, y_2) \end{array} \right), \end{array}$$

for every $i \in [n-1]$, $(\alpha, \beta) \in P_v^{\mathfrak{T}}$, $\gamma \in T$:

 $\forall y_1, y_2, p_1, p_2, \mathbf{0}, \mathbf{1}$

$$\left(\begin{array}{c} \text{TILE}_{i,\alpha}(\mathbf{0},\mathbf{1},p_1) \wedge \text{TILE}_{i,\beta}(\mathbf{0},\mathbf{1},p_2) \\ \wedge \text{VERT}_{i,\gamma}(\mathbf{0},\mathbf{1},p_1,p_2,y_1,y_2) \wedge \text{PAD}(y_1,y_2,\mathbf{0},\mathbf{1},p_1,p_2) \\ & \\ \downarrow \\ \text{VERT}_{i+1,\gamma}(\mathbf{0},\mathbf{1},p_1,p_2,y_1,y_2) \end{array}\right),$$

and, for all $(\alpha, \beta) \in P_v^{\mathfrak{T}} \cap (P_r^{\mathfrak{T}})^2, \ \gamma \in T$:

$$\begin{array}{c} \forall y_1, y_2, p_1, p_2, 0, 1 \\ \\ \begin{pmatrix} & \text{TILE}_{n,\alpha}(0, 1, p_1) \land \text{TILE}_{n,\beta}(0, 1, p_2) \\ & \land \text{VERT}_{n,\gamma}(0, 1, p_1, p_2, y_1, y_2) \land \text{PAD}(y_1, y_2, 0, 1, p_1, p_2) \\ & \downarrow \\ & \downarrow \\ & Q(0, 1, p_2, y_1, y_2) \end{array} \right)$$

The top row is verified in a similar fashion as the bottom row, except that the nullary falsity predicate is derived at the end of the verification process. We include the following sentences as conjuncts in Φ_2 . For all $\alpha \in P_{\ell}^{\mathfrak{T}} \cap P_t^{\mathfrak{T}}$, $\gamma \in T$:

$$\begin{array}{c} \forall y_1, y_2, p, \mathbf{0}, \mathbf{1} \\ \\ \begin{pmatrix} \underbrace{\mathrm{TILE}_{1,\alpha}(\mathbf{0}, \mathbf{1}, p) \land Q(\mathbf{0}, \mathbf{1}, p, y_1, y_2) \land \mathrm{PAD}(y_1, y_2, \mathbf{0}, \mathbf{1}, p) \\ \\ & \underbrace{ \downarrow} \\ \\ \mathrm{TOP}_{1,\gamma}(\mathbf{0}, \mathbf{1}, p, y_1, y_2) \end{array} \right), \end{array}$$

for every $i \in [n-1], \, \alpha, \beta \in P_t^{\mathfrak{T}}, \, \gamma \in T$:

$$\begin{array}{c} \forall y_1, y_2, p, \mathbf{0}, \mathbf{1} \\ \\ \begin{pmatrix} & \text{TILE}_{i,\alpha}(\mathbf{0}, \mathbf{1}, p) \land \text{TILE}_{i+1,\beta}(\mathbf{0}, \mathbf{1}, p) \\ & \land \text{TOP}_{i,\gamma}(\mathbf{0}, \mathbf{1}, p, y_1, y_2) \land \text{PAD}(y_1, y_2, \mathbf{0}, \mathbf{1}, p) \\ & & \downarrow \\ & \text{TOP}_{i+1,\gamma}(\mathbf{0}, \mathbf{1}, p, y_1, y_2) \end{pmatrix}, \end{array} \right),$$

and, for all $\alpha \in P_r^{\mathfrak{T}} \cap P_t^{\mathfrak{T}}$, $\gamma \in T$:

 $\forall y_1, y_2, p, 0, 1$

$$\left(\begin{array}{c} \underbrace{\mathrm{TILE}_{n,\alpha}(\mathbf{0},\mathbf{1},p)\wedge\mathrm{TOP}_{n,\gamma}(\mathbf{0},\mathbf{1},p,y_1,y_2)\wedge\mathrm{PAD}(y_1,y_2,\mathbf{0},\mathbf{1},p)}_{\downarrow\downarrow} \\ \downarrow \\ \end{array}\right).$$