Mahalanobis balancing: a multivariate perspective on approximate covariate balancing

Yimin Dai

School of Mathematics, Sun Yat-sen University, Guangzhou, China, 510275 and

Ying Yan

School of Mathematics, Sun Yat-sen University, Guangzhou, China, 510275

Abstract

In the past decade, various exact balancing-based weighting methods were introduced to the causal inference literature. It eliminates covariate imbalance by imposing balancing constraints in a certain optimization problem, which can nevertheless be infeasible when there is bad overlap between the covariate distributions in the treated and control groups or when the covariates are high-dimensional. Recently, approximate balancing was proposed as an alternative balancing framework. It resolves the feasibility issue by using inequality moment constraints instead. However, it can be difficult to select the threshold parameters. Moreover, moment constraints may not fully capture the discrepancy of covariate distributions. In this paper, we propose Mahalanobis balancing to approximately balance covariate distributions from a multivariate perspective. We use a quadratic constraint to control overall imbalance with a single threshold parameter, which can be tuned by a simple selection procedure. We show that the dual problem of Mahalanobis balancing is an ℓ_2 norm-based regularized regression problem, and establish interesting connection to propensity score models. We derive asymptotic properties, discuss the high-dimensional scenario, and make extensive numerical comparisons with existing balancing methods.

Keywords: causal inference, Mahalanobis distance, multivariate imbalance, overlap, propensity score

1 Introduction

Inference about causation gains increasing attention in medical science, economics, computer science, and many other disciplines. Propensity score (Rosenbaum & Rubin, 1983), the probability of receiving a treatment conditional on the covariates, plays a central role in causal inference for observational studies. There are several classes of propensity score-based methods in practitioners' toolkit, including matching, weighting, and subclassification. We focus on weighting in this article (Rosenbaum, 1987; Robins et al., 1994; Hirano et al., 2003). Inverse probability weighting and its doubly robust version are perhaps the most commonly used weighting methods. They explicitly estimate the propensity score via a parametric or nonparametric model (e.g. Hirano et al., 2003). However, as demonstrated by Kang & Schafer (2007) and related articles, inverse probability weighting is prone to misspecification of the propensity score model, and covariate balance may not be attained after reweighting.

In the past decade, numerous robust weighting methods have been proposed, which aim to directly balance covariates in the estimation procedure. Hainmueller (2012) introduced the entropy balancing method, which optimizes the entropy loss function subject to a set of balancing constraints. The balancing constraints enforce the weighted moments in the control group to be equal to the unweighted counterparts in the treated group, and thus finite-sample exact balance is achieved. The balanced weights are then used to reweight the subjects in the control group to produce a weighted estimator of the average treatment effect on the treated. This method has close connection to survey sampling, missing data, and machine learning (e.g. Dehejia & Wahba, 1999; Graham et al., 2012; Mohri et al., 2018). Zhao & Percival (2017) revealed that entropy balancing is doubly robust. Recently, entropy balancing has received great attention in applied areas. For example, it was adapted to integrate randomized clinical trial and observational study (Lee et al., 2023), to transport trial results to a target population (Josey et al., 2021a), and so forth. Imai & Ratkovic (2014) considered parametric propensity score model and solved the likelihood score function together with the balancing constraints simultaneously to produce covariate balancing propensity score. Fan et al. (2022) improved the covariate balancing propensity score method. Chan et al. (2016) constructed a class of calibration

weighting estimators by minimizing a distance measure subject to balancing constraints. Empirical likelihood and exponential tilting belong to this class. Yiu & Su (2018) proposed to eliminate the association between treatment and covariates in the weight construction procedure. Hazlett (2020) recommended to construct balancing constraints from the kernel viewpoint. Josey et al. (2021b) proposed a Bregman distance framework which unifies many existing methods.

A common feature of the aforementioned balancing methods is that the balancing conditions are directly imposed as equality moment constraints in a convex optimization problem. We call them *exact balancing* methods in this paper. Compared to inverse probability weighting, exact balancing is capable of producing stable weights and is more robust to model misspecification in many circumstances. Nevertheless, exact balancing performs poorly in the bad-overlap scenario where there is limited overlap between the covariate distributions in the treated and control groups, or in the high-dimensional scenario where the number of constraints is large. Even worse, the optimization problem may be infeasible in these difficult scenarios. We refer readers to a discussion of convex hull problems in Hayakawa et al. (2023), which implies infeasibility of exact balancing in the high-dimensional scenario.

To alleviate the infeasibility problem, Zubizarreta (2015) and Wang & Zubizarreta (2020) proposed an *approximate balancing* framework, i.e., the minimal dispersion approximately balancing weights (MDABW) method, which replaces the exact balancing conditions with less restrictive inequality constraints. The inequality constraints are obtained by controlling univariate dispersion of each covariate. Wang & Zubizarreta (2020) showed that the dual problem of MDABW is a weighted ℓ_1 norm-based regularized regression problem. There are several other robust balancing methods which may not suffer the infeasibility problem. For example, Wong & Chan (2018) directly minimized covariate functional imbalance. Zhao (2019) proposed a general balancing framework using tailored loss functions. Li et al. (2018) and Ma & Wang (2020) provided insights about robust weighting from different perspectives.

In general, covariate balancing becomes much more difficult in the high-dimensional scenario, because the overlap condition tends to be more restrictive as the dimensionality increases. We refer the readers to D'Amour et al. (2021) for a formal discussion of the overlap condition in the high-dimensional scenario. Regularization plays a key role in high-dimensional analysis, and it has been adapted to covariate balancing recently. Among others, Athey et al. (2018); Ning et al. (2020); Tan (2020a,b) addressed high-dimensional balancing problems using regularization techniques. We call them high-dimensional regularized balancing methods. Many of these methods involved outcome models and thus cannot be categorized as data preprocessing procedures. Interestingly, Athey et al. (2018); Ning et al. (2020); Tan (2020b) established tight connections of their methods to the MD-ABW method.

Now we discuss potential limitations of the MDABW method as an approximate balancing framework. The MDABW method controls the weighted absolute standardized mean difference (ASMD) for each covariate with a threshold parameter, where the ASMD is a univariate imbalance measure frequently used in observational studies. Approximate balance is achieved for each covariate separately by controlling the threshold parameters, and thus the MDABW method is a *univariate approximate balancing* framework. However, it can be difficult to select the threshold parameters simultaneously. For example, an attempt to reduce imbalance of some covariate by shrinking the corresponding threshold parameter may reversely enlarge imbalance of other covariates. This dilemma is not uncommon in the bad-overlap scenario. Second, even if univariate approximate balancing is achieved for all constraints separately, the overall imbalance is not necessarily under control. Moreover, the MDABW method requires that the number of constraints is much smaller than the sample size. Our numerical studies demonstrate that the MDABW method may perform unsatisfactorily in the bad-overlap and high-dimensional scenarios.

The limitation of univariate approximate balancing motivates the proposed multivariate approximate balancing method in this paper. We show that univariate approximate balancing can be greatly improved by monitoring and controlling overall covariate balance from the multivariate perspective. The concept of multivariate imbalance measure is widely adopted in the matching literature to quantify the discrepancy of the covariate distributions (e.g. Iacus et al., 2011, 2012; Diamond & Sekhon, 2013; Imbens & Rubin, 2015). However, it is largely neglected in the covariate balancing literature. In this paper, we highlight the importance of controlling multivariate imbalance.

The main contribution of this paper is that we propose Mahalanobis balancing as a

multivariate approximate balancing framework. We use a generalized version of Mahalanobis distance to measure multivariate imbalance of the covariates between groups. A sufficiently small value of the multivariate imbalance measure suggests that the covariates are approximately balanced from the multivariate perspective. The multivariate balancing condition is encoded as a single quadratic inequality constraint in a convex optimization problem with a pre-specified loss function (e.g., entropy, empirical likelihood). We study the primal optimization problem by utilizing the Fenchel duality theory (e.g. Bertsekas, 2016; Mohri et al., 2018; Mordukhovich & Nam, 2022). We show that the dual problem is an unconstrained regularized regression problem with an ℓ_2 norm regularizer. Off-the-shelf convex optimization algorithms can be employed to solve the dual problem. We adopt the BFGS quasi-Newton algorithm in this paper.

Mahalanobis balancing differs from the MDABW method by that multivariate imbalance, rather than univariate imbalance, is controlled. This simple idea has important consequences that make Mahalanobis balancing a highly competitive balancing method. First, a univariate threshold parameter is attached to the quadratic inequality constraint in Mahalanobis balancing, and thus the issue of tuning multiple or high-dimensional threshold parameters is resolved. Second, there are situations where univariate approximate balancing is achieved but the covariate distributions remain imbalanced after reweighting (e.g., Scenarios B, C, E in Section 4). The MDABW method does not perform satisfactorily in these situations, and treatment effect estimation can be considerably biased. It suggests that controlling univariate imbalance in reweighting can be insufficient to delineate multivariate imbalance and obtain valid treatment effect estimation. In comparison, Mahalanobis balancing greatly improves the performance of univariate approximate balancing in these situations, which confirms the importance of controlling multivariate imbalance. Third, Mahalanobis balancing is not restricted to the low-dimensional situation. In the Supplementary Material, we propose a high-dimensional version of Mahalanobis balancing where the idea of multivariate imbalance is further exploited. The simulations reveal that it is competitive to the high-dimensional regularized balancing methods (Athey et al., 2018; Ning et al., 2020; Tan, 2020b). In comparison, the MDABW method is restricted to the low-dimensional scenario.

The rest of the paper is organized as follows. In Section 2, we introduce necessary

notations, existing imbalance measures, and balancing methods. In Section 3, we present the proposed Mahalanobis balancing method, and discuss the high-dimensional situation. Asymptotic properties of Mahalanobis balancing are studied. In Section 4, we make extensive comparison of Mahalanobis balancing to the existing balancing methods in numerical studies. We draw conclusion in the last section. Extended discussion of the highdimensional setting, proofs, and additional numerical results are available in the online Supplementary Material. An R package of Mahalanobis balancing and numerical examples are available at Github via the link: https://github.com/yimindai0521/ACBalancing/.

2 Notation and Preliminary

2.1 The framework

Consider a simple random sample of n subjects from a population. Let T be a binary treatment indicator with T = 1 if the subject is treated and T = 0 otherwise. Let $X = (X_1, \dots, X_p)^{\top}$ be a p-dimensional vector of pre-treatment covariates. Let Y be the univariate outcome. The observed data are $\{(X_i, T_i, Y_i) : i = 1, \dots, n\}$, which are nindependent and identically distributed copies of the triplet (X, T, Y). Under the potential outcome framework (e.g. Imbens & Rubin, 2015), we define a pair of potential outcomes $\{Y(0), Y(1)\}$ for each subject if he were not treated or treated. The observed outcome is Y = (1 - T)Y(0) + TY(1).

We impose the following strong ignorability and overlap assumptions for identification and inference of the average treatment effect.

Assumption 1 (Strong Ignorability). $\{Y(0), Y(1)\} \perp T \mid X$.

Here, \perp denotes independence. Assumption 1 states that the potential outcomes $\{Y(0), Y(1)\}$ are independent of the treatment indicator T given pre-treatment covariates X, which implies that there are no unmeasured confounders. The probability $\pi(X) = \Pr(T = 1 \mid X)$ is the propensity score function. Assumption 1 implies $\pi(X)$ is a balancing score (Rosenbaum & Rubin, 1983) in the sense that $\{Y(0), Y(1)\} \perp T \mid \pi(X)$. Under Assumption 1, the average treatment effect is identifiable.

Moreover, to ensure that there are informative observations for treatment effect estimation, we impose the overlap assumption:

Assumption 2 (Overlap). 0 < Pr(T = 1 | X) < 1 for any X in its support.

In this paper, the causal estimand of interest is the average treatment effect (ATE), defined by $\tau = E\{Y(1) - Y(0)\} = \mu_1 - \mu_0$, where $\mu_j = E\{Y(j)\}, j = 0, 1$. The proposed method can be adapted to other causal estimands, e.g., average treatment effect on the treated (ATT). Under Assumption 1, $\tau = E\{\mu_1(X) - \mu_0(X)\}$, where $\mu_j(X) = E\{Y(j) \mid X\}$, j = 0, 1.

A generic reweighting scheme constructs weights $w = (w_1, \dots, w_n)^{\top}$ by exploiting the sampled data and possibly external information from the population, and then estimates the ATE by $\hat{\tau} = \sum_{i=1}^{n} T_i w_i Y_i - \sum_{i=1}^{n} (1 - T_i) w_i Y_i$. The covariate balancing methods introduced in Section 1 explicitly balance the covariates in the reweighting procedure and produce a set of balanced weights.

2.2 Imbalance measures

A set of pre-specified basis functions, denoted by $\phi_1(X), \dots, \phi_K(X)$, can be used instead of the original covariates X in the construction of balanced weights (e.g., Hainmueller, 2012; Imai & Ratkovic, 2014; Wang & Zubizarreta, 2020). In Section 3, we discuss the choice of the basis functions. Define $\Phi(X) = (\phi_1(X), \dots, \phi_K(X))^{\top}$ to be a vector of basis functions, which is an $\mathbb{R}^p \to \mathbb{R}^K$ feature mapping.

Assessment of covariate balance is crucial in the inference of causal effects. Many imbalance measures are proposed and widely used in the literature. The absolute standardized mean difference (ASMD) is the most popular univariate imbalance measure (e.g., Imbens & Rubin, 2015). The ASMD for the kth basis function $\phi_k(X)$ is expressed as

$$ASMD_{k} = \frac{\left|\phi_{1,k} - \phi_{0,k}\right|}{\sqrt{\left(s_{1,k}^{2} + s_{0,k}^{2}\right)/2}}, \quad k = 1, \cdots, K,$$
(1)

where $\bar{\phi}_{j,k} = \sum_{i=1}^{n} I(T_i = j)\phi_k(X_i)/n_j$ and $s_{j,k}^2 = \sum_{i=1}^{n} I(T_i = j)(\phi_k(X_i) - \bar{\phi}_{j,k})^2/(n_j - 1)$ are the treatment-specific sample mean and sample variance for ϕ_k , j = 0, 1. Here, $I(\cdot)$ is the indicator function, and n_1 and n_0 are the sample sizes of the treated and control groups, respectively. When the ASMD for a basis function does not exceed a pre-specified threshold parameter δ , then this basis function is considered to be approximately balanced. Common choices of δ are 0.1, 0.2 and 0.25 (e.g. Stuart et al., 2013; Imbens & Rubin, 2015; Xie et al., 2019). However, even when all basis functions are approximately balanced in the sense that the average of ASMD₁, \cdots , ASMD_K is smaller than the threshold, say $\delta = 0.2$, the difference in the outcomes as an unadjusted average treatment effect estimation can be still severely biased. This can be explained by that the ASMD is a univariate imbalance measure which does not fully characterize multivariate imbalance.

The targeted absolute standardized mean difference (TASMD), introduced by Chattopadhyay et al. (2020), is defined as

$$\mathrm{TASMD}_{j,k} = \frac{\left|\bar{\phi}_{j,k} - \bar{\phi}_k\right|}{s_k},$$

where $\bar{\phi}_k = \sum_{i=1}^n \phi_k(X_i)/n$ and $s_k^2 = \sum_{i=1}^n (\phi_k(X_i) - \bar{\phi}_k)^2/(n-1)$, $j = 0, 1; k = 1, \dots, K$. The ASMD quantifies the disparity between the treated and control groups, whereas the TASMD measure quantifies the disparity of each group in relation to a target population. In practical applications, we can employ the treatment-adjusted subgroup mean difference method to evaluate balance in relation to various target populations. Examples include the population of treated units when estimating the ATT, the entire population when estimating the ATE, or a specific population defined by its observed covariates when estimating the conditional average treatment effect (CATE).

Multivariate imbalance measure can be used to assess the imbalance for all covariates or basis functions simultaneously. The squared Mahalanobis distance (MD) (Imbens & Rubin, 2015) is a popular multivariate imbalance measure, given by

$$MD = (\bar{\Phi}_1 - \bar{\Phi}_0)^\top \left(\frac{\hat{\Sigma}_1 + \hat{\Sigma}_0}{2}\right)^{-1} (\bar{\Phi}_1 - \bar{\Phi}_0),$$

where $\bar{\Phi}_j = (\bar{\phi}_{j,1}, \dots, \bar{\phi}_{j,K})^{\top}$, j = 0, 1, and $\hat{\Sigma}_1$ and $\hat{\Sigma}_0$ are the sample covariance matrices of $\Phi(X)$ in the treated and control groups, respectively. The MD was used in Mahalanobis distance matching (e.g. Imbens & Rubin, 2015) to determine the closeness of subjects between the treated and control groups. Moreover, it was adapted to genetic matching (Diamond & Sekhon, 2013), which aims at minimizing a generalized version of

MD to optimize postmatching covariate balance. The ℓ_1 distance-based measure proposed by Iacus et al. (2011) is another example of multivariate imbalance measure, which motivates a class of matching methods with the monotonic imbalance bounding property. Iacus et al. (2012) derived coarsened exact matching from this class, and emphasized the importance of optimizing multivariate balance in matching. Zhu et al. (2018) proposed to use the Kernel distance as a multivariate imbalance measure. Huling & Mak (2020) studied energy distance-based covariate balancing.

2.3 Exact balancing v.s. approximate balancing

The existing covariate balancing methods achieve exact balance by imposing a set of equality balancing constraints

$$\sum_{i=1}^{n} w_i T_i \phi_k(X_i) = \bar{\phi}_k,$$
and
$$\sum_{i=1}^{n} w_i (1 - T_i) \phi_k(X_i) = \bar{\phi}_k, \quad k = 1, \cdots, K,$$
(2)

in the construction of the weights w_1, \dots, w_n , where $\bar{\phi}_k = \sum_{i=1}^n \phi_k(X_i)/n$. These constraints force the first moment of the basis functions to be exactly balanced after reweighting. Examples of exact balancing methods include entropy balancing (Hainmueller, 2012), covariate balancing propensity score (Imai & Ratkovic, 2014), and calibration weighting (Chan et al., 2016), among others.

Define a weighted version of the ASMD by

$$\text{ASMD}_{k}^{w} = \frac{\left|\sum_{i=1}^{n} w_{i} T_{i} \phi_{k}(X_{i}) - \sum_{i=1}^{n} w_{i}(1 - T_{i}) \phi_{k}(X_{i})\right|}{\sqrt{\left(s_{1,k}^{2} + s_{0,k}^{2}\right)/2}}, \quad k = 1, \cdots, K$$

The ASMD^w_k is a weighted univariate imbalance measure that quantifies remaining imbalance for the kth basis function ϕ_k after reweighting. It is used in this paper to compare the univariate balancing performance of weighting methods. Usually, we normalize the weights for each group such that $\sum_{i=1}^{n} T_i w_i = 1$ and $\sum_{i=1}^{n} (1 - T_i) w_i = 1$ before comparison. It is straightforward that the equality balancing constraints (2) lead to that ASMD^w_k = 0 for all $k = 1, \dots, K$. Therefore, finite-sample univariate exact balance is achieved by the exact balancing methods. In comparison, inverse probability weighting does not possess this attractive property.

When exact balancing is not attainable in the bad-overlap situation, the MDABW method (Zubizarreta, 2015; Wang & Zubizarreta, 2020) can be used instead. The balanced weights for the treated group $\{i : T_i = 1\}$ are the global minimum of the following optimization problem:

$$\begin{array}{ll}
\text{minimize} & \sum_{i=1}^{n} T_{i} f\left(w_{i}\right) \\
\text{subject to} & \left|\sum_{i=1}^{n} w_{i} T_{i} \phi_{k}(X_{i}) - \bar{\phi}_{k}\right| \leq \delta_{k}, k = 1, \cdots, K,
\end{array} \tag{3}$$

where $f(\cdot)$ is a pre-specified loss function, and $\delta_1, \dots, \delta_K \ge 0$ are the threshold parameters. Because the constraints are imposed separately for the basis functions, MDABW is a univariate approximate balancing method. The choices of $f(\cdot)$ is discussed in Section 3. Similarly, the weights for the control group $\{i : T_i = 0\}$ can be obtained by replacing T_i with $1 - T_i$ in the optimization problem (3).

If we normalize the basis function $\phi_k(X_i)$ to be $\phi_k(X_i)/\sqrt{(s_{1,k}^2 + s_{0,k}^2)/2}$ in the optimization problem (3), then $\text{ASMD}_k^w \leq 2\delta_k$ for $k = 1, \dots, K$. A small value of δ_k , say 0.1, indicates that the ϕ_k is approximately balanced. However, it is difficult to define optimality for all threshold parameters $(\delta_1, \dots, \delta_K)$ simultaneously in the bad-overlap setting, because decreasing univariate imbalance for some basis function may inevitably increase imbalance for other basis functions. Correlation of the basis functions are not taken into account in Problem (3). Tuning a large number of parameters is very time-consuming, and there is lack of guideline to tune these parameters simultaneously. These potential limitations motivate Mahalanobis balancing in the following section.

We define the following Mahalanobis imbalance measure (MIM) as a weighted version of the squared Mahalanobis distance:

$$MIM^{w} = \left\{ \sum_{i=1}^{n} w_{i} T_{i} \Phi(X_{i}) - \sum_{i=1}^{n} w_{i} (1 - T_{i}) \Phi(X_{i}) \right\}^{\top} \left(\frac{\hat{\Sigma}_{1} + \hat{\Sigma}_{0}}{2} \right)^{-1} \\ \times \left\{ \sum_{i=1}^{n} w_{i} T_{i} \Phi(X_{i}) - \sum_{i=1}^{n} w_{i} (1 - T_{i}) \Phi(X_{i}) \right\},$$
(4)

which can be used to measure the remaining overall imbalance after reweighting. Note that

 $ASMD_k^w = 0$ for all $k = 1, \dots, K$ is equivalent to $MIM^w = 0$. However, univariate approximate balance does not imply multivariate approximate balance, because it is possible that the $ASMD_k^w$ is small for all $k = 1, \dots, K$ but the MIM^w is large.

3 Proposed Methodology

3.1 Mahalanobis balancing

To alleviate the limitation of univariate approximate balancing, we propose the Mahalanobis balancing (MB) method as a multivariate approximate balancing framework, which directly controls multivariate imbalance to produce balanced weights. Specifically, Mahalanobis balancing obtains the balanced weights for the treated group by solving the convex optimization problem:

$$\begin{array}{ll}
\underset{w \in \mathbb{R}^{n}}{\operatorname{minimize}} : & \sum_{i=1}^{n} T_{i}f(w_{i}) \\
\text{subject to:} & \begin{cases} w_{i} \geq 0, & i \in \{j : T_{j} = 1\}; \\ \sum_{i=1}^{n} T_{i}w_{i}\{\Phi(X_{i}) - \bar{\Phi}\}^{\top}W\sum_{i=1}^{n} T_{i}w_{i}\{\Phi(X_{i}) - \bar{\Phi}\} \leq \delta, \end{cases}$$
(5)

where $f(\cdot)$ is a convex loss function, $\Phi(X)$ is a vector of basis functions defined in Section 2.2, $\bar{\Phi} = \sum_{i=1}^{n} \Phi(X_i)/n$, W is a user-specified $K \times K$ positive-definite weight matrix, and $\delta \geq 0$ is a univariate threshold parameter. The MB weights are obtained after normalization for the treated subjects, i.e., $w_i^{MB} = w_i / \sum_{j=1}^{n} T_j w_j$. Similarly, the MB weights for the control group can be obtained.

The role of each w_i is to down-weigh or up-weigh the deviation of $\Phi(X_i)$ from the pooledsample mean $\overline{\Phi}$. The matrix W weighs the relative importance of the basis functions $\phi_1(\cdot), \cdots, \phi_K(\cdot)$. In this paper, we consider two choices of W: (i) the diagonal matrix $W_1 = [\operatorname{diag}\{(\hat{\Sigma}_1 + \hat{\Sigma}_0)/2\}]^{-1}$, where its main diagonal contains the reciprocals of the diagonal elements of $(\hat{\Sigma}_1 + \hat{\Sigma}_0)/2$; (ii) $W_2 = \{(\hat{\Sigma}_1 + \hat{\Sigma}_0)/2\}^{-1}$. We use W_2 when the dimension Kis not large and the basis functions are not highly correlated. Otherwise, we use W_1 . We choose W in the data-preprocessing step to normalize the data, and thus treat it as fixed in the theoretical development.

For an arbitrary set of normalized weights for the treated subjects, we define the gen-

eralized Mahalanobis imbalance measure (GMIM) for the treated group:

$$GMIM_{1}^{w} = \left\{ \sum_{i=1}^{n} T_{i}w_{i}\Phi(X_{i}) - \bar{\Phi} \right\}^{\top} W \left\{ \sum_{i=1}^{n} T_{i}w_{i}\Phi(X_{i}) - \bar{\Phi} \right\}.$$
 (6)

Similarly, we define GMIM_0^w for the control group. The GMIM_1^w measures the residual multivariate difference between the basis functions in the treated group and the sample average $\overline{\Phi}$ after reweighting. Note that the term $\sum_{i=1}^n T_i w_i \{\Phi(X_i) - \overline{\Phi}\}^\top W \sum_{i=1}^n T_i w_i \{\Phi(X_i) - \overline{\Phi}\}$ in Problem (5) is the unnormalized version of the GMIM_1^w . Therefore, Problem (5) explicitly restricts the unnormalized residual multivariate imbalance by a single threshold parameter δ . When Problem (5) is feasible for $\delta = 0$, then MB reduces to exact balancing. When Problem (5) is infeasible for $\delta = 0$, we need to tune a positive value for δ . In Section 3.4, we propose to optimize post-weighting multivariate balance by monitoring GMIM_1^w in the selection procedure.

We remark that we prefer to minimize GMIM_1^w and GMIM_0^w separately rather than minimize the Mahalanobis imbalance measure MIM^w in equation (4). First, it allows us to easily obtain balanced weights from two separate optimization problems. Second, by minimizing GMIM_1^w and GMIM_0^w , the Mahalanobis imbalance measure MIM^w is under control. More importantly, compared to the MIM^w , the GMIM_1^w and GMIM_0^w are more relevant to the post-weighting multivariate balancing performance. In particular, even if the MIM^w is very small, it does not imply that the weighted basis functions $\sum_{i=1}^n T_i w_i \Phi(X_i)$ and $\sum_{i=1}^n (1 - T_i) w_i \Phi(X_i)$ are close to the sample average $\overline{\Phi}$. That is, it is possible that the weighted distributions in the two groups are close to each other, but meanwhile they are different from the underlying distribution in the target population so that the GMIM_1^w and GMIM_0^w are large. This phenomenon frequently occurs in the bad-overlap and highdimensional settings.

Next, we discuss the choice of the loss function $f(\cdot)$. One may use the entropy function $f(x) = x \log(x)$ (Hainmueller, 2012), the negative of empirical likelihood $f(x) = -\log(x)$, the quadratic function by Zubizarreta (2015), the distance measure by Chan et al. (2016), and the Bregman distance by Josey et al. (2021b), among others. We prefer the entropy function $f(x) = x \log(x)$ for its stable performance and theoretical properties.

The choice of the basis functions is important for the consistency of the balancing

methods (e.g. Zhao & Percival, 2017). In practise, we consider the first and second moments of X as the basis functions, and sometimes the interaction terms are included. Motivated by the theory of Reproducing Kernel Hilbert Space (RKHS) (Wong & Chan, 2018; Zhao, 2019; Hazlett, 2020), we also consider the second choice $\Phi(X) = (K(X, X_1), \dots, K(X, X_n))^{\top}$ when the dimension of the covariates p is large compared to the sample size n. Here, $K(\cdot, \cdot)$ is a pre-specified kernel function (e.g., Mohri et al., 2018), and the number of basis functions is K = n.

In the following, we study the dual of the primal optimization problem (5). By the Fenchel duality theory (Bertsekas, 2016; Mohri et al., 2018; Mordukhovich & Nam, 2022), we show that the dual problem is a regularized propensity score model with an ℓ_2 norm penalty. In Theorem 1, we formally establish the connection between Mahalanobis balancing and ℓ_2 shrinkage estimation of the propensity score model.

We define some additional notations. Apply Cholesky decomposition to the weight matrix W and write $W = (W^{1/2})^{\top}W^{1/2}$. Let $\|\theta\|_2 = \sqrt{\theta_1^2 + \cdots + \theta_K^2}$ be the ℓ_2 norm for an arbitrary K-dimensional vector $\theta = (\theta_1, \cdots, \theta_K)^{\top}$. The quadratic inequality constraint in Problem (5) can then be rewritten as $\|\sum_{i=1}^n T_i w_i W^{1/2}(\Phi(X_i) - \overline{\Phi})\|_2 \leq \sqrt{\delta}$. Let $\tilde{f}(p) =$ f(p) if $p \geq 0$, and $\tilde{f}(p) = +\infty$, otherwise. Let $\partial g(x)$ be the subgradient of the function $g(\cdot)$ at x in the domain of the function (Mohri et al., 2018). Define a convex subset $\mathcal{C} = \{u \in \mathbb{R}^K : \|u\|_2 \leq \sqrt{\delta}\}$. Define $I_{\mathcal{C}}(u) = 0$ if $u \in \mathcal{C}$, and $I_{\mathcal{C}}(u) = +\infty$ otherwise. Then, Problem (5) is equivalent to the following unconstrained primal optimization problem:

$$\underset{w \in \mathbb{R}^{n_1}}{\text{minimize}} : \sum_{i=1}^n T_i \tilde{f}(w_i) + I_{\mathcal{C}} \left(\sum_{i=1}^n T_i w_i W^{1/2}(\Phi(X_i) - \bar{\Phi}) \right).$$
(7)

Let $\tilde{f}^*(\cdot)$ be the conjugate function of $\tilde{f}(\cdot)$ (Bertsekas, 2016). The following theorem states the dual problem.

Theorem 1. The dual of Problem (7) is the following unconstrained optimization problem:

$$\underset{\theta \in \mathbb{R}^{K}}{\text{minimize}} : \sum_{i=1}^{n} T_{i} \tilde{f}^{*} \left(\theta^{\top} W^{1/2} (\Phi(X_{i}) - \bar{\Phi}) \right) + \sqrt{\delta} \|\theta\|_{2}.$$

$$(8)$$

Let $\hat{w} = \{\hat{w}_i : T_i = 1\}$ be the solution of the primal problem (7) and $\hat{\theta} = (\hat{\theta}_1, \cdots, \hat{\theta}_K)^{\top}$ be

the solution of the dual problem (8). The \hat{w}_i can be expressed as a function of $\hat{\theta}$:

$$\hat{w}_i = \partial(\tilde{f}^*) \left(\hat{\theta}^\top W^{1/2}(\Phi(X_i) - \bar{\Phi}) \right), \quad \text{for } i \in \{j : T_j = 1\}.$$

$$\tag{9}$$

In the rest of the paper, we consider the simpler situation that the loss function $f(\cdot)$ is differentiable and that it has non-negative domain as in Josey et al. (2021b). Entropy, KL divergence, empirical likelihood, logistic loss (Tan, 2020b) and many other choices satisfy this condition, whereas the quadratic function in Zubizarreta (2015) does not. This condition allows us to consider gradient instead of subgradient, and obtain $\hat{w}_i = (\tilde{f}^*)' \left(\hat{\theta}^\top W^{1/2}(\Phi(X_i) - \bar{\Phi})\right) = (\tilde{f}')^{-1}(\hat{\theta}^\top W^{1/2}(\Phi(X_i) - \bar{\Phi})),$ where $(\tilde{f}^*)'(\cdot)$ is the first derivative of $\tilde{f}^*(\cdot)$. For example, when we use the entropy $f(x) = x \log(x)$ as the loss function, the dual problem (8) becomes

$$\underset{\theta \in \mathbb{R}^{K}}{\text{minimize}} : \sum_{i=1}^{n} T_{i} \exp\left\{\theta^{\top} W^{1/2}(\Phi(X_{i}) - \bar{\Phi}) - 1\right\} + \sqrt{\delta} \|\theta\|_{2},$$

and the estimated balanced weight $\hat{w}_i = \exp\left\{\hat{\theta}^\top W^{1/2}(\Phi(X_i) - \bar{\Phi}) - 1\right\}$. The estimated MB weight, denoted by \hat{w}_i^{MB} , is then obtained by normalization:

$$\hat{w}_{i}^{MB} = \frac{\hat{w}_{i}}{\sum_{j:T_{j}=1} \hat{w}_{j}} = \frac{\exp\left\{\hat{\theta}^{\top} W^{1/2} \Phi(X_{i})\right\}}{\sum_{j:T_{j}=1} \exp\left\{\hat{\theta}^{\top} W^{1/2} \Phi(X_{j})\right\}}, \quad \text{for all } i \in \{j: T_{j}=1\}.$$
(10)

When the propensity score $\pi(x) = \Pr(T = 1 \mid x)$ follows the log model: $\log(\pi(x;\beta)) = \beta^{\top} \Phi(x)$, the inverse probability weight $1/\{n\pi(X_i;\beta)\}$ coincides with the expression of the unnormalized weight \hat{w}_i when $\Phi(\cdot)$ includes an intercept. Moreover, the normalized inverse probability weight $\pi^{-1}(X_i;\beta)/\{\sum_{j:T_j=1}\pi^{-1}(X_j;\beta)\}$ coincides with the expression of the MB weight \hat{w}_i^{MB} . There is a one-to-one correspondence of the dual parameter θ and the coefficient β in the propensity score model $\pi(x;\beta)$: $\beta = (W^{1/2})^{\top}\theta$. Therefore, solving the dual problem (8) is equivalent to fitting an ℓ_2 norm-based regularized generalized linear model with the logarithm as the link function.

When exact balancing is feasible, the balanced weights by entropy balancing (Hainmueller, 2012; Zhao & Percival, 2017) have the same expression as in equation (10), though parameter estimation is different because it does not involve regularization. In the bad-overlap

or high-dimensional situation, entropy balancing is infeasible. In contrast, there is no normalization constraint $\sum_{i=1}^{n} T_i w_i = 1$ in the primal problem (5). Instead, we normalize the weights after solving Problem (5). This two-step strategy guarantees that Problem (5) is always feasible for any $\delta > 0$ under the mild condition that W is positive-definite and $\phi_k(X_i)$ is bounded for any i and k.

Moreover, by regularizing the dual parameter θ with the ℓ_2 norm, Mahalanobis balancing stabilizes the estimated weights. The degree of regularization is determined by the tuning parameter δ , which controls the level of residual multivariate imbalance after reweighting. In contrast to entropy balancing which enforces finite-sample univariate exact balance, Mahalanobis balancing maintains finite-sample multivariate approximate balance.

When the loss function is the quadratic function by Zubizarreta (2015), the distance measure by Chan et al. (2016), or the Bregman distance by Josey et al. (2021b), the conjugate function can be calculated analogously but the expression can be complicated. We refer the readers to the monograph by Bertsekas (2016) for an extensive discussion of the conjugate functions. We prefer the entropy loss function $f(x) = x \log(x)$ in Mahalanobis balancing for its theoretical properties (Zhao & Percival, 2017) and stable performance in our numerical experience.

Intuitively, a larger value of δ results in more conservative θ value, which leads to more stable MB weights. Consequently, treatment effect estimation may be more biased but exhibit less variability. Hence, selection of δ is a key to the bias and variance trade-off for treatment effect estimation. In Section 3.2, we make comparison of the proposed method to existing balancing methods from the perspective of treatment effect estimation. In Section 3.3, we study the asymptotic properties. In Section 3.4, we discuss selection of δ in details.

3.2 A comparison with existing balancing methods

We compare Mahalanobis balancing to existing balancing methods from the outcome modelling perspective. Write

$$Y_i(1) = \mu_1(X_i) + \epsilon_{1i},$$

where $\mu_1(X_i) = E\{Y_i(1) \mid X_i\}$. Assume that the ϵ_{1i} are mutually independent with mean zero and finite variance, and that they are independent of the covariates. Let $\hat{w}_1, \dots, \hat{w}_n$ be

the normalized weights estimated by an arbitrary weighting method, and $\hat{\tau}_1 = \sum_{i=1}^n T_i \hat{w}_i Y_i$ be the resulting weighted estimator of $\tau_1 = E\{Y(1)\}$. It follows that (Wong & Chan, 2018)

$$\widehat{\tau}_{1} - \tau_{1} = \underbrace{\sum_{i=1}^{n} \left(T_{i} \widehat{w}_{i} - \frac{1}{n} \right) \mu_{1}(X_{i})}_{A_{1}} + \underbrace{\sum_{i=1}^{n} T_{i} \widehat{w}_{i} \epsilon_{1i}}_{A_{2}} + \underbrace{\frac{1}{n} \sum_{i=1}^{n} \mu_{1}(X_{i}) - E\{Y(1)\}}_{A_{3}}.$$

A good weighting method should minimize or control the magnitudes of the terms A_1 and A_2 . We first discuss the term A_1 . If the linear outcome model $\mu_1(X) = \beta_1^{\top} \Phi(X)$ holds (e.g., Athey et al., 2018), then $A_1 = \beta_1^{\top} (\sum_{i=1}^n T_i \hat{w}_i \Phi(X_i) - \bar{\Phi})$. The exact balancing methods (e.g., Hainmueller, 2012; Imai & Ratkovic, 2014; Chan et al., 2016; Fan et al., 2022) impose $\sum_{i=1}^{n} T_i \hat{w}_i \Phi(X_i) = \bar{\Phi}$, and thus A_1 is exactly zero. In the bad-overlap or high-dimensional situation, the term A_1 cannot be fully eliminated, and exact balancing is not applicable. The covariate functional balancing method (Wong & Chan, 2018) directly minimizes A_1^2 with the basis functions $\Phi(\cdot)$ restricted to the RKHS. The MD-ABW method (Zubizarreta, 2015; Wang & Zubizarreta, 2020) bounds the kth component of $|\sum_{i=1}^{n} T_i \hat{w}_i \Phi(X_i) - \bar{\Phi}|$ by a threshold parameter δ_k , $k = 1, \dots, K$. Therefore, $|A_1| \leq |\beta_1^{\top} \delta|$, where $\delta = (\delta_1, \cdots, \delta_K)^{\top}$, suggesting that A_1 is controlled by the MDABW method. However, it is difficult to tighten the upper-bound by tuning $(\delta_1, \dots, \delta_K)$ alone without modelling the outcomes. Note that $|A_1| \leq \|\beta_1^\top W^{-1/2}\|_2 \sqrt{\text{GMIM}_1^{\hat{w}}}$, where the term $\text{GMIM}_1^{\hat{w}}$ does not involve the unknown regression parameter β_1 . This inequality immediately suggests a simple guideline for the selection of the univariate threshold parameter δ in Mahalanobis balancing. A good choice of δ should make $\text{GMIM}_1^{\hat{w}}$ small enough so that the term A_1 is under control. We provide more details in Section 3.4.

Moreover, regardless of the linear outcome model assumption, the Cauchy-Schwarz inequality leads to that $|A_1| \leq \sqrt{\|\hat{w}\|_2^2 - 1/n} \sqrt{\sum_{i=1}^n \{\mu_1(X_i)\}^2}$ and $|A_2| \leq \|\hat{w}\|_2 \|\epsilon_1\|_2$, where $\|\hat{w}\|_2 = \sqrt{\sum_{i=1}^n T_i \hat{w}_i^2}$ and $\|\epsilon_1\|_2 = \sqrt{\sum_{i=1}^n T_i \epsilon_{1i}^2}$. Therefore, to minimize or control the magnitudes of the terms A_1 and A_2 , a good weighting method should control $\|\hat{w}\|_2$. Intuitively, it suggests that the weights should be stable and no extreme weights are allowed. The covariate functional balancing method (Wong & Chan, 2018) uses $\|\hat{w}\|_2^2$ as a regularizer. The approximately residual balancing method (Athey et al., 2018) minimizes a linear combination of $\|\sum_{i=1}^n T_i \hat{w}_i \Phi(X_i) - \bar{\Phi}_0\|_{\infty}^2$ and $\|\hat{w}\|_2^2$, where $\bar{\Phi}_0$ is the sample average of $\Phi(X_i)$ in the control group. Mahalanobis balancing stabilizes the estimated weights via the ℓ_2 norm regularizer. We show that $\|\hat{w}\|_2 = O_p(n^{-1/2})$ holds for Mahalanobis balancing in the Supplementary Material.

3.3 Asymptotic properties

The proposed Mahalanobis balancing method estimates the average treatment effect by

$$\hat{\tau}^{MB} = \sum_{i=1}^{n} T_i \hat{w}_i^{MB} Y_i - \sum_{i=1}^{n} (1 - T_i) \, \hat{w}_i^{MB} Y_i,$$

where the \hat{w}_i^{MB} are the estimated MB weights obtained by normalizing the weights \hat{w}_i in equation (9). In this subsection, we prove that this MB-based ATE estimator is doubly robust and semiparametrically efficient under mild regularity conditions. The proofs are given in the Supplementary Material.

Assumption 3. Assume the following conditions:

(3.1). The optimization problem $\min_{\theta \in \Theta} E[\tilde{f}^*(\theta^\top W^{1/2}(\Phi(X_i) - E\{\Phi(X)\}))]$ has a unique global minimizer for θ , where $\theta \in int(\Theta)$, and Θ is a compact parameter space for θ .

(3.2). $\delta = o(n)$.

(3.3). The conjugate function $\tilde{f}^*(\cdot)$ satisfies the property that if $\tilde{f}^*(\theta^\top W^{1/2}(\Phi(X) - E\{\Phi(X)\})) = C * \tilde{f}^{*'}(\theta^{*\top} W^{1/2}(\Phi(X) - E\{\Phi(X)\}))$ for some constant C for all X, then $\theta^* = \theta$.

(3.4). $E\{\exp(a\theta^{\top}W^{1/2}[\Phi(X) - E\{\Phi(X) \mid T = j\}])\} < \infty \text{ for all } \theta \in \Theta, \text{ where } a = 1, 2, 3, j = 0, 1.$ Moreover, $Var(\Phi(X)) < \infty$ and $Var(\Phi(X) \mid T) < \infty.$

(3.5). The noises $\epsilon_{1i} = Y_i(1) - E\{Y_i(1) \mid X_i\}$ are mutually independent and sub-Gaussian. Similarly, the noises $\epsilon_{0i} = Y_i(0) - E\{Y_i(0) \mid X_i\}$ are mutually independent and sub-Gaussian. All the noises are independent of X.

(3.6). $\eta \le \pi(X) \le 1 - \eta$, where $0 < \eta < 1/2$.

Assumption (3.1) is a standard requirement for consistency of the ATE estimator. Assumption (3.2) requires that the threshold parameter δ should not be large. Assumption (3.3) is satisfied by the common choices $f(x) = x \log(x)$ or $-\log(x)$. Assumptions (3.4) and (3.5) are mild conditions to control the moments of the covariates and the noises. Assumption (3.6) is the strict overlap assumption.

Theorem 2. Suppose that Assumptions 1-3 hold, then the MB-based ATE estimator $\hat{\tau}^{MB}$ is doubly robust in the sense that:

(i). If the propensity score satisfies $1/\pi(X) = C * \tilde{f}^{*'}(\theta^{*\top}W^{1/2}(\Phi(X) - \bar{\Phi}))$ for some $\theta^* \in int(\Theta)$ and some constant C for all X, then $\hat{\tau}^{MB} - \tau = O_p(n^{-1/2});$

(ii). If the conditional potential outcomes $E\{Y(1)|X\}$ and $E\{Y(0)|X\}$ are linear combinations of the basis functions $\Phi(X) = (\phi_1(X), \dots, \phi_K(X))^{\top}$, then $\widehat{\tau}^{MB} - \tau = O_p(n^{-1/2})$ when at least one of the following two statements is true: (a). The loss function is the entropy function; (b). The following conditions hold:

 $\sum_{i=1}^{n} I(T_i = j) (\hat{w_i}^{MB})^3 / (\sum_{i=1}^{n} I(T_i = j) (\hat{w_i}^{MB})^2)^{3/2} = o_p(1), \text{ and } \sum_{i=1}^{n} I(T_i = j) (\hat{w_i}^{MB})^2 = O_p(n^{-1}), \ j = 0, 1.$

Note that the condition $1/\pi(X) = C * \tilde{f}^{*'}(\theta^{*\top}W^{1/2}(\Phi(X) - \bar{\Phi}))$ implies that the unnormalized MB weight in equation (9) is proportional to the inverse weight $1/\{n\pi(X_i)\}$. When we use entropy as the loss function, then this assumption says that $1/\pi(X) = C * \exp(\theta^{*\top}\Phi(X))$ holds for some θ^* and constant C. The two conditions in statement (b) require that the weights should be smoothly distributed and no extremely large weights are allowed. That is, the weights should be stable. When the loss function is the entropy function, these two conditions are automatically satisfied under Assumption 3.

3.4 Tuning parameter selection

Recall that Problem (5) is always feasible for any $\delta > 0$ under mild conditions. Therefore, we are allowed to freely try a set of grid points and select the best δ that minimizes multivariate imbalance. In contrast to the MDABW method, no resampling is required in the selection procedure.

Now, we present the selection procedure in details. First, set up a set of positive values \mathcal{D} for δ selection. We use $\mathcal{D} = \{10^{-k} : k = 0, 1, \dots, 6\}$ in the simulations and application. In extensive numerical studies, we find that searching δ in this range leads to satisfactory performance of the MB method. More grid points can be added to \mathcal{D} to potentially better control multivariate imbalance, though we do not find notable improvement in our simulations.

The algorithm for selection of δ is as follows:

Algorithm 1. Selection of δ :

For each $\delta \in \mathcal{D}$:

Compute the dual parameter $\hat{\theta}$ by solving Problem (8); Compute the balanced weights \hat{w}_i using (9); Obtain the Mahalanobis balancing weights by normalization: $\hat{w}_i^{MB} = \hat{w}_i / \sum_{j:T_j=1} \hat{w}_j$; Calculate GMIM^w₁ in (6) using the \hat{w}_i^{MB} ; Output δ^* that minimizes GMIM^w₁.

Similarly, we select the optimal threshold parameter for the control group by minimizing GMIM_0^w . Let $(\hat{w}_1^{MB}, \dots, \hat{w}_n^{MB})$ be the resulting MB weights corresponding to the optimal threshold parameters in the two groups. The average treatment effect is then estimated by $\hat{\tau}^{MB} = \sum_{i=1}^n T_i \hat{w}_i^{MB} Y_i - \sum_{i=1}^n (1 - T_i) \hat{w}_i^{MB} Y_i$.

Interestingly, our empirical experience is that the ATE estimation is not quite sensitive to the choice of δ if it is small enough. In the Supplementary Material, the simulations reveal that the outputs of ATE estimation and imbalance measure are similar when $\delta \leq 10^{-2}$. Therefore, if computation cost is a concern, we suggest using a fixed small value for δ in practise, say $\delta = 10^{-4}$.

We offer some empirical guideline for evaluating whether the GMIM value is "small enough" for the MB method and other weighting methods. The concept that a threshold for GMIM is "small enough" is empirical, as analogous to the thresholds 0.1, 0.2 and 0.25 for ASMD (e.g. Stuart et al., 2013; Imbens & Rubin, 2015; Xie et al., 2019). Our guideline is that if GMIM ≤ 1 and GMIM/ $K \leq 0.01$, then the covariates are approximately balanced in the multivariate sense. The latter condition implies that the TASMD values are small. If either condition fails, then it implies that the weighting method cannot effectively balance covariates, and the corresponding treatment effect estimation may exhibit significant bias.

3.5 High-dimensional Mahalanobis balancing

In the high-dimensional setting where K is large compared to the sample size n, it becomes very difficult to simultaneously balance all basis functions even when each of the basis functions is approximately balanced in the univariate sense. The difficulty is easily seen from the multivariate perspective: when the basis functions are mutually independent, the squared Mahalanobis distance increases linearly with K, and hence multivariate imbalance can be hardly controlled for large K. Exact balancing is infeasible, and it becomes more difficult for univariate approximate balancing to obtain optimal threshold parameters.

Mahalanobis balancing is still feasible in the high-dimensional situation under the mild conditions that the weight matrix is positive-definite and the basis functions are bounded. Moreover, it still only needs to tune one threshold parameter. By minimizing multivariate imbalance to some extent, Mahalanobis balancing generally produces more balanced weights and less biased treatment effect estimation compared to univariate approximate balancing.

We provide asymptotic property for the MB-based ATE estimator in the high-dimensional situation. We choose the entropy function as the loss function.

Assumption 4. Assume the following conditions:

(4.1). The number of constraints $K/n \to \tau$, where $0 < \tau < 1$. (4.2). $|\theta^{\top}W^{1/2}(E\{\Phi(X)\} - E\{\Phi(X) \mid T\})| \leq \gamma \log(n)$ for all $\theta \in \Theta$, where $\gamma > 0$. (4.3). $\mu_j(X) = \beta_j^{\top} \Phi(X)$ with $\|\beta_j\|_2 = O(n^{\alpha})$, j = 0, 1, where α is a real number. (4.4) $\delta = O(n^{2s})$, where s is a real number.

Assumption (4.1) restricts the number of constraints K. It is allowed to increase with the sample size n, but it should not exceed n. Assumption (4.2) adds additional assumption on $W^{1/2}(E\{\Phi(X)\} - E\{\Phi(X) \mid T\})$. It is satisfied, for instance, when $\|\theta\|_2 \leq \gamma \log(n)$ and $\|E\{\Phi(X) \mid T\} - E\{\Phi(X)\}\|_2 \leq 1$, or when the θ is sparse. Assumption (4.3) imposes linear outcome models where the magnitude of the coefficients is not too large. Assumption (4.4) requires that δ should not be large.

Theorem 3. Suppose that Assumptions 1-4 hold, then $\hat{\tau}^{MB} - \tau = O_p(n^{-1/2}) + O_p(n^{s+\gamma+\alpha})$.

It suggests that δ should be small, so one may set s = 0 in Assumption (4.4). According to Theorem 3, given that $s + \gamma + \alpha \leq -1/2$, the MB-based ATE estimator $\hat{\tau}^{MB}$ is \sqrt{n} - consistent for τ . Moreover, this theorem illustrates the difficulty of ATE estimation in the high-dimensional setting. For example, when K = n and $\gamma > 0$, then Theorem 3 no longer guarantees the consistency of the MB-based ATE estimator. In the Supplementary Material, we propose a modified version of Mahalanobis balancing, and make extensive comparison of Mahalanobis balancing with existing regularized balancing methods in both sparse and non-sparse high-dimensional simulation settings.

3.6 Semiparametric efficiency

Next, we prove the semiparametric efficiency property for the MB-based ATE estimator.

Assumption 5. Assume the following conditions:

(5.1). Suppose that $1/\pi(X) = \tilde{f}^{*'}(m^*(X))$ for all X, where $m^*(\cdot) \in \mathcal{M}$ and \mathcal{M} is a set of smooth functions satisfying $\log n_{[]}\{\varepsilon, \mathcal{M}, L_2(P)\} \leq C_1(1/\varepsilon)^{1/k_1}$. Here, C_1 is a positive constant, $k_1 > 1/2$, and $n_{[]}\{\varepsilon, \mathcal{M}, L_2(P)\}$ denotes the covering number of \mathcal{M} by ε -brackets. (5.2). $\tilde{f}^{*'}(m^*(X))$ is Lipschitz in \mathcal{X} , where \mathcal{X} is the domain of covariate X.

(5.3). There exists a value θ^* such that $\sup_{x \in \mathcal{X}} \|\theta^{*\top} \left(\Phi(x) - \bar{\Phi}\right) - m^*(x)\|_2 = o_p(1).$

(5.4). The conditional potential outcomes $E\{Y(1)|X\}$ and $E\{Y(0)|X\}$ are linear combinations of the basis functions $\Phi(X) = (\phi_1(X), \cdots, \phi_K(X))^\top$.

(5.5). There exists a constant C_2 such that

$$\sup_{x \in \mathcal{X}} \| \left(\Phi(x) - \bar{\Phi} \right) \|_2 \le C_2 n^{1/2} \text{ and } E \left\{ \| \left(\Phi(x) - \bar{\Phi} \right) \|_2^2 \right\} \le C_2$$

(5.6). $s + \gamma + \alpha < -\frac{1}{2}$. (5.7). Let $g(X_i) = \frac{T_i}{\pi(X_i)} \left(\Phi(X_i) - \bar{\Phi} \right)$, then g(X) has independent σ^2 -sub-gaussian entries g_j with $\operatorname{Var}(g_j) > \tau^2$, where $\sigma = O(\operatorname{polyLog}(n))$ and $\tau = O\left(\frac{1}{\operatorname{polyLog}(n)}\right)$.

Our setting is different from Wang & Zubizarreta (2020), where they require the dimension K should be shrunk faster than the sample size n. Assumption (5.1) requires that the complexity of the function class is sufficiently smooth. Wang & Zubizarreta (2020) noted that the Hölder class with smoothness parameter v with v/K > 1/2 satisfies this condition (see also van der Vaart & Wellner, 1996; Fan et al., 2022). Assumption (5.2) bounds the second derivative of the function $\tilde{f}^*(\cdot)$. Assumption (5.3) requires that the $m^*(\cdot)$ can be approximated by the linear span of the basis functions. It is similar to Assumption 1.6 by Wang & Zubizarreta (2020). Assumption (5.4) is a standard condition for semiparametric efficiency. This condition is different from Wang & Zubizarreta (2020), since we discuss the weaker assumption that $K/n \rightarrow \tau > 0$. Assumption (5.5) is a standard technical assumption that restricts the magnitude of the basis functions. Assumption (5.6) corresponds to the level of sparsity. Assumption (5.7) provides a lower bound for the smallest eigenvalue for the empirical covariance matrix of $\Phi(X)$.

Theorem 4. Suppose that Assumptions 1-5 hold, then the MB-based ATE estimator reaches the semiparametric efficiency bound.

4 Numerical Studies

In this section, we compare Mahalanobis balancing to two classes of existing balancing methods for ATE estimation in numerical studies. The first class consists of exact balancing methods, including entropy balancing (EB) (Hainmueller, 2012) implemented by the R package WeightIt, covariate balancing propensity score (CBPS) (Imai & Ratkovic, 2014) implemented by the R package CBPS, and calibration weighting (CAL) (Chan et al., 2016) implemented by the R package ATE. The second class consists of univariate approximate balancing methods, among which the MDABW method (Wang & Zubizarreta, 2020) is implemented by the R package sbw. We also report the unadjusted ATE estimator using the simple difference in the outcomes (Unad) and the propensity score weighting estimator (PS) using logistic regression implemented by the R package WeightIt. In the end of this section, we discuss the kernel-based covariate balancing method (Wong & Chan, 2018).

For each method, we report the bias (Bias) of ATE estimation, Monte Carlo standard deviation (SD), root mean squared error (RMSE), Monte Carlo average of the mean of the TASMD_k, $k = 1, \dots, K$ (meanTASMD) as a univariate post-reweighting imbalance measure, and the generalized Mahalanobis imbalance measure (GMIM) GMIM = GMIM₁+ GMIM₀ with W_1 as the weight matrix to quantify residual multivariate imbalance.

We assess the performance of Mahalanobis balancing using W_1 (denoted by MB) or W_2 (denoted by MB2) as the weight matrix, respectively, and make comparison with exact balancing methods (EB, CBPS, CAL) and the MDABW method. We consider five

scenarios.

For each scenario, we conduct 1000 Monte Carlo simulations. The sample size is set to be n = 200 or n = 1000. Table 1 summarizes the outputs. The Supplementary Material provides additional information.

In Scenario A, we consider the situation where both propensity score and outcome models are misspecified. This setting similar to that of Kang & Schafer (2007). For each simulation, we first generate a standard normal random vector $Z = (Z_1, \ldots, Z_p)^{\top}$ with p = 10 for each observation, and then generate the covariates X from $X_1 = \exp(Z_1/2), X_2 =$ $Z_2/(1 + \exp(Z_1)), X_3 = (Z_1Z_3 + 0.6)^3, X_4 = (Z_2 + Z_4 + 20)^2, X_j = Z_j, j = 5, \cdots, 10.$ The potential outcomes are derived from linear regression utilizing the variables Z as follows: $(Y(1), Y(0)) = (210 + 13.7 \sum_{i=1}^{4} Z_i + \epsilon_1, 210 - 6.85 \sum_{i=1}^{4} Z_i + \epsilon_0)$, where ϵ_1 and ϵ_0 are independent standard normal variables which are independent of (T, Z). The true ATE is zero. The treatment indicator is generated from $T \sim$ Bernoulli $(\pi(Z))$, where $\pi(Z) = 1/(1 + \exp(0.5Z_1 + 0.1Z_4))$. The Z are latent variables and the X are observed. Hence, when one specifies a linear outcome model and a logistic propensity score model using the X as the covariates, both models are misspecified. In this scenario, $\Phi(X) = X$, and K = p = 10. Scenario A is a good-overlap setting, since the distributions of X are not quite different between the two treatment groups, and the exact balancing methods are applicable.

The MB and MB2 methods and other balancing methods show similar outputs. They all successfully remove covariate imbalance and obtain almost identical ATE estimates. The RMSEs of MB, MB2, CAL, and EB are somewhat smaller than those of MDABW and CBPS. We conclude that Mahalanobis balancing maintains the advantages of the exact balancing methods in Scenario A. Moreover, the computation cost of Mahalanobis balancing is less than 0.5% than that of MDABW. This is because MDABW needs 1000 bootstraps for each possible tuning parameter.

In Scenario B, we first generate $T \sim \text{Bernoulli}(0.5)$ for each observation. Set p = 10. If T = 1, the ten-dimensional covariates are generated from $X \sim N(1, \Sigma_1)$, where $Cov(X_j, X_k) = 2^{-I(j \neq k)}$; if T = 0, then the covariates are generated from $X \sim N(1, \Sigma_0)$, where $Cov(X_j, X_k) = I(j = k)$. This data generation procedure allows us to delineate the discrepancy of covariate distributions between the treated and control groups: the

mean and variance are the same, but the interaction terms are different. The potential outcomes are generated from $(Y(1), Y(0)) = (2\sum_{i=1}^{10} X_i + 2(X_1X_2 + X_2X_3 + \dots + X_9X_{10} + X_{10}X_1) + \epsilon_1, \sum_{i=1}^{10} X_i + (X_1X_2 + X_2X_3 + \dots + X_9X_{10} + X_{10}X_1) + \epsilon_0)$, where $\epsilon_j \sim N(0, 1)$, j = 0, 1. The true value of ATE is calculated via simulation. The first two moments of X are well balanced. In contrast, the interaction terms are strong confounders. We set $\Phi(X) = \{X_i, X_jX_k : 1 \le i \le 10, 1 \le j \le k \le 10\}$ and thus K = 65.

Because the covariance structures are quite dissimilar in the treated and control groups, both EB and CAL do not admit solutions, and thus Scenario B is a bad-overlap setting. In comparison, MB is able to approximately balance the interaction terms. The two imbalance measures of MB are substantially lower than those of MDABW and CBPS. Furthermore, MB significantly improves the performance of exact balancing and univariate approximate balancing for ATE estimation. It exhibits much smaller bias and RMSE than those of MD-ABW and CBPS. Nevertheless, MB2 performs unsatisfactorily because the weight matrix W_2 is unstable due to considerable correlation among the interaction terms, and its GMIM value is much larger than that of MB. We do not recommend MB2 in this situation.

In Scenario C, the covariate means are different in the two groups. Set p = 10. We first generate $T \sim \text{Bernoulli}(0.5)$ for each observation. If T = 1, the covariates are generated from $X \sim N(1, \Sigma_1)$; otherwise, $X \sim N(0, \Sigma_0)$. The covariance matrices Σ_1 and Σ_0 are the same as those in Scenario B. The potential outcomes are generated from (Y(1), Y(0)) = $(2\sum_{i=1}^{10} X_i + \epsilon_1, \sum_{i=1}^{10} X_i + \epsilon_0)$, where $\epsilon_j \sim N(0, 1)$, j = 0, 1. The true ATE is 5. In this scenario, we set $\Phi(X) = X$, and thus K = 10. Scenario C is a bad-overlap setting, because the covariate distributions are quite different such that EB and CAL are infeasible. MDABW exhibits much less bias compared to CBPS, but it has larger standard deviation. The proposed MB and MB2 methods have least biased ATE estimation, smallest RMSEs, and smallest GMIM values.

In Scenario D, the sample sizes are highly imbalanced. This situation is commonly seen in cohort studies where the exposure is rare and the size of the control group is very large. We fix the expected sample size of the treated group to be 50, and the control group has a sample size of roughly 950. Set $\Phi(X) = X$ and K = p = 10. The covariates are simulated by $X \sim N(1, I_{10\times10})$. The treatment indicator is simulated from $T \sim Bernoulli(\pi(X))$ with $\pi(X) = 1/\{1 + 19 \exp(\sum_{i=1}^{10} X_i - 10)\}$. The potential outcomes are simulated from $(Y(1), Y(0)) = (2 \sum_{i=1}^{10} X_i + \epsilon_1, \sum_{i=1}^{10} X_i + \epsilon_0)$, where $\epsilon_j \sim N(0, 1)$, j = 0, 1. The true ATE is 10. Again, this is a bad-overlap setting, where EB and CAL are infeasible. In contrast to Scenario C, the ATE estimate of CBPS is less biased than that of MDABW, and the RMSE is smaller. MB and MB2 greatly outperform MDABW and CBPS in terms of bias, RMSE, and GMIM.

In Scenario E, the distribution of the covariates is heavy-tailed. In specific, the covariate X is generated by $\log(X) \sim N(0, I_{5\times 5})$. The treatment indicator is generated from $T \sim Bernoulli(\pi(X))$, where $\pi(X) = 1/\{1+0.1 \exp(\sum_{i=1}^{5} X_i - 5)\}$. The potential outcomes are simulated from $(Y(1), Y(0)) = (2\sum_{i=1}^{5} X_i + \epsilon_1, \sum_{i=1}^{5} X_i + \epsilon_0)$, where $\epsilon_j \sim N(0, 1), j = 0, 1$. The true ATE is 8.24. Once again, this is a bad-overlap scenario, where the EB and CAL methods are not feasible. The CBPS estimate is less biased than that of MDABW, and the RMSE is smaller. Moreover, the MB and MB2 significantly outperform other methods.

We remark that although CBPS is an exact balancing method, it is still applicable to Scenarios B, C, D, and E, all of which are bad-overlap settings. This is because it employs the generalized method of moments for parameter estimation, which still works even when the moment constraints are not met exactly. In this sense, CBPS can be regarded as an approximate balancing method, though it does not control univariate dispersion of each covariate. The CBPS method has larger meanTASMD and GMIM values than the MDABW method in all scenarios. Scenarios B, C, D, and E suggest that the covariate distributions in each group (that is, the conditional distributions of X|T = 1 and X|T = 0) are still highly dissimilar to the distribution of X in the population after CBPS reweighting, leading to large estimation bias and RMSE values.

The GMIM recognizes post-weighting multivariate imbalance and is predictive of the performance of balancing methods for treatment effect estimation. Therefore, we recommend to use GMIM instead of meanTASMD to assess residual covariate imbalance in the bad-overlap settings.

We also implement the kernel-based covariate balancing method by Wong & Chan (2018), and the results are in column "KERNEL" in Table 1. In Scenario A where the model setup is similar to that in Wong & Chan (2018), the kernel-based covariate balancing method outperforms other methods. This result is not surprising because the nonparametric kernel method tends to be more robust to model misspecification. However, in all

the bad-overlap scenarios (Scenarios B, C, D, and E), the performance of the kernel-based method is not satisfactory. The slow convergence rate of the nonparametric method may explain the inferior performance of the kernel-based method in finite samples. Hence, while the kernel-based covariate balancing method may be preferred in some cases where model misspecification is a concern, it may not always be the optimal choice for achieving balance in observational studies especially when the overlap between two groups is poor, the sample sizes are imbalanced, or the covariates are heavy-tailed.

4.1 Application

We revisit a dataset from the National Supported Work program (Dehejia & Wahba, 1999). The National Supported Work program is a labor training program implemented in the 1970s by providing work experience to selected individuals. The data consist of a National Supported Work experimental group with sample size 185 and a nonexperimental comparison group from the Panel Study of Income Dynamics with sample size 429. We regard them as the treated and control groups, respectively. The outcome variable Y is the post-intervention earning measured in Year 1978.

The covariates include four numeric covariates age, education, earn1974, earn1975, four binary variables married, black, nodegree and hispanic, and sixteen interaction terms between the numeric covariates and the binary variables. In addition, we include the ratio education/age, which represents possible nonlinear effect of age at each level of education. Table 2 shows the results. Here, maxTASMD and medTASMD are the Monte Carlo averages of the maximum and medium of TASMD_k, $k = 1, \dots, K$, respectively. 500 bootstrap resamples are used to calculate the standard error for all methods.

This is a bad-overlap setting, since the EB and CAL methods fail to output ATE estimation. This can be explained by that the distribution of the covariate *education/age* is highly disimilar between the two groups. The CBPS method substantially reduces univariate and multivariate imbalance compared to the unadjusted method. The PS method has similar performance. The MDABW and KERNEL methods perform better than CBPS in the sense of smaller residual univariate and multivariate imbalance after reweighting. In comparison, the proposed Mahalanobis balancing methods produce most balanced weights as they have smallest TASMD and GMIM values, although they have larger standard error. The Mahalanobis balancing methods are recommended when balanced weights are in demand in data analysis.

5 Concluding Remarks

In this paper, we proposed Mahalanobis balancing. It produces balanced weights by solving a convex optimization problem with a quadratic constraint that represents multivariate imbalance. We discussed the high-dimensional setting. Different choices of the basis functions were examined in the simulations. We compared Mahalanobis balancing with the exact balancing, univariate approximate balancing, and high-dimensional regularized balancing methods in extensive numerical studies, and found that Mahalanobis balancing generally led to more balanced weights and less biased ATE estimation.

The proposed Mahalanobis balancing methods can be easily extended to the situation with multiple treatment arms. Other causal estimands, such as average treatment effect on the treated and average treatment effect on the control, can be also estimated using Mahalanobis balancing, though some modification is needed. For example, when the average treatment effect on the control is of interest, one needs to replace $\bar{\Phi}$ with the sample average of $\Phi(X)$ in the control group in Problem (5) to obtain Mahalanobis balancing weights.

Our method can be used to address the transportability and generalizability issues in data integration and data fusion. Recently, exact balancing methods were applied to these interesting problems (e.g. Lee et al., 2023; Josey et al., 2021b). Nevertheless, the trial population and the target population usually exhibit high degree of discrepancy, and there can be a large number of confounders. In these situations, exact balancing is infeasible and Mahalanobis balancing is recommended.

In the numerical studies, we used entropy as the loss function. One may further investigate the performance of Mahalanobis balancing using other loss functions (e.g. Chan et al., 2016; Josey et al., 2021b). Moreover, one may consider other multivariate imbalance measures instead of the Mahalanobis metric to improve robustness, e.g., energy distance (Huling & Mak, 2020) and kernel distance (Zhu et al., 2018). However, it is beyond the scope of this paper and we do not further pursue it here.

Supplementary Material

The online Supplementary Material includes discussion of the high-dimensional setting, proofs and additional simulation studies.

Acknowledgements

Ying Yan's research is supported by the National Natural Science Foundation of China (Grant No. 11901599).

References

- ATHEY, S., IMBENS, G. W. & WAGER, S. (2018). Approximate residual balancing: debiased inference of average treatment effects in high dimensions. J. R. Stat. Soc. Ser. B. Stat. Methodol. 80, 597–623.
- BERTSEKAS, D. P. (2016). Nonlinear Programming. Athena Scientific, 3rd ed.
- CHAN, K. C. G., YAM, S. C. P. & ZHANG, Z. (2016). Globally efficient non-parametric inference of average treatment effects by empirical balancing calibration weighting. J. R. Stat. Soc. Ser. B. Stat. Methodol. 78, 673–700.
- CHATTOPADHYAY, A., HASE, C. H. & ZUBIZARRETA, J. R. (2020). Balancing vs modeling approaches to weighting in practice. *Stat. Med.* **39**, 3227–3254.
- D'AMOUR, A., DING, P., FELLER, A., LEI, L. & SEKHON, J. (2021). Overlap in observational studies with high-dimensional covariates. J. Econometrics 221, 644–654.
- DEHEJIA, R. H. & WAHBA, S. (1999). Causal effects in nonexperimental studies: Reevaluating the evaluation of training programs. J. Amer. Statist. Assoc. 94, 1053–1062.
- DIAMOND, A. & SEKHON, J. S. (2013). Genetic matching for estimating causal effects: a general multivariate matching method for achieving balance in observational studies. *The Review of Economics and Statistics* **95**, 932–945.
- FAN, J., IMAI, K., LEE, I., LIU, H., NING, Y. & YANG, X. (2022). Optimal covariate balancing conditions in propensity score estimation. J. Bus. Econom. Statist. 41, 97–110.
- GRAHAM, B. S., CAMPOS DE XAVIER PINTO, C. & EGEL, D. (2012). Inverse probability tilting for moment condition models with missing data. *Rev. Econ. Stud.* **79**, 1053–1079.

- HAINMUELLER, J. (2012). Entropy balancing for causal effects: A multivariate reweighting method to produce balanced samples in observational studies. *Political analysis* 20, 25– 46.
- HAYAKAWA, S., LYONS, T. & OBERHAUSER, H. (2023). Estimating the probability that a given vector is in the convex hull of a random sample. *Probab. Theory Related Fields* **185**, 705–746.
- HAZLETT, C. (2020). Kernel balancing: A flexible non-parametric weighting procedure for estimating causal effects. *Statist. Sinica* **30**, 1155–1189.
- HIRANO, K., IMBENS, G. W. & RIDDER, G. (2003). Efficient estimation of average treatment effects using the estimated propensity score. *Econometrica* **71**, 1161–1189.
- HULING, J. D. & MAK, S. (2020). Energy balancing of covariate distributions. Arxiv.
- IACUS, S. M., KING, G. & PORRO, G. (2011). Multivariate matching methods that are monotonic imbalance bounding. J. Amer. Statist. Assoc. 106, 345–361.
- IACUS, S. M., KING, G. & PORRO, G. (2012). Causal inference without balance checking: coarsened exact matching. *Political Analysis* **20**, 1–24.
- IMAI, K. & RATKOVIC, M. (2014). Covariate balancing propensity score. J. R. Stat. Soc. Ser. B. Stat. Methodol. 76, 243–263.
- IMBENS, G. W. & RUBIN, D. B. (2015). Causal Inference in Statistics, Social, and Biomedical Sciences. Cambridge University Press.
- JOSEY, K. P., BERKOWITZ, S. A., GHOSH, D. & RAGHAVAN, S. (2021a). Transporting experimental results with entropy balancing. *Stat. Med.* 40, 4310–4326.
- JOSEY, K. P., JUAREZ-COLUNGA, E., YANG, F. & GHOSH, D. (2021b). A framework for covariate balance using bregman distances. *Scand. J. Stat.* 48, 790–816.
- KANG, J. D. & SCHAFER, J. L. (2007). Demystifying double robustness: A comparison of alternative strategies for estimating a population mean from incomplete data. *Statist. Sci.* 22, 523–539.
- LEE, D., YANG, S., DONG, L., WANG, X., ZENG, D. & CAI, J. (2023). Improving trial generalizability using observational studies. *Biometrics* **79**, 1213–1225.
- LI, F., MORGAN, K. L. & ZASLAVSKY, A. M. (2018). Balancing covariates via propensity score weighting. J. Amer. Statist. Assoc. 113, 390–400.

- MA, X. & WANG, J. (2020). Robust inference using inverse probability weighting. J. Amer. Statist. Assoc. 115, 1851–1860.
- MOHRI, M., ROSTAMIZADEH, A. & TALWALKAR, A. (2018). Foundations of Machine Learning. MIT press, 2nd ed.
- MORDUKHOVICH, B. S. & NAM, N. M. (2022). Convex Analysis and Beyond: Volume I: Basic Theory. Springer.
- NING, Y., PENG, S. & IMAI, K. (2020). Robust estimation of causal effects via a highdimensional covariate balancing propensity score. *Biometrika* **107**, 533–554.
- ROBINS, J., ROTNITZKY, A. & ZHAO, L. (1994). Estimation of regression coefficients when some regressors are not always observed. J. Amer. Statist. Assoc. 89, 846–866.
- ROSENBAUM, P. R. (1987). Model-based direct adjustment. J. Amer. Statist. Assoc. 82, 387–394.
- ROSENBAUM, P. R. & RUBIN, D. B. (1983). The central role of the propensity score in observational studies for causal effects. *Biometrika* **70**, 41–55.
- STUART, E. A., LEE, B. K. & LEACY, F. P. (2013). Prognostic score-based balance measures can be a useful diagnostic for propensity score methods in comparative effectiveness research. *Journal of Clinical Epidemiology* 66, S84–S90.
- TAN, Z. (2020a). Model-assisted inference for treatment effects using regularized calibrated estimation with high-dimensional data. Ann. Statist. 48, 811–837.
- TAN, Z. (2020b). Regularized calibrated estimation of propensity scores with model misspecification and high-dimensional data. *Biometrika* **107**, 137–158.
- VAN DER VAART, A. W. & WELLNER, J. A. (1996). Weak Convergence and Empirical Processes. Springer.
- WANG, Y. & ZUBIZARRETA, J. R. (2020). Minimal dispersion approximately balancing weights: asymptotic properties and practical considerations. *Biometrika* **107**, 93–105.
- WONG, R. K. & CHAN, K. C. G. (2018). Kernel-based covariate functional balancing for observational studies. *Biometrika* **105**, 199–213.
- XIE, Y., ZHU, Y., COTTON, C. A. & WU, P. (2019). A model averaging approach for estimating propensity scores by optimizing balance. *Stat. Methods Med. Res.* 28, 84–101.
- YIU, S. & SU, L. (2018). Covariate association eliminating weights: a unified weighting framework for causal effect estimation. *Biometrika* 105, 709–722.

- ZHAO, Q. (2019). Covariate balancing propensity score by tailored loss functions. Ann. Statist. 47, 965–993.
- ZHAO, Q. & PERCIVAL, D. (2017). Entropy balancing is doubly robust. J. Causal Inference 5, 20160010.
- ZHU, Y., SAVAGE, J. S. & GHOSH, D. (2018). A kernel-based metric for balance assessment. J. Causal Inference 6, 20160029.
- ZUBIZARRETA, J. R. (2015). Stable weights that balance covariates for estimation with incomplete outcome data. J. Amer. Statist. Assoc. 110, 910–922.

Ying Yan

Corresponding Author. yanying7@mail.sysu.edu.cn

School of Mathematics, Sun Yat-sen University, Guangzhou, China, 510275

	Unad	\mathbf{PS}	\mathbf{EB}	CBPS	\mathbf{CAL}	\mathbf{MB}	MB2	MDABW	KERNEL	
Scenario A with $(n, K) = (200, 10)$: both propensity score and outcome models are misspecified.										
Bias	-1.79	0.57	0.31	0.28	0.31	0.29	0.29	0.51	-0.03	
\mathbf{SD}	2.94	3.33	3.06	3.12	3.06	3.06	3.06	3.08	3.09	
RMSE	3.44	3.38	3.08	3.13	3.08	3.07	3.07	3.12	3.09	
meanTASMD	0.15	0.06	0.00	0.05	0.00	0.00	0.00	0.01	0.02	
GMIM	0.20	0.06	0.00	0.02	0.00	0.00	0.00	0.00	0.01	
Scenario B with	(n, K) =	(200, 6)	55): the	e confoun	ders incl	lude co	variate i	nteractions.		
Bias	6.93	4.14	-	3.72	-	-0.53	3.13	1.44	2.95	
\mathbf{SD}	4.07	7.49	-	5.17	-	1.00	2.58	1.38	2.11	
RMSE	8.04	8.56	-	6.37	-	1.13	4.05	1.99	3.62	
meanTASMD	0.21	0.33	-	0.23	-	0.04	0.13	0.12	0.12	
GMIM	2.09	7.36	-	2.94	-	0.12	0.81	0.55	225.50	
Scenario C with	(n,K) =	(200, 1	10: cor	variate me	eans are	very d	ifferent.			
Bias	15.00	8.71	-	7.05	-	0.69	0.92	3.31	8.94	
\mathbf{SD}	0.76	4.24	-	4.52	-	0.84	0.92	1.45	2.12	
RMSE	15.01	9.69	-	8.37	-	1.09	1.30	3.61	9.19	
meanTASMD	0.89	0.67	-	0.51	-	0.04	0.05	0.20	0.53	
GMIM	4.07	2.80	-	1.74	-	0.03	0.04	0.23	1.63	
Scenario D with	(n,K) =	(1000,	10): sat	ample size	es are ve	ery imb	alanced	across treatm	ent groups.	
Bias	-8.60	-1.60	-	-0.74	-	-0.25	-0.25	-1.37	-6.06	
\mathbf{SD}	0.31	2.11	-	0.41	-	0.44	0.44	0.77	2.32	
\mathbf{RMSE}	8.60	2.65	-	0.85	-	0.50	0.50	1.57	6.49	
meanTASMD	0.48	0.25	-	0.19	-	0.01	0.01	0.09	0.31	
GMIM	1.58	0.86	-	0.28	-	0.00	0.00	0.05	1.03	
	<pre> `</pre>	,								
Scenario E with	(n,K) =	(200, 5)	b): hear	vy-tailed o	covariat	es.				
Bias	-8.91	-3.40	-	-1.33	-	-0.66	-0.66	-2.55	-4.41	
SD	0.61	2.10	-	0.85	-	0.68	0.68	1.56	1.12	
RMSE	8.93	3.99	-	1.58	-	0.94	0.95	2.99	4.55	
meanTASMD	0.35	0.25	-	0.23	-	0.02	0.02	0.16	0.14	
GMIM	0.91	0.70	-	0.42	-	0.01	0.01	0.21	0.21	

Table 1: Simulation outputs in the low-dimensional settings.

Table 2: Data analysis of National Supported Work program.

	Unad	\mathbf{PS}	\mathbf{EB}	CBPS	\mathbf{CAL}	MB	MB2	MDABW	KERNEL
ATE	-606.09	635.29	-	1329.66	-	1529.37	1516.03	960.71	141.87
\mathbf{SE}	690.03	1368.81	-	1525.42	-	2044.52	2082.54	1231.93	1018.23
$\max TASMD$	1.31	0.35	-	0.38	-	0.01	0.02	0.12	0.26
medTASMD	0.34	0.21	-	0.19	-	0.00	0.02	0.08	0.12
meanTASMD	0.56	0.21	-	0.20	-	0.00	0.01	0.08	0.11
GMIM	3.38	0.49	-	0.30	-	0.00	0.01	0.05	0.21

Supplementary Material for "Mahalanobis balancing: a multivariate perspective on approximate covariate balancing"

Yimin Dai

School of Mathematics, Sun Yat-sen University, Guangzhou, China, 510275 and

Ying Yan*

School of Mathematics, Sun Yat-sen University, Guangzhou, China, 510275

A High-dimensional Mahalanobis balancing

Asymptotic theory of Mahalanobis balancing becomes much more difficult in the ultra high-dimensional setting with $K \gg n$. For example, the general high-dimensional regularized M-estimation theory (Wainwright, 2019) does not directly apply because the ℓ_2 norm regularizer in the dual problem (8) is not decomposable.

When the true propensity score model is not sparse, Mahalanobis balancing outperforms the high-dimensional regularized balancing methods (Athey et al., 2018; Ning et al., 2020; Tan, 2020a,b) and univariate approximate balancing (Wang and Zubizarreta, 2020) in the simulation studies. Nevertheless, when the true propensity score model is sparse, the performance of Mahalanobis balancing is not entirely satisfactory compared to highdimensional regularized balancing. This is not surprising because high-dimensional regularized balancing commonly selects a sparse subset of basis functions by ℓ_1 or elastic net regularization. In contrast, the ℓ_2 regularization in the dual problem (8) suggests that Mahalanobis balancing does not automatically perform variable selection.

In the sparse setting, we propose the high-dimensional Mahalanobis balancing method, which is a modified version of Mahalanobis balancing. It differs from Mahalanobis balancing by performing selection of basis functions before producing balanced weights.

^{*}Corresponding Author. yanying7@mail.sysu.edu.cn

Recall that we write $X = (X_1, \dots, X_p)^{\top}$. In the following high-dimensional setting, we use the trivial feature mapping: $\Phi(X) = X$, and thus K = p. In high-dimensional Mahalanobis balancing, we select a subset of the covariates X with dimension K_0 , where $K_0 \leq p$. The following algorithm gives a principled way for subset selection. Let $ASMD_j$ be the absolute standardized mean difference for X_j , $j = 1, \dots, p$. Rank X_1, \dots, X_p as $X_{(1)}, \dots, X_{(p)}$ such that $X_{(1)}$ has the largest ASMD value, $X_{(2)}$ has the second largest ASMD value, and so forth.

Algorithm S1. High-dimensional Mahalanobis balancing

For each step $j \in \{1, \dots, p\}$: Apply Algorithm 1 to obtain the MB weights \hat{w}_i^{*MB} using $\Phi(X) = (X_{(1)}, \dots, X_{(j)})^{\top}$; Calculate $GMIM_1^w$ in (6) using the \hat{w}_i^{MB} , and define $GMIM_{1,j}^w = GMIM_1^w/j$; Add $(j, GMIM_{1,j}^w)$ to the scatter plot; Observe whether there is a kink at $(j, GMIM_{1,j}^w)$: If no, let j = j + 1; If yes, stop and output $K_0 = j - 1$. If no kink is observed, output $K_0 = p$.

We then obtain $\Phi(X) = (X_{(1)}, \dots, X_{(K_0)})^{\top}$ and the MB weights at Step K_0 . Similarly, we obtain the MB weights for the control group. The average treatment effect is then estimated by the weighted difference of the outcomes.

The rationale of Algorithm S1 hinges on the adjusted multivariate imbalance measure $\text{GMIM}_{1,j}^w$. It represents the average contribution of the j most univariate imbalanced covariates to the residual multivariate imbalance after MB weighting. If it remains stable as j increases, then MB is capable of controlling multivariate imbalance. However, if there is a kink at Step j, it implies that adding the jth largest imbalanced covariate greatly increases the multivariate imbalance measure, implying that MB starts to lose control of overall imbalance at Step j. Therefore, we stop and choose the outputs at Step j - 1. The kink usually occurs when $K_0 = O(\sqrt{p})$ in the numerical studies. If there is no kink for all $j = 1, \dots, p$, it suggests that the performance of MB is acceptable even if all pcovariates are included. One may choose $K_0 = p$ in this situation, or $K_0 = \sqrt{p}$ if a small subset of covariates is preferred. Our numerical experience reveals that high-dimensional

	Unad	\mathbf{MB}	$\mathbf{kernelMB}$	hdMB	MDABW	RCAL1	RCAL2	ARB	hdCBPS
Scenario F with $(n, p) = (200, 100)$: sparse propensity score model.									
Bias	-3.44	-0.85	-0.92	-0.01	-2.13	-0.48	-0.47	-0.22	-0.19
\mathbf{SD}	0.36	0.30	0.29	0.32	1.11	0.32	0.31	0.27	0.37
RMSE	3.47	0.90	0.96	0.32	2.39	0.58	0.57	0.35	0.38
meanASMD	0.16	0.07	0.08	0.14	0.15	0.15	0.16	-	-
GMIM	4.35	0.53	0.71	$4.19(0.00)^{*}$	2.12	2.17	3.43	-	-
Scenario F with	(n, p) =	= (200.	500): sparse p	ropensity sco	re model.				
Bias	-3.46	-2.46	-2.49	0.00	-3.43	-0.76	-0.76	-0.43	-0.38
\mathbf{SD}	0.37	0.33	0.33	0.35	0.37	0.29	0.29	0.25	0.40
RMSE	3.48	2.48	2.52	0.35	3.45	0.81	0.81	0.48	0.55
meanASMD	0.12	0.10	0.10	0.22	0.12	0.12	0.13	-	-
GMIM	7.32	5.15	5.30	36.26(0.00)	7.28	7.36	7.43	-	-
Scenario G with	n (<i>n</i> . <i>p</i>) =	= (200.	100): dense p	ropensity scor	e model.				
Bias	-1.35	-0.35	-0.31	-0.55	-0.91	-0.76	-0.75	-0.54	-0.86
\mathbf{SD}	0.21	0.23	0.22	0.39	0.39	0.30	0.30	0.23	0.23
RMSE	1.36	0.42	0.38	0.68	0.99	0.82	0.81	0.59	0.89
meanASMD	0.20	0.07	0.08	0.16	0.15	0.20	0.20	-	-
GMIM	3.53	0.64	0.82	4.87(0.00)	1.86	3.03	3.20	-	-
Scenario G with $(n, n) = (200, 500)$; dense propensity score model									
Bias	-0.33	-0.23	-0.22	-0.27	-0.33	-0.32	-0.33	-0.27	-0.34
\mathbf{SD}	0.17	0.17	0.17	0.33	0.17	0.18	0.18	0.19	0.17
RMSE	0.37	0.29	0.28	0.43	0.37	0.37	0.37	0.33	0.39
meanASMD	0.12	0.10	0.10	0.22	0.12	0.18	0.13	-	-
GMIM	7.29	5.13	5.28	34.55(0.00)	7.26	7.38	7.52	-	-

Table S1: Simulation outputs in the high-dimensional settings.

* If we use $\Phi(X) = X$ to compute the GMIM for the hdMB method, then $\text{GMIM}^w = 4.19$; if we use $\Phi(X) = (X_{(1)}, \dots, X_{(K_0)})^{\top}$ with K_0 determined by Algorithm S1, then $\text{GMIM}^w = 0.00$.

Mahalanobis balancing substantially improves the performance of Mahalanobis balancing in the sparse setting.

Algorithm S1 tends to select covariates that exhibit large univariate imbalance. Algorithm S1 may not work well when the unselected imbalanced covariates are correlated with the outcomes. To alleviate this issue, one may construct a bias-corrected version for treatment effect estimation by augmenting the MB-based ATE estimator with the outcome models.

A.1 Simulation for the high-dimensional settings

In this subsection, we assess the performance of three Mahalanobis balancing methods in the high-dimensional setting, and compare them with high-dimensional regularized balancing methods, including two versions of regularized calibrated estimation (RCAL1 and RCAL2) (Tan, 2020a,b) implemented with the R package RCAL, approximately residual balancing (ARB) (Athey et al., 2018) implemented with the R package balanceHD, and high-dimensional covariate balancing propensity score (hdCBPS) (Ning et al., 2020) implemented with the R package CBPS. We also report the unadjusted ATE estimator using the simple difference in the outcomes (Unad).

The MB method was described in Section 4. We also consider kernel-based Mahalanobis balancing (kernelMB), where we use $\Phi(X) = (K(X, X_1), ..., K(X, X_n))^{\top}$ with $K(\cdot, \cdot)$ set to be the Gaussian kernel. High-dimensional Mahalanobis balancing (hdMB) was described in Section A, where $\Phi(X) = (X_{(1)}, \dots, X_{(K_0)})^{\top}$ and K_0 is selected by Algorithm S1. We compare them with four high-dimensional regularized balancing methods (RCAL1, RCAL2, ARB, hdCBPS). We do not report imbalance measures for ARB and hdCBPS, because they utilize outcome information in weight construction. For all methods except kernelMB, we set $\Phi(X) = X$ and thus K = p. In each scenario, we consider (n, p) = (200, 100) or (200, 500). Table S1 summarizes the outputs.

In Scenario F, we generate the covariates from $X \sim N(0, \Sigma)$ where $Cov(X_j, X_k) = 2^{-I(j \neq k)}$. The treatment assignment is generated by $T \sim Bernoulli(\pi(X))$ with $\pi(X) = 1/\{1 + \exp(X_1 + \sum_{j=2}^6 X_j/2)\}$. Therefore, treatment assignment is correlated with a sparse subset of covariates. The outcome is generated from $Y = T(\sum_{j=1}^5 X_j) + (1 - T)(\sum_{j=1}^5 X_j/2) + \epsilon$.

Both MB and kernelMB have low bias, small standard deviation, and small imbalance measures when (n, p) = (200, 100), implying that they are capable of approximately balancing covariates when the covariate dimension is substantially smaller than the sample size. When (n, p) = (200, 500), MB and kernelMB exhibit quite large GMIM values, and their biases are larger than those of the high-dimensional regularized balancing methods but are smaller than that of MDABW. The hdMB method shrinks the GMIM for the selected covariates to be zero, but its GMIM value is very large when all covariates are used to calculate this measure. It achieves lower bias and RMSE compared to MDABW and all high-dimensional regularized balancing methods (RCAL1, RCAL2, ARB, hdCBPS). In conclusion, the hdMB method is recommended over MB and kernelMB when the true propensity score model is sparse. It is competitive to the existing high-dimensional regularized balancing methods.

In Scenario G, the data generation procedure of the covariates is the same as the one

in Scenario F. The treatment index is generated by $T \sim Bernoulli(\pi(X))$ with $\pi(X) = 1/\{1 + \exp(X_1 + \sum_{j=2}^5 X_j/2 + 10 \sum_{j=6}^p X_j/p)\}$. Note that treatment assignment is correlated with all covariates, and thus the propensity score model is dense. The outcome is generated from $Y = T(10 \sum_{j=1}^p X_j/p) + (1 - T)(5 \sum_{j=1}^p X_j/p) + \epsilon$.

Both MB and kernelMB have lower bias and smaller standard deviation compared to hdMB and MDABW. The high-dimensional regularized balancing methods (RCAL1, RCAL2, ARB, hdCBPS) have larger biases than MB and kernelMB. Both MB and kernelMB produce much smaller GMIM values than other methods when (n, p) = (200, 100). The MB and kernelMB methods are recommended since they enjoy lowest biases, smallest RMSEs, and smallest GMIMs. The hdMB method is not recommended in this dense scenario.

A.2 Additional Simulation for Model Misspecification

We consider high-dimensional situation with (n, p) = (200, 100) when the outcome model is misspecified as a linear model. The data generation procedure of the covariates is the same as the one in Scenario F.

In the Scenario M1, we consider a sparse propensity score model

$$\pi(X_i) = \frac{1}{1 + \exp(X_{i1} + \sum_{j=2}^{6} X_{ij}/2)}$$

and a nonlinear outcome model

$$Y_i = 2T_i \left(\sum_{j=1}^6 X_{ij} + \sum_{j=1}^6 X_{ij}^2 \right) + (1 - T_i) \left(\sum_{j=1}^6 X_{ij} + \sum_{j=1}^6 X_{ij}^2 \right) + \epsilon_i,$$

where ϵ_i is a standard normal random variable.

In the Scenario M2, we consider a dense propensity score model

$$\pi(X_i) = \frac{1}{1 + \exp(X_{i1} + \sum_{j=2}^{5} X_{ij}/2 + \sum_{j=6}^{100} X_{ij}/10)},$$

Scenario M1	Bias	\mathbf{SD}	RMSE	maxASMD	meanASMD	medianASMD	GMIM
Unad	-3.45	0.50	3.49	1.54	0.16	0.10	3.22
\mathbf{MB}	-1.47	0.61	1.59	0.58	0.06	0.05	0.52
kernelMB	-1.52	0.56	1.62	0.80	0.08	0.06	0.71
hdMB	-0.72	0.73	1.03	1.25	0.14	0.10	4.19
MDABW	-2.50	0.96	2.68	1.54	0.14	0.11	2.12
RCAL1	-1.68	0.60	1.79	1.10	0.15	0.11	2.17
RCAL2	-1.67	0.60	1.78	1.49	0.15	0.11	2.76
ARB	-1.22	0.55	1.34	-	-	-	-
hdCBPS	-1.68	0.61	1.80	-	-	-	-
Scenario M2	Bias	SD	RMSE	maxASMD	meanASMD	medianASMD	GMIM
Scenario M2 Unad	Bias -1.35	SD 0.36	RMSE 1.40	maxASMD 1.30	meanASMD 0.20	medianASMD 0.17	GMIM 3.53
Scenario M2 Unad MB	Bias -1.35 -0.50	SD 0.36 0.48	RMSE 1.40 0.68	maxASMD 1.30 0.56	meanASMD 0.20 0.07	medianASMD 0.17 0.06	GMIM 3.53 0.64
Scenario M2 Unad MB kernelMB	Bias -1.35 -0.50 -0.35	SD 0.36 0.48 0.37	RMSE 1.40 0.68 0.51	maxASMD 1.30 0.56 0.74	meanASMD 0.20 0.07 0.08	medianASMD 0.17 0.06 0.07	GMIM 3.53 0.64 0.82
Scenario M2 Unad MB kernelMB hdMB	Bias -1.35 -0.50 -0.35 -0.70	SD 0.36 0.48 0.37 0.69	RMSE 1.40 0.68 0.51 0.98	maxASMD 1.30 0.56 0.74 1.28	meanASMD 0.20 0.07 0.08 0.16	medianASMD 0.17 0.06 0.07 0.11	GMIM 3.53 0.64 0.82 4.87
Scenario M2 Unad MB kernelMB hdMB MDABW	Bias -1.35 -0.50 -0.35 -0.70 -1.02	SD 0.36 0.48 0.37 0.69 0.51	RMSE 1.40 0.68 0.51 0.98 1.14	maxASMD 1.30 0.56 0.74 1.28 1.30	meanASMD 0.20 0.07 0.08 0.16 0.15	medianASMD 0.17 0.06 0.07 0.11 0.14	GMIM 3.53 0.64 0.82 4.87 1.86
Scenario M2 Unad MB kernelMB hdMB MDABW RCAL1	Bias -1.35 -0.50 -0.35 -0.70 -1.02 -1.20	SD 0.36 0.48 0.37 0.69 0.51 0.41	RMSE 1.40 0.68 0.51 0.98 1.14 1.27	maxASMD 1.30 0.56 0.74 1.28 1.30 0.93	meanASMD 0.20 0.07 0.08 0.16 0.15 0.20	medianASMD 0.17 0.06 0.07 0.11 0.14 0.18	GMIM 3.53 0.64 0.82 4.87 1.86 3.03
Scenario M2 Unad MB kernelMB hdMB MDABW RCAL1 RCAL2	Bias -1.35 -0.50 -0.35 -0.70 -1.02 -1.20 -1.19	SD 0.36 0.48 0.37 0.69 0.51 0.41 0.41	RMSE 1.40 0.68 0.51 0.98 1.14 1.27 1.26	maxASMD 1.30 0.56 0.74 1.28 1.30 0.93 1.24	meanASMD 0.20 0.07 0.08 0.16 0.20 0.20	medianASMD 0.17 0.06 0.07 0.11 0.14 0.18	GMIM 3.53 0.64 0.82 4.87 1.86 3.03 3.18
Scenario M2 Unad MB kernelMB hdMB MDABW RCAL1 RCAL2 ARB	Bias -1.35 -0.50 -0.35 -0.70 -1.02 -1.20 -1.19 -0.78	SD 0.36 0.48 0.37 0.69 0.51 0.41 0.41 0.45	RMSE 1.40 0.68 0.51 0.98 1.14 1.27 1.26 0.90	maxASMD 1.30 0.56 0.74 1.28 1.30 0.93 1.24 -	meanASMD 0.20 0.07 0.08 0.16 0.15 0.20 0.20	medianASMD 0.17 0.06 0.07 0.11 0.14 0.18 0.18 0.18 -	GMIM 3.53 0.64 0.82 4.87 1.86 3.03 3.18

Table S2: Model misspecification scenarios.

and a nonlinear outcome model

$$Y_i = T_i \left(\sum_{j=1}^{100} X_{ij} + \sum_{j=1}^{50} X_{ij}^2 \right) / 10 + (1 - T_i) \left(\sum_{j=1}^{100} X_{ij} + \sum_{j=1}^{50} X_{ij}^2 \right) / 20 + \epsilon_i.$$

Table S2 gives the results. We observe that MB and hdMB have best performance in Scenario M1, and kernelMB has best performance in Scenario M2. Mahalanobis balancing does not use information of the outcomes, and thus is robust in these scenarios.

B Lemma and Proof

Lemma 1 is similar to Lemma 8 by Athey et al. (2018), but the assumptions are different.

Lemma 1. (Weight Behaviour I) Suppose that $\sum_{\{i:T_i=1\}} w_i^3 / ||w||_2^3 = o_p(1)$ and that the ε_{1i} are sub-Gaussian. Then, as $n \to \infty$,

$$\frac{1}{\|w\|_2} \sum_{\{i:T_i=1\}} w_i \varepsilon_{1i} \stackrel{d}{\to} N\left(0, \sigma^2\right),$$

where $\sigma^2 = Var(\varepsilon_{1i})$.

Proof. Since the MB weights do not utilize the outcomes, the ε_{1i} are independent of the w_i given the X_i . Therefore,

$$E\{\sum_{\{i:T_i=1\}} w_i \varepsilon_{1i} \mid w_i, T_i = 1\} = 0 \text{ and } \operatorname{Var}\{\sum_{\{i:T_i=1\}} w_i \varepsilon_{1i} \mid w_i, T_i = 1\} = \sigma^2 \|w\|_2^2$$

Since the ε_{1i} are sub-Gaussian,

$$E\{\sum_{\{i:T_i=1\}} (w_i \varepsilon_{1i})^3 \mid w_i, T_i = 1\} \le C \sum_{\{i:T_i=1\}} w_i^3$$

for some positive constant C. Therefore,

$$\frac{E\{\sum_{\{i:T_i=1\}} (w_i \varepsilon_{1i})^3 \mid w_i, T_i = 1\}}{Var\{\sum_{\{i:T_i=1\}} w_i \varepsilon_{1i} \mid w_i, T_i = 1\}^{3/2}} = O\left(\sum_{\{i:T_i=1\}} w_i^3 / \|w\|_2^3\right) = o_p(1).$$

Hence, the Lyapunov condition is verified, and the proof is completed by applying Lyapunov's central limit theorem. $\hfill \Box$

The next lemma asserts that the MB weights are stable when the entropy loss function is employed.

Lemma 2. (Weight Behaviour II) Suppose that the entropy loss function is used. Assume that $E\{\exp(a\theta^{\top}W^{1/2}[\Phi(X) - E\{\Phi(X) \mid T = 1\}]) \mid T = 1\} \leq K_a$ for all $\theta \in \Theta$, where K_a is some positive constant, a = 1, 2, 3. Assume that $\theta^{\top}W^{1/2}(\Phi(X) - E\{\Phi(X) \mid T = 1\})$ is sub-Gaussian. Then the following properties hold for the MB weights:

$$\sum_{i:T_i=1} (\hat{w_i}^{MB})^a = O_p(n^{1-a}), \text{ for } a = 1, 2, 3;$$
$$\max_{i:T_i=1} \hat{w_i}^{MB} = O_p(n^{-1/2}).$$

Proof. When the entropy loss function is employed,

$$\begin{split} \hat{w}_{i}^{MB} &= \frac{\exp\left\{\hat{\theta}^{\top}W^{1/2}\Phi(X_{i})\right\}}{\sum_{j:T_{j}=1}\exp\left\{\hat{\theta}^{\top}W^{1/2}\Phi(X_{j})\right\}} \\ &= \frac{\exp\left\{\hat{\theta}^{\top}W^{1/2}\Phi(X_{i})\right\}}{n_{1}E\left[\exp\left\{\hat{\theta}^{\top}W^{1/2}\Phi(X)\right\} \mid T=1\right]}(1+o_{p}(1)) \\ &\leq \frac{\exp\left\{\hat{\theta}^{\top}W^{1/2}\Phi(X_{i})\right\}}{n_{1}\exp\left(\hat{\theta}^{\top}W^{1/2}E\left\{\Phi(X) \mid T=1\right\}\right)}(1+o_{p}(1)) \\ &= \frac{1}{n_{1}}\exp\left(\hat{\theta}^{\top}W^{1/2}\left[\Phi(X_{i}) - E\left\{\Phi(X) \mid T=1\right\}\right]\right)(1+o_{p}(1)) \\ &= O_{p}(n^{-1}). \end{split}$$

Therefore,

$$\begin{split} \sum_{i:T_i=1} (\hat{w}_i^{MB})^a &= \frac{1}{n_1^a} \sum_{i:T_i=1} \exp\left(a\hat{\theta}^\top W^{1/2} \left[\Phi\left(X_i\right) - E\left\{\Phi\left(X\right) \mid T=1\right\}\right]\right) (1+o_p(1)) \\ &= n_1^{1-a} E\left\{\exp(a\theta^\top W^{1/2} \left[\Phi(X) - E\left\{\Phi(X) \mid T=1\right\}\right]\right) \mid T=1\right\} (1+o_p(1)) \\ &= n_1^{1-a} K_a (1+o_p(1)) \\ &= O_p(n^{1-a}). \end{split}$$

Moreover, by the maximal inequality (e.g., Rigollet and Hütter, 2015), we obtain that

$$Pr\left(\max_{i:T_i=1}\frac{1}{n_1}\exp\left(\hat{\theta}^{\top}W^{1/2}(\Phi(X_i) - E\{\Phi(X) \mid T=1\})\right) \ge \frac{1}{n_1}\exp(t)\right) \le n_1\exp\left(-\frac{t^2}{2\sigma^2}\right).$$

Set $t = \sqrt{2}\sigma\log(n_1^{1/2}s)$, we have $\max_{i:T_i=1}\hat{w}_i^{MB} = O_p(n^{-1/2}).$

The following lemma shows the asymptotic property for the solution $\hat{\theta}$ of the MB dual problem.

Lemma 3. Assume that $1/\pi(X) = C * \tilde{f}^{*'}(\theta^{*\top}W^{1/2}(\Phi(X) - \bar{\Phi}))$ for some $\theta^* \in int(\Theta)$ and some constant C for all X. Suppose that Assumption 3 holds. Then $\hat{\theta}$ is consistent for θ^* . Moreover, $\hat{\theta}$ is asymptotically normal.

Proof. The first order optimality condition for the dual problem is

$$\sum_{i=1}^{n} T_{i} \tilde{f}^{*'} \left(\hat{\theta}^{\top} W^{1/2} \left(\Phi \left(X_{i} \right) - \bar{\Phi} \right) \right) \left(\Phi_{j} \left(X_{i} \right) - \bar{\Phi}_{j} \right) + \sqrt{\delta} \frac{\hat{\theta}_{j}}{||\hat{\theta}||_{2}} = 0, \quad j = 1, \cdots, K,$$

where $\Phi_j(X_i)$, $\bar{\Phi}_j$, and $\hat{\theta}_j$ are the *j*th components of $\Phi(X_i)$, $\bar{\Phi}$, and $\hat{\theta}$, respectively. Write

$$\frac{1}{n}\sum_{i=1}^{n}\psi(X_{i},T_{i};\theta) = \frac{1}{n}\sum_{i=1}^{n}T_{i}\tilde{f}^{*'}\left(\theta^{\top}W^{1/2}\left(\Phi\left(X_{i}\right) - E\{\Phi\left(X\right)\}\right)\right)\left(\Phi\left(X_{i}\right) - E\{\Phi\left(X\right)\}\right),$$

which is a set of K estimating functions. Note that

$$E \{ \psi(X_i, T_i; \theta) \} = E \{ E (\psi(X, T; \theta)) \mid X \}$$

= $E \left[\pi(X) \tilde{f}^{*'} \left(\theta^\top W^{1/2} \left(\Phi (X) - E \{ \Phi (X) \} \right) \right) W^{1/2} (\Phi (X) - E \{ \Phi (X) \}) \right].$

For the above conditional expectation to be zero, it must be true that

$$\pi(X)\tilde{f}^{*'}\left(\theta^{\top}W^{1/2}\left(\Phi\left(X\right)-E\{\Phi\left(X\right)\}\right)\right)$$

is a constant for any X. By the assumption that $1/\pi(X) = C * \tilde{f}^{*'}(\theta^{*\top}W^{1/2}(\Phi(X) - \bar{\Phi}))$, we obtain that θ^* is the unique solution of $E\{\psi(X_i, T_i; \theta)\} = 0$. Therefore, by the estimating equation theory (Van der Vaart, 1998), the solution of the estimating equations

$$\frac{1}{n}\sum_{i=1}^{n}\psi(X_i, T_i; \theta) = 0,$$

denoted by $\tilde{\theta}$, is asymptotically consistent for θ^* . Moreover, by the assumption that $\delta = o(n)$, we obtain $\frac{1}{n}\sqrt{\delta}\frac{\theta_j}{||\theta||_2} = o_p(n^{-1/2})$ for any $\theta \in int(\Theta)$. Therefore, the difference between $\hat{\theta}$ and $\tilde{\theta}$ is asymptotically negligible, and thus $\hat{\theta}$ is asymptotically consistent for θ^* . Furthermore, by Taylor expansion, we obtain that as $n \to \infty$,

$$\sqrt{n}(\hat{\theta} - \theta^*) \xrightarrow{d} N(0, \Sigma),$$

where
$$\Sigma = \left\{ E(\frac{\partial \psi}{\partial \theta^{\top}}) \right\}^{-1} E(\psi \psi^{\top}) \left\{ E(\frac{\partial \psi}{\partial \theta}) \right\}^{-1}$$
.

The following lemma asserts that the MB weight is close to the unknown inverse probability weight. The proof is similar to arguments by Lee et al. (2023).

Lemma 4. (Weight Behaviour III) Assume that $1/\pi(X) = C * \tilde{f}^{*'}(\theta^{*\top}W^{1/2}(\Phi(X) - \bar{\Phi}))$ for some $\theta^* \in int(\Theta)$ and some constant C for all X. Suppose that Assumption 3 holds. Then the following property holds:

$$n\hat{w}_i^{MB} = \frac{1}{\pi(X_i)} + O_p(n^{-\frac{1}{2}}).$$

Proof.

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^{n} T_i \tilde{f}^{*'} \left(\hat{\theta}^\top W^{1/2} \left(\Phi\left(X_i\right) - \bar{\Phi} \right) \right) &= \frac{1}{n} \sum_{i=1}^{n} T_i \tilde{f}^{*'} \left(\hat{\theta}^\top W^{1/2} \left(\Phi\left(X_i\right) - E\{\Phi\left(X\right)\} \right) \right) + O_p(n^{-\frac{1}{2}}) \\ &= \frac{1}{n} \sum_{i=1}^{n} T_i \tilde{f}^{*'} \left(\theta^{*\top} W^{1/2} \left(\Phi\left(X_i\right) - E\{\Phi\left(X\right)\} \right) \right) + O_p(n^{-\frac{1}{2}}) \\ &= \frac{1}{n} \sum_{i=1}^{n} \frac{T_i}{C\pi(X_i)} + O_p(n^{-\frac{1}{2}}) \\ &= \frac{1}{C} + o_p(1). \end{aligned}$$

Therefore,

$$n\hat{w}_{i}^{MB} = \frac{\tilde{f}^{*'}\left(\hat{\theta}^{\top}W^{1/2}\left(\Phi\left(X_{i}\right) - \bar{\Phi}\right)\right)}{\frac{1}{n}\sum_{i:T_{i}=1}\tilde{f}^{*'}\left(\hat{\theta}^{\top}W^{1/2}\left(\Phi\left(X_{i}\right) - \bar{\Phi}\right)\right)}$$
$$= \frac{\tilde{f}^{*'}\left(\theta^{*\top}W^{1/2}\left(\Phi\left(X_{i}\right) - E\{\Phi\left(X\right)\}\right)\right) + O_{p}(n^{-1/2})}{1/C + o_{p}(1)}$$
$$= \frac{1}{\pi(X_{i})} + O_{p}(n^{-\frac{1}{2}}).$$

B.1 Proof for Theorem 1

Proof. We utilize the Fenchel duality theorem (Mohri et al., 2018, Theorem B.39). Without loss of generability, suppose that subjects $i = 1, \dots, n_1$ are in the treated group, and

subjects $i = n_1 + 1, \dots, n$ are in the control group. The primal problem (7) is

$$\min_{w \in \mathbb{R}^{n_1}} : \sum_{i=1}^{n_1} \tilde{f}(w_i) + I_{\mathcal{C}}\left(\sum_{i=1}^{n_1} w_i W^{1/2}(\Phi(X_i) - \bar{\Phi})\right),$$

where $w = (w_1, \dots, w_{n_1})^{\top}$. Let $F(w) = \sum_{i=1}^{n_1} \tilde{f}(w_i)$. The conjugate function of F is given by

$$F^*(w) = \sup_{v} \left(\sum_{i=1}^{n_1} w_i v_i - F(v) \right)$$
$$= \sup_{v} \sum_{i=1}^{n_1} \left(w_i v_i - \tilde{f}(v_i) \right)$$
$$= \sum_{i=1}^{n_1} \sup_{v_i} (w_i v_i - \tilde{f}(v_i))$$
$$= \sum_{i=1}^{n_1} \tilde{f}^*(w_i).$$

Note that if the normalization constraint is added to the primal problem, then the conjugate function F^* is no longer additive. Let $g(\theta) = I_{\mathcal{C}}(\theta)$, for any $\theta \in \mathbb{R}^K$. The conjugate function of g is given by

$$g^{*}(\theta) = \sup_{u} \left(\sum_{k=1}^{K} \theta_{k} u_{k} - I_{\mathcal{C}}(u) \right)$$
$$= \sup_{\|u\|_{2} \le \sqrt{\delta}} \left(\sum_{k=1}^{K} \theta_{k} u_{k} \right)$$
$$= \sup_{\|u\|_{2} \le \sqrt{\delta}} \left(\|\theta\|_{2} \|u\|_{2} \right)$$
$$= \sqrt{\delta} \|\theta\|_{2}.$$

Define the mapping $A : \mathbb{R}^{n_1} \to \mathbb{R}^K$ such that $Aw = \sum_{i=1}^{n_1} w_i W^{1/2} (\Phi(X_i) - \overline{\Phi})$. Then A is a bounded linear map. Let A^* be the adjoint operator of A. Then for all $\theta = (\theta_1, \cdots, \theta_K)^\top \in \mathbb{R}^K$,

$$A^*\theta = (\sum_{k=1}^K \theta_k W^{1/2} (\Phi_k(X_1) - \bar{\Phi}_k), \cdots, \sum_{k=1}^K \theta_k W^{1/2} (\Phi_k(X_{n_1}) - \bar{\Phi}_k))^\top$$
$$= (\theta^\top W^{1/2} (\Phi(X_1) - \bar{\Phi}), \cdots, \theta^\top W^{1/2} (\Phi(X_{n_1}) - \bar{\Phi}))^\top.$$

Define $\theta_0 = Aw_0 = \sum_{i=1}^{n_1} W^{1/2} (\Phi(X_i) - \overline{\Phi})/n_1^a$, where $w_0 = (1/n_1^a, \cdots, 1/n_1^a)^\top$. Here, we choose *a* to be sufficiently large such that $\|\theta_0\|_2 < \sqrt{\delta}$. Since each component of

 w_0 is non-negative, $w_0 \in dom(F)$, where dom(F) denotes the domain of F. Therefore, $\theta_0 \in A(dom(F))$. Since $\|\theta_0\|_2 < \sqrt{\delta}$, we obtain that $g(\theta_0) = 0$ and g is continuous at θ_0 . Therefore, $\theta_0 \in A(dom(F)) \cap cont(g)$, implying that $A(dom(F)) \cap cont(g) \neq \emptyset$, where cont(g) is the set of continuous points of g. Therefore, the strong duality condition of the Fenchel duality theorem is verified. Moreover,

$$F(A^*\theta) + g^*(-\theta) = \sum_{i=1}^{n_1} \tilde{f}^*(\theta^\top W^{1/2}(\Phi(X_i) - \bar{\Phi})) + \sqrt{\delta} \|\theta\|_2.$$

The Fenchel duality theorem leads to that

$$\begin{array}{ll} \underset{w \in \mathbb{R}^{n_1}}{\text{minimize}} : & \sum_{i=1}^{n_1} \tilde{f}(w_i) + I_{\mathcal{C}} \left(\sum_{i=1}^{n_1} w_i W^{1/2}(\Phi(X_i) - \bar{\Phi}) \right) \\ = \underset{\theta \in \mathbb{R}^{K}}{\text{minimize}} : & \sum_{i=1}^{n_1} \tilde{f}^* \left(\theta^\top W^{1/2}(\Phi(X_i) - \bar{\Phi}) \right) + \sqrt{\delta} \|\theta\|_2. \end{array}$$

Furthermore, since the strong duality condition holds, equality of the Fenchel's inequality (Mohri et al., 2018, Prop. 38) holds. leading to that $A^*\hat{\theta}$ is a subgradient of F at \hat{w} . We consider the simpler situation that the loss function $f(\cdot)$ is differentiable and it has non-negative domain as in (Josey et al., 2021). Then, $A^*\hat{\theta} = F'(\hat{w})$, or equivalently,

$$\hat{\theta}^\top W^{1/2}(\Phi(X_i) - \bar{\Phi}) = \tilde{f}'(\hat{w}_i).$$

Therefore, $\hat{w}_i = (\tilde{f}')^{-1} (\hat{\theta}^\top W^{1/2} \left(\Phi(X_i) - \bar{\Phi} \right)) = (\tilde{f}^*)' (\hat{\theta}^\top W^{1/2} \left(\Phi(X_i) - \bar{\Phi} \right)), i = 1, \cdots, n_1,$ where the second equality holds by Fenchel's identity (Ryu and Yin, 2022; Mordukhovich and Nam, 2022).

	-	

B.2 Proof for Theorem 2

Proof. We only need to show that $\hat{\tau}_1^{MB} = \tau_1 + O_p(n^{-1/2})$, where $\hat{\tau}_1^{MB} = \sum_{i:T_i=1} \hat{w}_i^{MB} Y_i$. First, assume the propensity score satisfies that $1/\pi(X) = C * \tilde{f}^{*'}(\theta^{*\top}W^{1/2}(\Phi(X) - \bar{\Phi}))$ for some $\theta^* \in int(\Theta)$ and some constant C. By Lemma 3,

$$\sum_{i:T_i=1} \hat{w}_i^{MB} Y_i = \sum_{i:T_i=1} \left(\hat{w}_i^{MB} - \frac{1}{n\pi(X_i)} \right) Y_i + \sum_{i:T_i=1} \frac{Y_i}{n\pi(X_i)}$$
$$= \sum_{i:T_i=1} O_p(n^{-3/2}) O_p(1) + O_p(n^{-1/2}) + \tau_1.$$

Therefore, $\hat{\tau}_1 - \tau_1 = O_p(n^{-1/2}).$

Second, we assume that the conditional potential outcome $E\{Y(1)|X\} = \beta^{\top}W^{1/2}\Phi(X)$ with parameter β , $\sum_{i:T_i=1} (\hat{w_i}^{MB})^3 / \|\hat{w}^{MB}\|_2^3 = o_p(1)$, and $\|\hat{w}^{MB}\|_2^2 = O_p(n^{-1})$. Note that the following decomposition holds:

$$\sum_{i:T_i=1} \hat{w_i}^{MB} Y_i - \tau_1 = \beta^\top \sum_{i:T_i=1} \hat{w_i}^{MB} W^{1/2} (\Phi(X_i) - \bar{\Phi}) + \sum_{i:T_i=1} \hat{w_i}^{MB} \epsilon_{1i} + \frac{1}{n} \sum_{i=1}^n \beta^\top W^{1/2} \Phi(X_i) - \tau_1.$$

Let the \hat{w}_i be the unnormalized MB weights. By Assumption 3.4,

$$\sum_{i:T_i=1} \hat{w}_i = \sum_{i:T_i=1} \exp\left(\theta^\top W^{1/2}\left(\Phi(X_i) - \bar{\Phi}\right)\right) = O_p(n).$$

Therefore,

$$\begin{aligned} |\beta^{\top} \sum_{i:T_{i}=1} \hat{w}_{i}^{MB} W^{1/2}(\Phi(X_{i}) - \bar{\Phi})| &\leq \|\beta\|_{2} \|\sum_{i:T_{i}=1} \hat{w}_{i}^{MB} W^{1/2}(\Phi(X_{i}) - \bar{\Phi})\|_{2} \\ &= \|\beta\|_{2} \|\sum_{i:T_{i}=1} \hat{w}_{i} W^{1/2}(\Phi(X_{i}) - \bar{\Phi})\|_{2} / \sum_{i:T_{i}=1} \hat{w}_{i} \\ &\leq \|\beta\|_{2} \sqrt{\delta} / \sum_{i:T_{i}=1} \hat{w}_{i} \\ &= o_{p}(n^{-1/2}). \end{aligned}$$

By Lemma 1, we have $\frac{1}{\|\hat{w}^{MB}\|_2} \sum_{i:T_i=1} \hat{w}_i^{MB} \epsilon_{1i} = O_p(1)$. Therefore, by the assumption that $\|\hat{w}^{MB}\|_2^2 = O_p(n^{-1})$, we obtain $\sum_{i:T_i=1} \hat{w}_i^{MB} \epsilon_{1i} = O_p(\|\hat{w}^{MB}\|_2) = O_p(n^{-1/2})$. Finally, $\frac{1}{n} \sum_{i=1}^n \beta^\top W^{1/2} \Phi(X_i) - \tau_1 = O_p(n^{-1/2})$. Therefore, $\hat{\tau}_1 - \tau_1 = O_p(n^{-1/2})$. The double robustness property is proved.

B.3 Proof for Theorem 3

Proof. By the assumption for $\theta^{\top} W^{1/2} (E\{\Phi(X_i)\} - E\{\Phi(X_i) \mid T = 1\})$, we obtain $\exp(\theta^{\top} W^{1/2} (E\{\Phi(X_i)\} - E\{\Phi(X_i) \mid T = 1\})) = O_p(n^{\gamma})$. It follows that

Therefore,

$$|\beta^{\top} \sum_{i:T_i=1} \hat{w_i}^{MB} W^{1/2}(\Phi(X_i) - \bar{\Phi})| \le \|\beta\|_2 \|\sum_{i:T_i=1} \hat{w_i}^{MB} W^{1/2}(\Phi(X_i) - \bar{\Phi})\|_2 \le O_p(n^{s+\gamma+\alpha}).$$

By Lemmas 1 and 2, $\sum_{i:T_i=1} \hat{w}_i^{MB} \epsilon_{1i} = O_p(\|\hat{w}^{MB}\|_2) = O_p(n^{-1/2})$. Here, the condition $\sum_{\{i:T_i=1\}} w_i^3 / \|w\|_2^3 = o_p(1)$ holds. Moreover,

$$\frac{1}{n}\sum_{i=1}^{n}\beta^{\top}W^{1/2}\Phi(X_i) - \tau_1 = \frac{1}{n}\sum_{i=1}^{n}\mu_1(X_i) - \tau_1 = O_p(n^{-1/2}).$$

Therefore,

$$\begin{aligned} \hat{\tau_1}^{MB} &- \tau_1 &= \beta^\top \sum_{i:T_i=1} \hat{w}_i^{MB} W^{1/2} (\Phi(X_i) - \bar{\Phi}) + \sum_{i:T_i=1} \hat{w}_i^{MB} \epsilon_{1i} \\ &+ \frac{1}{n} \sum_{i=1}^n \beta^\top W^{1/2} \Phi(X_i) - \tau_1 \\ &= O_p(n^{-1/2}) + O_p(n^{s+\gamma+\alpha}). \end{aligned}$$

B.4 Proof of Theorem 4

Proof. To simplify the notations, let $\pi_i = \pi(X_i)$. Write $E\{Y(1)|X\} = \mu_1(X) = \beta^{\top} W^{1/2} \Phi(X)$ with parameter β . Then the following decomposition holds:

$$\begin{aligned} \hat{\tau}_1 - \tau_1 &= \sum_{i=1}^n T_i \hat{w}_i^{MB} Y_i - \tau_1 \\ &= \sum_{i=1}^n T_i \left(\hat{w}_i^{MB} - \frac{1}{n\pi_i} \right) \left(Y_i - \mu_1 \left(X_i \right) \right) + \sum_{i=1}^n \frac{T_i}{n\pi_i} \left(Y_i - \mu_1 \left(X_i \right) \right) \\ &+ \sum_{i=1}^n \left(T_i \hat{w}_i^{MB} - \frac{1}{n} \right) \beta^\top W^{1/2} \Phi \left(X_i \right) + \left(\frac{1}{n} \sum_{i=1}^n \mu_1 (X_i) - \tau_1 \right) \\ &= A_1 + R_1 + A_2 + R_2. \end{aligned}$$

Firstly, following the proof of theorem 2 in Wang and Zubizarreta (2020), we show that

$$\sup_{x \in \mathcal{X}} |\hat{w}_i^{MB} - \frac{1}{n\pi_i}| = o_p(1).$$

Lemma 5. There exists a global minimizer $\hat{\theta}$ such that

$$\left\|\hat{\theta} - \theta_{true}\right\|_2 = O_p\left(n^{-\frac{1}{4}+\epsilon}\right) \text{ for arbitrary } \epsilon > 0$$

for the objective function $G(\cdot)$:

$$G(\theta) := \sum_{i=1}^{n} T_i \tilde{f}^* \left(\theta^\top W^{1/2} \left(\Phi(X_i) - \bar{\Phi} \right) \right) + \delta^{\frac{1}{2}} ||\theta||_2,$$

To show that a minimizer Δ^* of $G(\theta_{true} + \Delta)$ exists in

$$\mathcal{C} = \left\{ \Delta \in \mathcal{R}^K : \|\Delta\|_2 \le C n^{s-1+\epsilon} \right\}$$

for some constant C, it suffices to show that

$$E\left\{\inf_{\Delta\in\mathcal{C}}G\left(\theta_{true}+\Delta\right)-G\left(\theta_{true}\right)>0\right\}\to 1, \text{ as } n\to\infty$$
(S1)

by the continuity of $G(\cdot)$.

To show (S1), we use mean value theorem: for some $\tilde{\theta}$ between θ_{true} and $\hat{\theta}$,

$$\begin{aligned} G\left(\theta_{true} + \Delta\right) &- G\left(\theta_{true}\right) \\ &= \sum_{i=1}^{n} T_{i} \left(\tilde{f}^{*} \left((\theta_{true} + \Delta)^{\top} W^{1/2} \left(\Phi\left(X_{i}\right) - \bar{\Phi}\right) \right) - \tilde{f}^{*} \left(\theta_{true}^{\top} W^{1/2} \left(\Phi\left(X_{i}\right) - \bar{\Phi}\right) \right) \right) \\ &+ \delta^{\frac{1}{2}} ||\theta_{true} + \Delta||_{2} - \delta^{\frac{1}{2}} ||\theta_{true}||_{2} \quad \left(||\theta_{true} + \Delta||_{2} - ||\theta_{true}||_{2} \ge -||\Delta||_{2} \right) \\ &\geq \Delta \cdot \sum_{i=1}^{n} T_{i} \tilde{f}^{*'} \left((\theta_{true} + \Delta)^{\top} W^{1/2} \left(\Phi\left(X_{i}\right) - \bar{\Phi} \right) \right) \left(\Phi\left(X_{i}\right) - \bar{\Phi} \right) \\ &+ \frac{1}{2} \sum_{i=1}^{n} T_{i} \tilde{f}^{*''} \left(\theta_{true}^{\top} W^{1/2} \left(\Phi\left(X_{i}\right) - \bar{\Phi} \right) \right) \left(\left(\Phi\left(X_{i}\right) - \bar{\Phi} \right)^{\top} \Delta \right)^{2} \\ &+ \frac{1}{6} \sum_{i=1}^{n} T_{i} \tilde{f}^{*'''} \left(\tilde{\theta}_{true}^{\top} W^{1/2} \left(\Phi\left(X_{i}\right) - \bar{\Phi} \right) \right) \left(\left(\Phi\left(X_{i}\right) - \bar{\Phi} \right)^{\top} \Delta \right)^{3} \\ &= \delta^{\frac{1}{2}} ||\Delta|| \quad (We treat \left(\Phi\left(X_{i}\right) - \bar{\Phi} \right)^{\top} \Delta \text{ as a univariate variable and apply mean way and ap$$

$$\begin{split} &-\delta^{\frac{1}{2}}||\Delta||_{2} \quad (\text{We treat } \left(\Phi\left(X_{i}\right)-\bar{\Phi}\right)^{\top}\Delta\text{ as a univariate variable and apply mean value theorem.})\\ &\geq \frac{1}{2}\sum_{i=1}^{n}T_{i}\tilde{f}^{*''}\left(\theta_{true}^{\top}W^{1/2}\left(\Phi\left(X_{i}\right)-\bar{\Phi}\right)\right)\left(\left(\Phi\left(X_{i}\right)-\bar{\Phi}\right)^{\top}\Delta\right)^{2}\\ &= \frac{1}{2}\sum_{i=1}^{n}T_{i}\tilde{f}^{*''}\left(\theta_{true}^{\top}W^{1/2}\left(\Phi\left(X_{i}\right)-\bar{\Phi}\right)\right)\left(\left(\Phi\left(X_{i}\right)-\bar{\Phi}\right)^{\top}\Delta\right)^{2}\\ &+ \frac{1}{6}\sum_{i=1}^{n}T_{i}\tilde{f}^{*''}\left(\tilde{\theta}_{true}^{\top}W^{1/2}\left(\Phi\left(X_{i}\right)-\bar{\Phi}\right)\right)\left(\left(\Phi\left(X_{i}\right)-\bar{\Phi}\right)^{\top}\Delta\right)^{3}\\ &- \left(\delta^{\frac{1}{2}}+||\sum_{i=1}^{n}T_{i}\tilde{f}^{*'}\left(\theta_{true}^{\top}W^{1/2}\left(\Phi\left(X_{i}\right)-\bar{\Phi}\right)\right)\left(\Phi\left(X_{i}\right)-\bar{\Phi}\right)||_{2}\right)||\Delta||_{2}. \quad (\text{Cauchy inequality}) \end{split}$$

Recall that the Bernstein's inequality for random matrices in Tropp et al. (2015) says the following. Let $\{M_k\}$ be a sequence of independent random matrices with dimensions $d_1 \times d_2$. Assume that $E(M_k) = 0$ and $||M_k||_2 \leq R_n$ almost surely. Define

$$\sigma_n^2 = \max\left\{ \left\| \sum_{k=1}^n E\left(M_k M_k^{\mathsf{T}}\right) \right\|_2, \left\| \sum_{k=1}^n E\left(M_k^{\mathsf{T}} M_k\right) \right\|_2 \right\}$$

Then for all $t \ge 0$,

$$\operatorname{pr}\left(\left\|\sum_{k=1}^{n} M_{k}\right\|_{2} \ge t\right) \le (d_{1}+d_{2}) \exp\left(-\frac{t^{2}/2}{\sigma_{n}^{2}+R_{n}t/3}\right).$$

Let $M_i = \frac{T_i}{\pi(X_i)} \left(\Phi(X_i) - \overline{\Phi} \right)$. We derive that

$$E(M_{i}) = E\left(\frac{T_{i}}{\pi(X_{i})}\left(\Phi(X_{i}) - \bar{\Phi}\right)\right) = E\left(E\left(\frac{T_{i}}{\pi(X_{i})}\left(\Phi(X_{i}) - \bar{\Phi}\right) \mid X_{i}\right)\right) = 0,$$

$$||M_{i}||_{2} = ||\frac{T_{i}}{\pi(X_{i})}\left(\Phi(X_{i}) - \bar{\Phi}\right)||_{2} \le ||\frac{T_{i}}{\pi(X_{i})}||_{2}||\left(\Phi(X_{i}) - \bar{\Phi}\right)||_{2} \le \frac{C_{2}}{\eta^{1/2}}n^{1/2},$$

$$E\left(||\sum_{i=1}^{n} M_{i}M_{i}^{\top}||_{2}\right) = E\left(||\sum_{i=1}^{n} \frac{T_{i}}{\pi(X_{i})^{2}}\left(\Phi(X_{i}) - \bar{\Phi}\right)\left(\Phi(X_{i}) - \bar{\Phi}\right)^{\top}||_{2}\right)$$

$$\le \sup_{x \in \mathcal{X}} \frac{T_{i}}{\pi(X_{i})^{2}}E\left(||\sum_{i=1}^{n} \left(\Phi(X_{i}) - \bar{\Phi}\right)\left(\Phi(X_{i}) - \bar{\Phi}\right)^{\top}||_{2}\right) = \frac{n}{\eta^{2}}E\left(||\left(\Phi(X_{i}) - \bar{\Phi}\right)\left(\Phi(X_{i}) - \bar{\Phi}\right)^{\top}||_{2}\right)$$

$$\le \frac{C_{2}n}{\eta^{2}},$$

$$E\left(||\sum_{i=1}^{n} M_{i}^{\top}M_{i}||_{2}\right) = E\left(||\sum_{i=1}^{n} \frac{T_{i}}{\pi(X_{i})^{2}}\left(\Phi(X_{i}) - \bar{\Phi}\right)^{\top}\left(\Phi(X_{i}) - \bar{\Phi}\right)||_{2}\right)$$

$$\le \frac{n}{\eta^{2}}E\left(||\left(\Phi(X_{i}) - \bar{\Phi}\right)^{\top}\left(\Phi(X_{i}) - \bar{\Phi}\right)||_{2}\right) \le \frac{C_{2}n^{3/2}}{\eta^{2}}.$$

Therefore,

$$\sigma_n^2 = \max\left\{ \left\| \sum_{i=1}^n E\left(M_i M_i^{\top} \right) \right\|_2, \left\| \sum_{i=1}^n E\left(M_i^{\top} M_i \right) \right\|_2 \right\} = \frac{C_2 n^{3/2}}{\eta^2},$$

and

$$\Pr\left(\left\|\sum_{i=1}^{n} M_{i}\right\|_{2} \ge t\right) \le (1+K) \exp\left(-\frac{t^{2}/2}{\frac{C_{2}n^{3/2}}{\eta^{2}} + C_{2}n^{1/2}t/3}\right).$$

Letting $t = O_p(n^{3/4+\epsilon})$ for arbitrary $\epsilon > 0$, we have

$$\begin{aligned} &||\sum_{i=1}^{n} T_{i}\tilde{f}^{*'}\left(\theta_{true}^{\top}W^{1/2}\left(\Phi\left(X_{i}\right)-\bar{\Phi}\right)\right)\left(\Phi\left(X_{i}\right)-\bar{\Phi}\right)||_{2} \\ &= ||\sum_{i=1}^{n} \frac{T_{i}}{\pi(X_{i})}\left(\Phi\left(X_{i}\right)-\bar{\Phi}\right)||_{2} \le C_{3}n. \end{aligned}$$

We have

$$\sum_{i=1}^{n} T_{i} \tilde{f}^{*''} \left(\theta_{true}^{\top} W^{1/2} \left(\Phi \left(X_{i} \right) - \bar{\Phi} \right) \right) \left(\left(\Phi \left(X_{i} \right) - \bar{\Phi} \right)^{\top} \Delta \right)^{2} \\ \geq \frac{1}{\eta} \sum_{i=1}^{n} \left(\frac{T_{i}}{\pi(X_{i})} \left(\Phi \left(X_{i} \right) - \bar{\Phi} \right)^{\top} \Delta \right)^{2} \\ = \frac{1}{\eta} n \Delta^{\top} \frac{1}{n} \sum_{i=1}^{n} \left(\frac{T_{i}}{\pi(X_{i})} \left(\Phi \left(X_{i} \right) - \bar{\Phi} \right) \right) \left(\frac{T_{i}}{\pi(X_{i})} \left(\Phi \left(X_{i} \right) - \bar{\Phi} \right) \right)^{\top} \Delta.$$

According to Section 3.3.1 in Lei et al. (2018), the smallest eigenvalue λ_1 of matrix $\frac{1}{n} \sum_{i=1}^{n} M_i M_i^{\top}$ satisfies that $\lambda_1 = O_p(\frac{1}{\log(n)^{\gamma}})$ for some $\gamma > 0$. Then,

$$n\Delta^{\top} \frac{1}{n} \sum_{i=1}^{n} \left(\frac{T_{i}}{\pi(X_{i})} \left(\Phi\left(X_{i}\right) - \bar{\Phi}\right)\right) \left(\frac{T_{i}}{\pi(X_{i})} \left(\Phi\left(X_{i}\right) - \bar{\Phi}\right)\right)^{\top} \Delta \ge n\lambda_{1} ||\Delta||_{2}^{2} = O_{p}\left(\frac{n}{\log(n)^{\gamma}}\right) ||\Delta||_{2}^{2}.$$
(S2)

Finally, we have

$$\frac{1}{6} \sum_{i=1}^{n} T_{i} \tilde{f}^{*'''} \left(\tilde{\theta}_{true}^{\top} W^{1/2} \left(\Phi \left(X_{i} \right) - \bar{\Phi} \right) \right) \left(\Delta^{\top} \left(\Phi \left(X_{i} \right) - \bar{\Phi} \right) \right)^{3} \\
\geq -\frac{1}{6} \sum_{i=1}^{n} T_{i} \tilde{f}^{*'''} \left(\tilde{\theta}_{true}^{\top} W^{1/2} \left(\Phi \left(X_{i} \right) - \bar{\Phi} \right) \right) ||\Delta||_{2}^{3} || \left(\Phi \left(X_{i} \right) - \bar{\Phi} \right) ||_{2}^{3} \\
\geq -\frac{1-\eta}{6} ||\Delta||_{2}^{3} \sum_{i=1}^{n} || \left(\Phi \left(X_{i} \right) - \bar{\Phi} \right) ||_{2}^{3} \\
= -\frac{1-\eta}{6} ||\Delta||_{2}^{3} O_{p} (K^{3/2} n^{-3/2}).$$
(S3)

Using (S2) and (S3), we conclude that

$$\begin{aligned} &G\left(\theta_{true} + \Delta\right) - G\left(\theta_{true}\right) \\ \geq &\frac{1}{\eta} n\lambda_1 ||\Delta||_2^2 - (\delta^{\frac{1}{2}} + O_p(n^{3/4+\epsilon}))||\Delta||_2 - \frac{1-\eta}{6} ||\Delta||_2^3 O_p(K^{3/2}n^{-3/2}) \\ = &O_p(\frac{n}{\log(n)^{\gamma}})||\Delta||_2^2 - O_p(n^{3/4+\epsilon})||\Delta||_2 - ||\Delta||_2^3 \\ = &O_p(\frac{n^{\frac{1}{2}+2\epsilon}}{\log(n)^{\gamma}}) - O_p(n^{1/2}) - O_p(n^{-3/4+3\epsilon}) \\ \geq &O_p(n^{\frac{1}{2}+1.5\epsilon}) - O_p(n^{1/2}) - O_p(n^{-3/4+3\epsilon}). \quad (\text{ since } O_p(n^{0.5\epsilon}/\log(n)^{\gamma} \to \infty \text{ for arbitrary } \epsilon > 0) \end{aligned}$$

For $\epsilon > 0$ such that $\frac{1}{2} + 1.5\epsilon > -3/4 + 3\epsilon$, Lemma 5 is proved.

Hence, we conclude that

$$\begin{split} \sup_{x \in \mathcal{X}} \left| \tilde{f}^{*'} \left(\hat{\theta}^{\top} W^{1/2} \left(\Phi \left(X_{i} \right) - \bar{\Phi} \right) \right) - \frac{1}{\pi(X_{i})} \right| \\ &= \sup_{x \in \mathcal{X}} \left| \tilde{f}^{*'} \left(\hat{\theta}^{\top} W^{1/2} \left(\Phi \left(X_{i} \right) - \bar{\Phi} \right) \right) - \tilde{f}^{*'}(m^{*}(X_{i})) \right| \\ &\leq \sup_{x \in \mathcal{X}} \left| \tilde{f}^{*'} \left(\hat{\theta}^{\top} W^{1/2} \left(\Phi \left(X_{i} \right) - \bar{\Phi} \right) \right) - \tilde{f}^{*'} \left(\theta^{\top}_{true} W^{1/2} \left(\Phi \left(X_{i} \right) - \bar{\Phi} \right) \right) \right| \\ &+ \sup_{x \in \mathcal{X}} \left| \tilde{f}^{*'} \left(\theta^{\top}_{true} W^{1/2} \left(\Phi \left(X_{i} \right) - \bar{\Phi} \right) \right) - \tilde{f}^{*'}(m^{*}(X_{i})) \right| \\ &= O\{ \sup_{x \in \mathcal{X}} \left| \left(\theta_{true} - \hat{\theta} \right)^{\top} W^{1/2} \left(\Phi \left(X_{i} \right) - \bar{\Phi} \right) \right\} + o_{p}(1) \\ &\leq O\left\{ \sup_{x \in \mathcal{X}} \left| \left(\theta_{true} - \hat{\theta} \right) \right| _{2} \right| |W^{1/2} \left(\Phi \left(X_{i} \right) - \bar{\Phi} \right) ||_{2} \right\} + o_{p}(1) \\ &= O_{p}\left\{ n^{s - \frac{1}{2} + \epsilon} \right\} + o_{p}(1) \\ &= o_{p}(1). \end{split}$$

Therefore, we prove that $\sup_{x \in \mathcal{X}} |\hat{w}_i - \frac{1}{\pi(X_i)}| = o_p(1).$

Following the empirical process arguments by Wang and Zubizarreta (2020) and Fan et al. (2022), it holds that $A_1 = o_p(n^{-1/2})$. Next,

$$A_{2} = \sum_{i=1}^{n} T_{i} \hat{w}_{i}^{MB} \left(\beta^{\top} W^{1/2} \Phi(X_{i}) - \frac{1}{n} \sum_{i=1}^{n} \beta^{\top} W^{1/2} \Phi(X_{i}) \right)$$

$$\leq \|\beta\|_{2} \|\hat{w}_{i}^{MB} W^{1/2} \left(\Phi(X_{i}) - \bar{\Phi} \right) \|_{2}$$

$$\leq O_{p}(n^{s+\gamma+\alpha}) = o_{p}(n^{-1/2}).$$

The R_1 and R_2 are regular and asymptotically linear, and they determine the asymptotic expansion of $\hat{\tau}_1 - \tau_1$. Similar expansion holds for $\hat{\tau}_0 - \tau_0$. Using these asymptotic expansions, it follows that the semiparametric efficiency bound for ATE estimation is attained.

C Additional Information for Numerical Studies

We provide more information about the numerical studies in the main article. In particular, we display the results of MB and MB2 by varying (δ_1, δ_0) , where δ_1 and δ_0 are the threshold parameters for the treated and control groups, respectively. The outputs show that both

Scenario A	Bias	\mathbf{SD}	RMSE	maxASMD	meanASMD	medASMD	GMIM
$\delta_1 = \delta_0 = 1$							
\mathbf{MB}	-0.25	3.16	3.17	0.11	0.01	0.01	0.00
MB2	-0.24	3.16	3.17	0.11	0.01	0.01	0.00
$\delta_1 = \delta_0 = 10^{-2}$							
\mathbf{MB}	-0.11	3.21	3.21	0.05	0.00	0.00	0.00
MB2	-0.11	3.21	3.21	0.05	0.00	0.00	0.00
$\delta_1 = \delta_0 = 10^{-4}$							
\mathbf{MB}	-0.10	3.22	3.22	0.03	0.00	0.00	0.00
$\mathbf{MB2}$	-0.10	3.22	3.22	0.03	0.00	0.00	0.00
$\delta_1 = \delta_0 = 10^{-6}$							
\mathbf{MB}	-0.10	3.22	3.22	0.03	0.00	0.00	0.00
MB2	-0.10	3.22	3.22	0.03	0.00	0.00	0.00
Scenario C	Bias	\mathbf{SD}	RMSE	maxASMD	meanASMD	medASMD	GMIM
$\delta_1 = \delta_0 = 10^{-2}$							
\mathbf{MB}	0.26	0.81	0.85	0.39	0.02	0.01	0.02
MB2	0.37	0.86	0.93	0.30	0.03	0.01	0.03
$\delta_1 = \delta_0 = 10^{-4}$							
\mathbf{MB}	0.21	0.80	0.83	0.36	0.02	0.00	0.02
MB2	0.26	0.85	0.89	0.27	0.02	0.00	0.02
Scenario D	Bias	\mathbf{SD}	RMSE	maxASMD	meanASMD	medASMD	GMIM
$\delta_1 = \delta_0 = 10^{-2}$							
\mathbf{MB}	-0.24	0.46	0.52	0.15	0.01	0.00	0.01
$\mathbf{MB2}$	-0.21	0.45	0.49	0.17	0.01	0.00	0.00
$\delta_1 = \delta_0 = 10^{-4}$							
\mathbf{MB}	-0.16	0.46	0.49	0.13	0.01	0.00	0.00
MB2	-0.15	0.45	0.47	0.17	0.01	0.00	0.00

Table S3: Performance of MB and MB2 by varying (δ_1, δ_0) in Scenarios A, C, D

MB and MB2 are not sensitive to the values of the threshold parameters if they are small enough. The results are summarized in Table S3.

D Alternative formulations of Mahalanobis balancing

We discuss the formulation of Mahalanobis balancing when the normalization constraint is added to Problem (5). Using the entropy function as the loss function, the optimization problem is

$$\begin{array}{ll}
\underset{w \in \mathbb{R}^{n_{1}}}{\text{minimize}} : & \sum_{i=1}^{n} T_{i} w_{i} \log(w_{i}) \\ \\
\text{subject to:} & \begin{cases} w_{i} \geq 0, & i \in \{j : T_{j} = 1\}; \\ \sum_{i=1}^{n} T_{i} w_{i} = 1; \\ \sum_{i=1}^{n} T_{i} \{w_{i} \Phi(X_{i}) - \bar{\Phi}\}^{\top} W \sum_{i=1}^{n} T_{i} \{w_{i} \Phi(X_{i}) - \bar{\Phi}\} \leq \delta. \end{cases}$$

By the Fenchel duality theory, the corresponding dual problem is

$$\underset{\theta \in \mathbb{R}^{K}}{\text{minimize}} : -\log\left[\sum_{i=1}^{n} T_{i} \exp\left\{\theta^{\top} W^{1/2} \Phi(X_{i})\right\}\right] + \theta^{\top} W^{1/2} \bar{\Phi} + \sqrt{\delta} \|\theta\|_{2}.$$
(S4)

The estimated balanced weight for subject i in the treated group is

$$\hat{w}_{i}^{MB} = \frac{\exp\left\{\hat{\theta}^{\top} W^{1/2} \Phi(X_{i})\right\}}{\sum_{j:T_{j}=1} \exp\left\{\hat{\theta}^{\top} W^{1/2} \Phi(X_{j})\right\}}, \text{ for all } i \in \{j: T_{j}=1\}.$$

where $\hat{\theta}$ is solved from the dual problem. When $\delta = 0$, the dual problem (S4) reduces to that by entropy balancing (Hainmueller, 2012; Zhao and Percival, 2017). Therefore, Mahalanobis balancing with the normalization constraint is a direct generalization of entropy balancing. The parameter δ can be more difficult to tune when the normalization constraint is inserted.

References

- ATHEY, S., G. W. IMBENS, AND S. WAGER (2018): "Approximate residual balancing: debiased inference of average treatment effects in high dimensions," J. R. Stat. Soc. Ser. B. Stat. Methodol., 80, 597–623.
- FAN, J., K. IMAI, I. LEE, H. LIU, Y. NING, AND X. YANG (2022): "Optimal covariate balancing conditions in propensity score estimation," J. Bus. Econom. Statist., 41, 97–110.
- HAINMUELLER, J. (2012): "Entropy balancing for causal effects: A multivariate reweighting method to produce balanced samples in observational studies," *Political analysis*, 20, 25–46.

- JOSEY, K. P., E. JUAREZ-COLUNGA, F. YANG, AND D. GHOSH (2021): "A framework for covariate balance using Bregman distances," *Scand. J. Stat.*, 48, 790–816.
- LEE, D., S. YANG, L. DONG, X. WANG, D. ZENG, AND J. CAI (2023): "Improving trial generalizability using observational studies," *Biometrics*, 79, 1213–1225.
- LEI, L., P. J. BICKEL, AND N. EL KAROUI (2018): "Asymptotics for high dimensional regression M-estimates: fixed design results," *Probab. Theory Related Fields*, 172, 983–1079.
- MOHRI, M., A. ROSTAMIZADEH, AND A. TALWALKAR (2018): Foundations of Machine Learning, MIT press, 2nd ed.
- MORDUKHOVICH, B. S. AND N. M. NAM (2022): Convex Analysis and Beyond: Volume I: Basic Theory, Springer.
- NING, Y., S. PENG, AND K. IMAI (2020): "Robust estimation of causal effects via a highdimensional covariate balancing propensity score," *Biometrika*, 107, 533–554.
- RIGOLLET, P. AND J. HÜTTER (2015): *High-Dimensional Statistics*, Lecture notes for the course 18.657 at MIT.
- RYU, E. K. AND W. YIN (2022): Large-Scale Convex Optimization: Algorithms & Analyses via Monotone Operators, Cambridge University Press.
- TAN, Z. (2020a): "Model-assisted inference for treatment effects using regularized calibrated estimation with high-dimensional data," Ann. Statist., 48, 811–837.
- (2020b): "Regularized calibrated estimation of propensity scores with model misspecification and high-dimensional data," *Biometrika*, 107, 137–158.
- TROPP, J. A. ET AL. (2015): "An introduction to matrix concentration inequalities," Foundations and Trends® in Machine Learning, 8, 1–230.
- VAN DER VAART, A. W. (1998): Asymptotic statistics, Cambridge university press.
- WAINWRIGHT, M. J. (2019): *High-Dimensional Statistics: A Non-Asymptotic Viewpoint*, Cambridge University Press.
- WANG, Y. AND J. R. ZUBIZARRETA (2020): "Minimal dispersion approximately balancing weights: asymptotic properties and practical considerations," *Biometrika*, 107, 93–105.

ZHAO, Q. AND D. PERCIVAL (2017): "Entropy balancing is doubly robust," J. Causal Inference, 5, 20160010.