A Boosted-DCA with Power-Sum-DC Decomposition for Linearly Constrained Polynomial Programs

Hu Zhang 🔸 Yi-Shuai Niu 💷

Received: date / Accepted: date

Abstract This paper proposes a novel Difference-of-Convex (DC) decomposition for polynomials using a power-sum representation, achieved by solving a sparse linear system. We introduce the Boosted DCA with Exact Line Search $(BDCA_e)$ for addressing linearly constrained polynomial programs within the DC framework. Notably, we demonstrate that the exact line search equates to determining the roots of a univariate polynomial in an interval, with coefficients being computed explicitly based on the power-sum DC decompositions. The subsequential convergence of BDCA_e to critical points is proven, and its convergence rate under the Kurdyka-Łojasiewicz property is established. To efficiently tackle the convex subproblems, we integrate the Fast Dual Proximal Gradient (FDPG) method by exploiting the separable block structure of the power-sum DC decompositions. We validate our approach through numerical experiments on the Mean-Variance-Skewness-Kurtosis (MVSK) portfolio optimization model and box-constrained polynomial optimization problems. Comparative analysis of BDCA_e against DCA, BDCA with Armijo line search, UDCA, and UBDCA with projective DC decomposition, alongside standard nonlinear optimization solvers FMINCON and FILTERSD, substantiates the efficiency of our proposed approach.

Keywords Power-sum DC decomposition \cdot Polynomial optimization \cdot Boosted DCA with exact line search \cdot FDGP method \cdot Portfolio optimization

Mathematics Subject Classification (2010) 90C26 · 90C30 · 91G10

Accepted by the Journal of Optimization Theory and Applications February 15, 2024. Communicated by Alper Yildirim.

Hu Zhang

School of Mathematical Sciences, Shanghai Jiao Tong University, China

Yi-Shuai Niu

Beijing Institute of Mathematical Sciences and Applications, Beijing, China niuyishuai@bimsa.cn; niuyishuai82@hotmail.com

1 Introduction

We are interested in solving the polynomial optimization problem with linear constraints:

$$\min_{\boldsymbol{x}\in\mathcal{P}} f(\boldsymbol{x}), \tag{POPL}$$

where f is a multivariate polynomial, and \mathcal{P} represents a nonempty convex polyhedral set (not necessarily bounded), defined by

$$\mathcal{P} := \{ \boldsymbol{x} \in \mathbb{R}^n : \langle \boldsymbol{a}_i, \boldsymbol{x} \rangle \le b_i, \langle \boldsymbol{p}_j, \boldsymbol{x} \rangle = q_j, i = 1, \dots, m, j = 1, \dots, r \}.$$

Here, $a_i, p_j \in \mathbb{R}^n$ and $b_i, q_j \in \mathbb{R}$. Throughout the paper, the polynomial function f is assumed to be lower bounded over \mathcal{P} .

This problem arises in many applications including, but not limited to, higher-order moment portfolio selection [33, 48, 49, 42], eigenvalue complementarity problem [28, 41, 38, 39], tensor complementarity problem [54], Euclidean distance matrix completion problem [9], copositivity of matrix [17, 20, 56], Boolean polynomial program [37], and bilinear matrix inequalities [16, 40], among other. Each of these applications underscores the essential and widespread utility of addressing the stated polynomial optimization problem with linear constraints, thereby highlighting the imperative nature of devising efficient and robust solutions for it.

The (POPL) is indeed a special class of the difference-of-convex (DC) program [24,25,31], defined as below:

$$\min_{\boldsymbol{x}\in\mathcal{P}}\{f(\boldsymbol{x}):=g(\boldsymbol{x})-h(\boldsymbol{x})\},\tag{P_{dc}}$$

where g and h are convex polynomials. There are some related works on DC decompositions for polynomials. The DC decomposition for quadratic functions has been studied in [27, 15, 29] based on the eigenvalue decomposition of real symmetric matrices. However, in the case of general multivariate polynomials, this technique is no longer available and the DC decomposition is more difficult to obtain (especially in dense polynomials). Niu et al. [49] proposed a DC decomposition for general polynomials in the form of $\frac{\sigma}{2} \| \boldsymbol{x} \|^2 - (\frac{\sigma}{2} \| \boldsymbol{x} \|^2 - f)$ where $\sigma > 0$ is obtained by estimating an upper bound of the spectral radius of the Hessian matrix of f over a compact convex set. This technique has been successfully applied to several real-world applications such as the higher-order moment portfolio optimization [49] and the eigenvalue complementarity problems [41, 38, 39]. Ahmadi et al. [1] explored DC decompositions for polynomials via algebraic relaxation techniques. They showed in [2] that convexity is different from sos-convexity, and characterized the discrepancy between convexity and sos-convexity in [3] for polynomials. Several DC decompositions are proposed in [1] by solving linear, second-order cone, or semidefinite programs. Meanwhile, Niu defined in [36] the so-called Difference-of-Convex-Sums-of-Squares (DC-SOS) decomposition for general polynomials and developed several practical DC-SOS decomposition techniques. Some of these decomposition techniques are also parallelizable. The DC decomposition technique proposed in this paper is indeed a special DC-SOS decomposition in form of power-sums of linear forms, namely the power-sum DC (PSDC) decomposition. We will show that this decomposition can be established by solving a sparse linear system, which is discussed in Section 3.

Concerning the solution method for DC programming formulation (P_{dc}) , the most popular method is DCA, introduced by Pham [50] in 1985 as an extension of the subgradient method, and extensively developed by Le Thi and Pham since 1994 (see, e.g., [29, 30, 31, 46, 47] and the references therein). The main idea of DCA is to minimize a sequence of convex majorizations of the DC objective function f by linearizing the second DC component h at the current iterate. The general convergence theorem of DCA ensures that every limit point of the generated sequence $\{x^k\}$ by DCA is a critical point of (P_{dc}) (see, e.g., [46,30]). In recent years, some accelerated DCAs are established. Artacho et al. [6] proposed a boosted DCA (BDCA) by incorporating DCA with an Armijo-type line search to potentially accelerate DCA under the smoothness assumption of both q and h. This method is extended to the non-smooth case of h component in [7]. Meanwhile, Niu et al. [42] further extended BDCA to the convex constrained DC program for both smooth and nonsmooth cases. The global convergence theorem is established under the Lojasiewicz subgradient inequality. Besides, some other accelerations based on the heavy-ball method [52] and Nesterov's extrapolation [35] are also proposed. The inertial DCA (InDCA) [44] is established by de Oliveira et al. as a heavy-ball type [52] accelerated DCA. Le Thi et al. proposed in [51] a Nesterov-type accelerated DCA (ADCA). Wen et al. [55] proposed a proximal DCA algorithm with Nesterov-type extrapolation [23]. In this paper, based on the BDCA framework for convex constrained DC program established in [42], we introduce a variant of BDCA with exact line search for (POPL).

Contribution. (i) We propose two special DC-SOS decompositions in the power-sum of linear forms (namely, the power-sum DC decomposition) for multivariate polynomials. These decompositions can be efficiently generated by solving sparse linear systems. (ii) We introduce BDCA with an exact line search (BDCA_e) for solving (POPL), wherein the line search entails computing roots of a univariate polynomial in an interval, and an upper bound for the initial step size of the line search is estimated. (iii) The convex subproblems required in BDCA_e are tackled by employing the Fast Dual Proximal Gradient (FDPG) method [12], which leverages the separable block structure of the power-sum DC decompositions. (iv) The convergence analysis of BDCA_e is established akin to the methodology in [42], and its convergence rate under the Kurdyka-Łojasiewicz property is substantiated.

We delineate three major advantages of our proposed power-sum DC decompositions. Firstly, they furnish effective DC decompositions for polynomials. Secondly, the convex subproblem in BDCA_e can be efficiently resolved in parallel due to the unique power-sum structure. Thirdly, the exact line search is simplified to the process of locating the roots of a univariate polynomial, with coefficients being computed explicitly based on the power-sum DC decompositions. These advantages are also regarded as major contributions of this paper.

The rest of this paper is organized as follows. In Section 2, we recall some notations and preliminaries required in the paper. In Section 3, we establish two power-sum DC decompositions to get two DC programming formulations for (POPL). The corresponding BDCA_e with the two DC formulations are proposed in Section 4. The convergence analysis and the convergence rate of BDCA_e are proved in Section 5. The FDPG method for efficiently solving the convex subproblem is discussed in Section 6. Some numerical experiments on the Mean-Variance-Skewness-Kurtosis portfolio optimization model and the polynomial optimization problem with box constraint are reported in Section 7. Some concluding remarks and important questions are summarized in the final section.

2 Notations and preliminaries

Throughout this paper, matrices and vectors are written in bold uppercase letters and lowercase letters (e.g., \boldsymbol{X} and \boldsymbol{x}) respectively. We use x_i to denote the *i*-th coordinate of the vector \boldsymbol{x} , and \boldsymbol{X}^{\top} and \boldsymbol{X}^{-1} to denote the transpose matrix and the inverse matrix of \boldsymbol{X} . Let \mathbb{R}^n be the *n*-dimensional Euclidean space endowed with the classical inner product $\langle \boldsymbol{x}, \boldsymbol{y} \rangle := \sum_{i=1}^n x_i y_i$ for $\boldsymbol{x}, \boldsymbol{y} \in$ \mathbb{R}^n , the induced Euclidean norm $\|\boldsymbol{x}\| := \sqrt{\langle \boldsymbol{x}, \boldsymbol{x} \rangle}$ and the ℓ_{∞} norm $\|\boldsymbol{x}\|_{\infty} :=$ $\max_{1 \leq i \leq n} |x_i|$. For $\boldsymbol{X} \in \mathbb{R}^{n \times m}$, $\|\boldsymbol{X}\|$ denotes the spectral norm of \boldsymbol{X} defined by $\|\boldsymbol{X}\| := \sqrt{\lambda_{\max}(\boldsymbol{X}^{\top}\boldsymbol{X})}$, where $\lambda_{\max}(\boldsymbol{X}^{\top}\boldsymbol{X})$ is the largest eigenvalue of the matrix $\boldsymbol{X}^{\top}\boldsymbol{X}$. Let $\boldsymbol{e} \in \mathbb{R}^n$ be the vector of ones and $\mathbb{N} := \{0, 1, \ldots\}$ be the set of natural numbers. The gradient of a differentiable function $f : \mathbb{R}^n \to \mathbb{R}$ at $\boldsymbol{x} \in \mathbb{R}^n$ is denoted by $\nabla f(\boldsymbol{x}) \in \mathbb{R}^n$.

A function $f : \mathbb{R}^n \to \mathbb{R}$ is convex if

$$f(\lambda \boldsymbol{x} + (1-\lambda)\boldsymbol{y}) \leq \lambda f(\boldsymbol{x}) + (1-\lambda)f(\boldsymbol{y}) \text{ for all } \boldsymbol{x}, \boldsymbol{y} \in \mathbb{R}^n \text{ and } \lambda \in (0,1),$$

and f is called ρ -strongly convex $(\rho > 0)$ if $f - \frac{\rho}{2} \| \cdot \|^2$ is convex.

A function $\psi:\mathbb{R}^n\to\mathbb{R}^n$ is called Lipschitz continuous, if there is some constant L>0 such that

$$\|\psi(\boldsymbol{x}) - \psi(\boldsymbol{y})\| \leq L \|\boldsymbol{x} - \boldsymbol{y}\|$$
 for all $\boldsymbol{x}, \boldsymbol{y} \in \mathbb{R}^n$

furthermore, ψ is said to be locally Lipschitz continuous if, for each $\boldsymbol{x} \in \mathbb{R}^n$, there exists a neighborhood \mathcal{U} of \boldsymbol{x} such that ψ restricted to \mathcal{U} is Lipschitz continuous.

For a proper closed function $f : \mathbb{R}^n \to (-\infty, \infty]$, the Fréchet subdifferential of f at $x_0 \in \text{dom} f := \{ x \in \mathbb{R}^n : f(x) < \infty \}$ is defined by

$$\partial^F f(\boldsymbol{x}_0) = \{ \boldsymbol{u} \in \mathbb{R}^n : \liminf_{\boldsymbol{h} \to \boldsymbol{0}} \frac{f(\boldsymbol{x}_0 + \boldsymbol{h}) - f(\boldsymbol{x}_0) - \langle \boldsymbol{u}, \boldsymbol{h} \rangle}{\|\boldsymbol{h}\|} \ge 0 \}$$

and if $\boldsymbol{x}_0 \notin \text{dom} f$, we set $\partial^F f(\boldsymbol{x}_0) = \emptyset$. A point $\boldsymbol{x}_0 \in \mathbb{R}^n$ is called a Fréchet critical point of f, if $0 \in \partial^F f(\boldsymbol{x}_0)$. The effective domain of $\partial^F f$ is defined by

$$\operatorname{dom}(\partial^F f) := \{ \boldsymbol{x} \in \operatorname{dom} f : \partial^F f(\boldsymbol{x}) \neq \emptyset \}.$$

Particularly, when f is convex, $\partial^F f$ coincides with the classical subdifferential in convex analysis, defined by

$$\partial f(\boldsymbol{x}_0) = \{ \boldsymbol{u} \in \mathbb{R}^n : f(\boldsymbol{x}) - f(\boldsymbol{x}_0) \geq \langle \boldsymbol{u}, \boldsymbol{x} - \boldsymbol{x}_0 \rangle, \forall \boldsymbol{x} \in \mathbb{R}^n \}.$$

Let $\Gamma_0(\mathbb{R}^n)$ denote by the set of all proper closed and convex functions from \mathbb{R}^n to $(-\infty, \infty]$. For a function $f \in \Gamma_0(\mathbb{R}^n)$, its conjugate function f^* is defined by

$$f^*(\boldsymbol{y}) := \sup_{\boldsymbol{x}} \{ \langle \boldsymbol{y}, \boldsymbol{x}
angle - f(\boldsymbol{x}) : \boldsymbol{x} \in \mathbb{R}^n \}, orall \boldsymbol{y} \in \mathbb{R}^n.$$

The proximal mapping of f at $x \in \mathbb{R}^n$ is given by

$$\operatorname{prox}_{f}(\boldsymbol{x}) := \operatorname{argmin}_{\boldsymbol{u}} \{ f(\boldsymbol{u}) + \frac{1}{2} \| \boldsymbol{x} - \boldsymbol{u} \|^{2} : \boldsymbol{u} \in \mathbb{R}^{n} \}.$$

We call that f is 'prox-friendly' if $\operatorname{prox}_f(x)$ is easy to compute. See a list of prox-friendly convex functions in [10, Chapter 7]. For a nonempty closed convex set \mathcal{C} of \mathbb{R}^n , the indicator function of \mathcal{C} is defined by

$$\chi_{\mathcal{C}}(\boldsymbol{x}) := \left\{egin{array}{cc} 0 & ext{if } \boldsymbol{x} \in \mathcal{C}, \ \infty & ext{otherwise} \end{array}
ight.$$

Note that $\chi_{\mathcal{C}}$ belongs to $\Gamma_0(\mathbb{R}^n)$ and $\partial \chi_{\mathcal{C}}(\boldsymbol{x}) = \mathcal{N}_{\mathcal{C}}(\boldsymbol{x})$ if $\boldsymbol{x} \in \mathcal{C}$, where $\mathcal{N}_{\mathcal{C}}(\boldsymbol{x})$ denotes the normal cone of \mathcal{C} at \boldsymbol{x} , defined by

$$\mathcal{N}_{\mathcal{C}}(\boldsymbol{x}) := \{ \boldsymbol{v} \in \mathbb{R}^n : \langle \boldsymbol{v}, \boldsymbol{y} - \boldsymbol{x} \rangle \leq 0, \forall \boldsymbol{y} \in \mathcal{C} \},$$

and $\mathcal{N}_{\mathcal{C}}(\boldsymbol{x}) = \emptyset$ if $\boldsymbol{x} \notin \mathcal{C}$.

A function $f : \mathbb{R}^n \to [-\infty, \infty]$ is called difference-of-convex (DC) if f = g - h with $g, h \in \Gamma_0(\mathbb{R}^n)$, where g and h are called DC components of f. By introducing $\chi_{\mathcal{P}}$ into the component g of the DC program (P_{dc}) as

$$\min_{\boldsymbol{x}}\{(g+\chi_{\mathcal{P}})(\boldsymbol{x})-h(\boldsymbol{x}):\boldsymbol{x}\in\mathbb{R}^n\},$$

a point $x^* \in \mathcal{P}$ is called a (DC) critical point of (P_{dc}) if and only if

$$0 \in \partial(g + \chi_{\mathcal{P}})(\boldsymbol{x}^*) - \partial h(\boldsymbol{x}^*).$$

If h is differentiable at $\boldsymbol{x} \in \mathbb{R}^n$, then one has

$$\partial^F f(\boldsymbol{x}) = \partial g(\boldsymbol{x}) - \nabla h(\boldsymbol{x}).$$

In particular, to the case where both g and h are polynomials and \mathcal{P} is a convex polyhedral set, then $x^* \in \mathcal{P}$ is a DC critical point of (P_{dc}) if and only if

$$abla h(x^*) -
abla g(x^*) \in \mathcal{N}_{\mathcal{P}}(x^*).$$

The cone of feasible directions of \mathcal{P} at $\bar{y} \in \mathcal{P}$ is defined as

$$\mathcal{D}(\bar{\boldsymbol{y}}) := \{ \boldsymbol{d} \in \mathbb{R}^n : \exists \delta > 0 \text{ such that } \bar{\boldsymbol{y}} + t \boldsymbol{d} \in \mathcal{P}, \forall t \in [0, \delta] \},\$$

and the active set at \bar{y} is defined by

$$\mathcal{A}(\bar{\boldsymbol{y}}) = \{i \in \{1, \dots, m\} : \langle \boldsymbol{a}_i, \bar{\boldsymbol{y}} \rangle = b_i\}.$$

Clearly,

$$\mathcal{D}(\bar{\boldsymbol{y}}) = \{ \boldsymbol{d} \in \mathbb{R}^n : \langle \boldsymbol{a}_i, \boldsymbol{d} \rangle \le 0, \langle \boldsymbol{p}_j, \boldsymbol{d} \rangle = 0, i \in \mathcal{A}(\bar{\boldsymbol{y}}), j = 1, \dots, r \}.$$

Let $\mathbb{H}_d[\mathbf{x}]$ denote by the set of all *n*-variable and *d*-degree homogeneous polynomials (forms) with coefficients in \mathbb{R} , defined by

$$\mathbb{H}_{d}[\boldsymbol{x}] := \{ f = \sum_{\boldsymbol{\alpha}} c_{\boldsymbol{\alpha}} \boldsymbol{x}^{\boldsymbol{\alpha}} : \boldsymbol{x} \in \mathbb{R}^{n}, c_{\boldsymbol{\alpha}} \in \mathbb{R}, \boldsymbol{\alpha} \in \mathcal{I}_{n,d} \}.$$

Here, $\mathcal{I}_{n,d}$ denotes the set encompassing all weak compositions of d into n parts, defined by $\mathcal{I}_{n,d} = \{ \boldsymbol{\alpha} = (\alpha_1, \ldots, \alpha_n) : \sum_{i=1}^n \alpha_i = d, \alpha_i \in \mathbb{N} \}$. The cardinality of this set is denoted by $s_{n,d}$, is $\binom{n+d-1}{d}$. Thus, the set of all real polynomials with degree up to d in n variables is defined by $\mathbb{R}_d[\boldsymbol{x}] := \bigcup_{k=0}^d \mathbb{H}_k[\boldsymbol{x}]$. The power-product matrix associated with $\mathcal{I}_{n,d}$ is given by

$$\widehat{\boldsymbol{V}}(n,d) := \left({d \choose \boldsymbol{\alpha}^{j}} (\boldsymbol{\alpha}^{i})^{\boldsymbol{\alpha}^{j}} \right)_{1 \leq i,j \leq s_{n,d}},$$

where $\binom{d}{\alpha^{j}} := \frac{d!}{\alpha_{1}^{j}!\cdots\alpha_{n}^{j}!}$ is the multinomial coefficient and $(\alpha^{i})^{\alpha^{j}} := \prod_{k=1}^{n} (\alpha_{k}^{i})^{\alpha_{k}^{j}}$. The power-product matrix $\widehat{V}(n, d)$ is nonsingular for all positive integers n and d, and is sparse if $n \gg d$. See [43] for more properties on the power-product matrix.

A form $f(\boldsymbol{x}) \in \mathbb{H}_d[\boldsymbol{x}]$ is said to have a *power-sum* representation, if there exist linear forms $L_i(\boldsymbol{x}) \in \mathbb{H}_1[\boldsymbol{x}]$ and scalars $\lambda_i \in \mathbb{R}$ (i = 1, ..., r) such that

$$f(oldsymbol{x}) = \sum_{i=1}^r \lambda_i L_i^d(oldsymbol{x}), orall oldsymbol{x} \in \mathbb{R}^n$$

A proper and closed function $f : \mathbb{R}^n \to (-\infty, \infty]$ is said to have the Kurdyka–Łojasiewicz (KL) property at $\boldsymbol{x}^* \in \operatorname{dom}(\partial^F f)$, if there exist some constant $\eta \in (0, \infty)$, a concave function $\varphi : [0, \eta] \to [0, \infty)$ and a neighborhood \mathcal{U} of \boldsymbol{x}^* , such that

- (i) φ belongs to the class \mathscr{C}^1 on $(0, \eta)$;
- (ii) $\varphi(0) = 0$ and $\varphi' > 0$ on $(0, \eta)$;
- (iii) $\forall \boldsymbol{x} \in \mathcal{U}$ with $f(\boldsymbol{x}) f(\boldsymbol{x}^*) \in (0, \eta)$, we have the KL inequality

$$\varphi'(f(\boldsymbol{x}) - f(\boldsymbol{x}^*)) \operatorname{dist}(0, \partial^F f(\boldsymbol{x})) \ge 1.$$

The class of functions endowed with the KL property is very ample. For example, all semi-algebraic and real subanalytic functions adhere to the KL inequality [14,26,32]. Consequently, the sum of polynomial functions and indicator functions associated with polyhedral sets also exhibits the KL property.

3 Power-Sum-DC decompositions of polynomials

3.1 Power-sum representation

Numerous studies have established the existence of a power-sum representation for homogeneous polynomials (see e.g., [13,53]). In the quadratic case, the power-sum representation is nothing but spectral decomposition. Reznick outlined a necessary and sufficient condition for constructing power-sum decompositions of binary forms in [53, section 5], providing a cornerstone for the subsequent formulation of an efficient algorithm to decompose binary polynomials into power-sums of linear forms with the minimal term count, as expounded in [18]. In the more general case, Biosca proved in [13] that the set of the *d*-th power of all linear forms $\{L^d : L \in \mathbb{H}_1[x]\}$ is a generating set for $\mathbb{H}_d[x]$. Nevertheless, Biosca's proof relied on induction, leaving a general method for identifying a finite generating set for any polynomial as a stillunresolved challenge. In our work, we unveil a finite generating set applicable to any polynomial $\varphi(x) \in \mathbb{H}_d[x]$. Furthermore, we elucidate that constructing a power-sum decomposition for $\varphi(x)$ becomes a tractable task through the solution of a sparse linear system. Lemma 1 summarizes the essential result.

Lemma 1 Let $\varphi(\boldsymbol{x})$ be a form in $\mathbb{H}_d[\boldsymbol{x}]$, expressed as $\varphi(\boldsymbol{x}) = \sum_{\boldsymbol{\alpha} \in \mathcal{I}_{n,d}} c_{\boldsymbol{\alpha}} \boldsymbol{x}^{\boldsymbol{\alpha}}$, where $\boldsymbol{c} := (c_{\boldsymbol{\alpha}^1}, \ldots, c_{\boldsymbol{\alpha}^{s_{n,d}}})^{\top}$. Then, a power-sum representation of $\varphi(\boldsymbol{x})$ is given by

$$\varphi(\boldsymbol{x}) = \sum_{\boldsymbol{\alpha} \in \mathcal{I}_{n,d}} \lambda_{\boldsymbol{\alpha}} \langle \boldsymbol{\alpha}, \boldsymbol{x} \rangle^d, \tag{1}$$

where $\boldsymbol{\lambda} = \left(\lambda_{\boldsymbol{\alpha}^1}, \dots, \lambda_{\boldsymbol{\alpha}^{s_{n,d}}}\right)^\top$ is the unique solution of the linear system

$$\boldsymbol{V}(n,d)\boldsymbol{\lambda} = \boldsymbol{c}.$$
 (2)

Proof Let $\boldsymbol{x} \in \mathbb{R}^n$, the multinomial equation reads as

$$\left(\sum_{i=1}^{n} x_i\right)^d = \sum_{\boldsymbol{\alpha} \in \mathcal{I}_{n,d}} \binom{d}{\boldsymbol{\alpha}} \boldsymbol{x}^{\boldsymbol{\alpha}}$$

Subsequently, utilizing the relationship

$$\langle oldsymbol{lpha}, oldsymbol{x}
angle^d = \sum_{oldsymbol{eta} \in \mathcal{I}_{n,d}} inom{d}{oldsymbol{eta}} oldsymbol{lpha}^{oldsymbol{eta}} oldsymbol{x}^{oldsymbol{eta}}, orall oldsymbol{lpha} \in \mathcal{I}_{n,d},$$

we deduce that

$$\widetilde{\mathcal{B}} = \widehat{V}(n,d)\mathcal{B},\tag{3}$$

where $\mathcal{B} = (\boldsymbol{x}^{\boldsymbol{\alpha}^1}, \boldsymbol{x}^{\boldsymbol{\alpha}^2}, \dots, \boldsymbol{x}^{\boldsymbol{\alpha}^{s_{n,d}}})^{\top}$ and $\widetilde{\mathcal{B}} = (\langle \boldsymbol{\alpha}^1, \boldsymbol{x} \rangle^d, \langle \boldsymbol{\alpha}^2, \boldsymbol{x} \rangle^d, \dots, \langle \boldsymbol{\alpha}^{s_{n,d}}, \boldsymbol{x} \rangle^d)^{\top}$. Given the nonsingularity of $\widehat{\boldsymbol{V}}(n, d)$ proved in [43], we can uniquely rewrite (3) as

$$\mathcal{B} = \widehat{\boldsymbol{V}}^{-1}(n,d)\widehat{\mathcal{B}}$$

Hence

$$\varphi(\boldsymbol{x}) = \boldsymbol{c}^{\top} \boldsymbol{\mathcal{B}} = \boldsymbol{c}^{\top} \widehat{\boldsymbol{V}}^{-1}(n, d) \widetilde{\boldsymbol{\mathcal{B}}} = \boldsymbol{\lambda}^{\top} \widetilde{\boldsymbol{\mathcal{B}}},$$

where $\boldsymbol{\lambda} = (\widehat{\boldsymbol{V}}^{-1}(n, d))^{\top} \boldsymbol{c}.$

Remark 1 As demonstrated above, a power-sum representation of forms can be obtained by solving the linear system as expressed in Equation (2). It's important to note that the matrix $\hat{V}(n,d)$ is asymmetric in general, and its size is $\binom{n+d-1}{d}^2$, which could be very large even if n and d are not too large. For instance, when d = 4 and n = 100, the magnitude of $\hat{V}(n,d)$ exceeds millions. However, owing to the sparsity of $\hat{V}(n,d)$ particularly when $n \gg d$, solving Equation (2) becomes remarkably efficient. In most real-world applications of polynomial optimization, the number of variables n significantly exceeds the degree d, hence the matrix $\hat{V}(n,d)$ is sparse. Fig. 1 provides some illustrative examples. More sparsity¹ properties of $\hat{V}(n,d)$ is referred to [43].



Fig. 1 The sparsity of the matrix $\hat{V}(n, d)$ changes with variable *n* from 2 to 16 for some fixed degree $d \in \{3, 4, 5, 6\}$. It is observed that, for n > 6, the sparsity of $\hat{V}(n, d)$ for all $d \in \{3, 4, 5, 6\}$ is more than 90%, and the sparsity goes to 1 as $n \to \infty$ for any fixed *d*.

3.2 Power-sum difference-of-convex decompositions

In this section, we establish two DC decompositions for any polynomial, namely the *termwise power-sum-DC* (T-PSDC) decomposition and the *homogenizing dehomogenizing power-sum-DC* (HD-PSDC) decomposition.

 $^{^1}$ The sparsity of $\pmb{M} \in \mathbb{R}^{n \times m}$ is defined by the ratio of zero entries to the total number of entries in the matrix

T-PSDC decomposition : We will consider two cases (the odd degree case and the even degree case) in the next proposition.

Proposition 1 Let $f(\mathbf{x}) \in \mathbb{H}_d[\mathbf{x}]$.

(i) If d is even, then a PSDC decomposition is

$$f(\boldsymbol{x}) = \sum_{\boldsymbol{\alpha} \in \mathcal{I}^+} \lambda_{\boldsymbol{\alpha}} \langle \boldsymbol{\alpha}, \boldsymbol{x} \rangle^d - \sum_{\boldsymbol{\alpha} \in \mathcal{I}^-} (-\lambda_{\boldsymbol{\alpha}}) \langle \boldsymbol{\alpha}, \boldsymbol{x} \rangle^d,$$
(4)

where $\mathcal{I}^+ = \{ \boldsymbol{\alpha} \in \mathcal{I}_{n,d} : \lambda_{\boldsymbol{\alpha}} > 0 \}$ and $\mathcal{I}^- = \{ \boldsymbol{\alpha} \in \mathcal{I}_{n,d} : \lambda_{\boldsymbol{\alpha}} < 0 \}.$ (ii) If d is odd, then a PSDC decomposition is

$$f(\boldsymbol{x}) = \sum_{\boldsymbol{\alpha} \in \mathcal{I}^+} \lambda_{\boldsymbol{\alpha}} \langle \boldsymbol{\alpha}, \hat{\boldsymbol{x}} \rangle^{d+1} - \sum_{\boldsymbol{\alpha} \in \mathcal{I}^-} (-\lambda_{\boldsymbol{\alpha}}) \langle \boldsymbol{\alpha}, \hat{\boldsymbol{x}} \rangle^{d+1},$$
(5)

where
$$\mathcal{I}^+ = \{ \boldsymbol{\alpha} \in \mathcal{I}_{n+1,d+1} : \lambda_{\boldsymbol{\alpha}} > 0 \}, \mathcal{I}^- = \{ \boldsymbol{\alpha} \in \mathcal{I}_{n+1,d+1} : \lambda_{\boldsymbol{\alpha}} < 0 \}$$
 and $\hat{\boldsymbol{x}} = \begin{pmatrix} \boldsymbol{x} \\ 1 \end{pmatrix}.$

Proof Case (i) is elucidated by Lemma 1. For the proof of case (ii), we multiply $f(\boldsymbol{x})$ with an additional variable x_h , yielding an even degree form $\tilde{f}(\boldsymbol{x}, x_h) = x_h f(\boldsymbol{x})$. Subsequently, case (ii) is deduced by applying case (i) to $\tilde{f}(\boldsymbol{x}, x_h)$ and setting $x_h = 1$.

For any polynomial $f(\mathbf{x}) \in \mathbb{R}_d[\mathbf{x}]$, we can express it as:

$$f(\boldsymbol{x}) = \sum_{k=0}^{d} f_k(\boldsymbol{x})$$

where each $f_k(\boldsymbol{x}) \in \mathbb{H}_k[\boldsymbol{x}]$ for $k = 0, 1, \ldots, d$. We can then apply Proposition 1 to each form $f_k(\boldsymbol{x}), k = 1, 2, \ldots, d$, obtaining a so-called *termwise power-sum-DC* (T-PSDC) decomposition of $f(\boldsymbol{x})$. The decomposition procedure is summarized in Algorithm 1.

HD-PSDC decomposition : An *n*-variate *d*-degree form can be dehomogenized to a polynomial in n-1 variables of degree at most *d* by assigning a constant value of 1 to one variable (cf. dehomogenization). Conversely, any polynomial can be transformed into a form by introducing a new variable x_h and adjusting each monomial with appropriate powers of x_h , as expressed by $f_H(\boldsymbol{x}, x_h) = x_h^d f(x_1/x_h, \ldots, x_n/x_h)$ (cf. homogenization). Consequently, we propose the homogenizing-dehomogenizing power-sum-DC (HD-PSDC) decomposition as encapsulated in Algorithm 2, founded on Proposition 2.

Proposition 2 Let $f(\mathbf{x}) \in \mathbb{R}_d[\mathbf{x}]$. One has the PSDC decomposition

$$f(\boldsymbol{x}) = \sum_{\boldsymbol{\alpha} \in \mathcal{I}^+} \lambda_{\boldsymbol{\alpha}} \langle \boldsymbol{\alpha}, \hat{\boldsymbol{x}} \rangle^{d_f} - \sum_{\boldsymbol{\alpha} \in \mathcal{I}^-} (-\lambda_{\boldsymbol{\alpha}}) \langle \boldsymbol{\alpha}, \hat{\boldsymbol{x}} \rangle^{d_f},$$
(6)

where $\mathcal{I}^+ = \{ \boldsymbol{\alpha} \in \mathcal{I}_{n+1,d_f} : \lambda_{\boldsymbol{\alpha}} > 0 \}, \mathcal{I}^- = \{ \boldsymbol{\alpha} \in \mathcal{I}_{n+1,d_f} : \lambda_{\boldsymbol{\alpha}} < 0 \}$ and $\underline{d_f = 2 \lceil \frac{d}{2} \rceil^2, \hat{\boldsymbol{x}} = \begin{pmatrix} \boldsymbol{x} \\ 1 \end{pmatrix}.}$

² $\left[\cdot\right]$ is the ceiling function of a number.

Algorithm 1: T-PSDC Decomposition

Input: $f_k(\boldsymbol{x}), k = 1, \dots, d$ (all homogeneous components of the polynomial $f(\boldsymbol{x})$); **Output:** DC components $g_k(\boldsymbol{x})$ and $h_k(\boldsymbol{x}), k = 1, \ldots, d$; 1 for k = 1, ..., d do if k is odd then 2 $\hat{n} \leftarrow n+1, \hat{d} \leftarrow k+1, \hat{x} \leftarrow \begin{pmatrix} x \\ 1 \end{pmatrix};$ 3 4 else $\hat{n} \leftarrow n, \hat{d} \leftarrow k, \hat{\boldsymbol{x}} \leftarrow \boldsymbol{x};$ 5 6 \mathbf{end} Rewrite $f_k(\boldsymbol{x}) = \sum_{\boldsymbol{\alpha} \in \mathcal{I}_{\hat{n},\hat{d}}} c_{\boldsymbol{\alpha}} \boldsymbol{x}^{\boldsymbol{\alpha}};$ 7 8 $\mathcal{I} \leftarrow \mathcal{I}_{\hat{n},\hat{d}};$ $\boldsymbol{c} \leftarrow$ the coefficients of $f_k(\boldsymbol{x})$; 9 Compute $\boldsymbol{\lambda}$ by solving linear equations: $\widehat{\boldsymbol{V}}^{\top}(\hat{n}, \hat{d})\boldsymbol{\lambda} = \boldsymbol{c}$; 10 Set $\mathcal{I}^+ = \{ \boldsymbol{\alpha} \in \mathcal{I} : \lambda_{\boldsymbol{\alpha}} > 0 \}, \mathcal{I}^- = \{ \boldsymbol{\alpha} \in \mathcal{I} : \lambda_{\boldsymbol{\alpha}} < 0 \};$ 11 Compute $g_k(\boldsymbol{x}) = \sum_{\boldsymbol{\alpha} \in \mathcal{I}^+} \lambda_{\boldsymbol{\alpha}} \langle \boldsymbol{\alpha}, \hat{\boldsymbol{x}} \rangle^{\hat{d}}$ and $h_k(\boldsymbol{x}) = \sum_{\boldsymbol{\alpha} \in \mathcal{I}^-} -\lambda_{\boldsymbol{\alpha}} \langle \boldsymbol{\alpha}, \hat{\boldsymbol{x}} \rangle^{\hat{d}}$; 12 13 end 14 return $g_k(x)$ and $h_k(x), k = 1, ..., d;$

Proof Let $f(\mathbf{x}) \in \mathbb{R}_d[\mathbf{x}]$. We will establish the result through two distinct cases:

(i) For an even d, we homogenize $f(\mathbf{x})$ to yield an (n+1)-variate polynomial of degree d as

$$f_H(\boldsymbol{x}, x_h) = x_h^d f(x_1/x_h, \dots, x_n/x_h)$$

Then, we apply Lemma 1 for $f_H(\boldsymbol{x}, x_h)$ and assign $x_h = 1$, resulting in the desired expression.

(ii) For an odd d, we homogenize $f(\mathbf{x})$ to attain an (n+1)-variate polynomial of degree d+1 as

$$f_H(\boldsymbol{x}, x_h) = x_h^{d+1} f(x_1/x_h, \dots, x_n/x_h).$$

Again, we apply Lemma 1 for $f_H(\mathbf{x}, x_h)$ and set $x_h = 1$, yielding the desired result.

Numerical test: We compare the proposed two PSDC decompositions and the DSOS-DC decomposition in [1]. All algorithms are implemented on MATLAB using the polynomial optimization library SPOT [34] and tested with the solver MOSEK [5] and MATLAB R2021a on a laptop equipped with Intel Core i5-1035G1 CPU 1.19GHz and 8GB RAM. The numerical results are summarized in Table 1.

We observed that for odd degree cases $d \in \{3, 5\}$, the total running time for T-PSDC and HD-PSDC are almost the same, whereas for even degree cases $d \in \{4, 6\}$, HD-PSDC performs about 2 times faster than T-PSDC. Moreover, T-PSDC and HD-PSDC are at least 3 times faster than DSOS-DC for all

Algorithm 2: HD-PSDC Decomposition

Input: polynomial $f(\boldsymbol{x})$; Output: DC components $g(\boldsymbol{x})$ and $h(\boldsymbol{x})$; 1 $d \leftarrow$ the degree of $f(\boldsymbol{x})$; 2 if d is odd then 3 $\mid d \leftarrow d+1$; 4 end 5 $n \leftarrow n+1, \hat{\boldsymbol{x}} \leftarrow \begin{pmatrix} \boldsymbol{x} \\ 1 \end{pmatrix}$; 6 Rewrite $f(\boldsymbol{x}) = \sum_{\boldsymbol{\alpha} \in \mathcal{I}_{n,d}} c_{\boldsymbol{\alpha}} \boldsymbol{x}^{\boldsymbol{\alpha}}$; 7 $\mathcal{I} \leftarrow \mathcal{I}_{n,d}$; 8 $\boldsymbol{c} \leftarrow$ the coefficients of $f(\boldsymbol{x})$; 9 Compute $\boldsymbol{\lambda}$ by solving the linear system: $\hat{\boldsymbol{V}}^{\top}(n,d)\boldsymbol{\lambda} = \boldsymbol{c}$; 10 Set $\mathcal{I}^+ = \{\boldsymbol{\alpha} \in \mathcal{I} : \lambda_{\boldsymbol{\alpha}} > 0\}, \mathcal{I}^- = \{\boldsymbol{\alpha} \in \mathcal{I} : \lambda_{\boldsymbol{\alpha}} < 0\}$; 11 Compute $g(\boldsymbol{x}) = \sum_{\boldsymbol{\alpha} \in \mathcal{I}^+} \lambda_{\boldsymbol{\alpha}} \langle \boldsymbol{\alpha}, \hat{\boldsymbol{x}} \rangle^d$ and $h(\boldsymbol{x}) = \sum_{\boldsymbol{\alpha} \in \mathcal{I}^-} -\lambda_{\boldsymbol{\alpha}} \langle \boldsymbol{\alpha}, \hat{\boldsymbol{x}} \rangle^d$; 12 return $g(\boldsymbol{x})$ and $h(\boldsymbol{x})$.

cases. The total running time increases slowly with respect to the increase of n, while sharply with respect to the increase of d. Hence, for higher degree cases with $d \ge 5$, all three methods suffer from the out-of-memory issue for some large n on MATLAB. Furthermore, the running time of HD-PSDC and T-PSDC mainly depends on the generating time of the power-sum matrix, which is approximately $10 \sim 100$ times greater than solving the sparse linear system.

3.3 DC reformulations of (POPL) leveraging PSDC decompositions

Using the PSDC decompositions T-PSDC and HD-PSDC, as described in Propositions 1 and 2, we transform (POPL) into two DC programs as follows:

Let $|\cdot|_p$ represent the ℓ_p norm. Then

(i) Using T-PSDC decomposition, problem (POPL) can be reformulated as

$$\min_{\boldsymbol{x}\in\mathcal{P}} f(\boldsymbol{x}) = \underbrace{\sum_{p=1}^{\left\lceil \frac{d}{2} \right\rceil} \|\boldsymbol{A}_{p}^{+}\boldsymbol{x} + \boldsymbol{b}_{p}^{+}\|_{2p}^{2p}}_{g_{1}(\boldsymbol{x})} - \underbrace{\sum_{p=1}^{\left\lceil \frac{d}{2} \right\rceil} \|\boldsymbol{A}_{p}^{-}\boldsymbol{x} + \boldsymbol{b}_{p}^{-}\|_{2p}^{2p}}_{h_{1}(\boldsymbol{x})}, \quad (T-P_{dc})$$

where $A_p^+ \in \mathbb{R}^{m_1 \times n}, A_p^- \in \mathbb{R}^{m_2 \times n}, b_p^+ \in \mathbb{R}^{m_1}, b_p^- \in \mathbb{R}^{m_2}$ with

$$m_1 + m_2 \le s_{n,2p} + s_{n+1,2p}$$

and $p = 1, \ldots, \lceil \frac{d}{2} \rceil$.

(ii) Using HD-PSDC decomposition, problem (POPL) can be reformulated as

$$\min_{\boldsymbol{x}\in\mathcal{P}} f(\boldsymbol{x}) = \underbrace{\|\boldsymbol{A}^{+}\boldsymbol{x} + \boldsymbol{b}^{+}\|_{d_{f}}^{d_{f}}}_{g_{2}(\boldsymbol{x})} - \underbrace{\|\boldsymbol{A}^{-}\boldsymbol{x} + \boldsymbol{b}^{-}\|_{d_{f}}^{d_{f}}}_{h_{2}(\boldsymbol{x})}, \quad (\text{HD-P}_{\text{dc}})$$

$ \begin{array}{c c c c c c c c c c c c c c c c c c c $														d d		
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	6	c	л	4	3	6	57	4	3	6	57	4	ω 	Ľ	'n	
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	εų	ċ	ŝ	0.	0.	0.03	0.02	0.00	0.00	0.05	0.04	0.01	0.01	T_{gen}		
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	75	0	70	28	74	0.01	0.01	0.00	0.01	0.02	0.02	0.00	0.00	T_{sol}	0.	
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	288	202	283	<u>ب</u>	5.	12.21	12.31	0.12	0.13	19.43	12.35	0.22	0.14	T_{gen}	1	
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	1.09	0.44	л 22	71	32	1.94	2.04	0.01	0.01	3.53	1.96	0.02	0.01	T_{sol}	UT UT	
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	1			39.	36.	945.31	926.45	1.65	1.65	1547.60	940.43	2.94	1.69	T_{gen}	24	
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$				82	75	34.73	35.27	0.08	0.07	59.82	35.01	0.13	0.07	T_{sol}	1	
$\begin{array}{c c c c c c c c c c c c c c c c c c c $				198	156	I	I	13.43	13.60	I	Ι	24.46	13.72	T_{gen}	3	
$\begin{array}{c c c c c c c c c c c c c c c c c c c $.58	.66	I	Ι	0.24	0.24	Ι	Ι	0.46	0.25	T_{sol}	ω	
$\begin{array}{c c c c c c c c c c c c c c c c c c c $				567.	508.	I	I	71.85	74.55	I	Ι	135.68	74.90	T_{gen}	42	
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$				10	64	I	Ι	0.80	0.77	Ι	Ι	1.53	0.77	T_{sol}		
$\begin{array}{c c c c c c c c c c c c c c c c c c c $				1218	1062	1	I	300.62	319.52	I	I	577.79	318.35	T_{gen}	5	
$\begin{array}{c c c c c c c c c c c c c c c c c c c $.55	.69	I	I	1.85	1.73	I	Ι	3.36	1.74	T_{sol}		
T _{sol} 3.29 6.70 - 3.24 3.59						I	I	1032.44	1120.31	I	I	2030.30	1121.43	T_{gen}	60	
						I	I	3.59	3.24	I	Ι	6.70	3.29	T_{sol}		

 $\{0, 15, \dots, 60\}$, and uniformly distributed random integer coefficients sampled in [-10, 10]. For each pair (n, d), we perform 10 runs of each algorithm and compare the average CPU time (in seconds). For T-PSDC and HD-PSDC, T_{gen} and T_{sol} denote the average running time for generating the matrix \hat{V} and for solving the linear system (2) respectively. For DSOS-DC, only the average total running time is reported.

where $d_f = 2\lceil \frac{d}{2} \rceil$ and $\mathbf{A}^+ \in \mathbb{R}^{m_1 \times n}, \mathbf{A}^- \in \mathbb{R}^{m_2 \times n}, \mathbf{b}^+ \in \mathbb{R}^{m_1}, \mathbf{b}^- \in \mathbb{R}^{m_2}$ with $m_1 + m_2 \le s_{n+1,d_f}.$

4 BDCA_e for solving $(T-P_{dc})$ and $(HD-P_{dc})$

Problem (T–P_{dc}) and (HD–P_{dc}) are a subclass of DC programming. The classical DCA [46] is viable and efficient for solving these problems. Meanwhile, Niu et al. proved in [42, Proposition 1] that the direction $d^k = y^k - x^k$ in DCA is a descent direction (namely, DC descent direction) of f at y^k if (1) $d^k \neq 0$, (2) d^k is a feasible direction of \mathcal{P} at y^k , and (3) h is strongly convex. In this scenario, executing a line search (either exact or inexact) along d^k will yield an additional reduction in the value of the objective function, thereby ensuring an acceleration of DCA. Subsequently, we introduce a Boosted DCA with Exact Line Search (BDCA_e) for problems $(T-P_{dc})$ and $(HD-P_{dc})$, as delineated in Algorithm 3.

Input: $\boldsymbol{x}^0 \in \mathcal{P}, \, \varepsilon > 0;$ for k = 0, 1, ... do 1 Step 1: Solving the subproblem 2 $oldsymbol{y}^k \in \operatorname*{argmin}_{oldsymbol{x} \in \mathcal{P}} \ g_i(oldsymbol{x}) - \langle
abla h_i(oldsymbol{x}^k), oldsymbol{x}
angle,$ (\mathbf{P}_k)

Algorithm 3: BDCA_e for solving problem $(T-P_{dc})$ (or $(HD-P_{dc})$)

where i = 1 for $(\mathbf{T}-\mathbf{P}_{dc})$ and i = 2 for $(\mathbf{H}\mathbf{D}-\mathbf{P}_{dc})$. **Step 2:** Set $d^k = y^k - x^k$. If $||d^k||/(1 + ||x^k||) < \varepsilon$, return x^k ; з

Step 3: Set $\mathcal{I}(\boldsymbol{d}^k) = \{i \in \{1, \dots, m\} : \langle \boldsymbol{a}_i, \boldsymbol{d}^k \rangle > 0\}$ and compute 4

 $\bar{t} = \min_{i \in \mathcal{I}(\boldsymbol{d}^k)} \frac{b_i - \langle \boldsymbol{a}_i, \boldsymbol{y}^k \rangle}{\langle \boldsymbol{a}_i, \boldsymbol{d}^k \rangle}.$

Step 4: If $\bar{t} > 0$, then d^k is a feasible direction, we compute 5 $t_k = \text{ExactLineSearch}(\boldsymbol{d}^k, \boldsymbol{y}^k, \bar{t})$ by Algorithm 4, Otherwise, set $t_k = 0$; Step 5: Set $\boldsymbol{x}^{k+1} = \boldsymbol{y}^k + t_k \boldsymbol{d}^k$; 6 7 end

Here are some comments on Algorithm 3:

(i) BDCA_e reduces to the classical DCA when excluding **Step 3–4** and setting $t_k = 0$ for all k in **Step 5**. Additionally, if the exact line search in **Step** 4 is replaced by the Armijo line search, then BDCA_e reduces to BDCA in [42].

- (ii) In **Step 1**, we leverage the fast dual-based proximal gradient (FDGP) method [12] to solve (P_k) by taking advantage of the power-sum structure, which will be discussed in Section 6.
- (iii) In Step 3–4, we introduce the exact line search to accelerate DCA, which will be discussed in the next two paragraphs. In particular, our exact line search amounts to computing the roots of the derivative of a univariate polynomial, whose coefficients can be computed explicitly by Equation (9) based on the power-sum formulation. Note that the assumption that f is lower bounded over \mathcal{P} guarantees the finiteness of t_k . Otherwise, when $\mathcal{I}(\mathbf{d}^k) = \emptyset$, then $\bar{t} = \min \emptyset = \infty$ by convention and it is possible to find $t_k = \infty$. In this case, (P_{dc}) has no optimal solution.

Algorithm 4: Exact line search

Input: A descent direction d^k , a point y^k and a bound \bar{t} ; **Output:** Optimal step size t_k ;

- 1 Compute $c_j (j = 0, \ldots, d_f)$ by Equation (9);
- 2 Set $\mathcal{R} \leftarrow \{t : \hat{f}'(t) = 0\} \cup \{0, \bar{t}\};$
- $s t_k \in \operatorname{argmin}_{t \in \mathcal{R}} \hat{f}(t);$

In the following, we will explore various facets of BDCA_e, encompassing: the verification of the feasibility of direction d^k at y^k over \mathcal{P} , the estimation of an upper bound for the line search step size, and the elucidation of the exact line search procedure.

Upper bound estimation for the line search step size : The next proposition shows that

$$\bar{t} = \min_{i \in \mathcal{I}(\boldsymbol{d}^k)} \left\{ \frac{b_i - \langle \boldsymbol{a}_i, \boldsymbol{y}^k \rangle}{\langle \boldsymbol{a}_i, \boldsymbol{d}^k \rangle} \right\}$$
(7)

is an upper bound for the line search step size t_k to ensure that $\boldsymbol{y}^k + t_k \boldsymbol{d}^k \in \mathcal{P}$, where $\mathcal{I}(\boldsymbol{d}^k) = \{i \in \{1, \ldots, m\} : \langle \boldsymbol{a}_i, \boldsymbol{d}^k \rangle > 0\}.$

Lemma 2 Let \mathbf{y}^k and \mathbf{d}^k be generated in Algorithm 3. Then $\mathbf{d}^k \in \mathcal{D}(\mathbf{y}^k)$ is equivalent to $\bar{t} > 0$. Moreover, if $t \in [0, \bar{t}]$, then $\mathbf{y}^k + t\mathbf{d}^k \in \mathcal{P}$, and if $t > \bar{t}$, then $\mathbf{y}^k + t\mathbf{d}^k \notin \mathcal{P}$.

Proof We first prove the following result:

$$\boldsymbol{d}^k \in \mathcal{D}(\boldsymbol{y}^k) \hspace{0.2cm} \Leftrightarrow \hspace{0.2cm} \min_{i \in \mathcal{I}(\boldsymbol{d}^k)} \{b_i - \langle \boldsymbol{a}_i, \boldsymbol{y}^k \rangle\} > 0.$$

If $\mathcal{I}(\boldsymbol{d}^k) = \emptyset$, the conclusion is obvious by the definition of the cone of feasible directions $\mathcal{D}(\boldsymbol{y}^k)$.

If $\mathcal{I}(\boldsymbol{d}^k) \neq \emptyset$, then for $i \in \{1, \dots, m\}$ and $j \in \{1, \dots, r\}$, we have

$$oldsymbol{d}^k \in \mathcal{D}(oldsymbol{y}^k) \ \Leftrightarrow \ \left\{ egin{array}{ll} \langle oldsymbol{a}_i,oldsymbol{y}^k
angle = b_i \Rightarrow \langle oldsymbol{a}_i,oldsymbol{d}^k
angle \leq 0 \ \langle oldsymbol{p}_j,oldsymbol{d}^k
angle = 0. \end{array}
ight.$$

Invoking the fact that $\langle \boldsymbol{p}_j, \boldsymbol{d}^k \rangle = \langle \boldsymbol{p}_j, \boldsymbol{y}^k \rangle - \langle \boldsymbol{p}_j, \boldsymbol{x}^k \rangle = q_j - q_j = 0$, we deduce

$$\boldsymbol{d}^{k} \in \mathcal{D}(\boldsymbol{y}^{k}) \iff \langle \boldsymbol{a}_{i}, \boldsymbol{y}^{k} \rangle = b_{i} \Rightarrow \langle \boldsymbol{a}_{i}, \boldsymbol{d}^{k} \rangle \leq 0 \iff \langle \boldsymbol{a}_{i}, \boldsymbol{d}^{k} \rangle > 0 \Rightarrow \langle \boldsymbol{a}_{i}, \boldsymbol{y}^{k} \rangle < b_{i}.$$

The last equivalence condition implies that the conclusion holds.

We now proceed to prove the main result as the following:

(i) If $\mathcal{I}(\boldsymbol{d}^k) = \emptyset$, we have $\langle \boldsymbol{a}_i, \boldsymbol{d}^k \rangle \leq 0$ for i = 1, 2, ..., m. Thus, $\forall t \in [0, \infty)$, we observe that the fact

$$\begin{cases} \langle \boldsymbol{a}_i, \boldsymbol{y}^k + t\boldsymbol{d}^k \rangle = \langle \boldsymbol{a}_i, \boldsymbol{y}^k \rangle + t \langle \boldsymbol{a}_i, \boldsymbol{d}^k \rangle \leq b_i, \\ \langle \boldsymbol{p}_j, \boldsymbol{y}^k + t\boldsymbol{d}^k \rangle = \langle \boldsymbol{p}_j, \boldsymbol{y}^k \rangle + t \langle \boldsymbol{p}_j, \boldsymbol{d}^k \rangle = q_j. \end{cases}$$

holds for i = 1, 2, ..., m and j = 1, 2, ..., r. Therefore, we have $\boldsymbol{y}^k + t\boldsymbol{d}^k \in \mathcal{P}$. (ii) If $\mathcal{I}(\boldsymbol{d}^k) \neq \emptyset$, invoking the fact that $\bar{t} = \min_{i \in \mathcal{I}(\boldsymbol{d}^k)} \{ \frac{b_i - \langle \boldsymbol{a}_i, \boldsymbol{y}^k \rangle}{\langle \boldsymbol{a}_i, \boldsymbol{d}^k \rangle} \} > 0$, hence, $\forall t \in [0, \bar{t}]$, we have

$$\langle oldsymbol{p}_j,oldsymbol{y}^k+toldsymbol{d}^k
angle=\langleoldsymbol{p}_j,oldsymbol{y}^k
angle+t\langleoldsymbol{p}_j,oldsymbol{d}^k
angle=q_j.$$

for j = 1, ..., r.

Next, we verify the result $\langle \boldsymbol{a}_i, \boldsymbol{y}^k + t\boldsymbol{d}^k \rangle \leq b_i$ for $i = 1, \ldots, m$ in two cases: **Case 1:** If $i \in \mathcal{I}(\boldsymbol{d}^k)$, it implies $\langle \boldsymbol{a}_i, \boldsymbol{d}^k \rangle > 0$, we get that

$$\langle \boldsymbol{a}_i, \boldsymbol{y}^k + t \boldsymbol{d}^k \rangle = \langle \boldsymbol{a}_i, \boldsymbol{y}^k \rangle + t \langle \boldsymbol{a}_i, \boldsymbol{d}^k \rangle \leq \langle \boldsymbol{a}_i, \boldsymbol{y}^k \rangle + \min_{i \in \mathcal{I}(\boldsymbol{d}^k)} \{ \frac{b_i - \langle \boldsymbol{a}_i, \boldsymbol{y}^k \rangle}{\langle \boldsymbol{a}_i, \boldsymbol{d}^k \rangle} \} \langle \boldsymbol{a}_i, \boldsymbol{d}^k \rangle \leq b_i.$$

Case 2: If $i \notin \mathcal{I}(\mathbf{d}^k)$, it implies $\langle \mathbf{a}_i, \mathbf{d}^k \rangle \leq 0$. Thus, for all $i \notin \mathcal{I}(\mathbf{d}^k)$, we have

$$\langle \boldsymbol{a}_i, \boldsymbol{y}^k + t \boldsymbol{d}^k \rangle = \langle \boldsymbol{a}_i, \boldsymbol{y}^k \rangle + t \langle \boldsymbol{a}_i, \boldsymbol{d}^k \rangle \leq b_i$$

Moreover, $\forall t_0 > \bar{t}$ (Here, we always assume that $\bar{t} < \infty$), we have some $i_0 \in \mathcal{I}(\mathbf{d}^k)$ such that

$$t_0 > \frac{b_{i_0} - \langle \boldsymbol{a}_{i_0}, \boldsymbol{y}^k \rangle}{\langle \boldsymbol{a}_{i_0}, \boldsymbol{d}^k \rangle}.$$

Therefore, we can deduce that

$$\langle \boldsymbol{a}_{i_0}, \boldsymbol{y}^k + t_0 \boldsymbol{d}^k \rangle = \langle \boldsymbol{a}_{i_0}, \boldsymbol{y}^k + \delta_0 \boldsymbol{d}^k \rangle = \langle \boldsymbol{a}_{i_0}, \boldsymbol{y}^k \rangle + \delta_0 \langle \boldsymbol{a}_{i_0}, \boldsymbol{d}^k \rangle > b_{i_0}$$

This implies $\boldsymbol{y}^k + t_0 \boldsymbol{d}^k \notin \mathcal{P}$.

L		
L		
-		1

Exact line search : The exact line search procedure in **Step 4** is detailed in Algorithm 4. This algorithm is founded on the minimization problem (ELS):

$$\min_{t>0} \{ f(\boldsymbol{y}^k + t\boldsymbol{d}^k) : \boldsymbol{y}^k + t\boldsymbol{d}^k \in \mathcal{P} \}.$$
(ELS)

Without loss of generality, we assume that $f(\boldsymbol{x})$ can be represented as Proposition 2, given by the equation:

$$f(\boldsymbol{x}) = \sum_{\boldsymbol{\alpha} \in \mathcal{I}_{n+1,d_f}} \lambda_{\boldsymbol{\alpha}} \langle \boldsymbol{\alpha}, \hat{\boldsymbol{x}} \rangle^{d_f}.$$
(8)

Subsequently, substituting $\hat{\boldsymbol{x}} = \hat{\boldsymbol{y}}^k + t\hat{\boldsymbol{d}}^k$ into (8) with $\hat{\boldsymbol{y}}^k = \begin{pmatrix} \boldsymbol{y}^k \\ 1 \end{pmatrix}$ and $\hat{\boldsymbol{d}}^k = \begin{pmatrix} \boldsymbol{d}^k \\ 0 \end{pmatrix}$, we obtain

$$\hat{f}(t) = f(\boldsymbol{y}^k + t\boldsymbol{d}^k) = \sum_{j=0}^{d_f} c_j t^j,$$

where the coefficients c_j are determined by

$$c_j = \sum_{i=1}^{s_{n+1,d_f}} \lambda_i {\binom{d_f}{j}} \langle \boldsymbol{\alpha}^i, \hat{\boldsymbol{y}}^k \rangle^{d_f - j} \langle \boldsymbol{\alpha}^i, \hat{\boldsymbol{d}}^k \rangle^j.$$
(9)

Combined with Lemma 2, problem (ELS) can be represented as the following univariate polynomial minimization problem:

$$\min_{t} \{ \tilde{f}(t) : 0 \le t \le \bar{t} \}.$$

$$\tag{10}$$

The minimization problem (10) is well-defined. Specifically, if the set $\mathcal{I}(\boldsymbol{d}^k)$ is non-empty, it implies that \bar{t} is finite, thus confirming the validity of the assertion. Conversely, if $\mathcal{I}(\boldsymbol{d}^k)$ is empty, then \bar{t} assumes an infinite value, and the assertion remains valid due to the property that $|\hat{f}(t)| \to \infty$ as $t \to \infty$ and the assumption that f is lower bounded over \mathcal{P} .

Remark 2 Contrary to the Armijo line search method elaborated upon in [6, 42], our suggested technique necessitates the computation of the parameter c_j as illustrated in Equation (9), and the resolution of Problem (10), diverging from the traditional computation of objective function values. The minimization of Problem (10) can be effectively executed by identifying the roots of the derivative $\hat{f}'(t)$ of $\hat{f}(t)$. Particularly, when $d_f \leq 5$, the roots of $\hat{f}'(t)$ can be efficiently computed using a root-finding formula, whereas for instances where $d_f > 5$, a numerical method is a viable alternative. This methodology allows us to derive an optimal step size for each iteration of the line search procedure.

5 Convergence analysis for $\mathrm{BDCA}_{\mathrm{e}}$

Assumption 1 The component h is ρ -strongly convex with $\rho > 0$.

Note that this assumption is easy to satisfy for problems $(\mathbf{T}-\mathbf{P}_{dc})$ and $(\mathbf{H}\mathbf{D}-\mathbf{P}_{dc})$. Indeed, for a DC function $f = \tilde{g} - \tilde{h}$, we can take $g := \tilde{g} + \frac{\rho}{2} \|\boldsymbol{x}\|^2$ and $h := \tilde{h} + \frac{\rho}{2} \|\boldsymbol{x}\|^2$ with a small $\rho > 0$ to ensure the strong convexity of h.

Lemma 3 [6,42] Under Assumption 1, let $\{\mathbf{y}^k\}$, $\{\mathbf{x}^k\}$ and $\{\mathbf{d}^k\}$ be the sequences generated by Algorithm 3 when applied to problem (\mathbf{P}_{dc}) , it holds that

$$f(\boldsymbol{y}^k) \le f(\boldsymbol{x}^k) - \rho \|\boldsymbol{d}^k\|^2.$$
(11)

Theorem 1 Under Assumption 1, if the sequence $\{x^k\}$ generated by $BDCA_e$ in Algorithm 3 when applied to problem (P_{dc}) is bounded, then

(i) The sequence $\{f(\boldsymbol{x}^k)\}$ is decreasing and converging to some finite f_{opt} . (ii) Every limit point of the sequence $\{\boldsymbol{x}^k\}$ is a critical point of problem (P_{dc}) . (iii) $\sum_{k=0}^{\infty} \|\boldsymbol{d}^k\|^2 < \infty$.

Proof (i)To simplify the notations, we use g (resp. h) instead of g_i (resp. h_i) for $i \in \{1, 2\}$ in Algorithm 3. Invoking **Step 4** and **Step 5** of Algorithm 3, we deduce that

$$f(\boldsymbol{x}^{k+1}) \le f(\boldsymbol{y}^k)$$

Combining the previous inequality and (11), we derive that

$$f(\boldsymbol{x}^{k+1}) \le f(\boldsymbol{x}^k) - \rho \| \boldsymbol{d}^k \|^2,$$
 (12)

as $\{f(\boldsymbol{x}^k)\}$ is decreasing and f is lower bounded over \mathcal{P} , we conclude (i). (ii) By the boundedness of $\{\boldsymbol{x}^k\}$, we know that convergent subsequences exist. Let \boldsymbol{x}^* be a limit point of $\{\boldsymbol{x}^k\}$ and $\{\boldsymbol{x}^{k_j}\}$ be a subsequence of $\{\boldsymbol{x}^k\}$ converging to \boldsymbol{x}^* . As $k_j \to \infty$, we infer from (12) that

$$\|\boldsymbol{y}^{k_j} - \boldsymbol{x}^{k_j}\| \to 0.$$

this, combined with the fact that $\|\boldsymbol{y}^{k_j} - \boldsymbol{x}^*\| \leq \|\boldsymbol{y}^{k_j} - \boldsymbol{x}^{k_j}\| + \|\boldsymbol{x}^{k_j} - \boldsymbol{x}^*\|$, yields $\boldsymbol{y}^{k_j} \to \boldsymbol{x}^*$. By invoking the fact that

$$oldsymbol{y}^{k_j} \in \operatorname*{argmin}_{oldsymbol{x}}\{g(oldsymbol{x}) - \langle
abla h(oldsymbol{x}^{k_j}), oldsymbol{x}
angle, oldsymbol{x} \in \mathcal{P} \},$$

we can conclude, for all $x \in \mathcal{P}$, that:

$$\langle \nabla g(\boldsymbol{y}^{k_j}) - \nabla h(\boldsymbol{x}^{k_j}), \boldsymbol{x} - \boldsymbol{y}^{k_j} \rangle \ge 0.$$

Taking $k_j \to \infty$, and noting that g and h are continuously differentiable, we obtain

$$\langle \nabla f(\boldsymbol{x}^*), \boldsymbol{x} - \boldsymbol{x}^* \rangle \geq 0$$

This implies that \boldsymbol{x}^* is a critical point for problem (P_{dc}). (iii)Summing inequality (12) from k = 0 to ∞ , we obtain

$$\sum_{k=0}^{\infty} \rho \|\boldsymbol{d}^k\|^2 \le \sum_{k=0}^{\infty} (f(\boldsymbol{x}^k) - f(\boldsymbol{x}^{k+1})) = f(\boldsymbol{x}^0) - \lim_k f(\boldsymbol{x}^{k+1}) < \infty,$$

therefore, we can conclue that $\sum_{k=0}^{\infty} \|\boldsymbol{d}^k\|^2 < \infty$.

Remark 3 Note that if we assume that the line search step size t_k is upper bounded for all $k \ge 0$, then we will have a stronger convergence result based on [42, Theorem 11] that the sequence $\{x^k\}$ in BDCA_e is also convergent.

By employing the following useful lemma, we can obtain the rate of convergence of the sequence $\{f(\boldsymbol{x}^k)\}$ generated by Algorithm 3.

Lemma 4 Let $\{s_k\}$ be a nonincreasing and nonnegative real sequence converging to 0. Suppose that there exist $\alpha \geq 0$ and $\beta > 0$ such that for all large enough k,

$$s_{k+1}^{\alpha} \le \beta(s_k - s_{k+1}).$$
 (13)

Then

(i) if α = 0, then the sequence {s_k} converges to 0 in a finite number of steps;
(ii) if α ∈ (0, 1], then the sequence {s_k} converges linearly to 0 with rate β/(1+β).
(iii) if α > 1, then the sequence {s_k} converges sublinearly to 0, i.e., there exists η > 0 such that

$$s_k \le \eta k^{\frac{1}{1-\alpha}} \tag{14}$$

for large enough k.

Proof (i) If $\alpha = 0$, then (13) implies that

$$0 \le s_{k+1} \le s_k - \frac{1}{\beta}.$$

It follows by $s_k \to 0$ and $\frac{1}{\beta} > 0$ that $\{s_k\}$ converges to 0 in a finite number of steps, and we can estimate the number of steps as:

$$0 \le s_{k+1} \le s_k - \frac{1}{\beta} \le s_{k-1} - \frac{2}{\beta} \le \dots \le s_N - \frac{k - N + 1}{\beta}.$$

Hence

$$k \le \beta s_N + N - 1.$$

(ii) If $\alpha \in (0, 1]$. Since $s_k \to 0$, we have that $s_k < 1$ for large enough k. Thus, $s_{k+1} \leq s_k < 1$, and it follows by (13) that

$$s_{k+1} \le s_{k+1}^{\alpha} \le \beta(s_k - s_{k+1})$$

for large enough k. Hence

$$s_{k+1} \le \left(\frac{\beta}{1+\beta}\right) s_k,$$

i.e., $\{s_k\}$ converges linearly to 0 with rate $\frac{\beta}{1+\beta}$ for large enough k. (iii) If $\alpha > 1$, then we have two cases:

Case 1: if $\{x^k\}$ converges in a finite number of steps, then the inequality (14) trivially holds.

Case 2: Otherwise, $s_k > 0$ for all $k \in \mathbb{N}$. Let $\phi(t) = t^{-\alpha}$ and $\tau > 1$.

(a) Suppose that $\phi(s_{k+1}) \leq \tau \phi(s_k)$. By the decreasing of $\phi(t)$ and $s_k \geq s_{k+1}$, we have

$$\phi(s_k)(s_k - s_{k+1}) \le \int_{s_{k+1}}^{s_k} \phi(t) \, \mathrm{d}t = \frac{1}{1 - \alpha} (s_k^{1 - \alpha} - s_{k+1}^{1 - \alpha}).$$

It follows from (13) that

$$\frac{1}{\beta} \le \phi(s_{k+1})(s_k - s_{k+1}) \le \tau \phi(s_k)(s_k - s_{k+1}) \le \frac{\tau}{\alpha - 1}(s_{k+1}^{1-\alpha} - s_k^{1-\alpha}).$$

Hence

$$s_{k+1}^{1-\alpha} - s_k^{1-\alpha} \ge \frac{\alpha - 1}{\beta\tau} \tag{15}$$

for large enough k.

(b) Suppose that $\phi(s_{k+1}) \ge \tau \phi(s_k)$. Taking $q := \tau^{-\alpha^{-1}} \in (0, 1)$, then

$$s_{k+1} \le q s_k.$$

Hence

$$s_{k+1}^{1-\alpha} \ge q^{1-\alpha} s_k^{1-\alpha}$$

It follows that $\exists N > 0, \forall k \ge N$:

$$s_{k+1}^{1-\alpha} - s_k^{1-\alpha} \ge (q^{1-\alpha} - 1)s_k^{1-\alpha} \ge (q^{1-\alpha} - 1)s_N^{1-\alpha}.$$

In both cases, there exist a constant $\zeta := \min\{(q^{1-\alpha}-1)s_N^{1-\alpha}, \frac{\alpha-1}{\beta\tau}\}$ such that for large enough k, we have

$$s_{k+1}^{1-\alpha} - s_k^{1-\alpha} \ge \zeta$$

Summing for k from N to M - 1 > N, we have

$$s_M^{1-\alpha} - s_N^{1-\alpha} \ge \zeta(M-N).$$

Then, by the decreasing of $t^{\frac{1}{1-\alpha}}$, we get

$$s_M \le \left(s_N^{1-\alpha} + \zeta(M-N)\right)^{\frac{1}{1-\alpha}}.$$

Hence, there exist some $\eta > 0$ such that

$$s_M < \eta M^{\frac{1}{1-\alpha}}$$

for large enough M, which completes the proof.

We prove the convergence rate of the sequence $\{f(\boldsymbol{x}^k)\}$ generated by BDCA_e as the sequence $\{\boldsymbol{x}^k\}$ has a limit point \boldsymbol{x}^* at which $\tilde{f} = f + \chi_{\mathcal{P}}$ satisfies the Kurdyka–Łojasiewicz (KL) property [22] and ∇h is locally Lipschitz continuous.

Theorem 2 Under the assumptions of Theorem 1, let $\{\boldsymbol{x}^k\}, \{\boldsymbol{y}^k\}$ be sequences generated by $BDCA_e$ from a starting point $\boldsymbol{x}^0 \in \mathcal{P}$, and \mathcal{V} be the set of limit points of $\{\boldsymbol{x}^k\}$. Suppose that $\tilde{f} := f + \chi_{\mathcal{P}}$ satisfies the KL property at $\boldsymbol{x}^* \in \mathcal{V}$ with $\varphi(s) = Ms^{1-\theta}$ for some M > 0 and $0 \le \theta < 1$. Then the following statements hold:

- (i) If $\theta = 0$, then the sequence $\{f(\boldsymbol{x}^k)\}$ converges to f_{opt} in a finite number of steps.
- (ii) If $\theta \in (0, 1/2]$, then the sequence $\{f(\boldsymbol{x}^k)\}$ converges linearly to f_{opt} , that is, there exist positive constants N_0, η_0 and 0 < q < 1 such that $f(\boldsymbol{x}^k) f_{opt} \leq \eta_0 q^k$ for all $k \geq N_0$.
- (iii) If $\theta \in (1/2, 1)$, then the sequence $\{f(\boldsymbol{x}^k)\}$ converges sublinearly to f_{opt} , that is, there exist positive constants η_0 and N_0 such that $f(\boldsymbol{x}^k) - f_{opt} \leq \eta_0 k^{-\frac{1}{2\theta-1}}$ for all $k \geq N_0$.

Proof We conclude that f shares the same value over \mathcal{V} . As f is continuously differentiable, this follows directly from

$$f_{opt} = \lim_{k \to \infty} f(\boldsymbol{x}^k) = \lim_{j \to \infty} f(\boldsymbol{x}^{k_j}) = f(\boldsymbol{x}^*),$$

where the subsequence $\{\boldsymbol{x}^{k_j}\} \to \boldsymbol{x}^* \in \mathcal{V}$.

Given that ∇h is locally Lipschitz continuous, for each $x^* \in \mathcal{V}$, there exist some constants $L_{x^*} \geq 0$ and $\delta'_{x^*} > 0$ satisfying

$$\|\nabla h(\boldsymbol{x}) - \nabla h(\boldsymbol{y})\| \le L_{\boldsymbol{x}^*} \|\boldsymbol{x} - \boldsymbol{y}\|, \quad \forall \boldsymbol{x}, \boldsymbol{y} \in \mathbb{B}(\boldsymbol{x}^*, \delta_{\boldsymbol{x}^*}').$$
(16)

Furthermore, as f satisfies the Kurdyka–Łojasiewicz property at $\boldsymbol{x}^* \in \mathcal{V}$, there exists some constant $\delta_{\boldsymbol{x}^*}' > 0$, $\forall \boldsymbol{x} \in \mathbb{B}(\boldsymbol{x}^*, \delta_{\boldsymbol{x}^*}')$ with $f(\boldsymbol{x}^*) < f(\boldsymbol{x}) < f(\boldsymbol{x}^*) + \eta$ such that the Kurdyka–Łojasiewicz inequality holds. Let $\delta_{\boldsymbol{x}^*} = \min\{\delta_{\boldsymbol{x}^*}', \delta_{\boldsymbol{x}^*}''\}$, we can construct an open cover of the set \mathcal{V} :

$$\mathcal{V} \subset \bigcup_{\boldsymbol{x}^* \in \mathcal{V}} \mathbb{B}(\boldsymbol{x}^*, \delta_{\boldsymbol{x}^*}/4).$$

Since the sequence $\{x^k\}$ is bounded, then \mathcal{V} is a compact set. Thus, a finite number of points $v_1, \ldots, v_l \in \mathcal{V}$ exist such that

$$\mathcal{V} \subset \bigcup_{i=1}^{l} \mathbb{B}(\boldsymbol{v}_i, \delta_{\boldsymbol{v}_i}/4).$$

Let $L = \max_{1 \le i \le l} L_{\boldsymbol{v}_i}$ and $\delta = \min_{1 \le i \le l} \delta_{\boldsymbol{v}_i}/2$. Invoking the fact from Theorem 1 that $\{\boldsymbol{x}^k\}$ and $\{\boldsymbol{y}^k\}$ share the same set of limit points \mathcal{V} , we obtain $N_1 > 0$ such that $\{\boldsymbol{x}^k\} \in \bigcup_{i=1}^l \mathbb{B}(\boldsymbol{v}_i, \delta_{\boldsymbol{v}_i}/2)$ and $\|\boldsymbol{x}^k - \boldsymbol{y}^k\| \le \delta$ for all $k \ge N_1$.

Thus, for all $k \geq N_1$, there exists some \boldsymbol{v}_i such that $\boldsymbol{x}^k, \boldsymbol{y}^k \in \mathbb{B}(\boldsymbol{v}_i, \delta_{\boldsymbol{v}_i})$. Combined with (16), this implies that for all $k > N_1$

$$\|\nabla h(\boldsymbol{x}^k) - \nabla h(\boldsymbol{y}^k)\| \le L_{\boldsymbol{v}_i} \|\boldsymbol{x}^k - \boldsymbol{y}^k\| \le L \|\boldsymbol{x}^k - \boldsymbol{y}^k\|.$$
(17)

On the other hand, since $\lim_{k\to\infty} f(\boldsymbol{x}^k) = f_{opt}, f(\boldsymbol{x}^k) \geq f_{opt}$, there exist $N_2 > 0$ and $\eta > 0$ such that $f_{opt} < f(\boldsymbol{x}^k) < f_{opt} + \eta$ for all $k \geq N_2$. Setting $N = \max\{N_1, N_2\}$ and for each $k \geq N$, we conclude that there exists some $\boldsymbol{v}_i \in \mathcal{V}$ such that $\boldsymbol{x}^k, \boldsymbol{y}^k \in \mathbb{B}(\boldsymbol{v}_i, \delta_{\boldsymbol{v}_i})$. Hence, for each $\boldsymbol{y}^k \in \mathbb{B}(\boldsymbol{v}_i, \delta_{\boldsymbol{v}_i})$ and $k \geq N$, we have

$$f_{opt} < f(\boldsymbol{y}^k) \le f(\boldsymbol{x}^k) < f_{opt} + \eta.$$

Since \tilde{f} satisfies the Kurdyka–Łojasiewicz property and $\tilde{f}(\boldsymbol{x}) = f(\boldsymbol{x})$ over \mathcal{P} , we obtain

$$\varphi'(\tilde{f}(\boldsymbol{y}^k) - \tilde{f}(\boldsymbol{x}^*)) \operatorname{dist}(0, \partial^F \tilde{f}(\boldsymbol{y}^k)) \ge 1.$$
(18)

Because $\boldsymbol{y}^k \in \operatorname{argmin}_{\boldsymbol{x}} \{ g(\boldsymbol{x}) - \langle \nabla h(\boldsymbol{x}^k), \boldsymbol{x} \rangle, \boldsymbol{x} \in \mathcal{P} \}$, we get that

$$abla h(oldsymbol{x}^k) =
abla g(oldsymbol{y}^k) + oldsymbol{u}_k, oldsymbol{u}_k \in N_\mathcal{P}(oldsymbol{y}^k),$$

using the previous equation, we obtain

$$abla h(oldsymbol{y}^k) -
abla h(oldsymbol{x}^k) =
abla h(oldsymbol{y}^k) -
abla g(oldsymbol{y}^k) - oldsymbol{u}_k \in -\partial^F \widetilde{f}(oldsymbol{y}^k).$$

Then it follows from (17) that

dist
$$(0, \partial^F \tilde{f}(\boldsymbol{y}^k)) \le \|\nabla h(\boldsymbol{y}^k) - \nabla h(\boldsymbol{x}^k)\| \le L \|\boldsymbol{x}^k - \boldsymbol{y}^k\|.$$
 (19)

Combining (18) and (19) with $\varphi(s) = Ms^{1-\theta}$, we can obtain for all $k \ge N$ that

$$(f(\boldsymbol{y}^k) - f_{opt})^{\theta} \le ML(1-\theta) \|\boldsymbol{x}^k - \boldsymbol{y}^k\|.$$
(20)

For all $k \ge N$, it follows from $f(\boldsymbol{y}^k) - f_{opt} \ge f(\boldsymbol{x}^{k+1}) - f_{opt} \ge 0$, (12) and (20) that

$$(f(\boldsymbol{x}^{k+1}) - f_{opt})^{2\theta} \leq (f(\boldsymbol{y}^k) - f_{opt})^{2\theta} \\ \leq (ML)^2 (1 - \theta)^2 \|\boldsymbol{x}^k - \boldsymbol{y}^k\|^2 \\ \leq \frac{(ML)^2 (1 - \theta)^2}{\rho} (f(\boldsymbol{x}^k) - f(\boldsymbol{x}^{k+1})) \\ = \frac{(ML)^2 (1 - \theta)^2}{\rho} ((f(\boldsymbol{x}^k) - f_{opt}) - (f(\boldsymbol{x}^{k+1}) - f_{opt})).$$

Hence, setting $s_k = f(\boldsymbol{x}^k) - f_{opt}$ and $\tau = \frac{(ML)^2(1-\theta)^2}{\rho} > 0$, we can obtain

$$s_{k+1}^{2\theta} \leq \tau(s_k - s_{k+1})$$
 for all $k \geq N$.

Invoking Lemma 4, we can conclude (i), (ii) and (iii) immediately.

6 An efficient method for subproblem (P_k)

In this section, we focus on how to solve subproblem (\mathbf{P}_k) by introducing separable block structure into the function $g(\mathbf{x})$ and leveraging the fast dualbased proximal gradient (FDGP) method [12]. FDGP amounts to computing the vector projection on the polyhedral set and evaluating a proximal mapping in each iteration. Particularly, evaluating the proximal mapping is equivalent to finding the unique root of a strictly convex univariate polynomial, which can be performed in parallel due to the separable block structure. The vector projection on the polyhedral set can also be computed efficiently by splitting the equality and inequality constraints.

Without loss of generality, let us consider the objective function of problem (HD–P_{dc}) with a ρ -strongly convex DC decomposition as

$$f(\boldsymbol{x}) = \underbrace{\|\boldsymbol{A}^{+}\boldsymbol{x} + \boldsymbol{b}^{+}\|_{d_{f}}^{d_{f}} + \frac{\rho}{2}\|\boldsymbol{x}\|^{2}}_{g(\boldsymbol{x})} - \underbrace{\|\boldsymbol{A}^{-}\boldsymbol{x} + \boldsymbol{b}^{-}\|_{d_{f}}^{d_{f}} + \frac{\rho}{2}\|\boldsymbol{x}\|^{2}}_{h(\boldsymbol{x})}.$$
 (21)

The convex subproblem (\mathbf{P}_k) reads

$$\min_{\boldsymbol{x}\in\mathcal{P}}\left\{\|\boldsymbol{A}^{+}\boldsymbol{x}+\boldsymbol{b}^{+}\|_{d_{f}}^{d_{f}}+\frac{\rho}{2}\|\boldsymbol{x}\|^{2}-\langle\nabla h(\boldsymbol{x}^{k}),\boldsymbol{x}\rangle\right\},$$
(22)

where $A^+ = (a_1^+, a_2^+, \dots, a_{m_1}^+)^\top$ and $b^+ = (b_1^+, b_2^+, \dots, b_{m_1}^+)^\top$.

Now, we will discuss the solution method for (\mathbf{P}_k) by supposing $\chi_{\mathcal{P}}$ to be prox-friendly or not respectively.

Case 1: $\chi_{\mathcal{P}}(\cdot)$ is prox-friendly. By incorporating $\chi_{\mathcal{P}}(x)$ into the objective function of (22), we get

$$\min_{\boldsymbol{x}\in\mathbb{R}^n} \{\psi(\boldsymbol{x}) + \varphi(\boldsymbol{A}^+\boldsymbol{x})\},\tag{P_f}$$

where

$$\begin{cases} \varphi(\boldsymbol{A}^{+}\boldsymbol{x}) = \sum_{i=1}^{m_{1}} \varphi_{i}(\langle \boldsymbol{a}_{i}^{+}, \boldsymbol{x} \rangle) \text{ with } \varphi_{i}(\langle \boldsymbol{a}_{i}^{+}, \boldsymbol{x} \rangle) = (\langle \boldsymbol{a}_{i}^{+}, \boldsymbol{x} \rangle + b_{i})^{d_{f}}, i = 1, \dots, m_{1}, \\ \psi(\boldsymbol{x}) = \frac{\rho}{2} \|\boldsymbol{x}\|^{2} - \langle \nabla h(\boldsymbol{x}^{k}), \boldsymbol{x} \rangle + \chi_{\mathcal{P}}(\boldsymbol{x}). \end{cases}$$

By introducing the linear constraint $A^+x - \zeta = 0$, the Lagrangian is

$$L(oldsymbol{x},oldsymbol{\zeta};oldsymbol{z})=\psi(oldsymbol{x})+arphi(oldsymbol{\zeta})-\langleoldsymbol{A}^+oldsymbol{x}-oldsymbol{\zeta},oldsymbol{z}
angle$$

with $\boldsymbol{z} = (z_1, z_2, \dots, z_{m_1})^{\top}$. Then the Lagrangian dual problem of (P_f) is given by

$$\min_{\boldsymbol{z}} \{ \Psi(\boldsymbol{z}) + \Phi(\boldsymbol{z}) \}, \tag{D}_{\mathrm{f}}$$

where $\Psi(\mathbf{z}) = \psi^*((\mathbf{A}^+)^\top \mathbf{z}), \Phi(\mathbf{z}) = \varphi^*(-\mathbf{z})$. Applying the renowned FISTA [11] to (D_f), we get the FDPG method for (P_f) described in Algorithm 5.

Some comments on Algorithm 5 are summarized below:

- (i) In the settings of **Input**, we choose the initial point $z^0 = -\nabla \varphi(A^+ x^k)$ by solving the KKT system with respect to $\boldsymbol{\zeta}$ and \boldsymbol{z} at \boldsymbol{x}^k .
- (ii) In **Step 1**, the projection on \mathcal{P} (e.g., simplex or box) is easy to compute by assuming that $\chi_{\mathcal{P}}(\cdot)$ is prox-friendly.
- (iii) In Step 2, the proximal operator of all strictly convex functions φ_i is computed by finding the unique real root of its derivative. This can be achieved through a closed-form solution using the root formula when $d_f \leq 5$, or through numerical computation using Newton's method when $d_f > 5$. Notably, this step can be performed in parallel.

The convergence result of the sequence $\{u^l\}$ is given bellow.

Lemma 5 (See [12]) Suppose that $\{u^l\}_{l\geq 0}$ is the sequence generated by Algorithm 5. Then for the unique optimal solution x_{opt} of problem (P_f) and any optimal solution z_{opt} of problem (D_f) and $l \geq 1$,

$$\|\boldsymbol{u}^{l} - \boldsymbol{x}_{opt}\| \le 2\sqrt{\frac{L}{\rho}} \frac{\|\boldsymbol{z}^{0} - \boldsymbol{z}_{opt}\|}{l+1}$$

Hence, by setting $L = \frac{\|\mathbf{A}^+\|^2}{\rho}$ in Algorithm 5, we can simplify from Lemma 5 that

$$\|\boldsymbol{u}^{l} - \boldsymbol{x}_{opt}\| \le 2 \frac{\|\boldsymbol{A}^{+}\|}{\rho} \frac{\|\boldsymbol{z}^{0} - \boldsymbol{z}_{opt}\|}{l+1}.$$
 (23)

Lemma 6 Let A^+ be defined in problem (HD-P_{dc}) and c be the coefficient vector of its objective function f(x). Then

$$\|\boldsymbol{A}^{+}\| \leq \|\boldsymbol{L}_{n,d_{f}}\| \|\widehat{\boldsymbol{V}}^{-1}(n,d_{f})\|^{\frac{1}{d_{f}}} \|\boldsymbol{c}\|_{\infty}^{\frac{1}{d_{f}}},$$
(24)

where $\boldsymbol{L}_{n,d_f} = (\boldsymbol{\alpha}^1, \dots, \boldsymbol{\alpha}^{s_{n,d_f}})^\top$ and $d_f = 2\lceil \frac{d}{2} \rceil$.

23

Proof Following from Proposition 2 and (HD–P_{dc}), let us denote by $\boldsymbol{\lambda} = (\boldsymbol{\widehat{V}}^{-1}(n, d_f))^{\top} \boldsymbol{c}$, then we have

$$egin{pmatrix} egin{pmatrix} egin{aligned} egin{aligned} egin{aligned} egin{pmatrix} egin{aligned} egin{pmatrix} eta^+ & eta^+ \\ egin{pmatrix} egin{pmatrix}$$

Then according to the Cauchy interlacing theorem [45], we get

$$\|oldsymbol{A}^+\|\leq\|egin{pmatrix}oldsymbol{A}^+&oldsymbol{b}^+\oldsymbol{A}^-&oldsymbol{b}^-\end{pmatrix}\|\leq\|oldsymbol{L}_{n,d_f}\|(\|oldsymbol{\lambda}\|_\infty)^{rac{1}{d_f}}$$

Combining with $\|\boldsymbol{\lambda}\|_{\infty} \leq \|\boldsymbol{\widehat{V}}^{-1}(n, d_f)\| \|\boldsymbol{c}\|_{\infty}$, we get the desired conclusion.

Lamma 6 establishes the connection between $\|\mathbf{A}^+\|$ and the coefficients c of $f(\mathbf{x})$. When n and d_f are given, then \mathbf{L}_{n,d_f} and $\widehat{\mathbf{V}}(n,d_f)$ are fixed, hence

$$\|\boldsymbol{A}^+\| = O\left(\|\boldsymbol{c}\|_{\infty}^{\frac{1}{d_f}}\right).$$

Case 2: $\chi_{\mathcal{P}}(\cdot)$ is not prox-friendly. We split \mathcal{P} into two parts: the inequalities \mathcal{P}_I and the equalities \mathcal{P}_E as

$$\mathcal{P}_I = \{oldsymbol{x}:oldsymbol{A}oldsymbol{x} \leq oldsymbol{b}\} \quad ext{ and } \quad \mathcal{P}_E = \{oldsymbol{x}:oldsymbol{P}oldsymbol{x} = oldsymbol{q}\}$$

with $\boldsymbol{A} = (\boldsymbol{a}_1, \dots, \boldsymbol{a}_m)^{\top}, \boldsymbol{b} = (b_1, \dots, b_m)^{\top}$ and $\boldsymbol{P} = (\boldsymbol{p}_1, \dots, \boldsymbol{p}_r)^{\top}, \boldsymbol{q} = (q_1, \dots, q_r)^{\top}$. Here, we assume that \boldsymbol{P} has full row rank (otherwise, we use Gauss's elimination to get an equivalence equality constraint in full row rank). Thus, the indicator function $\chi_{\mathcal{P}_I}$ can be rewritten as

$$\chi_{\mathcal{P}_I}(\boldsymbol{x}) = \chi_{\mathrm{Box}[-\infty,\boldsymbol{b}]}(\boldsymbol{A}\boldsymbol{x}),$$

and problem (22) reads

$$\min_{\boldsymbol{x}} \{ \psi(\boldsymbol{x}) + \varphi(\boldsymbol{A}^{+}\boldsymbol{x}) + \chi_{\text{Box}[-\infty,\boldsymbol{b}]}(\boldsymbol{A}\boldsymbol{x}) \}, \quad (P_{\text{nf}})$$

where

$$\begin{cases} \varphi(\boldsymbol{A}^{+}\boldsymbol{x}) = \sum_{i=1}^{m_{1}} \varphi_{i}(\langle \boldsymbol{a}_{i}^{+}, \boldsymbol{x} \rangle) \text{ with } \varphi_{i}(\langle \boldsymbol{a}_{i}^{+}, \boldsymbol{x} \rangle) = (\langle \boldsymbol{a}_{i}^{+}, \boldsymbol{x} \rangle + b_{i})^{d_{f}}, i = 1, \dots, m_{1}, \\ \psi(\boldsymbol{x}) = \frac{\rho}{2} \|\boldsymbol{x}\|^{2} - \langle \nabla h(\boldsymbol{x}^{k}), \boldsymbol{x} \rangle + \chi_{\mathcal{P}_{E}}(\boldsymbol{x}). \end{cases}$$

The FDPG can also be applied for solving (P_{nf}) as in Case 1. The differences include:

(i) In **Step 1**, we compute $w^l \to u^{l+\frac{1}{2}} \to u^l$ as

$$\begin{cases} \boldsymbol{u}^{l+\frac{1}{2}} = \frac{1}{\rho} \left[\nabla h(\boldsymbol{x}^{k}) + \begin{pmatrix} \boldsymbol{A}^{+} \\ \boldsymbol{A} \end{pmatrix}^{\top} \boldsymbol{w}^{l} \right]; \\ \boldsymbol{u}^{l} = \boldsymbol{u}^{l+\frac{1}{2}} - \boldsymbol{P}^{\top} (\boldsymbol{P} \boldsymbol{P}^{\top})^{-1} (\boldsymbol{P} \boldsymbol{u}^{l+\frac{1}{2}} - \boldsymbol{q}). \end{cases}$$

(ii) In Step 2, there are some extra proximal operators to compute as

$$v_{i}^{l} = \begin{cases} \operatorname{prox}_{L\varphi_{i}}((\boldsymbol{a}_{i}^{+})^{\top}\boldsymbol{u}^{l} - Lw_{i}^{l}), & i = 1, \dots, m_{1}; \\ \min\{\boldsymbol{a}_{i}^{\top}\boldsymbol{u}^{l} - Lw_{i}^{l}, b_{i}\}, & i = m_{1} + 1, \dots, m_{1} + m_{n} \end{cases}$$

(iii) In **Step 3**, the z^{l+1} is updated by:

$$oldsymbol{z}^{l+1} = oldsymbol{w}^l - rac{1}{L} \left[egin{pmatrix} oldsymbol{A}^+ \ oldsymbol{A} \end{pmatrix} oldsymbol{u}^l - oldsymbol{v}^l
ight].$$

Note that similar method can also be viable for solving the convex subproblem (P_k) for $(T-P_{dc})$.

7 Numerical experiment

In this section, we test BDCA_e against the classical DCA and BDCA [42] applied to our new PSDC decompositions, UDCA [49] and UBDCA [42] (with projective DC decomposition) as well as FILTERSD [21] and FMINCON for solving the higher-order moment portfolio optimization model and the polynomial optimization problem with box constraint. All experiments are performed on MATLAB R2021a with a laptop equipped with Intel Core i5-1035G1 CPU 1.19GHz and 8GB of RAM. According to Lemma 1, the gradient computation of any form $\varphi(\mathbf{x}) \in \mathbb{H}_d[\mathbf{x}]$ is given by

$$\nabla \varphi(\boldsymbol{x}) = d \sum_{\boldsymbol{\alpha} \in \mathcal{I}_{n,d}} \lambda_{\boldsymbol{\alpha}} \langle \boldsymbol{\alpha}, \boldsymbol{x} \rangle^{d-1} \boldsymbol{\alpha}.$$
 (25)

7.1 The Mean-Variance-Skewness-Kurtosis portfolio optimization problem

Let $\boldsymbol{r} = (r_1, r_2, \ldots, r_n)^{\top}$ be a random return vector of n risky assets, \boldsymbol{x} be the n-dimensional decision vector, where x_i is the percentage invested in the i-th risky asset, and \mathbb{E} be the expectation operator. The moments of the return are defined as the following: the mean of the return denoted by $\boldsymbol{\mu} \in \mathbb{R}^n$ and $\forall i \in \{1, \ldots, n\}, \mu_i = \mathbb{E}(r_i)$; the variance of the return denoted by $\boldsymbol{V} \in \mathbb{R}^{n^2}$ and $\forall (i, j) \in \{1, \ldots, n\}^2, V_{ij} = \mathbb{E}\left([r_i - \mathbb{E}(r_i)][r_j - \mathbb{E}(r_j)]\right)$; the skewness of the return denoted by $\boldsymbol{S} \in \mathbb{R}^{n^3}$ and $\forall (i, j, k) \in \{1, \ldots, n\}^3,$ $S_{ijk} = \mathbb{E}\left([r_i - \mathbb{E}(r_i)][r_j - \mathbb{E}(r_j)][r_k - \mathbb{E}(r_k)]\right)$; the kurtosis of the return denoted by $\boldsymbol{K} \in \mathbb{R}^{n^4}$ and $\forall (i, j, k, l) \in \{1, \ldots, n\}^4, K_{ijkl} = \mathbb{E}([r_i - \mathbb{E}(r_i)][r_j -$ $\mathbb{E}(r_j)[r_k - \mathbb{E}(r_k)][r_l - \mathbb{E}(r_l)]$). The mean-variance-skewness-kurtosis portfolio model can be studied as a weighted nonconvex polynomial minimization problem (see, e.g., [42,49]):

$$\min_{\boldsymbol{x}} \{ f(\boldsymbol{x}) : \boldsymbol{e}^{\top} \boldsymbol{x} = 1, \ \boldsymbol{x} \ge \boldsymbol{0}, \ \boldsymbol{x} \in \mathbb{R}^n \},$$
(MVSK)

where

$$f(\boldsymbol{x}) := -\omega_1 \sum_i \mu_i x_i + \omega_2 \sum_{i,j} V_{ij} x_i x_j - \omega_3 \sum_{i,j,k} S_{ijk} x_i x_j x_k + \omega_4 \sum_{i,j,k,l} K_{ijkl} x_i x_j x_k x_l$$

with the parameter $\boldsymbol{\omega} = (\omega_1, \omega_2, \omega_3, \omega_4) \in \mathbb{R}^4_+$ being the investor's preference. Note that for some large n (e.g., n = 100), the total number of monomials in kurtosis can reach up to millions. However, due to the symmetry of \boldsymbol{V} (resp. \boldsymbol{S} and \boldsymbol{K}), the computation complexity on \boldsymbol{V} (resp. \boldsymbol{S} and \boldsymbol{K}) can be reduced from n^2 (resp. n^3 and n^4) to $\binom{n+1}{2}$ (resp. $\binom{n+2}{3}$ and $\binom{n+3}{4}$), see e.g., [33,49]. Data generation: We test on the dataset collected from monthly records

Data generation: We test on the dataset collected from monthly records from 1995 to 2015 of 48 industry-sector portfolios of the U.S. stock market with the number of assets $n \in \{11, 16, 21, 26, 31, 36, 41, 46\}$. For each n, we test problems with three different investor's preferences, including riskseeking $\boldsymbol{\omega} = (10, 1, 10, 1)$, risk-aversing $\boldsymbol{\omega} = (1, 10, 1, 10)$ and risk-neutral $\boldsymbol{\omega} = (10, 10, 10, 10)$.

Setup: The initial points are generated using MATLAB function rand: we first set x0=rand(n,1), then compute x0=x0/sum(x0) to get a feasible initial point $x^0 \in \mathcal{P}$. The gradients required in DCA, BDCA and BDCA_e are computed by Equation (25) and the gradients required in UDCA, UBDCA, FILTERSD and FMINCON for model (MVSK) are computed by the next formula (see e.g., [8,49]):

$$abla f(oldsymbol{x}) = -\omega_1 oldsymbol{\mu} + 2\omega_2 oldsymbol{V} oldsymbol{x} - 3\omega_3 oldsymbol{S}(oldsymbol{x}\otimesoldsymbol{x}) + 4\omega_4 oldsymbol{K}(oldsymbol{x}\otimesoldsymbol{x}\otimesoldsymbol{x}).$$

We terminate DCA, BDCA and BDCA_e with PSDC decomposition when

$$\|\boldsymbol{y}^k - \boldsymbol{x}^k\| / (1 + \|\boldsymbol{x}^k\|) \le 10^{-3}$$

Algorithm 5 for (P_k) is terminated if

$$\|\boldsymbol{z}^{l+1} - \boldsymbol{z}^{l}\|/(1 + \|\boldsymbol{z}^{l}\|) \le 10^{-3}.$$

UDCA and UBDCA are terminated if

$$\|\boldsymbol{y}^k - \boldsymbol{x}^k\| / (1 + \|\boldsymbol{x}^k\|) \le 10^{-5}.$$

FILTERSD and FMINCON are terminated with default parameters. The initial step size for Armijo line search is computed as in [42] by

$$t = \min\left\{\min_{i \in \mathcal{I}(\boldsymbol{d}^k)} \{\frac{b_i - \langle \boldsymbol{a}_i, \boldsymbol{y}^k \rangle}{\langle \boldsymbol{a}_i, \boldsymbol{d}^k \rangle}\}, \frac{\sqrt{2}}{\|\boldsymbol{d}^k\|}\right\}$$

We reduce t by $t = \beta t$ with $\beta = 0.8$ until

$$f(\boldsymbol{x}^{k+1}) - f(\boldsymbol{y}^k) \le -\sigma t^2 \|\boldsymbol{d}^k\|^2$$

is verified, where $\sigma = 10^{-3}$ is used.

Numerical results : In Table 2–4, The objective values (obj), the CPU time (time in seconds), the number of iterations (iter), the average CPU time and the average number of iterations are reported by comparing BDCA_e, DCA and BDCA applied to $(T-P_{dc})$ and $(HD-P_{dc})$, UDCA and UBDCA with projective DC decomposition, and the solvers FILTERSD and FMINCON on problem (MVSK).

27

We observe that BDCA_e is as fast as BDCA for solving (HD–P_{dc}), and both are the fastest methods among the others. Regarding the objective values, FILTERSD often provides the best ones. BDCA_e for both (T–P_{dc}) and (HD–P_{dc}) shares the same objective values as FILTERSD in most of the cases (15/24) with the difference of order $O(10^{-5})$. Moreover, BDCA_e and BDCA for both (T–P_{dc}) and (HD–P_{dc}) almost always require fewer iterations and result in better objective values than UDCA and UBDCA. This implies that the quality of the proposed two PSDC decompositions is better than the projective DC decomposition. Concerning the CPU time, BDCA_e for (HD–P_{dc}) is almost as fast as BDCA (HD–P_{dc}) and FMINCON, and at least 2 times faster than DCA and UDCA. We conclude that BDCA_e and BDCA for (T–P_{dc}) and (HD–P_{dc}) are promising approaches for solving (MVSK).

Fig. 2 illustrates the log average CPU time along the number of assets n from the results in Table 2–4. We observe that the DCA, BDCA, and BDCA_e algorithms, when applied to (HD–P_{dc}), exhibit faster performance compared to their counterparts applied to (T–P_{dc}). Furthermore, UDCA has the worst performance over all methods. FILTERSD is the fastest method when $n \leq 40$. As n > 40, the best performance is given by BDCA_e for (HD–P_{dc}), which demonstrates that BDCA_e should be a promising approach on large-scale settings.



Fig. 2 Log average CPU time along *n* for results of DCA, BDCA and BDCA_e for $(T-P_{dc})$ and $(HD-P_{dc})$ as well as FILTERSD and FMINCON reported in Tables 2–4.

	1																										
aver	24	23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	x	7	6	ы	4	ယ	2				
age	46	46	46	41	41	41	36	36	36	31	31	31	26	26	26	21	21	21	16	16	16	11	11	11		n	
77	72	96	5 8	72	77	58 8	55 57	93	59	66	94	56	65	58 85	118	78	58 8	119	63	89	24	06	80	113	iter		
0.82	2.74	3.96	2.14	1.52	1.67	1.11	0.72	1.31	0.74	0.48	0.76	0.39	0.28	0.40	0.49	0.17	0.14	0.24	0.07	0.13	0.04	0.05	0.05	0.04	time	D	
	-1.145e-01	7.829e-04	-1.650e-01	-1.145e-01	1.409e-03	-1.650e-01	-1.057e-01	8.497 e-04	-1.649e-01	-1.142e-01	1.139e-03	-1.649e-01	-9.899e-02	2.689e-03	-1.047e-01	-1.079e-01	5.191 e-03	-1.302e-01	-1.075e-01	8.466e-03	-1.646e-01	-9.608e-02	7.378e-03	-8.705e-02	obj	CA	A
20	25	23	18	21	29	17	20	23	17	15	22	17	13	26	23	19	23	17	20	23	15	15	22	19	iter		lgorith
0.33	1.40	1.20	0.79	0.72	0.84	0.43	0.40	0.44	0.29	0.14	0.24	0.16	0.08	0.17	0.13	0.09	0.07	0.05	0.04	0.05	0.04	0.02	0.03	0.02	time	BD	ims for
	-1.146e-01	7.303 e-04	-1.650e-01	-1.145e-01	1.412e-03	-1.650e-01	-1.057e-01	7.928e-04	-1.649e-01	-1.143e-01	1.051 e-03	-1.649e-01	-9.898e-02	2.687 e-03	-1.047e-01	-1.079e-01	$5.194\mathrm{e}{-03}$	-1.302e-01	-1.076e-01	8.391 e-03	-1.646e-01	-9.608e-02	7.354 e-03	-8.705e-02	obj	CA	solving (T–P _{dc}
14	14	19	14	14	22	12	15	16	14	11	17	12	10	16	16	14	16	12	15	13	12	11	12	10	iter		
0.31	0.98	1.31	0.74	0.44	0.95	0.41	0.31	0.55	0.27	0.15	0.32	0.18	0.07	0.18	0.14	0.07	0.10	0.04	0.03	0.05	0.04	0.02	0.02	0.01	time	BD	
	-1.146e-01	7.138e-04	-1.650e-01	-1.145e-01	1.354e-03	-1.650e-01	-1.057e-01	7.838e-04	-1.649e-01	-1.143e-01	1.057e-03	-1.649e-01	-9.902e-02	2.649 e-03	-1.047e-01	-1.079e-01	5.192 e-03	-1.302e-01	-1.076e-01	8.391e-03	-1.646e-01	-9.609e-02	7.348e-03	-8.705e-02	obj	CA_e	

Tabl
00 N 3
Numerical 1
esults o
f DCA,
BDCA
an B
DCA,
with
with
$\rho = 1$
for problem (
(T-P _{dc})
; Bold
values
for
best
results i
n o
bjective
function.

Table 3 Numeric					N																							
al res	avera	24	3	22	21	0	19	8	7	16	5	14	ω Π	12	Ξ	10	9	x	7	6	CT	4	ယ	2	<u> </u>			
ults o	ıge	46	46	46	41	41	41	36	36	36	31	31	31	26	26	26	21	21	21	16	16	16	11	11	11		n	
f DCA,	76	72	96	58 8	72	77	58 8	55 57	93	59	66	94	56	65	$^{20}_{85}$	118	78	58	119	60	89	22	68	74	113	iter		
BDCA a	0.72	2.50	3.56	1.96	1.32	1.49	1.04	0.62	1.16	0.66	0.42	0.65	0.35	0.23	0.31	0.40	0.11	0.09	0.19	0.03	0.05	0.02	0.02	0.02	0.03	time	D	
nd BDCA, with a		-1.145e-01	7.829e-04	-1.650e-01	-1.145e-01	1.409e-03	-1.650e-01	-1.057e-01	8.497 e-04	-1.649e-01	-1.142e-01	1.139e-03	-1.649e-01	-9.899e-02	2.689e-03	-1.047e-01	-1.079e-01	5.191e-03	-1.302e-01	-1.075e-01	8.466e-03	-1.646e-01	-9.608e-02	7.378e-03	-8.705e-02	obj	CA	A
= 1 for	20	25	23	18	21	29	17	20	23	17	15	22	17	13	26	23	19	22	17	19	23	15	13	23	18	iter		lgorithi
problem	0.29	1.29	1.02	0.74	0.64	0.77	0.40	0.35	0.38	0.24	0.12	0.20	0.15	0.07	0.14	0.11	0.05	0.05	0.04	0.03	0.03	0.02	0.00	0.01	0.01	time	BL	ms for s
(HD–P _{dc}); Bold		-1.146e-01	7.303 e-04	-1.650e-01	-1.145e-01	1.412 e-03	-1.650e-01	-1.057e-01	7.928 e-04	-1.649e-01	-1.143e-01	1.051 e-03	-1.649e-01	-9.898e-02	2.687 e-03	-1.047e-01	-1.079e-01	5.194 e-03	-1.302e-01	-1.076e-01	8.391 e-03	-1.646e-01	-9.608e-02	7.355e-03	-8.705 e-02	obj	CA	olving (HD-P _d
values fo	14	16	20	14	13	22	12	15	16	14	11	17	12	10	19	16	14	15	12	14	13	13	9	13	6	iter		c
r best res	0.27	0.95	1.15	0.66	0.38	0.81	0.36	0.27	0.44	0.23	0.13	0.28	0.16	0.05	0.16	0.11	0.04	0.06	0.04	0.02	0.03	0.02	0.01	0.01	0.01	time	BD	
ults in objective f		-1.146e-01	7.138e-04	-1.650e-01	-1.145e-01	1.354e-03	-1.650e-01	-1.057e-01	7.838e-04	-1.649e-01	-1.143e-01	1.057e-03	-1.649e-01	-9.902e-02	2.647 e-03	-1.047e-01	-1.079e-01	5.190e-03	-1.302e-01	-1.076e-01	8.391e-03	-1.646e-01	-9.608e-02	7.348e-03	-8.705e-02	obj	CAe	

ര ധ
5
Iumerical
results o
of DC
A, 1
ЗDCA
and
BDC.
A _e w
$^{\mathrm{ith}} ho$
for
problem
(HD-P
<u>ас</u>);
Bold
values
or b
est r
esults i
n ol
) jective f
function.

29

			UDC/			UBI	OCA	F	ILTERSD	ч	MINCON
		lter	time	obj	iter	time	obj	time	obj	time	obj
$1 \mid 1$	1 _	404	0.01	-8.705e-02	25	0.00	-8.705e-02	0.00	-8.705e-02	0.01	-8.705 e-02
2 1	1 1	000	0.13	7.345e-03	51	0.02	7.343e-03	0.01	7.343e-03	0.02	7.346e-03
3 1	1 1	8698	0.06	-9.611e-02	65	0.01	-9.611e-02	0.00	-9.611e-02	0.01	-9.611e-02
4 1	6	84	0.00	-1.646e-01	26	0.00	-1.646e-01	0.01	-1.646e-01	0.03	-1.646e-01
5 1	6 6	983	0.42	8.410e-03	68	0.05	8.381e-03	0.03	8.377e-03	0.02	8.381e-03
6 1	6 4	065	0.21	-1.076e-01	$^{20}_{80}$	0.04	-1.076e-01	0.02	-1.076e-01	0.02	-1.076e-01
7 2	1 1	653	0.22	-1.302e-01	50	0.03	-1.302e-01	0.04	-1.302e-01	0.04	-1.302e-01
8 2	1 5	609	0.85	5.163e-03	69	0.03	5.136e-03	0.03	5.108e-03	0.03	5.113e-03
9 2	1 8	290	1.01	-1.078e-01	82	0.04	-1.079e-01	0.06	-1.079e-01	0.07	-1.079e-01
10 2	6 3	814	0.89	-1.047e-01	75	0.04	-1.047e-01	0.05	-1.047e-01	0.04	-1.047e-01
11 2	6 10	0382	2.64	3.018e-03	91	0.10	2.768e-03	0.07	2.610e-03	0.07	2.617e-03
12 2	6 0	422	2.05	-9.884e-02	79	0.04	-9.895e-02	0.06	-9.902e-02	0.08	-9.901e-02
13 3	1 2	114	1.00	-1.649e-01	75	0.07	-1.649e-01	0.16	-1.649e-01	0.09	-1.649e-01
14 3	1 1:	2294	6.18	1.771 e-03	104	0.13	1.595e-03	0.11	1.042 e-03	0.12	1.050e-03
15 3	1 8	182	3.77	-1.140e-01	113	0.15	-1.140e-01	0.10	-1.143e-01	0.14	-1.143e-01
16 3	6 4	129	3.95	-1.649e-01	100	0.22	-1.649e-01	0.16	-1.649e-01	0.22	-1.649e-01
17 3	6 9	511	9.32	1.604 e-03	114	0.28	1.589e-03	0.29	7.721e-04	0.27	7.816e-04
18 3	6 10	8880	10.38	-1.048e-01	124	0.48	-1.053e-01	0.48	-1.057e-01	0.37	-1.057e-01
19 4	1 4	934	8.00	-1.650e-01	114	0.41	-1.650e-01	0.95	-1.650e-01	0.31	-1.650e-01
20 4	1 1:	3390	22.20	3.186e-03	125	0.54	3.166e-03	0.79	1.349e-03	1.07	1.360e-03
21 4	1 1:	2452	20.06	-1.134e-01	148	2.07	-1.135e-01	0.56	-1.145e-01	0.72	-1.145e-01
22 4	6 7	394	18.74	-1.650e-01	132	0.77	-1.650e-01	1.18	-1.650e-01	0.78	-1.650e-01
23 4	6 9	783	25.16	2.354 e-03	137	0.98	2.302e-03	1.37	7.006e-04	1.58	7.126e-04
24 4	6 1!	5278	38.82	-1.122e-01	157	1.97	-1.125e-01	1.88	-1.146e-01	1.25	-1.146e-01
averag	e 6	861	7.35		93	0.35		0.35		0.31	
Table 4 Numeric	al resu	lts of U	DCA, UE	3DCA, FILTERSD	and FM	INCON fo:	r (MVSK); Bold	values fo	r best results in o	objective	function.

7.2 Polynomial optimization problem with box constraints

We consider the box-constrained polynomial optimization problem (see [4, 19]):

$$\min_{\boldsymbol{x}} \{ f(\boldsymbol{x}) : \boldsymbol{l} \le \boldsymbol{x} \le \boldsymbol{u}, \ \boldsymbol{x} \in \mathbb{R}^n \}$$
(P_{box})

where $f(\boldsymbol{x})$ is a multivariate polynomial on $\mathbb{R}_d[\boldsymbol{x}]$ and $\boldsymbol{l}, \boldsymbol{u}$ are lower and upper bound of \boldsymbol{x} . We compare BDCA_e against the classical DCA and BDCA [42] applied to T-PSDC and HD-PSDC decompositions, respectively, as well as FILTERSD [21] and FMINCON for solving (P_{box}).

Data generation: The objective functions $f(\boldsymbol{x})$ in our experiments are randomly generated in $\mathbb{R}_d[\boldsymbol{x}]$, whose coefficients are uniformly distributed in the interval [-1, 1], with a specified density parameter $den \in (0, 1]$. Additionally, we set $[\boldsymbol{l}, \boldsymbol{u}] = [-\boldsymbol{e}, \boldsymbol{e}]$.

Setup: The initial points are randomly generated by MATLAB function rand. We start by setting x0 = rand(n,1), and then transform it with x0 = -1 + 2 * x0 to obtain a feasible initial point $x^0 \in \mathcal{P}$.

The gradients required in DCA, BDCA and BDCA_e are computed by Equation (25) and the gradients required in FILTERSD and FMINCON are approximated by

$$\frac{\partial f}{\partial x_i} = (f(\boldsymbol{x} + \delta \boldsymbol{e}_i) - f(\boldsymbol{x} - \delta \boldsymbol{e}_i))/(2\delta), i = 1, \dots, n,$$

where $\delta = 10^{-3}$. We terminate DCA, BDCA and BDCA_e when

$$\|\boldsymbol{y}^{k} - \boldsymbol{x}^{k}\| / (1 + \|\boldsymbol{x}^{k}\|) \le 5 \times 10^{-4}$$

Algorithm 5 for (\mathbf{P}_k) is terminated if

$$\|\boldsymbol{z}^{l+1} - \boldsymbol{z}^{l}\|/(1 + \|\boldsymbol{z}^{l}\|) \le 5 \times 10^{-5}.$$

The setting for the Armijo line search is the same as in (MVSK).

Numerical results: For each triple (n, d, den) with $n \in \{10, 20, 30, 40, 50\}, d \in \{3, 4\}$ and $den \in \{0.1, 0.4, 0.7, 1\}$, we generate 10 independent trials and compare five methods (namely, DCA, BDCA, BDCA_e with HD-PSDC decompositions, FILTERSD and FMINCON) in terms of the average CPU time (in seconds) over the 10 trials. Figures 3 and 4 show the performance of the five compared methods with d = 3 and 4 for all n and den. One can see that, our BDCA_e always outperforms DCA and BDCA in terms of the log average CPU time. In particular, for even d as d = 4 and for each den, BDCA_e improves with respect to FILTERSD and FMINCON when n increases, and for $den \ge 40\%$, BDCA_e always outperforms FILTERSD and FMINCON when $n \ge 30$. However, when d is odd (e.g., d = 3), regardless of the values of n and den, FILTERSD and FMINCON always outperform the others in terms of the log average CPU time. This is likely due to that the proposed power-sum DC decompositions transform an odd-degree polynomial into an even-degree polynomial, potentially increasing



Fig. 3 Comparison of DCA, BDCA and BDCA_e applied to HD-PSDC decomposition with $\rho = 1$, as well as FILTERSD and FMINCON for solving (P_{box}) with $n \in \{10, 20, 30, 40, 50\}, d = 3, den \in \{0.1, 0.4, 0.7, 1\}.$



Fig. 4 Comparison of DCA, BDCA and BDCA_e applied to HD-PSDC decomposition with $\rho = 1$, as well as FILTERSD and FMINCON for solving (P_{box}) with $n \in \{10, 20, 30, 40, 50\}, d = 4, den \in \{0.1, 0.4, 0.7, 1\}.$

the overall computational time when solving (P_{box}) . Furthermore, we observe that BDCA_e showcases reductions in log average CPU time compared to both FILTERSD and FMINCON as *d* increases. This may be attributed to the fact that the computational cost required in FILTERSD and FMINCON escalates more rapidly than the expense incurred in solving a sequence of convex subproblems by leveraging the power-sum structure as the degree *d* increases. Additionally, it's noteworthy that BDCA_e achieves objective values nearly identical to those of other methods across all tested scenarios.

33

Table 5-6 summarizes the numerical results of BDCA_e, DCA, BDCA (with T-PSDC and HD-PSDC decompositions, respectively), FILTERSD and FMINCON with $n \in \{10, 20, 30, 40, 50\}$, d = 4 and $den \in [0.5, 1]$ for (P_{box}). Here, we generate for each n four instances with a random density in [0.5, 1]. We observe that DCA, BDCA and BDCA_e for $(HD-P_{dc})$ consistently outperform the corresponding methods for (T-P_{dc}). Specifically, BDCAe exhibits nearly identical average CPU time for both $(T-P_{dc})$ and $(HD-P_{dc})$, and remarkably, they are the fastest methods (at least twice faster than the other methods in terms of the average CPU time). Furthermore, in about half of the examined scenarios, BDCA_e for both PSDC decompositions achieves the best objective values. Fig. 5 depicts the average CPU time on a logarithmic scale as a function of n. Specifically, it shows the average CPU time over the four generated instances for each variable count, as presented in Table 5-6. We can see that $BDCA_e$ for both two PSDC decompositions gives the best performance when n > 25, which confirms again that BDCA_e performs better than the others for large-scale cases.



Fig. 5 Log average CPU time along n for results of DCA, BDCA and BDCA_e for (T–P_{dc}) and (HD–P_{dc}) as well as FILTERSD and FMINCON reported in Table 5 and Table 6.

functio	Table .
D.	z
	ume
	rical
	rest
	ılts c
	of DC
	ДА, Е
	3DC
	A an
	d BI
)CA _e
	(wit
	h T-
	PSD
	C de
	dutoc
	ositi
	on aj
	nd ρ
	= 1)
	for
	(P _{bo}
	<u>;</u> ; В
	old v
	alue
	s for
	best
	resu
	lts in
	ı obj
	ective

	10.39	35		28.67	75		78.93	375	ge	avera	
$-1.154 \mathrm{e}{+03}$	33.57	43	$-1.154 \mathrm{e}{+03}$	145.07	120	$-1.154 \mathrm{e}{+03}$	259.53	655	0.66	50	20
$-1.349 \mathrm{e}{+03}$	36.73	48	-1.342 e + 03	81.11	107	-1.337e+03	235.82	482	0.73	50	19
-1.108e+03	33.31	57	-1.108e + 03	113.73	139	-9.967 e + 02	240.01	673	0.58	50	18
-1.292 e+03	31.72	62	$-1.019 \mathrm{e}{+03}$	63.97	116	-1.018e + 03	264.68	635	0.52	50	17
$-7.274 \mathrm{e}{+02}$	11.56	42	$-7.274 \mathrm{e}{+02}$	39.82	123	$-7.274 \mathrm{e}{+02}$	111.63	540	0.52	40	16
-5.871 e+02	11.65	49	-5.871 e+02	23.06	107	-5.770 e+02	91.58	598	0.53	40	15
$-6.138 \mathrm{e}{+02}$	11.98	41	$-6.106 \mathrm{e}{+02}$	27.87	106	$-6.015 \mathrm{e}{+02}$	96.48	373	0.55	40	14
$-7.695 \mathrm{e}{+02}$	14.69	70	$-7.695 \mathrm{e}{+02}$	31.65	06	$-7.695 \mathrm{e}{+02}$	116.89	457	0.70	40	13
-3.578e+02	4.54	31	$-3.578 \mathrm{e}{+02}$	9.44	67	-3.578e + 02	36.11	335	0.70	30	12
$-4.465 \mathrm{e}{+02}$	5.22	32	$-4.465 \mathrm{e}{+02}$	10.13	61	$-4.446e{+}02$	30.30	209	0.99	30	11
$-4.805 \mathrm{e}{+02}$	5.00	32	$-4.805 \mathrm{e}{+02}$	10.93	78	$-4.805 \mathrm{e}{+02}$	37.15	290	0.94	30	10
$-3.660 \mathrm{e}{+02}$	3.94	31	$-3.660 \mathrm{e}{+02}$	7.93	65	-3.658e + 02	23.45	195	0.52	30	9
$-1.446 \mathrm{e}{+02}$	0.87	21	$-1.446 \mathrm{e}{+02}$	1.69	42	$-1.446 \mathrm{e}{+02}$	6.02	185	0.62	20	x
$-1.625 \mathrm{e}{+02}$	0.88	23	$-1.625 \mathrm{e}{+02}$	2.14	59	$-1.625 \mathrm{e}{+02}$	11.50	410	0.79	20	7
-1.114e+02	0.67	21	-1.107e+02	1.46	45	-1.107e + 02	5.85	382	0.57	20	6
-1.265e+02	0.84	36	-1.265 e+02	1.94	71	$-1.262 \mathrm{e}{+02}$	6.43	454	0.92	20	σ
$-2.495 \mathrm{e}{+01}$	0.16	11	$-2.494 \mathrm{e}{+01}$	0.34	24	$-2.494 \mathrm{e}{+01}$	1.27	146	0.77	10	4
$-3.983 \mathrm{e}{+01}$	0.22	11	$-3.983 \mathrm{e}{+01}$	0.38	19	$-3.983 \mathrm{e}{+01}$	0.83	57	0.59	10	ယ
$-3.097 \mathrm{e}{+01}$	0.17	16	$-3.097 \mathrm{e}{+01}$	0.45	33	$-3.097 \mathrm{e}{+01}$	1.19	154	0.96	10	2
$-2.622 \mathrm{e}{+01}$	0.14	17	$-2.622 \mathrm{e}{+01}$	0.35	31	$-2.622 \mathrm{e}{+01}$	1.88	275	0.98	10	
obj	time	iter	obj	time	iter	obj	time	iter			
CA_e	BD		CA	BD		A	DC		den	n	
			olving $(T-P_{dc})$	hms for s	Algorit						

in objective function.	of DCA, BDCA and BDCA _e (with HD-PSDC decomposition a in objective function.	, of DCA, BDCA and BDCA _e (with HD-PSDC decomposition and $\rho = 1$), ε in objective function.	of DCA, BDCA and BDCA _e (with HD-PSDC decomposition and $\rho = 1$), as well as FILTERSD and i in objective function.	Bold values for best results	Table 6Numerical results
l BDCA _e (with HD-	$1~{\rm BDCA_e}$ (with HD-PSDC decomposition a .	1 BDCA _e (with HD-PSDC decomposition and $\rho = 1$), ε	1 BDCA _e (with HD-PSDC decomposition and $\rho = 1$), as well as FILTERSD and :	in objective function	of DCA, BDCA and
	PSDC decomposition a	PSDC decomposition and $\rho = 1$), ε	PSDC decomposition and $\rho = 1$), as well as FILTERSD and i	•	l BDCA _e (with HD-

- 10	ŝ		חת	V		DF	5		סס	22	F.			TITINOON
	10		t	A		DL	UCA		ВД	CAe				
	it	er	time	obj	iter	time	obj	iter	time	obj	time	obj	time	obj
0 0.9	38 2	81	0.97	-2.622e+01	37	0.22	-2.622 e+01	17	0.08	-2.622e+01	0.02	-2.622e+01	0.07	-2.622 e + 01
0.9	96 1.	46	0.74	-3.097e+01	29	0.23	-3.097 e + 01	15	0.10	-3.097 e + 01	0.02	-3.097 e + 01	0.04	-2.155e+01
0.5	00	64	0.64	-3.983e + 01	18	0.21	$-3.983 \mathrm{e}{+01}$	11	0.14	$-3.983 \mathrm{e}{+01}$	0.01	-3.983 e+01	0.02	$-3.983 \mathrm{e}{+01}$
0.2	17 1	33	0.96	-2.495e+01	26	0.25	-2.495 e+01	11	0.10	-2.495 e+01	0.02	-2.495 e+01	0.07	$-2.495 \mathrm{e}{+01}$
0.9	92 4	45	5.19	-1.265e+02	70	1.45	-1.265e+02	36	0.65	-1.265e+02	0.94	-1.265e+02	1.20	$-1.386 \mathrm{e}{+02}$
0.2	57 4	30	4.33	-1.107e+02	45	1.10	-1.107e+02	21	0.52	-1.114e+02	0.40	-1.114e+02	0.55	$-1.227 \mathrm{e}{+02}$
0.0	79 4	25	7.49	-1.625e+02	57	1.49	$-1.625 \mathrm{e}{+02}$	23	0.60	-1.625 e+02	0.57	-1.625 e+02	0.94	$-1.625 \mathrm{e}{+02}$
0.0	52 1	72	4.03	-1.446e+02	42	1.19	$-1.446 \mathrm{e}{+02}$	21	0.60	$-1.446 \mathrm{e}{+02}$	0.38	-1.446e+02	0.61	-1.390 e+02
0.2	52 1:	86	20.93	-3.660e + 02	63	7.35	-3.660 e + 02	31	3.73	-3.660e + 02	4.18	-2.958e+02	6.57	$-4.037 \mathrm{e}{+02}$
0.9	94 2	86	25.74	-4.805e+02	66	7.93	$-4.805 \mathrm{e}{+02}$	30	3.77	$-4.805 \mathrm{e}{+02}$	5.48	-4.805 e+02	6.93	$-4.805 \mathrm{e}{+02}$
0.9	39 1	. 66	24.11	-4.446e+02	63	7.68	-4.465e+02	32	3.76	-4.465e+02	5.38	-4.465 e+02	5.86	$-4.933\mathrm{e}{+02}$
0.0	70 3	34	29.42	-3.578e + 02	66	7.96	-3.578e + 02	31	3.73	-3.578e + 02	4.64	-3.495 e+02	3.43	$-3.634 \mathrm{e}{+02}$
0.0	70 4.	53	91.54	-7.695e+02	88	24.70	-7.695 e+02	62	10.90	-7.695 e+02	30.04	-7.517e+02	14.46	$-7.695 \mathrm{e}{+02}$
0.0	95 99	32	77.69	-6.014e+02	92	23.84	-6.106e + 02	38	10.26	-6.138e + 02	21.83	-6.658e + 02	21.58	-6.094e + 02
0.2	55 57 57	75	76.55	-5.769e + 02	119	24.87	-5.871e + 02	52	10.10	-5.871e + 02	29.90	-5.871e+02	31.06	$-7.517 \mathrm{e}{+02}$
0.0	52	14 9	94.29	-7.274e+02	125	44.22	$-7.274 \mathrm{e}{+02}$	44	9.32	$-7.274 \mathrm{e}{+02}$	20.35	-7.274 e + 02	23.93	$-6.843 \mathrm{e}{+02}$
0.0	62 6	12 2	18.13	-1.014e+03	129	71.44	-1.019 e + 03	61	25.03	$-1.295 \mathrm{e}{+03}$	86.10	-9.023e+02	42.68	-1.145e+03
0.0	80 23	86 2	48.07	-1.108e+03	96	88.39	-1.108e + 03	52	33.27	-1.108e + 03	110.35	-1.108e+03	136.60	$-1.150 \mathrm{e}{+03}$
0.0	73 4	89 2	06.95	-1.337e+03	114	69.15	$-1.349\mathrm{e}{+03}$	57	34.30	$-1.349 \mathrm{e}{+03}$	118.62	-1.203e+03	80.83	-1.276e + 03
0.6	56 4	94 2	06.30	-1.154e+03	105	74.94	-1.154e + 03	55	29.47	-1.154e+03	111.47	-1.223e+03	83.72	$-1.275 \mathrm{e}{+03}$
rage	<u>ب</u>	48	57.20		73	22.93		35 57	9.02		27.53		23.06	
nuN	nerical	resul	ts of D	CA, BDCA a	nd BI		with HD-PSDC	4						
	Trage 0.00000000000000000000000000000000000	$\begin{array}{ c c c c c c }\hline i \\ \hline i \\ \hline 0 & 0.98 & 2 \\ 0 & 0.96 & 1 \\ 0 & 0.77 & 1 \\ 0 & 0.77 & 4 \\ 0 & 0.57 & 4 \\ 0 & 0.57 & 4 \\ 0 & 0.57 & 4 \\ 0 & 0.57 & 4 \\ 0 & 0.52 & 1 \\ 0 & 0.52 & 1 \\ 0 & 0.53 & 5 \\ 0 & 0.52 & 5 \\ 0 & 0.58 & 3 \\ 0 & 0.58 & 3 \\ 1 \\ \hline rage & 3 \\ \hline Numerical \\ \hline \end{array}$	$\begin{tabular}{ c c c c } \hline & \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \$	$\begin{tabular}{ l l l l l l l l l l l l l l l l l l l$	$\begin{array}{ $	$\begin{tabular}{ c c c c c c c c c c c c c c c c c c c$	$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	$ \begin{tabular}{ c c c c c c c c c c c c c c c c c c c$	$ \begin{tabular}{ c c c c c c c c c c c c c c c c c c c$	$ \begin{tabular}{ c c c c c c c c c c c c c c c c c c c$	$ \begin{tabular}{ c c c c c c c c c c c c c c c c c c c$	$ \begin{tabular}{ c c c c c c c c c c c c c c c c c c c$	$ \begin{tabular}{ c c c c c c c c c c c c c c c c c c c$	$ \begin{tabular}{ c c c c c c c c c c c c c c c c c c c$

8 Conclusions and future work

In this paper, we presented a novel DC-SOS decomposition technique based on the power-sum representation for polynomials. Then we proposed an accelerated DCA (BDCA_e) combining DCA with an exact line search along the DC descent direction for solving linearly constrained polynomial optimization problems. We proved that any limit point of the sequence $\{x^k\}$ generated by $BDCA_e$ is a critical point of (POPL) and established the convergence rate of the sequence $\{f(\boldsymbol{x}^k)\}$ under the Kurdyka-Łojasiewicz assumption. We applied the FDPG method by exploiting the power-sum-DC structure for efficiently solving the convex subproblems required in $BDCA_e$. The numerical experiments on the higher-order moment portfolio optimization model (MVSK) and the box-constrained polynomial optimization problem (P_{box}) have demonstrated that $\mathrm{BDCA}_{\mathrm{e}}$ is often comparable to <code>FILTERSD</code> and <code>FMINCON</code> in terms of the CPU time and objective values, and outperforms DCA, UDCA, BDCA and UBDCA on the tested dataset. Moreover, BDCA_e shows advantages in solving large-scale dense polynomial optimization problems. It's worth noting that there are three major benefits for our proposed power-sum DC decompositions. First, they provide good DC decompositions for polynomials which can be verified in numerical results that DCA with power-sum DC decompositions requires fewer iterations than DCA with the classical projective DC decomposition. Second, the convex subproblem in BDCA_e can be solved efficiently in parallel by FDGP as discussed in Section 6, which cannot be the case without the particular power-sum structure. Third, the exact line search in Section 4 amounts to finding the roots of a univariate polynomial whose coefficients are computed explicitly based on our power-sum decompositions, leading to efficient exact line search in some real-world applications.

Several pivotal questions warrant further exploration. First, the generation of power-sum DC decompositions with minimal square terms arises as a notable inquiry due to its crucial influence on the efficiency of DCA and its variants. Second, the pursuit of an adaptive methodology for updating the strongly convex parameter ρ remains essential to ensure robust numerical performance in BDCA_e. It's worth noting the intrinsic trade-off regarding the choice of modulus ρ concerning the overall performance of BDCA_e. A diminutive modulus ρ engenders a superior DC decomposition, thereby reducing the requisite iterations for DCA_e; however, as articulated by Lemma 5, a magnified ρ accelerates the convergence of FDPG in solving the convex subproblem. Therefore, the quest for an optimal ρ to assure peak performance is imperative.

Acknowledgements We extend our sincere gratitude to the anonymous referees for their insightful suggestions which significantly improved the quality of this manuscript.

Funding This work was supported by the Natural Science Foundation of China (Grant No: 11601327).

Data Availability All data generated and analysed during this study are available from the corresponding author on reasonable request.

Conflict of Interest The authors declare that they have no conflict of interest.

References

- 1. Ahmadi, A.A., Hall, G.: DC decomposition of nonconvex polynomials with algebraic techniques. Mathematical Programming **169**(1), 69–94 (2018)
- Ahmadi, A.A., Parrilo, P.A.: A convex polynomial that is not sos-convex. Mathematical Programming 135(1), 275–292 (2012)
- Ahmadi, A.A., Parrilo, P.A.: A complete characterization of the gap between convexity and sos-convexity. SIAM Journal on Optimization 23(2), 811–833 (2013)
- Anjos, M.F., Lasserre, J.B.: Handbook on Semidefinite, Conic and Polynomial Optimization, vol. 166. Springer Science & Business Media (2011)
- 5. ApS, M.: Mosek Optimization Toolbox for MATLAB. User's Guide and Reference Manual, Version 4 (2019)
- Artacho, F.J.A., Fleming, R.M., Vuong, P.T.: Accelerating the DC algorithm for smooth functions. Mathematical Programming 169(1), 95–118 (2018)
- Artacho, F.J.A., Vuong, P.T.: The boosted DC algorithm for nonsmooth functions. SIAM Journal on Optimization 30(1), 980–1006 (2020)
- Athayde, G.M.D., Flôres, R.G.: Incorporating skewness and kurtosis in portfolio optimization: A multidimensional efficient set. In: Advances in Portfolio Construction and Implementation, Quantitative Finance, pp. 243–257. Elsevier (2003)
- Bakonyi, M., Johnson, C.R.: The euclidian distance matrix completion problem. SIAM Journal on Matrix Analysis and Applications 16(2), 646–654 (1995)
- 10. Beck, A.: First-order Methods in Optimization. SIAM (2017)
- Beck, A., Teboulle, M.: A fast iterative shrinkage-thresholding algorithm for linear inverse problems. SIAM journal on imaging sciences 2(1), 183–202 (2009)
- Beck, A., Teboulle, M.: A fast dual proximal gradient algorithm for convex minimization and applications. Operations Research Letters 42(1), 1–6 (2014)
- Biosca, A.F.: Representation of a polynomial function as a difference of convex polynomials, with an application. In: Generalized Convexity and Generalized Monotonicity, Lecture Notes in Economics and Mathematical Systems, pp. 189–207. Springer (2001)
- Bolte, J., Daniilidis, A., Lewis, A.: The Łojasiewicz inequality for nonsmooth subanalytic functions with applications to subgradient dynamical systems. SIAM Journal on Optimization 17(4), 1205–1223 (2007)
- Bomze, I.M., Locatelli, M.: Undominated dc decompositions of quadratic functions and applications to branch-and-bound approaches. Computational Optimization and Applications 28(2), 227–245 (2004)
- Boyd, S., El Ghaoui, L., Feron, E., Balakrishnan, V.: Linear Matrix Inequalities in System and Control Theory. SIAM (1994)
- Brás, C., Eichfelder, G., Júdice, J.: Copositivity tests based on the linear complementarity problem. Computational Optimization and Applications 63(2), 461–493 (2016)
- Comon, P., Mourrain, B.: Decomposition of quantics in sums of powers of linear forms. Signal Processing 53(2-3), 93–107 (1996)
- De Klerk, E., Hess, R., Laurent, M.: Improved convergence rates for Lasserre-type hierarchies of upper bounds for box-constrained polynomial optimization. SIAM Journal on Optimization 27(1), 347–367 (2017)
- 20. Dür, M., Hiriart-Urruty, J.B.: Testing copositivity with the help of difference-of-convex optimization. Mathematical Programming **140**(1), 31–43 (2013)
- Fletcher, R.: Filtersd-a Library for Nonlinear Optimization Written in Fortran (2015). URL https://projects.coin-or.org/filterSD
- Frankel, P., Garrigos, G., Peypouquet, J.: Splitting methods with variable metric for Kurdyka–Lojasiewicz functions and general convergence rates. Journal of Optimization Theory and Applications 165(3), 874–900 (2015)

- Gotoh, J., Takeda, A., Tono, K.: DC formulations and algorithms for sparse optimization problems. Mathematical Programming 169(1), 141–176 (2018)
- Hartman, P.: On functions representable as a difference of convex functions. Pacific Journal of Mathematics 9(3), 707–713 (1959)
- Hiriart-Urruty, J.B.: Generalized differentiability, duality and optimization for problems dealing with differences of convex functions. In: Convexity and duality in optimization, Lecture Notes in Economics and Mathematical Systems, pp. 37–70. Springer (1985)
- Kurdyka, K., Parusinski, A.: wf-stratification of subanalytic functions and the Łojasiewicz inequality. Comptes rendus de l'Académie des sciences. Série 1, Mathématique 318(2), 129–133 (1994)
- Le Thi, H.A.: An efficient algorithm for globally minimizing a quadratic function under convex quadratic constraints. Mathematical programming 87(3), 401–426 (2000)
- Le Thi, H.A., Moeini, M., Pham, D.T., Judice, J.: A DC programming approach for solving the symmetric eigenvalue complementarity problem. Computational Optimization and Applications 51(3), 1097–1117 (2012)
- 29. Le Thi, H.A., Pham, D.T.: Solving a class of linearly constrained indefinite quadratic problems by DC algorithms. Journal of global optimization **11**(3), 253–285 (1997)
- Le Thi, H.A., Pham, D.T.: The DC (difference of convex functions) programming and DCA revisited with DC models of real world nonconvex optimization problems. Annals of operations research 133(1), 23–46 (2005)
- 31. Le Thi, H.A., Pham, D.T.: DC programming and DCA: thirty years of developments. Mathematical Programming **169**(1), 5–68 (2018)
- Lojasiewicz, S.: Sur La Géométrie Semi et Sous-analytique. In: Annales de l'institut Fourier, pp. 1575–1595 (1993)
- Maringer, D., Parpas, P.: Global optimization of higher order moments in portfolio selection. Journal of Global optimization 43(2), 219–230 (2009)
- 34. Megretski, A.: SPOT: Systems Polynomial Optimization Tools (2013)
- 35. Nesterov, Y.E.: A method for solving the convex programming problem with convergence rate $O(1/k^2)$. Dokl. Akad. Nauk SSSR **269**(3), 543–547 (1983)
- Niu, Y.S.: On difference-of-sos and difference-of-convex-sos decompositions for polynomials. arXiv :1803.09900 (2018)
- Niu, Y.S., Glowinski, R.: Discrete dynamical system approaches for boolean polynomial optimization. Journal of Scientific Computing 92(2), 1–39 (2022)
- Niu, Y.S., Júdice, J., Le Thi, H.A., Pham, D.T.: Solving the quadratic eigenvalue complementarity problem by DC programming. In: Modelling, Computation and Optimization in Information Systems and Management Sciences, Advances in Intelligent Systems and Computing, pp. 203–214. Springer (2015)
- Niu, Y.S., Júdice, J., Le Thi, H.A., Pham, D.T.: Improved dc programming approaches for solving the quadratic eigenvalue complementarity problem. Applied Mathematics and Computation 353, 95–113 (2019)
- Niu, Y.S., Pham, D.T.: DC programming approaches for BMI and QMI feasibility problems. In: Advanced Computational Methods for Knowledge Engineering, Advances in Intelligent Systems and Computing, pp. 37–63. Springer (2014)
- Niu, Y.S., Pham, D.T., Le Thi, H.A., Júdice, J.: Efficient DC programming approaches for the asymmetric eigenvalue complementarity problem. Optimization Methods and Software 28(4), 812–829 (2013)
- Niu, Y.S., Wang, Y.J., Le Thi, H.A., Pham, D.T.: Higher-order moment portfolio optimization via an accelerated difference-of-convex programming approach and sums-ofsquares. arXiv :1906.01509 (2019)
- 43. Niu, Y.S., Zhang, H.: Power-product matrix: nonsingularity, sparsity and determinant. Linear and Multilinear Algebra 0(0), 1–18 (2023)
- de Oliveira, W., Tcheou, M.P.: An inertial algorithm for dc programming. Set-Valued and Variational Analysis 27(4), 895–919 (2019)
- 45. Parlett, B.N.: The Symmetric Eigenvalue Problem. SIAM (1998)
- 46. Pham, D.T., Le Thi, H.A.: Convex analysis approach to DC programming: theory, algorithms and applications. Acta mathematica vietnamica 22(1), 289–355 (1997)
- Pham, D.T., Le Thi, H.A.: A DC optimization algorithm for solving the trust-region subproblem. SIAM Journal on Optimization 8(2), 476–505 (1998)

- Pham, D.T., Le Thi, H.A., Pham, V.N., Niu, Y.S.: DC programming approaches for discrete portfolio optimization under concave transaction costs. Optimization letters 10(2), 261–282 (2016)
- Pham, D.T., Niu, Y.S.: An efficient DC programming approach for portfolio decision with higher moments. Computational Optimization and Applications 50(3), 525–554 (2011)
- Pham, D.T., Souad, E.B.: Duality in DC (difference of convex functions) optimization. subgradient methods. In: Trends in Mathematical Optimization: 4th French-German Conference on Optimization, International Series of Numerical Mathematics, pp. 277– 293. Springer (1988)
- 51. Phan, D.N., Le, H.M., Le Thi, H.A.: Accelerated Difference of Convex functions Algorithm and its Application to Sparse Binary Logistic Regression. In: Proceedings of the 27th International Joint Conference on Artificial Intelligence and the 23rd European Conference on Artificial Intelligence, pp. 1369–1375 (2018)
- Polyak, B.T.: Some methods of speeding up the convergence of iteration methods. Ussr computational mathematics and mathematical physics 4(5), 1–17 (1964)
- 53. Reznick, B.A.: Sum of Even Powers of Real Linear Forms. American Mathematical Society (1992)
- Song, Y., Yu, G.: Properties of solution set of tensor complementarity problem. Journal of Optimization Theory and Applications 170(1), 85–96 (2016)
- 55. Wen, B., Chen, X., Pong, T.K.: A proximal difference-of-convex algorithm with extrapolation. Computational optimization and applications **69**(2), 297–324 (2018)
- You, Y., Niu, Y.S.: A refined inertial dc algorithm for dc programming. Optimization and Engineering 24(1), 65–91 (2023)