

A note on distinct differences in t -intersecting families

Jagannath Bhanja and Sayan Goswami

The Institute of Mathematical Sciences, A CI of Homi Bhabha National Institute, C.I.T.

Campus, Taramani, Chennai 600113, India

jbbhanja@imsc.res.in, sayangoswami@imsc.res.in

Abstract

For a family \mathcal{F} of subsets of $\{1, 2, \dots, n\}$, let $\mathcal{D}(\mathcal{F}) = \{F \setminus G : F, G \in \mathcal{F}\}$ be the collection of all (setwise) differences of \mathcal{F} . The family \mathcal{F} is called a t -intersecting family, if for some positive integer t and any two members $F, G \in \mathcal{F}$ we have $|F \cap G| \geq t$. The family \mathcal{F} is simply called intersecting if $t = 1$. Recently, Frankl proved an upper bound on the size of $\mathcal{D}(\mathcal{F})$ for the intersecting families \mathcal{F} . In this note we extend the result of Frankl to t -intersecting families.

2020 Mathematics Subject Classification: 05D05

Keywords: Setwise difference, t -intersecting family, Erdős-Ko-Rado theorem

1. Introduction

We denote the standard n -element set $\{1, 2, \dots, n\}$ by $[n]$, the set of all subsets of $[n]$ by $2^{[n]}$, and $\binom{[n]}{k}$ by the collection of all k -element subsets of $[n]$ for $0 \leq k \leq n$. We also use the standard notation $|S|$ for the cardinality of a set S and $\lfloor N \rfloor$ for the largest integer less than or equal to N .

A family \mathcal{F} of subsets of $[n]$ is said to be a t -intersecting family for some positive integer t , if $|F \cap G| \geq t$ for any two members $F, G \in \mathcal{F}$. A 1-intersecting family is simply called an intersecting family. The Erdős-Ko-Rado theorem [1] gives the tight upper bound on the size of t -intersecting families.

Theorem 1 (Erdős-Ko-Rado theorem for t -intersecting family). *There exists some $n_0(k, t)$ such that if $n \geq n_0(k, t)$ and $\mathcal{F} \subset \binom{[n]}{k}$ is t -intersecting, then*

$$|\mathcal{F}| \leq \binom{n-t}{k-t}.$$

The upper bound is tight for $n_0(k, t) = (t+1)(k-t+1)$; this was proved by Frankl [3] for $t \geq 15$ and for all t using a different technique by Wilson [6].

For a family \mathcal{F} , let

$$\mathcal{D}(\mathcal{F}) = \{F \setminus G : F, G \in \mathcal{F}\}$$

be the collection of all (setwise) differences of \mathcal{F} . Here we allow the empty set \emptyset to be in \mathcal{F} when $\mathcal{F} \neq \emptyset$. In this direction Marica and Schönheim [5] proved the following theorem.

Theorem 2 (Marica-Schönheim theorem). *For a nonempty family $\mathcal{F} \subset 2^{[n]}$ one has*

$$|\mathcal{D}(\mathcal{F})| \geq |\mathcal{F}|.$$

Recently, Frankl [2] proved the following upper bound for $\mathcal{D}(\mathcal{F})$ when \mathcal{F} is an intersecting family.

Theorem 3 (Frankl). *Suppose that $\mathcal{F} \subset \binom{[n]}{k}$ is an intersecting family with $n \geq k(k+3)$. Then*

$$|\mathcal{D}(\mathcal{F})| \leq \binom{n-1}{k-1} + \binom{n-1}{k-2} + \cdots + \binom{n-1}{0}.$$

For further developments in this direction see [4].

In this note we extend Theorem 3 to the t -intersecting families \mathcal{F} of $\binom{[n]}{k}$. More precisely we prove the following theorem.

Theorem 4. *Suppose that $\mathcal{F} \subset \binom{[n]}{k}$ is a t -intersecting family with $n \geq (k-t)(k+t+3) \cdot \lfloor \left(\frac{k+t+3}{k+t+1}\right)^{t-1} \rfloor$. Then*

$$|\mathcal{D}(\mathcal{F})| \leq \binom{n-t}{k-t} + \binom{n-t}{k-t-1} + \cdots + \binom{n-t}{0}.$$

2. Preliminaries

To prove Theorem 4 we require the following notions.

For a family $\mathcal{F} \subset 2^{[n]}$ and a non-negative integer ℓ let

$$\mathcal{D}^{(\ell)}(\mathcal{F}) = \left\{ D \in \binom{[n]}{\ell} : \exists F, G \in \mathcal{F}, F \setminus G = D \right\}.$$

Note that

$$|\mathcal{D}(\mathcal{F})| = \sum_{\ell \geq 0} |\mathcal{D}^{(\ell)}(\mathcal{F})|.$$

We call a family \mathcal{F} to be a t -star if there is some t -subset contained in every member of \mathcal{F} . Similarly, we call \mathcal{F} to be the full t -star all elements of $\binom{[n]}{k}$ containing a fix t -subset are the only elements of the family \mathcal{F} .

Lemma 1. *Suppose that the family $\mathcal{F} \subset \binom{[n]}{k}$ is t -intersecting and that $\mathcal{D}^{(k-t)}(\mathcal{F})$ has $k+t+2$ members that are pairwise disjoint. Then \mathcal{F} is a t -star.*

Proof. We can assume that $k \geq t+1$, as the case $k = t$ is trivial. Let $D_0, D_1, \dots, D_{k+t+1}$ be members of $\mathcal{D}^{(k-t)}(\mathcal{F})$ that are pairwise disjoint. As $D_i = F_i \setminus F'_i$ for some $F_i, F'_i \in \mathcal{F}$ and \mathcal{F} is t -intersecting, for each D_i there exist some subset X_i of $2^{[n]}$ with $|X_i| = t$ and $D_i \cup X_i \in \mathcal{F}$. Observe that each $x \in X_0$ can contain in at most one D_i , without loss of generality let $D_{k+2}, D_{k+3}, \dots, D_{k+t+1}$ be such D_i 's. Therefore, $(D_0 \cup X_0) \cap D_i = \emptyset$ for all $i = 1, 2, \dots, k+1$.

However, as $D_i \cup X_i \in \mathcal{F}$ and \mathcal{F} is a t -intersecting family, we have in particular that $|(D_0 \cup X_0) \cap (D_i \cup X_i)| \geq t$. But, D_i 's are pairwise disjoint and X_0 has nothing common with D_i for $1 \leq i \leq k+1$, the only possibility is that $X_i \subset (D_0 \cup X_0)$ for $1 \leq i \leq k+1$.

Further, as $|(D_1 \cup X_1) \cap (D_i \cup X_i)| \geq t$ for $1 \leq i \leq k+1$, we have $X_1 \cap D_i = \emptyset$, otherwise we would have $D_1 \cap D_i \neq \emptyset$ or $(D_0 \cup X_0) \cap D_1 \neq \emptyset$ giving contradictions. By similar argument $X_i \cap D_1 = \emptyset$. Thus, $X_1 = X_i$ for $1 \leq i \leq k+1$.

Our claim is that $X_1 \in F$ for all $F \in \mathcal{F}$. Suppose that $X_1 \not\subset F$ for some $F \in \mathcal{F}$. Then $|(D_i \cup X_i) \cap F| \geq t$ implies $|D_i \cap F| \neq \emptyset$ for all $1 \leq i \leq k+1$, which further implies $|\mathcal{F}| \geq k+1$, a contradiction. \square

If $\mathcal{F} \subset \binom{[n]}{k}$ is a t -star with $X \subset F$ for all $F \in \mathcal{F}$, then $\mathcal{D}^{(\ell)}(\mathcal{F}) \subset \binom{[n] \setminus X}{\ell}$ for all $0 \leq \ell \leq k-t$. Thus, for t -stars $|\mathcal{D}(\mathcal{F})| \leq \sum_{\ell=0}^{k-t} |\mathcal{D}^{(\ell)}(\mathcal{F})|$. The equality holds when \mathcal{F} is the full t -star.

Corollary 1. *It is sufficient to prove Theorem 4 for families \mathcal{F} with $\mathcal{D}^{(k-t)}(\mathcal{F})$ not containing $k+t+2$ pairwise disjoint members.*

3. The proof of Theorem 4

Proof of Theorem 4. Define $f(k, t) = (k-t)(k+t+3) \lfloor \left(\frac{k+t+3}{k+t+1} \right)^{t-1} \rfloor$. Choose a family $\mathcal{D} \subset \binom{[n]}{k-t}$ such that it has $(k+t+3) \lfloor \left(\frac{k+t+3}{k+t+1} \right)^{t-1} \rfloor$ members that are pairwise disjoint. The selection of \mathcal{D} has done in an uniform way. Then the expected value

$$\mathbb{E}(|\mathcal{D} \cap \mathcal{D}^{(k-t)}(\mathcal{F})|) = (k+t+3) \left\lfloor \left(\frac{k+t+3}{k+t+1} \right)^{t-1} \right\rfloor \frac{1}{\binom{n}{k-t}} |\mathcal{D}^{(k-t)}(\mathcal{F})|.$$

But by Corollary 1 we have

$$(k+t+3) \left\lfloor \left(\frac{k+t+3}{k+t+1} \right)^{t-1} \right\rfloor \frac{1}{\binom{n}{k-t}} |\mathcal{D}^{(k-t)}(\mathcal{F})| \leq k+t+1.$$

Thus,

$$\begin{aligned}
|\mathcal{D}^{(k-t)}(\mathcal{F})| &\leq \frac{(k+t+1)^t}{(k+t+3)^t} \binom{n}{k-t} \\
&= \frac{(k+t+1)^t}{(k+t+3)^t} \cdot \frac{n \cdot (n-1) \cdots (n-t+1)}{(n-k+t) \cdot (n-k+t-1) \cdots (n-k+1)} \cdot \binom{n-t}{k-t} \\
&< \frac{(k+t+1)^t}{(k+t+3)^t} \cdot \left(\frac{n}{n-k+1} \right)^t \cdot \binom{n-t}{k-t}.
\end{aligned}$$

As the function $(\frac{n}{n-k+1})^t$ is decreasing on n we see that $(\frac{n}{n-k+1})^t \leq (\frac{f(k,t)}{f(k,t)-k+1})^t \leq (\frac{k+t+2}{k+t+1})^t$. Therefore we have

$$\begin{aligned}
|\mathcal{D}^{(k-t)}(\mathcal{F})| &< \left(\frac{k+t+2}{k+t+3} \right)^t \binom{n-t}{k-t} \\
&< \frac{k+t+2}{k+t+3} \binom{n-t}{k-t} \\
&= \left(1 - \frac{1}{k+t+3} \right) \binom{n-t}{k-t}.
\end{aligned} \tag{1}$$

For all other values of $\ell = 0, 1, \dots, k-t-1$, as $\mathcal{F} \subset \binom{[n]}{k}$ is t -intersecting, we have

$$|\mathcal{D}^{(0)}(\mathcal{F})| \leq 1 = \binom{n-t}{0}, \tag{2}$$

and

$$|\mathcal{D}^{(\ell)}(\mathcal{F})| \leq \binom{n-t+1}{\ell} = \binom{n-t}{\ell} + \binom{n-t}{\ell-1} \tag{3}$$

for $\ell = 1, \dots, k-t-1$. Observe that

$$\sum_{\ell=0}^{k-t-2} \binom{n-t}{\ell} \leq (k-t-1) \binom{n-t}{k-t-2} < \frac{1}{k+t+3} \binom{n-t}{k-t}, \tag{4}$$

where the latter inequality can be easily verified by just expanding the binomial coefficients. Hence, combining the equations (1), (2), (3), and (4) we obtain

$$\begin{aligned}
|\mathcal{D}(\mathcal{F})| &= \sum_{\ell=0}^{k-t} |\mathcal{D}^{(\ell)}(\mathcal{F})| \\
&< \sum_{\ell=0}^{k-t-1} |\mathcal{D}^{(\ell)}(\mathcal{F})| + \left(1 - \frac{1}{k+t+3}\right) \binom{n-t}{k-t} \\
&= \sum_{\ell=0}^{k-t-1} \binom{n-t}{\ell} + \sum_{\ell=0}^{k-t-2} \binom{n-t}{\ell} + \left(1 - \frac{1}{k+t+3}\right) \binom{n-t}{k-t} \\
&< \sum_{\ell=0}^{k-t-1} \binom{n-t}{\ell} + \frac{1}{k+t+3} \binom{n-t}{k-t} + \left(1 - \frac{1}{k+t+3}\right) \binom{n-t}{k-t} \\
&= \sum_{\ell=0}^{k-t} \binom{n-t}{\ell}.
\end{aligned}$$

This completes the proof of the theorem. □

Acknowledgment

Both the authors wish to thank the Institute of Mathematical Sciences, Chennai for the financial support received through the institute postdoctoral program.

References

- [1] P. Erdős, C. Ko, and R. Rado, Intersection theorems for systems of finite sets, Q. J. Math. 12 (1961), 310–320.
- [2] P. Frankl, On the number of distinct differences in an intersecting family, Disc. Math. 344 (2) (2021), Paper No. 112210.
- [3] P. Frankl, The Erdős–Ko–Rado theorem is true for $n = ckt$, Combinatorics (Proc. Fifth Hungarian Colloq., Keszthely, 1976), Vol. I, 365–375, Colloq. Math. Soc. János Bolyai, 18, North–Holland, 1978.
- [4] P. Frankl, S. Kiselev, and A. Kupavskii, Best possible bounds on the number of distinct differences in intersecting families, European J. Combin. 107 (2023), Paper No. 103601.

- [5] J. Marica and J. Schönheim, Differences of sets and a problem of graham, *Canad. Math. Bull.* 12 (1969), 635–637.
- [6] R. M. Wilson, The exact bound in the Erdős–Ko–Rado theorem, *Combinatorica*, 4 (1984), 247–257.