# Maximum Independent Set when excluding an induced minor: $K_1 + tK_2$ and $tC_3 \uplus C_4$

Édouard Bonnet ⊠☆®

Univ Lyon, CNRS, ENS de Lyon, Université Claude Bernard Lyon 1, LIP UMR5668, France

Univ Lyon, CNRS, ENS de Lyon, Université Claude Bernard Lyon 1, LIP UMR5668, France

Univ Lyon, CNRS, ENS de Lyon, Université Claude Bernard Lyon 1, LIP UMR5668, France

Stéphan Thomassé ⊠

Univ Lyon, CNRS, ENS de Lyon, Université Claude Bernard Lyon 1, LIP UMR5668, France

Alexandra Wesolek 

□

□

Simon Fraser University, Burnaby, BC, Canada

#### Abstract

Dallard, Milanič, and Štorgel [arXiv '22] ask if for every class excluding a fixed planar graph H as an induced minor, Maximum Independent Set can be solved in polynomial time, and show that this is indeed the case when H is any planar complete bipartite graph, or the 5-vertex clique minus one edge, or minus two disjoint edges. A positive answer would constitute a far-reaching generalization of the state-of-the-art, when we currently do not know if a polynomial-time algorithm exists when H is the 7-vertex path. Relaxing tractability to the existence of a quasipolynomial-time algorithm, we know substantially more. Indeed, quasipolynomial-time algorithms were recently obtained for the t-vertex cycle,  $C_t$  [Gartland et al., STOC '21] and the disjoint union of t triangles,  $tC_3$  [Bonamy et al., SODA '23].

We give, for every integer t, a polynomial-time algorithm running in  $n^{O(t^5)}$  when H is the friend-ship graph  $K_1+tK_2$  (t disjoint edges plus a vertex fully adjacent to them), and a quasipolynomial-time algorithm running in  $n^{O(t^2 \log n)+t^{O(1)}}$  when H is  $tC_3 \uplus C_4$  (the disjoint union of t triangles and a 4-vertex cycle). The former extends a classical result on graphs excluding  $tK_2$  as an induced subgraph [Alekseev, DAM '07], while the latter extends Bonamy et al.'s result.

2012 ACM Subject Classification Theory of computation o Graph algorithms analysis

**Keywords and phrases** Maximum Independent Set, forbidden induced minors, quasipolynomial-time algorithms

**Funding** This work was supported by the ANR projects TWIN-WIDTH (ANR-21-CE48-0014) and Digraphs (ANR-19-CE48-0013).

Alexandra Wesolek: Supported by the Vanier Canada Scholarship Program.

Acknowledgements We thank Dibyayan Chakraborty for useful discussions.

### 1 Introduction

The MAX INDEPENDENT SET (MIS for short) problem asks, in its optimization form, for a largest *independent set* of its input graph G, i.e., a subset of pairwise non-adjacent vertices in G. In its decision form, the input is a graph G and an integer k, and the question is whether G admits an independent set of size at least k.

Besides the ubiquitous usefulness that such a fundamental problem has within combinatorial optimization, and notably in the areas of packing, scheduling, and coloring, MIS (or equivalently MAXIMUM CLIQUE, the same problem in the complement graph) has as wide a range of applications as possible, as evidenced for instance in map labeling [47],

coding theory [17], spatial scheduling [26], genetic analysis [1], information retrieval [10], macromolecular docking [28], and sociometry [27] (also see Butenko's thesis [16]). It is thus unfortunate that this problem is not only hard to solve but also very reluctant to being approximated. Indeed, the decision version of MIS is NP-complete [29] and W[1]-complete [25], while its optimization version cannot be approximated within ratio  $n^{1-\varepsilon}$  on n-vertex graphs, for any  $\varepsilon > 0$ , unless P=NP [35, 49].

In spite of this, theorists and practitioners have put a lot of effort in designing efficient algorithms for MIS. In parallel to generically solving MIS via integer programming, highly performant exact and heuristic MIS solvers have emerged in the recent years, based on diverse methods such as kernelization and evolutionary approaches [40], deep reinforcement learning [4], graph neural networks [44], and dataless training (where backpropagation is applied to a loss function based instead on the input) [9]. On the theory side, exact exponential algorithms have been developed for decades culminating in a running time below  $1.2^n n^{O(1)}$  [48].

Another approach is to try and exploit the structure that the input graphs may have. Indeed, in all the aforementioned applications, inputs are not uniformly sampled over all n-vertex graphs: They instead bare some structural properties, and in some cases, might avoid some specific patterns. Graph theory<sup>1</sup> offers two main notions of patterns or containment: the natural and straightforward subgraphs (obtained by removing vertices and edges), and the deeper minors (further allowing to contract edges). Both notions come with an induced variant, when edge removals are disallowed, bringing the number of containment types to four. It is then sensible to determine the patterns H whose absence makes MIS (more) tractable. It turns out that this question is completely settled for subgraphs and minors.

For the subgraph containment, the argument is the following. By the grid minor theorem [45], the class of graphs excluding H as a subgraph has bounded treewidth if (and only if) all the connected components of H are paths and subdivided claws (i.e., stars with three leaves); thus MIS can be solved in polynomial-time in this class, for instance by Courcelle's theorem [21]. If instead H has a connected component which is not a path nor a subdivided claw, MIS remains NP-complete since such a class either contains all subcubic graphs or the 2|V(H)|-subdivision of every graph, two families of graphs on which MIS is known to be NP-complete [43, 5, 8].

For minors, the dichotomy relies on the planarity of H. Indeed, if H is planar, then the class of graphs excluding H as a minor has bounded treewidth (again, mainly by the grid minor theorem), and MIS can be solved efficiently. If H is non-planar, then the H-minor-free graphs include all planar graphs for which MIS is known to be NP-complete [30].

The question is more intriguing for the induced containments, and the *induced subgraph* case has received a lot of attention. While it is known for a long time that if H is not the disjoint union of paths and subdivided claws, MIS remains NP-complete on graphs without H as an induced subgraph [43, 5], it has been conjectured that MIS is otherwise polynomial-time solvable. This has been proven when H is the 6-vertex path [34], a claw with exactly one edge subdivided [6, 41], or any disjoint union of claws [15]. The latter result extends a polynomial-time algorithm due to Alekseev when H is any disjoint union of edges [7]. The author indeed proves that the total number of maximal independent sets is polynomially bounded, and can be enumerated in polynomial time.

While we currently do not know of a polynomial-time algorithm when H is the 7-vertex path, Gartland and Lokshtanov [31] have obtained a quasipolynomial-time algorithm when

<sup>&</sup>lt;sup>1</sup> We refer the reader to Section 2 for the relevant background in graph theory.

H is  $P_t$ , the t-vertex path, for any positive integer t; also see [42]. Supporting the existence of at least a quasipolynomial-time algorithm when H is  $S_{i,j,k}$ , the claw whose three edges are subdivided i-1, j-1, and k-1 times, respectively, a quasipolynomial-time approximation scheme (QPTAS) [20] and a polynomial-time algorithm among bounded-degree graphs [2] have been proposed. The parameterized complexity of MIS when excluding a fixed induced subgraph has been studied [22, 12, 13], but the mere statement of which H make the problem fixed-parameter tractable (and which ones keep it W[1]-complete) is unclear [13].

We eventually arrive at the *induced minor* containment, the topic of the current paper. As for minors, the class of all graphs excluding a non-planar graph H as an induced minor contains all planar graphs; hence MIS remains NP-complete in such a class. However we do not know of a *planar* graph H, for which MIS remains NP-complete on H-induced-minor-free graphs. This has led Dallard, Milanič, and Štorgel [23] to ask if such classes exist:

▶ Question 1. Is it true that for every planar graph H, MAX INDEPENDENT SET can be solved in polynomial time in the class of graphs excluding H as an induced minor?

A first observation is that avoiding H as an induced minor implies avoiding it as an induced subgraph. Thus Question 1 is settled for  $P_6$ ,  $S_{1,1,2}$ , and  $tS_{1,1,1}$  (where tG denotes the disjoint of t copies of G). The same authors [23] further obtain a polynomial-time algorithm when H is  $K_5^-$  (the 5-vertex clique minus an edge),  $K_{2,t}$  (the bipartite complete graph with 2 vertices fully adjacent to t vertices), and  $W_4 = K_1 + C_4$  (a 4-vertex cycle  $C_4$  with a fifth vertex fully adjacent to the cycle). All three cases were shown by bounding the so-called tree-independence number (i.e., treewidth where bag size is replaced by independence number of the subgraph induced by the bag) [23], in which case a polynomial-time algorithm can be derived for MIS using the corresponding tree-decompositions [24]. They also show that this is as far as this sole technique can go: H-induced-minor-free graphs have bounded tree-independence number if and only if H is edgeless or an induced minor of  $K_5^-$ ,  $K_{2,t}$ , or  $W_4$  [23]. The framework of potential maximal cliques [14] and the container method has led to a polynomial-time algorithm when  $H = C_5$  [19, 3].

Question 1 a beautiful question, and, if true, a very difficult one. Indeed,  $H = P_7$ , the 7-vertex path, is a very simple planar graph for which we currently do not know such a polynomial-time algorithm. A natural relaxation of Question 1 is to only request a quasipolynomial-time algorithm:

▶ Question 2. Is it true that for every planar graph H, MAX INDEPENDENT SET can be solved in quasipolynomial time in the class of graphs excluding H as an induced minor?

We know somewhat more about Question 2. There is a quasipolynomial-time algorithm for MIS in  $C_t$ -induced-minor-free graphs [32], building upon the  $H = P_t$  case. Recently, Bonamy et al. [11] present a quasipolynomial-time algorithm when H is  $tC_3$ , i.e., the disjoint union of t triangles. See Table 1 for a summary of the introduction.

Expecting an affirmative solution to Question 1 or Question 2 may seem optimistic. However, as far as precise running time is concerned, we do know that for every planar H, MIS is probably not as difficult in H-induced-minor-free graphs as it is in general graphs. Indeed, Korhonen [39] describes a  $2^{O(n/\log^c n)}$ -time algorithm, for some constant c > 0, to solve MIS on n-vertex graphs excluding a fixed planar graph H as an induced minor. Assuming the Exponential-Time Hypothesis [36], such a running time is impossible in general graphs [37].

Our results. We make some progress regarding Questions 1 and 2. Our first contribution is, for every positive integer t, a polynomial-time algorithm when H is the  $friendship\ graph$ 

H exluded as	subgraph	minor	induced subgraph	induced minor
in P	$tS_{t,t,t}$	planar	$P_6, S_{1,1,2}, tS_{1,1,1}$	$K_5^-, K_{2,t}, W_4, C_5, K_1 + tK_2$
only known in QP	_	_	$P_t \ (t \geqslant 7)$	$C_t \ (t \geqslant 6), \ tC_3 \uplus C_4$
NP-c	$\neg t S_{t,t,t}$	non-planar	$\neg tS_{t,t,t}$	non-planar
open P vs NP-c	_	_	$P_7, \ldots$	$P_7, C_6, \ldots$
open QP vs NP-c	_	_	$S_{1,1,3}, S_{1,2,2} \dots$	$C_4 \uplus C_4, \ldots$

**Table 1** The complexity of MAX INDEPENDENT SET when H is excluded as one of the four main types of patterns. " $\neg tS_{t,t,t}$ " means that for no t does H is a subgraph of  $tS_{t,t,t}$ . Our results are framed.

 $K_1 + tK_2$  (also called *Dutch windmill graph* or fan), i.e., t independent edges universally linked to a 2t + 1-st vertex:

▶ **Theorem 1.** MAX INDEPENDENT SET can be solved in polynomial-time  $n^{O(t^5)}$  in n-vertex  $K_1 + tK_2$ -induced-minor-free graphs.

This extends Alekseev's result [7] for graphs excluding  $tK_2$  as an induced subgraph, or equivalently, as an induced minor. We indeed use this result to first derive a polynomial-time algorithm in subgraphs of  $K_1 + tK_2$ -induced-minor-free graphs G induced by vertices from a bounded number of breadth-first search (BFS) layers of G.

We then consider the connected components of our input graph G when deprived of a subset X of vertices inducing  $tK_2$  and, subject to that property, maximizing the size of the largest connected component in G-X. We show that, due to this careful selection of X, every component C of G-X admits an efficiently constructible path-decomposition  $\mathcal{P}$  with bounded adhesion (i.e., any two distinct bags have a bounded intersection), each bag of which is contained in a bounded number of consecutive BFS layers of C. Hence MIS can be solved efficiently within a bag, by our opening step (see previous paragraph). This part is quite technical, but mostly to justify the existence of  $\mathcal{P}$ . The algorithm itself remains simple.

Section 3 is then obtained by exhaustively finding X and guessing its intersection X' with a maximum independent set of G, and performing dynamic programming on the connected components of G - X, deprived of N(X'). The dynamic-programming table is filled via the efficient algorithm when handling an induced subgraph contained in few BFS layers.

Our second contribution is a quasipolynomial-time algorithm when H is,  $tC_3 \uplus C_4$ , the disjoint union of t triangles and a 4-vertex cycle:

▶ **Theorem 2.** MAX INDEPENDENT SET can be solved in quasipolynomial-time  $n^{O(t^2 \log n) + t^{O(1)}}$  in  $tC_3 \uplus C_4$ -induced-minor-free graphs.

We first perform a quasipolynomial branching step to get rid of holes of size at most 6 (i.e., induced cycles of length 4, 5, or 6). We then assume that the input graph G is not  $(t+2)C_3$ -induced-minor-free, for otherwise we conclude with Bonamy et al.'s algorithm [11]. Thus G, being  $tC_3 \uplus C_4$ -induced-minor-free, has to admit  $(t+2)C_3$  as an induced subgraph, i.e., a collection  $T_1, \ldots, T_{t+2}$  of t+2 pairwise vertex-disjoint and non-adjacent triangles. We define  $S_{i,j}$ , minimally separating  $T_i$  and  $T_j$  in the graph G deprived of the neighborhoods of the other triangles  $T_k$  (with  $k \neq i, k \neq j$ ).

We show that each  $S_{i,j}$  induces a clique. So does every intersection  $N_{i,j}$  of the neighborhood of two distinct triangles  $T_i, T_j$  of the collection (this is where getting rid of the holes of length at most 6 comes into play). We can therefore exhaustively guess the intersection of

a maximum independent set with the union of the sets  $S_{i,j}$  and  $N_{i,j}$  (for every  $i < j \in [t+2]$ ). We finally observe that  $G' = G - \bigcup_{i \neq j \in [t+2]} S_{i,j} \cup N_{i,j}$  is chordal, since the presence of a hole H in G' would imply the existence in G of t independent triangles in the non-neighborhood of H, a contradiction to the  $tC_3 \uplus C_4$ -induced-minor-freeness of G. We thus conclude by using a classic algorithm for MIS in chordal graphs [33, 46].

In Section 2 we introduce the relevant graph-theoretic background. In Section 3 we prove Theorem 1, and in Section 4 we prove Theorem 2.

#### 2 Preliminaries

If  $i \leq j$  are two integers, we denote by [i,j] the set of integers  $\{i,i+1,\ldots,j-1,j\}$ , and by [i], the set [1,i]. We denote by V(G) and E(G) the set of vertices and edges of a graph G, respectively. We denote by  $G_1 \simeq G_2$  the fact that the two graphs  $G_1$  and  $G_2$  are isomorphic, i.e., equal up to renaming their vertex set. For  $S \subseteq V(G)$ , the subgraph of G induced by G, denoted G[S], is obtained by removing from G all the vertices that are not in G (together with their incident edges). Then G - S is a short-hand for  $G[V(G) \setminus S]$ . G is an induced subgraph of G if there is an  $G \subseteq V(G)$  such that  $G[S] \subseteq H$ .

For G a graph and  $X \subseteq V(G)$ ,  $E_G(X)$  (or simply E(X)) is a short-hand for E(G[X]). For G a graph and  $X,Y \subseteq V(G)$  two disjoint sets,  $E_G(X,Y)$  denotes the set of edges of E(G) with one endpoint in X and the other endpoint in Y. We denote by  $N_G(v)$  and  $N_G[v]$ , the open, respectively closed, neighborhood of v in G. For  $S \subseteq V(G)$ , we set  $N_G(S) := \bigcup_{v \in S} N_G(v) \setminus S$  and  $N_G[S] := N_G(S) \cup S$ . We may omit the subscript if G is clear from the context. A connected component is a maximal connected induced subgraph.

Two cycles  $C_1, C_2$  are said *independent* if they are vertex-disjoint and there is no edge between  $C_1$  and  $C_2$ . A collection of cycles is *independent* if they are pairwise independent. Two vertex subsets  $X, Y \subseteq V(G)$  touch if  $X \cap Y \neq \emptyset$  or there is an edge  $uv \in E(G)$  with  $u \in X$  and  $v \in Y$ . Then two (or more) cycles are independent if and only if they do not touch. We say that  $X, Y \subseteq V(G)$  touch in Z if  $X \cap Y \cap Z \neq \emptyset$  or there is an edge  $uv \in E(G)$  with  $u \in X \cap Z$  and  $v \in Y \cap Z$ , or equivalently, if  $X \cap Z$  and  $Y \cap Z$  touch.

A graph H is a *induced minor* of a graph G if H can be obtained from G by a sequence of vertex deletions and edge contractions. A *minor* is the same but also allows edge deletions. Equivalently an induced minor H –with vertex set, say,  $\{v_1, \ldots, v_{V(H)}\}$  – of G can be defined as a vertex-partition  $B_1, \ldots, B_{|V(H)|}$  of an induced subgraph of G, such that every  $G[B_i]$  is connected and  $v_i v_j \in E(H)$  if and only if  $E_G(B_i, B_j) \neq \emptyset$  (i.e., when the disjoint sets  $B_i$  and  $B_j$  touch). Observe indeed that contracting each  $B_i$  into a single vertex (possible since each  $B_i$  induces a connected subgraph) results in H. A graph G (resp. a graph class) is said H-induced-minor-free if H is not an induced minor of G (resp. no graph of the class admits H as an induced minor).

We denote by  $C_{\ell}$  the  $\ell$ -vertex cycle, and by  $K_{\ell}$ , the  $\ell$ -vertex clique. A *hole* is an induced cycle of length at least four. A graph is *chordal* if it has no hole. For two disjoint sets  $X, Y \subseteq V(G)$  in a graph G, an (X,Y)-separator is a (possibly empty) set  $S \subseteq V(G) \setminus (X \cup Y)$  such that there is no path between X and Y in G - S. An (X,Y)-separator is *minimal* if no proper subset of it is itself an (X,Y)-separator.

The disjoint union  $G_1 \uplus G_2$  of two graphs  $G_1, G_2$  has vertex set  $V(G_1) \uplus V(G_2)$  and edge set  $E(G_1) \uplus E(G_2)$ , where  $V(G_1) \uplus V(G_2)$  presupposes that the vertices of  $G_1$  and  $G_2$  have disjoint labels. If  $t \geqslant 2$  is an integer and G a graph, tG is the graph  $G \uplus (t-1)G$ , and 1G is simply G. The join  $G_1 + G_2$  of two graphs  $G_1, G_2$  has vertex set  $V(G_1) \uplus V(G_2)$  and edge set  $E(G_1) \uplus E(G_2) \uplus \{uv : u \in V(G_1), v \in V(G_2)\}$ . In other words, the join of  $G_1$  and  $G_2$ 

is obtained from their disjoint union by adding all possible edges between  $G_1$  and  $G_2$ .

A breadth-first search (BFS) layering in G from a vertex  $v \in V(G)$  (or from a connected set  $S \subseteq V(G)$ ) is a partition of the remaining vertices into  $L_1, L_2, \ldots$  such that every vertex of  $L_i$  is at distance exactly i of v (or S). Such an  $L_i$  is called a BFS layer of G (from v, or from S). Note that there cannot be an edge in G between  $L_i$  and  $L_j$  if |i-j| > 1.

A path-decomposition  $\mathcal{P} = (B_1, \ldots, B_h)$  of a graph G is such that

6

- $\blacksquare$  for every  $e \in E(G)$ , there is some  $B_i$  that contains both endpoints of e, and
- whenever  $v \in B_i \cap B_j$  with i < j, v is also in all  $B_k$  with i < k < j.

The sets  $B_i$  (for  $i \in [h]$ ) are called the bags of  $\mathcal{P}$ , and the sets  $B_i \cap B_{i+1}$  (for  $i \in [h-1]$ ) the adhesions of  $\mathcal{P}$ . Path-decomposition  $\mathcal{P}$  has maximum adhesion p if all of its adhesions have size at most p. Note that the adhesion  $B_i \cap B_{i+1}$ , if disjoint from  $B_1 \cup B_h$ , is a vertex cutset disconnecting  $B_1$  from  $B_h$ .

## **3** Polynomial algorithm in $K_1 + tK_2$ -induced-minor-free graphs

We first show how to solve MAX INDEPENDENT SET in  $K_1 + tK_2$ -induced-minor-free graphs of bounded diameter. More generally, we show the following.

▶ Lemma 3. Let t, h be fixed non-negative integers. Let G be a  $K_1 + tK_2$ -induced-minor-free n-vertex graph, and  $L_0 \subseteq V(G)$  such that  $G[L_0]$  is connected. Let  $L_1, L_2, \ldots, L_h$  be such that every vertex of  $L_i$ , for any  $i \in [h]$ , is at distance exactly i of  $L_0$ . Then MAX INDEPENDENT SET can be solved in polynomial time  $n^{(2t-1)h+O(1)}$  on any induced subgraph of  $G[\bigcup_{1 \le i \le h} L_i]$ .

**Proof.** For every  $j \in [h]$ ,  $G[L_j]$  has no  $tK_2$  induced subgraph (or equivalently, induced minor). Indeed,  $G[\bigcup_{0 \le i \le j-1} L_i]$  is a connected graph, hence  $\bigcup_{0 \le i \le j-1} L_i$  can be contracted to a single vertex, and every vertex in  $L_j$  has at least one neighbor in  $L_{j-1}$ . Therefore a  $tK_2$  induced subgraph in  $G[L_j]$  would contradict the  $K_1 + tK_2$ -induced-minor-freeness of G.

Let H be any induced subgraph of  $G[\bigcup_{1 \leq i \leq h} L_i]$ . In particular  $H[L_j]$  has also no  $tK_2$  induced subgraph, for every  $j \in [h]$ . Hence, by a classical result of Alekseev [7],  $H[L_j]$  has at most  $n^{2t-1}$  maximal independent sets, which can be listed in time  $n^{2t+O(1)}$ .

We thus exhaustively list every h-tuple  $(I_1, \ldots, I_h)$  where, for every  $j \in [h]$ ,  $I_j$  is a maximal independent set of  $H[L_j]$ , in time  $n^{(2t-1)h+O(1)}$ . Note that if there is an edge in H between  $L_i$  and  $L_j$ , then  $|i-j| \leq 1$ . As each  $I_j$  (for  $j \in [h]$ ) is an independent set,  $H' = H[\bigcup_{j \in [h]} I_j]$  is a bipartite graph as witnessed by the bipartition  $(L_1 \cup L_3 \cup \ldots, L_2 \cup L_4 \cup \ldots)$ . A maximum independent set I can thus be computed in polynomial time in H'. Indeed, by the Kőnig-Egerváry theorem [38], finding a maximum independent set in a bipartite graph boils down to finding a maximum matching, which can be done in polynomial time (and now, even in almost linear time [18]) by solving a maximum flow problem. We output the largest independent set I found among every run.

The correctness of the algorithm is based on the observation that a maximum independent set  $I^*$  of H intersects every  $L_i$  (for  $i \in [h]$ ) at an independent set  $J_i$ , which by definition is contained in a maximal independent set  $I_i$  of  $H[L_i]$ . In the run when every maximal independent set  $I_i$  is guessed correctly, we obtain an independent set of the cardinality of  $I^*$ .

We say that G is reduced if it does not contain degree-1 vertices. If e = uv is an edge of G, let  $G \setminus e$  (resp.  $S \setminus e$ ,  $S \cup e$ , for some  $S \subseteq V(G)$ ) be the induced subgraph  $G[V(G) - \{u, v\}] = G - \{u, v\}$  (resp. the sets  $S \setminus \{u, v\}$ ,  $S \cup \{u, v\}$ ). More generally, for a

collection  $e_1 = u_1 v_1, \ldots, e_k = u_k v_k$  of edges of G, we denote by  $G \setminus \{e_1, \ldots, e_k\}$  the induced subgraph  $G[V(G) - \{u_1, v_1, \ldots, u_k, v_k\}] = G - \{u_1, v_1, \ldots, u_k, v_k\}.$ 

- ▶ Lemma 4. Let G be a reduced connected  $K_1 + tK_2$ -induced-minor-free n-vertex graph. Let  $X \subseteq V(G)$  maximize the size of a largest component of G' := G X, among those sets X such that  $G[X] \simeq tK_2$ . Then for any  $e \in E(G' N_G(X))$  contained in a connected component C of G',
- 1.  $C \setminus e$  is disconnected, and
- **2.** each connected component of  $C \setminus e$  contains a vertex in  $N_G(X)$ .

**Proof.** As G is reduced, every vertex of X has degree at least two (in G). Thus G' cannot be connected, for otherwise, contracting in G the set V(G') to a single vertex would form a  $K_1 + tK_2$  induced minor. We thus know that G' has at least two connected components.

Let  $C_1$  be a largest connected component of G'. Since G is connected, there exists a shortest path P in G from  $V(C_1)$  to  $V(G') \setminus V(C_1)$ . Say, that P ends in the connected component  $C_2 \neq C_1$  of G'. Path P has to have some internal vertices in X, but since  $G[X] \simeq tK_2$ , it follows that there is an edge  $e^*$  in E(X) (but not necessarily in P) incident to both  $N_G(V(C_1))$  and  $N_G(V(C_2))$ .

For every edge  $e \in E(G' - N_G(X))$  in component C (possibly  $C_1$ ),  $C \setminus e$  is disconnected. Indeed, for the sake of contradiction, suppose that  $C \setminus e$  is connected, and consider  $X' := (X \setminus e^*) \cup e$ . By assumption,  $G[X'] \cong tK_2$ . Furthermore, the connected component G - X' containing  $e^*$  is strictly larger than  $C_1$ , as it contains  $(V(C_1) \setminus e) \cup e^*$  and  $V(C_2) \setminus e$ , disjoint sets of size at least  $|V(C_1)|$  and at least one, respectively. This contradicts the maximality of X, and establishes the first item.

We now prove the second item, also by contradiction. Suppose that there is a connected component C' of  $C \setminus e$  that does not contain a vertex in  $N_G(X)$ . We will reach a contradiction by showing that C' contains an edge e' not incident to  $N_G(X)$ , and such that  $C \setminus e'$  is connected (and conclude in light of the previous paragraph).

Let  $L_i \subseteq V(C')$  be the *i-th neighborhood of e* in C', i.e., the vertices at distance *i* of one endpoint of *e*, and at least *i* of the other endpoint. We consider the *last layer*  $L_k$ , i.e., such that  $L_k \neq \emptyset$  and  $L_{k+1} = \emptyset$ . If  $L_k$  contains an edge e' then removing the endpoints of this edge does not disconnect C' (and hence C) since each vertex in  $L_k$  has a neighbor in  $L_{k-1}$  (with the convention that  $L_0$  consists of the endpoints of *e*) and  $G[L_0 \cup L_1 \cup \cdots \cup L_{k-1}]$  is connected.

If  $L_k$  does not contain an edge, then each vertex in  $L_k$  has two neighbors in  $L_{k-1}$ . This is because G is reduced, and by assumption that no vertex of C' has a neighbor in X. Hence, removing the endpoints of any edge e' incident to a vertex in  $L_k$  does not disconnect C' (nor C), since each vertex in  $L_k$  has at least one neighbor in  $L_{k-1}$  which is not an endpoint of e'. In either case, e' is an edge of  $C - N_G(X)$  that does not disconnect C, which we showed is not possible.

We now prove the main technical result of the section.

- ▶ Proposition 5. Let G be a reduced connected  $K_1 + tK_2$ -induced-minor-free n-vertex graph. Let  $X \subseteq V(G)$  maximize the size of a largest component of G' := G X, among those sets X such that  $G[X] \simeq tK_2$ . Then for every connected component C of G', a path-decomposition  $\mathcal{P}$  of C such that
- every bag of  $\mathcal{P}$  is contained in  $O(t^4)$  consecutive BFS layers, and
- every adhesion of  $\mathcal{P}$  is of size at most  $2t^2$ , can be computed in time  $n^{O(1)}$ .

**Proof.** Let v be a vertex in  $N_G(X) \cap V(C)$  and, for any positive integer s, let  $L_s$  be the vertices at distance s from v in C. Let q be the largest distance between v and a vertex of C. We set  $f(t) := t^2(6t^2 + 2) = O(t^4)$ . We will show that, for every  $s \in [q - f(t)]$ , there is a vertex cutset of size at most  $2t^2$  separating  $L_s$  from  $L_{s+f(t)}$ .

We show that any long-enough induced path (such as a shortest path from  $L_s$  to  $L_{s+f(t)}$ ) has some edges with both endpoints in  $V(C) \setminus N_G(X)$ .

 $\triangleright$  Claim 6. Any induced path P in C contains less than  $3t^2$  vertices in  $N_G(X)$ .


Proof. If there are  $3t^2$  vertices on P with a neighbor in X, then there are at least 3t vertices  $w_1, \ldots, w_{3t}$  on P that are neighbors of a fixed edge  $e \in E(X)$ . For each  $i \in [t]$ , contract every edge of P between  $w_{3i-2}$  and  $w_{3i-1}$  but one, say  $e_i$ . Contract e, and call z the resulting vertex. The vertex z and the t edges  $e_i$  contradict the fact that G is  $K_1 + tK_2$ -induced-minor-free.

<

For an edge  $e \in E(C)$  we denote by  $\operatorname{dist}(e, v)$  the length of a shortest path from an endpoint of e to v. We build a collection of paths  $P_i$  and edges  $e_i \in E(P_i)$ , for  $i = 1, 2, \ldots$ , while they are well-defined, in the following way.

Let  $P_1$  be a shortest path from  $L_s$  to  $L_{s+f(t)}$ . Let  $e_1 \in E(P_1)$  minimize  $e \mapsto \operatorname{dist}(e,v)$ , among those edges of  $P_1$  with both endpoints in  $V(C) \setminus N_G(X)$ . By Claim 6 there are less than  $6t^2$  edges on  $P_1$  with an endpoint in  $N_G(X)$ , hence  $\operatorname{dist}(e_1,v) \leqslant s+6t^2$ . We denote by  $Q_1$  the maximal subpath of  $P_1$  starting in  $L_s$  and not containing an endpoint of  $e_1$  (that is, stopping just before reaching an endpoint of  $e_1$ ). For the next iteration, we work in  $C \setminus e_1$  (recall that this stands for C deprived of the two endpoints of  $e_1$ ).

We now describe in general the *i*-th iteration. Let  $P_i$  be a shortest path from  $L_s$  to  $L_{s+f(t)}$  in  $C \setminus \{e_1, e_2, \ldots, e_{i-1}\}$ . Let  $e_i$  be the first edge of  $P_i$  (when starting from  $L_s$ ) such that  $\operatorname{dist}(e_i, v) \geqslant \operatorname{dist}(e_{i-1}, v) + 2$  and  $e_i$  has no endpoint in  $N_G(X)$ . Note that by Claim 6,  $\operatorname{dist}(e_i, v) \leqslant \operatorname{dist}(e_{i-1}, v) + 2 + 6t^2$ . Let  $Q_i$  be the maximal subpath of  $P_i$  starting in  $L_s$  and not containing an endpoint of  $e_i$ . See Figure 1.



**Figure 1** Illustration of X (for t = 6) and the first f(t) layers (from left to right) of the connected component C of G - X rooted at v. Vertices of C with a neighbor in X are filled. We represented the first three iterations:  $P_1, e_1$  (in red),  $P_2, e_2$  (in blue),  $P_3, e_3$  (in dark green). Not to clutter the figure, we do not represent all the edges, and we code the different labels: edges  $e_i$  are squiggly, and subpaths  $Q_i$  are thicker.

Let  $e_1, \ldots, e_k$  be the eventual collection of edges. In principle, the while loop stops when:

- (i)  $L_s$  is disconnected from  $L_{s+f(t)}$  in  $C \setminus \{e_1, \ldots, e_k\}$ , or
- (ii) there is no edge  $e_{k+1} \in E(P_{k+1})$  such that  $\operatorname{dist}(e_{k+1}, v) \geqslant \operatorname{dist}(e_k, v) + 2$  and  $e_{k+1}$  has no endpoint in  $N_G(X)$ .

 $\triangleright$  Claim 7. It holds that  $k \le t^2$ . In particular, as  $f(t) = t^2(6t^2 + 2)$ , (ii) cannot hold.

Proof. Assume for the sake of contradiction that  $k \ge t^2 + 1$ . By Lemma 4 (using both items), for each i there exists a vertex  $v_i \in V(C) \cap N_G(X)$  disconnected from v in  $C \setminus e_i$ . Since  $k \ge t^2 + 1$ , there are t + 1 vertices  $v_{a_1}, \ldots, v_{a_{t+1}}$  all adjacent to an endpoint of the same fixed edge  $e \in E(X)$ . Let C' be the component containing v in  $C \setminus \{e_{a_1}, \ldots, e_{a_t}\}$ , and let us contract (in G) set V(C') to a single vertex, say w. Since  $e_{a_{t+1}}$  is, by construction, an edge of C', vertex w is adjacent to an endpoint of e.

Observe also that, for every  $i \in [t]$ , each subpath  $Q_{a_i}$  is contained in C'. Vertex w is adjacent to an endpoint  $u_{a_i}$  of  $e_{a_i}$ , for every  $i \in [t]$ . Let  $C_{a_i}$  be the vertex set of the connected component of  $C \setminus e_{a_i}$  containing  $v_{a_i}$ . Let  $R_{a_i}$  be a path from  $u_{a_i}$  to  $v_{a_i}$  in the subgraph of G induced by  $C_{a_i}$  plus the endpoints of  $e_{a_i}$ . For each  $i \in [t]$ , if  $R_{a_i}$  has at least two edges, contract the edge  $wu_{a_i}$ , and all the edges of  $R_{a_i}$  but the last two; the last one being  $f_{a_i} = y_{a_i}v_{a_i}$ . Finally contract e, and the edge between e and w. We call the resulting vertex z.

We claim that z and the edges  $f_{a_i}$  make a  $K_1 + tK_2$  induced minor in G. One can see that z is adjacent to  $v_{a_i}$  via  $e \in E(X)$ , and to  $y_{a_i}$  via w and the path  $R_{a_i}$ . We shall justify that the edges  $f_{a_i}$  form an induced matching in G. Indeed, suppose there is an edge between some  $f_{a_i}$  and  $f_{a_j}$  with  $i \neq j \in [t]$ . Then in  $C \setminus e_{a_i}$ , there is a path from v to  $v_{a_i}$  via  $Q_j$ , a contradiction.

Claim 7 implies that the  $2k \leq 2t^2$  endpoints of  $e_1, \ldots, e_k$  form a vertex cutset disconnecting  $L_s$  from  $L_{s+f(t)}$ . We can now build the path-decomposition  $\mathcal{P}$ . Recall that the BFS search from v gives rise to q layers,  $L_1, \ldots, L_q$  (outside v).

For  $j \in [\lfloor q/f(t) \rfloor]$ , let  $S_j$  be the vertex cutset of size at most  $2t^2$  (and obtained as detailed above) disconnecting  $L_{(j-1)f(t)+1}$  from  $L_{jf(t)}$ . We denote by  $L_{s \to s'}$  the set  $\bigcup_{s \leqslant h \leqslant s'} L_h$ .

- Let  $B_1 \subseteq V(C)$  consist of v,  $S_1$ , plus all the vertices of connected components of  $C[L_{1\to f(t)}] S_1$  that do not intersect  $L_{f(t)}$ .
- For j going from 2 to  $\lfloor q/f(t) \rfloor 1$ , let  $B_j \subseteq V(C)$  consist of  $S_j \cup S_{j+1}$  plus the vertices not already present in one of  $B_1, \ldots, B_{j-1}$  of all the connected components of

$$C[L_{(i-1)}f(t)+1\to (i+1)}f(t)]-(S_i\cup S_{i+1})$$

that do not intersect  $L_{(j+1)f(t)}$ .

■ Let finally  $B_{\lfloor q/f(t)\rfloor} \subseteq V(C)$  consist of  $S_{\lfloor q/f(t)\rfloor}$  plus the vertices of all the connected components of  $C[L_{(\lfloor q/f(t)\rfloor-1)f(t)+1\to q}] - S_{\lfloor q/f(t)\rfloor}$  that intersect  $L_q$ .

Let  $\mathcal{P}$  be the path-decomposition  $(B_1, B_2, \ldots, B_{\lfloor q/f(t) \rfloor})$ .  $\mathcal{P}$  is indeed a path-decomposition, since our process entirely covers V(C), and by virtue of  $S_j$  separating  $L_{(j-1)f(t)+1}$  from  $L_{if(t)}$ . By construction,

- every bag intersects at most  $2f(t) = O(t^4)$  layers of BFS from vertex v, and
- for every  $j \in [\lfloor q/f(t) \rfloor 1]$ ,  $B_j \cap B_{j+1} = S_j$ , so every adhesion has size at most  $2t^2$ .

Finally note that once X is found, one can find the path-decomposition  $\mathcal{P}$  of C in time  $n^{O(1)}$ , since this only involves computing (at most n) shortest paths.

We can now wrap up, using Proposition 5 and Lemma 3.

▶ **Theorem 1.** MAX INDEPENDENT SET can be solved in polynomial-time  $n^{O(t^5)}$  in n-vertex  $K_1 + tK_2$ -induced-minor-free graphs.

**Proof.** Let G be our  $K_1+tK_2$ -induced-minor-free n-vertex input graph. As including vertices of degree 1 in the independent set is a safe reduction rule, we can assume that G is reduced. By dealing with the possibly several connected components of G separately, we can further

assume that G is connected. In time  $n^{O(t)}$  we find  $X \subseteq V(G)$  that maximizes the size of a largest component of G' := G - X, among those sets X such that  $G[X] \simeq tK_2$ .

We exhaustively guess the intersection X' of a fixed maximum independent set of G with the set X, with an extra multiplicative factor of  $2^{2t}$ . We are now left with solving MIS separately in  $C' := C - N_G(X')$  for each connected component C of G'. By Proposition 5, we obtain in time  $n^{O(1)}$  a path-decomposition  $\mathcal{P} = (B_1, \ldots, B_p)$  of C', such that

- every  $B_i$  (for  $i \in [p]$ ) is contained in  $O(t^4)$  consecutive BFS layers of C, and
- every adhesion  $A_i := B_i \cap B_{i+1}$  (for  $i \in [p-1]$ ) is of size at most  $2t^2$ .

Indeed, removing  $N_G(X')$  from C (and its path-decomposition) preserves those properties.

Let us define  $A_0, A_p$  to be empty. We proceed to the following dynamic programming. For  $i \in [0, p]$ , and for any  $S \subseteq A_i$ , T[i, S] is meant to eventually contain an independent set I of  $C'[\bigcup_{1 \le j \le i} B_j]$  of maximum cardinality among those such that  $I \cap B_i = S$ . We set  $T[0, \emptyset] = \emptyset$ , and observe that it is the only entry of the form  $T[0, \cdot]$ .

We fill this table by increasing value of  $i=1,2\ldots,p$ . Assume that all entries of the form  $T[i',\cdot]$  are properly filled for i'< i. For every  $S\subseteq A_i, T[i,S]$  is filled in the following way. For every  $S'\subseteq A_{i-1}$ , if  $S\cup S'$  is an independent set, we compute, by Lemma 3 (with  $L_1,\ldots,L_h$  being the  $O(t^4)$  consecutive BFS layers of C containing  $B_i$ , and  $L_0$  being the connected set, in C, formed by the union of all the previous layers), a maximum independent set  $I_i$  in  $C'[B_i]-N[S\cup S']$  in time  $n^{O(ht)}=n^{O(t^5)}$ . We finally set  $T[i,S]=T[i-1,S']\cup I_i\cup S$  for a run that maximizes the cardinality of  $T[i-1,S']\cup I_i$ .

It takes time  $p \cdot 2^{O(t^2)} \cdot n^{O(t^5)} = n^{O(t^5)}$  to completely fill T. Eventually  $T[p, \emptyset]$  contains a maximum independent set of C'. We return the union of X' and of the maximum independent sets of C' found for each connected component C of G'. The overall running time is  $n^{O(t^5)}$ .

# **4** Quasipolynomial algorithm in $tC_3 \uplus C_4$ -induced-minor-free graphs

We first show that the existence of a short hole allows for a quasipolynomial-time branching scheme in  $tC_3 \uplus C_4$ -induced-minor-free graphs.

▶ Lemma 8. Let G be an n-vertex  $tC_3 \uplus C_4$ -induced-minor-free graph, and some fixed constant  $\ell$ . While there is a hole H of length at most  $\ell$  and a  $tC_3$  induced subgraph in G, MAX INDEPENDENT SET admits a quasipolynomial branching step, running in  $n^{O_{\ell}(t \log n)}$ .

**Proof.** Let  $C(G) := \{X \in {V(G) \choose 3t} : G[X] \simeq tC_3\}$  be the collection of vertex subsets inducing t disjoint triangles, and assume  $\mu(G) := |C(G)| > 0$ , and H is a hole of G of length at most  $\ell$ . As G is  $tC_3 \uplus C_4$ -induced-minor-free, N[V(H)] intersects every  $X \in C(G)$ . In particular, there is a vertex  $v \in V(H)$  such that N[v] intersects at least a  $1/\ell$  fraction of the  $X \in C(G)$ .

We branch on two options: either we take v in an (initially empty) solution, and remove its closed neighborhood from G, or we remove v from G (without adding it to the solution). With the former choice, measure  $\mu$  drops by at least a  $1/\ell$  fraction (and the number of vertices of G decreases by at least 1), and with the latter choice, the number of vertices drops by 1. This branching is exhaustive. We simply need to argue about its running time.

Note that each option can be done at most n times, while the first option cannot be done more than  $\log_{\ell}(n^{3t}) = O_{\ell}(t\log n)$  times. Hence the branching tree has at most  $\binom{n}{\log_{\ell}(n^{3t})} = n^{O_{\ell}(t\log n)}$  leaves.

The previous lemma permits us to get rid of short holes, which turns out useful in some corner case.

▶ **Theorem 2.** MAX INDEPENDENT SET can be solved in quasipolynomial-time  $n^{O(t^2 \log n) + t^{O(1)}}$  in  $tC_3 \uplus C_4$ -induced-minor-free graphs.


**Proof.** We apply the quasipolynomial branching step of Lemma 8 with  $\ell = 6$ , until the input n-vertex graph G has no longer holes of length at most 6, or  $tC_3$  induced subgraph.

In time  $n^{O(t)}$ , we exhaustively look for a collection of pairwise vertex-disjoint and non-adjacent triangles  $T_1, T_2, \ldots, T_{t+2}$  in G. If such a collection does not exist, G is  $(t+2)C_3$ -induced-minor-free. Indeed, the absence of  $(t+2)C_3$  as an induced subgraph implies that at least one of the (t+2) independent cycles realizing a  $(t+2)C_3$  induced minor is of length at least four. This is ruled out by the assumption that G is  $tC_3 \uplus C_4$ -induced-minor-free. (Note here that only (t+1) independent cycles would suffice.)

We can thus assume that a collection  $T_1, T_2, \ldots, T_{t+2}$  exists, for otherwise, we can conclude with the quasipolynomial-time algorithm, running in  $n^{O(t^2 \log n) + t^{O(1)}}$ , of Bonamy et al. [11] for MAX INDEPENDENT SET in graphs with a bounded number of independent cycles (here,  $(t+2)C_3$ -induced-minor-free). In turn, as G contains a  $tC_3$  (even  $(t+2)C_3$ ) induced subgraph, we can, in light of the first paragraph, further assume that all the cycles of G have length either 3 or at least 7. We refer the reader to Figure 2 for a visual summary of the next two paragraphs.

For every pair  $T_i, T_j$  (with  $i < j \in [t+2]$ ), let  $S_{i,j}$  be a minimal  $(T_i, T_j)$ -separator in  $G - \bigcup_{k \in [t+2] \setminus \{i,j\}} N(T_k)$ . We first observe that  $S_{i,j}$  is a clique of G. Suppose, for the sake of contradiction, that  $u, v \in S_{i,j}$  are distinct and non-adjacent. Consider  $P_{u,i}, P_{v,i}, P_{u,j}, P_{v,j}$  four chordless paths between u and  $T_i, v$  and  $T_i, u$  and  $T_j, v$  and  $T_j$ , respectively, all of which intersect  $S_{i,j}$  exactly once. These paths exist since  $S_{i,j}$  is a minimal  $(T_i, T_j)$ -separator. Since  $P_{u,i}, P_{v,i}$  touch in  $T_i$ , and  $P_{u,j}, P_{v,j}$  touch in  $T_j$ , we can build a cycle C out of those four paths. There is by assumption no edge between u and v, nor between  $(P_{u,i} \cup P_{v,i}) \setminus \{u, v\}$  and  $(P_{u,j} \cup P_{v,j}) \setminus \{u, v\}$ . Therefore, a subset of V(C) induces a hole H. As C, hence H, does not intersect the neighborhood of  $\bigcup_{k \in [t+2] \setminus \{i,j\}} T_k$ , the set  $\{V(H)\} \cup \{T_k : k \in [t+2] \setminus \{i,j\}\}$  contradicts the  $tC_3 \uplus C_4$ -induced-minor-freeness of G.

Let  $N_{i,j}$  be the set  $N(T_i) \cap N(T_j)$  for each pair  $i < j \in [t+2]$ . Observe that the sets  $N_{i,j}$  need not be disjoint, and that when  $T_i$  and  $T_j$  are at distance at least 3 apart,  $N_{i,j}$  is empty. We notice that  $N_{i,j}$  is a clique, for otherwise we can exhibit an induced cycle of length 4, 5, or 6 in G (hence a hole of length at most 6).



**Figure 2** Illustration of the sets  $S_{i,j}$  and  $N_{i,j}$ , and the paths  $P_{u,i}$  (bottom) and  $P_{v,i}$  (top) in blue, and the paths  $P_{u,j}$  (bottom) and  $P_{v,j}$  (top) in red. A non-edge uv in  $S_{i,j}$  would imply, from these paths, that a hole exists in the non-neighborhood of the other triangles  $T_k$ , contradicting  $tC_3 \uplus C_4$ -minor-freeness. The absence of hole of length at most 6 implies that  $N_{i,j}$  is also a clique.

We claim that  $G' := G - (\bigcup_{i < j \in [t+2]} S_{i,j} \cup N_{i,j})$  is chordal. Indeed assume there is a hole H' in G'. The  $tC_3 \uplus C_4$ -induced-minor-freeness implies that H' intersects at least two sets

 $N[T_i]$  and  $N[T_j]$ . Thus, let P be a subpath of H' whose endpoints are in two distinct  $N(T_i)$  and  $N(T_j)$ , and no internal vertex of P lies in some  $N[T_k]$  with  $k \notin \{i, j\}$ . As G' does not include any vertex of  $\bigcup_{i' < j' \in [t+2]} N_{i',j'}$ , it further holds that no vertex of P (at all) lies in some  $N[T_k]$  with  $k \notin \{i, j\}$ . Therefore, path P contradicts that  $S_{i,j}$  separates  $T_i$  and  $T_j$  in  $G - \bigcup_{k \in [t+2] \setminus \{i,j\}} N(T_k)$ .

We can now describe the rest of the algorithm after the collection  $T_1, T_2, \ldots, T_{t+2}$  is found. We greedily compute the minimal separators  $S_{i,j}$ . We exhaustively try every subset  $S \subseteq \bigcup_{i < j \in [t+2]} S_{i,j} \cup N_{i,j}$  that is an independent set. Such a set S contains at most one vertex in each  $S_{i,j}$  and each  $N_{i,j}$ , as we have established that each  $S_{i,j}$  and each  $N_{i,j}$  form a clique. Hence there are  $n^{O(t^2)}$  such sets S. For each S, we compute a maximum independent set I in the chordal graph  $G - (N[S] \cup \bigcup_{i < j \in [t+2]} S_{i,j} \cup N_{i,j})$  in linear time (see [33, 46]). We finally output the set  $S \cup I$  maximizing  $|S \cup I|$ . Note that the overall running time is  $n^{O(t^2 \log n) + t^{O(1)}}$ .

#### References

- 1 Kuruvilla J. Abraham and Clara Diaz. Identifying large sets of unrelated individuals and unrelated markers. *Source code for biology and medicine*, 9(1):1–8, 2014.
- 2 Tara Abrishami, Maria Chudnovsky, Cemil Dibek, and Paweł Rzazewski. Polynomial-time algorithm for maximum independent set in bounded-degree graphs with no long induced claws. In Joseph (Seffi) Naor and Niv Buchbinder, editors, Proceedings of the 2022 ACM-SIAM Symposium on Discrete Algorithms, SODA 2022, Virtual Conference / Alexandria, VA, USA, January 9 12, 2022, pages 1448–1470. SIAM, 2022. doi:10.1137/1.9781611977073.61.
- 3 Tara Abrishami, Maria Chudnovsky, Marcin Pilipczuk, Paweł Rzążewski, and Paul D. Seymour. Induced subgraphs of bounded treewidth and the container method. In Dániel Marx, editor, Proceedings of the 2021 ACM-SIAM Symposium on Discrete Algorithms, SODA 2021, Virtual Conference, January 10 13, 2021, pages 1948–1964. SIAM, 2021. doi:10.1137/1.9781611976465.116.
- 4 Sungsoo Ahn, Younggyo Seo, and Jinwoo Shin. Learning what to defer for maximum independent sets. In *Proceedings of the 37th International Conference on Machine Learning, ICML 2020, 13-18 July 2020, Virtual Event*, volume 119 of *Proceedings of Machine Learning Research*, pages 134–144. PMLR, 2020. URL: http://proceedings.mlr.press/v119/ahn20a.html.
- 5 Vladimir E. Alekseev. The effect of local constraints on the complexity of determination of the graph independence number. *Combinatorial-algebraic methods in applied mathematics*, pages 3–13, 1982.
- 6 Vladimir E. Alekseev. Polynomial algorithm for finding the largest independent sets in graphs without forks. *Discrete Applied Mathematics*, 135(1-3):3–16, 2004. doi:10.1016/S0166-218X(02)00290-1.
- Vladimir E. Alekseev. An upper bound for the number of maximal independent sets in a graph. 17(4):355-359, 2007. URL: https://doi.org/10.1515/dma.2007.030 [cited 2023-02-03], doi:doi:10.1515/dma.2007.030.
- Paola Alimonti and Viggo Kann. Some APX-completeness results for cubic graphs. *Theoretical Computer Science*, 237(1):123 134, 2000. doi:https://doi.org/10.1016/S0304-3975(98) 00158-3.
- 9 Ismail R. Alkhouri, George K. Atia, and Alvaro Velasquez. A differentiable approach to the maximum independent set problem using dataless neural networks. *Neural Networks*, 155:168–176, 2022.
- J. Gary Augustson and Jack Minker. An analysis of some graph theoretical cluster techniques. Journal of the ACM (JACM), 17(4):571–588, 1970.
- Marthe Bonamy, Édouard Bonnet, Hugues Déprés, Louis Esperet, Colin Geniet, Claire Hilaire, Stéphan Thomassé, and Alexandra Wesolek. Sparse graphs with bounded induced cycle

- packing number have logarithmic treewidth. In *Proceedings of the 2023 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 3006–3028. SIAM, 2023.
- Édouard Bonnet, Nicolas Bousquet, Pierre Charbit, Stéphan Thomassé, and Rémi Watrigant. Parameterized complexity of independent set in H-free graphs. In Christophe Paul and Michał Pilipczuk, editors, 13th International Symposium on Parameterized and Exact Computation, IPEC 2018, August 20-24, 2018, Helsinki, Finland, volume 115 of LIPIcs, pages 17:1-17:13. Schloss Dagstuhl Leibniz-Zentrum für Informatik, 2018. doi:10.4230/LIPIcs.IPEC.2018.17.
- Édouard Bonnet, Nicolas Bousquet, Stéphan Thomassé, and Rémi Watrigant. When maximum stable set can be solved in FPT time. In Pinyan Lu and Guochuan Zhang, editors, 30th International Symposium on Algorithms and Computation, ISAAC 2019, December 8-11, 2019, Shanghai University of Finance and Economics, Shanghai, China, volume 149 of LIPIcs, pages 49:1-49:22. Schloss Dagstuhl Leibniz-Zentrum für Informatik, 2019. doi: 10.4230/LIPIcs.ISAAC.2019.49.
- Vincent Bouchitté and Ioan Todinca. Treewidth and minimum fill-in: Grouping the minimal separators. SIAM J. Comput., 31(1):212–232, 2001. doi:10.1137/S0097539799359683.
- Andreas Brandstädt and Raffaele Mosca. Maximum weight independent set for ℓclaw-free graphs in polynomial time. Discrete Applied Mathematics, 237:57–64, 2018. doi:10.1016/j.dam.2017.11.029.
- Sergiy Butenko. Maximum independent set and related problems, with applications. University of Florida, 2003.
- 17 Sergiy Butenko, Panos M. Pardalos, Ivan Sergienko, Vladimir Shylo, and Petro Stetsyuk. Finding maximum independent sets in graphs arising from coding theory. In Gary B. Lamont, Hisham Haddad, George A. Papadopoulos, and Brajendra Panda, editors, *Proceedings of the 2002 ACM Symposium on Applied Computing (SAC), March 10-14, 2002, Madrid, Spain*, pages 542–546. ACM, 2002. doi:10.1145/508791.508897.
- Li Chen, Rasmus Kyng, Yang P. Liu, Richard Peng, Maximilian Probst Gutenberg, and Sushant Sachdeva. Maximum flow and minimum-cost flow in almost-linear time. In 63rd IEEE Annual Symposium on Foundations of Computer Science, FOCS 2022, Denver, CO, USA, October 31 November 3, 2022, pages 612–623. IEEE, 2022. doi:10.1109/F0CS54457.2022.00064.
- Maria Chudnovsky, Marcin Pilipczuk, Michał Pilipczuk, and Stéphan Thomassé. On the maximum weight independent set problem in graphs without induced cycles of length at least five. SIAM J. Discret. Math., 34(2):1472–1483, 2020. doi:10.1137/19M1249473.
- 20 Maria Chudnovsky, Marcin Pilipczuk, Michał Pilipczuk, and Stéphan Thomassé. Quasi-polynomial time approximation schemes for the maximum weight independent set problem in H-free graphs. In Proceedings of the 2020 ACM-SIAM Symposium on Discrete Algorithms, SODA 2020, Salt Lake City, UT, USA, January 5-8, 2020, pages 2260–2278, 2020. doi: 10.1137/1.9781611975994.139.
- 21 Bruno Courcelle. The monadic second-order logic of graphs. i. recognizable sets of finite graphs. Inf. Comput., 85(1):12-75, 1990. doi:10.1016/0890-5401(90)90043-H.
- 22 Konrad K. Dabrowski, Vadim V. Lozin, Haiko Müller, and Dieter Rautenbach. Parameterized complexity of the weighted independent set problem beyond graphs of bounded clique number. J. Discrete Algorithms, 14:207–213, 2012. doi:10.1016/j.jda.2011.12.012.
- Clément Dallard, Martin Milanič, and Kenny Štorgel. Treewidth versus clique number. III. tree-independence number of graphs with a forbidden structure. *CoRR*, abs/2206.15092, 2022. arXiv:2206.15092, doi:10.48550/arXiv.2206.15092.
- 24 Clément Dallard, Martin Milanič, and Kenny Štorgel. Treewidth versus clique number. II. tree-independence number, 2021. URL: https://arxiv.org/abs/2111.04543, doi:10.48550/ ARXIV.2111.04543.
- Rodney G. Downey and Michael R. Fellows. Fixed-parameter tractability and completeness II: on completeness for W[1]. *Theor. Comput. Sci.*, 141(1&2):109–131, 1995. doi:10.1016/0304-3975(94)00097-3.

- 14
- 26 Duncan Eddy and Mykel J. Kochenderfer. A maximum independent set method for scheduling earth-observing satellite constellations. Journal of Spacecraft and Rockets, 58(5):1416–1429,
- 27 Elaine Forsyth and Leo Katz. A matrix approach to the analysis of sociometric data: preliminary report. Sociometry, 9(4):340-347, 1946.
- Eleanor J. Gardiner, Peter Willett, and Peter J. Artymiuk. Graph-theoretic techniques for macromolecular docking. J. Chem. Inf. Comput. Sci., 40(2):273-279, 2000. doi:10.1021/ ci990262o.
- 29 Michael R. Garey and David S. Johnson. Computers and Intractability: A Guide to the Theory of NP-Completeness. W. H. Freeman, 1979.
- Michael R. Garey, David S. Johnson, and Larry J. Stockmeyer. Some simplified NP-complete graph problems. Theor. Comput. Sci., 1(3):237-267, 1976. doi:10.1016/0304-3975(76) 90059-1.
- Peter Gartland and Daniel Lokshtanov. Independent set on  $P_k$ -free graphs in quasi-polynomial time. In Sandy Irani, editor, 61st IEEE Annual Symposium on Foundations of Computer Science, FOCS 2020, Durham, NC, USA, November 16-19, 2020, pages 613-624. IEEE, 2020. doi:10.1109/FOCS46700.2020.00063.
- Peter Gartland, Daniel Lokshtanov, Marcin Pilipczuk, Michał Pilipczuk, and Paweł Rzążewski. Finding large induced sparse subgraphs in  $C_{>t}$ -free graphs in quasipolynomial time. In Samir Khuller and Virginia Vassilevska Williams, editors, STOC '21: 53rd Annual ACM SIGACT Symposium on Theory of Computing, Virtual Event, Italy, June 21-25, 2021, pages 330-341. ACM, 2021. doi:10.1145/3406325.3451034.
- 33 Fănică Gavril. The intersection graphs of subtrees in trees are exactly the chordal graphs. Journal of Combinatorial Theory, Series B, 16(1):47–56, 1974.
- Andrzej Grzesik, Tereza Klimosová, Marcin Pilipczuk, and Michał Pilipczuk. Polynomial-time 34 algorithm for maximum weight independent set on  $P_6$ -free graphs. ACM Trans. Algorithms, 18(1):4:1-4:57, 2022. doi:10.1145/3414473.
- Johan Håstad. Clique is hard to approximate within  $n^{1-\varepsilon}$ . In Acta Mathematica, pages 35 627-636, 1996.
- 36 Russell Impagliazzo and Ramamohan Paturi. On the complexity of k-sat. J. Comput. Syst. Sci., 62(2):367-375, 2001. doi:10.1006/jcss.2000.1727.
- Russell Impagliazzo, Ramamohan Paturi, and Francis Zane. Which problems have strongly exponential complexity? J. Comput. Syst. Sci., 63(4):512-530, 2001. doi:10.1006/jcss.2001.
- 38 Denés Kőnig. Gráfok és mátrixok. matematikai és fizikai lapok, 38: 116-119, 1931.
- Tuukka Korhonen. Grid induced minor theorem for graphs of small degree. CoRR, abs/2203.13233, 2022. arXiv:2203.13233, doi:10.48550/arXiv.2203.13233.
- Sebastian Lamm, Peter Sanders, Christian Schulz, Darren Strash, and Renato F. Werneck. Finding near-optimal independent sets at scale. J. Heuristics, 23(4):207-229, 2017. doi: 10.1007/s10732-017-9337-x.
- Vadim V. Lozin and Martin Milanič. A polynomial algorithm to find an independent set of maximum weight in a fork-free graph. J. Discrete Algorithms, 6(4):595-604, 2008. doi: 10.1016/j.jda.2008.04.001.
- Marcin Pilipczuk, Michał Pilipczuk, and Paweł Rzążewski. Quasi-polynomial-time algorithm for independent set in  $P_t$ -free graphs via shrinking the space of induced paths. In Hung Viet Le and Valerie King, editors, 4th Symposium on Simplicity in Algorithms, SOSA 2021, Virtual Conference, January 11-12, 2021, pages 204-209. SIAM, 2021. doi:10.1137/1.9781611976496.
- 43 Svatopluk Poljak. A note on stable sets and colorings of graphs. Commentationes Mathematicae Universitatis Carolinae, 15(2):307–309, 1974.
- Thomas Pontoizeau, Florian Sikora, Florian Yger, and Tristan Cazenave. Neural maximum independent set. In Machine Learning and Principles and Practice of Knowledge Discovery in

- Databases International Workshops of ECML PKDD 2021, Virtual Event, September 13-17, 2021, Proceedings, Part I, volume 1524 of Communications in Computer and Information Science, pages 223–237. Springer, 2021. doi:10.1007/978-3-030-93736-2\\_18.
- **45** Neil Robertson and Paul D. Seymour. Graph minors. V. Excluding a planar graph. *J. Comb. Theory, Ser. B*, 41(1):92–114, 1986. doi:10.1016/0095-8956(86)90030-4.
- Donald J. Rose, Robert E. Tarjan, and George S. Lueker. Algorithmic aspects of vertex elimination on graphs. SIAM J. Comput., 5(2):266–283, 1976. doi:10.1137/0205021.
- Bram Verweij and Karen Aardal. An optimisation algorithm for maximum independent set with applications in map labelling. In Jaroslav Nesetril, editor, Algorithms ESA '99, 7th Annual European Symposium, Prague, Czech Republic, July 16-18, 1999, Proceedings, volume 1643 of Lecture Notes in Computer Science, pages 426-437. Springer, 1999. doi: 10.1007/3-540-48481-7\\_37.
- 48 Mingyu Xiao and Hiroshi Nagamochi. Exact algorithms for maximum independent set. *Inf. Comput.*, 255:126–146, 2017. doi:10.1016/j.ic.2017.06.001.
- David Zuckerman. Linear degree extractors and the inapproximability of max clique and chromatic number. *Theory of Computing*, 3(1):103–128, 2007. doi:10.4086/toc.2007.v003a006.