Solving constrained Procrustes problems: a conic optimization approach

Terézia Fulová^{1,*}, Mária Trnovská¹

^aFaculty of Mathematics, Physics and Informatics, Comenius University in Bratislava, Mlynska dolina F1, Bratislava, 84248, Slovakia

Abstract

Procrustes problems are matrix approximation problems searching for a transformation of the given dataset to fit another dataset. They find applications in numerous areas, such as factor and multivariate analysis, computer vision, multidimensional scaling or finance. The known methods for solving Procrustes problems have been designed to handle specific subclasses, where the set of feasible solutions has a special structure (e.g. a Stiefel manifold), and the objective function is defined using a specific matrix norm (typically the Frobenius norm). We show that a wide class of Procrustes problems can be formulated and solved as a (rank-constrained) semi-definite program. This includes balanced and unbalanced (weighted) Procrustes problems, possibly to a partially specified target, but also oblique, projection or two-sided Procrustes problems. The proposed approach can handle additional linear, quadratic, or semi-definite constraints and the objective function defined using the Frobenius norm but also standard operator norms. The results are demonstrated on a set of numerical experiments and also on real applications.

Keywords: Weighted Procrustes problems, Orthogonal Procrustes problems, Oblique Procrustes problems, Two-sided Procrustes problems, Rank-constrained semi-definite programming

^{*}Corresponding author

Email addresses: terezia.fulova@fmph.uniba.sk (Terézia Fulová), trnovska@fmph.uniba.sk (Mária Trnovská)

1. Introduction

In this paper, we study a general class of Procrustes problems (PPs), formulated as

$$\min_{\substack{\text{s.t.} X \in \mathcal{P},}} f(X) := \|\mathcal{L}(X)\|$$
(1)

where $X \in \mathbb{R}^{m \times n}$ is the matrix variable and $\mathcal{L} : \mathbb{R}^{m \times n} \to \mathbb{R}^{p \times q}$ is a linear map.

In the objective, we will consider several types of norms, listed in the table below:

matrix norm for $Y \in \mathbb{R}^{m \times n}$	notation	definition
l_1 norm	$ Y _1$	$\max_{1 \le j \le n} \sum_{i=1}^{m} Y_{ij} $
$l_{\infty} { m norm}$	$ Y _{\infty}$	$\max_{1 \le i \le m} \sum_{j=1}^n Y_{ij} $
l_2 (spectral norm)	$ Y _{2}$	maximal singular value of Y
Frobenius norm	$ Y _F$	$\sqrt{tr(YY^T)}$

Table 1: Matrix norms considered in the Procrustes problems (1).

The most commonly used matrix norm in the objective of the Procrustes problem (1) is the Frobenius norm (least-squares matrix approximation), see [1, 2, 3, 4, 5, 6, 7, 8]. A few authors (see [9, 10]), consider the l_1 norm, which is a robust alternative to the least squares. However, our conic approach also covers the l_{∞} norm and the spectral norm l_2 .

The classification of PPs as *balanced* or *unbalanced* can be made based on the dimension of the matrix variable, as described in [3, 2, 11]. Balanced Procrustes problems refer to cases where the matrix variable is squared (i.e. m = n), while unbalanced Procrustes problems refer to cases where $m \neq n$, indicating that the matrix variable is rectangular.

Another classification of PPs can be made regarding the choice of the linear map \mathcal{L} . Suppose $A \in \mathbb{R}^{p \times m}$, $B \in \mathbb{R}^{n \times q}$, $C \in \mathbb{R}^{p \times q}$ are the given data.¹ Then, we distinguish standard PPs [12, 3] if $\mathcal{L}(X) = C - AX$, weighted PPs [13, 14] if $\mathcal{L}(X) = C - AXB$, two-sided PPs [15] if $\mathcal{L}(X) = CX - XA$ and PPs to a partially specified target [9] if $\mathcal{L}(X) = W \circ (C - AXB)$ where \circ

¹Note that the general formulation (1) also covers the standard matrix approximation problems, that is, problems with $\mathcal{L}(X) = C - X$.

denotes the Hadamard (element-wise) product and the matrix W specifies the target, i.e.

$$W_{ij} = \begin{cases} 1 & \text{if } C_{ij} \text{ is specified,} \\ 0 & \text{otherwise.} \end{cases}$$

The feasible set \mathcal{P} in (1) is typically a matrix manifold, which encompasses cases such as orthogonal Procrustes problems (OPPs) [4, 6, 7, 1] and oblique Procrustes problems (ObPPs) [10, 16]. Some authors also consider other types of feasible sets, such as the set of positive semi-definite matrices [17]. In our paper, we allow for linear, semi-definite, quadratic, and rank constraints to define the feasibility set \mathcal{P} . This means that we cover all the aforementioned classes as well as other challenging cases that are difficult to handle using standard approaches. These may include orthogonal or oblique Procrustes problems with additional linear constraints or Procrustes problems minimizing other than the Frobenius norm of $\mathcal{L}(X)$.

The most well-known subclass of PPs is the class of orthogonal Procrustes problems (OPPs), where the matrix variable is assumed to be orthogonal or at least having orthogonal columns (rows). Since OPPs search for an orthogonal matrix, which maps the given set of data closest to another set of data (with respect to a given norm), they find application in numerous areas such as rigid body dynamics [18, 19], psychometrics [1, 5], multidimensional scaling [20], or global positioning system [21]. Furthermore, standard unbalanced OPPs map high-dimensional data (with dimension m) into a space with a lower dimension $n \ll m$. This applies, for example, in orthogonal least squares regression (OLSR), which may be used for feature extraction [2, 22].

In the standard balanced orthogonal Procrustes problem, the goal is to find an orthogonal matrix X that minimizes $||C - AX||_F$. In [1] it was shown that this problem has a closed-form solution, which can be obtained using the singular value decomposition. Later publications have focused on accelerating the computational time – e.g. in [3], a method based on eigenvalue decomposition is proposed.

Unlike balanced OPPs, there is no closed-form solution to the other classes of Procrustes problems known and therefore an algorithmic approach is required. Several algorithms have been proposed to solve unbalanced OPPs, using the structure of the Stiefel manifold. The approach introduced in [7] uses relaxation-based iterations. This method involves relaxing the orthogonality constraint on X and solving a sequence of relaxed sub-problems iteratively, until convergence is achieved. Another approach (see [5, 6]) uses the Newton-type method to update X in each iteration until a local optimum is reached. In addition to iterative methods, necessary and sufficient conditions for local optimality in unbalanced OPPs have been derived in [23]. These conditions provide insights into the properties of optimal solutions and can be used to guide the development of optimization algorithms for solving unbalanced OPPs.

The special case of unbalanced OPPs with n = 1 is known as the trustregion subproblem of the trust-region method in optimization [24]. This knowledge was used in [11] to design the successive projection method, where all but one column of X are fixed, and a trust-region sub-problem is solved in each iteration. In [22], an iterative algorithm based on the use of SVDs was introduced to solve the orthogonal least squares regression (OLSR), which has proven to be efficient in practice. More recently, in 2020, an eigenvalue-based approach was introduced in [2] that outperforms the successive projection method from [11]. Specifically, the authors proposed an iterative algorithm based on the self-consistent-field (SCF) iteration, which is an efficient method for solving eigenvector-dependent nonlinear eigenvalue problems.

Several algorithms have been developed for the class of weighted OPPs (where $B \neq I_n$) associated with the Frobenius norm. They are based on the extension of standard unconstrained optimization algorithms to the case of Stiefel manifolds [5, 25, 26]. In addition, in [13] and [14], an approach based on solving differential equations has been introduced for weighted OPPs. This approach has also been extended to solve weighted OPPs with the l_1 norm in the objective in [9]. Efforts to solve OPPs using a conic optimization approach have been made in the past. In [4], a relaxation-based approach was introduced, but it only addresses balanced OPPs with possible data uncertainties. Another SDP approach was used to solve unbalanced PPs in [8], where vectorization was used to obtain a semi-definite relaxation of the unbalanced OPP.

Procrustes problems defined over a feasible set given by quadratic constraints of the form $diag(X^TX) = \mathbf{1}_n$ are referred to as *oblique Procrustes* problems (ObPP). It is common to consider the Frobenius norm (see [27, 28]) or the l_1 norm in the objective (see [10, 29]). ObPPs arise in various applications, such as factor analysis [30] and shape analysis [31]. The ObPPs to a partially specified target and weighted ObPPs are also discussed in [10].

On the other hand, our conic approach also covers weighted OPPs and can handle additional linear and semi-definite constraints in the problem formulation. Moreover, unlike existing approaches, our approach handles various matrix norms in the objective, such as the l_1 norm and the spectral norm, which are robust with respect to outliers. This makes our approach more appropriate for problems such as the orthogonal least squares regression, as stated in [22].

A subclass of Procrustes problems defined over the cone of symmetric positive semi-definite matrices is known as *semi-definite Procrustes problems* (SDPP). This problem has been studied in several works (see [17, 32, 33], where the authors have formulated the necessary and sufficient conditions for the optimum and compared the performance of several numerical algorithms. An algorithm for solving the SDPP, based on computing the optimality conditions using specific singular value decompositions, has been designed in [33]. The SDPP's are recognized in numerous applications such as structural analysis [34], signal processing [35], and finance [36]. The problem of finding the nearest covariance matrix can be formulated as a special case of SDPP with m = n = p and $A = I_m$. This problem is commonly encountered when the initial estimate of a covariance matrix is non-positive semi-definite, which is common e.g. in foreign exchange markets [37, 36].

In this paper, we propose an approach based on reformulating the general Procrustes problem (1), where the set \mathcal{P} is assumed to be defined only using linear, semi-definite or (general) quadratic constraints, as a (rankconstrained) semi-definite program. We show that this approach covers a very wide class of Procrustes problems, including balanced and unbalanced weighted Procrustes problems to a partially specified target, with linear quadratic or semi-definite constraints. This includes orthogonal, oblique, semidefinite, or projection Procrustes problems. Also, our approach is not limited to a specific choice of matrix norm – it is suitable for the Frobenius norm as well as for the l_1, l_2 , or l_{∞} operator norms. To handle the reformulated problems, we combine a bisection method with known techniques, such as the log-det heuristic [37] and the so-called convex iteration algorithm [38], designed primarily for rank-constrained feasibility problems. The computational efficiency is demonstrated in several numerical examples, including a practical application. Furthermore, we compare the performance of the proposed bisection method to the existing methods for solving standard unbalanced OPPs and demonstrate its applicability to solve weighted OPPs and weighted ObPPs considering different matrix norms in the objective and also OPPs with additional linear constraints.

This paper is organized as follows: In Section 2, we introduce a reformulation of PPs of the form (1) into a (rank-constrained) semi-definite pro-

class	\mathbf{type}	norm	solution method	source	conic	
		Fro	SVD	[1]		
	balanced		eigenvalue dec.	[3]	\checkmark	
			semi-definite relaxation	[4]		
		Fro	relaxation-based	[7]		
			Newton-type	[5, 6]		
	unbalanced		successive proj.	[11]		
OPP	unbalanceu		OLSR	[22]	v	
			eigenvalue-based	[2]	1	
			SDP relaxation	[8]		
			Stiefel man.	[5, 25],		
	weighted	Fro	gradient	[26]	\checkmark	
			differential appr.	[13, 14]		
		l_1	differential appr.	[9]	\checkmark	
	l_2, l_∞				\checkmark	
		Fro	projection	[15]	.(
	standard	110	differential appr.	[28]	v	
ObPP		l_1	separation	[10, 29]	\checkmark	
	woighted	l_1	differential appr.	[10]	\checkmark	
	weighted	Fro, l_2 , l_{∞}			\checkmark	
		Fra	optimality	[17, 32],	(
SDPP		FIO	conditions	[33]	V	
		l_1, l_∞, l_2			\checkmark	
PPP					\checkmark	
add.c.					\checkmark	

Table 2: Solution methods for different classes of Procrustes problems. The last column indicates classes covered by the conic programming approach. The shortcut "add.c." stands for additional (linear, or semi-definite) constraints.

gram. Unlike other approaches, it applies to all subclasses of PPs that fit our program scheme. In the third section we describe several methods for solving rank-constrained feasibility problems and drawing from the existence of methods able to find solutions proving a lower and upper bound on the optimal value, we propose a bisection method for solving rank-constrained problems. The fourth section contains numerical results and the last section concludes.

2. Semi-definite programming approach to Procrustes problems

In this section, we show that the general Procrustes problem (1) can be represented as a rank-constrained semi-definite program, provided the feasible set \mathcal{P} is defined only using linear, semi-definite or (general) quadratic constraints. In the objective, we consider four different matrix norms: the l_1 norm, the l_2 norm, the l_{∞} norm and the Frobenius norm.

First, we reformulate problems of the form (1) into equivalent problems with a linear objective. The corresponding optimization problems are stated in the following theorem. The proof relies on auxiliary lemmas presented in Appendix A.

Proposition 1. The problem (1) can be equivalently reformulated as follows.

a) For the objective $\|\mathcal{L}(X)\|_F$, the problem (1) is equivalent to

$$\begin{array}{ll} \min & tr(Z) \\ s.t. & X \in \mathcal{P} \\ & \begin{pmatrix} I_q & \mathcal{L}(X)^T \\ \mathcal{L}(X) & Z \end{pmatrix} \succeq 0. \end{array}$$

$$(2)$$

b) For the objective $\|\mathcal{L}(X)\|_1$, the problem (1) is equivalent to

$$\begin{array}{ll} \min & t \\ s.t. & X \in \mathcal{P}, \\ & -S &\leq \mathcal{L}(X) &\leq S, \\ & S^T \mathbf{1}_p &\leq t \mathbf{1}_q. \end{array}$$

$$(3)$$

c) For the objective $\|\mathcal{L}(X)\|_{\infty}$, the problem (1) is equivalent to

$$\begin{array}{ll} \min & t \\ s.t. & X \in \mathcal{P}, \\ & -S & \leq \mathcal{L}(X) & \leq S, \\ & S\mathbf{1}_q & \leq t\mathbf{1}_p. \end{array}$$

$$(4)$$

d) For the objective $\|\mathcal{L}(X)\|_2$, the problem (1) is equivalent to

$$\begin{array}{l} \min \quad s \\ s.t. \quad X \in \mathcal{P} \\ \begin{pmatrix} sI_p & \mathcal{L}(X) \\ \mathcal{L}(X)^T & sI_q \end{pmatrix} \succeq 0. \end{array}$$

$$(5)$$

Proof. We will show that the optimal solution of the problem (1) (where the objective uses a specific matrix norm) defines an optimal solution of the corresponding problem (2)-(5), and vice versa.

a) Let \hat{X} be optimal for (1) with the Frobenius norm in the objective, and we define $\hat{Z} := \mathcal{L}(\hat{X})\mathcal{L}(\hat{X})^T$. Then, from Lemma 1 a) it follows that (\hat{X}, \hat{Z}) is feasible for (2). Clearly $tr(\hat{Z}) = f(\hat{X})$. Reversely, if (X^*, Z^*) is optimal for (2), then X^* is feasible for (1) with the Frobenius norm in the objective. From Lemma 1 a) and Lemma 3 we have that

$$tr(Z^*) \ge tr\left(\mathcal{L}(X^*)\mathcal{L}(X^*)^T\right) =$$
$$\|\mathcal{L}(X^*)\|_F = f(X^*).$$

Therefore

 $f(X^*) \le tr(Z^*) \le tr(\hat{Z}) = f(\hat{X}) \le f(X^*),$

where the second inequality follows from the optimality of Z^* for (2) and the last inequality follows from optimality of (\hat{X}, \hat{Z}) for (1). Therefore, $f(X^*) = f(\hat{X}) = \hat{t} = t^*$.

b) Let \hat{X} be optimal for (1) with the l_1 norm in the objective. Define $\hat{S} \in \mathbb{R}^{p \times q}$ such that $\hat{S}_{ij} := |\mathcal{L}(\hat{X})_{ij}|$ and $\hat{t} = \max_{j} \sum_{i=1}^{p} \hat{S}_{ij}$. Then $(\hat{X}, \hat{S}, \hat{t})$ is feasible for (3) and

$$f(\hat{X}) = \|\mathcal{L}(\hat{X})\|_1 = \max_j \sum_{i=1}^p |\mathcal{L}(\hat{X})_{ij}| = \hat{t}.$$

Reversely, if (X^*, S^*, t^*) is optimal for (3), then X^* is clearly feasible for (1). From the last constraint in (3) we have $t^* \geq \max_j \sum_{i=1}^p S^*_{ij}$, and from the second constraint we have $S^*_{ij} \geq |\mathcal{L}(X^*)_{ij}|$. Therefore

$$t^* \ge \max_j \sum_{i=1}^p S_{ij}^* = \|S^*\|_1 \ge \|\mathcal{L}(X^*)\|_1 = f(X^*).$$

To sum up, we have

$$f(X^*) \le t^* \le \hat{t} = f(\hat{X}) \le f(X^*)$$

where the second inequality follows from the optimality of t^* for (3), and the last inequality follows from the optimality of \hat{X} for (1). Therefore, $f(X^*) = f(\hat{X}) = \hat{t} = t^*$.

c) The proof is analogous to the proof of part b).

d) Let \hat{X} be optimal for (1) with the l_2 norm in the objective. Define $\hat{s} := \|\mathcal{L}(\hat{X})\|_2$. Then, if we denote $\lambda_{max}(.)$ the maximal eigenvalue of a given matrix, we get

$$(\hat{s})^2 = \lambda_{max} \big(\mathcal{L}(\hat{X}) \mathcal{L}(\hat{X})^T \big),$$

which is equivalent to

$$(\hat{s})^2 I_q - \mathcal{L}(\hat{X})^T \mathcal{L}(\hat{X}) \succeq 0.$$

The Schur complement property from Corollary 1 a) gives

$$\begin{pmatrix} \hat{s}I_p & \mathcal{L}(\hat{X}) \\ \mathcal{L}(\hat{X})^T & \hat{s}I_q \end{pmatrix} \succeq 0,$$

and hence (\hat{X}, \hat{s}) is feasible for (5) such that $\hat{s} = f(\hat{X})$. Reversely, if (X^*, s^*) is optimal for (5), then X^* is clearly feasible for (1). By applying Corollary 1 to the last constraint of (5) we obtain

$$(s^*)^2 I_q - \mathcal{L}(X^*)^T \mathcal{L}(X^*) \succeq 0$$

which is equivalent to

$$(s^*)^2 \ge \lambda_{max} \left(\mathcal{L}(X^*)^T \mathcal{L}(X^*) \right) = \| \mathcal{L}(X^*) \|_2^2.$$

Therefore

$$f(X^*) \le s^* \le \hat{s} = f(\hat{X}) \le f(X^*),$$

where the second inequality follows from the optimality of s^* for (5), and the last inequality follows from the optimality of \hat{X} for (1). Therefore, $f(X^*) = f(\hat{X}) = \hat{s} = s^*$.

Note that if \mathcal{P} is defined by linear, convex quadratic, or semi-definite constraints, the problems (2)-(5) are semi-definite programming problems. Furthermore, if \mathcal{P} is defined by non-convex quadratic constraints, the problems (2)-(5) are equivalent to rank-constrained SDP problems. The representations of quadratic constraints are listed in Table 3 and can be easily derived using the standard matrix analysis results; see Lemma 1, Lemma 2 in Appendix A.

Using the rank-constrained SDP representations in Table 3, we can also easily express quadratic constraints of the type

$$tr(X^TX) \le g, \ tr(X^TX) = g, \ tr(X^TX) \ge g$$

Constraint Representation

$$\begin{split} X^T X &\preceq G \quad V = \begin{pmatrix} I & X \\ X^T & G \end{pmatrix} \succeq 0 \\ X^T X &= G \quad V = \begin{pmatrix} I & X \\ X^T & G \end{pmatrix} \succeq 0 \quad rank(V) = m \\ X^T X \succeq G \quad V = \begin{pmatrix} I & X \\ X^T & Y \end{pmatrix} \succeq 0 \quad rank(V) = m, \, Y - G \succeq 0 \end{split}$$

Table 3: Quadratic constraints representation via semi-definite, linear, and rank constraints. Quadratic constraints are given by $m \times n$ matrix X and $n \times n$ matrix G.

or

$$diag(X^T X) \leq h, \ diag(X^T X) = h, \ tr(X^T X) \geq h$$

for scalar g and vector $h \in \mathbb{R}^n$. Note that non-convex matrix constraints include, e.g., orthogonal constraints $X^T X = I$ and projection constraints $X^T X = X$.

The advantage of the SDP reformulation of the Procrustes problems is that we can handle a wide class of problems with algorithms for rankconstrained SDP problems. We recall that the rank is a quasiconcave function on the cone of the positive semi-definite symmetric matrices, and its convex envelope is the trace function (see Theorem 1 in [39]). Using these results, several rank minimization heuristics and rank reduction algorithms were designed for solving rank-constrained SDP problems using only conic optimization tools. The most well-known heuristic is the so-called trace heuristic [37], which was upgraded into the so-called log-det heuristic in [40]. Besides these rank minimization heuristics, there are also several rank reduction algorithms that search for a lower-rank solution among feasible solutions starting from an initial solution of a higher rank. Such an algorithm is, e.g., the so-called convex iteration introduced in [38]. Moreover, thanks to the recent publication [41], even exact algorithms can be applied to solve rankconstrained problems.

3. Solving rank-constrained problems

In the previous section, we have shown that the Procrustes problem of the form (1) can be formulated as a rank-constrained semi-definite programming problem. The most common approach is to approximate the problem with its convex relaxation, which is obtained by omitting the rank constraint. This way we obtain a lower bound on the optimal value of (1). However, the contemporary interior-point methods for solving convex problems converge to the solution of the highest feasible rank (see [42]). Therefore, even if an optimal solution of the required rank exists in the set of optimal solutions of the convex relaxation, it is not guaranteed to be found. In this case, the rank reduction algorithm for solving the rank-constrained semi-definite problem can be applied, as proposed in [43].

However, typically an optimal solution of the required rank does not exist in the set of optimal solutions of the convex relaxation of (1). Since in this case the problem is way more complex, several heuristics have been designed to solve rank-constrained semi-definite *feasibility problems* of the form

find
$$X: X \in \mathcal{C}, rank(X) \le k,$$
 (6)

or rank minimization problems

$$\min_{X \in \mathcal{C}} rank(X) \tag{7}$$

where $\mathcal{C} \subseteq \mathcal{S}^n_+$ is a convex set. The well-known trace heuristic (see [37]) is based on the fact that trace(X) is the convex envelope of rank(X) on the set $\{X \in \mathcal{S}^n_+ \mid 0 \leq X \leq I\}$. To enhance the performance of the trace heuristic, the so-called log-det heuristic was proposed in [40]. The idea is to approximate the rank minimization problem (7) with

$$\min_{X \in \mathcal{C}} \log \det(X + \delta I), \tag{8}$$

which is then solved by an iterative method based on the first-order linear approximation.

An approach for solving the feasibility problem (6) was proposed in [38, §4.4.2]. It is based on iteratively solving two convex problems until convergence. In particular, in the *t*-th iteration we find

$$X_t := \underset{X \in \mathcal{C}}{\operatorname{argmin}} \ trace(U_{t-1}X) \tag{9}$$

and

$$U_t := \operatorname*{argmin}_{U:0 \preceq U \preceq I, trace(U) = n-k} trace(UX_t)$$
(10)

The algorithm starts with $U_0 = I$ and generates a sequence $\{U_t\}_{t=0}^T$ of the so-called direction matrices along with the sequence $\{X_t\}_{t=0}^T$ of approximate solutions of (6) such that the sequence $\{trace(X_tU_t)\}_{t=0}^T$ is non-increasing. It can be seen (e.g. in [44, §4.1]) that the objective of (10) represents the sum of n-k smallest eigenvalues of X_t . However, the sum of n-k smallest eigenvalues of X_t . However, the sum of n-k smallest eigenvalues of X_t equal to zero is equivalent to $rank(X_t) \leq k$, see Lemma 4 in Appendix A. Therefore, if the sequence $\{trace(X_tU_t)\}_{t=0}^\infty$ converges to zero, the algorithm converges to a rank-k-solution (with some accuracy).

In the more complex case, our aim is to solve a problem

$$\min \begin{array}{l} f(X) \\ X \in \mathcal{C}, \\ rank(X) \le k, \end{array}$$

$$(11)$$

where $\mathcal{C} \subseteq \mathcal{S}^n_+$ is a convex set. Assume that the problem (11) reaches an optimal solution X^* and that the optimal value is $f^* := f(X^*)$.

In the following, we propose a *bisection method* based on solving rankconstrained feasibility problems of the form

find
$$X$$

 $X \in \mathcal{C},$
 $rank(X) \le k,$
 $f(X) \le \gamma.$
(12)

Let $l, u \in \mathbb{R}$ be such that $f^* \in [l, u]$. The value l, satisfying $l \leq f^*$ can be found by solving the convex relaxation problem

$$\begin{array}{ll} \min & f(X) \\ & X \in \mathcal{C}, \end{array}$$
 (13)

and the value u, satisfying $f^* \leq u$ can be set as $u = f(\bar{X})$, where \bar{X} is a feasible solution of (11). Such a solution can be found e.g. by applying any heuristics mentioned above.²

We design a bisection algorithm to solve the rank-constrained problem (11) as follows:

²Note that if $\gamma = f(X_0)$ where X_0 is a solution of the convex relaxation (13), solving (12) finds an optimal solution of (13) if such a solution exists.

Algorithm 1: Bisection algorithm for solving rank-constrained problems

Input: Interval [l, u] containing f^* ; Accuracy constant $\delta > 0$; Initialize: $X_{\delta} \leftarrow \bar{X}$; while $|u - l| \ge \delta$ do $\begin{array}{c|c} \gamma \leftarrow \frac{l+u}{2};\\ \text{Solve (12) (or declare infeasibility);}\\ \text{if there exists a solution } X_{\gamma} \text{ of (12) then}\\ & X_{\delta} \leftarrow X_{\gamma};\\ & u \leftarrow \gamma;\\ & \text{else}\\ & \mid l \leftarrow \gamma;\\ & \text{end}\\ \end{array}$ end Output: $X^*_{\delta} := X^{\delta}$ (δ -optimal solution of (11) satisfying $rank(X^*_{\delta}) \le k$)

Proposition 2. The solution X^*_{δ} provided by Algorithm 1 satisfies

 $|f(X^*_{\delta}) - f^*| \le \delta.$

and it is obtained after $N = \left\lceil \log_2 \frac{|u-l|}{\rho} \right\rceil$. iterations.

Proof. Denote

$$\mathcal{P} = \{ X \in \mathcal{C} \mid rank(X) \le k \}$$

the set of feasible solutions of (11) and denote $X^i := X_{\delta}$ and $[l_i, u_i]$ the feasible solution and the corresponding interval in the *i*-th iteration, respectively. In each iteration of Algorithm 1 we either find $X_{\gamma} \in \mathcal{P}$ satisfying $f^* \leq f(X_{\gamma}) \leq \gamma$ or we find out that no such solution exists. In the latter case, we have that for all $X \in \mathcal{P}$ it holds $f(X) > \gamma$ and therefore $f^* = \inf_{X \in \mathcal{P}} \geq \gamma$. Therefore, in each iteration, the property $f^* \in [l_i, u_i]$ is satisfied. Our aim now is to show that in each iteration it holds $f(X^i) \in [l_i, u_i]$. Since at initialization, $X_{\delta} = \bar{X}$ is chosen so that $f(\bar{X}) = u$, the property is satisfied in the first iteration. Next, we show that if $f(X^i) \in [l_i, u_i]$, then $f(X^{i+1}) \in [l_{i+1}, u_{i+1}]$. If (12) is feasible, then X^{i+1} is updated to X_{γ} and hence $f(X^{i+1}) \leq \gamma = u_{i+1}$. Also, in this case $l_{i+1} = l_i \leq f^* \leq f(X^{i+1})$. On the other hand, if (12) is infeasible, we have that $X^{i+1} = X^i$ and $l_{i+1} = \gamma \leq f^* \leq f(X^i) \leq u_i$. Let $[l_N, u_N]$ be the final interval satisfying $u_N - l_N < \delta$. We have that the both values f^* and $f(X^*_{\delta})$ belong to $[l_N, u_N]$ and therefore

$$|f(X^*_{\delta}) - f^*| \le u_N - l_N < \delta.$$

During practical implementation, we search for an optimal solution with the required rank with respect to some accuracy. Since the variables are positive semi-definite, we define the ε -rank of a matrix $X \in S^n_+$ as follows:

$$\varepsilon$$
-rank $(X) = k \iff \lambda_1 > \varepsilon, ..., \lambda_k > \varepsilon, \lambda_{k+1} \le \varepsilon, ..., \lambda_n \le \varepsilon.$ (14)

Algorithm 1 is based on a method for the rank-constrained feasibility problem (12). For this purpose, any of the heuristics or methods mentioned at the beginning of this section can be applied.

4. Numerical results

The computations have been executed in MATLAB R2019a [45] on a laptop with the 11th Gen Intel(R) Core(TM) i7-1165G7 processor running at 2.80GHz. To solve SDP programs, we used CVX: a package for specifying and solving convex programs [46, 47]. The rank of a matrix was determined as the ε -rank according to (14) for $\varepsilon = 10^{-6}$. For some experiments, we determine an "empirical" ε as the (k + 1)-th eigenvalue of a solution. This value can be useful to analyze the performance of a particular algorithm or the quality of a found solution. The orthogonal matrix was generated using the build-in function RandOrth() from Matlab libraries and the oblique matrix was generated as $X_{oblique} = XDiag(diag(X^TX)^{\frac{1}{2}})$ where X is a randomly generated $m \times n$ matrix, as proposed in [28]. Then we generate problems with a zero optimal value by generating data matrices for the problem (1) and setting C = AXB, or with a non-zero optimal value, where we generated also a matrix $\Delta \in \mathbb{R}^{p \times q}$ from N(0, 1) to define $C = AXB + 0.5\Delta$, as suggested in [13].

In the following sets of experiments, we first demonstrate the applicability of the proposed conic approach to find a solution of an application of feature extraction with the Frobenius norm and even with the l_1 norm in the objective. Second, we demonstrate the versatility of the proposed conic approach in solving weighted OPPs and weighted ObPPs defined in terms of different norms. Then, we apply the proposed conic approach to find a permutation matrix that solves a system of linear equations, since such a problem can be formulated as a standard balanced OPP with additional linear constraints. Finally, we solve a graph isomorphism problem which can be formulated as a two-sided OPP. To assess the feasibility of the resulting solution X, we use the criteria $||X^TX - I_n||_F$ to verify orthogonality and $||diag(X^TX) - \mathbf{1}||_1$ to verify obliqueness.

4.1. Weighted OPPs and weighted ObPPs with the Frobenius, l_1 , l_2 and l_{∞} norm in the objective

This subsection focuses on weighted PPs of the form

$$\min_{\text{s.t.}} \|C - AXB\|$$

$$\text{s.t.} \quad X \in \mathcal{P},$$

$$(15)$$

where $B \neq I_n$ and $\mathcal{P} = \{X \in \mathbb{R}^{m \times n} | X^T X = I_n\}$ in case of OPPs or $\mathcal{P} = \{X \in \mathbb{R}^{m \times n} | diag(X^T X) = \mathbf{1}_n\}$ in case of ObPPs. Our aim is to demonstrate the versatility of the proposed conic approach by applying it to the weighted PPs regarding four matrix norms in the objective of (15), see Table 1.

It is important to note that there are several effective methods for solving weighted OPPs with the Frobenius norm in the objective, including the spectral projected gradient method [28, 25] which finds an optimal orthogonal or oblique solution within a few seconds also for large-size problems, as demonstrated by the experiments. However, unlike the approach in [28, 25] the conic approach can also be applied to weighted PPs with respect to l_1, l_2 or l_{∞} norm, see the results in Table 4 and Table 5.

For the case of l_1 norm in the objective, a differential approach was proposed in [9, 10] to solve weighted OPPs and weighted ObPPs with the l_1 norm in the objective. However, the performance of such an approach was illustrated on small examples, and the authors labeled this approach to be time-consuming since using build-in Matlab functions for ODE calculations. Regarding l_2 norm and l_{∞} norm in the objective of (15), there are no significant results in the literature (compare to Table 2).

The results obtained by the proposed bisection method applied to the rank-constrained SDP reformulation of (15) are summarized in Table 4 for weighted OPPs and Table 5 for weighted ObPPs. We can observe that, in all cases, orthogonal or oblique solutions were found by Algorithm 1 yielding only a slightly higher optimal value than the semi-definite relaxation. The results demonstrate that the conic approach is a possible computational tool for solving this class of OPPs, though being time-consuming. As expected, when solving weighted OPPs with the Frobenius norm in the objective, the conic approach cannot compete with the spectral projected gradient method [28, 25]. However, in the case of l_1 norm, it can be considered a reasonable alternative to the differential approach[9], and, to our best knowledge there are no alternative computational methods in the case of l_2, l_{∞} .

4.2. Standard balanced OPPs with additional linear constraints

In this subsection, we address the problem of finding a permutation matrix that minimizes the objective of the standard balanced OPP (17) with the l_1 norm in the objective. The l_1 norm is preferred since it enables handling matrices with many zero elements more effectively. It is worth noting that a permutation matrix is an orthogonal matrix with unit row and column sums, having elements of 0 or 1. Although the problem of finding such a matrix is an integer programming problem, we use the fact that a permutation matrix can be represented by an orthogonality constraint and constraints on its nonnegative elements. Therefore, the standard balanced OPP with additional linear constraints representing finding a permutation matrix can be formulated as follows

$$\min_{X \in \mathbb{R}^{n \times n}} \|C - AX\|
X^T X = I_n
X_{ij} \ge 0, \quad \forall i, j = 1, ..., n.$$
(16)

In the following set of experiments, we solve the OPP with the additional linear constraints of the form (16) generated for a random permutation matrix of size n. We focus on problems with a zero optimal value, which enables interpreting (16) as the problem of finding a permutation matrix that satisfies a linear system of equations.

Table 6 shows that a rank-*n* solution was obtained by the semi-definite relaxation of the proposed rank-constrained SDP reformulation of (16) in all cases. We test the optimality and feasibility of the found solutions of (16) using several criteria listed in the table. To conclude, the proposed conic approach was successful in finding a ε -optimal solution of all generated OPPs with additional linear constraints.

norm	criterion	SDP	Algorithm 1	Algorithm 1
		relaxation	(log-det)	(cvx.iter.)
	$\ C - AXB\ _F$	2.2556	2.3726	2.3379
Frob.	$\ X^T X - I_n\ _F$	1.0835	1.9947e-06	1.4682e-06
norm	ε -rank(V)	6.28	4	4
	time (s)	0.2642	47.5041	93.9932
	$\% \ (\varepsilon\text{-}rank(V) \le m)$	0	100	100
	$\ C - AXB\ _1$	3.6718	3.8106	3.8339
l_1	$\ X^T X - I_n\ _F$	1.0865	9.2466e-07	9.2440e-07
norm	ε -rank(V)	7.02	4	4
	time (s)	0.2489	40.6750	84.4602
	$\% \ (\varepsilon\text{-}rank(V) \le m)$	0	100	100
	$\ C - AXB\ _2$	1.7020	1.74875	1.7486
l_2	$\ X^T X - I_n\ _F$	1.1803	7.3899e-07	7.8219e-07
norm	ε -rank(V)	7.38	4	4
	time (s)	0.2807	32.6941	76.4913
	$\% (\varepsilon \text{-} rank(V) \le m)$	0	100	100
	$\ C - AXB\ _{\infty}$	1.3773	1.4848	1.4718
l_{∞}	$\ X^T X - I_n\ _F$	1.1161	1.4266e-06	9.8376e-07
norm	ε -rank(V)	6.76	4	4
	time (s)	0.2421	38.0758	72.5916
	$\% \ (\varepsilon\text{-}rank(V) \le m)$	0	100	100

Table 4: Results obtained by the proposed conic approach in solving weighted OPPs with the Frobenius norm, l_1 norm, l_{∞} norm, and spectral norm in the objective. Average values of optimal value, orthogonality criterion, ε -rank of the block matrix V, computation time and percentage of solutions having ε -rank equal to m = 4 obtained by semi-definite relaxation and Algorithm 1 in solving 100 generated problems of size (m, n, p, q) = (10, 4, 4, 3) with optimal value $f^* \neq 0$.

4.3. Application - Feature extraction

Consider the Yale data set³, consisting of 165 gray-scale images of 15 individuals, each having 11 images, representing different facial expressions or configurations. The task is to identify the most important facial features that predict the identity of the individual, such as the positions of certain

³Data sourced from https://www.kaggle.com/datasets/olgabelitskaya/yale-face-database

norm	criterion	SDP relaxation	Algorithm 1 (log-det)	Algorithm 1 (cvx.iter.)
	$\ C - AXB\ _F$	2.1410	2.1533	2.1658
Frob.	$\ diag(X^TX) - 1_n\ _1$	2.6696e-01	3.6455e-06	2.1883e-08
norm	ε -rank(V)	5.45	4	4
	time (s)	0.2624	37.6552	21.5325
	$\% \ (\varepsilon\text{-}rank(V) \le m)$	10	100	100
	$\ C - AXB\ _1$	3.3923	3.4455	3.3415
l_1	$\ diag(X^TX) - 1_n\ _1$	0.5372	5.2837e-07	5.2837e-07
norm	ε -rank(V)	7.40	4	4
	time (s)	0.2318	14.9645	38.4616
	$\% \ (\varepsilon\text{-}rank(V) \le m)$	5	100	100
	$\ C - AXB\ _2$	1.6164	1.6246	1.6215
l_2	$\ diag(X^TX) - 1_n\ _1$	1.1270	1.0289e-05	1.3706e-07
norm	ε -rank(V)	7.7	4	4
	time (s)	0.2337	15.4821	36.2344
	$\% \ (\varepsilon\text{-}rank(V) \le m)$	0	100	100
	$\ C - AXB\ _{\infty}$	0.9296	1.2757	1.3016
l_{∞}	$\ diag(X^TX) - 1_n\ _1$	0.3955	4.1862e-07	4.0443e-07
norm	ε -rank(V)	6.7	4	4
	time (s)	0.2302	16.2491	36.9748
	$\% \ (\varepsilon\text{-}rank(V) \le m)$	15	100	100

Table 5: Results obtained by the proposed conic approach in solving weighted ObPPs with the Frobenius norm, l_1 norm, l_{∞} norm, and spectral norm in the objective. Average values of optimal value, orthogonality criterion, ε -rank of the block matrix V, computation time and percentage of solutions having ε -rank equal to m = 4 obtained by the semi-definite relaxation and Algorithm 1 in solving 20 generated weighted ObPPs with the Frobenius norm, l_1 norm, l_{∞} norm and spectral norm in the objective of size (m, n, p, q) = (10, 4, 4, 3)with optimal value $f^* \neq 0$.

landmarks on the face or the intensities of certain regions. To achieve this, the orthogonal least squares regression can be used to extract the most informative features that are correlated with the identity labels of individuals.

Orthogonal least squares regression (OLSR) is a regression technique that involves finding an orthogonal transformation matrix $X \in \mathbb{R}^{m \times n}$ to project high-dimensional data (with dimension m) into a lower-dimensional space (with dimension $n \ll m$). Hence, it is formulated as a standard unbalanced

criterion $\setminus (m,n,p)$	(3,3,10)	(5,5,20)	(10, 10, 30)
$\ C - AX\ _1$	3.0607e-10	4.4304e-10	5.2880e-10
$\ X^T X - I_n\ _F$	2.3621e-11	2.4869e-11	2.3348e-11
ε -rank(V)	3	5	10
# of ε -rank(V) > m	0	0	0
$\ X1_n-1_n\ _1$	4.1927e-12	7.6000e-13	3.8232e-13
$\ X^T 1_n - 1_n\ _1$	4.1299r-12	8.3364e-13	1.8042e-13
$\ o_{max}-1_n\ _1$	1.4740e-11	2.4448e-11	3.4865e-11
$ z_{min} - 0_{n(n-1)} _1$	1.8849e-11	2.5154e-11	3.4811e-11
time (s)	0.1598	0.2892	0.5307
empirical ε	3.6912e-11	2.2944e-11	1.4833e-11

Table 6: Results obtained by the semi-definite relaxation in solving standard balanced OPPs with the l_1 norm in the objective and additional linear constraints that represent searching for permutation matrices. Average values of the listed criteria counted for 100 generated problems of each size. Here, o_{max} denotes a vector of n largest elements of X and z_{min} denotes a vector of n(n-1) smallest elements of X.

OPP of the form

$$\min_{\text{s.t.}} \|C - AX\|$$

$$\text{s.t.} \quad X^T X = I.$$

$$(17)$$

To perform feature extraction using orthogonal least squares regression, we follow the approach described in [2]. Consider a data set $S = [s_1, ..., s_p] \in \mathbb{R}^{m \times p}$, which contains p samples with m features drawn from n classes. In the Yale data set, we have p = 165 images (samples) with m = 256 features corresponding to n = 15 individuals. The task is to identify the most important facial features that predict the identity of the individual. Let $K = [k_1, ..., k_p] \in \mathbb{R}^{n \times p}$ be the class indicator matrix. This means that if the image s_i belongs to the j-th individual, then $k_i = e_j$, where $e_j \in \mathbb{R}^n$ is the j-th column of the standard basis. The model includes an orthogonal transformation matrix $X \in \mathbb{R}^{m \times n}$ and an associated bias $b \in \mathbb{R}^n$. Both Xand b are determined using the orthogonal least squares regression, which is formulated as an OPP of the form ([2])

min
$$h(X,b) = \|S^T X + \mathbf{1}_p b^T - K^T\|_F$$

 $X^T X = I,$ (18)

where $S \in \mathbb{R}^{m \times p}$ and $K \in \mathbb{R}^{n \times p}$ are the given data described above.

norm	criterion	OLSR method [22]	SDP relevation	log-det
		method [22]	тстахатюн	incuristic
Frobenius	$ C - AX _F$	2.9571	2.9570	2.9570
	$\ X^T X - I_n\ _F$	4.3586e-15	1.0000	1.1725e-08
	ε -rank(V)	256	258	256
	time (s)	0.4719	817.6205	1534.89
<i>l</i> ₁	$\ C - AX\ _1$	Х	7.2534	7.2534
	$\ X^T X - I_n\ _F$	х	0.3560	1.6922e-9
	ε -rank(V)	X	270	256
	time (s)	X	51.58	106.2779

Table 7: Results of the feature extraction application (17) applied to the Yalefaces data set. Comparison of the OLSR algorithm [22] and the proposed conic approach represented by the semi-definite relaxation, and the log-det heuristic (8) applied to find a rank-256 solution among optimal solutions of the semi-definite relaxation.

Since *b* can be expressed as $b = \frac{1}{p}(K\mathbf{1}_p - X^T S\mathbf{1}_p)$, the formulation (18) is simplified to the OPP of the form (17), where $A = (I_p - \frac{1}{p}\mathbf{1}_p\mathbf{1}_p^T)S^T \in \mathbb{R}^{p \times m}$ and $C = (I_p - \frac{1}{p}\mathbf{1}_p\mathbf{1}_p^T)K^T \in \mathbb{R}^{p \times n}$.

To solve the problem (17), we used an OLSR algorithm introduced in $[22]^4$. Our aim is to assess the precision of a solution found by solving the proposed rank-constrained SDP reformulation. In Table 7, our results are compared with those obtained by the OLSR algorithm. It can be seen that the log-det heuristic was able to find an orthogonal solution among the optimal solutions of the semi-definite relaxation. The comparison of computation time matches our expectations since the conic approach is based on solving large SDP problems, while the OLSR iteratively computes singular value decompositions.

As stated in [22], the l_1 norm is more suitable for feature extraction that the Frobenius norm, since it is robust to outliers. Nonetheless, the proposed conic approach, represented by the semi-definite relaxation and the modified log-det heuristic can handle this type of OPP. Moreover, the computational time is in this case much lower, which is related to the lower number of variables, (see Table 7).

⁴Accesible at https://github.com/StevenWangNPU/OLSR_NC2016

4.4. Application - Graph isomorphism problem

This subsection describes the graph isomorphism problem and its formulation in the form of a two-sided OPP. It is known that a graph is a set of vertices connected by edges. For the sake of simplicity, we only consider unweighted undirected graphs. Determining if two graphs are isomorphic or not is important in various fields such as chemistry, computer science, and data mining.

We can label the *n* vertices of a simple graph as 1, 2, ..., n. The graph can then be defined by its adjacency matrix $A \in \mathbb{R}^{n \times n}$, where each element $A_{ij} \in \{0, 1\}$ indicates whether vertices *i* and *j* are adjacent or not. The graphs *G* and \tilde{G} are isomorphic if and only if there is a permutation matrix $P \in \mathbb{R}^{n \times n}$ that satisfies

$$PA = \hat{A}P. \tag{19}$$

The task of the graph isomorphism problem is to determine whether the two given graphs are isomorphic, i.e. to verify whether there exists a permutation matrix satisfying (19).

The authors of [48] proposed a method for the graph isomorphism problem by formulating it as an integer linear program. Its LP relaxation, though being computationally efficient, lead to a solution with real-valued elements and therefore sophisticated rounding algorithms needed to be applied to get a permutation matrix. To avoid problems with rounding, we represent the permutation matrix as an orthogonal matrix having non-negative elements. Consequently, the graph isomorphism problem can be formulated as a socalled two-sided OPP (see [15]) of the following form

$$\min_{P \in \mathbb{R}^{n \times n}} \|PA - \tilde{A}P\|_{1}$$

$$P^{T}P = I,$$

$$P_{ij} \geq 0 \qquad \forall i, j = 1, ..., n.$$
(20)

Similarly to the previous subsection, the graph isomorphism problem (20) is formulated with the l_1 norm in the objective and the constraints of (20) define the set of permutation matrices $P \in \mathbb{R}^{n \times n}$.

After applying Proposition 1 and representation of the orthogonality constraint from Table 3, (20) can be rewritten as the following rank-constrained SDP problem

$$\begin{array}{rcl} \min & t \\ -S \leq PA - \tilde{A}P \leq S, \\ S^{T}\mathbf{1}_{n} \leq t\mathbf{1}_{n}, \\ P_{ij} \geq 0, \quad \forall i, j = 1, ..., n. \end{array}$$

$$V := \begin{bmatrix} I_{n} & P^{T} \\ P & I_{n} \end{bmatrix} \succeq 0, \\ rank(V) = n, \end{array}$$

$$\begin{array}{rcl} & (21) \\ \end{array}$$

where $P \in \mathbb{R}^{n \times n}, t \in \mathbb{R}, S \in \mathbb{R}^{n \times n}, V \in \mathbb{S}^{2n}$ are variables.

Furthermore, we can exploit the fact that two graphs represented by their adjacency matrices A and \tilde{A} are isomorphic if and only if P is a feasible solution of (20) with a zero optimal value. Therefore, in our numerical experiments, we solve the rank-constrained SDP problem (21) using the log-det heuristic (8) and the convex iteration (9),(10) to solve (12) where γ is a small positive constant. We solve (21) for four pairs of adjacency matrices of different sizes $n \in \{4, 6, 16, 25\}$ that represent pairs of isomorphic graphs, and we search for a permutation matrix that satisfies (19). The results summarized in Table 8 confirm that the found solutions are permutation matrices that can be considered ε -optimal solutions of the graph isomorphism problem (20)

5. Conclusion

In this paper we have demonstrated the applicability of the conic programming approach to a wide class of Procrustes problems related to four matrix norms. The proposed approach uses (but is not limited to) known heuristics for rank-constrained feasibility problems, in combination with a suitable bi-section algorithm. It has been shown that, in the case of specific sub-classes (such as orthogonal Procrustes problems with Frobenius norm), this approach cannot compete with algorithms based on singular-value decomposition or extensions of the standard gradient algorithms to the Stiefel manifolds. On the other hand, it can handle other sub-classes, for which no suitable methods are known. The applicability of the proposed approach was verified on two applications – by solving orthogonal least squares regression for feature extraction and the graph isomorphism problem. Furthermore, the techniques can be easily extended to the case of multiple norm criteria, e.g. appearing in the regularized Procrustes problems.

method	criterion	n = 6	n = 16	n = 25
	$\ PA - \tilde{A}P\ _1$	3.5187	6.0017	0.8683
	$ P^T P - I_n _F$	1.5643	2.4076	3.7997
	rank(V)	11	31	49
SDD	$\ P1_{n}-1_{n}\ _{1}$	1.4627e-11	1.1875e-10	2.0370e-12
rolayation	$\ P^T 1_n - 1_n\ _1$	1.4627e-11	1.1875e-10	2.0367e-12
Telaxation	$\ o_{max} - 1_n\ _1$	2.6188	5.8639	13.8898
	$ z_{min} - 0_{n(n-1)} _1$	2.6188	5.8639	13.8898
	time (s)	0.2827	0.8153	1.9829
	empirical ε	0.5958	1.6218	3.9723
	$\ PA - \tilde{A}P\ _1$	4.2576e-13	1.2472e-13	7.6111e-7
	$ P^T P - I_n _F$	1.4113e-7	1.2744e-7	6.1667e-6
	rank(V)	6	16	25
log-det	$\ P1_{n}-1_{n}\ _{1}$	1.3124e-12	5.5889e-10	9.6092e-9
heuristic	$\ P^T 1_n - 1_n\ _1$	1.3096e-12	5.5889e-10	9.6092e-9
(γ)	$\ o_{max}-1_n\ _1$	7.0571e-8	2.072e-7	1.4358e-5
	$\ z_{min} - 0_{n(n-1)}\ _1$	7.0570e-8	2.0716e-7	1.4348e-5
	time (s)	0.5958	1.6218	3.9723
	empirical ε	7.0564e-8	2.5362e-8	9.3755e-7
	$\ PA - \tilde{A}P\ _1$	2.4484e-14	7.1029e-15	4.6085e-15
	$ P^T P - I_n _F$	1.2035e-9	4.4462e-9	4.4291e-8
	rank(V)	6	16	25
$\begin{array}{c} \text{convex} \\ \text{iteration} \\ (\gamma) \end{array}$	$\ P1_n-1_n\ _1$	1.7152e-12	1.9921e-10	1.0913e-9
	$ P^T 1_n - 1_n _1$	1.7154e-12	1.9921e-10	1.0913e-9
	$\ o_{max}-1_n\ _1$	1.0254e-9	8.5908e-9	1.0850e-7
	$\ z_{min} - 0_{n(n-1)}\ _1$	1.0247e-9	8.3916e-9	1.0744e-7
	time (s)	0.7158	3.4213	11.5446
	empirical ε	$4.\overline{3997e}-10$	$7.\overline{2807e-10}$	4.5204e-9

Table 8: Results obtained by the semi-definite relaxation, the log-det heuristic and the convex iteration for $\gamma = 10^{-6}$ in solving two-sided OPPs of the form (21) representing the graph isomorphism problem (20).

A cknowledgement

The research was supported by the APVV-20-0311 project of the Slovak Research and Development Agency.

Appendix A. Auxilliary results

Lemma 1. Let

$$M = \begin{pmatrix} A & B \\ B^T & C \end{pmatrix}$$

be a symmetric matrix with square blocks A and C.

a) [49, §4.2.1], [50] If A is positive definite, then M is positive (semi)definite if and only if the Schur complement $C - B^T A^{-1}B$ is positive (semi)definite.

If C is positive definite, then M is positive (semi)definite if and only if the Schur complement $A - BC^{-1}B^{T}$ is positive (semi)definite.

b) [38, \S A.4], [50] If A is invertible, then

$$rankM = rank(A) + rank(C - B^{T}A^{-1}B).$$
(A.1)

Corollary 1. For $B \in \mathbb{R}^{p \times q}$ the following are equivalent:

a)
$$\begin{pmatrix} sI_p & B \\ B^T & s_qI \end{pmatrix} \succeq 0 \quad b)s^2I_p - BB^T \succeq 0, \quad c)s^2I_q - B^TB \succeq 0.$$

Proof. Note that if s = 0 in a), b) or c), then B = 0. In this case, the equivalence is trivial. The case s > 0 follows from Lemma 1 a).

Lemma 2. [51] Let $G \in \mathbb{S}^n$ and $X \in \mathbb{R}^{m \times n}$. Then

$$G = X^T X \quad \Leftrightarrow \quad \left[G \succeq X^T X \land rank \left(\left[\begin{array}{cc} I_m & X \\ X^T & G \end{array} \right] \right) = m \right].$$
(A.2)

Since the trace of a positive semi-definite matrix is non-negative, the following property holds:

Lemma 3. Let $X, Y \in S^n$. If $X \succeq Y$, then $tr(X) \ge tr(Y)$.

Lemma 4. [38, 50] Let $X \in \mathbb{S}^n_+$, $k \leq n$ and $\lambda_1 \geq \lambda_2 \geq ... \geq \lambda_n \geq 0$ be eigenvalues of X. Then it holds

$$\lambda_1 + \lambda_2 + \dots + \lambda_k = tr(X) \iff rank(X) \le k, \tag{A.3}$$

and

$$\lambda_{k+1} + \lambda_{k+2} + \dots + \lambda_n = 0 \quad \Leftrightarrow \quad rank(X) \le k. \tag{A.4}$$

References

- P. H. Schönemann, A generalized solution of the orthogonal procrustes problem, Psychometrika 31 (1) (1966) 1–10.
- [2] L.-H. Zhang, W. H. Yang, C. Shen, J. Ying, An eigenvalue-based method for the unbalanced procrustes problem, SIAM Journal on Matrix Analysis and Applications 41 (3) (2020) 957–983.
- [3] Y. Qiu, A. Wang, Solving balanced procrustes problem with some constraints by eigenvalue decomposition, Journal of computational and applied mathematics 233 (11) (2010) 2916–2924.
- [4] S. Ahmed, I. M. Jaimoukha, A relaxation-based approach for the orthogonal procrustes problem with data uncertainties, in: Proceedings of 2012 UKACC International Conference on Control, 2012, pp. 906–911. doi:10.1109/CONTROL.2012.6334752.
- [5] T. Viklands, P.-Å. Wedin, Algorithms for linear least squares problems on the stiefel manifold.
- [6] A. Edelman, T. A. Arias, S. Smith, The geometry of algorithms with orthogonality constraints, SIAM J. Matrix Anal. Appl. 20 (1998) 303– 353.
- [7] A. W. Bojanczyk, A. Lutoborski, The procrustes problem for orthogonal stiefel matrices, SIAM Journal on Scientific Computing 21 (4) (1999) 1291–1304. doi:10.1137/S106482759630992X.
- [8] Y. Ding, D. Ge, H. Wolkowicz, On equivalence of semidefinite relaxations for quadratic matrix programming, Mathematics of Operations Research 36 (1) (2011) 88–104.
- [9] N. T. Trendafilov, On the l_1 procrustes problem, Future Generation Computer Systems 19 (7) (2003) 1177–1186.
- [10] N. T. Trendafilov, G. Watson, The l_1 oblique procrustes problem, Statistics and Computing 14 (1) (2004) 39–51.
- [11] Z. Zhang, K. Du, Successive projection method for solving the unbalanced procrustes problem, Science in China Series A 49 (7) (2006) 971– 986.

- [12] P. Schönemann, A generalized solution of the orthogonal procrustes problem, Psychometrika 31 (1966) 1–10. doi:10.1007/BF02289451.
- [13] M. T. Chu, N. T. Trendafilov, The orthogonally constrained regression revisited, Journal of Computational and Graphical Statistics 10 (4) (2001) 746–771.
- [14] M. T. Chu, N. T. Trendafilov, On a differential equation approach to the weighted orthogonal procrustes problem, Statistics and Computing 8 (2) (1998) 125–133.
- [15] J. C. Gower, G. B. Dijksterhuis, Procrustes problems, Vol. 30, OUP Oxford, 2004.
- [16] J. C. Gower, G. B. Dijksterhuis, Procrustes problems, Vol. 30, OUP Oxford, 2004.
- [17] L.-E. Andersson, T. Elfving, A constrained procrustes problem, SIAM Journal on Matrix Analysis and Applications 18 (1) (1997) 124–139. doi:10.1137/S0895479894277545.
- [18] J. Berge, The rigid orthogonal procrustes rotation problem, Psychometrika 71 (2006) 201–205. doi:10.1007/s11336-004-1160-5.
- [19] I. Söderkvist, P.-Å. Wedin, On condition numbers and algorithms for determining a rigid body movement, BIT Numerical Mathematics 34 (3) (1994) 424–436.
- [20] M. A. Cox, T. F. Cox, Multidimensional scaling, in: Handbook of data visualization, Springer, 2008, pp. 315–347.
- [21] T. Bell, Global positioning system-based attitude determination and the orthogonal procrustes problem, Journal of Guidance, Control, and Dynamics 26 (5) (2003) 820–822. doi:10.2514/2.5117.
- [22] H. Zhao, Z. Wang, F. Nie, Orthogonal least squares regression for feature extraction, Neurocomputing 216 (2016) 200–207.
- [23] L. Eldén, H. Park, A procrustes problem on the stiefel manifold, Numerische Mathematik 82 (4) (1999) 599–619.

- [24] J. J. Moré, D. C. Sorensen, Computing a trust region step, SIAM Journal on scientific and statistical computing 4 (3) (1983) 553–572.
- [25] J. Francisco, T. Martini, Spectral projected gradient method for the procrustes problem, TEMA (São Carlos) 15 (2014) 83–96.
- [26] J. B. Francisco, F. S. V. Bazán, Nonmonotone algorithm for minimization on closed sets with applications to minimization on stiefel manifolds, Journal of Computational and Applied Mathematics 236 (10) (2012) 2717–2727.
- [27] J. Gower, Multivariate analysis. ordination, multidimensional scaling and allied topics, Handbook of Applicable Mathematics, VI.: Statistics (B).
- [28] N. T. Trendafilov, A continuous-time approach to the oblique procrustes problem, Behaviormetrika 26 (1999) 167–181.
- [29] C. Bogani, M. G. Gasparo, A. Papini, A gss method for oblique l_1 procrustes problems, in: Applied And Industrial Mathematics In Italy III, World Scientific, 2010, pp. 87–98.
- [30] S. A. Mulaik, Foundations of factor analysis, CRC press, 2009.
- [31] I. L. Dryden, K. V. Mardia, Statistical shape analysis: with applications in R, Vol. 995, John Wiley & Sons, 2016.
- [32] N. Gillis, P. Sharma, A semi-analytical approach for the positive semidefinite procrustes problem, Linear Algebra and its Applications 540 (2018) 112–137.
- [33] P. Jingjing, W. Qingwen, P. Zhenyun, C. Zhencheng, Solution of symmetric positive semidefinite procrustes problem, The Electronic Journal of Linear Algebra 35 (2019) 543–554.
- [34] J. E. Brock, Optimal matrices describing linear systems., AIAA Journal 6 (7) (1968) 1292–1296.
- [35] T. Suffridge, T. Hayden, Approximation by a hermitian positive semidefinite toeplitz matrix, SIAM journal on matrix analysis and applications 14 (3) (1993) 721–734.

- [36] A. Minabutdinov, I. Manaev, M. Bouev, Finding the nearest covariance matrix: the foreign exchange market case, Journal of Computational Finance 24 (2).
- [37] M. Fazel, H. Hindi, S. Boyd, Rank minimization and applications in system theory, in: Proceedings of the 2004 American control conference, Vol. 4, IEEE, 2004, pp. 3273–3278.
- [38] J. Dattorro, Convex optimization & Euclidean distance geometry, Lulu. com, 2010.
- [39] M. Fazel, H. Hindi, S. P. Boyd, A rank minimization heuristic with application to minimum order system approximation, in: Proceedings of the 2001 American Control Conference. (Cat. No. 01CH37148), Vol. 6, IEEE, 2001, pp. 4734–4739.
- [40] M. Fazel, H. Hindi, S. P. Boyd, Log-det heuristic for matrix rank minimization with applications to hankel and euclidean distance matrices, in: Proceedings of the 2003 American Control Conference, 2003., Vol. 3, IEEE, 2003, pp. 2156–2162.
- [41] D. Bertsimas, R. Cory-Wright, J. Pauphilet, Mixed-projection conic optimization: A new paradigm for modeling rank constraints, Operations Research 0 (0) (2021) null. doi:10.1287/opre.2021.2182.
- [42] Y. Ye, Interior Point Algorithms: Theory and Analysis, Vol. 51, 1998. doi:10.2307/254021.
- [43] A. Lemon, A. So, Y. Ye, Low-rank semidefinite programming: Theory and applications 2 (2016) 1–156. doi:10.1561/2400000009.
- [44] F. Alizadeh, Interior point methods in semidefinite programming with applications to combinatorial optimization, SIAM Journal on Optimization 5. doi:10.1137/0805002.
- [45] MATLAB, version R2019a, The MathWorks Inc., 2019.
- [46] M. Grant, S. Boyd, CVX: Matlab software for disciplined convex programming, version 2.1, http://cvxr.com/cvx (Mar. 2014).

- [47] M. Grant, S. Boyd, Graph implementations for nonsmooth convex programs, in: V. Blondel, S. Boyd, H. Kimura (Eds.), Recent Advances in Learning and Control, Lecture Notes in Control and Information Sciences, Springer-Verlag Limited, 2008, pp. 95–110, http://stanford.edu/~boyd/graph_dcp.html.
- [48] R. Takapoui, S. Boyd, Linear programming heuristics for the graph isomorphism problem, arXiv preprint arXiv:1611.00711.
- [49] A. Ben-Tal, A. Nemirovski, Lectures on modern convex optimization: analysis, algorithms, and engineering applications, SIAM, 2001.
- [50] F. Zhang, Matrix theory: basic results and techniques, Springer, 2011.
- [51] C. Sun, R. Dai, Rank-constrained optimization and its applications, Automatica 82 (2017) 128–136. doi:10.1016/j.automatica.2017.04.039.