Transversals via regularity

Yangyang Cheng^{*†}

Katherine Staden^{‡§}

June 7, 2023

Abstract

Given graphs G_1, \ldots, G_s all on the same vertex set and a graph H with $e(H) \leq s$, a copy of H is transversal or rainbow if it contains at most one edge from each G_c . We study the case when H is spanning and explore how the regularity blow-up method, that has been so successful in the uncoloured setting, can be used to find transversals. We provide the analogues of the tools required to apply this method in the transversal setting. Our main result is a blow-up lemma for transversals that applies to separable bounded degree graphs H.

Our proofs use weak regularity in the 3-uniform hypergraph whose edges are those xyc where xy is an edge in the graph G_c . We apply our lemma to give a large class of spanning 3-uniform linear hypergraphs H such that any sufficiently large uniformly dense *n*-vertex 3-uniform hypergraph with minimum vertex degree $\Omega(n^2)$ contains H as a subhypergraph. This extends work of Lenz, Mubayi and Mycroft.

1 Introduction

1.1 Transversal embedding

The problem of deciding whether an *n*-vertex graph G contains a given subgraph H is a central topic in graph theory. Since this problem is NP-complete, much of the research on this topic has focused on finding sufficient conditions on G that guarantee the presence of H. Given a graph parameter π , we seek the *best possible* bound $\pi_{H,n}$ such that if $\pi(G) > \pi_{H,n}$, then G contains a copy of H, whereas there are graphs G with $\pi(G) \leq \pi_{H,n}$ which are H-free.

In this paper, we investigate the generalisation of this problem to graph collections, also known as a multilayer graph. Here, we are given graphs G_1, \ldots, G_s on the same vertex set, where $s \ge e(H)$, and we seek a *transversal* copy of H, which is a copy of H containing at most one edge from each of the graphs G_1, \ldots, G_s . We often think of each G_c having the colour c, so a transversal copy of H is also called a *rainbow* copy.

Again, we seek a best possible condition on π which guarantees the existence of H. That is, we seek the minimum $\pi_{H,n}^s$ such that if G_1, \ldots, G_s are any graphs on the same vertex set of size n and $\pi(G_c) > \pi_{H,n}^s$ for all $1 \leq c \leq s$, then there is a transversal copy of H. Observe that we recover the original 'uncoloured' problem when $G_1 = \ldots = G_s$, so the transversal embedding problem is

^{*}Mathematical Institute, University of Oxford, Oxford, UK. Email: yangyang.cheng@maths.ox.ac.uk.

[†]Yangyang Cheng was supported by a PhD studentship of ERC Advanced Grant 883810.

[‡]School of Mathematics and Statistics, The Open University, Walton Hall, Milton Keynes, UK. Email: katherine.staden@open.ac.uk.

[§]Katherine Staden was supported by EPSRC Fellowship EP/V025953/1.

indeed a generalisation. Ideally, we take s = e(H) graphs in the collection and are most interested in determining $\pi_{H,n}^{\text{col}} := \pi_{H,n}^{e(H)}$.

There has been a lot of recent progress in this area. The central question about transversal embeddings, posed by Joos and Kim [25], is whether the addition of colours changes the answer to the problem. That is, given some graph parameter π , when do we have $\pi_{H,n} = \pi_{H,n}^{\text{col}}$? When equality does hold, we say that the embedding problem is *colour-blind*.

We already have a non-example of colour-blindness in the case of the triangle and the size parameter. Mantel's theorem from 1907 states that $e(G) > |n^2/4|$ guarantees that an *n*-vertex graph G contains a triangle, whereas the complete balanced bipartite graph shows that this is best possible. So $e_{K_{3,n}} = \lfloor n^2/4 \rfloor$. However, Aharoni, DeVos, de la Maza, Montejano and Šámal [2] proved that whenever graphs G_1, G_2, G_3 on the same set of n vertices satisfy $e(G_c) > an^2$ for all c = 1, 2, 3, 3where $a := \frac{26-2\sqrt{7}}{81} \approx 0.2557 > 1/4$, then there is a rainbow triangle, while there is an example which shows a cannot be decreased. That is, $e_{K_3,n}^{\text{col}} = \lfloor an^2 \rfloor$. The transversal embedding problem for larger cliques is still open.

When one is interested in embedding a spanning graph, it makes sense to consider minimum degree conditions rather than size conditions, as, for example, a graph could be very dense but contain an isolated vertex, and therefore not contain a copy of any given spanning graph with minimum degree at least 1. Let M_n and C_n be the perfect matching and Hamilton cycle on nvertices respectively. Joos and Kim [25] proved that $\delta_{C_n,n}^{\text{col}} = \delta_{C_n,n}$ and $\delta_{M_n,n}^{\text{col}} = \delta_{M_n,n}$ (which both have the common value = $\lfloor n/2 \rfloor - 1$; that is, Hamilton cycle embedding and matching embedding are both colour-blind with respect to minimum degree.

An easier question is to ask for approximately best possible conditions. For minimum degree, this means we would like to find δ_H and δ_H^{col} , where

- $\tilde{\delta}_H$ is the minimum δ such that for all $\alpha > 0$, any sufficiently large *n*-vertex graph G with $\delta(G) \ge (\delta + \alpha)n$ contains a copy of H; and • δ_{H}^{col} is the minimum δ such that for all $\alpha > 0$, any collection $G_1, \ldots, G_{e(H)}$ of sufficiently large
- *n*-vertex graphs with $\delta(G_c) \geq (\delta + \alpha)n$ for all $1 \leq c \leq e(H)$ contains a transversal copy of H.

So the results of [25] imply that $\tilde{\delta}_{C_n}^{\text{col}} = \tilde{\delta}_{M_n}^{\text{col}} = \frac{1}{2}$; this approximate version was earlier proved by Cheng, Wang and Zhao [11]. If $\tilde{\delta}_H = \tilde{\delta}_H^{\text{col}}$, then we say that *H* is approximately colour-blind.

Montgomery, Müvesser and Pehova [36] determined which F-factors are approximately colourblind for any given small graph F, and showed that spanning trees with maximum degree $o(\frac{n}{\log n})$ are approximately colour-blind. They observed that some spanning graphs are very far from being colour-blind: taking H to be the disjoint union of copies of $K_{2,3} \cup C_4$, it holds that $\widetilde{\delta}_H \leq \frac{4}{9}$ by a result of Kühn and Osthus [31], whereas $\widetilde{\delta}_{H}^{\text{col}} \geq \frac{1}{2}$. Gupta, Hamann, Müyesser, Parczyk and Sgueglia [23] showed that powers of Hamilton cycles are approximately colour-blind.

They also showed, improving results of Cheng, Han, Wang, Wang and Yang [10], that Hamilton ℓ -cycles in k-uniform hypergraphs are approximately 'd-colour-blind' for some range of ℓ , k and d, which means with respect to minimum d-degree, which we do not define here.

The aim of [36] was to provide a fairly general approach for transversal embedding problems that used the corresponding uncoloured embedding result as a black box. Roughly speaking, their approach is designed to work when the graph H to be embedded is made up of small almostdisconnected blocks: they prove results for F-factors, which are made up of vertex-disjoint copies of F, and for trees which are made up of an ordered sequence of small subtrees each sharing one vertex with a previous tree. Similarly, [23] gave a widely applicable sufficient condition for approximate colour-blindness, which applies to graphs obtained by 'cyclically gluing' copies of a smaller graph

(e.g. a Hamilton cycle is obtained by 'cyclically gluing' copies of an edge). Nevertheless, the H to which these two results apply are fairly specific, and in this paper we build on [36] to develop a method for H which are 'more connected' than the above.

Very recently, during the preparation of this paper, Chakraborti, Im, Kim and Liu [9] made important progress in this direction by proving a 'transversal bandwidth theorem'. A graph has bandwidth b if there is an ordering v_1, \ldots, v_n of its vertices such that $|i - j| \leq b$ whenever $v_i v_j$ is an edge. Their result states that if a bounded degree graph H of sublinear bandwidth has chromatic number k, then $\tilde{\delta}_H^{col} \leq 1/k$. This extends several of the above results, and is asymptotically best possible for many graphs H, as well as generalising the original bandwidth theorem of Böttcher, Schacht and Taraz [8] for single graphs. The proof uses similar ideas to the proof of the main result of our paper, a transversal blow-up lemma, which we introduce in the next section.

1.2 The regularity-blow-up method for transversals

The so-called 'regularity-blow-up method' has been employed to prove many results concerning the embedding of a given spanning subgraph H in a large graph G. Such proofs typically run along the following lines. Apply Szemerédi's regularity lemma [41] to obtain a constant-size 'reduced graph' Rof G, which approximates the structure of G: vertices of R correspond to disjoint vertex clusters in G, and edges of R correspond to regular pairs of vertex clusters, which informally means that the edges between them are randomlike and therefore easy to embed into. There may be a small number of 'exceptional' vertices which are not part of this structure. Next, find a suitable subgraph H' of R which is simpler than H, and often consists of many small connected components (whereas H could be connected). Next, embed small pieces of H which connect the components of H'. At this stage, some vertices may need to be moved from the structure into the exceptional set to make clusters balanced and to ensure regular pairs have sufficiently large minimum degree (they are *superregular*). Then incorporate the exceptional vertices: for each such vertex v find a component of H' where v has many suitable neighbours and a few vertices inside the H'-structure which can be used along with vto embed a small part of H. Finally apply the blow-up lemma of Komlós, Särkózy and Szemerédi [28] to embed most of H into H' allowing for the restrictions imposed by the initial embedding. The blow-up lemma states that, for the purposes of embedding a possibly spanning bounded degree graph, a regular pair behaves the same as a complete bipartite graph.

The primary goal of this paper is to provide the tools needed to apply the regularity-blow-up method to obtain transversal embeddings, following the same steps as above. The basic idea is that one can think of a graph collection $\mathbf{G} = (G_c : c \in \mathscr{C})$ with common vertex set V as the 3-uniform hypergraph $G^{(3)}$ with vertex and edge sets

$$V(G^{(3)}) = V \cup \mathscr{C} \quad \text{and} \quad E(G^{(3)}) = \{xyc : xy \in G_c\}.$$

This natural idea has been noticed and used before, for example in [25], where it was noted that a result of Aharoni, Georgakoupoulos and Sprüssel [3] on matchings in k-partite k-uniform hypergraphs can be used to find a transversal matching in a bipartite graph collection; and in [10], where the weak regularity lemma was applied to $G^{(3)}$ to find an almost spanning clique factor. In this paper, we take the idea further by using this perspective to provide the tools one would require to use the regularity-blow-up method for graph collections. To this end, we

- define regularity, superregularity, reduced graphs and provide a regularity lemma for graph collections;
- state some of the standard tools for regularity arguments, such as a 'slicing lemma' which

states that regularity is inherited in subpairs of regular pairs, degree inheritance in the reduced graph, and various embedding lemmas with target sets, candidate sets and prescribed colours;

• prove a blow-up lemma for 'separable graphs', which we define shortly.

The first item is mainly a convenient reformulation of weak regularity for 3-uniform hypergraphs and does not require any new ideas; nonetheless the statements are useful to have. The third (which builds upon the tools developed in the second item) is our main contribution, and the proof uses the original blow-up lemma combined with colour absorption ideas from [36].

We envisage that our blow-up lemma will be useful in solving various transversal embedding problems, making it a viable alternative to the commonly used absorption technique. Furthermore, there are cases where the blow-up lemma is a more appropriate tool. For instance, when attempting to embed a given transversal 2-factor (a spanning 2-regular subgraph) or, more generally, a graph with low bandwidth, the absorption technique is not as effective. In such situations, the blow-up lemma can still be employed for successful embedding. For example, our blow-up lemma can be used to embed powers of Hamilton cycles, providing an alternative proof of a result in [23]. Abbasi [1], proving a conjecture of El-Zahar [17], used the original blow-up lemma to obtain the best possible minimum degree bound in a large graph containing a given 2-factor; our blow-up lemma could be useful in proving a transversal version of this result. Another notable advantage of the blow-up lemma is its ability to aid in characterising extremal constructions. This is helpful for determining exact bounds in these types of problems, by utilising the stability method commonly employed in non-transversal graph embeddings. This natural problem was recently raised in [23]. We discuss applications to transversal embedding problems further in Section 6.

The secondary goal of the paper is to apply our transversal blow-up lemma to embeddings in uniformly dense 3-uniform hypergraphs. We introduce these 'randomlike' hypergraphs, and our results in this direction, in Section 1.5. Indeed, our blow-up lemma can be formulated as a blow-up lemma for 3-uniform hypergraphs. Consequently, there is also potential to employ this lemma in generalising results such as those of Kühn and Osthus [30] who studied the problem of embedding loose Hamiltonian cycles under vertex degree conditions.

All our results apply to 'separable graphs' with bounded maximum degree. An *n*-vertex graph H is μ -separable if there is $X \subseteq V(H)$ of size at most μn such that H - X consists of components of size at most μn . Separable graphs (with suitable small μ) include F-factors for fixed F, trees, 2-regular graphs, powers of a Hamilton cycle, and graphs of small bandwidth.

1.3 Transversals in uniformly dense graph collections

Our first main result concerns transversal embeddings of spanning graphs inside a quasirandom graph collection. A 'quasirandom' condition is one possessed by a random graph of a similar density. Our condition is that for every linear subset of vertices and linear subset of colours, the total number of edges in these colours spanned by the subset is large. A graph collection satisfying this condition is said to be *uniformly dense*. We also require that the number of edges of each colour and the total degree of each vertex is $\Omega(n^2)$, where the graph collection has a common vertex set of size n. This condition cannot be completely removed, since if a colour is not present or if a vertex is isolated, we certainly cannot find a transversal copy of a spanning graph H. A graph collection satisfying both conditions is *super uniformly dense*, in analogy with regular and superregular. We are interested in the following question: for which (spanning) graphs H must any super uniformly dense graph collection contain a transversal copy of H?

Definition 1.1 (Uniformly dense graph collection). Given $d, \eta > 0$, we say that a graph collection

 $G = (G_c : c \in \mathscr{C})$ with vertex set V of size n is (d, η) -dense if for all $A, B \subseteq V$ and $\mathscr{C}' \subseteq \mathscr{C}$, we have

$$\sum_{c \in \mathscr{C}'} e(G_c[A, B]) \ge d|\mathscr{C}'||A||B| - \eta n^3,$$

where $e(G_c[A, B])$ denotes the number of pairs $(x, y) \in A \times B$ for which $\{x, y\}$ is an edge of G_c . We informally refer to a graph collection G on n vertices that is (d, η) -dense for parameters $0 < 1/n \ll \eta \ll d \leq 1$ as uniformly dense, and if additionally there is $\alpha \gg \eta$ for which $\sum_{c \in \mathscr{C}} d_{G_c}(v) \geq \alpha |\mathscr{C}|n$ for all $v \in V$ and $e(G_c) \geq \alpha n^2$ for all $c \in \mathscr{C}$, then G is super uniformly dense.

Note that the condition is vacuous unless A, B and \mathscr{C}' are of linear size (in the number n of vertices). We prove that a transversal copy of a separable graph can be found in a super uniformly dense graph collection.

Theorem 1.2. For all Δ , d, δ , $\alpha > 0$, there are η , $\mu > 0$ and $n_0 \in \mathbb{N}$ such that the following holds for all integers $n \ge n_0$. Let $\mathbf{G} = (G_c : c \in \mathscr{C})$ be a (d, η) -dense graph collection on a common vertex set V of size n, where $|\mathscr{C}| \ge \delta n$, and suppose that for all $v \in V$ we have $\sum_{c \in \mathscr{C}} d_{G_c}(v) \ge \alpha |\mathscr{C}|n$, and for all $c \in \mathscr{C}$ we have $e(G_c) \ge \alpha n^2$. Then \mathbf{G} contains a transversal copy of any given μ -separable graph H on n vertices with $\Delta(H) \le \Delta$ and $e(H) = |\mathscr{C}|$.

This result is a consequence of a more general transversal blow-up lemma, which we state in the next section. First, we discuss possible extensions to other colour patterns. One can ask whether a graph collection contains a copy of H with a given colour pattern, extending the monochromatic case (a copy in a single graph G_c) and rainbow case (a copy with one edge from each G_c). For example, the following was shown in [36]. Not only does the same minimum degree $(\frac{2}{3} + o(1))3n$ which guarantees a triangle factor in a large graph with a vertex set of size 3n in fact guarantee a transversal copy when $|\mathscr{C}| = 3n$, it also guarantees a triangle factor where the *c*-th triangle lies inside G_c (has colour *c*), when $|\mathscr{C}| = n$.

Given $d, \eta > 0$, a (single) graph with a vertex set V of size n is (d, η) -dense if for all $A, B \subseteq V$, we have $e(G[A, B]) \ge d|A||B| - \eta n^2$. A collection of (d, η) -dense graphs on the same vertex set is a (d, η) -dense graph collection, but the converse does not hold. It is easy to see that, for any d > 0, as long as η and 1/n are sufficiently small, such a G contains $\Omega(n^3)$ triangles: the condition applied to A = B = V implies that there are $\Omega(n)$ vertices v with $d_G(v) \ge dn/2$; each one has $\Omega(n^2)$ edges in its neighbourhood.

However, there are uniformly dense graph collections with d fairly large which do not contain any monochromatic triangles, as the following example shows. (The example is essentially equivalent to one in the setting of uniformly dense hypergraphs – introduced in Section 1.5 – of Reiher, Rödl and Schacht [38], which itself has its roots in work of Erdős and Hajnal [19].) Form an auxiliary oriented 2-graph J by letting V, \mathscr{C} be disjoint, where, say, $|V| = |\mathscr{C}| = n$ is large, and first adding edges between every pair in V, and every pair in (V, \mathscr{C}) . Independently for each edge, choose an orientation uniformly at random. Add the pair xy to G_c with $x, y \in V$ and $c \in \mathscr{C}$ precisely when xycis a cyclic triangle. Then, with probability tending to 1 as $n \to \infty$, G is $(\frac{1}{4}, o(1))$ -dense. However, for every triple x, y, z of vertices in V and every colour $c \in \mathscr{C}$, there are at most two edges in $G_c[\{x, y, z\}]$.

This raises the question of which colour patterns of which graphs H one can expect to find in a (super) uniformly dense graph collection. If d is sufficiently large, then any bounded degree H with any colour pattern can be found; for those H which are not present when d is an arbitrary positive constant, how large must d be to guarantee such a copy of H? We will explore a generalisation of this question in Section 1.5.

1.4 A transversal blow-up lemma

In this section, we state a simplified version of our blow-up lemma for bipartite graphs. We defer the complete statement to Theorem 1.8 at the end of this section.

Our blow-up lemma has the advantages that its proof, sketched at the beginning of Section 4, is conceptually straightforward, following from the original blow-up lemma and a colour absorption tool introduced in [36]; and the lemma should be powerful enough for all of the transversal embedding applications we have in mind. This is since the usual regularity-blow-up method is almost always successfully applied to 'non-expanding' separable graphs, since one embeds them by using the blowup lemma on a series of small pieces. The disadvantage is that there does not seem to be a reason why the separability condition should be necessary.

Theorem 1.3 (Simplified transversal blow-up lemma). Let $0 < 1/n \ll \varepsilon, \mu \ll d, \delta, 1/\Delta \leq 1$. Let \mathscr{C} be a set of at least δn colours and let $\mathbf{G} = (G_c : c \in \mathscr{C})$ be a collection of bipartite graphs with the same vertex partition V_1, V_2 , where $n \leq |V_1| \leq |V_2| \leq n/\delta$, such that

• for all $V'_i \subseteq V_i$ for i = 1, 2 and $\mathscr{C}' \subseteq \mathscr{C}$ with $|V'_i| \ge \varepsilon |V_i|$ for i = 1, 2 and $|\mathscr{C}'| \ge \varepsilon |\mathscr{C}|$, we have that

$$\sum_{c \in \mathscr{C}'} e_{G_c}(V_1', V_2') \ge d|\mathscr{C}'||V_1'||V_2'|;$$

• for i = 1, 2 and every $v \in V_i$ we have $\sum_{c \in \mathscr{C}} d_{G_c}(v) \ge d|\mathscr{C}|n$ and for every $c \in \mathscr{C}$, we have $e(G_c) \ge dn^2$.

Let H be a μ -separable bipartite graph with parts of size $|V_1|, |V_2|$, and $|\mathcal{C}|$ edges and maximum degree Δ . Then G contains a transversal copy of H.

The theorem requires that the number of vertices and edges of H are comparable, so it does not apply to very sparse H. If one adds edges to H to obtain a suitable denser graph H', and duplicates colours until there are e(H') colours, the copy of H' produced by the theorem will not necessarily contain a transversal copy of H.

The general version of the theorem, Theorem 1.8, applies to a graph collection whose graphs have common parts V_1, \ldots, V_r , and a graph R with V(R) = [r] such that the bipartite condition in the theorem holds between all $ij \in E(R)$, and each ij has a dedicated set of colours \mathscr{C}_{ij} .

1.4.1 Rainbow blow-up lemmas

There are by now many blow-up lemmas for various settings which have been applied to many embedding problems. Of particular relevance to this paper are rainbow blow-up lemmas which apply to a single graph whose edges are coloured. We say that an edge-coloured graph G is k-bounded if no colour appears on more than k edges, and G is locally k-bounded if each vertex is incident to at most k edges of the same colour. Glock and Joos [22] proved a rainbow blow-up lemma for o(n)-bounded edge-colourings, which allows one to find a rainbow embedding of a given bounded degree graph H. Here, the number of colours is many times larger than e(H). Later, Ehard, Glock, and Joos [16] proved a similar lemma for locally O(1)-bounded colourings, which allows the number of colours to be (1 + o(1))e(H). Therefore it does not seem possible to use such blow-up lemmas for transversal embedding problems where we require exactly e(H) colours.

1.5 Applications to embedding in uniformly dense hypergraphs

In this section we revisit the hypergraph perspective and discuss the consequences of our work to embeddings in 3-uniform hypergraphs. 'Weak regularity' is the straightforward generalisation of Szemerédi regularity from (2-uniform hyper)graphs to k-uniform hypergraphs (hereafter referred to as k-graphs), and the 'weak regularity lemma' was proved by Chung [12] in much the same way as the original [41]. However, this lemma is not powerful enough to prove many of the analogues of graph results which one would like. In particular, there is no general counting lemma which guarantees that the number of copies of a given small graph F in a regularity partition is similar to what one would expect if pairs were replaced by uniform random hyperedges, with the same density. Thus the 'strong regularity lemma' was developed, which uses, as the name suggests, a more complicated and much stronger notion of regularity, and does have an associated counting lemma; this lemma is much more applicable than its weaker counterpart.

It was shown by Conlon, Hàn, Person and Schacht [13] and independently Kohayakawa, Nagle, Rödl and Schacht [27] that weak regularity is in fact strong enough to give a counting lemma for *linear* hypergraphs, which have the property that $|e \cap f| \leq 1$ for all distinct hyperedges e, f. As a transversal copy of a (simple) graph inside a graph collection G is a linear subhypergraph of the associated 3-graph $G^{(3)}$, weak regularity is an effective tool for transversal embedding problems. And conversely, the tools developed in this paper are useful for embedding 3-graphs.

We generalise the definition we gave in Section 1.3 for graphs.

Definition 1.4 (Uniformly dense k-graph). Let $d, \eta > 0$ and let $k \ge 2$ be an integer. A k-graph is (d, η) -dense if for all subsets $U_1, \ldots, U_k \subseteq V$, we have

$$e(U_1,\ldots,U_k) \ge d|U_1|\ldots|U_k| - \eta n^k,$$

where $e(U_1, \ldots, U_k)$ is the number of k-tuples $(x_1, \ldots, x_k) \in U_1 \times \ldots \times U_k$ for which $\{x_1, \ldots, x_k\}$ is a hyperedge. We informally refer to a k-graph G on n vertices that is (d, η) -dense for parameters $0 < 1/n \ll \eta \ll d \le 1$ as uniformly dense. If there is $\alpha \gg \eta$ for which additionally $d_G(v) \ge \alpha n^{k-1}$ for all $v \in V(G)$, then we say that G is super uniformly dense.

This is a type of quasirandomness, since a random k-graph of density at least d is (d, o(1))-dense with high probability. Several papers studying uniform density use the term 'quasirandom' instead.

Following a suggestion of Erdős and Sós [20], a systematic treatment of extremal problems in uniformly dense hypergraphs has been started by Rödl, Reiher and Schacht, in part due to the great difficulty of such problems in general hypergraphs. In [37], they fully answered the zero Turán density question for uniformly dense 3-graphs, i.e. for which F is the following true? For all d > 0, there exists $\eta > 0$ such that every sufficiently large (d, η) -dense 3-graph contains F as a subhypergraph. They showed that these F are precisely those with the following property. There is an ordering v_1, \ldots, v_r of V(F) and a colouring of the set of pairs of vertices contained in edges by red, blue and green, so that whenever $v_i v_j v_k \in E(F)$ with i < j < k, we have that $v_i v_j$ is red, $v_i v_k$ is blue and $v_j v_k$ is green. This set of F includes linear F and 3-partite F (Erdős [18] proved that a k-graph Fhas Turán density 0 if and only if F is k-partite), but is much richer than this: for example, the hypergraph obtained by removing one edge from the tight cycle on 5 vertices is such a hypergraph.

The simple argument given in Section 1.3 shows that for k = 2 and $\Delta > 0$, a sufficiently large uniformly dense 2-graph contains a copy (in fact many copies) of K_{Δ} . In contrast, for $k \ge 3$ there are very simple k-graphs F which require a fairly large density d to appear in any uniformly dense kgraph. Indeed, there is a $(\frac{1}{4}, o(1))$ -dense 3-graph in which $K_4^{(3)-}$, the (unique) 3-graph with 4 vertices and 3 edges, does not appear. (The example is closely related to the one given in Section 1.3.) In this paper, we are interested in which *spanning* hypergraphs appear in any super uniformly dense hypergraph. Note that we cannot remove 'super' since uniform density does not preclude the existence of isolated vertices.

Problem 1.5. Let $\mathcal{H} = (\mathcal{H}_{\ell} : \ell \in \mathbb{N})$ be a collection of k-graphs where $|V(\mathcal{H}_{\ell})| =: n_{\ell} \to \infty$. For which \mathcal{H} is the following true? For all $d, \alpha > 0$, there exist $\eta > 0$ and $\ell_0 \in \mathbb{N}$ such that for all integers $\ell > \ell_0$, every (d, η) -dense k-graph G on n_{ℓ} vertices with $d_G(v) \ge \alpha n_{\ell}^{k-1}$ for all $v \in V(G)$ contains \mathcal{H}_{ℓ} as a subhypergraph.

For k = 2, the blow-up lemma of Komlós, Sárközy and Szemerédi [28] implies that every uniformly dense graph contains as a subgraph any given spanning graph H with bounded degree and sublinear bandwidth (as observed by Glock and Joos, see Theorem 9.3 in [40]).

For general uniformities k, Lenz and Mubayi [34] proved that uniformly dense k-graphs contain an F-factor for any given fixed size linear F. In fact they found non-linear F for which uniform density is sufficient, and 'almost-linear' F for which it is not. Ding, Han, Sun, Wang and Zhou [14] completed this work for k = 3 by characterising those F for which uniform density guarantees an F-factor, and for general k they also obtained a characterisation for k-partite F. Lenz, Mycroft and Mubayi [35] showed that uniform density guarantees a loose cycle. To the best of our knowledge, there are no further results on embedding spanning F in uniformly dense hypergraphs of arbitrarily small positive density.

Our next main result generalises the result of [35] for k = 3 by providing a new family of spanning H which can be found in super uniformly dense 3-graphs. An expanded graph is obtained from a 2-graph by replacing every edge e = xy by some 3-edges xyc_1, \ldots, xyc_{t_e} , where all the new vertices c_1, \ldots, c_{t_e} are distinct, and the number t_e of new edges/vertices for the edge e depends on e. If every t_e is equal to t then we call this a *t*-expansion. For example, a loose 3-uniform cycle is an expanded cycle where each edge is replaced by one expanded edge, that is, a 1-expansion of a cycle. The 1-expansion of a (simple) graph H is linear and has |V(H)| + |E(H)| vertices.

Theorem 1.6. For all Δ , $d, \alpha > 0$ there are $\eta, \mu > 0$ and $n_0 \in \mathbb{N}$ such that the following holds for all integers $n \ge n_0$. Let G be a (d, η) -dense 3-graph on n vertices with $d_G(v) \ge \alpha n^2$ for all $v \in V(G)$. Then G contains a copy of the 1-expansion of any given μ -separable graph H with $\Delta(H) \le \Delta$ and $|V(H)| + |E(H)| \le n$.

Since a large 2-regular graph (a union of vertex-disjoint cycles) is o(1)-separable, the theorem implies that any super uniformly dense 3-graph G contains as a subhypergraph any 3-graph consisting of vertex-disjoint loose cycles.

Next, we reformulate Theorem 1.3 as a blow-up lemma for 3-graphs, which may be of independent interest.

Theorem 1.7 (Simplified weak hypergraph blow-up lemma). For all Δ , $d, \delta > 0$ there are $\varepsilon, \mu > 0$ and $n_0 \in \mathbb{N}$ such that the following holds for all integers $n \ge n_0$. Let G be a 3-partite 3-graph with parts V_1, V_2, V_3 where every $|V_i| =: n_i$ and $n \le n_1 \le n_2 \le n/\delta$ and $n_3 \ge \delta n$ such that G is a weakly (ε, d) -half-superregular triple, that is,

(i) for all $i \in [3]$ and $V'_i \subseteq V_i$ with $|V'_i| \ge \varepsilon |V_i|$, we have $e_G(V'_1, V'_2, V'_3) \ge d|V'_1||V'_2||V'_3|$, (ii) $d_G(v) \ge dn^2$ for all $v \in V(G)$.

Let H be a μ -separable bipartite 2-graph with parts of size n_1, n_2 , with n_3 edges and maximum degree Δ . Then G contains a copy of the 1-expansion of H, where the images of the new vertices lie in V_3 .

Weak superregularity and uniform density of 3-graphs and graph collections are closely connected. Observe that if G is $(d + \sqrt{\eta}, \eta)$ -dense, where $\eta \gg \varepsilon, \delta$, and V_1, V_2, V_3 is any partition with sizes as in the theorem, then the 3-partite subhypergraph G' of G induced on these parts satisfies (i). If $d_G(v) \ge \alpha n^2$ for all $v \in V$, and in addition the vertex partition is uniformly random, then with probability tending to 1 as $n \to \infty, G'$ is weakly $(\varepsilon, \min\{d, \alpha/2\})$ -half-superregular, say. Furthermore, if one takes any partition $V(G) = V \cup \mathscr{C}$ into parts which are not too small, then the graph collection G consisting of graphs G_c for $c \in \mathscr{C}$ with $V(G_c) = V$ and $E(G_c) = \{xy : x, y \in V \text{ and } xyc \in E(G)\}$ is (d, η) -dense. Again, if $d_G(v) \ge \alpha n^2$, a random such partition gives rise to a super uniformly dense graph collection.

As an example, any 3-graph G satisfying the hypotheses of the theorem with $n_1 = n_2$ and $n_3 = n_1 + n_2$ contains a copy of a loose Hamilton cycle. The full statement of a 3-graph version of our transversal blow-up lemma (Theorem 1.8) is stated in Theorem 6.1.

A much more general hypergraph blow-up lemma was established by Keevash [26] which applies to k-graphs satisfying a strong regularity property, which is designed to be used with the strong regularity lemma; this blow-up lemma applies to embed any bounded degree k-graph H. Nevertheless, it would be interesting to determine which graphs can be embedded into a weakly superregular triple. We discuss this further in Section 6.

There are many other notions of quasirandomness of hypergraphs, often related to the number of edges in/between subsets, that have been studied in the literature. We refer the reader to [4, 33, 43] for a detailed comparison of such notions; it turns out that equivalent notions of quasirandomness in graphs have inequivalent hypergraph analogues. The embedding of spanning structures has been studied in these various settings, for example tight Hamilton cycles in 3-graphs with a stronger quasirandomness condition than the one in this paper, in [5, 6, 21], and graphs of sublinear bandwidth in 'locally dense' 2-graphs, which is weaker than uniformly dense, in [40], and in 'inseparable' 2-graphs in [15]. While in this paper we ask which H are subhypergraphs of any quasirandom hypergraph G of positive density (where for us, 'quasirandom' means 'uniformly dense'), the results above mainly concern the generalisation of this question: given H, what minimum degree in a quasirandom hypergraph G is sufficient to imply the existence of H? There are various notions of degree in hypergraphs and the results mentioned in this paragraph consider several of them. Quasirandomness conditions that apply to sparse (hyper)graphs have also been well studied. Recently, Hàn, Han and Morris [24] extended the results of [34, 35] to the sparse regime.

1.6 The statement of our main result

The full statement of our transversal blow-up lemma is as follows. Its proof will follow from Theorem 3.5 which is a very similar statement written in the notation defined in Section 2.

Theorem 1.8 (Transversal blow-up lemma). For all $\nu, d, \delta, \Delta, r > 0$ where $r \ge 2$ is an integer, there exist $\varepsilon, \mu, \alpha > 0$ and $m_0 \in \mathbb{N}$ such that the following holds for all integers $m \ge m_0$.

Suppose that $G = (G_c : c \in \mathscr{C})$ is a graph collection with the following properties.

- There is a graph R with vertex set [r] and a partition $\mathscr{C} = \bigcup_{e \in E(R)} \mathscr{C}_e$ where $|\mathscr{C}_e| \ge \delta m$ for all $e \in E(R)$;
- for all $ij \in E(R)$ and $c \in \mathscr{C}_{ij}$, G_c is bipartite with parts V_i, V_j , and V is a vertex set of size n with partition $V = V_1 \cup \ldots \cup V_r$, where $m \leq |V_i| \leq m/\delta$ for each $i \in [r]$;
- for all $ij \in E(R)$,
 - $\text{ for all } V'_h \subseteq V_h \text{ for } h = i, j \text{ and } \mathscr{C}'_{ij} \subseteq \mathscr{C}_{ij} \text{ with } |V'_h| \ge \varepsilon |V_h| \text{ for } h = i, j \text{ and } |\mathscr{C}'_{ij}| \ge \varepsilon |\mathscr{C}_{ij}|,$

we have that

$$\sum_{c \in \mathscr{C}'_{ij}} e_{G_c}(V'_i, V'_j) \ge d|\mathscr{C}'_{ij}||V'_i||V'_j|;$$

- for h = i, j and every $v \in V_h$ we have $\sum_{c \in \mathscr{C}_{ij}} d_{G_c}(v) \ge d|\mathscr{C}_{ij}|m$ and for every $c \in \mathscr{C}_{ij}$, we have $e(G_c) \ge dm^2$.

Suppose that H is a graph with the following properties.

- $\Delta(H) \leq \Delta;$
- *H* is μ -separable;
- *H* has vertex partition $A_1 \cup \ldots \cup A_r$ such that $|A_i| = |V_i|$ for all $i \in [r]$ and for every $xy \in E(H)$ there is $ij \in E(R)$ such that $x \in A_i$ and $y \in A_j$;
- $e(H[A_i, A_j]) = |\mathscr{C}_{ij}|$ for all $ij \in E(R)$;
- for each $i \in [r]$, there is a set $U_i \subseteq A_i$ with $|U_i| \leq \alpha m$ and for each $x \in U_i$, a set $T_x \subseteq V_i$ with $|T_x| \geq \nu m$.

Then there is a transversal embedding of H inside G such that for every $i \in [r]$, every $x \in U_i$ is embedded inside T_x .

1.7 Notation and organisation

Notation. For reals x, a, b, we write $x = a \pm b$ if we have $a - b \leq x \leq a + b$. For any two constants $\alpha, \beta \in (0, 1)$, we write $\alpha \ll \beta$ if there exists a function $\alpha_0 = \alpha_0(\beta)$ such that the subsequent arguments hold for all $0 < \alpha \leq \alpha_0$. When we write multiple constants in a hierarchy, we mean that they are chosen from right to left. For any positive integer a, let $[a] := \{1, 2, \ldots, a\}$. Given a set X and a positive integer b, write $\binom{X}{b}$ to denote the set of b-element subsets of X.

Let $k \geq 2$ be an integer and let G = (V, E) be any k-graph, so each edge consists of k vertices. We use V(G) := V to denote its vertex set and E(G) := E to denote its edge set. Let v(G) := |V(G)| and e(G) := |E(G)| be their sizes. For any vertex $v \in V(G)$, let $N_G(v)$ be the *neighbourhood* of v, that is, the set of (k-1)-tuples of v that are incident to v and let $d_G(v) := |N_G(v)|$ be the *degree* of v. Sometimes we use d(v) for short if G is obvious from the text. We write $\delta(G) := \min_{v \in V(G)} d_G(v)$ and $\Delta(G) := \max_{v \in V(G)} d_G(v)$ for the minimum and maximum degree of G. For each vertex $v \in V(G)$ and subset $S \subseteq \binom{V(G)}{k-1}$, let $N_G(v, S) := N_G(v) \cap S$ and $d_G(v, S) := |N_G(v, S)|$. For any vertex set $U \subseteq V(G)$, let G[U] be the induced hypergraph of G on U, i.e. the hypergraph with vertex set U and all the edges of G with vertices only in U. Let $G - U := G[V(G) \setminus U]$. For a subhypergraph H of G, let $G \setminus H$ be the graph with vertex set V(G) and edge set $E(G) \setminus E(H)$. For any (not necessarily disjoint) vertex sets $U_1, \ldots, U_k \subseteq V(G)$, we write $G[U_1, \ldots, U_k]$ for the k-graph with vertex set $U \in V_i(G) \setminus U_i$. Let $e_G(U_1, \ldots, U_k)$ be the number of k-tuples $(x_1, \ldots, x_k) \in U_1 \times \ldots \times U_k$ for which $\{x_1, \ldots, x_k\} \in E_G(U_1, \ldots, U_k)$ (note that when these sets are not disjoint, edges may be counted more than once).

We say that a statement about a graph on n vertices holds with high probability if it holds with probability tending to 1 as $n \to \infty$. We use script letters (e.g. \mathscr{C}) to denote sets of colours and bold letters (e.g. G) to denote graph collections. It will be convenient to represent a graph collection in three equivalent ways:

- a collection $G = (G_c : c \in \mathscr{C})$ of graphs on the same vertex set V. We call \mathscr{C} the *colour set* of G;
- an edge-coloured graph \mathcal{G} on vertex set V with edge set $\bigcup_{c \in \mathscr{C}} G_c$, where xy has a multiset of

colours consisting of those $c \in \mathscr{C}$ for which $xy \in G_c$. We say that \mathcal{G} is the *edge-coloured graph* of G (but rarely use this representation);

• a 3-graph $G^{(3)}$ on vertex set $V \cup \mathscr{C}$ with edges xyc whenever $xy \in G_c$. We say that $G^{(3)}$ is the 3-graph of G.

Given two 2-graphs H and G, a graph homomorphism (from H to G) is a map $\phi: V(H) \to V(G)$ such that $\phi(x)\phi(y) \in E(G)$ whenever $xy \in E(H)$. An injective graph homomorphism is an embedding (of H into G). Given a graph H and a graph collection $\mathbf{G} = (G_c: c \in \mathscr{C})$ on vertex set V, a transversal embedding (of H into \mathbf{G}) is a pair $\tau: V(H) \to V$ and $\sigma: E(H) \to \mathscr{C}$ of injective maps such that $\tau(x)\tau(y) \in E(G_{\sigma(xy)})$ whenever $xy \in E(H)$.

Organisation. In Section 2, we define regularity for graph collections, state a regularity lemma for graph collections, define the 'template' graph collections to which our theory applies, and prove some of the basic tools one uses when applying the regularity method, such as a slicing lemma and a degree inheritance lemma. Section 3 contains some embedding lemmas which are the main ingredients of the proof of our transversal blow-up lemma, Theorem 3.5, but which are also useful tools when applying the method. In Section 4, we prove Theorem 3.5 (which readily implies Theorem 1.8) and in Section 5, we prove Theorems 1.2 and 1.6 on embeddings in uniformly dense graph collections and 3-graphs. Section 6 contains some concluding remarks.

2 The regularity-blow-up method for transversals

2.1 Regularity

Our notion of regularity for a graph collection G is essentially weak regularity for the 3-graph $G^{(3)}$ of G. We now define several related notions of weak regularity for k-graphs, which we will use for k = 2, 3. We usually omit weak and weakly since we will not use any stronger type of regularity.

Let G be a k-partite k-graph with classes V_1, \ldots, V_k , which we also denote as $(V_1, \ldots, V_k)_G$. We define the *density* of G to be

$$d_G(V_1,\ldots,V_k):=\frac{e(G)}{|V_1|\ldots|V_k|}.$$

Given $\varepsilon > 0$, we say that $(V_1, \ldots, V_k)_G$ is

• (weakly) ε -regular if for every subhypergraph $(V'_1, \ldots, V'_k)_G$ with $V'_i \subseteq V_i$ and $|V'_i| \ge \varepsilon |V_i|$ for all $i \in [k]$ we have

$$|d_G(V'_1,\ldots,V'_k)-d_G(V_1,\ldots,V_k)|<\varepsilon;$$

- (weakly) (ε, d) -regular if additionally $d_G(V_1, \ldots, V_k) \ge d$;
- (weakly) (ε , d)-superregular if it is (weakly) (ε , d)-regular and additionally $d_G(x) \ge d(|V_1| \dots |V_k|)/|V_i|$ for all $i \in [k]$ and $x \in V_i$;
- (weakly) (ε, d) -half-superregular if for every subhypergraph $(V'_1, \ldots, V'_k)_G$ with $V'_i \subseteq V_i$ and $|V'_i| \ge \varepsilon |V_i|$ for all $i \in [k]$ we have $d_G(V'_1, \ldots, V'_k) \ge d$ and $d_G(x) \ge d(|V_1| \ldots |V_k|)/|V_i|$ for all $i \in [k]$ and $x \in V_i$.

Our main results will use 'half-superregular' given its close connection to uniform density. The 2graph blow-up lemma [29] uses a similar notion, but nowadays 'superregular' is generally used instead. This makes little material difference: by definition, an (ε, d) -superregular hypergraph is $(\varepsilon, d - \varepsilon)$ half-superregular, and as shown by Rödl and Ruciński in the course of their alternative proof of the blow-up lemma [39], a half-superregular 2-graph contains a spanning superregular subgraph, with weaker parameters. This proof generalises easily to k-graphs, and therefore we postpone it to the appendix.

Lemma 2.1. Let $0 < 1/n \ll \varepsilon \ll \varepsilon' \ll d, \delta, 1/k \leq 1$ where $k \geq 2$ is an integer. Suppose that G is a k-partite k-graph with parts V_1, \ldots, V_k where $\delta n \leq |V_i| \leq n/\delta$ for all $i \in [k]$. If G is (ε, d) -half-superregular, then G contains a spanning subhypergraph G' that is $(\varepsilon', d^2/2)$ -superregular.

Regularity for a bipartite graph collection G is defined in terms of the 3-graph $G^{(3)}$ of G.

Definition 2.2 (Regularity, semi- and half-superregularity, superregularity). Suppose that G is a graph collection with colour set \mathscr{C} , where each G_c is bipartite with parts V_1, V_2 . Let $G^{(3)}$ be the 3-graph of G. We say that

• **G** is (ε, d) -regular if $G^{(3)}$ is (ε, d) -regular. That is, for all $V'_i \subseteq V_i$ with $|V'_i| \ge \varepsilon |V_i|$ for i = 1, 2and $\mathscr{C}' \subseteq \mathscr{C}$ with $|\mathscr{C}'| \ge \varepsilon |\mathscr{C}|$, we have

$$\left|\frac{\sum_{c\in\mathscr{C}'}e_{G_c}(V_1',V_2')}{|\mathscr{C}'||V_1'||V_2'|}-\frac{\sum_{c\in\mathscr{C}}e_{G_c}(V_1,V_2)}{|\mathscr{C}||V_1||V_2|}\right|<\varepsilon$$

- **G** is (ε, d) -semi-superregular if it is (ε, d) -regular and $d_{G^{(3)}}(v) = \sum_{c \in \mathscr{C}} d_{G_c}(v) \ge d|V_{3-i}||\mathscr{C}|$ for all $i \in [2]$ and $v \in V_i$.
- **G** is (ε, d) -superregular if $G^{(3)}$ is (ε, d) -superregular. That is, it is (ε, d) -semi-superregular and $d_{G^{(3)}}(c) = e(G_c) \ge d|V_1||V_2|$ for all $c \in \mathscr{C}$.
- \widetilde{G} is (ε, d) -half-superregular if $G^{(3)}$ is (ε, d) -half-superregular. That is, for all $V'_i \subseteq V_i$ with $|V'_i| \ge \varepsilon |V_i|$ for i = 1, 2 and $\mathscr{C}' \subseteq \mathscr{C}$ with $|\mathscr{C}'| \ge \varepsilon |\mathscr{C}|$, we have $\sum_{c \in \mathscr{C}'} e_{G_c}(V'_1, V'_2) \ge d|\mathscr{C}'||V'_1||V'_2|$ and $\sum_{c \in \mathscr{C}} d_{G_c}(v) \ge d|V_{3-i}||\mathscr{C}|$ for all i = 1, 2 and $v \in V_i$, and $e(G_c) \ge d|V_1||V_2|$ for all $c \in \mathscr{C}$.

Note that, if every G_c with $c \in \mathscr{C}$ is the same, then G is (ε, d) -regular if and only if G_c is (ε, d) -regular; and G is (ε, d) -superregular if and only if G_c is (ε, d) -superregular. The superregularity of G does not imply a minimum degree condition for any graph G_c in the collection, and indeed they could all have isolated vertices.

The following simple lemma shows that, in an (ε, d) -regular graph collection, most vertices – *typical* vertices – have large total degree (the sum of degrees over all colours) and typical colours have many edges.

Lemma 2.3 (Typical vertices and colours). Let $0 < \varepsilon \ll d \leq 1$, and let G be an (ε, d) -regular graph collection with colour set \mathscr{C} , where each G_c is bipartite with parts V_1, V_2 . Then the following hold:

(i) for every $i \in [2]$ and all but at most $\varepsilon |V_i|$ vertices $v \in V_i$ we have $\sum_{c \in \mathscr{C}} d_{G_c}(v) \ge (d-\varepsilon)|V_{3-i}||\mathscr{C}|$; (ii) for all but at most $\varepsilon |\mathscr{C}|$ colours $c \in \mathscr{C}$ we have $e(G_c) \ge (d-\varepsilon)|V_1||V_2|$.

Proof. For (i), let V'_1 be the set of vertices in V_1 without this property and suppose for a contradiction that $|V'_1| > \varepsilon |V_1|$. Then (ε, d) -regularity implies that $d_{G^{(3)}}(V'_1, V_2, \mathscr{C}) > d - \varepsilon$, but the definition of V'_1 implies that $e_{G^{(3)}}(V'_1, V_2, \mathscr{C}) \le |V'_1|(d - \varepsilon)|V_2||\mathscr{C}|$ and hence $d_{G^{(3)}}(V'_1, V_2, \mathscr{C}) \le d - \varepsilon$, a contradiction. By symmetry, the rest of (i) and (ii) are identical.

The following lemma is a standard tool, written in our graph collection notation.

Lemma 2.4 (Slicing lemma). Let $0 < 1/n \ll \varepsilon \ll \varepsilon' \ll \alpha \ll d \leq 1$, and let G be a graph collection with colour set \mathscr{C} , where each G_c is bipartite with parts V_1, V_2 each of size at least n, and let $V'_i \subseteq V_i$ for $i \in [2]$ and $\mathscr{C}' \subseteq \mathscr{C}$. Let $G' = (G_c[V'_1, V'_2] : c \in \mathscr{C}')$.

- (i) Suppose that **G** is (ε, d) -regular. Suppose $|V'_i| \ge \alpha |V_i|$ for $i \in [2]$ and $|\mathscr{C}'| \ge \alpha |\mathscr{C}|$. Then **G**' is $(\varepsilon/\alpha, d/2)$ -regular.
- (ii) Suppose that **G** is (ε, d) -superregular. Suppose $|V'_i| \ge (1-\alpha)|V_i|$ for $i \in [2]$ and $|\mathscr{C}'| > (1-\alpha)|\mathscr{C}|$. Then **G'** is $(2\varepsilon, d/2)$ -superregular.
- (iii) Suppose that G is (ε, d) -superregular. Given $n_1 \ge \alpha |V_1|, n_2 \ge \alpha |V_2|$ and $h \ge \alpha |\mathscr{C}|$, if $V'_i \subseteq V_i$ is a uniform random subset of size n_i for each i = 1, 2 and $\mathscr{C}' \subseteq \mathscr{C}$ is a uniform random subset of size h, then with high probability, G' is $(\varepsilon/\alpha, d^2/16)$ -superregular.

Proof. For (i), firstly we have $d_{G^{(3)}}(V'_1, V'_2, \mathscr{C}') \ge d - \varepsilon \ge d/2$. Secondly, given any subset $V''_i \subseteq V'_i$ with $|V''_i| \ge \varepsilon |V'_i|/\alpha \ge \varepsilon |V_i|$ where i = 1, 2 and any subset $\mathscr{C}'' \subseteq \mathscr{C}'$ with $|\mathscr{C}''| \ge \varepsilon |\mathscr{C}'|/\alpha \ge \varepsilon |\mathscr{C}|$, by regularity we have

$$\begin{aligned} |d_{G^{(3)}}(V_1'', V_2'', \mathscr{C}'') - d_{G^{(3)}}(V_1', V_2', \mathscr{C}')| \\ &\leq |d_{G^{(3)}}(V_1'', V_2'', \mathscr{C}'') - d_{G^{(3)}}(V_1, V_2, \mathscr{C})| + |d_{G^{(3)}}(V_1, V_2, \mathscr{C}) - d_{G^{(3)}}(V_1', V_2', \mathscr{C}')| \\ &\leq 2\varepsilon \leq \varepsilon/\alpha. \end{aligned}$$

For (ii), note that \mathbf{G}' is $(\varepsilon/(1-\alpha), d/2)$ -regular and thus $(2\varepsilon, d/2)$ -regular by (i). Let $G'_c := G_c[V'_1, V'_2]$ for each $c \in \mathscr{C}$. For each i = 1, 2 and $v \in V'_i$, we also have $\sum_{c \in \mathscr{C}'} d_{G'_c}(v) \ge d|V_{3-i}||\mathscr{C}| - 2\alpha|V_{3-i}||\mathscr{C}| \ge d|V'_{3-i}||\mathscr{C}'|/2$. Combining this with $e(G'_c) \ge d|V_1||V_2| - 2\alpha|V_1||V_2| \ge d|V_1||V_2|/2$ for each $c \in \mathscr{C}'$, we get that \mathbf{G}' is $(2\varepsilon, d/2)$ -superregular.

For (iii), similarly, we always have that \mathbf{G}' is $(\varepsilon/\alpha, d/2)$ -regular by (i). Let $G'_c := G_c[V'_1, V'_2]$ for all $c \in \mathscr{C}$. We only need to show that with high probability, for each i = 1, 2 and $v \in V'_i$, we have $\sum_{c \in \mathscr{C}'} d_{G'_c}(v) \ge d^2 |V'_{3-i}| |\mathscr{C}'| / 16$ and for each $c \in \mathscr{C}'$, we have $e(G'_c) \ge d^2 |V_1| |V_2| / 16$. Suppose that \mathscr{C}' has been chosen. Since \mathbf{G} is (ε, d) -superregular, for each $c \in \mathscr{C}'$, we have $e(G_c) \ge d|V_1| |V_2|$. Let $V_1^c := \{v \in V_1 : d_{G_c}(v) \ge d|V_2|/2\}$. Then we get $d|V_1| |V_2| \le |V_1^c| |V_2| + (|V_1| - |V_1^c|) d|V_2| / 2$ and thus $|V_1^c| \ge d|V_1| / (2 - d) \ge d|V_1| / 2$. A Chernoff bound implies that, with probability $1 - e^{-\Omega(n)}$, we have $|V'_1 \cap V_1^c| \ge dn_1 / 4$ and $|V'_2 \cap N_{G_c}(v)| \ge dn_2 / 4$ for each $v \in V_1^c$. Therefore, by a union bound, with high probability, we have $e(G'_c) \ge d^2 n_1 n_2 / 16$ for each $c \in \mathscr{C}'$. Similarly, with high probability, for each i = 1, 2 and $v \in V'_i$, we have $\sum_{c \in \mathscr{C}'} d_{G'_c}(v) \ge d^2 n_{3-i} h / 16$. It follows that \mathbf{G}' is $(\varepsilon/\alpha, d^2/16)$ -superregular with high probability. \Box

2.2 The regularity lemma for graph collections

We use the following version of the regularity lemma for graph collections, which is obtained by applying the degree version of the weak regularity lemma (Lemma A.2) to the 3-graph $G^{(3)}$ of G and cleaning up the clusters so that vertex clusters and colour clusters are separate. We postpone the derivation to the appendix.

Lemma 2.5 (Regularity lemma for graph collections). For all integers $L_0 \geq 1$ and every $\varepsilon, \delta > 0$, there is an $n_0 = n_0(\varepsilon, \delta, L_0)$ such that for every $d \in [0, 1)$ and every graph collection $\mathbf{G} = (G_c : c \in \mathscr{C})$ on vertex set V of size $n \geq n_0$ with $\delta n \leq |\mathscr{C}| \leq n/\delta$, there exists a partition of V into V_0, V_1, \ldots, V_L , of \mathscr{C} into $\mathscr{C}_0, \mathscr{C}_1, \ldots, \mathscr{C}_M$ and a spanning subgraph G'_c of G_c for each $c \in \mathscr{C}$ such that the following properties hold:

- (i) $L_0 \leq L, M \leq n_0 \text{ and } |V_0| + |\mathscr{C}_0| \leq \varepsilon n;$
- (*ii*) $|V_1| = \ldots = |V_L| = |\mathscr{C}_1| = \ldots = |\mathscr{C}_M| =: m;$
- $\begin{array}{l} (iii) \sum_{c \in \mathscr{C}} d_{G'_c}(v) > \sum_{c \in \mathscr{C}} d_{G_c}(v) (3d/\delta^2 + \varepsilon)n^2 \ for \ all \ v \in V \ and \ e(G'_c) > e(G_c) (3d/\delta^2 + \varepsilon)n^2 \\ for \ all \ c \in \mathscr{C}; \end{array}$
- (iv) if, for $c \in \mathcal{C}$, the graph G'_c has an edge with both vertices in a single cluster V_i for some $i \in [L]$, then $c \in \mathcal{C}_0$;

(v) for all triples $(\{h, i\}, j) \in {\binom{[L]}{2}} \times [M]$, we have that either $G'_c[V_h, V_i] = \emptyset$ for all $c \in \mathscr{C}_j$, or $G'_{hi,j} := (G'_c[V_h, V_i] : c \in \mathscr{C}_j)$ is (ε, d) -regular.

The sets V_i are called *vertex clusters* and the sets \mathscr{C}_j are called *colour clusters*, while V_0 and \mathscr{C}_0 are the exceptional vertex and colours sets respectively.

Definition 2.6 (Reduced graph collection). Given a graph collection $\boldsymbol{G} = (G_c : c \in \mathscr{C})$ on V and parameters $\varepsilon > 0, d \in [0, 1)$ and $L_0 \ge 1$, the reduced graph collection $\boldsymbol{R} = \boldsymbol{R}(\varepsilon, d, L_0)$, reduced 3-graph $R^{(3)} = R(\varepsilon, d, L_0)$ and the reduced edge-coloured graph $\mathcal{R} = \mathcal{R}(\varepsilon, d, L_0)$ of \boldsymbol{G} are defined as follows. Apply Lemma 2.5 to \boldsymbol{G} with parameters ε, d, L_0 to obtain \boldsymbol{G}' and a partition V_0, \ldots, V_L of V and $\mathscr{C}_0, \ldots, \mathscr{C}_M$ of \mathscr{C} where V_0, \mathscr{C}_0 are the exceptional sets and V_1, \ldots, V_L are the vertex clusters and $\mathscr{C}_1, \ldots, \mathscr{C}_M$ are the colour clusters. Then $\boldsymbol{R} = (R_1, \ldots, R_M)$ is a graph collection of M graphs each on the same vertex set [L], where, for $(\{h, i\}, j) \in {[L] \choose 2} \times [M]$, we have $hi \in R_j$ whenever $\boldsymbol{G}'_{hi,j}$ is (ε, d) -regular.

Also, $R^{(3)}$ is the 3-graph of **R** and \mathcal{R} is the reduced edge-coloured graph.

The next lemma (related to Lemma 5.5 in [32]) states that clusters inherit a minimum degree bound in the reduced graph from G.

Lemma 2.7 (Degree inheritance). Suppose p > 0, $L_0 \ge 1$ and $0 < 1/n \ll \varepsilon \le d \ll \delta, \gamma, p \le 1$. Let $\mathbf{G} = (G_c : c \in \mathscr{C})$ be a graph collection on a vertex set V of size n with $\delta(G_c) \ge (p + \gamma)n$ for all $c \in \mathscr{C}$ and $\delta n \le |\mathscr{C}| \le n/\delta$. Let $\mathbf{R} = \mathbf{R}(\varepsilon, d, L_0)$ be the reduced graph collection of \mathbf{G} on L vertices with M graphs. Then

(i) for every $i \in [L]$ there are at least $(1 - d^{1/4})M$ colours $j \in [M]$ for which $d_{R_j}(i) \ge (p + \gamma/2)L$; (ii) for every $j \in [M]$ there are at least $(1 - d^{1/4})L$ vertices $i \in [L]$ for which $d_{R_j}(i) \ge (p + \gamma/2)L$.

Proof. To prove (i), note that for all $v \in V \setminus V_0$ we have

$$\sum_{c \in \mathscr{C}} d_{G'_c - V_0}(v) \ge \sum_{c \in \mathscr{C}} d_{G_c}(v) - (3d/\delta^2 + \varepsilon)n^2 - \varepsilon |\mathscr{C}|s \ge \sum_{c \in \mathscr{C}} d_{G_c}(v) - 4dn^2/\delta^2$$

Let \mathscr{D}_v be the collection of colours c in $\mathscr{C} \setminus \mathscr{C}_0$ for which $d_{G'_c-V_0}(v) \geq d_{G_c}(v) - \sqrt{dn}$. Then

$$\sum_{c \in \mathscr{C}} d_{G'_c - V_0}(v) \le \sum_{c \in \mathscr{C}} d_{G_c}(v) - |\mathscr{C} \setminus \mathscr{D}_v| \sqrt{dn}$$

and therefore $|\mathscr{C} \setminus \mathscr{D}_v| \leq 4dn^2/(\delta^2 \sqrt{dn}) \leq d^{1/3}n/2$ by $d \ll \delta$, so $|\mathscr{D}_v| \geq |\mathscr{C}| - d^{1/3}n/2 - \varepsilon n \geq |\mathscr{C}| - d^{1/3}n$. We have $mM \leq |\mathscr{C}| \leq mM + \varepsilon n$ and $mL \leq n \leq mL + \varepsilon n$. Thus the number of clusters \mathscr{C}_j containing at least one colour of \mathscr{D}_v is at least

$$|\mathscr{D}_v|/m \ge M - d^{1/3}n/m \ge M - d^{1/3}L/(1-\varepsilon) \ge M - 2d^{1/3}M/\delta \ge (1-d^{1/4})M.$$

Now let $i \in [L]$ and $v \in V_i$. For each cluster \mathscr{C}_j as above, choose an arbitrary colour $c_j \in \mathscr{C}_j \cap \mathscr{D}_v$. Then the number of clusters V_h containing some $u \in N_{G'_{c_i}}(v)$ is at least

$$\frac{d_{G_{c_j}}(v) - \sqrt{dn}}{m} \ge \frac{(p + \gamma - \sqrt{d})n}{m} \ge (p + \gamma/2)L.$$

But then Lemma 2.5(v) implies that *i* is adjacent to each such V_h in R_j . So for every $i \in [L]$, $d_{R_i}(i) \ge (p + \gamma/2)L$ for at least $(1 - d^{1/4})M$ colours *j*. The proof of (ii) is similar and we omit it. \Box

2.3 Templates

We define the notion of a 'template', which is essentially a reduced graph in the transversal setting. We will use these as templates for embedding, in the same way that reduced graphs are used for embedding into a single graph.

Definition 2.8 (Templates). Let $0 < 1/m \le \varepsilon \le d, \delta \le 1$ be parameters and let $r \in \mathbb{N}$. Suppose that

- R is an r-vertex graph, with vertex set [r] unless otherwise specified,
- $\mathcal{V} = \{V_1, \ldots, V_r\}$ is a set of r disjoint vertex sets with $m \leq |V_j| \leq m/\delta$ for all $j \in [r]$, whose union we denote by V,
- $\mathscr{C} = \bigcup_{e \in E(R)} \mathscr{C}_e$ is a colour set where $|\mathscr{C}_e| \ge \delta m$ for all $e \in E(R)$,
- G is a graph collection with colour set \mathscr{C} where for each $c \in \mathscr{C}$, the graph G_c is the union of bipartite graphs G_c^e where G_c^e has parts V_i, V_j , over all e = ij for which $c \in \mathscr{C}_e$. For each $e \in E(R)$, let $G^e = (G_c^e : c \in \mathscr{C}_e)$.

We say that $\mathcal{F} = (\mathcal{V}, \mathscr{C}, \mathbf{G})$ is an *R*-template with parameters $(m, \varepsilon, d, \delta)$ if for every $e \in E(R)$, \mathbf{G}^e is (ε, d) -regular. If we replace regular with semi-superregular, it is a semi-super *R*-template; if we replace regular with half-superregular, it is a super *R*-template; if we replace regular with half-superregular, it is a half-super *R*-template. If $\mathscr{C} = \bigcup_{e \in E(R)} \mathscr{C}_e$ is a partition, we say that the template is rainbow.

A transversal embedding of a graph H inside \mathcal{F} is a copy of H with vertices in V such that for every edge e there is a distinct $c \in \mathscr{C}$ such that $e \in G_c$. That is, there exist injections $\tau : V(H) \to V$ and $\sigma : E(H) \to \mathscr{C}$ where $\tau(x)\tau(y) \in G_{\sigma(xy)}$ for all $xy \in E(H)$.

Note that the partition $\mathscr{C} = \bigcup_{e \in E(R)} \mathscr{C}_e$ is suppressed in the notation. Given a template \mathcal{F} , we explicitly use the notation in the definition unless otherwise specified. Observe that for an *R*-template $(\mathcal{V}, \mathscr{C}, \mathbf{G})$ with parameters $(m, \varepsilon, d, \delta)$ and v(R) = r,

$$rm \le |V| \le rm/\delta.$$

Given parameters $\beta, \varepsilon', d', \delta' > 0$ with $\beta \leq 1, \varepsilon' \geq \varepsilon, d' \leq d$ and $\delta' \leq \beta\delta$, any $(m, \varepsilon, d, \delta)$ template is also a $(\beta m, \varepsilon', d', \delta')$ template. We will often take *subtemplates* of templates, meaning that the new vertex clusters, colour clusters and graphs G_c are subsets/subgraphs of the originals. Some convenient notation for this is as follows: if $\mathcal{F} = (\mathcal{V}, \mathscr{C}, \mathbf{G})$ is a template, $\mathscr{C}' \subseteq \mathscr{C}$ and $\mathcal{V}' = \{V'_1, \ldots, V'_r\}$ where $V'_j \subseteq V_j$ for all $j \in [r]$, we say that $\mathcal{F}' = (\mathcal{V}', \mathscr{C}', \mathbf{G}')$ is the *subtemplate of* \mathcal{F} *induced by* $\mathcal{V}', \mathscr{C}'$ when each $\mathscr{C}'_e = \mathscr{C}_e \cap \mathscr{C}'$, and $(G')^e_c := G_c[V'_i, V'_j]$ is defined for each $c \in \mathscr{C}'_e$. We also say that \mathcal{F}' is obtained by deleting $\mathscr{C} \setminus \mathscr{C}'$ and $V \setminus V'$.

The following straightforward lemma shows that removing a small fraction of colours and vertices from a template produces a subtemplate with slightly weaker parameters, which remains super if the original template was super.

Lemma 2.9 (Template slicing). Let $0 < 1/m \ll \varepsilon \ll \varepsilon' \ll \alpha \ll d, \delta, 1/r, 1/k \leq 1$ where $r \geq 2$ is an integer. Let R be an r-vertex graph and let $\mathcal{F} = (\mathcal{V}, \mathcal{C}, \mathbf{G})$ be an R-template with parameters $(m, \varepsilon, d, \delta)$. Let $\mathcal{C}' \subseteq \mathcal{C}, V'_i \subseteq V_i$ for all $i \in [r]$ and let $\mathcal{F}' = (\mathcal{V}', \mathcal{C}', \mathbf{G}')$ be the subtemplate of \mathcal{F} induced by $\mathcal{V}', \mathcal{C}'$.

- (i) If every $|\mathscr{C}'_e| \geq \alpha |\mathscr{C}_e|/k$ and $\alpha |V_i| \leq |V'_i| \leq k\alpha |V_i|$, then \mathcal{F}' is a template with parameters $(\alpha m, \varepsilon/\alpha, d/2, \delta/k)$.
- (ii) If every $|\mathscr{C}_e \setminus \mathscr{C}'_e| \leq \alpha m$ and $|V_i \setminus V'_i| \leq \alpha m$, then \mathcal{F}' is a template with parameters $(m/2, 2\varepsilon, d/2, \delta/2)$. Moreover, if \mathcal{F} is super, then \mathcal{F}' is super.

- (iii) Given $\alpha |V_i| \leq n_i \leq k\alpha |V_i|$ for all $i \in [r]$ and $h_e \geq \alpha |\mathcal{C}_e|/k$ for all $e \in E(R)$, if V'_i is a uniform random subset of V_i of size n_i and \mathcal{C}'_e is a uniform random subset of \mathcal{C}_e of size h_e and \mathcal{F} is super, then with high probability, \mathcal{F}' is a super template with parameters $(\alpha m, \varepsilon / \alpha, d^2/16, \delta/k)$.
- (iv) Suppose \mathcal{F} is half-super. For all $c \in \mathscr{C}$, there is $G'_c \subseteq G_c$ such that, defining $\mathbf{G}' := (G'_c : c \in \mathscr{C})$, the template $(\mathcal{V}, \mathscr{C}, \mathbf{G}')$ is super with parameters $(m, \varepsilon', d^2/2, \delta)$.

Proof. For (i), we have $\alpha m \leq |V'_i| \leq \alpha m/(\delta/k)$ and $|\mathscr{C}'_e| \geq \alpha \delta m/k$. Also, by Lemma 2.4(i), for each $e \in E(R)$, $(\mathbf{G}')^e$ is $(\varepsilon/\alpha, d/2)$ -regular. Thus \mathcal{F}' is a template with parameters $(\alpha m, \varepsilon/\alpha, d/2, \delta/k)$.

For (ii), showing that \mathcal{F}' is a template with the given parameters is similar to (i). Note that if \mathcal{F} is super, then $(\mathbf{G}')^e$ is $(2\varepsilon, d/2)$ -superregular for each $e \in E(R)$ by Lemma 2.4(ii) and thus \mathcal{F}' is a super *R*-template with parameters $(m/2, 2\varepsilon, d/2, \delta/2)$.

For (iii), if \mathcal{F} is super, then \mathbf{G}^e is (ε, d) -superregular for each $e \in E(R)$. Thus by Lemma 2.4(iii), with high probability, $(\mathbf{G}')^e$ is $(\varepsilon/\alpha, d^2/16)$ -superregular for each $e \in E(R)$. Therefore, by (i), \mathcal{F}' is a super template with parameters $(\alpha m, \varepsilon/\alpha, d^2/16, \delta/k)$ with high probability.

Part (iv) follows immediately from Lemma 2.1.

Edges which lie in many graphs G_c are particularly useful for embedding and thus we define the (simple, uncoloured) *thick graph* consisting of all such edges.

Definition 2.10 (Thick graph of a template). Given $\lambda > 0$ and an *R*-template $\mathcal{F} = (\mathcal{V}, \mathscr{C}, \mathbf{G})$, let $T_{\mathcal{F}}^{\lambda}$ be the simple 2-graph with vertex set *V* such that $xy \in E(T_{\mathcal{F}}^{\lambda})$ whenever $xy \in G_c$ for at least $\lambda |\mathscr{C}_{ij}|$ colours $c \in \mathscr{C}_{ij}$ where $x \in V_i, y \in V_j$ and $ij \in E(R)$. We call $T_{\mathcal{F}}^{\lambda}$ the λ -thick graph of \mathcal{F} .

The following proposition states that the thick graph of a bipartite semi-super template has a spanning half-superregular subgraph. This is very useful for embedding as one can use the usual blow-up lemma to embed a bounded degree graph into the thick graph, and then greedily assign colours.

Proposition 2.11. Let $0 < 1/m \ll \varepsilon \ll \lambda \ll d, \delta, 1/r \leq 1$ where $r \geq 2$ is an integer. Let R be a graph on r vertices and let $\mathcal{F} = (\mathcal{V}, \mathscr{C}, \mathbf{G})$ be a semi-super R-template with parameters $(m, \varepsilon, d, \delta)$. Then for all $ij \in E(R)$, $T^{\lambda}_{\mathcal{F}}[V_i, V_j]$ is $(\varepsilon, d/2)$ -half-superregular.

Proof. Let $ij \in E(R)$ and let $U_i \subseteq V_i$ and $U_j \subseteq V_j$ be any two subsets such that $|U_i| \ge \varepsilon |V_i|$ and $|U_j| \ge \varepsilon |V_j|$. Let $\ell_{ij} := e(T^{\lambda}_{\mathcal{F}}[U_i, U_j])$. By regularity we have $|d_{G^{(3)}}(U_i, U_j, \mathscr{C}_{ij}) - d_{G^{(3)}}(V_i, V_j, \mathscr{C}_{ij})| \le \varepsilon$ and thus we get

$$(d-\varepsilon)|U_i||\mathcal{U}_j||\mathscr{C}_{ij}| \leq e_{G^{(3)}}(U_i,U_j,\mathscr{C}_{ij}) \leq \ell_{ij}|\mathscr{C}_{ij}| + \lambda(|U_i||U_j| - \ell_{ij})|\mathscr{C}_{ij}|.$$

Therefore, we have $\ell_{ij}/(|U_i||U_j|) \ge (d - \varepsilon - \lambda)/(1 - \lambda) \ge d/2$. Similarly, for any $x \in V_i$, let $\ell_x := d_{T^{\lambda}_{\mathcal{F}}}(x, V_j)$. By semi-superregularity, we have $\sum_{c \in \mathscr{C}_{ij}} d_{G_c}(x) \ge d|V_j||\mathscr{C}_{ij}|$. Then we get

$$d|V_j||\mathscr{C}_{ij}| \le \ell_x|\mathscr{C}_{ij}| + \lambda(|V_j| - \ell_x)|\mathscr{C}_{ij}|$$

and thus $\ell_x/|V_j| \ge (d-\lambda)/(1-\lambda) \ge d/2$. The half-superregularity then follows.

3 Embedding lemmas

In this section we state a series of embedding lemmas which we will combine to prove our transversal blow-up lemma, stated at the end of the section. These lemmas are also useful in their own right in applications.

Each lemma is of the following type: we are given an R-template and a bounded degree graph H whose partition matches the template. For a small number of vertices y in H we are given large target sets T_y where y must be embedded. The output of the lemma is a transversal embedding of H, which consists of an embedding τ of vertices and an embedding σ of colours, so $\tau(x)\tau(y) \in G_{\sigma(xy)}$ for all $xy \in E(H)$, and such that $\tau(y) \in T_y$ whenever there is a target set.

The first such lemma applies to embed a small graph H into a template. In fact we only embed some of H while finding large candidate sets for the rest of H. This means that we set as de some colours so that, for each unembedded vertex y and each of its embedded neighbours x, there is a distinct colour $\sigma(xy)$, and a large vertex set so that if the image of y is chosen in this set, then we can extend our embedding to a transversal embedding that uses the specified colours. That is, for every z in this set we have $\tau(x)z \in G_{\sigma(xy)}$. The proof embeds vertices one by one, at each step fixing the colours that will be used to future neighbours, and is similar to the 'partial embedding lemma', an uncoloured version, in [7].

Lemma 3.1 (Embedding lemma with target and candidate sets). Let $0 < 1/m \ll \varepsilon, \gamma \ll \nu' \ll$ $\nu, d, \delta, 1/\Delta, 1/r \leq 1$ where $r \geq 2$ is an integer.

- Let R be a graph on vertex set [r] such that $\mathcal{F} = (\mathcal{V}, \mathcal{C}, \mathbf{G})$ is an R-template with parameters $(m,\varepsilon,d,\delta).$
- Let H be a graph with $\Delta(H) \leq \Delta$ for which there is a graph homomorphism $\phi: V(H) \to V(R)$ such that $|\phi^{-1}(j)| \leq \gamma m$ for all $j \in [r]$, and suppose there is a partition $V(H) = X \cup Y$ where $E(H[Y]) = \emptyset.$
- For every $w \in V(H)$, suppose there is a set $T_w \subseteq V_{\phi(w)}$ with $|T_w| \ge \nu m$.

Then there are injective maps $\tau: X \to V$ and $\sigma: E(H) \to \mathscr{C}$ such that

- (i) $\tau(x) \in T_x$ for all $x \in X$;
- (ii) $\sigma(xy) \in \mathscr{C}_{\phi(x)\phi(y)}$ for all $xy \in E(H)$, and if $xx' \in E(H[X])$, then $\tau(x)\tau(x') \in E(G_{\sigma(xx')})$; (iii) for all $y \in Y$ there exists $C_y \subseteq V_{\phi(y)} \setminus \tau(X)$ such that $C_y \subseteq \bigcap_{x \in N_H(y) \cap X} N_{G_{\sigma(xy)}}(\tau(x)) \cap T_y$ and $|C_u| \geq \nu' m.$

Proof. Fix an ordering of V(H) where all vertices of X come before any vertex of Y, and for each $x \in X$, let $N^{<}(x)$ be the set of neighbours of x in H which appear before x in the ordering, and $N^{>}(x)$ the set of those which appear after. At step (x) (with $x \in X$), we will choose $\tau(x)$ and $\sigma(xy)$ for all $y \in N^{>}(x)$. Initially, define candidate sets $C_w := T_w$ for all $w \in V(H)$ and $C_{xy} := \mathscr{C}_{\phi(x)\phi(y)}$ for all $xy \in E(H)$. The following comprises step (x), which we perform for each $x \in X$ in order.

- (x,1) For all $y \in N^{>}(x)$, delete all vertices $v \in C_x$ with $\sum_{c \in C_{xy}} |N_{G_c}(v) \cap C_y| < (d-\varepsilon)|C_{xy}||C_y|$.
- (x,2) Choose $\tau(x) \in C_x$.
- (x,3) For all $u \in V(H)$, delete $\tau(x)$ from C_u .
- (x,4) For each $y \in N^{>}(x)$ in order:
- (x,y,4.1) delete all colours $c \in C_{xy}$ with $|N_{G_c}(\tau(x)) \cap C_y| < d|C_y|/2$,
- (x,y,4.2) choose $\sigma(xy) \in C_{xy}$,
- (x,y,4.3) for all $uv \in E(H)$, delete $\sigma(xy)$ from C_{uv} ,
- (x,y,4.4) delete all vertices $v \in C_y$ with $v \notin N_{G_{\sigma(xy)}}(\tau(x))$.

We claim that, at the end of the process, $|C_x|, |C_{xy}| \ge \nu' m$ for all $x \in V(H)$ and $xy \in E(H)$. First, let us see why the claim implies the lemma. Since candidate sets are never empty and every edge in H is incident to $X, \tau: X \to V$ and $\sigma: E(H) \to \mathscr{C}$ are defined, and by step (3) and (4.3), they are both injections. For (i), we choose $\tau(x)$ from C_x which is always a (non-empty) subset of T_x . For (ii), we choose $\sigma(xy)$ from C_{xy} which is always a (non-empty) subset of $\mathscr{C}_{\phi(x)\phi(y)}$. If $xx' \in E(H)$ where x' appears after x in the ordering, then we choose $\tau(x')$ from $C_{x'}$ and by step (x,x',4.4) we have $\tau(x') \in N_{G_{\sigma(xx')}}(\tau(x))$. For (iii), given the claim and the fact that Y comes after X in the ordering, it suffices to show that, for all $y \in Y$ and $x \in N^{<}(y)$, we have $C_y \subseteq N_{G_{\sigma(xy)}}(\tau(x))$. Noting that $x \in X$, this is a consequence of step (x,y,4.4).

It remains to prove the claim. We suppose that it is true up until step (x). The vertex candidate set C_x can only shrink at steps (x,1), (3) and at step (u,x,4.4) for $u \in N^{<}(x)$. Proposition 2.3 and the fact that, currently, $|C_x| \geq \nu' m > \varepsilon |V_{\phi(x)}|$, imply that at step (x,1), at most $\Delta \varepsilon |C_x|$ vertices are deleted. At step (3), at most |X| vertices are deleted. At step (u,x,4.4), the colour $\sigma(ux)$ was chosen from C_{ux} which has $|N_{G_{\sigma(ux)}}(\tau(u)) \cap C_x| \geq d|C_x|/2$ due to step (u,x,4.1). Thus C_x shrinks by a factor of at most d/2 at this step. Therefore

$$|C_x| \ge ((d/2)^{\Delta} - \Delta\varepsilon)|T_x| - |X| \ge ((d/2)^{\Delta} - \Delta\varepsilon)\nu - r\gamma)m \ge \nu(d/3)^{\Delta}m > \nu'm.$$

The colour candidate set C_{xy} , with x before y in the ordering, can only shrink at steps (x,y,4.1)and (4.3). At most e(H) colours are lost at step (4.3). At step (x,y,4.1), we chose $\tau(x)$ from C_x which, immediately after (x,1), satisfies $\sum_{c \in C_{xy}} |N_{G_c}(\tau(x)) \cap C_y| \ge (d-\varepsilon)|C_{xy}||C_y|$. Writing $A \subseteq C_{xy}$ for the subset of colours which are not deleted at step (x,y,4.1), we have $(d-\varepsilon)|C_{xy}||C_y| \le |A||C_y| + (d/2)(|C_{xy}| - |A|)|C_y|$ and thus $|A| \ge (d/2 - \varepsilon)|C_{xy}|$. Therefore

$$|C_{xy}| \ge (d/2 - \varepsilon)|\mathscr{C}_{\phi(x)\phi(y)}| - e(H) \ge (d/2 - \varepsilon)\delta m - \Delta r\gamma m \ge \delta dm/3 \ge \nu' m.$$

This completes the proof of the claim, and hence of the lemma.

The next ingredient is a version of the above lemma where we are embedding a small graph H such that a small fraction of its vertices have target sets, but additionally we now have a very small set of colours which must be used in the embedding (together with any other colours). These prescribed colours will be used on an induced matching M in H. This matching, together with its neighbours, will be embedded greedily and then Lemma 3.1 will apply to extend the embedding to the whole of H.

Lemma 3.2 (Embedding lemma with targets and prescribed colours). Let $0 < 1/m \ll \varepsilon \ll \alpha, \gamma \ll \lambda_1, \lambda_2 \ll \nu, d, \delta, 1/\Delta, 1/r \leq 1$ where $r \geq 2$ is an integer.

- Let R be a graph on vertex set [r] and let $(\mathcal{V}, \mathscr{C}, \mathbf{G})$ be an R-template with parameters $(m, \varepsilon, d, \delta)$.
- Let H be a graph with $\Delta(H) \leq \Delta$ for which there is a graph homomorphism $\phi: V(H) \to V(R)$ such that $|\phi^{-1}(j)| \leq \lambda_1 m$ for all $j \in [r]$ and $e(H[\phi^{-1}(i), \phi^{-1}(j)]) \geq \lambda_2 m$ for all $ij \in E(R)$.
- Suppose there is a set $W \subseteq V(H)$ with $|W| \leq \alpha m$, such that for all $w \in W$ there is a set $T_w \subseteq V_{\phi(w)}$ with $|T_w| \geq \nu m$.
- For each $e \in E(R)$, let $\mathscr{D}_e \subseteq \mathscr{C}_e$ be a set of at most γm colours and let $\mathscr{D} = \bigcup_{e \in E(R)} \mathscr{D}_e$, and suppose that $e(G_c[V_i, V_j]) \ge d|V_i||V_j|$ for all $c \in \mathscr{D}_{ij}$ and $ij \in E(R)$, and $|\mathscr{C}_e \setminus \mathscr{D}| \ge d|\mathscr{C}_e|$ for all $e \in E(R)$.

Then there are injective maps $\tau: V(H) \to V$ and $\sigma: E(H) \to \mathscr{C}$ such that

- (i) $\tau(x) \in V_{\phi(x)}$ for all $x \in V(H)$;
- (ii) $\tau(w) \in T_w$ for all $w \in W$;
- (iii) for all $xx' \in E(H)$ we have $\tau(x)\tau(x') \in E(G_{\sigma(xx')})$;
- (iv) $\mathscr{D} \subseteq \sigma(E(H)).$

Proof. Let e_1, \ldots, e_s be an arbitrary ordering of E(R). Let $\mathscr{D}'_{e_i} := \mathscr{D}_{e_i} \setminus (\mathscr{D}_{e_1} \cup \ldots \cup \mathscr{D}_{e_{i-1}})$ for all $i \in [s]$, so that the \mathscr{D}'_e over $e \in E(R)$ are pairwise disjoint, and their union equals \mathscr{D} . We claim that

for each $e_i = jj' \in E(R)$, we can choose a matching $M_{jj'} \subseteq H[\phi^{-1}(j), \phi^{-1}(j')]$ such that, writing $M := \bigcup_{e \in E(B)} M_e$, we have

- $V(M) \cap W = \emptyset$ and M is an induced matching in H such that for every $y \in V(H) \setminus V(M)$, there is $xx' \in E(M)$ such that $N_H(y, V(M)) \subseteq \{x, x'\}$;
- $|M_e| = |\mathscr{D}'_e|$ for each $e \in E(R)$.

We can do this greedily, as follows. Suppose we have found $M_{e_1}, \ldots, M_{e_{i-1}}$ with the required properties, for some $i \ge 1$. We show how to find M_{e_i} , where we write $e_i := jj'$. Vizing's theorem states that every graph J with maximum degree Δ can be properly edge-coloured with at most $\Delta + 1$ colours. Thus J contains a matching of size $[e(J)/(\Delta+1)]$. Thus $H[V_i, V_{i'}]$ contains a matching of size $[\lambda_2 m/(\Delta+1)]$. Now, from this matching, delete W and any vertex at distance at most two to any vertex in a previously found matching. There are at most r-1 neighbours h of j in R for which we have defined M_{jh} , and for each one we delete at most $(1 + \Delta + \Delta^2)v(M_{jh}) \leq 4\Delta^2\gamma m$ such vertices. The total number of deleted vertices is at most $|W| + 8r\Delta^2 \gamma m$, which leaves a matching M' of size at least $\lambda_2 m/(4\Delta)$. Now we will greedily choose a suitable submatching. Suppose we have chosen a subset $E(M^q) := \{x_1y_1, \ldots, x_{q-1}y_{q-1}\} \subseteq E(M')$ where $1 \leq q \leq |\mathscr{D}'_{jj'}|$ and the distance in $H \setminus M^q$ between any pair of vertices in $V(M^q)$ is at least three. To choose x_qy_q , we delete all vertices in V(M') at distance in $H \setminus M^q$ at most two. The number of deleted vertices is at most $(1 + \Delta + \Delta^2) 2q \leq 4\Delta^2 \gamma m < \lambda_2 m/(4\Delta)$. Thus we can always choose $x_q y_q$ from the remaining vertices of M'. We claim that $M^{q+1} := M^q \cup \{x_q y_q\}$ is a suitable extension of M^q . Indeed, x_q, y_q have no neighbours in M^q so M^{q+1} is induced, and if x_q shared a neighbour z with some other $y \in M^q$, then it would be at distance two from y, a contradiction. Thus we can obtain M_{e_i} and hence M_{e_1}, \ldots, M_{e_s} satisfying both required properties.

Let X := V(M) and $Y := N_H(X) \setminus X$. We will embed M into the template first, defining injections $\tau: X \to V$ and $\sigma: E(M) \cup E(H[X,Y]) \to \mathscr{D}$ so that $\sigma(M_e) = \mathscr{D}'_e$ for all $e \in E(R)$. We want to choose good images for the vertices $x \in X$ so that there are many choices for each neighbour $y \in N_H(x) \setminus X.$

Now we will embed M greedily, using the fact that $e(G_c) \ge d|V_j||V_{j'}|$ for all $c \in \mathscr{D}_{jj'}$ and $\gamma \ll d$. For each $p \le e(M)$, let $i \le s$ be such that $e(M_{e_1}) + ... + e(M_{e_{i-1}}) \le p < e(M_{e_1}) + ... + e(M_{e_i})$. Let $M_p := M_{e_1} \cup \ldots \cup M_{e_{i-1}} \cup M'_{e_i}$ where $e(M_p) = p$ and we have fixed an ordering of $E(M_{e_i})$ and M'_{e_i} is the (possibly empty) matching which consists of an initial segment. Let $X_p := V(M_p)$ and $Y_p := N_H(X_p) \setminus X_p$. Suppose we have found injections $\tau : X_p \to V$ and $\sigma : E(\widetilde{M}_p) \cup E(H[X_p, Y_p]) \to \mathscr{C}$ such that

- $\tau(x) \in V_{\phi(x)}$ for all $x \in X_p$;

- $\tau(x)\tau(y) \in G_{\sigma(xy)}$ for all $xy \in E(\widetilde{M}_p)$; $\sigma(E(M_{e_h})) = \mathscr{D}'_{e_h}$ for all h < i and $\sigma(E(M'_{e_i})) \subseteq \mathscr{D}'_{e_i}$; for all $y \in Y_p$ there exists $C_y \subseteq V_{\phi(y)}$ such that $C_y \subseteq \bigcap_{x \in N_H(y) \cap X_p} N_{G_{\sigma(xy)}}(\tau(x))$, and $|C_y| \ge C_{\phi(y)}$ $d|V_{\phi(y)}|/6.$

We want to extend τ and σ by embedding an edge xx' in $M_{e_i} \setminus M'_{e_i}$ with a colour $c^* \in \mathscr{D}'_{e_i}$ and choosing colours and candidate sets for its unembedded neighbours. So fix any such c^* and let $\sigma(xx') := c^*$. Let N(x), N(x') be the set of neighbours of x, x' respectively in $H \setminus M$. The first property of M implies that N(x), N(y), V(M) are pairwise disjoint for all $y \in V(M) \setminus \{x, x'\}$, and similarly for x'. However, we could have $N(x) \cap N(x') \neq \emptyset$ (for example if H is a union of vertex-disjoint triangles).

First, we will define $\tau(x)$ and colours and candidate sets for every vertex in N(x). Write $e_i = jj'$ and $U_h := V_h \setminus X_p$ for h = j, j'. We have $e(G_{c^*}[U_j, U_{j'}]) \ge d|V_j||V_{j'}| - 2\gamma m(|V_j| + |V_{j'}|) \ge d|V_j||V_{j'}|/2$.

$$Z(x) := \{ v \in U_j : d_{G_{c^*}}(v, U_{j'}) \ge d|U_{j'}|/4 \}.$$

A simple counting argument implies that $|Z(x)| \ge d|U_j|/4$. We will choose $\tau(x) \in Z(x)$ and candidate sets C_y for each $y \in N(x)$, as follows. Order N(x) as y_1, \ldots, y_ℓ , where $\ell \le \Delta$, and let a_1, \ldots, a_ℓ be such that $\phi(y_i) = a_i$, so $a_i \in [r]$. Suppose we have found, for $i \ge 0$,

- $c_i \in \mathscr{C}_{ja_i}$ such that c^*, c_1, \ldots, c_i are all distinct and disjoint from $\sigma(E(\widetilde{M}_p))$ and
- a set $Z_i(x) \subseteq Z(x)$ with $|Z_i(x)| \ge (d/6)^i |Z(x)|$ and $d_{G_{c_h}}(z, V_{a_h}) \ge d|V_{a_h}|/6$ for all $h \in [i]$ and $z \in Z_i(x)$.

Now, $\mathbf{G}^{ja_{i+1}}[Z_i(x), V_{a_{i+1}}]$ is $(\sqrt{\varepsilon}, d/2)$ -regular by Lemma 2.4(i) (the slicing lemma), so Lemma 2.3(ii) implies there are at least $(1 - \sqrt{\varepsilon})|\mathscr{C}_{ja_{i+1}}|$ colours $c \in \mathscr{C}_{ja_{i+1}}$ with $e(G_c[Z_i(x), V_{a_{i+1}}]) \geq (d/2 - \sqrt{\varepsilon})|Z_i(x)||V_{a_{i+1}}| \geq d|Z_i(x)||V_{a_{i+1}}|/3$. Let c_{i+1} be any such colour which does not lie in the set $\{c^*, c_1, \ldots, c_i\} \cup \sigma(E(\widetilde{M}_p))$, which exists since the number of available colours is at least $|\mathscr{C}_{ja_{i+1}} \setminus \mathscr{D}| - 1 - \Delta \geq d|\mathscr{C}_{ja_{i+1}}|/2 \geq d\delta m/2$. By a simple counting argument, the number of vertices $z \in Z_i(x)$ with $d_{G_{c_{i+1}}}(z, V_{a_{i+1}}) \geq d|V_{a_{i+1}}|/6$ is at least $d|Z_i(x)|/6$, and we let the set of these vertices be $Z_{i+1}(x)$. Thus we can complete the iteration and find $Z_\ell(x)$. Now let $\tau(x)$ be an arbitrary vertex of $Z_\ell(x)$, and for each $i \in [\ell]$, define the candidate set $C_{y_i} := N_{G_{c_i}}(\tau(x), V_{a_i})$ and $\sigma(xy_i) := c_i$.

Next, we will define $\tau(x')$ and colours and candidate sets for every vertex in N(x'). For each $y \in N(x) \cap N(x')$, the candidate set for y will be a subset of C_y . For this, let $Z(x') := N_{G_c^*}(\tau(x), U_{j'})$, so $|Z(x')| \ge d|U_{j'}|/4$ since $\tau(x) \in Z(x)$. Write $N(x') = \{w_1, \ldots, w_k\}$ and let b_1, \ldots, b_k be such that $\phi(w_i) = b_i$. For each $w_i \in N(x)$, we already defined C_{w_i} ; for the other w_i , let $C_{w_i} := V_{b_i}$. Using the fact that $\mathbf{G}^{j'b_i}[Z(x'), C_{w_i}]$ is $(\sqrt{\varepsilon}, d/2)$ -regular, exactly the same argument for N(x') as for N(x) implies that we can find

- for each $i \in [k], c'_i \in \mathscr{C}_{j'b_i}$ which are distinct and disjoint from $\{c^*, c_1, \ldots, c_\ell\} \cup \sigma(E(\widetilde{M}_p))$ and
- $Z_k(x') \subseteq Z(x')$ with $|Z_k(x')| \ge (d/6)^k |Z(x')|$ and $d_{G_{c'_i}}(z, C_{w_i}) \ge d|C_{w_i}|/6$ for all $i \in [k]$ and $z \in Z_k(x')$.

We let $\tau(x')$ be an arbitrary vertex of $Z_k(x')$ and for each $i \in [k]$, let $T_{w_i} := N_{G_{c'_i}}(\tau(x'), C_{w_i})$ and $\sigma(x'w_i) := c'_i$. Note that $\tau(x') \in Z(x') \subseteq N_{G_{c^*}}(\tau(x))$ so $\tau(x)\tau(x') \in G_{c^*} = G_{\sigma(xx')}$. For $y_i \in N(x) \setminus N(x')$, C_{y_i} has not changed since it was first defined, and we let $T_{y_i} := C_{y_i}$. We have $|T_y| \ge d^2 |V_{\phi(y)}|/36 \ge d^2 m/36$ for all $y \in N(x) \cup N(x')$.

Thus we can complete the iteration and embed the whole of M. This completes the required extension of τ and σ , so we have obtained $\tau : X \to V$ and $\sigma : E(M) \cup E(H[X,Y]) \to \mathscr{C}$, with $\sigma(E(M)) = \mathscr{D}$.

Finally, we extend the embedding to the whole of H by applying Lemma 3.1. Indeed, let \mathcal{F}' be the subtemplate of \mathcal{F} induced by $\mathcal{V}', \mathscr{C}'$, where $\mathcal{V}' = \{V'_1, \ldots, V'_r\}$ and $V'_i := V_i \setminus \tau(X)$, and $\mathscr{C}'_e := \mathscr{C}_e \setminus (\sigma(E(M) \cup E(H[X, Y])))$. Lemma 2.9(ii) implies that \mathcal{F}' is a template with parameters $(m/2, 2\varepsilon, d/2, \delta/2)$. For each $y \in Y$, we have $|T_y \cap V'_{\phi(y)}| \ge d^2|V_{\phi(y)}|/36 - |\phi^{-1}(\phi(y))| \ge d^2m/36 - \lambda_1 m \ge d^2m/37$. Similarly for each $w \in W$, we have $|T_w \cap V'_{\phi(w)}| \ge (\nu - \lambda_1)m \ge \nu m/2$. For all vertices y of H for which T_y is not defined, let $T_y := V_{\phi(y)}$. Thus we can apply Lemma 3.1 with parameters $m/2, 2\varepsilon, 2\lambda_1, \min\{\nu, d^2/37\}, d/2, \delta/2$ playing the roles of $m, \varepsilon, \gamma, \nu, d, \delta$, target sets T_y and $Y = \emptyset$ (so no candidate sets), to find the desired embedding. Most of the required properties are immediate but we justify why (iii) holds. If $xx' \in E(M)$ then we already showed that $\tau(x)\tau(x') \in G_{\sigma(xx')}$. If $xy \in E(H) \setminus E(M)$ where $x \in X$, we have $y \in Y$, and then the choice of M guaranteed that $N_H(y, X) \subseteq \{x, x'\}$ where $xx' \in E(M)$ and, if this neighbourhood equals $\{x\}$, we chose $\tau(y) \in T_y \subseteq N_{G_{\sigma(xy)}}(\tau(x))$, while if it equals $\{x, x'\}$ we chose $\tau(y) \in T_y$ and $T_y \subseteq N_{G_{\sigma(xy)}}(\tau(x)) \cap N_{G_{\sigma(x'y)}}(\tau(x'))$.

Let

If $xy \in E(H) - X$, then this follows from Lemma 3.1.

Next we state the usual blow-up lemma, which we will use in the proof of our transversal blow-up lemma. Note that the lemma is usually stated in terms of the stronger 'superregular' condition rather than 'half-superregular', but the same proof applies. Alternatively, one can apply Lemma 2.1.

Theorem 3.3 (Blow-up lemma [29]). Let $0 < 1/m \ll \varepsilon, \alpha \ll \nu, d, \delta, 1/\Delta, 1/r \le 1$.

- Let R be a graph on vertex set [r].
- Let G be a graph with vertex classes V_1, \ldots, V_r where $m \leq |V_i| \leq m/\delta$ for all $i \in [r]$, such that $G[V_i, V_j]$ is (ε, d) -half-superregular whenever $ij \in E(R)$.
- Let H be a graph with $\Delta(H) \leq \Delta$ for which there is a graph homomorphism $\phi: V(H) \to V(R)$ such that $|\phi^{-1}(j)| \leq |V_j|$ for all $j \in [r]$.
- For each $i \in [r]$, let $U_i \subseteq \phi^{-1}(i)$ with $|U_i| \leq \alpha m$ and suppose there is a set $T_x \subseteq V_i$ with $|T_x| \geq \nu m$ for all $x \in U_i$.

Then there is an embedding of H inside G such that for every $i \in [r]$, every $x \in U_i$ is embedded inside T_x .

The final embedding lemma that we prove in this section applies to embed a spanning graph H which consists of small components and such that a small fraction of its vertices have target sets. The template \mathcal{F} is required to be semi-super (every vertex has large total degree) and rainbow, since H could have many edges in every pair, and additionally \mathcal{F} must contain many more colours than required for a transversal embedding.

Lemma 3.4 (Embedding lemma with extra colours). Let $0 < 1/m \ll \varepsilon, \mu, \alpha \ll \beta \ll \nu, d, \delta, 1/\Delta, 1/r \le 1/2$ where r is an integer.

- Let R be an r-vertex graph and let $\mathcal{F} = (\mathcal{V}, \mathcal{C}, \mathbf{G})$ be a semi-super rainbow R-template with parameters $(m, \varepsilon, d, \delta)$.
- Let n := |V| and suppose that H is an n-vertex graph with $\Delta(H) \leq \Delta$ which is the union of vertex-disjoint components of size at most μn , and there is a graph homomorphism ϕ of H into R such that $|\phi^{-1}(j)| = |V_j|$ for all $j \in [r]$ and $e(H[\phi^{-1}(i), \phi^{-1}(j)]) \leq |\mathscr{C}_{ij}| \beta m$ for all $ij \in E(R)$.
- For each $i \in [r]$, let $U_i \subseteq \phi^{-1}(i)$ with $|U_i| \leq \alpha m$ and suppose there is a set $T_x \subseteq V_i$ with $|T_x| \geq \nu m$ for each $x \in U_i$.

Then there is a transversal embedding of H inside \mathcal{F} such that for every $i \in [r]$, every $x \in U_i$ is embedded inside T_x .

Proof. Choose new parameters $\mu', \gamma, \zeta, \lambda$ satisfying $\varepsilon, \mu, \alpha \ll \mu' \ll \gamma \ll \zeta \ll \lambda \ll \beta$. Note that n and m are similar in size, since

$$rm \le n \le rm/\delta,$$
 (1)

so in particular $\delta n/r \le m \le |V_j| \le m/\delta \le n/(r\delta)$ for each j.

We have that H is the vertex-disjoint union of connected components H_1, \ldots, H_t each of size at most μn . We will partition each $\phi^{-1}(j)$ into s parts of size roughly γm and one part of size roughly $\mu'm$ that respect components. For each $h \in [t]$ and $j \in [r]$, let $A_{hj} := V(H_h) \cap \phi^{-1}(j)$ and $a_{hj} := |A_{hj}|$. Let t^* be the largest integer such that $b_{0j} := |B_{0j}| \ge \mu'm$ for all $j \in [r]$, where $B_{0j} := A_{t^*+1,j} \cup \ldots \cup A_{tj}$. Obtain $s \in \mathbb{N}$ and $B_{ij} \subseteq \phi^{-1}(j)$ for $i \in [s], j \in [r]$ iteratively as follows. Let $\ell_0 := 0$ and $a_{0j} := 0$ for all $j \in [r]$ and do the following for $i \ge 1$. If

$$|\phi^{-1}(j)| - (a_{1j} + \ldots + a_{\ell_{i-1}j}) > q := 2(\Delta + 1)^{r-1} \gamma m$$

for all $j \in [r]$, let $B_{ij} := A_{\ell_{i-1}+1,j} \cup \ldots \cup A_{\ell_i j}$ where ℓ_i is the smallest integer so that $b_{ij} := |B_{ij}|$ is at least γm for all $j \in [r]$. Otherwise, set s := i, let $B_{sj} := A_{\ell_{s-1}+1,j} \cup \ldots \cup A_{t^*j}$, and let $b_{sj} := |B_{sj}|$. This process always terminates since we remove at least $r\gamma m$ vertices each time, so defines a partition $V_j = B_{0j} \cup B_{1j} \cup \ldots \cup B_{sj}$ for each $j \in [r]$. We have

$$b_{0j} + b_{1j} + \ldots + b_{sj} = |\phi^{-1}(j)| = |V_j|.$$

We claim that

$$\mu' m \leq b_{0j} \leq 2(\Delta+1)^{r-1} \mu' m \quad \text{for all } j \in [r],$$

$$\gamma m \leq b_{ij} \leq 2(\Delta+1)^{2r-2} \gamma m \quad \text{for all } i \in [s], j \in [r] \quad \text{and}$$

$$s \leq 1/(\delta\gamma).$$
(2)

To prove the claim, we first note the following. Since every H_h is a connected graph with $\Delta(H_h) \leq \Delta$,

$$|A_{hj}| \le |V(H_h)| \le (\Delta+1)^{r-1} |A_{hj'}| \quad \text{for every } j, j' \in [r] \text{ and } h \in [t].$$
(3)

(The second inequality can be proved by induction of the number of parts of H_h .)

By construction, for every $j \in [r]$ we have $b_{0j} \geq \mu'm$. There at least one $j' \in [r]$ for which $|B_{0j'}| \leq \mu'm + \mu n \leq 2\mu'm$, otherwise we would have chosen a larger t^* , and so the required upper bound on every b_{0j} follows from (3).

For the second part, suppose first that $i \in [s-1]$. Then every $b_{ij} \geq \gamma m$ by construction. Also there is some $j' \in [r]$ for which $b_{ij'} \leq \gamma m + \mu n \leq 3\gamma m/2$, otherwise ℓ_i would be smaller. Thus $b_{ij} \leq 3(\Delta + 1)^{r-1}\gamma m/2$ for all $j \in [r]$ by (3).

Now consider s = i. There is at least one $j^* \in [r]$ which caused the partitioning to stop during the s-th step due to

$$q \ge |\phi^{-1}(j^*)| - (a_{1j^*} + \ldots + a_{\ell_{s-1}j^*}) = |\phi^{-1}(j^*)| - (b_{1j^*} + \ldots + b_{(s-1)j^*}) = b_{0j^*} + b_{sj^*}.$$

Since the partitioning did not stop during step s - 1, we similarly have $b_{0j^*} + b_{(s-1)j^*} + b_{sj^*} > q$. Thus $b_{sj^*} \ge q - 2(\Delta + 1)^{r-1} \mu' m - 3(\Delta + 1)^{r-1} \gamma m/2 \ge \gamma m$. Altogether, $\gamma m \le b_{sj^*} \le q$.

For any $j \in [r]$ which did not cause the partitioning to stop, we have $b_{0j} + b_{sj} > q$ and hence $b_{sj} > q - 2(\Delta + 1)^{r-1} \mu' m \ge \gamma m$. Moreover, (3) implies that

$$b_{0j} + b_{sj} = a_{\ell_{s-1}+1,j} + \ldots + a_{tj} \le (\Delta + 1)^{r-1} (a_{\ell_{s-1}+1,j^*} + \ldots + a_{tj^*}) = (\Delta + 1)^{r-1} (b_{0j^*} + b_{sj^*}) \le (\Delta + 1)^{r-1} q.$$

Thus $\gamma m \leq b_{sj} \leq (\Delta + 1)^{r-1} q$.

Thus for every $j \in [r]$ we have $\gamma m \leq b_{sj} \leq (\Delta + 1)^{r-1}q$, and hence $\gamma m \leq b_{ij} \leq (\Delta + 1)^{r-1}q$ for all $i \in [s]$ and $j \in [r]$. For the third part, the lower bound implies that $s\gamma m \leq |V_j| \leq m/\delta$ and so $d \leq \delta/\gamma$, completing the proof of the claim that (2) holds. Another related estimate that will be useful later is

$$b_{0j} - sr\varepsilon^{1/3}m > \mu'm - \left(\frac{r|V_j|}{\gamma m}\varepsilon^{1/3}m\right) > \mu'm - \frac{r\varepsilon^{1/3}m}{\gamma} \ge \mu'm/2.$$
(4)

Let $B^i := \bigcup_{i \in [r]} B_{ij}$. We have

$$e(H[B^i]) \le \Delta |B^i| \le 2(\Delta+1)^{2r-1} r\gamma m \quad \text{for all } i \in [s] \quad \text{and}$$

$$e(H[B^0]) \le \Delta |B^0| \le 2(\Delta+1)^r r\mu' m.$$
(5)

In the next part of the proof, we will embed $H[B^1], \ldots, H[B^s]$ in turn. For this, we first partition each V_j into sets for each stage of the embedding, and find a buffer set of colours which will be used in the final part of the embedding. Given an edge xy of G, write $c(xy) := \{c \in \mathscr{C} : xy \in G_c\}$. For each $jj' \in E(R)$ and $x \in V_j$, define $M_{jj'}(x) := \{y \in V_{j'} : |c(xy)| \ge d|\mathscr{C}_{jj'}|/2\}$. We claim that each such set satisfies

$$|M_{jj'}(x)| \ge d|V_{j'}|/2$$

Indeed, semi-superregularity implies that

$$d|V_{j'}||\mathscr{C}_{jj'}| \le \sum_{c \in \mathscr{C}_{jj'}} d_{G_c}(x, V_{j'}) = \sum_{y \in V_{j'}} |c(xy)| \le |M_{jj'}(x)||\mathscr{C}_{jj'}| + |V_{j'}|d|\mathscr{C}_{jj'}|/2$$

as required.

For each $j \in [r]$, let $V_j^0 \cup V_j^1 \cup \ldots \cup V_j^s$ be a random partition of V_j into parts of size $b_{0j} - sr\varepsilon^{1/3}m, b_{1j} + r\varepsilon^{1/3}m, \ldots, b_{sj} + r\varepsilon^{1/3}m$ respectively. For every $i \in [s], j \in [r]$ and $x \in B_{ij}$, we will embed x into V_j^i , which has size slightly larger than necessary. For each $e \in E(R)$, let \mathscr{C}_e^0 be a uniform random subset of \mathscr{C}_e of size $\zeta |\mathscr{C}_e|$ and let $\widetilde{\mathscr{C}}_e := \mathscr{C}_e \setminus \mathscr{C}_e^0$.

By a Chernoff bound, using (2) and (4) to see that all parts are sufficiently large, we may assume that the following hold:

- (C1) for all $jj' \in E(R)$, $x \in V_j$ and $y \in M_{jj'}(x)$, we have $|c(xy) \cap \mathscr{C}^0_{jj'}| \ge d|\mathscr{C}^0_{jj'}|/4$;
- (C2) for all $jj' \in E(R)$ and $x \in V_j$ we have $|M_{jj'}(x) \cap V_{j'}^0| \ge d|V_{j'}^0|/4$;

(C3) for all $0 \le i \le s$ and $x \in V(B^i)$ we have $|T'_x| \ge \nu |V^i_{\phi(x)}|/2$, where $T'_x := T_x \cap V^i_{\phi(x)}$.

Indeed, each of these lower bounds is at most half the expectation of the quantity in question.

We iteratively construct the transversal embedding of $H[B^1 \cup \ldots \cup B^s]$. Suppose that we have obtained an embedding g_{i-1} for some $i \ge 1$, such that

- (i) g_{i-1} is a transversal embedding of $H[W_{i-1}]$, where $W_{i-1} := B^1 \cup \ldots \cup B^{i-1}$;
- (ii) $g_{i-1}(x) \in V_i^{i'}$ for every $x \in B_{i'j} \subseteq W_{i-1}$;
- (iii) for every $j\tilde{j}' \in E(R)$ and every embedded pair x, y of vertices where $xy \in E(H[V_j, V_{j'}])$, the colour of xy is in $\widetilde{\mathscr{C}}_{jj'}$ and these colours are distinct over the embedding; and
- (iv) if $x \in U_j \cap W_{i-1}$, then $g_{i-1}(x) \in T_x$ for every $j \in [r]$.

That is, g_{i-1} consists of injections $\tau : W_{i-1} \to V$ and $\sigma : E(H[W_{i-1}]) \to \mathscr{C}$ such that $\tau(x)\tau(y) \in G_{\sigma(xy)}$, with the stated properties.

Now we will extend g_{i-1} to a transversal embedding g_i of $W_i := W_{i-1} \cup B^i$ with the analogous properties for *i*. First observe that the vertices we are about to embed, that is, $B^i = W_i \setminus W_{i-1}$, have no previously embedded neighbours. Therefore it suffices to find a transversal embedding of B^i when B_{ij} is embedded into V_i^i for each $j \in [r]$ which uses unused colours.

Obtain \mathscr{C}_e^i from $\widetilde{\mathscr{C}}_e$ by deleting any colour we have used in the embedding g_{i-1} . Every colour cluster in the original template contains extra colours, so for every $jj' \in E(R)$, we have

$$|\mathscr{C}_{jj'}^{i}| \ge |\mathscr{C}_{jj'}| - \sum_{1 \le \ell \le i-1} e(H[B_{\ell j}, B_{\ell j'}]) - \zeta |\mathscr{C}_{jj'}| \ge (1-\zeta)|\mathscr{C}_{jj'}| - e(H[B^{i}]) \ge \beta m/2.$$
(6)

For each $j \in [r]$ and $j' \in N_R(j)$, let

$$V_j^{i,j',\text{bad}} := \{ v \in V_j^i : \sum_{c \in \mathscr{C}_{jj'}^i} d_{G_c}(v, V_{j'}^i) < 2d |V_{j'}^i| |\mathscr{C}_{jj'}^i|/3 \}.$$

Lemma 2.4(i) implies that $G_{jj'}^i := (G_c[V_j^i, V_{j'}^i] : c \in \mathscr{C}_{jj'}^i)$ is $(\sqrt{\varepsilon}, d/2)$ -regular and thus, by Lemma 2.3(i) and (2), we have that $|V_j^{i,j',\text{bad}}| < \sqrt{\varepsilon}|V_j^i| < \varepsilon^{1/3}m$. For each $i \in [s]$ and $j \in [r]$, obtain a subset

 Y_j^i of V_j^i by removing $V_j^{i,j',\text{bad}}$ for each $j' \in N_R(j)$ and enough additional arbitrary vertices so that $|Y_j^i| = |V_j^i| - r\varepsilon^{1/3}m = b_{ij}$. Let $\mathcal{Y}^i := \{Y_1^i, \dots, Y_r^i\}$ and let $\mathscr{C}^i := \bigcup_{e \in E(R)} \mathscr{C}_e^i$.

We claim that the subtemplate $\mathcal{F}_i := (\mathcal{Y}^i, \mathscr{C}^i, \mathbf{G}^i)$ of \mathcal{F} induced by $\mathcal{Y}^i, \mathscr{C}^i$ is a semi-super rainbow R-template with parameters $(\gamma m, \sqrt{\varepsilon}, d/2, 1/(2(\Delta+1)^{2r-2}))$. That vertex clusters have suitable sizes follows from (2), and that colour clusters have suitable sizes follows from (6). We have seen that $\mathbf{G}^i_{jj'}$ is $(\sqrt{\varepsilon}, d/2)$ -regular. Moreover, for all $jj' \in E(R)$ and $v \in Y^i_j$, since $v \notin V^{i,j',\text{bad}}_j$, we have

$$\sum_{c \in \mathscr{C}_{jj'}^i} d_{G_c}(v, Y_{j'}^i) \ge 2d|V_{j'}^i||\mathscr{C}_{jj'}^i|/3 - \varepsilon^{1/3}m|\mathscr{C}_{jj'}^i| > d|V_{j'}^i||\mathscr{C}_{jj'}^i|/2$$

so $G^i_{ij'}$ is $(\sqrt{\varepsilon}, d/2)$ -semi-superregular. This completes the proof of the claim.

Therefore, for each $jj' \in E(R)$, we can apply Proposition 2.11 to see that the λ -thick graph $T_{\mathcal{F}_i}^{\lambda}$ is such that $T_{jj'}^i := T_{\mathcal{F}_i}^{\lambda}[Y_j^i, Y_{j'}^i]$ is $(\sqrt{\varepsilon}, d/4)$ -half-superregular for all $jj' \in E(R)$. Apply Theorem 3.3 (the blow-up lemma) with target sets T'_x and parameters $\gamma m, \sqrt{\varepsilon}, \sqrt{\alpha}, \nu/2, d/4$ playing the roles of $m, \varepsilon, \alpha, \nu, d$ to embed $H[B^i]$ into $T_{jj'}^i$ such that for every $j \in [r]$, every $x \in U_j \cap B^i$ is mapped to $T'_x \subseteq V_j^i$, which is possible since, by (C3), $|T'_x| \geq \nu \gamma m/2$ and $|U_j \cap B^i| \leq \alpha m$. Since for each edge of $T_{ij'}^i$, the number of unused colour graphs it lies in is

$$\lambda |\mathscr{C}_{jj'}^i| \stackrel{(6)}{>} \lambda \beta m/2 > 3(\Delta + 1)^{2r-1} r \gamma m \stackrel{(5)}{>} e(H[B^i]),$$

we can greedily assign colours to the embedding. This completes the construction of g_i which satisfies (i)–(iv).

Thus we can obtain g_s with the required properties. At the end of this process, the unembedded part of H is $H[B^0]$, and the set of vertices of V_j which are not an image of any embedded vertex of His precisely $Z_j := V_j^0 \cup \bigcup_{i \in [s]} (V_j^i \setminus Y_j^i)$, which has size b_{0j} . We will use the colours in $\mathscr{C}^0 := \bigcup_{e \in E(R)} \mathscr{C}_e^0$ to embed each B_{0j} into Z_j .

We claim that the subtemplate $\mathcal{F}_0 := (\mathcal{Z}, \mathscr{C}^0, \mathbf{G}_0)$ of \mathcal{F} induced by $\mathcal{Z} := (Z_j : j \in [r]), \mathscr{C}^0$ is a semi-super rainbow template with parameters $(\mu'm, \sqrt{\varepsilon}, d^2/16, 1/(2(\Delta+1)^{r-1}))$. Indeed, (4) implies that vertex clusters have sizes satisfying $\mu'm \leq b_{0j} = |Z_j| \leq 2(\Delta+1)^{r-1}\mu'm$. Colour clusters have size $|\mathscr{C}_{jj'}^0| = \zeta|\mathscr{C}_{jj'}| \geq \zeta \delta m > \mu'm$. Lemma 2.4(i) implies that $\mathbf{G}_{jj'}^0 := (G_c[V_j^0, V_{j'}^0] : c \in \mathscr{C}_{jj'}^0)$ is $(\sqrt{\varepsilon}, d/2)$ -regular for every $jj' \in E(R)$. Additionally, for all $x \in Z_j \subseteq V_j$,

$$\sum_{c \in \mathscr{C}_{jj'}^0} d_{G_c}(x, Z_{j'}) = \sum_{y \in V_{j'}^0} |c(xy) \cap \mathscr{C}_{jj'}^0| \ge \sum_{y \in M_{jj'}(x) \cap V_{j'}^0} |c(xy) \cap \mathscr{C}_{jj'}^0| \stackrel{(C1), (C2)}{\ge} d|V_{j'}^0| / 4 \cdot d|\mathscr{C}_{jj'}^0| / 4$$
$$= d^2 |V_{j'}^0| |\mathscr{C}_{jj'}^0| / 16,$$

so the template is semi-super.

As before, we can apply Proposition 2.11 to see that the λ -thick graph $T^{\lambda}_{\mathcal{F}_0}$ is such that $T^0_{jj'} := T^{\lambda}_{\mathcal{F}^0}[Z^0_j, Z^0_{j'}]$ is $(\sqrt{\varepsilon}, d/4)$ -half-superregular for each $jj' \in E(R)$. We have $\alpha m < \sqrt{\alpha}|Z_j|$, and (C3) implies that $|T'_x| \ge \nu |V^0_{\phi(x)}|/2 \ge \nu |Z_{\phi(x)}|/3$ for all $x \in V(B^0)$ with a target set. Thus we can apply Theorem 3.3 (the blow-up lemma) with target sets T'_x and parameters $\mu'm, \sqrt{\varepsilon}, \sqrt{\alpha}, \nu/3, d/4$ playing the roles of $m, \varepsilon, \alpha, \nu, d$ to embed $H[B^0]$ into $T^0_{jj'}$ such that for every $j \in [r]$, every $x \in U_j$ is mapped to $T'_x \subseteq V^0_j$, which is possible by (C3). The number of colours on each edge of $T^0_{jj'}$ is at least

$$\lambda |\mathscr{C}_{jj'}^0| = \lambda \zeta |\mathscr{C}_{jj'}| \ge \lambda \zeta \delta m > 3(\Delta + 1)^r r \mu' m \stackrel{(5)}{>} e(H[B^0]).$$

Thus we can again greedily assign colours of $\mathscr{C}_{jj'}^0$ to obtain a transversal embedding of $H[B^0]$ using colours untouched by the previous embedding, which completes the transversal embedding of H. \Box

Finally we state a transversal blow-up lemma, which we prove in the next section by combining the embedding lemmas of this section. Its statement is slightly stronger than Theorem 1.8 since the template is super rather than half-super, but the proof of Theorem 1.8 is a simple matter of applying Lemma 2.9(iv) (essentially Lemma 2.1) first. The details can be found at the end of the next section.

Theorem 3.5 (Transversal blow-up lemma). Let $0 < 1/m \ll \varepsilon, \mu, \alpha \ll \nu, d, \delta, 1/\Delta, 1/r \le 1/2$ where r is an integer.

- Let R be a simple graph with vertex set [r] and let $\mathcal{F} = (\mathcal{V}, \mathscr{C}, \mathbf{G})$ be a rainbow super R-template with parameters $(m, \varepsilon, d, \delta)$.
- Suppose that H is a μ -separable graph with $\Delta(H) \leq \Delta$ and there is a graph homomorphism ϕ of H into R such that $|\phi^{-1}(j)| = |V_j|$ for all $j \in [r]$ and $e(H[\phi^{-1}(i), \phi^{-1}(j)]) = |\mathscr{C}_{ij}|$ for all $ij \in E(R)$.
- For each $i \in [r]$, let $U_i \subseteq \phi^{-1}(i)$ with $|U_i| \leq \alpha m$ and suppose there is a set $T_x \subseteq V_i$ with $|T_x| \geq \nu m$ for each $x \in U_i$.

Then there is a transversal embedding of H inside \mathcal{F} such that for every $i \in [r]$, every $x \in U_i$ is embedded inside T_x .

4 Proof of the transversal blow-up lemma

4.1 Sketch of the proof

The proof is a combination of the usual blow-up lemma applied to the thick graph, together with the rainbow partial embedding lemmas in the previous section and the colour absorbing approach pioneered by Montgomery, Müyesser and Pehova in [36].

We begin by outlining the steps of the proof in [36], which is a common sequence of steps for embeddings using absorption, and which we shall also use. The transversal bandwidth theorem in [9] also follows this general outline, as well as using the (usual) blow-up lemma as a key tool. Let $H = H_{\rm con} \cup H_{\rm abs} \cup H_{\rm app} \cup H_{\rm col} \cup H_{\rm vx}$ be a suitable partition, which will be carefully chosen. These parts will be embedded in turn, each time using new colours to extend the transversal embedding.

- Step 0. Embed a small 'connecting graph'. Embed $H_{\rm con}$ whose vertex set, as guaranteed by separability, is such that its removal disconnects H into very small components (Lemma 3.1).
- Step 1. Find a colour absorber. Embed H_{abs} into the edge-coloured graph \mathcal{G} of G, and find disjoint sets $\mathscr{A}, \mathscr{B} \subseteq \mathscr{C}$ of colours such that, given any $e(H_{abs}) |\mathscr{A}|$ colours in \mathscr{B} , we can choose the colours of the edges of H_{abs} using exactly those colours and the colours in \mathscr{A} .
- Step 2. Use most of the colours outside $\mathscr{A} \cup \mathscr{B}$. Embed H_{app} using most of the colours outside $\mathscr{A} \cup \mathscr{B}$ (Lemma 3.4).
- Step 3. Use the remaining colours outside $\mathscr{A} \cup \mathscr{B}$. Embed H_{col} using every unused colour in $\mathscr{C} \setminus (\mathscr{A} \cup \mathscr{B})$ as well as some colours of \mathscr{B} (Lemma 3.2).
- Step 4. Embed the remaining vertices of H using colours in \mathscr{B} . Embed H_{vx} using colours of \mathscr{B} (Lemma 3.4).
- Step 5. Use the colour absorber. The colours for H_{abs} will consist of \mathscr{A} along with the unused colours in \mathscr{B} .

We recall that in [36], these steps were used to find transversal embeddings of spanning trees, and F-factors. When embedding an F-factor where F is a small graph, one can embed each copy of F

in turn, ensuring that they are disjoint. Similarly, to embed a tree, one can embed small subtrees one by one, ensuring that the single vertex of the current tree that was already embedded matches up. The new difficulty in our setting (and in [9]) is that we would like to embed graphs which are more highly-connected. Nevertheless, they have the property that a small fraction of edges can be removed to produce a graph which consists of small (but still linear) connected components.

One difficulty is that we do not have a minimum degree condition. Every vertex has large total degree and every colour graph is dense, but there could be a small proportion of vertices which do not have any edges of a given colour.

For Step 1, we use the following lemma from [36] which is their key tool for colour absorption. Here we present a slightly weaker version with modified parameters for simplicity.

Lemma 4.1 ([36]). Let $0 < \lambda_1 \ll \lambda_2 \ll \lambda_3 < 1$ and let ℓ, m, n be integers with $\ell = \lambda_1 m$ and $1 \leq m \leq \lambda_3^2 n/8$. Suppose that $G = (U, \mathscr{C})$ is a bipartite graph with |U| = m and $|\mathscr{C}| = n$ such that $d(v, \mathscr{C}) \geq \lambda_3 n$ for each $v \in U$. Then there exist disjoint subsets $\mathscr{A}, \mathscr{B} \subseteq \mathscr{C}$ with $|\mathscr{A}| = m - \ell$ and $|\mathscr{B}| = \lambda_2 n$ such that, for every subset \mathscr{B}^0 of \mathscr{B} with size ℓ , we can find a perfect matching between U and $\mathscr{A} \cup \mathscr{B}^0$.

4.2 Proof of Theorem 3.5

Proof of Theorem 3.5. Without loss of generality, we may assume that $\varepsilon \ll \mu \ll \alpha$. We further define constants $\lambda_1, p_{\text{abs}}, p_{\text{col}}, p_{\text{vx}}, \lambda_3, \nu'$ so that altogether

$$0 < 1/m \ll \varepsilon \ll \mu \ll \alpha \ll \lambda_1 \ll p_{\text{abs}} \ll p_{\text{col}} \ll p_{\text{vx}} \ll \lambda_3 \ll \nu' \ll \nu, d, \delta, 1/\Delta, 1/r \le 1/2.$$

Let $p_{\text{app}} := 1 - (p_{\text{vx}} + p_{\text{abs}} + p_{\text{col}})$. Let R be a graph on vertex set [r], $\mathcal{F} = (\mathcal{V}, \mathscr{C}, \mathbf{G})$ a rainbow super template with parameters $(m, \varepsilon, d, \delta)$ and H a graph as in the statement. Let n := |V| be the number of vertices in the template, which equals v(H). Note that

$$rm \le n \le rm/\delta$$
, and
 $m \le |\phi^{-1}(i)| = |V_i| \le m/\delta$ for all $i \in [r]$, and $\delta m \le |\mathscr{C}_e| \le \Delta m/\delta$ for all $e \in E(R)$, (7)

where the final assertion follows from the fact that every $|\mathscr{C}_{ij}| = e(H[\phi^{-1}(i), \phi^{-1}(j)]) \leq \Delta |\phi^{-1}(i)| \leq \Delta m/\delta$.

Preparation of H. First we will choose a partition $H = H_{\rm con} \cup H_{\rm abs} \cup H_{\rm app} \cup H_{\rm vx} \cup H_{\rm col}$ such that each part H_{\circ} has roughly a given number $p_{\circ}n$ of vertices, and moreover $e(H_{\circ}[\phi^{-1}(i), \phi^{-1}(j)]) \approx p_{\circ}e(H[\phi^{-1}(i), \phi^{-1}(j)])$ for all $ij \in E(R)$ and $\circ \in \{\text{abs, app, vx, col}\}$. Since H is μ -separable, there is a set X of size at most μn such that H - X consists of disjoint components H_1, \ldots, H_t , each of size at most μn . For all $\ell \in [t]$ and $ij \in E(R)$, let $H_{ij}^{\ell} := H_{\ell}[\phi^{-1}(i), \phi^{-1}(j)]$ and let $n_j^{\ell} := |V(H_{\ell}) \cap \phi^{-1}(j)|$ and $h_{ij}^{\ell} := e(H_{ij}^{\ell})$. Let $H_{\rm con} := H[X]$. Note that

$$|X| = |V(H_{\rm con})| \le \mu n \le r\mu m/\delta \le \sqrt{\mu}m.$$
(8)

Independently, for each $\ell \in [t]$, add $H^{\ell} := \bigcup_{ij \in E(R)} H_{ij}^{\ell}$ to H_{\circ} with probability p_{\circ} for $\circ \in \{abs, app, vx, col\}$. Then, for all $i \in [r]$ we have $\mathbb{E}(|V(H_{\circ}) \cap \phi^{-1}(i)|) = p_{\circ}|V_i| \pm |X| \ge (p_{\circ} - \sqrt{\mu})m$, and for all $ij \in E(R)$ we have

$$\mathbb{E}(e(H_{\circ}[\phi^{-1}(i),\phi^{-1}(j)])) = p_{\circ}\sum_{\ell\in[t]}h_{ij}^{\ell} = p_{\circ}e(H[\phi^{-1}(i),\phi^{-1}(j)]) \pm \Delta|X| = p_{\circ}|\mathscr{C}_{ij}| \pm \Delta\sqrt{\mu}m$$
$$\geq (p_{\circ}\delta - \Delta\sqrt{\mu})m.$$

Now, $t \ge (1-\mu)/\mu \ge 1/(2\mu)$ and $e(H_{ij}^{\ell}) \le \Delta n_i^{\ell} \le \Delta \mu n$ for all $\ell \in [t]$ and $ij \in E(R)$, and each of these expectations is at least $\delta p_{abs}m/2$, so a Chernoff bound implies that these values are within a multiplicative factor of $(1 \pm \frac{1}{3})$ of their expectations with probability at least

$$1 - 4\left(r + \binom{r}{2}\right) \exp\left(\frac{-(\delta p_{\rm abs}m/2)^2/9}{\frac{1}{2\mu}(\Delta\mu n)^2}\right) \ge 1 - 4r^2 \exp\left(\frac{-p_{\rm abs}^2\delta^4}{18r\Delta\mu}\right) \ge \frac{1}{2}.$$

Thus we may assume that, for all $\circ \in \{abs, app, vx, col\},\$

$$n_i^{\circ} := |V(H_{\circ}) \cap \phi^{-1}(i)| = (1 \pm \frac{1}{2})p_{\circ}|V_i| \in [p_{\circ}m/2, 3p_{\circ}m/(2\delta)] \quad \text{for all } i \in V(R);$$
(9)

$$h_{ij}^{\circ} := e(H_{\circ}[\phi^{-1}(i), \phi^{-1}(j)]) = (1 \pm \frac{1}{2})p_{\circ}|\mathscr{C}_{ij}| \in [p_{\circ}\delta m/2, 3\Delta p_{\circ}m/(2\delta)] \quad \text{for all } ij \in E(R).$$
(10)

We write $H_{ij}^{\circ} := H_{\circ}[\phi^{-1}(i), \phi^{-1}(j)]$ for all $ij \in E(R)$. Every vertex $x \in V(H) \setminus V(H_{\text{con}})$ lies in some H_{\circ} where $\circ \in \{\text{abs, app, vx, col}\}$, and x can only have neighbours inside H_{\circ} or $V(H_{\text{con}})$. Note here that for each $\circ \in \{\text{abs, app, col, vx}\}$, H_{\circ} is the union of vertex-disjoint components of size at most $\mu n \leq (2\mu/p_{\circ})|V(H_{\circ})| \leq \sqrt{\mu}|V(H_{\circ})|$, by (9).

To find a transversal embedding of H, we need to define two injective maps $\tau : V(H) \to V$ and $\sigma : E(H) \to \mathscr{C}$ where $\tau(x)\tau(y) \in G_{\sigma(xy)}$ for all $xy \in E(H)$. For this, we will define $\tau(x)$ for every vertex in H_{con} , followed by every vertex in H_{abs} , H_{app} , H_{col} , H_{vx} . When defining τ for vertices in H_{\circ} , we simultaneously define σ for all future incident edges except for $\circ = \text{abs}$, for which colours are defined at the end. Whenever we define $\tau(y)$ for some y we always choose $\tau(y) \in V_{\phi(y)}$ and if y has a target set T_y , we ensure $\tau(y) \in T_y$. We also always choose $\sigma(xy) \in \mathscr{C}_{\phi(x)\phi(y)}$.

Step 0. We first embed H_{con} into \mathcal{F} by applying Lemma 3.1, as follows. Recall that $X = V(H_{\text{con}})$, and define $Y := N_H(X) \setminus X$. Let H'_{con} be the graph with vertex set $X \cup Y$ and edge set $E(H[X \cup Y]) \setminus E(H[Y])$. Then (8) implies that

$$|V(H'_{\rm con})| \le (\Delta+1)|V(H_{\rm con})| \stackrel{(8)}{\le} 2\Delta\sqrt{\mu}m \quad \text{and hence} \quad e(H'_{\rm con}) \le \mu^{1/3}m.$$
(11)

For each $w \in V(H'_{con})$ where T_w is not defined, let $T_w := V_{\phi(w)}$. Lemma 3.1 applied with $2\Delta\sqrt{\mu}$ playing the role of γ (and the other parameters the same) implies that there are injective maps $\tau : X \to V$ and $\sigma : E(H'_{con}) \to \mathscr{C}$ with $\tau(x)\tau(x') \in G_{\sigma(xx')}$ for all $xx' \in E(H_{con})$, and so that $\tau(x) \in T_x$ for all $x \in X$, and so that there are candidate sets C_y for each $y \in Y$ where $C_y \subseteq V_{\phi(y)} \setminus \tau(X)$ such that $C_y \subseteq \bigcap_{x \in N_H(y) \cap X} N_{G_{\sigma(xy)}}(\tau(x)) \cap T_y$ and $|C_y| \ge \nu' m$.

We update the target sets, by defining $T_y^1 := C_y \subseteq T_y$ for $y \in Y$ where this set is defined, and $T_y^1 := T_y$ if $y \notin V(H'_{\text{con}})$ and this set is defined. This updates target sets T_y^1 for all unembedded vertices y, and all target sets have size at least $\nu'm$. Note that for all $y \in Y$, if we choose $\tau(y) \in T_y^1$, then $\tau(x)\tau(y) \in G_{\sigma(xy)}$ for all $x \in N_H(y) \cap X$, so the colour $\sigma(xy)$ will be used as desired. For each $i \in [r]$, let U_i^1 be the set of vertices y in $\phi^{-1}(i)$ with a target set T_y^1 . We have

$$|U_i^1| \le |U_i| + |Y| \le \alpha m + 2\Delta \sqrt{\mu} m \le 2\alpha m.$$
(12)

This set will only shrink during the rest of the proof as vertices with target sets are embedded. Some target sets of vertices in U_i^1 will also shrink, but will remain large enough.

Let $n_i^{\text{con}} := |V(H_{\text{con}}) \cap \phi^{-1}(i)|$ for all $i \in [r]$ and $h_{ij}^{\text{con}} := e(H'_{\text{con}}[\phi^{-1}(i), \phi^{-1}(j)])$ for all $ij \in E(R)$. We have

$$|V_i| = n_i^{\text{con}} + n_i^{\text{abs}} + n_i^{\text{app}} + n_i^{\text{vx}} + n_i^{\text{col}} \quad \text{for all } i \in [r] \quad \text{and} |\mathscr{C}_{ij}| = e(H[\phi^{-1}(i), \phi^{-1}(j)]) = h_{ij}^{\text{con}} + h_{ij}^{\text{abs}} + h_{ij}^{\text{app}} + h_{ij}^{\text{vx}} + h_{ij}^{\text{col}} \quad \text{for all } ij \in E(R).$$
(13)

Let $\mathcal{F}' = (\mathcal{V}', \mathscr{C}', \mathbf{G}')$ be the subtemplate of \mathcal{F} obtained by deleting the vertices $\tau(X)$ and the colours $\sigma(E(H'_{\text{con}}))$ from \mathcal{F} , so, defining for all $i \in [r]$ and $e \in E(R)$

$$V'_i := V_i \setminus \tau(X), \quad \mathscr{C}'_e := \mathscr{C}_e \setminus \sigma(E(H'_{\text{con}})),$$

we have $(1 - \mu^{1/4})|\mathscr{C}_e| \stackrel{(7),(11)}{\leq} |\mathscr{C}'_e| \stackrel{(13)}{=} h_e^{\text{abs}} + h_e^{\text{app}} + h_e^{\text{col}} + h_e^{\text{vx}}.$

By Lemma 2.9(ii), \mathcal{F}' is a rainbow super *R*-template with parameters $(m/2, 2\varepsilon, d/2, \delta/2)$.

Step 1. Next we embed (the vertices but not the colours of) H_{abs} into \mathcal{F}' . For each $i \in [r]$, let $V_i^{abs} \subseteq V_i'$ be a uniform random subset of size n_i^{abs} . Lemma 2.9(iii) applied with $\mathcal{F}', p_{abs}/3$, 6 playing the roles of \mathcal{F}, α, k implies that the subtemplate \mathcal{F}_{abs} of \mathcal{F}' induced by $(V_i^{abs} : i \in [r])$ is a rainbow super *R*-template with parameters $(p_{abs}m/3, \sqrt{\varepsilon}, d^2/64, \delta/6)$. Furthermore, a Chernoff bound implies that we may assume $|T_y^1 \cap V_{\phi(y)}^{abs}| \ge 2\nu' n_i^{abs}/3 \ge \nu' |V_i^{abs}|/3$ for all $y \in V(H_{abs})$ with a target set T_y^1 .

Let $V_i'' := V_i' \setminus V_i^{\text{abs}}$. We have

$$|V_i''| \ge |V_i| - 2\Delta\sqrt{\mu}m - n_i^{\text{abs}} \stackrel{(9)}{\ge} (1 - 2p_{\text{abs}})|V_i| \ge m/2.$$
(14)

Lemma 2.9(ii) applied with $\mathcal{F}, 2p_{abs}$ playing the roles of \mathcal{F}, α implies that the subtemplate \mathcal{F}'' induced by V''_i is a rainbow super *R*-template with parameters $(m/2, 2\varepsilon, d/2, \delta/2)$, and a Chernoff bound implies that we may assume

$$|T_y^1 \cap V_{\phi(y)}''| \ge 2(\nu'm - n_i^{\text{abs}})/3 \ge \nu'm/2 \tag{15}$$

for all y with a target set T_y^1 .

We will embed H_{abs} into the λ_3 -thick graph $T_{\mathcal{F}_{abs}}^{\lambda_3}$. By Proposition 2.11, for every $ij \in E(R)$, $T_{\mathcal{F}_{abs}}^{\lambda_3}[V_i^{abs}, V_j^{abs}]$ is $(\sqrt{\varepsilon}, d^2/128)$ -half-superregular. Apply Theorem 3.3 (the blow-up lemma) with target sets $T_y^1 \cap V_{\phi(y)}^{abs}$ and parameters $\sqrt{\varepsilon}, \sqrt{\alpha}, \sqrt{\nu'}, d^2/128$ playing the roles of $\varepsilon, \alpha, \nu, d$ to find an embedding τ of H_{abs} into $T_{\mathcal{F}_{abs}}^{\lambda_3}$ such that $\tau(x) \in V_{\phi(x)}$ for all $x \in V(H_{abs})$ and $\tau(y) \in T_y^1$ for every $y \in U_i^1$ with $i \in [r]$. Note that we haven't yet defined $\sigma(e)$ for any $e \in E(H_{abs})$. However, by our definition, for every such e = xy we have $\tau(x)\tau(y) \in E(G_c)$ for at least $\lambda_3|\mathscr{C}'_e|$ colours $c \in \mathscr{C}'_e$. For each $e \in E(R)$, let $G_{abs,e}$ be the auxiliary bipartite graph with vertex classes $Z_e := \{\tau(x)\tau(y) : xy \in E(H_{abs}^e)\}$ and \mathscr{C}'_e , where $\{\tau(x)\tau(y), c\}$ is an edge whenever $\tau(x)\tau(y) \in E(G_c)$. Then (10) implies that $|Z_e| = h_e^{abs} \leq 3p_{abs}|\mathscr{C}_e|/2 \leq 2p_{abs}|\mathscr{C}'_e|$, and by construction, every $\tau(x)\tau(y)$ in Z_e has degree at least $\lambda_3|\mathscr{C}'_e|$ in $G_{abs,e}$. For each $e \in E(R)$, define constants ℓ_e and p_e via

$$\ell_{e} := \lambda_{1} h_{e}^{\text{abs}} \text{ and}$$

$$p_{e} |\mathscr{C}'_{e}| := (1 - p_{\text{abs}}) |\mathscr{C}'_{e}| - (h_{e}^{\text{abs}} - \ell_{e}) - h_{e}^{\text{app}} = h_{e}^{\text{vx}} + h_{e}^{\text{col}} - p_{\text{abs}} |\mathscr{C}'_{e}| + \lambda_{1} h_{e}^{\text{abs}}$$

$$\stackrel{(10)}{=} (1 \pm \frac{1}{4}) h_{e}^{\text{vx}} \in [p_{\text{vx}} |\mathscr{C}'_{e}| / 3, 3p_{\text{vx}} |\mathscr{C}'_{e}|].$$

$$(17)$$

Thus for each $e \in E(R)$ we can apply Lemma 4.1 with $G_{abs,e}, Z_e, \mathscr{C}'_e, \lambda_1, p_e, \lambda_3$ playing the roles of $G, U, \mathscr{C}, \lambda_1, \lambda_2, \lambda_3$ to obtain disjoint sets $\mathscr{A}_e, \mathscr{B}_e \subseteq \mathscr{C}'_e$ such that

- (Q1) $|\mathscr{A}_e| = h_e^{\text{abs}} \ell_e \overset{(10)}{\leq} 2p_{\text{abs}}|\mathscr{C}'_e|;$
- (Q2) $|\mathscr{B}_e| = p_e |\mathscr{C}'_e|;$
- (Q3) for any set $\mathscr{B}_e^0 \subseteq \mathscr{B}_e$ of size ℓ_e , there exists a colouring using colours from $\mathscr{A}_e \cup \mathscr{B}_e^0$ that makes the embedding τ of H_e^{abs} rainbow. That is, there is a bijection $\sigma_e : E(H_e^{\text{abs}}) \to \mathscr{A}_e \cup \mathscr{B}_e^0$ such that $\tau(x)\tau(y) \in G_{\sigma_e(xy)}$ for all $xy \in E(H_e^{\text{abs}})$.

Preparation for Steps 2–4. Recall that the template $\mathcal{F}'' = (\mathcal{V}'', \mathcal{C}', \mathbf{G}'')$ has vertex clusters $\mathcal{V}'' := (V''_i := V'_i \setminus V_i^{abs} : i \in [r])$, colour clusters $(\mathcal{C}'_e : e \in E(R))$ and graphs $(\mathbf{G}'')^{ij} = (G_c[V''_i, V''_j] : c \in \mathcal{C}'_{ij})$, and \mathcal{F}'' is a rainbow super *R*-template with parameters $(m/2, 2\varepsilon, d/2, \delta/2)$. Let $n_i^{colvx} := n_i^{col} + n_i^{vx}$ for each $i \in [r]$ and let $p_{colvx} := p_{col} + p_{vx}$. During the rest of the proof, for each $\delta \in \{app, colvx\}$ we will identify pairwise disjoint vertex sets V_i^{δ} in the vertex clusters $V_i'', i \in [r]$, so that

 $|V_i^\diamond| = n_i^\diamond, \quad \text{and} \quad V_i'' = V_i^{\text{app}} \cup V_i^{\text{colvx}}, \quad \text{where} \quad \mathcal{V}^\diamond := \{V_1^\diamond, \dots, V_r^\diamond\} \; \forall \diamond \in \{\text{app, colvx}\}.$

We will choose $\tau(x) \in V_{\phi(x)}^{\text{app}}$ for all $x \in V(H_{\text{app}})$ and $\tau(x) \in V_{\phi(x)}^{\text{colvx}}$ for all $x \in V(H_{\text{col}}) \cup V(H_{\text{vx}})$. We will identify pairwise disjoint colour sets \mathscr{C}° , so that

$$\mathscr{C}^{\circ} = \bigcup_{e \in E(R)} \mathscr{C}^{\circ}_{e} \ \forall \circ \in \{ \operatorname{app}, \operatorname{col}, \operatorname{vx} \}, \quad \mathscr{C}'_{e} = \bigcup_{\circ \in \{ \operatorname{app}, \operatorname{col}, \operatorname{vx} \}} \mathscr{C}^{\circ}_{e}$$

and for each $xy \in E(H_{\circ})$, we will ensure $\sigma(xy) \in G_c$ for some $c \in \mathscr{C}^{\circ}_{\phi(x)\phi(y)}$. Given such sets of vertices and colours for $\circ \in \{\text{app, col, vx}\}$, we define $\mathcal{F}_{\circ} := (\mathcal{V}^{\circ'}, \mathscr{C}^{\circ}, \mathbf{G}^{\circ})$ to be the subtemplate of \mathcal{F}'' induced by $\mathcal{V}^{\circ'}, \mathscr{C}^{\circ}$, where app' = app and vx' = col' = colvx. Note that these templates are always rainbow *R*-templates with pairwise disjoint sets of colours, but \mathcal{F}_{col} and \mathcal{F}_{vx} share the same vertex clusters.

We will now define the vertex sets $V_i^{\circ'}$ for $\circ' \in \{\text{app, colvx}\}$, but colour sets will be defined sequentially, given the colours used in each step. For each $i \in [r]$, do the following. For each $j \in N_R(i)$, recall that $(G_c[V_i'', V_j''] : c \in \mathscr{C}'_{ij})$ is $(2\varepsilon, d/2)$ -superregular. By (16) and (Q2), we have $|\mathscr{B}_{ij}| \geq p_{\text{vx}}|\mathscr{C}'_{ij}|/2$, so by Lemma 2.4(i), $(G_c[V_i'', V_j''] : c \in \mathscr{B}_{ij})$ is $(4\varepsilon/p_{\text{vx}}, d/4)$ -regular and hence $(\sqrt{\varepsilon}/r, d/4)$ -regular. By Lemma 2.3(i), there is a set $\overline{V_i^j} \subseteq V_i''$ with $|\overline{V_i^j}| \leq \sqrt{\varepsilon}|V_i''|/r$ such that

$$\sum_{c \in \mathscr{B}_{ij}} d_{G_c}(v) \ge d|V_j''||\mathscr{B}_{ij}|/5 \quad \text{for all } v \in V_i'' \setminus \overline{V}_i^j.$$
(18)

Let $\overline{V}_i := \bigcup_{j \in N_R(i)} \overline{V}_i^j$, so $|\overline{V}_i| \le \sqrt{\varepsilon} |V_i''|$, and let V_i^{colvx} be a uniform random subset of $V_i'' \setminus \overline{V}_i$ of size n_i^{colvx} , and let $V_i^{\text{app}} := V_i'' \setminus V_i^{\text{colvx}}$. So $|V_i^{\text{app}}| = n_i^{\text{app}}$. Let $T_y^2 := T_y^1 \cap V_{\phi(y)}^{\circ'(y)}$ for all $y \in U_i^1$ and $i \in [r]$ (i.e. those y for which T_y^1 has been defined), where $\circ'(y) = \text{app}$ if $y \in V(H_{\text{app}})$ and $\circ'(y) = \text{colvx}$ if $y \in V(H_{\text{col}}) \cup V(H_{\text{vx}})$.

For all $c \in \mathscr{C}'_{ij}$, the superregularity of our graph collections implies that

$$\begin{split} \mathbb{E}(e(G_c[V_i^{\text{colvx}}, V_j^{\text{colvx}}])) &\geq (n_i^{\text{colvx}}/|V_i''|)(n_j^{\text{colvx}}/|V_j''|)(e(G_c[V_i'', V_j'']) - \sqrt{\varepsilon}|V_i''||V_j''|) \\ &\geq dn_i^{\text{colvx}}n_j^{\text{colvx}}/3 \stackrel{(9)}{\geq} dp_{\text{vx}}^2 \delta^2 m^2/12. \end{split}$$

For all $i \in [r]$ and $y \in U_i^1$ we have

$$\mathbb{E}(|T_y^2|) \ge (n_i^{\circ'(y)}/|V_i''|)(|T_y^1 \cap V_i''| - \sqrt{\varepsilon}|V_i''|) \stackrel{(9),(15),(14)}{\ge} p_{\circ'(y)}\nu'm/5.$$

Further, for all $ij \in E(R)$ and $v \in V_i^{\text{colvx}}$, since $v \notin \overline{V}_i^j$ we have

$$\mathbb{E}\left(\sum_{c\in\mathscr{B}_{ij}} d_{G_c}(v, V_j^{\text{colvx}})\right) \ge (n_j^{\text{colvx}}/|V_j''|) \sum_{c\in\mathscr{B}_{ij}} (d_{G_c}(v, V_j'') - \sqrt{\varepsilon}|V_j''|) \stackrel{(18)}{\ge} dn_j^{\text{colvx}}|\mathscr{B}_{ij}|/11.$$

By Chernoff bounds, we may assume that each of the above quantities are close to their expectations, so

- (C1) $|T_y^2| \ge p_{o'(y)}\nu'm/6$ for all $y \in U_i^1$ and $i \in [r]$ (i.e. all those y which have a target set); (C2) $e(G_c[V_i^{\text{colvx}}, V_j^{\text{colvx}}]) \ge dn_i^{\text{colvx}} n_j^{\text{colvx}}/13$ for all $ij \in E(R)$ and $c \in \mathscr{C}'_{ij}$;
- (C3) $\sum_{c \in \mathscr{B}_{ij}} d_{G_c}(v, V_j^{\text{colvx}}) \ge dn_j^{\text{colvx}} |\mathscr{B}_{ij}|/12 \text{ for all } ij \in E(R) \text{ and } v \in V_i^{\text{colvx}}.$

Let $m' := p_{vx}m/8$. Now (12) and (C1) imply that, for all $i \in [r]$, the number $|U_i^1|$ of vertices $y \in \phi^{-1}(i)$ with a target set T_y^2 satisfy

$$|U_i^1| \le 2\alpha m \le \sqrt{\alpha} m', \quad \text{and} \quad |T_y^2| \ge p_{vx} \nu' m/6 > \nu' m'.$$
(19)

For all $i \in [r]$ we have $|V_i^{\text{colvx}}| = n_i^{\text{colvx}} = n_i^{\text{col}} + n_i^{\text{vx}}$, so

$$p_{\rm vx}|V_i''|/3 \stackrel{(14)}{\leq} p_{\rm vx}|V_i|/2 \stackrel{(9)}{\leq} |V_i^{\rm colvx}| \stackrel{(9)}{\leq} 2p_{\rm vx}|V_i| \stackrel{(14)}{\leq} 3p_{\rm vx}|V_i''|.$$
(20)

We claim that the following properties about templates \mathcal{F}_{\circ} with colour sets \mathscr{C}_{e}° for $\circ \in \{app, col, vx\}$ hold.

(F1) Let

$$\mathscr{C}_e^{\operatorname{app}} := \mathscr{C}'_e \setminus (\mathscr{A}_e \cup \mathscr{B}_e) \text{ for all } e \in E(R).$$

Then \mathcal{F}_{app} , with vertex clusters $(V_i^{app} : i \in [r])$, is a rainbow super *R*-template with parameters $(m/4, 4\varepsilon, d/4, \delta/4).$

(F2) Suppose

$$\mathscr{C}_e^{\operatorname{col}} = \mathscr{B}_e \cup \mathscr{D}_e$$
 where $\mathscr{D}_e \subseteq \mathscr{C}^{\operatorname{app}}$ for all $e \in E(R)$.

Then \mathcal{F}_{col} , with vertex clusters $(V_i^{colvx} : i \in [r])$, is a rainbow *R*-template with parameters $(m', \sqrt{\varepsilon}, d/4, \delta/24)$ with the property that $e(G_c[V_i^{colvx}, V_j^{colvx}]) \geq d|V_i^{colvx}||V_j^{colvx}|/22$ for all $c \in \mathscr{D}_e$ and $ij \in E(R)$.

(F3) Suppose

 $\mathscr{C}_e^{\mathrm{vx}} \subseteq \mathscr{B}_e$ and $|\mathscr{B}_e \setminus \mathscr{C}_e^{\mathrm{vx}}| \leq 2p_{\mathrm{col}}|\mathscr{C}_e'|$ for all $e \in E(R)$.

Then \mathcal{F}_{vx} , with colour clusters $(V_i^{colvx} : i \in [r])$, is a rainbow super *R*-template with parameters $(m', \sqrt{\varepsilon}, d/13, \delta/24).$

These properties follow from Lemma 2.9 applied to \mathcal{F}_{\circ} as a subtemplate of $\mathcal{F}'' = (\mathcal{V}'', \mathscr{C}', \mathbf{G}'')$ (which has parameters $(m/2, 2\varepsilon, d/2, \delta/2))$, as follows. First, (F1) holds by Lemma 2.9(ii) since \mathcal{F}_{app} is a small perturbation of \mathcal{F}'' . Indeed, (Q1) and (Q2) implies that $|\mathscr{C}'_e \setminus \mathscr{C}^{app}_e| \leq (2p_{abs} + 3p_{vx})|\mathscr{C}'_e| \leq (2p_{abs} + 3p_{vx})|$ $4p_{\rm vx}|\mathscr{C}'_e| \leq \sqrt{p_{\rm vx}}m/2$, and $|V''_i \setminus V^{\rm app}_i| = |V^{\rm colvx}_i| \leq 3p_{\rm vx}|V''_i| \leq \sqrt{p_{\rm vx}}m/2$ by (20), so we can apply the lemma with $2\varepsilon, \sqrt{p_{\rm vx}}, d/2$ playing the roles of ε, α, d to obtain (F1).

For (F2), (Q2) and (16) imply that $|\mathscr{C}_e^{col}| \ge |\mathscr{B}_e| = p_e |\mathscr{C}'_e| > p_{vx} |\mathscr{C}'_e|/4$. Together with (20), this means that we can apply Lemma 2.9(i) with parameters $m/2, 2\varepsilon, p_{\rm vx}/4, d/2, \delta/2, 12$ playing the roles of $m, \varepsilon, \alpha, d, \delta, k$ to see that \mathcal{F}_{col} is a rainbow *R*-template with parameters $(m', \sqrt{\varepsilon}, d/4, \delta/24)$. The second property follows from (C2) since $\mathscr{D}_e \subseteq \mathscr{C}_e^{\operatorname{app}} \subseteq \mathscr{C}'_e$.

For property (F3), we have $|\mathscr{C}_e^{\mathrm{vx}}| \geq |\mathscr{B}_e| - 2p_{\mathrm{col}}|\mathscr{C}'_e| \geq p_{\mathrm{vx}}|\mathscr{C}'_e|/4$ by (16). Together with (20), this means we can apply Lemma 2.9(i) with the same parameters as previously to see that \mathcal{F}_{vx} is a rainbow R-template with the given parameters. For superregularity, we have for all $e = ij \in E(R)$ and $v \in V_i^{\text{colvx}}$ that

$$\sum_{c \in \mathscr{C}_{ij}^{\mathrm{vx}}} d_{G_c}(v, V_j^{\mathrm{colvx}}) \ge \sum_{c \in \mathscr{B}_{ij}} d_{G_c}(v, V_j^{\mathrm{colvx}}) - 2p_{\mathrm{col}}|\mathscr{C}_{ij}'||V_j''| \overset{(7), (18), (C3)}{\ge} dn_j^{\mathrm{colvx}}|\mathscr{B}_{ij}|/12 - 2p_{\mathrm{col}}\Delta m^2/\delta^2$$
$$\ge dn_j^{\mathrm{colvx}}|\mathscr{C}_{ij}^{\mathrm{vx}}|/13,$$

and $e(G_c[V_i^{\text{colvx}}, V_j^{\text{colvx}}]) \ge dn_i^{\text{colvx}} n_j^{\text{colvx}}/13$ for all $c \in \mathscr{C}_{ij}^{\text{vx}}$ by (C2). Thus (F1)–(F3) all hold.

Now it is a matter of embedding each H_{\circ} into its corresponding template, using suitable (unused) colours at each step. The image of $V(H_{vx})$ in each V_i will be the remaining vertices of V_i^{colvx} after embedding H_{col} , which is most of this set since H_{col} is much smaller than H_{vx} .

Step 2. We embed H_{app} into \mathcal{F}_{app} using Lemma 3.4 (embedding lemma with extra colours) with parameters $m/4, 4\varepsilon, \sqrt{\mu}, \sqrt{\alpha}, 2p_{abs}\delta, \nu'/7, d/4, \delta/4$ playing the roles of $m, \varepsilon, \mu, \alpha, \beta, \nu, d, \delta$. For this, we recall that H_{app} is the union of components of size at most $\sqrt{\mu}|V(H_{app})|$, and there is a graph homomorphism from H_{app} into R. By (C1) and (F1), it now suffices to check that colour clusters have a suitable size to apply the lemma. Indeed, for all $e \in E(R)$ we have

$$\begin{aligned} |\mathscr{C}_{e}^{\mathrm{app}}| - h_{e}^{\mathrm{app}} &\stackrel{(F1)}{=} & |\mathscr{C}_{e}'| - |\mathscr{A}_{e}| - |\mathscr{B}_{e}| - h_{e}^{\mathrm{app}} \stackrel{(Q1),(Q2)}{=} |\mathscr{C}_{e}'| - (h_{e}^{\mathrm{abs}} - \ell_{e}) - p_{e}|\mathscr{C}_{e}'| - h_{e}^{\mathrm{app}} \stackrel{(16)}{=} p_{\mathrm{abs}}|\mathscr{C}_{e}'| \\ & \in & [p_{\mathrm{abs}}\delta m/2, p_{\mathrm{abs}}\Delta m/\delta]. \end{aligned}$$

$$(21)$$

Thus we can apply Lemma 3.4 to obtain $\tau(V(H^{app}))$ and $\sigma(E(H^{app}))$ where each y with a target set T_y^2 is embedded inside it.

For each $e \in E(R)$, let $\mathscr{D}_e := \mathscr{C}_e^{\operatorname{app}} \setminus \sigma(E(H_{\operatorname{app}}))$ and let $\mathscr{C}_e^{\operatorname{col}} := \mathscr{B}_e \cup \mathscr{D}_e$.

Step 3. We embed $H_{\rm col}$ into $\mathcal{F}_{\rm col}$ by applying Lemma 3.2 (embedding lemma with target sets and prescribed colours) with $m', \sqrt{\varepsilon}, \sqrt{\alpha}, \sqrt{p_{\rm abs}}, \sqrt{p_{\rm col}}, p_{\rm col}, \nu', d/4, \delta/24$ playing the roles of $m, \varepsilon, \alpha, \gamma, \lambda_1, \lambda_2, \nu, d, \delta$. To see that this is possible, we have $|V(H_{\rm col}) \cap \phi^{-1}(i)| = n_i^{\rm col} \leq 2p_{\rm col}m/\delta \leq \sqrt{p_{\rm col}}m'$. We also have

$$e(H_{\rm col}[\phi^{-1}(i),\phi^{-1}(j)]) = h_{ij}^{\rm col} \stackrel{(10)}{\geq} p_{\rm col}\delta m/2 \ge p_{\rm col}m'.$$

Equation (19) implies that the number of vertices with target sets and the size of these target sets are suitable. By (21), for every $e \in E(R)$ we have

$$|\mathscr{D}_e| = |\mathscr{C}_e^{\mathrm{app}}| - h_e^{\mathrm{app}} \le p_{\mathrm{abs}} \Delta m / \delta \le \sqrt{p_{\mathrm{abs}}} m'.$$
(22)

By (F2), it remains to check that $|\mathscr{C}_e^{col} \setminus \mathscr{D}| \ge d|\mathscr{C}_e^{col}|/4$. Since the original template was rainbow, $\mathscr{C}_e^{col} \setminus \mathscr{D} = \mathscr{C}_e^{col} \setminus \mathscr{D}_e = \mathscr{B}_e$. Equation (22) implies that $|\mathscr{B}_e| \ge |\mathscr{C}_e^{col}| - \sqrt{p_{abs}}m' \ge |\mathscr{C}_e^{col}|/2 \ge d|\mathscr{C}_e^{col}|/4$, as required. Thus we can apply Lemma 3.2 to obtain $\tau(V(H_{col}))$ and $\sigma(E(H_{col}))$ where each y with a target set T_y^2 (i.e. those in $U_{\phi(y)}^1$) is embedded inside it.

Let
$$\mathscr{C}_e^{\mathrm{vx}} := \mathscr{B}_e \setminus \sigma(E(H_{\mathrm{col}}))$$
 and $V_i^{\mathrm{vx}} := V_i^{\mathrm{colvx}} \setminus \tau(V(H_{\mathrm{col}}))$, so $V_i^{\mathrm{vx}} = n_i^{\mathrm{vx}}$.

Step 4. Let \mathcal{F}'_{vx} be the subtemplate of \mathcal{F}_{vx} induced by $(V_i^{vx} : i \in [r])$. It is a small perturbation: $|V_i^{colvx} \setminus V_i^{vx}| = n_i^{col} \leq 2p_{col}m/\delta \leq p_{vx}m'$, so Lemma 2.9(ii) with $m', \sqrt{\varepsilon}, p_{vx}, d/13, \delta/24$ playing the roles of $m, \varepsilon, \alpha, d, \delta$ implies that \mathcal{F}'_{vx} is a rainbow super template with parameters $(m'/2, 2\sqrt{\varepsilon}, d/26, \delta/48)$. Let $T_y^3 := T_y^2 \cap V_i^{vx}$ for all $i \in [r]$ and $y \in V_i^{colvx} \cap U_i^1$.

We embed H_{vx} into \mathcal{F}'_{vx} by applying Lemma 3.4 with parameters $\sqrt{\mu}, \sqrt{\alpha}, \lambda_1^2, \nu'/3$ playing the roles of μ, α, β, ν (and template parameters as above). For this, we recall that H_{vx} is the union of vertex disjoint components of size at most $\sqrt{\mu}|V(H_{vx})|$ and there is a graph homomorphism from H_{vx} into R. By (19), there are suitably few vertices with target sets T_y^3 , and all target sets are suitably large. By (F3), it now suffices to check that every $|\mathscr{C}_e^{vx}| - h_e^{vx}$ is large. We have

$$\begin{aligned} |\mathscr{C}_e^{\mathrm{vx}}| - h_e^{\mathrm{vx}} &= |\mathscr{B}_e| - h_e^{\mathrm{col}} - h_e^{\mathrm{vx}} \stackrel{(F1)}{=} |\mathscr{C}_e'| - |\mathscr{A}_e| - h_e^{\mathrm{app}} - h_e^{\mathrm{col}} - h_e^{\mathrm{vx}} = h_e^{\mathrm{abs}} - |\mathscr{A}_e| = \ell_e = \lambda_1 h_e^{\mathrm{abs}} \\ &\geq \lambda_1 p_{\mathrm{abs}} \delta m/2 \geq \lambda_1^2 m'/2. \end{aligned}$$

Thus we can apply Lemma 3.4 to obtain $\tau(V(H_{vx}))$ and $\sigma(E(H_{vx}))$ where each y with a target set T_y^2 is embedded inside it.

Let
$$\mathscr{C}_e^{\text{abs}} := \mathscr{A}_e \cup (\mathscr{C}_e^{\text{vx}} \setminus \sigma(E(H_{\text{vx}}))).$$

Step 5. Since $\mathscr{C}'_e = \mathscr{C}^{\mathrm{app}}_e \cup \mathscr{C}^{\mathrm{col}}_e \cup \mathscr{C}^{\mathrm{vx}}_e \cup \mathscr{C}^{\mathrm{abs}}_e$ is a disjoint union, and $\mathscr{A}_e \subseteq \mathscr{C}^{\mathrm{abs}}_e \subseteq \mathscr{A}_e \cup \mathscr{B}_e$, the set $\mathscr{C}^{\mathrm{abs}}_e \cap \mathscr{B}_e$ must have exactly the right size:

$$|\mathscr{C}_e^{\mathrm{abs}} \cap \mathscr{B}_e| = |\mathscr{C}_e'| - h_e^{\mathrm{app}} - h_e^{\mathrm{col}} - h_e^{\mathrm{vx}} - |\mathscr{A}_e| = |\mathscr{C}_e'| - (|\mathscr{C}_e'| - h_e^{\mathrm{abs}}) - |\mathscr{A}_e| = \ell_e.$$

Thus, by (Q3), for each $e \in E(R)$ we can find σ_e with image $\sigma_e(E(H_{abs})) = \mathscr{C}_e^{abs}$. Extending σ by all of these σ_e completes the transversal embedding.

4.3 Proof of Theorem 1.8

Proof of Theorem 1.8. We may assume without loss of generality that $1/m \ll \varepsilon, \mu, \alpha \ll \nu, d, \delta, 1/\Delta, 1/r \leq 1/2$. Choose an additional constant ε' with $\varepsilon \ll \varepsilon' \ll \nu, d, \delta, 1/\Delta, 1/r$ such that the conclusion of Lemma 2.1 holds with $\varepsilon, \varepsilon', d, \delta, 3$ playing the roles of $\varepsilon, \varepsilon', d, \delta, k$. We may further assume that the conclusion of Theorem 3.5 holds with $\varepsilon', d^2/2$ playing the roles of ε, d and the other parameters unchanged.

Let $G, \mathscr{C}, R, \mathcal{V} := (V_1, \ldots, V_r)$ be as in the statement of Theorem 1.8. Let $\phi : V(H) \to V(R)$ be such that $\phi(x) = j$ whenever $x \in A_j$. So ϕ is a graph homomorphism. Our hypothesis is that $\mathcal{F} = (\mathcal{V}, \mathscr{C}, \mathbf{G})$ is a rainbow half-super *R*-template with parameters $(m, \varepsilon, d, \delta)$. Lemma 2.9(iv) implies that for all $c \in \mathscr{C}$ there is $G'_c \subseteq G_c$ such that, writing $\mathbf{G}' := (G'_c : c \in \mathscr{C})$, the template $\mathcal{F}' = (\mathcal{V}, \mathscr{C}, \mathbf{G}')$ is super with parameters $(m, \varepsilon', d^2/2, \delta)$.

Apply Theorem 3.5 to \mathcal{F}' to obtain the required embedding.

5 Proof of Theorems 1.2 and 1.6

In this section we prove two applications of our transversal blow-up lemma, to super uniformly dense graph collections (Theorem 1.2) and super uniformly dense 3-graphs (Theorem 1.6). The latter is an easy consequence of the former.

Proof of Theorem 1.2. Let $\Delta, d, \delta, \alpha > 0$ be given and let $r := \Delta + 1$. Choose additional constants such that

$$0 < 1/n_0 \ll \eta, \mu \ll \varepsilon \ll \delta_1 \ll \delta_2 \ll \ldots \ll \delta_{\binom{r}{2}+1} \ll \nu' \ll \alpha \ll \delta, d, 1/\Delta,$$

where we are assuming without loss of generality that $\alpha \ll \delta, d/1/\Delta$, and so that the conclusion of Lemma 2.1 holds with $3, \eta^{1/4}, \varepsilon, \alpha^2/6$ playing the roles of $k, \varepsilon, \varepsilon', d$, and Lemma 3.1 holds with $n_0/r, \varepsilon, \sqrt{\delta_1}, \alpha^4/72$ playing the roles of $m, \varepsilon, \gamma, d$ and Theorem 3.5 holds with $\delta_2^2 n_0/2, \sqrt{\varepsilon}, 2\mu, \delta_1^{1/3}, \nu', \alpha^9, \delta/2$ playing the roles of $m, \varepsilon, \mu, \alpha, \nu, d, \delta$. Let $n \ge n_0$ be an integer and let $\mathbf{G} = (G_c : c \in \mathscr{C})$ be a graph collection on a vertex set V of size n and H a graph on n vertices satisfying the conditions of the theorem.

By assumption, $\delta n \leq e(H) = |\mathscr{C}| \leq \Delta n$. Let $m := \lfloor n/r \rfloor$. The Hajnal-Szemerédi theorem implies that there is a partition $V(H) = A_1 \cup \ldots \cup A_r$ into parts of size m and m+1 such that all edges go between different parts. Let $V = V_1 \cup \ldots \cup V_r$ be a random partition of V with $|V_i| = |A_i|$ for all $i \in [r]$, and let $\mathcal{V} := \{V_1, \ldots, V_r\}$. We claim that with high probability, the following hold for any $ij \in E(K_r) = {[r] \choose 2}$:

- (i) for all $V'_h \subseteq V_h$ with $|V'_h| \ge \eta^{1/4} |V_h|$ for h = i, j and $\mathscr{C}' \subseteq \mathscr{C}$ with $|\mathscr{C}'| \ge \eta^{1/4} |\mathscr{C}|$, we have $\sum_{c \in \mathscr{C}'} e_{G_c}(V'_i, V'_j) \ge d|\mathscr{C}'||V'_i||V'_j|/2;$
- (ii) for each labelling $\{g, h\} = \{i, j\}$ and $v \in V_g$, we have $\sum_{c \in \mathscr{C}} d_{G_c}(v, V_h) \ge \alpha^2 |V_h| |\mathscr{C}|/6$;
- (iii) $e(G_c) \ge \alpha^2 |V_i| |V_j|/6$ for all $c \in \mathscr{C}$.

For (i), we have

$$\eta n^3 < (d/2) \cdot \eta^{3/4} \delta n^3 / (2r)^2 < d\eta^{3/4} \delta n m^2 / 2 \le d(\eta^{1/4})^3 |\mathscr{C}| |V_i| |V_j| / 2 \le d|\mathscr{C}'| |V_i'| |V_j'| / 2.$$

The statement then follows directly from the definition of (d, η) -dense. To prove (ii), for each vertex $v \in V$, there are at least $\alpha |\mathscr{C}|/2$ colours $c \in \mathscr{C}$ for which $d_{G_c}(v) \geq \alpha n/2$, and a Chernoff-type bound implies that, with high probability $d_{G_c}(v, V_j) \geq \alpha |V_j|/3$. Part (iii) is proved similarly by swapping colour and vertex.

Parts (i)–(iii) imply that the 3-graph $G^{(3),ij}$ of $\mathbf{G}^{ij} := (G_c[V_i, V_j] : c \in \mathscr{C})$ is $(\eta^{1/4}, \alpha^2/6)$ -half-superregular. Lemma 2.1 and our choice of parameters implies that every $G^{(3),ij}$ contains a spanning subhypergraph $J^{(3),ij}$ which is $(\varepsilon, \alpha^4/72)$ -superregular. Thus $\mathcal{F} := (\mathcal{V}, \mathscr{C}, \mathbf{J})$ is a super K_r -template with parameters $(m, \varepsilon, \alpha^4/72, \delta)$, where \mathbf{J}^{ij} is the graph collection whose 3-graph is $J^{(3),ij}$ and $\mathscr{C}_{ij} = \mathscr{C}$ for all $ij \in E(K_r)$.

Let $d_{ij} := e(H[A_i, A_j])/n$ for all $ij \in E(K_r)$. We have $\sum_{ij} d_{ij} = e(H)/n \ge \delta$. Thus there is at least one ij such that $d_{ij} \ge \delta_{\binom{r}{2}+1}$. By the pigeonhole principle, there is $\ell \in \binom{r}{2}$ such that for all ij, either $d_{ij} \le \delta_{\ell}$, or $d_{ij} \ge \delta_{\ell+1}$. Let $P^{<} := \{ij \in \binom{[r]}{2} : d_{ij} \le \delta_{\ell}\}$. We will embed those sparse $H[A_i, A_j]$, with indices ij in $P^{<}$, using Lemma 3.1 (Embedding lemma with target and candidate sets), while the remaining (dense) pairs will be embedded using Theorem 3.5 (Transversal blow-up lemma) with target sets from the initial embedding of sparse pairs.

Let X be the union of non-isolated vertices in $H[A_i, A_j]$ over all $ij \in P^<$, let $Y := N_H(X) \setminus X$, and let $H^< := H[X \cup Y] \setminus H[Y]$, whose edge set is precisely those edges of H incident to X. We have $v(H^<) \leq 2\binom{r}{2}\delta_\ell n \leq \sqrt{\delta_\ell}m$. Apply Lemma 3.1 to \mathcal{F} with $m, \varepsilon, \sqrt{\delta_\ell}, \nu', 1, \alpha^4/72, \delta, \Delta, r$ playing the roles of $m, \varepsilon, \gamma, \nu', \nu, d, \delta, \Delta, r$, and $T_w := V_i$ for all $w \in V(H^<) \cap A_i$ and $i \in [r]$, to find injective maps $\tau : X \to V$ and $\sigma : E(H^<) \to \mathscr{C}$ such that $\tau(x)\tau(x') \in J_{\sigma(xx')}$ for all $xx' \in E(H[X])$, for all $i \in [r]$ we have $\tau(x) \in V_i$ for all $x \in A_i$, and for all $y \in Y \cap A_i$ there exists $C_y \subseteq V_i \setminus \tau(X)$ such that $C_y \subseteq \bigcap_{x \in N_{H^<}(y) \cap X} N_{G_{\sigma(xy)}}(\tau(x))$ and $|C_y| \geq \nu'm$.

Let $\mathscr{C}' := \mathscr{C} \setminus \sigma(E(H^{\leq}))$ and let $\mathcal{V}' := \{V'_1, \ldots, V'_r\}$ where $V'_i := V_i \setminus \tau(X)$ for each $i \in [r]$. By Lemma 2.9(ii) applied with $\alpha = \Delta \sqrt{\delta_\ell}$, the template $(\mathcal{V}', \mathscr{C}', \mathbf{J}')$ induced by $\mathcal{V}', \mathscr{C}'$ is super with parameters $(m/2, 2\varepsilon, \alpha^4/144, \delta/2)$. Let $H^{\geq} := H - X$. Note that H^{\geq} (with respect to its slightly smaller vertex set) is 2μ -separable. Now we randomly partition \mathscr{C}' into $\binom{r}{2}$ (some perhaps empty) parts $\mathscr{C}' = \bigcup_{ij \in E(K_r)} \mathscr{C}''_{ij}$ such that $|\mathscr{C}''_{ij}| = e(H^{\geq}[V'_i, V'_j])$ for all $ij \in \binom{[r]}{2}$. This is possible since $|\mathscr{C}'| = |\mathscr{C}| - e(H^{\leq}) = e(H) - e(H^{\leq}) = e(H^{\geq})$. Let $\mathcal{F}'' := (\mathcal{V}', \mathscr{C}'', \mathbf{J}')$, where each \mathscr{C}''_{ij} is as above (so $\mathscr{C}'' = \mathscr{C}'$). Each colour set is either large or empty, indeed, if $ij \in P^{\leq}$, then $\mathscr{C}''_{ij} = \emptyset$, but otherwise we have

$$|\mathscr{C}_{ij}''| \ge e(H[A_i, A_j]) - e(H^{<}) \ge \delta_{\ell+1}n - \Delta\sqrt{\delta_{\ell}}m \ge \delta_{\ell+1}n/2 \ge \delta_{\ell+1}|\mathscr{C}'|/(2\Delta) \ge \delta_{\ell+1}^2|\mathscr{C}'|.$$

Lemma 2.9(iii) applied with $\delta_{\ell+1}^2$, 1 playing the roles of α, k implies that \mathcal{F}'' is a rainbow super template with parameters ($\delta_{\ell+1}^2 m/2, \sqrt{\varepsilon}, \alpha^9, \delta/2$). Apply Theorem 3.5 (transversal blow-up lemma) with target sets C_y of size at least $\nu'm$ for the at most $\sqrt{\delta_\ell}m \leq \delta_\ell^{1/3}\delta_{\ell+1}^2m/2$ vertices $y \in Y$, and parameters $\delta_{\ell+1}^2m/2, \sqrt{\varepsilon}, 2\mu, \delta_1^{1/3}, \nu', \alpha^9, \delta/2$ playing the roles of $m, \varepsilon, \mu, \alpha, \nu, d, \delta$ to obtain a transversal embedding of $H^>$ inside \mathcal{F}'' such that every $y \in Y$ is embedded inside C_y . Together with the embedding of $H^<$, this gives a transversal embedding of H. Proof of Theorem 1.6. Let $\Delta, d, \alpha > 0$ be given. Without loss of generality, we may assume that $\alpha \ll d, 1/\Delta$. Choose constants $\eta', \mu, n_0 > 0$ such that the conclusion of Theorem 1.2 holds, applied with $\Delta, d, 1/5, \alpha^3$ playing the roles of $\Delta, d, \delta, \alpha$. Let $\eta := \eta'/(2\Delta)$.

Let $n \ge n_0$ be an integer and let G be a (d, η) -dense 3-graph on n vertices with $d_G(v) \ge \alpha n^2$ for all $v \in V(G)$.

Let *H* be a μ -separable graph with $\Delta(H) \leq \Delta$ and $v(H) + e(H) \leq n$. First, if *H* is small, we will enlarge it, as follows. If e(H) > n/4 - 1, let H' := H. Otherwise, obtain *H'* from *H* by successively adding an edge between isolated vertices until e(H) > n/4 - 1. Let *H'* be the obtained graph restricted to non-isolated vertices. We have $e(H') \leq n/4$, which implies that $v(H') + e(H') \leq 4e(H') \leq n$.

In both cases, we have $\Delta(H') \leq \Delta$, and H' is μ -separable; e(H') > n/4 - 1 and $v(H') \geq 3e(H')/\Delta > 3(n-4)/(4\Delta) > n/(2\Delta)$; and $v(H') + e(H') \leq n$.

Let V, \mathscr{C} be disjoint subsets of V(G) chosen uniformly at random subject to

$$|V| = v(H') > n/(2\Delta)$$
 and $|\mathscr{C}| = e(H') > n/4 - 1.$

Standard Chernoff-type bounds imply that, with high probability, for every $v \in V$ we have

$$\begin{split} e(G[\{v\},V,\mathscr{C}]) \geq \alpha^2 |V||\mathscr{C}|/6 \geq \alpha^2 n^2/(49\Delta) \geq \alpha^3 n^2 \quad \text{for all } v \in V \\ |\{xyc \in E(G) : x, y \in V\}| \geq \alpha^2 |V|^2/12 \geq \alpha^2 n^2/(48\Delta^2) \geq \alpha^3 n^2 \quad \text{for all } c \in \mathscr{C}. \end{split}$$

For each $c \in \mathscr{C}$, let G_c be the 2-graph with vertex set V and edge set $\{xy : x, y \in V \text{ and } xyc \in E(G)\}$, and let $\mathbf{G} := (G_c : c \in \mathscr{C})$. Thus $\sum_{c \in \mathscr{C}} d_{G_c}(v, V) = e(G[\{v\}, V, \mathscr{C}]) \geq \alpha^3 n^2$, and for every $c \in \mathscr{C}$ we have $e(G_c) \geq \alpha^3 n^2$. Since G is (d, η) -dense and $|V| \geq n/(2\Delta)$, we have that \mathbf{G} is (d, η') -dense. Theorem 1.2 applied with $\Delta, d, 1/5, \alpha^3$ playing the roles of $\Delta, d, \delta, \alpha$ implies that \mathbf{G} contains a transversal copy of H' and thus a transversal copy of H, that is, there are injective maps $\tau : V(H) \to V$ and $\sigma : E(H) \to \mathscr{C}$ so that $\tau(x)\tau(y) \in G_{\sigma(xy)}$ for every $xy \in E(H)$. Define $\rho : V(H) \cup E(H) \to V(G)$ by setting $\rho(x) := \tau(x)$ for $x \in V(H)$ and $\rho(e) := \sigma(e)$ for $e \in E(H)$. Then ρ is injective and $\rho(x)\rho(y)\rho(xy) \in E(G)$ for all $xy \in E(H)$; that is, ρ defines a copy of the 1-expansion of H in G, as required.

6 Concluding remarks

In this paper, we have proved a transversal blow-up lemma that embeds separable graphs into graph collections, which can be used to apply the regularity-blow-up method to transversal embedding problems. We conclude with some remarks on future directions.

Separability. In our proof of the transversal blow-up lemma, the separability condition is necessary because we need to divide the graph into linear-sized pieces and then use the blow-up lemma to embed them piece by piece. For this embedding process to work, the number of edges between different pieces should not be too large. We wonder if our transversal blow-up lemma can be generalised to embed any graph with bounded maximum degree. If such a version can be proven, it would directly generalise the original blow-up lemma for graphs (since a collection of identical superregular pairs is a superregular collection).

Future applications to transversal embedding. In a subsequent paper, we will utilise the transversal blow-up technique developed in this paper in combination with the absorption method to provide a new proof of the transversal version of the approximate Pósa-Seymour conjecture,

which was recently established in [23], and is a special case of the yet more recent main result of [9]. Additionally, we will prove a stability result for transversal Hamilton cycles that has not been demonstrated before.

However, using our method to obtain transversal versions of embedding results proved using the regularity blow-up method does not seem quite as straightforward as simply following the same proof. The regularity lemma produces some exceptional vertices which need to be incorporated into the structure built between vertex clusters: they are 'absorbed' by clusters. For transversal embedding of a spanning graph H, one needs to insert the exceptional vertices as well as some exceptional colours into the structure built between vertex clusters and colour clusters. Such an insertion may make some new colours exceptional. Right at the end of the process, inserting vertices and colours simultaneously becomes difficult, when there are no more 'spare colours'. Thus, we need to construct an absorption set for the remaining vertices and colours prior to embedding H.

Applications to hypergraph embedding. We state the full 3-graph version of our transversal blow-up lemma.

Theorem 6.1 (weak 3-graph blow-up lemma). Let $0 < 1/m \ll \varepsilon, \mu, \alpha \ll \nu, d, \delta, 1/\Delta, 1/r \le 1$.

- Let R be a 2-graph with vertex set [r].
- Let G be a 3-graph with parts V_1, \ldots, V_r and V_{ij} for $ij \in E(R)$ where $m \leq |V_i| \leq m/\delta$ for all $i \in [r]$, and $|V_{ij}| \geq \delta m$ for all $ij \in [r]$. Suppose that $G[V_i, V_j, V_{ij}]$ is weakly (ε, d) -(half-)superregular for all $ij \in E(R)$.
- Let H be a μ -separable 2-graph with $\Delta(H) \leq \Delta$ for which there is a graph homomorphism $\phi: V(H) \rightarrow V(R)$ with $|\phi^{-1}(i)| = |V_i|$ for all $i \in [r]$ and $e(H[\phi^{-1}(i), \phi^{-1}(j)]) = |V_{ij}|$ for all $i \in E(R)$.
- Suppose that for each $i \in [r]$ there is a set $U_i \subseteq \phi^{-1}(i)$ with $|U_i| \leq \alpha m$ and $T_x \subseteq V_{\phi(x)}$ with $|T_x| \geq \nu m$ for all $x \in U_i$.

Then G contains a copy of the 1-expansion of H, where each vertex $x \in V(H)$ is mapped to $V_{\phi(x)}$ and the new vertex for $xy \in E(H)$ is mapped to $V_{\phi(x)\phi(y)}$; and moreover, for every $i \in [r]$ every $x \in U_i$ is mapped to T_x .

This is a reformulation of Theorem 1.8 since for each $ij \in E(R)$, the 3-graph $G_{ij}^{(3)}$ of $(G_c : c \in \mathscr{C}_{ij})$ in Theorem 1.8 is weakly half-superregular, so we simply take $V_{ij} := \mathscr{C}_{ij}$. It would be interesting to extend Theorem 1.6 on embeddings in uniformly dense 3-graphs beyond 1-expansions of separable 2graphs. It is not clear which 3-graphs one should expect to be able to embed in a weakly superregular triple (or in a uniformly dense 3-graph). Even the case of 3-partite 3-graphs seems difficult; that is, to extend Theorem 1.7 (simplified weak hypergraph blow-up lemma).

Problem 6.2. Given $0 < 1/n \ll \varepsilon \ll d \ll \delta \leq 1$, which J are subhypergraphs of any weakly (ε, d) -superregular triple $G[V_1, V_2, V_3]$, where $\delta n \leq |V_1|, |V_2|, |V_3| \leq n/\delta$?

A 3-partite 3-graph J is equivalent to an edge-coloured bipartite 2-graph \mathcal{J} (where each edge can receive multiple colours) using our usual identification of one part with a set of colours. If J has $\Delta(J) \leq \Delta$, then $\Delta(\mathcal{J}) \leq \Delta$ and the colouring is Δ -bounded. (If J is linear, then the edge-colouring is proper.) Thus we would like to extend Theorem 1.3 (simplified weak transversal blow-up lemma). to find an embedding of \mathcal{J} with a given edge-colouring up to permutation of colours. If every colour only appears on a single edge, that is, J is an expansion of some 2-graph H where the 2-edge e is replaced by some bounded number $t_e \geq 1$ of 3-edges (which is not linear if some $t_e > 1$), it seems plausible that our techniques would work. However, the colour absorption we use from [36] does not seem to be able to deal with colours playing multiple roles. Let F be a 3-partite 3-graph of fixed size. Recall that an old result of Erdős [18] implies that any large 3-graph of positive density contains a copy of F. However, the results of [14] imply that any large uniformly dense 3-graph G (whose number of vertices is a multiple of v(F) and where every vertex sees a positive fraction of pairs) contains an F-factor if and only if there is a vertex $v^* \in V(F)$ such that for any two edges e, e' where e contains v^* and e' does not, e and e' share at most one vertex. Probably, the characterisation for weakly superregular triples with balanced classes is the same.

Another instructive case is the tight Hamilton cycle (whose number of vertices is divisible by 6). As a graph collection problem, this corresponds to finding a (2-graph) Hamilton cycle on 4n vertices with 2n colours, where, cyclically labelling the edges e_1, \ldots, e_{4n} , the consecutive edges $e_{2c-1}, e_{2c}, e_{2c+1}$ are all in G_c , for each $c \in [2n]$ where indices are taken modulo 4n. A construction in [34] (see also [35]) shows that a weakly superregular triple need not contain a tight Hamilton cycle, even if its density is close to $\frac{1}{8}$.

Example 6.3. Let V be a vertex set of size 6n, and let $V = V_1 \cup V_2 \cup V_3$ be a partition, where V_1, V_2, V_3 have equal size 2n, and let $X \subseteq V$ where $|X \cap V_1|$ is odd. Independently for each distinct $i, j \in [3]$ add uniform random edges between parts V_i, V_j with probability $\frac{1}{2}$, to obtain a 3-partite 2-graph J. Now form a 3-partite 3-graph G by, for each triple $abc \in A \times B \times C$, as follows:

- add *abc* to G if $|\{a, b, c\} \cap X|$ is even and $\{a, b, c\}$ spans a triangle in J.
- add *abc* to G if $|\{a, b, c\} \cap X|$ is odd and $\{a, b, c\}$ spans an independent set in J.

Suppose $v_1^1 v_2^1 v_3^1 \dots v_1^{2n} v_2^{2n} v_3^{2n}$ is a tight cycle H in J, where without loss of generality, $v_i^j \in V_i$ for all j. For any $1 \leq j < 2n$, Since $v_1^j v_2^j v_3^j, v_2^j v_3^j v_1^{j+1}$ are edges of H, we have that $|\{v_1^j, v_2^j, v_3^j\} \cap X|$, $|\{v_1^{j+1}, v_2^j, v_3^j\} \cap X|$ are both odd or both even. This implies that v_1^j, v_1^{j+1} are either both in X or neither in X. By considering the disjoint pairs $v_1^1 v_1^2, v_1^3 v_1^4, \dots, v_1^{2n-1} v_1^{2n}$, this is a contradiction to $|X \cap V_1|$ odd.

It can easily be checked using Chernoff bounds (see [34]) that, if every $|X \cap V_i|$ has size about n, then with high probability G is weakly $(\frac{1}{8}, o(1))$ -superregular.

References

- [1] Sarmad Abbasi. Spanning subgraphs of dense graphs and a combinatorial problem on strings. Rutgers The State University of New Jersey, School of Graduate Studies, 1998.
- [2] Ron Aharoni, Matt DeVos, Sebastián González Hermosillo de la Maza, Amanda Montejano, and Robert Šámal. A rainbow version of Mantel's theorem. *Advances in Combinatorics*, 2, 2020.
- [3] Ron Aharoni, Agelos Georgakopoulos, and Philipp Sprüssel. Perfect matchings in r-partite r-graphs. European Journal of Combinatorics, 30(1):39–42, 2009.
- [4] Elad Aigner-Horev, David Conlon, Hiêp Hàn, Yury Person, and Mathias Schacht. Quasirandomness in hypergraphs. *Electronic Journal of Combinatorics*, 25(3):P3.34, 2018.
- [5] Elad Aigner-Horev and Gil Levy. Tight Hamilton cycles in cherry-quasirandom 3-uniform hypergraphs. Combinatorics, Probability and Computing, 30(3):412–443, 2021.
- [6] Pedro Araújo, Simón Piga, and Mathias Schacht. Localised codegree conditions for tight Hamilton cycles in 3-uniform hypergraphs. arXiv:2005.11942, 2020.

- [7] Julia Böttcher, Mathias Schacht, and Anusch Taraz. Spanning 3-colourable subgraphs of small bandwidth in dense graphs. Journal of Combinatorial Theory, Series B, 98(4):752–777, 2008.
- [8] Julia Böttcher, Mathias Schacht, and Anusch Taraz. Proof of the bandwidth conjecture of Bollobás and Komlós. *Mathematische Annalen*, 343(1):175–205, 2009.
- [9] Debsoumya Chakraborti, Seonghyuk Im, Jaehoon Kim, and Hong Liu. A bandwidth theorem for graph transversals. arXiv:2302.09637, 2023.
- [10] Yangyang Cheng, Jie Han, Bin Wang, Guanghui Wang, and Donglei Yang. Rainbow Hamilton cycle in hypergraph systems. arXiv:2111.07079, 2021.
- [11] Yangyang Cheng, Guanghui Wang, and Yi Zhao. Rainbow pancyclicity in graph systems. *The Electronic Journal of Combinatorics*, 28(3), 2021.
- [12] Fan Chung. Regularity lemmas for hypergraphs and quasi-randomness. Random Structures & Algorithms, 2:241 – 252, 06 2007.
- [13] David Conlon, Hiêp Hàn, Yury Person, and Mathias Schacht. Weak quasi-randomness for uniform hypergraphs. *Random Structures & Algorithms*, 40(1):1–38, 2012.
- [14] Laihao Ding, Jie Han, Shumin Sun, Guanghui Wang, and Wenling Zhou. F-factors in quasirandom hypergraphs. Journal of the London Mathematical Society, 106(3):1810–1843, 2022.
- [15] Oliver Ebsen, Giulia S Maesaka, Christian Reiher, Mathias Schacht, and Bjarne Schülke. Embedding spanning subgraphs in uniformly dense and inseparable graphs. *Random Structures & Algorithms*, 57(4):1077–1096, 2020.
- [16] Stefan Ehard, Stefan Glock, and Felix Joos. A rainbow blow-up lemma for almost optimally bounded edge-colourings. In *Forum of Mathematics, Sigma*, volume 8, page e37, 2020.
- [17] Mohamed H El-Zahar. On circuits in graphs. Discrete mathematics, 50:227–230, 1984.
- [18] Paul Erdős. On extremal problems of graphs and generalized graphs. Israel Journal of Mathematics, 2(3):183–190, 1964.
- [19] Paul Erdős and András Hajnal. On Ramsey like theorems, problems and results. In Combinatorics (Proc. Conf. Combinatorial Math., Math. Inst., Oxford, 1972), pages 123–140. Citeseer, 1972.
- [20] Paul Erdős and Vera T. Sós. On Ramsey-Turán type theorems for hypergraphs. Combinatorica, 2:289–295, 1982.
- [21] Luyining Gan and Jie Han. Hamiltonicity in cherry-quasirandom 3-graphs. *European Journal* of Combinatorics, 102:103457, 2022.
- [22] Stefan Glock and Felix Joos. A rainbow blow-up lemma. Random Structures & Algorithms, 56(4):1031–1069, 2020.
- [23] Pranshu Gupta, Fabian Hamann, Alp Müyesser, Olaf Parczyk, and Amedeo Sgueglia. A general approach to transversal versions of Dirac-type theorems. arXiv:202209.09289, 2022.

- [24] Hiêp Hàn, Jie Han, and Patrick Morris. Factors and loose Hamilton cycles in sparse pseudorandom hypergraphs. Random Structures & Algorithms, 61(1):101–125, 2022.
- [25] Felix Joos and Jaehoon Kim. On a rainbow version of Dirac's theorem. Bulletin of the London Mathematical Society, 53(2):498–504, 2020.
- [26] Peter Keevash. A hypergraph blow-up lemma. Random Structures & Algorithms, 39(3):275–376, 2011.
- [27] Yoshiharu Kohayakawa, Brendan Nagle, Vojtěch Rödl, and Mathias Schacht. Weak hypergraph regularity and linear hypergraphs. *Journal of Combinatorial Theory, Series B*, 100(2):151–160, 2010.
- [28] János Komlós, Gábor N Sárközy, and Endre Szemerédi. Blow-up lemma. Combinatorica, 17:109– 123, 1997.
- [29] János Komlós, Gábor N Sárközy, and Endre Szemerédi. Proof of the Seymour conjecture for large graphs. Annals of Combinatorics, 2:43–60, 1998.
- [30] Daniela Kühn and Deryk Osthus. Loose hamilton cycles in 3-uniform hypergraphs of high minimum degree. Journal of Combinatorial Theory, Series B, 96(6):767–821, 2006.
- [31] Daniela Kühn and Deryk Osthus. The minimum degree threshold for perfect graph packings. Combinatorica, 29(1):65–107, 2009.
- [32] Daniela Kühn, Deryk Osthus, and Timothy Townsend. Fractional and integer matchings in uniform hypergraphs. *European Journal of Combinatorics*, 38:83–96, 2014.
- [33] John Lenz and Dhruv Mubayi. The poset of hypergraph quasirandomness. *Random Structures* & Algorithms, 46(4):762–800, 2015.
- [34] John Lenz and Dhruv Mubayi. Perfect packings in quasirandom hypergraphs I. Journal of Combinatorial Theory, Series B, 119:155–177, 2016.
- [35] John Lenz, Dhruv Mubayi, and Richard Mycroft. Hamilton cycles in quasirandom hypergraphs. Random Structures & Algorithms, 49(2):363–378, 2016.
- [36] Richard Montgomery, Alp Müyesser, and Yani Pehova. Transversal factors and spanning trees. Advances in Combinatorics, 2022.
- [37] Christian Reiher, Vojtěch Rödl, and Mathias Schacht. Hypergraphs with vanishing Turán density in uniformly dense hypergraphs. Journal of the London Mathematical Society, 97(1):77– 97, 2018.
- [38] Christian Reiher, Vojtěch Rödl, and Mathias Schacht. On a Turán problem in weakly quasirandom 3-uniform hypergraphs. Journal of the European Mathematical Society, 20(5):1139–1159, 2018.
- [39] Vojtěch Rödl and Andrzej Ruciński. Perfect matchings in ε -regular graphs and the blow-up lemma. *Combinatorica*, 19(3):437–452, 1999.
- [40] Katherine Staden and Andrew Treglown. The bandwidth theorem for locally dense graphs. In Forum of Mathematics, Sigma, volume 8, page e42, 2020.

- [41] Endre Szemerédi. Regular partitions of graphs. Technical report, Stanford Univ Calif Dept of Computer Science, 1975.
- [42] Timothy Townsend. Extremal problems on graphs, directed graphs and hypergraphs. PhD thesis, University of Birmingham, 2016.
- [43] Henry Towsner. σ -algebras for quasirandom hypergraphs. Random Structures & Algorithms, 50(1):114-139, 2017.

Appendix Α

Weak hypergraph regularity lemmas A.1

We first introduce the following special form of the weak hypergraph regularity lemma for k-partite k-graphs. Its proof follows easily from the original lemma of Chung [12], which is very similar to the 2-graph regularity lemma of Szemerédi [41].

Lemma A.1. For every $L_0, k \ge 1$ and every $\varepsilon, \delta > 0$, there is an $n_0 > 0$ such that for every kpartite k-graph G on $n \ge n_0$ vertices with parts V_1, \ldots, V_k where $\delta n \le |V_i| \le n/\delta$, there exists a refined partition $V_i = \bigcup_{j=1}^{t_i} V_{i,j}$ for each $i \in [k]$ such that the following properties hold:

- (i) $L_0 \leq t := t_1 + \ldots + t_k \leq n_0;$
- (ii) $||V_{i,j}| |V_{i',j'}|| \le 1$ for any $i, i' \in [k], j \in [t_i]$ and $j' \in [t_{i'}];$ (iii) all but at most εt^k k-tuples $(V_{1,j_1}, \ldots, V_{k,j_k})$ are ε -regular.

The following weak hypergraph regularity lemma [32] is proved in the same way as the original degree form of the regularity lemma (see [42]), which in turn can be derived from the standard regularity lemma via some cleaning.

Theorem A.2 (Degree form of the weak hypergraph regularity lemma). For all integers $L_0 \geq 1$ and every $\varepsilon > 0$, there is an $n_0 = n_0(\varepsilon, L_0)$ such that for every $d \in [0, 1)$ and for every 3-graph G = (V, E)on $n \ge n_0$ vertices there exists a partition of V into V_0, V_1, \ldots, V_L and a spanning subhypergraph G' of G such that the following properties hold:

- (i) $L_0 \leq L \leq n_0$ and $|V_0| \leq \varepsilon n$;
- (*ii*) $|V_1| = \ldots = |V_L| =: m;$
- (iii) $d_{G'}(v) > d_G(v) (d + \varepsilon)n^2$ for all $v \in V$;
- (iv) every edge of G' with more than one vertex in a single cluster V_i for some $i \in [L]$ has at least one vertex in V_0 ;
- (v) for all triples $\{h, i, j\} \in {[L] \choose 3}$, we have that $(V_h, V_i, V_j)_{G'}$ is either empty or (ε, d) -regular.

The proof of Lemma 2.5 (the regularity lemma for graph collections) is a routine but tedious consequence of this theorem applied to the 3-graph $G^{(3)}$ of a graph collection G.

Proof of Lemma 2.5. By increasing L_0 and decreasing ε, δ as necessary, we may assume without loss of generality that $0 < 1/L_0 \ll \varepsilon \ll \delta \ll 1$. Let $L'_0 := 2L_0/\delta$. Choose new constants ε', α , which we may assume satisfy $0 < 1/L'_0 \ll \varepsilon' \ll \alpha \ll \varepsilon$. Let n'_0 be obtained from Theorem A.2 applied with parameters ε', L'_0 . We may assume that $1/n'_0 \ll 1/L'_0$ by increasing n'_0 . Altogether,

$$0 < 1/n'_0 \ll 1/L'_0 \ll \varepsilon' \ll \alpha \ll \varepsilon \ll \delta \ll 1.$$

Let $n_0 := 2n'_0/\alpha$. Let $n \ge n_0$ be an integer and suppose that **G** is a graph collection on a vertex set V of size n and with colour set \mathscr{C} , where $\delta n \leq |\mathscr{C}| \leq n/\delta$. Let $d \in [0,1)$ and let $G^{(3)}$ be the 3-graph of **G** with vertex set $U := V \cup \mathscr{C}$. Theorem A.2 implies that there is a partition U_0, U_1, \ldots, U_K of U and a spanning subhypergraph $G^{(3)'}$ of $G^{(3)}$ such that

- (a) $L'_0 \le K \le n'_0$ and $|U_0| \le \varepsilon' |U|$; (b) $|U_1| = \ldots = |U_K| =: m'$;
- (c) $d_{G^{(3)'}}(u) > d_{G^{(3)}}(u) (2d + \varepsilon')|U|^2$ for all $u \in U$;
- (d) every edge of $G^{(3)'}$ with more than one vertex in a single cluster U_i has at least one vertex in $U_0;$

(e) for all triples $\{h, i, j\} \in {\binom{[K]}{3}}$, we have that $(U_h, U_i, U_j)_{G^{(3)'}}$ is either empty or $(\varepsilon', 2d)$ -regular. Partition each cluster $U_i, i \in [K]$, into $1/\alpha$ subclusters of size at most m so that all but at most two subclusters of U_i have size exactly m and the property that they lie entirely within V, or within \mathscr{C} . If a subcluster does not have this property, add it to U_0 . The new exceptional set has size at most $|U_0| + 2\alpha m' K$, and we let V_0 be its intersection with V, and \mathscr{C}_0 its intersection with \mathscr{C} . Relabel the subclusters so that those which are subsets of V are V_1, \ldots, V_L and those which are subsets of \mathscr{C} are $\mathscr{C}_1, \ldots, \mathscr{C}_M$. Let G'_c be the graph with vertex set V and edge set $\{xy : xyc \in G^{(3)'}\}$ for all $c \in \mathscr{C}$. We claim that the properties of the lemma are satisfied.

For (i), we have m'K < |U| so

$$|V_0| + |\mathscr{C}_0| \le |U_0| + 2\alpha m' K \stackrel{(a)}{\le} (\varepsilon' + 2\alpha)|U| \le (\varepsilon' + 2\alpha)(n + n/\delta) \le \varepsilon n.$$

Also, $L_0 \leq L'_0 \leq L'_0 \leq K/\alpha \leq L + M \leq 2K/\alpha \leq 2n'_0/\alpha = n_0$, proving the required upper bound for both L and M. Furthermore, $m(L+M) \leq n + |\mathscr{C}|$ so

$$L \ge \frac{n - |V_0|}{m} \ge \frac{(n - |V_0|)(L + M)}{n + |\mathscr{C}|} \ge \frac{(n - |V_0|)L'_0}{n + n/\delta} \ge \delta L'_0/2 = L_0$$

and similarly for M. Part (ii) follows by construction.

For (iii), we have $\sum_{c \in \mathscr{C}} d_{G'_c}(v) = d_{G^{(3)'}}(v)$ for all $v \in V$, and $e(G'_c) = d_{G^{(3)'}}(c)$ for all $c \in \mathscr{C}$, and similarly for G_c . Further, $(2d + \varepsilon')(n + |\mathscr{C}|)^2 < (2d + \varepsilon')(1 + 1/\delta)^2 n^2 \leq (3d/\delta^2 + 2\varepsilon'/\delta^2)n^2 < (3d/\delta^2 + 2\varepsilon'/\delta^2)n^2 > (3d/\delta^2 + 2\varepsilon'$ $(3d/\delta^2 + \varepsilon)n^2$. Thus (c) implies the required.

For (iv), if G'_c has an edge xy with $x, y \in V_i$ for some $i \in [L]$, then $x, y \in U_{i'}$ for the cluster $U_{i'}$ containing V_i . Since $xyc \in E(G^{(3)'})$, (d) implies that $c \in U_0$. Thus $c \in \mathscr{C}_0$, as required.

For (v), suppose that $(\{h, i\}, j) \in {\binom{[L]}{2}} \times M$ and $\mathbf{G}'_{hi,j}$ is non-empty. Let $h', i', j' \in [K]$ be such that $V_h \subseteq U_{h'}, V_i \subseteq U_{i'}$ and $\mathscr{C}_j \subseteq U_{j'}$. Since $G'_{hi,j}$ is non-empty, we have that $(U_{h'}, U_{i'}, U_{j'})_{G^{(3)'}}$ is non-empty. So h', i', j' are distinct by (d). Part (e) implies that $(U_{h'}, U_{i'}, U_{j'})_{G^{(3)'}}$ is $(\varepsilon', 2d)$ -regular. Therefore, for each $c \in U_{i'}$, letting J_c be the bipartite graph with partition $(U_{h'}, U_{i'})$ and edge set $\{xy: x \in U_{h'}, y \in U_{i'}, xyc \in G^{(3)'}\}$, we have that $J := (J_c: c \in U_{j'})$ is $(\varepsilon', 2d)$ -regular. Lemma 2.4(i) (the slicing lemma) implies that $G'_{hi,i}$ is $(\varepsilon'/\alpha, d)$ -regular and thus (ε, d) -regular. This completes the proof.

We conclude the appendix with a proof of Lemma 2.1, which states that a half-superregular (k-graph) k-tuple contains a spanning superregular subhypergraph. The proof is the same as the one in [39] for k = 2.

Proof of Lemma 2.1. Choose a new parameter ε'' such that $0 < \varepsilon \ll \varepsilon'' \ll \varepsilon'$. Apply Lemma A.1 to G with parameters $L_0 := 1, \varepsilon'', \delta$ to obtain $n_0 > 0$. Increasing n_0, n and decreasing ε if necessary, we may assume that $1/n \ll \varepsilon \ll 1/n_0 \ll \varepsilon''$. Now let $G = (V_1, \ldots, V_k)_G$ be an (ε, d) -half-superregular kgraph with $\delta n \leq |V_i| \leq n/\delta$ for all $i \in [k]$. Lemma A.1 implies that there is a refinement $V_i = \bigcup_{j=1}^{t_i} V_{i,j}$ for each $i \in [k]$ such that $t := t_1 + \ldots + t_k \leq n_0$, every pair of subparts differ in size by at most one, and all but at most $\varepsilon'' t^3$ k-tuples $(V_{1,j_1}, \ldots, V_{k,j_k})$ are ε'' -regular.

Obtain a spanning subhypergraph G' of G by, for each ε'' -regular triple (X_1, \ldots, X_k) of subparts, removing every k-edge in (X_1, \ldots, X_k) independently at random with probability 1 - d/d', where d'is the density of (X_1, \ldots, X_k) . (Note that by $|X_i| \ge |V_i|/n_0 \ge \varepsilon |V_i|$ for each $i \in [k]$, we have $d' \ge d$ due to half-superregularity.) We claim that, with high probability,

- $d_{G'}(v) \ge d^2(|V_1| \dots |V_k|)/(2|V_i|)$ for any $i \in [k]$ and $v \in V_i$,
- any ε'' -regular k-tuple (of subparts) in G is $2\varepsilon''$ -regular in G',
- any ε'' -regular k-tuple (of subparts) in G has density $d \pm 2\varepsilon''$ in G'.

This follows from Chernoff bounds.

Now we claim that, given that the above hold, G' is $(\varepsilon', d^2/2)$ -superregular. We only need to check that G' is $(\varepsilon', d^2/2)$ -regular. Let (A_1, \ldots, A_k) be any k-tuple where $A_i \subseteq V_i$ and $|A_i| \ge \varepsilon' |V_i|$ for all $i \in [k]$. For each $i \in [k]$, we have partition $A_i = \bigcup_{j=1}^{t_i} (A_i \cap V_{i,j})$. We say a part $A_{i,j} := A_i \cap V_{i,j}$ is big if $|A_i \cap V_{i,j}| \ge \varepsilon'' |V_{i,j}|$ and otherwise it is small. Note that in total we have at most $\varepsilon''(|V_1| + \ldots + |V_k|) \le \varepsilon'' kn/\delta$ vertices in small parts. Also note that we have at most $\varepsilon'' t^k$ non-regular k-tuples that contain in total at most $\varepsilon''(n/\delta)^k$ edges. Thus

$$e_{G'}(A_1,\ldots,A_k) = \sum_{\substack{(A_{1,j_1},\ldots,A_{k,j_k}) \text{ all big and} \\ (V_{1,j_1},\ldots,V_{k,j_k}) \in \mathcal{C}'' \text{ regular}}} e_{G'}(A_{1,j_1},\ldots,A_{k,j_k}) \pm \left(2\varepsilon''kn/\delta \cdot (n/\delta)^{k-1} + \varepsilon''(n/\delta)^k\right).$$

For every summand index (j_1, \ldots, j_k) , we have $e_{G'}(A_{1,j_1}, \ldots, A_{k,j_k}) = (d \pm 2\varepsilon'')|A_{1,j_1}| \ldots |A_{k,j_k}|$ by $2\varepsilon''$ -regularity. Thus

$$e_{G'}(A_1,\ldots,A_k) = (d \pm 2\varepsilon'')|A_1|\ldots|A_k| \pm 3\varepsilon'' kn^k/\delta^k = (d \pm \varepsilon')|A_1|\ldots|A_k|.$$

This implies that G' is $(\varepsilon', d^2/2)$ -superregular and thus we finish the proof.