NORMED LINEAR SPACES WHICH ARE ISOMETRIC TO ORDER UNIT SPACES

ANIL KUMAR KARN

ABSTRACT. In this paper, we consider the linear direct sum of a real normed linear space with an order unit space and with a base normed space to obtain respectively a new order unit space and a new base normed space. As a consequence, we find that an ℓ_1 -space may be shown to be an order unit space. (However, not in the natural order.) Dually, an ℓ_{∞} -space may be shown to be a base normed space. To understand this aberration, we characterize the real normed linear spaces which are isometrically isomorphic to order unit spaces. We prove that other than the classical case of ℓ_{∞} , ℓ_1 is also isometrically isomorphic to an order unit space whereas ℓ_2 is not isometrically isomorphic to any order unit space.

1. INTRODUCTION

Alaoglu theorem roughly says that every normed linear space can be isometrically embedded into C(K) for some suitable compact Haudorff space K. However, in case of an order unit space, this embedding enhances to an order embedding. Even otherwise, order units play a very significant role in functional analysis in general and operator algebras in particular. For example, they sit on the interface of the commutative and the non-commutative theory. Therefore, discovering more classes of order unit spaces is always desirable.

It is a folklore that we can adjoin an order unit to any real normed linear space. Probably, this construction is due to M. M. Day [2, Theorem 1.6.1 (Alternative proof)]. Recently, in [6] the author studied the notion of absolute value in such spaces. Since the adjoining of an order unit to a real normed linear space is a one dimensional extension, such spaces also have a base normed structure. This aspect was studied by the author in [7]. In this paper, we generalize this construction. We consider the (linear) direct sum of a real normed linear space either with an order unit space or with a base normed space to construct a

²⁰²⁰ Mathematics Subject Classification. Primary: 46B40; Secondary: 46B20.

Key words and phrases. Norming order unit, adjoining a normed linear space to an order unit space.

ANIL KUMAR KARN

new order unit space and a new base normed space respectively. It is interesting to note that in the case of an order unit space the direct sum turns out to be an ℓ_1 -sum and in the case of a base normed space, it is an ℓ_{∞} -sum. As a consequence, we find that an ℓ_1 -space may be shown to be an order unit space. (However, not in the natural order.) Dually, an ℓ_{∞} -space may be shown to be a base normed space.

Thus it will interesting to know under what condition a normed linear space is isometrically isomorphic to an order unit space. In this paper, we address this question. We introduce the notion of *norming order units* in a normed linear space. We observe that the standard order unit in an order unit space is a norming order unit. More importantly, we show that if a norm one element in a normed linear space is a norming order unit, then it determines a (unique) order structure under which the space is an order unit space and the order unit norm is equal to the given norm. We examine some known (ℓ_{∞}) as well as some unknown (ℓ_1) cases. We also find an example of a normed linear space that does not possess any norming order unit.

2. Adjoining normed linear spaces with ordered normed linear spaces

In this section, we consider the (linear) direct sum of an ordered linear space with a normed linear space and obtain a new ordered linear space under two suitable conditions. We begin with a base normed space which is easier to handle.

Let (V, V^+) be a real ordered vector space. A convex subset $B \subset V^+$ is called a *base* of V^+ , if $0 \notin B$ and for each $u \in V^+$ with $u \neq 0$, there is a unique $b \in B$ and a real number k > 0 such that u = kb. If V^+ is generating, then the set $K := \operatorname{co}(B \bigcup -B)$ is an absolutely convex and absorbing set in V which determines a seminorm $\|\cdot\|_B$ in V. This is called the *base seminorm* generated by B. When $\|\cdot\|_B$ is a norm, (V, V^+, B) is called a *base normed space*. For a detailed discussion on such spaces, one may refer to any standard book on ordered vector spaces. We follow [1, 2, 9].

Theorem 2.1. Let (V, V^+, B) be a base normed space and let X be any real normed linear space. Then $(V_X(:= V \oplus_{\infty} X), V_X^+, B_X)$ is a base normed space for the cone $V_X^+ := \{(u, x) : u \in V^+ \text{ and } ||u|| \ge ||x||\}$ and the base $B_X := B \times X_1$. (Here X_1 is the closed unit ball of X.)

Proof. Note that $(0,0) \in V_X^+$. Let $(u,x), (v,y) \in V_X^+$. Then $u, v \in V^+$ and we have $||u|| \ge ||x||$ and $||v|| \ge ||y||$. As V^+ is a cone, we have

 $u + v \in V^+$. Further, as the base norm is additive in V^+ , we get

$$||u + v|| = ||u|| + ||v|| \ge ||x|| + ||y|| \ge ||x + y||$$

Thus $(u+v, x+y) \in V_X^+$. Now, if $k \in \mathbb{R}^+$, then $ku \in V^+$ and we have $||ku|| = k||u|| \ge k||x|| = ||kx||$

$$||ku|| = k||u|| \ge k||x|| = ||kx||$$

Thus $k(u, x) \in V_X^+$ so that V_X^+ is a cone. Also, as B and X_1 are convex subsets of V^+ and X respectively, we get that B_X is convex. Also, if $(u, x) \in B_X$, then $u \in B$ and $x \in X_1$. Thus ||u|| = 1 and $||x|| \leq 1$ so that $(u, x) \in V_X^+$ and subsequently, B_X is a convex subset of V_X^+ . We show that B_X is a base for V_X^+ . For this, let $(u, x) \in V_X^+$ with $(u, x) \neq (0, 0)$. Then $u \in V^+$ with $||u|| \geq ||x||$ so that $u \neq 0$. Since B is a base for V^+ , there is a unique $b \in B$ such that u = ||u||b. Put $z = ||u||^{-1}x$. Then $(b, z) \in B_X$ and (u, x) = ||u||(b, z). Hence B_X is a base for V_X^+ .

By definition, the base (semi) norm corresponding to the base B_X is gauge of $co(B_X \cup -B_X)$. As $-X_1 = X_1$, we get that

$$\operatorname{co}(B_X \cup -B_X) = \operatorname{co}(B \cup -B) \times X_1 = V_1 \times X_1 = (V \oplus_{\infty} X)_1.$$

Thus the base norm on V_X determined by B_X is given by

$$||(v,x)||_{\infty} := \max\{||v||, ||x||\}$$

for any $(v, x) \in V_X$.

Now we turn to the dual notion. Let (V, V^+) be a real ordered vector space and let $e \in V^+$. We say that e is an *order unit* for V, if for every $v \in V$ there exists $\lambda \in \mathbb{R}$ with $\lambda > 0$ such that $\lambda e \pm v \in V^+$. In this case, e determines a semi-norm $\|\cdot\|_e$ given by

$$||v||_e := \inf\{\lambda > 0 : \lambda e \pm v \in V^+\}$$

for all $v \in V$.

We say that V^+ is proper, if $V^+ \cap -V^+ = \{0\}$. Further, we say that V^+ is Archimedean, if for any $v \in V$ with $\lambda u + v \in V^+$ for a fixed $u \in V$ and all $\lambda > 0$, we have $v \in V^+$. Now, an order unit space is a real ordered vector space with an order unit e such that V^+ is proper and Archimedean. (In this case, $\|\cdot\|_e$ is a norm on V such that V^+ is norm-closed.) For details, we refer to [1, 2, 9], or for that matter, any standard book on ordered vector space is an order unit space and vice versa.

We noticed that the adjoining a normed linear space to an order unit space is a little more complicated. Let X be a real normed linear space and let (V, V^+, e) be an order unit space. Then $V' \oplus_{\infty} X'$ is a base

normed space as described in Theorem 2.1. Thus the expected cone V_X^+ in $V \oplus_1 X$ should be described as

 $V_X^+ = \{(u, x) \in V \oplus_1 X : \langle (\phi, f), (u, x) \rangle \ge 0 \text{ for all } \phi \in S(V) \text{ and } f \in X_1' \}.$ Let $(u, x) \in V_X^+$. Then

$$\inf\{\phi(u): \phi \in S(V)\} + \inf\{f(x): f \in X_1'\} \ge 0.$$

Lemma 2.2. Let X be a real normed linear space and let (V, V^+, e) be an order unit space.

(1) For $x \in X$, we have

$$\inf\{f(x) : f \in X_1'\} = -\|x\|$$

(2) For $u \in V^+$, we have

$$\inf\{\phi(u):\phi\in S(V)\} = \|u\| - \|\|u\|e - u\|.$$

Proof. (1): Find $f_0 \in X'_1$ such that $||x|| = f_0(x)$. Then $f(x) \leq f_0(x)$ for all $f \in X'_1$. Since $-f \in X'_1$ whenever $f \in X'_1$, we get that

$$\inf\{f(x): f \in X_1'\} = -f_0(x) = -||x||.$$

(2): Since S(V) is weak*-compact, there exists $\phi_0 \in S(V)$ such that $\phi_0(u) \leq \phi(u)$ for all $\phi \in S(V)$. Thus

$$\phi(\|u\|e - u) = \|u\| - \phi(u) \le \|u\| - \phi_0(u) = \phi_0(\|u\|e - u)$$

for all $\phi \in S(V)$. Therefore,

$$|||u||e - u|| = \phi_0(||u||e - u) = ||u|| - \phi_0(u),$$

, or equivalently,

$$\inf\{\phi(u):\phi\in S(V)\}=\phi_0(u)=\|u\|-\|\|u\|e-u\|.$$

We deduce that the expected cone V_X^+ should thus be given by

$$V_X^+ = \{(u, x) \in V \oplus_1 X : u \in V^+ \text{ and } \|x\| \le \|u\| - \|\|u\|e - u\|\}.$$

The condition given in V_X^+ can be presented in a simpler form.

Lemma 2.3. Let (V, V^+, e) be an order unit space. Then for $u \in V$ and $k \in \mathbb{R}^+$, following two conditions are equivalent:

(1) $u \in V^+$ and $k \le ||u|| - |||u||e - u||$. (2) $ke \le u$.

Proof. First we assume that $u \in V^+$ and $k \le ||u|| - |||u||e - u||$. Then $|||u||e - u|| \le ||u|| - k$. Thus $0 \le ||u||e - u \le (||u|| - k)e$ so that $ke \le u$.

Conversely, we assume that $ke \leq u$. Then $u \in V^+$ with $k \leq ||u||$. Since $u \leq ||u||e$, we get $0 \leq ||u||e - u \leq ||u||e - ke$. Thus $|||u||e - u|| \leq ||u|| - k$, whence $k \leq ||u|| - |||u||e - u||$. **Theorem 2.4.** Let (V, V^+, e) be an order unit space and X is a real normed linear space. Then $(V \oplus_1 X, V_X^+, e_X)$ is an order unit space where $V_X^+ = \{(u, x) : ||x|| e \leq u\}$ and $e_X := (e, 0)$.

Proof. Let $(u, x) \in V_X^+$ and let α be a positive real number. Then $||x||e \leq u$ so that $||\alpha x||e \leq \alpha u$. Thus $\alpha(u, x) \in V_X^+$. Next, let $(u, x), (v, y) \in V_X^+$. Then $||x||e \leq u$ and $||y||e \leq v$. Thus $||x + y||e \leq (||x|| + ||y||)e \leq u + v$ so that $(u, x) + (v, y) \in V_X^+$ as $u + v \in V^+$. Hence V_X^+ is a cone.

Next, let $\pm(u, x) \in V_X^+$. Then $||x|| e \leq u$ and $||-x|| e \leq -u$. Thus $\pm u \in V^+$. Since V^+ is proper, we get u = 0 and consequently, x = 0. Thus V_X^+ is also proper.

Now, we show that e_X is an order unit for $V \oplus_1 X$. For this, let $(v, x) \in V \oplus_1 X$. Put k = ||v|| + ||x||. Since $||v|| e \pm v \ge 0$, we get $ke \pm v \ge ||x|| e$. Thus $ke_X \pm (u, x) \in V_X^+$. Therefore, e_X is an order unit for $V \oplus_1 X$. We also notice that $||(v, x)||_o \le ||(v, x)||_1$ for all $(v, x) \in V \oplus_1 X$ where $||\cdot||_o$ is the corresponding order unit (semi)norm on $V \oplus_1 X$.

Next, we prove that V_X^+ is Archimedean. Let $(v, x) \in V \oplus_1 X$ be such that $ke_X + (v, x) \in V_X^+$ for all k > 0. That is, $(ke + v, x) \in V_X^+$ for all k > 0. Then $||x||e \le ke + v$ for all k > 0. Now by the Archimedean property in V^+ , we get $||x||e \le v$, that is, $(v, x) \in V_X^+$. Therefore, V_X^+ is Archimedean.

Hence $(V \oplus_1 X, V_X^+, e_X)$ is an order unit space. Also, we have that the corresponding order unit norm $\|\cdot\|_o \leq \|\cdot\|_1$. We prove their equality. Let $(v, x) \in V \oplus_1 X$. Assume that $\lambda e_X \pm (v, x) \in V_X^+$, that is, $(\lambda e \pm v, \pm x) \in V_X^+$ for some $\lambda > 0$. Then $\|x\|_e \leq \lambda e \pm v$. Thus $\pm v \leq (\lambda - \|x\|)e$. It follows that $\|v\| \leq \lambda - \|x\|$, that is $\|(v, x)\|_1 \leq \lambda$. Hence $\|(v, x)\|_1 \leq \|(v, x)\|_o$. This completes the proof. \Box

Now we describe the order unit space (ℓ_1, e_1) as an order unit space by adjoining a normed linear space to an order unit space.

Example 2.5. Consider the order unit space $(\mathbb{R}, 1)$ and the normed linear space \mathbb{R} in their natural setting. Then by Theorem 2.4, ℓ_1^2 is an order unit space with the cone $(\ell_1^2)_{e_1^2}^+ := \{(\alpha, \beta) : \alpha \ge |\beta|\}$ and the order unit $e_1^2 := (1, 0)$. Next, consider the resulting order unit space $(\ell_1^2, (\ell_1^2)_{e_1^2}^+, e_1^2)$ and again the naturally normed \mathbb{R} . Again invoking Theorem 2.4, we now obtain the order unit space $(\ell_1^3, (\ell_1^3)_{e_1^3}^+, e_1^3)$ where $e_1^3 = (1, 0, 0)$ and $(\ell_1^3)_{e_1^3}^+ := \{(\alpha, \beta, \gamma) : (\alpha, \beta) \ge |\gamma| e_1^2$ in $(\ell_1^2)_{e_1^2}^+ \}$. If $(\alpha, \beta, \gamma) \in (\ell_1^3)_{e_1^3}^+$, then $|\gamma| e_1^2 \le (\alpha, \beta)$ in $(\ell_1^2)_{e_1^2}^+$ so that $(\alpha - |\gamma|, \beta) \in (\ell_1^2)_{e_1^2}^+$. Thus $|\beta| \le \alpha - |\gamma|$, that is, $||(\beta, \gamma)||_1 = |\beta|| + |\gamma| \le \alpha$. Now

ANIL KUMAR KARN

it follows that $(\ell_1^3, (\ell_1^3)_{e_1^3}^+, e_1^3)$ is also the oder unit obtained by adjoining the normed linear space ℓ_1^2 to the order unit space $(\mathbb{R}, 1)$. Also following the induction on n, we may conclude that for each $n \in \mathbb{N}$, $(\ell_1^{n+1}, (\ell_1^{n+1})_{e_1^{n+1}}^+, e_1^{n+1})$ can be obtained as the order unit space by adjoining \mathbb{R} to $(\mathbb{R}, 1)$ successively n-times or simply adjoining ℓ_1^n to $(\mathbb{R}, 1)$. As a result of induction on n, we can also conclude that $(\ell_1, (\ell_1)_{e_1^{n+1}}^+, e_1)$ can be obtained as the order unit space by adjoining \mathbb{R} to $(\mathbb{R}, 1)$ successively countably many times or simply adjoining ℓ_1 to $(\mathbb{R}, 1)$.

Remark 2.6. A dual statement may be made about ℓ_{∞} as a base normed space which we leave to the readers. Furthermore, when a (real) Hilbert space is adjoined to the order unit space (\mathbb{R} , 1), we get spin factors [3, Section 6.1]. (See also [6] where we have generalized this idea.)

Remark 2.7. By [8, Lemma 5.1], every 2-dimensional order unit space is isometrically isomorphic to ℓ_{∞}^2 . Here we construct a map to demonstrate that as a real normed linear space, ℓ_1^2 is isometrically isomorphic to the order unit space ℓ_{∞}^2 .

Consider \mathbb{R} as a real normed linear space with the standard norm as the modulus and also consider $(\mathbb{R}, 1)$ as the order unit space with its natural cone. Then $\mathbb{R} \oplus_1 \mathbb{R} \cong \ell_1^2$ and by Theorem 2.4, it is an order unit space with its cone $C = \{(\alpha, \beta) : |\beta| \le \alpha\}$ and the order unit $e_1 := (1, 0)$. Now define $\psi : \ell_1^2 \to \ell_\infty^2$ given by $\psi(\alpha, \beta) = (\alpha + \beta, \alpha - \beta)$ for all $\alpha, \beta \in \mathbb{R}$. Then ψ is a linear, isometric isomorphism. Also $\psi(C) = \{(x, y) : x, y \in [0, \infty)\}$ and $\psi(e_1) = (1, 1)$.

3. Norming order unit

When we think of an order unit space, generally ℓ_{∞} -type spaces appear in our minds. However, we have noted in the previous section that even ℓ_1 can be shown to be an order unit space, albeit not in its natural order. This leads to a curiosity: which normed linear space can be seen as order unit space. In this section, we try to address this question. Let us begin with a notion.

Definition 3.1. Let X be a non-zero real normed linear space and let $e \in X$ with ||e|| = 1. We say that e is a norming order unit for X, if $||2x - ||x||e|| \le ||x||$ whenever $||2x - \lambda e|| \le \lambda$ for some $\lambda \ge ||x||$.

Note that -e is a norming order unit for X whenever e is a norming order unit for X. More generally, if $T: X \to X$ is a surjective linear isometry, then T(e) is a norming order unit for X whenever e is a

6

norming order unit for X. The next result justifies the nomenclature "norming order unit".

Proposition 3.2. Let (V, e) be an order unit space and let $u \in V$. Then the following statements are equivalent:

- (1) $u \in V^+$.
- (2) $||2u \lambda e|| \le \lambda$ for any $\lambda \ge ||u||$.
- (3) $||2u \lambda e|| \le \lambda$ for some $\lambda \ge ||u||$.
- (4) ||2u ||u||e|| = ||u||.

Proof. Let $u \in V^+$ and assume that $\lambda \geq ||u||$. Then $0 \leq u \leq ||u||e \leq \lambda e$. Thus $\lambda e \pm (2u - \lambda e) \in V^+$ and we have $||2u - \lambda e|| \leq \lambda$. Conversely, assume that $||2u - \lambda e|| \leq \lambda$ for some $\lambda \geq ||u||$. Then $\lambda e \pm (2u - \lambda e) \in V^+$ so that $u \in V^+$. Hence (1), (2) and (3) are equivalent. Also then (4) implies (1). Finally, assume that $u \in V^+$ and let $||2u - ||u||e|| = \alpha$. Then $\alpha e \pm (2u - ||u||e) \in V^+$. Thus $0 \leq u \leq \frac{1}{2}(||u|| + \alpha)e$. It follows that $||u|| \leq \alpha$. By (2), we have $\alpha \leq ||u||$ which completes the proof. \Box

We show that whenever a real normed linear space possesses a norming order unit, it is isometric to an order unit space.

Theorem 3.3. Let $(X, \|\cdot\|)$ be a real normed linear space and assume that $e \in X$ with $\|e\| = 1$ be a norming order unit for X. Put

$$X_e^+ = \{ x \in X : \|2x - \|x\|e\| \le \|x\| \}.$$

Then X_e^+ is a proper and Archimedean cone in X and e is an order unit for (X, X_e^+) so that (X, X_e^+, e) is an order unit space. Moreover, e determines $\|\cdot\|$ as an order unit norm on X.

Proof. We have $0, e \in X_e^+$ so that $X_e^+ \neq \emptyset$. Let $x \in X_e^+$ and $\alpha \ge 0$. Then $||2x - ||x||e|| \le ||x||$. Thus

 $||2\alpha x - ||\alpha x||e|| = \alpha ||2x - ||x||e|| \le \alpha ||x|| = ||\alpha x||$

so that $\alpha x \in X_e^+$. Next, let $x, y \in X_e^+$. Then $||2x - ||x||e|| \le ||x||$ and $||2y - ||y||e|| \le ||y||$. Thus

$$\begin{aligned} \|2(x+y) - (\|x\| + \|y\|)e\| &\leq \|2x - \|x\|e\| + \|2y - \|y\|e\| \\ &\leq \|x\| + \|y\|. \end{aligned}$$

Since e is a norming order unit for X, we conclude that $||2(x+y) - ||x+y||e|| \le ||x+y||$. Thus $x+y \in X_e^+$ and X_e^+ is a cone. We observe that if $x \in X_e^+$, then for any $\lambda \ge ||x||$, we have $||2x - \lambda e|| \le \lambda$. In fact,

$$\begin{aligned} \|2x - \lambda e\| &= \|(2x - \|x\|e) - (\lambda - \|x\|)e\| \\ &\leq \|2x - \|x\|e\| + \|(\lambda - \|x\|)e\| \\ &\leq \|x\| + \lambda - \|x\| = \lambda. \end{aligned}$$

ANIL KUMAR KARN

Hence, $x \in X_e^+$ if and only if $||2x - \lambda e|| \le \lambda$ for any $\lambda \ge ||x||$.

Now, we show that e is an order unit for (X, X_e^+) . Let $x \in X$. Then $\|(\|x\|e \pm x) - \|x\|e\| = \|x\|$. Since $\|\|x\|e \pm x\| \le 2\|x\|$, we conclude that $\|x\|e \pm x \in X_e^+$ for all $x \in X$. Thus e is an order unit for X. We also deduce that $\|x\|_e \le \|x\|$ for all $x \in X$ where $\|\cdot\|_e$ is the order unit (semi)norm on X. We prove the equality. Let $x \in X$ and $\lambda \ge 0$ be such that $\lambda e \pm x \in X_e^+$. Since $\|\lambda e \pm x\| \le \lambda + \|x\|$, we have

$$||2(\lambda e \pm x) - (\lambda + ||x||)e|| \le \lambda + ||x||.$$

In other words, $||2x \pm (\lambda - ||x||)e|| \le \lambda + ||x||$. Thus

$$4\|x\| \le \|2x + (\lambda - \|x\|)e\| + \|2x - (\lambda - \|x\|)e\| \le 2(\lambda + \|x\|)$$

so $||x|| \leq \lambda$. Hence $||x|| \leq ||x||_e$ and consequently, $||x|| = ||x||_e$ for every $x \in X$. Further, if $\pm x \in X_e^+$ for some $x \in X$, then as above, we get $||x|| \leq 0$. Thus x = 0 so that X_e^+ is proper.

Finally, we prove that X_e^+ is Archimedean. Assume that $x \in X$ be such that $x + \lambda e \in X_e^+$ for all $\lambda > 0$. Since $||x + \lambda e|| \le ||x|| + \lambda$, we have $||2(x + \lambda e) - (||x|| + \lambda)e|| \le ||x|| + \lambda$ for all $\lambda > 0$. Letting $\lambda \to 0^+$, we get $||2x - ||x||e|| \le ||x||$. Thus $x \in X_e^+$. That is, X_e^+ is Archimedean.

The following observation is a direct consequence of Proposition 3.2 which is the converse of Theorem 3.3.

Corollary 3.4. Let (V, e) be an order unit space. Then e is a norming order unit for V with $V_e^+ = V^+$.

Corollary 3.5. A strictly convex real normed linear space of dimension more than 1 does not possess any norming order unit.

Proof. Let V be a strictly convex real normed linear space of dimension more than 1. Assume to the contrary that $e \in V$ with ||e|| = 1 is a norming order unit for V. Then (V, V_e^+, e) is an order unit space such that the norm on V is the order unit norm determined by e. Here $V_e^+ = \{u \in V : ||2u - ||u||e|| \le ||u||\}$. Let $u \in V_e^+$ with ||u|| = 1 and $u \neq e$. It follows from Proposition 3.2 that ||2u - e|| = 1. Put v :=2u - e. Then ||v|| = 1 with $v \neq e$. Since V is strictly convex, we must have $||\frac{1}{2}(e+v)|| < 1$. This leads to a contradiction as $\frac{1}{2}(e+v) = u$. \Box

In [8], the author has shown that the skeleton $S_V := \{u \in V^+ : \|u\| = 1 = \|e-u\|\}$ of an order unit space (V, V^+, e) describes the whole space. So it may be interesting to describe skeletal elements (that is, the elements in S_V) in terms of norming order units independently. We shall use ∞ -orthogonality for this purpose. Let us recall that $x, y \in$

8

X in a real Banach space X are said to *orthogonal* to each other, if $||x + ky|| = \max\{||x||, ||ky||\}$ for all real number k [5].

Proposition 3.6. Let X be a real normed linear space, $e \in X$ with ||e|| = 1 be a norming order unit for X and assume that $u \in X$. Then u is a skeletal element in the order unit space (X, X_e^+, e) if and only if ||u|| = ||e - u|| and ||2u - e|| = 1.

Proof. First, we assume that u is a skeletal element of (X, X_e^+, e) . That is, $u \in X_e^+$ and ||u|| = 1 = ||e - u||. Then $e - u \in X_e^+$. Since 1 = ||e|| = ||u + (e - u)||, we have $u \perp_{\infty} (e - u)$ by [5, Theorem 3.3]. Thus

$$||2u - e|| = ||u - (e - u)|| = \max\{||u||, ||e - u||\} = 1.$$

Conversely, we assume that ||u|| = ||e - u|| and ||2u - e|| = 1. Since e is the norming order unit for X_e^+ , we have $e \pm (2u - e) \in X_e^+$, or equivalently, $u, e - u \in X_e^+$. Thus $||u|| = ||e - u|| \le 1$. Also then

$$1 = \|2u - e\| = \|u - (e - u)\| \le \max\{\|u\|, \|e - u\|\} = \|u\|$$

so that ||u|| = ||e - u|| = 1. Thus u is a skeletal element in the order unit space (X, X_e^+, e) .

3.1. Some examples.

Example 3.7. Norming order units in ℓ_{∞} . Since the constant sequence $\langle 1 \rangle$ is the norming order unit for ℓ_{∞} together with its natural cone, and since $\langle \alpha_n \rangle \mapsto \langle \epsilon_n \alpha_n \rangle$ is a surjective isometry in ℓ_{∞} where $\epsilon_n \in \{1, -1\}$, we deduce that $\langle \epsilon_n \rangle$ is also a norming order unit for ℓ_{∞} for every choices of ϵ_n . We show that these are the only norming order units in ℓ_{∞} .

Let $e_0 = \langle \alpha_n \rangle \in \ell_{\infty}$ with $\|\langle \alpha_n \rangle\|_{\infty} = 1$. Without any loss of generality, we also assume that $\alpha_n \geq 0$ for every n. Thus $\sup_{n \in \mathbb{N}} \alpha_n = 1$. Let that $\alpha_{n_0} < 1$ for some $n_0 \in \mathbb{N}$. Put $\lambda_0 = \frac{2}{1+\alpha_{n_0}}$. Then $\lambda_0 > 1$. Consider $e_{n_0} \in \ell_{\infty}$ with 1 at n_0 -th place and 0 elsewhere. Then $\|e_{n_0}\|_{\infty} = 1$. Also

$$||2e_{n_0} - \lambda_0 e_0||_{\infty} = \max\{|2 - \lambda_0 \alpha_{n_0}|, \sup_{n \neq n_0} \alpha_n\} = \lambda_0.$$

However,

$$||2e_{n_0} - e_0||_{\infty} = \max\{|2 - \alpha_{n_0}|, \sup_{n \neq n_0} \alpha_n\} = 2 - \alpha_{n_0} > 1 = ||e_{n_0}||_{\infty}.$$

Thus e_0 fails to be a norming order unit for ℓ_{∞} .

Example 3.8. Norming order units in ℓ_1 . Let $e_0 = \langle \alpha_n \rangle \in \ell_1$ with $\|\langle \alpha_n \rangle\|_1 = 1$. Without any loss of generality, we can again assume that $\alpha_n \geq 0$ for every n. Then $\sum_{n \in \mathbb{N}} \alpha_n = 1$.

Case 1. Assume that $0 < \alpha_{n_0} < 1$ for some n_0 . Consider $e_{n_0} \in \ell_1$ with 1 at n_0 -th place and 0 elsewhere. Then $||e_{n_0}||_1 = 1$. Also

$$\begin{aligned} \left\| 2e_{n_0} - \frac{2}{\alpha_{n_0}} e_0 \right\|_1 &= \left(\frac{2}{\alpha_{n_0}} \right) \|\alpha_{n_0} e_{n_o} - e_0\|_1 \\ &= \left(\frac{2}{\alpha_{n_0}} \left(\sum_{n \neq n_0} \alpha_n \right) \right) \\ &= \left(\frac{2(1 - \alpha_{n_0})}{\alpha_{n_0}} \right) \\ &< \left(\frac{2}{\alpha_{n_0}} \right). \end{aligned}$$

However

$$\begin{aligned} \|2e_{n_0} - e_0\|_1 &= (2 - \alpha_{n_0}) + \left(\sum_{n \neq n_0} \alpha_n\right) \\ &= (2 - \alpha_{n_0}) + (1 - \alpha_{n_0}) \\ &> 1. \end{aligned}$$

Thus in this case, e_0 fails to be a norming order unit for ℓ_1 . So we assume now that $\alpha_n \in \{0, 1\}$.

Case 2. Assume that $\alpha_{n_0} = 1$ for some n_0 . Then $\alpha_n = 0$, if $n \neq n_0$ so that $e_0 = e_{n_0}$. We show that e_0 is a norming order unit for ℓ_1 . Let $u = \langle x_n \rangle \in \ell_1$ with $||u||_1 = 1$. Let $\lambda \ge 1$ be such that $||2u - \lambda e_0||_1 \le \lambda$ or equivalently, $|2x_{n_0} - \lambda| + 2\sum_{n \ne n_0} |x_n| \le \lambda$. Since $\sum_{n \ne n_0} |x_n| = 1 - |x_{n_0}|$, we have $|2x_{n_0} - \lambda| \le \lambda + 2|x_{n_0}| - 2$. In other words,

$$-(\lambda + 2|x_{n_0}| - 2) \le 2x_{n_0} - \lambda \le \lambda + 2|x_{n_0}| - 2,$$

whence $1 - |x_{n_0}| \le x_{n_0} \le \lambda + |x_{n_0}| - 1$. Thus $x_{n_0} \ge 0$ and consequently, $x_{n_0} \ge \frac{1}{2}$. Therefore,

$$||2u - e_0||_1 = (2x_{n_0} - 1) + 2\sum_{n \neq n_0} |x_n| = (2x_{n_0} - 1) + 2(1 - x_{n_0}) = 1.$$

Hence e_{n_0} is a norming order unit for ℓ_1 . In this way, we may conclude that $\{\pm e_n : n \in \mathbb{N}\}$ is the set of all norming order units for ℓ_1 .

We end the paper with the following conjecture.

Conjecture 3.9. Let (V, V^+, e) be an order unit space and assume that $u \in V$ with ||u|| = 1 is a norming order unit. Then there exists a surjective linear isometry $T: V \to V$ such that T(e) = u.

NORMING ORDER UNITS

Acknowledgements: The author was partially supported by Science and Engineering Research Board, Department of Science and Technology, Government of India sponsored Mathematical Research Impact Centric Support project (reference no. MTR/2020/000017).

References

- E. M. Alfsen, Compact convex sets and boundary integrals, Springer-Verlag, 1971.
- [2] G. J. O. Jameson, Ordered linear spaces, Lecture Notes in Mathematics, Springer-Verlag, 1970.
- [3] H. Hanche-Olsen and E. Störmer, *Jordan operator algebras*, Pitman Advanced Publishing Program, 1984.
- [4] R. V. Kadison, A representation theory for commutative topological algebras, Mem. Amer. Math. Soc., 7(1951).
- [5] A. K. Karn, Orthogonality in sequence spaces and its bearing on ordered Banach spaces, Anil K. Karn; Positivity, 18(2) (2014), 223-234.
- [6] A. K. Karn, A generalization of spin factors, Acta. Sci. Math. (Szeged)), 87 (2021), 551-569.
- [7] A. K. Karn, Centre of a compact convex set, Banach Journal of Mathematical Analysis, 16(4) (2022), Article 68, 19 pages.
- [8] A. K. karn, On the geometry of order unit spaces, Advances in Operator Theory 9, 28 (2024), 18 pages.
- [9] Y.-C. Wong and K.-F. Ng, *Partially ordered topological spaces*, Oxford University Press, Oxford, 1973.

School of Mathematical Sciences, National Institute of Science Education and Research Bhubaneswar, An OCC of Homi Bhabha National Institute, P.O. - Jatni, District - Khurda, Odisha - 752050, India.

Email address: anilkarn@niser.ac.in