# A FOOTNOTE TO A PAPER OF DEODHAR

DAVIDE FRANCO

ABSTRACT. Let  $X \subseteq G/B$  be a Schubert variety in a flag manifold and let  $\pi : \tilde{X} \to X$  be a Bott-Samelson resolution of X. In this paper we prove an effective version of the decomposition theorem for the derived pushforward  $R\pi_*\mathbb{Q}_{\tilde{X}}$ . As a by-product, we obtain recursive procedure to extract Kazhdan-Lusztig polynomials from the polynomials introduced by V. Deodhar in [6], which does not require prior knowledge of a minimal set. We also observe that any family of equivariant resolutions of Schubert varieties allows to define a new basis in the Hecke algebra and we show a way to compute the transition matrix, from the Kazhdan-Lusztig basis to the new one.

*Keywords*: Kazhdan-Lusztig polynomials, Intersection cohomology, Decomposition theorem, Schubert varieties, Bott-Samelson resolution, Hecke algebra.

MSC2010: Primary 14B05, 14M15; Secondary 14E15, 14F45, 32S20, 32S60, 58K15.

# 1. INTRODUCTION

As the title suggests, this work is a sort of appendix to [6]. In such a paper, Vinay Deodhar introduces a statistic, called *defect*, on the subexpressions of a given reduced expression of an element of a Coxeter group W (see also [2, Section 6.3.17]). Specifically, let  $w \in W$  be an element of length l(w) and let

$$w = s_1 \dots s_l, \quad l = l(w)$$

be a reduced expression of w. A subexpression  $\sigma = (\sigma_0, \ldots, \sigma_l)$  is a sequence of Coxeter group elements such that

$$\sigma_{i-1}^{-1}\sigma_i \in \{id, s_j\}, \quad \text{for all} \quad 1 \le j \le l.$$

Let  $\mathcal{S}$  be the set such sequences for the given reduced expression and let

$$\pi(\sigma) := \sigma_l, \quad \text{if} \quad \sigma = (\sigma_0, \dots, \sigma_l) \in \mathcal{S}.$$

For any subexpression  $\sigma = (\sigma_0, \ldots, \sigma_l) \in \mathcal{S}$ , Deodhar defines the *defect* of  $\sigma$  by

$$d(\sigma) := \#\{1 \le j \le l \mid \sigma_{j-1}^{-1} s_j < \sigma_{j-1}\}.$$

If one fix a reduced expression for all  $w \in W$  and consider  $v \in W$  such that  $v \leq w$  in the Bruhat order, then one can use the defect to define the following polynomial

(1) 
$$Q_{w,v} := \sum_{\sigma \in \mathcal{S}, \, \pi(\sigma) = v} q^{d(\sigma)} \in \mathbb{Z}[q]$$

In [6], Deodhar proves that the Kazhdan-Lusztig polynomial  $P_{w,v}$  admit a description as subsum of (1). More precisely, he gives a recursive algorithm for computing a minimal set

### DAVIDE FRANCO

 $E_{min} \subseteq \mathcal{S}$  such that

(2) 
$$P_{w,v} := \sum_{\sigma \in E_{min}, \, \pi(\sigma) = v} q^{d(\sigma)} \in \mathbb{Z}[q],$$

for any pair  $w, v \in W$ ,  $v \leq w$ . What is more, in [6] one can find a new basis of the Hecke algebra of W that is defined starting from the polynomials (1).

The main aim of this paper is to give a recursive procedure to extract Kazhdan-Lusztig polynomials  $P_{w,v}$  from polynomials  $Q_{w,v}$ , without going through the computation of the minimal set  $E_{min} \subseteq S$ , in the case W is the Weyl group of a semisimple connected algebraic group over  $\mathbb{C}$ . Instead, our approach is based on an effective version of the Beilinson-Bernstein-Deligne-Gabber decomposition theorem (BBDG for short) for the Bott-Samelson resolution. As a by-product of our analysis, we provide a recursive procedure to compute the change-of-basis matrix, from the Kazhdan-Lusztig basis of the Hecke algebra to the basis defined in [6].

The starting point of our analysis is Proposition 3.9 of [6], where the author shows that, when W is the Weyl group of a semisimple connected algebraic group, the polynomial (1) has a nice geometric interpretation as Poincaré polynomial of a suitable fiber of the *Bott-Samelson* resolution of the Schubert variety X(w) (see section (2) for a short review of some standard definitions and notations concerning Schubert varieties). More precisely, if we denote by Gan algebraic group with Weyl group W and Borel subgroup B, then to the chosen reduced expression  $w = s_1 \dots s_l$  it is also associated the Bott-Samelson resolution

$$\pi_w: X(w) \to X(w)$$

The smooth variety  $\tilde{X}(w)$  is defined as the subvariety of  $(G/B)^l$  consisting of *l*-tuples  $(g_1B, \ldots, g_lB)$  such that

$$g_{i-1}^{-1}g_i \in \overline{Bs_iB}, \quad 1 \le i \le l \quad (\text{by convention}, \ g_0 = 1),$$

and  $\pi_w$  is the projection on the last factor.

The polynomial (1) is the Poincaré polynomial of the fiber of  $\pi_w$  over the cell  $\Omega_v \subset X(w)$  associated to v:

(3) 
$$Q_{w,v} = \sum_{i} \dim H^{2i}(\pi_w^{-1}(x))q^i, \quad \forall x \in \Omega(v).$$

Our approach for extracting the Kazhdan-Lusztig polynomials from the polynomials defined in (1), is to prove an effective version of the BBDG decomposition theorem for the Bott-Samelson resolution and, more generally, for any equivariant resolution of a Schubert variety (in [9], [11], and [3] partial results in this direction were previously obtained). Specifically, let  $D_c^b(X)$  be the derived category of bounded complexes of constructible Q-vector sheaves on a Schubert variety  $X \subseteq G/B$ . The decomposition theorem applied to an equivariant resolution  $\pi : \tilde{X} \to X$ , states that the derived direct image  $R\pi_*\mathbb{Q}_{\tilde{X}}[\dim X]$  splits in  $D_c^b(X)$  as a direct sum of shifts of irreducible perverse sheaves on X. By [5, § 1.5], we have a non-canonical decomposition

(4) 
$$R\pi_*\mathbb{Q}_{\tilde{X}}[\dim X] \cong \bigoplus_{i\in\mathbb{Z}} \bigoplus_{j\in\mathbb{N}} IC(L_{ij})[-i],$$

where the summands are shifted intersection cohomology complexes of the semisimple local systems  $L_{ij}$ , each of which is supported on a suitable locally closed stratum of codimension j, usually called a *support* of the decomposition. The summand supported in the general point

is precisely the intersection cohomology of X. The supports appearing in the splitting (4) and the local systems  $L_{ij}$  are, generally, rather mysterious objects when  $j \ge 1$ .

Quite luckily, in our case a crucial simplification arises because all the local systems  $L_{ij}$  appearing in the decomposition (4) are trivial by an easy argument that is explained in Proposition 3.2. As a consequence, we have

(5) 
$$R\pi_* \mathbb{Q}_{\tilde{X}}[\dim X] \cong \bigoplus_{X(v) \subseteq X} \bigoplus_{\alpha \in \mathbb{Z}} IC_{X(v)}^{\oplus s_{v,\alpha}},$$

for suitable multiplicities  $s_{v,\alpha}$ . Since we have a splitting like (5) for any equivariant resolution  $\pi_w : \tilde{X}(w) \to X(w)$ , we can define a Laurent polynomial recording the contribution to the decomposition theorem for  $\pi_w$ , with support X(v):

$$D_{w,v}(t) := \sum_{\alpha \in \mathbb{Z}} s_{v,\alpha}^w \cdot t^\alpha \in \mathbb{Z}[t, t^{-1}],$$

for avery pair (v, w) in W, such that  $v \leq w$ , and for every resolution.

Now, assume to have fixed an equivariant resolution  $\pi_w : X(w) \to X(w)$  for every Schubert variety X(w) and assume that the cohomology of the fibers of  $\pi_w$  vanish in odd degrees (this property is satisfied by any reasonable resolution of Schubert varieties). Similarly as in (3), define the analogue Deodhar's polynomial  $Q_{w,v}$  as the Poincaré polynomial of the fibers of  $\pi_w$ over the cell  $\Omega(v)$ . The main results contained in this paper can be summarized as follows:

a) we set up an iterative procedure that allows to compute **both** the Kazhdan-Lusztig polynomials  $P_{w,v}$  and the Laurent polynomials  $D_{w,v}$  starting from Deodhar's polynomials  $Q_{w,v}$ ;

b) we observe that the polynomials  $Q_{w,v}$  allow to construct a new basis  $\{B_w \mid w \in W\}$  of the Hecke algebra;

c) we prove that the transition matrix, from the Kazhdan-Lusztig basis to the new basis  $\{B_w \mid w \in W\}$ , can be easily deduced from the Laurent polynomials  $D_{w,v}$  and does not require prior knowledge of the transition matrix from the Kazhdan-Lusztig basis to the standard one (compare with Remark 5.2).

### 2. NOTATIONS AND BASIC FACTS

In this section we review some basic facts concerning Buhat decomposition, Schubert varieties and combinatorics of subexpressions that are needed in the following.

(i) Let G be a semisimple connected algebraic group over  $\mathbb{C}$ . Let T and B be a maximal torus and a Borel subgroup of G, respectively. Denote by W be the Weyl group of G. If we consider

$$\{e_w \mid w \in W\} \subset G/B,$$

the set of fixed points for the torus action on G/B, then we have the Bruhat decomposition of G/B i.e. the disjoint union of Bruhat cells

$$G/B = \bigsqcup_{w \in W} \Omega(w), \quad \Omega(w) := Be_w.$$

For every  $w \in W$  the *Schubert variety* associated to w is defined as the Zariski closure of the corresponding Bruhat cell:

$$X(w) := \overline{\Omega(w)}$$

(ii) There is a partial order on the Weyl group W determined by the decomposition above. Specifically, for  $w_1, w_2 \in W$  we have

$$w_1 \ge w_2 \quad \Leftrightarrow \quad X(w_1) \supseteq X(w_2)$$

Furthermore, we have

$$X(w) = \bigcup_{v \le w} \Omega(v)$$

We borrow from Deodhar's paper [6] some crucial definitions concerning the combinatorics of subexpressions of a reduced word in a Coxeter group.

# **Definition 2.1.** [6, Def. 2.1 - 2.2]

(1) Let  $w \in W$  be an element of length l(w) and let

$$w = s_1 \dots s_l, \quad l = l(w)$$

be a reduced expression of w. A subexpression  $\sigma = (\sigma_0, \ldots, \sigma_l)$  is a sequence of Weyl group elements such that  $\sigma_0 = id$  and

$$\sigma_{j-1}^{-1}\sigma_j \in \{id, s_j\}, \text{ for all } 1 \le j \le l.$$

For any reduced word r, let  $S_r$  be the set of reduced expressions of r and let

$$\pi(\sigma) := \sigma_l, \quad \text{if} \quad \sigma = (\sigma_0, \dots, \sigma_l) \in \mathcal{S}_r.$$

(2) For any  $\sigma = (\sigma_0, \ldots, \sigma_l) \in S_r$  define the *defect* of  $\sigma$  by

$$d(\sigma) := \#\{1 \le j \le l \mid \sigma_{j-1}^{-1} s_j < \sigma_{j-1}\},\$$

If  $v \leq w$ , we let

(6) 
$$Q_{w,v} := \sum_{\sigma \in \mathcal{S}_r, \, \pi(\sigma) = v} q^{d(\sigma)} \in \mathbb{Z}[q].$$

From now on we assume that we have chosen a reduced expression for all  $w \in W$ , hence (6) provides a polynomial  $Q_{w,v} \in \mathbb{Z}[q]$  for any pair (w, v) such that  $v \leq w$ .

As explained in [6, Proposition 3.9], the polynomial above has a nice geometric interpretation as Poincaré polynomial of the *Bott-Samelson resolution*. We recall that to each reduced expression  $w = s_1 \dots s_l$ , it is also associated the *Bott-Samelson resolution* 

$$\pi_w: X(w) \to X(w)$$

The smooth variety  $\tilde{X}(w)$  is defined as the subvariety of  $(G/B)^l$  consisting of *l*-tuples  $(g_1B, \ldots, g_lB)$  such that

 $g_{i-1}^{-1}g_i \in \overline{Bs_iB}, \quad 1 \le i \le l \quad (\text{by convention}, g_0 = 1),$ 

and  $\pi_w$  is the projection on the last factor. It is clear that  $\pi_w$  is equivariant under the action of the Borel subgroup B.

By [6, Proposition 3.9], (6) is the Poincaré polynomial of the fiber of  $\pi_w$  over the cell  $\Omega(v) \subset X(w)$ :

(7) 
$$Q_{w,v} = \sum_{i} \dim H^{2i}(\pi_w^{-1}(x))q^i, \quad \forall x \in \Omega(v).$$

# 3. The Decomposition theorem

Formula (7) and Deodhar's paper suggest that there should be a closed relationship between the Poincaré polynomials of the stalk cohomology of the complex  $R\pi_*\mathbb{Q}_{\tilde{X}(w)}$  and the Kazhdan-Lusztig polynomials. The most important result concerning the complex  $R\pi_*\mathbb{Q}_{\tilde{X}(w)}$ and, in general, concerning the topology of proper algebraic map is the *decomposition theorem* of Beilinson, Bernstein, Deligne and Gabber, which we now recall.

In what follows, we shall work cohomology with  $\mathbb{Q}$ -coefficients and the self-dual perversity  $\mathfrak{p}$  (see [1, §2.1], and [12, p. 79]).

**Theorem 3.1.** (*Decomposition theorem* [5, 1.6.1]) Let  $f : X \to Y$  be a proper map of complex algebraic varieties. In  $D_c^b(Y)$ , the derived category of bounded complexes of constructible  $\mathbb{Q}$ -vector sheaves on Y, there is a non-canonical isomorphism

(8) 
$$Rf_*IC_X \cong \bigoplus_{\alpha \in \mathbb{Z}} {}^{\mathfrak{p}}\mathcal{H}^{\alpha}(Rf_*IC_X) \left[-\alpha\right].$$

Furthermore, the perverse sheaves  ${}^{\mathfrak{p}}\mathcal{H}^{\alpha}(Rf_*IC_X)$  are semisimple; i.e. there is a decomposition into finitely many disjoint locally closed and nonsingular subvarieties  $Y = \coprod S_{\beta}$  and a canonical decomposition into a direct sum of intersection complexes of semisimple local systems

(9) 
$${}^{\mathfrak{p}}\mathcal{H}^{\alpha}(Rf_*IC_X) \cong \bigoplus_{\beta} IC_{\overline{S_{\beta}}}(L_{\alpha,S_{\beta}})$$

Combining (8) and (9) we have

(10) 
$$Rf_*IC_X \cong \bigoplus_{\alpha \in \mathbb{Z}} {}^{\mathfrak{p}}\mathcal{H}^{\alpha}(Rf_*IC_X)[-\alpha] \cong \bigoplus_{\alpha \in \mathbb{Z}} \bigoplus_{\beta} IC_{\overline{S_{\beta}}}(L_{\alpha,S_{\beta}})[-\alpha],$$

which can be written in the form

$$Rf_*IC_X \cong \bigoplus_{\alpha \in \mathbb{Z}} {}^{\mathfrak{p}}\mathcal{H}^{\alpha}(Rf_*IC_X)[-\alpha] \cong \bigoplus_{\alpha \in \mathbb{Z}} \bigoplus_{S} {}^{\mathfrak{p}}\mathcal{H}^{\alpha}(Rf_*IC_X)_S[-\alpha],$$

where S, called a **support** of f, is any  $\overline{S_{\beta}}$  associated to a non-zero local system  $L_{\alpha,S_{\beta}}$  (see [14, Definition 9.3.41]). In the literature one can find different approaches to the Decomposition Theorem (see [1], [15], [5], [16]), which is a very general result but also rather implicit. On the other hand, there are many special cases for which the Decomposition Theorem admits a simplified and explicit approach. One of these is the case of varieties with isolated singularities. For instance, in the work [10], a simplified approach to the Decomposition Theorem for varieties with isolated singularities is developed, in connection with the existence of a *natural Gysin morphism*, as defined in [7, Definition 2.3] (see also [8] for other applications of the Decomposition Theorem to the Noether-Lefschetz Theory).

As remarked before, the Bott-Samelson resolution

$$\pi_w: X(w) \to X(w),$$

is equivariant under the action of the Borel subgroup B, hence  $\pi_w$  is stratified according to the Bruhat decomposition

$$X(w) = \bigsqcup_{v \le w} \Omega(v).$$

In this case, the supports of the decomposition theorem applied to the resolution  $\pi_w$  are the Schubert subvarieties

$$X(v) = \overline{\Omega}(v), \qquad v \le w$$

Furthermore, all local systems appearing in the decomposition theorem are trivial since the isotropy subgroup of each orbit  $\Omega(v)$  is connected [13, Remark 11.6.2].

We include in the following proposition a proof of these facts, although they are probably well-known, in the attempt of making the present paper reasonably self-contained and also because the simple argument is very close to the rest of the paper.

**Proposition 3.2.** Let  $\pi : \tilde{X} \to X$  be an equivariant resolution of a Schubert variety X = X(w) of dimension l. In the derived category  $D_c^b(X)$ , we have a splitting

$$R\pi_*(\mathbb{Q}_{\tilde{X}})[l] = \bigoplus_{v \le w} \bigoplus_{\alpha \in \mathbb{Z}} IC_{X(v)}^{\oplus s_{v,\alpha}}[-\alpha],$$

for suitable multiplicities  $s_{v,\alpha}$ . In other words, the supports of the decomposition theorem applied to the resolution  $\pi$  are the Schubert varieties contained in X and all local systems are trivial.

*Proof.* Let l := l(w), fix r such that  $-1 \le r \le l$  and define the following decreasing sequence of open sets of X

$$\mathcal{U}_r := X(w) \setminus \bigsqcup_{v \le w, l(v) \le r} \Omega(v).$$

Clearly we have  $\mathcal{U}_l = \emptyset$  and  $\mathcal{U}_{-1} = X$ . We are going to prove, by decreasing induction on r, that

(11) 
$$R\pi_*(\mathbb{Q}_{\tilde{X}})[l] \mid_{\mathcal{U}_r} = \bigoplus_{v \le w, \, r < l(v)} \bigoplus_{\alpha \in \mathbb{Z}} IC_{X(v)}^{\oplus s_{v,\alpha}}[-\alpha] \mid_{\mathcal{U}_r},$$

for suitable multiplicities  $s_{v,\alpha}$ .

Since  $\mathcal{U}_{l-1} = \Omega := \Omega(w)$  and since  $\pi$  is an isomorphism over  $\Omega$ , we have

 $R\pi_*\mathbb{Q}_{\tilde{X}}[l]\mid_{\Omega}\cong\mathbb{Q}_{\Omega}[l],$ 

hence the *base step* follows from the well known isomorphism

$$\mathbb{Q}_{\Omega}[l] \cong IC_X \mid_{\Omega}$$

(compare e.g. with [14, Definition 6.3.1]). As for the *inductive step*, let

$$\mathcal{D} := \bigsqcup_{v \le w, \, r = l(v)} \Omega(v).$$

By induction, we have

$$R\pi_*(\mathbb{Q}_{\tilde{X}})[l] \mid_{\mathcal{U}_r} = \bigoplus_{v \le w, \, r < l(v)} \bigoplus_{\alpha \in \mathbb{Z}} IC_{X(v)}^{\oplus s_{v,\alpha}}[-\alpha] \mid_{\mathcal{U}_r}$$

hence we deduce

(12) 
$$R\pi_*(\mathbb{Q}_{\tilde{X}})[l] \mid_{\mathcal{U}_{r-1}} = \mathcal{L} \mid_{\mathcal{D}} \bigoplus \bigoplus_{v \le w, r < l(v)} \bigoplus_{\alpha \in \mathbb{Z}} IC_{X(v)}^{\oplus s_{v,\alpha}}[-\alpha] \mid_{\mathcal{U}_{r-1}},$$

where  $\mathcal{L}$  gathers all the summands supported in  $\mathcal{U}_r^* = \bigsqcup_{v \le w, \, l(v) \le r} \Omega(v)$ 

$$\mathcal{L} := \bigoplus_{\alpha \in \mathbb{Z}} \bigoplus_{S \subseteq \mathcal{U}_r^*} {}^{\mathfrak{p}} \mathcal{H}^{\alpha}(R\pi_*(\mathbb{Q}_{\tilde{X}})[l])_S[-\alpha].$$

In the previous formula, the summand  ${}^{\mathfrak{p}}\mathcal{H}^{\alpha}(R\pi_*(\mathbb{Q}_{\tilde{X}})[l])_S$  denotes the S component of  ${}^{\mathfrak{p}}\mathcal{H}^{\alpha}(R\pi_*(\mathbb{Q}_{\tilde{X}})[l])$ in the decomposition by supports [4, Section 1.1]. By proper base change we also have

(13) 
$$R\pi_*(\mathbb{Q}_{\tilde{X}})[l] \mid_{\mathcal{D}} = \mathcal{L} \mid_{\mathcal{D}} \oplus \bigoplus_{v \le w, \, r < l(v)} \bigoplus_{\alpha \in \mathbb{Z}} IC_{X(v)}^{\oplus s_{v,\alpha}}[-\alpha] \mid_{\mathcal{D}}.$$

Since  $\pi : \tilde{X} \to X$  is equivariant and  $\mathcal{D}$  is a disjoint union of *B*-orbits, the dimension of the cohomology stalk  $\mathcal{H}^i R \pi_*(\mathbb{Q}_{\tilde{X}})[l]_x$  is independent of  $x \in \mathcal{D}$ , for all *i*. The same holds true also for all  $IC_{X(v)}[-\alpha] \mid_{\mathcal{D}}$ . Then (13) shows that the dimension of the cohomology stalk  $\mathcal{H}^i \mathcal{L}_x$  is independent of  $x \in \mathcal{D}$ , for all *i*. Thus  $\mathcal{L} \mid_{\mathcal{D}}$  is a direct sum of shifted local systems because, by [5, Remark 1.5.1], the perverse cohomology sheaves  ${}^{\mathfrak{p}}\mathcal{H}^i(\mathcal{L} \mid_{\mathcal{D}})$  concide, up to a shift, with the ordinary cohomology

$${}^{\mathfrak{p}}\mathcal{H}^{i}(\mathcal{L}\mid_{\mathcal{D}})\cong\mathcal{H}^{i-r}\mathcal{L}\mid_{\mathcal{D}}[r],\quad\forall i$$

We are done, because  $\mathcal{D} = \bigsqcup_{v \leq w, r = l(v)} \Omega(v)$  is a disjoint union of affine spaces of dimension r (compare e.g. with [13, Theorem 9.9.5 (i)]) so any local system on  $\mathcal{D}$  is trivial and (11) follows for the restriction to the open set  $\mathcal{U}_{r-1}$ .

### 4. A CONSEQUENCE OF THE DECOMPOSITION THEOREM

In this section we assume to have fixed an equivariant resolution  $\pi_w : \tilde{X}(w) \to X(w)$ , for any Schubert variety  $X(w), w \in W$ . As a consequence of Proposition 3.2, the decomposition theorem for  $\pi_w$  can be stated as

(14) 
$$(R\pi_w)_* \mathbb{Q}_{\tilde{X}(w)}[l(w)] = \bigoplus_{v \le w} \bigoplus_{\alpha \in \mathbb{Z}} IC_{X(v)}^{\oplus s_{v,\alpha}^w}[-\alpha],$$

for suitable multiplicities  $s_{v,\alpha}^w$ , where recall that  $l(w) = \dim X(w)$ .

Notations 4.1. For any pair (v, w) of permutations such that  $v \leq w$ , let

(15) 
$$D_{w,v}(t) := \sum_{\alpha \in \mathbb{Z}} s_{v,\alpha}^{w} \cdot t^{\alpha} \in \mathbb{Z}[t, t^{-1}]$$

be the Laurent polynomial recording the contribution to the decomposition theorem (14) coming from support X(v). Let moreover

(16) 
$$F_{w,v}(t) := \sum_{\alpha \in \mathbb{Z}} f_{w,v}^{l(w) + \alpha} t^{\alpha} \in \mathbb{Z}[t, t^{-1}], \quad f_{w,v}^i := \dim H^i(\pi_w^{-1}(x)), \ x \in \Omega(v),$$

be the shifted Poincaré polynomial of the fibers of  $\pi_w$  over  $\Omega(v)$ .

### DAVIDE FRANCO

**Remark 4.2.** Let X(w) be a Schubert variety and let  $\Omega(v) \subseteq X(w)$  be a Schubert cell. It is well known the dimensions of the stalks  $\mathcal{H}^{\alpha}(IC_{X(w)})_x$  do not depend on  $x \in \Omega(v)$ .

The Laurent polynomial encoding the dimensions of the stalks  $\mathcal{H}^{\alpha}(IC^{\bullet}_{\mathcal{S}_{\tau}})_x$  is the shifted Kazhdan-Lusztig polynomial:

$$H_{w,v}(t) := \sum_{\alpha \in \mathbb{Z}} h_{w,v}^{\alpha} t^{\alpha}, \quad h_{w,v}^{\alpha} := \dim \mathcal{H}^{\alpha}(IC_{X(w)})_x, \ x \in \Omega(v)$$

Recall that we have

(17) 
$$P_{w,v}(q) = q^{\frac{l(w)}{2}} H_{w,v}(\sqrt{q})$$

where  $P_{w,v}(q)$  is the corresponding Kazhdan-Lusztig polynomial (2) (compare e.g. with [2, Theorem 6.1.11]).

Before stating the main result of this section, let us introduce the truncation U and symmetrizing S operators:

$$U: \sum_{\alpha \in \mathbb{Z}} a_{\alpha} t^{\alpha} \in \mathbb{Z}[t, t^{-1}] \mapsto \sum_{\alpha \ge 0} a_{\alpha} t^{\alpha} \in \mathbb{Z}[t];$$
  
$$S: \sum_{\alpha \ge 0} a_{\alpha} t^{\alpha} \in \mathbb{Z}[t] \mapsto a_{0} + \sum_{\alpha > 0} a_{\alpha} (t^{\alpha} + t^{-\alpha}) \in \mathbb{Z}[t, t^{-1}].$$

**Theorem 4.3.** With notations as above, let  $u \leq w$  in W. Then we have the following recursive formulae for the computation of the Laurent polynomials  $D_{w,u}$  and the shifted Kazhdan-Lusztig polynomials  $H_{w,u}$ :

$$\begin{cases} D_{w,u} = S \circ U(R_{w,u}) \\ H_{w,u} = t^{-l(u)}(R_{w,u} - D_{w,u}) \end{cases}$$

where

$$R_{w,u} := t^{l(u)} \left( F_{w,u} - \sum_{u < v < w} D_{w,v} \cdot H_{v,u} \right)$$

*Proof.* For the sake of simplicity, in the proof we set  $\pi : \tilde{X} \to X$  instead of  $\pi_w : \tilde{X}(w) \to X(w)$ . By Proposition 3.2, we have

(18) 
$$R\pi_*\mathbb{Q}_{\tilde{X}}[l] = \bigoplus_{v \le w} \bigoplus_{\alpha \in \mathbb{Z}} IC_{X(v)}^{\oplus s_{w,\alpha}^w}[-\alpha].$$

Consider a cell  $\Omega(u) \subset X$ , take the stalk cohomology at  $x \in \Omega(u)$  and recall (15) and (16). From (18) we infer

(19) 
$$F_{w,u}(t) = \sum_{u \le v \le w} D_{w,v}(t) \cdot H_{v,u}(t) = D_{w,w}(t) \cdot H_{w,u}(t) + D_{w,u}(t) \cdot H_{u,u}(t) + \sum_{u < v < w} D_{w,v}(t) \cdot H_{v,u}(t).$$

Since the resolution  $\pi : \tilde{X} \to X$  is equivariant, it must be an isomorphism over  $\Omega(w)$  and we have  $D_{w,w}(t) = 1$  (recall (14) and (15)). Furthermore, from the well known isomorphism  $IC_{X(u)}|_{\Omega(u)} \cong \mathbb{Q}_{\Omega(u)}[l(u)]$  (compare e.g. with [14, Definition 6.3.1]) we deduce  $H_{u,u}(t) = t^{-l(u)}$ . Hence, from (20) we get

(20) 
$$F_{w,u}(t) = H_{w,u}(t) + t^{-l(u)} \cdot D_{w,u}(t) + \sum_{u < v < w} D_{w,v}(t) \cdot H_{v,u}(t).$$

The last equality can be written as

$$D_{w,u}(t) = t^{l(u)} \cdot \left( F_{w,u}(t) - \sum_{u < v < w} D_{w,v}(t) \cdot H_{v,u}(t) \right) - t^{l(u)} \cdot H_{w,u}(t) = R_{w,u} - t^{l(u)} \cdot H_{w,u}(t).$$

The support conditions for perverse sheaves imply that  $t^{l(u)} \cdot H_{w,u}(t)$  is concentrated in negative degrees (see [5, p. 552, equation 12]), thus

$$U(D_{w,u}(t)) = U(R_{w,u}(t)).$$

Finally, the Laurent polynomials  $D_{w,u}(t)$  are symmetric because of Hard-Lefschetz theorem (see [5, Theorem 1.6.3]), that is to say

$$D_{w,u}(t) = D_{w,u}(t^{-1}).$$

Therefore we have

(21) 
$$\begin{cases} D_{w,u}(t) = S \circ U(R_{w,u}(t)) \\ H_{w,u}(t) = t^{-l(u)}(R_{w,u}(t) - D_{w,u}(t)) \end{cases}$$

and the statement follows.

By (17), the last theorem provides an iterative procedure to compute Kazhdan-Lusztig polynomials  $P_{w,v}$  from Poincaré polynomials  $Q_{w,v}$ . To this end, let us introduce the following operators:

$$U_{\beta}: \sum_{\alpha \ge 0} c_{\alpha} t^{\alpha} \in \mathbb{Z}[t] \mapsto \sum_{\alpha \ge \beta} c_{\alpha} t^{\alpha} \in \mathbb{Z}[t], \ \forall \beta \ge 0.$$

Next statement follows from Theorem 4.3 and collects all informations we obtained until now.

**Corollary 4.4.** Assume to have fixed an equivariant resolution  $\pi_w : \tilde{X}(w) \to X(w)$ , for any Schubert variety X(w),  $w \in W$ . For any pair w, u in W such that  $u \leq w$ , let  $\tilde{F}_{w,u}$  be the Poincaré polynomials of the fiber  $\pi_w^{-1}(x)$ ,  $\forall x \in \Omega(u)$ . Then we have the following recursive formulae:

$$\begin{cases} \tilde{D}_{w,u} = t^{l(w)-l(u)} \circ S \circ t^{l(u)-l(w)} \circ U_{l(w)-l(u)}(\tilde{R}_{w,u}) \\ \tilde{H}_{w,u} = \tilde{R}_{w,u} - \tilde{D}_{w,u} \end{cases}$$

where

$$\tilde{D}_{w,u} := t^{l(w) - l(u)} D_{w,u}, \quad \tilde{H}_{w,u} := t^{l(w)} H_{w,u}, \quad \tilde{R}_{w,u} := \tilde{F}_{w,u} - \sum_{u < v < w} \tilde{D}_{w,v} \cdot \tilde{H}_{v,u}$$

*Proof.* From (16) we find  $\tilde{F}_{w,u} = t^{l(w)}F_{w,u}$ . Thus we have

$$\tilde{R}_{w,u} := \tilde{F}_{w,u} - \sum_{u < v < w} \tilde{D}_{w,v} \cdot \tilde{H}_{v,u} = t^{l(w)} F_{w,u} - \sum_{u < v < w} t^{l(w) - l(u)} D_{w,v} \cdot t^{l(v)} H_{v,u} = t^{l(w) - l(v)} R_{w,u},$$

and the statement straightforwardly follows just combining Theorem 4.3 with

$$t^{l(u)-l(w)} \circ U_{l(w)-l(u)} \circ t^{l(w)-l(v)} = U_0.$$

**Remark 4.5.** By (17), we have

$$P_{w,v}(q) = q^{\frac{l(w)}{2}} H_{w,v}(\sqrt{q}) = \tilde{H}_{w,v}(\sqrt{q}),$$

hence previous corollary provides an iterative procedure that allows to compute **both** the Kazhdan-Lusztig polynomials  $P_{w,v}$  and the Laurent polynomials  $D_{w,v}$ .

#### 5. Bases for the Hecke Algebra

As in the previous section we assume to have fixed an equivariant resolution  $\pi_w$  for any Schubert variety X(w),  $w \in W$  and we assume in addition that the cohomology of the fibers of  $\pi_w$  vanish in odd degrees (this property is satisfied by any reasonable resolution of Schubert varieties). As a consequence, we have that the coefficients of the polynomials  $\tilde{F}_{w,v}$  vanish in odd degree. Furthermore, from the relations

$$\begin{cases} \tilde{D}_{w,u} = t^{l(w)-l(u)} \circ S \circ t^{l(u)-l(w)} \circ U_{l(w)-l(u)}(\tilde{R}_{w,u}) \\ \tilde{R}_{w,u} := \tilde{F}_{w,u} - \sum_{u < v < w} \tilde{D}_{w,v} \cdot \tilde{H}_{v,u} \end{cases}$$

one deduces immediately that the same holds true for the polynomials  $D_{w,v}$ . We define

(22) 
$$Q_{w,v}(q) = \tilde{F}_{w,v}(\sqrt{q}) \in \mathbb{Z}[q],$$

(23) 
$$S_{w,v}(q) = D_{w,v}(\sqrt{q}) \in \mathbb{Z}[q]$$

for all  $v, w \in W$  such that  $v \leq w$ . Our aim in this section is to define a new basis for the Hecke algebra by means of the polynomials  $Q_{w,v}(q)$ . We start by recalling the definition of Hecke algebra.

Let  $\mathcal{H}$  be the *Hecke algebra* of W i.e. the algebra over  $\mathbb{Z}[q^{\frac{1}{2}}, q^{-\frac{1}{2}}]$  with basis elements  $\{T_w \mid w \in W\}$  and relations [2, Sec. 6.1]

$$\begin{cases} T_{s_i} T_w = T_{s_i w} & \text{if } l(s_i w) > l(w), \\ T_{s_i} T_{s_i} = (q-1) T_{s_i} + q T_{id}. \end{cases}$$

The Hecke algebra is also equipped with the Kazhdan-Lusztig basis  $\{C_w \mid w \in W\}$  where

(24) 
$$C_w = T_w + \sum_{v < w} P_{w,v} T_v$$

where  $P_{w,v} \in \mathbb{Z}[q]$  are the Kazhdan-Lusztig polynomials.

**Theorem 5.1.** For any  $w \in W$ , let

$$B_w = T_w + \sum_{v < w} Q_{w,v} T_v \in \mathcal{H}.$$

The polynomials  $S_{w,v}(q) \in \mathbb{Z}[q]$  are the coefficients of  $B_w$  with respect to the Kazhdan-Lusztig basis

$$B_w = C_w + \sum_{v < w} S_{w,v} C_v \in \mathcal{H}.$$

*Proof.* From (20) and recalling  $\tilde{F}_{w,u} = t^{l(w)}F_{w,u}$ , we get

(25) 
$$\tilde{F}_{w,u} = t^{l(w)} F_{w,u} = \sum_{u \le v \le w} t^{l(w) - l(v)} D_{w,v} \cdot t^{l(v)} H_{v,u} = \sum_{u \le v \le w} \tilde{D}_{w,v} \cdot \tilde{H}_{v,u},$$

where we have taken into account 4.4. Combining 4.5 with (22) and (23) and evaluating in  $t = \sqrt{q}$  the last equality, we get

(26) 
$$Q_{w,u} = \sum_{u \le v \le w} S_{w,v} \cdot P_{v,u} \in \mathbb{Z}[q].$$

Since the resolution  $\pi: \tilde{X} \to X$  is equivariant, it must be an isomorphism over  $\Omega(w)$ , so we have  $Q_{w,w} = 1$  and

$$B_w = T_w + \sum_{u < w} Q_{w,u} T_u = \sum_{u \le w} Q_{w,u} T_u = \sum_{u \le w} \left( \sum_{u \le v \le w} S_{w,v} \cdot P_{v,u} \right) T_u = \sum_{v \le w} S_{w,v} \left( \sum_{u \le v} P_{v,u} T_u \right).$$

Again, since  $\pi_w$  an isomorphism over  $\Omega(w)$  and we have  $D_{w,w} = S_{w,w} = 1$  (recall (14) and (15)) and the last sum can be written as

$$\sum_{u \le w} P_{w,u} T_u + \sum_{v < w} S_{w,v} \left( \sum_{u \le v} P_{v,u} T_u \right)$$

that coincides with

$$C_w + \sum_{v < w} S_{w,v} C_v$$

in view of (24).

**Remark 5.2.** Theorem above shows that the transition matrix, from the Kazhdan-Lusztig basis  $\{C_w \mid w \in W\}$  to the new one  $\{B_w \mid w \in W\}$ , is triangular with coefficients  $S_{w,v} \in \mathbb{Z}[q]$ . In this work we have set up an iterative procedure that allows the computation of such a matrix and which does not require prior knowledge of the transition matrix from the Kazhdan-Lusztig basis to the standard one  $\{T_w \mid w \in W\}$ .

#### Conflicts of interest

The author has no conflict of interest to declare that are relevant to this article.

#### DAVIDE FRANCO

#### References

- A. A. Beĭlinson, J. Bernstein, and P. Deligne, <u>Faisceaux pervers</u>, Analysis and topology on singular spaces, I (Luminy, 1981), Astérisque, vol. 100, Soc. Math. France, Paris, 1982, pp. 5–171.
- [2] Sara Billey and V. Lakshmibai, <u>Singular loci of Schubert varieties</u>, Progress in Mathematics, vol. 182, Birkhäuser Boston, Inc., Boston, MA, 2000.
- [3] Francesca Cioffi, Davide Franco, and Carmine Sessa, <u>An effective decomposition theorem for Schubert varieties</u>, J. Symbolic Comput. **121** (2024), 102238. DOI: 10.1016/j.jsc.2023.102238
- [4] Mark Andrea A. de Cataldo, <u>Hodge-theoretic splitting mechanisms for projective maps</u>, J. Singul. 7 (2013), 134-156.
- [5] Mark Andrea A. de Cataldo and Luca Migliorini, <u>The decomposition theorem</u>, perverse sheaves and the topology of algebraic maps, Bull. Amer. Math. Soc. (N.S.) 46 (2009), no. 4, 535–633.
- [6] Vinay V. Deodhar, <u>A combinatorial setting for questions in Kazhdan-Lusztig theory</u>, Geom. Dedicata 36 (1990), no. 1, 95–119.
- [7] Vincenzo Di Gennaro and Davide Franco, <u>On the existence of a Gysin morphism for the blow-up of an ordinary singularity</u>, Ann. Univ. Ferrara Sez. VII Sci. Mat. **63** (2017), no. 1, 75–86.
- [8] Vincenzo Di Gennaro and Davide Franco, Néron-Severi group of a general hypersurface, Commun. Contemp. Math., 19 (2017), no. 1, 1650004, 15 pp. DOI: 10.1142/S0219199716500048
- [9] Vincenzo Di Gennaro and Davide Franco, <u>On a resolution of singularities with two strata</u>, Results Math. 74 (2019), no. 3, Paper No. 115, 22 pp. DOI: 10.1007/s00025-019-1040-9
- [10] Vincenzo Di Gennaro and Davide Franco, <u>On the topology of a resolution of isolated singularities</u>, II, J. Singul. 20 (2020), 95–102.
- [11] Davide Franco, <u>Explicit decomposition Theorem for special Schubert varieties</u>, Forum Math. **32** (2020), no. 2, 447-470.
- [12] Mark Goresky and Robert MacPherson, Intersection homology. II, Invent. Math. 72 (1983), no. 1, 77–129.
- [13] Ryoshi Hotta, Kiyoshi Takeuchi, and Toshiyuki Tanisaki, <u>D-modules, perverse sheaves, and representation</u> <u>theory</u>, Progress in Mathematics, vol. 236, Birkhäuser Boston, Inc., Boston, MA, 2008, Translated from the 1995 Japanese edition by Takeuchi. MR 2357361
- [14] Laurenţiu G. Maxim, Intersection homology & perverse sheaves—with applications to singularities, Graduate Texts in Mathematics, vol. 281, Springer, Cham, 2019.
- [15] Morihiko Saito, Mixed Hodge modules, Proc. Japan Acad. Ser. A Math. Sci. 62 (1986), no. 9, 360–363.
- [16] Geordie Williamson, <u>The Hodge theory of the decomposition theorem</u>, no. 390, 2017, Séminaire Bourbaki. Vol. 2015/2016. Exposés 1104–1119, pp. Exp. No. 1115, 335–367.

UNIVERSITÀ DI NAPOLI "FEDERICO II", DIPARTIMENTO DI MATEMATICA E APPLICAZIONI "R. CACCIOP-POLI", VIA CINTIA, 80126 NAPOLI, ITALY.

Email address: davide.franco@unina.it