AN EQUIVARIANT COMPACTIFICATION FOR ADJOINT REDUCTIVE GROUP SCHEMES

SHANG LI

ABSTRACT. Wonderful compactifications of adjoint reductive groups over an algebraically closed field play an important role in algebraic geometry and representation theory. In this paper, we construct an equivariant compactification for adjoint reductive groups over arbitrary base schemes. Our compactifications parameterize classical wonderful compactifications of De Concini and Procesi as geometric fibers. Our construction is based on a variant of the Artin–Weil method of birational group laws. In particular, our construction gives a new intrinsic construction of wonderful compactifications. The Picard group scheme of our compactification in the study of torsors under reductive group schemes.

CONTENTS

1.	Introduction	1
2.	Recollection of wonderful compactifications	5
3.	Rational action on big cells	6
4.	Compactification for split adjoint reductive group schemes	11
5.	Divisors	20
6.	Descent	23
7.	Applications	28
References		29

1. INTRODUCTION

Reductive groups play a fundamental role both in algebraic geometry and number theory. Seeking for an equivariant compactification of reductive groups is beyond any doubt an attractive and important problem, see the ICM report [Spr06] for recent development on this topic. The wonderful compactification for adjoint semisimple groups over algebraically closed fields is a good solution to this problem. The wonderful compactifications have a wide range of applications, for instance, in the study of character sheaves (see, *loc.cit*, [Gin89] and [Lus04]) and in the study of local models of Shimura varieties (see [He13] and [PRS13, Section 8]).

In this article, for an adjoint reductive group scheme \mathbf{G} over a scheme S, we construct the following equivariant compactification for \mathbf{G} :

Theorem 1.1. (Corollary 6.4). There is a smooth projective S-scheme \mathcal{X} containing **G** as an open dense subscheme such that \mathcal{X} is equipped with a $(\mathbf{G} \times \mathbf{G})$ -action which extends the left and the right translations of **G** on itself. Each geometric fiber of \mathcal{X} is identified with the wonderful compactification of the corresponding geometric fiber of **G**. The boundary $\mathcal{X} \setminus \mathbf{G}$ is a smooth relative Cartier divisor with relative normal crossings to S.

Date: October 30, 2023.

In particular, if the base scheme S in Theorem 1.1 is the spectrum of an algebraically closed field, the scheme \mathcal{X} recovers the classical wonderful compactification of \mathbf{G} . Our construction of \mathcal{X} in Theorem 1.1 is intrinsic and functorial in the sense that we avoid embedding the group \mathbf{G} into an ambient projective space, and we use a sheaf-theoretic argument so that the formation of \mathcal{X} is compatible with any base change.

In [Ces22, Conjecture 6.2.3], Cesnavičius raises a question about how to equivariantly embed a reductive group scheme into a projective scheme in which the group is fiberwise dense. Theorem 1.1 gives a positive answer to this question for adjoint reductive group schemes. As a consequence of Theorem 1.1, we deduce the following lifting property of torsors:

Corollary 1.2. (Proposition 7.2). Let A be an isotrivial torsor under an isotrivial adjoint reductive group **G** over an affine semilocal scheme S. For $a \in A(Z)$ with Z a closed subscheme of S, there exists a finite étale cover \widetilde{S} of S, a morphism $\nu: Z \to \widetilde{S}$ and a section $\widetilde{a} \in A(\widetilde{S})$ whose ν -pullback is a.

Indeed, by a twist argument, we deduce from Theorem 1.1 an equivariant compactification containing the torsor A as a fiberwise dense open subscheme. Then we use the Bertini theorem to each closed fiber of the compactification of A to cut out the desired finite étale cover \tilde{S} . For a further consequence of Theorem 1.1 that concerns the techniques of equating reductive (not necessarily adjoint) group schemes, see Proposition 7.4.

1.1. Classical wonderful compactifications. Let us briefly review the theory of wonderful compactifications. Consider an adjoint reductive group G over an algebraically closed field k. The wonderful compactification X of G was introduced by De Concini and Procesi [DCP83] in characteristic zero, and by Strickland [Str87] in arbitrary characteristic, and it satisfies the following properties:

- X is a smooth projective variety containing G as an open dense subvariety such that X is equipped with an action of $G \times_k G$ that extends the action $(g_1, g_2) \cdot g = g_1 g g_2^{-1}$ of $G \times_k G$ on G;
- the boundary $X \setminus G$ is a union of $(G \times_k G)$ -stable smooth divisors X_i (i = 1, ..., l) with normal crossings;
- the closures of the $(G \times_k G)$ -orbits are the partial intersections of the X_i .

The idea of their constructions is to define X as the closure of G in a projective space constructed via representation theory. In [Bri03], Brion gave an algebro-geometric realization of X by taking the closure of G in the Hilbert scheme of a certain flag variety of $G \times G$.

To investigate the arithmetic aspects of the wonderful compactifications, a basic questions is to seek an integral model of wonderful compactifications. In [Fal97], Faltings sketched an integral model of wonderful compactifications (for more general symmetric varieties) over $\operatorname{Spec}(\mathbb{Z}_p)$, where p is the characteristic of the base field k. In an unpublished work, Gabber established the wonderful compactification for a Chevalley group over $\operatorname{Spec}(\mathbb{Z})$.

The main feature of the wonderful compactification X that we are going to make use of is that X contains an open subvariety $\overline{\Omega}$ which is usually called the big cell of X and is isomorphic to the affine space $U^- \times_k \mathbb{A}_k^l \times_k U^+$, where l is the rank of G and U^+ (resp., U^-) is the unipotent radical of a Borel subgroup (resp., an opposite Borel subgroup). The geometry of X is, in many respects, largely controlled by $\overline{\Omega}$ mainly because translations of $\overline{\Omega}$ under the action of $G \times_k G$ cover X. For instance, the smoothness of X follows from that of $\overline{\Omega}$ (see, for instance, [BK05, Theorem 6.1.8 (i)]). Therefore, we may seek to recover X as a quotient of $G \times_k \overline{\Omega} \times_k G$. One advantage of this is that $\overline{\Omega}$ is much simpler than X in the sense that it has explicit coordinates, and after shrinking if needed, the action $G \times_k G$ on $\overline{\Omega}$ can be defined over Spec(\mathbb{Z}). This suggests for us to define a birational action of $G \times_k G$ on $\overline{\Omega}$ over more general base schemes, here a birational action is an analogue of the notion of a birational group law that was first due to Weil in the setting of varieties, and later was generalized to schemes by Artin (see [ER15]).

1.2. The proof of Theorem 1.1. The overall strategy of the proof is that we first establish Theorem 1.1 for split groups and then make a descent argument.

1.2.1. Split case. Now assume that **G** splits over S with respect to a split maximal torus **T**, choose a Borel subgroup **B** containing **T**, and denote the corresponding based root datum of **G** by $(X, \Psi, \Delta, X^{\vee}, \Psi^{\vee}, \Delta^{\vee})$. Then we have two big cells, and an immersion between them modeling on classical wonderful compactifications:

$$\Omega_{\mathbf{G}} := \mathbf{U}^- \times_S \mathbf{T} \times_S \mathbf{U}^+ \longrightarrow \overline{\Omega} := \mathbf{U}^- \times_S \overline{\mathbf{T}} \times_S \mathbf{U}^+,$$

that is the identity on \mathbf{U}^+ and \mathbf{U}^- , and sends \mathbf{T} to $\overline{\mathbf{T}}$ via negative simple roots, where $\overline{\mathbf{T}}$ is the product $\prod_{\Delta} \mathbb{G}_a$. In Theorem 3.4, we obtain a rational action of $\mathbf{G} \times_S \mathbf{G}$ on $\overline{\Omega}$ which is a rational morphism

$$\pi \colon \mathbf{G} \times_S \overline{\Omega} \times_S \mathbf{G} \dashrightarrow \overline{\Omega}$$

such that for any $x \in S$, the geometric fiber $\pi_{\overline{k(x)}}$, on certain large enough open subscheme, coincides with the action of $(\mathbf{G})_{\overline{k(x)}} \times_{\overline{k(x)}} (\mathbf{G})_{\overline{k(x)}}$ on the big cell $\overline{\Omega}_{\overline{k(x)}}$ of the wonderful compactification of $\mathbf{G}_{\overline{k(x)}}$ over the algebraically closed field $\overline{k(x)}$. The construction of π proceeds as follows: we first produce a rational endomorphism f (resp., f') of $\overline{\Omega}$ that fiberwise is the conjugation by w_0 (resp., w_0^{-1}) where w_0 is the element of longest length in the Weyl group (Lemma 3.2). With f and f' in hand, to define the image of a section A of $\mathbf{G} \times_S \overline{\Omega} \times_S \mathbf{G}$, the key idea is to apply f to the component of A on U^- and that on U^+ to switch them into the correct place so that we could apply f' to "multiply" them, while over each geometric fiber the action of f and f' are offset, see the proof of Lemma 3.3 and Theorem 3.4 for details. The similar strategy is used to prove the Existence Theorem, see [GP11b, Exposé XXV].

Using the rational action π , we define the following equivalence relation (Definition 4.1) on $\mathbf{G} \times \overline{\Omega} \times \mathbf{G}$: for every S-scheme S' and (g_1, x, g_2) , $(g'_1, x', g'_2) \in (\mathbf{G} \times_S \overline{\Omega} \times_S \mathbf{G})(S')$, we say that (g_1, x, g_2) and (g'_1, x', g'_2) are equivalent if and only if, there exist an fppf cover $S'' \to S'$ and a section $(a_1, a_2) \in (\mathbf{G} \times_S \mathbf{G})(S'')$ such that $\pi(a_1g_1, x, a_2g_2)$ and $\pi(a_1g'_1, x', a_2g'_2)$ are both well defined and are equal over S''.

One advantage of the above definition is that, although π is merely defined over an open subscheme of $\mathbf{G} \times_S \overline{\Omega} \times_S \mathbf{G}$, one can use $\mathbf{G} \times_S \mathbf{G}$ -translations to bring two sections into the definition domain of π so that they can be compared via π . We can then define the compactification \mathcal{X} of \mathbf{G} as a sheaf by taking the quotient of $\mathbf{G} \times_S \overline{\Omega} \times_S \mathbf{G}$ with respect to this equivalence relation.

The proof of the fact that \mathcal{X} is a scheme involves two steps. We first reinterpret \mathcal{X} as a quotient of $\mathbf{G} \times_S \overline{\Omega} \times_S \mathbf{G}$ with respect to an fppf relation (Theorem 4.7), hence by a theorem due to Artin ([LMB00, Corollaire (10.4)]), \mathcal{X} is an algebraic space. Secondly, we endow \mathcal{X} with a group action of $\mathbf{G} \times_S \mathbf{G}$ (Definition-Proposition 4.9), so that the ($\mathbf{G} \times_S \mathbf{G}$)translations of the open subscheme $\overline{\Omega}$ of \mathcal{X} (Lemma 4.10) cover \mathcal{X} . Thus, the schematic nature of \mathcal{X} follows from a Zariski gluing of scheme-like open neighborhoods of points of \mathcal{X} , see [BLR90, Section 6.6, Theorem 2].

The scheme \mathcal{X} is projective over S (Corollary 4.15). Roughly the quasi-projectivity of \mathcal{X} comes from the relative ampleness of the boundary divisor $\mathcal{X}\setminus\overline{\Omega}$, see [BLR90, Section 6.6,

Theorem 2 (d)]. By a variant of the valuation criterion for properness and the Iwahori decomposition, we deduce the properness of \mathcal{X} by the toric structure of the closure of the maximal torus **T** in \mathcal{X} , see Theorem 4.14 for details.

To finish the proof of Theorem 1.1 in the split case, by the funtoriality of the definition of \mathcal{X} and the fiberwise condition on π , we deduce that the geometric fibers of \mathcal{X} conincide with the wonderful compactifications of the corresponding geometric fibers of \mathbf{G} , see Proposition 4.12.

1.2.2. Non-split case. Now we consider **G** as an adjoint reductive group scheme over S. By [GP11b, Exposé XXII, Corollaire 2.3], there exists an étale cover U of S and a split reductive S-group \mathbf{G}_0 , which is isomorphic to \mathbf{G} over U. Applying the split case of Theorem 1.1 to \mathbf{G}_0 to produce a compactification \mathcal{X}_0 , the rest of the proof is to descend $\widetilde{\mathcal{X}} := \mathcal{X}_0 \times_S U$ to get the sought \mathcal{X} for \mathbf{G} . Note that \mathbf{G} is descended from $\widetilde{\mathbf{G}} := \mathbf{G}_0 \times_S U$ via a Čech cocycle χ which is an element of $\operatorname{Aut}_{\mathbf{G}_0/S}(U \times_S U)$ together with a cocycle condition. Then by [GP11b, Exposé XXIV, Théoreme 1.3], χ is the composition of an inner automorphism δ and an outer automorphism η . The inner part δ naturally induces an automorphism of $\widetilde{\mathcal{X}}$ via the group action on $\widetilde{\mathcal{X}}$, and we show that the outer part η also induces an automorphism of $\widetilde{\mathcal{X}}$, see Lemma 6.1. Combining these two automorphisms on $\widetilde{\mathcal{X}}$ together, by the density of $\widetilde{\mathbf{G}}$ in $\widetilde{\mathcal{X}}$, we get a descent datum on $\widetilde{\mathcal{X}}$. To make this descent datum effective, we construct an ample divisor compatible with the descent datum, see Theorem 6.3. The group action and the fiberwise condition on $\widetilde{\mathcal{X}}$ also descend, see Corollary 6.4.

1.3. Divisors and relative Picard schemes. In Theorem 5.1, we show that the Picard scheme of \mathcal{X} in Theorem 1.1, when **G** is split, is a constant scheme associated to the free abelian group of rank equal to the rank of **G**. Roughly, base change reduces us to the case when the base S is Spec(\mathbb{Z}), and a fibral criterion [Gro67, Corollaire 17.9.5] further reduces us to the case when S is an algebraically closed field, which follows from the cooresponding result for the classical wonderful compactifications ([BK05, Lemma 6.19]).

We also show that when **G** is split, \mathcal{X} in Theorem 1.1 shares the same wonderful properties of classical wonderful compactifications as we recall in Section 1.1, see Proposition 5.2 and Theorem 5.3.

1.4. Notation and conventions. All rings are commutative and unital. For the n-fold product $X_1 \times X_2 \times ... \times X_n$ of schemes, for $\{i_1, i_2, ..., i_l\}$ which is a strict ascending sequence of $\{1, 2, ..., n\}$, we denote by $\operatorname{pr}_{i_1, i_2, ..., i_l}$ the projection from $X_1 \times X_2 \times ... \times X_n$ onto the product of the factors with indices $i_1, i_2, ..., i_l$.

We use dotted arrows to depict a rational morphism. For an S-rational morphism f between two schemes X and Y over a scheme S, a test S-scheme S' and a section $x \in X(S')$, we say that x is well defined with respect to f if the image of x in X lies in the definition domain of f. We denote by Dom(f) the definition domain of f. For the existence and the uniqueness of definition domain of a rational morphism, see [SGA3II, Exposé XVIII, Définition 1.5.1].

For an effective Cartier divisor \mathbf{D} of a scheme X, we write $\mathcal{L}_X(\mathbf{D})$ for the corresponding invertible sheaf of \mathbf{D} .

A reductive group scheme is assumed to be connected.

For a presheaf of groups \mathfrak{F} over a category \mathfrak{C} and $T \in ob(\mathfrak{C})$, denote by e the identity section of $\mathfrak{F}(T)$ if the object T is clear in the context.

For an affine scheme X and $f \in \Gamma(X, \mathcal{O}_X)$, we denote the closed (resp., open) subscheme of X defined by f as $V_X(f)$ (resp., $D_X(f)$).

1.5. Acknowledgements. I would like to thank my advisor Kestutis Česnavičius deeply for his instruction on mathematics and writing and his sustained encouragement. I am especially grateful to Ofer Gabber for sharing his insights that are crucial to this paper. I heartfully thank Alexis Bouthier for several very helpful conversations and suggestions. I thank Ning Guo, Arnab Kundu, Zhouhang Mao, Sheng Chen and Federico Scavia for helpful correspondence. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 851146).

2. Recollection of wonderful compactifications

In this section, we briefly review the classical theory of wonderful compactifications. We fix an algebraically closed field k, and if a fiber product is formed over k, we will omit the subscript k.

2.1. The group setup. Let G be an adjoint reductive group over k, whose rank is l, and let $G^{\rm sc}$ be its simply connected covering. Fix a Borel subgroup $\widetilde{B} \subset G^{\rm sc}$ that contains a maximal subtorus \widetilde{T} of \widetilde{G}^{sc} , then \widetilde{B} determines a system of simple roots $\Delta = \{\alpha_1, ..., \alpha_l\}$ for the root system Ψ of $G^{\rm sc}$ in the character lattice $X^*(\widetilde{T})$. We denote the resulting sets of positive roots and negative roots by Ψ^+ and Ψ^- . Let \widetilde{B}^- be the opposite Borel of \widetilde{B} such that $\widetilde{B}^- \cap \widetilde{B} = \widetilde{T}$, and let U^+ and U^- be the unipotent radicals of \widetilde{B} and \widetilde{B}^- . For a root $\alpha \in \Psi$, we write U_{α} for the root subgroup defined by α . Let T be the image of \widetilde{T} in G. We identify the set of roots of G^{sc} with that of G by the natural embedding $X^*(T) \hookrightarrow X^*(\widetilde{T})$.

We choose a character $\lambda \colon \widetilde{T} \to \mathbb{G}_m$ that is regular dominant in the sense that $\langle \lambda, \alpha^{\vee} \rangle > 0$ for every positive coroot α^{\vee} .

2.2. Construction of wonderful compactification. By [BK05, Lemma 6.1.1 and Remark 6.1.2], we can fix an algebraic finite-dimensional representation V_{λ} of $G^{\rm sc}$ such that

- λ is the highest weight of V_{λ} ;
- any weight of \widetilde{T} in the representation V_{λ} is lower than λ ;
- for a simple root $\alpha \in \Delta$, the weight space of $\lambda \alpha$ is nonzero.

By the adjointness of G and the regularity of λ , the group G is embedded into $\mathbb{P}(\text{End}(V_{\lambda}))$ in such a way that the following square commutes:



We define the wonderful compactification X of G to be the closure of G in $\mathbb{P}(\operatorname{End}(V_{\lambda}))$. The $G \times G$ -action on G naturally extends to X. By [BK05, Theorem 6.1.8 (iv)], the wonderful compactification X does not depend on the choice of λ and V_{λ} .

2.3. Big cells of wonderful compactifications. In order to study the geometry of the wonderful compactification X, we fix a basis $v_0, v_1, ..., v_n$ of V_{λ} consisting of eigenvectors of T such that v_0 has weight λ . Let $(v_i^*)_{0 \leq i \leq n}$ be the dual basis of $(v_i)_{0 \leq i \leq n}$ in the dual vector space V_{λ}^* . Consider the open subscheme

$$\overline{\Omega}(k) := \left\{ f \in X(k) \mid v_0^*(\widetilde{f}(v_0)) \neq 0 \text{ for some lift } \widetilde{f} \text{ of } f \text{ in } \operatorname{End}(V_\lambda) \right\} \subset X(k)$$

which is often referred as the big cell of X. By [BK05, Proposition 6.1.7], the big cell $\overline{\Omega}$ of X is isomorphic to $U^- \times \overline{T} \times U^+$, where $\overline{T} = \prod_{\Delta} \mathbb{G}_a$.

By [BK05, Theorem 6.1.8 (i)], the $(G \times G)$ -translations of the big cell $\overline{\Omega}$ cover X i.e.,

$$X = \bigcup_{(g_1, g_2) \in G \times G} (g_1, g_2) \cdot \overline{\Omega}.$$
 (1)

3. RATIONAL ACTION ON BIG CELLS

In this section, we define a rational action of a product of an adjoint semisimple group scheme G on an affine space which models a big cell of a wonderful compactification. Passing to each geometric fiber, this rational action reflects the group action on wonderful compactifications. We then study various properties of this rational action. These properties are crucial for our constructions in Section 4.

3.1. The setup. Let G be an adjoint split reductive group scheme over a scheme S, and choose a maximal split torus T as in [GP11b, Définition 1.13] and a Borel subgroup B containing T. In this section, if a fiber product is formed over S, the lower subscript S will be omitted. Assume that $\operatorname{rk}(T) = l$. We denote the Lie algebras of $T \subset B \subset G$ by $\mathfrak{t} \subset \mathfrak{b} \subset \mathfrak{g}$. Let $(X, \Psi, \Delta, X^{\vee}, \Psi^{\vee}, \Delta^{\vee})$ be the based root datum defined by the Borel B, and let Ψ^+ (resp., Ψ^-) be the set of positive (resp., negative) roots. Then we have the canonical decomposition of \mathfrak{g} into root spaces: $\mathfrak{g} = \bigoplus_{\alpha \in \Psi} \mathfrak{g}_{\alpha}$. We fix an enumeration $\{\alpha_1, ..., \alpha_l\}$ of Δ .

According to [GP11b, Exposé XXIII, Prosposition 6.2], we can fix a Chevalley system $(X_{\alpha} \in \Gamma(S, \mathfrak{g}_{\alpha})^{\times})_{\alpha \in \Psi}$ for G. Each X_{α} gives rise to an isomorphism of groups

$$p_{\alpha} \colon \mathbb{G}_{a,S} \longrightarrow U_{\alpha}, \ \alpha \in \Psi.$$

In the following, we identify the coordinate function of U_{α} with the coordinate function of $\mathbb{G}_{a,S}$ via p_{α} .

We define a morphism:

$$\nu \colon \Omega_G := U^- \times T \times U^+ \longrightarrow \overline{\Omega} := U^- \times \overline{T} \times U^+ \tag{2}$$

via

$$(u', t, u) \mapsto (u', \prod_{\alpha_i \in \Delta} -\alpha_i(t), u),$$

where $\overline{T} := \prod_{\Delta} \mathbb{G}_a$. In the rest of this paper, the subscheme \overline{T} is canonically identified with the subschem $\{e\} \times \overline{T} \times \{e\} \subset \overline{\Omega}$, and similarly for U^- and U^+ . By the adjointness of G, this ν is a momomorphism which is an immersion by appealing to [SGA3II, Exposé XVI, Corollaire 1.5 a)] or [Con14, Theorem 5.3.5]. In the following, we will implicitly view Ω_G as a subscheme of $\overline{\Omega}$ via ν . In particular, T acts on \overline{T} via ν . For each simple root $\alpha_i \in \Delta$, we denote by \mathbb{X}_i the coordinate of \mathbb{G}_a indexed by α_i in $\overline{\Omega}$.

3.2. The rational action. We want to define an action of $G \times G$ on $\overline{\Omega}$. In the case of classical wonderful compactification, we know that their product may not lie in $\overline{\Omega}$. Hence, instead of defining the action on the entire big cell $\overline{\Omega}$, we seek to take advantage of "rational action" which is, roughly, an action defined over a fiberwise dense open subscheme of $\overline{\Omega}$, and which is similar to the notion of a birational group law [SGA3II, Exposé XVIII, Définition 3.1] (see also [BLR90, Chapter 4]). We require that this "rational action" should coincide geometrically fiberwise with the group actions on wonderful compactifications. This is done step by step by the subsequent lemmas.

Lemma 3.1. Let $\alpha_i \in \Delta$ and let $n_{\alpha_i} = p_{\alpha_i}(1)p_{-\alpha_i}(-1)p_{\alpha_i}(1) \in \operatorname{Norm}_G(T)(S)$, which is a representative of the simple reflection s_i defined by α_i in the Weyl group W. There exists a morphism $f_i: V_i \longrightarrow \overline{\Omega}$ such that

- (1) $V_i = D_{\overline{\Omega}}(\mathbb{X}_i + xy)$, where x, y are the coordinates of U_{α_i} and $U_{\alpha_{-i}}$ in $\overline{\Omega}$;
- (2) f_i extends the restriction of the conjugation by n_{α_i} on G to $\Omega_G \bigcap V_i \subset G$;
- (3) f_i sends U_{α} to $U_{s_i(\alpha)}$, for $\alpha \in \Psi$.

Proof. By [GP11b, Exposé XXIII, Lemme 3.1.1 (iii)] and the definition of Chevalley system [GP11b, Exposé XXIII, Définition 6.1], for each root $b \neq \pm \alpha_i \in \Psi$, we have

$$\operatorname{Ad}(n_{\alpha_i})p_b(x) = p_{s_i(b)}(e_b x),$$

where $e_b = \pm 1$ are given in *loc. cit.* and $x \in \Gamma(S', \mathcal{O}_{S'})$ for a test S-scheme S'. By [GP11b, Exposé XXIII 1.2], the above equality holds for $b = \pm \alpha_i$ with $e_b = -1$. For a section

$$u := \left(\prod_{c \in \Psi^- - \{-\alpha_i\}} p_c(x_c) \cdot p_{-\alpha_i}(x), t, p_{\alpha_i}(y) \cdot \prod_{c \in \Psi^+ - \{\alpha_i\}} p_c(x_c)\right) \in \overline{\Omega}(S'),$$

where $t = (t_1, ..., t_l) \in \overline{T}(S')$, we define

$$V_i := D_{\overline{\Omega}}(\mathbb{X}_i + xy).$$

If $u \in V_i(S')$, we define f_i by sending u to

$$\left(\prod_{c\in\Psi^{-}-\{-\alpha_{i}\}}p_{s_{i}(c)}(e_{c}x_{c})\cdot p_{-\alpha_{i}}\left(\frac{-y}{D}\right),\ \alpha_{i}^{\vee}(D)\cdot t,\ p_{\alpha_{i}}\left(\frac{-x}{D}\right)\cdot\prod_{c\in\Psi^{+}-\{\alpha_{i}\}}p_{s_{i}(c)}(e_{c}x_{c})\right)$$
(3)

where $D = X_i + xy$.

It is clear that V_i contains U^- and U^+ . The verification of the second condition follows from Equation (2) and a computation in the group G:

$$\begin{aligned} \operatorname{Ad}(n_{\alpha_{i}})(\prod_{c\in\Psi^{-}-\{-\alpha_{i}\}}p_{c}(x_{c})\cdot p_{-\alpha_{i}}(x)\cdot t\cdot p_{\alpha_{i}}(y)\cdot\prod_{c\in\Psi^{+}-\{\alpha_{i}\}}p_{c}(x_{c})) \\ &=\prod_{c\in\Psi^{-}-\{-\alpha_{i}\}}p_{s_{i}(c)}(e_{c}x_{c})\cdot p_{\alpha_{i}}(-x)\cdot (t\cdot\alpha_{i}^{\vee}(\alpha_{i}(t))^{-1})\cdot p_{-\alpha_{i}}(-y)\cdot\prod_{c\in\Psi^{+}-\{\alpha_{i}\}}p_{s_{i}(c)}(e_{c}x_{c}) \\ &=\prod_{c\in\Psi^{-}-\{-\alpha_{i}\}}p_{s_{i}(c)}(e_{c}x_{c})\cdot p_{-\alpha_{i}}\left(\frac{-y}{D}\right)\cdot (t\cdot\alpha_{i}^{\vee}(D))\cdot p_{\alpha_{i}}\left(\frac{-x}{D}\right)\cdot\prod_{c\in\Psi^{+}-\{a\}}p_{s_{i}(c)}(e_{c}x_{c}), \end{aligned}$$

for $t \in T(S')$, where $D = \alpha_i(t)^{-1} + xy$ and [GP11b, Exposé XXII, Définitions 1.5 (b)] is used in the first equality and [GP11b, Exposé XXII, Notations 1.3] is used in the second equality. \Box

Lemma 3.2. Set $n_0 = n_{\alpha_1}...n_{\alpha_l} \in \operatorname{Norm}_G(T)(S)$ and $n_{\alpha_i} = p_{\alpha_i}(1)p_{-\alpha_i}(-1)p_{\alpha_i}(1)$. There exist two open dense subschemes V and V' of $\overline{\Omega}$ and two morphisms

$$f: V \longrightarrow \overline{\Omega} \quad and \quad f': V' \longrightarrow \overline{\Omega}$$

such that

- (1) V contains $\mathcal{N}^- \times \mathcal{T} \times \mathcal{N}^+$ for nonempty open subschemes $\mathcal{N}^- \subset U^-$ and $\mathcal{N}^+ \subset U^+$ and $\mathcal{T} \subset \overline{T}$ such that \mathcal{N}^- and \mathcal{N}^+ are stable under the conjugation by T and $(0)_{1 \leq i \leq l} \in \mathcal{T}$; similarly, V' contains $\mathcal{M}^- \times \mathcal{T}' \times \mathcal{M}^+$ for nonempty open subschemes $\mathcal{M}^- \subset U^-$ and $\mathcal{M}^+ \subset U^+$ and $\mathcal{T}' \subset \overline{T}$ such that \mathcal{M}^- and \mathcal{M}^+ are stable under the conjugation by Tand $(0)_{1 \leq i \leq l} \in \mathcal{T}'$;
- (2) f (resp., f') extends the conjugation by n_0 (resp., n_0^{-1}) on $\Omega_G \cap V$ and $\Omega_G \cap V'$;

(3) the open V (resp., V') contains U⁻, U⁺, and the morphisms f and f' takes U⁻ (resp., U⁺) to U⁺ (resp., U⁻).

Proof. We define \tilde{V}_i by decreasing induction on i with $\tilde{V}_l = V_l$ and $\tilde{V}_i = f_i^{-1}(\tilde{V}_{i+1})$ for $1 \leq i \leq l-1$ where V_i and f_i come from Lemma 3.1. Then, it suffices to take $V = \tilde{V}_1$ and $f = f_l \dots f_1|_V$. Note that $n_0^{-1} = n_{\alpha_l}^3 n_{\alpha_{l-1}}^3 \dots n_{\alpha_1}^3$ because of the equalities $n_{\alpha_i}^4 = e$ in [GP11b, Exposé XX, Théorème 3.1 (v)]. Hence, similarly we get V' and f'. Note that, by Lemma 3.1, the section $(u^-, (0)_{1 \leq i \leq l}, u^+) \in \overline{\Omega}$ where all coordinates of u^- and u^+ are nonzero lies in V and V' because by Equation (3) the images of $(u^-, (0)_{1 \leq i \leq l}, u^+)$ under f_i still have nonzero coordinates on each root subgroups. Hence (1) follows. The claim (3) follows from (3) of *loc. cit.*

The main purpose of Lemma 3.2 is to use the adjoint action of n_0 (resp., n_0^{-1}) to swap the positive and the negative root subgroups, which is an important operation in the following result, even if it is merely defined over an open subscheme.

Lemma 3.3. Consider the S-rational morphism

$$\Theta: U^+ \times \overline{T} \times U^- \dashrightarrow \overline{\Omega}$$

given by the group multiplication of G. Then $\text{Dom}(\Theta)$ contains $\{e\} \times \overline{T} \times \{e\}$.

Proof. Note that by the definition of Θ , $\text{Dom}(\Theta)$ already contains $\{e\} \times T \times \{e\}$. Hence it suffices to show that $\{e\} \times (0)_{1 \leq i \leq l} \times \{e\}$ lies in $\text{Dom}(\Theta)$. For this, we will construct an *S*-rational morphism $\Theta' : U^+ \times \overline{T} \times U^- \dashrightarrow \overline{\Omega}$ whose definition domain contains this special point such that $\Theta = \Theta'$.

We fix a section $(u_0^-, (0)_{1 \leq i \leq l}, v_0^+) \in \overline{\Omega}(S)$ where $u_0^- \in \mathcal{N}^-(S)$ and $v_0^+ \in \mathcal{N}^+(S)$ and $\mathcal{N}^$ and \mathcal{N}^+ are defined in Lemma 3.2 (1). For any test S-scheme S' and a section $(u^+, \overline{t}, v^-) \in (U^+ \times \overline{T} \times U^-)(S')$, we define $\Theta'(u^+, \overline{t}, v^-)$ step by step and along the way we shrink the definition domain of Θ' .

As the first step, by shrinking the definition domain of Θ' , we can have

$$u^{+}(u_{0}^{-})^{-1} = (\dot{u}^{-}, \dot{t}, \dot{u}^{+}) \in \Omega_{G}(S'),$$

$$(v_{0}^{+})^{-1}v^{-} = (\dot{v}^{-}, \dot{s}, \dot{v}^{+}) \in \Omega_{G}(S').$$

As the second step, we further shrink $Dom(\Theta')$ so that the section

$$f'(f(\dot{t}\dot{u}^+\dot{t}^{-1}), f(\dot{t}^{-1}u_0^-\dot{t}, \dot{t}^{-1}\overline{t}\dot{s}, \dot{s}^{-1}u_0^+\dot{s}), f(\dot{s}^{-1}\dot{v}^-\dot{s})) \in \overline{\Omega}(S')$$

is well-defined, and we denote it by $(\tilde{u}^-, \tilde{t}, \tilde{v}^+)$.

As the *third step*, we define

$$\Theta'(u^+, \overline{t}, v^-) = (\dot{u}^- \widetilde{u}^-, \widetilde{t}, \widetilde{v}^+ \dot{v}^+) \in \overline{\Omega}(S').$$

The fact that $\Theta'|_{U^+ \times T \times U^-}$ coincides with the group multiplication of G follows from the claim (2) of Lemma 3.2. Hence we have $\Theta' = \Theta$ as S-rational morphisms. By the choice of u_0^- and v_0^+ and Lemma 3.2 (1), we have $\{e\} \times (0)_{1 \le i \le l} \times \{e\} \in \text{Dom}(\Theta')$.

Now we are in the position to define an rational action of $G \times G$ on the big cell $\overline{\Omega}$. The following theorem is, in spirit, similar to [GP11b, Exposé XXV Proposition 2.9], and we base our proof on theirs.

Theorem 3.4. There exist an S-rational morphism $\pi: G \times \overline{\Omega} \times G \dashrightarrow \overline{\Omega}$ whose definition domain \mathcal{R} contains $\{e\} \times \overline{\Omega} \times \{e\}$ such that

(1) The restriction of π to the open subscheme $(G \times \Omega_G \times G) \cap \mathcal{R}$ is given by

$$(g_1, g_2, g_3) \xrightarrow{\circ} g_1 g_2 g_3^{-1}.$$

(2) For every $x \in S$, the geometric fiber $\pi_{\overline{k(x)}}$ agrees with the restriction of the action of $G_{\overline{k(x)}} \times G_{\overline{k(x)}}$ on the big cell $\overline{\Omega_{\overline{k(x)}}}$ of the wonderful compactification of $G_{\overline{k(x)}}$ over the algebraically closed field $\overline{k(x)}$.

(3) The restriction $\pi|_{\{e\}\times\overline{\Omega}\times\{e\}}$ is the projection onto the middle factor.

Proof. Note that (2) follows from (1) by the fiberwise density of Ω_G in $\overline{\Omega}$.

Since Ω_G is S-fiberwise dense in G, it suffices to define an S-rational morphism

$$\pi\colon\Omega_G\times\overline{\Omega}\times\Omega_G\dashrightarrow\overline{\Omega}$$

whose definition domain \mathcal{R} contains $\{e\} \times \overline{\Omega} \times \{e\}$. We will define our π step by step and shrink $\Omega_G \times \overline{\Omega} \times \Omega_G$ along the way to obtain the sought \mathcal{R} . Let S' be a test S-scheme. Consider an element

$$A := ((u_1^-, t_1, u_1^+), (u_0^-, t_0, u_0^+), (u_2^-, t_2, u_2^+)) \in (\Omega_G \times \overline{\Omega} \times \Omega_G)(S').$$

The principle behind our construction is that we need to "move" all components of A on U^- (resp., U^+ , \overline{T}) together to form that of $\pi(A)$.

As the first step, let V be $\sigma^{-1}(\Omega_G) \cap \Omega_G$ where σ is the inverse morphism of the group scheme G, and we define an S-fiberwise dense open subscheme \mathcal{R}_1 of $\Omega_G \times \overline{\Omega} \times \Omega_G$ as $\Omega_G \times \overline{\Omega} \times V$. If $A \in \mathcal{R}_1(S')$, we write $\sigma(u_2^-, t_2, u_2^+) = (\hat{u}_2^-, \hat{t}_2, \hat{u}_2^+) \in \Omega_G(S')$.

As the second step, we want to "exchange" the position of the negative part and the positive part of the pair (u_1^+, u_0^-) and (u_0^+, \hat{u}_2^-) . Although the reductive group G is not commutative, we can still achieve this by restricting to the fiberwise dense open subscheme $\mathcal{R}_2 \subset \mathcal{R}_1$ defined by the following condition:

$$A \in \mathcal{R}_2 \iff u_1^+ u_0^- \text{ and } u_0^+ \hat{u}_2^- \text{ lie in } \Omega_G(S').$$

If so, we write

$$u_1^+ u_0^- = (\dot{u}_1^-, \dot{t}_1, \dot{u}_0^+) \in \Omega_G(S')$$
(4)

and

$$u_0^+ \hat{u}_2^- = (\dot{u}_0^-, \dot{t}_2, \dot{u}_2^+) \in \Omega_G(S').$$
(5)

As the *third step*, we seek to "exchange" the position of the positive part and the negative part of the triple $(\dot{u}_0^+, t_0, \dot{u}_0^-)$. To simplify the notations, we do this for the triple $(\dot{t}_1\dot{u}_0^+\dot{t}_1^{-1}, \dot{t}_1t_0\dot{t}_2, \dot{t}_2^{-1}\dot{u}_0^-\dot{t}_2)$. To do this, we further restrict to the open subscheme $\mathcal{R} \subset \mathcal{R}_2$ defined by the following:

$$A \in \mathcal{R} \iff (\dot{t}_1 \dot{u}_0^+ \dot{t}_1^{-1}, \dot{t}_1 t_0 \dot{t}_2, \dot{t}_2^{-1} \dot{u}_0^- \dot{t}_2)) \in \mathrm{Dom}(\Theta)$$

where Θ is defined in Lemma 3.3. The open subscheme \mathcal{R} is S-fiberwise dense because of *loc. cit.* If $A \in \mathcal{R}(S')$, we write

$$\Theta(\dot{t}_1 \dot{u}_0^+ \dot{t}_1^{-1}, \dot{t}_1 t_0 \dot{t}_2, \dot{t}_2^{-1} \dot{u}_0^- \dot{t}_2) = (\ddot{u}^-, \ddot{t}, \ddot{u}^+) \in \overline{\Omega}(S').$$
(6)

Now we have all components of $\pi(A)$ on U^- , U^+ and \overline{T} in the "right" places. Hence we define the image of A under π to be

$$(u_1^{-}(t_1\dot{u}_1^{-}t_1^{-1})(t_1\ddot{u}^{-}t_1^{-1}), t_1\ddot{t}t_2, (t_2^{-1}\ddot{u}^{+}t_2)(t_2^{-1}\dot{u}_2^{+}t_2)\hat{u}_2^{+}) \in \overline{\Omega}(S').$$

$$(7)$$

Note that \mathcal{R} contains $\{e\} \times \Omega \times \{e\}$ because of *loc. cit.*

To show the claim (1), according to the definition of π , if $A \in (\Omega_G \times \Omega_G \times \Omega_G)(S') \bigcap \mathcal{R}(S')$, we have that

$$\pi(A) = u_1^- t_1 \dot{u}_1^- \ddot{u} \ddot{t} \ddot{u}^+ \dot{u}_2^+ t_2 \dot{u}_2^+ \tag{8}$$

holds in G(S'). Notice that Equation (4) – (6) give rise to

$$u_1^+ u_0^1 = \dot{u}_1^- \dot{t}_1 \dot{u}_0^+, \ u_0^+ u_2^- = \dot{u}_0^- \dot{t}_2 \dot{u}_2^+$$

$$\dot{t}_1 \dot{u}_0^+ \dot{t}_1^{-1} \dot{t}_1 t_0 \dot{t}_2 \dot{t}_2^{-1} \dot{u}_0^- \dot{t}_2 = \ddot{u}^- \ddot{t} \ddot{u}^+$$

in G(S'), where Lemma 3.3 is used to deduce the last equation. Also since σ is the inverse operation, combining the first step, we have

$$\hat{u}_2^- \hat{t}_2 \hat{u}_2^+ = (u_2^- t_2 u_2^+)^{-1}.$$

Substituting the above four formulas into Equation (8) in turn, after a computation, we have

$$\pi(A) = (u_1^- t_1 u_1^+) (u_0^- t_0 u_0^+) (u_2^- t_2 u_2^+)^{-1},$$

as desired. The claim (3) follows from the definition of π .

3.3. Properties of the rational action. We are going to show several properties of π . These properties are analogue of the conditions which is imposed on a birational group law to ensure the existence of a group scheme solution to the group law.

The following lemma gives a feature of π defined in Theorem 3.4: if a section of $\Omega_G \times \overline{\Omega} \times \Omega_G$ equals to another section, which is "of group nature", under the morphism π , then the section itself is "of group nature" as well.

Lemma 3.5. For an S-scheme S' and two sections (g_1, x, g_2) , $(b_1, e, b_2) \in (\Omega_G \times \overline{\Omega} \times \Omega_G)(S')$, if $\pi(g_1, x, g_2) = \pi(b_1, e, b_2)$, then $x \in \Omega_G(S')$.

Proof. We shall keep the notations in the proof of Theorem 3.4, and spell out (g_1, x, g_2) as

$$((u_1^-, t_1, u_1^+), (u_0^-, t_0, u_0^+), (u_2^-, t_2, u_2^+)).$$

Recall that $\overline{\Omega}$ is defined as the product $U^- \times \prod_{\Delta} \mathbb{G}_a \times U^+$ and Ω_G is embedded into $\overline{\Omega}$ via Equation (2). Hence, to show $x \in \Omega_G(S')$, it suffices to check that each component of t_0 is nonzero. Note that $\pi(b_1, e, b_2) \in \Omega_G(S')$, and by the definition of π i.e., Equation (7) and the equality $\pi(g_1, x, g_2) = \pi(b_1, e, b_2)$, each component of \ddot{t} is nonzero. By the definition of f (resp., f') in the proof of Lemma 3.2, f (resp., f') is the composition of f_i (i = 1, ..., l)(resp., f'_i) defined in Lemma 3.1, and by Equation (3), the only impact of f_i (resp., f'_i) on the coordinates of $\overline{T}(S')$ is multiplying nonzero scalars given by elements in T(S'). Therefore, by Equation (6), $\dot{t}_1 t_0 \dot{t}_2$ has nonzero components, so is t_0 , as desired.

The following lemma says that the rational action π of $G \times G$ on $\overline{\Omega}$ defined in Theorem 3.4 is associative.

Lemma 3.6. The following two S-rational morphisms

$$\phi \colon (G \times G) \times (G \times \overline{\Omega} \times G) \dashrightarrow \overline{\Omega},$$
$$((g_1, g_2), (g'_1, x, g'_2)) \mapsto \pi(g_1, \pi(g'_1, x, g'_2), g_2)$$

and

$$\psi \colon (G \times G) \times (G \times \Omega \times G) \dashrightarrow \Omega, ((g_1, g_2), (g'_1, x, g'_2)) \mapsto \pi(g_1g'_1, x, g_2g'_2)$$

coincide.

Proof. By (1) of Theorem 3.4, ϕ and ψ coincide over an S-fiberwise dense open subscheme.

The morphism π in Theorem 3.4 defines a (relative) rational morphism in the sense of [SGA3II, Exposé XVIII, Définition 1.5], which can be viewed as a rational action of $G \times G$ on $\overline{\Omega}$. In the following theorem, we prove that this rational action is "birational" and "strict", which is an analogue of the notion of a strict birational group law, see [BLR90, Chapter 5, Section 5.2, Definition 1]. We adopt the strategy of the proof of [Con14, Section 6, Proposition 6.3.13] into our context.

Lemma 3.7. The S-rational morphism

$$\Phi \colon G \times \overline{\Omega} \times G \dashrightarrow G \times \overline{\Omega} \times G,$$

$$(g_1, x, g_2) \mapsto (g_1, \pi(g_1, x, g_2), g_2),$$

is birational. Moreover, there exists an $\overline{\Omega}$ -dense open subscheme \mathcal{V} of the definition domain of Φ containing $\{e\} \times \overline{\Omega} \times \{e\}$ such that $\Phi(\mathcal{V})$ is $\overline{\Omega}$ -dense and the restriction $\Phi|_{\mathcal{V}}$ is an open immersion into $G \times \overline{\Omega} \times G$.

Proof. The rational morphism Φ is birational because, by Lemma 3.6, it has the following rational inverse:

$$\Psi \colon G \times \Omega \times G \dashrightarrow G \times \Omega \times G,$$

$$(g_1, x, g_2) \mapsto (g_1, \pi(g_1^{-1}, x, g_2^{-1}), g_2)$$

where Ψ is defined over an S-dense open subscheme of $G \times \overline{\Omega} \times G$ obtained by further shrinking the definition domain of Φ . Let \mathcal{U} and \mathcal{U}' be the definition domain of Φ and Ψ . Let

$$\mathcal{V} := \Phi^{-1}(\mathcal{U}') \bigcap \mathcal{U} \text{ and } \mathcal{V}' := \Psi^{-1}(\mathcal{U}) \bigcap \mathcal{U}'.$$

Note that $\Phi|_{\mathcal{V}}$ and $\Psi|_{\mathcal{V}'}$ are inverse to each other and by Theorem 3.4, \mathcal{V} and \mathcal{V}' contain $\{e\} \times \overline{\Omega} \times \{e\}$.

4. Compactification for split adjoint reductive group schemes

We shall keep the notations of Section 3. In this section, we first define a compactification of the split adjoint reductive group G as a sheaf. Then we prove that this compactification is in fact a $G \times G$ -projective scheme and contains G as an open dense subscheme. Moreover, we show that geometric fibers of this compactification are wonderful compactifications as we recall in Section 2.

4.1. An equivalence relation. Recall from Theorem 3.4 that we have the S-rational morphism $\pi: G \times \overline{\Omega} \times G \dashrightarrow \overline{\Omega}$. The following construction is inspired by [SGA3II, Exposé XVIII, 3.2.3].

Definition 4.1. We define a relation on $G \times \overline{\Omega} \times G$ as follows: for a test S-scheme S' and $(g_1, x, g_2), (g'_1, x', g'_2) \in (G \times \overline{\Omega} \times G)(S')$, we say that (g_1, x, g_2) and (g'_1, x', g'_2) are equivalent, if and only if, there exist an fppf cover $S'' \to S'$ and a section $(a_1, a_2) \in (G \times G)(S'')$ such that $\pi(a_1g_1, x, a_2g_2)$ and $\pi(a_1g'_1, x', a_2g'_2)$ are both well defined and are equal over S''. We write $(g_1, x, g_2) \sim (g'_1, x', g'_2)$ if $(g_1, x, g_2), (g'_1, x', g'_2) \in (G \times \overline{\Omega} \times G)(S')$ are equivalent.

Our next goal is to show that the relation in Definition 4.1 is an equivalence relation. As a preparation, we need the following lemma which is inspired by [SGA3II, Exposé XVIII, Lemme 3.3].

Lemma 4.2. Consider two sections

$$(g_1, x, g_2), (g'_1, x', g'_2) \in (G \times \overline{\Omega} \times G)(S)$$

together with $(a_1, a_2) \in (G \times G)(S_1)$, where S_1 is an fppf cover of S, such that

$$\pi(a_1g_1, x, a_2g_2) = \pi(a_1g_1', x', a_2g_2').$$

Then, for $(h_1, h_2) \in (G \times G)(S_2)$ where S_2 is an S-scheme, if $\pi(h_1g_1, x, h_2g_2)$ and $\pi(h_1g'_1, x', h_2g'_2)$ are well defined, then they are equal.

Proof. We have that

$$\pi(z_1g_1, x, z_2g_2) = \pi(z_1a_1^{-1}, \pi(a_1g_1, x, a_2g_2), z_2a_2^{-1})$$

= $\pi(z_1a_1^{-1}, \pi(a_1g'_1, x', a_2g'_2), z_2a_2^{-1})$
= $\pi(z_1g'_1, x', z_2g'_2)$

holds as an equality of S_1 -rational morphisms with respect to the variable (z_1, z_2) , where Lemma 3.6 is used in the first and the last equality. Hence by [SGA3II, Exposé XVIII, Proposition 1.6], $\pi(z_1g_1, x, z_2g_2) = \pi(z_1g'_1, x', z_2g'_2)$ holds as an equality of S-rational morphisms. Therefore $\pi(h_1g_1, x, h_2g_2) = \pi(h_1g'_1, x', h_2g'_2)$.

Lemma 4.3. The relation in Definition 4.1 is an equivalence relation.

Proof. It is clear that this relation is symmetric.

To show the reflexivity, for an S-scheme S' and $(g_1, x, g_2) \in (G \times \overline{\Omega} \times G)(S')$, we need to find a section $(a_1, a_2) \in (G \times G)(S'')$ for some S'' is an fppf cover of S' such that $\pi(a_1g_1, x, a_2g_2)$ is well defined. By Lemma 3.7, the definition domain \mathcal{U} of Φ is an $\overline{\Omega}$ -dense open subscheme. In the view of the stability of being relative schematic dense under base change (see [SGA3II, Exposé XVIII, Proposition 1.1 (2)]), the base change $\mathcal{U}' := \mathcal{U} \times_{\overline{\Omega}} S'$ of \mathcal{U} along x is an S'-dense open subscheme of $(G \times G) \times S'$. Then the image of the sought section (a_1, a_2) is required to lie in $((g_1^{-1}, g_2^{-1}) \cdot \mathcal{U}')$, which is an open dense subscheme of $G_{S'} \times G_{S'}$. Thus, the existence of such a section (a_1, a_2) over some fppf covering of S' is ensured by [SGA3II, Exposé XVIII, Proposition 1.7]. This proves the reflexivity.

To show the transitivity, we assume that, over S', we have

$$(g_1, x, g_2) \sim (g'_1, x', g'_2), \ (g'_1, x', g'_2) \sim (g''_1, x'', g''_2).$$

By Lemma 3.7 and [SGA3II, Exposé XVIII, Proposition 1.1 (i)], there exists an S-dense open subscheme \mathcal{V} of $G \times G$ such that for any section (h_1, h_2) of \mathcal{V} valued in some S-scheme, $\pi(h_1g_1, x, h_2g_2), \pi(h_1g'_1, x', h_2g'_2)$ and $\pi(h_1g''_1, x'', h_2g''_2)$ are all well defined. Such a section (h_1, h_2) exists by [SGA3II, Exposé XVIII, Proposition 1.7]. Then the transitivity follows from Lemma 4.2.

Lemma 4.4. Consider the following S-rational morphism

$$\phi \colon (G \times \overline{\Omega} \times G) \times (G \times G) \dashrightarrow \overline{\Omega},$$
$$(g_1, x, g_2, g'_1, g'_2) \longmapsto \pi(g'_1^{-1}, \pi(g_1, x, g_2), g'^{-1}).$$

Let \mathcal{U} be the definition domain of the rational morphism ϕ , and let Γ be the graph of the morphism $\phi|_{\mathcal{U}}$. We consider a section

$$(g_1, x, g_2, g'_1, g'_2, y) \in ((G \times \overline{\Omega} \times G) \times (G \times G) \times \overline{\Omega})(S).$$

If $(g_1, x, g_2, g'_1, g'_2, y) \in \Gamma(S)$, then $(g_1, x, g_2) \sim (g'_1, y, g'_2)$.

Proof. Assume that $(g_1, x, g_2, g'_1, g'_2, y) \in \Gamma(S)$ i.e., $y = \phi(g_1, x, g_2, g'_1, g'_2)$. Note that $\pi(g_1, x, g_2)$ may not be well defined because $\operatorname{pr}_{123}(\mathcal{U})$ is possibly larger than the definition domain of π . However, by Lemma 3.7 and [SGA3II, Exposé XVIII, Proposition 1.7], there exists an $(h_1, h_2) \in (G \times G)(S'')$, where S'' is an fppf cover of S, such that $\pi(h_1g_1, x, h_2g_2)$ and $\pi(h_1g'_1, y, h_2g'_2)$ are both well defined. Again, by the associativity of π (Lemma 3.6) and the definition of ϕ and by viewing $(g_1, x, g_2, g'_1, g'_2)$ as variables, we have the following equalities of S"-rational morphisms

$$\pi(h_1g'_1, y, h_2g'_2) = \pi(h_1g'_1, \phi(g_1, x, g_2, g'_1, g'_2), h_2g'_2)$$

= $\pi(h_1, \pi(g'_1, \phi(g_1, x, g_2, g'_1, g'_2), g'_2), h_2)$
= $\pi(h_1, \pi(g_1, x, g_2), h_2) = \pi(h_1g_1, x, h_2g_2).$

It follows that $\pi(h_1g'_1, y, h_2g'_2) = \pi(h_1g_1, x, h_2g_2).$

Lemma 4.5. The scheme Γ in Lemma 4.4 is flat and locally of finite presentation over $G \times \overline{\Omega} \times G$ with respect to pr_{123} and pr_{456} , where we implicitly identify the target of the projection pr_{456} with $G \times \overline{\Omega} \times G$ by switching the last two coordinates.

Proof. By definition, Γ fits into the following cartesian diagram



Therefore, Γ is of finite presentation over $\mathcal{U} \times \overline{\Omega}$ via $\operatorname{pr}_{12345} \times \operatorname{pr}_6$ because so is Δ . It follows that the morphism $\operatorname{pr}_{123}|_{\Gamma}$ is locally of finite presentation as well because so are G and $\overline{\Omega}$ over S. Observe also that Γ is isomorphic to $\operatorname{pr}_{12345}(\Gamma) = \mathcal{U}$ which is, by definition, an open subscheme of $(G \times \overline{\Omega} \times G) \times (G \times G)$. Combining the flatness of G and $\overline{\Omega}$ over S, we deduce that the morphism $\operatorname{pr}_{123}|_{\mathcal{U}}$ is flat.

To show that pr_{456} is flat and locally of finite presentation, we claim that Γ is symmetric under the permutation of components sending $(g_1, x, g_2, g'_1, g'_2, y)$ to $(g'_1, y, g'_2, g_1, g_2, x)$. Then the claim and a similar argument give rise to the fact that $\operatorname{pr}_{456}|_{\Gamma}$ is flat and locally of finite presentation.

To show the claim, it suffices to check that

$$\phi(g_1',\phi(g_1,x,g_2,g_1',g_2'),g_2',g_1,g_2)=x.$$

This follows from the associativity of π (Lemma 3.6) and a computation of rational morphisms.

4.2. Schematic nature of the compactification. Now we are in the position to construct a compactification of the group G as a sheaf. The following definition is inspired by the definition given in [SGA3II, Exposé XVIII, Page 387].

Definition 4.6. Thanks to Lemma 4.3, we define the compactification of the adjoint group scheme G to be the quotient sheaf \mathcal{X} of the scheme $G \times \overline{\Omega} \times G$ with respect to the equivalence relation constructed in Lemma 4.3. We denote the quotient morphism by $Q_G: G \times \overline{\Omega} \times G \to \mathcal{X}$.

We adapt the method due to Artin in the proof of [SGA3II, Exposé XVIII, Proposition 3.5] to deduce the following result which says that the equivalence relation of Lemma 4.3 can be encoded in a nice scheme.

Theorem 4.7. The quotient sheaf of the relation $\Gamma \rightrightarrows G \times \overline{\Omega} \times G$ is isomorphic to \mathcal{X} , where Γ is defined in Lemma 4.4, the two morphisms are pr_{123} and pr_{456} and we implicitly identify the target of the projection pr_{456} with $G \times \overline{\Omega} \times G$ by switching the last two coordinates.

Proof. We will verify that two sections of $G \times \overline{\Omega} \times G$ are equivalent if and only if they come from a section of Γ . As in Lemma 4.4, let \mathcal{U} be the definition domain of

$$\phi \colon (G \times \overline{\Omega} \times G) \times (G \times G) \dashrightarrow \overline{\Omega}$$

and Γ is the graph of $\phi|_{\mathcal{U}}$. Then we have the natural immersion

$$\Gamma \hookrightarrow (G \times \overline{\Omega} \times G) \times (G \times G) \times \overline{\Omega}$$

We fix a test S-scheme S' and two sections $(g_1, x, g_2), (g'_1, y, g'_2) \in (G \times \overline{\Omega} \times G)(S')$. Since the definition domain of $\phi_{S'}$ certainly contains $\mathcal{U} \times S'$, by applying Lemma 4.4 to $\phi_{S'}$, it follows that $(g_1, x, g_2, g'_1, g'_2, y) \in \Gamma(S')$ implies that $(g_1, x, g_2) \sim (g'_1, y, g'_2)$.

Conversely, suppose that, for a test S-scheme S',

 $(g_1, x, g_2), (g'_1, y, g'_2) \in (G \times \overline{\Omega} \times G)(S') \text{ and } (g_1, x, g_2) \sim (g'_1, y, g'_2),$

i.e., that there exist an fppf cover S'' over S' and a section $(a_1, a_2) \in (G \times G)(S'')$ such that $\pi(a_1g_1, x, a_2g_2) = \pi(a_1g'_1, y, a_2g'_2)$. We have to show that $(g_1, x, g_2, g'_1, g'_2, y) \in \Gamma(S')$. By [GP11a, Exposé IV, Proposition 6.3.1 (iii)], the scheme Γ represents a sheaf. Hence to check that $(g_1, x, g_2, g'_1, g'_2, y) \in \Gamma(S')$, it suffices to localize on S'. Note that $(g_1, x, g_2) \sim (g'_1, y, g'_2)$ remains true while passing to étale neighborhoods. We assume that S' is the spectrum of a strictly henselian local ring. Then consider the strict localization \widetilde{S} of S at the image of the geometric closed point of S'. By [Gro67, Proposition 18.8.8 (iii), (iv)], the morphism $\widetilde{S} \longrightarrow S$ is flat. Therefore, by [BLR90, 2.5, Proposition 6], the formation of Γ commutes with the base change to \widetilde{S} , hence, we may assume that S is also strictly henselian and the morphism $S' \to S$ is local.

We claim that there exist a flat and finitely presented morphism $S_0 \longrightarrow S$ and a section $(h_1, h_2) \in (G \times G)(S_0)$ such that $\pi(h_1g_1, x, h_2g_2)$ and $\pi(h_1g'_1, y, h_2g'_2)$ are well defined, here we implicitly pullback (h_1, h_2) to $(G \times G)(S' \times_S S_0)$.

Granted the claim, consider the S_0 -rational morphism

$$\varphi \colon (G \times \overline{\Omega} \times G) \times (G \times G) \dashrightarrow \overline{\Omega},$$
$$(a_1, \omega, a_2, a'_1, a'_2) \longmapsto \pi((h_1 a'_1)^{-1}, \pi(h_1 a_1, \omega, h_2 a_2), (h_2 a'_2)^{-1}).$$

By Lemma 3.6, two S_0 -rational morphisms φ and ϕ_{S_0} equal, where ϕ is defined in Lemma 4.4. By Lemma 4.2, the equivalence $(g_1, x, g_2) \sim (g'_1, y, g'_2)$ implies that

$$\pi(h_1g_1, x, h_2g_2) = \pi(h_1g_1', y, h_2g_2').$$

This means that $\pi((h_1g'_1)^{-1}, \pi(h_1g_1, x, h_2g_2), (h_2g'_2)^{-1})$ is well defined and equals to y. Therefore, since \mathcal{U} is the definition domain of ϕ , by [SGA3II, Exposé XVIII, Proposition 1.6], it follows that $(g_1, x, g_2, g'_1, g'_2) \in \mathcal{U}(S')$ and also $(g_1, x, g_2, g'_1, g'_2, y)$ lies in $\Gamma(S')$.

To show the claim above, by Lemma 3.7, we are left with showing that there is a flat and finitely presented morphism $S_0 \longrightarrow S$ and a S'-morphism $S_0 \times S' \longrightarrow (G \times G) \times S'$ landing in the S'-dense open subcheme

$$\mathcal{W}' \coloneqq ((g_1^{-1}, g_2^{-1}) \cdot \mathcal{V}_x) \bigcap ((g_1'^{-1}, g_2'^{-1}) \cdot \mathcal{V}_y) \bigcap ((G \times G) \times S') \subset (G \times G) \times S',$$

where $\mathcal{V}_x \subset (G \times G) \times S'$ (resp., $\mathcal{V}_y \subset (G \times G) \times S'$) is the base change of the open subscheme $\mathcal{V} \subset (G \times \overline{\Omega} \times G) \times S'$, which is obtained by applying Lemma 3.7 over S', along $x \in \overline{\Omega}(S')$ (resp., $y \in \overline{\Omega}(S')$). To do this, we adopt the strategy of the proof of [Gro67, Corollaire 17.16.2].

Let s' (resp., s) be the closed point of S' (resp., S). Since \mathcal{W}' is S'-dense, we can assume that the special fiber $\mathcal{W}'_{s'}$ contains an open subscheme of the form $\mathcal{W} \times_{k(s)} k(s')$ where \mathcal{W} is a nonempty open subscheme of $(G \times G)_s$. We choose a closed point $h \in \mathcal{W} \subset G \times G$. We can choose a system of parameters $(\tilde{t}_i)_{1 \leq i \leq n}$ in the local ring $\mathcal{O}_{(G \times G)_s,h}$, because the special fiber $(G \times G)_s$ is a regular scheme. Then we can find an affine open neighborhood V containing h and n sections $(t_i)_{1 \leq i \leq n} \subset \Gamma(V, \mathcal{O}_{G \times G})$ lifting $(\tilde{t}_i)_{1 \leq i \leq n}$. Let $V' \subset V$ be the closed subscheme cut out by $(t_i)_{1 \leq i \leq n}$, and, by [Gro65b, Théorème 11.3.8 b') and c)], after shrinking V' if needed, we can assume that V' is flat over S. Since $(\tilde{t}_i)_{1 \leq i \leq n}$ is a system of parameters, $\mathcal{O}_{V'_s,h}$ is artinian. Since h is a closed point in V'_s , it is isolated in V'_s . Hence $\operatorname{Spec}(\mathcal{O}_{V',h})$ is quasi-finite over S. As S is henselien, by the Zariski main theorem, the S-scheme $\operatorname{Spec}(\mathcal{O}_{V',h})$ is even finite, see [Gro67, Théorème 18.5.11 c')]. We take $\operatorname{Spec}(\mathcal{O}_{V',h})$ as S_0 and consider the natural S-morphism ϵ : $\operatorname{Spec}(\mathcal{O}_{V',h}) \to G \times G$.

Now we show that the S'-morphism $\epsilon \times \operatorname{Id}_{S'}: S_0 \times S' \to (G \times G) \times S'$ has image in \mathcal{W}' . Since ϵ sends the closed point of $\operatorname{Spec}(\mathcal{O}_{V',h})$ to $h \in \mathcal{W}$ and, by [Gro60, Proposition 2.4.4], ϵ preserves generalizations of the closed point of $\operatorname{Spec}(\mathcal{O}_{V',h})$, the image of the closed fiber ϵ_s lies in \mathcal{W} . Hence, the special fiber $(\epsilon \times \operatorname{Id}_{S'})_{s'}$ has image in $\mathcal{W} \times_{k(s)} k(s') \subset \mathcal{W}'$. Now since S' is henselian, by [Gro67, Proposition 18.5.9 (ii)], the S'-finite scheme $\operatorname{Spec}(S_0 \times S')$ decomposes as the disjoint union of some local schemes. We already know that the closed points of these local schemes are mapped into \mathcal{W}' by $\epsilon \times \operatorname{Id}_{S'}$. Hence, by [Gro60, Proposition 2.4.4], the whole space $\operatorname{Spec}(S_0 \times S')$ also lands in \mathcal{W}' , as desired. \Box

Corollary 4.8. The sheaf \mathcal{X} is an S-algebraic space.

Proof. By Lemma 4.5, the corollary then follows from a result due to Artin, which says that the quotient of an algebraic space with respect to an fppf relation is again an algebraic space, see [LMB00, Corollaire (10.4)]. \Box

Our next goal is to show that \mathcal{X} is, in fact, a scheme. To do this, we first endow \mathcal{X} with a group action of $G \times G$.

Definition-Proposition 4.9. For any S-scheme S', consider $g = (a_1, a_2) \in (G \times G)(S')$ and $x \in \mathcal{X}(S')$ represented by a section $(g_1, y, g_2) \in (G \times \overline{\Omega} \times G)(S'')$, where S'' is an fppf cover of S'. We define gx to be the section of $\mathcal{X}(S')$ represented by the section

$$(a_1g_1, y, a_2g_2) \in (G \times \overline{\Omega} \times G)(S'').$$

We denote by

$$\Theta \colon G \times \mathcal{X} \times G \longrightarrow \mathcal{X}$$

the resulting action of $G \times G$ on \mathcal{X} .

Proof. We need to check that the action of $G \times G$ on \mathcal{X} is well defined. Fix a test S-scheme S'. Let $g = (a_1, a_2) \in (G \times G)(S')$. Let (g_1, y, g_2) and (g'_1, y', g'_2) be two sections of $G \times \overline{\Omega} \times G$ valued in two different fppf covers S''_1 and S''_2 of S'. By pullback two sections to $S''_1 \times_{S'} S''_2$, we can assume that $S''_1 = S''_2$ which we rename to S''. By Lemma 3.7 and [SGA3II, Exposé XVIII, Propostion 1.7], we can find a section (b_1, b_2) of $G \times G$ valued in some fppf cover of S'' such that $(b_1a_1g_1, y, b_2a_2g_2)$ and $(b_1a_1g'_1, y', b_2a_2g'_2)$ are both well defined. Then we conclude by Lemma 4.2.

The following lemma is inspired by the fact that the group scheme that is associated to a birational group law on a scheme X contains X as an open dense subscheme ([SGA3II, Exposé XVIII, Théorème 3.7 (ii)]). The proof that we present here is, in spirit, close to [ER15, Lemma 3.13].

Lemma 4.10. The morphism $\mathbf{j} \colon \overline{\Omega} \to \mathcal{X}$ which takes a section $a \in \overline{\Omega}(S')$ to the equivalence class represented by (e, a, e) in $\mathcal{X}(S')$, is represented by an open immersion.

Proof. We first verify that **j** is a monomorphism. For an S-scheme S', let $a_1, a_2 \in \overline{\Omega}(S')$ be such that $\mathbf{j}(a_1) = \mathbf{j}(a_2)$. Then there exist an fppf cover S'' of S' and a section $(g_1, g_2) \in (G \times G)(S'')$ such that $\pi(g_1, a_1, g_2) = \pi(g_1, a_2, g_2)$. Then by Lemma 4.2, we get $a_1 = a_2$ in $\overline{\Omega}(S'')$, hence $a_1 = a_2$ in $\overline{\Omega}(S')$.

Now to show that **j** is an open immersion, we need to prove that, for a test S-scheme S', and a morphism $\alpha: S' \to \mathcal{X}$, there exist an open subscheme $U' \subset S'$ and a section of $\overline{\Omega}(U')$ such that the following diagram



is Cartesian.

We first produce the open subscheme U' and a section of $\overline{\Omega}(U')$. By fppf descent, without loss of generality, we can assume that $\alpha \in \mathcal{X}(S')$ is represented by a section $(g_1, x, g_2) \in (G \times \overline{\Omega} \times G)(S')$. We consider the following S'-rational morphism

$$\begin{aligned} f\colon (G\times G)\times S'\dashrightarrow \Omega\times S'\\ (a_1,a_2)\longmapsto \phi(a_1g_1,x,a_2g_2,a_1,a_2), \end{aligned}$$

where (a_1, a_2) is an arbitrary section of $G \times G$ over an S'-scheme and ϕ is defined in Lemma 4.4. Let \mathcal{V}' be the definition domain of the S'-rational morphism

$$(G \times G) \times S' \dashrightarrow \overline{\Omega} \times S'$$
$$(a_1, a_2) \longmapsto \pi(a_1g_1, x, a_2g_2).$$

Consider the S'-morphism

$$h: \mathcal{V}' \longrightarrow (G \times \overline{\Omega} \times G) \times (G \times G) \times S'$$
$$(a_1, a_2) \longmapsto (a_1g_1, x, a_2g_2, a_1, a_2).$$

Let $\mathcal{U} = h^{-1}(\text{Dom}(\phi) \times S')$ where ϕ is defined in Lemma 4.4. As G is flat and finitely presented over S, by [Gro65a, Théorème 2.4.6], the image of \mathcal{U} under the projection from $(G \times G) \times S'$ to S' is an open subscheme of S', denoted by U'. We claim that there is a morphism $f_{U'}$ that fits into a commutative diagram



in which $i_{U'}$ is the natural open immersions and $\operatorname{pr}_{U'}$ is the projection from \mathcal{U} to the last coordinate U'. To see this, by the definition of f and \mathcal{U} , we have $\mathbf{j} \circ \operatorname{pr}_{\overline{\Omega}} \circ f|_{\mathcal{U}} = \alpha \circ i_{U'} \circ \operatorname{pr}_{U'}$. Then, since \mathbf{j} is a monomorphism and $\mathcal{U} \to \mathcal{U}'$ is an fppf cover, $\operatorname{pr}_{\overline{\Omega}} \circ f|_{\mathcal{U}}$ descends to give the sought morphism $f_{U'}$.

To finish the proof, it suffices to check that the bottom-right square of the above diagram is cartesian. For this, let T be an S-scheme, and let $h_{S'}: T \to S'$ and $h_{\overline{\Omega}}: T \to \overline{\Omega}$ be two morphisms over S such that $\mathbf{j} \circ h_{\overline{\Omega}} = \alpha \circ h_{S'}$. Then we are left to show that there exists a unique morphism $h_{U'}: T \to U'$ that fits into a commutative diagram



As $i_{U'}$ is an open immersion, the uniqueness of $h_{U'}$ follows from [Sta, 01L7]. It suffices to show the existence of $h_{U'}$. By viewing $h_{\overline{\Omega}}$ as a section of $\overline{\Omega}(T)$, the equality $\mathbf{j} \circ h_{\overline{\Omega}} = \alpha \circ h_{S'}$ gives rise to $(e, h_{\overline{\Omega}}, e) \sim (g_1, x, g_2)$. This implies that there exists an S'-morphism $(c_1, c_2) \in (G \times G)(T')$ such that

$$\pi(c_1, h_{\overline{\Omega}}, c_2) = \pi(c_1g_1, x, c_2g_2),$$

where T' is an fppf cover of T. By the definition of \mathcal{U} and f and Theorem 4.7, we have

 $(c_1, c_2) \in \mathcal{U}(T')$ and $f(c_1, c_2) = h_{\overline{\Omega}}$.

Since $T' \to T$ is an fppf cover and U' is the image of \mathcal{U} in S', the morphism $\operatorname{pr}_{U'} \circ (c_1, c_2)$ descends to give the desired morphism $h_{U'}$.

Proposition 4.11. The algebraic space \mathcal{X} is a separated scheme over S.

Proof. By Lemma 4.10, the $\overline{\Omega}$ is an open subspace of \mathcal{X} . By the definition of the action of $G \times G$ on \mathcal{X} (Definition-Proposition 4.9), we have $\mathcal{X} = (G \times G) \cdot \overline{\Omega}$. Then the proposition follows from [BLR90, Section 6.6, Theorem 2 (a) and (b)] and the smoothness of $\overline{\Omega}$ over S. \Box

4.3. Relation with wonderful compactifications. We are going to show that the geometric fibers of \mathcal{X} are classical wonderful compactifications (as we review in Section 2.2). For this, recall that the wonderful compactification \mathbf{X}_s of $G_{\overline{k(s)}}$ contains an open subscheme called the big cell which is canonically isomorphic to $(\overline{\Omega})_{\overline{k(s)}}$, see, for instance, [BK05, Proposition 6.1.7].

Proposition 4.12. There is a $G \times G$ -equivariant open immersion $\mathbf{i}: G \to \mathcal{X}$ such that every geometric fiber of \mathcal{X} over a point $s \in S$ is identified with the wonderful compactification of $G_{\overline{k(s)}}$. Viewing $\overline{\Omega}$ as an open subscheme of \mathcal{X} via Lemma 4.10, the restriction of \mathbf{i} to Ω_G agrees with Equation (2). For a test S-scheme S', the S'-valued points of the image of \mathbf{i} are

$$Q_G(\bigcap_{\alpha \in \Delta} D_{G \times \overline{\Omega} \times G}(\mathbb{X}_\alpha)(S'))$$

where Q_G is the quotient map as in Definition 4.6 and $D_{G \times \overline{\Omega} \times G}(\mathbb{X}_{\alpha})$ is the principal open subscheme of $G \times \overline{\Omega} \times G$ defined by \mathbb{X}_{α} .

Proof. For a section $g \in G(S')$ where S' is an S-scheme, we define $\mathbf{i}: G \to \mathcal{X}$ by taking $\mathbf{i}(g)$ to the equivalence class of (g, e, e) in $\mathcal{X}(S')$. The fact that \mathbf{i} is $G \times G$ -equivariant follows from Theorem 3.4.

We claim that $G \cap \overline{\Omega} = \Omega_G$ and $\mathbf{i}|_{\Omega_G} \colon \Omega_G \to \overline{\Omega}$ coincides with Equation (2), here the intersection indicates the pullback of $\overline{\Omega}$, which is an open subscheme of \mathcal{X} (Lemma 4.10), along \mathbf{i} .

Granted the claim, as Ω_G is an open subscheme of G [GP11b, Exposé XXII, Proposition 4.1.2] and \mathcal{X} is covered by the translations of $\overline{\Omega}$ under the action of $G \times G$, it follows that **i** is an open immersion.

To prove the claim, we choose a section $g \in \Omega_G(S')$. By Definition-Proposition 4.9, $\mathbf{i}(g)$ is the section of \mathcal{X} represented by the section $(g, e, e) \in (G \times \overline{\Omega} \times G)(S')$. Note that, by Definition 4.1, $(g, e, e) \sim (e, \nu(g), e)$, where ν is Equation (2). Hence $\Omega_G \subset G \cap \overline{\Omega}$ and the restriction $\mathbf{i}|_{\Omega_G}$ coincides with ν . For the converse direction, take a section $\beta \in G(S') \cap \overline{\Omega}(S')$, after base change if needed, we assume that there is an fppf cover S'' of S' so that β is represented by two sections (e, y, e), (b_1, e, e) of $(G \times \overline{\Omega} \times G)(S'')$ satisfying $(e, y, e) \sim (b_1, e, e)$. Hence, by Lemma 3.5, y lies in $\Omega_G(S'')$, then $\beta \in \Omega_G(S')$ follows. This proves the claim.

For a point $s \in S$, by the functoriality of the definition of \mathcal{X} (Definition 4.6), the geometric fiber $\mathcal{X}_{\overline{k(s)}}$ of \mathcal{X} is the quotient sheaf $G_{\overline{k(s)}} \times (\overline{\Omega})_{\overline{k(s)}} \times G_{\overline{k(s)}} / \sim_s$, where \sim_s stands for the equivalence relation obtained by applying Definition 4.1 to $G_{\overline{k(s)}}$. To define a morphism from $\mathcal{X}_{\overline{k(s)}}$ to \mathbf{X}_s , by the universal property of sheafication, it suffices to define a morphism χ_s^+ at the level of presheaves. For a $\overline{k(s)}$ -scheme T and a section $(g_1, x, g_2) \in (G_{\overline{k(s)}} \times (\overline{\Omega})_{\overline{k(s)}} \times G_{\overline{k(s)}})(T)$, we define the image of (g_1, x, g_2) under χ_s^+ to be $g_1 x g_2^{-1}$ in $\mathbf{X}_s(T)$. The fact that χ_s is well defined follows from the fibral description of π in Theorem 3.4. The associated morphism from $\mathcal{X}_{\overline{k(s)}}$ to \mathbf{X}_s is denoted by χ_s . Note that, by Definition-Proposition 4.9, this χ_s is $G_{\overline{k(s)}} \times G_{\overline{k(s)}}$ equivariant. By Lemma 4.10 and the definition of χ_s , $\chi_s|_{(\overline{\Omega})_{\overline{k(s)}}}$ is an open immersion onto the big cell of \mathbf{X}_s . By the translations of $G_{\overline{k(s)}} \times G_{\overline{k(s)}}$, the χ_s is quasi-finite, hence by the Zariski main theorem [Gro61b, Corollaire 4.4.9], is an open immersion. By Equation (1), χ_s is an isomorphism.

The last part of the proposition follows essentially from the definition of ν and the equivariant action of $G \times G$.

Remark 4.13. In the proof of Proposition 4.12, we construct i via the left action of G on \mathcal{X} . In the view of the condition (1) of Theorem 3.4, it is equivalent to use the right action of G.

4.4. **Projectivity of compactification.** In Section 4.3, we see that the geometric fibers of \mathcal{X} over S are projective. However, this does not imply the projectivity of \mathcal{X} because the projectivity does not satisfy even Zariski descent, see [Har77, Chapter II, Exercise 7.13] for a counterexample.

In this part, we will show that \mathcal{X} is projective over S. We first use the toric structure inside \mathcal{X} to deduce the properness of \mathcal{X} .

Theorem 4.14. The scheme \mathcal{X} is proper over S.

Proof. We are going to verify the valuative criterion for properness [Gro61a, Théorème 7.3.8]. Hence, by base change, we can assume that the base scheme S is the spectrum of a discrete valuation ring R. Let K be the fraction field of R, and let η be the generic point of S.

Since, by Proposition 4.11, \mathcal{X} is separated over S, combining with [Rom13, Lemma 4.1.1], it suffices to show that for any $x \in G(K)$, there exists a section $y \in \mathcal{X}(R)$, such that the following diagram



commutes, where **i** is from Proposition 4.12. By the Iwahori–Cartan decomposition (see, [AHHL21, Remark 3.5, Theorem 4.1]), we have

$$G(K) = G(R)T(K)G(R).$$

Therefore, it is enough to show that any section $z \in T(K)$ can be lifted to an R section of \mathcal{X} . For this, let $Z \subset \mathcal{X}$ be the schematic closure of the generic fiber T_{η} in \mathcal{X} , we claim that Z is a proper toric scheme over S (see [KKMSD73, Chapter IV, § 3] for the theory of toric schemes over DVRs). Assuming the claim, the theorem follows by applying the valuative criterion for properness to Z to lift z to a section in Z(R).

To show the claim, denote by X (resp., X^{\vee}) the character lattice (resp., cocharacter lattice) of T, and by W the Weyl group of G. We denote by $N_G(T)$ the normalizer subgroup scheme of T in G. Note that $N_G(T)$ acts on T, hence, by the flatness of $N_G(T)$ over S ([Con14, Proposition 2.1.2]) and the proof of [GGMB14, Lemme 5.1.1], acts on Z as well. Also, \overline{T} is contained in Z. By Proposition 4.12, $T_{\eta} \subset T$ is embedded into \overline{T} via

$$t \mapsto \prod_{\alpha_i \in \Delta} \alpha_i(t)^{-1},$$

which is an affine embedding of torus defined by the rational polyhedral cone

$$\sigma := C_0 \times \mathbb{R}_{\geq 0} \subset X^{\vee}_{\mathbb{R}} \times \mathbb{R}_{\geq 0},$$

where $C_0 \subset X_{\mathbb{R}}^{\vee}$ is the negative Weyl chamber of G. Similarly, for $w \in W$, since the open immersion **i** in Proposition 4.12 is $(G \times G)$ -equivariant, the embedding from the generic torus T_{η} to $\dot{w} \cdot \overline{T} \cdot \dot{w}^{-1}$ is defined by the rational polyhedral cone

$$\sigma_w := C_w \times \mathbb{R}_{\geq 0} \subset X_{\mathbb{R}}^{\vee} \times \mathbb{R}_{\geq 0},$$

where $\dot{w} \in \mathcal{N}_G(T)(S)$ is a representative of w and $C_w = \{\gamma \in X_{\mathbb{R}}^{\vee} | \langle w\alpha_i, \gamma \rangle \leq 0 \}$ is the translation of the negative Weyl chamber C_0 under the action of w. As the fan of Weyl chambers is complete, we have that the fan defining the toric scheme $\bigcup_{w \in W} \dot{w} \cdot \overline{T} \cdot \dot{w}^{-1}$ which is $\bigcup_{w \in W} \sigma_w$, is also complete in the sense that

$$\bigcup_{w \in W} \sigma_w = X_{\mathbb{R}}^{\vee} \times \mathbb{R}_{\geq 0}.$$

Thus, by [KKMSD73, Chapter IV, §3, (I.), g)], the subscheme $\bigcup_{w \in W} \dot{w} \cdot \overline{T} \cdot \dot{w}^{-1}$ of Z is proper over S, hence equals Z, as desired.

Corollary 4.15. The scheme \mathcal{X} is projective over S.

Proof. First note that the scheme \mathcal{X} is the base change of a scheme \mathcal{X}_0 over $\operatorname{Spec}(\mathbb{Z})$ obtained by applying Definition 4.6 to the Chevalley group that shares the same root datum with G. Thus it suffices to show that \mathcal{X}_0 is projective. The normality of \mathbb{Z} allows us to utilize [BLR90, Section 6.6, Theorem 2 (d)] to get the quasi-projectivity of \mathcal{X}_0 . By [Sta, Lemma 0BCL (1)], the corollary follows from Theorem 4.14.

Remark 4.16. Our method of constructing the projective scheme \mathcal{X} containing G as an open subscheme fails for non adjoint reductive groups. A key feature of our approach is that for the adjoint group G, the set of simple roots forms a basis of the character space $X_{\mathbb{R}}$ and cuts out a rational cone C_0 (the negative Weyl chamber) in the cocharacter space $X_{\mathbb{R}}^{\vee}$ such that the translations of C_0 under the Weyl group form a complete fan in $X_{\mathbb{R}}^{\vee}$. In non adjoint case, there is no such a basis in the character space. In fact, since the action of the Weyl group on the cocharacter space is generated by simple reflections, the cone C_0 cut out by such a basis must lie in some Weyl chamber. Since the fan of the translates of C_0 is complete, the cone C_0 must be a Weyl chamber. Hence this basis is the translation of simple roots under some element in the Weyl group. This implies the root datum of G is adjoint.¹

Remark 4.17. It is also plausible to replace $G \times \overline{\Omega} \times G$ by $G \times \overline{T} \times G$ in the construction of \mathcal{X} , which is similar to [Gab12] where Gabber sketched a proper compactification for pseudoreductive groups of minimal type over fields. Another possible model for \mathcal{X} , which is similar to [BLR90, Chapter 5], is to realize it as the subsheaf of $\mathcal{R}(\overline{\Omega})$ which is the sheaf of rational endomorphisms of $\overline{\Omega}$.

5. Divisors

Divisors of a scheme often reflect many geometric properties of the scheme itself. In this section, we will study the divisors of \mathcal{X} constructed in Section 4 and their interactions to $G \times G$ -orbits of \mathcal{X} . In this section, we shall keep notations of Section 4.

We denote by G_0 the unique Chevalley group over $\text{Spec}(\mathbb{Z})$ which shares the same root datum with G, and denote by \mathcal{X}_0 the scheme obtained by applying Definition 4.6 to G_0 . The following result has been established for classical wonderful compactifications, see, for instance, [BK05, Chapter 6, Lemma 6.19], whose argument can be carried over to our situation with mild modifications, except that it is more natural to consider relative Picard schemes instead of absolute Picard groups.

Theorem 5.1. The complement $\mathcal{X}\setminus\overline{\Omega}$ is the union of the relative effective Cartier divisors $(See [Sta, 062T]) \overline{Bs_iB^-}$, where $s_i \in W(S)$ is the simple reflection defined by the simple root $\alpha_i \in \Delta$ and bar indicates schematic closure in \mathcal{X} . Moreover, the relative Picard functor $P_{\mathcal{X}/S}$, which is the fppf sheaf associated to the functor

$$(Sch/S)^{op} \longrightarrow Grp, \ T \longmapsto Pic(X \times_S T)$$

from the category of S-schemes to the category of groups, is represented by the constant group scheme $\underline{\mathbb{Z}}^{\oplus l}_{S}$.

Proof. As \mathcal{X} is separated by Proposition 4.11, together with the affineness of $\overline{\Omega}$, the open embedding $\mathbf{j} \colon \overline{\Omega} \hookrightarrow \mathcal{X}$ in Lemma 4.10 is affine. By [Gro67, Corollaire 21.12.7], $\mathcal{X} \setminus \overline{\Omega}$ is of pure codimension 1. We claim that $G \setminus \overline{\Omega}$ is schematically dense in $\mathcal{X} \setminus \overline{\Omega}$. In fact, schematic density can be fiberwise checked by [SGA3II, Exposé XVIII, Corollaire 1.3]. Hence, by base change to geometric fibers and Proposition 4.12, the claim follows from the same result for wonderful compactifications [BK05, Chapter 6, proof of Lemma 6.1.9]. Furthermore, by the Bruhat decomposition [GP11b, Exposé XXII, Théorème 5.7.4, (i)], combining with the fact that $G \cap \overline{\Omega} = \Omega_G$, we have

$$G \backslash \overline{\Omega} = \coprod_{w \in W(S) \backslash 1} B \cdot \dot{w} \cdot B^{-},$$

where $\dot{w} \in \operatorname{Norm}_G(T)(S)$ is a lifting of w. Therefore, after taking schematic closure, we obtain

$$\mathcal{X} \setminus \overline{\Omega} = \bigcup_{s_i \in W(S)} \overline{B \cdot \dot{s_i} \cdot B^-},$$

by the usual Bruhat order [Spr09, 8.5.4, Proposition 8.5.5] applied to each geometric fiber.

Let us first assume $S = \text{Spec}(\mathbb{Z})$. In this case, we rename \mathcal{X} to \mathcal{X}_0 . As now S is a Dedekind scheme, by [Har77, Chapter III, Proposition 9.7], these boundary divisors $\overline{B \cdot \dot{s}_i \cdot B^-}$ are flat over S because $\overline{B \cdot \dot{s}_i \cdot B^-}$ dominates S. As \mathcal{X}_0 is locally factorial, by [Har77, Chapter II, Proposition 6.11], $\overline{B \cdot \dot{s}_i \cdot B^-}$ is an effective Cartier divisor, hence is a relative effective

¹We thank Will Sawin for pointing this out to us.

Cartier divisor. For a point $s \in S$, the geometric fiber $(\overline{B \cdot \dot{s}_i \cdot B^-})_{\overline{k(s)}}$ certainly contains $\overline{B_{\overline{k(s)}} \cdot \dot{s}_i \cdot B^-_{\overline{k(s)}}}$, where the closure is taken in the wonderful compactification of $(G_0)_{\overline{k(s)}}$. By [Sta, 056Q], $(\overline{B \cdot \dot{s}_i \cdot B^-})_{\overline{k(s)}}$ is again an effective Cartier divisor. Note that $(\overline{B \cdot \dot{s}_i \cdot B^-})_{\overline{k(s)}}$ can not contain other boundary divisor $\overline{B_{\overline{k(s)}} \cdot \dot{s}_j \cdot B^-_{\overline{k(s)}}}$ for $j \neq i$ otherwise, by intersecting with the group $(G_0)_{\overline{k(s)}}$, a contradiction against the usual Bruhat decomposition [Spr09, 8.5.4] would arise. Therefore, we have

$$(\overline{B \cdot \dot{s_i} \cdot B^-})_{\overline{k(s)}} = \overline{B_{\overline{k(s)}} \cdot \dot{s_i} \cdot B_{\overline{k(s)}}^-}.$$
(9)

Take a Weil divisor D in \mathcal{X}_0 . As $\overline{\Omega}$ is an affine space over $\operatorname{Spec}(\mathbb{Z})$, by [GW20, Proposition 11.40], D is equivalent to a divisor spanned by these boundary divisors. If D is equivalent to a principal divisor associated to a rational function h in the function field $K(\mathcal{X}_0)$, after restricting to $\overline{\Omega}$ which is an affine space over $\operatorname{Spec}(\mathbb{Z})$, we have $h \in \mathbb{Z}^{\times}$, hence D itself is zero. Therefore $\operatorname{Pic}(\mathcal{X}_0)$ is freely generated by these boundary divisors.

Since, by Corollary 4.15, \mathcal{X}_0 is smooth and projective over $\operatorname{Spec}(\mathbb{Z})$ and by Proposition 4.12 the geometric fibers of $\mathcal{X}_0/\operatorname{Spec}(\mathbb{Z})$ are reduced and irreducible, according to the result [BLR90, § 8.2 Theorem 1] of Grothendieck, the relative Picard functor $P_{\mathcal{X}_0/\operatorname{Spec}(\mathbb{Z})}$ is representable by a seperated S-scheme, say $\operatorname{Pic}_{\mathcal{X}_0/S}$, which is locally of finite presentation over S. By Proposition 4.12, the generic fiber of \mathcal{X}_0 over S is geometrically integral, hence, by [GW20, Proposition 5.51], \mathbb{Q} is algebraically closed in the function field $K(\mathcal{X}_0)$. Then by [Gro61b, Corollaire 4.3.12], the pushforward of the structure sheaf $\mathcal{O}_{\mathcal{X}_0}$ to S equals to \mathcal{O}_S . By [BLR90, § 8.1, Proposition 4], $\operatorname{Pic}_{\mathcal{X}_0/S}(S)$ equals the Picard group $\operatorname{Pic}(\mathcal{X}_0)$, which is, as we have seen, the free abelian group $\mathbb{Z}^{\oplus l}$. Hence we can define a morphism

$$\theta: \underline{\mathbb{Z}}^{\oplus l}_S \longrightarrow \operatorname{Pic}_{\mathcal{X}_0/S}$$

by sending an *l*-tuple of integers (a_i) i = 1, ..., l to the divisor $\sum_{i=1}^{l} a_i (\overline{B \cdot \dot{s}_i \cdot B^-})$. By applying [BK05, Corollary 6.2.8] to the wonderful compactification $(\mathcal{X}_0)_{\overline{k(s)}}$ for a point $s \in S$, we have $H^1((\mathcal{X}_0)_{\overline{k(s)}}, \mathcal{O}_{(\mathcal{X}_0)_{\overline{k(s)}}}) = 0$. Combining this with [BLR90, § 8.4 Theorem 1 (b)], it follows that $(\operatorname{Pic}_{\mathcal{X}_0/S})_{\overline{k(s)}}$, which is isomorphic to $\operatorname{Pic}_{(\mathcal{X}_0)_{\overline{k(s)}}/\overline{k(s)}}$, has dimension zero, i.e., is disjoint union of points. By [BK05, Lemma 6.1.9], the underlying topological space of $(\operatorname{Pic}_{\mathcal{X}_0/S})_{\overline{k(s)}}$ is bijective to $\mathbb{Z}^{\oplus l}$ via the basis $\overline{B_{\overline{k(s)}} \cdot \dot{s}_i \cdot B_{\overline{k(s)}}^-}$. Hence, combining Equation (9), each geometric fiber of θ is an isomorphism, and then by [Gro67, Corollaire 17.9.5], θ itself is an isomorphism.

Back to the general base S, by Definition 4.6, \mathcal{X} is the base change of \mathcal{X}_0 along the natural morphism from S to $\operatorname{Spec}(\mathbb{Z})$. By a similar argument using the Bruhat decomposition as above, each boundary divisor $\overline{B \cdot \dot{s}_i \cdot B^-}$ descends to a boundary divisor of \mathcal{X}_0 , hence by [Sta, 056Q], is again a relative effective Cartier divisor. The relative Picard functor $P_{\mathcal{X}/S}$ is represented by the base change of $\operatorname{Pic}_{\mathcal{X}_0/\operatorname{Spec}(\mathbb{Z})}$ along the natural morphism from S to $\operatorname{Spec}(\mathbb{Z})$, which is simply the constant group scheme $\mathbb{Z}^{\oplus l}_{S}$.

For $s \in S$, by Proposition 4.12, the geometric fiber $\mathcal{X}_{\overline{k(s)}}$ is the wonderful compactification of $G_{\overline{k(s)}}$, and $\overline{\Omega}_{\overline{k(s)}}$ is the big cell of $\mathcal{X}_{\overline{k(s)}}$. We denote as h_i the element of $\overline{T}_{\overline{k(s)}}$ which has *i*-th coordinate 1, and 0 otherwise, here $i \in \{1, ..., l\}$. By [BK05, Theorem 6.1.8 (ii)], the complement $\mathcal{X}_{\overline{k(s)}} \setminus G_{\overline{k(s)}}$ is the union of prime divisors $\mathbf{X}_1, ..., \mathbf{X}_l$ with normal crossings, where \mathbf{X}_i is spanned by the element h_i under the action of $G_{\overline{k(s)}} \times_{\overline{k(s)}} G_{\overline{k(s)}}$. A relative version of this result is the following. **Proposition 5.2.** The complement $\mathcal{X} \setminus G$ is covered by $G \times G$ -invariant smooth relative effective Cartier divisors S_{α_i} with relative normal crossings to S^2 , for $\alpha_i \in \Delta$, such that after passing to each geometric fiber, S_{α_i} becomes the boundary divisor \mathbf{X}_i . For a test scheme S' over S, the set of S'-valued points of S_{α_i} is $Q_G(V_{G \times \overline{\Omega} \times G}(\mathbb{X}_{\alpha_i})(S'))$ where Q_G is the quotient map in Definition 4.6.

Proof. We first prove the proposition when $S = \text{Spec}(\mathbb{Z})$. By the same argument as in the proof of Theorem 5.1, $\mathcal{X} \setminus G$ is of pure codimension 1. Consider an irreducible component **C** of $\mathcal{X} \setminus G$. As $\mathcal{X} \setminus G$ is $G \times G$ -stable and G is connected, \mathbf{C} is $G \times G$ -stable. Since $\mathbf{C} = \bigcup_{a \in G \times G} g \cdot (\mathbf{C} \cap \overline{\Omega})$ and C is of codimension 1 in \mathcal{X} , $\mathbf{C} \cap \overline{\Omega}$ is of codimension 1 in the big cell $\overline{\Omega}$. Note that now the coordinate ring of $\overline{\Omega}$ is a polynomial ring over \mathbb{Z} which is a UFD, hence by the Krull's Hauptidealsatz (see, for instance, [Har77, Chapter I, Theorem 1.11A]), $\mathbf{C} \cap \overline{\Omega}$ is a hypersurface in $\overline{\Omega}$, i.e., defined by a single element of the coordinate ring of $\overline{\Omega}$. Notice that $\mathbf{C} \cap \overline{\Omega}$ is also $U^{-}T \times U^{+}$ -stable, hence it is nothing else but the hypersurface defined by the coordinate \mathbb{X}_{α_i} of the big cell $\overline{\Omega}$ for a simple root $\alpha_i \in \Delta$. Therefore, combining with the $G \times G$ action, we obtain the final statement of the proposition. Since \mathcal{X} is covered by the $G \times G$ translations of $\overline{\Omega}$ whose local rings now are all UFDs, by [Har77, Chapter II, Proposition 6.11], the prime Weil divisor C is locally principal. The assertion of the proposition about geometric fibers follows from the construction of the embedding \mathbf{i} as in the proof of Proposition 4.12. Finally, by [Sta, Lemma 062Y], the locally principal subscheme C in \mathcal{X} is a relative effective Cartier divisor because so are all its geometric fibers. The divisor $\mathcal{X} \setminus G$ is of relative normal crossings to S because $\mathcal{X} \setminus G \cap (g_1, g_2) \overline{\Omega}$ is of strict relative normal crossings in $(g_1, g_2) \overline{\Omega}$ for every $(g_1, g_2) \in (G \times G)(S)$.

Back to the original base S, note that by the functoriality of the construction of \mathcal{X} , \mathcal{X} comes from the base change of the scheme \mathcal{X}_0 obtained by applying Definition 4.6 to G_0 . By the functorial description of the boundary divisors of \mathcal{X}_0 in the previous case, we deduce that the boundary $\mathcal{X} \setminus G$ is covered by the base changes of the relative Cartier divisors in the boundary $\mathcal{X}_0 \setminus G_0$ along the morphism $S \longrightarrow \operatorname{Spec}(\mathbb{Z})$, which are still relative effective Cartier divisors by [Sta, Lemma 056Q]. The relative normal crossings to S of $\mathcal{X} \setminus G$ inherit from that of $\mathcal{X}_0 \setminus G_0$. \Box

The following result says that the equivariant compactification \mathcal{X} of the adjoint group scheme G is a schematic version of a wonderful variety in the sense of [Lun96]. Before proceeding to the theorem, we first clarify the notion of an orbit for a group scheme action, that is used in this paper. For an S-scheme Y acted on by a group scheme H over S and an S-valued point $x \in Y(S)$, we refer the subsheaf of Y spanned by x under the action of H as the orbit of x (compare with [MFK94, Definition 0.4]).

For a subset $I \subset \Delta$, let $z_I \in \mathcal{X}(S)$ be the section represented by

$$\dot{z}_I := (e, (e, \omega_I, e), e) \in (G \times \overline{\Omega} \times G)(S), \tag{10}$$

where $\omega_I = (\omega_1, \omega_2, ..., \omega_l) \in \prod_{\Delta} \mathbb{G}_m(S)$ such that $\omega_i = 0$ if $i \in I$ and $\omega_i = 1$ otherwise.

Theorem 5.3. Each $G \times G$ -orbit in \mathcal{X} in the above sense is represented by a scheme. For a $G \times G$ -orbit in \mathcal{X} , there is a unique subset I of Δ such that the schematic closure of the orbit

• $D = \bigcup_{i \in I} V_X(f_i);$

• for every $x \in \text{Supp}(D)$, X is smooth on x over S, and the closed subscheme $V((f_i)_{i \in I(x)}) \subset X$ is smooth over S of codimension |I(x)|, where $I(x) = \{i \in I | f_i(x) = 0\}$.

The divisor D is with relative normal crossings to S, if étale locally it is strictly with relative normal crossings.

²Following the terminology of [03, Exposé XIII, 2.1], for a scheme X over a scheme S, we say that a relative Cartier divisor $D \subset X$ is strictly with relative normal crossings to S, if there exists a finite family $(f_i \in \Gamma(X, \mathcal{O}_X))_{i \in I}$ such that

in \mathcal{X} is $\bigcap_{\alpha_i \in I} S_{\alpha_i}$, where S_{α_i} are the boundary divisors in Proposition 5.2 and the intersection is transversal.

Proof. For a section $\gamma \in \mathcal{X}(S)$ represented by a section

$$(u, (x, t, y), v) \in (G \times \overline{\Omega} \times G)(S'),$$

where S' is an fppf cover of S, by the definition of the group action on \mathcal{X} (Definition-Proposition 4.9),

$$\gamma = (u, v) \cdot (e, (x, t, y), e).$$

Furthermore, by Definition 4.1, we have $(e, (x, t, y), e) \sim (x, (e, t, e), y) = (x, y) \cdot (e, (e, t, e), e)$. By the adjointness of G, there exists a section $\dot{t} \in T(S')$ such that $t = \dot{t} \cdot \omega_I$ for some $I \subset \Delta$, then $(e, (e, t, e), e) = (\dot{t}, e) \cdot (e, (e, \omega_I, e), e)$. Therefore, the $G \times G$ -orbit \mathbf{O}_{γ} of γ is spanned by z_I under the action of $G \times G$, and \mathbf{O}_{γ} is, as a sheaf, isomorphic to

$$Q_G(\bigcap_{i\in I} V_{G\times\overline{\Omega}\times G}(\mathbb{X}_{\alpha_i}) \bigcap_{i\notin I} D_{G\times\overline{\Omega}\times G}(\mathbb{X}_{\alpha_i})),$$

where Q_G is the quotient map in Definition 4.6. Note that now \mathbf{O}_{γ} can be identified with the quotient sheaf of the subscheme $\bigcap_{i \in I} V_{G \times \overline{\Omega} \times G}(\mathbb{X}_{\alpha_i}) \bigcap_{i \notin I} D_{G \times \overline{\Omega} \times G}(\mathbb{X}_{\alpha_i})$ of $G \times \overline{\Omega} \times G$ with respect to the restriction of the equivalence relation in Theorem 4.7 to this subscheme, which is again an fppf relation. Thus by a result due to Artin [LMB00, Corollaire (10.4)], \mathbf{O}_I is an algebraic space over S. By intersecting the open subscheme $\overline{\Omega}$ of \mathcal{X} (Lemma 4.10) and its translations by $G \times G$, we see that \mathbf{O}_{γ} , Zariski locally on \mathcal{X} , is a subscheme of \mathcal{X} . By [Sta, 03M4], being an immersion for a morphism of algebraic spaces is Zariski local on target, hence \mathbf{O}_{γ} is a subscheme of \mathcal{X} . Observe that by the description of the open immersion from $\overline{\Omega}$ into \mathcal{X} in Lemma 4.10, $\mathbf{O}_{\gamma} \cap \overline{\Omega}$ is simply $\bigcap_{i \in I} V_{\overline{\Omega}}(\mathbb{X}_{\alpha_i}) \bigcap_{i \notin I} D_{\overline{\Omega}}(\mathbb{X}_{\alpha_i})$, which is schematic dense in $(\bigcap_{\alpha_i \in I} S_{\alpha_i}) \cap \overline{\Omega}$. Hence, by the translation of $G \times G$, the schematic closure of \mathbf{O}_{γ} in \mathcal{X} is $\bigcap_{\alpha_i \in I} S_{\alpha_i}$, here the intersection is transversal because so is it over $\overline{\Omega}$ (see the proof of Proposition 5.2).

6. Descent

An adjoint reductive group is descended from a split adjoint reductive group. In this section, we deduce an equivariant compactification of a general adjoint reductive group scheme by twisting the equivariant compactification (defined in Section 4) of a split form of this group.

6.1. **Descent datum of groups.** We consider an adjoint semisimple group scheme (not necessarily split) G over a scheme S. If the fiber product is formed over S, we will omit the subscript S. By [GP11b, Exposé XXII, Corollaire 2.3], this G splits after base change to an étale cover U of S, and, by [GP11b, Exposé XXV, Théorème 1.1], there exists a split adjoint group G_0 over S such that G is a form of G_0 . We fix a maximal split torus T of G_0 as in [GP11b, Exposé XXII, Définition 1.13] with $\operatorname{rk}(T) = n$, in a Borel subgroup B of G_0 , and denote by $(X, \Psi, \Delta, X^{\vee}, \Psi^{\vee}, \Delta^{\vee})$ the corresponding based root datum. We also fix a pinning $\{X_a\}_{a\in\Delta}$ of G_0 , where $X_a \in \Gamma(S, \mathfrak{g}_a)^{\times}$ and \mathfrak{g}_a is the weight subsheaf of the Lie algebra \mathfrak{g} of G_0 with respect to the weight a. By Proposition 4.12 and Corollary 4.15, we obtain a projective smooth $(G_0 \times G_0)$ -scheme \mathcal{X}_0 that equivariantly contains G_0 as an open dense subscheme.

Denote by U' (resp., U'') the 2-fold (resp., 3-fold) product of U over S. By [Con14, Example 7.1.4], the group G is descended from $G_0 \times U$ via a Čech 1-cocycle

$$\xi \in \mathbb{Z}^1(U/S, \operatorname{Aut}_S(G_0)).$$

More precisely, this ξ is an automorphism of the group scheme $\tilde{G} := G_0 \times U'$ over U' such that $\operatorname{pr}_{13}^*(\xi) = \operatorname{pr}_{12}^*(\xi) \operatorname{pr}_{23}^*(\xi)$. By [GP11b, Exposé XXIV, Théorème 1.3], we have the following short exact sequence of group schemes:

$$1 \longrightarrow \widetilde{G} \longrightarrow \operatorname{Aut}_{U'}(\widetilde{G}) \longrightarrow \operatorname{Out}_{U'}(\widetilde{G}) \longrightarrow 1.$$
(11)

Further, the induced pinning $\{\tilde{X}_a\}_{a\in\Delta}$ on \tilde{G} from the pinning $\{X_a\}_{a\in\Delta}$ splits the above short exact sequence and yields an isomorphism:

$$\operatorname{Aut}_{U'}(\widetilde{G}) \simeq \widetilde{G} \rtimes \operatorname{Out}_{U'}(\widetilde{G}).$$
 (12)

At the same time, since \tilde{G} is split, by *loc. cit.*, the outer automorphism group scheme $\operatorname{Out}_{U'}(\tilde{G})$ is identified with a constant group scheme $E_{U'}$, where E is the group of automorphisms of the based root datum $(X, \Psi, \Delta, X^{\vee}, \Psi^{\vee}, \Delta^{\vee})$.

For the element η , by [GP11b, Exposé XXIV, Théorème 1.3 (iii)], it is induced by an automorphism $(\rho, {}^t \rho)$ of the based root datum $(X, \Psi, \Delta, X^{\vee}, \Psi^{\vee}, \Delta^{\vee})$. More precisely, this ρ (resp., ${}^t \rho$) is an automorphism of the abelian group X (resp., X^{\vee}), and it induces a bijection on Ψ (resp., Ψ^{\vee}) and preserves simple roots (resp., simple coroots) (see [GP11b, Exposé XXI, Définition 6.1]). Now η is determined in the following way (see [GP11b, Exposé XXIII, proof of Théorème 4.1 and Exposé XXIII, 1.5]):

• ρ induces an automorphism ρ_T of T:

$$\rho_T = \mathcal{D}_{U'}(\rho) \colon T = \mathcal{D}_{U'}(X) \longrightarrow \mathcal{D}_{U'}(X) = T;$$

• For each simple root $\alpha \in \Delta$, set

$$n_{\alpha} = p_{\alpha}(1)p_{-\alpha}(-1)p_{\alpha}(1),$$

where $p_{\alpha} : \mathbb{G}_{a,U'} \to U_{\alpha}$ is given by the corresponding X_{α} in the pinning we fixed. Then ρ_T is extended to an automorphism ρ_N of $\operatorname{Norm}_G(T)$ by taking n_{α} to $n_{\rho(\alpha)}$.

• For each simple root $\alpha \in \Delta$, set

$$\rho_{\alpha} \colon U_{\alpha} \longrightarrow U_{\rho(\alpha)},$$
$$p_{\alpha}(x) \longmapsto p_{\rho(\alpha)}(x).$$

• For each root $r \in \Psi$, there exists a $n \in \operatorname{Norm}_{\widetilde{G}}(T)(U')$ such that $\overline{n}(r) \in \Delta$, \overline{n} is the image of n in the Weyl group W. Thus, by [GP11b, Exposé XXIII, Lemme 2.3.5], we have a morphism

$$\rho_r \colon U_r \longrightarrow U_{\rho(r)},$$

such that $\operatorname{Ad}(\rho_N(n))\rho_r(x) = \rho_{\overline{n}(r)}(\operatorname{Ad}(n)x)$, for any $x \in U_r(V), V \to U'$.

Then, modulo some technical arguments as in [GP11b, Exposé XXIII, proof of Théoreme 4.1], the above data glue together to yield the group automorphism η .

6.2. Descent datum for compactifications. Now we are in the position to define a descent datum on $\mathcal{X}_0 \times U'$. Since the construction in Definition 4.6 commutes with any base change, then $\mathcal{X}_0 \times U'$ is identified to the compactification of \tilde{G} , denoted by $\tilde{\mathcal{X}}$. By Equation (12), we can uniquely decompose ξ into the product of $\delta \in \tilde{G}(U')$ and $\eta \in \operatorname{Out}_{U'}(\tilde{G})(U')$.

We are going to define two automorphisms induced from δ and η on $\tilde{\mathcal{X}}$ respectively and then, take their composition as an action induced by ξ on $\tilde{\mathcal{X}}$.

For the element δ , we define its action on $\hat{\mathcal{X}}$ to be the conjugation action, see Definition-Proposition 4.9.

For the element η , inspired by the above explicit description of η , we define an automorphism $\rho_{\overline{T}}$ of $\overline{T} = \prod_{\Delta} (\mathbb{G}_a)$ to be the permutation of coordinates induced by ρ so that the following diagram



commutes, where two horizontal morphisms are the embedding of T induced by negative simple roots as Equation (2). As ρ preserves Δ , the η defines an automorphism \hat{H} of $\tilde{G} \times \overline{\Omega} \times \tilde{G}$ by applying $\rho_{\overline{T}}$ and ρ_r , $r \in \Psi$ to each component, where $\overline{\Omega} = U^+ \times \overline{T} \times U^+$ is over U'.

Lemma 6.1. The automorphism \hat{H} of $\tilde{G} \times \overline{\Omega} \times \tilde{G}$ preserves the relation defined in Definition 4.1.

Proof. In the viewing of Definition 4.1, it suffices to show that the following diagram of U'-rational morphisms



commutes, where the dotted vertical arrows indicate that π is a U'-rational morphism. Since \hat{H} is induced by η which is an automorphism of G, the commutativity of the diagram then follows from Theorem 3.4.

Thanks to Lemma 6.1, combining with Definition 4.6, \hat{H} induces an automorphism H of $\tilde{\mathcal{X}}$. We take H as the action induced by η on $\tilde{\mathcal{X}}$. We thus obtain an automorphism φ of $\tilde{\mathcal{X}}$ by composing H and the conjugation action of δ . We claim that φ is a descent datum for the projective scheme $\mathcal{X}_0 \times U'$. In fact, it suffices to check the cocycle condition on φ , which follows from the cocycle condition on ξ and the schematic density of G_0 in \mathcal{X}_0 .

6.3. Effectivity of descent datum. A descent datum for a projective scheme, in general, is not effective (see [Sta, 08KE]). In order to make our descent datum effective, we are going to construct an ample line bundle compatible with the descent datum φ . To do this, we need the following combinatorial lemma on root datum.

Lemma 6.2. For the based root datum $\mathbf{R} = (X, \Psi, \Delta, X^{\vee}, \Psi^{\vee}, \Delta^{\vee})$, there exists a linear combination $\gamma = \sum_{\alpha_i \in \Delta} n_i \alpha_i$ of simple roots with $n_i \in \mathbb{Z}_{\geq 0}$ such that

• γ is invariant under arbitrary automorphism of **R**;

 α_1

• γ is regular dominant in the sense that for any simple root $\alpha_i \in \Delta, <\gamma, \alpha_i^{\vee} >> 0$.

Proof. By [GP11b, Exposé XXI, Corollaire 7.1.6], the root datum \mathbf{R} decomposes into product of irreducible adjoint root data, hence we assume that \mathbf{R} is irreducible. As an irreducible adjoint root datum is determined by the corresponding Dynkin diagram ([GP11b, Exposé XXI, Corollaire 7.4.4]), we can analyse \mathbf{R} case by case.

• **R** is of type $A_n \ (n \ge 2)$.

$$\alpha_2 \quad \alpha_3 \quad \alpha_{n-2} \quad \alpha_{n-1} \quad \alpha_n$$

Suppose $\gamma = \sum_{i=1}^{n} m_i \alpha_i$ with $m_i \in \mathbb{Z}_{\geq 0}$. To make γ satisfy the requirements in the lemma, it suffices to impose the following three conditions on coefficients:

$$m_{i} = m_{n-i}, \text{ for all } 1 \leq i \leq \left[\frac{n}{2}\right];$$
$$< \sum_{i=1}^{n} m_{i} \alpha_{i}, \alpha_{j}^{\vee} >= 2m_{j} - m_{j-1} - m_{j+1} \geq 0, \text{ for all } 2 \leq j \leq \left[\frac{n}{2}\right];$$
$$< \sum_{i=1}^{n} m_{i} \alpha_{i}, \alpha_{1}^{\vee} >= 2m_{1} - m_{2}.$$

Notice that the above three requirements can be fulfilled for some m_i .

• **R** is of type D_n $(n \ge 5)$.



In the view of the first requirement in the lemma, we suppose that $\gamma = \sum_{i=1}^{n-2} m_i \alpha_i + m_{n-1}(\alpha_{n-1} + \alpha_n)$ with $m_i \in \mathbb{Z}_{\geq 0}$. To make γ regular dominant, it suffices to impose the following conditions on the coefficients:

$$\langle \gamma, \alpha_{1}^{\vee} \rangle = 2m_{1} - m_{2} > 0; \quad \langle \gamma, \alpha_{j}^{\vee} \rangle = 2m_{j} - m_{j+1} - m_{j-1}, \text{ for all } 2 \leqslant j \leqslant n-3;$$

$$\langle \gamma, \alpha_{n-2}^{\vee} \rangle = 2m_{n+1} - m_{n-3} - m_{n-1} - m_{n} > 0;$$

$$\langle \gamma, \alpha_{n-1}^{\vee} \rangle = 2m_{n-1} - m_{n-2} > 0; \quad \langle \gamma, \alpha_{n}^{\vee} \rangle = 2m_{n} - m_{n-2} > 0.$$

A simple computation yields a solution to the above inequalities.

• **R** is of type E_6 .



It suffices to take, for instance, $\gamma = 31(\alpha_1 + \alpha_6) + 60(\alpha_3 + \alpha_5) + 85\alpha_4 + 49\alpha_2$.

• \mathbf{R} is of type D_4 .



It suffices to take, for instance, $\gamma = 5\alpha_2 + 3(\alpha_1 + \alpha_3 + \alpha_4)$.

• **R** is of type A_1 , B_n , C_n , E_7 , E_8 , F_4 , G_2 . There is no nontrivial automorphism on Dynkin diagram of these types, hence on **R**. Therefore, by the adjointness, it suffices to choose any regular dominant character as γ .

Theorem 6.3. There exists an ample effective Cartier divisor \mathbf{D} on $\widetilde{\mathcal{X}}$ such that \mathbf{D} is stable under the automorphism φ of $\widetilde{\mathcal{X}}$, and \mathbf{D} descends to an ample effective Cartier \mathbf{D}_0 of \mathcal{X}_0 .

Proof. By Lemma 6.2, there is a regular dominant weight $\gamma = \sum_{i=1}^{n} n_i \alpha_i$ of the based root datum **R** such that γ is stable under the action ρ . Consider the effective Cartier divisor $\mathbf{D} := \sum_{i=1}^{n} n_i S_{\alpha_i}$, here S_{α_i} are the boundary effective Cartier divisor of $\hat{\mathcal{X}}$ from Proposition 5.2. By [Gro65a, Corollaire 9.6.5], being relative ample can be checked fiberwise. Hence, to show that **D** is U'-ample, we are left to check that, for each point $u \in U'$, the geometric fiber

 $\mathbf{D}_{\overline{u}} = \sum_{i=1}^{n} n_i (S_{\alpha_i})_{\overline{u}}$ is an ample divisor of the geometric fiber $\widetilde{\mathcal{X}}_{\overline{u}}$ which, by Proposition 4.12, is identified with the wonderful compactification of $\widetilde{G}_{\overline{u}}$. By Proposition 5.2, $(S_{\alpha_i})_{\overline{u}}$ is a boundary divisor in $\widetilde{\mathcal{U}} \setminus \widetilde{G}_{\overline{u}}$. By [BK05, Chapter 6, Proposition 6.1.11 (ii)], $(S_{\alpha_i})_{\overline{u}}$ corresponds to the line bundle $\mathcal{L}_{\widetilde{G}_{\overline{u}}}(\alpha_i)$. Thus we have that

$$\mathcal{L}_{\widetilde{G}_{\overline{u}}}(\mathbf{D}_{\overline{u}}) = \sum_{i=1}^{n} n_i (S_{\alpha_i})_{\overline{u}} = \sum_{i=1}^{n} n_i \mathcal{L}_{\widetilde{G}_{\overline{u}}}(\alpha_i) = \mathcal{L}_{\widetilde{G}_{\overline{u}}}(\sum_{i=1}^{n} n_i \alpha_i) = \mathcal{L}_{\widetilde{G}_{\overline{u}}}(\gamma).$$

As γ is regular dominant, by [BK05, Chapter 6, Proposition 6.1.11 (iii)], $\mathcal{L}_{\widetilde{G}_{\overline{u}}}(\gamma)$ is ample. Hence **D** is an ample effective Cartier divisor.

Now we show that **D** is stable under φ . Recall that φ is the composition of the conjugate action of an element of $\tilde{G}(U')$ and the automorphism H defined by the automorphism (f, f) of the based root datum **R**. By Proposition 5.2, each S_{α_i} is $(\tilde{G} \times \tilde{G})$ -stable, hence so is **D**. Moreover, by *loc. cit.*, we see that H takes S_{α_i} to $S_{\alpha_{f(i)}}$, and their coefficients n_i and $n_{f(i)}$ in **D** is designed to be equal, thus **D** is stable under H as well, as desired.

By applying Proposition 5.2 to \mathcal{X}_0 , we get divisors $S_{0 \alpha_i}$ on the boundary of G_0 in \mathcal{X}_0 . Notice that, by the functorial description of S_{α_i} in Proposition 5.2, each S_{α_i} descends to $S_{0 \alpha_i}$. Thus **D** descends to $\mathbf{D}_0 = \sum_{i=1}^n n_i S_{0 \alpha_i}$ which is an ample effective divisor by the same argument as above.

Corollary 6.4. The $(G_0 \times U) \times (G_0 \times U)$ -scheme $\mathcal{X}_0 \times U$ over U descends to a smooth projective $(G \times G)$ -scheme \mathcal{X} over S such that the geometric fibers of \mathcal{X} are the wonderful compactifications of the corresponding geometric fibers of G, and \mathcal{X} contains G as a fiberwise dense open subscheme. Moreover, the boundary $\mathcal{X} \setminus G$ is a smooth relative Cartier divisor with relative normal crossings to S.

Proof. By [BLR90, Section 6.1, Theorem 7], the descent datum φ of the pair

$$(\mathcal{X}_0 \times U, \mathcal{L}_{\mathcal{X}_0 \times U}(\mathbf{D}_0 \times U))$$

is effective, where \mathbf{D}_0 is given in Theorem 6.3. Thus $\mathcal{X}_0 \times U$ descends to a scheme \mathcal{X} over S, which is quasi-projective over S because $\mathcal{L}_{\mathcal{X}_0 \times U}(\mathbf{D}_0 \times U)$ descends to a relative ample line bundle on \mathcal{X} . By [Gro65a, Proposition (2.7.1) (vii)], this \mathcal{X} inherits properness from \mathcal{X}_0 , hence \mathcal{X} is projective over S.

In order to descend the action of $(G_0 \times U) \times (G_0 \times U)$ on the scheme $\mathcal{X}_0 \times U$, we have to show that the following diagram:

$$\begin{array}{c} \widetilde{G} \times \widetilde{\mathcal{X}} \times \widetilde{G} \xrightarrow{\widetilde{\Theta}} \widetilde{\mathcal{X}} \\ & \downarrow_{\xi \times \varphi \times \xi} & \downarrow^{\varphi} \\ \widetilde{G} \times \widetilde{\mathcal{X}} \times \widetilde{G} \xrightarrow{\widetilde{\Theta}} \widetilde{\mathcal{X}} \end{array}$$

commutes, where $\widetilde{\Theta}$ is the action of $\widetilde{G} \times \widetilde{G}$ on $\widetilde{\mathcal{X}}$ defined in Definition-Proposition 4.9. Recall that ξ (resp. φ) consists of an inner part and an outer part, hence it suffices to verify the above diagram separately. For the inner part, the commutativity follows from the group action law on \widetilde{X} ; for the outer part, it follows from the definition of φ . Then by [BLR90, Section 6.1, Theorem 6 (a)], this $\widetilde{\Theta}$ descends to a morphism $\Theta: G \times \mathcal{X} \times G \longrightarrow \mathcal{X}$ which is again a group action. In order to descend the open embedding from $G_0 \times U$ into $\mathcal{X}_0 \times U$, we need to show that the following diagram



commutes, where \tilde{i} is the open embedding defined in Proposition 4.12. The commutativity of the above diagram follows from the definition of φ . Again, by [BLR90, Section 6.1, Theorem 6 (a)], this \tilde{i} descends to a morphism $i: G \longrightarrow \mathcal{X}$ which is an open immersion by [Gro65a, Proposition 2.7.1(x)]. Finally, note that for any geometric point $\operatorname{Spec}(\overline{k(s)}) \to S$ of S, where $s \in S$, by [Gro67, Corollaire 17.16.2], there is a section of the base change $U \times_S \operatorname{Spec}(\overline{k(s)})$ over $\operatorname{Spec}(\overline{k(s)})$. Then the fiberwise claim for the point s is fulfilled via this section and Proposition 4.12.

By Proposition 5.2, the complement $\mathcal{X}_0 \setminus G_0$ is of relative normal crossing to S. Hence, since $\mathcal{X} \setminus G$ is descended from $(\mathcal{X}_0 \setminus G_0) \times U$ by the definition of being of relative normal crossings [03, Exposé XIII, 2.1], $\mathcal{X} \setminus G$ is of relative normal crossing to S as well.

Remark 6.5. In [Wed18], Wedhorn introduced the notion of a spherical space over an arbitrary base scheme as a generalization of a spherical variety over an algebraically closed field. More precisely, in loc. cit., a spherical space for a reductive group scheme G over a scheme S is defined to be a flat, separated, and finitely presented algebraic space X over S together with an action by G such that for any $s \in S$, $X_{\overline{k(s)}}$ is a spherical $G_{\overline{k(s)}}$ -variety. Since by Luna's result [Lun96] which says that every wonderful variety is spherical, Corollary 6.4 produces plenty of meaningful examples of spherical spaces.

7. Applications

In this section, we discuss applications of the compactification constructed in Section 6 on torsors of reductive group schemes.

The initial motivation of this paper comes from the following conjecture about equivariant compactification for reductive group schemes by Česnavičius:

Conjecture 7.1. ([Čes22, Conjecture 6.2.3]). For an isotrivial reductive group G over a Noetherian scheme S in the sense that there exists a finite étale cover S' of S such that $G \times_S S'$ splits, there exists a projective, finitely presented scheme X containing G as a fiberwise dense open subscheme, together with a left action of G on X that extends the left translation of G on itself.

If S is a field, then such an X can be obtained by simply taking schematic closure of G in some projective space. The main difficulty of Conjecture 7.1 lies in the fact that, in general, the schematic image of a morphism does not commute with nonflat base change.

Note that Corollary 6.4 gives an affirmative answer to Conjecture 7.1 for adjoint reductive group schemes without the isotriviality restriction, and also the left and the right Gtranslations are both extended. Notice also that the solution of Conjecture 7.1 for tori has been given, see [Čes22, Theorem 6.3.1].

The following result is established in [Ces22, Proposition 6.2.4] for isotrivial torsors of an isotrivial reductive group schemes under the assumption that Conjecture 7.1 holds.

Proposition 7.2. For an isotrivial torsor X under an isotrivial adjoint reductive group scheme G over a Noetherian scheme S, there exists a projective, finitely presented G-scheme \overline{X}/S

containing X as a fiberwise open dense subscheme such that the G-action on \overline{X} restricts to the G-action on X. If S is the spectrum of a semilocal Noetherian ring, then, for any section a from a closed subscheme $Z \subset S$ to X, there exists a finite étale cover \widetilde{S} of S, a morphism $\nu: Z \to \widetilde{S}$ and a section $\widetilde{a} \in X(\widetilde{S})$ whose ν -pullback is a.

Proof. The proposition follows from Corollary 6.4 and [Čes22, Proposition 6.2.4]. More precisely, the construction of \overline{X} is obtained by twisting the compactification \mathcal{X} of G given in Corollary 6.4. Let \overline{X} be the contracted product $X \times^G \mathcal{X}$, which is an algebraic space by [Sta, 06PH], and by étale descent for quasi-affine schemes [Sta, 0247], is a scheme. See [Čes22, proof of Proposition 6.2.4] for the projectivity of \overline{X} over S, in which the isotriviality condition is used. For the last part of the proposition, roughly, we use the Bertini theorem to cut the compactification \overline{X} to produce a finite étale cover of \overline{Z} , see [Čes22, Lemma 6.2.2] for details.

Remark 7.3. The subtlety of Proposition 7.2 lies in arranging the finiteness of \tilde{S} . Without the finiteness requirement, the second part of the proposition follows from [Čes22, Proposition 6.1.1 (a)].

Another application is the following trick of equating reductive group schemes, which is frequently used (in various forms) in the study of the Grothendieck–Serre conjecture (see [Čes22, Section 3.1] for a detailed survey of this conjecture). The following result is established in [Čes22, Proposition 6.2.5] under the assumption of Conjecture 7.1. A surprising point of the following proposition is that it works for reductive group schemes even if we only dispose of the compactifications for adjoint reductive group schemes.

Proposition 7.4. Let S be the spectrum of a Noetherian semilocal ring whose local rings are all geoemetrically unibranch, e.g., normal (see [Gro64, 23.2.1]), and let $Z \subset S$ be a closed subscheme. Assume that G_1 and G_2 are two reductive group schemes over S that have the same root datum over each geometric fiber, and that there is an isomorphism of group schemes $\varphi: G_1|_Z \to G_2|_Z$. Then, there are a finite étale cover \widetilde{S} of S, a morphism $c: Z \to \widetilde{S}$, and an isomorphism $\widetilde{\varphi}: G_1 \times_S \widetilde{S} \to G_2 \times_S \widetilde{S}$ that lifts φ along c.

Proof. The proposition follows from Corollary 6.4 and [Ces22, Proposition 6.2.5]. Roughly, by [GP11b, Exposé XXII, Corollaire 2.3], G_1 and G_2 split étale locally over S. Then thanks to the fiberwise assumption, G_1 and G_2 are étale locally isomorphic. Thus, the quotient, say $\overline{\mathbf{I}}$, of the isomorphism scheme $\mathbf{I} := \text{Isom}_{\text{grp}}(G_1, G_2)$ by the action of $(G_1)_{\text{ad}}$ is étale locally constant because, étale locally, $\overline{\mathbf{I}}$ is the outer automorphism scheme of a split reductive group scheme, which is, by [GP11b, Exposé XXIV, Théorème 1.3 (iii)], a constant group scheme. By descent [Gro65a, Proposition 2.6.1 (i)] and [Gro67, Corollaire 17.7.3 (ii)], $\overline{\mathbf{I}}$ is étale and surjective over S. By [SGA3II, Exposé X, Corollaire 5.14] and [Gro60, Corollaire 6.1.9], $\overline{\mathbf{I}}$ decomposes as a disjoint union of connected components, which are clopen, finite and étale over S. Now consider the section $\overline{\varphi} \in \overline{\mathbf{I}}(Z)$ induced by φ , and take \overline{Z} as the finite disjoint union of all connected components of $\overline{\mathbf{I}}$ that intersect the image of Z. Then \overline{Z} is finite étale over S, and the natural embedding of \overline{Z} into $\overline{\mathbf{I}}$, denoted by $\overline{\alpha}$, gives a lifting of $\overline{\varphi}$. By base changing the $(G_1)_{\text{ad}}$ -torsor \mathbf{I} over $\overline{\mathbf{I}}$ along $\overline{\alpha}$ and considering the schematic image of Z in \overline{Z} , the proposition then follows from Proposition 7.2.

References

[[]AHHL21] Jarod Alper, Jochen Heinloth, and Daniel Halpern-Leistner, Cartan-Iwahori-Matsumoto decompositions for reductive groups, Pure Appl. Math. Q. 17 (2021), no. 2, 593–604. MR4257594

- [BK05] Michel Brion and Shrawan Kumar, Frobenius splitting methods in geometry and representation theory, Progress in Mathematics, vol. 231, Birkhäuser Boston, Inc., Boston, MA, 2005. MR2107324
- [BLR90] Siegfried Bosch, Werner Lütkebohmert, and Michel Raynaud, Néron models, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 21, Springer-Verlag, Berlin, 1990. MR1045822
- [Bri03] Michel Brion, Group completions via Hilbert schemes, J. Algebraic Geom. 12 (2003), no. 4, 605–626. MR1993758
- [Čes22] Kęstutis Česnavičius, Problems about torsors over regular rings, Acta Math. Vietnam. 47 (2022), no. 1, 39–107. With an appendix by Yifei Zhao. MR4406561
- [Con14] Brian Conrad, *Reductive group schemes*, Autour des schémas en groupes. Vol. I, 2014, pp. 93–444. MR3362641
- [DCP83] C. De Concini and C. Procesi, Complete symmetric varieties, Invariant theory (Montecatini, 1982), 1983, pp. 1–44. MR718125
- [ER15] Bas Edixhoven and Matthieu Romagny, Group schemes out of birational group laws, Néron models, Autour des schémas en groupes. Vol. III, 2015, pp. 15–38. MR3525600
- [Fal97] Gerd Faltings, Explicit resolution of local singularities of moduli-spaces, J. Reine Angew. Math. 483 (1997), 183–196. MR1431845
- [Gab12] Ofer Gabber, On pesudo-reductive groups and compactification theorems, Oberwolfach Reports (2012), 2371–2374.
- [GGMB14] Ofer Gabber, Philippe Gille, and Laurent Moret-Bailly, Fibrés principaux sur les corps valués henséliens, Algebr. Geom. 1 (2014), no. 5, 573–612. MR3296806
 - [Gin89] Victor Ginsburg, Admissible modules on a symmetric space, 1989, pp. 9–10, 199–255. Orbites unipotentes et représentations, III. MR1021512
 - [GP11a] Philippe Gille and Patrick Polo (eds.), Schémas en groupes (SGA 3). Tome I. Propriétés générales des schémas en groupes, Documents Mathématiques (Paris) [Mathematical Documents (Paris)], vol. 7, Société Mathématique de France, Paris, 2011. Séminaire de Géométrie Algébrique du Bois Marie 1962–64. [Algebraic Geometry Seminar of Bois Marie 1962–64], A seminar directed by M. Demazure and A. Grothendieck with the collaboration of M. Artin, J.-E. Bertin, P. Gabriel, M. Raynaud and J-P. Serre, Revised and annotated edition of the 1970 French original. MR2867621
 - [GP11b] (ed.), Schémas en groupes (SGA 3). Tome III. Structure des schémas en groupes réductifs, Documents Mathématiques (Paris) [Mathematical Documents (Paris)], vol. 8, Société Mathématique de France, Paris, 2011. Séminaire de Géométrie Algébrique du Bois Marie 1962–64. [Algebraic Geometry Seminar of Bois Marie 1962–64], A seminar directed by M. Demazure and A. Grothendieck with the collaboration of M. Artin, J.-E. Bertin, P. Gabriel, M. Raynaud and J-P. Serre, Revised and annotated edition of the 1970 French original. MR2867622
 - [Gro60] A. Grothendieck, Éléments de géométrie algébrique. I. Le langage des schémas, Inst. Hautes Études Sci. Publ. Math. 4 (1960), 228. MR217083
 - [Gro61a] _____, Éléments de géométrie algébrique. II. Étude globale élémentaire de quelques classes de morphismes, Inst. Hautes Études Sci. Publ. Math. 8 (1961), 222. MR217084
 - [Gro61b] _____, Éléments de géométrie algébrique. III. Étude cohomologique des faisceaux cohérents. I, 1961. MR217085
 - [Gro64] _____, Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. I, Inst. Hautes Études Sci. Publ. Math. 20 (1964), 259. MR173675
 - [Gro65a] _____, Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. II, Inst. Hautes Études Sci. Publ. Math. **24** (1965), 231. MR199181
 - [Gro65b] _____, Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. III, Inst. Hautes Études Sci. Publ. Math. 24 (1965), 231. MR199181
 - [Gro67] _____, Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas IV, 1967. MR238860
 - [GW20] Ulrich Görtz and Torsten Wedhorn, Algebraic geometry I. Schemes—with examples and exercises, Springer Studium Mathematik—Master, Springer Spektrum, Wiesbaden, 2020. Second edition [of 2675155]. MR4225278
 - [Har77] Robin Hartshorne, Algebraic geometry, Graduate Texts in Mathematics, No. 52, Springer-Verlag, New York-Heidelberg, 1977. MR0463157

- [He13] Xuhua He, Normality and Cohen-Macaulayness of local models of Shimura varieties, Duke Math. J. 162 (2013), no. 13, 2509–2523. MR3127807
- [KKMSD73] G. Kempf, Finn Faye Knudsen, D. Mumford, and B. Saint-Donat, *Toroidal embeddings. I*, Lecture Notes in Mathematics, Vol. 339, Springer-Verlag, Berlin-New York, 1973. MR0335518
 - [LMB00] Gérard Laumon and Laurent Moret-Bailly, Champs algébriques, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 39, Springer-Verlag, Berlin, 2000. MR1771927
 - [Lun96] D. Luna, Toute variété magnifique est sphérique, Transform. Groups 1 (1996), no. 3, 249–258. MR1417712
 - [Lus04] G. Lusztig, Parabolic character sheaves. II, Mosc. Math. J. 4 (2004), no. 4, 869–896, 981. MR2124170
 - [MFK94] D. Mumford, J. Fogarty, and F. Kirwan, Geometric invariant theory, Third, Ergebnisse der Mathematik und ihrer Grenzgebiete (2) [Results in Mathematics and Related Areas (2)], vol. 34, Springer-Verlag, Berlin, 1994. MR1304906
 - [PRS13] Georgios Pappas, Michael Rapoport, and Brian Smithling, Local models of Shimura varieties, I. Geometry and combinatorics, Handbook of moduli. Vol. III, 2013, pp. 135–217. MR3135437
 - [Rom13] Matthieu Romagny, Models of curves, Arithmetic and geometry around Galois theory, 2013, pp. 149–170. MR3408164
 - [SGA3II] Schémas en groupes. II: Groupes de type multiplicatif, et structure des schémas en groupes généraux, Lecture Notes in Mathematics, Vol. 152, Springer-Verlag, Berlin-New York. Séminaire de Géométrie Algébrique du Bois Marie 1962/64 (SGA 3), Dirigé par M. Demazure et A. Grothendieck. MR0274459
 - [03] Revêtements étales et groupe fondamental (SGA 1), Documents Mathématiques (Paris) [Mathematical Documents (Paris)], vol. 3, Société Mathématique de France, Paris, 2003. Séminaire de géométrie algébrique du Bois Marie 1960–61. [Algebraic Geometry Seminar of Bois Marie 1960-61], Directed by A. Grothendieck, With two papers by M. Raynaud, Updated and annotated reprint of the 1971 original [Lecture Notes in Math., 224, Springer, Berlin; MR0354651 (50 #7129)]. MR2017446
 - [Spr06] Tonny A. Springer, Some results on compactifications of semisimple groups, International Congress of Mathematicians. Vol. II, 2006, pp. 1337–1348. MR2275648
 - [Spr09] T. A. Springer, *Linear algebraic groups*, second, Modern Birkhäuser Classics, Birkhäuser Boston, Inc., Boston, MA, 2009. MR2458469
 - [Sta] The Stacks Project Authors, Stacks Project.
 - [Str87] Elisabetta Strickland, A vanishing theorem for group compactifications, Math. Ann. 277 (1987), no. 1, 165–171. MR884653
 - [Wed18] Torsten Wedhorn, Spherical spaces, Ann. Inst. Fourier (Grenoble) **68** (2018), no. 1, 229–256. MR3795478

UNIVERSITÉ PARIS-SACLAY, LABORATOIRE DE MATHÉMATIQUES D'ORSAY, F-91405, ORSAY, FRANCE *Email address*: shang.li@universite-paris-saclay.fr