# A Discharging Method: Improved Kernels for Edge Triangle Packing and Covering

Zimo Sheng and Mingyu Xiao<sup>[0000-0002-1012-2373]</sup>\*

School of Computer Science and Engineering, University of Electronic Science and Technology of China, Chengdu, China shengzimo2016@gmail.com, myxiao@uestc.edu.cn

**Abstract.** EDGE TRIANGLE PACKING and EDGE TRIANGLE COVERING are dual problems extensively studied in the field of parameterized complexity. Given a graph G and an integer k, EDGE TRIANGLE PACKING seeks to determine whether there exists a set of at least k edge-disjoint triangles in G, while EDGE TRIANGLE COVERING aims to find out whether there exists a set of at most k edges that intersects all triangles in G. Previous research has shown that EDGE TRIANGLE PACKING has a kernel of  $(3 + \epsilon)k$  vertices, while EDGE TRIANGLE COVERING has a kernel of 6kvertices. In this paper, we show that the two problems allow kernels of 3kvertices, improving all previous results. A significant contribution of our work is the utilization of a novel discharging method for analyzing kernel size, which exhibits potential for analyzing other kernel algorithms.

# 1 Introduction

Preprocessing is a fundamental and commonly used step in various algorithms. However, most preprocessing has no theoretical guarantee on the quality. Kernelization, originating from the field of parameterized algorithms [1], now has been found to be an interesting way to analyze the quality of preprocessing. Consequently, kernelization has received extensive attention in both theoretical and practical studies.

Given an instance (I, k) of a problem, a kernelization (or a kernel algorithm) runs in polynomial time and returns an equivalent instance (I', k') of the same problem such that (I, k) is a yes-instance if and only if (I', k') is a yes-instance, where  $k' \leq k$  and  $|I'| \leq g(k)$  for some computable function g only of k. The new instance (I', k') is called a *kernel* and g(k) is the size of the kernel. If  $g(\cdot)$  is a polynomial or linear function, we classify the problem as having a polynomial or linear kernel, respectively.

EDGE TRIANGLE PACKING (ETP), to check the existence of k edge-disjoint triangles in a given graph G is NP-hard even on planar graphs with maximum degree 5 [2]. The optimization version of this problem is APX-hard on general graphs [3]. A general result of [4] leads to a polynomial-time  $(3/2+\epsilon)$  approximation algorithm for any constant  $\epsilon > 0$ . When the graphs are restricted to planar

<sup>\*</sup> Corresponding author

graphs, the result can be improved. A polynomial-time approximation scheme for the vertex-disjoint triangle packing problem on planar graphs was given by [5], which can be extended to ETP on planar graphs. In terms of parameterized complexity, a 4k-vertex kernel and an  $O^*(2^{\frac{9k}{2}\log k + \frac{9k}{2}})$ -time parameterized algorithm for ETP were developed in [6]. Later, the size of the kernel was improved to 3.5k [7]. The current best-known result is  $(3 + \epsilon)k$  [8], where  $\epsilon > 0$  can be any positive constant. On tournaments, there is also a kernel of 3.5k vertices [9].

Another problem considered in this paper is EDGE TRIANGLE COVERING (ETC). ETC is the dual problem of ETP, which is to check whether we can delete at most k edges from a given graph such that the remaining graph has no triangle. ETC is also NP-hard even on planar graphs with maximum degree 7 [10]. In terms of kernelization, a 6k-vertex kernel for ETC was developed [10]. On planar graphs, the result was further improved to  $\frac{114}{2}$  [10].

In this paper, we will deeply study the structural properties of EDGE TRI-ANGLE PACKING and EDGE TRIANGLE COVERING and give some new reduction rules by using a variant of crown decomposition. After that, we will introduce a new technology called the discharging method to analyze the size of problem kernels. Utilizing the new discharging method, we obtain improved kernel sizes of 3k vertices for both ETP and ETC. Notably, our results even surpass the previously best-known kernel size for ETC on planar graphs [10].

# 2 Preliminaries

Let G = (V, E) denote a simple and undirected graph with n = |V| vertices and m = |E| edges. A vertex is a *neighbor* of another vertex if there is an edge between them. The set of neighbors of a vertex v is denoted by N(v), and the degree of v is defined as d(v) = |N(v)|. For a vertex subset  $V' \subseteq V$ , we let  $N(V') = \bigcup_{v \in V'} N(v) \setminus V'$  and  $N[V'] = N(V') \cup V'$ . The subgraph induced by a vertex subset  $V' \subseteq V$  is denoted by G[V'] and the subgraph spanned by an edge set  $E' \subseteq E$  is denoted by G[E']. The vertex set and edge set of a graph H are denoted by V(H) and E(H), respectively.

A complete graph on 3 vertices is called a *triangle*. We will use vuw to denote the triangle formed by vertices v, u, and w. If there is a triangle vuw in G, we say that vertex v spans edge uw. An edge triangle packing in a graph is a set of triangles such that every two triangles in it have no common edge. The EDGE TRIANGLE PACKING problem (ETP) is defined below.

Edge Triangle Packing (ETP)	Parameter: k
<b>Input</b> : An undirected graph $G = (V,$	E), and an integer $k$ .
Question: Does there exist an edge	triangle packing of size at least $k$ in $G$ ?

An edge covers a triangle if it is contained in the triangle. An edge triangle covering in a graph is a set of edges S such that there is no triangle after deleting S from G. The EDGE TRIANGLE COVERING problem (ETC) is defined below.

Edge Triangle Covering (ETC) <b>Parameter:</b> $k$	
<b>Input</b> : An undirected graph $G = (V, E)$ , and an integer k.	
Question: Does there exist an edge triangle covering of size at most $k$ in $G$ ?	



Fig. 1. An illustration for the fat-head crown decomposition

## 3 Fat-Head Crown Decomposition

One important technique in this paper is based on a variant of the crown decomposition. Crown decomposition is a powerful technique for the famous VER-TEX COVER problem and it has been extended to solve several related problems [11,12,13,14]. Specifically, we employ a specific variant called the fat-head crown decomposition to tackle (ETP) [8]. This variant of the crown decomposition will also be applied in our algorithms for both ETP and ETC. To provide a comprehensive understanding, let us begin by introducing the definition of the fat-head crown decomposition.

A fat-head crown decomposition of a graph G = (V, E) is a triple (C, H, X) such that C and X form a partition of V and  $H \subseteq E$  is a subset of edges satisfying the following properties:

1. C is an independent set.

2. *H* is the set of edges spanned by at least one vertex in C.

3. No vertex in C is adjacent to a vertex in  $X \setminus V(H)$ .

4. There is an edge-disjoint triangle packing P of size |P| = |H| such that each triangle in P contains exactly one vertex in C and exactly one edge in H. The packing P is also called the *witness packing* of the fat-head crown decomposition.

An illustration of the fat-head crown decomposition is shown in Figure 1. To determine the existence of fat-head crown decompositions in a given graph structure, we present three lemmas.

**Lemma 1.** (Lemma 2 in [8]) Let G = (V, E) be a graph such that each edge and each vertex is contained in at least one triangle. Given a non-empty independent set  $I \subseteq V$  such that |I| > |S(I)|, where S(I) is the set of edges spanned by at least one vertex in I. A fat-head crown decomposition (C, H, X) of G with  $C \subseteq I$ and  $H \subseteq S(I)$  together with a witness packing P of size |P| = |H| > 0 can be found in polynomial time.

**Lemma 2.** Given a graph G = (V, E), a vertex set  $A \subseteq V$ , and an edge set  $B \subseteq E$ , where  $A \cap V(B) = \emptyset$ . There is a polynomial-time algorithm that checks whether there is a fat-head crown decomposition (C, H, X) such that  $\emptyset \neq C \subseteq A$  and  $H \subseteq B$  and outputs one if yes.

Proof. We begin by demonstrating that if there exists an edge  $uv \in E$  between two vertices u and  $v \in A$ , neither u nor v can belong to C in the fat-head crown decomposition. Suppose, without loss of generality, that u is in C. Since C is an independent set, v cannot be in C. Moreover, v cannot be in  $X \setminus V(H)$  since no vertex in C is adjacent to a vertex in  $X \setminus V(H)$ . Additionally, v cannot be in V(H) since  $A \cap V(B) = \emptyset$ . Thus, this scenario is impossible. Therefore, neither u nor v can be in C for any fat-head crown decomposition (C, H, X) with  $C \subseteq A$ and  $H \subseteq B$ . Consequently, we can exclude vertices in A that are adjacent to other vertices in A when computing the decomposition.

To facilitate the computation, we construct an auxiliary bipartite graph G' = (A', B', E'), where each vertex in A' corresponds to a vertex in A that is not adjacent to any other vertex in A, each vertex in B' corresponds to an edge in B, and an edge exists between  $w \in A'$  and  $e \in B'$  if and only if w spans e in G. Based on the construction of G', we deduce that a fat-head crown decomposition (C, H, X) exists in G if and only if there exist  $C' \subseteq A'$  and  $H' \subseteq B'$  in G' such that there is a matching of size |H'| in the induced bipartite graph  $G'[C' \cup H']$ .

The existence of the matching can be determined in  $O(mn^{1.5})$  time using the expansion lemma [15]. Thus, the algorithm can verify the existence of a fathead crown decomposition and output one if it exists, all within polynomial time complexity.

**Lemma 3.** If there is a fat-head crown decomposition (C, H, X) in G, then G has an edge-disjoint triangle packing (resp., edge triangle covering) of size k if and only if the graph G' has an edge-disjoint triangle packing (resp., edge triangle covering) of size k - |H|, where G' is the graph obtained from G by deleting vertex set C and deleting edge set H.

*Proof.* For a fat-head crown decomposition (C, H, X), there is witness packing P of size |P| = |H|.

We first show that there is an edge-disjoint triangle packing D of size k in Gif and only if there is an edge-disjoint triangle packing D' of size k - |H| in G'. We only prove the sufficient condition and the other direction is obvious since  $D' \cup P$  is a valid edge-disjoint triangle packing in G. Let  $D^* \subseteq D$  be the set of triangles containing either a vertex in C or an edge in H. For the former case, triangle T still contains one edge in H since any triangle containing a vertex in C must contain one edge in H. Thus, each triangle in  $D^*$  contains at least one edge in H, which means that  $|D^*| \leq |H|$ . Therefore,  $D - D^*$  is an edge-disjoint triangle packing of size at least k - |H| in graph G'.

Next, we prove that there is an edge triangle covering M of size k in G if and only if there is an edge triangle covering M' of size k-|H| in G'. For the necessary condition. We prove that  $M' \cup H$  is a valid edge triangle covering in G. Consider an arbitrary triangle T' in G. If  $V(T') \cap V(C) = \emptyset$  and  $E(T') \cap E(H) = \emptyset$ , then T' is covered by M' since M' is an edge triangle covering for G'. If  $V(T') \cap C \neq \emptyset$ , then T' is covered by H since vertices in C can only span edges in H by the definition of the fat-head crown decomposition. If  $E(T') \cap E(H) \neq \emptyset$ , then T'is covered by H. For any case, triangle T' is covered by at least one edge in  $M' \cup H$ . For the sufficient condition. Let  $M^* = M \cap E(P)$ . We know that  $|M^*| \ge |P| = |H|$  since P is an edge-disjoint triangle packing of size |H|. Thus,  $M - M^*$  is an edge triangle covering of size at most k - |H| in G'.  $\Box$ 

## 4 The Algorithms

In this section, we present our kernelization algorithms for the EDGE TRIAN-GLE PACKING (ETP) and EDGE TRIANGLE COVERING (ETC) problems. Our algorithms involve a set of reduction rules that are applied iteratively until no further reduction is possible. Each reduction rule is applied under the assumption that all previous reduction rules have already been applied and cannot be further applied to the current instance. A reduction rule is *correct* if the original instance (G, k) is a yes-instance if and only if the resulting instance (G', k') after applying the reduction rule is a yes-instance.

We have one algorithm for ETP and ETC, respectively. The two algorithms are similar. We will mainly describe the algorithm for ETP and introduce the difference for ETC. In total, we have nine reduction rules. The first four rules are simple rules to handle some special structures, while the remaining five rules are based on a triangle packing. Especially, the last rule will use the fat-head crown decomposition. We will show that the algorithms run in polynomial time.

#### 4.1 Simple Rules

**Reduction Rule 1** For ETP, if  $k \leq 0$ , then return 'yes' to indicate that the instance is a yes-instance; if k > 0 and the graph is empty, then return 'no' to indicate that the instance is a no-instance.

For ETC, if  $k \ge 0$  and the graph is empty, then return 'yes' to indicate that the instance is a yes-instance; if k < 0, then return 'no' to indicate that the instance is a no-instance.

**Reduction Rule 2** If there is a vertex or an edge not appearing in any triangle, then delete it from the graph.

**Reduction Rule 3** If there are 4 vertices  $u, v, w, x \in V$  inducing a complete graph (i.e., there are 6 edges  $uv, uw, ux, vw, vx, wx \in E$ ) such that none of the 6 edges is in a triangle except uwv, uvx, uwx, and vwx, then

- For ETP, delete the 6 edges uv, uw, ux, vw, vx and wx and let k = k 1;
- For ETC, delete the 6 edges uv, uw, ux, vw, vx and wx and let k = k 2.

The correctness of Reduction Rule 3 is based on the following observation. For ETP, any edge triangle packing can have at most one triangle containing some edge from these 6 edges and we can simply take one triangle from this local structure. For ETC, any edge triangle covering must contain at least two edges from these 6 edges and after deleting vu and wx, none of uw, ux, vw, and vx is contained in a triangle anymore.



Fig. 2. An Illustration for Reduction Rule 4

**Reduction Rule 4** If there is a vertex  $v \in V$  such that all edges incident to v can be partitioned into two parts  $E_1$  and  $E_2$  such no triangle in G contains an edge in  $E_1$  and an edge in  $E_2$ , then split v into two vertices v' and v'' such that all edges in  $E_1$  are incident on v' and all edges in  $E_2$  are incident on v''.

An illustration of Reduction Rule 4 is shown in Figure 2. This reduction rule will increase the number of vertices in the graph. However, this operation will simplify the graph structure and our analysis.

Lemma 4. Reduction Rule 4 is correct and can be executed in polynomial time.

*Proof.* First, we consider the correctness. Let G' = (V', E') be the graph after applying Reduction Rule 4 on a vertex v. We can establish a one-to-one mapping between the edges in E and the edges in E' by considering the vertices v' and  $v'' \in V'$  as  $v \in V$ . Three edges in E form a triangle in G if and only if the three corresponding edges in E' form a triangle in G' since there is no triangle in Gcontains an edge in  $E_1$  and an edge in  $E_2$ . Thus, an edge triangle packing of size k (resp., an edge triangle covering of size k) in G is also an edge triangle packing of size k (resp., an edge triangle covering of size k) in G'. This implies that Reduction Rule 4 is correct for both ETP and ETC.

We give a simple greedy algorithm to find the edge sets  $E_1$  and  $E_2$  for a given vertex v. Initially, let  $E_1$  contain an arbitrary edge e incident on v. We iteratively perform the following steps until no further updates occur: if there is a triangle containing an edge in  $E_1$  and an edge e' incident on v but not in  $E_1$ , then add edge e' to  $E_1$ . It is easy to see that all edges in  $E_1$  must be in the same part to satisfy the requirement. If  $E_1 \neq E$ , then we can split E to two parts  $E_1$  and  $E_2 = E \setminus E_1$ . Otherwise, the edges incident on v cannot be split.  $\Box$ 

#### 4.2 Adjustments Based on a Triangle Packing

After applying the first four rules, our algorithms will find a maximal edgedisjoint triangle packing S by using an arbitrary greedy method. This can be done easily in polynomial time. The following rules are based on the packing S. From now on, we let  $F = V \setminus V(S)$  denote the set of vertices not appearing in S and  $R = E \setminus E(S)$  denote the set of edges not appearing in S. We begin with the following trivial rule. **Reduction Rule 5** If |S| > k, for ETP, return 'yes' to indicate that the instance is a yes-instance; and for ETC, return 'no' to indicate that the instance is a no-instance.

The following three rules just update the packing S by replacing some triangles in it and do not change the graph. Illustrations of the three rules are shown in Figure 3.



Fig. 3. Illustrations of Reduction Rules 6-8

**Reduction Rule 6** If there is a triangle  $v_1v_2v_3 \in S$  such that there are at least two edge-disjoint triangles in the spanned graph  $G[R \cup \{v_1v_2, v_1v_3, v_2v_3\}]$ , then replace  $v_1v_2v_3$  with these triangles in S to increase the size of S by at least 1.

**Reduction Rule 7** If there are two edge-disjoint triangles  $v_1v_2v_3$  and  $v_4v_5v_6 \in S$  such that there are at least three edge-disjoint triangles in the spanned graph  $G[R \cup \{v_1v_2, v_1v_3, v_2v_3, v_4v_5, v_4v_6, v_5v_6\}]$ , then replace  $v_1v_2v_3$  and  $v_4v_5v_6$  with these triangles in S to increase the size of S by at least 1.

**Reduction Rule 8** If there are two edge-disjoint triangles  $v_1v_2v_3$  and  $v_4v_5v_6 \in S$  such that there are two edge-disjoint triangles  $v'_1v'_2v'_3$  and  $v'_4v'_5v'_6$  in the induced graph  $G[F \cup \{v_1, v_2, v_3, v_4, v_5, v_6\}]$  such that  $|\{v'_1, v'_2, v'_3\} \cup \{v'_4, v'_5, v'_6\}| > |\{v_1, v_2, v_3\} \cup \{v_4, v_5, v_6\}|$ , then replace triangles  $v_1v_2v_3$  and  $v_4v_5v_6$  with triangles  $v'_1v'_2v'_3$  and  $v'_4v'_5v'_6$  in S to increase the number of vertices appearing in S by at least 1.

Note that an application of Reduction Rules 6-8 will not change the structure of the graph. Thus, the first four reduction rules will not be applied after executing Reduction Rules 6-8.

#### 4.3 A Reduction Based on Fat-head Crown Decomposition

After Reduction Rule 8, we obtain the current triangle packing S. An edge in E(S) is called a *labeled edge* if it is spanned by at least one vertex in F. We let L denote the set of labeled edges.

We can find a fat-head crown decomposition (C, H, X) with  $C \subseteq V \setminus V(L)$ and  $H \subseteq L$  in polynomial time if it exists by Lemma 2. Moreover, we will apply the following reduction rule to reduce the graph, the correctness of which is based on Lemma 3.

**Reduction Rule 9** Use the algorithm in Lemma 2 to check whether there is a fat-head crown decomposition (C, H, X) such that  $\emptyset \neq C \subseteq V \setminus V(L)$  and  $H \subseteq L$ . If yes, then delete vertex set C and edge set H, and let k = k - |H|.

An instance is called *reduced* if none of the nine reduction rules can be applied to it. The corresponding graph is also called a reduced graph.

**Lemma 5.** For any input instance, the kernelization algorithms run in polynomial time to output a reduced instance.

*Proof.* Lemma 2 demonstrates that one execution of Reduction Rule 9 can be performed within polynomial time, while Lemma 4 shows that one execution of Reduction Rule 4 can also be accomplished in polynomial time. Additionally, each of the remaining reduction rules can be easily executed within polynomial time. Next, we only need to show that each reduction rule can be executed for at most polynomial times to complete our proof. Reduction Rules 1 and 5 can be applied for at most 1 time. Reduction Rule 2 can be applied for at most mtimes since it will delete at least one edge each time. Reduction Rule 4 can be applied for at most 2m times since it will increase one vertex each time and there are at most 2m vertices for the worst case. Reduction Rules 3 and 9 can be applied for at most k times since each of them will decrease k by at least 1. We consider Reduction Rules 6-8. For a fixed graph G = (V, E), Reduction Rules 6 and 7 can be continuously executed for at most k times and Reduction Rule 8 can be continuously executed for at most 3k times. Since Reduction Rule 9 can be applied for at most k times to change the graph, we know that Reduction Rules 6-8 can be applied for at most  $O(k^2)$  times. Thus, the algorithms run in polynomial times. 

# 5 Analysis Based on Discharging

Next, we use a discharging method to analyze the size of a reduced instance. Note that there is no significant difference between ETC and ETP in the analysis. We partition the graph into two parts: one part is the edge-disjoint triangle packing S after applying all the reductions; the other part is the set F of vertices not appearing in S. Before proceeding with the analysis, we will establish some properties that will be utilized.



Fig. 4. Three cases in Lemma 6

**Lemma 6.** Consider a reduced graph G = (V, E) with triangle packing S. For any triangle  $uvw \in S$ , at most one of  $\{uv, vw, uw\}$  is a labeled edge.

*Proof.* Assume to the contrary that there are two edges, say uv and vw are spanned by vertices in F. We show some contradiction.

If edges uv and vw are spanned by two different vertices  $x, x' \in F$  respectively, then Reduction Rule 6 could be applied (Case 1 in Figure 4). Therefore, edges uv and vw are spanned by the same vertex  $x \in F$ . Since Reduction Rule 3 is not applied on the four vertices  $\{u, v, w, x\}$ , we know that at least one edge in  $\{uw, uw, ux, vw, vx, wx\}$  is contained in a triangle other than uwv, uvx, uwx, and vwx. Due to symmetry, we only need to consider two edges vw and xw.

Assume that edge vw is contained in a triangle vwy, where  $y \notin \{u, x\}$ . If none of  $\{yv, yw\}$  appears in E(S), then Reduction Rule 6 could be applied to replace uvw with two triangles xvu and yvw in S. If at least one edge in  $\{yv, yw\}$ is contained in E(S), without loss of generality, assume  $yw \in E(S)$  and there is a triangle  $ywz \in S$ . For this case, Reduction Rule 8 could be applied to replace vuw and ywz with xvu and ywz (Case 2 in Figure 4).

Assume that edge xw is contained in a triangle xwy, where  $y \notin \{u, v\}$ . By the maximality of S, we know that at least one of  $\{xy, yw\}$  must appear in E(S). However, edge xy can not appear in E(S) since  $x \in F$ . We know that  $wy \in E(S)$ and there is a triangle  $wyz \in S$ . For this case, Reduction Rule 8 could be applied to replace vuw and ywz with xvu and wyz (Case 3 in Figure 4).

In any of these cases, we can find a contradiction to the fact that the graph is reduced.  $\hfill \Box$ 

A triangle  $uvw \in S$  is good if it contains a labeled edge and bad otherwise. By Lemma 6, we know that there is exactly one labeled edge in each good triangle. We let G' be the graph obtained by deleting the set L of labeled edges from G. Consider a good triangle uvw with labeled edge uv. If the two edges vw and wu are not in any triangle in G', we call the triangle *excellent*. Otherwise, we call the triangle *pretty-good*. We let  $S_1$  denote the set of excellent triangles,  $S_2$ denote the set of pretty-good triangles, and  $S_3$  denotes the set of bad triangles in S. The number of triangles in  $S_1$ ,  $S_2$  and  $S_3$  are denoted by  $k_1, k_2$ , and  $k_3$ , respectively. Let  $V_1 = V(S_1) \setminus (V(L) \cup V(S_2) \cup V(S_3))$  and  $V_2 = V(S_2) \setminus V(L)$ . See Figure 5 for an illustration of these concepts.

9

10 Zimo Sheng, Mingyu Xiao



Fig. 5. An illustration for triangles and vertices in G

#### 5.1 The analysis

The discharging method stands as a renowned technique in graph theory, finding its most notable application in the proof of the famous Four Color Theorem. In this section, we will use the discharging method to analyze the number of vertices present in  $S_1$ ,  $S_2$ ,  $S_3$ , and F. The idea of the method is as follows.

First, we initially assign some integer values to vertices, edges, and triangles in S. The total value assigned is at most 3k. Subsequently, we perform steps to update the values, where certain values on vertices, edges, and triangles are transformed into other vertices, edges, and triangles. In these steps, we never change the structure of the graph and the total value in the graph. After performing these transformations, we demonstrate that each vertex in the graph has a value of at least 1. Consequently, we conclude that the number of vertices in the graph is at most 3k.

**Initialization:** Assign value 3 to each edge in L and each triangle in  $S_3$ . Edges not in L, vertices, and triangles in  $S_1 \cup S_2$  are assigned a value of 0.

By Lemma 6, we know that each of excellent and pretty-good triangles contains exactly one labeled edge in L and each bad triangle in  $S_3$  contains no labeled edge. Thus, the total value in the graph is  $3k_1 + 3k_2 + 3k_3 \leq 3k$ .

**Step 1:** For each labeled edge in L, transform a value of 1 to each of its two endpoints; for each triangle in  $S_3$ , transform a value of 1 to each of its three vertices.

Figure 6 illustrates the transformation process of Step 1. After Step 1, each labeled edge has a value of 1, and all triangles have values of 0. Note that some vertices may have a value of 2 or more, as they may serve as endpoints of multiple labeled edges and can also be vertices in  $V(S_3)$ . However, vertices in  $F \cup V_1 \cup V_2$  still retain a value of 0.

A triangle component is a connected component in the graph H = (V(S), E(S)). For a vertex  $v \in V(S)$ , we let C(v) denote the set of vertices in the triangle component which contain v.

**Step 2:** For each triangle component in G, we iteratively transform a value of 1 from a vertex with a value of at least 2 to a vertex with a value of 0 in the same triangle component, where vertices in  $V_1$  have a higher priority to get the value.



Fig. 6. An illustration for Step 1

**Lemma 7.** After Step 2, each triangle component has at most one vertex with a value of 0. Moreover,

(i) For any triangle component containing a triangle in  $S_3$ , each vertex in the triangle component has a value of at least 1;

(ii) For any triangle component containing at least one triangle in  $S_2$ , if there is a vertex with a value of 0 in the triangle component, then the vertex must be a vertex in  $V_2$ .

Proof. Let Q be a triangle component with x triangles. Since Q is connected, it contains at most 2x + 1 vertices. Assume that among the x triangles, there are  $x_1$  triangles in  $S_1 \cup S_2$  and  $x_2$  triangles in  $S_3$ , where  $x_1 + x_2 = x$ . By the definition, we know that each triangle in  $S_1 \cup S_2$  contains a distinct labeled edge. According to the initialization of the assignment, we know that the total value is  $2x_1 + 3x_2 = 2x + x_2$ . It always holds that  $2x + 1 \leq (2x + x_2) + 1$ , and  $2x + 1 \leq 2x + x_2$  when  $x_2 \geq 1$ . Thus, Q has at most one vertex with a value of 0. When Q contains some triangles from  $S_3$ , i.e.,  $x_2 \geq 1$ , all vertices in Q will get a value of at least 1. The statement (ii) holds because vertices in  $V_1$  have a higher priority to receive the value in Step 2.

After Step 2, only vertices in F, some vertices in  $V_1$ , and some vertices in  $V_2$  have values of 0. We use the following lemma to transform some values to vertices in  $V_2$  with a value of 0.

**Lemma 8.** Consider two triangles vuw and  $vxy \in S_2$  sharing a common vertex v, where uv and  $vx \in L$ . If there is an edge  $wy \in E$ , then uv and vx are spanned by exactly one vertex in  $F \cup V_1 \setminus C(v)$ .

*Proof.* Since vu and vx are labeled edges in L, we know that each of vu and vx is spanned by at least one vertex in F. Assume to the contrary that there are two different vertices  $q_1$  and  $q_2 \in F \cup V_1 \setminus C(v)$  that span edges uv and vx, respectively. We show some contradiction. First, none edge in  $\{q_1u, q_1v, q_2v, q_2x\}$  belongs to E(S) since  $q_1$  and  $q_2$  are not part of C(v). Thus, Reduction Rule 7 could be applied to replace triangles uvw and vxy with triangles  $q_1uv, q_2vx$ , and vwy, a contradiction to the factor that the graph is reduced. See Figure 7 for an illustration of the proof.

**Step 3:** If there are two triangles vuw and  $vxy \in S_2$  sharing a common vertex v such that uv and vx are two labeled edges in L and there is an edge



Fig. 7. An illustration for Lemma 8



Fig. 8. An illustration for Step 3, where the number in parentheses next to each vertex represents the value of that vertex

 $wy \in E$ , we transform a value of 1 from edge vx to the unique vertex  $q_1 \in F$  spanning vx and transform a value of 1 from edge uv to the vertex with a value of 0 in C(v) if this vertex exists.

See Figure 8 for an illustration of Step 3. We have the following property.

#### **Lemma 9.** Every vertex in $V_2$ has a value of at least 1 after Step 3.

*Proof.* Assume to the contrary there is a vertex  $w \in V_2$  with a value of 0. We know that all vertices in  $C(w) \setminus \{w\}$  have a value of at least 1 by Lemma 7. Let  $wuv \in S_2$  be the triangle containing w, where uv is the labeled edge spanning by a vertex  $q \in F$ . As shown in Figure 9. At least one of uw and vw is in a triangle in graph G - L by the definition of  $S_2$ . Without loss of generality, we assume that uw is contained in a triangle uwx, where  $x \neq v$ . At least one of ux and wx is contained in a triangle in S otherwise Reduction Rule 6 could be applied.

**Case 1**: Edge ux is contained in a triangle  $uxy \in S$ . See Case 1 in Figure 9. The triangle uxy is not in  $S_3$  otherwise there is a contradiction that w would have a value of at least 1 by Lemma 7. Thus, triangle uxy must be in  $S_1 \cup S_2$  and there is exactly one of ux, uy, and xy is a labeled edge. Edge ux would not be a labeled edge since triangle uwx is contained in G - L. If xy is the labeled edge that is spanned by a vertex  $q' \in F$ , then Reduction Rule 8 could be applied to replace triangles uvw and uxy with triangles uvw and xyq', a contradiction to the factor that the graph is reduced. If uy is the labeled edge, then triangle  $uxy \in S_2$  since ux is contained in a triangle  $uxw \in G - L$ . For this case, the



Fig. 9. An illustration for Lemma 9

vertex w would have a value of at least 1 by Lemma 7. We can always find a contradiction.

**Case 2**: Edge wx is contained in a triangle  $wxz \in S$ . See Case 2 in Figure 9. If  $z \neq q$ , then Reduction Rule 8 could be applied to replace triangles uvw and wxz with triangles wxz and qvu, a contradiction to the factor that the graph is reduced. If z = q, then at least two edges in triangle vuw are spanned by vertices in F, a contradiction to Lemma 6.

In either case, a contradiction is reached, which implies that the assumption of a vertex  $w \in V_2$  having a value of 0 is incorrect. Therefore, every vertex in  $V_2$  has a value of at least 1 after Step 3.

After Step 3, all vertices with a value of 0 are in either F or  $V_1$ . We let  $V'_1$  denote the set of vertices with a value of 0 in  $V_1$ , F' denote the set of vertices with a value of 0 in F, and L' denote the set of edges with a value of 1 in L after Step 3. We give more properties.

## **Lemma 10.** Set $F' \cup V'_1$ is an independent set.

*Proof.* We prove that  $F \cup V_1$  is an independent set, which implies  $F' \cup V'_1$  is an independent set since  $F' \cup V'_1 \subseteq F \cup V_1$ . Assume to the contrary that there is an edge uv between two vertices in  $F \cup V_1$ . There is at least one triangle  $uvw \in G$  containing uv since Reduction Rule 2 has been applied. At least one of uv, vw, and uw must be in E(S) by the maximality of S.

If  $uv \in E(S)$ , we let uvx be the triangle in S containing uv. First, we know that u and  $v \in V_1$  since  $F \cap V(S) = \emptyset$ . By the definition of  $V_1$ , we get that none of u and v is contained in a triangle in  $S_3$  and none of u and v is an endpoint of a labeled edge. Thus, triangle uvx is not a triangle in  $S_3$  and it does not contain any labeled edge and then it is not a triangle in  $S_1 \cup S_2$ , which implies triangle uvx is not in S, a contradiction.

Otherwise, one of uw and vw, say uw, is contained in E(S). Let uwy be the triangle in S containing uw. We also have that  $u \in V_1$  since  $F \cap V(S) = \emptyset$ . By the definition of  $V_1$ , we know that u is not a vertex in a triangle in  $S_3$ , and then uwy is not a triangle in  $S_3$ . Thus, triangle uwy can only be in  $S_1 \cup S_2$ . Note that none of uv, vw and uw can be a labeled edge since u and  $v \in F \cup V_1$ . Thus, edge uw is still in a triangle uvw in G - L, and then uwy can not be a triangle in  $S_1$ .

14 Zimo Sheng, Mingyu Xiao



Fig. 10. Some cases in the proof of Lemma 12

However, triangle uvw can not be a triangle in  $S_2$  too since u is a vertex in  $V_1$ . We also get a contradiction that triangle uvw is not in S.

Hence, we have shown that no edge exists between any two vertices in  $F' \cup V'_1$ , which proves that  $F' \cup V'_1$  forms an independent set.

## **Lemma 11.** Vertices in F' only span edges in L'.

*Proof.* Assume to the contrary that there is a vertex  $z \in F'$  that spans an edge  $e \notin L'$ . Note that  $z \in F$  can only span edges in L. Thus,  $e \in L \setminus L'$ , i.e., the value of e is 0 now. Only Step 3 will create labeled edges with a value of 0. Therefore, edge e appears as a labeled edge in the graph structure described in Step 3. By Lemma 8, we know that z is the unique vertex in F spanning e and the value of z is at least 1 after Step 3, a contradiction to  $z \in F'$ . Thus, vertices F' can only span edges in L'.

**Lemma 12.** Vertices in  $V'_1$  only span edges in L'.

*Proof.* Let  $v' \in V'_1$  and Y be the triangle component containing v'. First, we show that  $N(v') \subseteq V(Y)$ .

We let  $E_1$  denote the set of edges with one endpoint being v' and the other endpoint being in  $N(v') \cap V(Y)$  and  $E_2$  denote the set of edges with one endpoint being v' and the other endpoint being in  $N(v') \setminus V(Y)$ . Assume that there are edges  $x'v' \in E_1$  and  $u'v' \in E_2$  forming a triangle x'v'u' in G and we show some contradiction. If  $x'v' \notin E(S)$ , then none of x'v', u'v', and x'u' is contained in E(S) since u' is not in the triangle component containing v'. which is a contradiction to the maximality of S. Otherwise  $x'v' \in E(S)$  and we let triangle  $x'v'y' \in S$  be the triangle containing x'v'. Since the value of  $v_1$  is 0, by Lemma 7, we know that Y could not contain any triangle in  $S_2 \cup S_3$  and then  $x'v'y' \in S_1$ . We know that x'y' is the labeled edge since  $v' \in V'_1$ . Let  $q' \in F$  be the vertex spanning x'y'. We can see that  $q' \neq u'$  otherwise all the three edges in triangle x'y'y' are labeled edges spanned by a vertex  $q' \in F$ , a contradiction to Lemma 6. However, for this case, Reduction Rule 6 could be applied to replace triangle x'v'y' with triangles q'x'y' and u'x'v' in S, a contradiction to the factor that the graph is reduced. Therefore, no triangle in G contains edges both from  $E_1$  and  $E_2$ . If both of  $E_1$  and  $E_2$  are not empty, then the condition of Reduction Rule 4 would hold, a contradiction. Since  $E_1$  can not be empty, we know that  $E_2$  is empty. Thus, we have  $N(v') \subseteq V(Y)$ . See Figure 10(a) for an illustration of the proof.

Assuming that v' spans an edge ab, we can conclude that both a and b are in V(Y) since N(v') is a subset of V(Y). Our goal is to prove that ab belongs to L. If  $ab \in E(S) \setminus L$ , then ab is contained in a triangle  $abd \in S_1$  since Y contains only triangles in  $S_1$ . We have  $d \neq v'$ , otherwise the triangle  $abv' \in S_1$  would have no labeled edge, a contradiction. The triangle  $abd \in S_1$  contains an edge ab which will be in a triangle abv' in G - L, which contradicts the definition of  $S_1$  (See Figure 10(b)). If  $ab \notin E(S)$ , then at least one of bv' and av', say bv', is contained in E(S) by the maximality of S. Let  $bv'c \in S_1$  be the triangle containing bv' where  $c \neq a$  since  $ab \notin E(S)$ . The triangle  $bcv' \in S_1$  contains an edge bv' which will be in a triangle abv' in G - L, which contradicts the definition of  $S_1$  (See Figure 10(c)). Thus, it holds that  $ab \in L$ . Any vertex  $v' \in V'_1$  only span edges  $ab \in L$ , where both a and b are in V(Y). Then, edge ab can only be a labeled edge in a triangle in  $S_1$ . Since all labeled edges in triangles in  $S_1$  are in L', we know that vertices in  $V'_1$  can only span edges in L'.

**Lemma 13.** After Step 3, it holds that  $|F' \cup V'_1| \le |L'|$ .

*Proof.* By Lemma 10, 11, and 12, we know that  $F' \cup V'_1$  is an independent set and any vertex  $v' \in F' \cup V'_1$  only span edges in L'. If  $|F' \cup V'_1| > |L'|$ , then by Lemma 1 there is a fat-head crown decomposition (C, H, X) of G with  $C \subseteq F' \cup V'_1$  and  $H \subseteq L'$ . Moreover, the fat-head crown decomposition can be detected by Lemma 2 and will be handled by Reduction Rule 9 since  $F' \cup V'_1 \subseteq F \cup V_1 \subseteq V \setminus V(L)$ and  $L' \subseteq L$ . Thus, we know the lemma holds.  $\Box$ 

**Step 4:** We transform value from edges in L' to vertices in  $F' \cup V'_1$  such that each vertex in  $F' \cup V'_1$  gets value at least 1 by Lemma 13.

After Step 4, each vertex in G has a value of at least 1. Since the total value in G is at most 3k, we can conclude that the graph has at most 3k vertices.

**Theorem 1.** EDGE TRIANGLE PACKING and EDGE TRIANGLE COVERING admit a kernel of at most 3k vertices.

#### 6 Conclusion

In this paper, we present simultaneous improvements in the kernel results for both EDGE TRIANGLE PACKING and EDGE TRIANGLE COVERING. Our approach incorporates two key techniques to achieve these enhancements. The first technique involves utilizing fat-head crown decomposition, which enables us to effectively reduce various graph structures. By applying this technique, we can simplify the problem instances. The second technique we introduce is the discharging method, which plays a crucial role in analyzing kernel size. This method is simple and intuitive, and we believe it has the potential to be applied to the analysis of other kernel algorithms.

Acknowledgments. The work is supported by the National Natural Science Foundation of China, under the grants 62372095 and 61972070.

### References

- M. Cygan, F. V. Fomin, L. Kowalik, D. Lokshtanov, D. Marx, M. Pilipczuk, M. Pilipczuk, S. Saurabh, Parameterized Algorithms, Springer, 2015.
- I. Holyer, The NP-completeness of some edge-partition problems, SIAM J. Comput. 10 (4) (1981) 713–717.
- 3. V. Kann, Maximum bounded h-matching is MAX snp-complete, Inf. Process. Lett. 49 (6) (1994) 309–318.
- 4. C. A. J. Hurkens, A. Schrijver, On the size of systems of sets every t of which have an sdr, with an application to the worst-case ratio of heuristics for packing problems, SIAM J. Discret. Math. 2 (1) (1989) 68–72.
- B. S. Baker, Approximation algorithms for np-complete problems on planar graphs, J. ACM 41 (1) (1994) 153–180.
- 6. L. Mathieson, E. Prieto-Rodriguez, P. Shaw, Packing edge disjoint triangles: A parameterized view, in: Parameterized and Exact Computation, First International Workshop, IWPEC 2004, Bergen, Norway, September 14-17, 2004, Proceedings, Vol. 3162 of Lecture Notes in Computer Science, Springer, 2004, pp. 127–137.
- Y. Yang, Towards optimal kernel for edge-disjoint triangle packing, Inf. Process. Lett. 114 (7) (2014) 344–348.
- W. Lin, M. Xiao, A (3+ε)k-vertex kernel for edge-disjoint triangle packing, Inf. Process. Lett. 142 (2019) 20–26.
- H. Yuan, Q. Feng, J. Wang, Improved kernels for triangle packing in tournaments, Sci. China Inf. Sci. 66 (5) (2023).
- D. Brügmann, C. Komusiewicz, H. Moser, On generating triangle-free graphs, Electron. Notes Discret. Math. 32 (2009) 51–58.
- F. K. H. A. Dehne, M. R. Fellows, F. A. Rosamond, P. Shaw, Greedy localization, iterative compression, modeled crown reductions: New FPT techniques, an improved algorithm for set splitting, and a novel 2k kernelization for vertex cover, in: Parameterized and Exact Computation, First International Workshop, IWPEC 2004, Bergen, Norway, September 14-17, 2004, Proceedings, Vol. 3162 of Lecture Notes in Computer Science, Springer, 2004, pp. 271–280.
- M. Xiao, S. Kou, Parameterized algorithms and kernels for almost induced matching, Theor. Comput. Sci. 846 (2020) 103–113.
- M. Xiao, S. Kou, Kernelization and parameterized algorithms for 3-path vertex cover, in: Theory and Applications of Models of Computation - 14th Annual Conference, TAMC 2017, Bern, Switzerland, April 20-22, 2017, Proceedings, Vol. 10185 of Lecture Notes in Computer Science, 2017, pp. 654–668.
- R. Cervený, P. Choudhary, O. Suchý, On kernels for d-path vertex cover, in: 47th International Symposium on Mathematical Foundations of Computer Science, MFCS 2022, August 22-26, 2022, Vienna, Austria, Vol. 241 of LIPIcs, Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2022, pp. 29:1–29:14.
- H. L. Bodlaender, F. V. Fomin, D. Lokshtanov, E. Penninkx, S. Saurabh, D. M. Thilikos, (meta) kernelization, J. ACM 63 (5) (2016) 44:1–44:69.