### ON THE CONVERGENCE OF ADAPTIVE APPROXIMATIONS FOR STOCHASTIC DIFFERENTIAL EQUATIONS

JAMES FOSTER\* AND ANDRAŽ JELINČIČ\*

Abstract. In this paper, we study numerical approximations for stochastic differential equations (SDEs) that use *adaptive* step sizes. In particular, we consider a general setting where decisions to reduce step sizes are allowed to depend on the future trajectory of the underlying Brownian motion. Since these adaptive step sizes may not be previsible, the standard mean squared error analysis cannot be directly applied to show that the numerical method converges to the solution of the SDE. Building upon the pioneering work of Gaines and Lyons, we shall instead use rough path theory to establish convergence for a wide class of adaptive numerical methods on general Stratonovich SDEs (with sufficiently smooth vector fields). To our knowledge, this is the first convergence guarantee that applies to standard solvers, such as the Milstein and Heun methods, with non-previsible step sizes. In our analysis, we require the sequence of adaptive step sizes to be nested and the SDE solver to have unbiased Lévy area terms in its Taylor expansion. We conjecture that for adaptive SDE solvers more generally, convergence is still possible provided the scheme does not introduce "Lévy area bias". We present a simple example where the step size control can skip over previously considered times, resulting in the numerical method converging to an incorrect limit (i.e. not the Stratonovich SDE). Finally, we conclude with an experiment demonstrating the accuracy of Heun's method and a newly introduced Splitting Path-based Runge-Kutta scheme (SPaRK) when used with adaptive step sizes.

Key words. Numerical methods, adaptive step size control, stochastic differential equations, rough path theory

AMS subject classifications. 60H35, 60L90, 65C30

1. Introduction. Stochastic differential equations (SDEs) have seen widespread use in the physical, engineering and social sciences for describing random systems. Examples of SDEs can be found within finance [3, 55, 62], biology [1, 4, 36, 67], physics [13, 42, 50, 63] and, more recently, in data science [8, 9, 12, 19, 33, 39, 46, 47, 65, 69]. However, just as for ordinary differential equations (ODEs), solutions of SDEs are rarely obtainable in closed form, and numerical methods are often required in practice.

In such applications, this is typically done through Monte Carlo simulation, where numerical solutions are computed using independently generated random variables (for example, corresponding to the increments of the underlying Brownian motion). By independently generating multiple numerical solutions, one can then use a Monte Carlo estimator to compute quantities relating to the average behaviour of the SDE. As motivation for this paper, we note that the *status quo* is to propagate numerical solutions of SDEs forwards in time using a fixed step size, whereas so-called "adaptive" step sizes have seen great success for discretizing ODEs (see [26, Section II.4] and [64]). We refer the reader to the book [28] for an accessible introduction to SDE numerics.



FIG. 1. In the Monte Carlo paradigm, Brownian motion is discretized and then mapped to a numerical solution of the SDE. Despite the random fluctuations of underlying Brownian motion, which can occasionally be very large, numerical methods for SDEs typically only use fixed step sizes.

<sup>\*</sup> University of Bath, Department of Mathematical Sciences. Email: { jmf68, aj2382}@bath.ac.uk 1

In this paper, we will study numerical methods with adaptive step size control for general Stratonovich SDEs of the form:

(1.1) 
$$dy_t = f(y_t)dt + g(y_t) \circ dW_t,$$

where  $y = \{y_t\}_{t \in [0,T]}$  is an  $\mathbb{R}^e$ -valued process,  $W = (W^1, \cdots, W^d) = \{W_t\}_{t \in [0,T]}$  is a *d*-dimensional Brownian motion, the vector fields  $f : \mathbb{R}^e \to \mathbb{R}^e$  and  $g = (g_1, \cdots, g_d)$ :  $\mathbb{R}^e \to \mathbb{R}^{e \times d}$  are sufficiently smooth and  $g(y_t) \circ dW_t$  is understood as  $\sum_{i=1}^d g_i(y_t) \circ dW_t^i$ .

In [21], Gaines and Lyons consider adaptive step size controls for SDE simulation where the current "candidate" step size  $h_k$  at time  $t_k$  may be halved depending on the increment of the underlying Brownian motion  $W_{t_k+h_k} - W_{t_k}$  sampled over  $[t_k, t_k + h_k]$ . This approach is intuitive since it can allow the numerical method for (1.1) to adapt to the "large" fluctuations of Brownian motion which, despite having a low probability, can significantly increase the overall approximation error. However, such adaptive step sizes are clearly *non-previsible* and therefore carry theoretical and practical challenges. To address these challenges, Gaines and Lyons made the following key contributions.

• Using Lévy's well-known construction of Brownian motion, a tree structure called the *Brownain tree* was proposed for the generation and storage of Brownian paths. The first level stores increments of Brownian motion over intervals of a fixed size, whereas subsequent levels are dynamically created whenever the Brownian motion is sampled at midpoints within the intervals. Thus, the finest discretization of the Brownian path, which is used to compute a numerical solution of the SDE (1.1), is stored at the lowest level of the tree. We illustrate a Brownian tree in Figure 2.



FIG. 2. If an SDE solver has an adaptive step size control that can only halve step sizes, then it naturally corresponds to a nested sequence of partitions with increments stored in a Brownian tree. Of course, as the mesh of these partitions tends to zero, we would hope the approximation converges.

In (scientific) machine learning, computing the gradient of a loss functions with respect to the parameters of a differential equation can be memory intensive [56]. As a consequence, memory-efficient extensions of the Brownian tree have been developed, such as the *Virtual Brownian Tree* [34, 46] and *Brownian Interval* [38].

• Separate from the practical challenge of implementing numerical methods for SDEs with adaptive step sizes, there is a "simple" theoretical question to be addressed: Do numerical methods for SDEs with non-previsible adaptive step sizes converge?

It was show in [21, Theorem 4.3] that pathwise convergence in the adaptive step size setting does still hold provided that the numerical solution  $Y = \{Y_k\}$  satisfies

$$Y_{k+1} = Y_k + f(Y_k)h_k + g(Y_k)W_k + \sum_{i,j=1}^d g'_j(Y_k)g_i(Y_k) \int_{t_k}^{t_{k+1}} \left(W_t^i - W_{t_k}^i\right) \circ dW_t^j + o(h_k)g_i(Y_k) \int_{t_k}^{t_{k+1}} \left(W_t^i - W_t^i\right) \circ dW_t^j + o(h_k)g_i(Y_k) \int_{t_k}^{t_k} \left(W_t^i - W_t^i\right) \otimes dW_t^j + o(h_k)g_i(Y_k) \int_{t_k}^{t_k} \left(W_t^i - W_t^i\right) \otimes dW_t^j + o(h_k)g_i(Y_k) \int_{t_k}^{t_k} \left(W_t^i - W_t^i\right) \otimes dW_t^j + o(h_k)g_i(Y_k) \otimes dW_t^j + o(h_k)g_i(Y_k)$$

where  $W_k := W_{t_{k+1}} - W_{t_k}$  and  $h_k := t_{k+1} - t_k$  denote the Brownian increment and step size used to propagate the numerical solution  $Y_k$  over the interval  $[t_k, t_{k+1}]$ .

• To motivate this pathwise convergence result, Gaines and Lyons present a simple counterexample [21, Section 4.1] demonstrating that the Euler-Maruyama method, which clearly does not satisfy the above condition, can converge to an incorrect (i.e. not Itô) solution of the SDE when the adaptive step sizes are non-previsible.

Therefore, it is proposed in [21] that second iterated integrals of Brownian motion should be finely discretized (with  $o(h_k)$  accuracy) to ensure the SDE solver converges. As a consequence, following [21], research has focused on either previsible step sizes [14, 29, 30, 41, 52], non-previsible adaptive step sizes for restricted classes of SDEs [5, 6, 18, 32, 34, 49, 57, 60] or adaptive high order schemes with iterated integrals [2].

In this paper, we argue that such integrals are not needed and traditional methods, such as the no-area Milstein and Heun methods, do converge with adaptive step sizes. To this end, we shall present an informal version of our main result, Theorem 2.19.

THEOREM 1.1 (Convergence of adaptive SDE approximations, informal version). Suppose that the vector fields of the SDE (1.1) are bounded and twice differentiable with bounded derivatives. In addition, we assume the second derivatives are  $\alpha$ -Hölder continuous with  $\alpha \in (0, 1)$ . We make the following assumptions about the SDE solver:

1. Let  $Y = \{Y_k\}$  denote a numerical solution of (1.1) where  $Y_0 := y_0$  and each step  $Y_k \mapsto Y_{k+1}$ , between times  $t_k$  and  $t_{k+1}$ , is a function of the (random) quantities:

(1.2) 
$$h_k = t_{k+1} - t_k, \ \mathcal{W}_k := \left\{ \int_{t_k}^{t_{k+1}} \left( \frac{t - t_k}{t_{k+1} - t_k} \right)^j dW_t \right\}_{0 \le j \le q}, \ z_k \in \mathbb{R}^m,$$

where  $h_k$  is the step size used,  $q \ge 0$  and the random vector  $z_k$  is assumed to be independent from the Brownian motion. For example,  $z_k$  could be the additional uniform random variable used within so-called "randomized" numerical methods [27, 31, 40, 51, 61]. In addition, we assume that each step of the method satisfies

(1.3) 
$$Y_{k+1} = Y_k + f(Y_k)h_k + g(Y_k)W_k + \sum_{i,j=1}^d g'_j(Y_k)g_i(Y_k) \mathbb{E}\bigg[\int_{t_k}^{t_{k+1}} (W_t^i - W_{t_k}^i) \circ dW_t^j \, \Big| \, \mathcal{W}_k\bigg] + R_k\big(Y_k, h_k, \mathcal{W}_k, z_k\big),$$

where the remainder  $R_k$  is assumed to be small so that  $R_k \sim o(h_k)$  almost surely.

2. Given an initial (random) partition  $D_0$  on [0,T], we define a sequence  $\{D_n\}$  of nested partitions inductively by

(1.4) 
$$D_{n+1} := \mathcal{R}\big(\mathcal{E}\big(D_n, W\big)\big).$$

where  $\mathcal{E}$  gives the required information (1.2) about W with respect to the partition,

$$\mathcal{E}(\lbrace t_k \rbrace, W) := \left\{ \begin{pmatrix} h_k = t_{k+1} - t_k, & z_k \in \mathbb{R}^m, \\ \left(h_k, \mathcal{W}_k, z_k\right) : & \mathcal{W}_k = \left\{ \int_{t_k}^{t_{k+1}} \left(\frac{t - t_k}{t_{k+1} - t_k}\right)^j dW_t \right\}_{0 \le j \le q} \right\},$$

and  $\mathcal{R}$  maps a given partition to a finer partition, that is  $D_n \subset \mathcal{R}(\mathcal{E}(D_n, W))$ . We also assume the sequence  $\{D_n\}$  satisfies  $\lim_{n\to\infty} \operatorname{mesh}(D_n) = 0$  almost surely.

The "refinement" map  $\mathcal{R}$  represents the adaptive step size control for the solver which, when given a partition  $D_n$ , may add steps based on only the information about the Brownian motion that was generated with respect to  $D_n$  (see Figure 2).

Then, letting  $Y^n$  denote the numerical solution computed over  $D_n = \{t_k^n\}_{k \ge 0}$ , we have

$$\sup_{k} \left\| Y_k^n - y(t_k^n) \right\|_2 \to 0,$$

as  $n \to 0$  almost surely, where  $\|\cdot\|_2$  denotes the standard Euclidean norm.

Remark 1.2. The condition (1.3) is much less restrictive than [21, Equation (4.6)] since it does not require the second iterated integrals of the Brownian motion to be finely discretized with o(h) accuracy. In practice, (1.3) can be verified as, for example,

(1.5) 
$$\mathbb{E}\left[\int_{t_k}^{t_{k+1}} \left(W_t^i - W_{t_k}^i\right) \circ dW_t^j \,\middle|\, W_k\right] = \frac{1}{2} W_k^i W_k^j$$

(1.6) 
$$\mathbb{E}\left[\int_{t_k}^{t_{k+1}} \left(W_t^i - W_{t_k}^i\right) \circ dW_t^j \,\Big|\, W_k, H_k\right] = \frac{1}{2} W_k^i W_k^j + H_k^i W_k^j - W_k^i H_k^j,$$

where  $H_k = \int_{t_k}^{t_{k+1}} \left(\frac{1}{2} - \frac{t}{h_k}\right) dW_t \sim \mathcal{N}(0, \frac{1}{12}h_k I_d)$  is independent of  $W_k$  [17, Theorem 5.4]. Intuitively, the fact that these approximations are conditional expectations enables us to employ certain martingale arguments for establishing convergence in our analysis. Otherwise, the O(h) local errors may propagate linearly and give a global O(1) error.

To support the above remark, we give a few examples of SDE solvers which satisfy the condition (1.3) and have an embedded method to enable adaptive step size control. Of course, these are not extensive and we hope that our general convergence analysis will enable the development of further solvers with non-previsible adaptive step sizes.

1.1. Examples of adaptive numerical methods. We introduce the notation

(1.7) 
$$F(y) := \begin{pmatrix} f(y) & g(y) \end{pmatrix}, \quad \overline{W}_t := \begin{pmatrix} t & W_t \end{pmatrix}^{\mathsf{T}},$$

so that the SDE (1.1) can be rewritten as a simpler "controlled" differential equation:

(1.8) 
$$dy_t = F(y_t) \circ d\overline{W}_t$$

To define our examples of adaptive numerical methods, we also introduce the notation

(1.9) 
$$\overline{W}_k := \begin{pmatrix} h_k & W_k \end{pmatrix}^{\mathsf{T}}, \qquad \overline{H}_k := \begin{pmatrix} 0 & H_k \end{pmatrix}^{\mathsf{T}}$$

where  $H_k := \int_{t_k}^{t_{k+1}} \left(\frac{1}{2} - \frac{t}{h_k}\right) dW_t = \frac{1}{h_k} \int_{t_k}^{t_{k+1}} (W_t - W_{t_k} - \frac{t - t_k}{h_k} W_k) dt \sim \mathcal{N}(0, \frac{1}{12} h_k I_d)$  is the "space-time" Lévy area of Brownian motion on  $[t_k, t_{k+1}]$ , see [20, Definition 3.5].

We now present a few straightforward numerical methods that are compatible with the class of adaptive step size controls satisfying the conditions of Theorem 1.1.

Due to their popularity in ODE numerics, we propose embedded methods which output  $Y_{k+1}$  alongside another approximation  $\tilde{Y}_{k+1}$  from  $(Y_k, h_k, \mathcal{W}_k, z_k)$  at each step (often by reusing vector field evaluations). The difference between  $Y_{k+1}$  and  $\tilde{Y}_{k+1}$  may then serve as a proxy for the local error and used by the adaptive step size controller. For example, the decision to halve a step size  $h_k$  can simply be made if  $||Y_{k+1} - \tilde{Y}_{k+1}||_2$ is above a user-specified tolerance. In particular, this should improve accuracy when "large" Brownian increments  $W_k$  are generated (which happens with low probability).

Whilst [21] showed that the Euler-Maruyama method may fail to converge to the Itô SDE solution when used with an adaptive step size, our main result establishes the convergence of the following "increment only" Milstein method introduced in [22].

DEFINITION 1.3 (No-area Milstein method with embedded Euler for Itô SDEs). We define a numerical solution  $\{Y_k\}$  for the below SDE (understood in the Itô sense)

(1.10) 
$$dy_t = f(y_t)dt + g(y_t)dW_t,$$

by  $Y_0 := y_0$  and, for  $k \ge 0$ , we define  $Y_{k+1}$  and the embedded approximation  $\widetilde{Y}_{k+1}$  as

(1.11) 
$$\widetilde{Y}_{k+1} := Y_k + f(Y_k)h_k + g(Y_k)W_k$$

(1.12) 
$$Y_{k+1} := Y_k + f(Y_k)h_k + g(Y_k)W_k + \frac{1}{2}g'(Y_k)g(Y_k)(W_k^{\otimes 2} - h_kI_d)$$

Remark 1.4. To apply our main result, we rewrite (1.10) as a Stratonovich SDE:

$$dy_t = \bar{f}(y_t)dt + g(y_t) \circ dW_t,$$

where  $\tilde{f}(y) := f(y) - \frac{1}{2} \sum_{i=1}^{d} g'_i(y) g_i(y)$  is obtained via the Itô-Stratonovich correction. Then the no-area Milstien method given by (1.12) simplifies and can be expressed as

$$Y_{k+1} := Y_k + \widetilde{f}(Y_k)h_k + g(Y_k)W_k + \frac{1}{2}g'(Y_k)g(Y_k)W_k^{\otimes 2}.$$

By equations (1.3) and (1.5), with  $R_k = 0$ , it is clear the above has the required form.

To our knowledge, the following are the first embedded stochastic Runge-Kutta methods for general SDEs (see [59, 60] for examples with scalar or additive noise). We first consider a multidimensional extension of the Heun scheme proposed in [58].

DEFINITION 1.5 (Heun's method with embedded Euler-Maruyama). Using the notation in (1.7) and (1.9), we define a numerical solution  $\{Y_k\}$  for the SDE (1.8) as

(1.13) 
$$\widetilde{Y}_{k+1} := Y_k + F(Y_k)\overline{W}_k$$

(1.14) 
$$Y_{k+1} := Y_k + \frac{1}{2} \left( F(Y_k) + F(\widetilde{Y}_{k+1}) \right) \overline{W}_k,$$

for  $k \ge 0$  with  $Y_0 := y_0$ .

In addition, we shall introduce a new stochastic Runge-Kutta method that uses space-time Lévy area  $H_k$  to achieve a slightly better local accuracy than traditional "increment only" schemes, including Heun's method (see Appendix A for details). For additive noise SDEs, the proposed scheme becomes high order and achieves an  $O(h^{1.5})$  strong convergence rate (i.e. when the noise vector fields  $g_i$  are constant). This scheme was inspired by a recent path-based splitting method [16, Example 1.3]. DEFINITION 1.6 (Splitting Path Runge-Kutta (SPaRK) with embedded Heun). We define a numerical solution  $\{Y_k\}$  for the SDE (1.8) by  $Y_0 := y_0$  and, for all  $k \ge 0$ ,

$$Y_{k+\frac{1}{2}} := Y_k + F(Y_k) \left(\frac{1}{2}\overline{W}_k + \sqrt{3}\overline{H}_k\right), \qquad Z_{k+1} := Y_k + F(Y_{k+\frac{1}{2}})\overline{W}_k,$$
  
1.15) 
$$\widetilde{Y}_{k+1} := Y_k + \frac{1}{2} (F(Y_k) + F(Z_{k+1}))\overline{W}_k,$$

$$(1.16) \quad Y_{k+1} := Y_k + F\left(Y_k\right) \left(a\overline{W}_k + \overline{H}_k\right) + bF\left(Y_{k+\frac{1}{2}}\right)\overline{W}_k + F\left(Z_{k+1}\right) \left(a\overline{W}_k - \overline{H}_k\right),$$

where  $a := \frac{3-\sqrt{3}}{6}$  and  $b := \frac{\sqrt{3}}{3}$ .

Further details on this stochastic Runge-Kutta method is given in Appendix A. In particular, the improved local accuracy of (1.16) over Heun's method is detailed. However, the formulation of the above (embedded) Runge-Kutta methods is only one component of the SDE solver – the other being an algorithm for controlling step sizes.

For example, we could consider the following simple adaptive step size control, though we emphasise that our analysis (Theorem 2.19) holds more generally, provided the associated sequence of partitions is nested and has its mesh size converging to zero.

DEFINITION 1.7 (A simple step size control for embedded Runge-Kutta methods). Let  $h_{init}, C > 0$  be fixed user-specified constants (corresponding to the initial step size and local error tolerance). Then we depth-first recursively construct a Brownian tree, with the n-th level containing Brownian information on intervals of size  $h_{init}/2^n$ , until

(1.17) 
$$\left\|Y_{k+1} - \widetilde{Y}_{k+1}\right\|_2 \leqslant C\sqrt{h_k},$$

holds for all  $k \ge 0$ , where the numerical solution  $Y = \{Y_k\}_{k\ge 0}$  is constructed using the information of the Brownian motion provided by the leaves of the Brownian tree.

Remark 1.8. In SDE numerics, it is common to consider mean squared errors. Informally speaking, in the constant step size regime, a local mean squared error of  $O(h^p)$  gives a global mean squared error of  $O(h^{p-1})$  (see [50, Theorem 1.1] for details). Therefore, if we would like the global error to be less than  $\varepsilon > 0$ , then we would expect local mean squared errors to be at most  $O(\varepsilon h)$ . This motivates the estimator (1.17) since we expect  $||Y_{k+1} - \tilde{Y}_{k+1}||_2^2$  to be a good proxy for the local mean squared error.

**1.2.** Organisation of the paper. In Section 2, we will establish our main convergence result (Theorem 2.19) for a wide class of adaptive SDEs approximations. The central idea in our proof methodology is to employ *rough path theory* and first show that piecewise polynomial approximations of Brownian motion converge in the *p*-variation metric along a sequence of nested partitions satisfying certain properties. This argument is a natural extension of the analysis presented in [68, Section 13.3.2], which presents similar results for piecewise linear approximations of Brownian motion that are defined along a sequence of partitions that are both nested and deterministic.

Once we have convergence in *p*-variation for piecewise polynomial approximations, we can then apply the well-known Universal Limit Theorem from rough path theory to show that the corresponding sequence of Controlled Differential Equations (CDEs) converges to the solution of the (Stratonovich) SDE. We show that this convergence is preserved even when small o(h) local perturbations are added to the CDE solutions. By considering the local Taylor expansion of the CDEs between discretization times, we obtain the condition (1.3) from Theorem 1.1 for the numerical method to satisfy.

(

We conclude Section 2 by showing the no-area Milstein, Heun and Splitting Path Runge-Kutta (SPaRK) methods in Section 1.1 satisfy the convergence condition (1.3).

In Section 3, we give an example demonstrating that without nested partitions, an adaptive step size control can induce a non-zero bias which prevents convergence. Here, the decision to reduce a step size is based on whether it increases the value of the numerical solution. In particular, step sizes can "skip over" known values of W. Ultimately, this leads to an approximation with an incorrect mean, even in the limit.

In Section 4, we present an experiment comparing methods with different step size controls for the well-known SABR<sup>1</sup> model used in mathematical finance [7, 11, 25, 44]. The SABR model describes the evolution of an interest (or exchange) rate  $S = \{S_t\}$ whose local volatility  $\sigma = \{\sigma_t\}$  is also stochastic. It is given by the following Itô SDE:

(1.18) 
$$dS_t = \sqrt{1 - \rho^2} \sigma_t (S_t)^\beta dW_t^1 + \rho \sigma_t (S_t)^\beta dW_t^2,$$
$$d\sigma_t = \alpha \sigma_t dW_t^2,$$

where W is a standard two-dimensional Brownian motion and  $\alpha, \beta, \rho$  are parameters. For simplicity, we will set  $\alpha = 1$  and  $\beta = \rho = 0$ . In Stratonovich form, (1.18) becomes

(1.19) 
$$dS_t = \sigma_t \circ dW_t^1,$$
$$d\sigma_t = -\frac{1}{2}\sigma_t dt + \sigma_t \circ dW_t^2.$$

Despite its simple form, the SDE (1.19) is challenging to accurately discretize since its Taylor expansion contains the non-Gaussian stochastic integral of  $W^1$  against  $W^2$ ,

(1.20) 
$$S_{t+\Delta t} = S_t + \sigma_t \left( W_{t+\Delta t}^1 - W_t^1 \right) + \sigma_t \int_t^{t+\Delta t} \left( W_u^2 - W_t^2 \right) \circ dW_u^1 + R_{\Delta t}(t),$$

where the remainder is  $R_{\Delta t}(t) = \sigma_t \int_t^{t+\Delta t} \left(e^{-\frac{1}{2}(u-t)+W_u^2-W_t^2} - \left(1+W_u^2-W_t^2\right)\right) \circ dW_u^1$ . Therefore by (1.20), approximating  $S_{t+\Delta t}$  using one step of Heun's method, with a sufficiently small step size, will result in a mean squared error of  $\frac{1}{4}\sigma_t^2(\Delta t)^2 + O((\Delta t)^3)$ . As a consequence, we would expect that the overall accuracy of a numerical method can be improved by decreasing  $\Delta t$  if  $\sigma_t$  when large and increasing  $\Delta t$  when  $\sigma_t$  is small. More specifically, in our experiment, we observe that previsible step sizes of the form  $\Delta t = \log(1 + C\sigma_t^{-2})$  give a faster convergence rate which is linear instead of  $O(\sqrt{\Delta t})$ . We also demonstrate that the same first order convergence rate can be observed when local mean squared errors are estimated via an embedded Runge-Kutta method and used by a standard "Proportional-Integral" (PI) adaptive step size controller [6, 32].

In Section 5, we conclude the paper and discuss potential areas of future research. Among the areas of future research, we would like to extend our convergence analysis to a multidimensional extension of the stochastic Runge-Kutta method proposed in [58] (which can be viewed as a "Heun's method for Itô SDEs"). This would then enable general Itô SDEs to be solved with non-previsible step sizes, but without explicitly requiring vector field derivatives – which is the case for the no-area Milstein method.

**1.3. Notation.** In this subsection, we summarise the notation used in the paper. We use  $\|\cdot\|_2$  to denote the usual Euclidean norm. For a sequence of random variables  $\{X_n\}$  in a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ , we say  $X_n \to X$  in  $L^2(\mathbb{P})$  if  $\mathbb{E}_{\mathbb{P}}[\|X_n - X\|_2^2] \to 0$ . For  $w : \mathbb{R}_+ \to \mathbb{R}_+$ , we say that  $w(h) \sim o(h)$  if  $\frac{w(h)}{h} \to 0$  as  $h \to 0$  and  $w(h) \sim O(h^p)$  with p > 0, if there exists a constant C such that  $w(h) \leq Ch^p$  for sufficiently small h.

 $<sup>^{1}</sup>$ Stochstic Alpha-Beta-Rho

Throughout,  $W = \{W_t\}_{t \in [0,T]}$  will be a standard *d*-dimensional Brownian motion. For a sequence of (random) nested partitions  $\{D_n\}$  of [0,T], we use  $\widetilde{W}^n = \{\widetilde{W}^n_t\}_{t \in [0,T]}$  to denote a piecewise polynomial approximation of W defined with respect to  $D_n$ . Most notably, this can be the usual piecewise linear discretization (see [20] for details). We write increments of the paths as  $W_{s,t} := W_t - W_s$  and  $\widetilde{W}^n_{s,t} := \widetilde{W}^n_t - \widetilde{W}^n_s$  for  $s \leq t$ .

We will denote SDE solutions by  $y = \{y_t\}_{t \in [0,T]}$  and CDE solutions by  $\tilde{x}^n$  or  $\tilde{y}^n$ . Each partition can be expressed as  $D_n = \{0 = t_0^n < t_1^n < \cdots < t_{K_n-1}^n < t_{K_n}^n = T\}$  or  $\{D_n\} = \{h_k^n\}_{0 \leq k < K_n}$  where  $K_n \geq 1$  and  $h_k^n := t_{k+1}^n - t_k^n$  represents the step size. We often use the shorthand notation  $W_k := W_{t_{k+1}} - W_{t_k}$  when defining numerical methods – which are denoted by  $Y = \{Y_k\}_{k \geq 0}$ . Similarly, we write  $H_k \sim \mathcal{N}(0, \frac{1}{12}h_k I_d)$  for the space-time Lévy area of Brownian motion on  $[t_k, t_{k+1}]$ , see [20, Definition 3.5].

Given a random vector X, we let  $\sigma(X)$  denote the  $\sigma$ -algebra generated by X. Filtrations are denoted by  $\{\mathcal{F}_n\}$  or  $\{\mathcal{G}_n\}$  and give information about W and partitions. Blackboard bold symbols lie in  $\mathbb{R}^{d \times d}$ , such as iterated integrals of Brownian motion  $\mathbb{W}_{s,t} = \left\{\int_s^t W_{s,u}^i \circ dW_u^j\right\}_{1 \le i,j \le d}$ . Bold symbols lie in the tensor algebra  $\mathbb{R} \oplus \mathbb{R}^d \oplus \mathbb{R}^{d \times d}$ , such as the *Stratonovich enhanced* Brownian motion  $\mathbf{W} = \{\mathbf{W}_{s,t}\} = \{(1, W_{s,t}, \mathbb{W}_{s,t})\}$ .

We use  $\otimes$  to denote the usual tensor product with  $a \otimes b := \{a_i b_j\}_{1 \leq i,j \leq d} \in \mathbb{R}^{d \times d}$ for vectors  $a = \{a_i\}_{i=1}^d$  and  $b = \{b_i\}_{i=1}^d$  in  $\mathbb{R}^d$ . In particular, we can use this notation to define the iterated integrals  $\mathbb{W}_{s,t} := \int_s^t W_{s,u} \otimes \circ dW_u$  and  $\widetilde{\mathbb{W}}_{s,t}^n := \int_s^t \widetilde{W}_{s,u}^n \otimes d\widetilde{W}_u^n$ . In addition, there will be also notation for rough paths  $\boldsymbol{X}$  and  $\boldsymbol{Y}$  (see Definition 2.9). This includes the  $\alpha$ -Hölder norm  $\|\boldsymbol{X}\|_{\alpha-\text{Höl};[s,t]}$ , the  $\alpha$ -Hölder metric  $d_{\alpha-\text{Höl};[s,t]}(\boldsymbol{X}, \boldsymbol{Y})$ and the *p*-variation metric  $d_{p-\text{var};[0,\mathrm{T}]}(\boldsymbol{X}, \boldsymbol{Y})$ , where  $\alpha \in (\frac{1}{3}, \frac{1}{2})$  and  $p = \frac{1}{\alpha} \in (2, 3)$ . For a vector  $\boldsymbol{X} = (1, X, \mathbb{X}) \in \mathbb{R} \oplus \mathbb{R}^{d} \oplus \mathbb{R}^{d \times d}$ , we also define  $\|\boldsymbol{X}\| := \max(\|X\|_2, \|\mathbb{X}\|_2^{\frac{1}{2}})$ .

2. Main result. To begin, we will first establish the "rough path" convergence of piecewise polynomial approximations along nested partitions to Brownian motion. However, unlike [68, Section 13.3.2], which our analysis is based on, we will consider a sequence of nested partitions that may depend non-trivially on the Brownian motion. Since they correspond to an adaptive solver and we wish to use martingale arguments, we construct the next partition as a function of only the information generated about the Brownian motion with respect to the current partition. We note that this strategy for showing convergence was originally proposed in the doctoral thesis [15, Chapter 6]. That said, in this paper, we consider a more general class of adaptive approximations as we allow dependence on random variables independent of the Brownian motion. To define the corresponding sequence of partitions, we make the following definitions:

DEFINITION 2.1 (Enhancement function). For a d-dimensional Brownian motion W and a partition  $D = \{h_k\}$  of [0,T], we consider the functions  $\mathcal{E}_W$  and  $\mathcal{E}_Z$  given by

$$(2.1) \qquad \mathcal{E}_{\mathcal{W}}(\{h_k\}, W) := \left\{ \left(h_k, \mathcal{W}_k\right) : \mathcal{W}_k = \left\{ \int_{t_k}^{t_{k+1}} \left(\frac{t - t_k}{t_{k+1} - t_k}\right)^j dW_t \right\}_{0 \le j \le q} \right\},$$

$$(2.2) \qquad \mathcal{E}_{\mathcal{Z}}(\{h_k\}, W) := \left\{ \left(h_k, z_k\right) : z_k \in \mathbb{R}^m \right\},$$

where  $q \in \{0, 1 \cdots\}$  and  $z_k$  is a random vector in  $\mathbb{R}^m$ , with  $m \ge 1$ , that only depends on the Brownian motion W through the information contained in  $\sigma(\mathcal{E}_{\mathcal{W}}(\{h_k\}, W))$ . By combining functions (2.1) and (2.2), we define the "enhancement" function  $\mathcal{E}$  as

(2.3) 
$$\mathcal{E}(\{h_k\}, W) := \mathcal{E}_{\mathcal{W}}(\{h_k\}, W) \cup \mathcal{E}_{\mathcal{Z}}(\{h_k\}, W).$$

DEFINITION 2.2 (Refinement function associated with an adaptive SDE solver). Let  $\mathcal{D}$  be the set of partitions on [0,T], which we can express in terms of step sizes as

$$\mathcal{D} := \left\{ \{h_k\}_{0 \le k < N} : h_k > 0, \sum_{k=0}^{N-1} h_k = T \text{ and } N \ge 1 \right\},\$$

where  $\{h_k\}$  corresponds to the partition  $\{0 = t_0 < \cdots < t_N = T\}$  with  $h_k = t_{k+1} - t_k$ . Similarly, we define the set of enhanced partitions on [0, T] as

$$\widetilde{\mathcal{D}} := \left\{ \left\{ \left(h_k, w_k\right) \right\}, \left\{ \left(h_k, z_k\right) \right\} : \left\{h_k\right\} \in \mathcal{D}, w_k \in (\mathbb{R}^d)^{\geq 1} \text{ and } z_k \in \mathbb{R}^m \right\},\right.$$

where  $(\mathbb{R}^d)^{\geq 1}$  is a Euclidean space  $\mathbb{R}^{pd}$  with  $p \geq 1$ . A refinement function  $\mathcal{R}$  is a map

$$\mathcal{R}: \widetilde{\mathcal{D}} \to \mathcal{D},$$

such that the times  $\{t_k\}$  associated with the partition  $\{h_k\}$  are contained in  $\mathcal{R}(\{h_k, \cdot\})$ . That is,  $\mathcal{R}(\{h_k, \cdot\})$  is a refinement of the input partition  $\{h_k\}$  for any  $\{w_k\}$  and  $\{z_k\}$ .

DEFINITION 2.3 (Partitions and filtrations defined using the refinement function). Let  $\mathcal{E}$  and  $\mathcal{R}$  denote the enhancement and refinement functions given by Definitions 2.1 and 2.2. Given a standard Brownian motion  $W = \{W_t\}_{t \in [0,T]}$  and a random initial partition  $D_0$ , we will define a sequence  $\{D_n\}$  of nested partitions inductively by

(2.4) 
$$D_{n+1} := \mathcal{R}\big(\mathcal{E}\big(D_n, W\big)\big).$$

We define a sequence of  $\sigma$ -algebras  $\{\mathcal{F}_n\}$  by  $\mathcal{F}_0 := \sigma(\mathcal{E}(D_0, W))$  and for each  $n \ge 0$ ,

(2.5) 
$$\mathcal{F}_{n+1} := \sigma\Big(\mathcal{F}_n \cup \mathcal{E}\big(D_{n+1}, W\big)\Big).$$

Since  $\{\mathcal{F}_n\}$  is clearly an increasing sequence of  $\sigma$ -algebras, it is therefore a filtration. We will also define another filtration  $\{\mathcal{G}_n\}$  by  $\mathcal{G}_0 := \sigma(\mathcal{E}_{\mathcal{Z}}(D_0, W))$  and for each  $n \ge 0$ ,

(2.6) 
$$\mathcal{G}_{n+1} := \sigma \Big( \mathcal{G}_n \cup \mathcal{E}_{\mathcal{Z}} \big( D_{n+1}, W \big) \Big).$$

Having defined a sequence of partitions and the associated filtrations, we now show the limiting filtration contains all the information about the Brownian motion. To do so, we first identify  $\mathcal{F}_n$  as just the  $\sigma$ -algebra containing information about Wwith respect to  $D_n$ , with the additional independent random vectors contained in  $\mathcal{G}_n$ .

PROPOSITION 2.4 (Filtration captures path information on the finest partition). Let  $\{\mathcal{F}_n\}$  and  $\{\mathcal{G}_n\}$  be the filtrations given by (2.5) and (2.6) in Definition 2.3. Then

(2.7) 
$$\mathcal{F}_n = \sigma\Big(\mathcal{E}_{\mathcal{W}}\big(D_n, W\big) \cup \mathcal{G}_n\Big).$$

*Proof.* We assume (2.7) for some n and aim to show  $\mathcal{F}_{n+1} = \sigma(\mathcal{E}(D_{n+1}, W) \cup \mathcal{G}_n)$ . Since  $D_{n+1} = \mathcal{R}(\mathcal{E}(D_n, W))$ , we see that  $D_{n+1}$  is a refinement of the partition  $D_n$ . Therefore, by merging a finite number of subintervals in  $D_{n+1}$ , we can obtain  $D_n$ . Suppose we merge the subintervals [s, u] and [u, t]. Then, for  $j \in \{0, \dots, q\}$ , we have

$$\begin{split} \int_{s}^{t} \left(\frac{r-s}{t-s}\right)^{j} dW_{r} &= \int_{s}^{u} \left(\frac{r-s}{t-s}\right)^{j} dW_{t} + \int_{u}^{t} \left(\frac{r-s}{t-s}\right)^{j} dW_{r} \\ &= \left(\frac{u-s}{t-s}\right)^{j} \int_{s}^{u} \left(\frac{r-s}{u-s}\right)^{j} dW_{r} + \left(\frac{t-u}{t-s}\right)^{j} \int_{u}^{t} \left(\frac{r-s}{t-u}\right)^{j} dW_{r} \\ &= \left(\frac{u-s}{t-s}\right)^{j} \int_{s}^{u} \left(\frac{r-s}{u-s}\right)^{j} dW_{r} \\ &+ \left(\frac{t-u}{t-s}\right)^{j} \sum_{l=0}^{j} {j \choose l} \left(\frac{u-s}{t-u}\right)^{j-l} \int_{u}^{t} \left(\frac{r-u}{t-u}\right)^{l} dW_{r}. \end{split}$$

In other words, the integrals  $\{\int_{s}^{t} \left(\frac{r-s}{t-s}\right)^{j} dW_{r} : 0 \leq j \leq q\}$  can be recovered from  $\{\int_{s}^{u} \left(\frac{r-s}{u-s}\right)^{j} dW_{r}\}$  and  $\{\int_{u}^{t} F\left(\frac{r-u}{t-u}\right)^{j} dW_{r}\}$ . Moreover, since  $D_{n+1}$  is a refinement of  $D_{n}$ , we can recover the information contained in  $\sigma(\mathcal{E}_{\mathcal{W}}(D_{n},W))$  from  $\sigma(\mathcal{E}_{\mathcal{W}}(D_{n+1},W))$ . Hence, the  $\sigma$ -algebras depending on W (i.e. without  $\mathcal{G}_{n}$ ) would lead to a filtration as

(2.8) 
$$\sigma(\mathcal{E}_{\mathcal{W}}(D_n, W)) \subseteq \sigma(\mathcal{E}_{\mathcal{W}}(D_{n+1}, W))$$

Thus, by "expanding"  $\mathcal{F}_{n+1}$  using its definition and the induction hypothesis, we have

$$\mathcal{F}_{n+1} = \sigma\left(\mathcal{F}_n \cup \mathcal{E}(D_{n+1}, W)\right)$$
  
=  $\sigma\left(\left(\mathcal{E}_{\mathcal{W}}(D_n, W) \cup \mathcal{G}_n\right) \cup \left(\mathcal{E}_{\mathcal{W}}(D_{n+1}, W) \cup \mathcal{E}_{\mathcal{Z}}(D_{n+1}, W)\right)\right)$   
=  $\sigma\left(\mathcal{E}_{\mathcal{W}}(D_{n+1}, W) \cup \mathcal{G}_n \cup \mathcal{E}_{\mathcal{Z}}(D_{n+1}, W)\right)$   
=  $\sigma\left(\mathcal{E}_{\mathcal{W}}(D_{n+1}, W) \cup \mathcal{G}_{n+1}\right).$ 

The result now follows by induction as  $\mathcal{F}_0 := \sigma(\mathcal{E}(D_0, W)), \mathcal{G}_0 := \sigma(\mathcal{E}_{\mathcal{Z}}(D_0, W)).$ 

PROPOSITION 2.5. Let  $\{D_n\}$  be the sequence of partitions given by Definition 2.3 and suppose that  $\operatorname{mesh}(D_n) \to 0$  almost surely. Then  $\mathcal{F}_{\infty} := \bigcup_{n \ge 0} \mathcal{F}_n$  is given by

(2.9) 
$$\mathcal{F}_{\infty} = \sigma\Big(\big\{W_t : t \in [0,T]\big\} \cup \mathcal{G}_{\infty}\Big),$$

where  $\mathcal{G}_{\infty} := \bigcup_{n \ge 0} \mathcal{G}_n$ .

*Proof.* Since the enhancement function produces  $\mathcal{W}_k = \{\int_{t_k}^{t_{k+1}} \left(\frac{t-t_k}{t_{k+1}-t_k}\right)^j dW_t\},\$ we see that  $\mathcal{W}_k$  contains the Brownian motion's increment  $\int_{t_k}^{t_{k+1}} 1 dW_t = W_{t_{k+1}} - W_{t_k}.$ 

For any fixed  $t \in [0, T]$ , we can find the largest time point  $t_n \in D_n$  such that  $t_n \leq t$ . As  $0 \leq t - t_n < \operatorname{mesh}(D_n)$ , it follows from the assumption that  $t_n \to t$  almost surely. Moreover, since Brownian motion is continuous almost surely, we have  $W_{t_n} \to W_t$ almost surely and we can thus obtain  $W_t$  from  $\mathcal{F}_\infty$ . That is,  $\sigma(W_t : t \in [0, T]) \subseteq \mathcal{F}_\infty$ . It is also clear from the definitions of the filtrations that  $\mathcal{G}_n \subseteq \mathcal{F}_n$  and thus  $\mathcal{G}_\infty \subseteq \mathcal{F}_\infty$ .

On the other hand, since W has finite p-variation almost surely, where p > 2, it follows that the Riemann-Stieltjes integrals  $\left\{\int_{t_k}^{t_{k+1}} \left(\frac{t-t_k}{t_{k+1}-t_k}\right)^j dW_t\right\}$  exist and are uniquely determined by W. Thus, we have  $\mathcal{F}_n \subseteq \sigma\left(\{W_t\}_{t \in [0,T]} \cup \mathcal{G}_{\infty}\right)$ , as required.

Having now established the above key properties of the filtration  $\mathcal{F} = \{\mathcal{F}_n\}_{n \ge 0}$ , we will proceed to show that the corresponding sequence of (adaptive) approximations  $\{\widetilde{W}^n\}_{n \ge 0}$  converges to Brownian motion on the interval [0, T] in a rough path sense. **2.1. Rough path convergence of piecewise polynomial approximations.** In order to show that our approximations of Brownian motion converge in a rough path sense (i.e. *p*-variation), we will first perform a standard mean squared analysis.

THEOREM 2.6 (Nested piecewise unbiased approximations of Brownian motion). Let  $\{D_n\}$  denote the sequence of partitions and  $\{\mathcal{F}_n\}$  the corresponding filtration given by Definition 2.3. Moreover, we will assume (2.7) and mesh $(D_n) \to 0$  almost surely. We define a sequence of approximations  $\{\widetilde{W}^n\}$  of Brownian motion as

(2.10) 
$$\widetilde{W}_t^n := \mathbb{E} \Big[ W_t \,|\, \mathcal{F}_n \Big].$$

Then for  $t \in [0,T]$ ,  $\widetilde{W}_t^n \to W_t$  almost surely and

(2.11) 
$$\mathbb{E}\Big[\big(\widetilde{W}_t^n - W_t\big)^2\Big] \to 0.$$

*Proof.* As  $\mathcal{F}_n$  is a filtration, it follows that  $\{\widetilde{W}^n\} = \{\mathbb{E}[W | \mathcal{F}_n]\}$  is a martingale. Moreover, each  $\widetilde{W}^n$  is square-integrable as by Jensen's inequality and the tower law,

$$\mathbb{E}\Big[\big(\widetilde{W}_t^n\big)^2\Big] = \mathbb{E}\Big[\mathbb{E}\big[W_t \,|\, \mathcal{F}_n\big]^2\Big] \leqslant \mathbb{E}\Big[\mathbb{E}\big[W_t^2 \,|\, \mathcal{F}_n\big]\Big] = \mathbb{E}\big[W_t^2\big] = t.$$

Therefore, by Doob's martingale convergence theorem, we have that for each  $t \in [0, T]$ ,

(2.12) 
$$\mathbb{E}[W_t | \mathcal{F}_n] \to \mathbb{E}[W_t | \mathcal{F}_\infty].$$

almost surely and in  $L^2(\mathbb{P})$ . Since mesh $(D_n) \to 0$  almost surely, we can see that  $\sigma(W_t : t \in [0,T]) \subseteq \mathcal{F}_{\infty}$  by Proposition 2.4. The result now directly follows by (2.12). It is worth noting that any additional information given by  $\mathcal{G}_n$  is independent of W, and therefore will have no effect on the conditional expectations within our proof.  $\Box$ 

Remark 2.7. If q = 0, then  $\mathcal{W}_n = \{W_{t_{n+1}} - W_{t_n}\}$  and each  $\widetilde{W}^n$  is the piecewise linear path that agrees with the Brownian motion W at the discretization points  $\{t_n\}$ given by the partition  $D_n$ . More generally,  $\widetilde{W}^n$  is a piecewise degree q polynomial that matches the "moment" information of W on  $D_n$  (see [20, Theorem 2.4] for details).

In order to utilise results from rough path theory, we extend the above theorem to include the second iterated integrals of Brownian motion (or equivalent Lévy areas).

THEOREM 2.8 (Convergence of nested unbiased approximations for Lévy area). Let  $\{\widetilde{W}^n\}$  denote the sequence of piecewise polynomial approximations for W given by Theorem 2.6. With the same definitions and assumptions used in Propositions 2.4 and 2.5, we consider the following sequence of approximate Lévy areas  $\{\widetilde{A}^n_{s,t}\}$  on [s,t].

$$\widetilde{A}_{s,t}^{n} := \left\{ \int_{s}^{t} \left( \widetilde{W}_{s,r}^{n} \right)^{i} d\left( \widetilde{W}_{r}^{n} \right)^{j} - \int_{s}^{t} \left( \widetilde{W}_{s,r}^{n} \right)^{j} d\left( \widetilde{W}_{r}^{n} \right)^{i} \right\}_{1 \leq i,j \leq d},$$

where  $0 \leq s \leq t \leq T$  and  $\widetilde{W}_{s,r}^n := \widetilde{W}_r^n - \widetilde{W}_s^n$  denotes the path increment of  $\widetilde{W}^n$ . Then there exists a random variable  $\widetilde{A}_{s,t}$  such that  $\widetilde{A}_{s,t}^n \to \widetilde{A}_{s,t}$  almost surely and in  $L^2(\mathbb{P})$ . Furthermore  $\widetilde{A}_{s,t}$  coincides almost surely with the "Lévy area" of Brownian motion,

$$(2.13) A_{s,t} := \left\{ \int_s^t W_{s,r}^i \circ dW_r^j - \int_s^t W_{s,r}^i \circ dW_r^j \right\}_{1 \le i,j \le d}$$

where  $W_{s,r} := W_r - W_s$ .

*Proof.* When i = j, we see that  $(\tilde{A}_{s,t}^n)^{i,j} = A_{s,t}^{i,j} = 0$  and there is nothing to prove. Since  $(\tilde{A}_{s,t}^n)^{i,j} = -(\tilde{A}_{s,t}^n)^{j,i}$  and  $A_{s,t}^{i,j} = -A_{s,t}^{j,i}$ , it suffices to consider the case i < j. Let  $n \ge 0$  be fixed and suppose  $t_k$  and  $t_{k+1}$  are adjacent points in the partition  $D_n$ . We first note the following four properties about Brownian motion:

- 1. Brownian motion has independent increments that are normally distributed. Thus  $\{W_{t_k,t}\}_{t \in [t_k, t_{k+1}]}$  is independent of  $\{W_t\}_{t \in [0, t_k]}$  and  $\{W_{t_{k+1},t}\}_{[t_{k+1},T]}$ .
- 2. Brownian motion has independent coordinate processes. That is, when  $i \neq j$ ,  $W^i$  and  $W^j$  are independent.
- 3. For c > 0, the process  $t \mapsto \frac{1}{c} W_{c^2 t}$  has the same law as a Brownian motion.
- 4. For a fixed  $q \ge 0$ , the process  $\{W_t \mathbb{E}[W_t | \{\int_0^1 s^l dW_s\}_{0 \le l \le q}]\}_{t \in [0,1]}$  is a centered Gaussian process on [0, 1] and is independent of  $\{\int_0^1 s^j dW_s\}_{0 \le j \le q}$ . This is a direct consequence of a polynomial expansion of Brownian motion (see, for example, [20, Theorem 2.2] or [24, Proposition 1.3] for such details).

Since the information currently known about W is given in  $\mathcal{F}_n = \sigma(\mathcal{E}(D_n, W) \cup \mathcal{G}_n)$ , and  $D_n$  contains no further subintervals of  $[t_k, t_{k+1}]$  for adjacent times  $t_k, t_{k+1} \in D_n$ , it follows from the first and fourth properties of Brownian motion that

$$\widetilde{W}_{t,t_k}^n = \mathbb{E}\bigg[W_{t_k,t} \,\Big| \, \int_{t_k}^{t_{k+1}} t^l \, dW_t, \, 0 \leqslant l \leqslant q \bigg],$$

for  $t \in [t_k, t_{k+1}]$ . In addition, it directly follows from the last three properties that the *i*-th and *j*-th coordinates of

$$\begin{cases} \frac{1}{\sqrt{t_{k+1} - t_k}} \left( W_{t_k, (t_{k+1} - t_k)t} - \mathbb{E} \left[ W_{t_k, (t_{k+1} - t_k)t} \right| \int_0^1 \left( (t_{k+1} - t_k)t \right)^l dW_{(t_{k+1} - t_k)t}, 0 \le l \le m \right] \right) \end{cases}_{t \in [0, 1]},$$

are independent centered Gaussian processes. Hence, the *i*-th and *j*-th coordinates of

$$\left\{\frac{1}{\sqrt{t_{k+1}-t_k}}\Big(W_{t_k,t(t_{k+1}-t_k)}-\widetilde{W}^n_{t_k,t(t_{k+1}-t_k)}\Big)\right\}_{t\in[0,1]},$$

are independent centered Gaussian processes. As before, we note that any additional information in  $\mathcal{G}_n$  is independent of W and does not affect conditional expectations. Therefore, it follows that  $\{W_{t_k,t}^i\}_{t \in [t_k,t_{k+1}]}$  and  $\{W_{t_k,t}^j\}_{t \in [t_k,t_{k+1}]}$  are  $\mathcal{F}_n$ -conditionally independent processes and

$$\widetilde{W}_t^n = \mathbb{E}\big[W_{t_k,t} | \mathcal{F}_n\big].$$

Thus for a sequence of uniform partitions  $\{\Delta_m\}$  of [s, t] with mesh $(\Delta_m) \to 0$ , we have

(2.14) 
$$\mathbb{E}\left[\sum_{t_l\in\Delta_m} W_{s,t_l}^i W_{t_l,t_{l+1}}^j \middle| \mathcal{F}_n\right] = \sum_{t_l\in\Delta_m} \mathbb{E}\left[W_{s,t_l}^i \middle| \mathcal{F}_n\right] \mathbb{E}\left[W_{t_l,t_{l+1}}^j \middle| \mathcal{F}_n\right] = \sum_{t_l\in\Delta_m} \left(\widetilde{W}_{s,t_l}^n\right)^i \left(\widetilde{W}_{t_l,t_{l+1}}^n\right)^j.$$

Since Itô integrals can be defined as the limit of their Riemann sum approximations, with convergence taking place is in the standard  $L^2(\mathbb{P})$  sense, it follows that

$$\mathbb{E}\bigg[\bigg(\mathbb{E}\bigg[\sum_{t_l\in\Delta_m} W_{s,t_l}^i W_{t_l,t_{l+1}}^j \Big| \mathcal{F}_n\bigg] - \mathbb{E}\bigg[\int_s^t W_{s,u}^i \, dW_u^j \Big| \mathcal{F}_n\bigg]\bigg)^2\bigg] \\ \leqslant \mathbb{E}\bigg[\bigg(\mathbb{E}\bigg[\sum_{t_l\in\Delta_m} W_{s,t_l}^i W_{t_l,t_{l+1}}^j\bigg] - \mathbb{E}\bigg[\int_s^t W_{s,u}^i \, dW_u^j\bigg]\bigg)^2\bigg] \to 0,$$

by Jensen's inequality and the tower property. Since  $W^i$  and  $W^j$  are independent, we also observe that the Itô and Stratonovich formulations of the integral coincide. That is,  $\int_s^t W_{s,u}^i dW_u^j = \int_s^t W_{s,u}^i \circ dW_u^j$ . Hence, by taking the limit of (2.14), we have

(2.15) 
$$\mathbb{E}\left[\int_{s}^{t} W_{s,u}^{i} \circ dW_{u}^{j} \middle| \mathcal{F}_{n}\right] = \int_{s}^{t} \left(\widetilde{W}_{s,u}^{n}\right)^{i} d\left(\widetilde{W}_{u}^{n}\right)^{j}.$$

As before, since  $\{F_n\}$  is a filtration, it follows that the sequence (2.15) is a martingale. Thus, by applying Doob's martingale convergence theorem to these integrals, we have

(2.16) 
$$\int_{s}^{t} \left(\widetilde{W}_{s,u}^{n}\right)^{i} d\left(\widetilde{W}_{u}^{n}\right)^{j} \to \mathbb{E}\left[\int_{s}^{t} W_{s,u}^{i} \circ dW_{u}^{j} \middle| \mathcal{F}_{\infty}\right] = \int_{s}^{t} W_{s,u}^{i} \circ dW_{u}^{j},$$

as  $n \to \infty$  almost surely and in  $L^2(\mathbb{P})$ .

We now introduce the notions of rough paths and p-variation. These are the two key ingredients in the Universal Limit Theorem [48], which establishes SDE (or CDE) solutions as continuous functions of the Brownian motion (or the driving rough path).

DEFINITION 2.9. A rough path is a continuous function  $\mathbf{X} : \Delta_T \to \mathbb{R} \oplus \mathbb{R}^d \oplus \mathbb{R}^{d \times d}$ where  $\Delta_T := \{(s,t) : 0 \leq s \leq t \leq T\}$ . The rough path  $\mathbf{X} = (1, X, \mathbb{X})$  is said to be  $\alpha$ -Hölder continuous if  $\|\mathbf{X}\|_{\alpha - H\"{o}l;[0,T]} < \infty$  where the norm  $\|\cdot\|_{\alpha - H\"{o}l;[s,t]}$  is defined as

(2.17) 
$$\|\boldsymbol{X}\|_{\alpha-H\ddot{o}l;[s,t]} := \max\left(\sup_{s\leqslant u < v\leqslant t} \frac{\|X_{u,v}\|_2}{|v-u|^{\alpha}}, \sup_{s\leqslant u < v\leqslant t} \frac{\|\mathbb{X}_{u,v}\|_2}{|v-u|^{2\alpha}}\right)$$

is finite. Similarly, we define the  $\alpha$ -Hölder metric between rough paths X and Y as

(2.18) 
$$d_{\alpha-H\ddot{o}l;[s,t]}(\boldsymbol{X},\boldsymbol{Y}) := \max\left(\sup_{s\leqslant u < v\leqslant t} \frac{\|X_{u,v} - Y_{u,v}\|_2}{|v-u|^{\alpha}}, \sup_{s\leqslant u < v\leqslant t} \frac{\|\mathbb{X}_{u,v} - \mathbb{Y}_{u,v}\|_2}{|v-u|^{2\alpha}}\right).$$

For  $p \in [2,3)$ , we define the p-variation metric between rough paths X and Y as

(2.19) 
$$d_{p\text{-}var;[0,T]}(\boldsymbol{X},\boldsymbol{Y}) := \max\left(\left(\sup_{\mathcal{D}}\sum_{k} \|X_{t_{k},t_{k+1}} - Y_{t_{k},t_{k+1}}\|_{2}^{p}\right)^{\frac{1}{p}}, \\ \left(\sup_{\mathcal{D}}\sum_{k} \|\mathbb{X}_{t_{k},t_{k+1}} - \mathbb{Y}_{t_{k},t_{k+1}}\|_{2}^{\frac{p}{2}}\right)^{\frac{2}{p}}\right).$$

Remark 2.10. In the literature, the paths X and Y would also be referred to as p-rough paths where  $p \in [2,3)$ . Although there do exist p-rough paths with  $p \ge 3$  (such as fractional Brownian motion with Hurst parameter  $H \le \frac{1}{3}$ ), for the purposes of this paper, it will be sufficient to only consider the above setting where  $p \in (2,3)$ .

Remark 2.11. From the above definitions, it is straightforward to show that

$$d_{\frac{1}{\alpha}-\operatorname{var};[0,\mathrm{T}]}(\boldsymbol{X},\boldsymbol{Y}) \leq \max(T^{\alpha},T^{2\alpha}) d_{\alpha-\operatorname{Höl};[0,\mathrm{T}]}(\boldsymbol{X},\boldsymbol{Y}).$$

DEFINITION 2.12. We define a "homogeneous" norm  $\|\cdot\|$  for path increments as

(2.20) 
$$\|\boldsymbol{X}\| := \max\left(\|\boldsymbol{X}\|_2, \|\boldsymbol{X}\|_2^{\frac{1}{2}}\right),$$

where  $\mathbf{X} = (1, X, \mathbb{X}) \in \mathbb{R} \oplus \mathbb{R}^d \oplus \mathbb{R}^{d \times d}$  and  $\|\cdot\|_2$  denotes the standard Euclidean norm.

Now we have introduced some key definitions from rough path theory, we can strengthen the pointwise and mean squared convergence established in Theorem 2.6. To do so, we will largely follow the rough path analysis detailed in [68, Section 13.3.2].

THEOREM 2.13 (Rough path convergence of nested unbiased approximations). Let  $\widetilde{\boldsymbol{W}}^n = \{\widetilde{\boldsymbol{W}}^n_{s,t}\}_{0 \leq s \leq t \leq T}$  be the rough path defined by the piecewise polynomial  $\widetilde{W}^n$  as

$$\widetilde{\boldsymbol{W}}_{s,t}^{n} := \left(1, \widetilde{W}_{s,t}^{n}, \widetilde{\mathbb{W}}_{s,t}^{n}\right),$$

where  $\widetilde{W}_{s,t}^n := \left\{ (\widetilde{W}_t^n)^i - (\widetilde{W}_s^n)^i \right\}_{1 \leq i \leq d}$  and  $\widetilde{W}_{s,t}^n := \left\{ \int_s^t (\widetilde{W}_{s,u}^n)^i d(\widetilde{W}_u^n)^j \right\}_{1 \leq i,j \leq d}$ . Then

(2.21) 
$$d_{\alpha-H\"ol;[0,T]}(\widetilde{\boldsymbol{W}}^{n},\boldsymbol{W}) \to 0,$$

as  $n \to \infty$  almost surely, where  $\mathbf{W} = \{\mathbf{W}_{s,t}\}_{0 \leq s \leq t \leq T}$  is the "Stratonovich enhanced" Brownian motion given by

$$\boldsymbol{W}_{s,t} := \left(1, W_{s,t}, \mathbb{W}_{s,t}\right),$$

with  $W_{s,t} := \left\{ W_t^i - W_s^i \right\}_{1 \leqslant i \leqslant d} \in \mathbb{R}^d$  and  $\mathbb{W}_{s,t} := \left\{ \int_s^t W_{s,u}^i \circ dW_u^j \right\}_{1 \leqslant i,j \leqslant d} \in \mathbb{R}^{d \times d}$ .

*Proof.* Let  $\beta \in (\alpha, \frac{1}{2})$  denote a fixed constant. Then by [68, Corollary 13.14 (i)], there exists c > 0 (depending on  $\beta$  and T) such that

$$\mathbb{E}\bigg[\exp\left(c\|\boldsymbol{W}\|_{\beta-\mathrm{H\"ol};[0,\mathrm{T}]}^{2}\right)\bigg] < \infty$$

In particular, this implies that there exists a non-negative random variable  $C_1$  with  $\|C_1\|_{L^2(\mathbb{P})} < \infty$  so that, for almost all  $\omega \in \Omega$ , the increments of the path  $W(\omega)$  satisfy

$$\left\|\boldsymbol{W}_{s,t}\right\|^2 \leqslant C_1 |t-s|^{2\beta},$$

for  $0 \leq s < t \leq T$  where  $\|\cdot\|$  is the norm previously given by (2.20) in Definition 2.12. Hence, there exists a non-negative random variable  $C_2$  with  $\|C_2\|_{L^2(\mathbb{P})} < \infty$  and

(2.22) 
$$-C_2|t-s|^{2\beta} \leqslant \int_s^t W_{s,u}^i \circ dW_u^j \leqslant C_2|t-s|^{2\beta}.$$

Since  $\mathbb{E}[C_2|\mathcal{F}_n]$  is a martingale, we can apply Doob's maximal inequality to give

(2.23) 
$$\left\| \sup_{0 \le m \le n} \left( \mathbb{E} \left[ C_2 \left| \mathcal{F}_m \right] \right) \right\|_{L^2(\mathbb{P})} \le 2 \left\| \mathbb{E} \left[ C_2 \left| \mathcal{F}_n \right] \right\|_{L^2(\mathbb{P})} \le 2 \| C_2 \|_{L^2(\mathbb{P})}.$$

15

Taking the expectation of the inequality (2.22) conditional on the  $\sigma$ -algebra  $\mathcal{F}_n$  and applying equation (2.15) gives

(2.24) 
$$-C_3|t-s|^{2\beta} \leqslant \int_s^t \left(\widetilde{W}_{s,u}^n\right)^i d\left(\widetilde{W}_u^n\right)^j \leqslant C_3|t-s|^{2\beta},$$

where  $C_3 := \sup_{n \ge 0} \left( \mathbb{E} [C_2 | \mathcal{F}_n] \right)$  is finite almost surely due to the upper bound (2.23). Therefore, by setting  $C_4 := d^2 C_3$ , we have

(2.25) 
$$\sup_{n \ge 0} \left\| \int_{s}^{t} \left( \widetilde{W}_{s,u}^{n} \right) \otimes d\left( \widetilde{W}_{u}^{n} \right) \right\|_{2} \leq C_{4} |t-s|^{2\beta}.$$

By a similar martingale argument, the exists a positive random variable  $C_5$  such that

$$\sup_{n \ge 0} \left\| \widetilde{W}_{s,t}^n \right\|_2 \le C_5 |t-s|^{\beta},$$

where  $C_5$  is finite almost surely. Thus, letting  $C_6 := \max\left(\sqrt{C_1}, \sqrt{C_4}, C_5\right)$ , we have

(2.26) 
$$\left\|\boldsymbol{W}_{s,t}\right\| \leq C_6 |t-s|^{\beta},$$

(2.27) 
$$\left\|\widetilde{\boldsymbol{W}}_{s,t}^{n}\right\| = \max\left(\left\|\widetilde{W}_{s,t}^{n}\right\|_{2}, \left\|\int_{s}^{t}\left(\widetilde{W}_{s,u}^{n}\right)\otimes d\left(\widetilde{W}_{u}^{n}\right)\right\|_{2}^{\frac{1}{2}}\right) \leqslant C_{6}|t-s|^{\beta},$$

for  $n \ge 0$ . Letting  $\mathbb{W}_{s,t}$  and  $\widetilde{\mathbb{W}}_{s,t}^n$  denote the second iterated integrals of W and  $\widetilde{W}^n$  over the interval [s, t], we note that

$$\frac{\left\|W_{s,t} - \widetilde{W}_{s,t}^{n}\right\|_{2}}{|t-s|^{\alpha}} \leqslant \left(\frac{\left\|W_{s,t} - \widetilde{W}_{s,t}^{n}\right\|_{2}}{|t-s|^{\beta}}\right)^{\frac{\alpha}{\beta}} \left(\sup_{0 \leqslant s \leqslant t \leqslant T} \left\|W_{s,t} - \widetilde{W}_{s,t}^{n}\right\|_{2}\right)^{1-\frac{\alpha}{\beta}},$$
$$\frac{\left\|\mathbb{W}_{s,t} - \widetilde{W}_{s,t}^{n}\right\|_{2}}{|t-s|^{2\alpha}} \leqslant \left(\frac{\left\|\mathbb{W}_{s,t} - \widetilde{W}_{s,t}^{n}\right\|_{2}}{|t-s|^{2\beta}}\right)^{\frac{\alpha}{\beta}} \left(\sup_{0 \leqslant s \leqslant t \leqslant T} \left\|\mathbb{W}_{s,t} - \widetilde{W}_{s,t}^{n}\right\|_{2}\right)^{1-\frac{\alpha}{\beta}}.$$

By (2.26) and (2.27), there exists an almost surely finite random variable  $C_7 > 0$ , which can bound the first terms on the right hand sides.

(2.28) 
$$\frac{\|W_{s,t} - \widetilde{W}_{s,t}^n\|_2}{|t-s|^{\alpha}} \leq C_7 \left(\sup_{0 \leq s \leq t \leq T} \|W_{s,t} - \widetilde{W}_{s,t}^n\|_2\right)^{1-\frac{\alpha}{\beta}},$$

(2.29) 
$$\frac{\left\|\mathbb{W}_{s,t} - \mathbb{W}_{s,t}^n\right\|_2}{|t-s|^{2\alpha}} \leq C_7 \left(\sup_{0 \leq s \leq t \leq T} \left\|\mathbb{W}_{s,t} - \widetilde{\mathbb{W}}_{s,t}^n\right\|_2\right)^{1-\frac{1}{\beta}}$$

Recall that by Theorems 2.6 and 2.8, it was shown that  $\widetilde{\boldsymbol{W}}_{s,t}^{n}$  converges to  $\boldsymbol{W}_{s,t}$  almost surely and in  $L^{2}(\mathbb{P})$  as  $n \to \infty$ . Therefore, it immediately follows from the uniform Hölder bounds (2.26) and (2.27), that any subsequence of  $\{(W - \widetilde{W}^{n}, W - \widetilde{W}^{n})\}_{n \geq 0}$  is bounded and uniformly equicontinuous in (s,t) and converges to zero almost surely. Hence, by the Arzelà–Ascoli theorem, there will exist a further subsequence which converges uniformly to zero. Hence, the right hand sides of both (2.28) and (2.29) will converge to zero almost surely, leading to  $d_{\alpha-\text{Höl};[0,\text{T}]}(\widetilde{\boldsymbol{W}}^{n}, \boldsymbol{W}) \to 0$  as required.  $\Box$ 

COROLLARY 2.14 (Rough path convergence of nested unbiased approximations). Let  $\boldsymbol{W}$  and  $\widetilde{\boldsymbol{W}}^n$  denote the rough path lifts of W and its approximation  $\widetilde{W}^n$ . Then

(2.30) 
$$d_{p\text{-}var;[0,T]}(\widetilde{\boldsymbol{W}}^{n},\boldsymbol{W}) \to 0,$$

as  $n \to \infty$  almost surely.

*Proof.* The result follows as  $d_{\frac{1}{\alpha}-\operatorname{var};[0,T]}(\cdot,\cdot) \leq \max(T^{\alpha}, T^{2\alpha}) d_{\alpha-\operatorname{Höl};[0,T]}(\cdot,\cdot).$   $\Box$ 

### J. FOSTER AND A. JELINČIČ

2.2. Convergence of "polynomial driven" CDEs with small local errors. Since we have shown that  $\{\widetilde{\boldsymbol{W}}^n\}$  converges to the Stratonovich Brownian motion  $\boldsymbol{W}$ , we can apply the Universal Limit Theorem from rough path theory [48, Theorem 5.3] to show that the associated sequence of CDEs (with the *n*-th CDE driven by  $\widetilde{\boldsymbol{W}}^n$ ) converges to the CDE driven by  $\boldsymbol{W}$  – which is simply just the Stratonovich SDE (1.1). To this end, we use the following definition to impose smoothness on the vector fields.

DEFINITION 2.15. We say that a function  $f : \mathbb{R}^e \to \mathbb{R}^e$  is  $\operatorname{Lip}(\gamma)$  with  $\gamma > 1$ if it is bounded with  $[\gamma]$  bounded derivatives, the last being Hölder continuous with exponent  $(\gamma - [\gamma])$ . We say that f is  $\operatorname{Lip}(1)$  if it is bounded and Lipschitz continuous.

THEOREM 2.16 (Rough path convergence of piecewise polynomial driven CDEs). Suppose that the vector fields f and  $\{g_i\}$  of the SDE (1.1) are  $\operatorname{Lip}(\gamma)$  with  $\gamma \in (2,3)$ . Let  $\{D_n\}$  denote a sequence of nested partitions obtained from the Brownian motion using an enhancement function  $\mathcal{E}$ , with  $q \ge 0$  fixed, and a refinement function  $\mathcal{R}$ , according to (2.4). For each (random) partition  $D_n$ , let  $\widetilde{W}^n$  be the associated piecewise degree q polynomial approximation that matches the increments and integrals in (2.1). Let y be the solution of the SDE (1.1) and, for  $n \ge 0$ , we define  $\widetilde{y}^n$  as the solution of

(2.31) 
$$d\widetilde{y}_t^n = f(\widetilde{y}_t^n)dt + g(\widetilde{y}_t^n)d\widetilde{W}_t^n,$$
$$\widetilde{y}_0^n = y_0.$$

Then, letting y and  $\tilde{y}^n$  denote the unique (Stratonovich) lifts of y and  $\tilde{y}^n$ , we have

$$(2.32) d_{p-var;[0,T]}(\widetilde{\boldsymbol{y}}^n, \boldsymbol{y}) \to 0$$

as  $n \to \infty$  almost surely, where  $p \in (2, \gamma)$ .

*Proof.* The result follows from the universal limit theorem and Corollary 2.14. □

Remark 2.17. Technically speaking, we define the rough path  $\boldsymbol{y} = (1, y, y)$  as the limit of  $\{\tilde{\boldsymbol{y}}^n\}$  and, by the universal limit theorem,  $\boldsymbol{y}$  can be taken to be the solution of the rough differential equation driven by Stratonovich enhanced Brownian motion. In particular, from  $\boldsymbol{y}$ , we can define a process  $y = \{y_t\}$  as  $y_t = y_0 + y_{0,t}$  for  $t \in [0,T]$ , which coincides almost surely with the usual solution of the Stratonovich SDE (1.1).

In practice, it is unlikely that each polynomial driven CDE can be solved exactly. However, if our numerical method approximates the CDE (2.31) with  $o(h_k)$  accuracy, then we would still expect convergence. For example, Heun's method (1.14), is clearly obtained by discretizing the "piecewise linear" CDE using a second order ODE solver.

To make this precise, we follow the error analysis presented in [21, Section 4.2]. However, since rough path theory was formulated after [21], we can employ a result known as *Davie's estimate* to establish the following Lipschitz property of CDE flows.

THEOREM 2.18 (CDEs driven by Brownian polynomials produce Lipschitz flows). As before, let  $\{\widetilde{W}^n\}$  be the piecewise polynomial approximations of W corresponding to the sequence of nested partitions  $\{D_n\}$ . For each  $n \ge 0$ , we consider two solutions  $\widetilde{x}^n$  and  $\widetilde{y}^n$  of the CDE (2.31), driven by the same  $\widetilde{W}^n$ , but with different initial values. Then there exists a positive random variable L with  $L < \infty$  almost surely, such that

(2.33) 
$$\|\widetilde{x}_t^n - \widetilde{y}_t^n\|_2 \leq L \|\widetilde{x}_s^n - \widetilde{y}_s^n\|_2,$$

for all  $0 \leq s \leq t \leq T$  and  $n \geq 0$ .

17

*Proof.* By applying Davie's estimate [68, Theorem 10.29] to the CDE solutions,  $\tilde{x}^n$  and  $\tilde{y}^n$ , using the control function w(s,t) := t - s and  $\alpha = \frac{1}{n} \in (\frac{1}{3}, \frac{1}{2})$ , we obtain

(2.34) 
$$d_{\alpha-\mathrm{H\"ol};[\mathbf{s},\mathbf{t}]}(\widetilde{x}^n,\widetilde{y}^n) \leq C\nu \exp\left(C\nu^p(t-s)\right) \|\widetilde{x}^n_s - \widetilde{y}^n_s\|_2,$$

where  $\nu$  is the so-called "Lip( $\gamma$ ) norm" of CDE vector fields rescaled by  $\|\widetilde{W}^n\|_{\alpha-\text{H\"ol};[s,t]}$ (as we need to scale the driving path by  $\|\widetilde{W}^n\|_{\alpha-\text{H\"ol};[s,t]}^{-1}$  to apply [68, Theorem 10.29]), and the constant C only depends on  $(p, \gamma)$ , which are fixed and satisfy 2 . $The result now follows from (2.34) as <math>\{\|\widetilde{W}^n\|_{\alpha-\text{H\"ol};[0,T]}\}$  can be uniformly bounded by an almost surely finite positive random variable (see the proof of Theorem 2.13).

Using the Lipschitz estimate (2.33), we now adapt the proof of [21, Theorem 4.3] to give our main result on pathwise convergence for adaptive approximations close to polynomial driven CDEs – such as Heun's method (1.14) and the SRK method (1.16).

THEOREM 2.19 (Pathwise convergence of strong adaptive SDE approximations). For each partition  $D_n$ , appearing in Theorems 2.16 and 2.18, let  $Y^n = \{Y_k^n\}$  be a numerical solution of (1.1) computed at times  $\{0 = t_0^n < t_1^n < \cdots < t_{K_n}^n = T\} \subseteq D_n$ , with the initial condition  $Y_0^n := y_0$  such that, for all  $k \in \{0, 1, \cdots, K_n - 1\}$ , we have

(2.35) 
$$\left\|Y_{k+1}^n - \Phi_{t_k^n, t_{k+1}^n}^n(Y_k^n)\right\|_2 \le w(t_k^n, t_{k+1}^n),$$

where  $\Phi_{a,b}^n(y) \in \mathbb{R}^e$  denotes the solution  $\widetilde{x}_b^n$  at time b of the following CDE on [a, b],

(2.36) 
$$d\widetilde{x}_{t}^{n} = f(\widetilde{x}_{t}^{n})dt + g(\widetilde{x}_{t}^{n})d\widetilde{W}_{t}^{n},$$
$$\widetilde{x}_{a}^{n} := y,$$

driven by the same piecewise polynomial  $\{\widetilde{W}_t^n\}_{t\in[0,T]}$  considered in Theorem 2.16 and

(2.37) 
$$\sum_{k=0}^{K_n-1} w(t_k^n, t_{k+1}^n) \to 0,$$

as  $n \to \infty$  almost surely. Then the numerical solutions  $\{Y^n\}$  converges pathwise, i.e.

(2.38) 
$$\sup_{0 \le k \le K_n} \|Y_k^n - y_{t_k^n}\|_2 \to 0,$$

as  $n \to \infty$  almost surely.

*Proof.* Since  $\Phi_{t_k^n, t_k^n}^n(Y_k^n) = Y_k^n$  and  $(t_0^n, Y_0^n) = (0, y_0)$  for all  $n \ge 0$ , by expressing  $Y_k^n - \Phi_{0, t_k^n}^n(y_0)$  as a telescoping sum and applying the triangle inequality, we obtain

$$\left\|Y_{k}^{n}-\Phi_{0,t_{k}^{n}}^{n}(y_{0})\right\|_{2} \leqslant \sum_{i=0}^{k-1} \left\|\Phi_{t_{i+1}^{n},t_{k}^{n}}^{n}(Y_{i+1}^{n})-\Phi_{t_{i}^{n},t_{k}^{n}}^{n}(Y_{i}^{n})\right\|_{2}.$$

As  $\Phi^n$  denotes the solution of a differential equation, it has a natural flow property,  $\Phi^n_{s,t} = \Phi^n_{u,t} \circ \Phi^n_{s,u}$  for  $u \in [s,t]$ . In particular, by applying this to  $\Phi^n_{t^n_t,t^n_t}(Y^n_i)$ , we have

$$\left\|Y_{k}^{n}-\Phi_{0,t_{k}^{n}}^{n}(y_{0})\right\|_{2} \leqslant \sum_{i=0}^{k-1} \left\|\Phi_{t_{i+1}^{n},t_{k}^{n}}^{n}(Y_{i+1}^{n})-\Phi_{t_{i+1}^{n},t_{k}^{n}}^{n}\left(\Phi_{t_{i}^{n},t_{i+1}^{n}}^{n}(Y_{i}^{n})\right)\right\|_{2}.$$

Thus, applying the Lipschitz estimate (2.33) from Theorem 2.18 to each  $\Phi_{t_{i+1},t_{k}}^{n}$  gives

$$\begin{aligned} \left\|Y_{k}^{n} - \Phi_{0,t_{k}^{n}}^{n}(y_{0})\right\|_{2} &\leqslant \sum_{i=0}^{k-1} L \left\|Y_{i+1}^{n} - \Phi_{t_{i}^{n},t_{i+1}^{n}}^{n}(Y_{i}^{n})\right\|_{2} \\ &\leqslant L \sum_{i=0}^{k-1} w(t_{i}^{n},t_{i+1}^{n}), \end{aligned} \qquad [by (2.35)]. \end{aligned}$$

By the assumption (2.37) that the "error bounds"  $\{w(t_i^n, t_{i+1}^n)\}$  are small, we see that

(2.39) 
$$\sup_{0 \le k \le K_n} \left\| Y_k^n - \Phi_{0,t_k^n}^n(y_0) \right\|_2 \le L \sum_{i=0}^{K_n-1} w(t_i^n, t_{i+1}^n) \to 0.$$

as  $n \to \infty$  almost surely. The result now immediately follows by Theorem 2.13, which established the rough path convergence of  $\{\Phi_{0,t}^n(y_0)\}$  to the Stratonovich solution.

Remark 2.20. If the local errors can be bounded by  $w(t_k^n, t_{k+1}^n) = \widetilde{w}(t_{k+1}^n - t_k^n)$  for some function  $\widetilde{w}$  with  $\widetilde{w}(h) \sim o(h)$ , then  $\sum_{k=0}^{K_n-1} w(t_k^n, t_{k+1}^n) \to 0$  is trivially satisfied. This will be the case for standard numerical methods, such as Heun's method (1.14), where under suitable regularity conditions, we have  $\widetilde{w}(h) \sim O(h^{\frac{3}{2}})$  in an  $L^2(\mathbb{P})$  sense.

**2.3.** Numerical methods for SDEs viewed as approximations of CDEs. To conclude the section, we will argue why the convergence criterion (2.35) is satisfied the no-area Milstein, Heun and stochastic Runge-Kutta methods given in Section 1.1. By considering Taylor expansions, we can show these numerical methods accurately approximate the CDE (2.36) so that our main result (Theorem 2.19) can be applied.

THEOREM 2.21 (Taylor expansion of CDEs driven by "Brownian" polynomials). Consider the controlled differential equation considered in Theorem 2.36 and given by

(2.40) 
$$d\widetilde{x}_t^n = f(\widetilde{x}_t^n)dt + g(\widetilde{x}_t^n)d\widetilde{W}_t^n$$

where  $f : \mathbb{R}^e \to \mathbb{R}^e$  and  $g : \mathbb{R}^e \to \mathbb{R}^{e \times d}$  are bounded twice continuously differentiable vector fields with bounded derivatives. Then the solution of (2.40) can be expanded as

$$\widetilde{x}_{t_{k+1}}^n = \widetilde{x}_{t_k}^n + f(\widetilde{x}_{t_k}^n)h_k + g(\widetilde{x}_{t_k}^n)W_k + g'(\widetilde{x}_{t_k}^n)g(\widetilde{x}_{t_k}^n)\int_{t_k}^{t_{k+1}}\int_{t_k}^t d\widetilde{W}_s^n \otimes d\widetilde{W}_t^n + R,$$

where, for any fixed  $\alpha \in (0, \frac{1}{2})$ , there an almost surely finite random variable C > 0such that the remainder term R satisfies  $||R||_2 \leq Ch_k^{3\alpha}$ , and hence  $o(h_k)$  when  $\alpha \geq \frac{1}{3}$ .

*Proof.* The solution of the CDE (2.40) can be expressed in integral form as

$$\widetilde{x}_{t_{k+1}}^n = \widetilde{x}_{t_k}^n + \int_{t_k}^{t_{k+1}} f(\widetilde{x}_t^n) dt + \int_{t_k}^{t_{k+1}} g(\widetilde{x}_t^n) d\widetilde{W}_t^n,$$

and, given a differentiable function  $F \in \mathcal{C}^1(\mathbb{R}^e)$ , we also have the following chain rule:

$$F(\widetilde{x}_t^n) = F(\widetilde{x}_{t_k}^n) + \int_{t_k}^t F'(\widetilde{x}_s^n) d\widetilde{x}_s^n.$$

Therefore, using the above equations, we can derive a Taylor expansion for the CDE.

$$\begin{split} \widetilde{x}_{t_{k+1}}^n &= \widetilde{x}_{t_k}^n + \int_{t_k}^{t_{k+1}} f(\widetilde{x}_{t_k}^n) + \int_{t_k}^t f'(\widetilde{x}_s^n) d\widetilde{x}_s^n dt + \int_{t_k}^{t_{k+1}} g(\widetilde{x}_{t_k}^n) + \int_{t_k}^t g'(\widetilde{x}_s^n) d\widetilde{x}_s^n d\widetilde{W}_t^n \\ &= \widetilde{x}_{t_k}^n + f(\widetilde{x}_{t_k}^n) h_k + g(\widetilde{x}_{t_k}^n) (\widetilde{W}_{t_{k+1}}^n - \widetilde{W}_{t_k}^n) \\ &+ \int_{t_k}^{t_{k+1}} \int_{t_k}^t f'(\widetilde{x}_s^n) f(\widetilde{x}_s^n) \, ds \, dt + \int_{t_k}^{t_{k+1}} \int_{t_k}^t f'(\widetilde{x}_s^n) g(\widetilde{x}_s^n) \, d\widetilde{W}_s^n \, dt \\ &+ \int_{t_k}^{t_{k+1}} \int_{t_k}^t f'(\widetilde{x}_s^n) g(\widetilde{x}_s^n) \, ds \, d\widetilde{W}_t^n + \int_{t_k}^{t_{k+1}} \int_{t_k}^t g'(\widetilde{x}_s^n) g(\widetilde{x}_s^n) \, d\widetilde{W}_s^n \, d\widetilde{W}_t^n \\ &= \widetilde{x}_{t_k}^n + f(\widetilde{x}_{t_k}^n) h_k + g(\widetilde{x}_{t_k}^n) W_k + g'(\widetilde{x}_{t_k}^n) g(\widetilde{x}_{t_k}^n) \int_{t_k}^{t_{k+1}} \int_{t_k}^t d\widetilde{W}_s^n \otimes d\widetilde{W}_t^n + R, \end{split}$$

where the remainder term R is given by

$$\begin{split} R &:= \int_{t_k}^{t_{k+1}} \int_{t_k}^t \left( g'(\widetilde{x}_s^n) g(\widetilde{x}_s^n) - g'(\widetilde{x}_{t_k}) g(\widetilde{x}_{t_k}^n) \right) d\widetilde{W}_s^n d\widetilde{W}_t^n \\ &+ \int_{t_k}^{t_{k+1}} \int_{t_k}^t f'(\widetilde{x}_s^n) g(\widetilde{x}_s^n) d\widetilde{W}_s^n dt + \int_{t_k}^{t_{k+1}} \int_{t_k}^t f'(\widetilde{x}_s^n) g(\widetilde{x}_s^n) ds d\widetilde{W}_t^n \\ &+ \int_{t_k}^{t_{k+1}} \int_{t_k}^t f'(\widetilde{x}_s^n) f(\widetilde{x}_s^n) ds dt. \end{split}$$

As  $\widetilde{W}^n$  is a polynomial on  $[t_k, t_{k+1}]$ , we can apply the chain rule to rewrite any " $d\widetilde{W}_t^n$ " integral as an " $(\widetilde{W}_t^n)'dt$ " integral. This allows the first integral to be estimated as

$$\begin{split} \left\| \int_{t_k}^{t_{k+1}} \int_{t_k}^t \left( g'(\widetilde{x}_s^n) g(\widetilde{x}_s^n) - g'(\widetilde{x}_{t_k}) g(\widetilde{x}_{t_k}^n) \right) d\widetilde{W}_s^n d\widetilde{W}_t^n \right\|_2 \\ & \leq \int_{t_k}^{t_{k+1}} \int_{t_k}^t \left\| \left( g'(\widetilde{x}_s^n) g(\widetilde{x}_s^n) - g'(\widetilde{x}_{t_k}) g(\widetilde{x}_{t_k}^n) \right) \left( \widetilde{W}_s^n \right)' \left( \widetilde{W}_t^n \right)' \right\|_2 ds dt \\ & \leq \int_{t_k}^{t_{k+1}} \int_{t_k}^t \left\| g'(\widetilde{x}_s^n) g(\widetilde{x}_s^n) - g'(\widetilde{x}_{t_k}) g(\widetilde{x}_{t_k}^n) \right\| \left\| \left( \widetilde{W}_s^n \right)' \right\|_2 \left\| \left( \widetilde{W}_t^n \right)' \right\|_2 ds dt \\ & \leq \frac{1}{2} h_k^2 \| g'(\cdot) g(\cdot) \|_{\operatorname{Lip}(1)} \sup_{t \in [t_k, t_{k+1}]} \| \widetilde{x}_t^n - \widetilde{x}_{t_k}^n \|_2 \sup_{t \in [t_k, t_{k+1}]} \| \left( \widetilde{W}_t^n \right)' \|_2^2. \end{split}$$

We estimate the term involving  $\tilde{x}^n$  using the integral form of the CDE solution as

$$\begin{aligned} \|\widetilde{x}_t^n - \widetilde{x}_{t_k}^n\|_2 &\leqslant \int_{t_k}^t \|f(\widetilde{x}_s^n)\|_2 ds + \int_{t_k}^t \|g(\widetilde{x}_s^n)\|_2 \|(\widetilde{W}_s^n)'\|_2 ds \\ &\leqslant \Big(\|f\|_{\infty} + \|g\|_{\infty} \sup_{t \in [t_k, t_{k+1}]} \|(\widetilde{W}_t^n)'\|_2\Big) h_k. \end{aligned}$$

Due to the  $\alpha$ -Hölder continuity of W and Brownian scaling, we see that  $\{\widetilde{W}_t^n\}_{t \in [t_k, t_{k+1}]}$  is a linear function of  $\mathcal{W}_k$  that scales as  $O(h_k^{\alpha})$ . For example, in the low-degree cases,

$$\widetilde{W}_{t_k,t}^n = \frac{t - t_k}{h_k} W_k, \qquad (\text{degree 1})$$
$$\widetilde{W}_{t_k,t}^n = \frac{t - t_k}{h_k} W_k + \frac{6(t_{k+1} - t)(t - t_k)}{h_k^2} H_k. \quad (\text{degree 2})$$

Thus, since  $\widetilde{W}^n$  is a polynomial in t, the derivative  $(\widetilde{W}^n)'$  scales as  $O(h_k^{\alpha-1})$  and so

$$\begin{split} & \left\| \int_{t_{k}}^{t_{k+1}} \int_{t_{k}}^{t} \left( g'(\widetilde{x}_{s}^{n}) g(\widetilde{x}_{s}^{n}) - g'(\widetilde{x}_{t_{k}}) g(\widetilde{x}_{t_{k}}^{n}) \right) d\widetilde{W}_{s}^{n} d\widetilde{W}_{t}^{n} \right\|_{2} \\ & \leq \frac{1}{2} h_{k}^{3} \|g'g\|_{\operatorname{Lip}(1)} \Big( \|f\|_{\infty} + \|g\|_{\infty} \sup_{t \in [t_{k}, t_{k+1}]} \left\| \left(\widetilde{W}_{t}^{n}\right)' \right\|_{2} \Big) \sup_{t \in [t_{k}, t_{k+1}]} \left\| \left(\widetilde{W}_{t}^{n}\right)' \right\|_{2}^{2} \sim O(h^{3\alpha}). \end{split}$$

As  $(\widetilde{W}^n)' \sim O(h_k^{\alpha-1})$ , it is straightforward to show the middle terms in R are  $O(h^{\alpha+1})$ . Finally, the fourth term in R is clearly  $O(h_k^2)$  due to the boundedness of f and f'.

Remark 2.22. By equation (2.15), which was shown in the proof of Theorem 2.8, we see that the polynomial iterated integrals are optimal unbiased approximations as

(2.42) 
$$\int_{t_k}^{t_{k+1}} \int_{t_k}^t d\widetilde{W}_s^n \otimes d\widetilde{W}_t^n = \mathbb{E}\bigg[\int_{t_k}^{t_{k+1}} W_{t_k,t} \otimes \circ dW_t \,\Big|\, \mathcal{W}_k\bigg].$$

Therefore, it is clear that the convergence condition (1.3) originally presented in the introduction simply follows from Theorems 2.19 and 2.21 along with equation (2.42).

*Remark* 2.23. Whilst a general formula for polynomial approximations of second iterated integrals (or equivalent Lévy areas) is given in [17, Theorem 5.4], we only use

(2.43) 
$$\mathbb{E}\left[\int_{t_k}^{t_{k+1}} W_{t_k,t} \otimes \circ dW_t \,\Big|\, W_k\right] = \frac{1}{2} W_k^{\otimes 2},$$

(2.44) 
$$\mathbb{E}\left[\int_{t_k}^{t_{k+1}} W_{t_k,t} \otimes \circ dW_t \,\Big|\, W_k, H_k\right] = \frac{1}{2} W_k^{\otimes 2} + H_k \otimes W_k - W_k \otimes H_k.$$

With the Taylor expansion given by Theorem 2.21, it is now straightforward to see why our analysis is applicable to the numerical methods discussed in Section 1.1.

# • The "no-area" Milstein method (1.12)

As discussed in Remark 1.4, by using the Itô-Stratonovich correction, the no-area Milstein method can be rewritten as the following method for Stratonovich SDEs:

$$Y_{k+1} := Y_k + \widetilde{f}(Y_k)h_k + g(Y_k)W_k + \frac{1}{2}g'(Y_k)g(Y_k)W_k^{\otimes 2}$$

where  $\tilde{f}(\cdot) = f(\cdot) - \frac{1}{2} \sum_{i=1}^{d} g'_i(\cdot) g_i(\cdot)$  is the drift for the SDE in Stratonovich form. By (2.43), it is clear that the no-area Milstein method falls under our framework.

• Heun's method (1.14)

Our strategy is simply to show that Heun's method is close to no-area Milstein. Using the Taylor expansion  $F(b) = F(a) + F'(a)c + \int_0^1 (1-r)F''(a+rc)(c,c)dr$  for  $a, b, c \in \mathbb{R}^e$  where c = b - a, and the notation  $F''(\cdot)(u \otimes v) \equiv F''(\cdot)(u, v)$ , we have

$$Y_{k+1} = Y_k + \frac{1}{2} \left( F(Y_k) + F(\widetilde{Y}_{k+1}) \right) \overline{W}_k$$
  

$$= Y_k + \frac{1}{2} F(Y_k) \overline{W}_k + \frac{1}{2} \left( F(Y_k) + F'(Y_k) (\widetilde{Y}_{k+1} - Y_k) + R \right) \overline{W}_k$$
  

$$= Y_k + F(Y_k) \overline{W}_k + \frac{1}{2} \left( F'(Y_k) F(Y_k) \overline{W}_k + R \right) \overline{W}_k$$
  

$$= Y_k + F(Y_k) \overline{W}_k + F'(Y_k) F(Y_k) \left( \frac{1}{2} \overline{W}_k^{\otimes 2} \right) + \frac{1}{2} R \overline{W}_k,$$
  

$$= \mathbb{E} \left[ \int_{t_k}^{t_{k+1}} (\overline{W}_t - \overline{W}_{t_k}) \otimes \circ d\overline{W}_t | \overline{W}_k \right]$$

where the remainder term R is given by

$$R = \int_0^1 (1-r)F'' (Y_k + (1-r)F(Y_k)\overline{W}_k) dr (F(Y_k)\overline{W}_k)^{\otimes 2}.$$

Since W is  $\alpha$ -Hölder continuous for  $\alpha \in (0, \frac{1}{2})$ , we can employ a similar argument as the proof of Theorem 2.21 to show that  $R \sim O(h_k^{2\alpha})$ , and thus  $R\overline{W}_k \sim O(h_k^{3\alpha})$ . Therefore, as the vector fields and their derivatives are bounded, by taking  $\alpha > \frac{1}{3}$ , we see that Heun's method has the desired expansion (up to a  $o(h_k)$  remainder).

## • SPaRK method $(1.16)^2$

Taylor expanding this numerical method is slightly more involved, but results in

$$\begin{aligned} Y_{k+1} &= Y_k + F(Y_k) \left( a \overline{W}_k + \overline{H}_k \right) + b F(Y_{k+\frac{1}{2}}) \overline{W}_k + F(Z_{k+1}) \left( a \overline{W}_k - \overline{H}_k \right) \\ &= Y_k + F(Y_k) \left( a \overline{W}_k + \overline{H}_k \right) + b \left( F(Y_k) + F'(Y_k) \left( Y_{k+\frac{1}{2}} - Y_k \right) + R_1 \right) \right) \overline{W}_k \\ &+ \left( F(Y_k) + F'(Y_k) \left( Z_{k+1} - Y_k \right) + R_2 \right) \left( a \overline{W}_k - \overline{H}_k \right) \\ &= Y_k + F(Y_k) \overline{W}_k + b F'(Y_k) F(Y_k) \left( \frac{1}{2} \overline{W}_k + \sqrt{3} \overline{H}_k \right) \otimes \overline{W}_k + b R_1 \overline{W}_k \\ &+ F'(Y_k) F(Y_k) \left( \overline{W}_k \otimes \left( a \overline{W}_k - \overline{H}_k \right) \right) + (R_2 + R_3) \left( a \overline{W}_k - \overline{H}_k \right) \\ &= Y_k + F(Y_k) \overline{W}_k + F'(Y_k) F(Y_k) \left( \frac{1}{2} \overline{W}_k^{\otimes 2} + \overline{H}_k \otimes \overline{W}_k - \overline{W}_k \otimes \overline{H}_k \right) + R, \\ &= \mathbb{E} \Big[ \int_{t_k}^{t_{k+1}} (\overline{W}_t - \overline{W}_{t_k}) \otimes o d \overline{W}_t | \overline{W}_k, \overline{H}_k \Big] \end{aligned}$$

where R is given by  $R := bR_1\overline{W}_k + (R_2 + R_3)(a\overline{W}_k - \overline{H}_k)$  with  $a := \frac{3-\sqrt{3}}{6}, b := \frac{\sqrt{3}}{3},$ 

$$R_{1} = \int_{0}^{1} (1-r)F''(Y_{k} + (1-r)(Y_{k+\frac{1}{2}} - Y_{k}))dr(Y_{k+\frac{1}{2}} - Y_{k})^{\otimes 2},$$
  

$$R_{2} = \int_{0}^{1} (1-r)F''(Y_{k} + (1-r)(Z_{k+1} - Y_{k}))dr(Z_{k+1} - Y_{k})^{\otimes 2},$$
  

$$R_{3} = (F(Y_{k+\frac{1}{2}}) - F(Y_{k}))\overline{W}_{k}.$$

Recall that for  $\alpha \in (0, \frac{1}{2})$ , Brownian motion is  $\alpha$ -Hölder continuous almost surely. That is, there exists an almost surely finite random variable C > 0 such that  $\|W_{s,t}\|_2 \leq C|t-s|^{\alpha}$ . Hence, space-time Lévy area is also  $\alpha$ -Hölder continuous as,

$$\|H_{s,t}\|_{2} = \left\|\frac{1}{t-s}\int_{s}^{t}W_{s,u}du - \frac{1}{2}W_{s,t}\right\|_{2} \leq C\left(\frac{1}{1+\alpha} + \frac{1}{2}\right)|t-s|^{\alpha}.$$

Using the  $\alpha$ -Hölder estimates for  $W_k$  and  $H_k$  along with the boundedness of F, F'and F'', it is straightforward to estimate each remainder term as  $||R_i||_2 \leq C_i h_k^{2\alpha}$ . We can produce estimates for  $R_1, R_2 \sim O(h_k^{2\alpha})$  using similar arguments as before. The third term  $R_3$  can simply be estimated using the Lipschitz continuity of F as

$$\begin{split} \|R_3\|_2 &\leq \|F(Y_{k+\frac{1}{2}}) - F(Y_k)\|_2 \|\overline{W}_k\|_2 \\ &\leq \|F\|_{\text{Lip}(1)} \|Y_{k+\frac{1}{2}} - Y_k\|_2 \|\overline{W}_k\|_2 \\ &\leq \|F\|_{\text{Lip}(1)} \|F\|_{\infty} \Big(\frac{1}{2} \|\overline{W}_k\|_2 + \sqrt{3} \|\overline{H}_k\|_2 \Big) \|\overline{W}_k\|_2. \end{split}$$

Thus, it follows that the remainder R is  $O(h_k^{3\alpha})$ , which is  $o(h_k)$  if we take  $\alpha > \frac{1}{3}$ .

<sup>&</sup>lt;sup>2</sup>Splitting Path Runge-Kutta method

3. Counterexample involving non-nested step sizes. In this brief section, we will present a simple counterexample showing that non-nested adaptive step sizes can potentially prevent the approximation converging to the desired SDE solution. The idea is simply that by selectively "skipping over" values of the Brownian path, we can induce a local O(h) bias with a positive expectation. These local biases can then propagate linearly to produce a global O(1) bias, also with positive expectation. We illustrate this phenomenon using Theorem 3.1, which details the counterexample.

THEOREM 3.1. For  $N \ge 1$ , we will define an approximation  $\{(X_k, Y_k)\}_{0 \le k \le N}$  of

(3.1)  
$$dx_t = dW_t^1,$$
$$dy_t = x_t \circ dW_t^2,$$
$$(x_0, y_0) = 0,$$

over [0,T] by  $(X_0,Y_0) := 0$  and for each  $k \ge 0$ , the numerical solution is propagated using either one step or two steps of Heun's method (depending on which is largest),

$$\begin{aligned} X_{k+1} &:= X_k + W_{t_{k+1}}^1 - W_{t_k}^1, \\ Y_{k+1} &:= \max\left(Y_k + \frac{1}{2} \left(W_{t_k}^1 + W_{t_{k+1}}^1\right) \left(W_{t_{k+1}}^2 - W_{t_k}^2\right), \\ Y_k + \frac{1}{2} \left(W_{t_k}^1 + W_{t_{k+\frac{1}{2}}}^1\right) W_{t_k, t_{k+\frac{1}{2}}}^2 + \frac{1}{2} \left(W_{t_{k+\frac{1}{2}}}^1 + W_{t_{k+1}}^1\right) W_{t_{k+\frac{1}{2}}, t_{k+1}}^2 \right), \end{aligned}$$

where  $t_k := kh$  with  $h = \frac{T}{N}$  denoting the coarsest step size. Then, at time T, we have

$$\mathbb{E}[y_T] = 0, \quad \mathbb{E}[Y_N] = \frac{1}{8}T.$$

*Proof.* Since the x-component of the SDE is solved exactly by Heun's method, we have  $X_k = W_{t_k}^1$  for all  $k \ge 0$ . Rearranging the second part of the y-component gives

$$\begin{split} Y_{k} &+ \frac{1}{2} \Big( W_{t_{k}}^{1} + W_{t_{k+\frac{1}{2}}}^{1} \Big) \Big( W_{t_{k+\frac{1}{2}}}^{2} - W_{t_{k}}^{2} \Big) + \frac{1}{2} \Big( W_{t_{k+\frac{1}{2}}}^{1} + W_{t_{k+1}}^{1} \Big) \Big( W_{t_{k+1}}^{2} - W_{t_{k+\frac{1}{2}}}^{2} \Big) \\ &= Y_{k} + \frac{1}{2} \Big( W_{t_{k}}^{1} + W_{t_{k+1}}^{1} \Big) \Big( W_{t_{k+1}}^{2} - W_{t_{k}}^{2} \Big) \\ &+ \frac{1}{2} \Big( W_{t_{k+\frac{1}{2}}}^{1} - W_{t_{k}}^{1} \Big) \Big( W_{t_{k+1}}^{2} - W_{t_{k+\frac{1}{2}}}^{2} \Big) - \frac{1}{2} \Big( W_{t_{k+1}}^{1} - W_{t_{k+\frac{1}{2}}}^{1} \Big) \Big( W_{t_{k+\frac{1}{2}}}^{2} - W_{t_{k}}^{2} \Big) \Big) \end{split}$$

Since the last line gives the difference between one and two steps of Heun's method, it is enough for us to compute its expected value (conditional on being non-negative).

$$\mathbb{E}[Y_{k+1}] = \mathbb{E}\left[Y_k + \frac{1}{2} (W_{t_k}^1 + W_{t_{k+1}}^1) (W_{t_{k+1}}^2 - W_{t_k}^2) + \max\left(0, \frac{1}{2} (W_{t_k, t_{k+\frac{1}{2}}}^1 W_{t_{k+\frac{1}{2}}, t_{k+1}}^2 - W_{t_{k+\frac{1}{2}}, t_{k+1}}^1 W_{t_k, t_{k+\frac{1}{2}}}^2)\right)\right]$$
$$= \mathbb{E}[Y_k] + \frac{1}{4} \mathbb{E}\left[|W_{t_k, t_{k+\frac{1}{2}}}^1 W_{t_{k+\frac{1}{2}}, t_{k+1}}^2 - W_{t_{k+\frac{1}{2}}, t_{k+1}}^1 W_{t_k, t_{k+\frac{1}{2}}}^2|\right]$$
$$= \mathbb{E}[Y_k] + \frac{1}{8} h \mathbb{E}\left[|Z_1 Z_2 - Z_3 Z_4|\right],$$

where  $Z_1, Z_2, Z_3, Z_4 \sim N(0, 1)$  are independent standard normal random variables. The result now follows as the expectation given above equals 1 by Theorem C.2.

For a counterexample using nested step sizes and the Euler-Maruyama method, which does not satisfy the convergence condition (1.3) in general, see [21, Section 4.1].

4. Numerical example. In this section, we will consider the well-known SABR model used in mathematical finance [7, 11, 25, 44]. The model describes the evolution of an interest (or exchange) rate S whose volatility  $\sigma$  is also stochastic. It is given by

(4.1) 
$$dS_t = \sqrt{1 - \rho^2} \sigma_t (S_t)^\beta dW_t^1 + \rho \sigma_t (S_t)^\beta dW_t^2,$$
$$d\sigma_t = \alpha \sigma_t dW_t^2,$$

where W is a standard two-dimensional Brownian motion and  $\alpha$ ,  $\beta$ ,  $\rho$  are parameters. For simplicity, we set  $\alpha = 1$  and  $\beta = \rho = 0$ . In Stratonovich form, (4.1) then becomes

(4.2) 
$$dS_t = \exp(\nu_t) \circ dW_t^1,$$
$$d\nu_t = -\frac{1}{2}dt + dW_t^2,$$

with  $\nu_t := \log(\sigma_t)$ . We will also use  $S_0 = \nu_0 = 0$  and a fixed time horizon of T = 10. As discussed in Section 1.2, it may be difficult to accurately simulate the SDE (4.2) since the noise vector fields are non-commutative. That is, second iterated integrals of Brownian motion (or Lévy areas) will appear within stochastic Taylor expansions.

By Itô's isometry, we can show that the local error for Heun's method is given by

$$\mathbb{E}\left[\left(S_{t+h} - \left(S_t + \frac{1}{2}\sigma_t \left(1 + e^{-\frac{1}{2}h + (W_{t+h}^2 - W_t^2)}\right) \left(W_{t+h}^1 - W_t^1\right)\right)\right)^2\right] = \frac{1}{4}\sigma_t^2 h(e^h - 1)$$

Thus, by setting this proportional to h, we arrive at the following previsible step size,

(4.3) 
$$h(\nu_t) := \log(1 + C \exp(-2\nu_t)),$$

where C > 0 is a user-specified constant. In our experiments, we found that (4.3) gave the best choice of previsible step size – even for the Euler-Maruyama method. A simple lower bound for the "average step size"  $\mathbb{E}[h(\nu_t)]$  is derived in Appendix D.

For the non-previsible adaptive step size, we use the Proportional-Integral (PI) controller for commutative SDEs that is detailed in [32] and already implemented in the popular ODE/SDE package Diffrax [37]. This computes a new step size  $h_{k+1}$  as

$$h_{k+1} = h_k \left( \operatorname{Fac}_{\max} \wedge \left( \operatorname{Fac}_{\min} \vee \left( \frac{\operatorname{Fac} \cdot C}{e(Y_k, h_k, \mathcal{W}_k)} \right)^{K_I} \left( \frac{e(Y_{k-1}, h_{k-1}, \mathcal{W}_{k-1})}{e(Y_k, h_k, \mathcal{W}_k)} \right)^{K_P} \right) \right),$$

where {Fac<sub>max</sub>, Fac<sub>min</sub>} are the maximum and minimum factors  $h_k$  can change by, Fac is a safety factor, { $K_I, K_P$ } are the "integral" and "proportional" coefficients, C > 0 is the absolute tolerance and  $e(Y_k, h_k, W_k)$  denotes the local error estimator  $||Y_{k+1} - \tilde{Y}_{k+1}||_2$  obtained by computing approximations  $Y_{k+1}\tilde{Y}_{k+1}$  from  $(Y_k, h_k, W_k)^3$ . The first three parameters are set to their default values in Diffrax, Fac<sub>max</sub> = 10, Fac<sub>min</sub> = 0.2, Fac = 0.9 and as recommended by [32], we choose  $K_I = 0.3, K_P = 0.1$ . In the experiment, we compare the following numerical methods and step size controls:

- The Euler-Maruyama method, with constant and previsible step sizes.
- Heun's method (Definition 1.5), with constant, previsible and PI step sizes.
- The SPaRK method (Definition 1.6) with constant, previsible and PI step sizes.

<sup>&</sup>lt;sup>3</sup>In our case, the pair  $(Y_{k+1}, \tilde{Y}_{k+1})$  will be obtained from an embedded Runge-Kutta method. Whilst it performed worse in our experiment, one can also compare one step and two half-steps of the same method. Of course, the numerical solution should then be propagated using the two half-steps.

For each of these methods, we will compute the following strong error estimator:

DEFINITION 4.1 (Monte Carlo estimator for strong / root mean squared errors). Let  $\{Y_{t_k}^C\}_{0 \leq t_k \leq T}$  be a numerical solution of SDE (4.1), computed over [0,T] with  $Y_0^C := y_0$ , where C denotes the user-specified parameter(s) used to determine steps. For  $N \ge 1$ , we consider N independent samples of  $Y^C$ , denoted by  $\{\{Y_{t_k,i}^C\}_{k\ge 0}\}_{1\le i\le N}$ .

Then, for a given C and N, we will define the strong error estimator  $S_{N,C}$  as

(4.4) 
$$S_{N,C} := \max_{0 \le k \le 64} \sqrt{\frac{1}{N} \sum_{i=1}^{N} \left\| Y_{k\Delta t,i}^{C} - Y_{k\Delta t,i}^{fine} \right\|_{2}^{2}}$$

where  $\Delta t := \frac{1}{64}T$  and  $\{Y_{t_k,i}^{fine}\}_{0 \leq t_k \leq T}$  denotes a "fine" numerical approximation that is computed using the same SDE solver – but which uses a finer step size of  $h^{\text{fine}} = \frac{1}{8}h$ .

Remark 4.2. A crucial aspect of the error estimator (4.4) is that  $Y^C$  and  $Y^{\text{fine}}$ are computed with respect to the same Brownian motion. Fortunately, this can be achieved in a convenient manner by Diffrax's (single-seed) Virtual Brownian Tree [34].

The estimated convergence rates for the different methods and step size controls are given in Figure 3 and Python code for reproducing these results can be found at github.com/andyElking/Adaptive\_SABR.





FIG. 3. Estimated convergence rates for the different numerical methods and step size controls. Errors for the constant and variable step size SDE solvers were estimated using N = 50,000 and N = 20,000 samples. Since the Euler-Maruyama, Heun and SPaRK methods use 1,2 and 3 vector field evaluations, the x-axis is obtained as the average number of steps multiplied by these numbers.

As expected, the constant step size solvers exhibit a convergence rate of  $O(\sqrt{h})$ , but with SPaRK being the most accurate method. However, with variable step sizes, the methods can exhibit convergence that is linear in the (expected) number of steps. Perhaps this is not surprising for the previsible step sizes, which were model-specific and derived to achieve the desired mean squared error condition. On the other hand, for the PI adaptive SDE solvers, we have used the recommended parameter settings. Since only SPaRK showed linear convergence with the PI adaptive step sizes, it may be the case that Heun is a more effective embedded method than Euler-Maruyama.

It is important to note that the PI controller does not satisfy the conditions of Theorem 2.19, as it can skip over previously evaluated values of the Brownian motion. Since we still observe convergence, we conjecture that the embedded Heun and SPaRK methods do not introduce an  $O(h_k)$  "Lévy area bias" after each adaptive step size  $h_k$ .

5. Conclusion and future work. The primary contribution of this paper shows that standard SDE solvers, such as the Heun and no-area Milstein methods, converge when applied with non-previsible adaptive step sizes (under mild conditions). This enables us to use embedded methods, which are popular within ODE numerics, to establish local errors for an adaptive step size controller in the general SDE setting.

We also introduce a numerical method called "SPaRK" which uses space-time Lévy area to achieve smaller local errors than standard "increment only" SDE solvers. This stochastic Runge-Kutta method is compatible with the proposed adaptive step size theory and we empirically demonstrated its efficacy on a two-dimensional SDE. The results of this paper may also lead to further questions for adaptive SDE solvers:

• When is it acceptable for step sizes to "skip" information of the Brownian motion?

For example, in our experiment, we observed that the Heun and SPaRK methods both appeared to converge with a Proportional-Integral (PI) step size controller.

• Do numerical methods that "weakly" approximate Lévy area terms, or otherwise introduce local unbiasaed O(h) perturbations, converge with adaptive step sizes?

Examples of such schemes include high order weak approximations [35, 53, 54, 66] and the below extension of Heun's method for general Itô SDEs introduced in [58]:

DEFINITION 5.1 (Heun's method with embedded Euler-Maruyama for Itô SDEs). We define a numerical solution  $Y = \{Y_k\}$  for the Itô SDE (1.10) by  $Y_0 := y_0$  and, for  $k \ge 0$ , we define  $(Y_{k+1}, \tilde{Y}_{k+1})$  similar to Heun's method (see Definition 1.5) as

$$Z_{k+1} = Y_k + F(Y_k) \left(\overline{W}_k + \overline{S}_k\right)$$

(5.1) 
$$\widetilde{Y}_{k+1} := Y_k + F(Y_k)\overline{W}_k,$$

(5.2) 
$$Y_{k+1} = Y_k + \frac{1}{2}F(Y_k)\left(\overline{W}_k + \overline{S}_k\right) + \frac{1}{2}F(Z_{k+1})\left(\overline{W}_k - \overline{S}_k\right),$$

where  $\overline{S}_k := (0 \quad \sqrt{h_k} S_k)^{\top}$  with the random vector  $S_k \sim Uniform(\{-1,1\}^d)$  assumed to be independent of the Brownian motion W.

- Is it possible to obtain convergence rates for non-previsible adaptive SDE solvers?
- Similarly, can we prove linear convergence for the previsible solvers in Section 4?
- Can the regularity of the drift vector field f(·) be weakened from Lip(γ) to Lip(1)?
   This may require a Davie's estimate for mixed regularity (1, p)-rough paths [23, 45].
- Does the Lévy construction detailed in Appendix B extend to an explicit algorithm for generating N terms in the polynomial expansion of Brownian motion? [20, 24]

This would allow adaptive solvers to use accurate Lévy area approximations [17].

Acknowledgements. We appreciate the support from the University of Bath, the DataSig group under EPSRC grant EP/S026347/1 and the Alan Turing Institute. We would also like to thank Terry Lyons and Harald Oberhauser for supervising the first author as a graduate student and having countless interesting discussions about stochastic numerics, rough paths and (non-previsible) adaptive step sizes for SDEs.

#### J. FOSTER AND A. JELINČIČ

### REFERENCES

- L. ALLEN, An Introduction to Stochastic Processes with Applications to Biology, Chapman & Hall/CRC, second ed., 1991.
- C. BAYER, S. BRENEIS, AND T. LYONS, An Adaptive Algorithm for Rough Differential Equations, https://arxiv.org/abs/2307.12590, (2023).
- [3] D. BRIGO AND F. MERCURIO, Interest Rate Models Theory and Practice, Springer, 2006.
- [4] A. BROWNING, D. WARNE, K. BURRAGE, R. BAKER, AND M. SIMPSON, Identifiability analysis for stochastic differential equation models in systems biology, Journal of the Royal Society Interface, 17 (2020).
- [5] P. M. BURRAGE AND K. BURRAGE, A Variable Stepsize Implementation for Stochastic Differential Equations, SIAM Journal on Scientific Computing, 24 (2003), pp. 848–864.
- [6] P. M. BURRAGE, R. HERDIANA, AND K. BURRAGE, Adaptive stepsize based on control theory for stochastic differential equations, Journal of Computational and Applied Mathematics, 170 (2004), pp. 317–336.
- [7] N. CAI, Y. SONG, AND N. CHEN, Exact Simulation of the SABR Model, Operational Research, 65 (2017), pp. 837–1113.
- [8] T. CHEN, E. B. FOX, AND C. GUESTRIN, Stochastic Gradient Hamiltonian Monte Carlo, Proceedings of the 31st International Conference on Machine Learning, (2014).
- [9] X. CHENG, N. S. CHATTERJI, P. L. BARTLETT, AND M. I. JORDAN, Underdamped Langevin MCMC: A non-asymptotic analysis, Proceedings of the 31st Conference On Learning Theory, Volume 75 of Proceedings of Machine Learning Research, (2018).
- [10] G. M. CICUTA AND M. L. MEHTA, Probability density of determinants of random matrices, Journal of Physics A: Mathematical and General, 33 (2000).
- [11] Z. CUI, J. L. KIRKBY, AND D. NGUYEN, A General Valuation Framework for SABR and Stochastic Local Volatility Models, SIAM Journal on Financial Mathematics, 9 (2018), pp. 520–563.
- [12] T. DOCKHORN, A. VAHDAT, AND K. KREIS, Score-Based Generative Modeling with Critically-Damped Langevin Diffusion, Proceedings of the International Conference on Learning Representations, (2022).
- [13] R. DU TOIT STRAUSS AND F. EFFENBERGER, A Hitch-hiker's Guide to Stochastic Differential Equations: Solution Methods for Energetic Particle Transport in Space Physics and Astrophysics, Space Science Reviews, 212 (2017), pp. 151–192.
- [14] W. FANG AND M. B. GILES, Adaptive Euler-Maruyama method for SDEs with nonglobally Lipschitz drift, Annals of Applied Probability, 30 (2020), pp. 526–560.
- [15] J. FOSTER, Numerical approximations for stochastic differential equations, PhD thesis, University of Oxford, 2020.
- [16] J. FOSTER, G. DOS REIS, AND C. STRANGE, High order splitting methods for SDEs satisfying a commutativity condition, SIAM Journal on Numerical Analysis, 62 (2024), pp. 500–532.
- [17] J. FOSTER AND K. HABERMANN, Brownian bridge expansions for Lévy area approximations and particular values of the Riemann zeta function, Combinatorics, Probability and Computing, 32 (2023), pp. 370–397.
- [18] J. FOSTER, T. LYONS, AND V. MARGARINT, An Asymptotic Radius of Convergence for the Loewner Equation and Simulation of SLE Traces via Splitting, Journal of Statistical Physics, 189 (2022).
- [19] J. FOSTER, T. LYONS, AND H. OBERHAUSER, The shifted ODE method for underdamped Langevin MCMC, https://arxiv.org/abs/2101.03446, (2021).
- [20] J. FOSTER, H. OBERHAUSER, AND T. LYONS, An optimal polynomial approximation of Brownian motion, SIAM Journal on Numerical Analysis, 58 (2020), pp. 1393–1421.
- [21] J. G. GAINES AND T. LYONS, Variable Step Size Control in the Numerical Solution of Stochastic Differential Equations, SIAM Journal on Applied Mathematics, 57 (1997), pp. 1455–1484.
- [22] M. B. GILES AND L. SZPRUCH, Antithetic multilevel Monte Carlo estimation for multidimensional SDEs without Lévy area simulation, Annals of Applied Probability, 24 (2014), pp. 1585–1620.
- [23] L. G. GYURKÓ, Differential Equations Driven by Π-Rough Paths, Proceedings of the Edinburgh Mathematical Society, 59 (2016), pp. 741–758.
- [24] K. HABERMANN, A semicircle law and decorrelation phenomena for iterated Kolmogorov loops, Journal of the London Mathematical Society, 103 (2021), pp. 558–586.
- [25] P. S. HAGAN, D. KUMAR, A. S. KESNIEWSKI, AND D. E. WOODWARD, Managing Smile Risk, Wilmott Magazine, 1 (2002), pp. 84–108.
- [26] E. HAIRER, W. WANNER, AND S. P. NØRSETT, Solving Ordinary Differential Equations I: Nonstiff Problems, Springer, second ed., 1993.

- [27] Y. HE, K. BALASUBRAMANIAN, AND M. A. ERDOGDU, On the Ergodicity, Bias and Asymptotic Normality of Randomized Midpoint Sampling Method, Advances in Neural Information Processing Systems, (2020).
- [28] D. HIGHAM AND P. KLOEDEN, An Introduction to the Numerical Simulation of Stochastic Differential Equations, Society for Industrial and Applied Mathematics, 2021.
- [29] H. HOEL AND S. RAGUNATHAN, Higher-order adaptive methods for exit times of Itô diffusions, IMA Journal of Numerical Analysis, (2023).
- [30] N. HOFMANN, T. MÜLLER-GRONBACH, AND K. RITTER, Step size control for the uniform approximation of systems of stochastic differential equations with additive noise, Annals of Applied Probability, 10 (2000), pp. 616–633.
- [31] Z. HU, F. HUANG, AND H. HUANG, Optimal Underdamped Langevin MCMC Method, Advances in Neural Information Processing Systems, (2021).
- [32] S. ILIE, K. R. JACKSON, AND W. H. ENRIGHT, Adaptive time-stepping for the strong numerical solution of stochastic differential equations, Numerical Algorithms, 68 (2014), pp. 791–812.
- [33] Z. ISSA, B. HORVATH, M. LEMERCIER, AND C. SALVI, Non-adversarial training of Neural SDEs with signature kernel scores, Advances in Neural Information Processing Systems, (2023).
- [34] A. JELINČIČ, J. FOSTER, AND P. KIDGER, Single-seed generation of Brownian paths and integrals for adaptive and high order SDE solvers, https://arxiv.org/abs/2405.06464, (2024).
- [35] A. JELINČIČ, J. TAO, W. F. TURNER, T. CASS, J. FOSTER, AND H. NI, Generative Modelling of Lévy Area for High Order SDE Simulation, https://arxiv.org/abs/2308.02452, (2023).
- [36] S. JHA AND C. LANGMEAD, Exploring behaviors of stochastic differential equation models of biological systems using change of measures, BMC Bioinformatics, 13 (2012).
- [37] P. KIDGER, On Neural Differential Equations, PhD thesis, University of Oxford, https://arxiv.org/abs/2202.02435, (2022). The Diffrax software package is available at https://github.com/patrick-kidger/diffrax.
- [38] P. KIDGER, J. FOSTER, X. LI, AND T. LYONS, Efficient and Accurate Gradients for Neural SDEs, Advances in Neural Information Processing Systems, (2021).
- [39] P. KIDGER, J. FOSTER, X. LI, H. OBERHAUSER, AND T. LYONS, Neural SDEs as Infinite-Dimensional GANs, Proceedings of the 38th International Conference on Machine Learning, (2021).
- [40] R. KRUSE AND Y. WU, A randomized Milstein method for stochastic differential equations with non-differentiable drift coefficients, Discrete and Continuous Dynamical Systems – Series B, 24 (2019), pp. 3475–3502.
- [41] H. LAMBA, J. C. MATTINGLY, AND A. M. STUART, An adaptive Euler-Maruyama scheme for SDEs: convergence and stability, IMA Journal of Numerical Analysis, 27 (2007), pp. 479– 506.
- [42] B. LEIMKUHLER AND C. MATTHEWS, Molecular Dynamics: With Deterministic and Stochastic Numerical Methods, Interdisciplinary Applied Mathematics, Springer, 2015.
- [43] B. LEIMKUHLER, C. MATTHEWS, AND M. V. TRETYAKOV, On the long-time integration of stochastic gradient systems, Proceedings of the Royal Society A, 470 (2014).
- [44] A. LEITAO, L. A. GRZELAK, AND C. W. OOSTERLEE, On an efficient multiple time step Monte Carlo simulation of the SABR model, Quantitative Finance, 17 (2017), pp. 1549–1565.
- [45] A. LEJAY AND N. VICTOIR, On (p,q)-rough paths, Journal of Differential Equations, 225 (2006), pp. 103–133.
- [46] X. LI, T.-K. L. WONG, R. T. Q. CHEN, AND D. DUVENAUD, Scalable Gradients for Stochastic Differential Equations, Proceedings of the 23rd International Conference on Artificial Intelligence and Statistics, (2020).
- [47] X. LI, D. WU, L. MACKEY, AND M. A. ERDOGDU, Stochastic Runge-Kutta Accelerates Langevin Monte Carlo and Beyond, Advances in Neural Information Processing Systems, (2019).
- [48] T. J. LYONS, M. CARUANA, T. LÉVY, AND J. PICARD, Differential equations driven by rough paths, Ecole d'été de Probabilités de Saint-Flour, 34 (2004), pp. 1–93.
- [49] S. MAUTHNER, Step size control in the numerical solution of stochastic differential equations, Journal of Computational and Applied Mathematics, 100 (1998), pp. 93–109.
- [50] G. N. MILSTEIN AND M. V. TRETYAKOV, Stochastic Numerics for Mathematical Physics, Springer, 2004.
- [51] P. M. MORKISZ AND P. PRZYBYŁOWICZ, Randomized derivative-free Milstein algorithm for efficient approximation of solutions of SDEs under noisy information, Journal of Computational and Applied Mathematics, 383 (2021).
- [52] A. NEUENKIRCH, M. SZÖLGYENYI, AND L. SZPRUCH, An Adaptive Euler-Maruyama Scheme for Stochastic Differential Equations with Discontinuous Drift and its Convergence Analysis, SIAM Journal on Numerical Analysis, 57 (2019), pp. 378–403.
- [53] M. NINOMIYA AND S. NINOMIYA, A new higher-order weak approximation scheme for stochastic

differential equations and the Runge-Kutta method, Finance and Stochastics, 13 (2009), pp. 415–443.

- [55] B. ØKSENDAL, Stochastic Differential Equations: An Introduction with Applications, Springer, Berlin, sixth ed., 2003.
- [56] C. RACKAUCKAS, Y. MA, J. MARTENSEN, C. WARNER, K. ZUBOV, R. SUPEKAR, D. SKINNER, A. RAMADHAN, AND A. EDELMAN, Universal Differential Equations for Scientific Machine Learning, https://arxiv.org/abs/2001.04385, (2021).
- [57] C. RACKAUCKAS AND Q. NIE, Adaptive methods for stochastic differential equations via natural embeddings and rejection sampling with memory, Discrete and Continuous Dynamical Systems – Series B, 22 (2017), pp. 2731–2761.
- [58] A. J. ROBERTS, Modify the improved Euler scheme to integrate stochastic differential equations, https://arxiv.org/abs/1210.0933, (2012).
- [59] A. RÖSSLER, Embedded Stochastic Runge-Kutta Methods, Proceedings in Applied Mathematics & Mechanics, 2 (2003), pp. 461–462.
- [60] F. SAMMÜLLER AND M. SCHMIDT, Adaptive Brownian Dynamics, The Journal of Chemical Physics, 155 (2021).
- [61] R. SHEN AND Y. T. LEE, The Randomized Midpoint Method for Log-Concave Sampling, Advances in Neural Information Processing Systems, (2019).
- [62] S. SHREVE, Stochastic Calculus for Finance II: Continuous-Time Models, Springer, 2004.
- [63] K. SOBCZYK, Stochastic Differential Equations: With Applications to Physics and Engineering, Mathematics and its Applications, Springer, 1991.
- [64] G. SÖDERLIND, Automatic control and adaptive time-stepping, Numerical Algorithms, 31 (2002), pp. 281–310.
- [65] Y. SONG, J. SOHL-DICKSTEIN, D. P. KINGMA, A. KUMAR, S. ERMON, AND B. POOLE, Score-Based Generative Modeling through Stochastic Differential Equations, Proceedings of the International Conference on Learning Representations, (2021).
- [66] D. TALAY, Second-order discretization schemes of stochastic differential systems for the computation of the invariant law, Stochastics and Stochastic Reports, 29 (1990), pp. 13–36.
- [67] F. VADILLO, Comparing stochastic Lotka-Volterra predator-prey models, Applied Mathematics and Computation, 360 (2020), pp. 181–189.
- [68] N. B. VICTOIR AND P. K. FRIZ, Multidimensional Stochastic Processes as Rough Paths: Theory and Applications, Cambridge University Press, 2010.
- [69] M. WELLING AND Y. W. TEH, Bayesian Learning via Stochastic Gradient Langevin Dynamics, Proceedings of the 28th International Conference on Machine Learning, (2011).
- [70] A. WINKELBAUER, Moments and Absolute Moments of the Normal Distribution, https://arxiv.org/abs/1209.4340, (2014).

Appendix A. Improved local accuracy of the SPaRK method (1.16). In this section, we will detail certain properties of the newly introduced stochastic Runge-Kutta method (1.16) that make it an appealing alternative to Heun's method.

We first recall the definition of this method (using the notation from Section 1.1),

$$Y_{k+\frac{1}{2}} := Y_k + F(Y_k) \left(\frac{1}{2}\overline{W}_k + \sqrt{3}\,\overline{H}_k\right), \qquad Z_{k+1} := Y_k + F(Y_{k+\frac{1}{2}})\overline{W}_k,$$
  
(A.1) 
$$Y_{k+1} := Y_k + F(Y_k) \left(a\overline{W}_k + \overline{H}_k\right) + bF(Y_{k+\frac{1}{2}})\overline{W}_k + F(Z_{k+1}) \left(a\overline{W}_k - \overline{H}_k\right),$$

with  $a = \frac{3-\sqrt{3}}{6}$  and  $b = \frac{\sqrt{3}}{3}$ . In Section 2.3, the scheme (A.1) was Taylor expanded as

(A.2) 
$$Y_{k+1} = Y_k + f(Y_k)h_k + g(Y_k)W_k + g'(Y_k)g(Y_k)\left(\frac{1}{2}W_k^{\otimes 2} + H_k \otimes W_k - W_k \otimes H_k\right) + o(h_k).$$

By comparing (A.2) with the Taylor expansion of the SDE solution started from  $Y_k$ , we see that the leading term in the expansion of the one-step mean squared error is

$$\begin{split} & \mathbb{E}\bigg[ \left\| g'(Y_k) g(Y_k) \bigg( \int_{t_k}^{t_{k+1}} W_{t_k,t} \otimes \circ dW_t - \left( \frac{1}{2} W_k^{\otimes 2} + H_k \otimes W_k - W_k \otimes H_k \right) \right) \right\|_2^2 \bigg] \\ &= \sum_{i \neq j} \mathbb{E} \bigg[ \left\| g'_j(Y_k) g_i(Y_k) \right\|_2^2 \bigg] \underbrace{\mathbb{E} \bigg[ \left( \int_{t_k}^{t_{k+1}} W_{t_k,t}^i \circ dW_t^j - \left( \frac{1}{2} W_k^i W_k^j + H_k^i W_k^j - W_k^i H_k^j \right) \right)^2 \bigg]}_{= \frac{1}{12} h_k^2 \text{ by } [17, \text{ Theorem 5.4] and Brownian scaling}} \\ &= \frac{1}{12} h_k^2 \sum_{i \neq j} \mathbb{E} \bigg[ \left\| g'_j(Y_k) g_i(Y_k) \right\|_2^2 \bigg]. \end{split}$$

On the other hand, two half steps of Heun's method on  $[t_k, t_{k+1}]$  can be expanded as

$$\begin{split} Y_{k+1} &= Y_{k+\frac{1}{2}} + \frac{1}{2} f(Y_{k+\frac{1}{2}}) h_k + g(Y_{k+\frac{1}{2}}) W_{k+\frac{1}{2}} + g'(Y_{k+\frac{1}{2}}) g(Y_{k+\frac{1}{2}}) \left(\frac{1}{2} W_{k+\frac{1}{2}}^{\otimes 2}\right) + o(h_k) \\ &= Y_k + f(Y_k) h_k + g(Y_k) \left(W_k + W_{k+\frac{1}{2}}\right) \\ &+ g'(Y_k) g(Y_k) \left(\frac{1}{2} W_k^{\otimes 2} + W_k \otimes W_{k+\frac{1}{2}} + \frac{1}{2} W_{k+\frac{1}{2}}^{\otimes 2}\right) + o(h_k), \end{split}$$

where we now use the notation  $W_k := W_{t_{k+\frac{1}{2}}} - W_{t_k}$  and  $W_{k+\frac{1}{2}} := W_{t_{k+1}} - W_{t_{k+\frac{1}{2}}}$ . For  $i \neq j$ , we have  $\int_s^t W_{s,u}^i \circ dW_u^j = \int_s^t W_{s,u}^i dW_u^j$ , and so applying Itô's isometry gives

$$\begin{split} & \mathbb{E}\bigg[\bigg(\int_{t_{k}}^{t_{k+1}} W_{t_{k},t}^{i} \circ dW_{t}^{j} - \bigg(\frac{1}{2}W_{k}^{i}W_{k}^{j} + W_{k}^{i}W_{k+\frac{1}{2}}^{j} + \frac{1}{2}W_{k+\frac{1}{2}}^{i}W_{k+\frac{1}{2}}^{j}\bigg)\bigg)^{2}\bigg] \\ &= \mathbb{E}\bigg[\bigg(\int_{t_{k}}^{t_{k+\frac{1}{2}}} W_{t_{k},t}^{i} \circ dW_{t}^{j} - \frac{1}{2}W_{k}^{i}W_{k}^{j}\bigg)^{2} + \bigg(\int_{t_{k+\frac{1}{2}}}^{t_{k+1}} W_{t_{k+\frac{1}{2}},t}^{i} \circ dW_{t}^{j} - \frac{1}{2}W_{k+\frac{1}{2}}^{i}W_{k+\frac{1}{2}}^{j}\bigg)^{2}\bigg] \\ &= \frac{1}{4}\bigg(\frac{1}{2}h_{k}\bigg)^{2} + \frac{1}{4}\bigg(\frac{1}{2}h_{k}\bigg)^{2} = \frac{1}{8}h_{k}^{2}. \end{split}$$

Hence, the leading term in the mean squared error of the two-step Heun's method is

$$\frac{1}{8}h_k^2 \sum_{i \neq j} \mathbb{E}\Big[ \left\| g_j'(Y_k) g_i(Y_k) \right\|_2^2 \Big].$$

Thus, provided the step size is sufficiently small, we see that one step of the stochastic Runge-Kutta method (1.16) is more accurate than two steps of Heun's method (1.14). In addition, two steps of Heun's method uses an extra evaluation of the vector fields.

It is also known that Heun's method achieves second order weak convergence for SDEs with additive noise [43] (that is, when  $g(y) = \sigma$  for a constant matrix  $\sigma \in \mathbb{R}^{e \times d}$ ),

$$dy_t = f(y_t)dt + \sigma dW_t$$

where, for the purposes of this section, we will assume that f is sufficiently smooth. For additive noise SDEs, the Splitting Path Runge-Kutta (SPaRK) method becomes

$$\begin{split} Y_{k+\frac{1}{2}} &:= Y_k + \frac{1}{2}f(Y_k)h_k + \sigma\bigg(\frac{1}{2}W_k + \sqrt{3}H_k\bigg), \qquad Z_{k+1} := Y_k + f(Y_{k+\frac{1}{2}})h_k + \sigma W_k, \\ Y_{k+1} &:= Y_k + \bigg(\frac{3-\sqrt{3}}{6}f(Y_k) + \frac{\sqrt{3}}{3}f(Y_{k+\frac{1}{2}}) + \frac{3-\sqrt{3}}{6}f(Z_{k+1})\bigg)h_k + \sigma W_k. \end{split}$$

Using the classical Taylor theorem  $f(b) = f(a) + f'(a)(b-a) + \frac{1}{2}f''(a)(b-a)^{\otimes 2} + R$ with  $R := \frac{1}{2} \int_0^1 (1-t)^2 f'''(a+t(b-a))(b-a)^{\otimes 3} dt$ , we can Taylor expand  $Y_{k+1}$  to give

$$\begin{split} Y_{k+1} &= Y_k + f(Y_k)h_k + \sigma W_k + \frac{\sqrt{3}}{3}f'(Y_k) \bigg(\frac{1}{2}f(Y_k)h_k + \sigma\bigg(\frac{1}{2}W_k + \sqrt{3}H_k\bigg)\bigg)h_k \\ &\quad + \frac{3-\sqrt{3}}{6}f'(Y_k) \bigg(f(Y_{k+\frac{1}{2}})h_k + \sigma W_k\bigg)h_k \\ &\quad + \frac{\sqrt{3}}{3}\bigg(\frac{1}{2}f''(Y_k)\bigg(\frac{1}{2}f(Y_k)h_k + \sigma\bigg(\frac{1}{2}W_k + \sqrt{3}H_k\bigg)\bigg)^{\otimes 2}\bigg)h_k \\ &\quad + \frac{3-\sqrt{3}}{6}\bigg(\frac{1}{2}f''(Y_k)\bigg(f(Y_{k+\frac{1}{2}})h_k + \sigma W_k\bigg)^{\otimes 2}\bigg)h_k + O(h_k^{\frac{5}{2}}) \\ &= Y_k + f(Y_k)h_k + \sigma W_k + f'(Y_k)\sigma\bigg(\frac{1}{2}W_k + H_k\bigg)h_k + \frac{1}{2}f'(Y_k)f(Y_k)h_k^2 + O(h_k^{\frac{5}{2}}) \\ &\quad + \frac{1}{2}f''(Y_k)\sigma^{\otimes 2}\bigg(\frac{6-\sqrt{3}}{12}W_k^{\otimes 2} + \frac{1}{2}W_k \otimes H_k + \frac{1}{2}H_k \otimes W_k + \sqrt{3}H_k^{\otimes 2}\bigg)h_k. \end{split}$$

We note that in the above Taylor expansion, we have  $\left(\frac{1}{2}W_k + H_k\right)h_k = \int_{t_k}^{t_{k+1}} W_{t_k,t} dt$ and  $\mathbb{E}\left[\left(\frac{6-\sqrt{3}}{12}W_k^{\otimes 2} + \frac{1}{2}W_k \otimes H_k + \frac{1}{2}H_k \otimes W_k + \sqrt{3}H_k^{\otimes 2}\right)h_k\right] = \frac{1}{2}h_k^2 = \mathbb{E}\left[\int_{t_k}^{t_{k+1}} W_{t_k,t}^{\otimes 2} dt\right].$ 

Thus, by the standard mean squared error analysis of Milstein and Tretyakov [50], we see that stochastic Runge-Kutta method will converge strongly with rate  $O(h^{\frac{3}{2}})$ . Similarly, it is also clear that the method will achieve second order weak convergence. We refer the reader to [16] for more details on the analysis of commutative noise SDEs. For non-additive noise SDEs, it is straightforward to see that the proposed stochastic Runge-Kutta method exhibits the same strong convergence rates as Heun's method.

	Heun's method One step Two steps		Splitting Path Runge-Kutta	
Vector field evaluations	2	4	3	
Gaussian random vectors	1	2	2	
Coefficient in front of leading term $h_k^2 \sum_{i \neq j} \mathbb{E} \Big[ \ g'_j(Y_k)g_i(Y_k)\ _2^2 \Big]$ in expansion of mean squared error	$\frac{1}{4}$	$\frac{1}{8}$	$\frac{1}{12}$	
Convergence rates for SDEs with	Strong	Weak	Strong	Weak
general noise	$O(h^{\frac{1}{2}})$	O(h)	$O(h^{\frac{1}{2}})$	O(h)
commutative noise $(g'_i(y)g_j(y) = g'_j(y)g_i(y))$	O(h)	O(h)	O(h)	O(h)
additive noise $(g(y) \equiv \sigma \in \mathbb{R}^{e \times d})$	O(h)	$O(h^2)$	$O(h^{\frac{3}{2}})$	$O(h^2)$

To conclude, we summarise the properties of these numerical methods in Table 1.

Table 1

Properties of Heun's method (1.14) and the proposed SPaRK method (1.16).

Appendix B. Generating increments and integrals of Brownian motion. In this section, we will briefly outline how increments  $W_{s,t}$  and space-time Lévy areas  $H_{s,t} := \frac{1}{t-s} \int_s^t W_{s,u} du$  of Brownian motion can be generated for adaptive SDE solvers.

We emphasise that the results detailed in this section are already well-established in the SDE numerics literature [15, 34] and are only included here for completeness.

To begin, we note that for any  $u \in [s, t]$ , a path increment  $W_{s,u} := W_u - W_s$  can be generated conditional on  $W_{s,t}$  using standard properties of the Brownian bridge. When  $u = \frac{1}{2}(s+t)$ , this leads to well-known *Lévy construction of Brownian motion*.

THEOREM B.1 (Conditional distribution of Brownian increments). For  $s \leq u \leq t$ ,

$$W_{s,u} = \frac{u-s}{t-s} W_{s,t} + B_{s,u}, \qquad W_{u,t} = \frac{t-u}{t-s} W_{s,t} - B_{s,u},$$

where the Brownian bridge increment  $B_{s,u} \sim \mathcal{N}\left(0, \frac{(u-s)(t-u)}{t-s}I_d\right)$  is independent of  $W_{s,t} \sim \mathcal{N}\left(0, (t-s)I_d\right)$ . When  $u = \frac{1}{2}(s+t)$  is the midpoint of [s,t], this simplifies to

$$W_{s,u} = \frac{1}{2}W_{s,t} + B_{s,u}, \qquad W_{u,t} = \frac{1}{2}W_{s,t} - B_{s,u}$$

where  $B_{s,u} \sim \mathcal{N}(0, \frac{1}{4}(t-s)I_d)$ .

*Proof.* Let  $B_{s,u} := W_{s,u} - \frac{u-s}{t-s}W_{s,t}$ . The covariance between  $B_{s,u}$  and  $W_{s,t}$  is

$$Cov(B_{s,u}, W_{s,t}) = Cov(W_{s,u}, W_{s,t}) - \frac{u-s}{t-s} Cov(W_{s,t}, W_{s,t})$$
$$= (u-s) - \frac{u-s}{t-s}(t-s) = 0.$$

Since  $B_{s,u}$  and  $W_{s,t}$  depend on the same Brownian motion, they are jointly normal, and therefore independent. The variance of  $B_{s,u}$  is straightforward to compute.

#### J. FOSTER AND A. JELINČIČ

Extending the theorem given above to include both increments and space-time Lévy areas of the Brownian motion is more involved, but based on a similar argument (which also uses that uncorrelated jointly normal random variables are independent).

THEOREM B.2 (Conditional distribution of increments and space-time Lévy areas of Brownian motion within an interval, [15, Theorems 6.1.4 and 6.1.6]). For  $s \leq u \leq t$ ,

$$\begin{split} W_{s,u} &= \frac{u-s}{h} W_{s,t} + \frac{6(u-s)(t-u)}{h^2} H_{s,t} + \frac{2(a+b)}{h} X_1 \\ W_{u,t} &= \frac{t-u}{h} W_{s,t} - \frac{6(u-s)(t-u)}{h^2} H_{s,t} - \frac{2(a+b)}{h} X_1 \\ H_{s,u} &= \frac{(u-s)^2}{h^2} H_{s,t} - \frac{a}{u-s} X_1 + \frac{c}{u-s} X_2, \\ H_{u,t} &= \frac{(u-s)^2}{h^2} H_{s,t} - \frac{b}{t-u} X_1 - \frac{c}{t-u} X_2, \end{split}$$

where  $W_{s,t} \sim \mathcal{N}(0, hI_d), H_{s,t} \sim \mathcal{N}\left(0, \frac{1}{12}hI_d\right)$  and  $X_1, X_2 \sim \mathcal{N}(0, I_d)$  are independent, h = t - s denotes the length of the interval and the coefficients  $a, b, c \in \mathbb{R}$  are given by

$$\begin{split} a &:= \frac{(u-s)^{\frac{7}{2}}(t-u)^{\frac{1}{2}}}{2h\sqrt{(u-s)^3 + (t-u)^3}}, \qquad b := \frac{(u-s)^{\frac{1}{2}}(t-u)^{\frac{7}{2}}}{2h\sqrt{(u-s)^3 + (t-u)^3}}, \\ c &:= \frac{\sqrt{3}(u-s)^{\frac{3}{2}}(t-u)^{\frac{3}{2}}}{6\sqrt{(u-s)^3 + (t-u)^3}}. \end{split}$$

When  $u = \frac{1}{2}(s+t)$  is the midpoint of the interval [s,t], these formulae simplify to

$$\begin{split} W_{s,u} &= \frac{1}{2} W_{s,t} + \frac{3}{2} H_{s,t} + Z_{s,u}, \qquad W_{u,t} = \frac{1}{2} W_{s,t} - \frac{3}{2} H_{s,t} - Z_{s,u}, \\ H_{s,u} &= \frac{1}{4} H_{s,t} - \frac{1}{2} Z_{s,u} + \frac{1}{2} N_{s,t}, \qquad H_{u,t} = \frac{1}{4} H_{s,t} - \frac{1}{2} Z_{s,u} - \frac{1}{2} N_{s,t}, \end{split}$$

where  $W_{s,t} \sim \mathcal{N}(0, hI_d), Z_{s,u} \sim \mathcal{N}(0, \frac{1}{16}hI_d)$  and  $H_{s,t}, N_{s,t} \sim \mathcal{N}(0, \frac{1}{12}hI_d)$  are all independent Gaussian random vectors.



FIG. 4. Illustration of the Brownian increments and areas in Theorem B.2 (diagram from [15])

Appendix C. Norm of the determinant for a Gaussian random matrix. In this section, we present the following known result, which was helpful in Section 3.

THEOREM C.1  $(L^1(\mathbb{P}) \text{ norm of the determinant for a Gaussian random matrix}).$ For  $n \ge 1$ , let  $A_n = \{a_{ij}\}_{1 \le i,j \le n}$  be an  $n \times n$  matrix with independent entries  $a_{ij} \sim \mathcal{N}(0,1)$  and let  $Z \sim \mathcal{N}(0,1)$  denote a standard normal random variable. Then, (C.1)  $\mathbb{E}[|\det A_n|] = \mathbb{E}[|Z|^n].$ 

*Proof.* Let  $p_n$  denote the probability density function of  $|\det A_n|$ . Then the Mellin transform of  $p_n$  is given by [10, Equation (2.3)], but with  $a = \frac{1}{2}$  and  $\beta = 1$ . That is,

$$\mathcal{M}_n(s) := \int_0^\infty x^{s-1} p_n(x) dx = \left(\frac{1}{2}\right)^{-\frac{1}{2}n(s-1)} \prod_{j=1}^n \left(\frac{\Gamma\left(\frac{1}{2}(s-1) + \frac{1}{2}j\right)}{\Gamma\left(\frac{1}{2}j\right)}\right),$$

where  $\Gamma$  is the standard Gamma function. Since  $\mathbb{E}[|\det A_n|] = \int_0^\infty x p_n(x) dx$ , we have

$$\mathbb{E}[|\det A_n|] = \left(\frac{1}{2}\right)^{-\frac{1}{2}n} \prod_{j=1}^n \left(\frac{\Gamma(\frac{1}{2} + \frac{1}{2}j)}{\Gamma(\frac{1}{2}j)}\right)$$
$$= \left(\frac{1}{2}\right)^{-\frac{1}{2}n} \frac{\Gamma(\frac{1}{2} + \frac{1}{2}n)}{\Gamma(\frac{1}{2})}$$
$$= 2^{\frac{1}{2}n} \frac{\Gamma(\frac{1}{2}(n+1))}{\sqrt{\pi}},$$

which is the *n*-th moment of a half-normal distribution (see [70, Equation (18)]).  $\Box$ 

In particular, when n = 2, we obtain the result used in the proof of Theorem 3.1.

THEOREM C.2. Let  $A, B, C, D \sim \mathcal{N}(0, 1)$  be independent random variables. Then (C.2)  $\mathbb{E}[|AD - BC|] = 1.$ 

Remark C.3. Although (C.1) immediately follows from [10, Equation (2.3)] and [70, Equation (18)], it was difficult for the authors to find it stated in the literature. Hence, (C.1) may be a lesser-known (but elegant) fact about the normal distribution.

Appendix D. A lower bound for the SABR-specific previsible step size. By the construction of the previsible step size (4.3), applying Heun's method to the SABR model (4.2) with step size  $h \equiv h(\nu_t)$  gives a local mean-squared error of  $\frac{1}{4}Ch$ . Since the local errors propagate linearly, the global mean-squared error will be O(C). On the other hand, we can also consider the "average step size" taken by the method.

THEOREM D.1. Let  $\nu_t := -\frac{1}{2}t + W_t$  denote the log-volatility component of (4.2) and let  $h(\nu_t) := \log(1 + Ce^{-2\nu_t})$  be the previsible step size control with C > 0. Then (D.1)  $\mathbb{E}[h(\nu_t)] \ge \log(1 + Ce^t).$ 

*Proof.* We first note that the previsible step size control  $x \mapsto h(x)$  is convex as  $\frac{d^2}{dx^2}h(x) = \frac{d^2}{dx^2}(\log(1+Ce^{-2x})) = \frac{4Ce^{2x}}{(C+e^{2x})^2} > 0.$  Thus, by Jensen's inequality, we have

$$\mathbb{E}[h(\nu_t)] \ge h\left(\mathbb{E}[\nu_t]\right) = h\left(-\frac{1}{2}t\right) = \log\left(1 + Ce^t\right).$$

Hence, if C is sufficiently small, the average step size at any  $t \in [0, T]$  is O(C), and results in a global  $L^2(\mathbb{P})$  error of  $O(\sqrt{C})$ . This mimics the  $O(\sqrt{h})$  strong convergence of SDE solvers with a constant step size. However, since (D.1) increases with time, we can see that the variable step size methods tend to take increasingly larger steps.