# The diagrammatic presentation of equations in categories

Kevin Arlin

James Fairbanks Tim Hosgood

Evan Patterson

17 January 2024

#### Abstract

A lift of a diagram  $D: J \rightarrow X$  in a category against a discrete opfibration  $\pi: E \rightarrow X$  can be interpreted as presenting a solution to a system of equations in X presented by D. With this interpretation in mind, it is natural to ask if there is a notion of equivalence of diagrams  $D \simeq D'$  that precisely captures the idea of the two diagrams "having the same solutions". We give such a definition, and then show how the localisation of the category of all diagrams in X along such equivalences is isomorphic to the localisation of the slice category Cat/X along the class of initial functors. Finally, we extend this result to the 2-categorical setting, proving the analogous statement for any locally presentable 2-category in place of Cat.

#### Contents

1	Introduction	1
2	Diagram categories	5
3	Initial and relatively initial functors	14
4	Localisations of categories of diagrams	20
5	The general 2-categorical story	32
References		41

# **1** Introduction

A diagram in a category can be viewed as a presentation of a system of equations given by equating parallel paths in its image. For example, we can present the equation

$$y^2 = x^3 + x + 1$$

describing an elliptic curve in an affine plane  $\mathbb{A}^2$  over a field k by the diagram

$$\mathbb{A}^2 \underbrace{\overset{y^2}{\overbrace{x^3+x+1}}}_{x^3+x+1} k$$
 .

What is crucial here is that this diagram *does not commute*. The locus on which the equation  $y^2 = x^3 + x + 1$  holds, which for a geometer is the variety determined by the system, is given categorically by the equalizer; in general, the limit of a diagram in a category may be viewed as the object of all solutions to the system of equations the diagram presents.

From previous work [PBHF22], we are motivated by partial differential equations, a field in which mathematicians must in general study only points, or families of points, in the solution space. An individual solution is given in familiar categories by a cone over the diagram tipped by a terminal object or monoidal unit. More generally, a family of solutions indexed by an arbitrary object x is given by a cone over the diagram tipped by x. Thus we might, for instance, obtain a two-parameter family of solutions to a system of differential equations using a cone tipped by  $\mathbb{R}^2$ .

The categorical framing allows us to generalize yet further. Letting  $D : J \to X$  be a diagram in a category X, we shift perspective to consider the choice of a cone over D tipped by x as a *lift* of the functor D into the coslice category x/X:



We are led to consider generalized solutions via lifting against functors into X. For instance, if the spaces  $\mathbb{R}^n$  are in our category X then, by considering lifts against the canonical functor  $\coprod_n(\mathbb{R}^n/\mathsf{X}) \to \mathsf{X}$ , we can reify the concept of "an *n*-parameter family of solutions for some *n*." If the interval [0, 1] is available in X then, by the same token, by lifting against the projection  $C \to X$ , where *C* is the coequalizer of the canonical functors  $\mathbb{R}^1 \times [0, 1]/\mathsf{X} \to \mathbb{R}^1/\mathsf{X}$ , we reify the concept of "a homotopy equivalence class of one-parameter families of solutions." It is thus rather natural to close the family of coslice projections  $x/\mathsf{X} \to \mathsf{X}$  under arbitrary colimits and consider notions of solution resulting from lifting against all of the resulting functors into X. The result of this colimit closure is precisely the class of discrete opfibrations over X. (We recall the definition of discrete opfibration below in Definition 2.6.) For reasons that will become clear following Definition 2.9 below, it is infelicitous to continue generalizing to lifts against arbitrary functors over X, and so we have found our basic setting. This was already the setting that emerged from a recent study in applied category theory by Patterson, Baas, Hosgood, and Fairbanks [PBHF22].

To summarize thus far: we are interested in lifts of diagrams  $D: J \to X$  against discrete opfibrations  $\pi: \mathsf{E} \to \mathsf{X}$ . We take as our central question under what circumstances such a diagram D admits "the same" lifts against all such discrete opfibrations—in equational terms, the same solutions for every way of interpreting the variables when the diagram is viewed as a system of equations—as another diagram  $D': J' \to X$ . Since we want to allow the shape of the diagram to vary, we are led to consider the category  $Diag_{\leftarrow}(X)$ of all diagrams in X. (See Section 2 for the details on this category; note for now that it is not simply the slice Cat/X but incorporates a lax aspect in its morphisms.) The category  $Diag_{\leftarrow}(X)$ , as a suitable domain for a global limit functor, goes all the way back to Eilenberg and Mac Lane [EM45]. It is nevertheless not very extensively studied (though see [GV77] for a characterization of  $Diag_{\leftarrow}(X)$  as a lax cocompletion and [Mes23] for a description of colimits in  $Diag_{\leftarrow}(X)$ ), and the notion of limit is more often formalized "locally," one diagram shape at a time. The diagram category has nonetheless arisen repeatedly in the short history of applied category theory thus far, both in the current context and in the modeling of functorial data migrations [Spi23]; in both cases at the heart of the usefulness of  $Diag_{\leftarrow}(X)$  is the fact that a diagram in X gives a convenient presentation of a discrete opfibration over X—for computational purposes, importantly, a presentation that might well be finite even when X is not.

Any morphism  $D \to D'$  in  $\text{Diag}_{\leftarrow}(X)$  induces a function from lifts of D against a discrete opfibration  $\pi$  to lifts of D' against  $\pi$ , and we can use this canonical function to define a class of *weak equivalences* on  $\text{Diag}_{\leftarrow}(X)$  as those morphisms which induce bijections of lifts against arbitrary discrete opfibrations over X. Such maps will be viewed as isomorphisms of the systems of equations presented by the two diagrams. A very similar class of weak equivalences was studied in Section 9 of [PBHF22].

The primary work of this paper, leading to Theorem 4.6, is to compute the localization  $Diag_{\leftarrow}(X)[W^{-1}]$ , where W is the class of weak equivalences described above. The *localization* (Definition 4.1) of a category at some class of morphisms universally adjoins inverses to those morphisms [GZ67]. In particular, the localization  $Diag_{\leftarrow}(X)[W^{-1}]$ , universally turning the equivalences of systems of equations into isomorphisms, can be thought of as "the category of systems of equations in X." From a theoretical point of view, we obtain a completely satisfying answer:

# The category $Diag_{\leftarrow}(X)[W^{-1}]$ of systems of equations in a category X is equivalent to the full subcategory of the slice Cat/X spanned by the discrete opfibrations.

It was not at all obvious a priori to these authors that this theorem should hold, although it is reads quite naturally after the fact. In particular, the proof includes some technical work in establishing that the morphisms in the localization can all be represented by *strict* maps in Cat/X.

So, the motto is that a "a system of equations in X is a discrete opfibration over X." That said, we are especially interested in computationally tractable descriptions of

this localization for future work in algorithmic rewriting for PDEs. The description of the localization in terms of discrete opfibrations (or, equivalently, copresheaves) is not appropriate for this purpose, due to the loss of the opportunity for finite presentation of objects mentioned above. We thus expect to find more applications for the description of the localization as the category whose objects are diagrams in X and whose morphisms  $D \rightarrow D''$  are the zigzags  $D \leftarrow D' \rightarrow D''$  with the reversed arrow determined by an initial functor between the domains of D and of D'. Initial functors (Definition 5.5) can be described either as functors, restriction along which does not change the limit of a diagram, or as those left orthogonal to discrete opfibrations. The other key aspect of the proof of our main theorem, in addition to the strictification mentioned above, is to show that inverting diagram morphisms given by initial functors suffices to invert all weak equivalences—even though not nearly all weak equivalences are themselves given by initial morphisms.

Our first main result, Theorem 4.6, is not sufficiently general to apply to most systems of interest in physics and engineering, including those in [PBHF22]. The reason is that diagrams in a category without extra structure can only describe equations involving unary operations, whereas equations in applied mathematics tend to involve operations or differential operators of higher arity. Thus, one often prefers to take diagrams in a categorical structure going beyond a bare category.

For instance, as simple an equation as B(x, y) = 0, where B is a bilinear form on a vector space V, is most naturally modelled by considering the symmetric monoidal or multicategorical structure on vector spaces. We generally prefer the latter, though we address both, so that we would draw this equation diagrammatically like as

$$(V,V) \underbrace{\overset{0}{\overbrace{\phantom{a}}}_{B} k \ .$$

For various types of equations, we might prefer to work in multicategories, perhaps symmetric or cartesian or equipped with coproducts, or the analogous monoidal, symmetrical monoidal, cartesian monoidal, or distributive monoidal categories. Motivated by these many possibilities, the final Section 5 generalizes our main result from the 2-category of categories to an arbitrary locally presentable 2-category. This setting directly includes the multicategorical examples and indirectly includes the others, as we will explain later using standard tricks from 2-monad theory.

To even state the broadened result, the concepts of discrete opfibration and initial functor must be defined in a general 2-category. These concepts sometimes generalize in more than one useful way; for our purposes, we propose a definition of discrete opfibration in an arbitrary 2-category that is inspired by Riehl and Verity's treatment of formal  $(\infty$ -)category theory in a virtual equipment [RV22]. This definition reduces to the standard one in the 2-category of categories. We then *define* the initial morphisms to be the class of morphisms left orthogonal to discrete opfibrations. Thus, what was a theorem

for categories, the existence of Street and Walters' *comprehensive factorization* [SW73], becomes essentially a definition, using standard features of factorization systems in locally presentable categories.

With these definitions and a limited amount of groundwork, the proofs in Section 4 go through essentially without change to establish a wide generalization of our main theorem:

For any locally presentable 2-category  $\mathcal{K}$  containing an object x, the category of discrete opfibrations over x, as a full subcategory of the slice  $\mathcal{K}/x$ , is equivalent to the localization of the diagram category  $\text{Diag}_{\leftarrow}(x)$  at a class of weak equivalences defined just as in the case  $\mathcal{K} = \text{Cat.}$ 

The results of this paper illustrate how relatively deep categorical mathematics, including localizations, 2-dimensional monad theory, and comprehensive factorization systems, can be used to give a novel notion of sameness for systems of equations. In future work, we intend to make use of these results to implement algorithms for the simplification of such systems that would be essentially indescribable without the categorical formalism.

**Conventions** We write categories in sans-serif, e.g. X and J, functors and objects of an arbitrary 2-category in serif capitals, e.g. D and X, morphisms of a category or a 2-category in serif lower case, e.g. f and g, and natural transformations and other 2-morphisms in Greek script.

While we would philosophically lean toward diagrammatic order of composition, we cannot bring ourselves to apply functions or functors on the left. So, in the Cat-centric first four sections of the paper, the juxtaposition FG means to apply F after G, in the classical order. In Section 5, however, we will be working in an abstract 2-category with no ability to apply 1-morphisms to elements of their domain, and so we will use the diagrammatic order of composition. We highlight the intent to compose diagrammatically with  $\cdot$  but sometimes suppress the  $\cdot$  in more complicated composites when no ambiguity can arise. We make no notational distinction between horizontal and vertical composition of 2-morphisms, taking the perspective that both are just special cases of the fundamental composition operation—that of pasting—on a 2-category.

**Acknowledgments** Arlin, Fairbanks, and Patterson acknowledge support from DARPA Award HR00112220038. Hosgood thanks Evan Cavallo and Ivan di Liberti for helpful conversations.

# 2 Diagram categories

Let X be an arbitrary category.

**Definition 2.1.** The (oplax) *category of diagrams in* X, or *diagram category*, denoted  $Diag_{\rightarrow}(X)$ , is defined as follows:

- the objects are diagrams D: J → X indexed by a small category J, which we denote by (J, D) (or, more often, just D when J is implicit);
- the morphisms  $(J, D) \rightarrow (K, E)$  are pairs  $(R, \rho)$ , where  $R: J \rightarrow K$  is a functor and  $\rho: D \Rightarrow ER$  is a natural transformation. (See Figure 1.)

Dually, we define the contravariant category of diagrams  $Diag_{\leftarrow}(X)$  as  $Diag_{\rightarrow}(X^{op})$ . Thus  $Diag_{\leftarrow}(X)$  has the same objects as  $Diag_{\rightarrow}(X)$ , but have morphisms  $(J, D) \rightarrow (K, E)$  given by pairs  $(R, \rho)$ , where  $R: K \rightarrow J$  is a functor and  $\rho: DR \Rightarrow E$  is a natural transformation. (See Figure 1).

We say that a morphism  $(R, \rho)$  in  $\text{Diag}_{\rightarrow}(X)$  or  $\text{Diag}_{\leftarrow}(X)$  is *pseudo*<sup>1</sup> if  $\rho$  is a natural isomorphism and is *strict* if  $\rho$  is an identity; we write  $\text{Diag}_{\rightarrow}^{\text{ps}}(X)$  and  $\text{Diag}_{\rightarrow}^{\text{strict}}(X)$  for the wide subcategories consisting of pseudo and strict morphisms, respectively.



Figure 1: *Left:* a morphism in  $Diag_{\rightarrow}(X)$ . *Right:* a morphism in  $Diag_{\leftarrow}(X)$ .

**Remark 2.2.** The construction of the diagram categories is functorial: given a functor  $F: X \to Y$ , we get a functor  $\text{Diag}_{\to}(F): \text{Diag}_{\to}(X) \to \text{Diag}_{\to}(Y)$  by post-composition with F, sending (J, D) to  $(J, F \circ D)$  and acting on morphisms  $(R, \rho)$  by whiskering with  $\rho$ . The same is true for  $\text{Diag}_{\leftarrow}$ .

It is possible to give  $Diag_{\rightarrow}$  and  $Diag_{\leftarrow}$  a universal property as a kind of categorified slice category. Much as an ordinary slice category may be seen as the limit of a span in the 2-category Cat, weighted by a certain span  $\mathbb{1} \to \mathbb{2} \leftarrow \mathbb{1}$  of categories, so can the diagram category be seen as the limit of a span in V weighted by the same span  $\mathbb{1} \to \mathbb{2} \leftarrow \mathbb{1}$  viewed in V, where V is the category OpLaxGray of 2-categories enriched with the oplax Gray tensor product. We shall make no use of this fact, but point it out as an amusing reflection on the depth of vision of Eilenberg and Mac Lane's original paper, and perhaps as motivation for the work of this paper in getting *out* of the relatively mysterious diagram category and into the more familiar slice Cat/X.

<sup>&</sup>lt;sup>1</sup>One could also use the terminology "*strong*" but we opt for convention most common in the 2-categorical literature.

Note that the wide subcategory  $\text{Diag}^{\mathrm{strict}}_{\to}(X)$  consisting only of strict morphisms is precisely the slice category Cat/X, and similarly for  $\text{Diag}^{\mathrm{strict}}_{\leftarrow}(X)$  and  $(\text{Cat}/X)^{\mathrm{op}}$ . Thus  $\text{Diag}^{\mathrm{strict}}_{\leftarrow}(X) \cong \text{Diag}^{\mathrm{strict}}_{\to}(X)^{\mathrm{op}}$ . We can generalise this correspondence to the case of pseudo morphisms using the following definition.

**Definition 2.3.** The *mate* of a morphism  $(R, \rho): D \to E$  in  $\text{Diag}_{\to}^{ps}(X)$  is the morphism  $(R, \rho^{-1}): E \to D$  in  $\text{Diag}_{\leftarrow}^{ps}(X)$ . We also refer to the inverse operation, which sends a morphism  $(S, \sigma)$  in  $\text{Diag}_{\leftarrow}^{ps}(X)$  to the morphism  $(S, \sigma^{-1})$  in  $\text{Diag}_{\to}^{ps}(X)$ , as the *mate*.

It is immediately observed that the identity-on-objects functor  $\text{Diag}_{\rightarrow}^{ps}(X) \rightarrow \text{Diag}_{\leftarrow}^{ps}(X)^{op}$ that sends a morphism  $(R, \rho) \colon D \rightarrow E$  to its mate  $(R, \rho^{-1}) \colon E \rightarrow D$  induces an isomorphism of categories.

This gives us one general motivation for studying diagram categories: they are extensions of the slice category Cat/X that can describe more sensitive 2-categorical information. Objects of the diagram categories are also useful as presentations for discrete (op)fibrations, as we shall see. Furthermore, diagram categories are fundamental even at the 1-categorical level, as shown by the following lemma.

**Lemma 2.4.** Let X be a complete category. Then giving a choice of limit for every diagram in X it is equivalent to giving a functor

lim: 
$$Diag_{\leftarrow}(X) \longrightarrow X$$
.

The dual statement for cocomplete categories also holds, giving a functor colim:  $Diag_{\rightarrow}(X) \rightarrow X$ .

*Proof.* An early version of this statement appears in Eilenberg and Mac Lane's original paper on category theory [EM45, §23]; a proof can also be found in [PBHF22, Proposition 4.2].  $\Box$ 

The interest in  $\text{Diag}_{\rightarrow}(X)$  and  $\text{Diag}_{\leftarrow}(X)$  in [PBHF22] was motivated by how diagrams can be used to present systems of equations, and how lifts of diagrams then describe solutions to these systems. We will briefly recall the relevant definitions, but we refer the interested reader to *loc.cit.* for a more detailed discussion.

**Definition 2.5.** Given a diagram  $D: J \to X$  and a functor  $\pi: E \to X$ , a *lift of* D *along*  $\pi$  is a functor  $\overline{D}: J \to E$  making the triangle

$$\begin{array}{c} \mathsf{E} \\ \overline{D}, \overline{\mathcal{A}} \\ \downarrow^{\pi} \\ \mathsf{J} \xrightarrow{D} \mathsf{X} \end{array}$$

commutes. We write  $\operatorname{Lift}^{\pi} D$  for the set of lifts of D along  $\pi$ .

There is indeed a natural notion of *category* of lifts along  $\pi$ , where morphisms lie over  $id_D$ , but for our purposes  $\pi$  will always be a discrete opfibration and thus the category of lifts will be discrete.

While not all of the following definitions are standard, they are consistent with the usual ones: a functor is a discrete opfibration (in the usual sense) precisely when it is a discrete opfibration at every morphism in X.

**Definition 2.6.** Let  $\pi: \mathsf{E} \to \mathsf{X}$  be a functor. For any object  $x \in \mathsf{X}$ , we write  $\mathsf{E}_x$  for the *fibre of*  $\mathsf{E}$  *over* x, defined as the set of all objects  $\overline{x} \in \mathsf{E}$  such that  $\pi(\overline{x}) = x$ .

We say that  $\pi$  is a *discrete opfibration at a morphism*  $f: x \to y$  in X if, for every  $\overline{x} \in \mathsf{E}_x$ , there exists a unique object  $\overline{y} \in \mathsf{E}$  and a unique morphism  $\overline{f}: \overline{x} \to \overline{y}$  such that  $\pi(\overline{f}) = f$ ; we refer to  $\overline{f}$  as the (unique) *lift* of the pair  $(\overline{x}, f)$ .

Similarly,  $\pi$  is said to be a *discrete fibration at* f if  $\pi^{op} \colon E^{op} \to X^{op}$  is a discrete opfibration at f. If  $\pi$  is both a discrete fibration at f and a discrete opfibration at f, then we say that it is a *discrete bifibration at* f.



Figure 2: A discrete opfibration  $\pi$  at  $f: x \to y$ . As the picture suggests, the codomain of the lift  $\overline{f}$  is not given, but is instead part of the existence and uniqueness statement.

**Remark 2.7.** An important consequence of the uniqueness of lifts is that discrete opfibrations give *functorial* lifts: given morphisms  $f: x \to y$  and  $g: y \to z$  in X and an object  $\overline{x}$  in  $\mathbb{E}_x$ , we can lift  $(\overline{x}, f)$  to get  $\overline{f}: \overline{x} \to \overline{y}$ , and then lift  $(\overline{y}, g)$  to get  $\overline{g}: \overline{y} \to \overline{z}$ ; by uniqueness of lifts, it must be the case that  $\overline{g} \circ \overline{f} = \overline{g \circ f}$ , since the former also defines a lift of  $g \circ f: x \to z$ . This is at the heart of the proof that the Grothendieck construction gives as equivalence between the category of copresheaves on X and the category of discrete opfibrations over X.

**Remark 2.8.** We have so far used the word "lift" to mean two seemingly different things: in Definition 2.5 it means a functor making a triangle commute, and in Definition 2.6 it means an object or morphism in the fibre of a functor. But the former is actually an example of the latter, since finding a lift of  $D: J \to X$  along  $\pi: E \to X$  is exactly finding an object in the fibre of  $(\pi \circ -): [J, E] \to [J, X]$  over D. With this apology made, we shall feel free to use the same overline notation for lifts in both senses. The perspective of diagrams presenting systems of equations leads to a question, however, that remains largely invisible in standard algebraic notation for equations: there can exist diagrams that are not isomorphic but which nonetheless present systems of equations that have "the same" sets of solutions. The authors of [PBHF22] propose a definition of a *weak equivalence* of diagrams, which aims to rectify this situation by defining two diagrams to be weakly equivalent exactly when there is a pseudo map between them inducing a bijection between sets of lifts against all discrete opfibrations. As discussed in the introduction, these lifting sets are interpreted as the sets of solutions to the systems of equations presented by the two diagrams.

In this paper we somewhat weaken this definition, allowed for weak equivalences that aren't necessarily pseudo, and avoiding dependence on the mate construction in the definition. In the below definition we consider only weak equivalences in  $\text{Diag}_{\leftarrow}(X)$ , as it is this category whose morphisms induce functions between lifting sets against discrete opfibrations.

**Definition 2.9.** We say that a morphism  $(R, \rho) \colon D \to E$  in  $\text{Diag}_{\leftarrow}(X)$  is a *weak equivalence* if, for every discrete opfibration  $\pi \colon \mathsf{E} \to \mathsf{X}$ , the map  $R_* \colon \text{Lift}^{\pi} D \to \text{Lift}^{\pi} E$  induced by  $(R, \rho)$  (see below) is a bijection.

The map  $R_*$  is given by applying the discrete opfibration property of  $\pi$  to find a lift of  $\rho$  whose codomain is the image of a lift of D, as we now explain in more detail, following [PBHF22, Theorem 5.2]. A slightly different proof is given at a higher level of abstraction in Section 5.

For any  $k \in K$ , the natural transformation  $\rho$  gives us a morphism

$$\rho_k \colon DR(k) \longrightarrow E(k)$$

in X. If we have some lift  $\overline{D}: J \to E$  of  $D: J \to X$  along  $\pi$ , then we have the necessary data to apply the discrete opfibration property: there exists a unique lift  $\overline{\rho_k}$  of  $\rho_k$ , which has a codomain we shall name  $\overline{E(k)}$ , as seen here:

$$\overline{D}R(k) \xrightarrow{\overline{\rho_k}} \overline{E(k)}$$

$$\downarrow^{\pi}$$

$$DR(k) \xrightarrow{\rho_k} E(k)$$

With that done, we define the functor  $\overline{E} \colon \mathsf{K} \to \mathsf{E}$  on objects by  $\overline{E}(k) = \overline{E(k)}$ . Then, given a morphism  $g \colon k \to k'$  in K, we can apply the discrete opfibration property again: there exists a unique lift  $\overline{E(g)}$  of domain  $\overline{E}(k)$ . We call its codomain  $\widetilde{E(k')}$ :

$$\overline{E}(k) \xrightarrow{\overline{E(g)}} \widetilde{E(k')}$$

$$\downarrow^{\pi}$$

$$E(k) \xrightarrow{E(g)} E(k')$$

In order to define  $\overline{E}(g) = \overline{E(g)}$ , we need to know that  $E(k') = \overline{E}(k')$ . This follows directly from discrete opfibrations having functorial lifting (Remark 2.7), but for clarity we explain the details in this case. We already know that  $\overline{\rho_k}$  is the unique lift of  $\rho_k$  with domain  $\overline{DR}(k)$ . Similarly,  $\overline{E(g)}$  is by definition the unique lift of E(g) with domain  $\overline{E}(k)$ .

Together, this implies that  $\overline{E(g)} \circ \overline{\rho_k}$ , which has codomain  $\overline{E(k')}$ , is the unique lift of  $E(g) \circ \rho_k$  with domain  $\overline{D}R(k)$ . But the naturality of  $\rho$  says that  $E(g) \circ \rho_k = \rho_{k'} \circ DR(g)$ , and by a similar argument as above, the unique lift of  $\rho_{k'} \circ DR(g)$  with domain  $\overline{D}R(k)$  has codomain  $\overline{E}(k)$ . Thus  $\widetilde{E(k')} = \overline{E(k')}$ , as desired.

**Remark 2.10**. Note that this argument breaks down if R goes in the other direction: morphisms in  $Diag_{\rightarrow}(X)$  do not necessarily induces maps of lifts against discrete opfibrations. For this reason, here and in [PBHF22] we have been primarily interested in the category  $Diag_{\leftarrow}(X)$ .

In the above construction of  $R_*$ : Lift<sup> $\pi$ </sup>  $D \to \text{Lift}^{\pi} E$ , we have actually also constructed a natural transformation  $\overline{\rho} \colon \overline{D}R \Rightarrow \overline{E}$  by setting  $\overline{\rho}_k = \overline{\rho_k}$ , which gives us a morphism  $(R, \overline{\rho}) \colon \overline{D} \to \overline{E}$  in  $\text{Diag}_{\leftarrow}(\mathsf{E})$ . In fact, by the uniqueness of the maps constructed throughout, we have actually established the following result.

**Lemma 2.11** ([PBHF22, Theorem 5.2]). The functor  $Diag_{\leftarrow}$  preserves discrete opfibrations. That is, let  $\pi : E \to X$  be a discrete opfibration. Then the post-composition functor

$$Diag_{\leftarrow}(\pi): Diag_{\leftarrow}(E) \longrightarrow Diag_{\leftarrow}(X)$$

is also a discrete opfibration.

We can even express the definition of a weak equivalence in  $\text{Diag}_{\leftarrow}(X)$  purely in terms of properties of the functor  $\text{Diag}_{\leftarrow}(\pi)$ , avoiding explicit mention of lifts of diagrams altogether. To do so, we need to give an alternative formulation of discrete opfibrations.

**Definition 2.12.** Let  $\pi: E \to X$  be a discrete opfibration at some morphism  $f: x \to y$  in X. The discrete opfibration property of  $\pi$  induces a function between fibres which we denote by

$$f_* \colon \mathsf{E}_x \longrightarrow \mathsf{E}_y$$
$$\overline{x} \longmapsto \overline{y}$$

that sends  $\overline{x}$  to the codomain  $\overline{y}$  of the unique lift of f with domain  $\overline{x}$ .

Dually, if  $\pi$  is a discrete fibration at f, then it induces a function

$$f^* \colon \mathsf{E}_y \longrightarrow \mathsf{E}_x$$
$$\overline{y} \longmapsto \overline{x}$$

that sends  $\overline{y}$  to the domain of the unique lift  $\overline{f}$  of f with codomain  $\overline{y}$ .

┛

**Lemma 2.13.** Let  $\pi: E \to X$  be a discrete opfibration at some morphism  $f: x \to y$  in X. Then  $\pi$  is a discrete fibration (and thus a discrete bifibration) at f if and only if the induced map  $f_*: E_x \to E_y$  is a bijection.

*Proof.* If  $\pi$  is also a discrete fibration at f, then we have the induced map  $f^* \colon \mathsf{E}_y \to \mathsf{E}_x$  which sends  $\overline{y}$  to the domain of the corresponding unique lift of f. But the uniqueness of lifts implies that  $f^*f_* = \mathrm{id}_{\mathsf{E}_x}$  and  $f_*f^* = \mathrm{id}_{\mathsf{E}_y}$  and so  $f_*$  is a bijection.

Conversely, if  $f_*$  is a bijection, then define  $f^* \colon \mathsf{E}_y \to \mathsf{E}_x$  by  $f^*(\overline{y}) = f_*^{-1}(\overline{y})$ . Then  $f^*$  gives us the necessary domain for the discrete fibration property: let  $\overline{y} \in \mathsf{E}_y$  and set  $\overline{x} = f^*(\overline{y})$ ; since  $f_*(\overline{x}) = \overline{y}$  by construction, the discrete opfibration property gives a unique lift  $\overline{f} \colon \overline{x} \to \overline{y}$ .

**Corollary 2.14.** A morphism  $(R, \rho)$  in  $Diag_{\leftarrow}(X)$  is a weak equivalence if and only if, for every discrete opfibration  $\pi \colon E \to X$ , the functor  $Diag_{\leftarrow}(\pi) \colon Diag_{\leftarrow}(E) \to Diag_{\leftarrow}(X)$  is a discrete fibration (and thus a discrete bifibration, by Lemma 2.11) at  $(R, \rho)$ .

*Proof.* This follows simply by unravelling the definitions. First of all, to combine the notation of Definitions 2.9 and 2.12, notice that the objects of  $\text{Diag}_{\leftarrow}(\mathsf{E})_D$  are exactly the elements of  $\text{Lift}^{\pi} D$ , and a morphism  $(R, \rho) \colon D \to E$  in  $\text{Diag}_{\leftarrow}(\mathsf{X})$  induces the function on fibres

$$R_*: \operatorname{Diag}_{\leftarrow}(\mathsf{E})_D \longrightarrow \operatorname{Diag}_{\leftarrow}(\mathsf{E})_E$$
$$\overline{D} \longmapsto \operatorname{cod}(R, \overline{\rho})$$

where  $\overline{\rho}$  is constructed as in the proof of Lemma 2.11, and cod denotes the codomain. Note that this map  $R_*$  is exactly the map  $R_*$  in Definition 2.9.

Then, by Lemma 2.13, we know that  $\pi$  is a discrete fibration at  $(R, \rho)$  if and only if this map  $R_*$  is bijective, but this is exactly the the definition of a weak equivalence.

**Example 2.15.** Every discrete opfibration over X can be constructed out of the "representable" discrete opfibrations  $\pi_x : x/X \to X$ , where x is an object of X, via colimits. As described in the introduction, lifting a diagram D along  $\pi_x$  can be interpreted as giving an x-parameterized family of solutions to the system of equations presented by D. Since this is the most familiar notion of solution to a system of equations, it is natural to ask whether our notion of weak equivalence depends on the use of arbitrary discrete opfibrations, and in particular on whether there are non-weakly-equivalent diagrams admitting a map that does induce a bijection on lifts against representable discrete opfibrations.

Indeed, this can happen, intuitively because we are mapping *into* discrete opfibrations when constructing lifts but discrete opfibrations are built from representables via *colimits*, not via limits. For a minimal example, let X be the arrow category  $2 = (0 \rightarrow 1)$ . Then representable discrete opfibrations are given by the identity and by the inclusion  $1: \mathbb{1} \rightarrow$ X. Every morphism of diagrams induces a bijection on lifts against the identity; for the nontrivial representable discrete opfibration, a lift of  $D: J \rightarrow X$  along 1 exists if and only if D is constant at 1, in which case this lift is unique. Thus morphisms of diagrams in X which are either both constant at 1 or neither constant at 1 induce bijections on lifts against representable discrete opfibrations. However, there is a unique diagram morphism  $1 \rightarrow 1 + 1$  determined by the unique functor  $1 + 1 \rightarrow 1$ , and it certainly does not induce a bijection on lifts against 1 + 1 itself, since 1 + 1 has four lifts along itself but 1 has only two lifts along 1 + 1.

We now turn to the other diagram category,  $Diag_{\rightarrow}(X)$ . As mentioned above, it is not necessarily the case that a morphism in  $Diag_{\rightarrow}(X)$  induces a map on lifts along a discrete opfibration, so we give a name to those that do.

**Definition 2.16.** We say that a morphism  $(R, \rho)$  in  $\text{Diag}_{\rightarrow}(X)$  is a *weak equivalence* if, for any discrete opfibration  $\pi: E \rightarrow X$ , the functor  $\text{Diag}_{\rightarrow}(\pi): \text{Diag}_{\rightarrow}(E) \rightarrow \text{Diag}_{\rightarrow}(X)$  is a discrete opfibration at  $(R, \rho)$ .

The dual version of Lemma 2.11 is true in a limited sense: if we consider only pseudo morphisms, then  $Diag_{\rightarrow}$  sends discrete opfibrations to discrete fibrations. More precisely, we have the following lemma.

**Lemma 2.17.** Let  $\pi : \mathsf{E} \to \mathsf{X}$  be a discrete opfibration. Then

$$\operatorname{Diag}^{\operatorname{ps}}_{\to}(\pi) \colon \operatorname{Diag}^{\operatorname{ps}}_{\to}(\mathsf{E}) \longrightarrow \operatorname{Diag}^{\operatorname{ps}}_{\to}(\mathsf{X})$$

is a discrete fibration.

*Proof.* The idea of the proof is rather simple — take the mate of a morphism in  $\text{Diag}_{\rightarrow}^{\text{ps}}(X)$  and then apply the proof of Lemma 2.11.

We need to show that, given some (pseudo) morphism  $(S, \sigma): (D, \mathsf{J}) \to (E, \mathsf{K})$  in  $\operatorname{Diag}_{\to}^{\operatorname{ps}}(\mathsf{X})$  and a lift  $\overline{E}$  of E along  $\pi$ , there exists a unique lift  $(S, \overline{\sigma}): (\overline{D}, \mathsf{J}) \to (\overline{E}, \mathsf{K})$  in  $\operatorname{Diag}_{\to}^{\operatorname{ps}}(\mathsf{E})$  of  $(S, \sigma)$ . For any  $j \in \mathsf{J}$ , we have the morphism

$$\sigma_j \colon D(j) \longrightarrow ES(j)$$

in X. By hypothesis,  $\sigma$  is a natural isomorphism, which means that  $\sigma_j$  is an isomorphism, and thus we have an inverse morphism

$$\sigma_j^{-1} \colon ES(j) \longrightarrow D(j).$$

But now we are in exactly the same situation as in the proof of Lemma 2.11, just with the roles of D and E swapped, and so we can apply exactly the same argument to obtain

$$(S, \overline{\sigma^{-1}}) \colon (\overline{E}, \mathsf{K}) \longrightarrow (\overline{D}, \mathsf{J})$$

in  $\text{Diag}_{\leftarrow}^{\text{ps}}(\mathsf{E})$ . Since  $\overline{\sigma^{-1}}$  thus defined is a natural isomorphism, we can again take its inverse in order to obtain

$$(S,\overline{\sigma})\colon (\overline{D},\mathsf{J})\longrightarrow (\overline{E},\mathsf{K})$$

in  $Diag^{ps}_{\rightarrow}(E)$ , as desired.

We now know that, if  $\pi: E \to X$  is a discrete opfibration, then  $\text{Diag}_{\leftarrow}(\pi)$  (and thus  $\text{Diag}_{\leftarrow}^{\text{ps}}(\pi)$ , as we will explain in the proof of Corollary 2.19) is also a discrete opfibration, and  $\text{Diag}_{\rightarrow}^{\text{ps}}(\pi)$  is a discrete fibration. It is natural to then ask if  $\text{Diag}_{\leftarrow}^{\text{ps}}(\pi)$  being a discrete fibration is equivalent to  $\text{Diag}_{\rightarrow}^{\text{ps}}(\pi)$  being a discrete opfibration: do weak equivalences in  $\text{Diag}_{\leftarrow}^{\text{ps}}(X)$  and weak equivalences in  $\text{Diag}_{\rightarrow}^{\text{ps}}(X)$  coincide, up to the mate isomorphism? The following lemma and its corollary provides an affirmative answer to this question.

**Lemma 2.18.** Let  $\pi: \mathsf{E} \to \mathsf{X}$  be a discrete opfibration, and let  $(R, \rho): D \to E$  be a morphism in  $\mathsf{Diag}^{\mathrm{ps}}_{\leftarrow}(\mathsf{X})$ . Then  $\mathsf{Diag}^{\mathrm{ps}}_{\leftarrow}(\pi)$  is a discrete fibration at  $(R, \rho)$  if and only if  $\mathsf{Diag}^{\mathrm{ps}}_{\to}(\pi)$  is a discrete opfibration at  $(R, \rho^{-1})$ .

*Proof.* By Lemma 2.13, we know that  $\text{Diag}^{ps}_{\leftarrow}(\pi)$  is a discrete fibration at  $(R, \rho)$  if and only if  $(R, \rho)_*$ :  $\text{Lift}^D \pi \to \text{Lift}^E \pi$  is a bijection. The dual statement of the same lemma tells us that  $\text{Diag}^{ps}_{\to}(\pi)$  is a discrete opfibration at  $(R, \rho^{-1})$  if and only if  $(R, \rho^{-1})^*$ :  $\text{Lift}^D \pi \to \text{Lift}^E \pi$  is a bijection. Thus we need only show that  $(R, \rho)_*$  and  $(R, \rho^{-1})^*$  are the same function.

By definition, given  $\overline{D} \in \operatorname{Lift}^D \pi$ , the image  $(R, \rho)_*(\overline{D})$  is the codomain of the unique lift of  $(R, \rho)$  with domain  $\overline{D}$ , while  $(R, \rho^{-1})^*(\overline{D})$  is the domain of the unique lift of  $(R, \rho^{-1})$ with codomain  $\overline{D}$ . Since the discrete opfibration  $\pi$  is conservative, in a lift  $(R, \overline{\rho}) : \overline{D} \to \overline{E}$ of  $(R, \rho)$  we can be sure that  $\overline{\rho}$  is invertible. Thus the mate  $(R, \overline{\rho}^{-1}) : \overline{E} \to \overline{D}$  of the unique such lift must also be the unique lift determining  $(R, \rho^{-1})^*$ . Thus  $(R, \rho)_*$  and  $(R, \rho^{-1})^*$ are the same function, as desired.

**Corollary 2.19.** A morphism  $(R, \rho): D \to E$  in  $\text{Diag}_{\leftarrow}^{ps}(X)$  is a weak equivalence in the sense of Definition 2.9 if and only if its mate  $(R, \rho^{-1}): E \to D$  in  $\text{Diag}_{\rightarrow}^{ps}(X)$  is a weak equivalence in the sense of Definition 2.16.

*Proof.* We know from Corollary 2.14 that  $(R, \rho)$  is a weak equivalence if and only if, for any discrete opfibration  $\pi \colon \mathsf{E} \to \mathsf{X}$ , the discrete opfibration  $\operatorname{Diag}_{\leftarrow}(\pi) \colon \operatorname{Diag}_{\leftarrow}(\mathsf{E}) \to \operatorname{Diag}_{\leftarrow}(\mathsf{X})$  is also a discrete fibration at  $(R, \rho)$ . But any lift  $(R, \overline{\rho})$  of  $(R, \rho)$  along  $\pi$  will be a pseudo morphism in  $\operatorname{Diag}_{\leftarrow}(\mathsf{E})$ , since discrete opfibrations lift isomorphisms to isomorphisms (a consequence of functorial lifting, as in Remark 2.7) and so  $\overline{\rho}$  will be a natural isomorphism since  $\rho$  is. This means that  $\operatorname{Diag}_{\leftarrow}(\pi)$  is a discrete fibration at  $(R, \rho)$  if and only if  $\operatorname{Diag}_{\leftarrow}^{\operatorname{ps}}(\pi)$  is a discrete fibration at  $(R, \rho)$ . But then Lemma 2.18 tells us that this is equivalent to  $\operatorname{Diag}_{\to}^{\operatorname{ps}}(\pi)$  being a discrete opfibration at  $(R, \rho^{-1})$ , which is exactly the definition of  $(R, \rho^{-1})$  being a weak equivalence.

It turns out that there is a readily verifiable sufficient condition on the functor R that ensures that a morphism  $(R, \rho)$  in  $\text{Diag}_{\rightarrow}$  is a weak equivalence. That condition is *initiality*, which we now review.

### **3** Initial and relatively initial functors

In this section we will recall the definition of an initial functor in the context of orthogonal factorisation systems, and then explain how this relates to the weak equivalences from the previous section. This motivates a generalisation of initiality, leading to the definition of relatively initial morphisms. Note that throughout this section we will be working exclusively with  $Diag_{\rightarrow}$ , not  $Diag_{\leftarrow}$ .

Informally, a *zigzag* between two objects x and x', denoted  $x \leftrightarrow x'$ , is a finite sequence of morphisms  $x = x_0 \leftrightarrow x_1 \leftrightarrow x_2 \leftrightarrow \ldots \leftrightarrow x_{n+1} = x'$  in which each arrow is allowed to point in either direction (" $\leftrightarrow \in \{\leftarrow, \rightarrow\}$ "). More formally:

**Definition 3.1.** A *zigzag* between objects x and x' in a category X is a finite sequence of spans  $\{x_i \leftarrow y_i \rightarrow x_{i+1}\}_{i=0,\dots,n}$  in X such that  $x_0 = x$  and  $x_{n+1} = x'$ .

We can recover the informal definition of a zigzag from the formal one by taking some of the legs of the spans  $x_i \leftarrow y_i \rightarrow x_{i+1}$  to be identities, so that the directions of the arrows don't appear to strictly alternate.

**Definition 3.2.** A functor  $R: J \to K$  is *initial* if, for every object  $k \in K$ , the comma category R/k is *non-empty* and *connected*, that is, for every  $k \in K$ ,

- (non-empty) there exists at least one object  $j \in J$  along with a morphism  $f : Rj \to k$  in K; and
- (connected) given any two objects (j, f) and (j', f') in R/k, there exists a zigzag  $(j, f) \nleftrightarrow (j', f')$  in R/k.

More abstractly, a category X is *connected* if the right adjoint  $\pi_0$ : Cat  $\rightarrow$  Set to the chaotic/codiscrete category functor Set  $\rightarrow$  Cat sends X to a point.

A key way of thinking about initial functors is through following characterization, that restricting along an initial functor does not modify a diagram's limit.

**Lemma 3.3.** Let  $R: J \to K$  be a functor. Then R is initial if and only if, for any category X and any functor  $D: J \to X$ , the canonical morphism

$$\lim_{\mathsf{K}} D \longrightarrow \lim_{\mathsf{J}} (DR)$$

is an isomorphism whenever the limits involved exist. Furthermore, it is sufficient to establish this in the case that X = Set.

Proof. See, for example, [Rie14, Lemma 8.3.4].

Two particularly useful sufficient conditions for initiality are the following.

**Lemma 3.4.** Let  $L: J \to K$  be a left adjoint. Then L is initial.

*Proof.* One way of defining what it means for L to be a left adjoint is that the comma category L/k has a terminal object  $\varepsilon_k : LRk \to k$  for every k. Any category with a terminal object is *a fortiori* connected.

**Lemma 3.5.** Let  $R: J \to K$  be a full and essentially surjective functor. Then R is initial.

Proof. This is [PBHF22, Lemma 9.8].

The fact that a morphism  $(R, \rho)$  in  $\text{Diag}_{\rightarrow}(X)$  is a weak equivalence if R is initial is related to how initial functors and discrete opfibrations interact with each other. We can make this more precise by recalling the so-called comprehensive factorization system.

**Definition 3.6.** Let L and R be classes of morphisms in a category C. We say that (L, R) is an *orthogonal factorisation system* if

• every morphism  $f: c \to d$  in C factors as  $f = r \circ l$  for some morphism  $l \in L$  and  $r \in R$ , and the factorisation is unique up to unique isomorphism:



- L and R contains all isomorphisms;
- L and R are closed under composition.

We refer to L as the *left part* of the factorisation system, and to R as the *right part*.

**Remark 3.7.** A well-known equivalent definition says that classes of morphisms (L, R) in a category C form an orthogonal factorization system if:

- Every morphism in C factors in *some* way as an L-map followed by an R-map; and
- L and R are *orthogonal* in the sense that, whenever a solid square as below is given with ℓ ∈ L and r ∈ R, there exists a unique diagonal morphism indicated by the dashed arrow making both triangles commute.



The proof is straightforward. Assuming the assumptions just listed, if a morphism  $f: c_1 \rightarrow d_2$  factors in two ways as an  $\ell$  followed by an r, then that produces an orthogonality

situation in two different ways, producing arrows  $d_1 \leftrightarrows c_2$  which are mutually inverse isomorphisms by the usual arguments involving universal properties.



Conversely, assuming the conditions in Definition 3.6, when given an orthogonality situation as above, one factors the horizontal arrows to produce two factorizations of the path around the square and then uses essential uniqueness of the factorization to produce the desired diagonal filler.

The orthogonal factorisation system most relevant to us is the comprehensive factorisation [SW73], which will prove useful in the comparison between initial functors and weak equivalences in Cat/X in Section 4.

**Lemma 3.8.** The category Cat admits an orthogonal factorisation system (L, R), known as the comprehensive factorisation system, where L is the class of initial functors and R is the class of discrete opfibrations.

*Proof.* This is [SW73, Theorems 3 and 4], but it is interesting to note that it also follows from the proof of Lemma 3.12 (see [PBHF22, Theorem 9.12]). The idea of the factorization is to use the Yoneda embedding to interpret R as a diagram of copresheaves on K and take the colimit of this diagram, followed by the Grothendieck construction, to get the canonical discrete opfibration over K through which R factors.

The following lemma provides the link between initial functors and weak equivalences in  $Diag_{\rightarrow}$ , keeping Corollary 2.19 in mind.

**Lemma 3.9.** Let  $(R, \rho)$  be a morphism in  $\text{Diag}_{\rightarrow}(X)$ . If R is an initial functor, then  $(R, \rho)$  is a weak equivalence.

*Proof.* This is a special case of Lemma 3.12 stated below.

Although Lemma 3.9 tells us that initiality is a *sufficient* condition for a morphism to be a weak equivalence, it is not *necessary*, as demonstrated by Example 3.13 below. Indeed, this condition does not even make reference to the diagrams themselves, nor to the transformation  $\rho$ , so it is not surprising that it is too strong.

This defect is partially remedied by a more subtle version of initiality that does take into account both of these aspects.

**Definition 3.10.** Let  $(R, \rho)$ :  $(J, D) \to (K, E)$  be a morphism in  $\text{Diag}_{\to}(X)$ , and let k be an object of K. Then we define the *relative comma category*  $(R, \rho)/k$  as follows:

- its objects are pairs (j, f), where  $j \in J$  and  $f \colon Rj \to k$  in K;
- its morphisms  $(j,f) \to (j',f')$  are morphisms  $h \colon j \to j'$  in J such that the diagram



┛

commutes.

We sometimes write the objects (j,f) simply as  $Rj\xrightarrow{f}k.$ 

The relative comma category  $(R, \rho)/k$  has the same objects as the comma category R/k, but has, in general, more morphisms. Indeed, any morphism  $(j, f) \rightarrow (j', f')$  in R/k, given by the data of a morphism  $h: j \rightarrow j'$  in J, defines a morphism  $(j, f) \rightarrow (j', f')$  in  $(R, \rho)/k$ , since the naturality of  $\rho: D \Rightarrow ER$  implies that the diagram



commutes; however, the fact that the diagram



commutes (which is the case if h defines a morphism in  $(R, \rho)/k$ ) does not imply that



commutes. The latter holds only under futher assumptions, such as E being faithful and  $\rho_i$  being an epimorphism.

**Definition 3.11.** Let  $(R, \rho)$ :  $(J, D) \to (K, E)$  be a morphism in  $\text{Diag}_{\to}(X)$ . We say that the functor R is *initial relative to* the transformation  $\rho$  (or that the diagram morphism  $(R, \rho)$  is *relatively initial*) if, for all  $k \in K$ , the relative comma category  $(R, \rho)/k$  is non-empty and connected.

Relative initiality is a proper generalisation of initiality: a functor  $R: J \to K$  is initial if and only if the diagram morphism  $(R, id_R): (J, R) \to (K, id_K)$  is relatively initial, whereas there exist morphisms  $(R, \rho)$  that are relatively initial but where the functor R is not initial, such as in Example 3.13. A relatively initial morphism is still sufficient to give a weak equivalence, as the following lemma shows.

**Lemma 3.12.** Let  $(R, \rho)$  be a relatively initial morphism in  $\text{Diag}_{\rightarrow}(X)$ . Then  $(R, \rho)$  is a weak equivalence.

Proof. This is [PBHF22, Theorem 9.12].

Relatively initial morphisms satisfy some standard useful properties: they are closed under composition, and every isomorphism is relatively initial. Furthermore, relatively initial morphisms preserve limits, in much the same sense that initial functors do. For more details, see [PBHF22, Proposition 9.13 and Lemma 9.14].

**Example 3.13**. Let J be the walking arrow, or the category with two objects and one non-identity morphism, and let K be the category with two objects and two parallel non-identity morphisms:

$$\mathsf{J} = \left\{ 0 \stackrel{\alpha}{\rightarrow} 1 \right\} \quad \text{and} \quad \mathsf{K} = \left\{ 0 \stackrel{\alpha}{\Rightarrow} 1 \right\}.$$

Let  $f: x \to y$  be a morphism in some category X, and define diagrams  $D: J \to X$  and  $E: K \to X$  by

$$D(\mathsf{J}) = \left\{ x \xrightarrow{f} y \right\}$$
 and  $E(\mathsf{K}) = \left\{ x \xrightarrow{f} y \right\}.$ 

If we define  $R: J \to K$  to be the identity-on-objects functor implied by the naming of the morphisms (i.e. R sends the unique morphism  $\alpha: 0 \to 1$  in J to the morphism  $\alpha: 0 \to 1$  in J), then this defines a strict morphism of diagrams  $(R, id): (J, D) \to (K, E)$ in  $\text{Diag}_{\to}(X)$ . Then the following three things are true:

- 1. *R* is not initial; but
- 2. (R, id) is a weak equivalence; and
- 3. *R* is initial with respect to id.

 $\square$ 

The second point follows from the third point combined with Lemma 3.12, so it remains only to justify the two concerning initiality.

That R is not initial follows by considering the comma category R/1, which is nonempty but not connected: there is no morphism  $\sigma: 0 \to 0$  in K such that  $\alpha \sigma = \beta$  or  $\alpha = \beta \sigma$ , so we can never hope to find a zigzag between  $\alpha: R0 \to 1$  and  $\beta: R0 \to 1$ .

However, (R, id) is *relatively* initial, since the identity morphism  $id: 0 \to 0$  in J gives a morphism between  $\alpha: R0 \to 1$  and  $\beta: R0 \to 1$ , since the diagram



commutes. This example illustrates how relatively initiality is "initiality evaluated in X."

Importantly, although relative initiality is strictly more general than initiality and is still a sufficient condition for a morphism of diagrams to be a weak equivalence, it is also still *not necessary*.

**Lemma 3.14.** Let  $(R, \rho)$  be a morphism in  $\text{Diag}_{\rightarrow}(*)$ , where \* denotes the terminal category. Then  $(R, \rho)$  is a weak equivalence if and only if R induces a bijection on connected components, and is relatively initial if and only if it further satisfies the condition that every R/k is non-empty.

*Proof.* Since \* is the terminal category, there is an isomorphism  $\text{Diag}_{\rightarrow}(*) \cong \text{Cat.}$  So, a diagram  $D: J \to *$  is just the information of the indexing category J, and we can view any functor  $R: J \to K$  as a morphism (R, id) in  $\text{Diag}_{\rightarrow}(*)$ .

Now, a discrete opfibration over \* is precisely a discrete category, i.e., a set. This means that, given a discrete opfibration  $\pi: E \to *$ , a lift of D along  $\pi$  is simply an arbitrary function of sets  $\pi_0 J \to E$ , where  $\pi_0 J$  denotes the set of connected components of J. So asking for a morphism  $(R, \operatorname{id}): J \to K$  to be a weak equivalence is thus precisely asking that, for any set E and any function  $\overline{D}: \pi_0 J \to E$ , there exists a unique function  $\overline{E}: \pi_0 K \to E$  such that  $\overline{D} = \overline{E} \circ \pi_0 R$ . In particular, this must be true for the set  $E = \pi_0 J$  and the function  $\overline{D} = \operatorname{id}_{\pi_0 J}$ . But this then says that  $\pi_0 R : \pi_0 J \to \pi_0 K$  is a split monomorphism with unique splitting, and thus  $\pi_0 R$  is a bijection. Conversely, if  $\pi_0 R$  is a bijection, then  $(R, \operatorname{id})$  is a weak equivalence: given any function  $\overline{D}: \pi_0 J \to E$ , we obtain a lift  $\overline{E}: \pi_0 K \to E$  simply by  $\overline{D}$  with the inverse of  $\pi_0 R$ .

Now consider what it means for a morphism (R, id):  $J \to K$  to be relatively initial: the relative comma categories (R, id)/k need to be non-empty and connected for all  $k \in K$ . But the morphisms in these relative comma categories are just the morphisms in J, since the diagram in Definition 3.10 consists of objects and morphisms in X, which is here the trivial category \* and so this diagram commutes for any choice of morphism in J. In other words, (j, f) and (j', f') are in the same connected component of (R, id)/k if and only if j and j' are in the same connected component of J. So if  $\pi_0$  is a bijection *and* every (R, id)/k is non-empty, then R is relatively initial. Conversely, if R is relatively initial then every (R, id)/k is non-empty by definition, and thus  $\pi_0 R$  is surjective; but  $\pi_0 R$  is also injective, since if j and j' are in the same connected component of J then Rj and Rj' are in the same connected component of K.

# 4 Localisations of categories of diagrams

Motivated by the idea that diagrams present systems of equations, we have so far proposed a notion of equivalence between diagrams, which we call a weak equivalence in  $Diag_{\leftarrow}(X)$  (Definition 2.9); we showed that this is essentially equivalent to being a weak equivalence in  $Diag_{\rightarrow}(X)$ , at least for pseudo morphisms (Corollary 2.19); and we saw that initiality and, more generally, relative initiality are sufficient (Lemma 3.12) but not necessary (Example 3.13 and Lemma 3.14) conditions to have a weak equivalence.

Now, if we want to take seriously the consideration that two diagrams in the category X should be "the same" whenever they are weakly equivalent, then we need to formally invert the weak equivalences, i.e. freely turn them into isomorphisms. The tool to achieve this is known as *localisation* [GZ67].

**Definition 4.1.** Let W be a class of morphisms in a category C. The *localisation of* C *along* W is, if it exists, the data of a category  $C[W^{-1}]$  along with a functor  $Q: C \to C[W^{-1}]$  such that

- i. for all morphisms  $f \in W$ , the morphism Qf in  $C[W^{-1}]$  is an isomorphism;
- ii. any functor  $F: C \to D$  such that Ff is an isomorphism for all  $f \in W$  factors uniquely through  $Q: C \to C[W^{-1}]$ .

The universal property of the localisation makes  $C[W^{-1}]$  the initial category in which all morphisms in W become invertible.

The localisation is in general very complicated. One condition on the class of weak equivalences that begins to make the localisation more manageable without being too stringent is as follows.

**Definition 4.2.** Let C be a category and let W be a class of morphisms in C. We say that W satisfies 2-out-of-3 if, for any pair of composable morphisms f, g in C, if any two of f, g, and gf are in W then so too is the third.

**Example 4.3**. The prototypical class of morphisms that satisfies 2-out-of-3 is the class of isomorphisms in any category.

Since weak equivalences in  $Diag_{\leftarrow}$  are defined in terms of families of bijections, they also satisfy 2-out-of-3. The same thus holds, by Corollary 2.19, for weak equivalences in  $Diag_{\rightarrow}^{ps}$ .

**Remark 4.4.** Example 3.13 shows that initial functors do *not* satisfy 2-out-of-3, in contrast to weak equivalences.

Indeed, consider the unique identity-on-objects functor  $S \colon \mathsf{K} \to \mathsf{J}$ , which sends both  $\alpha$  and  $\beta$  to  $\alpha$ . Then one can show that S is initial, and  $SR = \mathrm{id}_{\mathsf{J}}$  is initial since identity functors always are, but we already know that R is not initial. In particular, this shows that the localisation of  $(\mathsf{Cat}/\mathsf{X})$  along the class of initial functors contains isomorphisms that do not arise directly from initial functors. This is an important point to keep in mind.

We now return to the specific case of categories of diagrams. As a preliminary remark, recall that strict diagram categories are isomorphic to slices of Cat:

┛

$$\operatorname{Diag}_{\rightarrow}^{\operatorname{strict}}(\mathsf{X}) \cong \operatorname{Cat}/\mathsf{X} \cong \operatorname{Diag}_{\leftarrow}^{\operatorname{strict}}(\mathsf{X})^{\operatorname{op}}.$$

Thus, we can talk about morphisms in Cat/X as weak equivalences in  $Diag_{\rightarrow}(X)$  or  $Diag_{\leftarrow}(X)$  by regarding them as morphisms in the relevant diagram category.

**Definition 4.5.** Write N for the class of morphisms in Cat/X given by initial functors,  $W_{\rightarrow}$  for the class of weak equivalences in  $Diag_{\rightarrow}(X)$ , and  $W_{\leftarrow}$  for the class of weak equivalences in  $Diag_{\leftarrow}(X)$ .

We also write  $W_{\rightarrow}^{\text{strict}} \subset W_{\rightarrow}$  and  $W_{\leftarrow}^{\text{strict}} \subset W_{\leftarrow}$  for the subclasses of strict morphisms. By the remark above, we can regard  $W_{\rightarrow}^{\text{strict}}$  and  $W_{\leftarrow}^{\text{strict}}$  as classes of weak equivalences in Cat/X and (Cat/X)<sup>op</sup>, respectively.

Our original motivation suggests that we are interested in the category  $\text{Diag}_{\leftarrow}(X)[W_{\leftarrow}^{-1}]$ . We know that initiality is a sufficient condition for being a weak equivalences, so this category admits a canonical map from  $(\text{Cat}/X)[N^{-1}]$ . The main theorem of this paper says that, in fact, this map is an isomorphism.

**Theorem 4.6.** The inclusion of  $Cat/X^{op}$  into  $Diag_{\leftarrow}(X)$  induces an isomorphism

$$(\mathsf{Cat}/\mathsf{X})^{\mathrm{op}}[\mathsf{N}^{-1}] \xrightarrow{\cong} \mathsf{Diag}_{\leftarrow}(\mathsf{X})[\mathsf{W}_{\leftarrow}^{-1}]$$

from the localisation of Cat/X<sup>op</sup> at the class N of initial functors to the localisation of  $Diag_{\leftarrow}(X)$  at the class  $W_{\leftarrow}$  of weak equivalences.

**Remark 4.7.** Some remarks on the nature of the category on the left-hand side of the isomorphism in Theorem 4.6 are in order. First, since Cat/X admits a lifted comprehensive factorization system, where, as in Cat, the left part consists of initial functors and

the right part of discrete opfibrations, we can express  $Cat/X[N^{-1}]$  more simply as the category of discrete opfibrations over X. Indeed, any diagram  $D: J \to X$  factors as  $\pi \circ i$ , where  $\pi$  is a discrete opfibration and i is initial, which means that once i is inverted, D becomes isomorphic to the discrete opfibration  $\pi$ . Furthermore, any morphism in Cat/X between discrete opfibrations is itself a discrete opfibration: if  $D, D': J, J' \to X$  are discrete opfibrations and  $R: J \to J'$  satisfies  $D' \circ R = D$ , then if R were not itself a discrete opfibration then factoring the initial part out of R would give a factorization of D with nontrivial initial part, which contradicts the essential uniqueness of the comprehensive factorization.

The conclusion is that  $Cat/X[N^{-1}]$  is equivalent to the category Dopf/X, where Dopf is the wide subcategory of Cat spanned by the discrete opfibrations. It is well-known that, via the Grothendieck construction, Dopf/X is itself equivalent to the category of copresheaves  $X \rightarrow Set$ . Thus the category of systems of equations in X (localised along initial functors) can also be viewed simply as the category of copresheaves on X; in particular, this category forms a topos, which demonstrates the exceptionally rich structure emerging from the categorification process relative to the traditional conception of equations.

That said, we do not prefer to think of  $Cat/X[N^{-1}]$  as the category of copresheaves on X, since the objects of this category are difficult to specify in a finitary way in the case where X is not itself finitely presented (which is our baseline). Instead, we prefer to consider how to express  $Cat/X[N^{-1}]$  most simply without changing the objects, since diagrams in X are much more amenable to finite specification in practical cases. To that end, we observe that the hom-set  $Cat/X[N^{-1}](D: J \to X, D': J' \to X)$  between any two diagrams is in bijection with the hom-set Dopf/X( $\pi: J \to X, \pi': J' \to X$ ), where  $\pi$  and  $\pi'$ are the discrete opfibrations generated by D and D' (respectively) via the comprehensive factorization system. This bijection is naturally induced by composition with  $i, i': J \rightarrow J$  $J, J' \rightarrow \overline{J'}$  respectively, which are the initial parts of the same comprehensive factorisations. Thus, each morphism in Cat/X[N<sup>-1</sup>]( $D: J \to X, D': J' \to X$ ) may be canonically written as the composite of i, a discrete opfibration over X, and  $(i')^{-1}$ ; composing the first two morphisms in Cat/X, we obtain a unique expression of such a morphism as a zigzag  $D \to \pi' \leftarrow D'$ : i'. Again, we shall often prefer not to calculate  $\pi'$  explicitly, so that the real import of this argument is that every morphism in  $Cat/X[N^{-1}](D: J \to X, D': J' \to X)$ may be written in *some* manner as a zigzag  $D \rightarrow D'' \leftarrow D'$ , where the reversed arrow is determined by an initial functor between the domains of D'' and D'. It is this expression that we expect to find most useful for applications. ∟

The rest of this section will be dedicated to a proof of Theorem 4.6, which we break down into four steps:

Step 1. In Cat/X, inverting initial functors is the same as inverting weak equivalences:

$$(\mathsf{Cat}/\mathsf{X})[\mathsf{N}^{-1}] \xrightarrow{\cong} (\mathsf{Cat}/\mathsf{X})[(\mathsf{W}_{\rightarrow}^{\mathrm{strict}})^{-1}]$$

Step 2. Inverting weak equivalences in Cat/X commutes with taking the opposite category:

 $(\mathsf{Cat}/\mathsf{X})[(\mathsf{W}_{\rightarrow}^{\mathrm{strict}})^{-1}]^{\mathrm{op}} \xleftarrow{\cong} (\mathsf{Cat}/\mathsf{X})^{\mathrm{op}}[(\mathsf{W}_{\leftarrow}^{\mathrm{strict}})^{-1}]$ 

Step 3. Inverting (strict) weak equivalences in  $(Cat/X)^{op}$  is the same as inverting them in all of  $Diag_{\leftarrow}(X)$ :

$$(\mathsf{Cat}/\mathsf{X})^{\mathrm{op}}[(\mathsf{W}_{\leftarrow}^{\mathrm{strict}})^{-1}] \xrightarrow{\cong} \mathsf{Diag}_{\leftarrow}(\mathsf{X})[(\mathsf{W}_{\leftarrow}^{\mathrm{strict}})^{-1}]$$

Step 4. In  $Diag_{\leftarrow}(X)$ , inverting strict weak equivalences is the same as inverting all weak equivalences:

$$\mathsf{Diag}_{\leftarrow}(\mathsf{X})[(\mathsf{W}_{\leftarrow}^{\mathrm{strict}})^{-1}] \xrightarrow{\cong} \mathsf{Diag}_{\leftarrow}(\mathsf{X})[\mathsf{W}_{\leftarrow}^{-1}].$$

The composition of these four isomorphisms will give us a proof of Theorem 4.6.

#### Step 1. Initial functors and weak equivalences in Cat/X

The key technical observation for this step is the following.

**Lemma 4.8.** Let  $P : E \to X$  and  $Q : F \to X$  be discrete opfibrations. Suppose we have  $R : E \to F$  such that  $R: (E, P) \to (F, Q)$  is a weak equivalence in Cat/X. Then R is an isomorphism of categories.

*Proof.* Since R is a weak equivalence, it induces a bijection  $R_*$ : Lift<sup> $\pi$ </sup>  $P \to$  Lift<sup> $\pi$ </sup> Q for every discrete optibration  $\pi$  over X. In particular we may consider  $\pi = Q$  and consider the lift id<sub>F</sub> of Q through itself, shown below:

$$\begin{array}{c}
\mathsf{E} \xrightarrow{R} \mathsf{F} \\
s & \downarrow Q \\
\mathsf{F} \xrightarrow{\mathrm{id}_{\mathsf{F}}} & \downarrow Q \\
\mathsf{F} \xrightarrow{Q} & \mathsf{X}
\end{array}$$

Then, since  $R_*$  is a bijection, we can find a unique S such that  $RS = id_F$  (and, in this case redundantly, PR = Q.) Thus R is a split epimorphism and S is a split monomorphism. Now, since R and  $id_F$  are weak equivalences, so is S by two-out-of-three, so that we can repeat the same argument with S in place of R to show that S is also a split epimorphism. Therefore, S is an isomorphism, and finally so is R.

We can now apply this lemma to prove the following.

**Lemma 4.9.** Let  $L_N: \operatorname{Cat}/X \to (\operatorname{Cat}/X)[N^{-1}]$  be the localisation functor that inverts the class of initial functors. Then  $L_N$  sends weak equivalences to isomorphisms.

*Proof.* Let  $R: (J, D) \to (K, E)$  be a weak equivalence in Cat/X.

Using the comprehensive factorisation system (Lemma 3.8), we can factor E as PI, with  $I: \mathsf{K} \to \mathsf{L}$  initial and  $P: \mathsf{L} \to \mathsf{X}$  a discrete opfibration. This gives us the commutative diagram



Since I is initial, it is a weak equivalence, by Lemma 3.9, and R is a weak equivalence by assumption; thus the composite IR is also a weak equivalence. It thus suffices to show that the composite IR is initial, since then  $(IR)^{-1}I$  will be an inverse for R in  $(Cat/X)[N^{-1}]$ .

If we again apply the comprehensive factorisation system to factor  $D: J \to X$  as QJ, with  $J: J \to M$  initial and  $Q: M \to X$  a discrete opfibration, then we obtain the commutative square

$$\begin{array}{c}
\mathsf{J} \xrightarrow{IR} \mathsf{L} \\
\mathsf{J} \downarrow \qquad \qquad \downarrow P \\
\mathsf{M} \xrightarrow{Q} \mathsf{X}
\end{array}$$

The unique lifting property of orthogonal factorisation systems tells us that the square above has a unique diagonal filler  $F: M \to L$ . In particular, we find that IR = FJ as morphisms  $D \to P$  in Cat/X:



Finally, since IR is a weak equivalence, and since J is initial and is thus a weak equivalence, applying 2-out-of-3 again tells us F is also a weak equivalence. We can now apply Lemma 4.8, which tells us that  $F: (M, Q) \to (L, P)$ , as a weak equivalence between two discrete opfibrations, is an isomorphism. But then IR = FJ is the composite of an initial functor J with an isomorphism F, and so is also initial, as was to be shown.

**Corollary 4.10**. The canonical functor

$$(\mathsf{Cat}/\mathsf{X})[\mathsf{N}^{-1}] \longrightarrow (\mathsf{Cat}/\mathsf{X})[(\mathsf{W}_{\rightarrow}^{\mathrm{strict}})^{-1}]$$

is an isomorphism.

*Proof.* We know from Lemma 3.9 that  $N \subseteq W_{\rightarrow}^{\text{strict}}$ , so inverting  $W_{\rightarrow}^{\text{strict}}$  will in particular invert N. Conversely, Lemma 4.9 tells us that inverting N also inverts  $W_{\rightarrow}^{\text{strict}}$  in the process. All together then, inverting N is equivalent to inverting  $W_{\rightarrow}^{\text{strict}}$ .

#### Step 2. Weak equivalences in Cat/X and (Cat/X)<sup>op</sup>

In Corollary 2.19, we showed that taking mates induces a correspondence between weak equivalences in  $\text{Diag}^{\text{ps}}_{\rightarrow}(X)$  and weak equivalences in  $\text{Diag}^{\text{ps}}_{\leftarrow}(X)$ . In fact, the mate actually gives an isomorphism of categories

$$\mathsf{Diag}^{\mathrm{ps}}_{\to}(\mathsf{X}) \cong \mathsf{Diag}^{\mathrm{ps}}_{\leftarrow}(\mathsf{X})^{\mathrm{op}}.$$

Corollary 2.19 then tells us that this induces an isomorphism of wide subcategories between pseudo morphisms that are weak equivalences in  $\text{Diag}^{ps}_{\rightarrow}(X)$  and pseudo morphisms that are weak equivalences in  $\text{Diag}^{ps}_{\leftarrow}(X)$ .

It follows immediately from the definition of localisation of categories that an isomorphism  $(C, A) \cong (D, B)$  of pairs of a category and a wide subcategory induces an isomorphism  $C[A^{-1}] \cong D[B^{-1}]$  of localisations. Restricting from pseudo morphisms to strict morphisms, this means that we have proved the following.

Corollary 4.11. The canonical functor

$$(\mathsf{Cat}/\mathsf{X})^{\operatorname{op}}[(\mathsf{W}_{\leftarrow}^{\operatorname{strict}})^{-1}] \longrightarrow (\mathsf{Cat}/\mathsf{X})[(\mathsf{W}_{\rightarrow}^{\operatorname{strict}})^{-1}]^{\operatorname{op}}$$

is an isomorphism.

#### Step 3. Weak equivalences in Cat/X, strict weak equivalences in $Diag_{\leftarrow}(X)$

The key technical observations for this step are the following lemmas, where  $D: J \rightarrow X$  is an object of Cat/X throughout.

Given a cospan  $F : A \to C \leftarrow B : G$ , we shall use the below notations for the comma category and the components of its canonical cone:

$$\begin{array}{c} F/G \xrightarrow{\pi_{/G}} \mathsf{B} \\ \pi_{F/\downarrow} \xrightarrow{\omega_{F/G}} & \downarrow_{G} \\ \mathsf{A} \xrightarrow{F} \mathsf{C} \end{array}$$

In the case that  $F = id_{\mathsf{C}}$ , we'll write  $\mathsf{C}/G$  and  $\pi_{\mathsf{C}/}$ , and similarly if  $G = id_{\mathsf{C}}$ .

For clarity and compatibility with the 2-categorical section later on, we work mainly with the universal property of the comma category: namely, that for any cospan (F,G) as above and for any D, the hom-category Cat(D, F/G) is canonically isomorphic to the category whose objects are triples  $(A: D \to A, B: D \to B, \gamma: FA \to GB)$  and whose morphisms  $(A, B, \gamma) \to (A', B', \gamma')$  are pairs of natural transformations  $(\alpha: A \to A', \beta: B \to B')$  such that the square below commutes:

$$FA \xrightarrow{\gamma} GB$$

$$F\alpha \downarrow \qquad \qquad \qquad \downarrow G\beta$$

$$FA' \xrightarrow{\gamma'} GB'$$

Specifically, this isomorphism is induced by sending a functor  $T: D \to F/G$  to the triple  $(\pi_{F/T}, \pi_{/G}T, \omega_{F/G}T)$  and a natural transformation  $\tau: T \Rightarrow T': D \to F/G$  to the pair  $(\pi_{F/T}, \pi_{/G}\tau)$ .

**Lemma 4.12.** The canonical inclusion of Cat/X into Diag $_{\rightarrow}(X)$  has a right adjoint given on objects by sending  $D: J \rightarrow X$  to the canonical projection functor  $\pi_{X/}: X/D \rightarrow X$ .

Dually, the canonical inclusion of  $(Cat/X)^{op}$  into  $Diag_{\leftarrow}(X)$  has a left adjoint given on objects by sending  $D: J \to X$  to the canonical projection functor  $\pi_{/X}: D/X \to X$ .

*Proof.* We want to construct a natural isomorphism

$$\mathsf{Diag}_{\to}(\mathsf{X})\left(\mathsf{J}\xrightarrow{D}\mathsf{X},\mathsf{K}\xrightarrow{E}\mathsf{X}\right)\cong\mathsf{Cat}/\mathsf{X}\left(\mathsf{J}\xrightarrow{D}\mathsf{X},(\mathsf{X}/E)\xrightarrow{\pi_{\mathsf{X}/}}\mathsf{X}\right).$$

By the universal property described just above the statement of the lemma, to give a functor  $S: J \to (X/E)$  it is equivalent to specify the functors  $\pi_{X/S}: J \to X$  and  $\pi_{/E}S: J \to K$ , and the natural transformation  $\omega_{X/E}S: \pi_{X/S} \Rightarrow E\pi_{/E}S$ . Such an S gives a morphism in Cat/X if and only if  $\pi_{X/S} = D$ , so that a morphism in Cat/X( $D, \pi_{X/}$ ) is precisely given by a functor  $R = \pi_{/E}S: J \to K$  and a natural transformation  $D \Rightarrow ER$ , which is the same as the data of a morphism on the left-hand side.

**Lemma 4.13.** Consider the functor  $\iota_{/D} : J \to X/D$  determined by

$$\pi_{\mathbf{X}/\ell/D} = D, \quad \pi_{/D}\ell_{/D} = \mathrm{id}_{\mathbf{J}}, \quad \omega_{\mathbf{X}/D}\ell_{/D} = \mathrm{id}_{D}.$$

Then  $\iota_{/D}$  is right adjoint to  $\pi_{/D}$ .

Dually,  $\pi_{D/}: D/X \to J$  has a left adjoint  $\iota_{D/}$  that is a monomorphism that splits  $\pi_{D/}$  and defines a strict morphism of diagrams  $(J, D) \to (D/X, \pi_{/X})$ .

Before we begin the proof, note that by definition,  $\iota_{/D}$  is a split monomorphism splitting  $\pi_{/D}$ . Furthermore,  $\iota_{/D}$  defines a strict morphism  $(J, D) \rightarrow (X/D, \pi_{X/})$  in Cat/X.

*Proof.* Define the counit  $\pi_{/D}\iota_{/D} \to \operatorname{id}_J$  to be the identity of  $\operatorname{id}_J$ , since  $\pi_{/D}$  splits  $\iota_{/D}$ . Next, the unit must be a map  $\eta : \operatorname{id}_{X/D} \to \iota_{/D}\pi_{/D}$ . First define  $\pi_{/D}\eta$  as  $\operatorname{id}_{\pi_{/D}}$ . For  $\pi_{X}/\eta$ , we need a natural transformation  $\pi_{X/} \to \pi_{X/}\iota_{/D}\pi_{/D} = D\pi_{/D}$ ; we are given such a natural transformation in  $\omega_{X/D}$ . This determines a morphism  $\eta$  as desired since  $\omega_{X/D} \operatorname{id}_{X/D} = \omega_{X/D}$  while  $\omega_{X/D}\iota_{/D} \circ \pi_{/D} = \operatorname{id}_{\pi_{/D}}$ .

The triangle identities will now follow once we check that  $\pi_{/D}\eta$  and  $\eta\iota_{/D}$  are isomorphisms, see [RV22, B.4.2]. But we defined  $\pi_{/D}\eta$  as  $\mathrm{id}_{\pi/D}$ . Meanwhile,  $\eta\iota_{/D} : \iota_{/D} \Rightarrow \iota_{/D} \circ \pi_{/D} \circ \iota_{/D} : J \to X/D$  will be an isomorphism as soon as its whiskerings with  $\pi_{/D}$  and  $\pi_{X/}$  are so. Again, the whiskering with  $\pi_{/D}$  is invertible by assumption; finally  $\pi_{X/}\eta\iota_{/D} = \omega_{X/D}\iota_{/D} = \mathrm{id}_{D}$ , and the claim is proven.

Of these two adjoint pairs,  $\pi_{/J} \dashv \iota_{/J}$  and  $\iota_{J/} \dashv \pi_{J/}$ , we are interested in the latter. This is because, although  $\iota_{/J}$  and  $\iota_{J/}$  both define strict morphisms of diagrams, it is only the latter which is a *left* adjoint and thus initial (Lemma 3.4). This allows us to prove the following.

**Corollary 4.14.** Every canonical projection functor  $\pi_{D/}: D/X \to J$  is a weak equivalence, as is its left adjoint  $\iota_{D/}: J \to D/X$ .

*Proof.* Firstly, the fact that  $\iota_{D/}$  is a weak equivalence follows from the fact that it is a left adjoint, thus initial (Lemma 3.4), thus a weak equivalence (Lemma 3.9. Then, to show that  $\pi_{D/}$  is a weak equivalence, note that  $\pi_{D/}\iota_{D/} = \operatorname{id}_J$ . But  $\operatorname{id}_J$  is a weak equivalence, so 2-out-of-3 tells us that  $\pi_{D/}$  is also a weak equivalence.

The above lemmas allow us to prove the main result of this step using the following approach.

Consider the commutative diagram

where L and M are the canonical localisation functors, i is the inclusion, and i' is the inclusion induced by i and functoriality of localisation. We want to show that i' is actually an isomorphism of categories, and we will do so by constructing a functor  $P: \operatorname{Diag}_{\leftarrow}(X) \to (\operatorname{Cat}/X)^{\operatorname{op}}[W_{\operatorname{strict}}^{-1}]$  such that the diagram



commutes. We will then show how this implies that P induces a functor P' that is inverse to i', i.e. such that

also commutes.

**Lemma 4.15.** There exists a functor P:  $Diag_{\leftarrow}(X) \rightarrow (Cat/X)^{op}[(W_{\leftarrow}^{strict})^{-1}]$  such that Diagram (B) commutes.

*Proof.* We proceed in three steps: first, we construct P as a morphism of graphs (i.e. without checking for functoriality); second, we show that P is indeed functorial; finally, we show that P does indeed make Diagram (B) commute. The main idea of the proof is fairly simple: define P using the canonical factorisation given by the unit of the adjunction  $\iota_{1/} \dashv \pi_{1/}$  from Lemma 4.12.

Step 1: P as a graph morphism

On *objects*, we define  $P: \text{Diag}_{\leftarrow}(X) \to (\text{Cat}/X)^{\text{op}}[(W_{\leftarrow}^{\text{strict}})^{-1}]$  to be the identity, sending  $D: J \to X$  to itself. Now, given a *morphism*  $(R, \rho): (J, D) \to (K, E)$  in  $\text{Diag}_{\leftarrow}(X)$ , we want to define the associated morphism  $P(R, \rho)$  in Cat/X. As promised, we start by factoring  $(R, \rho)$  through the unit of the adjunction  $\iota_{J/} \dashv \pi_{J/}$  from Lemma 4.12, via the equality



Since  $\iota_{D/}$  defines a strict morphism  $D \leftarrow \pi_{/X}$  (Lemma 4.13) that is also a weak equivalence (Corollary 4.14), it is in  $W_{\leftarrow}^{\text{strict}}$  and thus inverted by M. The fact that  $\iota_{D/}$  is also split by  $\pi_{D/}$  (Lemma 4.13) then implies that  $M(\iota_{D/}, \text{id})^{-1} = M(\pi_{D/}, \omega)$ , since splittings are preserved by all functors and isomorphisms can be split only by their inverse.

Putting this all together and using also the facts that the image of i consists of the strict morphisms in  $Diag_{\leftarrow}(X)$  and that Diagram (A) commutes, we see that<sup>2</sup>

$$M(R,\rho) = M(\widehat{\rho}, \mathrm{id})M(\pi_{D/}, \omega)$$
  
=  $M(\widehat{\rho}, \mathrm{id})M(\iota_{D/}, \mathrm{id})^{-1}$   
=  $Mi(\widehat{\rho})Mi(\iota_{D/})^{-1}$   
=  $i'L(\widehat{\rho})i'L(\iota_{D/})^{-1}$ .

With this in mind, if i' is really to turn out as an isomorphism with M = i'P, we are forced to finish the definition of P as:

$$P: \operatorname{Diag}_{\leftarrow}(\mathsf{X}) \longrightarrow (\operatorname{Cat}/\mathsf{X})^{\operatorname{op}}[(\mathsf{W}_{\leftarrow}^{\operatorname{strict}})^{\operatorname{op}}]$$
$$(\mathsf{J}, D) \longmapsto (\mathsf{J}, D)$$
$$(R, \rho) \longmapsto L(\widehat{\rho})L(\iota_D/)^{-1}.$$

<sup>&</sup>lt;sup>2</sup>Recall that we are working in  $\text{Diag}_{\leftarrow}(X)$ , so although  $R = \pi_{D/} \circ \hat{\rho}$  as a *functor*, as a *diagram morphism* we write  $(R, \rho) = (\hat{\rho}, \text{id}) \circ (\pi_{D/}, \omega)$ .

#### Step 2: Checking that P is a functor

When  $(R, \rho) = (id_J, id_D)$  is an identity morphism in  $Diag_{\leftarrow}(X)$ , the associated functor  $\hat{\rho}$  is simply  $\iota_{J/}$  itself, which shows that P preserves identities.

Now consider a composable pair of morphisms in  $Diag_{\leftarrow}(X)$ , as displayed below.



By definition, we must compare the two right-hand sides below:

$$P(S,\sigma)P(R,\rho) = L(\widehat{\sigma})L(\iota_{E/})^{-1}L(\widehat{\rho})L(\iota_{D/})^{-1}$$
  
$$P((S,\sigma)(R,\rho)) = P(RS,\sigma\rho) = L(\widehat{\sigma\rho}) \circ L(\iota_{D/})^{-1}.$$

Cancelling the isomorphism  $L(\iota_{D/})^{-1},$  it suffices to show that

$$L(\widehat{\sigma})L(\iota_{E/})^{-1}L(\widehat{\rho}) = L(\widehat{\sigma\rho}).$$

Working now in Cat/X, rather than its opposite, we have the four functors



and we want to show that the two paths from L to D/X, when we invert  $\iota_{K/}$ , are the same. For this, it suffices to construct a morphism between these two cospans in Cat/X, i.e. a functor  $T: E/X \to D/X$  such that



commutes. Indeed, from  $T\iota_{E/} = \hat{\rho}$  we will conclude  $L(\hat{\rho})L(\iota_{E/})^{-1} = L(T)$  and thus, from  $T\hat{\sigma} = \widehat{\sigma\rho}$ , we'll find  $L(\hat{\rho})L(\iota_{E/})^{-1}L(\hat{\sigma}) = L(T)L(\hat{\sigma}) = L(\widehat{\sigma\rho})$ , as desired.

We now define the necessary functor T as induced by the lax natural transformation



of spans, where unlabelled arrows are identities. Explicitly this means we have

$$\pi_{D/T} = R\pi_{E/}, \quad \pi_{/\mathsf{X}}T = \pi_{/\mathsf{X}}, \quad \alpha_{D/\mathsf{X}}T = \alpha_{E/\mathsf{X}}\rho,$$

as pictured below.

$$E/\mathsf{X} \xrightarrow{T} \underbrace{\longrightarrow} D/\mathsf{X}$$

$$\pi_{E/} \downarrow \xrightarrow{\alpha_{E/X}} \operatorname{id}_{\mathsf{X}} \xrightarrow{\alpha_{D/X}} \downarrow \pi_{D/}$$

$$\mathsf{K} \xrightarrow{\Lambda \rho}_{R} J$$

To see that T indeed gives a morphism of cospans, note firstly that  $T\hat{\sigma}\colon\mathsf{L}\to D/\mathsf{X}$  is induced by the pasting



Thus we have  $\pi_{D/T}\hat{\sigma} = RS, \pi_{/X}T\hat{\sigma} = F$ , and  $\alpha_{D/X} * (T\hat{\sigma}) : DRS \Rightarrow F = \sigma(\rho S)$ , which is precisely the definition of  $\widehat{\sigma\rho}$ .

The other commutative triangle is similar. We see that  $T\iota_{E/}$  is induced from the pasting



which also yields  $\hat{\rho}$ . Thus *P* is indeed a functor.

Step 3: P lifts M across i' and extends L along i

Finally we need to show that the functor  $P: \text{Diag}_{\leftarrow}(X) \to (\text{Cat}/X)^{\text{op}}[(W_{\leftarrow}^{\text{strict}})^{-1}]$  thus defined does indeed make Diagram (B) commute. We defined P exactly to get i'P = M, so it remains only to check that Pi = L.

If  $R: (\mathsf{J}, D) \to (\mathsf{K}, E)$  is a morphism in  $(\mathsf{Cat}/\mathsf{X})^{\mathrm{op}}$ , then by definition of P we have

$$P(i(R)) = P(R, \mathrm{id}_E)$$
  
=  $L(\widehat{\mathrm{id}_E})L(\iota_{D/})^{-1}.$ 

So we want to show that  $L(R) = L(\widehat{id_E})L(\iota_D)^{-1}$ , or, equivalently, that

$$L(R)L(\iota_D) = L(\widehat{\mathrm{id}_E}).$$

But  $L(R)L(\iota_{D/}) = L(R\iota_{D/})$ , and so it suffices to show that (switching to Cat from  $(Cat/X)^{\text{op}}$ ) we have  $\iota_{D/R} = \widehat{\operatorname{id}_E}$  as functors  $K \to D/X$ .

We know that  $id_E$  is determined by E, R, and  $id_E$ . Meanwhile  $\iota_{D/R}$  is determined by the following diagram:



Since DR = E, we see  $\widehat{\mathrm{id}_E} = \iota_{D/R}$ , as desired.

Corollary 4.16. The canonical functor

$$(\mathsf{Cat}/\mathsf{X})^{\mathrm{op}}[(\mathsf{W}_{\leftarrow}^{\mathrm{strict}})^{-1}] \longrightarrow \mathsf{Diag}_{\leftarrow}(\mathsf{X})[(\mathsf{W}_{\leftarrow}^{\mathrm{strict}})^{-1}]$$

is an isomorphism.

*Proof.* First of all, note that this canonical functor is exactly the functor i' in Diagram (B). Since P lands in  $(Cat/X)^{op}[(W_{\leftarrow}^{strict})^{-1}]$ , by definition it sends every morphism in  $W_{\leftarrow}^{strict}$  to an isomorphism; by the universal property of localisations, it thus factors as P = P'M, as in Diagram (C). It is then purely a matter of diagram chasing and abstract nonsense (which we spell out below) to show that this P' is the inverse to i', witnessing it as an isomorphism.

We need to show that P'i' and i'P' are both identity functors, but, again by the universal property of localisations, it suffices to show that P'i'L = L and i'P'M = M. By the commutativity of Diagram (B) and the factorisation P = P'M, we see that P'i'L = L

P'Mi = Pi; the factorisation P = P'M also implies that i'P'M = i'P. In summary then, we need to show that Pi - L

$$i' P = M$$

but both of these follow immediately from the commutativity of Diagram (B) (Lemma 4.15).  $\hfill \Box$ 

#### Step 4. Strict weak equivalences and weak equivalences in $Diag_{\leftarrow}(X)$

This step is handled rather anticlimactically relative to the previous one.

**Corollary 4.17**. The canonical functor

$$\mathsf{Diag}_{\leftarrow}(\mathsf{X})[(\mathsf{W}_{\leftarrow}^{\mathrm{strict}})^{-1}] \longrightarrow \mathsf{Diag}_{\leftarrow}(\mathsf{X})[\mathsf{W}_{\leftarrow}^{-1}]$$

is an isomorphism.

*Proof.* Recall the canonical factorisation



from the proof of Lemma 4.15. By Corollary 4.14,  $\pi_{D/}: D/X \to J$  is a weak equivalence, so 2-out-of-3 for weak equivalences says that, if  $(R, \rho)$  is a weak equivalence, then so too is  $(\hat{\rho}, id)$ . Since  $(\hat{\rho}, id)$  is strict, to show that inverting strict weak equivalences inverts R, it suffices to show that inverting strict weak equivalences inverts  $(\pi_{D/}, \omega)$ . But this follows immediately from the fact that  $\pi_{D/}$  is split by  $\iota_{D/}$ , which is strict (Lemma 4.13) and a weak equivalence (Corollary 4.14).

## 5 The general 2-categorical story

We now recapitulate the argument of Section 4 in a more abstract setting. To start, we recall how completeness and cocompleteness work for 2-categories.

Let  $\mathcal{K}$  denote an arbitrary finitely bicomplete 2-category. We take limits and colimits in 2-categories in the enriched sense, so that, for instance, for a product we require a 2-natural *isomorphism* of *categories*  $\mathcal{K}(A, B \times C) \cong \mathcal{K}(A, B) \times \mathcal{K}(A, C)$ . Thus a pair of 1morphisms into B and C lift fully uniquely to a 1-morphism into  $B \times C$ , and a 2-morphism between  $f, g: A \to B \times C$  is uniquely determined by its whiskerings with the projections to B and C. Finite completeness, in the enriched sense, amounts to the assumption that  $\mathcal{K}$  has finite products and equalisers as well as *cotensors* by the interval category  $2 = 0 \rightarrow 1$ . The cotensor  $b^2$  is a representation of the arrow category of a hom-category:  $\mathcal{K}(A, B^2) \cong$  $\mathcal{K}(A, B)^2$ . Thus a 1-morphism  $\alpha \colon A \to B^2$  corresponds uniquely to a 2-morphism between 1-morphisms from A to B, which, by abuse of notation, we'll also denote by  $\alpha \colon f \Rightarrow g \colon A \to B$ . Then a 2-morphism  $\alpha \Rightarrow \beta \colon A \to B^2$  corresponds to a commutative square in  $\mathcal{K}(A, B)$  as below:

$$\begin{array}{ccc} f & \stackrel{\alpha}{\longrightarrow} & g \\ \downarrow & & \downarrow \\ h & \stackrel{\beta}{\longrightarrow} & k \end{array}$$

Dually,  $\mathcal{K}$  is finitely cocomplete when it has finite coproducts, coequalisers, and *tensors* by 2, which by definition satisfy  $\mathcal{K}(A \otimes 2, B) \cong \mathcal{K}(A, B)^2$ .

Example 5.1. Many 2-categories of interest are finitely bicomplete. For example:

- Cat is finitely bicomplete, with tensors given by the cartesian product and cotensors by the internal hom.
- The 2-categories of multicategories, symmetric multicategories, cartesian multicategories, polycategories, etc. are also finitely bicomplete. Indeed, all these categories are finitely locally presentable, since they are the categories of models of essentially algebraic theories.

For instance, if X is a multicategory, then the cotensor  $X^2$  is the multicategory whose objects are the *unary* morphisms of X and whose multimorphisms  $X^2((f), g)$  are commutative squares of the form

$$\begin{array}{ccc} (x) & \longrightarrow z \\ (f) \downarrow & & \downarrow \\ (y) & \longrightarrow w \end{array}$$

It is interesting to note that the 2-category of multicategories is *not* monoidal closed in any obvious way specializing to its cotensoring by categories.

• The 2-category of strict algebras for a reasonable 2-monad on a reasonable 2category (say, an accessible 2-monad on a locally presentable 2-category, see [BKP89, Theorem 3.8]) and *strict* morphisms is finitely bicomplete. For instance, this includes the 2-category of monoidal categories and strict monoidal functors.

We are, naturally, more interested in 2-categories like that of monoidal categories and strong monoidal functors, cartesian categories and functors preserving finite products in the usual sense, multicategories with finite products, and other examples of 2-categories of algebras and *pseudo* morphisms over a 2-monad. Such 2-categories generally possess only *flexible* limits and *shrinkable* colimits, in the sense of [BLV23].

For instance, there is no initial object in the 2-category of monoidal categories, nor any equaliser of the two maps of cartesian categories from the terminal category to the isomorphism category  $\mathbb{I}$ . We shall later handle these cases by showing that we can cover our intended applications while considering only the strict morphisms.

We now define a notion of discrete opfibration in an arbitrary 2-category that is generally quite useful.

**Definition 5.2.** A morphism  $p: E \to B$  in a 2-category  $\mathcal{K}$  is called a *discrete opfibration* if, given a 2-morphism  $\alpha: f \Rightarrow g: X \to B$  and a 1-morphism  $\overline{f}: X \to E$  with  $\overline{f} \cdot p = f$ , there exists a unique 2-morphism  $\overline{\alpha}: \overline{f} \Rightarrow \overline{g}: X \to E$  with  $\overline{\alpha} \cdot p = \alpha$ .

If  $\mathcal{K}$  admits tensors with 2, then this definition can be re-expressed as the existence, given the solid square below, of a unique 1-morphism with the signature of the dotted arrow making both triangles commute:



where the map  $i_0: X \to X \otimes 2$  is induced by the inclusion  $0: \mathbb{1} \to 2$  of the initial object into 2.

In the language of factorisation systems, discrete opfibrations are thus precisely the morphisms orthogonal on the right to every morphism  $i_0: X \to X \otimes 2$ . A few remarks on this definition are in order.

- **Remark 5.3.** i. It is easy to check that a discrete opfibration of categories in the ordinary sense (Definition 2.6) induces a discrete opfibration in the abstract sense above (Definition 5.2).
  - ii. We do not claim that Definition 5.2 is the best notion of discrete opfibration for all purposes and all 2-categories K. A key 2-category of interest to us is that of multicategories, and there there exists a natural stricter notion of discrete opfibration, allowing for lifts against multimorphisms with a lifted domain. However, the notion used here suffices to prove our main theorem for multicategories. We thus will be able to describe the localization of the category of diagrams in a multicategory at weak equivalences defined with respect to this broader notion of discrete opfibration; we leave open the question of whether there are interesting examples of diagrams in a multicategory only weakly equivalent with respect to the stricter discrete opfibrations.

iii. While this notion of discrete opfibration for multicategories captures at least everything we intend to capture, for other 2-categories it is well-known that discrete opfibrations capture very *little* of interest. If  $\mathcal{K} = V$ -Cat for a non-cartesian symmetric monoidal V, then, in general, discrete opfibrations do not fully capture V-functors into V, which are instead modelled using discrete op-*co*fibrations, and therefore discrete opfibrations and thus the result to be proven are of limited interest; in short, the story of a diagram in such a V-category as a system of equations to be solved by lifting against a discrete opfibration does not go through.

Let us exhibit the key general example of discrete opfibrations.

**Proposition 5.4.** Given any cospan  $1 \xrightarrow{x} A \xleftarrow{f} B$  in a finitely complete 2-category  $\mathcal{K}$ , the projection from the comma object  $x/f \to B$  is a discrete opfibration.

*Proof.* Recall that the comma object is the terminal inhabitant of the right-hand side of the following situation:



Given maps from C and  $C \otimes 2$  as shown, we must show there exists a unique lift  $C \otimes 2 \rightarrow x/f$ . The given information amounts to a 2-morphism  $f : f_0 \Rightarrow f_1 : C \rightarrow A$  and a 2-morphism  $t :!_C \cdot x \Rightarrow f_0 \cdot g : C \rightarrow B$ , where  $!_C : C \rightarrow 1$  is the unique map to the terminal object. The desired map will be uniquely determined, by the universal property of x/f, by a 2-morphism  $!_{C\otimes 2} \cdot x \Rightarrow f \cdot g : C \otimes 2 \rightarrow B$ . Such a 2-morphism is itself given, by the universal property of the tensor product, by a choice of 2-morphism  $u :!_C \cdot x \Rightarrow f_1 \cdot g$  such that the square below commutes. But such a choice is visibly unique since the top leg of the square is an identity.

$$\begin{array}{cccc} !_C \cdot x & \stackrel{\mathrm{id}}{\longrightarrow} & !_C \cdot x \\ \downarrow & & \downarrow u \\ g \cdot f_0 & \stackrel{}{\xrightarrow{}} g \cdot g \rightarrow & f_1 \cdot g \end{array}$$

This result generalizes the analogous fact for coslice categories, where  $f = id_B$ . However, the sense in which x/f generalizes a coslice category can be slightly unintuitive, so let us explain it in more detail in the case of multicategories. The terminal multicategory 1 has a single object  $\star$  and a single *n*-ary morphism  $(\star) \to \star$  for every *n*. So if *X* is a multicategory, a multifunctor  $x: 1 \to X$  picks out a *monoid* in *X*, not just an object. If m is a monoid in X, then the objects of the coslice multicategory m/X are unary morphisms  $m \to a$ , and its multimorphisms  $(m/X)((m \to a), m \to b)$  are given by morphisms  $(a) \to b$  such that a square involving the monoid operation of m commutes. In contrast, if a is a mere object of a multicategory, it is represented by a functor from the multicategory  $1_{\perp}$  with one object and no non-unary morphisms, and then the analogous comma multicategory a/X also has no non-unary morphisms.

A particularly important example for applications is when X is a multicategory of vector spaces, vector bundles, or sheaves of vector spaces on some manifold. In any such case the standard notion of solution of a (system of) equation(s) corresponds to lifting against the codomain projection out of  $\mathbb{R}/X$ , where  $\mathbb{R}$  represents respectively the one-dimensional real vector space, the rank-one trivial bundle, or the sheaf of continuous real-valued functions. In every case  $\mathbb{R}$  is a monoid and so the full multicategorical slice exists, allowing us to continue with the story of solving equations in terms of lifting against discrete opfibrations.

From any notion of discrete opfibration, we immediately get a corresponding notion of initiality:

**Definition 5.5.** A morphism  $f: X \to Y$  in a 2-category  $\mathcal{K}$  is called *initial* if it is left orthogonal to all discrete opfibrations: thus if, given the solid square below in which p is a discrete opfibration, there exists a unique diagonal lift as indicated, making both triangles commute:



We are aiming to construct a comprehensive factorisation system on  $\mathcal{K}$ , and this property must hold for any putative class of initial maps — to ensure that this is the case, we have simply baked it in to the definition. While this is less satisfying than the independent definition for initial functors between categories, we can nevertheless show that initial 1-morphisms have a familiar sufficient classification:

#### **Proposition 5.6.** Every left adjoint in a 2-category $\mathcal{K}$ is initial.

*Proof.* Let  $\ell: X \to Y$  be a left adjoint in  $\mathcal{K}$ , so we can choose  $r: Y \to X$ ,  $\eta: 1_X \Rightarrow \ell \cdot r$ , and  $\varepsilon: r \cdot \ell \Rightarrow 1_Y$  satisfying the usual triangle equations. Consider also a discrete opfibration  $p: E \to B$  in  $\mathcal{K}$ , and a commutative square

$$\begin{array}{ccc} X & \stackrel{f}{\longrightarrow} & E \\ \ell \downarrow & & \downarrow^{p} \\ Y & \stackrel{g}{\longrightarrow} & B \end{array}$$

We are going to construct unique a 1-morphism  $Y \to E$  factoring  $\ell$  through f, and factoring g through p.

Uniqueness. Suppose we have two diagonal fillers k, k', so that



In particular then, we have  $\ell k = \ell k' = f$ , and thus  $r\ell k = r\ell k' = rf$ . Since kp = k'p = g, we have  $\varepsilon kp = \varepsilon k'p = \varepsilon g$ :  $rlg \Rightarrow g$ . Since p is a discrete opfibration, there can exist only one 2-morphism  $\alpha$  of domain rf such that  $\alpha p = \varepsilon g$ . We have two candidates  $\varepsilon k : rf \Rightarrow k$  and  $\varepsilon k' : rf \Rightarrow k'$  for such a 2-morphism, which implies that  $\varepsilon k = \varepsilon k'$ . In particular, the codomains of these 2-morphisms must be equal, whence k = k'.

*Existence.* There is a 1-morphism  $Y \to E$  ready to hand, namely,  $r \cdot f$ . Unfortunately, this is not quite sufficient, since the composite  $\ell r f$  need not coincide with f, nor  $rfp = r\ell g$  with g. Noting the irresistible fact that rf is correct "modulo" the unit and counit of the adjunction, however, we look to deform rf into the desired filler.

To construct the filler, we apply the discrete opfibration property to the following situation:



This produces a 2-morphism  $\alpha: rf \Rightarrow k: Y \to E$  such that  $\alpha p = \varepsilon g$ . In particular, the codomain kp of  $\alpha p$  must coincide with the codomain g of  $\varepsilon g$ , and so we see that k already satisfies the desideratum kp = g.

Now consider the following diagram:



By hypothesis,  $\eta f p = \eta \ell g$  and  $\ell \alpha p = \ell \varepsilon g$ . That is, the diagram commutes serially.

By the triangle identities, the vertical composite  $(\eta \ell g) \cdot (\ell \varepsilon g)$  is equal to  $\mathrm{id}_{\ell g}$ . In other words, the vertical composite in the upper half of the diagram is a lift of  $\mathrm{id}_{\ell g} = \mathrm{id}_{fp}$  along p with domain f. But  $\mathrm{id}_f$  itself is another such lift, and so, by the discrete opfibration property, we conclude that  $\mathrm{id}_f = (\eta f) \cdot (\ell \alpha)$ . In particular, this implies that  $\ell k = f$ , and since we already had kp = g, we have established that k gives the desired diagonal lifting.

If X is an object of any 2-category  $\mathcal{K}$ , we can define the categories  $\text{Diag}_{\rightarrow}(X)$  and  $\text{Diag}_{\leftarrow}(X)$  of diagrams in X, as well as their pseudo and strict variants, just as we did for  $X \in \text{Cat}$ . We highlight that  $\text{Diag}_{\rightarrow}(X)$  is merely a category, not an object of  $\mathcal{K}$ . The formal properties of these categories are quite similar in this generality to what we saw in Section 2. For instance:

**Lemma 5.7.** For any discrete opfibration  $p: E \to X$  in a finitely bicomplete 2-category  $\mathcal{K}$ , there is an induced discrete opfibration  $\text{Diag}_{\leftarrow}(p): \text{Diag}_{\leftarrow}(E) \to \text{Diag}_{\leftarrow}(X)$  of categories.

*Proof.* The functor itself is defined simply by whiskering, so we have only to show it is a discrete opfibration.

As in the case that  $\mathcal{K} = \text{Cat}$ , let  $d: J \to X$  and  $d': J \to X$  be diagrams in X, and let  $(r, \rho): d \to d'$  be a morphism in  $\text{Diag}_{\leftarrow}(X)$ . Let  $\overline{d}$  be a lift of d along p, so that we have the diagram



We need to show that there exists a unique lift  $\overline{\rho} \colon r\overline{d} \to \overline{d'}$  of  $\rho$ .

But we have given a 2-morphism  $\rho : rd \Rightarrow d'$  into X together with a lift rd of its domain, so the unique existence of  $\overline{\rho}$  is a direct application of the definition of discrete opfibrations in  $\mathcal{K}$ .

Given this result, we can define weak equivalences in  $X \in \mathcal{K}$  just as we did in  $X \in \mathsf{Cat.}$ 

**Definition 5.8.** A morphism  $(r, \rho): d \to d'$  in  $\text{Diag}_{\leftarrow}(X)$  is a *weak equivalence* if the function  $\overline{d} \mapsto \overline{d'}$  constructed above is a bijection for every discrete opfibration p, and similarly for pseudo morphisms in  $\text{Diag}_{\rightarrow}(X)$ .

As before, it is straightforward to establish that pseudo morphisms in  $\text{Diag}_{\leftarrow}(X)$  are weak equivalences if and only if their mates are also weak equivalences in  $\text{Diag}_{\rightarrow}(X)$ .

The diagram categories also continue to enjoy the following property:

**Lemma 5.9.** Let  $(r, \rho)$ :  $(J, d) \to (J', d')$  be a morphism in  $\text{Diag}_{\to}(X)$  such that  $r: J \to J'$  is initial. Then  $(r, \rho)$  is a weak equivalence.

*Proof.* Given a lift  $\overline{d}$  of d along some discrete opfibration  $p: E \to X$ , we need to show that there exists a unique lift  $\overline{d'}$  of d' along p.

First, we lift  $\rho$  itself. Indeed, since we're given a 2-morphism  $\rho: d \Rightarrow rd'$  into X and a lift  $\overline{d}$  of its domain, there is a unique  $\overline{\rho}: \overline{d} \Rightarrow s$  over  $\rho$ .



But now we have a square with edges r, d', p, and s, and the definition of initiality says precisely that there is a unique factorisation  $\overline{d'}: J' \to E$  of s through r which lifts d' along p.

As a final preparation for the main result of this section, we explain sufficient conditions under which  $\mathcal{K}$  admits a comprehensive factorisation system. Since initial morphisms are left orthogonal to discrete opfibrations by definition, any factorization of a morphism into an initial followed by a discrete opfibration will be unique up to unique isomorphism; but to show that such factorizations exist, we require some stronger assumptions on  $\mathcal{K}$ , namely, local presentability.

Recall that local presentability of a 2-category is closely related, but not equivalent, to local presentability of its underlying 1-category. To wit, a 2-category  $\mathcal{K}$  is locally presentable as a 2-category when it is cocomplete and admits a small set G of objects which jointly detects isomorphisms (not just equivalences) and which are finitely presentable in the 2-categorical sense. This means that the representable 2-functor corresponding to any object  $G \in G$  sends filtered colimits in  $\mathcal{K}$  to filtered colimits in Cat. Since the forgetful functor Cat  $\rightarrow$  Set preserves filtered colimits, a locally presentable 2-category is locally presentable as a 1-category. The converse holds in the context of 2-categories already assumed cocomplete, but not in general.

In any event, all of our 2-categories of interest, amounting to the algebras and strict morphisms for accessible 2-monads on categories, multicategories, or symmetric multicategories, are locally presentable, and indeed every specific example of interest is locally *finitely* presentable.

**Lemma 5.10.** If  $\mathcal{K}$  is locally presentable, then it admits a factorisation system whose left class is the class of initial morphisms and whose right class is the class of discrete opfibrations.

*Proof.* This argument is well-known to experts, particularly in the unenriched case, but we give the argument for convenience. Let  $\mathcal{K}$  be locally  $\lambda$ -presentable. The class of discrete opfibrations, having been defined in terms of a right orthogonality property, is closed

under all limits in the arrow 2-category  $\mathcal{K}^2$ . Furthermore, any morphism  $i_0: J \to J \otimes 2$  may be written as a  $\lambda$ -filtered colimit of the  $i_0$ s corresponding to  $\lambda$ -presentable objects.

This implies that a morphism  $p: E \to B$  in  $\mathcal{K}$  is a discrete opfibration if and only if it has the discrete opfibration property with respect to the morphisms  $i_0: J \to J \otimes 2$ such that J is  $\lambda$ -presentable. Indeed, the class of morphisms left orthogonal to p is closed under *all* colimits in the arrow category, not just  $\lambda$ -filtered ones.

Now we can see that discrete opfibrations are closed under  $\lambda$ -filtered colimits in  $\mathcal{K}^2$ , since, given a square

$$J \longrightarrow E \\ \downarrow^{i_0} \qquad {}^{p} \downarrow \\ J \otimes 2 \longrightarrow B$$

with  $J \lambda$ -presentable and p a  $\lambda$ -filtered colimit of discrete opfibrations, the whole square can be factored through one of the members of the  $\lambda$ -filtered colimit.

Since the class M of discrete opfibrations is closed in  $\mathcal{K}^2$  under limits and  $\lambda$ -filtered colimits, we conclude that M is a reflective subcategory of  $\mathcal{K}^2$ . It is then straightforward to check that the unit of the reflection gives the desired factorisation.

We are now in a position to reiterate the proof of Theorem 4.6 in the current more general situation.

**Theorem 5.11.** Given a locally presentable 2-category  $\mathcal{K}$  containing an object X, let  $\mathbb{N}$  be the class of morphisms in  $\mathcal{K}/X$  determined by an initial morphism in  $\mathcal{K}$ . If  $\mathbb{W}_{\leftarrow}$  denotes the weak equivalences in  $\text{Diag}_{\leftarrow}(X)$ , then the canonical functor  $(\mathcal{K}/X)[\mathbb{N}^{-1}] \to \text{Diag}_{\leftarrow}(X)[\mathbb{W}_{\leftarrow}^{-1}]$  is an isomorphism.

*Proof.* We follow the same four steps as in Section 4 for the proof of Theorem 4.6.

Step 1 really applies to any factorisation system in any category. In particular, since  $\mathcal{K}$  is locally presentable, there is a comprehensive factorisation system, and so Step 1 goes through as desired.

Step 2 is handled as in Section 4.

For Step 3, we first note that the properties given in Lemma 4.12 follow just from the universal property of the comma object, and not on its concrete description in Cat. The same is true for Lemma 4.13. Once more, the proof of Lemma 4.15 makes no use of the fact that, there, we had  $\mathcal{K} = \text{Cat.}$ 

For Step 4, again, there is nothing new to say.

To close, we explain why the above result, which computes the localisations of the diagram categories in a locally presentable 2-category, actually can be extended to handle every 2-categories like that of categories with chosen finite products and functors preserving these up to isomorphism, which is by no means locally presentable.

In all these examples, we are interested in the 2-category of algebras for an accessible 2-monad T on a locally presentable 2-category  $\mathcal{K}$  (namely,  $\mathcal{K}$  is either Cat or some flavor of multicategory.) As is well-known, in this case, the 2-category  $T-\mathcal{A}lg_{\text{strict}}$  of T-algebras and *strict* morphisms is again locally presentable. Thus Theorem 5.11 applies perfectly well to  $T-\mathcal{A}lg_{\text{strict}}$ .

Really, we are interested in T-Alg itself, where the morphisms may only be pseudo. However, under the present assumptions there is a left 2-adjoint to the inclusion of T- $Alg_{\text{strict}}$  into T-Alg, sending an algebra A to an algebra A' such that the category of strict morphisms  $A' \to B$  is isomorphic to the category of pseudo morphisms  $A \to B$ . (See [BKP89, Theorem 3.13].) Thus, when considering the category of diagrams  $d: J \to X$  in an object X of T-Alg, we can replace d with its strict adjunct  $\overline{d}: J' \to X$  and proceed using the results for T- $Alg_{\text{strict}}$ . As a practical matter, we will often have specified d to be strict in any case. Therefore the results in the strict case suffice to classify the objects of the diagram category even in the pseudo case.

# References

[BKP89]	R. Blackwell, G.M. Kelly, and A.J. Power. "Two-dimensional monad theory". Journal of Pure and Applied Algebra 59.1 (1989), pp. 1–41. ISSN: 0022-4049. DOI: https://doi.org/10.1016/0022-4049(89)90160-6.	
[BLV23]	John Bourke, Stephen Lack, and Lukáš Vokřínek. "Adjoint functor theorems for homotopically enriched categories". <i>Advances in Mathematics</i> 412 (2023). DOI: https://doi.org/10.1016/j.aim.2022.108812.	
[EM45]	Samuel Eilenberg and Saunders Mac Lane. "General theory of natural equivalences". <i>Transactions of the American Mathematical Society</i> 58 (1945), pp. 231–294. DOI: 10.1090/S0002-9947-1945-0013131-6.	
[GV77]	René Guitart and Luc Van den Bril. "Décompositions et Lax-complétions". fr. <i>Cahiers de topologie et géométrie différentielle</i> 18.4 (1977), pp. 333–407.	
[GZ67]	Peter Gabriel and Michel Zisman. <i>Calculus of fractions and homotopy theory</i> . Springer, 1967. DOI: 10.1007/978-3-642-85844-4.	
[Mes23]	Luca Mesiti. Colimits in 2-dimensional slices. 2023. arXiv: 2305.01494 [math.CT]	
[PBHF22]	Evan Patterson, Andrew Baas, Timothy Hosgood, and James Fairbanks. "A diagrammatic view of differential equations in physics". <i>Mathematics in Engineering</i> 5.2 (2022), pp. 1–59. DOI: 10.3934/mine.2023036.	
[Rie14]	Emily Riehl. <i>Categorical homotopy theory</i> . Cambridge University Press, 2014. DOI: 10.1017/CB09781107261457.	
[RV22]	Emily Riehl and Dominic Verity. <i>Elements of</i> $\infty$ -category theory. Cambridge University Press, 2022. DOI: 10.1017/9781108936880.	

- [Spi23] David I. Spivak. Functorial aggregation. 2023. arXiv: 2111.10968 [math.CT].
- [SW73] Ross Street and R.F.C. Walters. "The comprehensive factorization of a functor". *Bulletin of the American Mathematical Society* 79 (1973), pp. 936–941. DOI: 10.1090/S0002-9904-1973-13268-9.