Two step estimations via the Dantzig selector for models of stochastic processes with high-dimensional parameters

Kou Fujimori¹ and Koji Tsukuda²

¹Faculty of Economics and Law, Shinshu University. ²Faculty of Mathematics, Kyushu University.

Abstract

We consider the sparse estimation for stochastic processes with possibly infinitedimensional nuisance parameters, by using the Dantzig selector which is a sparse estimation method similar to Z-estimation. When a consistent estimator for a nuisance parameter is obtained, it is possible to construct an asymptotically normal estimator for the parameter of interest under appropriate conditions. Motivated by this fact, we establish the asymptotic behavior of the Dantzig selector for models of ergodic stochastic processes with high-dimensional parameters of interest and possibly infinitedimensional nuisance parameters. Applications to ergodic time series models including integer-valued autoregressive models and ergodic diffusion processes are presented.

1 Introduction

High-dimensional modeling in statistics has been attracted much attention over two decades. Specifically, the sparse estimation methods for high-dimensional parameter have been studied and commonly used in various fields. Most of sparse estimation procedures such as the Lasso (Tibshirani, 1996) are constructed by adding penalty terms to contrast functions in order to induce the sparsity of the parameters. On the other hand, the Dantzig selector proposed by Candes and Tao (2007) is defined as inequality constrained optimization problems. In this paper, we focus on the Dantzig selector type estimator for models of stochastic processes in semiparametric setting.

Let us begin with the following conditional heteroskedastic linear regression model:

$$Y_i = \beta^\top Z_i + \epsilon_i, \quad i = 1, \dots, n$$

where $\beta \in \mathbb{R}^p$ is a regression coefficient, $Z_i, i = 1, ..., n$ are \mathbb{R}^p -valued covariates and $\epsilon_i, i = 1, ..., n$ are independent random variables with $\mathbb{E}[\epsilon_i | Z_i] = 0$, a.s. and we put $\operatorname{Var}[\epsilon_i | Z_i] = \sigma^2(Z_i)$. The Dantzig selector type estimator $\hat{\beta}_D$ for β is then defined as follows:

$$\hat{\beta}_D := \arg\min_{\beta \in \mathcal{C}} \|\beta\|_1, \quad \mathcal{C} := \{\beta \in \mathbb{R}^p : \|\psi_n^{(1)}(\beta)\|_\infty \le \lambda\},\$$

where $\lambda \geq 0$ is a tuning parameter and

$$\psi_n^{(1)}(\beta) := \frac{1}{n} \sum_{i=1}^n Z_i (Y - \beta^\top Z_i)$$

is a score function based on the least squares method. When p is large and the true value β_0 is sparse, $\hat{\beta}_D$ behaves better than the ordinary least squares estimator under some technical conditions, such as restricted eigenvalue condition for Hessian matrix and sub-Gaussian property of the noise distribution; see, e.g., Bickel et al. (2009) for details. Using $\hat{\beta}_D$, we can consider the estimator \hat{T}_D for true support index set $T_0 := \{j : \beta_{0j} \neq 0\}$, which enables us to consider the dimension reduction. On the other hand, we can also consider the following score function:

$$\Psi_n(\beta, \sigma^2) = \frac{1}{n} \sum_{i=1}^n \frac{Z_i(Y_i - Z_i^\top \beta)}{\sigma^2(Z_i)}$$

Then, if we can construct a consistent estimator $\hat{\sigma}^2(\cdot)$ of $\sigma^2(\cdot)$ under some conditions, the estimator $\tilde{\beta}$ which is a solution to the following equation;

 $\Psi_n(\beta, \hat{\sigma}^2) \approx 0$

is more efficient than ordinary least squares in low-dimensional settings; see, e.g., Robinson (1987). Therefore, even in high-dimensional, but sparse settings, we may construct more efficient estimator by the following steps: First, construct an initial estimator and consider the variable selection via the Dantzig selector based on the score function $\psi_n^{(1)}$ for a high-dimensional parameter of interest. Second, plugging a consistent estimator for nuisance parameters in the score function Ψ_n , we construct the Z-estimator of the unknown parameter, which may be more efficient than the ordinary least squares estimator. Inspired by this ideas, we deal with semiparametric models of stochastic processes and time series in high-dimensional settings.

In Section 3, we consider the following conditionally heteroskedastic model of 1-dimensional time series with high-dimensional parameters:

$$X_t = S(\alpha^{\top} \phi(\mathbf{X}_{t-1}), \beta^{\top} Z_{t-1}) + u_t, \quad t \in \mathbb{Z},$$

where

$$\operatorname{E}[u_t|\mathcal{F}_{t-1}] = 0, \quad \operatorname{E}[u_t^2|\mathcal{F}_{t-1}] = \sigma^2(\boldsymbol{X}_{t-1};h),$$

 $X_{t-1} = (X_{t-1}, \ldots, X_{t-d}), \phi$ is a multi-dimensional functional, $\{Z_t\}$ is a high-dimensional covariate process, $\{\mathcal{F}_t\}$ is a filtration defined by $\mathcal{F}_t = \sigma(X_s, Z_s, s \leq t)$, and h is a possibly infinite-dimensional nuisance parameter. There are several previous works in which sparse estimation problems for time series models are considered. For example, Basu and Michailidis (2015) considered Lasso type estimators for a stationary Gaussian time series model in high-dimensional settings. Wong et al. (2020) also considered the

Lasso for a stationary non-Gaussian time series model in high-dimensional settings. In this paper, we consider the sparse estimation problems for conditionally heteroskedastic models and construct an asymptotically normal estimator for non-zero components of the true value of $\theta^{\top} = (\alpha^{\top}, \beta^{\top})$ based on the theory of Z-estimation and variable selection. As an example, we consider the higher order Integer-valued autoregressive model, discussed by, Al-Osh and Alzaid (1987), Du and Li (1991), Latour (1997, 1998) and so on. Moreover, we discuss an application to the estimation problems of Hawkes processes.

In Section 4, we consider models of diffusion processes with high-dimensional covari-There are some previous works in which sparse estimation problems for models ates. of diffusion processes in high-dimensional settings are discussed. Periera and Ibrahimi (2014), Gaïffas and Matulewicz (2019) and Ciołek et al. (2020) considered the (adaptive) Lasso or the Dantzig selector for drift parameters of linear models of diffusion processes such as Ornstein–Uhlenbeck processes under high-dimensional settings based on continuous observations, which do not deal with estimation problems of diffusion coefficients. Fujimori (2019) proposed the Dantzig selector for drift parameters of linear models of diffusion processes when the diffusion coefficient is a finite dimensional parameter based on discrete observations. In this paper, we focus on seimiparametric models of diffusion processes including the possibly infinite-dimensional parameter h in the diffusion coefficient, based on discrete observations and construct an asymptotically normal estimator for non-zero components of the true value of $\theta^{\top} = (\alpha^{\top}, \beta^{\top})$. As an example, we consider the high-dimensional Ornstein–Uhlenbeck processes as well as Gaïffas and Matulewicz (2019) and Ciołek et al. (2020), and discuss a sparse estimation problem based on discrete observations.

There are three major contributions in this paper. First, we provide sufficient conditions to derive the rate of convergence of the Dantzig selector for relatively general models including models of diffusion processes and time series under conditionally heteroskedasticity with possibly infinite-dimensional parameter. Second, we derive an asymptotically normal estimator for non-zero components of the high-dimensional parameter based on the theory of Z-estimators under sparse settings. Finally, we apply the general theory to the estimation problems for the integer autoregressive model of large order and the linear model of diffusion processes based on discrete observations. We verify some sufficient conditions to derive the rate of convergence of the Dantzig selector, such as some moment conditions and matrix conditions under some regularity conditions.

The rest of this paper is organized as follows. In Section 2, we present a sufficient conditions to derive the rate of convergence and selection consistency of the Dantzig selector in general settings. After the variable selection, we apply the general theory for Z-estimation with possibly infinite-dimensional nuisance parameter, which is originally established by Nishiyama (2009). See also van der Vaart and Wellner (1996) for the asymptotic theory of Z-estimation. The results for the ergodic time series are provided in Section 3, which includes a concrete example for integer-valued autoregressive models and numerical simulations. In Section 4, we present results for ergodic diffusion processes. Some proofs and technical lemmas are provided in Section 5. Especially, we use the maximal inequality provided in Nishiyama (2022) to verify one of the sufficient conditions to derive the rate of convergence of the Dantzig selector in Sections 3 and 4.

Throughout this paper, we denote by $||v||_q$ the l_q norm of a vector v and $||A||_q$ the operator norm of a matrix A for every $q \ge 1$.

For a vector $v \in \mathbb{R}^p$ and an index set $T \subset \{1, 2, \ldots, p\}$, we denote the sub-vector of v restricted by T, by v_T . Similarly, for a $p \times p$ matrix A and index sets $T, T' \subset \{1, 2, \ldots, p\}$, we denote the sub-matrix of A restricted by T and T', by $A_{T,T'} = (A_{ij})_{i \in T, j \in T'}$. For a random variable X and $q \geq 1$, we write $||X||_{L^q}$ for the L^q -norm of X if it exists; $||X||_{L^q} = \mathbb{E}[|X|^q]^{1/q}$. Moreover, we assume that all random elements are measurable in this paper.

2 Two step estimation for general models

Let (Ω, \mathcal{F}, P) be a probability space. Let $\theta \in \Theta \subset \mathbb{R}^p$ be an unknown parameter of interest and $h \in H$ a possibly infinite-dimensional nuisance parameter where H is a metric space equipped with a metric d_H . Consider the following random maps:

$$\psi_n^{(1)}: \Theta \to \mathbb{R}^p, \quad \tilde{\psi}_n^{(1)}: \Theta \to \mathbb{R}^p,$$
$$\Psi_n: \Theta \times H \to \mathbb{R}^p, \quad \tilde{\Psi}_n: \Theta \times H \to \mathbb{R}^p,$$

where n is a number of observations. The random maps $\psi_n^{(1)}$ and Ψ_n are corresponding to some score functions, and $\tilde{\psi}_n^{(1)}$ and $\tilde{\Psi}_n$ are their compensators, respectively. We suppose that the true values $\theta_0 \in \Theta$ and $h_0 \in H$ satisfy that

$$\tilde{\psi}_n^{(1)}(\theta_0) \approx 0, \quad \tilde{\Psi}_n(\theta_0, h_0) \approx 0.$$

We are interested in the estimation problem for $\theta_0 = (\theta_{01}, \ldots, \theta_{0p})^{\top}$ under the following high-dimensional and sparse settings;

- (i) The dimension p of θ possibly tends to ∞ as $n \to \infty$.
- (ii) Let T_0 be the support index set of θ_0 , i.e., $T_0 = \{j : \theta_{0j} \neq 0\}$. The sparsity s, which is the cardinality of T_0 , is smaller than n and p.

We describe the precise condition for s in the subsequent sections which discuss the applications to models of diffusion processes and time series models. We first construct an estimator for T_0 to achieve the dimension reduction. To do this, we consider the following Dantzig selector type estimator $\hat{\theta}_n^{(1)}$ for θ_0 , based on $\psi_n^{(1)}$:

$$\hat{\theta}_n^{(1)} := \arg\min_{\theta \in \mathcal{C}_n} \|\theta\|_1, \quad \mathcal{C}_n = \{\theta \in \Theta : \|\psi_n^{(1)}(\theta)\|_{\infty} \le \lambda_n\},\$$

where λ_n is a tuning parameter. Using $\hat{\theta}_n^{(1)}$, we define \hat{T}_n as follows:

$$\hat{T}_n := \{ j : |\hat{\theta}_{nj}^{(1)}| > \tau_n \},\$$

where τ_n is a threshold, which is a tuning parameter possibly depending on n, p and s. For every index set $T \subset \{1, \ldots, p\}, C_T$ be a set defined by

$$C_T := \{ v \in \mathbb{R}^p : \| v_{T^c} \|_1 \le \| v_T \|_1 \}.$$

Consider a $p \times p$ matrix $V_n^{(1)}$ which satisfies that for every $v \in C_{T_0}$ and $\theta \in \Theta$,

$$|v^{\top}[\psi_n^{(1)}(\theta) - \psi_n^{(1)}(\theta_0)]| \ge |v^{\top}V_n^{(1)}v|,$$
(2.1)

For the matrix $V_n^{(1)}$, we define the following quantity as well as Huang et al. (2013):

$$F_{\infty}(T_0, V_n^{(1)}) := \inf_{v \in C_{T_0} \setminus \{0\}} \frac{|v^\top V_n^{(1)} v|}{\|v_{T_0}\|_1 \|v\|_{\infty}}.$$
(2.2)

See also Bühlmann and van de Geer (2011) for such matrix factors. We assume the following conditions.

Assumption 2.1. (i) It holds that

$$P\left(\|\psi_n^{(1)}(\theta_0)\|_{\infty} > \lambda_n\right) \to 0, \quad n \to \infty$$

(ii) There exists a positive constant δ such that

$$P(F_{\infty}(T_0, V_n^{(1)}) > \delta) \to 1, \quad n \to \infty.$$

If $\psi_n^{(1)}(\theta_0)$ is a terminal value of square integrable martingale, we can verify the condition (i) by using, e.g., stochastic maximal inequality provided in Nishiyama (2022). The condition (ii) strongly depends on the model assumption such as stationarity, tail property, dependence structure of the processes, and the regularity condition of the information matrix. Therefore, in Sections 3 and 4, we provide the proof of the condition (i) for relatively general models of stochastic processes, while the condition (ii) is verified for more concrete models such as stationary integer-valued autoregressive models and Ornstein–Uhlenbeck processes.

Under Assumption 2.1, we establish the following theorem.

Theorem 2.2. Let Assumption 2.1 hold. Then, it holds that

$$P\left(\|\hat{\theta}_n^{(1)} - \theta_0\|_{\infty} > \frac{2}{\delta}c_n\right) \to 0, \quad n \to \infty,$$
(2.3)

where $c_n := \|\psi_n^{(1)}(\hat{\theta}_n^{(1)}) - \psi_n^{(1)}(\theta_0)\|_{\infty}$. Especially, it holds that

$$\|\theta_n^{(1)} - \theta_0\|_{\infty} = O_p(\lambda_n), \quad n \to \infty.$$

Moreover, if the threshold τ_n satisfies that $4\lambda_n/\delta < \tau_n < \inf_{j \in T_0} |\theta_{0j}|/2$, then it holds that

 $\mathbf{P}(\hat{T}_n = T_0) \to 1, \quad n \to \infty.$ (2.4)

Proof. For the proof of (2.3), it suffices to show that

$$\|\hat{\theta}_n^{(1)} - \theta_0\|_{\infty} \le \frac{2}{\delta}c_n$$

under the event that

$$\{\|\psi_n^{(1)}(\theta_0)\|_{\infty} \le \lambda_n\} \cap \left\{F_{\infty}(T_0, V_n^{(1)}) > \delta\right\}.$$

Put $v = \hat{\theta}_n^{(1)} - \theta_0$. Since $\hat{\theta}_n^{(1)} = \theta_0 + v$ is a minimizer of l_1 norm, we can show that

$$\begin{split} 0 \geq \|\theta_0 + v\|_1 - \|\theta_0\|_1 &= \sum_{j \in T_0^c} |v_{T_{0j}^c}| + \sum_{j \in T_0} (|\theta_{0j} + v_{T_{0j}}| - |\theta_{0j}|) \\ \geq \sum_{j \in T_0^c} |v_{T_{0j}^c}| + \sum_{j \in T_0} (|\theta_{0j}| - |v_{T_{0j}}| - |\theta_{0j}|) \\ \geq \sum_{j \in T_0^c} |v_{T_0^cj}| - \sum_{j \in T_0} |v_{T_{0j}}| \\ &= \|v_{T_0^c}\|_1 - \|v_{T_0}\|_1, \end{split}$$

which means $v \in C_{T_0}$. Moreover, we have

$$\|v\|_1 = \|v_{T_0^c}\|_1 + \|v_{T_0}\|_1 \le 2\|v_{T_0}\|_1.$$

Noting that $c_n := \|\psi_n^{(1)}(\hat{\theta}_n^{(1)}) - \psi_n^{(1)}(\theta_0)\|_{\infty}$, we have $|v^{\top}[\psi_n^{(1)}(\hat{\theta}_n^{(1)}) - \psi_n(\theta_0)]| \le \|v\|_1 c_n.$

Furthermore, it follows from the definitions of $V_n^{(1)}$ and $F_{\infty}(T_0, V_n^{(1)})$ that

$$|v^{\top}[\psi_n^{(1)}(\hat{\theta}_n^{(1)}) - \psi_n(\theta_0)]| \ge |v^{\top}V_n^{(1)}v| \ge F_{\infty}(T_0, V_n^{(1)}) ||v_{T_0}||_1 ||v||_{\infty} > \delta ||v_{T_0}||_1 ||v||_{\infty}.$$

Combining these facts, we have

$$\delta \|v_{T_0}\|_1 \|v\|_{\infty} \le \|v\|_1 c_n \le 2 \|v_{T_0}\|_1 c_n$$

It implies

$$\|v\|_{\infty} \le \frac{2}{\delta}c_n,$$

which concludes the proof of (2.3). Moreover, by using the triangle inequality and the definition of the $\hat{\theta}_n^{(1)}$, we have

$$c_n = \|\psi_n^{(1)}(\hat{\theta}_n^{(1)}) - \psi_n^{(1)}(\theta_0)\|_{\infty} \le 2\lambda_n,$$

which implies $||v||_{\infty} = O_p(\lambda_n)$.

Note that $\tau_n > 4\lambda_n/\delta \ge 2c_n/\delta$. To prove (2.4), it suffices to show that

$$\{\|\hat{\theta}_n^{(1)} - \theta_0\|_{\infty} \le \tau_n\} \subset \{\hat{T}_n = T_0\}.$$

Therefore, we consider the following two cases under the event $\{\|\hat{\theta}_n^{(1)} - \theta_0\|_{\infty} \le \tau_n\}$.

(i) Suppose that $j \in T_0$. Then, it holds that

$$|\theta_{0j}| - |\hat{\theta}_{nj}^{(1)}| \le |\hat{\theta}_{nj}^{(1)} - \theta_{0j}| \le \tau_r$$

under the event $\{\|\hat{\theta}_n^{(1)} - \theta_0\|_{\infty} \leq \tau_n\}$. Since $\tau_n < \inf_{j \in T_0} |\theta_{0j}|/2$, we have

$$|\hat{\theta}_{nj}| \ge |\theta_{0j}| - \tau_n > \tau_n,$$

which implies $j \in \hat{T}_n$, i.e., $T_0 \subset \hat{T}_n$.

(ii) Suppose that $j \in T_0^c$, then it holds that

$$|\hat{\theta}_{nj}^{(1)} - \theta_{0j}| = |\hat{\theta}_{nj}^{(1)}| \le \tau_n,$$

which implies $j \in \hat{T}_n^c$, i.e., $T_0^c \subset \hat{T}_n^c$.

It follows from (i) and (ii) that $\hat{T}_n = T_0$.

Remark 2.3. If we choose the tuning parameter λ_n satisfying that $\lambda_n \to 0$ as $n \to \infty$, then $\hat{\theta}_n^{(1)}$ is a consistent estimator in l_{∞} norms.

Hereafter, to discuss the asymptotically normal estimator for non-zero components of θ_0 , we assume that s is independent of n and p. For every index set T, we consider the following random maps restricted by T:

$$\Psi_{nT}:\Theta_T\times H\to \mathbb{R}^{|T|},\quad \tilde{\Psi}_{nT}:\Theta_T\times H\to \mathbb{R}^{|T|},$$

where Θ_T is a set of sub-vectors of Θ restricted by T. Let \hat{h}_n be an estimator of $h \in H$ such that $d_H(\hat{h}_n, h_0) = o_p(1)$ as $n \to \infty$. Then, we consider the new estimator $\tilde{\theta}_n$ for the parameter θ of interest, with help of \hat{h}_n and \hat{T}_n , as a solution to the following equations:

$$\Psi_{n\hat{T}_n}(\tilde{\theta}_{n\hat{T}_n},\hat{h}_n)\approx 0,\quad \tilde{\theta}_{n\hat{T}_n^c}=0.$$

Then, we establish the asymptotic normality of θ_n , which follows from Theorem 2.1 of Nishiyama (2009).

Theorem 2.4. Let Assumption 2.1 hold. Assume that s is independent of n and p. Suppose that there exist a sequence $r_n \uparrow \infty$, some fixed point (θ_0, h_0) and an invertible matrix V_{θ_0,h_0} which satisfy the following conditions.

(i) There exists a neighborhood $U \subset \Theta_{T_0} \times H$ of (θ_{0T_0}, h_0) such that

$$r_n(\Psi_{nT_0} - \Psi_{nT_0}) \rightarrow^d Z \quad in \ \ell^{\infty}(U),$$

where almost all paths $(\theta, h) \rightsquigarrow Z(\theta, h)$ are continuous with respect to $\rho = \|\cdot\|_2 \lor d_H$.

(ii) For given random sequence $(\hat{\theta}_{nT_0}, \hat{h}_n)$, it holds that

$$\tilde{\Psi}_{nT_0}(\hat{\theta}_{nT_0}, \hat{h}_n) - \tilde{\Psi}_{nT_0}(\theta_{T_0}, h_0) - V_{\theta_0, h_0}(\hat{\theta}_{nT_0} - \theta_{0T_0}) = o_p(r_n^{-1} + \|\hat{\theta}_{nT_0} - \theta_{0T_0}\|_2)$$

and that

$$\begin{aligned} \|\hat{\theta}_{nT_0} - \theta_{0T_0}\|_2 &\lor d_H(\hat{h}_n, h_0) = o_p(1), \quad \Psi_{nT_0}(\hat{\theta}_{nT_0}, \hat{h}_n) = o_p(r_n^{-1}), \\ \tilde{\Psi}_{nT_0}(\theta_{0T_0}, h_0) = o_p(r_n^{-1}) \end{aligned}$$

as $n \to \infty$.

Then, it holds that

$$r_n(\hat{\theta}_{n\hat{T}_n} - \theta_{0T_0}) \mathbf{1}_{\{\hat{T}_n = T_0\}} \to^d - V_{\theta_0,h_0} Z(\theta_{0T_0},h_0), \quad n \to \infty.$$

Proof. Note that

$$r_n(\hat{\theta}_{n\hat{T}_n} - \theta_{0T_0})\mathbf{1}_{\{\hat{T}_n = T_0\}} = r_n(\hat{\theta}_{nT_0} - \theta_{0T_0})\mathbf{1}_{\{\hat{T}_n = T_0\}}.$$

and that

$$1_{\{\hat{T}_n=T_0\}} \to^p 1, \quad n \to \infty$$

by Theorem 2.1. It follows from Theorem 2.1 of Nishiyama (2009) that

$$r_n(\hat{\theta}_{nT_0} - \theta_{0T_0}) \to^d - V_{\theta_0,h_0} Z(\theta_{0T_0},h_0), \quad n \to \infty.$$

Then, using the Slutsky lemma, we obtain the conclusion.

Remark 2.5. In Theorem 2.2, we establish the rate of convergence and the selection consistency of the Dantzig selector when s is allowed to diverge under appropriate conditions. On the other hand, to apply Theorem 2.4 directly, the sparsity s should be fixed. To consider the asymptotic normality in the case where s can be diverge, we should consider another type of asymptotic theory. See, e.g., Chernozhukov et al. (2021) for the Gaussian approximation for time series models in such a high-dimensional regime.

3 Applications to time series models

3.1 Models of ergodic time series with high-dimensional parameters

Let us consider the following time series model:

$$X_t = S(\alpha^{\top} \tilde{\phi}(X_{t-1}, \dots, X_{t-d_1}), \beta^{\top} Z_{t-1}) + u_t, \quad t \in \mathbb{Z}, \ d_1 \ge 1,$$

where S is a measurable and twice continuously differentiable function, $\alpha \in \mathbb{R}^{p_1}$, $\tilde{\phi} : \mathbb{R}^{d_1} \to \mathbb{R}^{p_1}$ is a measurable function, $\beta \in \mathbb{R}^{p_2}$, $\{Z_t\}_{t \in \mathbb{Z}}$ is an \mathbb{R}^{p_2} -valued covariate process and

 $\{u_t\}_{t\in\mathbb{Z}}$ is a square integrable martingale difference sequence with respect to the filtration $\{\mathcal{F}_t\}$ defined by

$$\mathcal{F}_t = \sigma(X_s, Z_s; s \le t), \quad t \in \mathbb{Z}.$$

We suppose that

$$\mathbb{E}[u_t^2|\mathcal{F}_{t-1}] = \tilde{\sigma}^2(X_{t-1}, \dots, X_{t-d_2}; h), \quad t \in \mathbb{Z}, \quad d_2 \ge 1,$$

where $\tilde{\sigma}$ is a measurable function and h is a possibly infinite-dimensional unknown parameter. Letting $d := d_1 \vee d_2$ and changing the domain of $\tilde{\phi}$ and $\tilde{\sigma}$, we write

$$X_t = S(\alpha^{\top} \phi(\boldsymbol{X}_{t-1}), \beta^{\top} Z_{t-1}) + u_t, \quad \mathbf{E}[u_t^2 | \mathcal{F}_{t-1}] = \sigma^2(\boldsymbol{X}_{t-1}; h),$$

where $\mathbf{X}_{t-1} = (X_{t-1}, \ldots, X_{t-d})$, without loss of generality. Let $\theta_0 = (\alpha_0^{\top}, \beta_0^{\top})^{\top}$ be the true value of $\theta = (\alpha^{\top}, \beta^{\top})^{\top}$, $\Theta = \Theta_{\alpha} \times \Theta_{\beta} \subset \mathbb{R}^{p_1+p_2}$ a parameter space for θ , and H a metric space equipped with a metric d_H . Put $T_{10} := \{j : \alpha_{0j} \neq 0\}, T_{20} := \{p_1 + j : \beta_{0j} \neq 0\}$, and $T_0 := T_{10} \cup T_{20}$. Our aim is to estimate $\theta = (\alpha^{\top}, \beta^{\top})^{\top}$ based on the observation X_{1-d_1}, \ldots, X_n and Z_0, \ldots, Z_n . Put $p = p_1 + p_2$, $s = s_1 + s_2$, where s_1 and s_2 are the cardinalities of T_{10} and T_{20} , respectively.

We first construct the estimator $\hat{\theta}_n^{(1)}$ for θ by the following Dantzig selector type estimator:

$$\hat{\theta}_n^{(1)} := \arg\min_{\theta \in \mathcal{C}_n} \|\theta\|_1, \quad \mathcal{C}_n := \{\theta \in \mathbb{R}^p : \|\psi_n^{(1)}(\theta)\|_{\infty} \le \lambda_n\},\$$

where

$$\psi_n^{(1)}(\theta) = \frac{1}{n} \sum_{t=1}^n \frac{\partial}{\partial \theta} S(\alpha^\top \phi(\mathbf{X}_{t-1}), \beta^\top Z_{t-1}) \{ X_t - S(\alpha^\top \phi(\mathbf{X}_{t-1}), \beta^\top Z_{t-1}) \}$$

and λ_n is a tuning parameter. Moreover, we define the following estimator \hat{T}_n for T_0 :

$$\hat{T}_n := \{ j : |\hat{\theta}_{nj}^{(1)}| > \tau_n \},$$

where τ_n is a threshold. For the second step, we construct a consistent estimator for h, by using $\hat{\theta}_n^{(1)}$. For example, if h is a finite-dimensional parameter, we can construct an estimator as a minimizer of the following function;

$$\mathcal{B}_{n}(h) = \frac{1}{n} \sum_{t=1}^{n} \left| |X_{t} - S(\hat{\alpha}_{n}^{(1)\top} \phi(\boldsymbol{X}_{t-1}), \hat{\beta}_{n}^{(1)\top} Z_{t-1})|^{2} - \sigma^{2}(\boldsymbol{X}_{t-1}; h) \right|^{2}.$$
(3.1)

If h is an infinite-dimensional parameter, we can use, e.g., spline method. Finally, using \hat{T}_n and \hat{h}_n , we consider the estimator $\hat{\theta}_n^{(2)}$ for θ as a solution to the following equation:

$$\Psi_{n\hat{T}_n}(\theta_{\hat{T}_n},\hat{h}_n)=0,\quad \tilde{\theta}_{n\hat{T}_n^c}=0,$$

where

$$\Psi_{nT}(\theta_T, \hat{h}_n) = \frac{1}{n} \sum_{t=1}^n \frac{\frac{\partial}{\partial \theta_T} S(\alpha_{T_1}^\top \phi(\mathbf{X}_{t-1})_{T_1}, \beta_{T_2}^\top Z_{t-1T_2})}{\sigma^2(\mathbf{X}_{t-1}; \hat{h}_n)} \cdot \{X_t - S(\alpha_{T_1}^\top \phi(\mathbf{X}_{t-1})_{T_1}, \beta^\top Z_{t-1T_2})\}$$

for every $T = T_1 \cup T_2$. Define the Hessian matrices $V_n^{(1)}(\theta)$ and $V_n^{(2)}(\theta, h)$ as follows:

$$V_{n}^{(1)}(\theta) := \frac{1}{n} \sum_{t=1}^{n} \frac{\partial^{2}}{\partial \theta \partial \theta^{\top}} S(\alpha^{\top} \phi(\mathbf{X}_{t-1}), \beta^{\top} Z_{t-1}) \left\{ X_{t} - S(\alpha^{\top} \phi(\mathbf{X}_{t-1}), \beta^{\top} Z_{t-1}) \right\}$$
$$- \frac{1}{n} \sum_{t=1}^{n} \frac{\partial}{\partial \theta} S(\alpha^{\top} \phi(\mathbf{X}_{t-1}), \beta^{\top} Z_{t-1}) \frac{\partial}{\partial \theta^{\top}} S(\alpha^{\top} \phi(\mathbf{X}_{t-1}), \beta^{\top} Z_{t-1}),$$

$$V_n^{(2)}(\theta,h) := \frac{1}{n} \sum_{t=1}^n \frac{\partial^2}{\partial \theta \partial \theta^{\top}} \frac{S(\alpha^{\top} \phi(\mathbf{X}_{t-1}), \beta^{\top} Z_{t-1})}{\sigma^2(\mathbf{X}_{t-1};h)} \left\{ X_t - S(\alpha^{\top} \phi(\mathbf{X}_{t-1}), \beta^{\top} Z_{t-1}) \right\} \\ - \frac{1}{n} \sum_{t=1}^n \frac{\partial}{\partial \theta} \frac{S(\alpha^{\top} \phi(\mathbf{X}_{t-1}), \beta^{\top} Z_{t-1})}{\sigma(\mathbf{X}_{t-1};h)} \frac{\partial}{\partial \theta^{\top}} \frac{S(\alpha^{\top} \phi(\mathbf{X}_{t-1}), \beta^{\top} Z_{t-1})}{\sigma(\mathbf{X}_{t-1};h)}.$$

To establish the asymptotic behavior of the estimators $\hat{\theta}_n^{(1)}$ and $\hat{\theta}_n^{(2)}$, we assume the following conditions.

Assumption 3.1. (i) There exists a constant $K_{p,s}$ possibly depending on p and s such that

$$\mathbb{E}\left[\left\|\frac{\partial}{\partial\theta}S(\alpha_0(\phi(\boldsymbol{X}_{t-1}),\beta_0^{\top}Z_{t-1})\right\|_{\infty}^2\sigma^2(\boldsymbol{X}_{t-1};h_0)\right] \leq K_{p,s}.$$

(ii) It holds that

$$c := \inf_{x \in \mathbb{R}} \inf_{h \in H} \sigma^2(x; h) > 0$$

There exists a integrable function Λ_σ such that

$$|\sigma^2(x;h_1) - \sigma^2(x;h_2)| \le \Lambda_\sigma(x) d_H(h_1,h_2)$$

for every $h_1, h_2 \in H$.

(iii) It holds that

$$\log p = o(n), \quad n \to \infty.$$

The tuning parameter λ_n satisfies that

$$\frac{K_{p,s}}{\lambda_n}\sqrt{\frac{\log p}{n}} \to 0, \quad n \to \infty.$$

The sufficient condition for $\|\psi_n^{(1)}(\theta_0)\|_{\infty}$ to obtain the rate of convergence of $\hat{\theta}_n^{(1)}$ can be verified as follows.

Proposition 3.2. Under Assumption 3.1, it holds that

$$\mathbf{P}(\|\psi_n^{(1)}(\theta_0)\|_{\infty} > \lambda_n) \to 0, \quad n \to \infty.$$

See Subsection 5.2 for the proof. Note that we can also use the Bernstein inequality for martingales to deduce a non-asymptotic inequality; see, e.g., Freedman (1975) for details. On the other hand, as the condition for $V_n^{(1)}$ strongly depends on the model structure such as stationarity or tail property of the process, it is difficult to verify it in general settings. Therefore, we assume the following condition.

Assumption 3.3. There exists a constant $\delta > 0$ such that

$$P\left(\inf_{0\neq v\in C_{T_0}}\inf_{\theta\in\Theta}\frac{|v^{\top}V_n^{(1)}(\theta)v|}{\|v_{T_0}\|_1\|v\|_{\infty}} > \delta\right) \to 1, \quad n \to \infty.$$
(3.2)

Note that the left-hand side of (3.2) may depend on n, p and s. We ensure this condition for integer-valued autoregressive models with large order in Subsection 3.2.

Then, we obtain the following rate of convergence.

Theorem 3.4. Let Assumptions 3.1 and 3.3 hold. Then, it holds that

$$P\left(\|\hat{\theta}_n^{(1)} - \theta_0\|_{\infty} > \frac{2}{\delta}c_n\right) \to 0, \quad n \to \infty,$$
(3.3)

where $c_n := \|\psi_n^{(1)}(\hat{\theta}_n^{(1)}) - \psi_n^{(1)}(\theta_0)\|_{\infty}$. Especially, it holds that

$$\|\hat{\theta}_n^{(1)} - \theta_0\|_{\infty} = O_p(\lambda_n), \quad n \to \infty.$$

Moreover, if the threshold τ_n satisfies that $4\lambda_n/\delta < \tau_n < \inf_{j \in T_0} |\theta_{0j}|/2$, it holds that

$$P(\hat{T}_n = T_0) \to 1, \quad n \to \infty.$$
(3.4)

Using \hat{T}_n , we can construct estimators $\tilde{\theta}_n^{(j)}$, j = 1, 2 as solutions to the following estimating equations:

$$\psi_{n\hat{T}_n}^{(1)}(\theta_{\hat{T}_n}) = 0, \quad \tilde{\theta}_{n\hat{T}_n^c} = 0$$

and

$$\Psi_{n\hat{T}_n}(\theta_{\hat{T}_n},\hat{h}_n)=0,\quad \tilde{\theta}_{n\hat{T}_n^c}=0,$$

respectively, where for every $T = T_1 \cup T_2$,

$$\psi_{nT}^{(1)}(\theta_T) = \frac{1}{n} \sum_{t=1}^n \frac{\partial}{\partial \theta_T} S(\alpha_{T_1}^\top \phi(\mathbf{X}_{t-1})_{T_1}, \beta_{T_2}^\top Z_{t-1,T_2}),$$

$$\Psi_{nT}(\theta_T, \hat{h}_n) = \frac{1}{n} \sum_{t=1}^n \frac{\frac{\partial}{\partial \theta_T} S(\alpha_{T_1}^\top \phi(\mathbf{X}_{t-1})_{T_1}, \beta_{T_2}^\top Z_{t-1})}{\sigma^2(\mathbf{X}_{t-1}; \hat{h}_n)} \{ X_t - S(\alpha_{T_1}^\top \phi(\mathbf{X}_{t-1})_{T_1}, \beta^\top Z_{t-1}) \}.$$

To establish the asymptotic normality of $\tilde{\theta}_n^{(j)}$, j = 1, 2, we assume the following conditions which correspond to conditions B1–B6 of Nishiyama (2009).

Assumption 3.5. (i) The sparsity s is independent of n and p. The parameter space $\Theta_{T_0} \subset \mathbb{R}^s$ is compact and the true value θ_{0T_0} is an interior point of Θ_{T_0} . There exists a measurable function Λ_1 on $\mathbb{R} \times \mathbb{R}^{s_2}$ such that

$$S(\alpha_{T_{10}}^{\top}\phi(x)_{T_{10}},\beta_{T_{20}}^{\top}z) - S(\alpha_{0T_{10}}^{\top}\phi(x)_{T_{10}},\beta_{0T_{20}}^{\top}z) = \frac{\partial}{\partial\theta_{T_0}}S(\alpha_{0T_{10}}^{\top}\phi(x)_{T_{10}},\beta_{0T_{20}}^{\top}z)(\theta_{T_0} - \theta_{0T_0}) + \Lambda_1(x,z)\epsilon(x,z;\theta_{T_0},\theta_{0T_0})$$

for every $x \in \mathbb{R}, z \in \mathbb{R}^{s_2}$ and $\theta \in \Theta$ where ϵ is a measurable function satisfying that

$$\sup_{x \in \mathbb{R}, z \in \mathbb{R}^{s_2}} |\epsilon(x, z; \theta_{T_0}, \theta_{0T_0})| = o(||\theta_{T_0} - \theta_{0T_0}||_2), \quad \theta_{T_0} \to \theta_{0T_0}.$$

(ii) There exists a measurable function Λ_2 on $\mathbb{R} \times \mathbb{R}^{s_2}$ such that

$$\sup_{\theta \in \Theta_{T_0}} |S(\alpha_{T_{10}}^\top \phi(x)_{T_{10}}, \beta_{T_{20}}^\top z)| \le \Lambda_2(x, z);$$
$$\sup_{\theta \in \Theta_{T_0}} \left\| \frac{\partial}{\partial \theta_{T_0}} S(\alpha_{T_{10}}^\top \phi(x)_{T_{10}}, \beta_{T_{20}}^\top z) \right\| \le \Lambda_2(x, z);$$
$$\sigma^2(x; h_0) \le \Lambda_2(x, z);$$

$$\left\| \frac{\partial}{\partial \theta_{T_0}} S(\alpha_{1T_{10}}^\top \phi(x)_{T_{10}}, \beta_{1T_{20}}^\top z) - \frac{\partial}{\partial \theta_{T_0}} S(\alpha_{2T_{10}}^\top \phi(x_1)_{T_{10}}, \beta_{2T_{20}}^\top z_1) \right\|_{2}$$

 $\leq \Lambda_2(x, z) \|\theta_{1T_0} - \theta_{2T_0}\|_{2}, \quad \forall \theta_{1T_0}, \theta_{2T_0} \in \Theta_{T_0};$
 $\|\sigma^2(x; h_1) - \sigma^2(x; h_2)\| \leq \Lambda_2(x, z) d_H(h_1, h_2), \quad \forall h_1, h_2 \in H.$

(iii) The process $\{(X_t, Z_{tT_{20}}^{\top})^{\top}\}_{t\geq 0}$ is ergodic under the true value (θ_0, h_0) in the sense that for d with an invariant measure μ_{d+s_2} such that for every μ_{d+s_2} -integrable function g, it holds that

$$\frac{1}{n} \sum_{t=1}^{n} g(X_{t-1}, \dots, X_{t-d}, Z_{tT_{20}}) \to^{p} \int_{\mathbb{R}^{d} \times \mathbb{R}^{s_{2}}} g(x_{1}, \dots, x_{d}, z) \mu_{d+s_{2}}(dx_{1}, \dots, dx_{d}dz),$$

as $n \to \infty$. We also assume that

$$\int_{\mathbb{R}^d \times \mathbb{R}^{s_2}} |\Lambda_i(x,z)|^5 \mu_{d+s_2}(dxdz) < \infty, \quad \int_{\mathbb{R}} |x_0|^4 \mu_1(dx_0) < \infty.$$

(iv) The following $s \times s$ matrices $\mathcal{I}^{(1)}(\theta_0, h_0)$, $\mathcal{J}^{(1)}(\theta_0)$ and $\mathcal{I}^{(2)}(\theta_0, h_0)$ are positive definite:

$$\mathcal{I}^{(1)}(\theta_{0},h_{0}) = \int_{\mathbb{R}^{d}\times\mathbb{R}^{s_{2}}} \frac{\partial}{\partial\theta} S(\alpha_{0T_{1}0}\phi(x)_{T_{10}},\beta_{T_{20}}^{\top}z) \frac{\partial}{\partial\theta^{\top}} S(\alpha_{0T_{1}0}\phi(x)_{T_{10}},\beta_{T_{20}}^{\top}z) \\ \cdot \sigma^{2}(x;h_{0})\mu_{d+s_{2}}(dxdz),$$
$$\mathcal{J}^{(1)}(\theta_{0}) = \int_{\mathbb{R}^{d}\times\mathbb{R}^{s_{2}}} \frac{\partial}{\partial\theta} S(\alpha_{0T_{1}0}\phi(x)_{T_{10}},\beta_{T_{20}}^{\top}z) \frac{\partial}{\partial\theta^{\top}} S(\alpha_{0T_{1}0}\phi(x)_{T_{10}},\beta_{T_{20}}^{\top}z)\mu_{d+s_{2}}(dxdz)$$

and

$$\mathcal{I}^{(2)}(\theta_{0},h_{0}) = \int_{\mathbb{R}^{d}\times\mathbb{R}^{s_{2}}} \frac{\frac{\partial}{\partial\theta_{T_{0}}} S(\alpha_{0T_{10}}^{\top}\phi(x)_{T_{10}},\beta_{0T_{20}}^{\top}z) \frac{\partial}{\partial\theta_{T_{0}}^{\top}} S(\alpha_{0T_{10}}^{\top}\phi(x)_{T_{10}},\beta_{0T_{20}}^{\top}z)}{\sigma^{2}(x;h_{0})} \mu_{d+s_{2}}(dxdz).$$

(v) For every $\epsilon > 0$,

$$\inf_{\substack{\theta_{T_0}: \|\theta_{T_0} - \theta_{0T_0}\|_2 > \epsilon}} \left\| \int_{\mathbb{R}^d \times \mathbb{R}^{s_2}} \frac{\frac{\partial}{\partial \theta_{T_0}} S(\alpha_{T_0}^\top \phi(x)_{T_{10}}, \beta_{T_{20}}^\top z)}{\sigma^2(x; h_0)} \right\|_2 \\ \left[S(\alpha_{T_{10}}^\top \phi(x)_{T_{10}}, \beta_{T_{20}}^\top z) - S(\alpha_{0T_{10}}^\top \phi(x)_{T_{10}}, \beta_{0T_{20}}^\top z) \right] \mu_{d+s_2}(dxdz) \right\|_2 > 0.$$

(vi) The metric entropy condition for (H, d_H) is satisfied:

$$\int_0^1 \sqrt{\log N(H, d_H, \epsilon)} d\epsilon < \infty.$$

(vii) The estimator \hat{h}_n is consistent estimator of h_0 , i.e., $d_H(\hat{h}_n, h_0) = o_p(1)$ as $n \to \infty$.

Some discussions about the metric entropy condition (vi) and examples satisfying such conditions are provided in, e.g., van der Vaart and Wellner (1996) and Nishiyama (2009). Sufficient conditions for the consistency of \hat{h}_n is described in Theorem 5.4 of Nishiyama (2009) when \hat{h}_n is given as a solution to $\mathcal{B}_n(h) = 0$, where $\mathcal{B}_n(h)$ is defined in (3.1). Under above assumptions, we can apply Theorem 2.4 to establish the asymptotic normality of $\tilde{\theta}_n^{(j)}$ as follows.

Theorem 3.6. Under Assumptions 3.1, 3.3, and 3.5,

$$\sqrt{n} \left(\tilde{\theta}_{n\hat{T}_{n}}^{(j)} - \theta_{0T_{0}} \right) \mathbf{1}_{\left\{ \hat{T}_{n} = T_{0} \right\}} \to^{d} N_{s}(0, \Sigma^{(j)}), \quad n \to \infty$$

for j = 1, 2, where

$$\Sigma^{(1)} = \mathcal{J}^{(1)}(\theta_0)^{-1} \mathcal{I}^{(1)}(\theta_0, h_0) \mathcal{J}^{(1)}(\theta_0)^{-1}$$

and

$$\Sigma^{(2)} = \mathcal{I}^{(2)}(\theta_0, h_0)^{-1}.$$

Proof. It follows from Lemmas 5.1, 5.2, and Theorem 5.3 of Nishiyama (2009) that

$$\sqrt{n}\left(\tilde{\theta}_{nT_0}^{(j)} - \theta_{0T_0}\right) \to^d N_s(0, \Sigma^{(j)}), \quad n \to \infty.$$

Combining this fact with (3.4) and Theorem 2.4, we obtain the conclusion.

Remark 3.7. We can show that $\Sigma^{(1)} - \Sigma^{(2)}$ is non-negative definite, which implies $\tilde{\theta}_n^{(2)}$ is asymptotically more efficient than $\tilde{\theta}_n^{(1)}$.

3.2 Example: Integer-valued autoregressive model

As an example, let us consider the following model:

$$X_t = \sum_{i=1}^p \alpha_i \circ X_{t-i} + \epsilon_t, \quad t \in \mathbb{Z},$$
(3.5)

where $\alpha_i \ge 0, i = 1, ..., p$, and, for a non-negative integer-valued random variable X_{t-1} ,

$$\alpha_i \circ X_{t-1} = \begin{cases} \sum_{j=1}^{X_{t-1}} \xi_j^{(i,t)} & X_{t-1} > 0\\ 0 & X_{t-1} = 0, \end{cases}$$

with i.i.d. non-negative integer-valued random sequences $\{\xi_j^{(i,t)}\}_{j\in\mathbb{N}} \ i = 1, \ldots, p, \ t \in \mathbb{Z}$ which is independent over i and t with $\operatorname{E}[\xi_j^{(i,t)}] = \alpha_i$, $\operatorname{Var}[\xi_j^{(i,t)}] = \beta_i$ for every $i = 1, \ldots, p$ and $t \in \mathbb{Z}$, and $\{\epsilon_t\}_{t\in\mathbb{Z}}$ is an i.i.d. non-negative integer-valued random sequence which is independent of $X_s, s < t$ with $\operatorname{E}[\epsilon_t] = \mu_{\epsilon}$ and $\operatorname{Var}[\epsilon_t] = \sigma_{\epsilon}^2$. Suppose that for every $i \neq i'$, $\{\xi_j^{(i,t)}\}_{j\in\mathbb{N}}$ and $\{\xi_j^{(i',t)}\}_{j\in\mathbb{N}}$ are independent. Let $\{\mathcal{F}_t\}_{t\in\mathbb{Z}}$ be a filtration defined by

$$\mathcal{F}_t = \sigma(X_s; s \le t), \quad t \in \mathbb{Z},$$

 $\theta = (\mu_{\epsilon}, \alpha_1, \dots, \alpha_p)^{\top}$ and $h = (\sigma_{\epsilon}^2, \beta_1, \dots, \beta_p)^{\top}$. We define an $\{\mathcal{F}_t\}_{t \in \mathbb{Z}}$ -martingale $\{u_t\}_{t \in \mathbb{Z}}$ as follows:

$$u_t := X_t - \mathbb{E}[X_t | \mathcal{F}_{t-1}], \quad t \in \mathbb{Z}.$$

Noting that

$$\mathbf{E}[X_t|\mathcal{F}_{t-1}] = \mu_{\epsilon} + \sum_{i=1}^p \alpha_i X_{t-i},$$

we have the following representation:

$$X_t = S(\alpha^{\top} \phi(\mathbf{X}_{t-1}), \mu_{\epsilon}) + u_t,$$

where

$$S(\alpha^{\top}\phi(\boldsymbol{X}_{t-1}),\mu_{\epsilon}) = \mu_{\epsilon} + \sum_{i=1}^{p} \alpha_{i}X_{t-i} \quad \alpha = (\alpha_{1},\ldots,\alpha_{p})^{\top},$$

and

$$\sigma^2(\boldsymbol{X}_{t-1};h) = \sigma_{\epsilon}^2 + \sum_{i=1}^p \beta_i X_{t-1}.$$

Let θ_0 and h_0 be the true values of θ and h, respectively. Let us assume that $\theta_{0i} \neq 0$ when $i \leq s$ and $\theta_{0i} = 0$ when i > s for some s > 1. Since $\{\xi_j^{(i,t)}\}$ and $\{\epsilon_t\}$ are non-negative integer-valued random sequences, $\mathbf{E}[\xi_j^{(i,t)}] = \alpha_i = 0$ or $\mathbf{E}[\epsilon_t] = \mu_{\epsilon} = 0$ implies that these random variables are degenerate, i.e., $\operatorname{Var}[\xi_j^{(i,t)}] = \beta_i = 0$ or $\operatorname{Var}[\epsilon_t] = \sigma_{\epsilon}^2 = 0$. Therefore, we can see $T_0 = \{j : \theta_{0j} \neq 0\} \supset \{j : h_{0j} \neq 0\}$, which means the sparsity of h_0 is at most s.

Our aim is to consider the order selection and to construct an asymptotically good estimator of the θ through the Dantzig selector based on the observations $X_t, t = 1 - p, \ldots, n$. The score functions and the Hessian matrices are given as follows:

$$\psi_n^{(1)}(\theta) = \frac{1}{n} \sum_{t=1}^n Y_{t-1} \left\{ X_t - \theta^\top Y_{t-1} \right\},$$
$$V_n^{(1)}(\theta) = -\frac{1}{n} \sum_{t=1}^n Y_{t-1} Y_{t-1}^\top,$$
$$\Psi_n(\theta, h) = \frac{1}{n} \sum_{t=1}^n \frac{Y_{t-1}}{h^\top Y_{t-1}} \left\{ X_t - \theta^\top Y_{t-1} \right\},$$

and

$$V_n(\theta, h) = -\frac{1}{n} \sum_{t=1}^n \frac{Y_{t-1} Y_{t-1}^\top}{h^\top Y_{t-1}},$$

where $Y_{t-1} = (1, X_{t-1}, \dots, X_{t-p})$. Let $V^{(1)} = \mathbb{E}[Y_0 Y_0^{\top}]$. We assume the following conditions for $\{X_t\}_{t \in \mathbb{Z}}$.

Assumption 3.8. (i) It holds that

$$\sum_{i=1}^{p} \alpha_{0i} < 1$$

(ii) For every $t \in \mathbb{Z}$, X_t is \mathcal{F}_{t-1} conditionally sub-Poissonian, that is,

$$\mathbb{E}[\exp(uX_t)|\mathcal{F}_{t-1}] \le \exp(\mu_{t-1}(e^u - 1)), \quad u > 0,$$

where

$$\mu_{t-1} = E[X_t | \mathcal{F}_{t-1}] = \theta_0^{\top} Y_{t-1}$$

The condition (i) implies that there exists a strictly stationary and ergodic solution $\{X_t\}_{t\in\mathbb{Z}}$ to (3.5). The condition (ii) is easily verified when, e.g., $\{\xi_j^{(i,t)}\}$ for all $i = 1, \ldots, p$ and $\{\epsilon_t\}$ are Poisson or Bernoulli sequences. See, e.g., Ahle (2022) for tail bounds of sub-Poissonian random variables, Du and Li (1991), Neumann (2011), Doukhan et al. (2012) and Doukhan et al. (2022), for details of the fundamental properties of integer-valued time series.

Now, we introduce the τ -mixing coefficient to apply the concentration inequality established by Merlevède et al. (2011). For any \mathbb{R}^d -valued random variable X on a probability space (Ω, \mathcal{F}, P) with $||X||_{L^1} < \infty$, and any σ -field $\mathcal{M} \subset \mathcal{F}$, let $P_{X|\mathcal{M}}$ be a conditional distribution of X given \mathcal{M} and P_X be the distribution of X. Then, we consider the following coefficient:

$$\tau(\mathcal{M}, X) = \left\| \sup_{f \in \Lambda_1(\mathbb{R}^d)} \int f(x) \operatorname{P}_{X|\mathcal{M}}(dx) - \int f(x) \operatorname{P}_X(dx) \right\|_{L^1},$$

where

$$\Lambda_1(\mathbb{R}^d) = \left\{ f : \mathbb{R}^d \to \mathbb{R} \left| \sup_{x \neq y} \frac{|f(x) - f(y)|}{\|x - y\|_1} \le 1 \right\}.$$

The τ -mixing coefficients $\tau(i)$ of a sequence $\{X_t\}$ are then given by

$$\tau(i) = \sup_{k \ge 0} \max_{1 \le l \le k} \sup \left\{ \tau(\mathcal{M}_j, (X_{j_1}, \dots, X_{j_l})) | j + i \le j_1 \le \dots \le j_l \right\},\$$

where $\mathcal{M}_j = \sigma(X_t, t \leq j)$. We call $\{X_t\}$ is τ -weakly dependent if $\tau(l) \to 0$ as $l \to \infty$. See, e.g., Doukhan et al. (2012) for the detail of the τ -coefficients. Especially, noting that the true order of the INAR $\{X_t\}$ is s, we have the following lemma.

Lemma 3.9. Suppose that Assumption 3.8 holds.

(i) There exist constants a, c > 0 and $\gamma_1 > 0$ such that

$$\tau(l) \le a \exp(-cl^{\gamma_1}), \quad l \in \mathbb{N}.$$

Moreover, it holds that

$$\gamma := \left(\frac{1}{\gamma_1} + 2\right)^{-1} < 1.$$

(ii) It holds that

$$\beta(l) \le as \exp(-cl^{\gamma_1}), \quad l \in \mathbb{N}.$$

Here, $\beta(l), l \in \mathbb{N}$ is a β -mixing coefficient, i.e.,

$$\beta(l) := \sup \frac{1}{2} \sum_{i=1}^{I} \sum_{j=1}^{J} | P(A_i \cap B_j) - P(A_i) P(B_j) |, \quad l \in \mathbb{N},$$

where the supremum is taken over all pair of partitions $\{A_i\}_{1 \leq i \leq I} \subset \sigma(X_u, u \leq t)$ and $\{B_j\}_{1 \leq j \leq J} \subset \sigma(X_u, u \geq t+l)$ for every $I, J \in \mathbb{N}$. See Theorems 1 and 2 of Doukhan et al. (2012) for the proof. We consider the following conditions.

Assumption 3.10. (i) There exists a positive constant f_{∞} such that

$$F_{\infty}(T_0; V) > f_{\infty}$$

(ii) The matrices

$$V_{T_0,T_0} = \mathbf{E}[Y_{0T_0}Y_{0T_0}^{\top}], \quad \mathcal{I}^{(2)}(\theta_0,h_0) = \mathbf{E}\left[\frac{Y_{0T_0}Y_{0T_0}}{h_{0T_0}^{\top}Y_{0T_0}}\right]$$

are invertible.

(iii) The order p of integer-valued autoregressive model and the sparsity s of the true value θ_0 , and tuning parameter λ_n satisfy that

$$s = O(n^{\zeta}), \quad ps \log p = o(n), \quad \frac{1}{\lambda_n} \sqrt{\frac{ps \log p}{n}} = o(1), \quad n \to \infty,$$

where $\zeta \in [0, 2/11 \land \gamma/(1 + \gamma))$ is a constant.

Let K_X be a constant defined by

$$K_X = \frac{2\mu_{0\epsilon} + 1}{2(1 - \sum_{i=1}^p \alpha_{0i})}.$$

Then, we have the following proposition.

Proposition 3.11. Let Assumptions 3.8 and 3.10 hold.

(i) For every $q \ge 1$, it holds that

$$||X_t||_{L^q} \le qK_X, \quad t \in \mathbb{Z}.$$

Moreover, it holds that

$$\mathbf{P}(\|\psi_n^{(1)}(\theta_0)\|_{\infty} > \lambda_n) \le \frac{12\sqrt{6}\|h_0\|_{\infty}^{1/2} K_X^{3/2}}{\lambda_n} \sqrt{\frac{ps\log(1+p)}{n}}$$

(ii) It holds that

$$\sup_{v \in C_{T_0} \cap \mathbb{S}_p} \sup_{q \ge 1} \frac{\mathrm{E}[|v^{\top} Y_{t-1}|^q]^{1/q}}{q} \le K_s, \quad t \in \mathbb{Z},$$

where

$$K_s = 2\sqrt{s}(1+K_X)$$

and $\mathbb{S}_p \subset \mathbb{R}^p$ is a unit sphere centered 0.

(iii) For every $0 < \eta < f_{\infty}/2$, it holds that

$$\mathbf{P}\left(F_{\infty}(T_0, V_n^{(1)}) > f_{\infty} - 2\eta\right) \to 1, \quad n \to \infty,$$

where

$$F_{\infty}(T_0, V_n^{(1)}) = \inf_{v \in C_{T_0} \setminus \{0\}} \frac{|v^\top V_n^{(1)} v|}{\|v_{T_0}\|_1 \|v\|_{\infty}}.$$

See Subsection 5.2 for the proof.

- **Remark 3.12.** (i) Drost et al. (2008) shows that if $\{\xi_j^{(i,t)}\}_{j\in\mathbb{N}}$, $i = 1, \ldots, p$ and $\{\epsilon_t\}_{t\in\mathbb{Z}}$ satisfy $\mathbb{E}[|\xi_j^{(i,t)}|^q] < \infty$ and $\mathbb{E}[|\epsilon_t|^q] < \infty$, it holds that $\mathbb{E}[|X_t|^q] < \infty$ under Assumption 3.10-(i). On the other hand, we proved the sub-exponential property of INAR under the sub-Poissonian condition in Proposition 3.11-(ii), which may be a new result.
 - (ii) Note that we can consider heavier tailed time series such that Assumption 3.10-(iv) is not satisfied. For such cases, we can use a self-weighted score function such as

$$\psi_{nw}^{(1)}(\theta) = \frac{1}{n} \sum_{t=1}^{n} w_{t-1} Y_{t-1} \left\{ X_t - \theta^\top Y_{t-1} \right\},\,$$

where w_t 's are non-negative \mathcal{F}_t -measurable random variables for every $t \in \mathbb{Z}$ satisfying

$$\sup_{v \in C_{T_0} \cap \mathbb{S}_p} \sup_{q \ge 1} \frac{\mathrm{E}[|v^\top w_{t-1} Y_{t-1}|^q]^{1/q}}{q} \le K_s, \quad t \in \mathbb{Z}.$$

for some $K_s > 0$.

As for the nuisance parameter h, we can construct an estimator \hat{h}_n by a solution to the following estimating equation:

$$\varphi_{n\hat{T}_{n}}(h_{\hat{T}_{n}}) = \frac{1}{n} \sum_{t=1}^{n} Y_{t-1\hat{T}_{n}} \left\{ |X_{t} - \hat{\theta}_{n\hat{T}_{n}}^{(1)\top} Y_{t-1\hat{T}_{n}}|^{2} - h_{\hat{T}_{n}}^{\top} Y_{t-1\hat{T}_{n}} \right\} = 0, \quad h_{\hat{T}_{n}^{c}} = 0,$$

where

$$\hat{T}_n = \{j : |\hat{\theta}_{nj}^{(1)}| > \tau_n\}, \quad \tau_n \in \left(\frac{4\lambda_n}{\delta}, \frac{\inf_{j \in T_0} |\theta_{0j}|}{2}\right)$$

Here, we consider the restriction by \hat{T}_n since $\{j : h_{0j} \neq 0\} \subset T_0$. Note that this equation can be solved as follows

$$\hat{h}_{n\hat{T}_{n}} = \left(\frac{1}{n}\sum_{t=1}^{n}Y_{t-1\hat{T}_{n}}Y_{t-1\hat{T}_{n}}^{\top}\right)^{-1}\frac{1}{n}\sum_{t=1}^{n}(X_{t}-\hat{\theta}_{n\hat{T}_{n}}^{(1)\top}Y_{t-1\hat{T}_{n}})^{2}Y_{t-1\hat{T}_{n}}, \quad \hat{h}_{n\hat{T}_{n}^{c}} = 0,$$

which is consistent when the Dantzig selector type estimator $\hat{\theta}_n^{(1)}$ and \hat{T}_n are consistent. Therefore, the two step estimator $\tilde{\theta}_n$ is defined as a solution to the following equations:

$$\Psi_{n\hat{T}_{n}}\left(\theta_{\hat{T}_{n}},\hat{h}_{n\hat{T}_{n}}\right) = \frac{1}{n}\sum_{t=1}^{n}\frac{Y_{t-1\hat{T}_{n}}}{\hat{h}_{n\hat{T}_{n}}^{\top}Y_{t-1\hat{T}_{n}}}\left\{X_{t}-\theta_{\hat{T}_{n}}^{\top}Y_{t-1\hat{T}_{n}}\right\} = 0, \ \theta_{\hat{T}_{n}^{c}} = 0.$$

In conclusion, under Assumption 3.10, we can apply the estimation procedure described in the previous subsection and obtain an asymptotically normal estimator when s is independent of n and p.

- **Remark 3.13.** (i) When $\{\xi_j^{(i,t)}\}$, i = 1, ..., p and $\{\epsilon_t\}$ are Poisson sequences, we have $h = \theta$. Therefore, we can plug $\hat{\theta}_n^{(1)}$ itself in Ψ_n when $\hat{\theta}_n^{(1)}$ is consistent.
 - (ii) Here, we consider the INAR (p) model with large p as an example. However, as well as Wang (2020), we may also apply the general theory to INAR models with high-dimensional covariates under appropriate conditions.

3.3 Application to the Hawkes processes

The Poisson INAR (p) process is known to be a discrete approximation of the Hawkes process. More precisely, let us consider the Hawkes process N with the intensity

$$\lambda(t) = \eta + \int_{\mathbb{R}} a(t-s) dN_s,$$

where $\eta > 0$ and $h : \mathbb{R} \to \mathbb{R}_+$ is a non-negative and continuous function with a(t) = 0 for $t \leq 0$. We assume that the support of a equals to $(0, \tau]$ for some constant $\tau > 0$ and that

$$\int_{\mathbb{R}} a(t)dt < 1$$

According to Kirchner (2016), there exist constants $\delta > 0$ and $\tilde{K} < 1$ such that for any $\Delta \in (0, \delta)$,

$$K^{(\Delta)} = \Delta \sum_{k=1}^{\infty} h(k\Delta) = \Delta \sum_{k=1}^{p} h(k\Delta) \le \tilde{K} < 1,$$

where $p \ge \lceil \tau/\Delta \rceil$. Let $\{X_t^{(\Delta)}\}$ be an INAR(p) process given by

$$X_t^{(\Delta)} = \sum_{i=1}^p \alpha_i^{(\Delta)} \circ X_{t-i}^{(\Delta)} + \epsilon_t^{(\Delta)},$$

where $\{\epsilon_t^{(\Delta)}\}\$ is an i.i.d. $Poisson(\Delta \eta)$ sequence and $\alpha_i^{(\Delta)} = \Delta a(i\Delta)$. We define the counting process $N^{(\Delta)}$ as follows:

$$N^{(\Delta)}(A) = \sum_{k:k\Delta \in A} X_k^{(\Delta)}, \quad A \in \mathcal{B}_b(\mathbb{R}), \quad \Delta \in (0,\delta),$$

where $\mathcal{B}_b(\mathbb{R})$ is a family of bounded Borel sets in \mathbb{R} . Then, Kirchner (2016) shows that

$$N^{(\Delta)} \to^d N, \quad \Delta \to 0.$$

Consider that we observe a Hawkes process $\{N_t\}_{t\in[0,T]}$. Fix a small constant $\Delta = \lceil T/n \rceil$ for $n \in \mathbb{N}$. We can consider the *n*-partitions of (0,T]:

$$(0,T] = \bigcup_{k=0}^{n-1} (k\Delta, (k+1)\Delta].$$

Then, we can see that if Δ is sufficiently small,

$$N((k\Delta, (k+1)\Delta]) \approx X_k^{(\Delta)},$$

where $\{X_k^{(\Delta)}\}$ is a INAR(*p*) process with sufficiently large *p*. Kirchner (2017) considered conditionally least squares estimator based on this approximation. On the other hand, we can also apply the estimation method discussed in Subsection 3.2. Let τ be a constant such that $\tau \leq p\Delta \leq T$ and $s = \lceil \tau/\Delta \rceil$. Then, $\alpha^{(\Delta)} = (\alpha_1^{(\Delta)}, \ldots, \alpha_p^{(\Delta)})^{\top}$ satisfies that

$$\alpha^{(\Delta)} = (\alpha_1^{(\Delta)}, \dots, \alpha_s^{(\Delta)}, 0, \dots, 0)^\top \in \mathbb{R}^p.$$

Therefore, we can estimate $\alpha^{(\Delta)}$ and the support $\tau = s\Delta$ by using the Dantzig selector and proposed estimator.

3.4 Simulation studies

We consider the sparse estimation of the regression coefficients of INAR(p) model presented in the previous subsection.

$$X_t = \sum_{i=1}^p \alpha_i \circ X_{t-i} + \epsilon_t, \quad t \in \mathbb{Z},$$

where the true value is given by

$$\alpha_0 = (0.25, 0.25, 0.2, 0.2, 0, \dots, 0) \in \mathbb{R}^p, \quad \mu_\epsilon = 0.3.$$

The scenario of the simulation study is given as follows.

Case 1. The order p of the INAR model is 10, and the number n of the observation points is 1000 or 2000.

Case 2. p = 20, n = 1000 or n = 2000.

Table 1: Performance of the estimators, 500 replications, Case 1.

	(n,p) = (1000,10)	(n,p) = (2000,10)
Mean of $\ \hat{\alpha}_n^{(1)} - \alpha_0\ _{\infty}$	0.067	0.049
Mean of $\ \hat{\alpha}_n^{(1)} - \alpha_0\ _2$	0.099	0.071
Proportion of $\hat{T}_n = T_0$	0.940	0.990
Mean of $\ \tilde{\alpha}_n - \alpha_0\ _{\infty}$	0.048	0.033
Mean of $\ \tilde{\alpha}_n - \alpha_0\ _2$	0.063	0.043
<i>p</i> -value of the Royston test for $\tilde{\alpha}_{nT_0}$	< 0.01	0.92

Table 2: Performance of the estimators, 500 replications, Case 2.

	(n,p) = (1000,20)	(n,p) = (2000,20)
Mean of $\ \hat{\alpha}_n^{(1)} - \alpha_0\ _{\infty}$	0.069	0.049
Mean of $\ \hat{\alpha}_n^{(1)} - \alpha_0\ _2$	0.102	0.073
Proportion of $\hat{T}_n = T_0$	0.940	0.992
Mean of $\ \tilde{\alpha}_n - \alpha_0\ _{\infty}$	0.050	0.033
Mean of $\ \tilde{\alpha}_n - \alpha_0\ _2$	0.065	0.044
<i>p</i> -value of the Royston test for $\tilde{\alpha}_{nT_0}$	< 0.01	0.65

Our aim is order selection via the Dantzig selector and to construct a good estimator for α . We choose the tuning parameter λ_n by a cross-validation method while we put the threshold $\tau_n = 2\sqrt{\log p/n}$, which is determined by taking into account of the rate of $\|\psi_n^{(1)}(\theta_0)\|_{\infty}$. To construct the Dantzig selector for α , we apply the gds method in the package hdme in R; see Sørensen (2019) to the centered process $\{X_t\}_{t\in\mathbb{Z}}$.

Tables 1 and 2 illustrate the average of l_{∞} and l_2 errors of the estimators, empirical probability of the success of the selection and the *p*-value of the Royston test in the package MVN in R (see Korkmaz et al. (2014) and Mecklin and Mundfrom (2005)), for the multivariate normality of the two step estimator restricted by the true support index set, over 500 replications. For both cases, the Dantzig selector works well in terms of errors and selection. Moreover, the two step estimator $\tilde{\alpha}_n$ has smaller errors than $\hat{\alpha}_n^{(1)}$ and seems to satisfy the normality when *n* is large.

4 Applications to models of diffusion processes

4.1 Diffusion processes with high-dimensional covariate

Let us consider the following model of the stochastic differential equation:

$$X_t = X_0 + \int_0^t S(\alpha^\top \phi(X_s), \beta^\top Z_s) ds + \int_0^t \sigma(X_s; h) dW_s, \quad t \ge 0,$$

where $\{W_t\}_{t\geq 0}$ is a 1-dimensional standard Brownian motion, $\phi: \mathbb{R} \to \mathbb{R}^{p_1}$ is a measurable function, $\{Z_t\}_{t\geq 0}$ is a p_2 -dimensional process, which is regarded as a covariate vector, $\alpha \in \mathbb{R}^{p_1}$ is an autoregressive coefficient, $\beta \in \mathbb{R}^{p_2}$ is a regression coefficient for covariate process $\{Z_t\}_{t\geq 0}$, S is a measurable and twice continuously defirentiable function with respect to α and β , and σ is a measurable function. We consider the estimation problem for $\theta = (\alpha^{\top}, \beta^{\top})^{\top} \in \mathbb{R}^p$, where $p = p_1 + p_2$, with the existence of a possibly infinitedimensional nuisance parameter $h \in H$. Let (θ_0, h_0) be a true value of (θ, h) , $\Theta =$ $\Theta_{\alpha} \times \Theta_{\beta} \subset \mathbb{R}^{p_1+p_2}$ a parameter space for θ and H a metric space equipped with a metric d_H . We observe the processes $\{X_t\}_{t\geq 0}, \{Y_t\}_{t\geq 0}$ and $\{Z_t\}_{t\geq 0}$ at n+1 equidistant time points $0 =: t_0^n < t_1^n < \cdots < t_n^n$. Put $T_{10} = \{j: \alpha_{0j} \neq 0\}, T_{20} = \{p_1 + j: \beta_{0j} \neq 0\}$ and $T_0 = T_{10} \cup T_{20}$. Assume that the true value θ_0 is high-dimensional and sparse, that is, ppossibly tends to ∞ as $n \to \infty$, and the number $s = s_1 + s_2$ of nonzero coefficients of θ_0 is relatively small, where s_1 is the sparsity for α_0 and s_2 is the sparsity for β_0 .

Consider the following score function for θ :

$$\psi_n^{(1)}(\theta) = \frac{1}{n\Delta_n} \sum_{k=1}^n \frac{\partial}{\partial \theta} S(\alpha^\top \phi(X_{t_{k-1}^n}), \beta^\top Z_{t_{k-1}^n}) \\ \cdot \{X_{t_{k^n}} - X_{t_{k-1}^n} - S(\alpha^\top \phi(X_{t_{k-1}^n}), \beta^\top Z_{t_{k-1}^n})\Delta_n\},$$

where $\Delta_n = t_n^n/n = |t_k^n - t_{k-1}^n|$, k = 1, ..., n. Then, we define the following estimator for θ_0 and T_0 , respectively;

$$\hat{\theta}_n^{(1)} := \arg\min_{\theta \in \mathcal{C}_n} \|\theta\|_1, \quad \mathcal{C}_n := \{\theta \in \Theta : \|\psi_n^{(1)}(\theta)\|_\infty \le \lambda_n\},\$$

and

$$\hat{T}_n = \{j : |\hat{\theta}_{nj}^{(1)}| > \tau_n\},\$$

where τ_n is a threshold.

We estimate θ based on the quasi-likelihood function given by

$$L_{n}(\theta,h) = \prod_{i=1}^{n} \frac{1}{\sqrt{2\pi\sigma^{2}(X_{t_{i-1}^{n}};h)|t_{i}^{n}-t_{i-1}^{n}|}} \\ \cdot \exp\left(-\frac{|X_{t_{i}^{n}}-X_{t_{i-1}^{n}}-S(\alpha^{\top}\phi(X_{t_{i-1}^{n}}),\beta^{\top}Z_{t_{i-1}^{n}})|t_{i}^{n}-t_{i-1}^{n}||^{2}}{2\sigma^{2}(X_{t_{i-1}^{n}};h)|t_{i}^{n}-t_{i-1}^{n}|}\right).$$

Let $\Psi_n(\theta; h)$ be the derivative of $\log L_n(\theta; h)/T_n$ with respect to θ , i.e.,

$$\Psi_n(\theta,h) = \frac{1}{n\Delta_n} \sum_{k=1}^n \frac{\frac{\partial}{\partial \theta} S(\alpha^\top \phi(X_{t_{k-1}^n}), \beta^\top Z_{t_{k-1}^n})}{\sigma^2(X_{t_{k-1}^n}; \hat{h}_n)} \\ \cdot \{X_{t_{k^n}} - X_{t_{k-1}^n} - S(\alpha^\top \phi(X_{t_{k-1}^n}), \beta^\top Z_{t_{k-1}^n}) \Delta_n\}.$$

As for the nuisance parameter h, we can construct a consistent estimator \hat{h}_n . When H is finite-dimensional space, see, e.g., Yoshida (1992), and Kessler (1997) while see, e.g., Hoffmann (2001) for the case where H is infinite-dimensional space. Then, we define the estimator $\tilde{\theta}_n$ for θ as a solution to the following equations:

$$\Psi_{n\hat{T}_n}(\theta_{\hat{T}_n},\hat{h}_n)=0,\quad \theta_{\hat{T}_n^c}=0,$$

where for every index set $T = T_1 \cup T_2$,

$$\Psi_{nT}(\theta_{T}, \hat{h}_{n}) = \frac{1}{n\Delta_{n}} \sum_{k=1}^{n} \frac{\frac{\partial}{\partial\theta_{T}} S(\alpha_{T_{1}}^{\top} \phi_{T_{1}}(X_{t_{k-1}}), \beta_{T_{2}}^{\top} Z_{t_{k-1}}^{n} T_{2})}{\sigma^{2}(X_{t_{k-1}}; \hat{h}_{n})} \cdot \{X_{t_{k}n} - X_{t_{k-1}}^{n} - S(\alpha_{T_{1}}^{\top} \phi_{T_{1}}(X_{t_{k-1}}), \beta_{T_{2}}^{\top} Z_{t_{k-1}}^{n} T_{2})\Delta_{n}\}.$$

Let $V_n^{(1)}(\cdot)$ and $V_n(\cdot, \cdot)$ be derivatives of $\psi_n^{(1)}$ and Ψ_n with respect to θ , respectively, i.e.,

$$V_{n}^{(1)}(\theta) = \frac{1}{n\Delta_{n}} \sum_{k=1}^{n} \frac{\partial^{2}}{\partial\theta\partial\theta^{\top}} S(\alpha^{\top}\phi(X_{t_{k-1}^{n}}), \beta^{\top}Z_{t_{k-1}^{n}})$$

$$\cdot \{X_{t_{k}^{n}} - X_{t_{k-1}^{n}} S(\alpha^{\top}\phi(X_{t_{k-1}^{n}}), \beta^{\top}Z_{t_{k-1}^{n}})\Delta_{n}\}$$

$$- \frac{1}{n\Delta_{n}} \sum_{k=1}^{n} \frac{\partial}{\partial\theta} S(\alpha^{\top}\phi(X_{t_{k-1}^{n}}), \beta^{\top}Z_{t_{k-1}^{n}}) \frac{\partial}{\partial\theta^{\top}} S(\alpha^{\top}\phi(X_{t_{k-1}^{n}}), \beta^{\top}Z_{t_{k-1}^{n}})\Delta_{n}$$

and

$$V_{n}(\theta,h) = \frac{1}{n\Delta_{n}} \sum_{k=1}^{n} \frac{\frac{\partial^{2}}{\partial\theta\partial\theta^{\top}} S(\alpha^{\top}\phi(X_{t_{k-1}^{n}}), \beta^{\top}Z_{t_{k-1}^{n}})}{\sigma^{2}(X_{t_{k-1}^{n}};h)} \\ \cdot \{X_{t_{k}^{n}} - X_{t_{k-1}^{n}} S(\alpha^{\top}\phi(X_{t_{k-1}^{n}}), \beta^{\top}Z_{t_{k-1}^{n}})\Delta_{n}\} \\ - \frac{1}{n\Delta_{n}} \sum_{k=1}^{n} \frac{\frac{\partial}{\partial\theta} S(\alpha^{\top}\phi(X_{t_{k-1}^{n}}), \beta^{\top}Z_{t_{k-1}^{n}})}{\sigma^{2}(X_{t_{k-1}^{n}};h)} \frac{\frac{\partial}{\partial\theta^{\top}} S(\alpha^{\top}\phi(X_{t_{k-1}^{n}}), \beta^{\top}Z_{t_{k-1}^{n}})}{\sigma^{2}(X_{t_{k-1}^{n}};h)}\Delta_{n}.$$

Let $C_{T_0} := \{v \in \mathbb{R}^p : \|v_{T_0^c}\|_1 \le \|v_{T_0}\|_1\}$. Hereafter, we consider the following conditions. **Assumption 4.1.** (i) For every $x_1, x_2 \in \mathbb{R}$ and $z_1, z_2 \in \mathbb{R}^{p_2}$, there exists a constant $L_S > 0$ such that

$$\left| S(\alpha_0^{\top} \phi(x_1), \beta_0^{\top} z_1) - S(\alpha_0^{\top} \phi(x_2), \beta_0^{\top} z_2) \right| \le L_S s(|x_1 - x_2| + ||z_1 - z_2||_{\infty}).$$

(ii) The function S is twice continuously differentiable with respect to α and β . Moreover, there exists a positive constant K_p possibly depending on p such that

$$\sup_{t \ge 0} \mathbf{E} \left[|S(\alpha_0^{\top} \phi(X_t), \beta_0^{\top} Z_t)|^4 \right]^{1/4} \le K_p,$$
$$\sup_{t \ge 0} \mathbf{E} \left[\left\| \frac{\partial}{\partial \alpha} S(\alpha_0^{\top} \phi(X_t), \beta_0^{\top} Z_t) \right\|_{\infty}^4 \right]^{1/4} \le K_p,$$
$$\sup_{t \ge 0} \mathbf{E} \left[\left\| \frac{\partial}{\partial \beta} S(\alpha_0^{\top} \phi(X_t), \beta_0^{\top} Z_t) \right\|_{\infty}^4 \right]^{1/4} \le K_p$$

and

$$\sup_{t \ge 0} \mathbf{E} \left[|\sigma(X_t; h_0)|^4 \right]^{1/4} \le K_p.$$

(iii) It holds that

$$c := \inf_{x \in \mathbb{R}} \inf_{h \in H} \sigma^2(x; h) > 0.$$

(iv) The processes $\{X_t\}_{t\geq 0}$ and $\{Z_t\}_{t\geq 0}$ satisfy that

$$\mathbb{E}\left[\sup_{t\in[t_{i-1}^n,t_i^n]}|X_t-X_{t_{i-1}^n}|^k\right] + \mathbb{E}\left[\sup_{t\in[t_{i-1}^n,t_i^n]}\|Z_t-Z_{t_{i-1}^n}\|_{\infty}^k\right] \le D_k\Delta_n^{k/2},$$

where $D_k > 0$ is a constant depending on k and p.

(v) It holds that

$$\Delta_n \asymp n^{-\zeta}, \quad \log p = o\left((n\Delta_n)^{1/2}\right)$$

as $n \to \infty$, where $\zeta \in (1/2, 1)$. Moreover, the tuning parameter λ_n is a sequence such that

$$\max\left\{\frac{K_p s \Delta_n^{1/2} D_2^{1/2}}{\lambda_n}, \frac{K_p^2}{\lambda_n} \sqrt{\frac{\log(1+p)}{n\Delta_n}}\right\} \to 0, \quad n \to \infty.$$

The condition for $\psi_n^{(1)}(\theta_0)$ can be verified as follows.

Proposition 4.2. Under Assumption 4.1, it holds that

$$P(\|\psi_n^{(1)}(\theta_0)\|_{\infty} > \lambda_n) \to 0, \quad n \to \infty.$$

$$(4.1)$$

See Subsection 5.3 for the proof. On the other hand, the condition for $V_n^{(1)}$ strongly depends on the model structure, it is difficult to provide some sufficient conditions. We thus introduce the following condition.

Assumption 4.3. There exists a constant $\delta > 0$ such that

$$P\left(\inf_{0\neq v\in C_{T_0}}\inf_{\theta\in\Theta}\frac{|v^{\top}V_n^{(1)}(\theta)v|}{\|v_{T_0}\|_1\|v\|_{\infty}} > \delta\right) \to 1, \quad n \to \infty.$$
(4.2)

Note that the left-hand side of (4.2) may depend on n, p and s. We ensure this condition for Ornstein–Uhlenbeck processes in Subsection 4.2. The following theorem follows from Theorem 2.2 and Proposition 4.2.

Theorem 4.4. Let Assumptions 4.1 and 4.3 hold. Then, it holds that

$$P\left(\|\hat{\theta}_n^{(1)} - \theta_0\|_{\infty} > \frac{2}{\delta}c_n\right) \to 0, \quad n \to \infty,$$
(4.3)

where $c_n := \|\psi_n^{(1)}(\hat{\theta}_n^{(1)}) - \psi_n^{(1)}(\theta_0)\|_{\infty}$. Especially, it holds that

$$\|\hat{\theta}_n^{(1)} - \theta_0\|_{\infty} = O_p(\lambda_n), \quad n \to \infty.$$

Moreover, if the threshold τ_n satisfies that $4\lambda_n/\delta < \tau_n < \inf_{j \in T_0} |\theta_{0j}|/2$, it holds that

$$P(\hat{T}_n = T_0) \to 1, \quad n \to \infty.$$
(4.4)

Using \hat{T}_n , we can define the estimator $\hat{\theta}_n^{(2)}$ as a solution to the following equations;

$$\Psi_{n\hat{T}_n}(\theta_{\hat{T}_n},\hat{h}_n)=0,\quad \theta_{\hat{T}_n^c}=0,$$

where for every $T = T_1 \cup T_2$,

$$\Psi_{nT}(\theta_T, h) = \frac{1}{n\Delta_n} \sum_{k=1}^n \frac{\frac{\partial}{\partial \theta_T} S(\alpha_{T_1}^\top \phi(X_{t_{k-1}})_{T_1}, \beta_{T_2}^\top Z_{t_{k-1}}T_2)}{\sigma^2(X_{t_{k-1}}; h)} \\ \cdot \left\{ X_{t_k}^n - X_{t_{k-1}}^n - S(\alpha_{T_1}^\top \phi(X_{t_{k-1}})_{T_1}, \beta_{T_2}^\top Z_{t_{k-1}}T_2) \right\}.$$

We assume the following conditions in order to establish the asymptotic behavior of the estimator $\hat{\theta}_n^{(2)}$.

Assumption 4.5. (i) The sparsity *s* is independent of *n* and *p*. The parameter space $\Theta_{T_0} \subset \mathbb{R}^s$ is compact and the true value θ_{0T_0} is an interior point of Θ_{T_0} . There exists a measurable function Λ_1 on $\mathbb{R} \times \mathbb{R}^{s_2}$ such that

$$S(\alpha_{T_{10}}^{\top}\phi(x)_{T_{10}},\beta_{T_{20}}^{\top}z) - S(\alpha_{0T_{10}}^{\top}\phi(x)_{T_{10}},\beta_{0T_{20}}^{\top}z) \\ = \frac{\partial}{\partial\theta_{T_0}}S(\alpha_{0T_{10}}^{\top}\phi(x)_{T_{10}},\beta_{0T_{20}}^{\top}z)(\theta_{T_0} - \theta_{0T_0}) + \Lambda_1(x,z)\epsilon(x,z;\theta_{T_0},\theta_{0T_0})$$

for every $x \in \mathbb{R}, z \in \mathbb{R}^{s_2}$ and $\theta \in \Theta$ where ϵ is a measurable function satisfying that

$$\sup_{x \in \mathbb{R}, z \in \mathbb{R}^{s_2}} |\epsilon(x, z; \theta_{T_0}, \theta_{0T_0})| = o(\|\theta_{T_0} - \theta_{0T_0}\|_2), \quad \theta_{T_0} \to \theta_{0T_0}.$$

(ii) There exists a constant K_{T_0} such that

$$\sup_{\theta_{T_0} \in \Theta_{T_0}} |S(\alpha_{T_{10}}^{\top} \phi(x_1)_{T_{10}}, \beta_{T_{20}}^{\top} z_1) - S(\alpha_{T_{10}}^{\top} \phi(x_2)_{T_{10}}, \beta_{T_{20}}^{\top} z_2)| \leq K_{T_0} \|y_1 - y_2\|_2;$$

$$\sup_{\theta_{T_0} \in \Theta_{T_0}} \|\frac{\partial}{\partial \theta_{T_0}} S(\alpha_{T_{10}}^{\top} \phi(x_1)_{T_{10}}, \beta_{T_{20}}^{\top} z_1) - \frac{\partial}{\partial \theta_{T_0}} S(\alpha_{T_{10}}^{\top} \phi(x_2)_{T_{10}}, \beta_{T_{20}}^{\top} z_2)\| \leq K_{T_0} \|y_1 - y_2\|_2;$$

$$\sup_{h \in H} |\sigma^2(x_1; h) - \sigma^2(x_2; h)| \leq K_{T_0} |x_1 - x_2|,$$
where $\alpha_{T_0}^{\top} = (x_1, x_1^{\top}), i = 1, 2$

where $y_i^{\top} = (x_i, z_i^{\top}), i = 1, 2.$

(iii) There exists a measurable function Λ_2 on $\mathbb{R} \times \mathbb{R}^{s_2}$ such that

$$\begin{aligned} \|\frac{\partial}{\partial\theta_{T_0}} S(\alpha_{1T_{10}}^{\top}\phi(x)_{T_{10}},\beta_{1T_{20}}^{\top}z) - \frac{\partial}{\partial\theta_{T_0}} S(\alpha_{2T_{10}}^{\top}\phi(x_1)_{T_{10}},\beta_{2T_{20}}^{\top}z_1)\|_2 \\ &\leq \Lambda_2(x,z) \|\theta_{1T_0} - \theta_{2T_0}\|_2, \quad \forall \theta_{1T_0}, \theta_{2T_0} \in \Theta_{T_0}; \\ |\sigma^2(x;h_1) - \sigma^2(x;h_2)| &\leq \Lambda_2(x,z) d_H(h_1,h_2), \quad \forall h_1,h_2 \in H. \end{aligned}$$

(iv) It holds that

$$\sup_{t \ge 0} \mathbf{E} \left[|\Lambda_1(X_t, Z_t)|^8 + |\Lambda_2(X_t, Z_t)|^8 + |X_t|^4 \right] < \infty.$$

(v) The process $\{(X_t, Z_{tT_{20}}^{\top})^{\top}\}_{t \ge 0}$ is ergodic with an invariant measure μ_0 , i.e., for every μ_0 -integrable function g, it holds that

$$\frac{1}{T}\int_0^T g(X_t, Z_{tT_{20}})dt \to^p \int_{\mathbb{R} \times \mathbb{R}^{s_2}} g(x, z)\mu_0(dxdz), \quad T \to \infty.$$

(vi) The following $s \times s$ matrix $\mathcal{I}(\theta_0, h_0)$ is positive definite:

$$\mathcal{I}(\theta_0, h_0) = \int_{\mathbb{R} \times \mathbb{R}^{s_2}} \frac{\frac{\partial}{\partial \theta_{T_0}} S(\alpha_{0T_{10}}^\top \phi(x)_{T_{10}}, \beta_{0T_{20}}^\top z) \frac{\partial}{\partial \theta_{T_0}^\top} S(\alpha_{0T_{10}}^\top \phi(x)_{T_{10}}, \beta_{0T_{20}}^\top z)}{\sigma^2(x; h_0)} \mu_0(dxdz).$$

(vii) It holds that

$$\inf_{\substack{\theta_{T_0}: \|\theta_{T_0} - \theta_{0T_0}\|_{2} > \epsilon}} \left\| \int_{\mathbb{R} \times \mathbb{R}^{s_2}} \frac{\frac{\partial}{\partial \theta_{T_0}} S(\alpha_{T_0}^{\top} \phi(x)_{T_{10}}, \beta_{T_{20}}^{\top} z)}{\sigma^2(x; h_0)} \right\|_{2} \\ \left[S(\alpha_{T_{10}}^{\top} \phi(x)_{T_{10}}, \beta_{T_{20}}^{\top} z) - S(\alpha_{0T_{10}}^{\top} \phi(x)_{T_{10}}, \beta_{0T_{20}}^{\top} z) \right] \mu(dxdz) \right\|_{2} > 0$$

for every $\epsilon > 0$.

(viii) The metric entropy condition for (H, d_H) is satisfied:

$$\int_0^1 \sqrt{\log N(H, d_H, \epsilon)} d\epsilon < \infty.$$

(ix) The estimator \hat{h}_n is a consistent estimator of h_0 , i.e., $d_H(\hat{h}_n, h_0) = o_p(1)$ as $n \to \infty$.

Note that Assumption 4.5 corresponds to conditions A1-A8 of Nishiyama (2009). The sufficient condition for (ix) is described in Theorem 4.4 of Nishiyama (2009) when \hat{h}_n is a minimizer of

$$\mathcal{A}_n(h) = \frac{1}{n} \sum_{k=1}^n \left| \frac{|X_{t_k^n} - X_{t_{k-1}^n}|^2}{\Delta_n} - \sigma^2(X_{t_{k-1}^n};h) \right|^2.$$

Finally, we obtain the following results.

Theorem 4.6. Under Assumptions 4.1, 4.3 and 4.5, it holds that

$$\sqrt{n\Delta_n} \left(\tilde{\theta}_{n\hat{T}_n} - \theta_{0T_0} \right) \mathbf{1}_{\left\{ \hat{T}_n = T_0 \right\}} \to^d N_s(0, \mathcal{I}^{-1}), \quad n \to \infty.$$

Proof. It follows from Lemma 4.1 and Theorem 4.3 of Nishiyama (2009) that

$$\sqrt{n\Delta_n} \left(\tilde{\theta}_{nT_0} - \theta_{0T_0} \right) \to^d N_s(0, \mathcal{I}^{-1}), \quad n \to \infty.$$

Combining this fact with Theorems 2.4 and 4.4, we obtain the conclusion.

4.2 Example: A linear model of Gaussian diffusion processes

Let us consider the p+1-dimensional Ornstein–Uhlenbeck process $\{Y_t\}$ which is a solution to the following equation:

$$Y_t = Y_0 + \int_0^t A Y_s ds + \Sigma \boldsymbol{W}_t, \quad t \ge 0,$$

where $A, \Sigma \in \mathbb{R}^{(p+1) \times (p+1)}$ are unknown matrices and $\{W_t\}_{t\geq 0}$ is a p+1-dimensional standard Brownian motion. Put $Y_t = (X_t, Z_t^{\top})^{\top}$, $W_t = (W_t, W_t^1, \dots, W_t^p)^{\top}$ for every $t \geq 0$. When $\Sigma = \text{diag}(\sigma, \sigma_1, \dots, \sigma_p)$, it holds that

$$X_t = X_0 + \int_0^t \{ \alpha X_s + \beta^\top Z_s \} ds + \sigma W_t, \quad t \ge 0,$$

where $\theta^{\top} := (\alpha, \beta^{\top})$ is the first row of the matrix A. Let $\theta_0^{\top} = (\alpha_0, \beta_0^{\top})$ and σ_0 be true values of θ and σ . We can consider the following score function and Hessian matrix:

$$\psi_n^{(1)}(\theta) = \frac{1}{n\Delta_n} \sum_{k=1}^n Y_{t_{k-1}^n} \left\{ X_{t_k^n} - X_{t_{k-1}^n} - \theta^\top Y_{t_{k-1}^n} \right\}, \quad V_n^{(1)} = -\frac{1}{n\Delta_n} \sum_{k=1}^n Y_{t_{k-1}^n} Y_{t_{k-1}^n}^\top \Delta_n.$$

We assume the following conditions.

Assumption 4.7. (i) The matrix A is diagonalizable and the real parts of its eigenvalues $\varsigma_p \leq \ldots \leq \varsigma_1$ are all negative. Moreover, the initial value $Y_0 \sim N_{p+1}(0, V)$.

(ii) There exist positive constants f_{∞} and K_p ,

$$F_{\infty}(T_0, V) > f_{\infty}, \quad ||V^*V||_2 \le K_p$$

where V^* is the Hermitian adjoint of V.

(iii) The matrix $\mathcal{I}(\theta_0, \sigma_0) = V_{T_0, T_0} / \sigma_0^2$ is invertible.

Condition (i) implies that $\{Y_t\}$ is strictly stationary Gaussian process with covariance function;

$$C(s,t) := \operatorname{Cov}(Y_s, Y_t) = \begin{cases} \exp((s-t)A)V & 0 \le t \le s < \infty \\ V \exp((t-s)A^{\top}) & 0 \le s < t < \infty. \end{cases}$$

See Chapter 5.6 of Karatzas and Shreve (1991) for details of condition (i). Especially, it holds that \sim

$$\sum_{l=0}^{\infty} \exp(l\Delta_n \varsigma_1) < \infty$$

under Assumption 4.7. Then, we have the following result.

Proposition 4.8. Let Assumption 4.7 hold.

(i) It holds that

$$\mathbf{E}[\|Y_0\|_{\infty}^4]^{1/4} < K_p.$$

(ii) There exists a constant $\eta \in (0, f_{\infty}/2)$ such that

$$P\left(F_{\infty}(T_0, V_n^{(1)}) > f_{\infty} - 2\eta\right) \to 1, \quad n \to \infty.$$

See Subsection 5.3 for the proof. The assertion (i) implies

$$\mathbf{P}(\|\psi_n^{(1)}(\theta_0)\|_{\infty} > \lambda_n) \to 0, \quad n \to \infty,$$

when λ_n satisfies that

$$\max\left\{\frac{K_p\Delta_n^{1/2}D_2^{1/2}}{\lambda_n}, \frac{K_p^2}{\lambda_n}\sqrt{\frac{\log(1+p)}{n\Delta_n}}\right\} \to 0, \quad n \to \infty.$$

Note that the process $\{Y_{tT_0}\}$ is ergodic with an invariant measure under Condition (i) of Assumption 4.7. As for the diffusion coefficient σ , we can construct a consistent estimator by classical methods, see, e.g., Yoshida (1992). In conclusion, under Assumption 4.7, we can apply the estimation procedure described in the previous subsection and obtain an asymptotically normal estimator.

5 Proofs

5.1 Stochastic maximal inequalities and concentration inequalities

In this section, we introduce three useful lemmas which are used to derive the error bounds of the Dantzig selector.

The following maximal inequality for square integrable martingales is provided in Nishiyama (2022).

Lemma 5.1 (Theorem A1.1.6 of Nishiyama (2022)). Let p be any positive integer. Let a p-dimensional martingale difference sequence $\xi = (\xi^1, \ldots, \xi^p)^\top$ on a stochastic basis $\boldsymbol{B} = (\Omega, \mathcal{F}, \mathrm{P}, \{\mathcal{F}_k\}_{k\geq 0})$ such that $\mathrm{E}[(\xi_k^i)^2] < \infty$ for all i, k given. Then, it holds for any finite stopping time T that

$$\mathbf{E}\left[\max_{1\leq i\leq p}\left|\sum_{k=1}^{T}\xi_{k}^{i}\right|\right] \leq 2\sqrt{2}\sqrt{E\left[\sum_{k=1}^{T}\max_{1\leq i\leq p}(\xi_{k}^{i})^{2}\right]}\sqrt{\log(1+p)}$$

and that

$$\mathbf{E}\left[\max_{1\leq i\leq p}\left|\sum_{k=1}^{T}\xi_{k}^{i}\right|\right] \leq 2\sqrt{2}\sqrt{\mathbf{E}\left[\sum_{k=1}^{T}\mathbf{E}\left[\max_{1\leq i\leq p}(\xi_{k}^{i})^{2}|\mathcal{F}_{k-1}\right]\right]}\sqrt{\log(1+p)}.$$

The inequality given by the following lemma is a variant of the Hanson–Wright inequality for Gaussian random variables, provided by Wong et al. (2020). See also Rudelson and Vershynin (2013) for details.

Lemma 5.2 (Lemma 11 of Wong et al. (2020)). Let $Y \sim N_n(\mathbf{0}, Q)$ be an *n*-dimensional random vector. Then, there exists a universal constant c > 0 such that for any $\eta > 0$,

$$P\left(\frac{1}{n}\left|\|\boldsymbol{Y}\|_{2}^{2} - E[\|\boldsymbol{Y}\|_{2}^{2}]\right| > \eta \|Q\|_{2}\right) \le 2\exp\left(-cn\min\{\eta,\eta^{2}\}\right).$$

The last lemma gives a concentration inequality for non-Gaussian time series satisfying geometric β -mixing condition, which is essentially equivalent to the Theorem 1 of Merlevède et al. (2011).

Lemma 5.3. Let $\{X_t\}_{t\in\mathbb{Z}}$ be an \mathbb{R} -valued τ -mixing zero mean strictly stationary process which satisfies that

$$\tau(l) \le a \exp(-cl^{\gamma_1}), \quad l \in \mathbb{N}$$

and

$$\sup_{q \ge 1} \operatorname{E}[|X_t|^q]^{1/q} q^{-1/\gamma_2} \le K, \quad \forall t \in \mathbb{Z}$$

for some constants a, c, γ_1, γ_2 , and K > 0. Let

$$\gamma := \left(\frac{1}{\gamma_1} + \frac{1}{\gamma_2}\right)^{-1} < 1.$$

Then, for every n > 4 and $\epsilon > 1/n$, it holds that

$$P\left(\left|\frac{1}{n}\sum_{t=1}^{n}X_{t}\right| > \epsilon\right) \leq n \exp\left(-\frac{(\epsilon n)^{\gamma}}{K^{\gamma}C_{1}}\right) + \exp\left(-\frac{\epsilon^{2}n^{2}}{K^{2}C_{2}\{1+n\mathcal{V}\}}\right) + \exp\left(-\frac{\epsilon^{2}n}{K^{2}C_{3}}\right),$$

$$(5.1)$$

where

$$\mathcal{V} = \sup_{M>0} \sup_{i>0} \left(\operatorname{Var}[\varphi_M(X_i)] + 2\sum_{j>i} |\operatorname{Cov}(\varphi_M(X_i), \varphi_M(X_j))| \right),$$
(5.2)

with

$$\varphi_M(x) = (x \wedge M) \lor (-M), \quad x \in \mathbb{R}, \quad M > 0$$

and $C_k, k = 1, 2, 3$ are constants depending only on a, c, γ_1 , and γ_2 .

See also Wong et al. (2020) for the proof.

5.2 Proofs for Section 3

Proof of Proposition 3.2. Let us first note that

$$\psi_n(\theta_0) = \frac{1}{n} \sum_{t=1}^n \frac{\partial}{\partial \theta} S(\alpha_0(\phi(\boldsymbol{X}_{t-1}), \beta_0^\top Z_{t-1}) u_t.$$

It follows from Lemma 5.1 that

$$\begin{split} \mathbf{E}\left[\|\psi_n(\theta_0)\|_{\infty}\right] &= \mathbf{E}\left[\max_{1\leq i\leq p}\left|\sum_{t=1}^n\xi_t^i\right|\right] \\ &\leq 2\sqrt{2}\sqrt{\mathbf{E}\left[\sum_{t=1}^n\max_{1\leq i\leq p}(\xi_t^i)^2\right]}\sqrt{\log(1+p)}, \end{split}$$

where

$$\xi_t^i = \frac{1}{n} \frac{\partial}{\partial \theta_i} S(\alpha_0(\phi(\boldsymbol{X}_{t-1}), \beta_0^\top Z_{t-1}) u_t.$$

We have

$$\begin{split} \mathbf{E}\left[\sum_{t=1}^{n}\max_{1\leq i\leq p}(\xi_{t}^{i})^{2}\right] &= \frac{1}{n^{2}}\sum_{t=1}^{n}\mathbf{E}\left[\left\|\frac{\partial}{\partial\theta}S(\alpha_{0}(\phi(\boldsymbol{X}_{t-1}),\beta_{0}^{\top}\boldsymbol{Z}_{t-1})\right\|_{\infty}^{2}\mathbf{E}[u_{t}^{2}|\mathcal{F}_{t-1}]\right] \\ &= \frac{1}{n^{2}}\sum_{t=1}^{n}\mathbf{E}\left[\left\|\frac{\partial}{\partial\theta}S(\alpha_{0}(\phi(\boldsymbol{X}_{t-1}),\beta_{0}^{\top}\boldsymbol{Z}_{t-1})\right\|_{\infty}^{2}\sigma^{2}(\boldsymbol{X}_{t-1};h_{0})\right] \\ &\leq \frac{K_{p,s}}{n}. \end{split}$$

Therefore, we obtain

$$\mathbb{E}\left[\|\psi_n(\theta_0)\|_{\infty}\right] \le 2\sqrt{2}K_{p,s}\sqrt{\frac{\log(1+p)}{n}},$$

which ends the proof.

Proof of Proposition 3.11. (i) Under Condition (iv), it follows from Theorem 1 of Ahle (2022) that

$$\mathbb{E}[X_t^q | \mathcal{F}_{t-1}] \le (\theta^\top Y_{t-1} + q/2)^q,$$

hence that

$$\mathbb{E}[X_t^q] \le \mathbb{E}[(\theta^\top Y_{t-1} + q/2)^q].$$

Using the triangle inequality, we have

$$\|X_{t}\|_{L^{q}} \leq \|\theta^{\top}Y_{t-1} + q/2\|_{L^{q}}$$

$$\leq \|\mu_{\epsilon} + \sum_{i=1}^{p} \alpha_{i}X_{t-i}\|_{L^{q}} + q/2$$

$$\leq \mu_{\epsilon} + \sum_{i=1}^{p} \alpha_{i}\|X_{t-i}\|_{L^{q}} + q/2.$$

Since $\{X_t\}_{t\in\mathbb{Z}}$ is a stationary process, we have

$$||X_t||_{L^q} \le qK_X.$$

As well as the proof of Proposition 3.2, we have

$$\mathbf{P}(\|\psi_n^{(1)}(\theta_0)\|_{\infty}) \leq \frac{2\sqrt{2}}{\lambda_n} \sqrt{\sum_{t=1}^n \mathbf{E}\left[\max_{1 \le i \le p} |\xi_t^i|^2\right]} \sqrt{\log(1+p)},$$

where

$$\xi_t^i = \frac{1}{n} Y_{t-1}^i u_t.$$

Since the sparsity of h_0 is at most s, we have

$$E\left[\max_{1 \le i \le p} |\xi_t^i|^2\right] = \frac{1}{n^2} E\left[||Y_{t-1}||_{\infty}^2 E[u_t^2|\mathcal{F}_{t-1}]\right]$$

$$= \frac{1}{n^2} E\left[||Y_{t-1}||_{\infty}^2 \sigma^2(\boldsymbol{X_{t-1}};h_0)\right]$$

$$= \frac{1}{n^2} E\left[||Y_{t-1}||_{\infty}^2(h_0^\top Y_{t-1})\right]$$

$$\le \frac{1}{n^2} E\left[||Y_{t-1}||_{\infty}^2||h_0||_1||Y_{t-1}||_{\infty})\right]$$

$$\le \frac{s||h_0||_{\infty}}{n^2} E\left[||Y_{t-1}||_{\infty}^3\right]$$

Since $\{X_t\}_{t\in\mathbb{Z}}$ is a stationary process, we have

$$E \left[\|Y_{t-1}\|_{\infty}^{3} \right] = E \left[\left\{ \max_{1 \le i \le p} |X_{t-i}| + 1 \right\}^{3} \right]$$

$$\leq 4 E \left[\max_{1 \le i \le p} |X_{t-i}|^{3} \right] + 4$$

$$\leq 4p E[|X_{0}|^{3}] + 4$$

$$\leq 4p \cdot 3^{3} K_{X}^{3} + 4,$$

which implies the conclusion.

(ii) For every $v = (v_0, v_1, \dots, v_p)^\top \in C_{T_0} \cap \mathbb{S}_p$, note that

$$\begin{aligned} \|v\|_1 &\leq 2 \|v_{T_0}\|_1 \\ &\leq 2\sqrt{s} \|v_{T_0}\|_2 \\ &\leq 2\sqrt{s}. \end{aligned}$$

Therefore, we have

$$\|v^{\top}Y_{t-1}\|_{L^{q}} \leq \left\|v_{0} + \sum_{j=1}^{p} v_{j}X_{t-j}\right\|_{L^{q}}$$
$$\leq |v_{0}| + \sum_{j=1}^{p} |v_{j}|qK_{X}$$
$$\leq \|v\|_{1}q(1+K_{X})$$
$$\leq 2\sqrt{s}\|v\|_{2}q(1+K_{X})$$
$$\leq 2\sqrt{s}q(1+K_{X}),$$

which completes the proof for $K_s = 2\sqrt{s}(1 + K_X)$.

(iii) Let $W_n = V_n^{(1)} - V$. Since we have

$$F_{\infty}(T_{0}, V_{n}^{(1)}) = \inf_{v \in C_{T_{0}} \setminus \{0\}} \frac{|v^{\top} V_{n}^{(1)} v|}{\|v_{T_{0}}\|_{1} \|v\|_{\infty}}$$

$$= \inf_{v \in C_{T_{0}} \setminus \{0\}} \frac{|v^{\top} V v|}{\|v_{T_{0}}\|_{1} \|v\|_{\infty}} - \sup_{v \in C_{T_{0}} \setminus \{0\}} \frac{|v^{\top} W_{n} v|}{\|v_{T_{0}}\|_{1} \|v\|_{\infty}},$$

it suffices to show that for every $0 < \eta < f_{\infty}/2$,

$$P\left(\sup_{v\in C_{T_0}\cap\mathbb{S}_p}|v^{\top}W_nv|>\eta\right)\to 0, \quad n\to\infty.$$
(5.3)

For every $v \in C_{T_0} \cap \mathbb{S}_p$, define that

$$Z_t(v) := |v^{\top} Y_{t-1}|^2 - \mathbb{E}[|v^{\top} Y_{t-1}|^2].$$

Then, we can observe that

$$v^{\top}W_n v = \frac{1}{n} \sum_{t=1}^n Z_t(v).$$

Since $v^{\top}Y_{t-1}$ satisfies that

$$\sup_{q \ge 1} \mathbf{E}[|v^{\top}Y_{t-1}|^q]^{1/q} q^{-1} \le K_s,$$

we have

$$\begin{aligned} \|Z_t(v)\|_{L^q} &= \left\| \|v^\top Y_{t-1}\|^2 - \mathbf{E}[|v^\top Y_{t-1}|^2] \right\|_{L^q} \\ &\leq \left\| \|v^\top Y_{t-1}\|^2 \right\|_{L^q} + \mathbf{E}[|v^\top Y_{t-1}|^2] \\ &= \mathbf{E}[|v^\top Y_{t-1}|^{2q}]^{1/q} + 4K_s^2 \\ &= \left(\mathbf{E}[|v^\top Y_{t-1}|^{2q}]^{1/2q} \right)^2 + 4K_s^2 \\ &\leq 4q^2 K_s^2 + 4K_s^2, \quad q \ge 1, \end{aligned}$$

which implies that there exists a constant $\tilde{K}>0$ such that

$$\sup_{q \ge 1} \mathbf{E}[|Z_t(v)|^q]^{1/q} q^{-2} \le \tilde{K}s.$$

By Lemma 3.9, we have

$$\beta(l) \le as \exp(-cl^{\gamma_1}), \quad l \in \mathbb{N}.$$

Then, as well as Wong et al. (2020), we can evaluate the constant \mathcal{V} appeared in (5.1) as follows.

$$\mathcal{V} \leq \|Z_1(v)\|_{L^2}^2 + 4\|Z_1(v)\|_{L^4}^2 \sum_{k=0}^{\infty} \sqrt{\beta(k)}$$

$$\leq \|Z_1(v)\|_{L^2}^2 + 4\|Z_1(v)\|_{L^4}^2 \sqrt{as} \sum_{k=0}^{\infty} \exp\left(-\frac{1}{2}ck^{\gamma_1}\right)$$

$$\leq \tilde{C}_2 s^{5/2}$$

for some constant $\tilde{C}_2 > 0$ depending on a, c, and \tilde{K} . It follows from Lemma 5.3 that

$$P\left(\left|\frac{1}{n}\sum_{t=1}^{n} Z_t(v)\right| > \eta\right) \leq n \exp\left(-\frac{(\eta n)^{\gamma}}{\tilde{K}^{\gamma} s^{\gamma} C_1}\right) + \exp\left(-\frac{\eta^2 n^2}{\tilde{K}^2 s^2 C_2 \{1 + n \tilde{K}^2 s^{5/2}\}}\right) + \exp\left(-\frac{\eta^2 n}{\tilde{K}^2 s^2 C_3}\right)$$

where

$$\gamma = \left(\frac{1}{\gamma_1} + 2\right)^{-1} = \frac{\gamma_1}{1 + 2\gamma_1},$$

 C_1 and C_2 are positive constants depending only on c appeared in the mixing condition (iii) of Assumption 3.8. Therefore, as well as Proof of Proposition 3.1 in Basu and Michailidis (2015) and the ϵ -net argument provided in Vershynin (2018),

we have

$$\begin{split} \mathbf{P}\left(\sup_{v\in C_{T_0}\cap\mathbb{S}_p}|v^{\top}W_nv| > \eta\right) &\leq \mathbf{P}\left(9\sup_{v\in\mathcal{B}_0(2s)}|v^{\top}W_nv| > \eta\right) \\ &\leq \mathbf{P}\left(\sup_{v\in\mathcal{B}_0(2s)}|v^{\top}W_nv| > \frac{\eta}{9}\right) \\ &\leq \left(\begin{array}{c}p\\2s\end{array}\right)6^{2s}\left\{n\exp\left(-\frac{(\eta n)^{\gamma}}{(9\tilde{K})^{\gamma}s^{\gamma}C_1}\right) \\ &+\exp\left(-\frac{\eta^2n^2}{81^2\tilde{K}^2s^2C_2\{1+\tilde{K}ns^{5/2}\}}\right) \\ &+\exp\left(-\frac{\eta^2n}{81^2\tilde{K}^2s^2C_3}\right)\right\} \\ &\leq \exp\left(2s\log 6p + \log n - \frac{(\eta n)^{\gamma}}{(9\tilde{K})^{\gamma}s^{\gamma}C_1}\right) \\ &+\exp\left(2s\log 6p - \frac{\eta^2n^2}{81^2\tilde{K}^2s^2C_2\{1+\tilde{K}ns^{5/2}\}}\right) \\ &+\exp\left(2s\log 6p - \frac{\eta^2n}{81^2\tilde{K}^2s^2C_3}\right) \\ &\to 0, \quad n \to \infty, \end{split}$$

where $\mathcal{B}_0(2s) = \{v \in \mathbb{R}^p : \sum_{j=1}^p \mathbb{1}_{\{v_j \neq 0\}} \leq 2s\}$. This completes the proof.

5.3 Proofs for Section 4

Proof of Proposition 4.2. Since it holds that

$$P(\|\psi_n^{(1)}(\theta_0)\|_{\infty} > \lambda_n) \le \frac{1}{\lambda_n} \{ E[\|A_n\|_{\infty}] + E[\|B_n\|_{\infty}] \},\$$

where

$$A_{n} = \frac{1}{n\Delta_{n}} \sum_{k=1}^{n} \frac{\partial}{\partial \theta} S(\alpha_{0}^{\top} \phi(X_{t_{k-1}}), \beta_{0}^{\top} Z_{t_{k-1}}) \\ \cdot \int_{t_{k-1}}^{t_{k}^{n}} \{ S(\alpha_{0}^{\top} \phi(X_{s}), \beta_{0}^{\top} Z_{s}) - S(\alpha_{0}^{\top} \phi(X_{t_{k-1}}), \beta_{0}^{\top} Z_{t_{k-1}}) \} ds$$

and

$$B_n = \frac{1}{n\Delta_n} \sum_{k=1}^n \frac{\partial}{\partial \theta} S(\alpha_0^\top \phi(X_{t_{k-1}^n}), \beta_0^\top Z_{t_{k-1}^n}) \int_{t_{k-1}^n}^{t_k^n} \sigma(X_s; h_0) dW_s,$$

it suffices to show that

$$\mathbf{E}\left[\|A_n\|_{\infty}\right] = o(\lambda_n), \quad n \to \infty$$

and that

$$\mathbb{E}\left[\left\|B_n\right\|_{\infty}\right] = o(\lambda_n), \quad n \to \infty.$$

Under our assumptions, we have

$$\begin{split} \mathbf{E} \left[\|A_{n}\|_{\infty} \right] &\leq \frac{1}{n\Delta_{n}} \sum_{k=1}^{n} \mathbf{E} \left[\left\| \frac{\partial}{\partial \theta} S(\alpha_{0}^{\top} \phi(X_{t_{k-1}}), \beta_{0}^{\top} Z_{t_{k-1}}) \right\|_{\infty} \\ & \cdot \left| \int_{t_{k-1}^{n}}^{t_{k}^{n}} \{ S(\alpha_{0}^{\top} \phi(X_{s}), \beta_{0}^{\top} Z_{s}) - S(\alpha_{0}^{\top} \phi(X_{t_{k-1}}), \beta_{0}^{\top} Z_{t_{k-1}^{n}}) \} ds \right| \right] \\ &\leq \frac{1}{n\Delta_{n}} \sum_{k=1}^{n} \mathbf{E} \left[\left\| \frac{\partial}{\partial \theta} S(\alpha_{0}^{\top} \phi(X_{t_{k-1}}), \beta_{0}^{\top} Z_{t_{k-1}^{n}}) \right\|_{\infty} \\ & \cdot \int_{t_{k-1}^{n}}^{t_{k}^{n}} \left| S(\alpha_{0}^{\top} \phi(X_{s}), \beta_{0}^{\top} Z_{s}) - S(\alpha_{0}^{\top} \phi(X_{t_{k-1}^{n}}), \beta_{0}^{\top} Z_{t_{k-1}^{n}}) \right| ds \right] \\ &\leq \frac{L_{S}s}{n\Delta_{n}} \sum_{k=1}^{n} \mathbf{E} \left[\left\| \frac{\partial}{\partial \theta} S(\alpha_{0}^{\top} \phi(X_{t_{k-1}^{n}}), \beta_{0}^{\top} Z_{t_{k-1}^{n}}) \right\|_{\infty} \\ & \cdot \int_{t_{k-1}^{n}}^{t_{k}^{n}} \left\{ |X_{s} - X_{t_{k-1}^{n}}| + \|Z_{s} - Z_{t_{k-1}^{n}}\|_{\infty} \right\} ds \right] \\ &\leq \frac{L_{S}s}{n\Delta_{n}} \sum_{k=1}^{n} \mathbf{E} \left[\left\| \frac{\partial}{\partial \theta} S(\alpha_{0}^{\top} \phi(X_{t_{k-1}^{n}}), \beta_{0}^{\top} Z_{t_{k-1}^{n}}) \right\|_{\infty}^{2} \right]^{1/2} \\ & \cdot \mathbf{E} \left[\left\| \int_{t_{k-1}^{n}}^{t_{k}^{n}} \left\{ |X_{s} - X_{t_{k-1}^{n}}| + \|Z_{s} - Z_{t_{k-1}^{n}}\|_{\infty} \right\} ds \right|^{2} \right]^{1/2} \\ &\leq 2^{1/2} L_{S} D_{2}^{1/2} sK_{p} \Delta_{n}^{1/2}, \end{split}$$

which implies

$$\frac{1}{\lambda_n} \operatorname{E}[\|A_n\|_{\infty}] \le \frac{sK_p \Delta_n^{1/2}}{\lambda_n} \cdot 2L_S D_2^{1/2} \to 0, \quad n \to \infty.$$

It follows from Lemma 5.1 that

$$E\left[\|B_n\|_{\infty}\right] = E\left[\max_{1 \le i \le p} \left|\sum_{k=1}^{n} \xi_k^i\right|\right]$$
$$\leq 2\sqrt{2}\sqrt{E\left[\sum_{k=1}^{n} \max_{1 \le i \le p} \left|\xi_k^i\right|^2\right]}\sqrt{\log(1+p)},$$

where

$$\xi_k^i = \frac{1}{n\Delta_n} \frac{\partial}{\partial \theta_i} S(\alpha_0^\top \phi(X_{t_{k-1}^n}), \beta_0^\top Z_{t_{k-1}^n}) \int_{t_{k-1}^n}^{t_k^n} \sigma(X_s; h_0) dW_s, \quad i = 1, \dots, p.$$

We have

$$\begin{split} & \mathbf{E}\left[\sum_{k=1}^{n} \max_{1 \leq i \leq p} \left|\xi_{k}^{i}\right|^{2}\right] \\ & \leq \frac{1}{n^{2}\Delta_{n}^{2}}\sum_{k=1}^{n} \mathbf{E}\left[\left\|\frac{\partial}{\partial\theta}S(\alpha_{0}^{\top}\phi(X_{t_{k-1}}),\beta_{0}^{\top}Z_{t_{k-1}})\right\|_{\infty}^{2} \mathbf{E}\left[\left\{\int_{t_{k-1}}^{t_{k}^{n}}\sigma(X_{s};h_{0})dW_{s}\right\}^{2}\middle|\mathcal{F}_{t_{k-1}}^{n}\right]\right] \\ & = \frac{1}{n^{2}\Delta_{n}^{2}}\sum_{k=1}^{n} \mathbf{E}\left[\left\|\frac{\partial}{\partial\theta}S(\alpha_{0}^{\top}\phi(X_{t_{k-1}}),\beta_{0}^{\top}Z_{t_{k-1}})\right\|_{\infty}^{2} \mathbf{E}\left[\left\{\int_{t_{k-1}^{n}}^{t_{k}^{n}}\sigma^{2}(X_{s};h_{0})ds\right\}\middle|\mathcal{F}_{t_{k-1}}^{n}\right]\right] \\ & \leq K_{p}^{2}\frac{1}{n^{2}\Delta_{n}^{2}}\sum_{k=1}^{n} \mathbf{E}\left[\left\{\int_{t_{k-1}^{n}}^{t_{k}^{n}}\sigma^{2}(X_{s};h_{0})ds\right\}^{2}\right]^{1/2} \\ & \leq \frac{1}{n\Delta_{n}}K_{p}^{4}. \end{split}$$

Therefore, we obtain

$$\mathbb{E}\left[\|B_n\|_{\infty}\right] \le 2\sqrt{2}K_p^2 \sqrt{\frac{\log(1+p)}{n\Delta_n}}.$$

This completes the proof.

Proof of Proposition 4.8. The assertion (i) follows from Assumption 4.1-(ii). To prove the assertion (ii), it suffices to prove that there exists a constant $0 < \eta < f_{\infty}/2$ such that

$$P\left(\sup_{v\in C_{T_0}\cap\mathbb{S}_p}|v^{\top}W_nv|>\eta\right)\to 0, \quad n\to\infty,$$
(5.4)

where $W_n = V_n^{(1)} - V$. Let Y(n) be the matrix defined by

$$Y(n) = (Y_0, \dots, Y_{t_{n-1}^n}).$$

Then, it holds that

$$V_n^{(1)} = \frac{1}{n} Y(n) Y(n)^{\top}, \quad V = \frac{1}{n} \mathbb{E}[Y(n) Y(n)^{\top}].$$

For every $v \in C_{T_0} \cap \mathbb{S}_p$, we have

$$v^{\top}W_n v = \frac{1}{n} \{ \|Y(n)^{\top}v\|_2^2 - \mathbb{E}[\|Y(n)^{\top}v\|_2^2] \}.$$

Noting that $Y(n)^{\top}v$ is a centered Gaussian random variable with covariance matrix

$$Q_n = \begin{pmatrix} v^\top \operatorname{E}[Y_0 Y_0^\top] v & \cdots & v^\top \operatorname{E}[Y_0 Y_{t_{n-1}}^\top] v \\ \vdots & \ddots & \vdots \\ v^\top \operatorname{E}[Y_{t_{n-1}}^n Y_0^\top] v & \cdots & v^\top \operatorname{E}[Y_{t_{n-1}}^n Y_{t_{n-1}}^\top] v \end{pmatrix},$$

we can use the Hanson–Wright inequality (see Rudelson and Vershynin (2013) for details) to deduce that for every $\xi > 0$,

$$P\left(\frac{1}{n}\left|\|Y(n)^{\top}v\|_{2}^{2} - E[\|Y(n)^{\top}v\|_{2}^{2}]\right| > \xi \|Q_{n}\|_{2}\right) \le 2\exp(-cn(\xi \wedge \xi^{2})).$$

From a similar calculation to the proof of Lemma 10 in Supplementary material of Wong et al. (2020), we have

$$||Q_n||_2 \le \max_{0\le i\le n-1} \sum_{l=0}^{n-1} ||\Sigma_Y((l-i)\Delta_n)||_2,$$

where

$$\Sigma_Y((l-i)\Delta_n) = \operatorname{Cov}(Y_{t_l^n}, Y_{t_i^n}).$$

From Assumption 4.7, it holds that

$$\|\Sigma_Y((l-i)\Delta_n)\|_2 = \|\exp(|l-i|\Delta_n A)V\|_2.$$

Since A is diagonalizable, there exists a non-singular matrix P with $||P||_2 = 1$ such that

$$P^{\top}AP = D,$$

where D is a diagonal matrix consists of eigenvalues of A. Therefore, we have

$$\begin{aligned} \|\Sigma_Y((l-i)\Delta_n)\|_2 &= \|\exp(|l-i|\Delta_n A)V\|_2 \\ &\leq \|\exp(|l-i|\Delta_n D)\|_2 \|V\|_2 \\ &\leq \exp(|l-i|\Delta_n\varsigma_1)K_p, \end{aligned}$$

which implies

$$\|Q_n\|_2 \leq \sum_{l=0}^{\infty} \exp(l\Delta_n \varsigma_1) K_p.$$

Then, for

$$\xi = \frac{\eta}{\sum_{l=0}^{\infty} \exp(l\Delta_n \varsigma_1) K_p},$$

we have

$$\mathbf{P}(|v^{\top}W_n v| > \eta) \le 2\exp(-cn(\xi \wedge \xi^2)).$$

Therefore, as well as Proof of Proposition 3.1 in Basu and Michailidis (2015) and the ϵ -net argument provided in Vershynin (2018), we have

$$\begin{aligned} \mathbf{P}\left(\sup_{v\in C_{T_0}\cap\mathbb{S}_p}|v^{\top}W_nv| > \eta\right) &\leq & \mathbf{P}\left(9\sup_{v\in\mathcal{B}_0(2s)}|v^{\top}W_nv| > \eta\right) \\ &\leq & \mathbf{P}\left(\sup_{v\in\mathcal{B}_0(2s)}|v^{\top}W_nv| > \frac{\eta}{9}\right) \\ &\leq & \left(\begin{array}{c}p\\2s\end{array}\right)6^{2s} \cdot 2\exp(-cn(\xi/9 \wedge \xi^2/9^2)) \\ &\leq & 2\exp(2s\log 6p - cn(\xi/9 \wedge \xi^2/9^2)) \\ &\to & 0, \quad n \to \infty, \end{aligned}$$

which ends the proof.

Acknowledgements

This work was supported by JSPS KAKENHI Grant Numbers 21K13271(K.F.), 21K13836 (K.T.).

References

- Ahle, T. D. (2022). Sharp and simple bounds for the raw moments of the binomial and Poisson distributions. *Statist. Probab. Lett.* 182, Paper No. 109306, 5.
- Al-Osh, M. A. and A. A. Alzaid (1987). First-order integer-valued autoregressive (INAR(1)) process. J. Time Ser. Anal. 8(3), 261–275.
- Basu, S. and G. Michailidis (2015). Regularized estimation in sparse high-dimensional time series models. Ann. Statist. 43(4), 1535–1567.
- Bickel, P. J., Y. Ritov, and A. B. Tsybakov (2009). Simultaneous analysis of lasso and Dantzig selector. Ann. Statist. 37(4), 1705–1732.
- Bühlmann, P. and S. van de Geer (2011). *Statistics for high-dimensional data*. Springer Series in Statistics. Springer, Heidelberg. Methods, theory and applications.
- Candes, E. and T. Tao (2007). The Dantzig selector: statistical estimation when p is much larger than n. Ann. Statist. 35(6), 2313–2351.
- Chernozhukov, V., W. Karl Härdle, C. Huang, and W. Wang (2021). LASSO-driven inference in time and space. Ann. Statist. 49(3), 1702–1735.

- Ciołek, G., D. Marushkevych, and M. Podolskij (2020). On Dantzig and Lasso estimators of the drift in a high dimensional Ornstein-Uhlenbeck model. *Electron. J. Stat.* 14(2), 4395–4420.
- Doukhan, P., K. Fokianos, and X. Li (2012). On weak dependence conditions: the case of discrete valued processes. Statist. Probab. Lett. 82(11), 1941–1948.
- Doukhan, P., A. Leucht, and M. H. Neumann (2022). Mixing properties of non-stationary INGARCH(1,1) processes. *Bernoulli* 28(1), 663–688.
- Drost, F. C., R. van den Akker, and B. J. M. Werker (2008). Note on integer-valued bilinear time series models. *Statist. Probab. Lett.* 78(8), 992–996.
- Du, J. G. and Y. Li (1991). The integer-valued autoregressive (INAR(p)) model. J. Time Ser. Anal. 12(2), 129–142.
- Freedman, D. A. (1975). On tail probabilities for martingales. Ann. Probability 3, 100–118.
- Fujimori, K. (2019). The Dantzig selector for a linear model of diffusion processes. Stat. Inference Stoch. Process. 22(3), 475–498.
- Gaïffas, S. and G. Matulewicz (2019). Sparse inference of the drift of a high-dimensional Ornstein-Uhlenbeck process. J. Multivariate Anal. 169, 1–20.
- Hoffmann, M. (2001). On estimating the diffusion coefficient: parametric versus nonparametric. Ann. Inst. H. Poincaré Probab. Statist. 37(3), 339–372.
- Huang, J., T. Sun, Z. Ying, Y. Yu, and C.-H. Zhang (2013). Oracle inequalities for the LASSO in the Cox model. Ann. Statist. 41(3), 1142–1165.
- Karatzas, I. and S. E. Shreve (1991). Brownian motion and stochastic calculus (Second ed.), Volume 113 of Graduate Texts in Mathematics. Springer-Verlag, New York.
- Kessler, M. (1997). Estimation of an ergodic diffusion from discrete observations. Scand. J. Statist. 24(2), 211–229.
- Kirchner, M. (2016). Hawkes and INAR(∞) processes. Stochastic Process. Appl. 126(8), 2494–2525.
- Kirchner, M. (2017). An estimation procedure for the Hawkes process. Quant. Finance 17(4), 571–595.
- Korkmaz, S., D. Goksuluk, and G. Zararsiz (2014). Mvn: An r package for assessing multivariate normality. R J. 6, 151.
- Latour, A. (1997). The multivariate GINAR(p) process. Adv. in Appl. Probab. 29(1), 228–248.

- Latour, A. (1998). Existence and stochastic structure of a non-negative integer-valued autoregressive process. J. Time Ser. Anal. 19(4), 439–455.
- Mecklin, C. J. and D. J. Mundfrom (2005). A Monte Carlo comparison of the type I and type II error rates of tests of multivariate normality. J. Stat. Comput. Simul. 75(2), 93–107.
- Merlevède, F., M. Peligrad, and E. Rio (2011). A Bernstein type inequality and moderate deviations for weakly dependent sequences. *Probab. Theory Related Fields* 151(3-4), 435–474.
- Neumann, M. H. (2011). Absolute regularity and ergodicity of Poisson count processes. Bernoulli 17(4), 1268–1284.
- Nishiyama, Y. (2009). Asymptotic theory of semiparametric Z-estimators for stochastic processes with applications to ergodic diffusions and time series. Ann. Statist. 37(6A), 3555–3579.
- Nishiyama, Y. (2022). Martingale methods in statistics, Volume 170 of Monographs on Statistics and Applied Probability. CRC Press, Boca Raton, FL.
- Periera, J. B. A. and M. Ibrahimi (2014). Support recovery for the drift coefficient of high-dimensional diffusions. *IEEE Trans. Inform. Theory* 60(7), 4026–4049.
- Robinson, P. M. (1987). Asymptotically efficient estimation in the presence of heteroskedasticity of unknown form. *Econometrica* 55(4), 875–891.
- Rudelson, M. and R. Vershynin (2013). Hanson-Wright inequality and sub-Gaussian concentration. *Electron. Commun. Probab.* 18, no. 82, 9.
- Sørensen, Ø. (2019). hdme: High-dimensional regression with measurement error. The Journal of Open Source Software 4 (37).
- Tibshirani, R. (1996). Regression shrinkage and selection via the lasso. J. Roy. Statist. Soc. Ser. B 58(1), 267–288.
- van der Vaart, A. W. and J. A. Wellner (1996). *Weak convergence and empirical processes*. Springer Series in Statistics. Springer-Verlag, New York. With applications to statistics.
- Vershynin, R. (2018). *High-dimensional probability*, Volume 47 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, Cambridge. An introduction with applications in data science, With a foreword by Sara van de Geer.
- Wang, X. (2020). Variable selection for first-order Poisson integer-valued autoregressive model with covariables. Aust. N. Z. J. Stat. 62(2), 278–295.

- Wong, K. C., Z. Li, and A. Tewari (2020). Lasso guarantees for β -mixing heavy-tailed time series. Ann. Statist. 48(2), 1124–1142.
- Yoshida, N. (1992). Estimation for diffusion processes from discrete observation. J. Multivariate Anal. 41(2), 220–242.