INTERNATIONAL JOURNAL OF GEOMETRY Vol. 3 (2013), No. 1, 5 - 14



FROM TWO SIMPLE PROBLEMS TO THE CONNECTION OF SPECIAL POINTS

NGUYEN TRAN HUU THINH

Abstract. Both the USA TST 2008 [1] and the ELMO Shortlist 2013 [2] suggested two issues that are connected to fixed points. These problems provide a strong linkage between the various attributes of specific points in a triangle. In this article, we will first investigate various theorems concerning the fixed points that have been presented, and then we will demonstrate how those points are connected to a few triangle centers.

1. INTRODUCTION

The fixed-point problems have been seen in a great number of international mathematical olympiad competitions [4]. The generalization of these problems in this area, as well as the features of these problems, leave behind a plethora of notable results [10]. In this work, we talk about some of those problems and show the relationship to a few of different kinds of unique triangle centers.

In USA Team Selection Test 2008 [1], there is a fixed-point problem as following:

Problem 1. Given triangle ABC, centroid G. Let P be a variable point on segment BC. Let Q, R be points on AC, AB such that $PQ \parallel AB, PR \parallel AC$. Prove that (AQR) passes through a fixed point X such that (AB, AG) = (AX, AC).

Proof. Let ω_1 be the circle which passes A, B and tangents to AC. Let ω_2 be the circle which passes A, C and tangents to AB. Let X be intersection of ω_1 and ω_2 , so X is fixed. Since two triangle ABX and AXC are homothetic, hence $\frac{AX}{BX} = \frac{AC}{AB}$. On the other hand, $\frac{AC}{AB} = \frac{PR}{BR} = \frac{AQ}{BR}$ so $\frac{AX}{BX} = \frac{AQ}{BR}$, also (BR, BX) = (AX, AX) hence two triangle BRX and AQX are homothetic, therefore (RB, RX) = (QA, QX), following that A, Q, X, R are cyclic. Moreover, we have

$$\frac{\sin(AB, AX)}{\sin(CA, CX)} = \frac{\sin(AB, AX)}{\sin(BA, BX)} = \frac{BX}{AX} = \frac{AB}{AC}$$

Keywords and phrases: Fixed point, Triangle, Brocard point, Hagge circle (2020)Mathematics Subject Classification: 51M04, 51-08

Received: 1.01.2013. In revised form: 13.06.2013. Accepted: 10.09.2013.



FIGURE 1. Proof of Problem 1

so AX is the symmetrian line of triangle ABC, which means (AB, AG) = (AX, AC). The problem has been solved.

With this configuration, ELMO Shortlist 2013 [2] provide the following problem using the antiparallel lines instead of parallel ones:

Problem 2. Given triangle ABC and a centroid G. Let P be a variable point on segment BC. Let Q, R respectively be points on AC, AB such that PQ, PR are antiparallel lines of AB, AC. Prove that (AQR) passes through a fixed point.



FIGURE 2. Proof of Problem 2

Proof. Let γ_1 be the circles passes A, B and tangents to BC. Let γ_2 be the circle passes A, C and tangents to BC. Let Y be intersection of γ_1 and γ_2 ,

so Y is fixed. Let K be intersection of BQ and CR. We hav

$$(RK, RA) = (PC, PA) = (PB, PA) = (QK, QA) \pmod{\pi}.$$

Hence four points A, R, K, Q are cyclic. Therefore:

$$(KB, KC) = (KR, KQ) = (AR, AQ) = (AR, AY) + (AY, AQ)$$
$$= (BY, BC) + (CB, CY)$$
$$= (YB, YC) \pmod{\pi}.$$

So four points B, K, Y, C are cyclic. We have

$$(KR, KY) = (KY, KC) = (BY, BC) = (AR, AY) \pmod{\pi}.$$

Hence Y is on $(ARK) \equiv (AQR)$. Moreover, by using the power of point's properties, we have the radical axis of γ_1 and γ_2 passes their common tangents, following that AY passes BC at its midpoint or AY is the median line of triangle ABC.

The subsequent sections will go into the various qualities associated with the aforementioned pair of fixed points, which are closely linked to several notable triangle centers.

2. PROPERTIES RELATED TO THE FIRST PROBLEM

Two above problems appeared in many contests [1, 2] and journals [4] in the world, but it seems like there are just few properties about. This article will put forward some results about two fixed points mentioned above. We denotes A_X, A_Y be that two points in the previous problems.

Theorem 2.1. A_X is projection point of the circumcenter of triangle ABC on the A-symmetry and line of that triangle.



FIGURE 3. Proof of Theorem 2.1

Proof. Let A_S be the second intersection of AA_X and (ABC). From Problem 1, we have two triangle ABA_X and AA_XC are homothetic hence $A_XA^2 = A_XB \cdot A_XC$. Moreover, we have

$$(A_XB, A_XA_S) = (A_XB, AA_X) = (AA_X, A_XC) = (A_XA_S, A_XC),$$
$$(A_SB, A_SA_X) = (CB, CA) = (CB, CA_X) + (CA_X, CA)$$
$$= (CB, CA_X) + (AA_S, AB)$$
$$= (CB, CA_X) + (CA_S, CB)$$
$$= (CA_S, CA_X).$$

So two triangle $A_X B A_S$ and $A_X A_S C$ are homothetic. Therefore $A_X A_S^2 = A_X B \cdot A_X C$. Hence $A_X A = A_X A_S$ or A_X is midpoint of $A A_S$, following that A_X is projection point of the circumcenter of triangle ABC on the A-symmetry A-symmetry of that triangle.

Theorem 2.2. Brocard circle [14] is the circle which passes two Brocard points Z_1, Z_2 [12, 13], Lemoine point L [16] and circumcenter O of triangle ABC. Let A_X be the fixed point mentioned in Problem 1. The construction of B_X and C_X is performed in a similar manner, but with the utilization of vertices B and C. Then A_X , B_X and C_X lie on Brocard circle [14] of triangle ABC.

Remark 2.1. $A_X B_X C_X$ is the second Brocard triangle of triangle ABC.

Lemma 2.1. (Mannheim's theorem) Given triangle ABC and three points L, M, N respectively be on BC, CA, AB. Let A', B', C' respectively be three points that lie on (AMN), (BNL), (CLM) such that AA', BB', CC' are concurrent at K. Prove that four points A', B', C', K are cyclic.



FIGURE 4. Proof of Lemma 2.1

Proof. Let *P* be Miquel point of triangle *LMN* with respect to triangle *ABC*. Let *Q* be intersection of *PB'* and *(AMN)*. Applying Reim's theorem to *(AMN)* and *(BNL)* we have $AQ \parallel BB'$. Through angle chasing:

$$(KA, KB') = (AA', AQ) = (PA', PB') \pmod{\pi}$$

Therefore A', P, B', K are cyclic. Similarly, B', P, C', K are cyclic. Hence four points A', B', C', K are cyclic.

Lemma 2.2. Given triangle ABC inscribed in (O) and a point P is not this triangle's vertex. Lines AP, BP, CP respectively cut (ABC) at S, T, U. Let X, Y, Z be midpoints of segments AS, BT, CU, respectively. Prove that X, Y, Z lie on (OP) (a circle with center O and radius OP).

Proof. We have X, Y, Z respectively be projection of O on AS, BT, CU. Through angle chasing:

$$(XP, XO) = (YP, YO) = (ZP, ZO) = \frac{\pi}{2}.$$

Therefore X, Y, Z lie on (OP). The lemma has been proved. Back to Theorem 2.2,



FIGURE 5. Proof of Theorem 2.2

Proof. Applying Lemma 2.1, to triangle ABC with AA_X, BB_X, CC_X are concurrent at L and Z_1 is intersection of $(ABB_X), (BCC_X), (CAA_X)$ we have L, Z_1 lie on $(A_X B_X C_X)$. Similarly, Z_2 lies on $(A_X B_X C_X)$. From Lemma 2.2, we have A_X, B_X, C_X lie on OL. The theorem has been proved.

Theorem 2.3. Ray AA_Y cuts (ABC) at A_M . Prove that BA_YCA_M is parallelogram.



FIGURE 6. Proof of Theorem 2.3

Proof. Using the properties of A_Y at Problem 2, we have

 $(BA_Y, BC) = (AB, AA_M) = (CB, CA_M).$

Therefore $A_YB \parallel A_MC$. Similarly, $A_YC \parallel A_MC$ so BA_YCA_M is a parallelogram. The problem has been solved.

Corollary 2.1. A_Y is symmetric with A_S through BC.

Theorem 2.4. A_X is the isogonal conjugate of A_Y with respect to triangle ABC.



FIGURE 7. Proof of Theorem 2.4

Proof. Because AA_X and AA_Y respectively be symmedian line and median line at vertex A are two isogonal conjugate lines so we have to prove that BA_X and BA_Y are two isogonal conjugate ones. Through angle chasing,

$$(BA_X, BA) = (AA_X, AC) = (AB, AA_Y) = (BC, BA_Y),$$

we have BA_X and BA_Y are isogonal conjugate, therefore A_X is the isogonal conjugate of A_Y with respect to triangle ABC.

Theorem 2.5. Given triangle ABC, centroid G. Let M_{AB} , M_{BC} , M_{CA} respectively be midpoints of BC, CA, AB. Let A_G be midpoint of AA_Y . Prove that A_G lies on $(M_{AB}M_{BC}M_{CA})$.



FIGURE 8. Proof of Theorem 2.5

Proof. Because $M_{AB}A_G$ is the midline of triangle ABA_Y so we have: $(M_{AB}A, M_{AB}A_G) = (BA, BA_Y)$ Draw altitude CH_C of triangle ABC. Through angle chasing, $(M_{BC}H_C, M_{BC}A) = (M_{BC}H_C, BA) + (BA, CB)$ $+ (CB, CA) + (CA, M_{BC}A)$ $= 2(BA, CB) + (CB, CA) + (CA, M_{BC}A)$ $= (BA, CB) + (AB, AM_{BC})$ $= (BA, BA_Y) + (BA_Y, CB) + (CB, CA_M)$ $= (BA, BA_Y)$ $= (M_{AB}A, M_{AB}A_G).$

Hence A_G lies on $(M_{AB}H_CM_{BC}) \equiv (M_{AB}M_{BC}M_{CA})$. We have done.

Theorem 2.6. Draw parallelogram $M_{AB}A_GM_{CA}A_{GY}$. Then A_{GY} is midpoint of AA_M .



FIGURE 9. Proof of Theorem 2.6

Proof. Let A_N be midpoint of $M_{AB}M_{CA}$. Because $M_{AB}A_GM_{CA}A_{GY}$ and $AM_{AB}M_{BC}M_{CA}$ are parallelogram, we can easily see that A_N is midpoint of both A_GA_{GY} and AM_{BC} . We have

$$\overline{AA_{GY}} = \overline{AA_N} + \overline{A_NA_{GY}} = \overline{A_NM_{BC}} + \overline{A_GA_N}$$
$$= \overline{A_GM_{BC}} = \overline{A_GA_Y} + \overline{A_YM_{BC}}$$
$$= \frac{1}{2}\left(\overline{AA_Y} + \overline{A_YA_M}\right) = \frac{1}{2}\overline{AA_M}.$$

Therefore A_{GY} is midpoint of AA_M .

Corollary 2.2. $A_X A_{GY} \parallel BC$.

Corollary 2.3. A_X is symmetric with A_G through $M_{AB}M_{CA}$.

Theorem 2.7. A_Y is the Anticomplement point of A_{GY} with respect to triangle ABC.

Proof. From these above properties, we have:

$$\overline{GA_{GY}} = \overline{GM_{BC}} - \overline{A_{GY}M_{BC}} = 2\overline{A_NG} - \overline{A_GA_Y} \\
= \overline{2A_NG} - \overline{A_GA_N} - \overline{A_NA_Y} = \overline{A_NA_Y} + 2\overline{A_YG} - \overline{A_NA_{GY}} \\
= \overline{A_{GY}A_Y} + 2\overline{A_YG} = \overline{A_{GY}G} + \overline{A_YG}.$$

Hence $\overline{A_YG} = 2\overline{GA_{GY}}$ therefore A_Y is the Anticomplement point of A_{GY} with respect to triangle ABC.

Theorem 2.8. The first Brocard triangle is a triangle which has three vertices respectively be intersection points of lines AZ_1 , BZ_1 , CZ_1 , AZ_2 , BZ_2 , CZ_2 . Let Z_A, Z_B, Z_C be three vertices of this triangle. Prove that $Z_A Z_B Z_C$ is inscribed in $(A_X B_X C_X)$. From two simple problems to the connection of special points



FIGURE 10. Proof of Theorem 2.8

$$\begin{aligned} & \textbf{Proof. Let } (AZ_1, AB) = (AC, AZ_2) = \alpha. \text{ Through angle chasing,} \\ & (A_XZ_1, A_XZ_2) = (A_XZ_1, A_XC) + (A_XC, A_XA) + (A_XA, A_XZ_2) \\ & = (AZ_1, AC) + (A_XA, A_XB) + (A_XA, Z_2A) + (Z_2A, A_XZ_2) \\ & = (AZ_1, AC) + (A_XA, A_XB) + (A_XA, Z_2A) + (BA, BA_X) \\ & = (AZ_1, AC) + (A_XA, A_XB) + (A_XA, Z_2A) + (AC, AA_X) \\ & = (AZ_1, AB) + (AB, AA_X) + (A_XA, A_XB) + (A_XA, Z_2A) \\ & = (AZ_1, AB) + (BA, A_XB) + (A_XA, Z_2A) \\ & = (AZ_1, AB) + (AC, AA_X) + (AA_X, AZ_2) \\ & = (AZ_1, AB) + (AC, AA_Z) + (AA_X, AZ_2) \\ & = (AZ_1, AB) + (AC, AZ_2) \\ & = (AZ_1, AB) + (AC, AZ_2) \\ & = 2\alpha \pmod{\pi}. \end{aligned}$$

On the other hand, we have

$$(Z_A Z_1, Z_A Z_2) = (Z_A Z_1, CB) + (CB, Z_A Z_2) = \alpha + \alpha = 2\alpha.$$

Therefore Z_A lies on $(Z_1A_XZ_2) \equiv (A_XB_XC_X)$. Similarly, Z_B, Z_C lie on $(A_XB_XC_X)$. WE have done.

Corollary 2.4. Z_1 is symmetric with Z_2 through OL and $(OZ_1, OZ_2) = 2\alpha$.

Theorem 2.9. OA_X , CZ_2 cut ω_1 at O_W , C_W . Then $O_W C_W \parallel OZ_A$.

Lemma 2.3. Let BZ_1 cut ω_2 at B_W . Prove that $B_W C_W \parallel BC$.

Proof. We have $(BB_W, BC) = (CA, CZ_1) = (B_WA, B_WZ_1)$ so $AB_W \parallel BC$. Similarly, $AC_W \parallel BC$ so $B_WC_W \parallel BC$. We have done. Back to Theorem 2.9,

Proof. By Lemma 2.3 and $(BC, BZ_A) = (CZ_A, CB)$, we have BCB_WC_W is an isosceles trapezoid or OZ_A is perpendicular to B_WC_W . On the other

Nguyen Tran Huu Thinh



FIGURE 11. Proof of Theorem 2.9

hand, by Theorem 2.2, we have $(A_X A, A_X O_W) = (C_W A, C_W O_W) = \frac{\pi}{2}$. Therefore $O_W C_W$ is perpendicular to $B_W C_W$. So $O_W C_W \parallel OZ_A$.

Theorem 2.10. $A_X Z_A$, $B_X Z_B$, $C_X Z_C$ are concurrent at G.

Lemma 2.4. Given triangle ABC. On the plane containing this triangle, choose A', B', C' such that the triangles A'CB, B'AC, C'BA are homothetic. We construct the parallelogram BA'CD. Prove that A'CDB is also a parallelogram.



FIGURE 12. Proof of Lemma 2.4

Proof. We have

$$\frac{C'B}{C'D} = \frac{C'B}{AB'} = \frac{C'A}{B'C} = \frac{B'D}{B'C}$$

and

$$\begin{split} (C'B,C'D) &= (C'B,C'A) + (C'A,C'D) \\ &= (B'A,B'C) + (B'D,B'A) = (B'D,B'C) \end{split}$$

Therefore triangle C'BD and triangle B'DC are homothetic. Hence

$$\frac{BD}{CD} = \frac{C'B}{B'D} = \frac{C'B}{C'A} = \frac{A'B}{A'C}.$$

On the other hand,

$$(DB, DC) = (DB, DC') + (DC', DB') + (DB', DC)$$

= (DC, B'C) + (B'A, DB') + (DB', DC)
= (B'A, B'C)
= (A'C, AB).

So triangle DBC and triangle A'CB are homothetic. Therefore, $BD \parallel A'C$ and $CD \parallel A'B$. We conclude that A'CDB is a parallelogram.

Lemma 2.5. G is the centroid of triangle $Z_A Z_B Z_C$.



FIGURE 13. Proof of Lemma 2.5

Proof. Let Z'_A be symmetric point with Z_A through BC. Since triangle BZ_AC is a isosceles triangle at Z_A , $BZ_ACZ'_A$ is a parallelogram. Applying Lemma 2.4 for triangle ABC, we have $AZ_CZ'_AZ_B$ is a parallelogram so AZ'_A passes midpoint N_{BC} of Z_BZ_C . In triangle $AZ_AZ'_A$, the two median lines AM_{BC} and Z_AN_{BC} intersect with each other at this triangle's centroid so that this point is also the centroid G of triangle ABC and triangle $Z_AZ_BZ_C$.

Lemma 2.6. Given triangle ABC and two disjoint arbitrary points P, Q. Let X, Y, Z respectively be projection of P on the lines which passes Q and is perpendicular to BC, CA, AB. Let X', Y', Z' respectively be projection of Q on AP, BP, CP. Prove that XX', YY' and ZZ' are concurrent.



FIGURE 14. Proof of Lemma 2.6

Proof. Noted that X, Y, Z, X', Y', Z' lie on a circle with diameter PQ and $AB \parallel PZ$, $AC \parallel PY$ so we have

$$(AB, AC) = (ZP, PY) = (XY, XZ).$$

Similarly, (BC, BA) = (YZ, YX) and (CA, CB) = (ZX, ZY) so two triangles XYZ and ABC are homothetic. Let P' be the image of P through the homothety transformation which turns triangle ABC into triangle XYZ. Let P^* be the isogonal conjugate point of P in triangle XYZ. We have

$$(XX', XZ) = (PX', PZ) = (AP, AB) = (XP^*, XZ).$$

Therefore XX' passes P^* . Similarly, YY' and ZZ' passes P^* , which solves the lemma.



FIGURE 15. Proof of Theorem 2.10

Back to Theorem 2.10,

Proof. By applying Lemmas 2.5 and 2.6 for triangle $Z_A Z_B Z_C$, we solves the problem.

By Corollary 2.3, the Brocard circle is definitely the Hagge circle [5] of G in triangle $M_{BC}M_{CA}M_{AB}$. We will discuss some properties related to this issue and use these ones to prove Theorem 2.10.

Lemma 2.7 (Hagge circle). Given triangle ABC, orthocenter H and a point P lies inside (ABC). Rays AP, BP, CP cuts (ABC) at the second points A_1 , B_1 , C_1 . Let A_2 , B_2 , C_2 respectively be the symmetry points of A_1 , B_1 , C_1 through A, B, C. Prove that A_2 , B_2 , C_2 and H are cyclic.

Remark 2.2. The circle mentioned above called Hagge circle \mathcal{H}_P of point P with respect to triangle ABC.



FIGURE 16. Proof of Lemma 2.7

Proof. Let P' be the isogonal conjugate of P in triangle ABC. AP', BP', CP' cut (ABC) respectively at A'_1 , B'_1 , C'_1 . Let B'_2 , C'_2 respectively be symmetry points of B_2 , C_2 through midpoint of BC. Set O is the center of (ABC) with diameter AA'. It is easy to see that A_2 , B_2 , C_2 are symmetry points of A'_1 , B'_2 , C'_2 through midpoint of BC. We have $CB_2AB'_1$ and $CB_2BB'_2$ are parallelograms so the quadrilateral $ABB'_2B'_1$ is also a parallelogram. We conclude that midpoint M_1 of AB'_2 is the projection of O on BP'. Similarly, midpoint M_2 of AC'_2 is the projection of O on CP'. Let M_3 be midpoint of AA'_1 . Then triangle $M_1M_2M_3$ is inscribed in a circle with diameter OP'. We use two geometric transformations respectively with the first one (1) is the homothety transformation \mathbf{H}^2_A with center A and ratio 2; and the later (2) is the symmetric one \mathbf{S}_{MBC} through midpoint M_{BC} of BC, to points M_3 , M_1 , M_2 , O such that

$$\mathbf{S}_{M_{BC}} \circ \mathbf{H}_A^2 \colon M_3, M_1, M_2, O \mapsto A_2, B_2, C_2, H.$$

Since M_1, M_2, M_3, O lie on a same circle so A_2, B_2, C_2, H are also cyclic.

Lemma 2.8. Given triangle ABC and two points S and S' are isogonal conjugate. Let X, X' respectively be intersection of AS, AS' with (ABC). Let V be intersection of AS' and BC. Prove that



FIGURE 17. Proof of Lemma 2.8

Proof. We have

So two triangles V

$$(CV, CX') = (AB, AX') = (AX, AC),$$

 $(X'V, X'C) = (XA, XC)$
 $(CX' \text{ and } CAX \text{ are homothetic. On the other hand,}$

$$(S'X', S'C) = (AS', AC) + (CA, CS')$$

= $(AB, AS) + (CS, CB)$
= $(CB, CX) + (CS, CB)$
= $(CS, CX),$

$$(X'S', X'C) = (XS, XC).$$

Therefore two triangles CX'S' and SXC are homothetic. So,

$$\overline{X'V} \cdot \overline{XA} = \overline{X'C} \cdot \overline{XC} = \overline{X'S'} \cdot \overline{XS}.$$

Hence, $\frac{\overline{XA}}{\overline{XS}} = \frac{\overline{X'S'}}{\overline{X'V}}$ or $\frac{\overline{AS}}{\overline{SX}} = \frac{\overline{S'V}}{\overline{VX'}}$.

Lemma 2.9. Let A_3 , B_3 , C_3 respectively be intersection of \mathcal{H}_P and AH, BH, CH. Prove that A_2A_3 , B_2B_3 , C_2C_3 are concurrent at P.

From two simple problems to the connection of special points



FIGURE 18. Proof of Lemma 2.9

Proof. We construct diameter HH' of \mathcal{H}_P . From Lemma 2.7, we have $\mathbf{S}_{M_{BC}} \circ \mathbf{H}^2_A : O, P' \mapsto H, H'.$

Therefore $OP' \parallel HH'$ and $\overline{H'H} = 2\overline{OP'}$ so H' is the Anticomplement point of P' in triangle ABC. On the other hand, $A_3H' \parallel BC$ and $BC \perp AH$ hence AA' = 2d(P', BC). Using Lemma 2.8, we have

$$\frac{\overline{AP}}{\overline{PA_1}} = \frac{\mathrm{d}(P', BC)}{\mathrm{d}(A_1', BC)} = \frac{\mathrm{d}(P', BC)}{\mathrm{d}(A_1, BC)} = \frac{\overline{AA_3}}{\overline{A_2A_1}}$$

This means A_2A_3 passes P. Similarly, B_2B_3 , C_2C_3 passes P. We have solved the lemma.

Back to Theorem 2.10,



FIGURE 19. Proof of Theorem 2.10

Proof. By Corollary 2.3, the Brocard cirlce is the Hagge circle of G with respect to triangle $M_{BC}M_{CA}M_{AB}$. By applying Lemma 2.7 and 2.9 for triangle $M_{BC}M_{CA}M_{AB}$, we have the result.

3. Properties related to the second problem

Theorem 3.1. A_Y is the projection of orthocenter H on the A-median line of triangle ABC.



FIGURE 20. Proof of Theorem 3.1

Proof. Let AA_O be diameter of (ABC). Since $BHCA_O$ is a parallelogram, M_{BC} is the midpoint of A_OH . On the other hand, M_{BC} is also the mid point of A_YA_M . By Theorem 2.3, $HA_YA_OA_M$ is a parallelogram therefore $A_YH \parallel A_OA_M$. Moreover, $A_OA_M \perp AA_M$ hence A_Y is the projection of H on the A-median line.

Corollary 3.1. Let B_Y and C_Y be constructed in a similar manner as A_Y , but with respect to B and C respectively. Let $H_A H_B H_C$ be the Orthocentroidal triangle with H_A , H_B , H_C representively be projection of G on AH, BH, CH. Then two triangles $A_Y B_Y C_Y$ and $H_A H_B H_C$ are inscribed in the circle with diameter GH which called Orthocentroidal circle.

Corollary 3.2. The Hagge circle of the Lemoine point L in triangle ABC is the Orthocentroidal circle of this triangle.

Theorem 3.2. The radical line of $(A_X B_X C_X)$ and $(A_Y B_Y C_Y)$ passes midpoint X (39) of two Brocard points.

We can see the proof of this property here [9] using Barycentric coordinate.

Theorem 3.3. A_YH_A , B_YH_B , C_YH_C are concurrent at L.

From two simple problems to the connection of special points



FIGURE 21. Demonstration of Corollary 3.2



FIGURE 22. Demonstration of Theorem 3.2

Lemma 3.1. L is also the Lemoine point of triangle $H_A H_B H_C$.

Proof. By Corollary 3.2, the Orthocentroidal circle of triangle ABC is the Hagge circle of L with respect to this triangle. Moreover, by Corollary 2.1, triangle $A_Y B_Y C_Y$ is the Hagge triangle of L, which means A_Y , B_Y , C_Y are constructed in the same way of A_2 , B_2 , C_2 in Lemma 2.7, but with respect to L. We then apply Lemma 2.9 to completely solve this result. Back to Theorem 3.3,

Proof. By applying Lemma 2.7, 2.9 and 3.1 for triangle ABC with Lemoine point L and the Orthocentroidal circle, we can prove the theorem.



FIGURE 23. Proof of Theorem 3.3

Theorem 3.4. The perpendicular bisector of AXA_{GY} cuts AL at L_N . Then $AXA_Y \parallel L_NM_{BC}$.



FIGURE 24. Proof of Theorem 3.4

Proof. By Corollary 2.2, L_N also lies on the perpendicular bisector of $M_{AB}M_{CA}$ containing A_N so $A_NL_N \parallel A_YA_S$. Therefore

 $\overline{AL_N} \cdot \overline{AA_Y} = \overline{AA_N} \cdot \overline{AA_S} = \overline{AM_{BC}} \cdot \overline{AA_X}$ so $\overline{\frac{AA_X}{AL_N}} = \overline{\frac{AA_Y}{AM_{BC}}}$. Hence, $A_X A_Y \parallel L_N M_{BC}$. **Theorem 3.5.** $(A_X A_{GY} L_N)$ is tangent to (BOC).



FIGURE 25. Proof of Theorem 3.5

Proof. By Theorem 2.1, A_X lies on (BOC). Let Ad be a ray which belongs to tangent line at A of $(A_X A_{GY} L_N)$ such that Ad and B are on the same side of the plane separated by AL. By Theorem 2.1, $A_X A_S$ is the A_X -bisector of triangle $A_X BC$. We have

$$(A_X d, A_X B) = (A_X d, A_X A_S) + (A_X A_S, A_X B)$$
$$= (A_X A_S, A_X A_{GY}) + (A_X C, A_X A_S)$$
$$= (A_X C, A_X A_{GY})$$
$$= (CB, CA_X).$$

So Ad is the tangent line of $(A_X BC) \equiv (BOC)$. We have done.

Theorem 3.6. Let A_{LX} be projection of H on AL. Then $M_{BC}A_Y = M_{BC}A_{LX}$.



FIGURE 26. Proof of Theorem 3.6

Proof. Draw altitudes BH_{CA} and CH_{AB} of triangle ABC. It can be seen that $A, H, H_{CA}, H_{AB}, A_Y, A_{LX}$ lie on the circle with diameter AH. Since M_{BC} is the intersection of two tangent lines at H_{CA} and H_{AB} of (AH) so $AH_{AB}A_YH_{BC}$ is a harmonic quadrilateral. Since AL is a median line of triangle $AH_{CA}H_{AB}$ hence $H_{CA}H_{AB} \parallel A_YA_{LX}$. Moreover, as M lies on the perpendicular bisector of $H_{CA}H_{AB}$, which is also the perpendicular bisector of A_YA_{LX} , so that $M_{BC}A_Y = M_{BC}A_{LX}$.

Corollary 3.3. $(A_Y M_{BC} A_{LX})$ is tangent to the Euler circle of triangle ABC.

Theorem 3.7. Let L_{BC} be intersection of AL and BC. Then $A_Y L_{BC}$ is a symmetrian line of triangle $A_Y BC$.



FIGURE 27. Proof of Theorem 3.7

Proof. Since AL_{BC} is a symmetrian line of triangle ABC so

$$\frac{\overline{L_{BC}B}}{\overline{L_{BC}C}} = -\left(\frac{AB}{AC}\right)^2.$$

On the other hand,

$$\left(\frac{AB}{AC}\right)^2 = \left(\frac{AB}{M_{BC}B} \cdot \frac{M_{BC}C}{AC}\right)^2 = \left(\frac{A_MC}{A_MM_{BC}} \cdot \frac{A_MM_{BC}}{A_MB}\right)^2 = \left(\frac{A_YB}{A_YC}\right)^2$$
So

$$\frac{\overline{L_{BC}B}}{\overline{L_{BC}C}} = -\left(\frac{A_YB}{A_YC}\right)^2.$$

We conclude that $A_Y L_{BC}$ is a symmetrian line of triangle $A_Y BC$.

Theorem 3.8. $(A_Y M_{BC} A_{LX})$ is tangent to (BHC).

Lemma 3.2. Given triangle ABC. On segment BC, we choose D, E such that AD, AE are isogonal conjugate of each other in this triangle. Prove that (ABC) is tangent to (ADE).



FIGURE 28. Proof of Lemma 3.2

Proof. The tangent line at A of (ABC) cuts line BC at F. Then (AF, AD) = (AF, AB) + (AB, AD) = (CA, CB) + (EA, CA) = (EA, CB)

so AF is tangent to (ADE). Thus, (ABC) is tangent to (ADE). Back to Theorem 3.8,



FIGURE 29. Proof of Theorem 3.8

Proof. Draw the altitude AH_{BC} of triangle ABC. We have that H_{BC} and A_{LX} lie on the circle with diameter HL_{BC} . Therefore,

 $(L_{BC}A_{LX}, L_{BC}M_{BC}) = (HA_{LX}, HH_{BC}) = (A_YA_{LX}, A_YA).$

So $A_Y, A_{LX}, L_{BC}, M_{BC}$ are cyclic. Using Theorem 3.7, $A_Y L_{BC}$ is a symmedian line of triangle $A_Y BC$. We use Lemma 3.2 to conclude that $(A_Y M_{BC} A_{LX})$ touches (BHC).

Theorem 3.9. The rays BA_Y , CA_Y cuts (ABC) at A_{BY} , A_{CY} . Let A'_{BY} , A'_{CY} be symmetry points of A_{BY} , A_{CY} through CA, AB respectively. Then A'_{BY} and A'_{CY} lie on AL.



FIGURE 30. Proof of Theorem 3.9

Proof. We have

$$(AA'_{BY}, AC) = (AC, AA_{BY}) = (BC, BA_{BY})$$
$$= (CB, CA_M) = (AB, AA_M)$$
$$= (AL, AC).$$

Thus AL passes A'_{BY} . Similarly, AL passes A'_{CY} .

Theorem 3.10. A_{LX} is midpoint of $A'_{BY}A'_{CY}$.

Proof. Rays BH and CH cut (ABC) respectively at H_B and H_C . We have

$$H_BA_{CY}, H_BB) = (CA_{CY}, CB) = (HA_Y, HH_B)$$

so $H_BA_{CY} \parallel HA_Y$, where $HA_Y \perp AG$. Simiarly $H_CA_{BY} \parallel HA_Y$, where $HA_Y \perp AG$ hence $H_BA_{CY}H_CA_{BY}$ is a isosceles trapezoid. We conclude that $H_CA_{CY} = H_BA_{BY}$. Using symmetry transformation through AB and AC, we have $\mathbf{S}_{AB} \colon H, A_{CY} \mapsto H_C, A'_{CY}$ and $\mathbf{S}_{AC} \colon H, A_{BY} \mapsto H_B, A'_{BY}$. Therefore $HA'_{BY} = H_BA_{BY} = H_CA_{CY} = HA'_{CY}$, which means triangle $HA'_{BY}A'_{CY}$ is isosceles at H. Moreover, A_{LX} is the projection of H on $A'_{BY}A'_{CY}$ so A_{LX} is midpoint of $A'_{BY}A'_{CY}$.

4. CONCLUSION

This article is a discussion on the properties associated with a pair of distinct triangle centers. Further investigations can be conducted in this domain to delve deeper into the subject matter and uncover additional distinctive characteristics pertaining to triangles.

References

- [1] AoPS site Admins, USA Team Selection Test 2008, available at https://artofproblemsolving.com/community/c4638.
- [2] Allen Liu, ELMO Shortlist 2013, Problem G3, available at https://web.evanchen.cc/exams/ELMO-2013-SL.pdf.
- [3] Clark Kimberling, *Encyclopedia Of Triangle Centers*, available at http://faculty.evansville.edu/ck6/encyclopedia/ETC.html
- [4] IMO Geometry Blogspot, IMO 1959 2021 116p, available at https://imogeometry.blogspot.com/p/1.html.
- [5] Jay Warendorff, *Hagge circle*, Wolfram Demonstrations Project, available at https://demonstrations.wolfram.com/TheHaggeCircle.
- [6] Jean-Louis Ayme, Brocard points properties, available at http://jl.ayme.pagesperso-orange.fr/vol7.html.
- [7] Jean-Louis Ayme, Hagge P-circle, available at http://jl.ayme.pagesperso-orange.fr/vol5.html.
- [8] Jean-Louis Ayme, A circle through six points problem, available at http://jl.ayme.pagesperso-orange.fr/vol9.html.
- [9] Jean-Louis Ayme, Brocard and Hagge circle, available at https://artofproblemsolving.com/community/c6h320075p1719735.
- [10] Nguyen Van Linh, *Circle passes through fixed points*, Nguyen Van Linh's Wordpress, available at

https://nguyenvanlinh.files.wordpress.com/2015/04/circles-pass-through-fixed-point.pdf.

- [11] TelvCohl, Some properties of Hagge circle, available at https://artofproblemsolving.com/community/c6h624925.
- [12] Weisstein, Eric W., First Brocard point, From MathWorld-A Wolfram Web Resource, available at https://mathworld.wolfram.com/FirstBrocardPoint.html.
- [13] Weisstein, Eric W., *Second Brocard point*, From MathWorld-A Wolfram Web Resource, available at http://mathworld.wolfram.com/SecondBrocardPoint.html.
- [14] Weisstein, Eric W., Brocard circle, From MathWorld-A Wolfram Web Resource, available at http://mathworld.wolfram.com/SecondBrocardPoint.html.
- [15] Weisstein, Eric W., Orthocentroidal circle, From MathWorld-A Wolfram Web Resource, available at http://mathworld.wolfram.com/SecondBrocardPoint.html.
- [16] Weisstein, Eric W., Lemoine point, From MathWorld-A Wolfram Web Resource, available at https://mathworld.wolfram.com/LemoinePoint.html.

FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, UNI-VERSITY OF SCIENCE, VIETNAM NATIONAL UNIVERSITY HO CHI MINH CITY, 227 NGUYEN VAN CU, WARD 4, DISTRICT 5, HO CHI MINH CITY

E-mail address: 21c29018@student.hcmus.edu.vn