Stabilizer entropies are monotones for magic-state resource theory

Lorenzo Leone^{1,*} and Lennart Bittel^{1,*}

¹Dahlem Center for Complex Quantum Systems, Freie Universität Berlin, 14195 Berlin, Germany

We establish the monotonicity of stabilizer entropies for $\alpha \geq 2$ within the context of magic-state resource theory restricted to pure states. Additionally, we show that linear stabilizer entropies serve as strong monotones. Furthermore, we provide the extension to mixed states of stabilizer entropies as magic monotones through convex roof constructions.

Introduction.— The inception of magic-state resource theory can be traced back to the seminal work of Bravvi and Kitaev [1]. They showed the feasibility of applying non-Clifford gates to a target state through stabilizer operations - i.e., Clifford operations and measurements leveraging a single-qubit ancilla state prepared in a nonstabilizer (or 'magic') state. Building upon this fundamental result, magic-state resource theory was then formalized [2]. The core principle hinges on the following dichotomy: while stabilizer operations are easily implemented in a fault-tolerant fashion (i.e., transversally) in many schemes [3], achieving fault-tolerance for nonstabilizer operations proves to be challenging [4]. As a result, stabilizer operations are regarded as the free operations within the resource theory framework, with stabilizer states designated as the corresponding *free states*, while non-Clifford operations and nonstabilizer states are considered resourceful. The task at hand involves utilizing nonstabilizer resource states and exclusively employing stabilizer operations to implement non-Clifford unitary gates fault-tolerantly, a process commonly referred to as magic-state distillation.

Stabilizer operations are not only non-universal, but have also been shown to be efficiently classically simulatable [5]. Nonstabilizer states become hard to simulate classically [6]. Hence, beyond the scope of magic-state distillation, magic-state resource theory is also employed to quantify the hardness of classically simulating quantum states using stabilizer formalism. From this perspective, nonstabilizer states are regarded as resourceful since they enable quantum computations that surpass the capabilities of classical computers.

From the technical point of view, the construction of a quantum resource theory is straightforward; given the set of free states \mathcal{T} uniquely defining the resource theory, free operations are quantum channels (completely positive, trace preserving and linear operations on a quantum system) leaving \mathcal{T} invariant. The challenge lies in selecting a *resource monotone*, akin to a "thermometer", to accurately quantify the resource amount in a given state $\psi \notin \mathcal{T}$, represented by a positive scalar function. While universal resource monotones (independent from the specific resource theory) do exist [7], they are impractical to compute and experimentally measure in real scenarios. For a resource theory to be applicable in practice, such as in quantum many-body physics and quantum computation, finding a resource-specific measure is an extremely important and nontrivial task.

Stabilizer entropies (SEs) indexed by a α -Rényi index have recently been proposed to probe nonstabilizerness in multiqubit quantum states [8]. They have garnered particular attention due to their analytical [9–11] and numerical [12, 13] computability as well as experimental measurability [14, 15]. Thanks to their computability advantages, stabilizer entropies have significantly advanced the study of the resource theory of magic within the context of many-body physics, allowing numerical simulations up to ~ 100 qubits [12, 13], which was previously limited to only a few bodies [16]. Beside, through stabilizer entropies, magic-state resource theory found connection to a number of topics in quantum physics, including quantum chaos and scrambling [8], as well as in quantum information with applications in cryptography [17], fidelity estimation [18], Pauli sampling [19], and classical shadows [20].

However, a counterexample to the monotonicity of stabilizer entropies with Rényi index $\alpha < 2$ under stabilizer protocols was provided [21]. Since then, the question of whether stabilizer entropies with $\alpha \geq 2$, when restricted to pure states, serve as a monotone for magic-state resource theory has remained unanswered. Given the significance of stabilizer entropies across various contexts, the question "are stabilizer entropies good monotones for magic-state resource theory?" has garnered increasing attention.

In this work, we demonstrate that stabilizer entropies with integer Rényi index $\alpha \geq 2$ are monotones for magicstate resource theory when restricted to pure states. We also show that their linearized versions – linear stabilizer entropies – serve as *strong* monotones. Moreover, we extend stabilizer entropies to mixed states while ensuring monotonicity under stabilizer protocols. See Table I for a summary. These results conclusively validate the use of stabilizer entropies to quantify "magic" in multiqubit quantum systems.

	Magic Monotone	Strong Magic Monotone
$M_{\alpha < 2}$	×	×
$M_{\alpha \ge 2}$	\checkmark	×
$M_{\alpha<2}^{\rm lin}$	×	×
$M_{\alpha \ge 2}^{\text{lin}}$	\checkmark	1

TABLE I. Summary of the results. M_{α} and M_{α}^{lin} represent the stabilizer entropy and the linear stabilizer entropy respectively (see Definition 4). Both are extended to mixed states.

Setup.— Throughout the paper we consider the Hilbert space \mathcal{H}_n of n qubits and denote as $d_n = 2^n$ its dimension. A natural operator basis is given by Pauli operators $P \in \mathbb{P}_n$, i.e. n-fold tensor products of ordinary Pauli matrices I, X, Y, Z. The subgroup of unitary matrices that maps Pauli operators to Pauli operators is known as the Clifford group. Stabilizer states, denoted as $|\sigma\rangle$ in this work, are pure states obtained from $|0\rangle^{\otimes n}$ with the action of unitary Clifford operators. Throughout the work, we will use ψ, ϕ to denote pure states, while ρ to denote (possibly mixed) general states.

Magic-state resource theory.— A resource theory is uniquely characterized by its set of free states. The set of free states for magic-state resource theory is given by the convex hull of pure stabilizer states, hereby denoted as STAB := $\{\sum_i p_i |\sigma_i\rangle \langle \sigma_i| : p_i \ge 0, \sum_i p_i = 1\}$.

Free operations are the *stabilizer protocols*, that are completely positive and trace preserving maps that leave invariant the set STAB. Denoting as \mathcal{S} the set of stabilizer protocols, every operation $\mathcal{E} \in \mathcal{S}$ can be built out of the following elementary operations: (i) Clifford unitaries; (ii) partial trace; (iii) measurements in the computational basis; (iv) composition with ancillary qubits in $|0\rangle$; the above operations conditioned on (v) measurement outcomes and (vi) classical randomness. Since these operations (and combination thereof) return different states based on conditioned operations and non-deterministic strategies, the most general form of a stabilizer protocol applied on a state ρ is $\mathcal{E}(\rho) = \{(p_i, \rho_i)\}$, i.e. a collection of n_i qubit states ρ_i with probabilities p_i . A stabilizer protocol \mathcal{E} is deterministic if $\mathcal{E}(\rho) = \tilde{\rho}$, i.e. a unique quantum state is reached with unit probability.

Given stabilizer protocols, one can define monotones for magic-state resource theory.

Definition 1 (Magic monotone). A monotone \mathcal{M} is a real-valued function for all $n \in \mathbb{N}$ qubit systems (or collection thereof) such that: (i) $\mathcal{M}(\rho) = 0$ if and only $\rho \in \text{STAB}$; (ii) \mathcal{M} is nonincreasing under stabilizer protocols S, i.e.

$$\mathcal{M}(\mathcal{E}(\rho)) \le \mathcal{M}(\rho), \quad \forall \mathcal{E} \in \mathcal{S}.$$
 (1)

As customary in resource theories [7], there are monotones that always work regardless of the specific resource theory, which includes (but is not limited to) the robustness of magic [22] or the relative entropy of magic [23]. In this regard, the challenge has always been to find resource monotones that meaningfully apply to the specific resource theory at hand, e.g., von Neumann entanglement entropy for entanglement. In magic-state resource theory, such a role is played by the stabilizer Rényi entropy [8]. As we shall see, the stabilizer entropy is naturally defined for pure states. We thus define a monotone for magic-state resource theory restricted to pure states.

Definition 2 (Pure-state magic monotone). A pure-state magic monotone is a real-valued function for all $n \in \mathbb{N}$

qubit systems such that: (i) $\mathcal{M}(\psi) = 0$ if and only if $|\psi\rangle$ is a pure stabilizer state; (ii) for every pair $(|\psi\rangle, \mathcal{E})$, where $\mathcal{E} \in S$ a stabilizer protocol, obeying $\mathcal{E}(|\psi\rangle\langle\psi|) = |\phi\rangle\langle\phi|$, it holds that:

$$\mathcal{M}(\phi) \le \mathcal{M}(\psi) \,. \tag{2}$$

In words, a pure-state magic monotone is a monotone when restricted to *deterministic* pure-state stabilizer protocols, i.e., those mapping pure states to pure states. Note that, as stabilizer protocols consist of multiple elementary operations, the aforementioned definition does not preclude intermediate states from becoming mixed, provided the final state remains pure. Clearly, being a pure-state monotone is a weaker condition than being a monotone and, as a matter of fact, Definition 1 implies Definition 2. However, deterministic pure-state stabilizer protocols have their own relevance: a famous example is provided by the magic-state injection, i.e., $\mathcal{E}(|T\rangle \otimes |\psi\rangle) = |0\rangle \otimes T |\psi\rangle$ where $|T\rangle = T |+\rangle$ is the Tstate and $T \coloneqq \text{diag}(1, e^{i\pi/4})$ being the T-gate [24]. As a last notion, let us introduce the concept of strong monotonicity for pure states. An analogous definition for mixed states follows.

Definition 3 (Strong pure-state magic monotone). Let \mathcal{M} a pure-state magic monotone. Let $|\psi\rangle$ be a pure state and consider the collection of pure states $\mathcal{E}(\psi) = \{(p_i, |\phi_i\rangle)\}$, obtained after applying a stabilizer protocol $\mathcal{E} \in \mathcal{S}$. \mathcal{M} is a strong monotone if

$$\mathcal{M}(\psi) \ge \sum_{i} p_i \mathcal{M}(\phi_i) \,. \tag{3}$$

Informally, the notion of strong monotonicity says that, on average, the resource of magic cannot increase after non-deterministic stabilizer protocols. Now that we have all the necessary definitions and tools, the next section reveals the main findings of our work.

Stabilizer entropies as magic monotones.— In this section, we establish the monotonicity of stabilizer entropies under stabilizer protocols in various flavors. Let us first define SEs. Given a pure state $|\psi\rangle$, the quantity $d_n^{-1}|\langle\psi|P|\psi\rangle|^2$ forms a probability distribution on the Pauli group \mathbb{P}_n , known also as *characteristic distribution* [25]. SEs are defined, up to an offset, as α -Rényi entropies of this probability distribution.

Definition 4 (α -stabilizer entropy). The α -Rényi stabilizer entropy of a pure quantum state $|\psi\rangle$ on n qubits reads

$$M_{\alpha}(\psi) \coloneqq \frac{1}{1-\alpha} \log P_{\alpha}(\psi), \ P_{\alpha}(\psi) \coloneqq \frac{1}{d_n} \sum_{P \in \mathbb{P}_n} |\langle \psi | P | \psi \rangle|^{2\alpha}$$

where $P_{\alpha}(\psi)$ is referred to as stabilizer purity. Linear stabilizer entropies are defined as $M_{\alpha}^{\text{lin}}(\psi) \coloneqq 1 - P_{\alpha}(\psi)$.

Stabilizer entropies exhibit two key properties required for magic-state resource theory: (i) they are minimized, i.e., are zero, only for stabilizer states and (ii) are invariant under the action of Clifford unitary operators. Additionally, SEs exhibit numerous useful properties and serve as lower bounds for key monotones within magic state resource theory, including the robustness of magic [22], minrelative entropy of magic [23], stabilizer nullity [26] and stabilizer extent [27]. For a comprehensive list, please refer to Appendix A. Operationally, SE quantifies the spread of a state in the operator basis of Pauli operators and, as such, can also be interpreted as the entropy of Pauli tomography [28].

However, mere invariance under Clifford operators does not suffice to establish stabilizer entropies as effective monotones for magic-state resource theory, given the significantly broader scope of stabilizer protocols. In fact, in a recent study (Ref. [21]), a specific counterexample to deterministic pure-state stabilizer protocols involving conditioned Clifford operations has been presented for $\alpha < 2$. Whether stabilizer entropies serve as monotones for larger values of α has remained an open question, and given the extensive list of useful features, it is indeed significant. We demonstrate below that stabilizer entropies function as monotones for deterministic stabilizer protocols for any $\alpha \geq 2$, thereby qualifying as pure-state magic monotones according to Definition 2.

Theorem 1. Stabilizer entropies M_{α} are pure-state magic monotones for every integer $\alpha \geq 2$.

Proof sketch. First of all, it is crucial to note that a deterministic pure-state stabilizer protocol only necessitates the initial and final states to be pure. We consider the following decomposition of a pure state $|\psi\rangle = \sqrt{p} |0\rangle \otimes |\phi_1\rangle + \sqrt{1-p} |1\rangle \otimes |\phi_2\rangle$. The key step of the proof is showing that $M_{\alpha}(\psi) \geq \min\{M_{\alpha}(\phi_1), M_{\alpha}(\phi_2)\}$. From this simple observation, for a general stabilizer protocol $\mathcal{E} \in \mathcal{S}$, it holds that

$$M_{\alpha}(\psi) \ge \inf_{(q_{ij},\phi_{ij})} \left\{ \min_{ij} M_{\alpha}(\phi_{ij}) : p_i \rho_i = \sum_j q_{ij} \phi_{ij} \right\},\$$

where the infimum is over the possible pure state convex decompositions of $p_i \rho_i$ where $\mathcal{E}(\psi) = \{(p_i, \rho_i)\}$. By requiring \mathcal{E} to be a deterministic pure-state stabilizer protocol, the result just follows. The complete proof can be found in Appendix C.

In fact, we can present an even stronger result. We show that linear stabilizer entropies are strong monotones according to Definition 3.

Theorem 2. Linear stabilizer entropies M_{α}^{lin} are strong magic monotones for every integer $\alpha \geq 2$.

Proof sketch. To show the strong pure-state monotonicity, we prove the more general result from which Theorem 3 follows. Starting from a pure state ψ , for all stabilizer operations $\mathcal{E} \in \mathcal{S}$ such that $\mathcal{E}(\psi) = \{(p_i, \rho_i)\}$, it holds that

$$M_{\alpha}^{\mathrm{lin}}(\psi) \ge \inf_{(q_{ij},\phi_{ij})} \left\{ \sum_{ij} q_{ij} M_{\alpha}^{\mathrm{lin}}(\phi_{ij}) : p_i \rho_i = \sum_j q_{ij} \phi_{ij} \right\}.$$

Similarly to Theorem 1, the crucial step is to consider an arbitrary pure state $|\psi\rangle = \sqrt{p} |0\rangle \otimes |\phi_1\rangle + \sqrt{1-p} |1\rangle \otimes |\phi_2\rangle$ and show that $M_{\alpha}^{\text{lin}}(\psi) \geq p M_{\alpha}^{\text{lin}}(\phi_1) + (1-p) M_{\alpha}^{\text{lin}}(\phi_2)$. With the restriction imposed on the collection $\mathcal{E}(\psi)$ to consist solely of pure states, the result follows.

According to Theorem 2, linear stabilizer entropies are powerful magic monotones defined through entropies of the characteristic distribution. In [21], it has been shown that stabilizer entropies M_{α} violate the strong monotonicity condition. However, since the two quantities are related to each other through a concave function, as $M_{\alpha} = \frac{1}{1-\alpha} \log(1-M_{\alpha}^{\text{lin}})$, ultimately this discrepancy is due to the Jensen's inequality which is equivalent to the difference between arithmetic mean and geometric mean. Let us present a counterexample for M_{α} for $\alpha \geq 2$. We consider the state $|\psi\rangle \propto |0\rangle^{\otimes n} + |1\rangle \otimes |\phi\rangle$ with $|\phi\rangle$ a Haar random state on n-1 qubits. We have $M_{\alpha}(|\psi\rangle) = O(1)$, as $|\psi\rangle$ has a non-vanishing overlap to the stabilizer state $|0\rangle^{\otimes n}$ (see Appendix A). While $M_{\alpha}(|\phi\rangle) = \Omega(n)$ (with overwhelming probability [17]). As such, there exist n_0 such that $\forall n \geq n_0$ one has $M_{\alpha}(|\psi\rangle) \leq \frac{1}{2}M_{\alpha}(|\phi\rangle)$, thus violating strong monotonicity of M_{α} for some sufficiently large n.

However, as a corollary of Theorem 1, one can see that stabilizer entropies obey a weaker condition compared to strong pure-state monotonicity. Given a nondeterministic stabilizer protocol that, starting from a pure state $|\psi\rangle$, yields a collection of pure states $\mathcal{E}(\psi) =$ $\{(p_i, |\phi_i\rangle)\}$, it follows that $M_{\alpha}(\psi) \geq \min_i M_{\alpha}(\phi_i)$ for every integer $\alpha \geq 2$.

That being said, while strong monotonicity is indeed a desirable trait with clear operational significance, it is worth noting that it is not an essential requirement for effective monotones. Min-relative entropy of magic [23], which is a natural monotone being the (negative log) value of the overlap with the closest stabilizer state, does not obey strong monotonicity either (the counterexample shown above holds true). Consequently, the adherence of linear stabilizer entropies to strong monotonicity can be regarded as a distinctive and advantageous feature.

Extension to mixed states. — While stabilizer entropies find their natural definition within the realm of pure states, when extending magic-state resource theory to mixed states, a diverse landscape emerges. This becomes evident when considering that magic-state ancillas are often assumed to be noisy, prompting interest in extracting a single clean non-Clifford unitary from many copies of these noisy ancillas [1]. To conclude, we leverage our findings to extend stabilizer entropies to mixed states, ensuring adherence to the monotonicity criterion outlined in Definition 1.

Definition 5 (Convex-roof extension of SEs). Let $C := \{(p_i, \rho_i)\}$ be a collection of n_i qubit quantum states. Extended stabilizer entropies are defined as

$$\widehat{M}_{\alpha}(\mathcal{C}) \coloneqq \frac{1}{1-\alpha} \log \widehat{P}_{\alpha}(\mathcal{C}), \quad \widehat{M}_{\alpha}^{\mathrm{lin}}(\mathcal{C}) \coloneqq 1 - \widehat{P}_{\alpha}(\mathcal{C}) \quad (4)$$

$$\widehat{P}_{\alpha}(\mathcal{C}) \coloneqq \sup_{(q_{ij},\phi_{ij})} \left\{ \sum_{ij} q_{ij} P_{\alpha}(\phi_{ij}) \right\}$$

The sup is over the possible convex pure-state decompositions $p_i \rho_i = \sum_j q_{ij} \phi_{ij}$ for $\{(p_i, \rho_i)\} \in C$.

It is worth noting that both the *extended* stabilizer entropy and linear stabilizer entropy reduce to the entropies defined on pure states in Definition 4. The following theorem establishes the monotonicity of the above convex roof constructions.

Theorem 3. Extended stabilizer entropies \widehat{M}_{α} and linear stabilizer entropies $\widehat{M}_{\alpha}^{\text{lin}}$ are monotones for every integer $\alpha \geq 2$. Moreover, $\widehat{M}_{\alpha}^{\text{lin}}$ is a strong monotone.

Proof. Both results for extended stabilizer entropy and linear stabilizer entropy follows by showing that, given a collection $C = \{(p_i, \rho_i)\}$ of n_i qubit states and an elementary stabilizer protocol $\mathcal{E} \in \mathcal{S}$, then:

$$\widehat{P}_{\alpha}(\mathcal{E}(\mathcal{C})) \ge \widehat{P}_{\alpha}(\mathcal{C}) \,. \tag{5}$$

The steps follow from the proof of Theorem 2. $\hfill \Box$

The above result extends stabilizer entropies, along with some of their amenable properties, to the realm of mixed states, thus unlocking the exploration of the magicstate resource theory in many-body physics in more intricate situations. Finding the optimal decomposition may prove challenging in practice, whereas even sub-optimal ones can provide useful upper-bounds for mixed states. While the convex-roof extension in Definition 5 reduces to the stabilizer entropy for ρ being pure, it is worth noting that it represents just one valid extension to mixed states. An alternative approach could involve considering the extremum of stabilizer entropies across potential purifications, effectively emulating the entanglement of purification [29].

Discussion.— In this work, we proved that stabilizer entropies serve as effective monotones for magic-state resource theory. Furthermore, their linearized counterparts exhibit strong monotonicity, meaning they do not, on average, increase following a non-deterministic protocol. Additionally, as a result of our investigation, we provided a convex-roof extension of stabilizer entropies to mixed states, ensuring monotonicity under stabilizer protocols and paving the way for further exploration of magic-state resource theory beyond pure states.

While our results solidify the utility of stabilizer entropies in quantifying "magic" within multiqubit systems, there remain intriguing open questions regarding their properties, which remain mostly unexplored. For instance, despite robust metrics such as robustness of magic, stabilizer extent, and stabilizer nullity directly quantifying the cost of classical simulations via stabilizer formalism, there lacks a simulation algorithm directly assessed by stabilizer entropy costs. Furthermore, the interconversion rates between different resource states under the lens of stabilizer entropies have not been investigated. In this context, the maximal resourceful state according to SEs remains unknown. While it is evident that the upper bound $M_2 < \log \frac{d_n+1}{2}$ cannot be achieved by any characteristic distribution (for $n \neq 1, 3$), a tighter bound and the state saturating it are yet to be determined.

In conclusion, this paper provides a definitive answer to the open question of whether stabilizer entropies serve as monotones for magic-state resource theory or merely act as weak detectors of nonstabilizerness. With this clarification, there arises a compelling need for further exploration of stabilizer entropies across diverse domains, ranging from many-body physics to classical simulation via stabilizer formalism, and within the realms of theoretical resource theory.

Acknowledgments.— We acknowledge an inspiring discussion with Ludovico Lami. We are grateful to Salvatore F.E. Oliviero for the careful reading of the manuscript. We thank Alioscia Hamma and Francesco A. Mele for their valuable comments and discussion. We thank Ryuji Takagi for having pointed out a mistake in an earlier version. L.L. is funded through the Munich Quantum Valley project (MQV-K8) by Bayerisches Staatsministerium für Wissenschaft und Kunst. L.B. has been funded by DFG (FOR 2724, CRC 183), the Cluster of Excellence MATH+ and the BMBF (MuniQC-Atoms).

- * Contributed equally. {lorenzo.leone, l.bittel}@fu-berlin.de
- S. Bravyi and A. Kitaev, Physical Review A 71, 022316 (2005).
- [2] V. Veitch, S. A. Hamed Mousavian, D. Gottesman, and J. Emerson, New Journal of Physics 16, 013009 (2014).
- [3] E. T. Campbell and D. E. Browne, Physical Review Letters 104, 030503 (2010).
- [4] B. Eastin and E. Knill, Phys. Rev. Lett. 102, 110502 (2009).
- [5] D. Gottesman, "The Heisenberg Representation of Quantum Computers," (1998), arxiv:quant-ph/9807006.
- [6] S. Aaronson and D. Gottesman, Physical Review A 70, 052328 (2004).
- [7] E. Chitambar and G. Gour, Review of Modern Physics 91, 025001 (2019).
- [8] L. Leone, S. F. E. Oliviero, and A. Hamma, Physical Review Letters 128, 050402 (2022).
- [9] G. Passarelli, R. Fazio, and P. Lucignano, "Nonstabilizerness of permutationally invariant systems," (2024), arXiv:2402.08551 [quant-ph].
- [10] S. F. E. Oliviero, L. Leone, and A. Hamma, Phys. Rev. A 106, 042426 (2022).
- [11] T. Haug and L. Piroli, Phys. Rev. B 107, 035148 (2023).
- [12] G. Lami and M. Collura, Phys. Rev. Lett. 131, 180401 (2023).
- [13] P. S. Tarabunga, E. Tirrito, M. C. Bañuls, and M. Dalmonte, "Nonstabilizerness via matrix product states in the pauli basis," (2024), arXiv:2401.16498 [quant-ph].
- [14] S. F. E. Oliviero, L. Leone, A. Hamma, and S. Lloyd, npj Quantum Inf 8, 1 (2022).

(2023).

arxiv:2305.19152 [quant-ph]. [16] M. Heinrich and D. Gross, Quantum **3**, 132 (2019/april).

bilizer entropies for quantum computers,"

- [17] A. Gu, L. Leone, S. Ghosh, J. Eisert, S. Yelin, and
- Y. Quek, "A Little Magic Means a Lot," (2023), arXiv:2308.16228 [quant-ph].
- [18] L. Leone, S. F. E. Oliviero, and A. Hamma, Phys. Rev. A 107, 022429 (2023).
- [19] M. Hinsche, M. Ioannou, S. Jerbi, L. Leone, J. Eisert, and J. Carrasco, "Efficient distributed inner product estimation via pauli sampling," (In preparation).
- [20] R. Brieger, M. Heinrich, I. Roth, and M. Kliesch, "Stability of classical shadows under gate-dependent noise," (2023), arXiv:2310.19947 [quant-ph].
- [21] T. Haug and L. Piroli, "Stabilizer entropies and nonstabilizerness monotones," (2023), arxiv:2303.10152 [condmat, physics:quant-ph].
- [22] M. Howard and E. Campbell, Physical Review Letters 118, 090501 (2017).

- [23] Z.-W. Liu and A. Winter, PRX Quantum 3, 020333 (2022).
- [24] X. Zhou, D. W. Leung, and I. L. Chuang, Physical Review A 62 (2000), 10.1103/physreva.62.052316.
- [25] H. Zhu, R. Kueng, M. Grassl, and D. Gross, "The Clifford Group Fails Gracefully to Be a Unitary 4-Design," (2016), 1609.08172 [quant-ph].
- [26] M. Beverland, E. Campbell, M. Howard, and V. Kliuchnikov, Quantum Science and Technology 5, 035009 (2020).
- [27] S. Bravyi, D. Browne, P. Calpin, E. Campbell, D. Gosset, and M. Howard, Quantum 3, 181 (2019).
- [28] L. Leone, S. F. E. Oliviero, and A. Hamma, "Learning t-doped Stabilizer States," (2023), arXiv:2305.15398 [quant-ph].
- [29] B. M. Terhal, M. Horodecki, D. W. Leung, and D. P. Di-Vincenzo, Journal of Mathematical Physics 43, 4286–4298 (2002).
- [30] S. Bravyi and D. Gosset, Physical Review Letters 116, 250501 (2016).

Appendix A: Stabilizer entropy

In this section, we list the properties of stabilizer entropies explored in a number of works. First of all, let us recall the definition. Let ψ be a pure state. Given the characteristic distribution $d_n^{-1} \operatorname{tr}^2(P\psi)$, stabilizer entropies are defined

$$M_{\alpha}(\psi) = \frac{1}{1-\alpha} \log \frac{1}{d_n} \sum_{P \in \mathbb{P}_n} \operatorname{tr}^{2\alpha}(P\psi).$$
(A1)

It obeys the following properties:

- $M_{\alpha}(\sigma) = 0$ if and only if σ is a pure stabilizer state;
- it is invariant under unitary Clifford operations C, i.e. $M_{\alpha}(C\psi C^{\dagger}) = M_{\alpha}(\psi);$
- it is additive $M_{\alpha}(\psi \otimes \phi) = M_{\alpha}(\psi) + M_{\alpha}(\phi);$
- ordered $M_{\alpha}(\psi) \ge M_{\beta}(\psi)$ if $\alpha < \beta$;
- it is bounded $M_{\alpha} \leq \log d_n$ and, for every $\alpha \geq 2$, the tighter bound $M_{\alpha} \leq \log(d_n + 1) 1$ holds
- it lower bounds the stabilizer nullity ν [26], $M_{\alpha}(\psi) \leq \nu(\psi)$. Given a state $|\psi\rangle$, the stabilizer nullity is proportional to the dimension $\nu(\psi) = n \dim(G)$ of the largest Abelian subgroup G of the Pauli group that stabilizes $|\psi\rangle$, i.e., $\forall P \in G$ then $P |\psi\rangle = |\psi\rangle$.
- for $\alpha \geq 2$, it lower bounds the min-relative entropy of magic [23], i.e. $M_{\alpha}(\psi) \leq \frac{2\alpha}{\alpha-1}D_{\min}(\psi) \coloneqq -\frac{2\alpha}{\alpha-1}\log\max_{\sigma}|\langle\psi|\sigma\rangle|^2$ [21];
- for $\alpha \geq \frac{1}{2}$, it lower bounds the Robustness of magic [8], i.e. $M_{\alpha} \leq 2\log \mathcal{R}(\psi) \coloneqq 2\min_x \{\sum_i |x_i| : \rho = \sum_i x_i \sigma_i, \sigma_i \in \text{STAB}\}$ [22];
- for $\alpha \geq 2$, it lower bounds the stabilizer extent [30], i.e. $M_{\alpha} \leq 4 \log \xi(\psi) \coloneqq \min\{\sum_{i} |x_{i}| : |\psi\rangle = \sum_{i} x_{i} |\sigma_{i}\rangle, |\sigma_{i}\rangle \in \text{STAB}\};$
- for $\alpha = 1$, it obeys a Fannes-type inequality [17]. For any two states $|\psi\rangle$ and $|\phi\rangle$, it holds that

$$|M_{1}(\psi) - M_{1}(\phi)| \leq \begin{cases} \|\psi - \phi\|_{1} \log(d_{n}^{2} - 1) + H_{\text{bin}}[\|\psi - \phi\|_{1}] & \text{for } \|\psi - \phi\|_{1} \leq 1/2, \\ \|\psi - \phi\|_{1} \log(d_{n}^{2} - 1) + 1 & \text{for } \|\psi - \phi\|_{1} > 1/2 \end{cases}$$
(A2)

• for every odd integer $\alpha \ge 2$, it can be measured efficiently. In particular $P_{\alpha}(\psi)$ (stabilizer purity) can be measured using $O(n\epsilon^{-2})$ many samples of the state $|\psi\rangle$ and consuming two copies of the state at a time [15].

Appendix B: Preliminary results

Lemma 1. Consider $|\psi\rangle = \sqrt{p} |0\rangle \otimes |\phi_1\rangle + \sqrt{1-p} |1\rangle \otimes |\phi_2\rangle$ and $|\phi_1\rangle, |\phi_2\rangle \in \mathbb{C}^{2\otimes(n-1)}$. For any integer $\alpha \geq 2$, it holds that

$$P_{\alpha}(\psi) \leq \sum_{i=0}^{\alpha} {\binom{2\alpha}{2i}} p^{2i} (1-p)^{2(\alpha-i)} P_{\alpha}^{\frac{i}{\alpha}}(\phi_1) P_{\alpha}^{\frac{\alpha-i}{\alpha}}(\phi_2) + 2^{2\alpha-1} p^{\alpha} (1-p)^{\alpha} \sqrt{P_{\alpha}(\phi_1)} P_{\alpha}(\phi_2) \,. \tag{B1}$$

Alternatively, the sum can be performed to obtain

$$P_{\alpha}(\psi) \leq \sum_{\pm} \frac{1}{2} \left(p P_{\alpha}^{1/2\alpha}(\phi_1) \pm (1-p) P_{\alpha}^{1/2\alpha}(\phi_2) \right)^{2\alpha} + 2^{2\alpha-1} p^{\alpha} (1-p)^{\alpha} \sqrt{P_{\alpha}(\phi_1) P_{\alpha}(\phi_2)} \,. \tag{B2}$$

Proof. First of all, we can express $P_{\alpha}(\psi)$ as

$$P_{\alpha}(\psi) = \frac{1}{d_n} \sum_{P} \operatorname{tr}^{2\alpha}(I \otimes P\psi) + \operatorname{tr}^{2\alpha}(Z \otimes P\psi) + \operatorname{tr}^{2\alpha}(X \otimes P\psi) + \operatorname{tr}^{2\alpha}(Y \otimes P\psi)$$
(B3)

$$= \frac{1}{d_n} \sum_{P} \left(p \operatorname{tr}(P\phi_1) + (1-p) \operatorname{tr}(P\phi_2) \right)^{2\alpha} + \frac{1}{d_n} \sum_{P} \left(p \operatorname{tr}(P\phi_1) - (1-p) \operatorname{tr}(P\phi_2) \right)^{2\alpha}$$
(B4)

+
$$\frac{1}{d_n} \sum_{P} (\operatorname{Re}[2\sqrt{p(1-p)}\langle\phi_2|P|\phi_1\rangle])^{2\alpha} + \frac{1}{d_n} \sum_{P} (\operatorname{Im}[2\sqrt{p(1-p)}\langle\phi_2|P|\phi_1\rangle])^{2\alpha},$$
 (B5)

where the sum runs over the Pauli group on n-1 qubits and $d_n := 2^n$. Let us first elaborate on (B4) and bound it as

$$\sum_{P} \left(p \operatorname{tr}(P\phi_{1}) + (1-p) \operatorname{tr}(P\phi_{2}) \right)^{2\alpha} + \sum_{P} \left(p \operatorname{tr}(P\phi_{1}) - (1-p) \operatorname{tr}(P\phi_{2}) \right)^{2\alpha}$$
(B6)
= $2 \sum_{P} \sum_{i=0}^{\alpha} \binom{2\alpha}{2i} (p \operatorname{tr}(P\phi_{1}))^{2i} ((1-p) \operatorname{tr}(P\phi_{2}))^{2(\alpha-i)}$
 $\leq d_{n} \sum_{i=0}^{\alpha} \binom{2\alpha}{2i} p^{2i} (1-p)^{2(\alpha-i)} P_{\alpha}^{\frac{i}{\alpha}}(\phi_{1}) P_{\alpha}^{\frac{\alpha-i}{\alpha}}(\phi_{2}).$ (B7)

In Eq. (B7) we used Hölder's inequality $\sum_{l} x_{l}^{i} y_{l}^{\alpha-i} \leq ||x||_{\alpha/i} ||y||_{\alpha/(\alpha-i)}$ with $i/\alpha + (\alpha - i)/\alpha = 1$. Moreover, we used the definition of stabilizer purity $P_{\alpha}(\phi_{i}) = \frac{1}{d_{n-1}} \sum_{P} \operatorname{tr}^{2\alpha}(P\phi_{i})$. Let us now analyze the term in Eq. (B5):

$$\sum_{P} (\operatorname{Re}[2\sqrt{p(1-p)}\langle\phi_{2}|P|\phi_{1}\rangle])^{2\alpha} + \sum_{P} (\operatorname{Im}[2\sqrt{p(1-p)}\langle\phi_{2}|P|\phi_{1}\rangle])^{2\alpha}$$

$$\leq 2^{2\alpha}p^{\alpha}(1-p)^{\alpha}\sum_{P} \operatorname{tr}^{\alpha}(P\phi_{1}P\phi_{2})$$
(B8)

$$\leq 2^{2\alpha} p^{\alpha} (1-p)^{\alpha} \sqrt{\sum_{P} \operatorname{tr}^{2\alpha}(P\phi_1)} \sqrt{\sum_{P} \operatorname{tr}^{2\alpha}(P\phi_2)}$$
(B9)

$$\leq d_n 2^{2\alpha - 1} p^{\alpha} (1 - p)^{\alpha} \sqrt{P_{\alpha}(\phi_1) P_{\alpha}(\phi_2)}.$$
 (B10)

In Eq. (B8) we used that for any complex number x it holds that $\operatorname{Re}(x)^{2\alpha} + \operatorname{Im}(x)^{2\alpha} \leq |x|^{2\alpha}$. For the above chain of inequalities to hold, we need to show Eq. (B9), that is

$$\sum_{P} \operatorname{tr}^{\alpha}(P\phi_1 P\phi_2) \le \sqrt{\sum_{P} \operatorname{tr}^{2\alpha}(P\phi_1)} \sqrt{\sum_{P} \operatorname{tr}^{2\alpha}(P\phi_2)} \,. \tag{B11}$$

To show Eq. (B11), we can first expand each (n-1) qubit state ϕ_i in the Pauli basis as $\phi_i = \frac{1}{d_{n-1}} \sum_P \operatorname{tr}(P\phi_i)P$ for i = 1, 2, and write

$$\sum_{P} \operatorname{tr}^{\alpha}(P\phi_{1}P\phi_{2}) = \frac{1}{d_{n-1}^{\alpha}} \sum_{P} \sum_{P_{1},\dots,P_{\alpha}} \prod_{i=1}^{\alpha} [\operatorname{tr}(P_{i}\phi_{1})\operatorname{tr}(P_{i}\phi_{2})] \prod_{i=1}^{\alpha} \Omega(P,P_{i}), \quad (B12)$$

where, in Eq. (B12), we defined the following function over Pauli operators:

$$\Omega(P,Q) \coloneqq \frac{1}{d_{n-1}} \operatorname{tr} \left(P Q P^{\dagger} Q^{\dagger} \right) = (-1)^{\delta_{QP \neq PQ}}, \qquad (B13)$$

which captures the commutation relation between P and Q. Notice that the following identity holds

$$\Omega(P,Q_1)\Omega(P,Q_2) = \Omega(P,Q_1Q_2) = d_{n-1}^{-1} \operatorname{tr}(PQ_1Q_2PQ_2Q_1), \quad \forall P,Q_1,Q_2 \in \mathbb{P}_{n-1}.$$
(B14)

Starting from Eq. (B12), we can use multiple times the above identity and write:

$$\sum_{P} \prod_{i=1}^{\alpha} \Omega(P, P_i) = \sum_{P} \Omega(P, P_1 \cdots P_{\alpha}) = \frac{1}{d_{n-1}} \sum_{P} \operatorname{tr} \left(PP_1 \cdots P_{\alpha} P(P_1 \cdots P_{\alpha})^{\dagger} \right) = d_{n-1}^2 \delta_{P_1 P_2 \dots P_{\alpha} \propto 1} .$$
(B15)

The last equality follows from Haar integration and the fact that the Pauli group forms a 1-design. Plugging the identity in Eq. (B15) back to Eq. (B12), we arrive to

$$\sum_{P} \operatorname{tr}^{\alpha}(P\phi_{1}P\phi_{2}) = \frac{1}{d_{n-1}^{\alpha-2}} \sum_{P_{1},\dots,P_{\alpha}} \delta_{P_{1}P_{2}\dots P_{\alpha} \propto \mathbb{1}} \operatorname{tr}(P_{1}\phi_{1}) \operatorname{tr}(P_{1}\phi_{2}) \cdots \operatorname{tr}(P_{\alpha}\phi_{1}) \operatorname{tr}(P_{\alpha}\phi_{2})$$
(B16)

$$\leq \frac{1}{d_{n-1}^{\alpha-2}} \sqrt{\sum_{P_1,\dots,P_{\alpha}} \delta \prod_{i=1}^{\alpha} \operatorname{tr}^2(P_i \phi_1)} \sqrt{\sum_{P_1,\dots,P_{\alpha}} \delta \prod_{i=1}^{\alpha} \operatorname{tr}^2(P_i \phi_2)}$$
(B17)

$$= \sqrt{\frac{1}{d_{n-1}^{\alpha}} \sum_{P,P_1,\dots,P_{\alpha}} \prod_{i=1}^{\alpha} \operatorname{tr}^2(P_i \phi_1) \prod_{i=1}^{\alpha} \Omega(P,P_i)} \sqrt{\frac{1}{d_{n-1}^{\alpha}} \sum_{P,P_1,\dots,P_{\alpha}} \prod_{i=1}^{\alpha} \operatorname{tr}^2(P_i \phi_2) \prod_{i=1}^{\alpha} \Omega(P,P_i)}$$
(B18)

$$= \sqrt{\sum_{P} \operatorname{tr}^{\alpha}(P\phi_1 P\phi_1)} \sqrt{\sum_{P} \operatorname{tr}^{\alpha}(P\phi_2 P\phi_2)}$$
(B19)

$$= \sqrt{\sum_{P} \operatorname{tr}^{2\alpha}(P\phi_1)} \sqrt{\sum_{P} \operatorname{tr}^{2\alpha}(P\phi_2)}.$$
(B20)

In Eq. (B17), we used the Cauchy–Schwarz inequality for the $d_{n-1}^{2\alpha}$ -dimensional vectors with components (for i = 1, 2)

$$\delta_{P_1P_2\cdots P_\alpha \propto \mathbb{1}} \operatorname{tr}(P_1\phi_i) \times \operatorname{tr}(P_2\phi_i) \times \ldots \times \operatorname{tr}(P_\alpha\phi_i), \quad P_1,\ldots,P_\alpha \in \mathbb{P}.$$
(B21)

Moreover, we used a short notation for $\delta \equiv \delta_{P_1P_2\cdots P_{\alpha} \propto 1}$ and we used $\delta^2 = \delta$. In Eq. (B18) we applied the identity in Eq. (B15) again. Finally, in Eq. (B20) we used that $\operatorname{tr}(P\phi_i P\phi_i) = \operatorname{tr}^2(P\phi_i)$ for i = 1, 2. This proves Eq. (B11). Plugging everything back to Eq. (B4) and (B5), we have

$$P_{\alpha}(\psi) \leq \sum_{i=0}^{\alpha} \binom{2\alpha}{2i} p^{2i} (1-p)^{2(\alpha-i)} P_{\alpha}^{\frac{i}{\alpha}}(\phi_1) P_{\alpha}^{\frac{\alpha-i}{\alpha}}(\phi_2) + 2^{2\alpha-1} p^{\alpha} (1-p)^{\alpha} \sqrt{P_{\alpha}(\phi_1)} P_{\alpha}(\phi_2),$$
(B22)

which concludes the proof of Eq. (B1). To obtain Eq. (B2) it is sufficient to perform the sum.

Corollary 1. Consider $|\psi\rangle = \sqrt{p} |0\rangle \otimes |\phi_1\rangle + \sqrt{1-p} |1\rangle \otimes |\phi_2\rangle$ and $|\phi_1\rangle, |\phi_2\rangle \in \mathbb{C}^{2\otimes (n-1)}$. For any integer $\alpha \geq 2$, it follows $M_{\alpha}(\psi) > \min\{M_{\alpha}(\phi_1), M_{\alpha}(\phi_2)\}.$ (B23)

Proof. Starting from Eq. (B1) in Lemma 1, we can bound

$$P_k(\psi) \le \left(\sum_{i=0}^{\alpha} \binom{2\alpha}{2i} p^{2i} (1-p)^{2(\alpha-i)} + 2^{2\alpha-1} p^{\alpha} (1-p)^{\alpha}\right) \max\{P_{\alpha}(\phi_1), P_{\alpha}(\phi_2)\}.$$
(B24)

The coefficient reads

$$\sum_{i=0}^{\alpha} \binom{2\alpha}{2i} p^{2i} (1-p)^{2(\alpha-i)} + 2^{2\alpha-1} p^{\alpha} (1-p)^{\alpha} = \frac{1}{2} \left((2p-1)^{2\alpha} + p^{\alpha} (4-4p)^{\alpha} + 1 \right) \le 1.$$
 (B25)

Therefore, one has $P_k(\psi) \leq \max\{P_\alpha(\phi_1), P_\alpha(\phi_2)\}$ which implies the statement (see Definition 4).

Corollary 2. Consider $|\psi\rangle = \sqrt{p} |0\rangle \otimes |\phi_1\rangle + \sqrt{1-p} |1\rangle \otimes |\phi_2\rangle$ and $|\phi_1\rangle, |\phi_2\rangle \in \mathbb{C}^{2\otimes (n-1)}$. For any integer $\alpha \geq 2$, it holds that

$$P_{\alpha}(\psi) \le p P_{\alpha}(\phi_1) + (1-p) P_{\alpha}(\phi_2).$$
(B26)

Proof. Let us start from Eq. (B1) in Lemma 1, which we display below for convenience.

$$P_{\alpha}(\psi) \leq \sum_{i=0}^{\alpha} {\binom{2\alpha}{2i}} p^{2i} (1-p)^{2(\alpha-i)} P_{\alpha}^{\frac{i}{\alpha}}(\phi_1) P_{\alpha}^{\frac{\alpha-i}{\alpha}}(\phi_2) + 2^{2\alpha-1} p^{\alpha} (1-p)^{\alpha} \sqrt{P_{\alpha}(\phi_1)} P_{\alpha}(\phi_2) \,. \tag{B27}$$

First of all, note that $P_{\alpha}^{\frac{i}{\alpha}}(\phi_1)P_{\alpha}^{\frac{\alpha-i}{\alpha}}(\phi_2) \leq \frac{i}{\alpha}P_{\alpha}(\phi_1) + \frac{\alpha-i}{\alpha}P_{\alpha}(\phi_2)$ for the geometric-arithmetic mean inequality. We can perform the sum over the index *i* easily. Since the expression is symmetric in $(p, P_{\alpha}(\phi_1)) \leftrightarrow (1 - p, P_{\alpha}(\phi_2))$, without loss of generality, let us pose $p \ge 1/2$. We thus get:

$$P_{\alpha}(\psi) \leq \frac{1}{2}p[(2p-1)^{2\alpha-1}+1]P_{\alpha}(\phi_1) - \frac{1}{2}(1-p)[(2p-1)^{2\alpha-1}-1]P_{\alpha}(\phi_2) + 2^{2\alpha-1}p^{\alpha}(1-p)^{\alpha}\sqrt{P_{\alpha}(\phi_1)P_{\alpha}(\phi_2)}.$$
 (B28)

Then, to show the statement, it is sufficient to show that the right hand side of Eq. (B28) is upper bounded by $pP_{\alpha}(\phi_1) + (1-p)P_{\alpha}(\phi_2)$. Taking the difference between the two terms, we need to show

$$F(x) = -f(p)x^{2} + g(p)x - h(p) \le 0, \quad x \ge 0,$$
(B29)

for every $1/2 \le p \le 1$, where $f(p) = \frac{p}{2}(1 - (2p - 1)^{2\alpha - 1})$, $h(p) = \frac{1-p}{2}((2p - 1)^{2\alpha - 1} + 1)$ and $g(p) = 2^{2\alpha - 1}p^{\alpha}(1 - p)^{\alpha}$. Note that we defined $x \coloneqq \sqrt{\frac{P_{\alpha}(\phi_1)}{P_{\alpha}(\phi_2)}}$. For x = 0 and $x \to \infty$ it readily holds that $F(x) \le 0$ since $f(p), h(p) \ge 0$. Maximizing over x, we get F'(x) =

g(p) - 2f(p)x, and substituting $x(p) = \frac{g(p)}{2f(p)}$ into F(x), we get

$$F_{\alpha}(x(p)) = \frac{(1-p)}{2} \left(-[1+(2p-1)^{2\alpha-1}] + \frac{[4p(1-p)]^{2\alpha-1}}{1-(2p-1)^{2\alpha-1}} \right).$$
(B30)

We are left to show that $F_{\alpha}(x(p)) \leq 0$ for all $1/2 \leq p \leq 1$. Except for p = 1 for which $F_{\alpha}(x(1)) = 0$ and the result readily follows, this is equivalent to show that $F_{\alpha}(p) \leq 0$, where

$$\tilde{F}_{\alpha}(p) = -[1 - (2p - 1)^{2(2\alpha - 1)}] + [4p(1 - p)]^{2\alpha - 1}.$$
(B31)

Noting that \tilde{F}_{α} is a monotonically decreasing function in α (being the sum of two monotonically decreasing functions. Indeed note that $4p(1-p) \leq 1$, it is sufficient to show the statement for $\alpha = 2$. We have

$$\tilde{F}_2(p) = [4p(1-p)]^3 - [1 - (2p-1)^6] = -12p(1-p)(2p-1)^2 \le 0,$$
(B32)

which concludes the proof.

Appendix C: Proof of main results: Theorem 1, Theorem 2 and Theorem 3

Proof of Theorem 1. Let us recall the definition of pure-state magic monotone. It must hold that for every pair $(|\psi\rangle, \mathcal{E})$ where $\mathcal{E} \in \mathcal{S}$ obeying $\mathcal{E}(|\psi\rangle\langle\psi|) = |\phi\rangle\langle\phi|$, then $M_{\alpha}(\psi) \geq M_{\alpha}(\phi)$. To show pure-state monotonicity of stabilizer entropies with $\alpha > 2$, we make use of Corollary 1. First of all, any stabilizer protocol can be decomposed as a combination of elementary operations, i.e., (i) Clifford unitaries; (ii) partial trace; (iii) measurements in the computational basis; (iv) composition with ancillary qubits in $|0\rangle$; the above operations conditioned on (v) measurement outcomes and on (vi) classical randomness.

It is crucial to note that a deterministic pure-state stabilizer protocol only necessitates the initial and final states to be pure. However, given that a general stabilizer protocol consists of, let us say, N elementary operations, denoted as $\mathcal{E} = \mathcal{E}_N \circ \mathcal{E}_{N-1} \circ \cdots \circ \mathcal{E}_1$, there is no obligation for each step to maintain a pure state, nor must it be deterministic. Therefore, given a collection of (possibly mixed) states $\{(p_i, \rho_i)\}$, we define

$$M_{\alpha}^{\min}(\{(p_i, \rho_i\}) \coloneqq \inf_{(q_{ij}, \phi_{ij})} \left\{ \min_{ij} M_{\alpha}(\phi_{ij}) : p_i \rho_i = \sum_j q_{ij} \phi_{ij}, \forall i \right\},$$
(C1)

which is a possible extension of the stabilizer entropy M_{α} to a collection of states. To clarify the notation: given all the states ρ_i ; for each convex decomposition $\rho_i = \sum_j q_{ij} \phi_{ij}$, we choose the minimum $\min_{ij} M_\alpha(\phi_{ij})$ and then optimize over all the possible convex decompositions $\inf_{(q_{ij},\phi_{ij})}$. Therefore, after the optimization, we can describe any state of the

system by a collection of pure states $\{(p_i q_{ij}, \phi_{ij})\}$, which describes the optimal decomposition of $\{(p_i, \rho_i)\}$ with respect to M_{α} and, after the optimization, we have $M_{\alpha}^{\min}(\{(p_i, \rho_i)\}) = \min_{ij}\{M_{\alpha}(\phi_{ij})\}$.

The strategy of the proof is as follows: we first demonstrate that M_{α}^{\min} is a magic monotone according to Definition 1 in full generality. Then, by restricting deterministic stabilizer protocols, we establish that M_{α} is a pure-state magic monotone, given that (crucially) $M_{\alpha}(\phi) = M_{\alpha}^{\min}(\phi)$ for every pure state ϕ .

With this tool at hand, let us first show that, starting from a pure state ϕ , for every elementary stabilizer operation $\mathcal{E}(\cdot)$ it holds that

$$M_{\alpha}(\phi) \ge M_{\alpha}^{\min}(\mathcal{E}(\phi)). \tag{C2}$$

- Clifford unitaries and appending stabilizer ancillas. Eq. (C2) holds trivially because M_{α} is invariant under unitary Clifford transformations and it is additive, see Appendix A.
- Measurement in the computational basis. This follows directly from Corollary 1. Since we map the state $|\psi\rangle = \sqrt{p} |0\rangle \otimes |\phi_1\rangle + \sqrt{1-p} |1\rangle \otimes |\phi_2\rangle$ to the collection of pure states $\{(p, \phi_1), (1-p, \phi_2)\}$, for which $M_{\alpha}(\psi) \geq \min\{M_{\alpha}(\phi_1), M_{\alpha}(\phi_2)\} \geq M_{\alpha}^{\min}(\mathcal{E}(\psi))$. If the post measurement state is maintained after measurement, the same follows because $M_{\alpha}(\phi_{i+1}) = M_{\alpha}(|i\rangle\langle i| \otimes \phi_{i+1})$ for i = 0, 1.
- Partial trace and dephasing. Both dephasing and partial trace can be simulated by a forgetful measurement. We can therefore select $\{(p, \phi_1), (1 p, \phi_2)\}$ for partial trace or $\{(p, |0\rangle\langle 0| \otimes \phi_1), (1 p, |1\rangle\langle 1| \otimes \phi_2)\}$ as our convex mixtures to upper-bound the infimum. This allows us to always maintain a pure state collection $\{(p_i, \phi_i)\}$ for every operation.
- Conditional operations based on measurement outcomes or classical randomness. In general, any elementary conditional operation acts as

$$\phi \mapsto \{(p_i, \phi_i)\} \mapsto \{(p_i, \mathcal{E}_i(\phi_i))\},\tag{C3}$$

where the first map comes from the result of a computational basis measurement on ϕ or a classical randomness splitting, i.e. $\phi \mapsto \{(p, \phi), (1 - p, \phi)\}$. Indeed note that this is the most general form of classical randomness: it includes both conditional preparation of stabilizer states and conditional operations. We have already seen that $M_{\alpha}(\phi) \ge M_{\alpha}^{\min}(\{(p_i, \phi_i)\})$ for a measurement. While, trivially, $M_{\alpha}(\phi) \ge M_{\alpha}(\phi)$ for classical randomness. Moreover, for every $i, M_{\alpha}(\phi_i) \ge M_{\alpha}^{\min}(\mathcal{E}_i(\phi_i))$. Hence:

$$M_{\alpha}(\phi) \ge M_{\alpha}^{\min}(\{(p_i, \phi_i)\}) = \min_i M_{\alpha}(\phi_i) \ge \min_i M_{\alpha}^{\min}(\mathcal{E}_i(\phi_i)) \ge M_{\alpha}^{(\min)}(\{p_i, \mathcal{E}_i(\phi_i)\}).$$
(C4)

The second equality follows by definition, while the last inequality follows because we have chosen a particular decomposition, not necessarily the infimum. The result just follows for conditional operations.

Now, we generalize the result to collections of mixed states and show that:

$$M_{\alpha}^{\min}(\{(p_i, \rho_i)\}) \ge M_{\alpha}^{\min}(\mathcal{E}(\{(p_i, \rho_i)\})).$$
(C5)

Let $\{(q_{ij}, \phi_{ij})\}$ be the optimal decomposition of $\{(p_i, \rho_i)\}$ into an collection of pure states. Consider an elementary stabilizer operation \mathcal{E} . We have already shown that, for every state ϕ_{ij} of such a collection, it holds that

$$M_{\alpha}(\phi_{ij}) \ge M_{\alpha}^{\min}(\mathcal{E}(\phi_{ij})).$$
(C6)

One can also verify that $\{(q_{ij}, \mathcal{E}(\phi_{ij}))\}$ is a potentially mixed state collection of $\mathcal{E}(\{(p_i, \rho_i)\})$. It therefore upper bounds the optimal value. We therefore have:

$$M_{\alpha}^{\min}(\mathcal{E}(\{(p_i, \rho_i)\})) \le \min_{ij} M_{\alpha}^{\min}(\mathcal{E}(\phi_{ij})) \le \min_{ij} M_{\alpha}(\phi_{ij}) = M_{\alpha}^{\min}(\{(p_i, \rho_i)\}).$$
(C7)

The first inequality arises since we selected a particular decomposition, i.e., $\{(q_{ij}, \mathcal{E}(\phi_{ij}))\}$, which is not necessarily the infimum; the last equality follows from $\{(q_{ij}, \phi_{ij})\}$ being a optimal decomposition of $\{(p_i, \rho_i)\}$. Since Eq. (C5) holds for all elementary stabilizer operations, it also holds for all composite stabilizer operations. To conclude the proof, consider a deterministic stabilizer protocol \mathcal{E} that, starting from $|\psi\rangle\langle\psi|$ returns a pure state $|\phi\rangle\langle\phi|$. We therefore have $M_{\alpha}(\psi) \geq M_{\alpha}^{\min}(\phi) = M_{\alpha}(\phi)$, which concludes the proof. Moreover, if the stabilizer protocol \mathcal{E} is non-deterministic and yields a collection of pure-states $\mathcal{E}(\psi) = \{(p_i, \phi_i)\}$, we have $M_{\alpha}(\psi) \geq \min_i M_{\alpha}(\phi_i)$.

Proof of Theorem 3. Let us first recall the definition of extended stabilizer purity \hat{P}_{α} on a collection of (possibly mixed) states $\{(p_i, \rho_i)\}$:

$$\widehat{P}_{\alpha}(\{(p_i,\rho_i\}) \coloneqq \sup_{(q_{ij},\phi_{ij})} \left\{ \sum_{ij} q_{ij} P_{\alpha}(\phi_{ij}) : p_i \rho_i = \sum_j q_{ij} \phi_{ij}, \forall i \right\}.$$
(C8)

That is, given the collection of states, we first compute the average stabilizer purity P_{α} defined on pure states in Definition 4 with respect to a convex decomposition of $p_i \rho_i$ and then take the superior over all the possible convex decompositions. We will use the same strategy used for the proof of Theorem 1: we first show that for every pure state ϕ and elementary stabilizer operation \mathcal{E} , the following holds:

$$P_{\alpha}(\phi) \le \widehat{P}_{\alpha}(\mathcal{E}(\phi)) \,. \tag{C9}$$

- Clifford unitaries and appending stabilizer ancillas. Eq. (C9) follows because P_{α} is invariant under Clifford unitaries and it is multiplicative, see Appendix A.
- Measurement in the computational basis. This follows directly from Corollary 2. Since we map the state $|\phi\rangle = \sqrt{p} |0\rangle \otimes |\phi_1\rangle + \sqrt{1-p} |1\rangle \otimes |\phi_2\rangle$ to the collection of pure states $\{(p, \phi_1), (1-p, \phi_2)\}$, for which $P_{\alpha}(\phi) \leq pP_{\alpha}(\phi_1) + (1-p)P_{\alpha}(\phi_2)$ holds. If the post measurement state is maintained after measurement, the same follows since $P_{\alpha}(\phi_{i+1}) = P_{\alpha}(|i\rangle\langle i| \otimes \phi_{i+1})$ for i = 0, 1.
- Partial trace and dephasing. Both dephasing and partial trace can be simulated by a forgetful measuremnt. We can therefore select $\{(p, \phi_1), (1 p, \phi_2)\}$ for partial trace or $\{(p, |0\rangle\langle 0| \otimes \phi_1), (1 p, |1\rangle\langle 1| \otimes \phi_2)\}$ as our convex mixture to lower-bound the superior. This allows us to always maintain a pure state collection $\{(p_i, \phi_i)\}$ for every operation.
- Conditional operations based on measurement outcomes or classical randomness. In general, any elementary conditional operation acts as

$$\phi \mapsto \{(p_i, \phi_i)\} \mapsto \{(p_i, \mathcal{E}_i(\phi_i))\},\tag{C10}$$

where the first map comes from the result of a computational basis measurement on ϕ or a classical randomness splitting, i.e. $\phi \mapsto \{(p, \phi), (1 - p, \phi)\}$. Indeed note that this is the most general form of classical randomness: it includes both conditional preparation of stabilizer states and conditional operations. We have already seen that $P_{\alpha}(\phi) \leq \hat{P}_{\alpha}\{(p_i, \phi_i)\}) = \sum_i p_i P_{\alpha}(\phi_i)$ for a measurement. While, somewhat trivially, it holds that $P_{\alpha}(\phi) \leq$ $pP_{\alpha}(\phi)+(1-p)P_{\alpha}(\phi)$ for classical randomness. Moreover we have also shown that, for every i, $P_{\alpha}(\phi_i) \leq \hat{P}_{\alpha}(\mathcal{E}_i(\phi_i))$. Hence:

$$P_{\alpha}(\phi) \leq \widehat{P}_{\alpha}(\{(p_i, \phi_i)\}) = \sum_{i} p_i P_{\alpha}(\phi_i) \leq \sum_{i} p_i \widehat{P}_{\alpha}(\mathcal{E}_i(\phi_i)) \leq \widehat{P}_{\alpha}(\{(p_i, \mathcal{E}_i(\phi_i))\}).$$
(C11)

The first equality follows by definition: note that for a collection of pure states, no optimization is needed. The last inequality follows because we have chosen a particular decomposition, not necessarily the superior. The result just follows for conditional operations.

Now, we generalize the above result to collections of mixed states and show that for every collection $\{(p_i, \rho_i)\}$ and elementary stabilizer operation \mathcal{E} , it holds that:

$$\widehat{P}_{\alpha}(\{(p_i, \rho_i)\}) \le \widehat{P}_{\alpha}(\mathcal{E}(\{(p_i, \rho_i)\})).$$
(C12)

Let $\{(q_{ij}, \phi_{ij})\}$ be the optimal decomposition of $\{(p_i, \rho_i)\}$ into an collection of pure states. Consider an elementary stabilizer protocol \mathcal{E} . We have already shown that, for every pure state ϕ_{ij} in such a collection, it holds that

$$\widehat{P}_{\alpha}(\mathcal{E}(\phi_{ij})) \ge P_{\alpha}(\phi_{ij}).$$
(C13)

One can also verify that $\{(q_{ij}, \mathcal{E}(\phi_{ij}))\}$ is a potentially mixed state collection of $\mathcal{E}(\{(p_i, \rho_i)\})$. Therefore, we can write the following chain of inequalities:

$$\widehat{P}_{\alpha}(\mathcal{E}(\{(p_i,\rho_i)\})) \ge \sum_{ij} q_{ij} \widehat{P}_{\alpha}(\mathcal{E}(\phi_{ij})) \ge \sum_{ij} q_{ij} P_{\alpha}(\phi_{ij}) = \widehat{P}_{\alpha}(\{(p_i,\rho_i)\}).$$
(C14)

where the first inequality arises since we selected a particular decomposition, i.e., $\{(q_{ij}, \mathcal{E}(\phi_{ij}))\}$, which is not necessarily the superior. The last equality follows because $\{(q_{ij}, \mathcal{E}(\phi_{ij}))\}$ is an optimal decomposition. Since this holds for all elementary stabilizer operations, Eq. (C12) holds every stabilizer protocol.

Lastly, the monotonicity of the extended stabilizer entropy and linear stabilizer entropy readily follows from Eq. (C12). What's more, the above theorem also shows that linear stabilizer entropies are strong monotones not only restricted to pure-states.

Proof of Theorem 2. The proof of Theorem 2 readily follows from Theorem 3. To see this, let us recall the definition of pure-state strong monotone. Starting from a pure state ϕ , for any non-deterministic stabilizer protocol \mathcal{E} ending in a collection $\mathcal{E} : \phi \mapsto \{(p_i, \phi_i)\}$ of pure states ϕ_i , it must hold that $\mathcal{M}(\phi) \geq \sum_i p_i \mathcal{M}(\phi_i)$. Given the definition of extended linear stabilizer entropy, we have shown that

$$M_{\alpha}^{\mathrm{lin}}(\phi) = \widehat{M}_{\alpha}^{\mathrm{lin}}(\phi) \ge \widehat{M}_{\alpha}^{\mathrm{lin}}(\{(p_i, \phi_i)\}) = \sum_i p_i M_{\alpha}^{\mathrm{lin}}(\phi_i) \,. \tag{C15}$$

The first equality follows because the two definitions adhere for pure states. The inequality follows because $\widehat{M}_{\alpha}^{\text{lin}}$ is a magic monotone thanks to Theorem 3, and the last equality follows by definition (see Definition 5). Note indeed that there is no optimization over convex decompositions being ϕ_i pure states. This concludes the proof.