

Optimizing the diffusion coefficient of overdamped Langevin dynamics

T. Lelièvre¹, G. Pavliotis², G. Robin³, R. Santet¹ and G. Stoltz¹

1: CERMICS, École des Ponts, Marne-la-Vallée, France & MATHERIALS project-team, Inria Paris, France

2: Department of Mathematics, Imperial College, London, United-Kingdom

3: CNRS & Laboratoire de Mathématiques et Modélisation d'Évry, Évry, France

May 2, 2024

Abstract

Overdamped Langevin dynamics are reversible stochastic differential equations which are commonly used to sample probability measures in high-dimensional spaces, such as the ones appearing in computational statistical physics and Bayesian inference. By varying the diffusion coefficient, there are in fact infinitely many overdamped Langevin dynamics which are reversible with respect to the target probability measure at hand. This suggests to optimize the diffusion coefficient in order to increase the convergence rate of the dynamics, as measured by the spectral gap of the generator associated with the stochastic differential equation. We analytically study this problem here, obtaining in particular necessary conditions on the optimal diffusion coefficient. We also derive an explicit expression of the optimal diffusion in some appropriate homogenized limit. Numerical results, both relying on discretizations of the spectral gap problem and Monte Carlo simulations of the stochastic dynamics, demonstrate the increased quality of the sampling arising from an appropriate choice of the diffusion coefficient.

1 Introduction

Predicting properties of materials and macroscopic physical systems in the framework of statistical physics [8], using for instance molecular dynamics [30, 68, 44, 49, 5] and obtaining the distribution of parameter values in Bayesian inference [60], both require the sampling of probability measures in high-dimensional spaces. Methods of choice to sample such probability measures rely on stochastic dynamics, in particular Markov Chain Monte Carlo (MCMC) methods. The convergence of these methods may however be quite slow because the target measure is typically concentrated on a few high probability modes separated by low probability regions, or because the Hessian of the corresponding potential, see (1), is badly conditioned. Various algorithms have been proposed over the years to improve the sampling of such probability measures, for instance using importance sampling strategies, and/or interacting replicas; see for instance the extensive review [41] in the context of molecular dynamics.

Sampling Boltzmann–Gibbs measures. We focus in this work on overdamped Langevin dynamics, a popular choice in molecular dynamics, computational statistics [62], and also machine learning (see for instance [20]). This dynamics is ergodic (and, in fact, reversible) with respect to the Boltzmann–Gibbs distribution, which is the probability measure with density

$$\mu(q) = Z^{-1} e^{-V(q)}, \quad Z = \int_{\mathcal{Q}} e^{-V} < +\infty, \quad (1)$$

where $V \in \mathcal{C}^\infty(\mathcal{Q})$. In molecular dynamics, the measure μ is the distribution sampled by the possible states of a system in the canonical thermodynamic ensemble. In that case, the system is

at thermal equilibrium with a heat bath of fixed temperature T , which is taken into account by considering βV instead of V in (1) with $\beta = 1/(k_B T)$, where k_B is the Boltzmann constant (in this work, we simply set $\beta = 1$). To simplify the presentation, and because periodic boundary conditions are commonly used for molecular dynamics simulations, we restrict ourselves to the case when the configuration space \mathcal{Q} is the d -dimensional torus \mathbb{T}^d (with $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ the one dimensional torus). Many arguments in our work can be extended, at least formally, to systems defined in unbounded configuration spaces.

The overdamped Langevin dynamics is given by the following stochastic differential equation (SDE):

$$dq_t = -\nabla V(q_t) dt + \sqrt{2} dW_t, \quad (2)$$

where $(W_t)_{t \geq 0}$ is a standard d -dimensional Brownian motion. This dynamics enjoys many nice properties. In particular, it is reversible with respect to the Gibbs measure (1), and, under appropriate s on the potential energy function V (see for instance [7]), its marginal law in time converges exponentially fast to the target distribution (1) in various norms and distances, e.g. total variation, relative entropy (Kullback–Leibler), $L^2(\mu)$, or weighted L^∞ norms. For example, when the overdamped Langevin dynamics in \mathbb{R}^d is considered, it is sufficient for the potential V to be smooth and convex at infinity to ensure the existence of Poincaré and logarithmic Sobolev inequalities for the Gibbs measure which, in turn, leads to exponentially fast convergence to the target distribution in $L^2(\mu)$ and in relative entropy, respectively. This class of potentials includes many potentials of interest, for example multistable potentials. Naturally, in the nonconvex case, the convergence rate typically scales very badly with respect to the dimension d (and the temperature). In this work, we will consider convergence in $L^2(\mu)$, *i.e.* the χ^2 -divergence. This is the relevant setting *e.g.* for obtaining error estimates on trajectory averages, relying on the Central Limit Theorem.

It is important to note that the constant in the Poincaré and in the logarithmic Sobolev inequalities is a property of the Gibbs measure. In particular, the rate of convergence to the target measure for the overdamped Langevin dynamics (2) depends only on the potential V , and convergence can be very slow when entropic or energetic barriers are present. It is therefore natural to consider alternatives to the overdamped Langevin dynamics that include “hyperparameters” that can be tuned to improve the rate of convergence to equilibrium and reduce the asymptotic variance.

There are in fact infinitely many overdamped Langevin dynamics which admit (1) as invariant probability measure; see [54, Section 4.6]. A comprehensive list of stochastic dynamics that can be used in order to sample from a Gibbs measure can be found in [25]. For instance, dynamics of the form

$$dq_t = (-\nabla V(q_t) + \gamma(q_t)) dt + \sqrt{2} dW_t,$$

where the vector field $\gamma : \mathbb{T}^d \rightarrow \mathbb{R}^d$ is smooth and such that

$$\operatorname{div}(\gamma\mu) = 0,$$

leave invariant the target probability measure with density (1); and is ergodic, but not reversible, with respect to this probability measure under certain s on V, γ . The hope is that the added drift term γ that renders the dynamics nonreversible, can accelerate convergence, measured in the $L^2(\mu)$ sense, and also reduce the asymptotic variance. This approach was extensively studied in the past decade [39, 40, 46, 58, 59, 24, 26]. For example, for Gaussian targets, the optimal nonreversible perturbation γ can be obtained in an algorithmic manner [46]. The presence of the divergence-drift term γ can lead to computational difficulties, since the resulting dynamics can be stiff. Appropriate numerical methods need to be developed to address this issue [24, 26]. In fact, the discretized nonreversible dynamics can perform worse than the reversible dynamics if the stiffness issue is not addressed adequately [71, 51].

Optimizing reversible dynamics for better sampling. Another alternative, which is the focus of this work, is to stay within the class of reversible dynamics, but with a space-dependent

diffusion matrix. For a given diffusion matrix $\mathcal{D} \in \mathcal{C}^1(\mathbb{T}^d, \mathcal{S}_d^+)$ (with \mathcal{S}_d^+ the set of real, symmetric, positive matrices of size $d \times d$), the associated overdamped Langevin dynamics reads

$$dq_t = (-\mathcal{D}(q_t)\nabla V(q_t) + \operatorname{div} \mathcal{D}(q_t)) dt + \sqrt{2} \mathcal{D}^{1/2}(q_t) dW_t, \quad (3)$$

where $\mathcal{D}^{1/2}$ is defined by functional calculus, and $\operatorname{div} \mathcal{D}$ is the vector whose i -th component is the divergence of the i -th column of the matrix $\mathcal{D} = [\mathcal{D}_1, \dots, \mathcal{D}_d]$, *i.e.* $\operatorname{div} \mathcal{D} = (\operatorname{div} \mathcal{D}_1, \dots, \operatorname{div} \mathcal{D}_d)^\top$. Modulating the diffusion to improve the sampling was explored to some extent both in the computational statistics literature [61] and in molecular dynamics [59, 1]; and also for simulated annealing [28].

The goal of this paper is to optimize the diffusion matrix in order to accelerate convergence. Intuitively, it seems relevant to accelerate the diffusion in regions of low probability under the target measure (equivalently, regions of high values of the energy function V), in order to more efficiently and rapidly find another mode, and slow down the diffusion in regions of high probability since these are the zones where sampling should be favored. Clearly, in order for this optimization problem to make sense, appropriate constraints on the magnitude of the diffusion matrix need to be introduced. These constraints on the diffusion matrix and their effects on the solution of the associated constrained optimization problem are discussed in detail later on.

The $L^2(\mu)$ convergence rate of the overdamped Langevin diffusion (3) is given by the spectral gap of the infinitesimal generator. The latter operator acts on test functions φ as

$$\mathcal{L}_{\mathcal{D}}\varphi = (-\mathcal{D}\nabla V + \operatorname{div}(\mathcal{D}))^\top \nabla \varphi + \mathcal{D} : \nabla^2 \varphi. \quad (4)$$

It is standard to show (see, *e.g.* [46, 7, 49]) that, $\mathcal{L}_{\mathcal{D}}$ admits a positive spectral gap $\Lambda(\mathcal{D})$ if and only if, for any initial condition $f(0) = \mu_0/\mu \in L^2(\mu)$, and denoting by $f(t)\mu$ the probability density function of the process q_t at time t ,

$$\forall t \geq 0, \|f(t) - 1\|_{L^2(\mu)} \leq e^{-\Lambda(\mathcal{D})t} \|f(0) - 1\|_{L^2(\mu)}, \quad (5)$$

where $\|\cdot\|_{L^2(\mu)}$ is the norm on $L^2(\mu)$, see Section 2.1 for precise definitions. For a fixed target probability density μ (*i.e.* a fixed potential energy function V), the generator $\mathcal{L}_{\mathcal{D}}$, and thus the spectral gap $\Lambda(\mathcal{D})$, are parameterized by the diffusion matrix \mathcal{D} .

The aim of this work is to compute, explicitly or numerically, the optimal diffusion matrix leading to the largest spectral gap, and thus to the largest convergence rate. For reversible Markov chains on discrete spaces, this question was explored in [16, 15]. A subtle issue in this endeavour is the normalization of the diffusion matrix \mathcal{D} . Indeed, the convergence rate in (5) can trivially be increased by a factor $\alpha \geq 1$ upon multiplying \mathcal{D} by α . In fact, it seems advantageous to let α go to infinity. The catch is of course that one should compare dynamics which are defined on similar timescales. From a practical point of view, one way of setting a timescale is to consider discretizations of overdamped Langevin dynamics. Larger diffusion matrices require smaller values of the time step. The issue of normalization is discussed more thoroughly in Section 2.2.

The optimization of the diffusion matrix is related to a body of literature on accelerating the convergence of Langevin-like dynamics through particular choices for the diffusion matrix \mathcal{D} . This is very much related to the optimization of the damping coefficient [19] or of the mass matrix in the Langevin Monte Carlo method, a.k.a. as preconditioning, see [18, 10, 32, 45, 34, 21]. One example of such methods is the Riemannian manifold Langevin Monte Carlo method, introduced in [32], and which reduces to (3) in the so-called overdamped limit. There, for strictly convex potentials, the diffusion matrix \mathcal{D} is chosen as the inverse of the Fisher–Rao metric tensor associated with the target measure, namely the inverse of the Hessian of V . From a functional analysis point of view, this can be understood, for a problem set on \mathbb{R}^d , from the Brascamb-Lieb inequality [37, 64]:

$$\operatorname{Var}_\mu(h) \leq \int_{\mathbb{R}^d} \langle \nabla h, (\operatorname{Hess}(V))^{-1} \nabla h \rangle \mu(q) dq,$$

for an arbitrary observable $h \in \mathcal{C}^1$ with $\operatorname{Var}_\mu(h) < +\infty$. In particular, when the diffusion matrix is chosen to be the inverse of the Hessian of V , this leads to a spectral gap for the dynamics (3)

that is independent of V and of the number of dimensions and is thus “optimal” in this sense. We emphasize that, although relevant, this unconstrained optimization problem for the diffusion matrix is quite different from the one we consider in this paper: we optimize the spectral gap for the dynamics on the torus, with no convexity s on V and with constraints on the magnitude of \mathcal{D} .

At the time we completed this work, we became aware of the preprint [23] where the authors consider a very similar problem, that they solve using different techniques, namely Stein Kernels and moment measures. More precisely, by using an appropriate transportation map, they transform the nonconvex log-probability to a convex one, for which the Brascamp–Lieb inequality yields the optimal diffusion, for a well-chosen normalizing constraint. We intend to further explore the links between this recent preprint and ours in future works.

Let us also mention results obtained via the theory of partial differential equations for eigenvalue problems associated with operators of the form $-\operatorname{div}(\mathcal{D}\nabla\cdot)$, see [38, Chapter 10]. The main difference with such results and ours is that we consider the Laplace operators on the weighted space $L^2(\mu)$ as we are interested in sampling the nontrivial probability measure μ .

Main contributions and extensions. The main contributions of this work are the following.

- First, we formulate the optimization of the convergence rate of the Langevin dynamics (3) with respect to \mathcal{D} as a convex optimization program, for which the well-posedness is guaranteed by a simple adaptation of standard results in the literature [38]. We also discuss some important properties of the resulting optimal diffusion matrix, such as its formal characterization using the Euler–Lagrange equation as well as its positivity. Note that, for Markov chains, it was already observed in [16] that maximizing the spectral gap can be formulated as a convex optimization problem.
- Second, we propose, in low dimensional scenarios, a numerical procedure combining a finite element parametrization and an optimization algorithm, to compute the optimal diffusion matrix in practice. The numerical procedure is illustrated on simple one-dimensional examples. All the methods and experiments are provided in open source Python and Julia codes, available at https://github.com/rsantet/Optimal_Overdamped_Langevin_Diffusion_Python and https://github.com/rsantet/Optimal_Overdamped_Langevin_Diffusion_Julia.
- Our third main contribution is to study the behavior of the optimal diffusion matrix in the homogenized limit where the target measure is associated with a highly oscillating potential. In a one-dimensional setting, we show that the optimal diffusion matrix has an analytical expression, proportional to the inverse of the target density, which is in accordance with various previous heuristics (as in [61]). This analytical solution can be used as an initialization guess for the optimization algorithm mentioned above, or as a proxy for the optimal diffusion matrix which does not require costly convex optimization procedures.
- Our last main contribution is to propose a sampling algorithm based on a simple Random Walk Metropolis Hastings algorithm with a proposal variance depending on the current state, which takes advantage of the precomputed optimal diffusion matrix. We show that the associated Markov chain converges to the Langevin dynamics with optimal diffusion matrix in the diffusive limit. We also present some numerical experiments illustrating the behavior of the complete procedure on simple examples, and highlighting in particular that the dynamics is less metastable when using the optimal diffusion matrix or its homogenized approximation. We show that the homogenization approximation yields similar results as the optimal diffusion matrix, which makes it a good choice in practice since it has an analytical expression.

We believe that this work calls for various perspectives and extensions, in particular:

- A key issue remains to find a relevant normalization constraint e.g. based on a numerical criterion, as discussed in Remark 3;

- In various contexts, we restrict ourselves to isotropic diffusion matrices or even to the one-dimensional setting (in particular for numerical simulations). Considering a larger class of positive definite symmetric matrices may be important to tackle anisotropic targets;
- In order to apply the methodology to actual systems of interest, one needs to adapt our numerical approach to higher dimensional scenarios, even in the case of a scalar diffusion coefficient, i.e. when the diffusion matrix is a multiple of the identity. It is indeed impossible to numerically optimize a diffusion matrix which genuinely depends on the full configuration q using discretization techniques such as finite elements, because of the curse of dimensionality of the corresponding approximation space. Apart from resorting to parsimonious functional representations of the diffusion matrix (which usually presuppose some form of smoothness), it seems appropriate to look for diffusion matrices parameterized by a low dimensional function $\xi(q)$, a so-called reaction coordinate in molecular dynamics [47], which summarizes some important information on the system. In this context, the potential energy function is replaced by the free energy associated with ξ , and the optimization problem can again be reformulated in a low dimensional setting where finite element methods can be used.
- In this paper we consider the problem of maximizing the spectral gap. The self-adjointness of the generator implies that maximizing the spectral gap is typically equivalent to minimizing the asymptotic variance, uniformly over the space of $L^2(\mu)$ observables [6]. We will explore the combination of spectral gap maximization and asymptotic variance minimization in a future work. In particular, our aim will be to minimize an appropriate cost functional that measures the true cost of the actual computation, by using the approach developed in [33].
- The objective in this paper is to optimize the convergence in χ^2 -divergence (see (5)). A natural question would be to consider convergence in other distances, such as relative entropy with respect to the target distribution (which amounts to optimizing Logarithmic Sobolev Inequality constants).

Outline of the work. We formally define the maximization of the spectral gap in Section 2 and make precise the normalization of the diffusion matrix. We then study the well posedness of the maximization problem in Section 3, and characterize the optimal diffusion matrix. We next present in Section 4 how to numerically solve the problem, using in particular a finite element method to discretize the diffusion matrix. As an alternative to a full-scale numerical simulations, or in order to start the optimization procedure with a relevant initial guess, we consider in Section 5 the optimal diffusion matrix arising in the homogenization limit where the given probability density is periodically replicated with a decreasing spatial period. We finally demonstrate in Section 6 with Monte Carlo simulations that an optimized diffusion matrix is beneficial for the convergence of the dynamics. In the appendices we present the proofs of various technical results.

2 Formulation of the optimization problem

We present in this section the problem of interest, namely the maximization of the spectral gap of the generator of the Langevin dynamics (3) with invariant measure μ given by (1). We start by precisely formulating the optimization problem in Section 2.1. Then, in Section 2.2, we discuss normalization constraints which are required for the well-posedness of the problem. For simplicity, we restrict ourselves to the case when the diffusion process under consideration takes as input a configuration in the d -dimensional torus \mathbb{T}^d . Various results can easily be adapted to more general domains.

2.1 Maximizing the spectral gap of Langevin dynamics

As discussed in the introduction around (5), the rate of convergence of the dynamics (3) towards the invariant measure μ is governed by the spectral gap of the operator $\mathcal{L}_{\mathcal{D}}$ (see *e.g.* [7, Chapter 4]).

In order to precisely define this quantity in our context, we work on the Hilbert space $L^2(\mu)$. For a given diffusion matrix $\mathcal{D} : \mathbb{T}^d \rightarrow \mathcal{S}_d^+$, a simple computation shows that the generator $\mathcal{L}_{\mathcal{D}}$, considered as an operator on $L^2(\mu)$, can be written as

$$\mathcal{L}_{\mathcal{D}} = -\nabla^* \mathcal{D} \nabla = - \sum_{i,j=1}^d \partial_{q_j}^* \mathcal{D}_{i,j} \partial_{q_i}, \quad (6)$$

where A^* is the $L^2(\mu)$ -adjoint of a closed operator A . In particular, $\partial_{q_i}^* = -\partial_{q_i} + \partial_{q_i} V$. The quadratic form associated with $\mathcal{L}_{\mathcal{D}}$ is easily obtained from the expression (6) of the generator: formally, for any $u : \mathbb{T}^d \rightarrow \mathbb{R}$,

$$\langle u, \mathcal{L}_{\mathcal{D}} u \rangle_{L^2(\mu)} = - \int_{\mathbb{T}^d} \nabla u(q)^\top \mathcal{D}(q) \nabla u(q) \mu(q) dq.$$

The right-hand side of the above equality is nonpositive, but possibly infinite at this level of generality. It is finite for instance when $\mathcal{D} \in L^\infty(\mathbb{T}^d, \mathcal{S}_d^+)$ and $u \in H^1(\mu)$, the subspace of $L^2(\mu)$ composed of functions in $L^2(\mu)$ whose (distributional) partial derivatives with respect to q_i also belong to $L^2(\mu)$. We restrict for simplicity to this setting in the sequel, and consider in particular diffusion matrices with bounded coefficients.

Since 0 is an eigenvalue of $\mathcal{L}_{\mathcal{D}}$ whose associated eigenvectors are constant functions, we introduce the following subspace to define the spectral gap:

$$H_0^1(\mu) = \left\{ u \in H^1(\mu) \mid \int_{\mathbb{T}^d} u(q) \mu(q) dq = 0 \right\}.$$

We also define $H_0^1(\mathbb{T}^d)$ as the space $H_0^1(\mathbf{1})$. The spectral gap is finally defined as

$$\Lambda(\mathcal{D}) = \inf_{u \in H_0^1(\mu) \setminus \{0\}} \frac{\int_{\mathbb{T}^d} \nabla u(q)^\top \mathcal{D}(q) \nabla u(q) \mu(q) dq}{\int_{\mathbb{T}^d} u(q)^2 \mu(q) dq}. \quad (7)$$

For a bounded positive definite diffusion matrix \mathcal{D} satisfying $\mathcal{D}(q) \geq c \text{Id}_d$ almost everywhere for some $c > 0$ (the inequality being understood in the sense of symmetric matrices), standard arguments from calculus of variations show that the infimum is attained for an eigenvector $u_{\mathcal{D}}$ associated with the second eigenvalue of $-\mathcal{L}_{\mathcal{D}}$ (ranked by increasing values), *i.e.* with the first nonzero eigenvalue of $-\mathcal{L}_{\mathcal{D}}$; and that $\Lambda(\mathcal{D})$ is in fact the first nonzero eigenvalue of $-\mathcal{L}_{\mathcal{D}}$ (see for instance [38, Chapter 1]).

Remark 1 (Degenerate diffusion matrices). *If there exists $q \in \mathbb{T}^d$ such that $\mathcal{D}(q)$ is rank deficient (i.e. $\xi^\top \mathcal{D}(q) \xi = 0$ for some $\xi \in \mathbb{R}^d \setminus \{0\}$), the domain of the quadratic form associated with $\mathcal{L}_{\mathcal{D}}$ can in fact be larger than $H_0^1(\mu)$; see [72]. In this case, $\Lambda(\mathcal{D})$ is in general no longer equal to the spectral gap of the operator $\mathcal{L}_{\mathcal{D}}$, but is larger than the spectral gap (and hence provides a too optimistic convergence rate) since the infimum is taken on a set of functions not sufficiently large. See [38, Remark 1.2.3] for a discussion on sufficient conditions for the operator $\mathcal{L}_{\mathcal{D}}$ to have a discrete spectrum even in the presence of degeneracies of the diffusion matrix.*

Remark 2 (Unbounded diffusion matrices). *We emphasize the fact that we will restrict ourselves to bounded diffusion matrices; otherwise the upper semicontinuity of the mapping $\mathcal{D} \mapsto \Lambda(\mathcal{D})$ is not clear (see Lemma 3). The latter property is crucial to ensure that the maximization problem (8) below is well posed.*

We propose to maximize the spectral gap of the overdamped Langevin dynamics with respect to the diffusion matrix \mathcal{D} . This corresponds here to solving the following optimization problem:

$$\Lambda^* = \sup_{\mathcal{D} \in \mathfrak{D}} \Lambda(\mathcal{D}), \quad (8)$$

where \mathfrak{D} is a subset of bounded measurable functions from \mathbb{T}^d to \mathcal{S}_d^+ . The choices we consider in the sequel for \mathfrak{D} take into account that the diffusion needs to be normalized in some way since, for any $\mathcal{D} : \mathbb{T}^d \rightarrow \mathcal{S}_d^+$,

$$\forall t \geq 0, \quad \Lambda(t\mathcal{D}) = t\Lambda(\mathcal{D}), \quad (9)$$

so that a maximization over a set containing a line $\mathbb{R}\mathcal{D}$ with $\Lambda(\mathcal{D}) > 0$ would lead to $\Lambda^* = +\infty$.

Remark 3 (A numerical motivation for normalizing the diffusion matrix). *The need for a normalization can also be motivated by numerical considerations. For instance, an Euler–Maruyama discretization of (3) with a time step $\Delta t > 0$ reads*

$$q^{n+1} = q^n + \Delta t [-\mathcal{D}\nabla V + \operatorname{div} \mathcal{D}] (q^n) + \sqrt{2\Delta t} \mathcal{D}^{1/2}(q^n) G^n, \quad (10)$$

where $(G^n)_{n \geq 0}$ is a family of independent and identically distributed standard d -dimensional Gaussian random variables. It is apparent from the above formula that the behavior of the numerical scheme actually depends on $\Delta t \mathcal{D}$ (and the divergence of this quantity), and not on \mathcal{D} or Δt alone, so that multiplying \mathcal{D} by a constant can be equivalently seen as multiplying the time step by the same constant. In this context, one should think of normalization criteria as ways to fix the amount of numerical error involved in the discretization, for instance by fixing the average rejection probability in a Metropolis–Hastings scheme whose proposal is provided by (10). The normalization we discuss in Section 2.2 is conceptually simpler than such a criterion, and not related to a specific numerical scheme. We intend to further study this perspective in following works.

2.2 Normalization constraint

We make precise in this section the normalization we consider for diffusion matrices, in view of the discussion after (8). In essence, we choose a (weighted) Lebesgue norm to be smaller than 1.

A quantity which naturally appears in the mathematical formulation of the problem is the product $\mathcal{D}(q)e^{-V(q)}$, present for instance in the numerator of the Rayleigh ratio (7). Note also that the divergence of this product is the drift in (3) (up to the factor $e^{-V(q)}$). This motivates normalizing the product $\mathcal{D}e^{-V}$ (which is proportional to $\mathcal{D}\mu$) rather than \mathcal{D} itself.

To formulate precisely the normalization we consider, it is convenient to introduce, for $a, b \geq 0$, the set

$$\mathcal{M}_{a,b} = \left\{ M \in \mathcal{S}_d^+ \mid \forall \xi \in \mathbb{R}^d, \ a|\xi|^2 \leq \xi^\top M \xi \leq \frac{1}{b}|\xi|^2 \right\},$$

with $|\cdot|$ the Euclidean norm on \mathbb{R}^d , and the convention that the right-hand side of the last inequality is $+\infty$ when $b = 0$. Considering $a, b > 0$ is useful to study homogenization limits, as we do in Section 5. We will always require that

$$ab \leq 1,$$

in order for $\mathcal{M}_{a,b}$ to be nonempty. We endow $\mathcal{M}_{a,b}$ with a norm $|\cdot|_{\mathbb{F}}$, which is any matrix norm compatible with the order on symmetric positive matrices, *i.e.* $|M_1|_{\mathbb{F}} \leq |M_2|_{\mathbb{F}}$ when $0 \leq M_1 \leq M_2$ in the sense of symmetric matrices. One such norm is the Frobenius norm.

We next denote by $L_V^p(\mathbb{T}^d, \mathcal{M}_{a,b})$ for $p \in [1, +\infty]$ the Banach space of measurable functions \mathcal{D} from \mathbb{T}^d to \mathcal{S}_d^+ such that $\mathcal{D}(q)e^{-V(q)} \in \mathcal{M}_{a,b}$ for almost every $q \in \mathbb{T}^d$ and $\|\mathcal{D}\|_{L_V^p} < +\infty$, with the norm

$$\|\mathcal{D}\|_{L_V^p} = \left(\int_{\mathbb{T}^d} |\mathcal{D}(q)|_{\mathbb{F}}^p e^{-pV(q)} dq \right)^{1/p}, \quad (11)$$

when $p < +\infty$, and $\|\mathcal{D}\|_{L_V^\infty} = \|\mathcal{D}|_{\mathbb{F}} e^{-V}\|_{L^\infty(\mathbb{T}^d)}$ when $p = +\infty$. Let us emphasize that the L_V^p -norm is in fact the usual L^p -norm applied to the product $|\mathcal{D}|_{\mathbb{F}} e^{-V}$. Other choices of normalization could be considered, see Remark 8 in Appendix D.4. One motivation for (11) is that it leads to simple expressions for optimal diffusion matrices in the homogenized limit (see Section 5.2.2).

Simply considering a normalization constraint in L_V^∞ -norm leads to a trivial solution for the maximization of the spectral gap (which simply saturates the constraint), as shown in the following result (see related discussions in [38, Section 10.1]).

Lemma 1. Fix $a \geq 0$ and $b > 0$ such that $ab \leq 1$. Then, the spectral gap $\Lambda(\mathcal{D})$ is maximized on $L_V^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$ by the diffusion matrix $\mathcal{D}^*(q) = b^{-1}e^V \text{Id}_d$.

Proof. Note first that, for any $\mathcal{D} \in L_V^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$, it holds $\mathcal{D}(q) \leq b^{-1}e^{V(q)} \text{Id}_d$ in the sense of symmetric matrices for almost all $q \in \mathbb{T}^d$. Therefore, for any $u \in H_0^1(\mu)$,

$$\frac{\int_{\mathbb{T}^d} \nabla u(q)^\top \mathcal{D}(q) \nabla u(q) \mu(q) dq}{\int_{\mathbb{T}^d} u(q)^2 \mu(q) dq} \leq \frac{\int_{\mathbb{T}^d} \nabla u(q)^\top \mathcal{D}^*(q) \nabla u(q) \mu(q) dq}{\int_{\mathbb{T}^d} u(q)^2 \mu(q) dq}.$$

Passing to the infimum over u on the left-hand side first, then on the right-hand side, leads to $\Lambda(\mathcal{D}) \leq \Lambda(\mathcal{D}^*)$ for all $\mathcal{D} \in L_V^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$. Moreover, it obviously holds that $b^{-1}e^V \text{Id}_d \in L_V^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$, which concludes the proof. \square

In view of Lemma 1, we focus in the remainder of this work on a constraint on the L_V^p norm of the diffusion matrix for $1 \leq p < +\infty$, and therefore consider the subsets

$$\mathfrak{D}_p^{a,b} = \left\{ \mathcal{D} \in L_V^\infty(\mathbb{T}^d, \mathcal{M}_{a,b}) \left| \int_{\mathbb{T}^d} |\mathcal{D}(q)|_{\mathbb{F}}^p e^{-pV(q)} dq \leq 1 \right. \right\}. \quad (12)$$

In order for these sets to be nonempty, the parameter $a \geq 0$ should be small enough.

Lemma 2. Fix $1 \leq p < +\infty$, and $a, b \geq 0$. The set $\mathfrak{D}_p^{a,b}$ is nonempty if and only if

$$ab \leq 1, \quad a \leq a_{\max} := \frac{1}{|\text{Id}_d|_{\mathbb{F}}}. \quad (13)$$

Proof. Note that any element of $\mathfrak{D}_p^{a,b}$ is lower bounded by $ae^V \text{Id}_d$. Therefore, the set is nonempty if and only if it contains $ae^V \text{Id}_d$. The equality

$$\int_{\mathbb{T}^d} |ae^V \text{Id}_d|_{\mathbb{F}}^p e^{-pV} = a^p |\text{Id}_d|_{\mathbb{F}}^p$$

then gives the claimed result. \square

We can now state the maximization problem which is considered in the remainder of this work. Fix $p \in [1, +\infty)$. For any $a \in [0, a_{\max}]$ and $b > 0$ such that $ab \leq 1$,

$$\text{find } \mathcal{D}^* \in \mathfrak{D}_p^{a,b} \text{ such that } \Lambda(\mathcal{D}^*) = \sup_{\mathcal{D} \in \mathfrak{D}_p^{a,b}} \Lambda(\mathcal{D}). \quad (14)$$

3 Theoretical analysis of the spectral gap optimization

We present in this section some theoretical results on the mathematical analysis of the optimization problem (14). We start by well-posedness results in Section 3.1, and we then formally characterize in Section 3.2 the optimal diffusion matrix.

3.1 Well posedness of the optimization problem

The aim of this section is to show that the maximization of the spectral gap $\Lambda(\mathcal{D})$ on the set $\mathfrak{D}_p^{a,b}$ defined in (12) is a well-posed problem. This is easily done by adapting arguments from [38]. Some results rely on the fact that the probability measure μ satisfies a Poincaré inequality. Recall that the potential V is assumed to be $\mathcal{C}^\infty(\mathbb{T}^d)$ in all this work. This implies in particular that the potential V is bounded and that the density μ is bounded from below by a positive constant and

bounded from above on \mathbb{T}^d . The measure with density (1) therefore satisfies a Poincaré inequality, *i.e.* there exists $C_\mu > 0$ such that

$$\inf_{u \in H_0^1(\mu)} \left\{ \frac{\int_{\mathbb{T}^d} |\nabla u(q)|^2 \mu(q) dq}{\int_{\mathbb{T}^d} u(q)^2 \mu(q) dq} \right\} = \frac{1}{C_\mu}. \quad (15)$$

In addition, there exists $u_\mu \in H_0^1(\mu) \setminus \{0\}$ such that the infimum (15) is attained, *i.e.*

$$\frac{\int_{\mathbb{T}^d} |\nabla u_\mu(q)|^2 \mu(q) dq}{\int_{\mathbb{T}^d} u_\mu(q)^2 \mu(q) dq} = \frac{1}{C_\mu}.$$

Let us first prove that the function Λ is well-defined on $\mathfrak{D}_p^{a,b}$ for $a, b \geq 0$. On the one hand, $\Lambda(\mathcal{D}) \geq 0$ (since \mathcal{D} is positive semi-definite on $\mathfrak{D}_p^{a,b}$). On the other hand, for a fixed function $u \in \mathcal{C}^\infty(\mathbb{T}^d)$ satisfying $\int_{\mathbb{T}^d} u(q) \mu(q) dq = 0$ and $\int_{\mathbb{T}^d} u(q)^2 \mu(q) dq = 1$, the function $|\nabla u(q)|^2$ is bounded on \mathbb{T}^d . Since there exists a constant $K \in \mathbb{R}_+$ such that $0 \leq \xi^\top \mathcal{D}(q) \xi \leq K |\mathcal{D}(q)|_F |\xi|^2$ (the constant K being 1 for the Frobenius norm), it holds (recall that Z is defined in (1))

$$\begin{aligned} \Lambda(\mathcal{D}) &\leq Z^{-1} K \|\nabla u\|_{L^\infty(\mathbb{T}^d)}^2 \int_{\mathbb{T}^d} |\mathcal{D}(q)|_F e^{-V(q)} dq \\ &\leq Z^{-1} K \|\nabla u\|_{L^\infty(\mathbb{T}^d)}^2 \left(\int_{\mathbb{T}^d} |\mathcal{D}(q)|_F^p e^{-pV(q)} dq \right)^{1/p} < +\infty, \end{aligned}$$

using the Hölder's inequality on \mathbb{T}^d . This shows that $\Lambda(\mathcal{D})$ is well-defined and that there exists $C > 0$ such that, for any $a, b \geq 0$ and any $\mathcal{D} \in \mathfrak{D}_p^{a,b}$, one has

$$0 \leq \Lambda(\mathcal{D}) \leq C.$$

Moreover, for any $1 \leq p \leq \infty$, the subset $\mathfrak{D}_p^{a,b}$ is weakly closed for the norm L_V^p defined in (11) (this follows directly from the weak closedness of standard Lebesgue spaces).

A first useful property to prove the well posedness of the maximization problem (14) is the following classical result on the concavity of the functional Λ , directly obtained from the fact that $\Lambda(\mathcal{D})$ is an infimum of linear functionals in \mathcal{D} (see for instance [38, Theorem 10.1.1]).

Lemma 3. *The functional $\mathcal{D} \mapsto \Lambda(\mathcal{D})$ is concave on the set of bounded, measurable functions $\mathbb{T}^d \rightarrow \mathcal{S}_d^+$.*

The following theorem, proved in Appendix A.1 states the well-posedness of the optimization problem we consider. Note that we assume that $b > 0$, as the upper-semicontinuity of Λ , a crucial argument for the result to hold, is not clear in the case $b = 0$.

Theorem 4. *Fix $p \in [1, +\infty)$. For any $a \in [0, a_{\max}]$ and $b > 0$ such that $ab \leq 1$, there exists a solution $\mathcal{D}^* \in \mathfrak{D}_p^{a,b}$ to (14). In addition, the optimal diffusion matrix \mathcal{D}^* satisfies the following properties:*

(i) *For any open set $\Omega \subset \mathbb{T}^d$, there exists $q \in \Omega$ such that $\mathcal{D}^*(q) \neq 0$;*

(ii) $\int_{\mathbb{T}^d} |\mathcal{D}^*(q)|_F^p e^{-pV(q)} dq = 1.$

Property (ii) states that, as expected in view of (9), the optimal diffusion matrix saturates the constraint. Property (i) is useful only for $a = 0$. It guarantees that \mathcal{D}^* cannot be identically 0 on open sets, but a priori $\mathcal{D}^*(q)$ need not be positive definite, and could vanish on sets of measure 0. If \mathcal{D}^* cancels at some point, it may be optimal in the sense of our functional Λ but not necessarily

in the sense of the spectral gap; see Remark 1. We discuss in Section 3.2 the positive definite character of \mathcal{D}^* . In addition, from a numerical perspective, we prove in Section 4.1.2 that the optimal diffusion matrix is bounded from below by a positive constant when the minimization is performed over the intersection of $\mathfrak{D}_p^{a,b}$ and a finite dimensional linear space. This lower bound is numerically observed to decrease in certain cases when the spatial discretization is refined (see Figure 3 below).

3.2 Characterization of uniformly positive definite maximizers \mathcal{D}^*

The objective of this section is to formally characterize the maximizers \mathcal{D}^* using the Euler–Lagrange condition satisfied by critical points of the functional Λ . When the optimal diffusion is uniformly positive definite, we first show in Section 3.2.1 that $\Lambda(\mathcal{D}^*)$ is degenerate. Therefore, the mapping $\mathcal{D} \mapsto \Lambda(\mathcal{D})$ is not necessarily differentiable at \mathcal{D}^* . In order to obtain an Euler–Lagrange equation characterizing \mathcal{D}^* , we circumvent this difficulty in Section 3.2.2 by relying on a smooth approximation of the definition of Λ . The theoretical results obtained in this section will be corroborated by numerical experiments in Section 4.

3.2.1 The eigenvalue $\Lambda(\mathcal{D}^*)$ is degenerate

Let us consider the solution \mathcal{D}^* to the problem (14) for $p \in (1, +\infty)$,

$$a = 0,$$

and $b > 0$. Note that we exclude the case $p = 1$ as the map $M \mapsto |M|_{\mathbb{F}}$ is not differentiable at any rank-deficient $M \in \mathcal{M}_{0,b}$. By Theorem 4, there exists a matrix valued function \mathcal{D}^* which is a maximizer of Λ over the constrained set $\mathfrak{D}_p^{0,b}$. In the remainder of this section, we make two s on \mathcal{D}^* :

$$\exists b_+ > b, \quad \mathcal{D}^*(q) e^{-V(q)} \leq \frac{1}{b_+} \text{Id}_d \text{ for a.e. } q \in \mathbb{T}^d. \quad (16)$$

and

$$\exists c > 0, \quad \mathcal{D}^*(q) \geq c \text{Id}_d \text{ for a.e. } q \in \mathbb{T}^d. \quad (17)$$

From (16) and (17), the constraints associated to the lower bound and upper bound in $\mathfrak{D}_p^{0,b}$ are not active at the critical point \mathcal{D}^* . A standard argument based on the Euler–Lagrange equation satisfied by the minimizer of the Rayleigh quotient in (7) shows that there exists a function $u_{\mathcal{D}^*} \in H_0^1(\mu)$ such that $\|u_{\mathcal{D}^*}\|_{L^2(\mu)} = 1$ and $\Lambda(\mathcal{D}^*) = \int_{\mathbb{T}^d} \nabla u_{\mathcal{D}^*}(q)^\top \mathcal{D}^*(q) \nabla u_{\mathcal{D}^*}(q) \mu(q) dq$.

In our numerical experiments, we always observe that (16) is satisfied up to choosing b sufficiently small. As explained below, (17) is in particular natural for potentials which are $1/k$ -periodic with $k \in \mathbb{N} \setminus \{0\}$, and k sufficiently large to approach the homogenized limit, for which the optimal diffusion is indeed uniformly positive definite; see Section 5.

We show in this section that, when (17) holds, $\Lambda(\mathcal{D}^*)$ is a degenerate eigenvalue of the operator $-\mathcal{L}_{\mathcal{D}^*}$ (under some regularity s in the case $d = 1$). The proof proceeds by contradiction and relies on the fact that, when $\Lambda(\mathcal{D}^*)$ is non-degenerate and (17) holds, one can write the Euler–Lagrange equation satisfied for \mathcal{D}^* . This leads to a characterization of the optimal diffusion matrix which contradicts (17). In order to state the result, we need to distinguish for technical reasons the cases $d = 1$ and $d > 1$.

Lemma 5. *Choose the Frobenius norm for the matrix norm $|\cdot|_{\mathbb{F}}$, consider $p \in (1, +\infty)$ and $a = 0$. Assume that the maximizer \mathcal{D}^* solution to (14) satisfies (16). Assume moreover that $\mathcal{D}^* \in \mathcal{C}^0(\mathbb{T}, \mathbb{R}_+)$ when $d = 1$. If (17) holds, then the optimal spectral gap $\Lambda(\mathcal{D}^*)$ is a degenerate eigenvalue of the diffusion operator $-\mathcal{L}_{\mathcal{D}^*}$.*

Proof. Suppose that $\mathcal{D}^* \geq c \text{Id}_d$ for $c > 0$. We denote by $u_{\mathcal{D}^*}$ an eigenvector satisfying $-\mathcal{L}_{\mathcal{D}^*} u_{\mathcal{D}^*} = \Lambda(\mathcal{D}^*) u_{\mathcal{D}^*}$. Observe first that, for any $\delta \mathcal{D} \in L^\infty(\mathbb{T}^d, \mathcal{S}_d)$, where \mathcal{S}_d is the set of real, symmetric, matrices of size $d \times d$ (note that $\delta \mathcal{D}$ does not output a positive matrix a priori), the matrix valued

function $\mathcal{D}^* + t\delta\mathcal{D}$ has values in the space of symmetric positive definite matrices when $|t|$ is sufficiently small.

The proof proceeds by contradiction. If the eigenvalue $\Lambda(\mathcal{D}^*)$ is non-degenerate, the eigenvalue $\Lambda(\mathcal{D}^* + t\delta\mathcal{D})$ remains non-degenerate and isolated for $|t|$ sufficiently small, and it is possible to choose the eigenvector $u_{\mathcal{D}^* + t\delta\mathcal{D}}$ associated with this eigenvalue so that $\|u_{\mathcal{D}^* + t\delta\mathcal{D}}\|_{L^2(\mu)} = 1$ and the mappings $t \mapsto u_{\mathcal{D}^* + t\delta\mathcal{D}}$ and $t \mapsto \Lambda(\mathcal{D}^* + t\delta\mathcal{D})$ are analytic in an open neighborhood of $t = 0$ (see, e.g., Theorem II.6.1 and the discussion in Section VII.3.1 of [42]). The optimal diffusion matrix \mathcal{D}^* is therefore characterized by the following Euler–Lagrange equation: there exists a Lagrange multiplier $\gamma \in \mathbb{R}$ such that

$$\forall \delta\mathcal{D} \in L^\infty(\mathbb{T}^d, \mathcal{S}_d), \quad \left. \frac{d}{dt} \Lambda(\mathcal{D}^* + t\delta\mathcal{D}) \right|_{t=0} + \gamma \left. \frac{d}{dt} \Phi_p(\mathcal{D}^* + t\delta\mathcal{D}) \right|_{t=0} = 0, \quad (18)$$

where Φ_p encodes the normalization constraint (11):

$$\Phi_p(\mathcal{D}) = \int_{\mathbb{T}^d} |\mathcal{D}(q)|_{\mathbb{F}}^p e^{-pV(q)} dq. \quad (19)$$

Note that there are no constraints in (18) related to positivity or boundedness because of (16)–(17).

Relying on the choice of the Frobenius norm for $|\cdot|_{\mathbb{F}}$ to compute the differential of Φ_p , (18) can be rewritten as: for any $\delta\mathcal{D} \in L^\infty(\mathbb{T}^d, \mathcal{S}_d)$,

$$\int_{\mathbb{T}^d} \delta\mathcal{D}(q) : (\nabla u_{\mathcal{D}^*} \otimes \nabla u_{\mathcal{D}^*}) \mu(q) dq = p\gamma \int_{\mathbb{T}^d} |\mathcal{D}^*(q)|_{\mathbb{F}}^{p-2} \mathcal{D}^*(q) : \delta\mathcal{D}(q) e^{-pV(q)} dq,$$

where $:$ and \otimes are respectively the double contraction and outer product operators: for any $M_1, M_2 \in \mathbb{R}^{d \times d}$ and $\xi, \zeta \in \mathbb{R}^d$,

$$M_1 : M_2 = \text{Tr}(M_1^T M_2) = \sum_{i,j=1}^d [M_1]_{i,j} [M_2]_{i,j}, \quad [\xi \otimes \zeta]_{i,j} = \xi_i \zeta_j.$$

To compute the derivative of $\Lambda(\mathcal{D}^* + t\delta\mathcal{D})$ around $t = 0$, as in the Hellmann–Feynman theorem of quantum mechanics, we have used that $du_{\mathcal{D}^* + t\delta\mathcal{D}}/dt$ is orthogonal to $u_{\mathcal{D}^* + t\delta\mathcal{D}}$ in $L^2(\mu)$ (which follows from the normalization condition $\|u_{\mathcal{D}^* + t\delta\mathcal{D}}\|_{L^2(\mu)} = 1$), and the equality

$$\nabla^*(\mathcal{D}^* \nabla u_{\mathcal{D}^*}) = \Lambda(\mathcal{D}^*) u_{\mathcal{D}^*}, \quad (20)$$

where we recall that ∇^* is the adjoint of ∇ on $L^2(\mu)$. We therefore deduce from the Euler–Lagrange condition (18) that the optimal diffusion matrix \mathcal{D}^* satisfies

$$\mathcal{D}^*(q) = \gamma_p |\mathcal{D}^*(q)|_{\mathbb{F}}^{2-p} e^{(p-1)V(q)} \nabla u_{\mathcal{D}^*}(q) \otimes \nabla u_{\mathcal{D}^*}(q), \quad (21)$$

where the constant $\gamma_p > 0$ is determined by the constraint $\Phi_p(\mathcal{D}^*) = 1$.

When $d \geq 2$, we immediately obtain a contradiction with the $\mathcal{D}^* \geq c \text{Id}_d$ since (21) implies that $\mathcal{D}^*(q)$ is rank deficient (in fact, rank 1) for a.e. $q \in \mathbb{T}^d$.

When $d = 1$, (21) simplifies (after some straightforward manipulations) as

$$\mathcal{D}^*(q) = \tilde{\gamma}_p e^{V(q)} |u'_{\mathcal{D}^*}(q)|^{2/(p-1)}, \quad (22)$$

where the constant $\tilde{\gamma}_p > 0$ is determined by $\Phi_p(\mathcal{D}^*) = 1$. We now show that $u'_{\mathcal{D}^*}$ cancels on \mathbb{T} , thus contradicting the fact that $\mathcal{D}^*(q) \geq c > 0$ for all $q \in \mathbb{T}$ (the inequality is indeed valid for all $q \in \mathbb{T}$, and not up to a set of Lebesgue measure 0 since \mathcal{D}^* is assumed to be continuous). To do so, we start by noticing that the continuity of \mathcal{D}^* implies that the eigenfunction $u_{\mathcal{D}^*}$ belongs to $\mathcal{C}^1(\mathbb{T}, \mathbb{R})$ (see [38, Remark 1.2.11] and [31, Theorem 9.15]), so that $u'_{\mathcal{D}^*}$ is continuous on \mathbb{T} . Since $u_{\mathcal{D}^*}$ is periodic on \mathbb{T} , its derivative $u'_{\mathcal{D}^*}$ cancels at least once on \mathbb{T} . In view of (22), the diffusion \mathcal{D}^* then also vanishes at this point. The contradiction finally allows to obtain the desired result for $d = 1$ as well. \square

Remark 4. Let us write more precisely the eigenproblem characterizing the second eigenpair in the situation where the Euler–Lagrange equation (21) holds, which is for instance the case when the second eigenvalue is nondegenerate. Since $|\cdot|_F$ is the Frobenius norm, (21) implies that $|\mathcal{D}^*(q)|_F = \gamma_p e^{V(q)} |\nabla u_{\mathcal{D}^*}(q)|^{2/(p-1)}$ for some parameter $\gamma_p > 0$ determined by the normalization condition, so that $\mathcal{D}^*(q) = \gamma_p e^{V(q)} |\nabla u_{\mathcal{D}^*}(q)|^{-2(p-2)/(p-1)} \nabla u_{\mathcal{D}^*}(q) \otimes \nabla u_{\mathcal{D}^*}(q)$. In view of (20) and the fact that, for any vector valued function φ , it holds $\nabla^* \varphi = -\mu^{-1} \operatorname{div}(\mu \varphi)$, it then follows that

$$-\gamma_p \operatorname{div} \left(|\nabla u_{\mathcal{D}^*}|^{2/(p-1)} \nabla u_{\mathcal{D}^*} \right) = \Lambda(\mathcal{D}^*) e^{-V} u_{\mathcal{D}^*}. \quad (23)$$

Up to a weight factor on the right-hand side, Equation (23) is an eigenvalue problem for the $2p/(p-1)$ -Laplacian (see, e.g. [52]). This equation is non-linear. The existing works on eigenvalue problems for nonlinear operators as the one appearing in (23) suggest that the solutions should be only $\mathcal{C}^{1,\alpha}$ (see again [52]). Singularities in the derivatives are indeed numerically observed in the case of vanishing optimal diffusion coefficients \mathcal{D}^* , see e.g. Figures 3b and 4a.

3.2.2 Euler–Lagrange equation for \mathcal{D}^*

Under (17), the first nonzero eigenvalue for the operator associated with the maximizer \mathcal{D}^* is degenerate, and one cannot rely on the Euler–Lagrange equation (21). Yet, a characterization of the optimal diffusion matrix can still be formally recovered using a smooth-min approximation. This leads to a generalization of the formula (21), which is corroborated by numerical experiments, see Section 4.2. The positivity of \mathcal{D}^* is discussed at the end of this section.

Smooth-min approximation of the maximization problem. Our approach relies on the fact that, for a positive α , the map $\mathfrak{m}_\alpha : \mathbb{R}^n \rightarrow \mathbb{R}$ defined by

$$\mathfrak{m}_\alpha(x_1, \dots, x_n) = \frac{\sum_{i=1}^n x_i e^{-\alpha x_i}}{\sum_{i=1}^n e^{-\alpha x_i}}$$

converges to the min function when $\alpha \rightarrow +\infty$, i.e.

$$\forall (x_1, \dots, x_n) \in \mathbb{R}^n, \quad \mathfrak{m}_\alpha(x_1, \dots, x_n) \xrightarrow{\alpha \rightarrow +\infty} \min_{1 \leq i \leq n} x_i. \quad (24)$$

We adapt this definition to obtain a smooth approximation of the spectral gap $\Lambda(\mathcal{D})$, from which one can compute the gradient, and exchange the derivative/limit when $\alpha \rightarrow +\infty$ in order to obtain a formal characterization of the optimal diffusion matrix similar to (21).

We first construct the counterpart of the map \mathfrak{m}_α in our framework. In order to do so, we fix $\mathcal{D} \in L_V^p(\mathbb{T}^d, \mathcal{M}_{a,b})$ such that $\mathcal{D} \geq c \operatorname{Id}_d$ for some $c > 0$ (see (17)). We denote by $g_\alpha : x \mapsto e^{-\alpha x}$ and $h_\alpha : x \mapsto x e^{-\alpha x}$. By functional calculus, the maps g_α and h_α can be generalized to functions of the (positive) self-adjoint operator $-\mathcal{L}_{\mathcal{D}}$. With an abuse of notation, we write in this case $g_\alpha(\mathcal{D})$ and $h_\alpha(\mathcal{D})$ instead of $g_\alpha(-\mathcal{L}_{\mathcal{D}})$ and $h_\alpha(-\mathcal{L}_{\mathcal{D}})$. Since \mathcal{D} is bounded and uniformly bounded from below by a positive constant (in the sense of symmetric matrices), the positive operator $-\mathcal{L}_{\mathcal{D}}$ has discrete spectrum, its minimum eigenvalue being simple and equal to 0, see Section 2.1. We therefore write

$$-\mathcal{L}_{\mathcal{D}} = \sum_{i \geq 1} \lambda_i P_i, \quad (25)$$

where $\lambda_1 = 0 < \lambda_2 < \dots < \lambda_i < \dots$ are the distinct eigenvalues of $-\mathcal{L}_{\mathcal{D}}$ and P_i is the orthogonal projection onto $\operatorname{Ker}(-\mathcal{L}_{\mathcal{D}} - \lambda_i \operatorname{Id})$. The multiplicity of λ_i is denoted by $N_i = \dim \operatorname{Ran}(P_i) \geq 1$. For $\alpha > 0$, we define

$$f_\alpha(\mathcal{D}) = \frac{\operatorname{Tr}(h_\alpha(\mathcal{D}))}{\operatorname{Tr}(g_\alpha(\mathcal{D})) - 1} = \frac{\operatorname{Tr}((-\mathcal{L}_{\mathcal{D}}) e^{\alpha \mathcal{L}_{\mathcal{D}}})}{\operatorname{Tr}(e^{\alpha \mathcal{L}_{\mathcal{D}}}) - 1}, \quad (26)$$

where the traces are taken on $L^2(\mu)$ (in case the traces are taken on another space than $L^2(\mu)$, this will be made precise with a subscript) and the operators $g_\alpha(\mathcal{D}), h_\alpha(\mathcal{D})$ are defined by functional calculus, *i.e.*

$$\begin{cases} g_\alpha(\mathcal{D}) = \sum_{i \geq 1} g_\alpha(\lambda_i) P_i, \\ h_\alpha(\mathcal{D}) = \sum_{i \geq 2} h_\alpha(\lambda_i) P_i, \end{cases}$$

where we used that $h_\alpha(\lambda_1) = 0$. The ratio (26) is well-defined in view of Lemma 23 in Appendix B.2 and the fact that $\text{Tr}(g_\alpha(\mathcal{D})) > 1$. Formally, since $\lambda_1 = 0$ and $N_1 = 1$,

$$f_\alpha(\mathcal{D}) = \frac{\sum_{i \geq 2} N_i \lambda_i e^{-\alpha \lambda_i}}{\sum_{i \geq 2} N_i e^{-\alpha \lambda_i}},$$

so that, again formally (compare with (24)),

$$f_\alpha(\mathcal{D}) = \frac{\lambda_2 \left(1 + \frac{1}{N_2 \lambda_2} \sum_{i \geq 3} N_i \lambda_i e^{-\alpha(\lambda_i - \lambda_2)} \right)}{1 + \frac{1}{N_2} \sum_{i \geq 3} N_i e^{-\alpha(\lambda_i - \lambda_2)}} \xrightarrow{\alpha \rightarrow +\infty} \lambda_2 = \min_{i \geq 2} \lambda_i. \quad (27)$$

This limit is justified after the statement of Lemma 23 in Appendix B.2 using the dominated convergence theorem. This shows that the quantity $f_\alpha(\mathcal{D})$ is an approximation of $\Lambda(\mathcal{D})$.

This motivates considering the following approximation of the problem (14): fix $p \in [1, +\infty)$. For any $a \in [0, a_{\max}]$ and $b > 0$ such that $ab \leq 1$,

$$\text{find } \mathcal{D}^{\star, \alpha} \in \mathfrak{D}_p^{a, b} \text{ such that } f_\alpha(\mathcal{D}^{\star, \alpha}) = \sup_{\mathcal{D} \in \mathfrak{D}_p^{a, b}} f_\alpha(\mathcal{D}). \quad (28)$$

From (27), we expect that $\mathcal{D}^{\star, \infty}$ (which is the limit of $\mathcal{D}^{\star, \alpha}$ as $\alpha \rightarrow +\infty$ whenever it exists) is the optimal diffusion \mathcal{D}^\star solution to (14): this will be corroborated by numerical experiments in Section 4.2. One can observe that, for a fixed \mathcal{D} , $f_\alpha(\mathcal{D})$ is a convex combination of the eigenvalues λ_i , which implies that $f_\alpha(\mathcal{D}) \geq \Lambda(\mathcal{D})$. Therefore, it holds $f_\alpha(\mathcal{D}^{\star, \alpha}) \geq \Lambda(\mathcal{D}^\star)$. One can also show that the map $\alpha \mapsto f_\alpha(\mathcal{D})$ is nonincreasing, either by computing its derivative and using the Cauchy–Schwarz inequality, or by observing that f_α is the opposite of the derivative of the logarithm of the partition function $\alpha \mapsto \sum_{i \geq 2} N_i e^{-\alpha \lambda_i}$, and using the fact that the second derivative of the logarithm of a partition function is only composed of a variance term and is therefore nonnegative. These computations can be made rigorous using the theory developed in Appendix B.

As in the previous section, we assume that there exists a solution to (28) for $a = 0$ and some $b > 0$, which satisfies (see (16)-(17))

$$\exists b_+ > b, \quad \mathcal{D}^{\star, \alpha}(q) e^{-V(q)} \leq \frac{1}{b_+} \text{Id}_d \text{ for a.e. } q \in \mathbb{T}^d, \quad (29)$$

and

$$\exists c > 0, \quad \mathcal{D}^{\star, \alpha}(q) \geq c \text{Id}_d \text{ for a.e. } q \in \mathbb{T}^d. \quad (30)$$

In particular, the constraints associated to the lower bound and the upper bound in $\mathfrak{D}_p^{0, b}$ are not active at the critical point $\mathcal{D}^{\star, \alpha}$. In our numerical experiments, we observe that (29) is always satisfied for b sufficiently small. Moreover, we use (30) below for technical reasons (in order to rigorously define f_α and prove that it is differentiable). However, we will observe numerically that the characterization that we finally obtain on $\mathcal{D}^{\star, \infty}$ (by sending $\alpha \rightarrow +\infty$, see Equation (36) below) seems to also hold even when (30) is not satisfied.

Characterization of the optimal diffusion matrix for a finite value of α . The map f_α defined by (26) is smooth over the set of functions $\mathcal{D} \in L_V^p(\mathbb{T}^d, \mathcal{M}_{a,b})$ such that $\mathcal{D} \geq c \text{Id}_d$ for some $c > 0$, and therefore its differential can be computed even when some eigenvalues are degenerate. The main result of this section is the computation of the differentials of the map g_α and h_α , whose proof is postponed to Appendix B.1. Note that the derivatives of the maps g_α and h_α , namely the maps g'_α and h'_α defined by

$$\forall x \in \mathbb{R}, \forall \alpha > 0, \quad g'_\alpha(x) = -\alpha e^{-\alpha x}, \quad h'_\alpha(x) = (1 - \alpha x) e^{-\alpha x},$$

can also be generalized to functions of the self-adjoint operator $-\mathcal{L}_\mathcal{D}$, and we then write $g'_\alpha(\mathcal{D}) = g'_\alpha(-\mathcal{L}_\mathcal{D})$ and $h'_\alpha(\mathcal{D}) = h'_\alpha(-\mathcal{L}_\mathcal{D})$.

Proposition 6. Fix $\alpha > 0$ and $\mathcal{D} \in L_V^p(\mathbb{T}^d, \mathcal{M}_{a,b})$ such that $\mathcal{D} \geq c \text{Id}_d$ for some $c > 0$. Then there exists $K \in \mathbb{R}_+$ such that, for any $\delta\mathcal{D} \in L^\infty(\mathbb{T}^d, \mathcal{S}_d)$ satisfying $\|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} \leq c/2$, the operators $g_\alpha(\mathcal{D} + \delta\mathcal{D})$, $g_\alpha(\mathcal{D})$, $g'_\alpha(\mathcal{D})\mathcal{L}_{\delta\mathcal{D}}$ and the operators $h_\alpha(\mathcal{D} + \delta\mathcal{D})$, $h_\alpha(\mathcal{D})$, $h'_\alpha(\mathcal{D})\mathcal{L}_{\delta\mathcal{D}}$ are trace-class on $L^2(\mu)$ and

$$\begin{cases} |\text{Tr}(g_\alpha(\mathcal{D} + \delta\mathcal{D})) - \text{Tr}(g_\alpha(\mathcal{D})) - \text{Tr}(g'_\alpha(\mathcal{D})(-\mathcal{L}_{\delta\mathcal{D}}))| \leq K \|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)}^2, \\ |\text{Tr}(h_\alpha(\mathcal{D} + \delta\mathcal{D})) - \text{Tr}(h_\alpha(\mathcal{D})) - \text{Tr}(h'_\alpha(\mathcal{D})(-\mathcal{L}_{\delta\mathcal{D}}))| \leq K \|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)}^2. \end{cases} \quad (31)$$

This result immediately implies a differentiability result for the map f_α .

Corollary 7. Fix $\alpha > 0$ and $\mathcal{D} \in L_V^p(\mathbb{T}^d, \mathcal{M}_{a,b})$ such that $\mathcal{D} \geq c \text{Id}_d$ for some $c > 0$. Then there exists $K \in \mathbb{R}_+$ such that, for any $\delta\mathcal{D} \in L^\infty(\mathbb{T}^d, \mathcal{S}_d)$ satisfying $\|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} \leq c/2$,

$$|f_\alpha(\mathcal{D} + \delta\mathcal{D}) - f_\alpha(\mathcal{D}) - \mathcal{F}_\alpha(\mathcal{D})(\delta\mathcal{D})| \leq K \|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)}^2,$$

where

$$\mathcal{F}_\alpha(\mathcal{D})(\delta\mathcal{D}) = \frac{\text{Tr}(h'_\alpha(\mathcal{D})(-\mathcal{L}_{\delta\mathcal{D}})) [\text{Tr}(g_\alpha(\mathcal{D})) - 1] - \text{Tr}(h_\alpha(\mathcal{D})) \text{Tr}(g'_\alpha(\mathcal{D})(-\mathcal{L}_{\delta\mathcal{D}}))}{(\text{Tr}(g_\alpha(\mathcal{D})) - 1)^2}.$$

Straightforward computations lead to

$$\mathcal{F}_\alpha(\mathcal{D})(\delta\mathcal{D}) = \sum_{i \geq 1} \left[\frac{\mathcal{G}_\alpha(1 - \alpha\lambda_i) + \alpha\mathcal{H}_\alpha}{\mathcal{G}_\alpha^2} e^{-\alpha\lambda_i} \right] \text{Tr}(P_i(-\mathcal{L}_{\delta\mathcal{D}})),$$

with

$$\mathcal{G}_\alpha = \sum_{j \geq 2} N_j e^{-\alpha\lambda_j}, \quad \mathcal{H}_\alpha = \sum_{j \geq 2} N_j \lambda_j e^{-\alpha\lambda_j}. \quad (32)$$

We now formally give an explicit formula of the optimal diffusion matrix $\mathcal{D}^{*,\alpha}$ solution to (28), following the same computations as in the proof of Lemma 5. This requires to introduce a Hilbert basis $(e_k)_{k \geq 1}$ composed of $L^2(\mu)$ normalized eigenvectors of $-\mathcal{L}_\mathcal{D}$, namely

$$\forall k, \ell \geq 1, \quad -\mathcal{L}_\mathcal{D} e_k = \lambda_k e_k, \quad \langle e_k, e_\ell \rangle_{L^2(\mu)} = \delta_{k,\ell},$$

where δ is the Kronecker symbol and the eigenvalues $\lambda_1 = 0 \leq \lambda_2 \leq \dots \leq \lambda_k \leq \dots$ are ordered and counted with their multiplicity:

$$\forall i \geq 1, \quad \forall k \in \left\{ 1 + \sum_{j=1}^{i-1} N_j, \dots, \sum_{j=1}^i N_j \right\}, \quad \lambda_k = \lambda_i.$$

We rewrite the spectral decomposition (25) of $-\mathcal{L}_\mathcal{D}$ in terms of this given Hilbert basis and with possibly degenerate eigenvalues: for a given smooth function φ ,

$$-\mathcal{L}_\mathcal{D} \varphi = \sum_{k \geq 1} \lambda_k \langle \varphi, e_k \rangle_{L^2(\mu)} e_k.$$

Exploiting the Euler–Lagrange equation for the objective function f_α , namely using that $\mathcal{F}_\alpha(\mathcal{D}^{\star,\alpha})$ is proportional to the differential of the constraint function (19):

$$\forall \delta \mathcal{D} \in L^\infty(\mathbb{T}^d, \mathcal{S}_d), \quad \mathcal{F}_\alpha(\mathcal{D}^{\star,\alpha})(\delta \mathcal{D}) \propto \int_{\mathbb{T}^d} |\mathcal{D}^{\star,\alpha}(q)|_{\mathbb{F}}^{p-2} e^{-pV(q)} \mathcal{D}^{\star,\alpha}(q) : \delta \mathcal{D}(q) dq.$$

Since

$$P_i = \sum_{k=1+\sum_{l=1}^{i-1} N_l}^{\sum_{l=1}^i N_l} \langle e_k, \cdot \rangle_{L^2(\mu)} e_k$$

and

$$\begin{aligned} \text{Tr} \left(\langle e_k, \cdot \rangle_{L^2(\mu)} e_k \nabla^* \delta \mathcal{D} \nabla \right) &= \int_{\mathbb{T}^d} e_k(q) [\nabla^* \delta \mathcal{D}(q) \nabla] e_k(q) \mu(q) dq \\ &= \int_{\mathbb{T}^d} \nabla e_k(q)^\top \delta \mathcal{D}(q) \nabla e_k(q) \mu(q) dq \\ &= \int_{\mathbb{T}^d} \delta \mathcal{D}(q) : (\nabla e_k(q) \otimes \nabla e_k(q)) \mu(q) dq, \end{aligned}$$

one obtains the following characterization of the optimal diffusion matrix for a fixed value of α :

$$\mathcal{D}^{\star,\alpha}(q) = \gamma_\alpha |\mathcal{D}^{\star,\alpha}(q)|_{\mathbb{F}}^{2-p} e^{(p-1)V(q)} \sum_{k \geq 2} \left[\frac{\mathcal{G}_\alpha(1 - \alpha \lambda_{k,\alpha}) + \alpha \mathcal{H}_\alpha}{\mathcal{G}_\alpha^2} e^{-\alpha \lambda_{k,\alpha}} \right] \nabla e_{k,\alpha}(q) \otimes \nabla e_{k,\alpha}(q), \quad (33)$$

with $\gamma_\alpha \in \mathbb{R}$ such that $\Phi_p(\mathcal{D}^{\star,\alpha}) = 1$, and where $(\lambda_{k,\alpha})_{k \geq 1}, (e_{k,\alpha})_{k \geq 1}$ are the eigenelements of the operator $-\mathcal{L}_{\mathcal{D}^{\star,\alpha}}$. Since $\mathcal{D}^{\star,\alpha}$ is a maximizer of f_α , the eigenvalues and eigenvectors also depend on α , and the definitions of the quantities \mathcal{G}_α and \mathcal{H}_α first defined in (32) and appearing in (33) are changed accordingly:

$$\mathcal{G}_\alpha = \sum_{j \geq 2} e^{-\alpha \lambda_{j,\alpha}}, \quad \mathcal{H}_\alpha = \sum_{j \geq 2} \lambda_{j,\alpha} e^{-\alpha \lambda_{j,\alpha}}.$$

Note that we used that $\nabla e_{1,\alpha} = 0$ to start the sum at $k = 2$ in (33) since the eigenspace associated to $\lambda_{1,\alpha} = 0$, which is one-dimensional, is spanned by constant functions. As a consequence, one obtains in the case $d = 1$ (recalling that we assume (30) so that the power $1/(p-1)$ is well defined)

$$\mathcal{D}^{\star,\alpha}(q) = \tilde{\gamma}_\alpha e^{V(q)} \left(\sum_{k \geq 2} \left[\frac{\mathcal{G}_\alpha(1 - \alpha \lambda_{k,\alpha}) + \alpha \mathcal{H}_\alpha}{\mathcal{G}_\alpha^2} e^{-\alpha \lambda_{k,\alpha}} \right] |e'_{k,\alpha}(q)|^2 \right)^{1/(p-1)}, \quad (34)$$

with $\tilde{\gamma}_\alpha > 0$. This formula is corroborated by numerical examples below, see Figure 2b.

Formal characterization in the limit $\alpha \rightarrow +\infty$. We now formally compute the limit when $\alpha \rightarrow +\infty$ in (33). For notational simplicity, we consider the case when the limiting problem obtained for $\alpha \rightarrow +\infty$ has a leading nonzero eigenvalue with a degeneracy of at most 2. It is straightforward to adapt the argument to degeneracies of arbitrary order. We make the following s:

- all the eigenelements converge when $\alpha \rightarrow +\infty$: for any $k \geq 1$, there exists $\lambda_{k,\infty} \in \mathbb{R}_+$ such that $\lambda_{k,\alpha} \xrightarrow{\alpha \rightarrow +\infty} \lambda_{k,\infty}$, and there exists $e_{k,\infty} \in H_0^1(\mu)$ such that $e_{k,\alpha} \xrightarrow{\alpha \rightarrow +\infty} e_{k,\infty}$ in $H_0^1(\mu)$;
- if $k \geq 4$, then $\lambda_{k,\infty} > \lambda_{3,\infty}$ (the eigenspace associated to the eigenvalue $\lambda_{2,\infty}$ is either 1 or 2 dimensional);
- when $\lambda_{2,\infty} = \lambda_{3,\infty}$, we further assume that the quantity $\alpha(\lambda_{3,\alpha} - \lambda_{2,\alpha})$ converges to $\eta \in \mathbb{R}_+$ when $\alpha \rightarrow +\infty$.

In the case when $\lambda_{2,\infty} < \lambda_{3,\infty}$, the limit when $\alpha \rightarrow +\infty$ of the sum on the right-hand side of (33) can be computed as

$$\sum_{k \geq 2} \left[\frac{\mathcal{G}_\alpha(1 - \alpha\lambda_{k,\alpha}) + \alpha\mathcal{H}_\alpha}{\mathcal{G}_\alpha^2} e^{-\alpha\lambda_{k,\alpha}} \right] \nabla e_{k,\alpha} \otimes \nabla e_{k,\alpha} \quad (35)$$

$$\underset{\alpha \rightarrow +\infty}{\sim} \frac{e^{-\alpha\lambda_{2,\infty}}(1 - \alpha\lambda_{2,\infty}) + \alpha\lambda_{2,\infty}e^{-\alpha\lambda_{2,\infty}}}{e^{2\alpha\lambda_{2,\infty}}} e^{-\alpha\lambda_{2,\infty}} \nabla e_{2,\infty} \otimes \nabla e_{2,\infty} = \nabla e_{2,\infty} \otimes \nabla e_{2,\infty}.$$

The characterization of the optimal diffusion matrix (33) becomes in the limit $\alpha \rightarrow +\infty$:

$$\mathcal{D}^{*,\infty}(q) = \gamma_\infty |\mathcal{D}^{*,\infty}(q)|_{\mathbb{F}}^{2-p} e^{(p-1)V} \nabla e_{2,\infty} \otimes \nabla e_{2,\infty},$$

with $\gamma_\infty > 0$ such that $\Phi(\mathcal{D}_{p,\infty}^*) = 1$, which simplifies in the case $d = 1$ to

$$\mathcal{D}^{*,\infty}(q) = \tilde{\gamma}_\infty e^{V(q)} |e'_{2,\infty}(q)|^{2/(p-1)},$$

with $\tilde{\gamma}_\infty > 0$. We therefore recover the characterizations (21) and (22) of Section 3.2.1.

When $\lambda_{2,\infty} = \lambda_{3,\infty}$, the factor in front of $\nabla e_{2,\alpha} \otimes \nabla e_{2,\alpha}$ in (35) is asymptotically equivalent to

$$\begin{aligned} & \frac{e^{-\alpha\lambda_{2,\alpha}}}{(e^{-\alpha\lambda_{2,\alpha}} + e^{-\alpha\lambda_{3,\alpha}})^2} [(e^{-\alpha\lambda_{2,\alpha}} + e^{-\alpha\lambda_{3,\alpha}})(1 - \alpha\lambda_{2,\alpha}) + \alpha(\lambda_{2,\alpha}e^{-\alpha\lambda_{2,\alpha}} + \lambda_{3,\alpha}e^{-\alpha\lambda_{3,\alpha}})] \\ &= \frac{1}{(1 + e^{-\alpha(\lambda_{3,\alpha} - \lambda_{2,\alpha})})^2} \left[(1 + e^{-\alpha(\lambda_{3,\alpha} - \lambda_{2,\alpha})})(1 - \alpha\lambda_{2,\alpha}) + \alpha\lambda_{2,\alpha} + \alpha\lambda_{3,\alpha}e^{-\alpha(\lambda_{3,\alpha} - \lambda_{2,\alpha})} \right] \\ &= \frac{1}{(1 + e^{-\alpha(\lambda_{3,\alpha} - \lambda_{2,\alpha})})^2} \left[1 + e^{-\alpha(\lambda_{3,\alpha} - \lambda_{2,\alpha})} + \alpha(\lambda_{3,\alpha} - \lambda_{2,\alpha})e^{-\alpha(\lambda_{3,\alpha} - \lambda_{2,\alpha})} \right] \\ &\xrightarrow{\alpha \rightarrow +\infty} \frac{1 + e^{-\eta} + \eta e^{-\eta}}{(1 + e^{-\eta})^2}, \end{aligned}$$

while the factor in front of $\nabla e_{3,\alpha} \otimes \nabla e_{3,\alpha}$ is asymptotically equivalent to $\frac{e^{-\eta}(1 + e^{-\eta} - \eta)}{(1 + e^{-\eta})^2}$. Therefore, one obtains the following characterization of the optimal diffusion matrix in the limit $\alpha \rightarrow +\infty$:

$$\mathcal{D}^{*,\infty}(q) = \gamma_\infty |\mathcal{D}^{*,\infty}(q)|_{\mathbb{F}}^{2-p} e^{(p-1)V} \left[\nabla e_{2,\infty} \otimes \nabla e_{2,\infty} + \frac{e^{-\eta}(1 + e^{-\eta} - \eta)}{1 + e^{-\eta} + \eta e^{-\eta}} \nabla e_{3,\infty} \otimes \nabla e_{3,\infty} \right], \quad (36)$$

with $\gamma_\infty \in \mathbb{R}$. In the case $d = 1$, and when the diffusion coefficient is positive, the formula (36) simplifies to

$$\mathcal{D}^{*,\infty}(q) = \tilde{\gamma}_\infty e^{V(q)} \left(|e'_{2,\infty}(q)|^2 + \frac{e^{-\eta}(1 + e^{-\eta} - \eta)}{1 + e^{-\eta} + \eta e^{-\eta}} |e'_{3,\infty}(q)|^2 \right)^{1/(p-1)}, \quad (37)$$

with $\tilde{\gamma}_\infty > 0$.

Formulas (36)-(37) suggest that, when the first nonzero eigenvalue $\lambda_{2,\infty}$ is degenerate, one should take into account all the eigenvectors associated with it, and reweight their contributions to obtain the analytical expression of the optimal diffusion matrix. The correctness of formula (37) is demonstrated on two numerical examples in Section 4.2, see Figures 1b and 2e below.

In the case $d = 1$, it is clear from (37) that if the value η satisfies $1 + e^{-\eta} - \eta = 0$, then the optimal coefficient vanishes where $e'_{2,\infty}$ vanishes. In that case, the (30) is not satisfied. However, we still believe that the formula (37) holds, from two numerical experiments in Section 4.2, see Figure 5 below. The particular value of η that satisfies $1 + e^{-\eta} - \eta = 0$ is

$$\eta^* = 1 + W\left(\frac{1}{e}\right) \approx 1.27846, \quad (38)$$

where W is the Lambert W function. In fact, a necessary condition (actually sufficient if $e'_{2,\infty}$ and $e'_{3,\infty}$ do not vanish at the same points) for (37) to define a positive definite diffusion coefficient is $\eta \in [0, \eta^*)$. In practice, we only observed numerical values of η in this range.

4 Numerical methods for optimizing the spectral gap

To solve the optimization problems (14) and (28) in practice, we introduce two finite-dimensional parametrizations: one for the diffusion matrix \mathcal{D} , and one for the eigenfunction u . We consider for simplicity the case of isotropic diffusion matrices

$$\mathcal{D}(q) = \mathcal{D}(q)\text{Id}_d. \quad (39)$$

In Section 4.1, we introduce the numerical methods we used to solve (14) and (28). We then present some numerical results obtained with our approach in Section 4.2.

4.1 Discretizations of the optimization problems (14) and (28)

We introduce a piecewise constant parametrization for \mathcal{D} in Section 4.1.1; and next a \mathbb{P}_1 finite element parametrization for the eigenfunctions u in Section 4.1.2, in the one dimensional case $d = 1$ for simplicity of exposition. The optimization problem then boils down to a constrained generalized eigenvalue problem, see Section 4.1.3. We describe in Section 4.1.4 how to solve the latter problem.

4.1.1 Parametrization of the diffusion matrix

In practice, we parametrize the scalar valued function \mathcal{D} in (39) using a finite-dimensional vector subspace of $L^p(\mathbb{T}^d, \mathbb{R})$. More precisely, we introduce a set of N non-negative functions $\{\psi_1, \dots, \psi_N\} \subset L^p(\mathbb{T}^d, \mathbb{R})$, and consider functions of the generic form

$$\mathcal{D}_D(q) = \sum_{n=1}^N D_n \psi_n(q), \quad D = (D_1, \dots, D_N) \in \mathbb{R}^N. \quad (40)$$

Nonnegativity conditions for \mathcal{D}_D may be cumbersome to write when the functions ψ_n have supports which overlap in a nontrivial way, as is the case for instance for (tensor products of) trigonometric functions. This motivates us to choose

$$\psi_n(q) = \mathbf{1}_{K_n}(q), \quad (41)$$

the indicator function of a domain $K_n \subset \mathbb{T}^p$, with the condition that (K_1, \dots, K_N) forms a partition of \mathbb{T}^p (typically, the sets K_n are rectangles obtained from a Cartesian mesh). In order for the function (40) to be in the set (12) for the choice (41), the components $(D_n)_{1 \leq n \leq N}$ should belong to the set

$$\delta_p^{a,b} = \left\{ D \in \left[ae^{V_+(K_1)}, \frac{1}{b} e^{V_-(K_1)} \right] \times \dots \times \left[ae^{V_+(K_N)}, \frac{1}{b} e^{V_-(K_N)} \right] \left| \sum_{n=1}^N \omega_{p,n} D_n^p \leq 1 \right. \right\}, \quad (42)$$

where

$$V_-(E) = \inf_E V, \quad V_+(E) = \sup_E V, \quad \omega_{p,n} = \int_{K_n} e^{-pV} |\text{Id}_d|_{\mathbb{F}}^p > 0.$$

We assume in the sequel that the conditions

$$ab \leq \min_{1 \leq n \leq N} e^{[V_-(K_n) - V_+(K_n)]}, \quad a \leq \left(\sum_{n=1}^N \omega_{p,n} e^{pV_+(K_n)} \right)^{-1/p}, \quad (43)$$

hold, so that the set $\delta_p^{a,b}$ is not empty. These conditions can be ensured by choosing $a, b \geq 0$ sufficiently small.

The maximization problem (14) can then be approximated by the finite dimensional maximization problem

$$\text{find } D^* \in \delta_p^{a,b} \text{ such that } \Lambda(\mathcal{D}_{D^*} \text{Id}_d) = \sup_{D \in \delta_p^{a,b}} \Lambda(\mathcal{D}_D \text{Id}_d). \quad (44)$$

The well-posedness of this optimization problem is guaranteed by the following result.

Proposition 8. Fix $N \geq 1$ and $p \in [1, +\infty)$, and consider a partition (K_1, \dots, K_N) of \mathbb{T}^d and $a, b \geq 0$ such that (43) holds. There exists a solution D^* to (44).

Note that the value $b = 0$ is allowed for N finite, in contrast to Theorem 4, since the inequality in (42) ensures that all the components $(D_n)_{1 \leq n \leq N}$ are bounded from above by $(\min_{1 \leq n \leq N} \omega_{p,n})^{-1/p}$.

Proof. The function $D \mapsto \Lambda(\mathcal{D}_D \text{Id}_d)$ is upper-semicontinuous on $\delta_p^{a,b}$ by arguments similar to the ones used in Appendix A.1 (and whatever the choice of the norm on \mathbb{R}^N). In addition, $\delta_p^{a,b}$ is closed and bounded, hence compact, which, combined with the upper-semicontinuity of the function to maximize, guarantees the existence of a maximum and thus proves the result. \square

Likewise, the maximization problem (28) can be approximated by the finite dimensional maximization problem

$$\text{find } D^{\star, \alpha} \in \delta_p^{a,b} \text{ such that } f_\alpha(\mathcal{D}_{D^{\star, \alpha}} \text{Id}_d) = \sup_{D \in \delta_p^{a,b}} f_\alpha(\mathcal{D}_D \text{Id}_d).$$

The well-posedness of this optimization problem can be obtained by similar arguments.

Notice that one can prove that any solution D^* of (44) is positive when the domains K_n have nonempty interiors (see Appendix A.2 for the proof).

Proposition 9. Fix $N \geq 1$ and $p \in [1, +\infty)$, and consider a partition (K_1, \dots, K_N) of \mathbb{T}^d for which the domains $(K_n)_{1 \leq n \leq N}$ have nonempty interiors. Assume that $a, b \geq 0$ satisfy (43), and denote by $D^* = (D_1^*, \dots, D_N^*) \in \delta_p^{a,b}$ a solution to the maximization problem (44). Then,

$$\forall n \in \{1, \dots, N\}, \quad D_n^* > 0.$$

Note however that this result does not give any information on the behavior of the lower bound on the components of D^* in the limit of a vanishing mesh size. The numerical results we present (see for instance Figure 3) indicate that the lower bound can indeed converge to 0 as $N \rightarrow +\infty$.

4.1.2 Finite element approximation of the eigenvalue problem

We discuss in this section how to numerically approximate $\Lambda(\mathcal{D})$ for a given diffusion matrix \mathcal{D} . For simplicity of exposition, we restrict the presentation to the one dimensional case $d = 1$, the extension to higher dimensional situations posing no difficulties in principle (but numerically viable only for $d \leq 3$ in terms of computational costs).

The minimization in (7) is performed over a linear space of dimension I obtained by a \mathbb{P}_1 finite element discretization over a regular mesh with nodes $q_i = i/I$ for $i \in \{0, \dots, I-1\}$ with $q_{-1} \equiv q_{I-1}$, $q_0 \equiv q_I$ and $q_1 \equiv q_{I+1}$ to comply with periodic boundary conditions. The continuous basis functions $(\varphi_i)_{i=1, \dots, I}$ have support on $[q_{i-1}, q_{i+1}]$ and are piecewise affine:

$$\varphi_i(q) = \begin{cases} I(q - q_{i-1}) & \text{if } q \in [q_{i-1}, q_i], \\ I(q_{i+1} - q) & \text{if } q \in [q_i, q_{i+1}]. \end{cases}$$

We denote by $S_I = \text{Span}(\varphi_1, \dots, \varphi_I)$, and by

$$S_{I,0} = \left\{ u \in S_I \left| \int_{\mathbb{T}} u(q) \mu(q) dq = 0 \right. \right\}.$$

We can then introduce the following mapping, which corresponds to a finite element approximation of $\Lambda(\mathcal{D}_D)$:

$$\lambda_{N,I}(D) = \inf_{u \in S_{I,0} \setminus \{0\}} \frac{\int_{\mathbb{T}} \mathcal{D}_D(q) u'(q)^2 \mu(q) dq}{\int_{\mathbb{T}} u(q)^2 \mu(q) dq}. \quad (45)$$

Let us recall the standard approach to solve (45). Upon writing $u = \sum_{i=1}^I U_i \varphi_i$, the optimization problem (45) can be rewritten as

$$\inf_{U \in \mathbb{R}^I \setminus \{0\}} \frac{U^\top A(D) U}{U^\top B U}, \quad A(D) = \sum_{n=1}^N D_n A_n,$$

where the matrices A_1, \dots, A_N and $B \in \mathbb{R}^{I \times I}$ have entries

$$[A_n]_{i,j} = \int_{K_n} \varphi'_j(q) \varphi'_i(q) e^{-V(q)} dq, \quad B_{i,j} = \int_{\mathbb{T}} \varphi_j(q) \varphi_i(q) e^{-V(q)} dq. \quad (46)$$

The Euler–Lagrange condition for the minimization problem (45) is therefore the following finite dimensional eigenvalue problem:

$$A(D)U = \sigma B U, \quad U^\top B U = 1. \quad (47)$$

Since $A(D)$ and B are real symmetric matrices and B is positive definite, the eigenvalue problem (47) admits $0 \leq \sigma_1(D) \leq \sigma_2(D) \leq \dots \leq \sigma_I(D)$ real and nonnegative eigenvalues (counted without multiplicities), with associated B -orthonormal eigenvectors $\{U_1(D), U_2(D), \dots, U_I(D)\}$. In addition, $\sigma_1(D) = 0$ and $U_1(D)$ is proportional to $(1, \dots, 1)$.

4.1.3 The discrete optimization problems

The optimization problem (14) is finally approximated by replacing $\Lambda(\mathcal{D}_D)$ in (44) by $\lambda_{N,I}(D)$: fix $p \in [1, +\infty)$. For any $a, b \geq 0$ such that (43) holds,

$$\text{find } D^* \in \delta_{a,b}^p \text{ such that } \sigma_2(D^*) = \sup_{D \in \delta_{a,b}^p} \sigma_2(D) = \lambda_{N,I}^*, \quad (48)$$

where $\sigma_2(D)$ is the first nonzero (second-smallest) eigenvalue of the problem (47).

The following proposition guarantees the existence of a minimizer to (48), which is moreover positive under some compatibility on the meshes used to discretize the diffusion coefficient and the eigenvalue problem.

Proposition 10. *Fix $p \in [1, +\infty)$. Consider $a, b \geq 0$ such that (43) holds. Then, there exists $D^* \in \delta_{a,b}^p$ such that*

$$\sigma_2(D^*) = \sup_{D \in \delta_{a,b}^p} \sigma_2(D) = \lambda_{N,I}^* > 0.$$

Assume moreover that, for all $n \in \{1, \dots, N\}$, there exists $i \in \{0, \dots, I-1\}$ such that $[q_{i-1}, q_{i+1}] \subset \overline{K_n}$. Then, any maximizer satisfies $\min_{1 \leq n \leq N} D_n^ > 0$.*

Proof. As already mentioned in the proof of Proposition 8, the set $\delta_{a,b}^p$ is compact. Thus, to prove the existence of a solution, it suffices to show that the application $D \mapsto \sigma_2(D)$ is upper-semicontinuous; in fact, we prove that it is continuous. Indeed, the application $M \in \mathcal{S}_I(\mathbb{R}) \mapsto \sigma_2(M)$ mapping any symmetric matrix to its second-smallest eigenvalue is continuous on $\mathcal{S}_I(\mathbb{R})$ (see, e.g., [11, Corollary III.2.6]). On the other hand, $D \mapsto A(D)$ is also continuous. By composition, $D \mapsto \sigma_2(D)$ is continuous on $\delta_{a,b}^p$, which proves the existence of a maximizer.

In addition, the Poincaré inequality (15) guarantees with the choice $D = \gamma \mathbb{1}_N$ (where $\mathbb{1}_N$ is the N -dimensional vector whose components are all equal to 1, and $\gamma > 0$ is a normalization constant chosen such that $D \in \delta_{a,b}^p$) that

$$\lambda_{N,I}^* \geq \sigma_2(\gamma \mathbb{1}_N) \geq \frac{\gamma}{C_\mu} > 0. \quad (49)$$

To prove that $D_n^* > 0$ for all $n \in \{1, \dots, N\}$, we proceed by contradiction. Assume that there exists $n \in \{1, \dots, N\}$ such that $D_n^* \leq 0$. Consider $i \in \{0, \dots, I-1\}$ such that $[q_{i-1}, q_{i+1}] \subset \overline{K_n}$. Setting $\phi_i = \varphi_i - \int_{\mathbb{T}} \varphi_i \mu$ (so that $\phi_i \in S_{I,0}$), we obtain

$$\sigma_2(D^*) \leq \frac{\int_{\mathbb{T}} D^*(q) |\nabla \phi_i(q)|^2 \mu(q) dq}{\int_{\mathbb{T}} \phi_i(q)^2 \mu(q) dq} = D_n^* \frac{\int_{K_n} |\nabla \phi_i(q)|^2 \mu(q) dq}{\int_{\mathbb{T}} \phi_i(q)^2 \mu(q) dq} \leq 0,$$

which contradicts (49) and thus proves the result. \square

The optimization problem (28) is approximated by: fix $p \in [1, +\infty)$. For any $a, b \geq 0$ such that (43) holds,

$$\text{find } D^{*,\alpha} \in \delta_{a,b}^p \text{ such that } F_\alpha(D^{*,\alpha}) = \sup_{D \in \delta_p^{a,b}} F_\alpha(D) = \lambda_{N,I}^{*,\alpha}, \quad (50)$$

where

$$F_\alpha(D) = \frac{H_\alpha(D)}{G_\alpha(D)}, \quad (51)$$

and

$$G_\alpha(D) = \sum_{i=2}^I e^{-\alpha \sigma_i(D)}, \quad H_\alpha(D) = \sum_{i=2}^I \sigma_i(D) e^{-\alpha \sigma_i(D)}. \quad (52)$$

The well-posedness of the optimization problem (50) (without the positivity of the optimizer) can be obtained by similar arguments. Using similar (and even simpler) arguments as the ones presented after (28), one can show that

$$\forall \alpha > 0, \quad \lambda_{N,I}^{*,\alpha} \geq \lambda_{N,I}^*,$$

and the map $\alpha \mapsto F_\alpha(D)$ is nonincreasing on \mathbb{R}_+^* . However, the monotonicity of the map $\alpha \mapsto \sigma_2(D^{*,\alpha})$ is not clear: it is observed numerically that this map is nondecreasing, see Figures 2c and 4b. Finally, let $(\alpha_n)_{n \geq 0}$ be a sequence of \mathbb{R}_+^* such that $\alpha_n \rightarrow +\infty$ when $n \rightarrow +\infty$. The sequence of maximizers $(D^{*,\alpha_n})_{n \geq 0}$ is bounded on $\delta_p^{a,b}$ so that, up to a subsequence (not indicated in the notations), there exists $D^{*,\infty} \in \delta_p^{a,b}$ such that

$$D^{*,\alpha_n} \xrightarrow{n \rightarrow +\infty} D^{*,\infty}.$$

By definition, it holds

$$F_{\alpha_n}(D^{*,\alpha_n}) = \lambda_{N,I}^{*,\alpha_n} \geq \lambda_{N,I}^* \geq \sigma_2(D^{*,\infty}).$$

By continuity of the maps $D \mapsto \sigma_i(D)$, it is then easy to show that $F_{\alpha_n}(D^{*,\alpha_n}) \rightarrow \sigma_2(D^{*,\infty})$, so that $\lambda_{N,I}^* = \sigma_2(D^{*,\infty})$ and $D^{*,\infty}$ solves (48). Moreover, $\lambda_{N,I}^{*,\alpha} \rightarrow \lambda_{N,I}^*$ as $\alpha \rightarrow +\infty$. The latter convergence is illustrated in Section 4.2.

In conclusion, solving (50) for large values of α should yield a tight upper bound on $\lambda_{N,I}^*$. In practice, solving (50) for small values of α already yields a good approximation in the situations we have considered.

4.1.4 Practical aspects of the implementation

We discuss in this section how to numerically solve (14), approximated as (48). The numerical solution of the smooth-min approximation (28), approximated as (50), is made precise in Appendix C.1.

To solve (48) in practice, we used either (i) Sequential Least Squares Quadratic Programming algorithm (SLSQP), which operates through linearization of the optimality conditions (see for instance [14, Section 15.1] for an introduction to such methods), relying on the implementation

available in SciPy [69]; or (ii) the IPOPT solver [70] within the Julia programming language, using the JuMP framework [27]. In order to use these methods, we need to compute the gradient of the objective function and the constraint, which we discuss next.

A first task is to approximate the integrals for the matrix elements in (46) and the weight factors $\omega_{p,n}$ in (42) by some quadrature method. In our numerical results, we set $N = I$ and $K_n = [(n-1)/N, n/N]$ for any $n \in \{1, \dots, N\}$, so that, for example, $[A_n]_{i,j}$ can be approximated by (with periodic boundary conditions for the indices)

$$N e^{-V(\frac{n-1}{N})} (\delta_{i,n} \delta_{i,j} + \delta_{i,n-1} \delta_{i,j} - \delta_{i,n} \delta_{i,j+1} - \delta_{i,n-1} \delta_{i,j-1}), \quad (53)$$

where δ is the Kronecker delta function, and $B_{i,j}$ can be approximated by

$$e^{-V(\frac{i-1}{N})} \frac{4\delta_{i,j} + \delta_{i,j-1} + \delta_{i,j+1}}{6N}. \quad (54)$$

We refer to Appendix C.2 for more details in order to obtain (53)-(54). Likewise, the weights $\omega_{p,n}$ can be approximated by

$$\frac{1}{N} e^{-pV(\frac{n-1}{N})}. \quad (55)$$

To run the optimization algorithms, we next need the gradient of the constraint

$$D \mapsto \sum_{n=1}^N \omega_{p,n} D_n^p,$$

which is readily computed. A more subtle point is to compute the gradient of the target function $\lambda_{N,I}(D) = \sigma_2(D)$ with the notation introduced in Section 4.1.2. It is readily computed assuming that the second eigenvalue $\sigma_2(D)$ is simple as the following classical result shows. Recall that the matrix B is symmetric positive definite, so that the matrix $B^{-1/2}$ is well-defined.

Lemma 11. *Consider $D^0 \in \mathbb{R}^N$ such that the second-smallest eigenvalue $\sigma_2(D^0)$ of the symmetric positive matrix $B^{-1/2} A(D^0) B^{-1/2}$ is simple. Then, $D \mapsto \sigma_2(D)$ is differentiable around D^0 , and the components of its gradient are*

$$\forall n \in \{1, \dots, N\}, \quad \frac{\partial \sigma_2}{\partial D_n}(D^0) = U(D^0)^\top A_n U(D^0), \quad (56)$$

where $U(D^0)$ is an eigenvector normalized as in (47).

Proof. The proof involves manipulations similar to the ones used to prove Lemma 5. When $\sigma_2(D^0)$ is simple, σ_2 is smooth in an open neighborhood of D^0 , and $D \mapsto U(D)$ can also be constructed to be smooth, so that, by taking the partial derivative with respect to D_n of the first condition in (47),

$$\frac{\partial A}{\partial D_n}(D) U(D) + A(D) \frac{\partial U}{\partial D_n}(D) = \frac{\partial \sigma_2}{\partial D_n}(D) B U(D) + \sigma_2(D) B \frac{\partial U}{\partial D_n}(D).$$

The desired result follows by multiplying this equality by $U(D)^\top$ on the left, using the normalization $U(D)^\top B U(D) = 1$ and that $U(D)^\top A(D) = \sigma_2(D) U(D)^\top B$ by transposing the relation (47) as the matrices $A(D)$ and B are real symmetric. \square

Using the approximation considered in (53) (recalling that $I = N$ in this case), the formula approximating the gradient (56) reads: for any $n \in \{1, \dots, N\}$,

$$\begin{aligned} \frac{\partial \sigma_2}{\partial D_n}(D) &\approx N e^{-V(\frac{n-1}{N})} (U(D)_{n-1}, U(D)_n)^\top \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} (U(D)_{n-1}, U(D)_n) \\ &= N e^{-V(\frac{n-1}{N})} [U(D)_n - U(D)_{n-1}]^2. \end{aligned} \quad (57)$$

Remark 5. Lemma 11 provides a formula for the gradient of σ_2 around points where this function is differentiable, namely for $D \in \mathbb{R}^N$ such that $\sigma_2(D)$ is simple. However, formula (56) makes sense even if $\sigma_2(D)$ is degenerate, although the resulting quantity is not necessarily the gradient of σ_2 in this situation. In practice, we use this formula for the gradient for any D , including if degeneracy happens. The downside is that the convergence may be slower around degeneracy points.

4.2 Numerical results

We apply the optimization procedure described in Section 4.1 to one-dimensional examples on the torus, with a normalization constraint corresponding to $p = 2$ and a mesh characterized by $N = I = 1000$ (recall that N is the dimension of the approximation space for the diffusion coefficient \mathcal{D} , and that I is the number of basis functions for the \mathbb{P}_1 finite element method). In particular, we choose $K_n = [(n-1)/N, n/N]$ for $n \in \{1, \dots, N\}$. Note that we used $N = I$ for simplicity in the implementation of A_n (see (53)) in order to compute the gradient of σ_2 (see Lemma 11) even though the in Proposition 10 to ensure positivity does not hold in that case. Similar results were obtained using $I = kN$ with $k \geq 2$ which is a sufficient condition for the positivity result to hold. The integrals for the matrix elements in (46) are approximated by (53)-(54), and the weights $\omega_{p,n}$ are approximated by (55). More details on the numerical approximations used are given in Appendix C. We relax the upper bound constraint on the diffusion coefficient by setting $b = 0$.

In this section, we name diffusion coefficients either the finite-dimensional vectors $D \in \mathbb{R}^N$ or the functions defined on \mathbb{T} by (40). In each figure, the target distribution is μ defined by (1); the optimal diffusion coefficient D^* is obtained numerically by solving (48), while the optimal diffusion coefficient $D^{*,\alpha}$ is obtained by solving (50); the constant diffusion coefficient is $D_{\text{cst}} = \gamma \mathbb{1}_N$ with

$$\gamma = \left(\sum_{n=1}^N \omega_{p,n} \right)^{-1/p},$$

so that $\sum_{n=1}^N \omega_{p,n} D_{\text{cst},n}^p = 1$; and the homogenized diffusion coefficient corresponds to the finite-dimensional representation of the proxy $\mathcal{D}_{\text{hom}}^*$ for the optimal diffusion coefficient obtained in a periodic homogenization limit, whose expression is given by (74) (see Section 5 for the presentation of the homogenized limit):

$$\forall n \in \{1, \dots, N\}, \quad D_{\text{hom},n}^* = \mathcal{D}_{\text{hom}}^* \left(\frac{n-1}{N} \right). \quad (58)$$


The initial condition for the diffusion coefficient in the optimization procedure solving either (48) or (50) is set to D_{hom}^* .

We denote by $D^{*,\infty}$ the numerical approximation of (37). This vector is defined componentwise as follows: for $n \in \{1, \dots, N\}$,

$$D_n^{*,\infty} = \tilde{\gamma}_\infty e^{V((n-1)/N)} \left(|U_{2,n} - U_{2,n-1}|^2 + \frac{e^{-\eta}(1 + e^{-\eta} - \eta)}{1 + e^{-\eta} + \eta e^{-\eta}} |U_{3,n} - U_{3,n-1}|^2 \right)^{1/(p-1)}, \quad (59)$$

where U_2 (respectively U_3) is a normalized eigenvector associated with $\sigma_2(D^*)$ (respectively $\sigma_3(D^*)$) and $\tilde{\gamma}_\infty$ is such that $\sum_{n=1}^N \omega_{p,n} (D_n^{*,\infty})^p = 1$. Note that periodic boundary conditions are imposed in (59), e.g. $U_{2,0} = U_{2,N}$. The scalar η is an approximation of η (see (37)), which is defined by $\eta = \lim_{\alpha \rightarrow +\infty} \alpha(\sigma_3(D^{*,\alpha}) - \sigma_2(D^{*,\alpha}))$, where $\sigma_2(D^{*,\alpha}) = \lambda_{N,I}^{*,\alpha}$ and $\sigma_3(D^{*,\alpha})$ are the leading two nonzero eigenvalues using the maximizer $D^{*,\alpha}$ solving (50).

We present the numerical results by first considering cases where $a = 0$ in Section 4.2.1, and then situations where the lower bound a is positive (see Section 4.2.2). The results for $a = 0$ are further distinguished depending on the value of the approximation of η among the cases $\eta = 0$, $\eta \in (0, \eta^*)$ and $\eta = \eta^*$. We finally draw conjectures from our numerical observations and discuss convergence issues in Section 4.2.3.



(a) Various diffusion coefficients and target distribution μ . The solid curve in red is the optimal diffusion coefficient D^* . (b) Comparison with the formal Euler-Lagrange characterizations (37) (approximated by (59)) for the case $\eta = 0$.

Figure 1: Results of the optimization procedure on the one-dimensional four-well potential $V(q) = \cos(8\pi q)$.

4.2.1 Results for $a = 0$

A case where $\eta = 0$. We plot in Figure 1a various diffusion coefficients as well as the target distribution for a four-well potential. We first observe that the optimal diffusion coefficient D^* takes larger values in regions where the target distribution is small. The analytical proxy D_{hom}^* has the same general shape as D^* . In fact, the two coefficients are almost superimposed. This observation is formalized in Section 5.3 below.

The numerical values of the spectral gaps obtained for D^* , D_{hom}^* and D_{cst} are respectively 30.24, 30.19 and 14.70. This simple example thus demonstrates the improvement obtained by optimizing the diffusion coefficient in terms of the spectral gap of the operator, and thus the enhanced convergence rate to equilibrium (further demonstrated by Monte Carlo simulations in Section 6). It also confirms the relevance of the homogenized proxy D_{hom}^* derived in Section 5, which can be computed explicitly and yields a spectral gap comparable to the one obtained through the optimization procedure.

The optimization procedure leads to an optimal diffusion coefficient which is uniformly lower bounded on the torus. In particular, we checked that the minimum of the optimal diffusion coefficient does not decrease as the mesh gets finer. A different behavior is observed in other numerical examples, see Figures 3a and 4a below.

The first three nonzero eigenvalues obtained for the optimal diffusion coefficient are $\sigma_2(D^*) = \lambda_{N,I}^* = 30.23813406$, $\sigma_3(D^*) = 30.23813407$, and $\sigma_4(D^*) = 86.06$. This means that the second eigenvalue can be considered degenerate, which is in accordance with Lemma 5 (we expect that both eigenvalues are equal in the limit $I, N \rightarrow +\infty$). Figure 1b displays the optimal diffusion coefficient D^* and the numerical approximation $D^{*,\infty}$ of (37) given by (59). We have observed that the two eigenvalues $\sigma_2(D^{*,\alpha})$ and $\sigma_3(D^{*,\alpha})$ are essentially the same (up to numerical precision) even for very small values of α , the values of $\alpha(\sigma_3(D^{*,\alpha}) - \sigma_2(D^{*,\alpha}))$ when $\alpha \in [1, 7]$ is at most of order 2.5×10^{-5} . We therefore consider that $\eta = 0$. With this choice, the two coefficients D^* and $D^{*,\infty}$ are superimposed, as predicted by (37). This confirms the relevance of the smooth-min approach used in Section 3.2.2.

A case where $\eta \in (0, \eta^*)$. We plot in Figure 2 the numerical results for a one-well potential using the value $\alpha = 1$. In Figure 2a, we plot the maximizer $D^{*,\alpha}$. The optimal diffusion coefficient D^* obtained by solving (48) is very similar, see Figures 2b and 2e). The numerical values of the spectral gaps obtained for $D^{*,\alpha}$, D_{hom}^* and D_{cst} are respectively 36.75, 32.43 and 30.47. Note that

the optimal spectral gap $\lambda_{N,I}^*$ is 36.88, and that $\lambda_{I,N}^{*,\alpha} = 36.94$. This shows that the map f_α is a good approximation of the objective function Λ , and that the diffusion coefficient obtained when solving (50) yields a spectral gap comparable to the one obtained when solving (48).

The diffusion coefficient $D^{*,\alpha}$ is uniformly lower bounded on \mathbb{T} . However, Lemma 5 does not apply since $D^{*,\alpha}$ is not optimal in the sense of maximizing the map Λ . Indeed, we check that the first three nonzero eigenvalues obtained for $D^{*,\alpha}$ are 36.75, 37.24 and 138.98. However, the optimal diffusion coefficient D^* obtained by solving (48) yields a degenerate eigenvalue (we obtain $\sigma_2(D^*) \approx \sigma_3(D^*) \approx 36.88$) which is in accordance with the result of Lemma 5. This means that checking for eigenvalue degeneracy when solving (50) does not yield information on the eigenvalue degeneracy of the optimal diffusion coefficient D^* . We check in Figure 2b that the characterization of the optimal diffusion coefficient using a finite value of α , *i.e.* formula (34), is valid, using similar approximations as for (59): more precisely, for $n \in \{1, \dots, N\}$, the n -th component of the discrete approximation of (34) is given by


$$\tilde{\gamma}_\alpha e^{V((n-1)/N)} \left(\sum_{k=2}^N \left[\frac{G_\alpha(D^{*,\alpha})(1 - \alpha\sigma_k(D^{*,\alpha})) + \alpha H_\alpha(D^{*,\alpha})}{G_\alpha(D^{*,\alpha})^2} e^{-\alpha\sigma_k(D^{*,\alpha})} \right] |U_{k,\alpha,n} - U_{k,\alpha,n-1}|^2 \right)^{1/(p-1)},$$

where G_α, H_α are defined in (52), $(U_{k,\alpha})_{2 \leq k \leq N}$ are the normalized eigenvectors associated with the eigenvalues $(\sigma_k(D^{*,\alpha}))_{2 \leq k \leq N}$ and $\tilde{\gamma}_\alpha$ is a normalizing constant. Note that this formula requires all the eigenvalues which can be computationally heavy.


We again seek to compute the value of η in order to compare the optimal diffusion with the formal Euler–Lagrange formula (37). We plot in Figure 2c the two leading nonzero eigenvalues using the maximizer $D^{*,\alpha}$ for increasing values of α . The behavior is very different from the previous case: the eigenvalues seem to converge to the same value but are quite different, even for rather large values of α . In Figure 2d, we plot the behavior of $\alpha(\sigma_3(D^{*,\alpha}) - \sigma_2(D^{*,\alpha}))$ as α increases, and fit the data with a function $\alpha \mapsto K/\alpha + \eta$, parameterized by $K, \eta \in \mathbb{R}$, using the method `curve_fit` from the `scipy.optimize` module. This gives a limiting value $\eta \approx -0.51$. The comparison between D^* and $D^{*,\infty}$ (as in (59)) is given in Figure 2e: the curves are superimposed, which again confirms that the formal discussion in Section 3.2.2 is relevant.

Cases where $\eta = \eta^*$. We plot in Figure 3a the numerical results for a double-well potential. We observe that the optimal diffusion coefficient displays two singularities at two points where it (almost) vanishes, around $q \approx 0.37$ and $q \approx 0.88$. We zoom around the point $q \approx 0.88$ in Figure 3b for various meshes and observe that, as the mesh gets finer ($I = N$ increases), the minimum value of the optimal diffusion coefficient gets smaller. The analytical proxy D_{hom}^* has the same general shape as D^* but displays no singularity, which may have numerical advantages. The numerical values of the spectral gaps obtained for D^* , D_{hom}^* and D_{cst} are respectively 11.23, 10.58 and 0.81. Other numerical properties of the optimal diffusion coefficient were difficult to obtain for this particular example, as the fact that the optimal diffusion almost vanishes makes the optimization either unstable or harder to converge, see Remark 6 below.


We present in Figure 4a the numerical results obtained for another double-well potential. As in the previous example, this diffusion coefficient almost vanishes and two singularities can be observed around $q \approx 0.25$ and $q \approx 0.75$. The numerical values of the spectral gap obtained for D^* , D_{hom}^* and D_{cst} are respectively 22.84, 21.18 and 8.46. The numerical convergence was easier to obtain for this example. In particular, we were able to compute the value of η by solving (50) for various values of α . We present in Figure 4b the behavior of the two leading nonzero eigenvalues for the values of α for which the optimization problems converged. We also fit the data similarly to what was done in Figure 2d to extract the limiting value $\eta \approx 1.276$, which is very close to the critical value η^* introduced in Section 3.2.2, see (38). This corroborates our claim around (38) that when $\eta = \eta^*$, the optimal diffusion vanishes at some points. We further back this statement by presenting in Figure 5 the results obtained with (37) (approximated by (59)) and $\eta = \eta^*$ (the value of η^* is approximated using the `lambertw` function of the `scipy.special` module). In particular, we see in Figure 5c that the diffusion given by the Euler–Lagrange formula with $\eta = \eta^*$ indeed vanishes.




(a) Various diffusion coefficients and target distribution μ . The solid curve in red is the optimal diffusion coefficient $D^{*,\alpha}$ with $\alpha = 1.0$.




(b) Comparison with the Euler-Lagrange equation (34) using $\alpha = 1.0$. The solid curve in red is the optimal diffusion coefficient D^* .



(c) Second and third eigenvalues associated with the maximizer $D^{*,\alpha}$ as a function of α .




(d) $\alpha(\sigma_3(D^{*,\alpha}) - \sigma_2(D^{*,\alpha}))$ as a function of α . The fit is given by $K/\alpha + \eta$ with $K \approx -0.027$ and $\eta \approx 0.51$.



(e) Comparison with the formal Euler-Lagrange characterizations (37) (approximated by (59)) for the case $\eta \in (0, \eta^*)$. The solid curve in red is the optimal diffusion coefficient D^* .

Figure 2: Results of the optimization procedure on the one-dimensional one-well potential $V(q) = \cos(2\pi q)$.



(a) Various diffusion coefficients and target distribution μ . The solid curve in red is the optimal diffusion coefficient D^* . (b) Zoom on the optimal diffusion coefficient D^* for increasing values of $I = N$.

Figure 3: Results of the optimization procedure on the one-dimensional double-well potential $V(q) = \sin(4\pi q)(2 + \sin(2\pi q))$.

Lower bound a	0.0	0.2	0.4	0.6	0.8	1.0
Spectral gap	11.227	11.226	11.208	11.145	10.983	10.572

Table 1: Spectral gap associated to optimal diffusion coefficients obtained with the optimization procedure using different lower bounds a in the same setting as Figure 6.

Lastly, note that the singularities observed in Figures 3b and 4a are consistent with the discussion in Remark 4. Indeed, when $\eta = \eta^*$, Equation (21) holds, and we thus expect the optimal diffusion to be non-smooth.

4.2.2 Results for $a > 0$

We now present results of the optimization procedure for the double-well potential energy function $V(q) = \sin(4\pi q)(2 + \sin(2\pi q))$ when imposing various lower bounds $a \in \{0, 0.2, 0.4, 0.6, 0.8, 1\}$. In Figure 6a, we plot the optimal diffusion coefficients. The case $a = 0$ has already been covered in Figure 3. It is the only one where the optimal diffusion coefficient almost vanishes on \mathbb{T} . When $a = 1$, the optimal diffusion coefficient is fully determined by the normalization constraint, and happens to coincide with the optimal diffusion coefficient in the homogenized limit D_{hom}^* . We highlight in Figure 6b how the lower bounds a affects the shape of the optimal diffusion coefficient: we plot the optimal diffusion coefficients divided by D_{hom}^* , and observe that the pointwise lower bound a is saturated on larger intervals as a increases to 1. We present in Table 1 the values of the spectral gap for the same values of the lower bound a . Note that the spectral gap for $a = 0.2$ is almost the same as the one for $a = 0$. This suggests that imposing a (small) positive lower bound may be useful in practice, especially in view of the interest of using a positive diffusion in Markov Chain Monte Carlo algorithms. This point is further discussed below in Section 6.2 (see Figure 10c).

4.2.3 Numerical conjectures and final remarks

In view of the results above, we conjecture that in all situations, in dimension 1, for the optimal diffusion coefficient:

- The first three nonzero eigenvalues associated with the operator $-\mathcal{L}_{\mathcal{D}^*}$ satisfy $\lambda_2 = \lambda_3 < \lambda_4$.




Figure 4: Results of the optimization procedure on the one-dimensional double-well potential $V(q) = \cos(4\pi q)$.



- The independent and normalized eigenvectors e_2 and e_3 (associated to $\lambda_2 = \lambda_3$) have derivatives that do not vanish at the same points (see (37)), which implies that $\eta \in [0, \eta^*]$, and there are then two situations:
 - either $\eta = \eta^*$, which is equivalent to the fact that the optimal diffusion vanishes at some points;
 - or $\eta < \eta^*$, which is equivalent to the fact that the optimal diffusion is uniformly positive.

It would be nice to prove such a conjecture and obtain conditions on V allowing to decide which situation is observed in practice. For example, we will see in the next section that when V is periodic with a period $1/k$ with an integer $k \in \mathbb{N} \setminus \{0\}$, the optimal diffusion converges rapidly to the optimal diffusion for the homogenized limit, which does not vanish.

Remark 6. *We have checked that all the numerical results reported in this section are converged up to numerical precision (with a convergence threshold of 10^{-15}). For some potentials, we were not able to get converged results, in the sense that the convergence threshold was not attained before the maximum number of iterations, set to 1000 in our examples. In some cases, the algorithm even stopped before the maximum number of iterations, because it was unable to find a descent direction compatible with the constraints. In practice, numerical convergence was harder to obtain when the optimal diffusion coefficient was almost zero at some points.*


5 Optimal diffusion matrix in the homogenized limit

A drawback of the optimization method described in Section 4 is the computational burden associated with the calculation of the optimal diffusion matrix in practice, which requires solving a high-dimensional eigenvalue problem. To obtain an approximation of the optimal diffusion, which can be used for instance as a good initial guess to the optimization procedure, we propose a simple model for the effective diffusion of the optimized overdamped Langevin dynamics. This model is obtained by studying the asymptotic behavior of the optimal diffusion matrix in the homogenized limit of periodic potentials with infinitely small spatial periods. The limit is explicit either for linear constraints, or for generic L_V^p constraints in the one dimensional case $d = 1$, and displays a simple dependence on the target density μ . The form of this dependence is backed up by prior results in the literature [61].




(a) $V(q) = \sin(4\pi q)(2 + \sin(2\pi q))$, using in (37) the value $\eta = \eta^*$ given by (38).

(b) $V(q) = \cos(4\pi q)$, using in (37) the value $\eta = \eta^*$ given by (38).




(c) Zoom around $q \approx 0.75$ for $V(q) = \cos(4\pi q)$, displaying the coefficient given by (37) with $\eta = \eta^*$ and $\eta = \eta$.

Figure 5: Comparison with the formal Euler-Lagrange characterizations (37) (approximated by (59)) for the cases $\eta = \eta^*$. The solid curves in red are the optimal diffusion coefficients D^* .



(a) Optimal diffusion coefficients for various lower bounds a .



(b) Optimal diffusion coefficients normalized by D_{hom}^* .

Figure 6: Results of the optimization procedure for various lower bounds a and $V(q) = \sin(4\pi q)(2 + \sin(2\pi q))$.

In order to present our results, we start by introducing in Section 5.1 the setting of homogenization theory, in particular the notion of H -convergence, and give some useful and standard results, adapted to our framework of periodic boundary conditions. Next, in Section 5.2, we study the homogenized limit of the diffusion matrices and then explicitly optimize the spectral gap of the homogenized limit for one-dimensional systems or linear constraints. Finally, in Section 5.3, we study the commutation of the homogenization and optimization procedures, by showing that the diffusion matrix obtained by optimizing first and then taking the homogenized limit coincides with the optimal diffusion matrix for the homogenized problem.

5.1 Definitions and compactness results

We present in this section compactness results for sequences of matrices $(\mathcal{A}^k)_{k \geq 1} \subset L^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$. Let us emphasize that we restrict ourselves to $a, b > 0$ in all this section. We start by recalling the notion of H -convergence in the setting of \mathbb{Z}^d -periodic functions. Compared to the classical H -convergence presented in the literature (see for instance [2, Chapter 1] and references therein, as well as [13]), the main difference is that we use periodic boundary conditions rather than Dirichlet boundary conditions.

Definition 1 (H -convergence). *Fix $a, b > 0$, and consider a sequence $(\mathcal{A}^k)_{k \geq 1} \subset L^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$. The sequence $(\mathcal{A}^k)_{k \geq 1}$ is said to H -converge to $\bar{\mathcal{A}} \in L^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$ (denoted by $\mathcal{A}^k \xrightarrow{H} \bar{\mathcal{A}}$) if, for any $f \in H^{-1}(\mathbb{T}^d)$ such that $\langle f, \mathbf{1} \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)} = 0$, the sequence $(u^k)_{k \geq 1} \subset H^1(\mathbb{T}^d)$ of solutions to*

$$\begin{cases} -\operatorname{div}(\mathcal{A}^k \nabla u^k) = f & \text{on } \mathbb{T}^d, \\ \int_{\mathbb{T}^d} u^k(q) dq = 0, \end{cases} \quad (60)$$

satisfies, in the limit $k \rightarrow \infty$,

$$\begin{cases} u^k \rightharpoonup u \text{ weakly in } H^1(\mathbb{T}^d), \\ \mathcal{A}^k \nabla u^k \rightharpoonup \bar{\mathcal{A}} \nabla u \text{ weakly in } L^2(\mathbb{T}^d)^d, \end{cases}$$

where $u \in H^1(\mathbb{T}^d)$ is the solution of the homogenized problem

$$\begin{cases} -\operatorname{div}(\bar{\mathcal{A}} \nabla u) = f & \text{on } \mathbb{T}^d, \\ \int_{\mathbb{T}^d} u(q) dq = 0. \end{cases} \quad (61)$$

Notice that the functions u^k in (60) and u in (61) are indeed well-defined since $\langle f, \mathbf{1} \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)} = 0$ (in application of the Lax–Milgram theorem, see Lemma 27 in Appendix D.2). The next theorem, proved for completeness in Appendix D.1, states a compactness result for sequences of periodic functions.

Theorem 12 (Murat–Tartar). *Fix $a, b > 0$, and consider a sequence $(\mathcal{A}^k)_{k \geq 1} \subset L^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$. Up to extraction of a subsequence, $(\mathcal{A}^k)_{k \geq 1}$ H -converges to a matrix $\bar{\mathcal{A}} \in L^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$ (non constant in general).*

The following theorem, adapted from [2, Theorem 1.3.18] (and proved for completeness in Appendix D.4.1, see Remark 9), states an H -convergence result for sequences of matrices defined through periodic homogenization. It is a particular case of Theorem 12 specific to periodic functions.

Theorem 13. *Fix $a, b > 0$, and let $\mathcal{A} \in L^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$. For $k \geq 1$, consider the sequence $(\mathcal{A}_{\#,k})_{k \geq 1}$ defined as*

$$\forall q \in \mathbb{T}^d, \quad \mathcal{A}_{\#,k}(q) = \mathcal{A}(kq).$$

The sequence $(\mathcal{A}_{\#,k})_{k \geq 1}$ H -converges to the constant matrix $\bar{\mathcal{A}} \in \mathcal{M}_{a,b}$ with entries

$$\forall 1 \leq i, j \leq d, \quad \bar{\mathcal{A}}_{ij} = \int_{\mathbb{T}^d} (e_i + \nabla w_i(q))^\top \mathcal{A}(q) (e_j + \nabla w_j(q)) dq, \quad (62)$$

where (e_1, \dots, e_d) is the canonical basis of \mathbb{R}^d and $\{w_i\}_{1 \leq i \leq d} \subset H^1(\mathbb{T}^d)$ is the family of unique solutions to the problems

$$\begin{cases} -\operatorname{div}(\mathcal{A}(e_i + \nabla w_i)) = 0, \\ \int_{\mathbb{T}^d} w_i(q) dq = 0. \end{cases} \quad (63)$$

The last theorem, proved in Appendix D.2 by a straightforward adaptation of [2, Theorem 1.3.16], states that H -convergence implies the convergence of the sequence of spectral gaps.

Theorem 14. Fix $a, b > 0$ and let $(\mathcal{A}^k)_{k \geq 1} \subset L^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$ be a sequence H -converging to a homogenized matrix $\bar{\mathcal{A}} \in L^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$. Let $(\rho^k)_{k \geq 1} \subset L^\infty(\mathbb{T}^d)$ be a sequence of positive functions converging weakly-* in $L^\infty(\mathbb{T}^d)$ to a limiting function ρ , and such that there exists $\rho_-, \rho_+ > 0$ for which $0 < \rho_- \leq \rho^k(q) \leq \rho_+ < +\infty$ for all $k \geq 1$ and for almost all $q \in \mathbb{T}^d$. Let λ^k be the smallest nonzero eigenvalue and u^k an associated normalized eigenvector of the spectral problem

$$\begin{cases} -\operatorname{div}(\mathcal{A}^k(q) \nabla u^k(q)) = \lambda^k \rho^k(q) u^k(q), \\ \int_{\mathbb{T}^d} u^k(q)^2 dq = 1. \end{cases}$$

Then,

$$\lim_{k \rightarrow +\infty} \lambda^k = \bar{\lambda}$$

and, up to a subsequence, $(u^k)_{k \geq 1}$ converges weakly in $H^1(\mathbb{T}^d)$ to u , where $(\bar{\lambda}, u)$ is an eigenpair of the homogenized operator:

$$\begin{cases} -\operatorname{div}(\bar{\mathcal{A}} \nabla u(q)) = \bar{\lambda} \rho(q) u(q), \\ \int_{\mathbb{T}^d} u(q)^2 dq = 1, \end{cases}$$

and $\bar{\lambda}$ is the smallest nonzero eigenvalue of this homogenized eigenvalue problem.

5.2 Optimization of the periodic homogenization limit

We still fix $a, b > 0$, and apply the results of Section 5.1 to a particular class of sequences of matrix valued-functions $(\mathcal{A}_{\#,k})_{k \geq 1}$, defined for $\mathcal{D} \in \mathfrak{D}_p^{a,b}$ by decreasing the period of the \mathbb{Z}^d -periodic matrix

$$\mathcal{A}(q) = \mathcal{D}(q) \exp(-V(q)) \quad (64)$$

to obtain the $(\mathbb{Z}/k)^d$ -periodic function

$$\mathcal{A}_{\#,k}(q) = \mathcal{D}_{\#,k}(q) \exp(-V_{\#,k}(q)), \quad \mathcal{D}_{\#,k}(q) = \mathcal{D}(kq), \quad V_{\#,k}(q) = V(kq). \quad (65)$$

Note that $\mathcal{A}_{\#,k}(q) \in \mathcal{M}_{a,b}$ for almost every $q \in \mathbb{T}^d$ and for all $k \geq 1$. The periodization procedure (65) allows us to prove in Section 5.2.1 the H -convergence of the sequence of matrices, and the convergence of the associated eigenvalues and eigenvectors (see Corollary 15). We then provide in Section 5.2.2 explicit expressions for diffusion matrices \mathcal{D} which maximize the spectral gap of the homogenized problem; see in particular Theorem 17 for the one dimensional case $d = 1$.

5.2.1 Periodic homogenization of the spectral gap problem

We consider the process $(q_t^k)_{t \geq 0}$ defined on \mathbb{T}^d by the dynamics

$$dq_t^k = [-\mathcal{D}_{\#,k}(q_t^k) \nabla V_{\#,k}(q_t^k) + \operatorname{div}(\mathcal{D}_{\#,k}(q_t^k))] dt + \sqrt{2} \mathcal{D}_{\#,k}^{1/2}(q_t^k) dW_t.$$

The generator associated to the dynamics acts on test functions $u : \mathbb{T}^d \rightarrow \mathbb{R}$ as

$$\mathcal{L}_k u = \exp(V_{\#,k}) \operatorname{div} [\mathcal{D}_{\#,k} \exp(-V_{\#,k}) \nabla u].$$

Since the eigenfunctions associated with the first eigenvalue 0 are constant functions, the Rayleigh–Ritz principle implies that the spectral gap of \mathcal{L}_k considered as an operator on $L^2(Z^{-1}e^{-V_{\#,k}})$ (recall that Z is defined in (1)) writes (recalling the notation (65)):

$$\Lambda_{\#,k}(\mathcal{D}) = \min_{u \in H^1(\mathbb{T}^d) \setminus \{0\}} \left\{ \frac{\int_{\mathbb{T}^d} \nabla u(q)^\top \mathcal{A}_{\#,k}(q) \nabla u(q) dq}{\int_{\mathbb{T}^d} u^2(q) e^{-V_{\#,k}(q)} dq} \mid \int_{\mathbb{T}^d} u e^{-V_{\#,k}} = 0 \right\}. \quad (66)$$

Let us emphasize that the test functions to find the minimum on the right-hand side of the previous equality are \mathbb{Z}^d -periodic, while the diffusion and the potential are $(\mathbb{Z}/k)^d$ -periodic. Note also that the minimization over the functions u is performed on $H^1(\mathbb{T}^d)$, which is equivalent in fact to the functional space $H^1(Z^{-1}e^{-V_{\#,k}})$ since \mathbb{T}^d is compact and V is smooth. This allows to work with a fixed functional space for the test functions u whatever the value of k .

The spectral gap $\Lambda_{\#,k}(\mathcal{D})$ is the smallest nonzero solution λ^k of the following eigenvalue problem posed on \mathbb{T}^d : find $(u^k, \lambda^k) \in H^1(\mathbb{T}^d) \setminus \{0\} \times (0, +\infty)$ such that

$$-\operatorname{div} [\mathcal{D}_{\#,k} \exp(-V_{\#,k}) \nabla u^k] = \lambda^k \exp(-V_{\#,k}) u^k, \quad \int_{\mathbb{T}^d} u^k e^{-V_{\#,k}} = 0.$$

By applying Theorem 14 to the sequence $(\mathcal{A}_{\#,k})_{k \geq 1}$, we therefore obtain the following result on the convergence of the sequence of spectral gaps $(\Lambda_{\#,k}(\mathcal{D}))_{k \geq 1}$ for a given diffusion $\mathcal{D} \in \mathfrak{D}_p^{a,b}$.

Corollary 15. *Fix $p \in [1, +\infty)$, $a, b > 0$ satisfying (13) and $\mathcal{D} \in \mathfrak{D}_p^{a,b}$. Then, $\mathcal{A}_{\#,k} \xrightarrow{H} \overline{\mathcal{A}}$ with $\overline{\mathcal{A}}$ defined in (62). Defining $\overline{\mathcal{D}} = Z^{-1} \overline{\mathcal{A}}$, the sequence $(\Lambda_{\#,k}(\mathcal{D}))_{k \geq 1}$ converges as $k \rightarrow \infty$ towards*

$$\Lambda_{\text{hom}}(\mathcal{D}) := \inf_{u \in H_0^1(\mathbb{T}^d) \setminus \{0\}} \frac{\int_{\mathbb{T}^d} \nabla u(q)^\top \overline{\mathcal{D}} \nabla u(q) dq}{\int_{\mathbb{T}^d} u^2(q) dq}.$$

In the next section, we seek to maximize the spectral gap $\Lambda_{\text{hom}}(\mathcal{D})$ with respect to the baseline diffusion matrix \mathcal{D} from which $\overline{\mathcal{D}}$ arises as the H -limit of $\mathcal{D}_{\#,k}(q) \exp(-V_{\#,k}(q))/Z$. There is of course a huge body of literature on optimizing homogenized coefficients, for example in the context of shape optimization of materials, typically by considering laminated structures which would correspond here to situations where the diffusion can only take two values; see for instance [65, 35, 9, 4] and [38, Chapter 10].

It is useful to provide a more explicit expression of the limiting matrix $\overline{\mathcal{D}}$ in order to maximize $\Lambda_{\text{hom}}(\mathcal{D})$. Consider $\xi = \xi_1 e_1 + \dots + \xi_d e_d \in \mathbb{R}^d$ and note that

$$\xi^\top \overline{\mathcal{D}} \xi = Z^{-1} \int_{\mathbb{T}^d} (\xi + \nabla w_\xi(q))^\top \mathcal{A}(q) (\xi + \nabla w_\xi(q)) dq,$$

where $w_\xi = \xi_1 w_1 + \dots + \xi_d w_d$ and the family $(w_i)_{1 \leq i \leq d}$ is defined in (63). Recalling (64), it follows that

$$\begin{aligned} \xi^\top \overline{\mathcal{D}} \xi &= \int_{\mathbb{T}^d} (\xi + \nabla w_\xi(q))^\top \mathcal{D}(q) (\xi + \nabla w_\xi(q)) \mu(q) dq \\ &= \int_{\mathbb{T}^d} (\xi^\top \mathcal{D}(q) \xi + 2\xi^\top \mathcal{D}(q) \nabla w_\xi(q) + \nabla w_\xi(q)^\top \mathcal{D}(q) \nabla w_\xi(q)) \mu(q) dq. \end{aligned} \quad (67)$$

By the weak formulation of (63), it holds, for any $v \in H^1(\mathbb{T}^d)$,

$$\int_{\mathbb{T}^d} \xi^\top \mathcal{D}(q) \nabla v(q) \mu(q) dq = - \int_{\mathbb{T}^d} \nabla w_\xi(q)^\top \mathcal{D}(q) \nabla v(q) \mu(q) dq. \quad (68)$$

Upon replacing v by w_ξ , we obtain with (67) that

$$\xi^\top \overline{\mathcal{D}} \xi = \xi^\top \left(\int_{\mathbb{T}^d} \mathcal{D}(q) \mu(q) dq \right) \xi - \int_{\mathbb{T}^d} \nabla w_\xi(q)^\top \mathcal{D}(q) \nabla w_\xi(q) \mu(q) dq. \quad (69)$$

5.2.2 Optimization of the homogenized limit

We now focus on the optimization with respect to $\mathcal{D} \in \mathfrak{D}_p^{a,b}$ of the homogenized limit $\Lambda_{\text{hom}}(\mathcal{D})$ given in Corollary 15, under some normalization constraint:

$$\Lambda_{\text{hom}}^* = \sup_{\mathcal{D} \in \mathfrak{D}_p^{a,b}} \Lambda_{\text{hom}}(\mathcal{D}). \quad (70)$$

This optimization problem is well posed by arguments similar to the ones used in Section 3.1, and using the fact that the mapping $\mathcal{D} \mapsto \overline{\mathcal{D}}$ in (69) is continuous on $\mathfrak{D}_p^{a,b}$ (relying on the continuity of the mapping $L^\infty(\mathbb{T}^d, \mathcal{M}_{a,b}) \ni \mathcal{A} \mapsto w_i \in H^1(\mathbb{T}^d)$ where w_i is the solution of (63)). In the most general case, one of the difficulties is to understand how the constraint $\mathcal{D} \in \mathfrak{D}_p^{a,b}$ homogenizes into a constraint on $\overline{\mathcal{D}}$. We start by analyzing a specific example of constrained set with linear constraints, which is not the constraint $\|\mathcal{D}\|_{L_V^1} \leq 1$ with the norm defined in (11). The interest of the linear constraint we consider is that the optimization problem becomes trivial. We next derive a similar result for the constraint $\|\mathcal{D}\|_{L_V^p} \leq 1$, but for the one-dimensional case $d = 1$. Both results hold provided $a, b > 0$ are sufficiently small so that the optimal diffusion matrix automatically satisfies the pointwise lower and upper bounds involving a and b .

Linear constraint. We start by considering the following constraint on \mathcal{D} , for a given constant matrix $M \in \mathcal{S}_d^{++}$ (with \mathcal{S}_d^{++} the set of real, symmetric, positive, definite matrices of sizes $d \times d$):

$$\int_{\mathbb{T}^d} \mathcal{D}(q) \mu(q) dq = M. \quad (71)$$

Lemma 16. *Consider $M \in \mathcal{S}_d^{++}$, with $M_-, M_+ > 0$ such that $M_- \text{Id}_d \leq M \leq M_+^{-1} \text{Id}_d$. Fix $a, b > 0$ such that*

$$0 < a \leq M_- Z, \quad 0 < b \leq \frac{1}{M_+ Z}. \quad (72)$$

Then, the diffusion $\mathcal{D}_{\text{hom}}^(q) = M/\mu(q)$ is a solution to the optimization problem*

$$\max \left\{ \Lambda_{\text{hom}}(\mathcal{D}) \mid \mathcal{D} \in L_V^\infty(\mathbb{T}^d, \mathcal{M}_{a,b}), \int_{\mathbb{T}^d} \mathcal{D}(q) \mu(q) dq = M \right\}.$$

Proof. In view of (69), for any $\mathcal{D} \in L_V^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$ satisfying (71), it holds

$$\forall \xi \in \mathbb{R}^d, \quad \xi^\top \overline{\mathcal{D}} \xi \leq \xi^\top \left(\int_{\mathbb{T}^d} \mathcal{D}(q) \mu(q) dq \right) \xi = \xi^\top M \xi.$$

Thus, for any $\mathcal{D} \in L_V^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$ satisfying the constraint (71),

$$\inf_{u \in H_0^1(\mathbb{T}^d) \setminus \{0\}} \frac{\int_{\mathbb{T}^d} \nabla u(q)^\top \overline{\mathcal{D}} \nabla u(q) dq}{\int_{\mathbb{T}^d} u^2(q) dq} \leq \inf_{u \in H_0^1(\mathbb{T}^d) \setminus \{0\}} \frac{\int_{\mathbb{T}^d} \nabla u(q)^\top M \nabla u(q) dq}{\int_{\mathbb{T}^d} u^2(q) dq}.$$

Now, note that $\mathcal{D}(q) = M/\mu(q)$ belongs to $L_V^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$ in view of the choice (72), and satisfies $\overline{\mathcal{D}} = M$ in view of (69) since $w_\xi = 0$ for any $\xi \in \mathbb{R}^d$ by (63). This diffusion matrix therefore maximizes the limiting spectral gap in view of the above inequality, which concludes the proof. \square

L_V^p constraint in the one-dimensional case. We now restrict ourselves to the case where $d = 1$. In this situation, $\Lambda_{\text{hom}}(\mathcal{D})$ is proportional to the nonnegative real number $\overline{\mathcal{D}}$. Thus, maximizing $\Lambda_{\text{hom}}(\mathcal{D})$ boils down to maximizing $\overline{\mathcal{D}} \geq 0$. In the one-dimensional case, denoting w_ξ as $w_{\mathcal{D}}$ (we omit the irrelevant parameter ξ , but make the dependence on \mathcal{D} explicit), equation (63) on $w_{\mathcal{D}}$ can be rewritten as

$$[e^{-V} \mathcal{D}(1 + w'_{\mathcal{D}})]' = 0, \quad (73)$$

and the expression (69) simplifies as

$$\overline{\mathcal{D}} = \int_{\mathbb{T}^d} \mathcal{D}(q) (1 - w'_{\mathcal{D}}(q)^2) \mu(q) dq,$$

while the normalization constraint $\|\mathcal{D}\|_{L_V^p} \leq 1$ simply reads

$$\int_{\mathbb{T}^d} \mathcal{D}(q)^p e^{-pV(q)} dq \leq 1.$$

The optimization problem (70) is therefore equivalent to

$$\max_{\mathcal{D} \in L_V^p(\mathbb{T}, \mathbb{R}_+)} \left\{ \int_{\mathbb{T}^d} \mathcal{D}(q) (1 - w'_{\mathcal{D}}(q)^2) \mu(q) dq \mid a \leq \mathcal{D}e^{-V} \leq b^{-1}, \int_{\mathbb{T}^d} \mathcal{D}(q)^p e^{-pV(q)} dq \leq 1 \right\}.$$

This optimization problem admits an explicit solution provided $a, b \leq 1$, as made precise in the following result.

Theorem 17. *Fix $p \in [1, +\infty)$, and consider $a, b \in (0, 1]$. Then, a maximizer for the spectral gap in the homogenized limit (70) is*

$$\mathcal{D}_{\text{hom}}^*(q) = e^{V(q)}. \quad (74)$$

Note that the maximizer $\mathcal{D}_{\text{hom}}^*$ is independent of p due to our specific choice of normalization, and that no extra normalization factor is needed.

Proof. Equation (73) can be integrated as $w'_{\mathcal{D}} + 1 = K\mathcal{D}^{-1}e^V$ for some constant $K \in \mathbb{R}$, so that (up to an irrelevant additive constant, set to 0 for simplicity)

$$w_{\mathcal{D}}(q) = -q + K \int_0^q \frac{e^V}{\mathcal{D}}.$$

The constant K is chosen to ensure the periodicity of the function. This gives

$$w_{\mathcal{D}}(q) = -q + \left(\int_{\mathbb{T}} \frac{e^V}{\mathcal{D}} \right)^{-1} \int_0^q \frac{e^V}{\mathcal{D}},$$

so that, by (69) and (68) (with $\xi = 1$ in this one-dimensional context, and $v = w_\xi = w_{\mathcal{D}}$),

$$\overline{\mathcal{D}} = \int_{\mathbb{T}} \mathcal{D} [1 - (w'_{\mathcal{D}})^2] \mu = \int_{\mathbb{T}} \mathcal{D} (1 + w'_{\mathcal{D}}) \mu = \left(\int_{\mathbb{T}} e^{-V} \right)^{-1} \left(\int_{\mathbb{T}} \mathcal{D}^{-1} e^V \right)^{-1}. \quad (75)$$

The maximization of the effective diffusion $\overline{\mathcal{D}}$ with respect to \mathcal{D} therefore amounts to solving the following optimization problem:

$$\min \left\{ \int_{\mathbb{T}} \mathcal{D}^{-1} e^V \mid a \leq \mathcal{D}e^{-V} \leq b^{-1}, \int_{\mathbb{T}} \mathcal{D}(q)^p e^{-pV(q)} dq \leq 1 \right\}.$$

The Euler-Lagrange equation associated with the minimization problem with the normalization constraint only (forgetting the pointwise upper and lower bounds), assuming that the constraint is saturated at its value 1, implies that $\mathcal{D}^{-2}e^V$ is proportional to $\mathcal{D}^{p-1}e^{-pV}$, i.e. \mathcal{D} is proportional to e^V ; and in fact equal to the latter function in view of the normalization constraint. The pointwise upper and lower bounds are then automatically satisfied given our choice for a, b , which allows to conclude the proof. \square

The extension of this result to the multi-dimensional case $d \geq 1$ is not as clear and we leave this open question for future work.

5.3 Homogenization of the optimal diffusion

We consider in this section the H -limit of diffusion matrices which maximize the spectral gap for potentials with decreasing spatial periodicity. This corresponds to first performing the optimization of the spectral gap, then the homogenization of the associated maximizers; whereas the analysis of Section 5.2.1 relied on first homogenizing the spectral gap problem, then optimizing it in Section 5.2.2.

For $k \geq 1$, define the constrained set associated with the oscillating potential $V_{\#,k}$:

$$\mathfrak{D}_{\#,k,p}^{a,b} = \left\{ \mathcal{D} \in L_{V_{\#,k}}^\infty(\mathbb{T}^d, \mathcal{M}_{a,b}) \left| \int_{\mathbb{T}^d} |\mathcal{D}(q)|_{\mathbb{F}}^p e^{-pV_{\#,k}(q)} dq \leq 1 \right. \right\}.$$

Note that, if $\mathcal{D} \in \mathfrak{D}_p^{a,b}$, then $\mathcal{D}_{\#,k} \in \mathfrak{D}_{\#,k,p}^{a,b}$, where we recall that $\mathcal{D}_{\#,k}$ is the periodized diffusion $\mathcal{D}_{\#,k}(q) = \mathcal{D}(kq)$. Of course, the set $\mathfrak{D}_{\#,k,p}^{a,b}$ contains more functions than those obtained by periodization. Let us indeed emphasize that the diffusion matrices in $\mathfrak{D}_{\#,k,p}^{a,b}$ are a priori only \mathbb{Z}^d -periodic. For $\mathcal{D} \in \mathfrak{D}_{\#,k,p}^{a,b}$ a \mathbb{Z}^d -periodic diffusion, consider the spectral gap associated with the oscillating potential $V_{\#,k}$:

$$\Lambda^k(\mathcal{D}) = \min_{u \in H^1(\mathbb{T}^d) \setminus \{0\}} \left\{ \frac{\int_{\mathbb{T}^d} \nabla u(q)^\top \mathcal{D}(q) \nabla u(q) e^{-V_{\#,k}(q)} dq}{\int_{\mathbb{T}^d} u^2(q) e^{-V_{\#,k}(q)} dq} \left| \int_{\mathbb{T}^d} u e^{-V_{\#,k}} = 0 \right. \right\}. \quad (76)$$

The optimal diffusion matrix corresponding to (76) satisfies the following optimization problem:

$$\Lambda^{k,\star} = \max_{\mathcal{D} \in \mathfrak{D}_{\#,k,p}^{a,b}} \Lambda^k(\mathcal{D}). \quad (77)$$

Note that the minimization problem in (76) is different from the one in (66) since the diffusion matrix \mathcal{D} is not $(\mathbb{Z}/k)^d$ -periodic, but only \mathbb{Z}^d -periodic a priori. It turns out however that the optimization problem (77) admits a $(\mathbb{Z}/k)^d$ -periodic solution, as made precise in the following lemma, proved in Appendix D.3.

Lemma 18. *Fix $p \in [1, +\infty)$ and consider $a \in [0, a_{\max}]$ and $b > 0$ such that $ab \leq 1$. For $k \in \mathbb{N}_+$, there exists $\mathcal{D}^{k,\star} \in \mathfrak{D}_p^{a,b}$ such that, denoting by $\mathcal{D}_{\#,k}^{k,\star}(q) = \mathcal{D}^{k,\star}(kq)$,*

$$\Lambda^k(\mathcal{D}_{\#,k}^{k,\star}) = \Lambda_{\#,k}(\mathcal{D}^{k,\star}) = \Lambda^{k,\star}.$$

Lemma 18 shows that, while solving the optimization problem (77) (well posed in application of Theorem 4), we may restrict ourselves to $(\mathbb{Z}/k)^d$ -periodic diffusion matrices. Theorem 19 below, proved in Appendix D.4, states that the limit $\bar{\Lambda}^\star$ when $k \rightarrow +\infty$ of the sequence $(\Lambda^{k,\star})_{k \geq 1}$ is in fact equal to the limit $\Lambda_{\text{hom}}^\star$ defined in (70), showing the commutation of homogenization and optimization procedures.

Theorem 19. *Fix $p \in [1, +\infty)$ and consider $a \in (0, a_{\max}]$ and $b > 0$ such that $ab \leq 1$. Then the sequence $(\Lambda^{k,\star})_{k \geq 1}$ converges to $\Lambda_{\text{hom}}^\star$ as $k \rightarrow \infty$.*

Theorem 19 suggests that, in the homogenized limit, the optimal diffusion matrix yields the same spectral gap as its proxy obtained by solving (70) – which, we recall, can be analytically written out in the simple cases discussed in Section 5.2.2.

Numerical results. We present two one-dimensional numerical examples illustrating the convergence towards the homogenized regime when the frequency k goes to infinity. The space discretizations of the optimization problem (77), the optimization algorithms, the notation and the hyperparameters are the same as in Section 4. The mesh size used to obtain these results is refined as k increases, using $N = I = 200k$. Denote by $D^{k,\star}$ the maximizer obtained by solving




Figure 7: Comparison of the two procedures: homogenize then optimize, versus optimize then homogenize. The left and right columns correspond to two different target distributions. The figures in the second row illustrate the convergence of the sequence $(\lambda_{N,I}^{k,*})_{k \geq 1}$ towards Λ_{hom}^* .

the approximation of the optimal spectral gap (77) for a given frequency $k \geq 1$ and by $\lambda_{N,I}^{k,*}$ the corresponding optimal value. The optimal homogenized coefficient is $\bar{D}_{\text{hom}}^* = 1/Z$, which is easily obtained from the formula (75) together with the expression of the optimal diffusion coefficient in the homogenized limit (74). Therefore, the value of the homogenized limit of the spectral gap is the product of the first non-zero eigenvalue of $-\Delta$ on \mathbb{T} with \bar{D}_{hom}^* , which yields $\Lambda_{\text{hom}}^* = 4\pi^2/Z$.

The convergence of $\lambda_{N,I}^{k,*}$ towards Λ_{hom}^* is illustrated in Figure 7. In particular, we represent on the first right and left plots the optimal diffusion coefficients $D^{k,*}$ for increasing values of k , along with the diffusion coefficient D_{hom}^* . Note that we always observe that the diffusion $D^{k,*}$ is $1/k$ -periodic, and we only plot its trace on $[0, 1/k]$ (and stretch it on $[0, 1]$) so that optimal diffusion coefficients for various values of k can be compared: more precisely, the function plotted for the optimal diffusion coefficient for the frequency $k \geq 1$ is

$$\forall q \in \mathbb{T}, \quad \mathcal{D}^{k,*}(q) = D_{\lfloor \frac{Nq}{k} \rfloor + 1}^{k,*}.$$

We observe that, for these simple one-dimensional examples, the optimal and homogenized diffusion coefficients coincide almost perfectly for relatively small values of the frequency, namely already for $k = 3$. We also observe that the optimal diffusion coefficient displays more regularity for $k \geq 3$: it cancels at two points for $k = 1$ (and for $k = 2$ for the numerical example presented in Figure 7a) where the derivative has a singularity, but remains uniformly positive and smooth for $k \geq 3$.

6 Application to sampling algorithms

We present in this section an application of the optimization procedure to accelerate sampling algorithms. We focus on the Random Walk Metropolis–Hastings (RWMH) algorithm [53, 36] to unbiasedly sample from a Boltzmann–Gibbs target distribution. As we show in Section 6.1, this algorithm provides a consistent discretization of the continuous dynamics (3). More precisely, we show that the RWMH with proposals constructed using a space dependent diffusion converges, in the limit $\Delta t \rightarrow 0$, to a Langevin diffusion with space dependent diffusion given by the proposal variance. Notice that we do not discuss ergodicity of such dynamics, see [50] for results in that direction. We next present in Section 6.2 numerical experiments demonstrating that optimizing the diffusion leads to a more efficient sampling of the target measure. We numerically compare three variants of RWMH corresponding to three different choices for the variance of the proposal. The first variant is a simple RWMH algorithm with constant proposal variance; the two others have a space-dependent proposal variance $\Delta t \mathcal{D}$, with \mathcal{D} given by (i) the pre-computed optimal diffusion matrix and (ii) the explicit homogenized limit of Section 5.2.2.

Remark 7 (On the choice of the dynamics). *We consider a simple Random-Walk algorithm with moves of size $O(\sqrt{\Delta t})$ rather than a Metropolization of a Euler–Maruyama discretization of the overdamped Langevin dynamics (the SmartMC method [63] in molecular dynamics, known as MALA [62] in computational statistics). Indeed the latter dynamics have rejection probabilities of order $\Delta t^{3/2}$ for constant diffusion matrices, but a much larger rejection probability of order $\sqrt{\Delta t}$ for genuinely position dependent diffusion matrices [29] (see also Lemma 34). In contrast, the rejection probability is of order $\sqrt{\Delta t}$ in all cases for Random Walk Metropolis–Hastings. We therefore choose RWMH in order to have algorithms with rejection probabilities of the same order of magnitude, which allows for a fairer comparison of the dynamics. Let us mention that numerical schemes leading to a rejection probability scaling as $\Delta t^{3/2}$ for Metropolizations of overdamped Langevin dynamics with position dependent diffusion matrices have recently been proposed in [48]. We also refer to [55] for discussions about efficient discretizations of the overdamped Langevin dynamics with position dependent diffusions, using time-rescaling transforms.*

6.1 Consistency of the Random Walk Metropolis–Hastings algorithm

We describe in this section the RWMH algorithm we consider, and then provide consistency results of the method in the limit $\Delta t \rightarrow 0$.

Description of the Metropolis algorithm. The Random Walk Metropolis–Hastings algorithm [53, 36] is obtained by first proposing moves constructed from the current configuration by adding a Gaussian increment, and then accepting or rejecting the proposal move according to a Metropolis criterion. More precisely, starting from the current configuration $q_{\Delta t}^i$ for $i \geq 0$ and using a discretization time step $\Delta t > 0$, a new configuration is proposed as

$$\tilde{q}_{\Delta t}^{i+1} = q_{\Delta t}^i + \sqrt{2\Delta t} \mathcal{D}^{1/2}(q_{\Delta t}^i) G^{i+1}, \quad (78)$$

where $(G^i)_{i \geq 1}$ is a sequence of independent and identically distributed (i.i.d.) normal random variables. The transition kernel $\mathcal{T}(q_{\Delta t}^i, q_{\Delta t}^{i+1})$ associated with this transition has density

$$\mathcal{T}(q, q') = \left(\frac{1}{4\pi\Delta t} \right)^{d/2} \det \mathcal{D}(q)^{-1/2} \exp \left(-\frac{1}{4\Delta t} (q' - q)^\top \mathcal{D}(q)^{-1} (q' - q) \right).$$

Note that the reverse move associated with the proposal (78) is

$$q_{\Delta t}^i = \tilde{q}_{\Delta t}^{i+1} - \sqrt{2\Delta t} \mathcal{D}^{1/2}(\tilde{q}_{\Delta t}^{i+1}) \tilde{G}^{i+1}, \quad \tilde{G}^{i+1} = \mathcal{D}^{-1/2}(\tilde{q}_{\Delta t}^{i+1}) \mathcal{D}^{1/2}(q_{\Delta t}^i) G^{i+1}.$$

This allows to compute the Metropolis ratio as

$$\begin{aligned} R_{\Delta t}(q_{\Delta t}^i, G^{i+1}) &= \min \left\{ 1, \frac{\mu(\tilde{q}_{\Delta t}^{i+1}) \mathcal{T}(\tilde{q}_{\Delta t}^{i+1}, q_{\Delta t}^i)}{\mu(q_{\Delta t}^i) \mathcal{T}(q_{\Delta t}^i, \tilde{q}_{\Delta t}^{i+1})} \right\} \\ &= \min \left\{ 1, \left(\frac{\det \mathcal{D}(q_{\Delta t}^i)}{\det \mathcal{D}(\tilde{q}_{\Delta t}^{i+1})} \right)^{1/2} e^{-[V(\tilde{q}_{\Delta t}^{i+1}) - V(q_{\Delta t}^i)] - (|\tilde{G}^{i+1}|^2 - |G^{i+1}|^2)/2} \right\}. \end{aligned} \quad (79)$$

Accepting the proposal with probability $R_{\Delta t}(q_{\Delta t}^i, G^{i+1})$ using a sequence of i.i.d. random variables $(U^i)_{i \geq 1}$ with uniform law on $[0, 1]$, independent of $(G^i)_{i \geq 1}$, then leads to setting the new configuration as

$$q_{\Delta t}^{i+1} = q_{\Delta t}^i + \sqrt{2\Delta t} \mathcal{D}^{1/2}(q_{\Delta t}^i) G^{i+1} \mathbb{1}_{\{U^{i+1} \leq R_{\Delta t}(q_{\Delta t}^i, G^{i+1})\}}. \quad (80)$$

Weak error estimates and pathwise weak convergence. The following result, similar to [29, Lemma 4], implies that the numerical scheme (80) is weakly consistent. The proof is omitted since it exactly follows the proof of [29, Lemma 4] as a particular case when the drift term is 0.

Lemma 20. *For any $\varphi \in \mathcal{C}^\infty(\mathbb{T}^d)$, there exists $K \in \mathbb{R}_+$ and $\Delta t^* > 0$ such that*

$$\mathbb{E}_q(\varphi(q_{\Delta t}^1)) = \varphi(q) + \Delta t \mathcal{L}_{\mathcal{D}} \varphi(q) + \Delta t^{3/2} r_{\varphi, \Delta t}(q),$$

with $\sup_{q \in \mathbb{T}^d} |r_{\varphi, \Delta t}(q)| \leq K$ for any $0 < \Delta t \leq \Delta t^*$.

We next state a weak consistency result of the Metropolis scheme with the baseline continuous dynamics (3) at the level of the paths of the process. Define the rescaled, linearly interpolated continuous-time process $(Q_t^{\Delta t})_{t \geq 0}$ by

$$Q_t^{\Delta t} = \left(\left\lceil \frac{t}{\Delta t} \right\rceil - \frac{t}{\Delta t} \right) q_{\Delta t}^{\lfloor \frac{t}{\Delta t} \rfloor} + \left(\frac{t}{\Delta t} - \left\lfloor \frac{t}{\Delta t} \right\rfloor \right) q_{\Delta t}^{\lceil \frac{t}{\Delta t} \rceil}, \quad (81)$$

with $Q_0^{\Delta t} = q_0$ given. We can then state the following pathwise consistency result, proved in Appendix E.

Theorem 21. *Assume that $\mathcal{D} \in \mathcal{C}^2(\mathbb{T}^d, \mathcal{S}_d^{++})$, and denote by $P_{\Delta t}$ the law of the process $(Q_t^{\Delta t})_{t \geq 0}$ on $\mathcal{C}^0([0, \infty), \mathbb{T}^d)$, starting from a given initial condition $Q_0^{\Delta t} = q_0$. Then, $P_{\Delta t}$ converges weakly to P as $\Delta t \rightarrow 0$, where P is the law of the unique solution to the stochastic differential equation*

$$dQ_t = (-\mathcal{D}(Q_t) \nabla V(Q_t) + \operatorname{div} \mathcal{D}(Q_t)) dt + \sqrt{2} \mathcal{D}^{1/2}(Q_t) dW_t, \quad (82)$$

where $(W_t)_{t \geq 0}$ is a standard d -dimensional Brownian motion, with initial condition $Q_0 = q_0$.

Note that we assume that $\mathcal{D} \in \mathcal{C}^2(\mathbb{T}, \mathcal{S}_d^{++})$ for some technical estimates, in particular those of Lemma 34.

6.2 Sampling experiments

In this section, we illustrate the benefits of using a position dependent diffusion coefficient for sampling a Boltzmann–Gibbs target distribution. We consider a linear interpolation of the optimal diffusion coefficient D^* (introduced in Section 4) obtained with $N = I = 1000$ and $a = b = 0$, the diffusion coefficient for the homogenized problem D_{hom}^* (see Section 5 and equation (74)), and the constant diffusion coefficient D_{cst} (see Section 4.2).

We choose the one-dimensional double well potential $V(q) = \sin(4\pi q)(2 + \sin(2\pi q))$ defined on \mathbb{T} (the associated Gibbs distribution is displayed in Figure 3a). The potential exhibits a global minimum on \mathbb{T} at $x_0 \approx 0.36544$ and a local minimum at $x_1 \approx 0.89714$.

Diffusion coefficient	Constant	Homogenized	Optimal
Spectral gap	0.81	10.6	11.2
M–H rejection probability (%)	3.72	4.00	6.42

Table 2: Spectral gap and Metropolis–Hastings rejection probability for the simulations in Figure 8.

Sample trajectories. We show in Figure 8 sample trajectories following the discretization (80) for each diffusion coefficient. In order to compare the simulations, the same Gaussian variables $(G^i)_{i \geq 1}$ are used in (78). They are run with time step $\Delta t = 10^{-4}$ for $N_{\text{it}} = 10^6$ iterations, with the same initial condition $q_0 = 0$. This time step is chosen so that the rejection probability in the Metropolis–Hastings procedure is of order 5%, see Table 2 for the values of the rejection probabilities in each case, as well as Figure 11b for a study of the scaling of the rejection probability as a function of the time step. We observe that the dynamics using a constant diffusion coefficient is stuck longer in a well, whereas the dynamics using either the optimal or homogenized diffusion coefficient transitions faster between metastable states.

In Figure 8d, we compute the mean squared distance (MSD) averaged over $N_{\text{sim}} = 1000$ realizations of the dynamics for the various optimal diffusion coefficients using a time step $\Delta t = 10^{-4}$ and $N_{\text{it}} = 10^6$ iterations. Each realization starts at position $q_0 = 0$. Therefore, the averaged MSD at iteration $n \geq 1$ is

$$\forall 1 \leq n \leq N_{\text{it}}, \quad (\text{MSD})_n = \frac{1}{N_{\text{sim}}} \sum_{i=1}^{N_{\text{sim}}} \frac{1}{n} \sum_{j=1}^n \left| q_{\Delta t}^{i,j} - q_0 \right|^2,$$

where $q_{\Delta t}^{i,j}$ is the position of the i -th realization of the dynamics at iteration j . We observe more displacement when using either the optimal or the homogenized diffusion coefficient, which confirms that the associated dynamics diffuse more over the configuration space than the dynamics using a constant diffusion coefficient. The fact that the diffusion coefficient $\mathcal{D}_{\text{hom}}^*$ (approximated by (58)) leads to a larger effective diffusion should not come as a surprise since it is optimized to maximize the MSD of the underlying continuous dynamics, see [29, Section 3.5.2].

Convergence of the empirical law towards the Gibbs measure. In order to further compare the various diffusion coefficients and their respective effects on the sampling efficiency, we numerically compute the weighted L^2 error (χ^2 divergence) between the empirical distribution of the discretized process and the Gibbs measure, see (83) below. As pointed out in (5), for the limiting continuous-in-time dynamics (82), this error converges exponentially fast towards 0, at a rate given by the spectral gap of the generator of the dynamics.


More precisely, we set $\Delta t = 10^{-6}$ and consider an initial distribution μ^0 made of $N_{\text{samples}} = 10^4$ samples generated independently and uniformly over $[0, 1]$:

$$\mu^0 = \frac{1}{N_{\text{samples}}} \sum_{i=1}^{N_{\text{samples}}} \delta_{q_{\Delta t}^{i,0}}, \quad q_{\Delta t}^{i,0} \sim \mathcal{U}([0, 1]).$$


Each sample is then independently updated using the RWMH algorithm presented in Section 6.1, so that the empirical distribution μ^n at time $n\Delta t$ is

$$\mu^n = \frac{1}{N_{\text{samples}}} \sum_{i=1}^{N_{\text{samples}}} \delta_{q_{\Delta t}^{i,n}}.$$


In practice, the weighted L^2 error appearing in the left-hand side of (5) is approximated by discretizing the interval $[0, 1]$ into N_{bins} bins of equal sizes: for $1 \leq k \leq N_{\text{bins}}$, we denote




(a) Constant diffusion coefficient.



(b) Optimal diffusion coefficient, $a = 0$.



(c) Homogenized diffusion coefficient.



(d) Squared distance over iterations.

Figure 8: Sample trajectories of the RWMH algorithm for a periodic, one-dimensional potential with double well $V(q) = \sin(4\pi q)(2 + \sin(2\pi q))$. Notice that we plot the trajectories over the unfolded configuration space \mathbb{R} instead of the torus $\mathbb{T} = \mathbb{R}/\mathbb{Z}$.

by $B_k = [(k-1)/N_{\text{bins}}, k/N_{\text{bins}})$ the k -th bin. The weighted L^2 error is then approximated as

$$\left(\frac{1}{N_{\text{bins}}} \sum_{k=1}^{N_{\text{bins}}} \frac{(\hat{\mu}_k^n - \hat{\mu}_k)^2}{\hat{\mu}_k} \right)^{1/2} \quad (83)$$

where $\hat{\mu}_k^n, \hat{\mu}_k$ are proportional to the fraction of samples in the k -th bin, and normalized as

$$\frac{1}{N_{\text{bins}}} \sum_{k=1}^{N_{\text{bins}}} \hat{\mu}_k^n = \frac{1}{N_{\text{bins}}} \sum_{k=1}^{N_{\text{bins}}} \hat{\mu}_k = 1.$$

In particular, in our case,

$$\forall 1 \leq k \leq N_{\text{bins}}, \quad \hat{\mu}_k = N_{\text{bins}} \frac{e^{-V((k-1)/N_{\text{bins}})}}{\sum_{j=1}^{N_{\text{bins}}} e^{-V((j-1)/N_{\text{bins}})}},$$

approximates the Gibbs distribution, and

$$\forall 1 \leq k \leq N_{\text{bins}}, \quad \hat{\mu}_k^n = \frac{N_{\text{bins}}}{N_{\text{samples}}} \left| \left\{ 1 \leq i \leq N_{\text{samples}}, q_{\Delta t}^{i,n} \in B_k \right\} \right|,$$

approximates the empirical distribution μ^n , where $|A|$ denotes the cardinal of a set A . We run the RWMH algorithm for $N_{\text{it}} = 10^6$ iterations and use $N_{\text{bins}} = 100$ to approximate the weighted L^2 error. The results are averaged over 10 simulations.

The error between the empirical distribution and the Gibbs distribution as a function of the physical time (*i.e.* the number of iterations multiplied by the time step) is presented in Figure 9. This confirms that the use of the optimal diffusion or the homogenized coefficient greatly improves the convergence of the law of the process towards equilibrium. The exponential rate of convergence, corresponding to the dotted lines, is quickly reached - although this regime is not present for a long time for the optimal and homogenized diffusion coefficients since the weighted L^2 error converges in a rather short time towards a small limiting value, which is nonzero because of statistical and numerical errors mainly related to the finiteness of N_{bins} and N_{samples} (this is why we only show the results up to the physical time 0.25). Recall that the spectral gaps for the optimal, homogenized and constant diffusion coefficients are respectively 11.23, 10.58 and 0.81. This is in agreement with the numerical results of Figure 9 which show a slightly lower error for the optimal diffusion coefficient than for the homogenized diffusion coefficient, and a significantly lower error than for the constant diffusion coefficient.

Spatial analysis of the rejection probability of the Metropolis–Hastings procedure.

We compute the average rejection probability of the Metropolis–Hastings procedure for a fixed time step $\Delta t = 0.01$. We discretize the torus with a set of points $q^i = i/1000, 0 \leq i < 1000$. For each point q^i , we propose 10^5 moves following the rule (78), and count how many proposals are accepted when performing the Metropolis–Hastings procedure, *i.e.* computing (79) and comparing its value to a random number uniformly distributed over $[0, 1]$.

We represent on Figure 10, for various diffusion coefficients, the three following functions: the target distribution, the diffusion coefficient and the average rejection probability. We discarded the error bars as they were too small to be even noticed in the plot.

For the constant diffusion coefficient, for which the results are presented in Figure 10a, the proposals are most likely rejected when q is at the bottom of a well. When using either the optimal diffusion coefficient or the homogenized diffusion coefficient, the behavior is the opposite: the proposals are most likely accepted when q is located at the bottom of a well and/or when the gradient of the potential energy function is small. This is shown in Figures 10b and 10d. Notice that for the case of the optimal diffusion, higher rejection probabilities are observed where the optimal diffusion coefficient almost vanishes. This is because of the ratio of determinant factors appearing in the denominator of (79): even though both terms are of similar orders,




Figure 9: Weighted L^2 error (in logarithmic scale) between the empirical distribution and the Gibbs distribution over time for various diffusion coefficients.

they are almost zero, which causes numerical instabilities. We plot in Figure 10c the average rejection probabilities when using the optimal diffusion coefficients obtained with the optimization procedure and imposing various lower bounds $a \in \{0.2, 0.4, 0.6, 0.8, 1\}$. We notice that, even for small (but positive) values of a , the two singularities disappear. This suggests that imposing a positive lower bound in the optimization procedure can yield an optimal diffusion coefficient which makes the numerical computations more stable, and therefore produces a better numerical sampling.

Transition times. We compute the mean transition time from the deepest well to any of its two nearby periodized copies. More precisely, we consider the initial condition $q^0 = x_0 \approx 0.36544$ and compute the average physical time it takes for the system to cross $x_0 - 1$ or $x_0 + 1$. This average is computed over $N_{\text{transitions}} = 10^5$ transitions. We show in Figure 11 the mean transition time with respect to the time step Δt , as well as the rejection probability of the Metropolis–Hastings procedure. The rejection probability due to the Metropolis–Hastings procedure scales as $O(\sqrt{\Delta t})$; see [29, Section 4.7] and Lemma 34. We see that the mean transition time between the two metastable states is much smaller when using either the optimal diffusion coefficient or the homogenized diffusion coefficient than with the constant diffusion coefficient. For a time step $\Delta t = 5 \times 10^{-5}$, the numerical values of the mean transition times when using D^* , D_{hom}^* and D_{cst} are respectively 2.37, 1.77 and 17.78. This shows that the optimal diffusion coefficients obtained by our numerical procedure or the homogenized diffusion coefficient can be used to faster sample the configurational space, especially in the case of metastable dynamics.




Figure 10: Spatial analysis of the Metropolis–Hastings rejection probability for the periodic, one-dimensional double well potential $V(q) = \sin(4\pi q)(2 + \sin(2\pi q))$. On Figures 10a, 10b and 10d, the left y -axis refers to the diffusion coefficient and the target density values, and the right y -axis refers to the average rejection probability values.




Figure 11: Transition times with RWMH for the three optimal diffusion coefficients with respect to the time step Δt .

A Proof of optimization results

A.1 Proof of Theorem 4

The fact that the optimization problem is well posed follows by a straightforward adaptation of [38, Theorem 10.3.1]. Recall that, by Lemma 3, the application $\mathcal{D} \mapsto \Lambda(\mathcal{D})$ is concave. Moreover, the function Λ is upper-semicontinuous for the weak-* L_V^∞ topology induced on $\mathfrak{D}_p^{a,b}$ for any $a \geq 0$, $b > 0$ and $p \geq 1$ since it is defined in (7) as the infimum over $u \in H_0^1(\mu) \setminus \{0\}$ of the functions

$$\mathcal{D} \mapsto \frac{\int_{\mathbb{T}^d} \nabla u(q)^\top \mathcal{D}(q) \nabla u(q) \mu(q) dq}{\int_{\mathbb{T}^d} u(q)^2 \mu(q) dq},$$

which are continuous for the weak-* L_V^∞ topology. We finally note that the set $\mathfrak{D}_p^{a,b}$ is nonempty by Lemma 2 and compact for the weak-* L_V^∞ topology. This already shows that there exists a solution to (14). Note that the L_V^p norm in item (ii) of Theorem 4 is necessarily saturated for the maximizer in view of (9).

To prove item (i), we rely on the following lemma, which shows that the optimal diffusion matrix $\mathcal{D}^*(q)$ cannot be degenerate on an open set when $a = 0$ (for $a > 0$, the diffusion is of course nondegenerate).

Lemma 22. *Fix $a = 0$. For any open set $\Omega \subset \mathbb{T}^d$, there exists $q \in \Omega$ such that $\mathcal{D}^*(q) \neq 0$.*

Proof. We proceed by contradiction. Note first that $\mathcal{D}(q) = c_{p,b} \text{Id}_d$ is in the constrained set $\mathfrak{D}_p^{0,b}$ for

$$c_{p,b} = \min \left\{ \frac{1}{|\text{Id}_d|_F} \left(\int_{\mathbb{T}^d} e^{-pV(q)} dq \right)^{-1/p}, \frac{1}{b} \right\}.$$

Then (recalling the constant C_μ in (15)),

$$\Lambda(\mathcal{D}^*) \geq \Lambda(c_{p,b} \text{Id}_d) = c_{p,b} \inf_{u \in H_0^1(\mu) \setminus \{0\}} \frac{\int_{\mathbb{T}^d} |\nabla u(q)|^2 \mu(q) dq}{\int_{\mathbb{T}^d} u(q)^2 \mu(q) dq} = \frac{c_{p,b}}{C_\mu} > 0. \quad (84)$$

Assume that there exists an open set $\Omega \subset \mathbb{T}^d$ such that $\mathcal{D}^*(q) = 0$ for all $q \in \Omega$. We can then consider a nonzero function $u_\Omega \in \mathcal{C}^\infty(\mathbb{T}^d)$ such that $\text{supp}(u_\Omega) \subset \Omega$, which leads to

$$0 \leq \Lambda(\mathcal{D}^*) \leq \frac{\int_{\mathbb{T}^d} \nabla u_\Omega(q)^\top \mathcal{D}^*(q) \nabla u_\Omega(q) \mu(q) dq}{\int_{\mathbb{T}^d} u_\Omega(q)^2 \mu(q) dq} = 0.$$

The latter equality contradicts (84) and thus provides the desired result. \square

A.2 Proof of Proposition 9

There exists $\gamma > 0$ such that $\gamma \mathbb{1}_N \in \delta_p^{a,b}$ (where $\mathbb{1}_N$ is the N -dimensional vector whose components are all equal to 1), and thus, since $\mathcal{D}_{\mathbb{1}_N}(q) = 1$ for any $q \in \mathbb{T}^d$, and in view of the Poincaré inequality (15),

$$\Lambda(\mathcal{D}_{\mathbb{1}_N} \text{Id}_d) \geq \gamma \Lambda(\mathcal{D}_{\mathbb{1}_N} \text{Id}_d) \geq \frac{\gamma}{C_\mu}.$$

Consider $n \in \arg\min_{m \in \{1, \dots, N\}} D_{p,m}^*$. Since K_n has a nonempty interior, there exists an open ball \mathcal{B}_n such that $\overline{\mathcal{B}_n} \subset K_n$. We consider the function v_n defined on \mathcal{B}_n as a normalized

eigenfunction associated with the first eigenvalue of the Laplace problem with Dirichlet boundary conditions:

$$-\Delta v_n = \lambda_{1,n} v_n \text{ on } \mathcal{B}_n, \quad v_n = 0 \text{ on } \partial \mathcal{B}_n, \quad \int_{\mathcal{B}_n} v_n^2 = 1. \quad (85)$$

Note that $\lambda_{1,n} > 0$. The function v_n is then extended by 0 on $\mathbb{T}^d \setminus \mathcal{B}_n$. It is easy to check that $v_n \in H^1(\mu)$. We finally define $u_n \in H_0^1(\mu)$ as

$$u_n = v_n - \int_{\mathcal{B}_n} v_n \mu.$$

In view of (85), and denoting by $\mu_+ = \max_{\mathbb{T}^d} \mu < +\infty$,

$$\int_{\mathbb{T}^d} \mathcal{D}_{D^*} |\nabla u_n|^2 \mu = D_{p,n}^* \int_{\mathcal{B}_n} |\nabla u_n|^2 \mu \leq \mu_+ D_{p,n}^* \int_{\mathcal{B}_n} |\nabla v_n|^2 = \mu_+ \lambda_{1,n} D_{p,n}^*.$$

Therefore, since $\|u_n\|_{L^2(\mu)}^2 > 0$ (otherwise v_n would be constant on \mathcal{B}_n , which is impossible by (85)),

$$\Lambda(\mathcal{D}_{D^*} \text{Id}_d) = \inf_{u \in H_0^1(\mathbb{T}^d) \setminus \{0\}} \frac{\int_{\mathbb{T}^d} \mathcal{D}_{D^*}(q) |\nabla u(q)|^2 \mu(q) dq}{\int_{\mathbb{T}^d} u(q)^2 \mu(q) dq} \leq \frac{\int_{\mathbb{T}^d} \mathcal{D}_{D^*}(q) |\nabla u_n(q)|^2 \mu(q) dq}{\int_{\mathbb{T}^d} u_n(q)^2 \mu(q) dq} \leq \frac{\mu_+ \lambda_{1,n} D_{p,n}^*}{\|u_n\|_{L^2(\mu)}^2}.$$

This allows to conclude that $D_n^* \geq \gamma \|u_n\|_{L^2(\mu)}^2 / (\mu_+ \lambda_{1,n} C_\mu) > 0$, which leads to the claimed result.

B Characterization of the optimal diffusion

Proposition 6 is proved in Appendix B.1 and relies on technical results which are proved in Appendix B.2.

Let us introduce some notation and recall standard results for trace-class operators that will be useful below. For two Banach spaces E, F , we denote by $\mathcal{B}(E, F)$ the Banach space of bounded linear operators from E to F . We denote by $\|A\|_1 = \text{Tr}(A^* A)^{1/2}$ the trace norm of an operator A on $L^2(\mu)$. Note that if, for any $s \in [0, 1]$, the operator $A(s)$ is a trace-class operator on $L^2(\mu)$ and there exist $K \in \mathbb{R}_+$ such that $\|A(s)\|_1 \leq K$ for any $s \in [0, 1]$, then

$$\left\| \int_0^1 A(s) ds \right\|_1 \leq \int_0^1 \|A(s)\|_1 ds \leq K. \quad (86)$$

We also recall that if A is bounded on $L^2(\mu)$ and B is trace-class on $L^2(\mu)$, the operators AB and BA are trace-class on $L^2(\mu)$ and

$$\|AB\|_1 \leq \|A\|_{\mathcal{B}(L^2(\mu))} \|B\|_1, \quad \|BA\|_1 \leq \|A\|_{\mathcal{B}(L^2(\mu))} \|B\|_1 \quad (87)$$

We refer to [56, 66] for more detailed results on trace-class operators.

B.1 Proof of Proposition 6

To prove (31), we first assume that $\delta \mathcal{D} \in \mathcal{C}^3(\mathbb{T}^d, \mathcal{S}_d)$ with $\|\delta \mathcal{D}\|_{L^\infty(\mathbb{T}^d)} \leq c/2$, and show the inequality (31) with a bound $\|\delta \mathcal{D}\|_{L^\infty(\mathbb{T}^d)}^2$ on the right-hand side. The desired result for any $\delta \mathcal{D} \in L^\infty(\mathbb{T}^d, \mathcal{S}_d)$ such that $\|\delta \mathcal{D}\|_{L^\infty(\mathbb{T}^d)} \leq c/2$ is then obtained by a density argument.

We perform the computations for the map g_α , a similar reasoning leading to the result for the map h_α . In this proof, the letter K denotes a generic nonnegative finite constant which may change from line to line.

Consider $\delta\mathcal{D} \in \mathcal{C}^3(\mathbb{T}^d, \mathcal{S}_d)$ such that $\|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} \leq c/2$. Then $\mathcal{D} + \delta\mathcal{D} \geq (c/2)\text{Id}_d$ so that the operator $\mathcal{L}_{\mathcal{D}+\delta\mathcal{D}}$ is negative and self-adjoint on $L^2(\mu)$. We can therefore use Duhamel's formula twice to obtain

$$\begin{aligned}
& g_\alpha(\mathcal{D} + \delta\mathcal{D}) - g_\alpha(\mathcal{D}) \\
&= e^{\alpha\mathcal{L}_{\mathcal{D}+\delta\mathcal{D}}} - e^{\alpha\mathcal{L}_{\mathcal{D}}} = \left[e^{s\alpha\mathcal{L}_{\mathcal{D}+\delta\mathcal{D}}} e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}} \right]_{s=0}^{s=1} \\
&= \alpha \int_0^1 e^{s\alpha\mathcal{L}_{\mathcal{D}+\delta\mathcal{D}}} \mathcal{L}_{\delta\mathcal{D}} e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}} ds \\
&= \alpha \int_0^1 \left(e^{s\alpha\mathcal{L}_{\mathcal{D}}} + \left[e^{s(1-u)\alpha\mathcal{L}_{\mathcal{D}}} e^{su\alpha\mathcal{L}_{\mathcal{D}+\delta\mathcal{D}}} \right]_{u=0}^{u=1} \right) \mathcal{L}_{\delta\mathcal{D}} e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}} ds \\
&= \alpha \int_0^1 e^{s\alpha\mathcal{L}_{\mathcal{D}}} \mathcal{L}_{\delta\mathcal{D}} e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}} ds + \alpha^2 \int_0^1 \int_0^1 e^{s(1-u)\alpha\mathcal{L}_{\mathcal{D}}} \mathcal{L}_{\delta\mathcal{D}} e^{su\alpha\mathcal{L}_{\mathcal{D}+\delta\mathcal{D}}} \mathcal{L}_{\delta\mathcal{D}} e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}} s du ds \\
&= \alpha \int_0^1 e^{s\alpha\mathcal{L}_{\mathcal{D}}} \mathcal{L}_{\delta\mathcal{D}} e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}} ds + \alpha^2 \int_0^1 \int_0^s e^{(s-u)\alpha\mathcal{L}_{\mathcal{D}}} \mathcal{L}_{\delta\mathcal{D}} e^{u\alpha\mathcal{L}_{\mathcal{D}+\delta\mathcal{D}}} \mathcal{L}_{\delta\mathcal{D}} e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}} du ds, \quad (88)
\end{aligned}$$

where we performed the change of variables $u \leftarrow su$ for the last equality.

By Lemma 23 below, the operators $g_\alpha(\mathcal{D} + \delta\mathcal{D})$ and $g_\alpha(\mathcal{D})$ are trace-class on $L^2(\mu)$. The operator defined by the first term on the right-hand side of (88) is linear in $\delta\mathcal{D}$. We now show that it is trace-class on $L^2(\mu)$ and that there exists $K \in \mathbb{R}_+$ such that

$$\left\| \alpha \int_0^1 e^{s\alpha\mathcal{L}_{\mathcal{D}}} \mathcal{L}_{\delta\mathcal{D}} e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}} ds \right\|_1 \leq K \|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)}. \quad (89)$$

For any $s \in [0, 1]$, let $T_s = e^{s\alpha\mathcal{L}_{\mathcal{D}}} \mathcal{L}_{\delta\mathcal{D}} e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}}$. We need to distinguish between the two cases $0 \leq s \leq 1/2$ and $1/2 \leq s \leq 1$: when $0 \leq s \leq 1/2$, the operator $e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}}$, possibly multiplied by any polynomial of $\mathcal{L}_{\mathcal{D}}$, is trace-class on $L^2(\mu)$ uniformly in s thanks to Lemma 23, while the operator $e^{s\alpha\mathcal{L}_{\mathcal{D}}}$ is bounded on $L^2(\mu)$, so that (87) can be used. A similar result is obtained in the case $1/2 \leq s \leq 1$ by switching $e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}}$ and $e^{s\alpha\mathcal{L}_{\mathcal{D}}}$ in the reasoning.

More precisely, for $s \in [0, 1/2]$, we rewrite T_s as

$$T_s = \underbrace{e^{s\alpha\mathcal{L}_{\mathcal{D}}}}_{=:A_1} \underbrace{\mathcal{L}_{\delta\mathcal{D}}(\text{Id} - \mathcal{L}_{\mathcal{D}})^{-1}}_{=:A_2} \underbrace{(\text{Id} - \mathcal{L}_{\mathcal{D}})e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}}}_{=:A_3}.$$

Note that $\|A_1\|_{\mathcal{B}(L^2(\mu))} \leq 1$, and A_2 is bounded on $L^2(\mu)$ by Lemma 26 (see (96) below). The operator A_3 is trace-class on $L^2(\mu)$ by Lemma 23, so that T_s is trace-class on $L^2(\mu)$ with trace norm uniformly bounded over $s \in [0, 1/2]$. In view of (87) and (96), there exists $K \in \mathbb{R}_+$ such that

$$\forall s \in [0, 1/2], \quad \|T_s\|_1 \leq K \|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)}.$$

In the case $1/2 \leq s \leq 1$, we rewrite T_s as

$$T_s = \underbrace{e^{s\alpha\mathcal{L}_{\mathcal{D}}}(\text{Id} - \mathcal{L}_{\mathcal{D}})}_{=:A_3} \underbrace{(\text{Id} - \mathcal{L}_{\mathcal{D}})^{-1}\mathcal{L}_{\delta\mathcal{D}}}_{=:A_2} \underbrace{e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}}}_{=:A_1},$$

and proceed with a similar argument as above. In view of (86), the operator defined by the first term on the right-hand side of (88) is therefore trace-class on $L^2(\mu)$ and (89) holds.

We now show that the operator defined by the second term on the right-hand side of (88) is trace-class on $L^2(\mu)$, and that its trace norm is bounded from above by $\|\delta\mathcal{D}\|_{\mathcal{C}^3(\mathbb{T}^d)}^2$. First, note that the operator $(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^2 e^{\theta\mathcal{L}_{\mathcal{D}}}$ is trace-class on $L^2(\mu)$ for any $\theta > 0$. Indeed, let us rewrite this operator as $(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^2 (\text{Id} - \mathcal{L}_{\mathcal{D}})^{-2} (\text{Id} - \mathcal{L}_{\mathcal{D}})^2 e^{\theta\mathcal{L}_{\mathcal{D}}}$, and let us set $T = (\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^2 (\text{Id} - \mathcal{L}_{\mathcal{D}})^{-2}$. Lemma 25 then implies that

$$0 \leq T^* T \leq (\text{Id} - \mathcal{L}_{\mathcal{D}})^{-2} \left(\text{Id} - \frac{\|\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} + \|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)}}{c} \mathcal{L}_{\mathcal{D}} \right)^4 (\text{Id} - \mathcal{L}_{\mathcal{D}})^{-2}. \quad (90)$$

Recall that if $A \leq B$ as symmetric operators on a Hilbert space, then $C^*AC \leq C^*BC$ for any bounded operator C on the same Hilbert space. The operator on the right-hand side of (90) is therefore bounded on $L^2(\mu)$ by functional calculus, so that T is also bounded on $L^2(\mu)$. Lemma 23 then implies that the operators $(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^2 e^{\theta\mathcal{L}_{\mathcal{D}}}$ are trace-class on $L^2(\mu)$, and more precisely that for any $\underline{\theta} > 0$, there exists $K \in \mathbb{R}_+$ (which depends on $\underline{\theta}$) such that

$$\forall \theta > \underline{\theta}, \quad \|(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^2 e^{\theta\mathcal{L}_{\mathcal{D}}}\|_1 \leq K. \quad (91)$$

For any $s \in [0, 1]$ and $u \in [0, s]$, let $T_{s,u} = e^{\alpha(s-u)\mathcal{L}_{\mathcal{D}}} \mathcal{L}_{\delta\mathcal{D}} e^{u\alpha\mathcal{L}_{\mathcal{D}+\delta\mathcal{D}}} \mathcal{L}_{\delta\mathcal{D}} e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}}$. We need to distinguish between three cases: if $0 \leq s \leq 1/2$, the operator $e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}}$ is trace-class on $L^2(\mu)$ uniformly in s and the other operators are bounded on $L^2(\mu)$; if $1/2 \leq s \leq 1$, then the operator $e^{\alpha(s-u)\mathcal{L}_{\mathcal{D}}}$ is trace-class on $L^2(\mu)$ uniformly in u, s for $0 \leq u \leq s/2$, while it is the operator $e^{u\alpha\mathcal{L}_{\mathcal{D}+\delta\mathcal{D}}}$ which is trace-class on $L^2(\mu)$ uniformly in u, s for $s/2 \leq u \leq s$.

If $s \in [0, 1/2]$, we rewrite $T_{s,u}$ as

$$T_{s,u} = \underbrace{e^{(s-u)\alpha\mathcal{L}_{\mathcal{D}}}}_{=:A_1} \underbrace{\mathcal{L}_{\delta\mathcal{D}}(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^{-1}}_{=:A_2} \underbrace{e^{u\alpha\mathcal{L}_{\mathcal{D}+\delta\mathcal{D}}}}_{=:A_3} \underbrace{(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})\mathcal{L}_{\delta\mathcal{D}}(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^{-2}}_{=:A_4} \underbrace{(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^2 e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}}}_{=:A_5}.$$

It holds $\|A_1\|_{\mathcal{B}(L^2(\mu))} \leq 1$ and $\|A_3\|_{\mathcal{B}(L^2(\mu))} \leq 1$. The operators A_2 and A_4 are bounded on $L^2(\mu)$ by Lemma 26, while the operator A_5 is trace-class on $L^2(\mu)$ by (91). Moreover, in view of (87), (91), (96) and (98), there exists $K \in \mathbb{R}_+$ such that, for any $s \in [0, 1/2]$ and any $u \in [0, s]$,

$$\|T_{s,u}\|_1 \leq K \|\delta\mathcal{D}\|_{\mathcal{C}^3(\mathbb{T}^d)}^2. \quad (92)$$

If $s \in [1/2, 1]$ and $u \in [0, s/2]$, we rewrite $T_{s,u}$ as

$$T_{s,u} = \underbrace{e^{(s-u)\alpha\mathcal{L}_{\mathcal{D}}}(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^2(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^{-2}}_{=:A_5} \underbrace{\mathcal{L}_{\delta\mathcal{D}}(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})}_{=:A_4} \underbrace{e^{u\alpha\mathcal{L}_{\mathcal{D}+\delta\mathcal{D}}}}_{=:A_3} \underbrace{(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^{-1}\mathcal{L}_{\delta\mathcal{D}}}_{=:A_2} \underbrace{e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}}}_{=:A_1},$$

and if $s \in [1/2, 1]$ and $u \in [s/2, s]$, we rewrite $T_{s,u}$ as

$$T_{s,u} = \underbrace{e^{(s-u)\alpha\mathcal{L}_{\mathcal{D}}}}_{=:A_1} \underbrace{\mathcal{L}_{\delta\mathcal{D}}(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^{-1}}_{=:A_2} \underbrace{(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})e^{u\alpha\mathcal{L}_{\mathcal{D}+\delta\mathcal{D}}}}_{=:A_5} \underbrace{(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})}_{=:A_4} \underbrace{(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^{-1}\mathcal{L}_{\delta\mathcal{D}}}_{=:A_3} \underbrace{e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}}}_{=:A_3}.$$

We then perform manipulations similar to the ones leading to (92) in the case $0 \leq s \leq 1/2$.

In view of (86), the operator in (88) is therefore trace-class on $L^2(\mu)$, and there exists $K \in \mathbb{R}_+$ such that

$$\left\| \alpha^2 \int_0^1 \int_0^s e^{\alpha(s-u)\mathcal{L}_{\mathcal{D}}} \mathcal{L}_{\delta\mathcal{D}} e^{u\alpha\mathcal{L}_{\mathcal{D}+\delta\mathcal{D}}} \mathcal{L}_{\delta\mathcal{D}} e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}} du ds \right\|_1 \leq K \|\delta\mathcal{D}\|_{\mathcal{C}^3(\mathbb{T}^d)}^2.$$

We can at this stage take the trace in the equality (88). In particular, using the linearity and the cyclicity of the trace, the trace of the operator defined by the first term on the right-hand side of (88) is

$$\begin{aligned} \text{Tr} \left(\alpha \int_0^1 e^{s\alpha\mathcal{L}_{\mathcal{D}}} \mathcal{L}_{\delta\mathcal{D}} e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}} ds \right) &= \alpha \int_0^1 \text{Tr} \left(e^{s\alpha\mathcal{L}_{\mathcal{D}}} \mathcal{L}_{\delta\mathcal{D}} e^{(1-s)\alpha\mathcal{L}_{\mathcal{D}}} \right) ds \\ &= \alpha \int_0^1 \text{Tr} \left(e^{\alpha\mathcal{L}_{\mathcal{D}}} \mathcal{L}_{\delta\mathcal{D}} \right) ds \\ &= \alpha \text{Tr} \left(e^{\alpha\mathcal{L}_{\mathcal{D}}} \mathcal{L}_{\delta\mathcal{D}} \right) = \text{Tr} (g'_\alpha(\mathcal{D}) (-\mathcal{L}_{\delta\mathcal{D}})). \end{aligned}$$

This shows that (31) hold with the bound $\|\delta\mathcal{D}\|_{\mathcal{C}^3(\mathbb{T}^d)}^2$ on the right-hand side.

Using the density of $\mathcal{C}^3(\mathbb{T}^d, \mathcal{S}_d)$ in $L^\infty(\mathbb{T}^d, \mathcal{S}_d)$, we now prove that (31) holds. It suffices to show that the trace of the operator defined by the second term on the right-hand side of (88) is bounded from above by $\|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)}^2$. We compute, using Fubini's theorem,

$$\begin{aligned} & \text{Tr} \left(\alpha^2 \int_0^1 \int_0^s e^{\alpha(s-u)\mathcal{L}_\mathcal{D}} \mathcal{L}_\delta \mathcal{D} e^{u\alpha\mathcal{L}_\mathcal{D}+\delta\mathcal{D}} \mathcal{L}_\delta \mathcal{D} e^{(1-s)\alpha\mathcal{L}_\mathcal{D}} du ds \right) \\ &= \alpha^2 \int_0^1 \int_0^s \text{Tr} \left(e^{(s-u)\alpha\mathcal{L}_\mathcal{D}} \mathcal{L}_\delta \mathcal{D} e^{u\alpha\mathcal{L}_\mathcal{D}+\delta\mathcal{D}} \mathcal{L}_\delta \mathcal{D} e^{(1-s)\alpha\mathcal{L}_\mathcal{D}} \right) du ds \\ &= \alpha^2 \int_0^1 \int_0^s \text{Tr} \left(e^{(1-u)\alpha\mathcal{L}_\mathcal{D}} \mathcal{L}_\delta \mathcal{D} e^{u\alpha\mathcal{L}_\mathcal{D}+\delta\mathcal{D}} \mathcal{L}_\delta \mathcal{D} \right) du ds \\ &= \alpha^2 \int_0^1 \int_u^1 \text{Tr} \left(e^{u\alpha\mathcal{L}_\mathcal{D}+\delta\mathcal{D}} \mathcal{L}_\delta \mathcal{D} e^{(1-u)\alpha\mathcal{L}_\mathcal{D}} \mathcal{L}_\delta \mathcal{D} \right) ds du \\ &= \alpha^2 \int_0^1 (1-u) \text{Tr} \left(e^{u\alpha\mathcal{L}_\mathcal{D}+\delta\mathcal{D}} \mathcal{L}_\delta \mathcal{D} e^{(1-u)\alpha\mathcal{L}_\mathcal{D}} \mathcal{L}_\delta \mathcal{D} \right) du. \end{aligned}$$

Denote by $V_u = e^{u\alpha\mathcal{L}_\mathcal{D}+\delta\mathcal{D}} \mathcal{L}_\delta \mathcal{D} e^{(1-u)\alpha\mathcal{L}_\mathcal{D}} \mathcal{L}_\delta \mathcal{D}$. If $u \in [0, 1/2]$, we rewrite V_u as

$$V_u = \underbrace{e^{u\alpha\mathcal{L}_\mathcal{D}+\delta\mathcal{D}}}_{=:A_1} \underbrace{\mathcal{L}_\delta \mathcal{D} (\text{Id} - \mathcal{L}_\mathcal{D})^{-1}}_{=:A_2} \underbrace{(\text{Id} - \mathcal{L}_\mathcal{D}) e^{(1-u)\alpha\mathcal{L}_\mathcal{D}} (\text{Id} - \mathcal{L}_\mathcal{D})}_{=:A_3} \underbrace{(\text{Id} - \mathcal{L}_\mathcal{D})^{-1} \mathcal{L}_\delta \mathcal{D}}_{=:A_4}.$$

It holds $\|A_1\|_{\mathcal{B}(L^2(\mu))} \leq 1$, the operators A_2 and A_4 are bounded on $L^2(\mu)$ by Lemma 26, while the operator A_3 is trace-class on $L^2(\mu)$ by Lemma 23. In view of (87) and (96), there exists $K \in \mathbb{R}_+$ such that, for all $u \in [0, 1/2]$,

$$\|V_u\|_1 \leq K \|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)}^2.$$

If $u \in [1/2, 1]$, using the cyclicity of the trace, we write

$$\text{Tr}(V_u) = \text{Tr} \left(\underbrace{(\text{Id} - \mathcal{L}_\mathcal{D}) e^{u\alpha\mathcal{L}_\mathcal{D}+\delta\mathcal{D}} (\text{Id} - \mathcal{L}_\mathcal{D})}_{=:A_1} \underbrace{(\text{Id} - \mathcal{L}_\mathcal{D})^{-1} \mathcal{L}_\delta \mathcal{D}}_{=:A_2} \underbrace{e^{(1-u)\alpha\mathcal{L}_\mathcal{D}}}_{=:A_3} \underbrace{\mathcal{L}_\delta \mathcal{D} (\text{Id} - \mathcal{L}_\mathcal{D})^{-1}}_{=:A_4} \right),$$

and proceed as above. The inequality

$$\left| \int_0^1 (1-u) \text{Tr}(V_u) du \right| \leq \int_0^1 |\text{Tr}(V_u)| du \leq \int_0^1 \|V_u\|_1 du,$$

then implies (31), which concludes the proof.

B.2 Technical results

The following lemma (whose proof is given in Appendix B.2.1) shows that f_α is well-defined whenever $\alpha > 0$ and $\mathcal{D} \in L_V^p(\mathbb{T}^d, \mathcal{M}_{a,b})$ satisfies $\mathcal{D} \geq c\text{Id}_d$ for some $c > 0$.

Lemma 23. *Fix $\mathcal{D} \in L_V^p(\mathbb{T}^d, \mathcal{M}_{a,b})$ such that $\mathcal{D} \geq c\text{Id}_d$ for some $c > 0$. Then, for any $n \in \mathbb{N}$ and $\theta > 0$, the operators $(\mathcal{L}_\mathcal{D})^n e^{\theta\mathcal{L}_\mathcal{D}}$ and $(\text{Id} - \mathcal{L}_\mathcal{D})^n e^{\theta\mathcal{L}_\mathcal{D}}$ are trace-class on $L^2(\mu)$. Moreover, for any $n \in \mathbb{N}$ and $\underline{\theta} > 0$, there exists $\theta^* > 0$ and $K \in \mathbb{R}_+$ (which both depend on n and $\underline{\theta}$) such that*

$$\forall \theta \geq \underline{\theta}, \quad \|(\text{Id} - \mathcal{L}_\mathcal{D})^n e^{\theta\mathcal{L}_\mathcal{D}}\|_1 \leq K. \quad (93)$$

Using the notation of Section 3.2.2, the limit in (27) is therefore obtained with the dominated convergence theorem, using that $e^{-\alpha(\lambda_i - \lambda_2)} \leq e^{-(\lambda_i - \lambda_2)}$ for any $\alpha \geq 1$ and that the sums $\sum_{i \geq 3} N_i \lambda_i e^{-\lambda_i}$ and $\sum_{i \geq 3} N_i e^{-\lambda_i}$ are finite since the operators $g_1(\mathcal{D})$ and $h_1(\mathcal{D})$ are trace-class on $L^2(\mu)$.

The three following lemmas are useful to bound various terms when identifying the differentials of $\mathcal{D} \mapsto g_\alpha(\mathcal{D})$ and $\mathcal{D} \mapsto h_\alpha(\mathcal{D})$. More precisely, the following lemma allows to compare the operators $\mathcal{L}_\mathcal{D}$ and $\mathcal{L}_{\delta\mathcal{D}}$ as a function of $\|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)}$.

Lemma 24. Fix $\mathcal{D} \in L_V^p(\mathbb{T}^d, \mathcal{M}_{a,b})$ such that $\mathcal{D} \geq c \text{Id}_d$ for some $c > 0$. Then for any $\delta\mathcal{D} \in L^\infty(\mathbb{T}^d, \mathcal{S}_d)$,

$$\frac{1}{c} \|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} \mathcal{L}_{\mathcal{D}} \leq \mathcal{L}_{\delta\mathcal{D}} \leq -\frac{1}{c} \|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} \mathcal{L}_{\mathcal{D}}. \quad (94)$$

The result is an immediate consequence of the inequality

$$\delta\mathcal{D} \leq \|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} \text{Id}_d \leq \frac{1}{c} \|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} \mathcal{D}.$$

Likewise, the following lemma (proved in Appendix B.2.2) allows to compare the operators $\mathcal{L}_{\mathcal{D}}$ and $\mathcal{L}_{\mathcal{D}+\delta\mathcal{D}}$ for a small enough perturbation $\delta\mathcal{D}$.

Lemma 25. Fix $\mathcal{D} \in L_V^p(\mathbb{T}^d, \mathcal{M}_{a,b})$ such that $\mathcal{D} \geq c \text{Id}_d$ for some $c > 0$, and consider $\delta\mathcal{D} \in L^\infty(\mathbb{T}^d, \mathcal{S}_d)$ such that $\|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} \leq c/2$. Then,

$$\frac{\|\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} + \|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)}}{c} \mathcal{L}_{\mathcal{D}} \leq \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}} \leq \frac{c}{2\|\mathcal{D}\|_{L^\infty(\mathbb{T}^d)}} \mathcal{L}_{\mathcal{D}} \leq 0. \quad (95)$$

Lastly, the following result (proved in Appendix B.2.3) shows that some operators appearing in the proof of Proposition 6 are bounded on $L^2(\mu)$, with an explicit bound on $\delta\mathcal{D}$ when needed. Note that the bound (98) is not written in terms of the L^∞ norm of $\delta\mathcal{D}$, but rather as a function of the \mathcal{C}^3 norm of $\delta\mathcal{D}$. This may be a technical limitation due to our technique of proof, which involves commutators, but the estimate is sufficient for our purposes.

Lemma 26. Fix $\mathcal{D} \in L_V^p(\mathbb{T}^d, \mathcal{M}_{a,b})$ such that $\mathcal{D} \geq c \text{Id}_d$ for some $c > 0$, and consider $\delta\mathcal{D} \in \mathcal{C}^3(\mathbb{T}^d, \mathcal{S}_d)$ such that $\|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} \leq c/2$. Then the following statements hold:

- The operators $\mathcal{L}_{\delta\mathcal{D}}(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^{-1}$ and $\mathcal{L}_{\delta\mathcal{D}}(\text{Id} - \mathcal{L}_{\mathcal{D}})^{-1}$ are bounded on $L^2(\mu)$. More precisely, there exists $K \in \mathbb{R}_+$ such that

$$\|\mathcal{L}_{\delta\mathcal{D}}(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^{-1}\|_{\mathcal{B}(L^2(\mu))} + \|\mathcal{L}_{\delta\mathcal{D}}(\text{Id} - \mathcal{L}_{\mathcal{D}})^{-1}\|_{\mathcal{B}(L^2(\mu))} \leq K \|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)}; \quad (96)$$

- The operator $(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^2 e^{\theta\mathcal{L}_{\mathcal{D}}}$ is bounded on $L^2(\mu)$ for any $\theta > 0$. More precisely, for any $\underline{\theta} > 0$, there exists $K_{\underline{\theta}} \in \mathbb{R}_+$ such that, for all $\theta > \underline{\theta}$,

$$\|(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^2 e^{\theta\mathcal{L}_{\mathcal{D}}}\|_{\mathcal{B}(L^2(\mu))} \leq K_{\underline{\theta}}; \quad (97)$$

- The operator $(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})\mathcal{L}_{\delta\mathcal{D}}(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^{-2}$ is bounded on $L^2(\mu)$. More precisely, there exists $K \in \mathbb{R}_+$ such that

$$\|(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})\mathcal{L}_{\delta\mathcal{D}}(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^{-2}\|_{\mathcal{B}(L^2(\mu))} \leq K \|\delta\mathcal{D}\|_{\mathcal{C}^3(\mathbb{T}^d)}, \quad (98)$$

where $\|\delta\mathcal{D}\|_{\mathcal{C}^3(\mathbb{T}^d)} = \sup_{1 \leq i, j \leq d} \|\delta\mathcal{D}_{i,j}\|_{\mathcal{C}^3(\mathbb{T}^d)}$.

B.2.1 Proof of Lemma 23

We first show that the operator $e^{\theta\mathcal{L}_{\mathcal{D}}}$ is trace-class on $L^2(\mu)$ for $\theta > 0$. Using the lower bound on \mathcal{D} , it holds $\mathcal{L}_{\mathcal{D}} \leq c\mathcal{L}_{\text{Id}_d}$ on $L^2(\mu)$. It therefore suffices to show that the operator $e^{\theta\mathcal{L}_{\text{Id}_d}}$ is trace-class on $L^2(\mu)$. Define the unitary transformation

$$U: \begin{cases} L^2(\mu) \rightarrow L^2(\mathbb{T}^d), \\ f \mapsto f e^{-V/2}. \end{cases} \quad (99)$$

Straightforward computations show that

$$U\mathcal{L}_{\text{Id}_d}U^* = \Delta + W \leq 0 \quad (100)$$

on $L^2(\mathbb{T}^d)$, where Δ is the Laplacian operator with periodic boundary conditions on $L^2(\mathbb{T}^d)$, and the smooth function

$$W := \frac{1}{2}\Delta V - \frac{1}{4}|\nabla V|^2 \quad (101)$$

is identified with a multiplication operator on $L^2(\mathbb{T}^d)$. This implies that

$$0 \leq \text{Tr}(e^{\theta \mathcal{L}_D}) \leq \text{Tr}(e^{c\theta \mathcal{L}_{\text{Id}_d}}) = \text{Tr}_{L^2(\mathbb{T}^d)}(e^{c\theta(\Delta+W)}) \leq e^{c\theta \mathcal{W}} \text{Tr}_{L^2(\mathbb{T}^d)}(e^{c\theta \Delta}),$$

where $\mathcal{W} = \max_{q \in \mathbb{T}^d} |W(q)|$. Since the operator $e^{c\theta \Delta}$ is trace-class on $L^2(\mathbb{T}^d)$ for any $\theta > 0$ (see for instance [57, Theorem XIII.76]), the operator $e^{\theta \mathcal{L}_D}$ is trace-class on $L^2(\mu)$ for any $\theta > 0$.

To conclude the proof, we rely on the fact that, for $\underline{\theta} > 0$ and $n \in \mathbb{N}$ fixed, there exists $\theta^* > 0$ and $K > 0$ such that $(-x)^n e^{\theta x} \leq K e^{\theta^* x}$ for any $x \in \mathbb{R}_-$ and any $\theta \geq \underline{\theta}$. This implies that the operator $(\mathcal{L}_D)^n e^{\theta \mathcal{L}_D}$ is trace-class on $L^2(\mu)$ for any $n \in \mathbb{N}$. It follows that the operator $(\text{Id} - \mathcal{L}_D)^n e^{\theta \mathcal{L}_D}$ is also trace-class on $L^2(\mu)$ as it is a linear combination of the operators $(\mathcal{L}_D)^k e^{\theta \mathcal{L}_D}$ for $0 \leq k \leq n$. This also implies that the estimate (93) holds, and concludes the proof.

B.2.2 Proof of Lemma 25

Since $\frac{c}{2}\text{Id}_d \leq \mathcal{D} + \delta \mathcal{D} \leq (\|\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} + \|\delta \mathcal{D}\|_{L^\infty(\mathbb{T}^d)}) \text{Id}_d$, it holds

$$(\|\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} + \|\delta \mathcal{D}\|_{L^\infty(\mathbb{T}^d)}) \mathcal{L}_{\text{Id}_d} \leq \mathcal{L}_{\mathcal{D} + \delta \mathcal{D}} \leq \frac{c}{2} \mathcal{L}_{\text{Id}_d} \leq 0. \quad (102)$$

Using now that $c \text{Id}_d \leq \mathcal{D} \leq \|\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} \text{Id}_d$, it holds $\|\mathcal{D}\|_{L^\infty(\mathbb{T}^d)}^{-1} \mathcal{D} \leq \text{Id}_d \leq c^{-1} \mathcal{D}$ so that

$$c^{-1} \mathcal{L}_D \leq \mathcal{L}_{\text{Id}_d} \leq \|\mathcal{D}\|_{L^\infty(\mathbb{T}^d)}^{-1} \mathcal{L}_D. \quad (103)$$

Combining (102) with (103) implies (95), which concludes the proof.

B.2.3 Proof of Lemma 26

We prove the first item. Denote by $T = \mathcal{L}_{\delta \mathcal{D}}(\text{Id} - \mathcal{L}_D)^{-1}$. Using Lemma 24, it holds

$$0 \leq T^* T \leq \frac{1}{c^2} \|\delta \mathcal{D}\|_{L^\infty(\mathbb{T}^d)}^2 (\text{Id} - \mathcal{L}_D)^{-1} \mathcal{L}_D^2 (\text{Id} - \mathcal{L}_D)^{-1}.$$

The operator on the right-hand side of the inequality is bounded on $L^2(\mu)$ since

$$\|(\text{Id} - \mathcal{L}_D)^{-1} \mathcal{L}_D^2 (\text{Id} - \mathcal{L}_D)^{-1}\|_{\mathcal{B}(L^2(\mu))} \leq \sup_{\lambda \geq 0} \frac{\lambda^2}{(1 + \lambda)^2} = 1.$$

Therefore, the operator T is bounded on $L^2(\mu)$. Now let $S = \mathcal{L}_{\delta \mathcal{D}}(\text{Id} - \mathcal{L}_{\mathcal{D} + \delta \mathcal{D}})^{-1}$. Using Lemma 25, it holds

$$0 \leq SS^* \leq \mathcal{L}_{\delta \mathcal{D}} \left(\text{Id} - \frac{c}{2\|\mathcal{D}\|_{L^\infty(\mathbb{T}^d)}} \mathcal{L}_D \right)^{-2} \mathcal{L}_{\delta \mathcal{D}}.$$

The operator $\mathcal{L}_{\delta \mathcal{D}} \left(\text{Id} - \frac{c}{2\|\mathcal{D}\|_{L^\infty(\mathbb{T}^d)}} \|\mathcal{D}\|_{L^\infty(\mathbb{T}^d)}^{-1} \mathcal{L}_D \right)^{-1}$ is bounded on $L^2(\mu)$ using a similar argument as above, with an upper bound proportional to $\|\delta \mathcal{D}\|_{L^\infty(\mathbb{T}^d)}$. This implies that the operator S is bounded on $L^2(\mu)$. The estimate (96) easily follows.

We next prove the second item. Denote now by $T_\theta = (\text{Id} - \mathcal{L}_{\mathcal{D} + \delta \mathcal{D}})^2 e^{\theta \mathcal{L}_D}$. In view of Lemma 25,

$$0 \leq T_\theta^* T_\theta \leq e^{\theta \mathcal{L}_D} \left(\text{Id} - c^{-1} (\|\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} + \|\delta \mathcal{D}\|_{L^\infty(\mathbb{T}^d)}) \mathcal{L}_D \right)^4 e^{\theta \mathcal{L}_D}.$$

The operators $(\mathcal{L}_{\mathcal{D}})^i e^{\theta \mathcal{L}_{\mathcal{D}}}$ for $0 \leq i \leq 2$ are bounded on $L^2(\mu)$ by functional calculus. This implies that the operator T_{θ} is bounded on $L^2(\mu)$, and the uniform estimate (97) is obtained in the same manner as in the end of the proof of Lemma 23.

We finally prove the third item. Let $T = (\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})\mathcal{L}_{\delta\mathcal{D}}(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^{-2}$. Consider $\kappa > 0$ (a parameter which will be made precise below) and rewrite the operator T as

$$T = \underbrace{(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})(\kappa\text{Id} - \mathcal{L}_{\text{Id}_d})^{-1}}_{:=A_1} \underbrace{(\kappa\text{Id} - \mathcal{L}_{\text{Id}_d})\mathcal{L}_{\delta\mathcal{D}}(\kappa\text{Id} - \mathcal{L}_{\text{Id}_d})^{-2}}_{:=A_2} \underbrace{(\kappa\text{Id} - \mathcal{L}_{\text{Id}_d})^2(\text{Id} - \mathcal{L}_{\mathcal{D}+\delta\mathcal{D}})^{-2}}_{:=A_3}. \quad (104)$$

The operators A_1 and A_3 are bounded on $L^2(\mu)$. We prove this statement for the operator A_1 , the proof for A_3 being similar. Using (102), it holds

$$0 \leq A_1^* A_1 \leq (\kappa\text{Id} - \mathcal{L}_{\text{Id}_d})^{-1} \left(\text{Id} - \left(\|\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} + \|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} \right) \mathcal{L}_{\text{Id}_d} \right)^2 (\kappa\text{Id} - \mathcal{L}_{\text{Id}_d})^{-1},$$

and the operator on the right-hand side of the previous inequality is bounded on $L^2(\mu)$ since

$$\begin{aligned} & \left\| (\kappa\text{Id} - \mathcal{L}_{\text{Id}_d})^{-2} \left(\text{Id} - \left(\|\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} + \|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} \right) \mathcal{L}_{\text{Id}_d} \right)^2 \right\|_{\mathcal{B}(L^2(\mu))} \\ & \leq \left(\sup_{\lambda \geq 0} \frac{1 + \left(\|\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} + \|\delta\mathcal{D}\|_{L^\infty(\mathbb{T}^d)} \right) \lambda}{\kappa + \lambda} \right)^2 < +\infty. \end{aligned}$$

In view of (104), it therefore suffices to show that the operator $(\kappa\text{Id} - \mathcal{L}_{\text{Id}_d})\nabla^*\delta\mathcal{D}\nabla(\kappa\text{Id} - \mathcal{L}_{\text{Id}_d})^{-2}$ is bounded on $L^2(\mu)$, or, equivalently, that the operator

$$U(\kappa\text{Id} - \mathcal{L}_{\text{Id}_d})\nabla^*\delta\mathcal{D}\nabla(\kappa\text{Id} - \mathcal{L}_{\text{Id}_d})^{-2}U^* = U(\kappa\text{Id} - \mathcal{L}_{\text{Id}_d})U^* (U\nabla^*\delta\mathcal{D}\nabla U^*) U(\kappa\text{Id} - \mathcal{L}_{\text{Id}_d})^{-2}U^*, \quad (105)$$

is bounded on $L^2(\mathbb{T}^d)$, where U denotes the unitary transformation defined by (99). Using (100), it holds

$$U(\kappa\text{Id} - \mathcal{L}_{\text{Id}_d})U^* = \kappa\text{Id} - \Delta - W, \quad U(\kappa\text{Id} - \mathcal{L}_{\text{Id}_d})^{-2}U^* = (\kappa\text{Id} - \Delta - W)^{-2}.$$

where W is defined in (101). Straightforward computations lead to

$$U\nabla^* = \left(-\nabla + \frac{1}{2}\nabla V \right) U, \quad \nabla U^* = U^* \left(\nabla + \frac{1}{2}\nabla V \right).$$

Therefore, the operator in (105) can be rewritten as

$$(\kappa\text{Id} - \Delta - W) \left[\sum_{1 \leq i, j \leq d} \left(-\partial_{q_j} + \frac{1}{2}\partial_{q_j} V \right) \delta\mathcal{D}_{i,j} \left(\partial_{q_i} + \frac{1}{2}\partial_{q_i} V \right) \right] (\kappa\text{Id} - \Delta - W)^{-2}.$$

We now choose κ such that $\kappa - W \geq 1$ (which is possible since W is smooth on the compact space \mathbb{T}^d), so that $\kappa\text{Id} - \Delta - W \geq \text{Id}$ on $L^2(\mathbb{T}^d)$. The operators $(\kappa\text{Id} - \Delta - W)(\text{Id} - \Delta)^{-1}$ and $(\text{Id} - \Delta)^2(\kappa\text{Id} - \Delta - W)^{-2}$ are then bounded on $L^2(\mathbb{T}^d)$, so that it suffices to show that the operator

$$(\text{Id} - \Delta) \left[\sum_{1 \leq i, j \leq d} \left(-\partial_{q_j} + \frac{1}{2}\partial_{q_j} V \right) \delta\mathcal{D}_{i,j} \left(\partial_{q_i} + \frac{1}{2}\partial_{q_i} V \right) \right] (\text{Id} - \Delta)^{-2}$$

is bounded on $L^2(\mathbb{T}^d)$ to conclude.

For $k \geq 1$, denote by $H^k(\mathbb{T}^d)$ the Sobolev space of square integrable functions on \mathbb{T}^d with all (distributional) derivatives up to order k belonging to $L^2(\mathbb{T}^d)$. Using the Fourier series characterization of Sobolev spaces and the fact that V is smooth, it holds

$$\left\{ \begin{array}{l} (\text{Id} - \Delta)^{-2} \in \mathcal{B}(L^2(\mathbb{T}^d), H^4(\mathbb{T}^d)), \\ \forall 1 \leq i, j \leq d, \quad \partial_{q_i} + \frac{1}{2} \partial_{q_i} V \in \mathcal{B}(H^4(\mathbb{T}^d), H^3(\mathbb{T}^d)), \\ \forall 1 \leq i, j \leq d, \quad -\partial_{q_j} + \frac{1}{2} \partial_{q_j} V \in \mathcal{B}(H^3(\mathbb{T}^d), H^2(\mathbb{T}^d)), \\ \text{Id} - \Delta \in \mathcal{B}(H^2(\mathbb{T}^d), L^2(\mathbb{T}^d)). \end{array} \right.$$

To obtain (98), it therefore suffices that $\delta \mathcal{D}_{i,j}$, seen as a multiplication operator on $H^3(\mathbb{T}^d)$, is bounded uniformly in $i, j \in \{1, \dots, d\}$ with $\|\delta \mathcal{D}_{i,j}\|_{\mathcal{B}(H^3(\mathbb{T}^d))} \leq K_{i,j} \|\delta \mathcal{D}\|_{\mathcal{C}^3(\mathbb{T}^d)}$ for a constant $K_{i,j} \geq 0$. This is easily seen to hold using the Leibniz rule since $\delta \mathcal{D} \in \mathcal{C}^3(\mathbb{T}^d, \mathcal{S}_d)$. Thus, the estimate (98) holds, which concludes the proof of Lemma 26.

C Numerical optimization implementation details

We set $I = N$ in order to ease computations. In particular, the nodes for the basis functions $q_i = i/I$ for $i \in \{0, \dots, I-1\}$ are such that $K_n = [q_{n-1}, q_n]$ for $n \in \{1, \dots, N\}$. Recall that periodic boundary conditions are imposed, *e.g.* $q_{-1} = q_{I-1}$.

C.1 Numerical optimization of the smooth-min approximation

In order to solve (50), we need to compute the gradient of F_α defined in (51) with respect to D . The expressions of the gradients of the eigenvalues with respect to D are similar to the expressions obtained in the proof of Lemma 11: if the maps $D \mapsto \sigma_i(D)$ with $i \in \{2, \dots, I\}$ are differentiable, one finds that

$$\forall n \in \{1, \dots, N\}, \quad \frac{\partial \sigma_i}{\partial D_n}(D) = U_i(D)^\top A_n U_i(D),$$

where $U_i(D)$ is an eigenvector associated with $\sigma_i(D)$ which satisfies $U_i(D)^\top B U_i(D) = 1$. Likewise, using the approximation (53), and similarly to (57), it holds

$$\forall n \in \{1, \dots, N\}, \quad \frac{\partial \sigma_i}{\partial D_n}(D) \approx N e^{-V(\frac{n-1}{N})} [U_i(D)_n - U_i(D)_{n-1}]^2.$$

Then, it holds

$$\frac{\partial G_\alpha}{\partial D_n}(D) = -\alpha \sum_{i=2}^I e^{-\alpha \sigma_i(D)} U_i(D)^\top A_n U_i(D),$$

and

$$\frac{\partial H_\alpha}{\partial D_n}(D) = \sum_{i=2}^I (1 - \alpha \sigma_i(D)) e^{-\alpha \sigma_i(D)} U_i(D)^\top A_n U_i(D),$$

from which the gradient of F_α is easily computed.

C.2 Approximation of the integral elements

Approximation of A_n . Fix $n \in \{1, \dots, N\}$. On K_n , the only basis functions which are nonzero are φ_n and φ_{n-1} . Therefore, the only nonzero entries for A_n correspond to the (row, column) pairs $(n-1, n-1)$, $(n-1, n)$, $(n, n-1)$ and (n, n) . For example, the entry (n, n) is approximated as

$$[A_n]_{n,n} = \int_{\frac{n-1}{N}}^{\frac{n}{N}} (\varphi'_n(q))^2 e^{-V(q)} dq \approx e^{-V(\frac{n-1}{N})} \int_{\frac{n-1}{N}}^{\frac{n}{N}} (\varphi'_n(q))^2 dq = N e^{-V(\frac{n-1}{N})},$$

while the entry $(n, n-1)$ is approximated as $[A_n]_{n,n-1} \approx -Ne^{-V(\frac{n-1}{N})}$. The formula (53) then follows.

Approximation of B . Fix $(i, j) \in \{1, \dots, N\}^2$. Similar approximations as above lead to

$$\begin{aligned}
B_{i,j} &= \int_0^1 \varphi_i(q) \varphi_j(q) e^{-V(q)} dq = \sum_{n=1}^N \int_{\frac{n-1}{N}}^{\frac{n}{N}} \varphi_i(q) \varphi_j(q) e^{-V(q)} dq \\
&\approx \sum_{n=1}^N e^{-V(\frac{n-1}{N})} \int_{\frac{n-1}{N}}^{\frac{n}{N}} \varphi_i(q) \varphi_j(q) dq \\
&= \frac{1}{6N} \sum_{n=1}^N e^{-V(\frac{n-1}{N})} (2\delta_{j,n-1}\delta_{i,j} + \delta_{j,n-1}\delta_{i,j+1} + 2\delta_{j,n}\delta_{i,j} + \delta_{j,n}\delta_{i,j-1}) \\
&= \frac{1}{6N} e^{-V(\frac{i-1}{N})} (2\delta_{i,j} + \delta_{i,j+1}) + \frac{1}{6N} e^{-V(\frac{i}{N})} (2\delta_{i,j} + \delta_{i,j-1}) \\
&\approx \frac{1}{6N} e^{-V(\frac{i-1}{N})} (4\delta_{i,j} + \delta_{i,j+1} + \delta_{i,j-1}),
\end{aligned}$$

which is exactly (54).

D Proof of homogenization results

We fix $a, b > 0$ in all this section. The proofs of the results in Appendices D.1 to D.2 are given for completeness, as they are obtained by straightforward modifications of standard results of the literature on homogenization problems with Dirichlet boundary conditions [2, 13].

We start by stating a result which will be repeatedly used in the following, and whose proof is based on the Lax-Milgram lemma (by an easy adaptation of [2, Lemma 1.3.21]).

Lemma 27. *Fix $a, b > 0$, and consider $f \in H^{-1}(\mathbb{T}^d)$ such that $\langle f, \mathbf{1} \rangle_{H^{-1}(\mathbb{T}^d), H_0^1(\mathbb{T}^d)} = 0$ and $\mathcal{A} \in L^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$. Then, there exists a unique solution $u \in H^1(\mathbb{T}^d)$ to the problem*

$$\begin{cases} -\operatorname{div} \mathcal{A} \nabla u = f & \text{in } H^{-1}(\mathbb{T}^d), \\ \int_{\mathbb{T}^d} u = 0. \end{cases}$$

Moreover, there exists $C > 0$ (which does not depend on u nor f) such that

$$\|u\|_{H^1(\mathbb{T}^d)} \leq C \|f\|_{H^{-1}(\mathbb{T}^d)}. \quad (106)$$

D.1 Proof of Theorem 12

The proof boils down to adapting the proof of [2, Theorem 1.2.16], which states a similar compactness result for equations with Dirichlet boundary conditions, to our case with periodic boundary conditions. To do so, we start by considering a modified problem on an extended domain with Dirichlet boundary conditions, and apply [2, Theorem 1.2.16] to obtain the H -convergence for this modified problem. We then use a result on the independence of H -convergence on boundary conditions, stated e.g. in [2, Proposition 1.2.19], to obtain the H -convergence in the case of periodic boundary conditions.

Fix $f \in H^{-1}(\mathbb{T}^d)$ such that $\langle f, \mathbf{1} \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)} = 0$. Consider the open set $\Omega = (-1, 2)^d$ and denote by $\tilde{\mathcal{A}}^k$ and \tilde{f} the extensions to Ω of \mathcal{A}^k and f obtained by periodicity. Consider $\tilde{u}_0^k \in H^1(\Omega)$

the solution to

$$\begin{cases} -\operatorname{div}(\tilde{\mathcal{A}}^k \nabla \tilde{u}_0^k) = \tilde{f} & \text{on } \Omega, \\ \tilde{u}_0^k = 0 & \text{on } \partial\Omega. \end{cases}$$

In view of [2, Theorem 1.2.16], and upon extraction of a subsequence (not explicitly indicated), there exists $\tilde{\mathcal{A}} \in L^\infty(\Omega, \mathcal{M}_{a,b})$ such that

$$\begin{cases} \tilde{u}_0^k \rightharpoonup \tilde{u}_0 & \text{weakly in } H^1(\Omega), \\ \tilde{\mathcal{A}}^k \nabla \tilde{u}_0^k \rightharpoonup \tilde{\mathcal{A}} \nabla \tilde{u}_0 & \text{weakly in } L^2(\Omega)^d, \end{cases}$$

where \tilde{u}_0 is the solution of the homogenized problem

$$\begin{cases} -\operatorname{div}(\tilde{\mathcal{A}} \nabla \tilde{u}_0) = \tilde{f} & \text{on } \Omega, \\ \tilde{u}_0 = 0 & \text{on } \partial\Omega. \end{cases}$$

We now turn back to our original problem with periodic boundary conditions. Denote by $u^k \in H^1(\mathbb{T}^d)$ the unique solution (thanks to Lemma 27) to

$$\begin{cases} -\operatorname{div}(\mathcal{A}^k \nabla u^k) = f & \text{on } \mathbb{T}^d, \\ \int_{\mathbb{T}^d} u^k(q) dq = 0. \end{cases} \quad (107)$$

Since $\mathcal{A}^k \in L^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$ and u^k has average 0 with respect to the Lebesgue measure on \mathbb{T}^d , standard estimates based on the Poincaré–Wirtinger inequality show that the sequence $(u^k)_{k \geq 1}$ is uniformly bounded in $H^1(\mathbb{T}^d)$, hence weakly converges in $H^1(\mathbb{T}^d)$ to a function u , up to the extraction of a subsequence. This extracted subsequence also strongly converges in $L^2(\mathbb{T}^d)$. In fact, as made precise at the end of the proof, we will prove that the whole sequence converges. With some abuse of notation, we therefore still denote the extracted subsequence as $(u^k)_{k \geq 1}$. Note that, in particular, $\int_{\mathbb{T}^d} u(q) dq = 0$.

Denoting by \tilde{u}^k, \tilde{u} the extensions to Ω of u^k, u obtained by periodicity, we claim that

$$\tilde{\mathcal{A}}^k \nabla \tilde{u}^k \rightharpoonup \tilde{\mathcal{A}} \nabla \tilde{u} \quad \text{weakly in } L^2_{\text{loc}}(\Omega)^d. \quad (108)$$

In view of [2, Proposition 1.2.19], it suffices to this end to check that $\tilde{u}^k \rightharpoonup \tilde{u}$ weakly in $H^1_{\text{loc}}(\Omega)$ and $-\operatorname{div}(\tilde{\mathcal{A}}^k \nabla \tilde{u}^k) = \tilde{f} \in H^{-1}_{\text{loc}}(\Omega)$. To prove the first statement, we combine the weak convergence of $(u^k)_{k \geq 1}$ in $H^1(\mathbb{T}^d)$ with an argument based on a partition of unity. Fix $\tilde{\phi} \in H^{-1}(\Omega)$ with compact support K in Ω . Thanks to the compactness of the support and for $\varepsilon \in (0, 1/3)$ fixed, there exists a finite number L of points $(x_\ell)_{1 \leq \ell \leq L}$ such that $K \subset \bigcup_{\ell=1}^L \mathcal{B}(x_\ell, \varepsilon)$, where $\mathcal{B}(x_\ell, \varepsilon)$ denotes the open ball of radius ε centered at x_ℓ . Upon reducing ε , it can be assumed that $\overline{\mathcal{B}(x_\ell, \varepsilon)} \subset \Omega$ for all $\ell \in \{1, \dots, L\}$. Note that, for all $\ell \in \{1, \dots, L\}$, the mapping $i_\ell : \mathcal{B}(x_\ell, \varepsilon) \rightarrow \mathbb{T}^d$ defined as $i_\ell(x) = x \bmod \mathbb{Z}^d$ is injective. In addition, there exists an associated partition of unity, namely a family of nonnegative smooth functions $(\chi_\ell)_{1 \leq \ell \leq L}$ such that $\operatorname{supp}(\chi_\ell) \subset \overline{\mathcal{B}(x_\ell, \varepsilon)}$ for all $\ell \in \{1, \dots, L\}$, and $\sum_{\ell=1}^L \chi_\ell = 1$ on K . We use the decomposition $\tilde{\phi} = \sum_{\ell=1}^L \tilde{\phi}_\ell$, where $\tilde{\phi}_\ell = \tilde{\phi} \chi_\ell$ has support on $\overline{\mathcal{B}(x_\ell, \varepsilon)}$. In particular, for all $k \geq 1$,

$$\langle \tilde{\phi}, \tilde{u}^k \rangle_{H^{-1}(\Omega), H^1(\Omega)} = \sum_{\ell=1}^L \langle \tilde{\phi}_\ell, \tilde{u}^k \rangle_{H^{-1}(\Omega), H^1(\Omega)}.$$

Define next $\phi_\ell \in H^{-1}(\mathbb{T}^d)$ from $\tilde{\phi}_\ell$ as

$$\forall v \in H^1(\mathbb{T}^d), \quad \langle \phi_\ell, v \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)} = \langle \tilde{\phi}, \chi_\ell(v \circ i_\ell) \rangle_{H^{-1}(\Omega), H^1(\Omega)}.$$

In particular, $\langle \tilde{\phi}_\ell, \tilde{u}^k \rangle_{H^{-1}(\Omega), H^1(\Omega)} = \langle \phi_\ell, u^k \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)}$ for all $k \geq 1$ and $\langle \tilde{\phi}_\ell, \tilde{u} \rangle_{H^{-1}(\Omega), H^1(\Omega)} = \langle \phi_\ell, u \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)}$, so that

$$\begin{aligned} \langle \tilde{\phi}, \tilde{u}^k \rangle_{H^{-1}(\Omega), H^1(\Omega)} &= \sum_{\ell=1}^L \langle \phi_\ell, u^k \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)} \\ &\xrightarrow{k \rightarrow +\infty} \sum_{\ell=1}^L \langle \phi_\ell, u \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)} = \sum_{\ell=1}^L \langle \tilde{\phi}_\ell, \tilde{u} \rangle_{H^{-1}(\Omega), H^1(\Omega)} = \langle \tilde{\phi}, \tilde{u} \rangle_{H^{-1}(\Omega), H^1(\Omega)}, \end{aligned}$$

which allows to conclude that $\tilde{u}^k \rightharpoonup \tilde{u}$ weakly in $H_{\text{loc}}^1(\Omega)$. A similar argument, with the same partition of unity, can be used to prove that $-\text{div}(\tilde{\mathcal{A}}^k \nabla \tilde{u}^k) = \tilde{f}$ in $H_{\text{loc}}^{-1}(\Omega)$ for all $k \geq 1$.

For any $\Phi \in L^2(\mathbb{T}^d)^d$, define the function $\Phi^\dagger : \Omega \rightarrow \mathbb{R}$ obtained by extending Φ by 0 outside of the set $(0, 1)^d$ identified with \mathbb{T}^d . More precisely, $\Phi^\dagger(q) = \mathbf{1}_{(0,1)^d}(q)\Phi(i(q))$, where $i : (0, 1)^d \rightarrow \mathbb{T}^d$ is the canonical injection. Note that $\Phi^\dagger \in L^2(\Omega)^d$ has compact support in Ω , and, using (108),

$$\begin{aligned} \int_{\mathbb{T}^d} \Phi(q)^\top \mathcal{A}^k(q) \nabla u^k(q) dq &= \int_{\Omega} \Phi^\dagger(q)^\top \tilde{\mathcal{A}}^k(q) \nabla \tilde{u}^k(q) dq \\ &\xrightarrow{k \rightarrow +\infty} \int_{\Omega} \Phi^\dagger(q)^\top \hat{\mathcal{A}}(q) \nabla \tilde{u}(q) dq = \int_{\mathbb{T}^d} \Phi(q)^\top \bar{\mathcal{A}}(q) \nabla u(q) dq, \end{aligned}$$

where $\bar{\mathcal{A}} = \hat{\mathcal{A}} \circ i^{-1}$. This shows that $\mathcal{A}^k \nabla u^k$ weakly converges to $\bar{\mathcal{A}} \nabla u$ in $L^2(\mathbb{T}^d)^d$. In view of (107),

$$\begin{cases} -\text{div}(\bar{\mathcal{A}} \nabla u) = f & \text{in } H^{-1}(\mathbb{T}^d), \\ \int_{\mathbb{T}^d} u(q) dq = 0. \end{cases}$$

Thus, u is uniquely defined, and the whole sequences $(u^k)_{k \geq 1}$ and $(\mathcal{A}^k \nabla u^k)_{k \geq 1}$ weakly converge respectively to u in $H^1(\mathbb{T}^d)$ and $\bar{\mathcal{A}} \nabla u$ in $L^2(\mathbb{T}^d)^d$, which provides the desired convergence in the sense of Definition 1.

D.2 Proof of Theorem 14

The proof of Theorem 14 is an adaptation of the proof of [2, Theorem 1.3.16]. The first step is to show that $(u^k)_{k \geq 1}$ and $(\lambda^k)_{k \geq 1}$ converge respectively (up to extraction) to an eigenvector and eigenvalue of the problem

$$\begin{cases} -\text{div}(\bar{\mathcal{A}} \nabla u(q)) = \bar{\lambda} \rho(q) u(q) & \text{on } \mathbb{T}^d, \\ \int_{\mathbb{T}^d} u(q)^2 dq = 1. \end{cases} \quad (109)$$

The second step is to prove that $\bar{\lambda}$ is indeed the smallest nonzero eigenvalue of (109).

Convergence to a solution of the problem (109). Let us first prove that, up to the extraction of a subsequence, the sequences $(\lambda^k)_{k \geq 1}$ and $(u^k)_{k \geq 1}$ respectively converge in \mathbb{R}_+ and weakly in $H^1(\mathbb{T}^d)$. By the min-max principle, using the fact that the eigenfunctions of the operator $-\text{div}(\mathcal{A}^k \nabla \cdot)$ associated with the eigenvalue 0 are constant functions, the second-smallest eigenvalue λ^k satisfies

$$\lambda^k = \min_{v \in H^1(\mathbb{T}^d) \setminus \{0\}} \left\{ \frac{\int_{\mathbb{T}^d} \nabla v(q)^\top \mathcal{A}^k(q) \nabla v(q) dq}{\int_{\mathbb{T}^d} v^2(q) \rho^k(q) dq} \mid \int_{\mathbb{T}^d} v \rho^k = 0 \right\}.$$

By , $a\text{Id}_d \leq \mathcal{A}^k \leq b^{-1}\text{Id}_d$ and $0 < \rho_- \leq \rho^k \leq \rho_+$, so that, for all $v \in H^1(\mathbb{T}^d)$,

$$\frac{a}{\rho_+} \frac{\int_{\mathbb{T}^d} |\nabla v(q)|^2 dq}{\int_{\mathbb{T}^d} v^2(q) dq} \leq \frac{\int_{\mathbb{T}^d} \nabla v(q)^\top \mathcal{A}^k(q) \nabla v(q) dq}{\int_{\mathbb{T}^d} \rho^k(q) v^2(q) dq} \leq \frac{1}{b\rho_-} \frac{\int_{\mathbb{T}^d} |\nabla v(q)|^2 dq}{\int_{\mathbb{T}^d} v^2(q) dq},$$

which implies

$$0 < \frac{4\pi^2 a}{\rho_+} \leq \lambda^k \leq \frac{4\pi^2}{b\rho_-} \quad (110)$$

since the first nonzero eigenvalue of the operator $-\Delta$ on \mathbb{T}^d is $4\pi^2$. This also implies

$$a \|\nabla u^k\|_{L^2(\mathbb{T}^d)}^2 \leq \int_{\mathbb{T}^d} \nabla u^k(q)^\top \mathcal{A}^k(q) \nabla u^k(q) dq = \lambda^k \int_{\mathbb{T}^d} \rho^k(q) u^k(q)^2 dq \leq \frac{4\pi^2 \rho_+}{b\rho_-},$$

since u^k is normalized in $L^2(\mathbb{T}^d)$. The sequences $(\lambda^k)_{k \geq 1}$ and $(u^k)_{k \geq 1}$ are therefore bounded respectively in \mathbb{R}_+ and $H^1(\mathbb{T}^d)$.

Upon extraction (not indicated explicitly in the notation), there exists a subsequence such that $\lambda^k \rightarrow \bar{\lambda}$, and $u^k \rightharpoonup u$ weakly in $H^1(\mathbb{T}^d)$ and strongly in $L^2(\mathbb{T}^d)$. In particular, $\|u\|_{L^2(\mathbb{T}^d)} = 1$. Since $\lambda^k \rho^k u^k$ is the product of two converging subsequences, respectively for the weak- $*$ L^∞ topology and for the strong L^2 topology, it converges to $\bar{\lambda} \rho u$ weakly in $L^2(\mathbb{T}^d)$, and thus strongly in $H^{-1}(\mathbb{T}^d)$. Note also that

$$\lambda^k \int_{\mathbb{T}^d} \rho^k u^k = \langle -\text{div}(\mathcal{A}^k \nabla u^k), \mathbf{1} \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)} = 0,$$

so that, since $\lambda^k \neq 0$ by (110) and $\rho^k u^k$ weakly converges to ρu in $L^2(\mathbb{T}^d)$,

$$0 = \int_{\mathbb{T}^d} \rho^k u^k = \langle \rho^k u^k, \mathbf{1} \rangle_{L^2(\mathbb{T}^d)} \xrightarrow{k \rightarrow +\infty} \int_{\mathbb{T}^d} \rho u.$$

Since ρu has integral 0, the solution v^k to the following problem is uniquely defined by Lemma 27:

$$-\text{div}(\mathcal{A}^k \nabla v^k) = \bar{\lambda} \rho u, \quad \int_{\mathbb{T}^d} v^k = 0.$$

By definition of the H -convergence $\mathcal{A}^k \xrightarrow[k \rightarrow +\infty]{H} \bar{\mathcal{A}}$, the sequence $(v^k)_{k \geq 1}$ converges weakly in $H^1(\mathbb{T}^d)$ to $v \in H^1(\mathbb{T}^d)$, and $\mathcal{A}^k \nabla v^k \rightharpoonup \bar{\mathcal{A}} \nabla v$ weakly in $L^2(\mathbb{T}^d)^d$, with v the unique solution in $H^1(\mathbb{T}^d)$ to

$$-\text{div}(\bar{\mathcal{A}} \nabla v) = \bar{\lambda} \rho u, \quad \int_{\mathbb{T}^d} v = 0. \quad (111)$$

As

$$-\text{div} \left(\mathcal{A}^k \nabla \left(u^k - v^k - \int_{\mathbb{T}^d} u^k \right) \right) = \lambda^k \rho^k u^k - \bar{\lambda} \rho u \xrightarrow[k \rightarrow +\infty]{} 0$$

strongly in $H^{-1}(\mathbb{T}^d)$, it follows by (106) that $u^k - v^k - \int_{\mathbb{T}^d} u^k \rightarrow 0$ strongly in $H^1(\mathbb{T}^d)$. Therefore, since $(u^k)_{k \geq 1}$ weakly converges to u in $H^1(\mathbb{T}^d)$, we obtain that $v^k \rightarrow u - \int_{\mathbb{T}^d} u$ weakly in $H^1(\mathbb{T}^d)$. By uniqueness of the weak limit in $H^1(\mathbb{T}^d)$, it holds $v = u - \int_{\mathbb{T}^d} u$. By plugging this equality in (111), we can finally conclude that $(u^k)_{k \geq 1}$ converges weakly in $H^1(\mathbb{T}^d)$ to a solution u of (109).

Proving that $\bar{\lambda}$ is the smallest nonzero eigenvalue. To conclude the proof, we show that $\bar{\lambda}$ is the smallest nonzero eigenvalue of (109), proceeding by contradiction. Assume that there exist $0 < \hat{\lambda} < \bar{\lambda}$ and $\hat{u} \in H^1(\mathbb{T}^d)$ such that

$$\begin{cases} -\text{div}(\bar{\mathcal{A}}(q) \nabla \hat{u}(q)) = \hat{\lambda} \rho(q) \hat{u}(q) & \text{on } \mathbb{T}^d, \\ \int_{\mathbb{T}^d} \hat{u}(q)^2 dq = 1. \end{cases}$$

In particular,

$$\int_{\mathbb{T}^d} \rho \widehat{u} = 0. \quad (112)$$

For $k \geq 1$, define $w^k \in H^1(\mathbb{T}^d)$ as the unique solution to (well-defined by Lemma 27)

$$-\operatorname{div}(\mathcal{A}^k \nabla w^k) = \widehat{\lambda} \rho \widehat{u}, \quad \int_{\mathbb{T}^d} w^k = 0. \quad (113)$$

The sequence $(w^k)_{k \geq 1}$ is bounded in $H^1(\mathbb{T}^d)$ by (106), and hence weakly converges in $H^1(\mathbb{T}^d)$ and strongly converges in $L^2(\mathbb{T}^d)$ up to extraction of a subsequence (not explicitly indicated). The limit \widetilde{u} satisfies, by definition of H -convergence,

$$\begin{cases} -\operatorname{div}(\overline{\mathcal{A}}(q) \nabla \widetilde{u}(q)) = \widehat{\lambda} \rho(q) \widehat{u}(q) & \text{on } \mathbb{T}^d, \\ \int_{\mathbb{T}^d} \widetilde{u}(q) dq = 0. \end{cases} \quad (114)$$

The function $\widehat{u} - \int_{\mathbb{T}^d} \widehat{u}$ also satisfies (114), and is therefore equal to \widetilde{u} by uniqueness of the solution to (114). The second condition in (114) then implies that

$$\int_{\mathbb{T}^d} \widehat{u} = - \frac{\int_{\mathbb{T}^d} \rho \widetilde{u}}{\int_{\mathbb{T}^d} \rho}. \quad (115)$$

Let us next introduce $\widetilde{w}^k = w^k - \int_{\mathbb{T}^d} \rho^k w^k / \int_{\mathbb{T}^d} \rho^k$. Note that $\int_{\mathbb{T}^d} \rho^k \widetilde{w}^k = 0$ by construction and $\widetilde{w}^k \rightarrow \widetilde{u} + \int_{\mathbb{T}^d} \widehat{u} = \widehat{u}$ strongly in $L^2(\mathbb{T}^d)$ as $k \rightarrow +\infty$ in view of (115). Then, using first the definition of λ^k , and next (113),

$$\begin{aligned} \lambda^k &\leq \frac{\int_{\mathbb{T}^d} \nabla \widetilde{w}^k(q)^\top \mathcal{A}^k(q) \nabla \widetilde{w}^k(q) dq}{\int_{\mathbb{T}^d} \widetilde{w}^k(q)^2 \rho^k(q) dq} = \frac{\int_{\mathbb{T}^d} \nabla w^k(q)^\top \mathcal{A}^k(q) \nabla w^k(q) dq}{\int_{\mathbb{T}^d} \widetilde{w}^k(q)^2 \rho^k(q) dq} = \widehat{\lambda} \frac{\int_{\mathbb{T}^d} w^k(q) \widehat{u}(q) \rho(q) dq}{\int_{\mathbb{T}^d} \widetilde{w}^k(q)^2 \rho^k(q) dq} \\ &\xrightarrow{k \rightarrow +\infty} \widehat{\lambda} \frac{\int_{\mathbb{T}^d} \widetilde{u}(q) \widehat{u}(q) \rho(q) dq}{\int_{\mathbb{T}^d} \widehat{u}(q)^2 \rho(q) dq} = \widehat{\lambda} \frac{\int_{\mathbb{T}^d} \widehat{u}(q)^2 \rho(q) dq}{\int_{\mathbb{T}^d} \widehat{u}(q)^2 \rho(q) dq} = \widehat{\lambda}, \end{aligned}$$

where we used (112) in the last but one equality. We therefore obtain that $\bar{\lambda} \leq \widehat{\lambda}$, which contradicts the .

The argument above can in fact be applied to any converging subsequence of the original sequence $(\lambda^k)_{k \geq 1}$, henceforth leading to the uniqueness of the limit and the convergence of the whole sequence $(\lambda^k)_{k \geq 1}$, which completes the proof.

D.3 Proof of Lemma 18

Consider a maximizer $\mathcal{D}_0 \in \mathfrak{D}_{\#,k,p}^{a,b}$ of (77) (which indeed exists in view of Theorem 4), and $u_0 \in H^1(\mathbb{T}^d) \setminus \{0\}$ a minimizer associated with $\Lambda^k(\mathcal{D}_0)$ in (76). For $\ell = (\ell_1, \dots, \ell_d) \in \{0, \dots, k-1\}^d$, introduce

$$\mathcal{D}_\ell(q) = \mathcal{D}_0 \left(q + \frac{\ell}{k} \right), \quad u_\ell(q) = u_0 \left(q + \frac{\ell}{k} \right).$$

For these two functions we have, using the change of variable $Q = q + \ell/k$ and the \mathbb{Z}^d -periodicity of V ,

$$\begin{aligned} \frac{\int_{\mathbb{T}^d} \nabla u_\ell(q)^\top \mathcal{D}_\ell(q) \nabla u_\ell(q) e^{-V_{\#,k}(q)} dq}{\int_{\mathbb{T}^d} u_\ell(q)^2 e^{-V_{\#,k}(q)} dq} &= \frac{\int_{\mathbb{T}^d} \nabla u_0\left(q + \frac{\ell}{k}\right)^\top \mathcal{D}_0\left(q + \frac{\ell}{k}\right) \nabla u_0\left(q + \frac{\ell}{k}\right) e^{-V(kq)} dq}{\int_{\mathbb{T}^d} u_0\left(q + \frac{\ell}{k}\right)^2 e^{-V(kq)} dq} \\ &= \frac{\int_{\mathbb{T}^d} \nabla u_0(Q)^\top \mathcal{D}_0(Q) \nabla u_0(Q) e^{-V(kQ-\ell)} dQ}{\int_{\mathbb{T}^d} u_0(Q)^2 e^{-V(kQ-\ell)} dQ} = \Lambda^k(\mathcal{D}_0). \end{aligned}$$

Thus, for all $\ell \in \{0, \dots, k-1\}^d$, the function \mathcal{D}_ℓ also satisfies $\Lambda^k(\mathcal{D}_\ell) = \Lambda^{k,*}$. Define now the diffusion matrix

$$\mathcal{D}_0^k(q) = \frac{1}{k^d} \sum_{\ell_1=0}^{k-1} \cdots \sum_{\ell_d=0}^{k-1} \mathcal{D}_\ell(q).$$

By concavity of the application $\mathcal{D} \mapsto \Lambda^k(\mathcal{D})$ (see Lemma 3),

$$\Lambda^k(\mathcal{D}_0^k) \geq \frac{1}{k^d} \sum_{\ell_1=0}^{k-1} \cdots \sum_{\ell_d=0}^{k-1} \Lambda^k(\mathcal{D}_\ell) = \Lambda^{k,*}.$$

Since $\mathcal{D}_0^k \in \mathfrak{D}_{\#,k,p}^{a,b}$ by the triangle inequality, \mathcal{D}_0^k also satisfies $\Lambda^k(\mathcal{D}_0^k) = \Lambda^{k,*}$. In addition, \mathcal{D}_0^k is $(\mathbb{Z}/k)^d$ -periodic by construction. There exists therefore $\mathcal{D}^{k,*} \in \mathfrak{D}_p^{a,b}$ such that $\mathcal{D}_0^k(q) = \mathcal{D}^{k,*}(kq)$, which concludes the proof.

D.4 Proof of Theorem 19

The following lemma, proved in Appendix D.4.1, shows that in the particular case where one considers a sequence of diffusion matrices $(\mathcal{A}^k)_{k \geq 1}$ such that \mathcal{A}^k is $(\mathbb{Z}/k)^d$ -periodic for any $k \geq 1$, the H -limit $\bar{\mathcal{A}}$ is in fact a constant matrix. It also includes comparison results between weak-* and homogenized limits, which are needed in the proof of Theorem 19. This result is an adaptation to the case of periodic boundary conditions of standard results for divergence operators with Dirichlet boundary conditions (see for instance [2, Lemmas 1.3.13 and 1.3.14]); see Appendix D.4.1 for the proof.

Lemma 28. *Let $a, b > 0$ and $(\mathcal{A}^k)_{k \geq 1} \subset L^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$ be a sequence of diffusion matrices such that \mathcal{A}^k is $(\mathbb{Z}/k)^d$ -periodic for any $k \geq 1$. Then, $(\mathcal{A}^k)_{k \geq 1}$ H -converges (up to the extraction of a subsequence) to a constant matrix $\bar{\mathcal{A}} \in \mathcal{M}_{a,b}$.*

Moreover, if $(\mathcal{B}^k)_{k \geq 1} \subset L^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$ is another sequence of diffusion matrices such that \mathcal{B}^k is $(\mathbb{Z}/k)^d$ -periodic for any $k \geq 1$, and

$$\forall k \geq 1, \quad \forall \xi \in \mathbb{T}^d, \quad \xi^\top \mathcal{A}^k \xi \leq \xi^\top \mathcal{B}^k \xi,$$

then the homogenized limits are also ordered, that is, denoting by $\bar{\mathcal{B}} \in \mathcal{M}_{a,b}$ the H -limit of $(\mathcal{B}^k)_{k \geq 1}$,

$$\forall \xi \in \mathbb{T}^d, \quad \xi^\top \bar{\mathcal{A}} \xi \leq \xi^\top \bar{\mathcal{B}} \xi.$$

Furthermore, if

$$\begin{aligned} \mathcal{A}^k &\rightharpoonup \mathcal{A}_+ \text{ weakly-* in } L^\infty(\mathbb{T}^d, \mathcal{M}_{a,b}), \\ (\mathcal{A}^k)^{-1} &\rightharpoonup \mathcal{A}_-^{-1} \text{ weakly-* in } L^\infty(\mathbb{T}^d, \mathcal{M}_{a,b}), \end{aligned}$$

then it holds

$$\forall \xi \in \mathbb{T}^d, \quad \xi^\top \mathcal{A}_- \xi \leq \xi^\top \bar{\mathcal{A}} \xi \leq \xi^\top \mathcal{A}_+ \xi.$$

For fixed $k \geq 1$, recalling the notation $\mathcal{D}_{\#,k}(q) = \mathcal{D}(kq)$ for any $\mathcal{D} \in \mathfrak{D}_p^{a,b}$, we obtain by definition of the diffusion matrix $\mathcal{D}^{k,\star}$ in Lemma 18 that

$$\forall \mathcal{D} \in \mathfrak{D}_p^{a,b}, \quad \Lambda^k(\mathcal{D}_{\#,k}^{k,\star}) \geq \Lambda^k(\mathcal{D}_{\#,k}). \quad (116)$$

Fix now $\mathcal{D} \in \mathfrak{D}_p^{a,b}$, and introduce the sequences $(\mathcal{A}^{k,\star})_{k \geq 1}$ and $(\mathcal{A}^k)_{k \geq 1}$ defined as

$$\mathcal{A}^{k,\star} = \mathcal{D}_{\#,k}^{k,\star} e^{-V_{\#,k}}, \quad \mathcal{A}^k = \mathcal{D}_{\#,k} e^{-V_{\#,k}}.$$

By Lemma 28, upon extraction a subsequence (still indexed by k with some abuse of notation), the sequence $(\mathcal{A}^{k,\star})_{k \geq 1}$ H -converges to a constant homogenized matrix $\overline{\mathcal{A}}^\star \in \mathcal{M}_{a,b}$. By Theorem 13, the sequence $(\mathcal{A}^k)_{k \geq 1}$ also H -converges to a constant matrix $\overline{\mathcal{A}} \in \mathcal{M}_{a,b}$. In addition, in view of the definitions (66) and (76),

$$\Lambda^k(\mathcal{D}_{\#,k}^{k,\star}) = \Lambda_{\#,k}(\mathcal{D}^{k,\star}), \quad \Lambda^k(\mathcal{D}_{\#,k}) = \Lambda_{\#,k}(\mathcal{D}).$$

Now, by Theorem 14 and Corollary 15, H -convergence implies the convergence of the associated spectral gap, and thus

$$\Lambda_{\#,k}(\mathcal{D}^{k,\star}) \xrightarrow[k \rightarrow +\infty]{} \overline{\Lambda}^\star = \min_{u \in H_0^1(\mathbb{T}^d) \setminus \{0\}} \frac{\int_{\mathbb{T}^d} \nabla u^\top \overline{\mathcal{A}}^\star \nabla u}{\int_{\mathbb{T}^d} u^2 \int_{\mathbb{T}^d} e^{-V}}, \quad \Lambda_{\#,k}(\mathcal{D}) \xrightarrow[k \rightarrow +\infty]{} \Lambda_{\text{hom}}(\mathcal{D}).$$

Thus, passing to the limit $k \rightarrow \infty$ in (116), we obtain that $\overline{\Lambda}^\star \geq \Lambda_{\text{hom}}(\mathcal{D})$. Since this result holds for any $\mathcal{D} \in \mathfrak{D}_p^{a,b}$, we finally obtain that

$$\overline{\Lambda}^\star \geq \max_{\mathcal{D} \in \mathfrak{D}_p^{a,b}} \Lambda_{\text{hom}}(\mathcal{D}) = \Lambda_{\text{hom}}^\star,$$

where we recall that $\Lambda_{\text{hom}}^\star$ is defined in (70).

To obtain the equality $\overline{\Lambda}^\star = \max_{\mathcal{D} \in \mathfrak{D}_p^{a,b}} \Lambda_{\text{hom}}(\mathcal{D})$, it therefore suffices to find a diffusion matrix $\widehat{\mathcal{D}} \in \mathfrak{D}_p^{a,b}$ such that $\Lambda_{\text{hom}}(\widehat{\mathcal{D}}) = \overline{\Lambda}^\star$. The natural candidate is $\widehat{\mathcal{D}}(q) = e^{V(q)} \overline{\mathcal{A}}^\star$. Note first that $\widehat{\mathcal{A}}_{\#,k} = \widehat{\mathcal{D}}_{\#,k} e^{-V_{\#,k}} = \overline{\mathcal{A}}^\star$ for any $k \geq 1$, so that the sequence $(\widehat{\mathcal{A}}_{\#,k})_{k \geq 1}$ H -converges to $\overline{\mathcal{A}}^\star$, and therefore $\Lambda_{\text{hom}}(\widehat{\mathcal{D}}) = \overline{\Lambda}^\star$ by Corollary 15. To conclude the proof, it suffices to check that $\widehat{\mathcal{D}} \in \mathfrak{D}_p^{a,b}$. Note first that, since $\mathcal{D}^{k,\star} \in \mathfrak{D}_p^{a,b}$, it holds for any $\xi \in \mathbb{R}^d$ and almost all $q \in \mathbb{T}^d$,

$$a|\xi|^2 \leq e^{-V_{\#,k}(q)} \xi^\top \mathcal{D}_{\#,k}^{k,\star}(q) \xi \leq \frac{1}{b} |\xi|^2.$$

By Lemma 28, the above inequality passes to the H -limit, so that,

$$\forall \xi \in \mathbb{R}^d, \quad a|\xi|^2 \leq \xi^\top \overline{\mathcal{A}}^\star \xi \leq \frac{1}{b} |\xi|^2.$$

Plugging $\widehat{\mathcal{D}}(q) = e^{V(q)} \overline{\mathcal{A}}^\star$ in the last inequality allows to conclude that $\widehat{\mathcal{D}} \in L_V^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$. Let us next check that

$$\|\widehat{\mathcal{D}}\|_{L_V^p}^p = \int_{\mathbb{T}^d} |\widehat{\mathcal{D}}(q)|_{\text{F}}^p e^{-pV(q)} dq \leq 1.$$

In view of Lemma 28, it holds $0 \leq \overline{\mathcal{A}}^\star \leq \mathcal{A}_+$ in the sense of symmetric matrices for any weak-* limit \mathcal{A}_+ of (a subsequence of) $(\mathcal{A}^{k,\star})_{k \geq 1}$. By lower semicontinuity of the norm, and since the norm $|\cdot|_{\text{F}}$ is compatible with the order on symmetric positive matrices and weak-* $L^\infty(\mathbb{T}^d)$ convergence implies weak $L^p(\mathbb{T}^d)$ convergence,

$$\|\widehat{\mathcal{D}}\|_{L_V^p}^p = |\overline{\mathcal{A}}^\star|_{\text{F}}^p \leq \int_{\mathbb{T}^d} |\mathcal{A}_+(q)|_{\text{F}}^p dq \leq \liminf_{k \rightarrow +\infty} \int_{\mathbb{T}^d} |\mathcal{A}^{k,\star}(q)|_{\text{F}}^p dq = \liminf_{k \rightarrow +\infty} \|\mathcal{D}^{k,\star}\|_{L_V^p}^p \leq 1, \quad (117)$$

which gives the claimed bound and completes the proof of Theorem 19.

Remark 8. *The convergence result (117) in fact holds for other choices of normalizations, in particular normalizations for the diffusion matrix of the form*

$$\int_{\mathbb{T}^d} |\mathcal{D}(q)|_{\mathbb{F}}^p e^{-pV(q)} \omega(q) dq$$

for $\omega \in L^\infty(\mathbb{T}^d)$ with $\omega \geq 0$. Therefore, Theorem 19 can be adapted for other normalizations than the one introduced in (12).

D.4.1 Proof of Lemma 28

Theorem 12 guarantees the H -convergence of $(\mathcal{A}^k)_{k \geq 1}$, up to the extraction of a subsequence, towards a general diffusion matrix $\overline{\mathcal{A}} \in L^\infty(\mathbb{T}^p, \mathcal{M}_{a,b})$. Thus, we only need to show that $\overline{\mathcal{A}}$ is constant. To do so, we follow the proof of [2, Theorem 1.3.18]. Since \mathcal{A}^k is $(\mathbb{Z}/k)^d$ -periodic, there exists a \mathbb{Z}^d -periodic matrix-valued function $\mathfrak{A}^k \in L^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$ such that $\mathcal{A}^k(q) = \mathfrak{A}^k(kq)$. Fix a smooth function φ defined on \mathbb{T}^d , and consider the sequence of oscillating test functions:

$$\tilde{w}^k(q) = \varphi(q) + \frac{1}{k} \sum_{i=1}^d w_i^k(kq) \partial_{q_i} \varphi(q), \quad (118)$$

where $w_i^k \in H^1(\mathbb{T}^d)$ is the periodic function uniquely defined (in view of Lemma 27) by the following equation on \mathbb{T}^d :

$$-\operatorname{div}(\mathfrak{A}^k(e_i + \nabla w_i^k)) = 0, \quad \int_{\mathbb{T}^d} w_i^k = 0. \quad (119)$$

The choice (118) is suggested in the proof of [3, Theorem 1.3.1] and was used for instance in the proof of [43, Proposition 7]. By construction, it holds

$$\nabla \tilde{w}^k(q) = \sum_{i=1}^d \partial_{q_i} \varphi(q) (e_i + \nabla w_i^k(kq)) + \frac{1}{k} \sum_{i=1}^d w_i^k(kq) \nabla \partial_{q_i} \varphi(q). \quad (120)$$

The sequence $(w_i^k)_{k \geq 1}$ is uniformly bounded in $H^1(\mathbb{T}^d)$. By (119), w_i^k solves

$$-\operatorname{div}(\mathfrak{A}^k \nabla w_i^k) = \operatorname{div}(\mathfrak{A}^k e_i), \quad \int_{\mathbb{T}^d} w_i^k = 0.$$

Since $\langle \operatorname{div}(\mathfrak{A}^k e_i), 1 \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)} = 0$, (106) implies that there exists a constant $C > 0$ (which does not depend on k) such that

$$\|w_i^k\|_{H^1(\mathbb{T}^d)} \leq C \|\operatorname{div}(\mathfrak{A}^k e_i)\|_{H^{-1}(\mathbb{T}^d)}. \quad (121)$$

Since $\mathfrak{A}^k \in L^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$ with $b > 0$, one easily bounds the right-hand side of (121) by a constant which does not depend on k .

The sequence $(\tilde{w}^k)_{k \geq 1}$ weakly converges in $H^1(\mathbb{T}^d)$ to φ . Since $\|w_i^k(k \cdot)\|_{L^2(\mathbb{T}^d)} = \|w_i^k\|_{L^2(\mathbb{T}^d)}$ and knowing that $(w_i^k)_{k \geq 1}$ is uniformly bounded in $L^2(\mathbb{T}^d)$, it is clear in view of (118) that $(\tilde{w}^k)_{k \geq 1}$ strongly converges to φ in $L^2(\mathbb{T}^d)$. As for the weak $L^2(\mathbb{T}^d)$ convergence of the gradient, we use Lemma 29 in Appendix D.4.2: the sequence $(e_j \cdot \nabla w_i^k)_{k \geq 1}$ is uniformly bounded in $L^2(\mathbb{T}^d)$ for any $1 \leq j \leq d$, and it holds

$$\int_{\mathbb{T}^d} e_j \cdot \nabla w_i^k(q) dq = 0.$$

Therefore, the sequence $(e_j \cdot \nabla w_i^k(k \cdot))_{k \geq 1}$ weakly converges in $L^2(\mathbb{T}^d)$ to 0, so that $(\nabla \tilde{w}^k)_{k \geq 1}$ weakly converges in $L^2(\mathbb{T}^d)^d$ to $\sum_{i=1}^d \partial_{q_i} \varphi e_i = \nabla \varphi$.

The sequence $(\mathcal{A}^k(e_i + \nabla w_i^k(k\cdot)))_{k \geq 1}$ weakly converges in $L^2(\mathbb{T}^d)^d$ to a constant vector \mathfrak{F}_i . Since $\|\nabla w_i^k(k\cdot)\|_{L^2(\mathbb{T}^d)} = \|\nabla w_i^k\|_{L^2(\mathbb{T}^d)}$, and knowing that $(\nabla w_i^k)_{k \geq 1}$ is uniformly bounded in $L^2(\mathbb{T}^d)$, the sequence $(\mathcal{A}^k(e_i + \nabla w_i^k(k\cdot)))_{k \geq 1}$ is uniformly bounded in $L^2(\mathbb{T}^d)^d$. Therefore, we can introduce, up to extraction, the weak limit \mathfrak{F}_i of this sequence in $L^2(\mathbb{T}^d)^d$. Let us show that this weak limit is constant using again Lemma 29. Indeed, for $1 \leq j \leq d$, let $a_k^j = e_j \cdot \mathfrak{A}^k(e_i + \nabla w_i^k) \in L^2(\mathbb{T}^d)$. The family $(a_k^j)_{k \geq 0}$ is uniformly bounded in $L^2(\mathbb{T}^d)$ so that, using the Cauchy-Schwarz inequality,

$$\sup_{k \geq 1} \int_{\mathbb{T}^d} |a_k^j| \leq \sup_{k \geq 1} \|a_k^j\|_{L^2(\mathbb{T}^d)} < +\infty.$$

Moreover, up to extraction, there exists $\rho_j \in \mathbb{R}$ such that $\int_{\mathbb{T}^d} a_k^j$ converges to ρ_j . This allows us to conclude that $\mathfrak{F}_{i,j}$, the j -th component of \mathfrak{F}_i , is constant. By multiplying (120) by \mathcal{A}^k , we therefore obtain that the sequence $(\mathcal{A}^k \nabla \tilde{w}^k)_{k \geq 1}$ weakly converges to $\mathfrak{F} \nabla \varphi$ in $L^2(\mathbb{T}^d)^d$.

A preliminary computation. We show a convergence result that will be used multiple times in the proof. This can be seen as the counterpart of the celebrated div-curl lemma in our context. Let $(u^k)_{k \geq 1}$ be a sequence converging weakly in $H^1(\mathbb{T}^d)$ and strongly in $L^2(\mathbb{T}^d)$ to $u \in H^1(\mathbb{T}^d)$. It then holds

$$\langle -\operatorname{div}(\mathcal{A}^k \nabla \tilde{w}^k), u^k \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)} \xrightarrow{k \rightarrow +\infty} \langle -\operatorname{div}(\mathfrak{F} \nabla \varphi), u \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)}, \quad (122)$$

where the constant matrix \mathfrak{F} has entries $\mathfrak{F}_{i,j}$. Indeed, using (120), it holds

$$\int_{\mathbb{T}^d} \mathcal{A}^k \nabla \tilde{w}^k \cdot \nabla u^k = \sum_{i=1}^d \int_{\mathbb{T}^d} (e_i + \nabla w_i^k(k\cdot)) \partial_{q_i} \varphi \cdot \mathcal{A}^k \nabla u^k + \frac{1}{k} \sum_{i=1}^d \int_{\mathbb{T}^d} w_i^k(k\cdot) \nabla \partial_{q_i} \varphi \cdot \mathcal{A}^k \nabla u^k. \quad (123)$$

The second term on the right-hand side of the previous equality converges to 0 as $k \rightarrow +\infty$ since $(\mathcal{A}^k)_{k \geq 1}$ is uniformly bounded in $L^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$ with $b > 0$, $(\nabla u^k)_{k \geq 1}$ weakly converges in $L^2(\mathbb{T}^d)^d$ to ∇u and the sequence $(w_i^k(k\cdot))_{k \geq 1}$ is uniformly bounded in $L^2(\mathbb{T}^d)$. For the first term, we use an integration by parts and (119) to write

$$\begin{aligned} \sum_{i=1}^d \int_{\mathbb{T}^d} (e_i + \nabla w_i^k(k\cdot)) \partial_{q_i} \varphi \cdot \mathcal{A}^k \nabla u^k &= - \sum_{i=1}^d \int_{\mathbb{T}^d} \mathcal{A}^k (e_i + \nabla w_i^k(k\cdot)) \cdot (\nabla \partial_{q_i} \varphi) u^k \\ &\xrightarrow{k \rightarrow +\infty} - \sum_{i=1}^d \int_{\mathbb{T}^d} \mathfrak{F}_i \cdot (\nabla \partial_{q_i} \varphi) u = - \sum_{i,j=1}^d \int_{\mathbb{T}^d} \mathfrak{F}_{i,j} (\partial_{q_i, q_j}^2 \varphi) u = \int_{\mathbb{T}^d} \mathfrak{F} \nabla \varphi \cdot \nabla u. \end{aligned} \quad (124)$$

This shows that (122) holds.

The matrix \mathfrak{F} is equal to $\overline{\mathcal{A}}$. Let us next identify the weak limit \mathfrak{F} . Fix $u \in H^1(\mathbb{T}^d)$, and introduce $f = -\operatorname{div}(\overline{\mathcal{A}} \nabla u) \in H^{-1}(\mathbb{T}^d)$. Note that $\langle f, \mathbf{1} \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)} = 0$. Introduce next the solution u^k (again well-defined by Lemma 27) of

$$-\operatorname{div}(\mathcal{A}^k \nabla u^k) = f, \quad \int_{\mathbb{T}^d} u^k = 0.$$

By definition of H -convergence, $u^k \rightharpoonup u - \int_{\mathbb{T}^d} u$ weakly in $H^1(\mathbb{T}^d)$, and strongly in $L^2(\mathbb{T}^d)$ upon extraction. Testing against \tilde{w}^k , it holds

$$\langle f, \tilde{w}^k \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)} = \int_{\mathbb{T}^d} \nabla \tilde{w}^k \cdot \mathcal{A}^k \nabla u^k = \langle -\operatorname{div}(\mathcal{A}^k \nabla \tilde{w}^k), u^k \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)}. \quad (125)$$

By (122), it holds

$$\int_{\mathbb{T}^d} \nabla \tilde{w}^k \cdot \mathcal{A}^k \nabla u^k \xrightarrow{k \rightarrow +\infty} \int_{\mathbb{T}^d} \mathfrak{F} \nabla \varphi \cdot \nabla \left(u - \int_{\mathbb{T}^d} u \right) = \int_{\mathbb{T}^d} \mathfrak{F} \nabla \varphi \cdot \nabla u.$$

As \tilde{w}^k weakly converges in $H^1(\mathbb{T}^d)$ to φ , it follows, by passing to the limit $k \rightarrow +\infty$ in (125), that

$$\langle f, \varphi \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)} = \int_{\mathbb{T}^d} \mathfrak{F} \nabla u \cdot \nabla \varphi.$$

By definition of f ,

$$\langle f, \varphi \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)} = \int_{\mathbb{T}^d} \overline{\mathcal{A}} \nabla u \cdot \nabla \varphi.$$

As the previous equalities hold true for any smooth function φ , we obtain that $\overline{\mathcal{A}} \nabla u = \mathfrak{F} \nabla u$, which indeed allows us to conclude that $\overline{\mathcal{A}} = \mathfrak{F}$ since $u \in H^1(\mathbb{T}^d)$ was arbitrary. This shows finally that the homogenized matrix is constant.

Remark 9 (The periodic case). *In the case when $\mathcal{A}^k(q) = \mathfrak{A}(kq)$ (which corresponds to the setting of Theorem 13), no extraction is needed for the sequence of functions $(\mathcal{A}^k(e_i + \nabla w_i^k(\cdot)))_{k \geq 1}$ to weakly converge in $L^2(\mathbb{T}^d)$. This is due to Lemma 29, and the fact that the functions w_i^k obtained from (119) are independent of k , and actually correspond to the functions w_i defined in (63). Therefore, there is a unique possible limiting constant matrix \mathfrak{F} , with*

$$\mathfrak{F}_i = \int_{\mathbb{T}^d} \mathcal{A}(e_i + \nabla w_i) = \sum_{j=1}^d e_j \int_{\mathbb{T}^d} e_j \cdot \mathcal{A}(e_i + \nabla w_i) = \sum_{j=1}^d e_j \int_{\mathbb{T}^d} (e_j + \nabla w_j) \cdot \mathcal{A}(e_i + \nabla w_i),$$

where we used (119) for the last equality. This allows us to recognize the matrix in (62) applied to e_i . Since all converging subsequences have the same limit, this shows in fact that the full sequence $(\mathcal{A}^k)_{k \geq 1}$ converges, as stated in Theorem 13.

We now show the second part of Lemma 28, whose proof is an adaptation of [2, Lemma 1.3.13]. Define the sequence of oscillating test functions corresponding to the sequence $(\mathcal{B}^k)_{k \geq 0}$, namely

$$\tilde{v}^k(q) = \varphi(q) + \frac{1}{k} \sum_{i=1}^d v_i^k(kq) \partial_{q_i} \varphi(q),$$

where v_i^k is such that

$$-\operatorname{div}(\mathfrak{B}^k(e_i + \nabla v_i^k)) = 0, \quad \int_{\mathbb{T}^d} v_i^k = 0,$$

with $\mathfrak{B}^k \in L^\infty(\mathbb{T}^d, \mathcal{M}_{a,b})$ such that $\mathcal{B}^k(q) = \mathfrak{B}^k(kq)$ for almost all $q \in \mathbb{T}^d$ (this is the counterpart of (118)-(119) with \mathcal{A}^k replaced by \mathcal{B}^k). The sequence $(\tilde{v}^k)_{k \geq 0}$ then weakly converges in $H^1(\mathbb{T}^d)$ to φ in the same way as the sequence $(\tilde{w}^k)_{k \geq 0}$ weakly converges in $H^1(\mathbb{T}^d)$ to φ . Since for almost all $q \in \mathbb{T}^d$, the matrix \mathcal{A}^k is symmetric and positive, it holds

$$\mathcal{A}^k \nabla \tilde{w}^k \cdot \nabla \tilde{w}^k - 2\mathcal{A}^k \nabla \tilde{w}^k \cdot \nabla \tilde{v}^k + \mathcal{A}^k \nabla \tilde{v}^k \cdot \nabla \tilde{v}^k = \mathcal{A}^k (\nabla \tilde{w}^k - \nabla \tilde{v}^k) \cdot (\nabla \tilde{w}^k - \nabla \tilde{v}^k) \geq 0.$$

Using that $\mathcal{A}^k \leq \mathcal{B}^k$, this implies that

$$\mathcal{A}^k \nabla \tilde{w}^k \cdot \nabla \tilde{w}^k - 2\mathcal{A}^k \nabla \tilde{w}^k \cdot \nabla \tilde{v}^k + \mathcal{B}^k \nabla \tilde{v}^k \cdot \nabla \tilde{v}^k \geq 0. \quad (126)$$

We now pass to the limit $k \rightarrow +\infty$. To do so, consider $\psi \in C^\infty(\mathbb{T}^d)$ with $\psi \geq 0$. Multiplying (126) by ψ and integrating over \mathbb{T}^d , it holds

$$\int_{\mathbb{T}^d} \mathcal{A}^k \nabla \tilde{w}^k \cdot \nabla \tilde{w}^k \psi - 2 \int_{\mathbb{T}^d} \mathcal{A}^k \nabla \tilde{w}^k \cdot \nabla \tilde{v}^k \psi + \int_{\mathbb{T}^d} \mathcal{B}^k \nabla \tilde{v}^k \cdot \nabla \tilde{v}^k \psi \geq 0. \quad (127)$$

For any $k \geq 0$, it holds $\tilde{w}^k \psi \in H^1(\mathbb{T}^d)$. Therefore, one obtains

$$\int_{\mathbb{T}^d} \mathcal{A}^k \nabla \tilde{w}^k \cdot \nabla \tilde{w}^k \psi = \langle -\operatorname{div}(\mathcal{A}^k \nabla \tilde{w}^k), \tilde{w}^k \psi \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)} - \int_{\mathbb{T}^d} \mathcal{A}^k \nabla \tilde{w}^k \cdot \nabla \psi \tilde{w}^k. \quad (128)$$

The sequence $(\tilde{w}^k \psi)_{k \geq 0}$ converges to $\varphi \psi$, weakly in $H^1(\mathbb{T}^d)$ and strongly in $L^2(\mathbb{T}^d)$, so that, using (122), it holds

$$\langle -\operatorname{div}(\mathcal{A}^k \nabla \tilde{w}^k), \tilde{w}^k \psi \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)} \xrightarrow{k \rightarrow +\infty} \langle -\operatorname{div}(\bar{\mathcal{A}} \nabla \varphi), \varphi \psi \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)}. \quad (129)$$

As $(\mathcal{A}^k \nabla \tilde{w}^k)_{k \geq 0}$ weakly converges in $L^2(\mathbb{T}^d)^d$ to $\bar{\mathcal{A}} \nabla \varphi$, and the sequence $(\tilde{w}^k \nabla \psi)_{k \geq 0}$ strongly converges to $\varphi \nabla \psi$ in $L^2(\mathbb{T}^d)^d$, one obtains

$$\int_{\mathbb{T}^d} \mathcal{A}^k \nabla \tilde{w}^k \cdot \nabla \psi \tilde{w}^k \xrightarrow{k \rightarrow +\infty} \int_{\mathbb{T}^d} \bar{\mathcal{A}} \nabla \varphi \cdot \nabla \psi \varphi. \quad (130)$$

Combining (129) and (130) in (128), it holds

$$\int_{\mathbb{T}^d} \mathcal{A}^k \nabla \tilde{w}^k \cdot \nabla \tilde{w}^k \psi \xrightarrow{k \rightarrow +\infty} \langle -\operatorname{div}(\bar{\mathcal{A}} \nabla \varphi), \varphi \psi \rangle_{H^{-1}(\mathbb{T}^d), H^1(\mathbb{T}^d)} - \int_{\mathbb{T}^d} \bar{\mathcal{A}} \nabla \varphi \cdot \nabla \psi \varphi = \int_{\mathbb{T}^d} \bar{\mathcal{A}} \nabla \varphi \cdot \nabla \varphi \psi.$$

Using a similar reasoning to identify the limits of the two other terms in (127) (using the convergence result (122) for the sequence $(\mathcal{B}^k)_{k \geq 1}$), it holds

$$\int_{\mathbb{T}^d} (\bar{\mathcal{A}} \nabla \varphi \cdot \nabla \varphi - 2\bar{\mathcal{A}} \nabla \varphi \cdot \nabla \varphi + \bar{\mathcal{B}} \nabla \varphi \cdot \nabla \varphi) \psi \geq 0.$$

Since $\psi \geq 0$ is arbitrary, this implies the following pointwise inequality:

$$0 \leq \bar{\mathcal{A}} \nabla \varphi \cdot \nabla \varphi - 2\bar{\mathcal{A}} \nabla \varphi \cdot \nabla \varphi + \bar{\mathcal{B}} \nabla \varphi \cdot \nabla \varphi = (\bar{\mathcal{B}} - \bar{\mathcal{A}}) \nabla \varphi \cdot \nabla \varphi.$$

Since φ is an arbitrary smooth function, this shows that $\bar{\mathcal{A}} \leq \bar{\mathcal{B}}$.

We finally show the third part of Lemma 28, which is an adaptation of [2, Lemma 1.3.14]. Using the coercivity and symmetry of \mathcal{A}^k , it holds

$$\mathcal{A}^k \nabla \tilde{w}^k \cdot \nabla \tilde{w}^k - 2\mathcal{A}^k \nabla \tilde{w}^k \cdot \nabla \varphi + \mathcal{A}^k \nabla \varphi \cdot \nabla \varphi = \mathcal{A}^k (\nabla \tilde{w}^k - \nabla \varphi) \cdot (\nabla \tilde{w}^k - \nabla \varphi) \geq 0.$$

Passing to the limit in the previous inequality using similar arguments as above, one obtains

$$\bar{\mathcal{A}} \nabla \varphi \cdot \nabla \varphi - 2\bar{\mathcal{A}} \nabla \varphi \cdot \nabla \varphi + \mathcal{A}_+ \nabla \varphi \cdot \nabla \varphi \geq 0,$$

which implies that $\mathcal{A}_+ \geq \bar{\mathcal{A}}$. Likewise, denoting by $\sigma = \mathcal{A}_- \nabla \varphi$,

$$\mathcal{A}^k \nabla \tilde{w}^k \cdot \nabla \tilde{w}^k - 2\nabla \tilde{w}^k \cdot \sigma + (\mathcal{A}^k)^{-1} \sigma \cdot \sigma = (\mathcal{A}^k)^{-1} (\mathcal{A}^k \nabla \tilde{w}^k - \sigma) \cdot (\mathcal{A}^k \nabla \tilde{w}^k - \sigma) \geq 0.$$

Passing to the limit $k \rightarrow +\infty$ yields

$$\bar{\mathcal{A}} \nabla \varphi \cdot \nabla \varphi - 2\nabla \varphi \cdot \mathcal{A}_- \nabla \varphi + \nabla \varphi \cdot \mathcal{A}_- \nabla \varphi \geq 0.$$

This shows that $\bar{\mathcal{A}} \geq \mathcal{A}_-$, which concludes the proof.

D.4.2 A technical result of periodic averaging

We conclude this section by stating and proving the following technical result, obtained by a simple adaptation of [2, Lemma 1.3.19].

Lemma 29. Consider a sequence of functions $(a_k)_{k \geq 1} \subset L^2(\mathbb{T}^d)$ such that

$$\int_{\mathbb{T}^d} a_k \xrightarrow[k \rightarrow +\infty]{} \varrho \in \mathbb{R}, \quad \sup_{k \geq 1} \int_{\mathbb{T}^d} |a_k| < +\infty. \quad (131)$$

Then the sequence $(a_k(k \cdot))_{k \geq 1}$ weakly converges in $L^2(\mathbb{T}^d)$ to the constant function ϱ .

Proof. The proof is a simple extension of [2, Lemma 1.3.19]. Consider $\phi \in \mathcal{C}^0(\mathbb{T}^d)$. We decompose \mathbb{T}^d as the union of k^d cubes $q_i^k + (\mathbb{T}/k)^d$ for $1 \leq i \leq k^d$, so that

$$\int_{\mathbb{T}^d} a_k(kq)\phi(q) dq = \sum_{i=1}^{k^d} \int_{(\mathbb{T}/k)^d} a_k(k(q_i^k + Q)) \phi(q_i^k + Q) dQ.$$

For a given $1 \leq i \leq k^d$, by the periodicity of a_k ,

$$\begin{aligned} & \left| \int_{(\mathbb{T}/k)^d} a_k(k(q_i^k + Q)) \phi(q_i^k + Q) dQ - \frac{1}{k^d} \phi(q_i^k) \int_{\mathbb{T}^d} a_k \right| \\ & \leq \int_{(\mathbb{T}/k)^d} |a_k(k(q_i^k + Q))| |\phi(q_i^k + Q) - \phi(q_i^k)| dQ \\ & \leq \frac{1}{k^d} \left(\int_{\mathbb{T}^d} |a_k| \right) \max_{|q - q'| \leq 1/k} |\phi(q') - \phi(q)|. \end{aligned}$$

Therefore,

$$\left| \int_{\mathbb{T}^d} a_k(kq)\phi(q) dq - \sum_{i=1}^{k^d} \frac{1}{k^d} \phi(q_i^k) \int_{\mathbb{T}^d} a_k \right| \leq \left(\int_{\mathbb{T}^d} |a_k| \right) \max_{|q - q'| \leq 1/k} |\phi(q') - \phi(q)|,$$

the right-hand side of the above inequality going to 0 as $k \rightarrow +\infty$ since ϕ is continuous on the compact set \mathbb{T}^d , hence uniformly continuous. Moreover,

$$\sum_{i=1}^{k^d} \frac{1}{k^d} \phi(q_i^k) \int_{\mathbb{T}^d} a_k \xrightarrow[k \rightarrow +\infty]{} \varrho \int_{\mathbb{T}^d} \phi$$

by results on the convergence of Riemann sums. This shows finally that

$$\int_{\mathbb{T}^d} a_k(kq)\phi(q) dq \xrightarrow[k \rightarrow +\infty]{} \varrho \int_{\mathbb{T}^d} \phi.$$

The claimed result follows by the density of $\mathcal{C}^0(\mathbb{T}^d)$ in $L^2(\mathbb{T}^d)$. \square

E Proof of Theorem 21

The proof consists of three main steps. For the ease of the presentation, we consider the specific case of processes associated with a sequence of time steps $(\Delta t_n)_{n \geq 1}$, which we further assume to be $\Delta t_n = 1/n$ in all this section, the proof in the general case following by straightforward modifications.

We first show in Proposition 30 that the sequence $\{P_{1/n}\}_{n \in \mathbb{N}^*}$ is tight in the family of probability measures on $\mathcal{C}^0([0, \infty), \mathbb{T}^d)$. By Prokhorov's theorem, $\{P_{1/n}\}_{n \in \mathbb{N}^*}$ then admits a weakly convergent subsequence. We next show that the weak limit solves the martingale problem for the limiting diffusion process. Finally, the uniqueness of the martingale problem implies that the weak limit is the law of the solution of the SDE defined in (80).

Let us now state more precisely the above results, starting with the tightness of the sequence $\{P_{1/n}\}_{n \in \mathbb{N}^*}$ (proved in Appendix E.1, with some technical results postponed to Appendix E.3).

Proposition 30 (Tightness). *The sequence $\{P_{1/n}\}_{n \in \mathbb{N}^*}$ is tight in the family of probability measures on $\mathcal{C}^0([0, \infty), \mathbb{T}^d)$.*

Proposition 30 and Prokhorov's theorem guarantee that $\{P_{1/n}\}_{n \in \mathbb{N}^*}$ admits a weakly converging subsequence P_* . To conclude the proof, we identify P_* as the law of the solution to the SDE (82), with generator (4), by considering the martingale problem.

Proposition 31 (Limit martingale problem). *Any weak limit P_* of the sequence $\{P_{1/n}\}_{n \in \mathbb{N}^*}$ is a solution to the martingale formulation of the stochastic differential equation (82). More precisely, for any \mathcal{C}^∞ function $\varphi : \mathbb{T}^d \rightarrow \mathbb{R}$, let us define the process $(M_t)_{t \geq 0}$ by:*

$$\forall t \geq 0, \quad M_t = \varphi(Q_t^*) - \varphi(q_0) - \int_0^t (\mathcal{L}_{\mathcal{D}}\varphi)(Q_s^*) ds$$

where $(Q_t^*)_{t \geq 0}$ is the canonical process on $\mathcal{C}^0([0, \infty), \mathbb{T}^d)$. Then, $(M_t)_{t \geq 0}$ is a P_* -martingale.

The proof of this result can be read in Appendix E.2 (with, again, some technical results postponed to Appendix E.3). The uniqueness of the solution to the martingale problem then implies that P_* is the law of the process (82).

To prove the results and for the sake of conciseness, we use the following notation, specific to our context. Let $f : \mathbb{T}^d \times \mathbb{R}^d \times [0, 1] \rightarrow \mathbb{R}$ be a measurable function. For any couple of independent random variables (G, U) where G is a centered Gaussian vector with identity covariance and U is uniformly distributed on $[0, 1]$, we say that

$$f(q, G, U) = O(n^{-a})$$

for some $a \geq 0$ if there exists an integer $k_f \in \mathbb{N}$ such that, for any $b \in \mathbb{N}$, there is $K_{b,f} \in \mathbb{R}_+$ for which, almost surely,

$$\forall q \in \mathbb{T}^d, \quad \mathbb{E}(|f(q, G, U)|^b | G) \leq \frac{K_{b,f}}{n^{ab}} (1 + |G|^{k_f})^b. \quad (132)$$

In particular, $f(q, G, U) = O(n^{-a})$ implies that, for all $b \in \mathbb{N}$, there exists a constant $C_{f,b} > 0$ such that

$$\mathbb{E}(|f(q, G, U)|^b) \leq \frac{C_{f,b}}{n^{ab}}.$$

E.1 Proof of Proposition 30

Recall that, by definition, the sequence $\{P_{1/n}\}_{n \in \mathbb{N}^*}$ of probability measures on $\mathcal{C}^0([0, \infty), \mathbb{T}^d)$ is tight if, for any $\varepsilon > 0$, there exists a relatively compact set $K \subseteq \mathcal{C}^0([0, \infty), \mathbb{T}^d)$ such that $\sup_{n \in \mathbb{N}^*} P_{1/n}(K^c) \leq \varepsilon$ (see, e.g., [12, Section 5]). The characterization of relatively compact subsets of $\mathcal{C}^0([0, \infty), \mathbb{T}^d)$ by the Arzelà–Ascoli theorem (see, e.g. [12, Theorem 7.2]) allows to derive conditions for the tightness of $\{P_{1/n}\}_{n \in \mathbb{N}^*}$ (see [12, Theorem 7.3]). We consider the following pair of conditions to get tightness:

- (i) For any $\varepsilon > 0$, there exists a compact set $\mathcal{K} \subseteq \mathbb{T}^d$ such that

$$\limsup_{n \in \mathbb{N}^*} P_{1/n} \left(Q_0^{1/n} \notin \mathcal{K} \right) \leq \varepsilon.$$

- (ii) For any $T \in \mathbb{R}_+$ and $\varepsilon > 0$,

$$\lim_{\delta \rightarrow 0} \limsup_{n \in \mathbb{N}^*} P_{1/n} \left(\sup_{\substack{s, t \in [0, T] \\ |s-t| \leq \delta}} |Q_s^{1/n} - Q_t^{1/n}| \geq \varepsilon \right) = 0.$$

The first condition holds trivially since the initial condition $Q_0^{1/n} = q_0$ is fixed. To prove the second condition, we start by observing that, for all $n \in \mathbb{N}^*$,

$$\sup_{\substack{s, t \in [0, T] \\ |s-t| \leq \delta}} |Q_s^{1/n} - Q_t^{1/n}| \leq 3 \max_{0 \leq k \leq \lfloor T/\delta \rfloor} \sup_{t \in [0, \delta]} |Q_{k\delta+t}^{1/n} - Q_{k\delta}^{1/n}|.$$

By Lemma 33 in Appendix E.3, for any $k \leq \lfloor T/\delta \rfloor$,

$$\begin{aligned} & \sup_{\substack{s, t \in [0, T] \\ |s-t| \leq \delta}} |Q_s^{1/n} - Q_t^{1/n}| \\ & \leq 3 \max \left(\max_{0 \leq k \leq \lfloor T/\delta \rfloor} \max_{\lfloor nk\delta \rfloor \leq \ell \leq \lceil n(k+1)\delta \rceil} |q_{1/n}^\ell - q_{1/n}^{\lfloor nk\delta \rfloor}|, \max_{0 \leq k \leq \lfloor T/\delta \rfloor} \max_{\lceil nk\delta \rceil \leq \ell \leq \lceil n(k+1)\delta \rceil} |q_{1/n}^\ell - q_{1/n}^{\lceil nk\delta \rceil}| \right). \end{aligned}$$

Thus,

$$\begin{aligned} \mathbb{P} \left(\sup_{\substack{s, t \in [0, T] \\ |s-t| \leq \delta}} |Q_s^{1/n} - Q_t^{1/n}| \geq \varepsilon \right) & \leq \underbrace{\mathbb{P} \left(\max_{0 \leq k \leq \lfloor T/\delta \rfloor} \max_{\lfloor nk\delta \rfloor \leq \ell \leq \lceil n(k+1)\delta \rceil} |q_{1/n}^\ell - q_{1/n}^{\lfloor nk\delta \rfloor}| \geq \frac{\varepsilon}{3} \right)}_{(a)} \\ & \quad + \underbrace{\mathbb{P} \left(\max_{0 \leq k \leq \lfloor T/\delta \rfloor} \max_{\lceil nk\delta \rceil \leq \ell \leq \lceil n(k+1)\delta \rceil} |q_{1/n}^\ell - q_{1/n}^{\lceil nk\delta \rceil}| \geq \frac{\varepsilon}{3} \right)}_{(b)}. \end{aligned} \quad (133)$$

The two terms (a) and (b) are controlled using similar arguments; for simplicity we make precise only the control of (a). We start by writing, for $\ell \geq \lfloor nk\delta \rfloor$,

$$q_{1/n}^\ell - q_{1/n}^{\lfloor nk\delta \rfloor} = \sum_{i=\lfloor nk\delta \rfloor}^{\ell-1} \Delta_n(q_{1/n}^i, G^{i+1}, U^{i+1}),$$

where (in view of (80) and recalling the notation $R_{\Delta t}$ from (79))

$$\Delta_n(q, G, U) = \sqrt{\frac{2}{n}} \mathcal{D}(q)^{1/2} G \mathbb{1}_{\{U \leq R_{1/n}(q, G)\}}. \quad (134)$$

By Lemma 35 in Appendix E.3, the following decomposition holds:

$$q_{1/n}^\ell - q_{1/n}^{\lfloor nk\delta \rfloor} = \sum_{i=\lfloor nk\delta \rfloor}^{\ell-1} M_n(q_{1/n}^i, G^{i+1}, U^{i+1}) + \sum_{i=\lfloor nk\delta \rfloor}^{\ell-1} T_n(q_{1/n}^i), \quad (135)$$

with $(M_n(q_{1/n}^i, G^{i+1}, U^{i+1}))_{i \geq 0}$ a sequence of martingale increments satisfying $M_n(q, G, U) = O(n^{-1/2})$ and $T_n(q) = O(n^{-1})$. Explicit expressions for $M_n(q_{1/n}^i, G^{i+1}, U^{i+1})$ and $T_n(q_{1/n}^i)$ are given in Lemma 35. From (135), we obtain

$$\begin{aligned} & \max_{0 \leq k \leq \lfloor T/\delta \rfloor} \max_{\lfloor nk\delta \rfloor \leq \ell \leq \lceil n(k+1)\delta \rceil} |q_{1/n}^\ell - q_{1/n}^{\lfloor nk\delta \rfloor}| \\ & \leq \max_{0 \leq k \leq \lfloor T/\delta \rfloor} \max_{\lfloor nk\delta \rfloor \leq \ell \leq \lceil n(k+1)\delta \rceil} \left| \sum_{i=\lfloor nk\delta \rfloor}^{\ell-1} M_n(q_{1/n}^i, G^{i+1}, U^{i+1}) \right| \\ & \quad + \max_{0 \leq k \leq \lfloor T/\delta \rfloor} \max_{\lfloor nk\delta \rfloor \leq \ell \leq \lceil n(k+1)\delta \rceil} \left| \sum_{i=\lfloor nk\delta \rfloor}^{\ell-1} T_n(q_{1/n}^i) \right|, \end{aligned}$$

so that

$$\begin{aligned}
& \mathbb{P} \left(\max_{0 \leq k \leq \lfloor T/\delta \rfloor} \max_{\lfloor nk\delta \rfloor \leq \ell \leq \lfloor n(k+1)\delta \rfloor} \left| q_{1/n}^\ell - q_{1/n}^{\lfloor nk\delta \rfloor} \right| \geq \frac{\varepsilon}{3} \right) \\
& \leq \underbrace{\mathbb{P} \left(\max_{0 \leq k \leq \lfloor T/\delta \rfloor} \max_{\lfloor nk\delta \rfloor \leq \ell \leq \lfloor n(k+1)\delta \rfloor} \left| \sum_{i=\lfloor nk\delta \rfloor}^{\ell-1} M_n(q_{1/n}^i, G^{i+1}, U^{i+1}) \right| \geq \frac{\varepsilon}{6} \right)}_I \\
& \quad + \underbrace{\mathbb{P} \left(\max_{0 \leq k \leq \lfloor T/\delta \rfloor} \max_{\lfloor nk\delta \rfloor \leq \ell \leq \lfloor n(k+1)\delta \rfloor} \left| \sum_{i=\lfloor nk\delta \rfloor}^{\ell-1} T_n(q_{1/n}^i) \right| \geq \frac{\varepsilon}{6} \right)}_{II}.
\end{aligned}$$

We next bound each of the terms on the right-hand side of the above inequality. We use C to denote a constant which can change from line to line.

Control of I . The term I is a sum of martingale increments, and can therefore be bounded using martingale inequalities. Consider to this end the sequence $(S_k^j)_{\ell \geq 0}$ for $0 \leq k \leq \lfloor T/\delta \rfloor$, defined as $S_k^0 = 0$ and

$$\forall 1 \leq j \leq \lfloor n(k+1)\delta \rfloor - \lfloor nk\delta \rfloor, \quad S_k^j = \sum_{i=\lfloor nk\delta \rfloor}^{\lfloor nk\delta \rfloor + j - 1} M_n(q_{1/n}^i, G^{i+1}, U^{i+1}).$$

For $j > \lfloor n(k+1)\delta \rfloor - \lfloor nk\delta \rfloor$, we set the increments to 0, which yields $S_k^j = S_k^{\lfloor n(k+1)\delta \rfloor - \lfloor nk\delta \rfloor}$. The sequence $(S_k^j)_{j \geq 1}$ is a martingale. We start by applying the union bound and Markov's inequality, which yields

$$\begin{aligned}
\mathbb{P} \left(\max_{0 \leq k \leq \lfloor T/\delta \rfloor} \max_{j \geq 1} |S_k^j| \geq \frac{\varepsilon}{6} \right) &= \mathbb{P} \left(\max_{0 \leq k \leq \lfloor T/\delta \rfloor} \max_{j \geq 1} |S_k^j|^4 \geq \frac{\varepsilon^4}{6^4} \right) \\
&\leq \sum_{k=0}^{\lfloor T/\delta \rfloor} \mathbb{P} \left(\max_{j \geq 1} |S_k^j|^4 \geq \frac{\varepsilon^4}{6^4} \right) \leq \frac{6^4}{\varepsilon^4} \sum_{k=0}^{\lfloor T/\delta \rfloor} \mathbb{E} \left(\max_{j \geq 1} |S_k^j|^4 \right). \tag{136}
\end{aligned}$$

We are then in position to resort to the martingale inequality from [17, Theorem 1.1], recalled here for convenience.

Theorem 32 (Theorem 1.1 in [17]). *Suppose that $X = (X_1, X_2, \dots)$ is a martingale, and introduce (with the convention $X_0 = 0$)*

$$X^* = \sup_{n \geq 1} |X_n|, \quad S(X) = \left(\sum_{k=1}^{+\infty} (X_k - X_{k-1})^2 \right)^{1/2}.$$

Consider a convex function $\Phi : [0, \infty) \rightarrow [0, \infty)$ such that $\Phi(0) = 0$, satisfying the following growth condition: There exists $c \in \mathbb{R}_+$ such that

$$\forall u > 0, \quad \Phi(2u) \leq c\Phi(u).$$

Set $\Phi(\infty) = \lim_{u \rightarrow \infty} \Phi(u)$. Then there exists $C_-, C_+ \in \mathbb{R}_+$ such that

$$C_- \mathbb{E}[\Phi(S(X))] \leq \mathbb{E}[\Phi(X^*)] \leq C_+ \mathbb{E}[\Phi(S(X))].$$

Applying Theorem 32 to the sequence $(S_k^j)_{j \geq 0}$ with $\Phi(u) = u^4$, and using a discrete Cauchy-Schwarz inequality, it follows that $\forall 0 \leq k \leq \lfloor T/\delta \rfloor$,

$$\begin{aligned} \mathbb{E} \left(\max_{j \geq 1} |S_k^j|^4 \right) &\leq C_+ \mathbb{E} \left[\left(\sum_{i=\lfloor nk\delta \rfloor}^{\lceil n(k+1)\delta \rceil - 1} \left| M_n \left(q_{1/n}^i, G^{i+1}, U^{i+1} \right) \right|^2 \right)^2 \right] \\ &\leq C_+(n\delta + 1) \sum_{i=\lfloor nk\delta \rfloor}^{\lceil n(k+1)\delta \rceil - 1} \mathbb{E} \left[\left| M_n \left(q_{1/n}^i, G^{i+1}, U^{i+1} \right) \right|^4 \right]. \end{aligned}$$

Since $M_n \left(q_{1/n}^i, G^{i+1}, U^{i+1} \right) = O(n^{-1/2})$ with bounds uniform in i , it follows that

$$\forall 0 \leq k \leq \lfloor T/\delta \rfloor, \quad \mathbb{E} \left(\max_{1 \leq j \leq \lceil n\delta \rceil} |S_k^j|^4 \right) \leq C \frac{(n\delta + 1)^2}{n^2}.$$

Combining the latter inequality with (136), we finally obtain that

$$\mathbb{P} \left(\max_{0 \leq k \leq \lfloor T/\delta \rfloor} \max_{1 \leq j \leq \lceil n\delta \rceil} |S_\ell| \geq \frac{\varepsilon}{6} \right) \leq C \left(\frac{T}{\delta} + 1 \right) \frac{(n\delta + 1)^2}{n^2 \varepsilon^4}.$$

The control of I is concluded by noticing that $\lim_{\delta \rightarrow 0} \limsup_{n \rightarrow +\infty} (T/\delta + 1)(n\delta + 1)^2 n^{-2} = 0$.

Control of II. Using Markov's inequality, then a Cauchy-Schwarz inequality,

$$\begin{aligned} \mathbb{P} \left(\max_{0 \leq k \leq \lfloor T/\delta \rfloor} \max_{\lfloor nk\delta \rfloor \leq \ell \leq \lceil n(k+1)\delta \rceil} \left| \sum_{i=\lfloor nk\delta \rfloor}^{\ell-1} T_n(q_{1/n}^i) \right| \geq \frac{\varepsilon}{6} \right) \\ \leq \frac{36}{\varepsilon^2} \mathbb{E} \left[\max_{0 \leq k \leq \lfloor T/\delta \rfloor} \max_{\lfloor nk\delta \rfloor \leq \ell \leq \lceil n(k+1)\delta \rceil} \left| \sum_{i=\lfloor nk\delta \rfloor}^{\ell-1} T_n(q_{1/n}^i) \right|^2 \right] \\ \leq \frac{36(n\delta + 1)}{\varepsilon^2} \mathbb{E} \left[\max_{0 \leq k \leq \lfloor T/\delta \rfloor} \max_{\lfloor nk\delta \rfloor \leq \ell \leq \lceil n(k+1)\delta \rceil} \sum_{i=\lfloor nk\delta \rfloor}^{\ell-1} \left| T_n(q_{1/n}^i) \right|^2 \right] \\ \leq \frac{36(n\delta + 1)}{\varepsilon^2} \mathbb{E} \left[\max_{0 \leq k \leq \lfloor T/\delta \rfloor} \sum_{i=\lfloor nk\delta \rfloor}^{\lceil n(k+1)\delta \rceil - 1} \left| T_n(q_{1/n}^i) \right|^2 \right] \\ \leq \frac{36(n\delta + 1)}{\varepsilon^2} \sum_{k=0}^{\lfloor T/\delta \rfloor} \sum_{i=\lfloor nk\delta \rfloor}^{\lceil n(k+1)\delta \rceil - 1} \mathbb{E} \left[\left| T_n(q_{1/n}^i) \right|^2 \right]. \end{aligned} \tag{137}$$

Yet, by Lemma 35, $T_n(q_{1/n}^i) = O(n^{-1})$ uniformly in i , so that

$$\sum_{i=\lfloor nk\delta \rfloor}^{\lceil n(k+1)\delta \rceil - 1} \mathbb{E} \left[\left| T_n(q_{1/n}^i) \right|^2 \right] \leq \frac{C(n\delta + 1)}{n^2}.$$

Plugging the last inequality in (137), we obtain that

$$\mathbb{P} \left(\max_{0 \leq k \leq \lfloor T/\delta \rfloor} \max_{\lfloor nk\delta \rfloor \leq \ell \leq \lceil n(k+1)\delta \rceil} \left| \sum_{i=\lfloor nk\delta \rfloor}^{\ell-1} T_n(q_{1/n}^i) \right| \geq \frac{\varepsilon}{6} \right) \leq C \left(\frac{T}{\delta} + 1 \right) \frac{(n\delta + 1)^2}{n^2 \varepsilon^2}.$$

The control of II is concluded by noticing that $\lim_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} C(T/\delta + 1)(n\delta + 1)^2 n^{-2} = 0$.

Conclusion of the proof. Combining the control of the two terms I and II , we conclude that

$$\lim_{\delta \rightarrow 0} \limsup_{n \rightarrow +\infty} \mathbb{P} \left(\max_{0 \leq k \leq \lfloor T/\delta \rfloor} \max_{\lfloor nk\delta \rfloor \leq \ell \leq \lceil n(k+1)\delta \rceil} \left| q_{1/n}^\ell - q_{1/n}^{\lfloor nk\delta \rfloor} \right| \geq \frac{\varepsilon}{3} \right) = 0.$$

It can similarly be proved that

$$\lim_{\delta \rightarrow 0} \limsup_{n \rightarrow +\infty} \mathbb{P} \left(\max_{0 \leq k \leq \lfloor T/\delta \rfloor} \max_{\lceil nk\delta \rceil \leq \ell \leq \lfloor n(k+1)\delta \rfloor} \left| q_{1/n}^\ell - q_{1/n}^{\lceil nk\delta \rceil} \right| \geq \frac{\varepsilon}{3} \right) = 0.$$

In view of (133), we finally obtain that

$$\lim_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} \mathbb{P} \left(\sup_{\substack{s, t \in [0, T] \\ |s-t| \leq \delta}} \left| Q_s^{1/n} - Q_t^{1/n} \right| \right) = 0,$$

which concludes the proof of Proposition 30.

E.2 Proof of Proposition 31

For $n \in \mathbb{N}^*$, let us denote by \mathcal{L}^n the operator defined by: for any bounded measurable function $\phi : \mathbb{T}^d \rightarrow \mathbb{R}$,

$$(\mathcal{L}^n \phi)(q) = \mathbb{E} \left[\phi(q_{1/n}^1) \mid q_{1/n}^0 = q \right] - \phi(q). \quad (138)$$

For a given function $\varphi \in \mathcal{C}^3(\mathbb{T}^d)$, we start by showing in Appendix E.2.1 that the sequence of functions $(n\mathcal{L}^n \varphi)_{n \geq 1}$ converges uniformly towards $\mathcal{L}_{\mathcal{D}} \varphi$ as $n \rightarrow \infty$, in the sense of uniform convergence of continuous functions over the compact set \mathbb{T}^d . We then write the martingale problem associated to \mathcal{L}^n in Appendix E.2.2, and prove that the limit P_∞ solves the martingale problem for the limit generator $\mathcal{L}_{\mathcal{D}}$.

E.2.1 Convergence of the sequence of functions $(n\mathcal{L}^n \varphi)_{n \geq 1}$

In order to determine the limit of the sequence of functions $(n\mathcal{L}^n \varphi)_{n \geq 1}$, we work out a Taylor expansion of the function (138). We expand to this end $\varphi(q_{1/n}^1)$ at second order around q , seeing $q_{1/n}^1 - q$ as a perturbation along the direction $\mathcal{D}(q)^{1/2} G^1$ with step size $\sqrt{2/n} \mathbb{1}_{\{U^1 \leq R_{1/n}(q, G^1)\}} = O(n^{-1/2})$. Writing G, U instead of G^1, U^1 to alleviate the notation,

$$\begin{aligned} \varphi \left(q + \sqrt{\frac{2}{n}} \mathbb{1}_{\{U \leq R_{1/n}(q, G)\}} \mathcal{D}(q)^{1/2} G \right) &= \varphi(q) + \sqrt{\frac{2}{n}} \mathbb{1}_{\{U \leq R_{1/n}(q, G)\}} \nabla \varphi(q)^\top \mathcal{D}(q)^{1/2} G \\ &\quad + \frac{1}{n} \mathbb{1}_{\{U \leq R_{1/n}(q, G)\}} G^\top \mathcal{D}(q)^{1/2} \nabla^2 \varphi(q) \mathcal{D}(q)^{1/2} G + O(n^{-3/2}), \end{aligned}$$

where the remainder (obtained for instance with a Taylor expansion with exact remainder) is uniformly controlled using derivatives of φ of order at most 3, by which we mean that the remainder satisfies an inequality such as (132), with a bound $K_{b,f} \leq c_{b,f} \|\varphi\|_{\mathcal{C}^3(\mathbb{T}^d)}$. Subtracting $\varphi(q)$ and taking the expectation with respect to U ,

$$\begin{aligned} \mathbb{E} \left[\varphi \left(q + \sqrt{\frac{2}{n}} \mathbb{1}_{\{U \leq R_{1/n}(q, G)\}} \mathcal{D}(q)^{1/2} G \right) \middle| G \right] - \varphi(q) &= \sqrt{\frac{2}{n}} R_{1/n}(q, G) \nabla \varphi(q)^\top \mathcal{D}(q)^{1/2} G \\ &\quad + \frac{1}{n} R_{1/n}(q, G) G^\top \mathcal{D}(q)^{1/2} \nabla^2 \varphi(q) \mathcal{D}(q)^{1/2} G + O(n^{-3/2}). \end{aligned}$$

In view of the expansion of the acceptance rate provided by Lemma 34 (with the analytical expression (140) for $\zeta(q, G)$), and further taking the expectation with respect to G , we obtain

$$(\mathcal{L}^n \varphi)(q) = -\frac{2}{n} \nabla \varphi(q)^\top \mathcal{D}(q)^{1/2} \mathbb{E}[G \max(0, \zeta(q, G))] + \frac{1}{n} \mathcal{D}(q) : \nabla^2 \varphi(q) + O(n^{-3/2}),$$

with a remainder term which is uniformly bounded in $q \in \mathbb{T}^d$. A simple expression for the expectation

$$\mathbb{E}[G \max(0, \zeta(q, G))] = \int_{\mathbb{R}^d} g \max(0, \zeta(q, g)) \frac{e^{-g^2/2}}{(2\pi)^{d/2}} dg$$

can be derived as in [29] using the anti-symmetry of the function ζ . Indeed, for all $q \in \mathbb{T}^d$ and for all $g \in \mathbb{R}^d$, a direct inspection of (140) reveals that $\zeta(q, g) = -\zeta(q, -g)$, so that $g\zeta(q, g) = -g\zeta(q, -g)$. Therefore, using [29, Lemma 10] to obtain the second line,

$$\begin{aligned} \mathbb{E}[G \max(0, \zeta(q, G))] &= \int_{\{g \in \mathbb{R}^d \mid \zeta(q, g) \geq 0\}} g \zeta(q, g) \frac{e^{-|g|^2/2}}{(2\pi)^{d/2}} dg = \frac{1}{2} \int_{\mathbb{R}^d} g \zeta(q, g) \frac{e^{-|g|^2/2}}{(2\pi)^{d/2}} dg \\ &= \frac{1}{2} \left[\mathcal{D}(q)^{1/2} \nabla V(q) - \mathcal{D}(q)^{-1/2} \operatorname{div}(\mathcal{D})(q) \right]. \end{aligned}$$

We conclude that

$$2\nabla\varphi(q)^\top \mathcal{D}(q)^{1/2} \mathbb{E}[G \min(0, \zeta(q, G))] = (\mathcal{D}(q) \nabla V(q) - \operatorname{div}(\mathcal{D})(q))^\top \nabla\varphi(q).$$

This ensures finally that $n(\mathcal{L}^n \varphi)(q) \rightarrow (\mathcal{L}_{\mathcal{D}} \varphi)(q)$ uniformly in $q \in \mathbb{T}^d$ as $n \rightarrow +\infty$.

E.2.2 Limit martingale problem

The argument is standard, and we refer for example to [67, Section 11.2] and [22, Theorem 5.8] for details. Let us however outline the reasoning for the sake of completeness.

For any $n \geq 1$ and any test function $\varphi \in \mathcal{C}^3(\mathbb{T}^d)$, the definition of \mathcal{L}^n ensures that

$$\varphi(q_{1/n}^k) - \varphi(q_{1/n}^0) - \frac{1}{n} \sum_{i=1}^{k-1} (n\mathcal{L}^n \varphi)(q_{1/n}^i)$$

is a martingale with respect to $P_{1/n}$. Now, using the definition (81),

$$\begin{aligned} \varphi(q_{1/n}^k) - \varphi(q_{1/n}^0) - \frac{1}{n} \sum_{i=1}^{k-1} (n\mathcal{L}^n \varphi)(q_{1/n}^i) \\ &= \varphi(Q_{k/n}^{1/n}) - \varphi(Q_0^{1/n}) - \frac{1}{n} \sum_{i=1}^{k-1} (n\mathcal{L}^n \varphi)(Q_{i/n}^{1/n}) \\ &= \varphi(Q_{k/n}^{1/n}) - \varphi(Q_0^{1/n}) - \frac{1}{n} \sum_{i=1}^{k-1} (\mathcal{L}_{\mathcal{D}} \varphi)(Q_{i/n}^{1/n}) + \frac{1}{n} \sum_{i=1}^{k-1} (n\mathcal{L}^n \varphi - \mathcal{L}_{\mathcal{D}} \varphi)(Q_{i/n}^{1/n}). \end{aligned}$$

The last term converges to 0 as $n \rightarrow +\infty$ in view of the uniform convergence $n(\mathcal{L}^n \varphi)(q) \rightarrow (\mathcal{L}_{\mathcal{D}} \varphi)(q)$. The first sum can be rewritten as

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^{k-1} (\mathcal{L}_{\mathcal{D}} \varphi)(Q_{i/n}^{1/n}) &= \sum_{i=1}^{k-1} \int_{i/n}^{(i+1)/n} (\mathcal{L}_{\mathcal{D}} \varphi)(Q_{i/n}^{1/n}) dt \\ &= \int_{1/n}^{k/n} (\mathcal{L}_{\mathcal{D}} \varphi)(Q_t^{1/n}) dt + O\left(\frac{1}{n}\right). \end{aligned}$$

For $t > 0$ fixed, we conclude by letting $n \rightarrow \infty$ with $k = \lfloor nt \rfloor$ that

$$\varphi(Q_t^*) - \varphi(Q_0^*) - \int_0^t (\mathcal{L}_{\mathcal{D}} \varphi)(Q_s^*) ds$$

is a martingale process for any weak limit P_* of $\{P_{1/n}\}_{n \geq 1}$, which concludes the proof.

E.3 Technical results

We gather in this section some technical results used in Appendices E.1 and E.2.

Lemma 33. *For any $\delta > 0$, $n \in \mathbb{N}$ and $k \in \mathbb{N}$, it holds*

$$\begin{aligned} & \sup_{t \in [0, \delta]} \left| Q_{k\delta+t}^{1/n} - Q_{k\delta}^{1/n} \right| \\ & \leq \max \left\{ \max_{[nk\delta] \leq \ell \leq \lceil n(k+1)\delta \rceil} \left| q_{1/n}^\ell - q_{1/n}^{\lfloor nk\delta \rfloor} \right|, \max_{\lceil nk\delta \rceil \leq \ell \leq \lceil n(k+1)\delta \rceil} \left| q_{1/n}^\ell - q_{1/n}^{\lceil nk\delta \rceil} \right| \right\}. \end{aligned}$$

Proof. Note first that, since $Q_{k\delta}^{1/n}$ linearly interpolates between $q_{1/n}^{\lfloor nk\delta \rfloor}$ and $q_{1/n}^{\lceil nk\delta \rceil}$,

$$\sup_{t \in [0, \delta]} \left| Q_{k\delta+t}^{1/n} - Q_{k\delta}^{1/n} \right| \leq \max \left\{ \sup_{s \in [k\delta, (k+1)\delta]} \left| Q_s^{1/n} - q_{1/n}^{\lfloor nk\delta \rfloor} \right|, \sup_{s \in [k\delta, (k+1)\delta]} \left| Q_s^{1/n} - q_{1/n}^{\lceil nk\delta \rceil} \right| \right\}. \quad (139)$$

We next decompose the supremums over $s \in [k\delta, (k+1)\delta]$ into supremums over intervals of length $1/n$, by noting that

$$[k\delta, (k+1)\delta] \subset \bigcup_{\ell = \lfloor nk\delta \rfloor}^{\lceil n(k+1)\delta \rceil - 1} \left[\frac{\ell}{n}, \frac{\ell+1}{n} \right].$$

For instance,

$$\sup_{s \in [k\delta, (k+1)\delta]} \left| Q_s^{1/n} - q_{1/n}^{\lfloor nk\delta \rfloor} \right| \leq \max_{[nk\delta] \leq \ell \leq \lceil n(k+1)\delta \rceil - 1} \sup_{\tau \in [0, 1/n]} \left| Q_{\ell/n+\tau}^{1/n} - q_{1/n}^{\lfloor nk\delta \rfloor} \right|.$$

Since $Q_{\ell/n+\tau}^{1/n}$ linearly interpolates between $q_{1/n}^\ell$ and $q_{1/n}^{\ell+1}$, it holds

$$\sup_{\tau \in [0, 1/n]} \left| Q_{\ell/n+\tau}^{1/n} - q_{1/n}^{\lfloor nk\delta \rfloor} \right| \leq \max \left\{ \left| q_{1/n}^\ell - q_{1/n}^{\lfloor nk\delta \rfloor} \right|, \left| q_{1/n}^{\ell+1} - q_{1/n}^{\lfloor nk\delta \rfloor} \right| \right\}.$$

The claimed conclusion then follows from a similar bound for the second maximum over s on the right-hand side of (139). \square

The following lemma provides an expansion of the acceptance rate in inverse powers of n .

Lemma 34 (Expansion of $R_{1/n}(q, G)$). *For any $q \in \mathbb{T}^d$ and $G \in \mathbb{R}^d$, it holds*

$$R_{1/n}(q, G) = 1 - \sqrt{\frac{2}{n}} \max(0, \zeta(q, G)) + O(n^{-1}),$$

with

$$\begin{aligned} \zeta(q, G) = & \nabla V(q)^\top \mathcal{D}(q)^{1/2} G + \frac{1}{2} \text{Tr} \left(\mathcal{D}(q)^{-1} \left[\mathcal{D}\mathcal{D}(q) \cdot \mathcal{D}(q)^{1/2} G \right] \right) \\ & - \frac{1}{2} G^\top \mathcal{D}(q)^{-1/2} \left[\mathcal{D}\mathcal{D}(q) \cdot \mathcal{D}(q)^{1/2} G \right] \mathcal{D}(q)^{-1/2} G, \end{aligned} \quad (140)$$

where $\mathcal{D}\mathcal{D}(q)$ is the differential matrix of $q \mapsto \mathcal{D}(q)$, i.e., for any $h \in \mathbb{R}^d$, $\mathcal{D}\mathcal{D}(q) \cdot h$ is the matrix with entries $\nabla \mathcal{D}_{ij}(q)^\top h$.

Proof. The computations are similar to the ones in [29, Section 4.7]. To control remainder terms, we use the fact that the functions at hand are sufficiently smooth, the diffusion matrix \mathcal{D} is positive and the domain \mathbb{T}^d is compact. The magnitude of the acceptance probability in the Metropolis algorithm reads

$$R_{1/n}(q, G) = \min \left\{ 1, \exp \left[-\alpha \left(q, \Phi_{1/n}(q, G) \right) \right] \right\}, \quad \Phi_{1/n}(q, G) = q + \sqrt{\frac{2}{n}} \mathcal{D}(q)^{1/2} G,$$

with

$$\begin{aligned}\alpha(q, \tilde{q}) &= \frac{1}{2} (\ln \det[\mathcal{D}(\tilde{q})] - \ln \det[\mathcal{D}(q)]) + [V(\tilde{q}) - V(q)] \\ &\quad + \frac{n}{4} \left[\left| \mathcal{D}(\tilde{q})^{-1/2}(q - \tilde{q}) \right|^2 - \left| \mathcal{D}(q)^{-1/2}(\tilde{q} - q) \right|^2 \right].\end{aligned}$$

First,

$$V(\Phi_{1/n}(q, G)) - V(q) = \sqrt{\frac{2}{n}} \nabla V(q)^\top \mathcal{D}(q)^{1/2} G + O(n^{-1}). \quad (141)$$

We next consider the terms involving $\det(\mathcal{D})$. Since \mathcal{D} is \mathcal{C}^2 on \mathbb{T}^d ,

$$\mathcal{D}(\Phi_{1/n}(q, G)) = \mathcal{D}(q) + \sqrt{\frac{2}{n}} \mathbb{D}\mathcal{D}(q) \cdot \mathcal{D}(q)^{1/2} G + O(n^{-1}).$$

It then holds, using the fact that \mathcal{D} is uniformly bounded from below by a positive constant,

$$\begin{aligned}\det[\mathcal{D}(\Phi_{1/n}(q, G))] &= \det[\mathcal{D}(q)] \det\left(\text{Id} + \sqrt{\frac{2}{n}} \mathcal{D}(q)^{-1} \mathbb{D}\mathcal{D}(q) \cdot \mathcal{D}(q)^{1/2} G + O(n^{-1})\right) \\ &= \det[\mathcal{D}(q)] \left(1 + \sqrt{\frac{2}{n}} \text{Tr}\left(\mathcal{D}(q)^{-1} [\mathbb{D}\mathcal{D}(q) \cdot \mathcal{D}(q)^{1/2} G]\right) + O(n^{-1})\right),\end{aligned}$$

so that

$$\ln \det[\mathcal{D}(\Phi_{1/n}(q, G))] - \ln \det[\mathcal{D}(q)] = \sqrt{\frac{2}{n}} \text{Tr}\left(\mathcal{D}(q)^{-1} [\mathbb{D}\mathcal{D}(q) \cdot \mathcal{D}(q)^{1/2} G]\right) + O(n^{-1}). \quad (142)$$

We finally turn to the term $|\mathcal{D}(\Phi_{1/n}(q, G))^{-1/2}(q - \Phi_{1/n}(q, G))|^2 - |\mathcal{D}(q)^{-1/2}(\Phi_{1/n}(q, G) - q)|^2$. We use the expansion $\mathcal{D}(q+x)^{-1} = \mathcal{D}(q)^{-1} - \mathcal{D}(q)^{-1} [\mathbb{D}\mathcal{D}(q) \cdot x] \mathcal{D}(q)^{-1} + O(|x|^2)$ (using again the uniform positive lower bound on \mathcal{D}) to write

$$\begin{aligned}&\left| \mathcal{D}(\Phi_{1/n}(q, G))^{-1/2}(q - \Phi_{1/n}(q, G)) \right|^2 - \left| \mathcal{D}(q)^{-1/2}(\Phi_{1/n}(q, G) - q) \right|^2 \\ &= (\Phi_{1/n}(q, G) - q)^\top [\mathcal{D}(\Phi_{1/n}(q, G))^{-1} - \mathcal{D}(q)^{-1}] (\Phi_{1/n}(q, G) - q) \\ &= -\left(\frac{2}{n}\right)^{3/2} G^\top \mathcal{D}(q)^{-1/2} [\mathbb{D}\mathcal{D}(q) \cdot \mathcal{D}(q)^{1/2} G] \mathcal{D}(q)^{-1/2} G + O(n^{-2}).\end{aligned}$$

Combining (141) and (142) with the latter equality leads to $\alpha(q, \Phi_{1/n}(q, G)) = (2/n)^{1/2} \zeta(q, G) + O(n^{-1})$, with $\zeta(q, G)$ given by (140). The desired conclusion then follows from the inequality $\max(x, 0) - \max(x, 0)^2/2 \leq 1 - \min(1, e^{-x}) \leq \max(x, 0)$, obtained by distinguishing the cases $x \leq 0$ and $x \geq 0$. \square

The last result makes precise the martingale decomposition of the increments $\Delta_n(q, G, U)$ defined in (134).

Lemma 35. *Fix $q \in \mathbb{T}^d$. The increment $\Delta_n(q, G, U)$ writes*

$$\Delta_n(q, G, U) = M_n(q, G, U) + T_n(q),$$

where $M_n(q, G, U) = \Delta_n(q, G, U) - \mathbb{E}[\Delta_n(q, G, U)]$ and $T_n(q) = \mathbb{E}[\Delta_n(q, G, U)]$, the latter expectations being with respect to realizations of the independent random variables G and U , with G a d -dimensional standard Gaussian random variable and U a uniform random variable on $[0, 1]$. Moreover,

$$\mathbb{E}[M_n(q, G, U)] = 0, \quad M_n(q, G, U) = O(n^{-1/2}), \quad T_n(q) = O(n^{-1}), \quad (143)$$

the last two estimates holding uniformly in $q \in \mathbb{T}^d$.

Proof. The first property of (143) holds by construction, so it suffices to prove the two other estimates. For the last one, we note that, by definition,

$$T_n(q) = \sqrt{\frac{2}{n}} \mathcal{D}(q)^{1/2} \mathbb{E} [GR_{1/n}(q, G)] = \sqrt{\frac{2}{n}} \mathcal{D}(q)^{1/2} \mathbb{E} [G(R_{1/n}(q, G) - 1)],$$

so that the conclusion follows from Lemma 34 and the fact that \mathcal{D} , its derivatives and its inverse, are uniformly bounded in q . The other estimate is obtained by noting that

$$M_n(q, G, U) = \sqrt{\frac{2}{n}} \mathcal{D}(q)^{1/2} \left(G \mathbb{1}_{\{U \leq R_{1/n}(q, G)\}} - \mathbb{E} [GR_{1/n}(q, G)] \right),$$

so that, in view of Lemma 34,

$$\forall u \in [0, 1], \quad |M_n(q, G, u)| \leq \frac{C}{\sqrt{n}} |\mathcal{D}(q)|_F^{1/2} |G| + O(n^{-1}).$$

This concludes the proof. \square

Acknowledgements. We thank Grégoire Allaire, Benjamin Bogosel, Andrew Duncan, Federico Ghimenti, Frédéric Legoll, Mathieu Lewin, Danny Perez, Samuel Power, Frédéric van Wijland and Jonathan Weare for stimulating discussions. The works of T.L., R.S. and G.S. benefit from fundings from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (project EMC2, grant agreement No 810367), and from the Agence Nationale de la Recherche through the grants ANR-19-CE40-0010-01 (QuAMProcs) and ANR-21-CE40-0006 (SINEQ). G.P. is partially supported by an ERC-EPSRC Frontier Research Guarantee through Grant No. EP/X038645, ERC through Advanced Grant No. 247031. This project was initiated as T.L. was a visiting professor at Imperial College of London (ICL), with a visiting professorship grant from the Leverhulme Trust. The Department of Mathematics at ICL and the Leverhulme Trust are warmly thanked for their support.

References

- [1] A. Abdulle, G. A. Pavliotis, and G. Vilmart. Accelerated convergence to equilibrium and reduced asymptotic variance for Langevin dynamics using Stratonovich perturbations. *Comptes Rendus Mathématique*, 357(4):349–354, 2019.
- [2] G. Allaire. *Shape Optimization by the Homogenization Method*, volume 146 of *Applied Mathematical Sciences*. Springer-Verlag New York, 2002.
- [3] G. Allaire. *Introduction to homogenization theory*. CEA-EDF-Inria school on homogenization (December 2010), 2010. <http://www.cmap.polytechnique.fr/~allaire/homog/lect1.pdf>.
- [4] G. Allaire, P. Geoffroy-Donders, and O. Pantz. Topology optimization of modulated and oriented periodic microstructures by the homogenization method. *Comput. Math. Appl.*, 78(7):2197–2229, 2019.
- [5] M. P. Allen and D. J. Tildesley. *Computer Simulation of Liquids*. Oxford University Press, Inc., 2nd edition, 2017.
- [6] C. Andrieu and S. Livingstone. Peskun-Tierney ordering for Markovian Monte Carlo: beyond the reversible scenario. *Ann. Statist.*, 49(4):1958–1981, 2021.
- [7] D. Bakry, I. Gentil, and M. Ledoux. *Analysis and Geometry of Markov Diffusion Operators*. Grundlehren der mathematischen Wissenschaften, Vol. 348. Springer, 2014.
- [8] R. Balian. *From Microphysics to Macrophysics. Methods and Applications of Statistical Physics*, volume I - II. Springer, 2007.

- [9] C. Barbarosie and A.-M. Toader. Shape and topology optimization for periodic problems. II. Optimization algorithm and numerical examples. *Struct. Multidiscip. Optim.*, 40(1-6):393–408, 2010.
- [10] C. H. Bennett. Mass tensor molecular dynamics. *Journal of Computational Physics*, 19(3):267–279, 1975.
- [11] R. Bhatia. *Matrix Analysis*. Graduate Texts in Mathematics. Springer, New York, NY, 1997.
- [12] P. Billingsley. *Convergence of Probability Measures*. Wiley Series in Probability and Statistics: Probability and Statistics. John Wiley & Sons, Inc., New York, second edition, 1999.
- [13] X. Blanc and C. Le Bris. *Homogenization Theory for Multiscale Problems*. Modeling, Simulation and Applications. Springer Cham, 2023.
- [14] J.-F. Bonnans, J. C. Gilbert, C. Lemarechal, and C. A. Sagastizábal. *Numerical Optimization: Theoretical and Practical Aspects*. Universitext. Springer, 2nd edition, 2006.
- [15] S. Boyd, P. Diaconis, P. Parrilo, and L. Xiao. Fastest mixing Markov chain on graphs with symmetries. *SIAM Journal on Optimization*, 20(2):792–819, 2009.
- [16] S. Boyd, P. Diaconis, and L. Xiao. Fastest mixing Markov chain on a graph. *SIAM Review*, 46(4):667–689, 2004.
- [17] D. L. Burkholder, B. J. Davis, and R. F. Gundy. Integral inequalities for convex functions of operators on martingales. In *Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability, Volume 2: Probability Theory*, pages 223–240. University of California Press, 1972.
- [18] F. Casas, J. M. Sanz-Serna, and L. Shaw. Split Hamiltonian Monte Carlo revisited. *Stat. Comput.*, 32(86), 2022.
- [19] M. Chak, N. Kantas, T. Lelièvre, and G. Pavliotis. Optimal friction matrix for underdamped langevin sampling. *ESAIM: Mathematical Modelling and Numerical Analysis*, 57(6):3335–3371, 2023.
- [20] T. Chen, E. Fox, and C. Guestrin. Stochastic Gradient Hamiltonian Monte Carlo. In E. P. Xing and T. Jebara, editors, *Proceedings of the 31st International Conference on Machine Learning*, volume 32 of *Proceedings of Machine Learning Research*, pages 1683–1691, Beijing, China, 22–24 Jun 2014. PMLR.
- [21] Y. Chen, D. Z. Huang, J. Huang, S. Reich, and A. M. Stuart. Gradient flows for sampling: Mean-field models, Gaussian approximations and affine invariance. *arXiv preprint*, 2302.11024, 2023.
- [22] F. Comets and T. Meyre. *Calcul stochastique et modèles de diffusions : cours et exercices corrigés*. Dunod, Paris, 2006.
- [23] T. Cui, X. Tong, and O. Zahm. Optimal Riemannian metric for Poincaré inequalities and how to ideally precondition Langevin dynamics. *arXiv preprint*, 2404.02554, 2024.
- [24] A. Duncan, T. Lelièvre, and G. Pavliotis. Variance reduction using nonreversible Langevin samplers. *J. Stat. Phys.*, 163(3):457–491, 2016.
- [25] A. B. Duncan, N. Nüsken, and G. A. Pavliotis. Using Perturbed Underdamped Langevin Dynamics to Efficiently Sample from Probability Distributions. *J. Stat. Phys.*, 169(6):1098–1131, 2017.
- [26] A. B. Duncan, G. A. Pavliotis, and K. C. Zygalakis. Nonreversible Langevin samplers: Splitting schemes, analysis and implementation. *arXiv preprint*, 1701.04247, 2017.

- [27] I. Dunning, J. Huchette, and M. Lubin. Jump: A modeling language for mathematical optimization. *SIAM Review*, 59(2):295–320, 2017.
- [28] H. Fang, M. Qian, and G. Gong. An improved annealing method and its large-time behavior. *Stochastic Process. Appl.*, 71(1):55–74, 1997.
- [29] M. Fathi and G. Stoltz. Improving dynamical properties of stabilized discretizations of overdamped Langevin dynamics. *Numer. Math.*, 136(2):545–602, 2017.
- [30] D. Frenkel and B. Smit. *Understanding Molecular Simulation*. Academic Press, 2nd edition, 2002.
- [31] D. Gilbarg and N. S. Trudinger. *Elliptic Partial Differential Equations of Second Order*, volume 224 of *Classics in Mathematics*. Springer, Berlin, Heidelberg, 2001.
- [32] M. Girolami and B. Calderhead. Riemann manifold Langevin and Hamiltonian Monte Carlo methods. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 73(2):123–214, 2011.
- [33] P. W. Glynn and W. Whitt. The asymptotic efficiency of simulation estimators. *Oper. Res.*, 40(3):505–520, 1992.
- [34] J. Goodman and J. Weare. Ensemble samplers with affine invariance. *Commun. Appl. Math. Comput. Sci.*, 5(1):65–80, 2010.
- [35] J. Haslinger and J. Dvořák. Optimum composite material design. *RAIRO Modél. Math. Anal. Numér.*, 29(6):657–686, 1995.
- [36] W. K. Hastings. Monte Carlo sampling methods using Markov chains and their applications. *Biometrika*, 57(1):97–109, 1970.
- [37] B. Helffer. Remarks on decay of correlations and Witten Laplacians, Brascamp-Lieb inequalities and semiclassical limit. *J. Funct. Anal.*, 155(2):571–586, 1998.
- [38] A. Henrot. *Extremum Problems for Eigenvalues of Elliptic Operators*. Birkhäuser Verlag, Basel - Boston - Berlin, 2006.
- [39] C.-R. Hwang, S.-Y. Hwang-Ma, and S. J. Sheu. Accelerating Gaussian diffusions. *Ann. Appl. Probab.*, 3:897–913, 1993.
- [40] C.-R. Hwang, S.-Y. Hwang-Ma, and S. J. Sheu. Accelerating diffusions. *Ann. Appl. Probab.*, 15:1433–1444, 2005.
- [41] J. Hénin, T. Lelièvre, M. R. Shirts, O. Valsson, and L. Delemotte. Enhanced sampling methods for molecular dynamics simulations. *Living Journal of Computational Molecular Science*, 4(1):1583, 2022.
- [42] T. Kato. *Perturbation Theory for Linear Operators*, volume 132 of *Grundlehren der mathematischen Wissenschaften*. Springer, 1976.
- [43] C. Le Bris, F. Legoll, and F. Thomines. Rate of convergence of a two-scale expansion for some “weakly” stochastic homogenization problems. *Asymptot. Anal.*, 80(3-4):237–267, 2012.
- [44] B. Leimkuhler and C. Matthews. *Molecular Dynamics: With Deterministic and Stochastic Numerical Methods*. Springer, 2015.
- [45] B. Leimkuhler, C. Matthews, and J. Weare. Ensemble preconditioning for Markov chain Monte Carlo simulation. *Stat. Comput.*, 28:277–290, 2018.
- [46] T. Lelièvre, F. Nier, and G. A. Pavliotis. Optimal non-reversible linear drift for the convergence to equilibrium of a diffusion. *Journal of Statistical Physics*, 152(2):237–274, 2013.

- [47] T. Lelièvre, M. Rousset, and G. Stoltz. *Free-energy Computations: A Mathematical Perspective*. Imperial College Press, 2010.
- [48] T. Lelièvre, R. Santet, and G. Stoltz. Unbiasing Hamiltonian Monte Carlo algorithms for a general Hamiltonian function. *arXiv preprint*, 2303.15918, 2023.
- [49] T. Lelièvre and G. Stoltz. Partial differential equations and stochastic methods in molecular dynamics. *Acta Numerica*, 25:681–880, 2016.
- [50] S. Livingstone. Geometric ergodicity of the Random Walk Metropolis with position-dependent proposal covariance. *Mathematics*, 9(4), 2021.
- [51] J. Lu and K. Spiliopoulos. Analysis of multiscale integrators for multiple attractors and irreversible Langevin samplers. *Multiscale Model. Simul.*, 16(4):1859–1883, 2018.
- [52] A. Lê. Eigenvalue problems for the p-Laplacian. *Nonlinear Analysis: Theory, Methods & Applications*, 64(5):1057–1099, 2006.
- [53] N. Metropolis, A. W. Rosenbluth, M. N. Rosenbluth, A. H. Teller, and E. Teller. Equation of state calculations by fast computing machines. *The Journal of Chemical Physics*, 21(6):1087–1092, 1953.
- [54] G. A. Pavliotis. *Stochastic processes and applications*, volume 60 of *Texts in Applied Mathematics*. Springer, New York, 2014. Diffusion processes, the Fokker-Planck and Langevin equations.
- [55] D. Phillips, C. Matthews, and B. Leimkuhler. Numerical methods with coordinate transforms for efficient Brownian dynamics simulations. *arXiv preprint*, 2307.02913, 2023.
- [56] M. Reed and B. Simon. *Methods of Modern Mathematical Physics. I. Functional Analysis*. Academic Press, New York-London, 1972.
- [57] M. Reed and B. Simon. *Methods of Modern Mathematical Physics. IV. Analysis of Operators*. Academic Press, New York-London, 1978.
- [58] L. Rey-Bellet and K. Spiliopoulos. Irreversible Langevin samplers and variance reduction: A large deviations approach. *Nonlinearity*, 28:2081–2103, 2015.
- [59] L. Rey-Bellet and K. Spiliopoulos. Improving the convergence of reversible samplers. *J. Stat. Phys.*, 164(3):472–494, 2016.
- [60] C. P. Robert. *The Bayesian Choice*. Springer Texts in Statistics. Springer, New York, second edition, 2007.
- [61] G. O. Roberts and O. Stramer. Langevin diffusions and Metropolis-Hastings algorithms. *Methodology and Computing in Applied Probability*, 4:337–357, 2002.
- [62] G. O. Roberts and R. L. Tweedie. Exponential convergence of Langevin distributions and their discrete approximations. *Bernoulli*, 2(4):341–363, 1996.
- [63] P. J. Rossky, J. D. Doll, and H. L. Friedman. Brownian dynamics as smart Monte Carlo simulation. *J. Chem. Phys.*, 69(10):4628–4633, 1978.
- [64] A. Saumard and J. A. Wellner. Log-concavity and strong log-concavity: a review. *Stat. Surv.*, 8:45–114, 2014.
- [65] O. Sigmund. Materials with prescribed constitutive parameters: An inverse homogenization problem. *Internat. J. Solids Structures*, 31(17):2313–2329, 1994.
- [66] B. Simon. *Trace ideals and their applications*, volume 120 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, second edition, 2005.

- [67] D. W. Stroock and S. R. S. Varadhan. *Multidimensional Diffusion Processes*. Classics in Mathematics. Springer-Verlag, Berlin, 2006.
- [68] M. Tuckerman. *Statistical Mechanics: Theory and Molecular Simulation*. Oxford University Press, 2010.
- [69] P. Virtanen, R. Gommers, T. E. Oliphant, M. Haberland, T. Reddy, D. Cournapeau, E. Burovski, P. Peterson, W. Weckesser, J. Bright, S. J. van der Walt, M. Brett, J. Wilson, K. J. Millman, N. Mayorov, A. R. J. Nelson, E. Jones, R. Kern, E. Larson, C. J. Carey, Í. Polat, Y. Feng, E. W. Moore, J. VanderPlas, D. Laxalde, J. Perktold, R. Cimrman, I. Henriksen, E. A. Quintero, C. R. Harris, A. M. Archibald, A. H. Ribeiro, F. Pedregosa, P. van Mulbregt, and SciPy 1.0 Contributors. SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. *Nature Methods*, 17:261–272, 2020.
- [70] A. Wächter and L. T. Biegler. On the implementation of an interior-point filter line-search algorithm for large-scale nonlinear programming. *Mathematical Programming*, 106(1):25–57, Mar 2006.
- [71] B. J. Zhang, Y. M. Marzouk, and K. Spiliopoulos. Geometry-informed irreversible perturbations for accelerated convergence of Langevin dynamics. *Stat. Comput.*, 32(5):Paper No. 78, 22, 2022.
- [72] V. V. Zhikov. Weighted Sobolev spaces. *Sbornik: Mathematics*, 189(8):1139–1170, 1998.