

On the rectilinear crossing number of complete balanced multipartite graphs and layered graphs *

Ruy Fabila-Monroy^{†‡} Rosna Paul^{§¶} Jenifer Viafara-Chanchi^{†**}
 Alexandra Weinberger^{§¶†‡}

April 23, 2024

Abstract

A rectilinear drawing of a graph is a drawing of the graph in the plane in which the edges are drawn as straight-line segments. The rectilinear crossing number of a graph is the minimum number of pairs of edges that cross over all rectilinear drawings of the graph. Let $n \geq r$ be positive integers. The graph K_n^r , is the complete r -partite graph on n vertices, in which every set of the partition has at least $\lfloor n/r \rfloor$ vertices. The layered graph, L_n^r , is an r -partite graph on n vertices, in which for every $1 \leq i \leq r-1$, all the vertices in the i -th partition are adjacent to all the vertices in the $(i+1)$ -th partition. In this paper, we give upper bounds on the rectilinear crossing numbers of K_n^r and L_n^r .

1 Introduction

Let G be a graph on n vertices and let D be a drawing of G . The crossing number of D is the number, $\text{cr}(D)$, of pairs of edges that cross in D . The *crossing number* of G is the minimum crossing number, $\text{cr}(G)$, over all drawings of G in the plane. A *rectilinear drawing* of G is a drawing of G in the plane in which its vertices are points in general position, and its edges are drawn as straight-line segments joining these points. The *rectilinear crossing number* of G , is the minimum crossing number, $\overline{\text{cr}}(G)$, over all rectilinear drawings of G in the plane. Computing crossing and rectilinear crossing numbers of graphs are important problems in Graph Theory and Combinatorial Geometry. For a comprehensive review of the literature on crossing numbers, we refer the reader to Schaefer's book [2].

Most of the research on crossing numbers have been focused around the complete graph, K_n , and the complete bipartite graph $K_{n,m}$. For the complete graph, Hill [3] gave the following drawing of K_n ; see Figure 1 for an example. Place half of the vertices equidistantly on the top circle of a cylinder, and the other half equidistantly on the bottom circle. Join the vertices with geodesics on the cylinder. Hill showed that the

*A preliminary version of this work has been presented at EGC'23[1]

[†]Departamento de Matemáticas, CINVESTAV

[‡]ruyfabila@math.cinvestav.edu.mx

[§]Institute for Software Technology, Graz University of Technology, Graz, Austria

[¶]Supported by the Austrian Science Fund (FWF) grant W1230.

[‡]ropaul@ist.tugraz.at

^{**}viafara@math.cinvestav.mx

^{††}Department of Mathematics and Computer Science, FernUniversität in Hagen, Hagen, Germany

^{‡‡}alexandra.weinberger@fernuni-hagen.de

This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 734922.






Figure 1: An example of Hill's drawings of K_8 , where here for convenience only the edges of one vertex are drawn. Left: The drawing on a cylinder. Right: An equivalent representation of Hill's drawings via concentric cycles.

following number, $H(n)$, is the crossing number of this drawing, and it is now conjectured to be optimal. Let

$$H(n) := \frac{1}{4} \left\lfloor \frac{n}{2} \right\rfloor \left\lfloor \frac{n-1}{2} \right\rfloor \left\lfloor \frac{n-2}{2} \right\rfloor \left\lfloor \frac{n-3}{2} \right\rfloor.$$

Conjecture 1 (Harary-Hill [4])

$$\text{cr}(K_n) = H(n).$$

For the complete bipartite graph, Zarankiewicz gave a rectilinear drawing with the following number, $Z(n, m)$, as crossing number of this drawing, and it is now conjectured to be optimal. Let

$$Z(n, m) := \left\lfloor \frac{n}{2} \right\rfloor \left\lfloor \frac{n-1}{2} \right\rfloor \left\lfloor \frac{m}{2} \right\rfloor \left\lfloor \frac{m-1}{2} \right\rfloor$$

and

$$Z(n) := Z(n, n).$$

Conjecture 2 (Zarankiewicz [5])

$$\text{cr}(K_{n,m}) = Z(n, m).$$

The number $Z(n, m)$ is also conjectured to be the general optimal crossing number, directly implying the following conjecture.

Conjecture 3

$$\overline{\text{cr}}(K_{n,m}) = \text{cr}(K_{n,m}).$$

Much less is known for the rectilinear crossing number of the complete graph. For $n \geq 10$, it is known that

$$\text{cr}(K_n) < \overline{\text{cr}}(K_n).$$

In contrast to the case of the complete bipartite graph, there is no conjectured value for $\overline{\text{cr}}(K_n)$, nor drawings conjectured to be optimal. The best bounds to date are

$$0.379972 \binom{n}{4} < \overline{\text{cr}}(K_n) < 0.380445 \binom{n}{4} + O(n^3).$$

The lower bound is due to Ábrego, Fernández-Merchant, Leaños, and Salazar [6], and the upper bound to Aichholzer, Duque, Fabila-Monroy, García-Quintero, and Hidalgo-Toscano [7]. It is known that

$$\lim_{n \rightarrow \infty} \frac{\overline{\text{cr}}(K_n)}{\binom{n}{4}} = \bar{q},$$

for some positive constant \bar{q} ; this constant is known as the *rectilinear crossing constant*.

Let K_{n_1, n_2, \dots, n_r} be the complete r -partite graph with n_i vertices in the i -th set of the partition; and let K_n^r be the complete balanced r -partite graph in which there are at least $\lfloor n/r \rfloor$ vertices in every partition set. Harborth [8] gave a drawing that provides an upper bound for $\text{cr}(K_{n_1, n_2, \dots, n_r})$; and gave an explicit formula for this number. He claims that for the case of $r = 3$, his drawing can be made rectilinear. More recently, Gethner, Hogben, Lidický, Pfender, Ruiz and Young [9] independently studied the problem of the crossing number and rectilinear crossing number of complete balanced r -partite graphs. For $r = 3$, they obtain the same bound as Harborth; and their drawing is rectilinear.

Let r be a positive integer and let n be a multiple of r . The *layered graph*, L_n^r , is the graph defined as follows. Its vertex set is partitioned into sets V_1, \dots, V_r , each consisting of n/r vertices. We call the set V_i , the i -th layer of L_n^r . The edge set of L_n^r is given by

$$\{uv : u \in L_i \text{ and } v \in L_{i+1}, \text{ for } i = 1, \dots, r-1\};$$

that is, the edges are exactly all possible edges between vertices on consecutive layers.

In this paper, we mainly focus on the rectilinear crossing numbers of K_n^r and L_n^r . If n is fixed and r tends to n , then K_n^r tends to K_n . We believe that studying the rectilinear crossing number of K_n^r might shed some light on how optimal rectilinear drawings of K_n look like.

This paper is organized as follows. In Section 2, we give a general technique to obtain non-rectilinear and rectilinear drawings of a given graph G on n vertices. It simply consists of mapping randomly the vertices of G to optimal drawings of K_n . We show how this technique upper bounds $\text{cr}(K_n^r)$ and $\overline{\text{cr}}(K_n^r)$. The bounds obtained in this way are very close to being optimal. However, for the layered graphs this technique gives rather poor upper bounds. In Section 3, we give a technique where we given a specific drawing of a graph, we use this drawing as a “seed” to produce larger drawings by replacing each vertex u with a cluster of collinear vertices S_u arbitrarily close to u . In the new drawing two vertices in different clusters S_u and S_v are adjacent whenever u and v are adjacent in the original drawing. We call the new larger drawing a “planted drawing”. The conjectured crossing optimal drawings of $K_{n,n}$ and $K_{n,n,n}$ mentioned above are actually planted drawings with drawings of $K_{2,2}$ and $K_{2,2,2}$ as seeds, respectively. However, we show that there is no rectilinear drawing of K_4 or K_8^4 that can be the seed of a crossing optimal planted drawing of K_n^4 . For the layered graph, we give a rectilinear planar drawing of L_{2r}^r . When used as a seed this drawing produces a planted drawing of L_n^r , with significantly smaller crossing number, than those produced by the random embedding technique. The proofs of many of our results are long and technical; for the sake of clarity, we have relocated most of the proofs and constructions to an appendix.

2 Random Embeddings into Drawings of K_n with Small Crossing Number

Suppose that we have a drawing (that can be rectilinear but doesn't have to be) D' of K_n . If $\text{cr}(D')$ is small, it might be a good idea to use this drawing to produce a drawing of a graph G on n vertices. Let D be the drawing of G that is produced by mapping the vertices of G randomly to the vertices of D' , and where the edges are drawn as their corresponding edges of D' . We call D a *random embedding* of G into D' .

In every 4-tuple of vertices of D' , there are three pairs of independent edges, which could cross. Of these three pairs at most one pair is crossing. For every pair of independent edges of G , we have a possible crossing in D ; thus, the probability that this pair of edges is mapped to a pair of crossing edges is equal to

$$\frac{1}{3} \cdot \frac{\text{cr}(D')}{\binom{n}{4}}.$$

By defining, for every pair of independent edges of G , an indicator random variable with value equal to one if the edges cross and zero otherwise, we obtain the following result where $||G||$ is the number of edges in G and $d(v)$ is the degree of a vertex v of G .

Lemma 4

$$E(\text{cr}(D)) = \frac{cr(D')}{3\binom{n}{4}} \left(\binom{\|G\|}{2} - \sum_{v \in V(G)} \binom{d(v)}{2} \right).$$

Complete r -partite Graphs

For an upper bound on the crossing number of K_n^r , we use Lemma 4 and Hill's drawing of K_n .

Theorem 5 *Suppose that n is a multiple of r . Let D be a random embedding of K_n^r into Hill's drawing of K_n . Then,*

$$\text{cr}(K_n^r) \leq E(\text{cr}(D)) \leq \frac{1}{16} \left(\frac{r-1}{r} \right)^2 \left(\frac{n^4}{4} - \frac{3n^3}{2} \right) + O(n^2).$$

In [9], the authors obtain the same bound on $\text{cr}(K_n^r)$ by considering a random mapping of the vertices of K_n^r into a sphere, and then joining the corresponding vertices with geodesics. This type of drawing is called a *random geodesic spherical drawing*. In 1965, Moon [10], showed that the expected number of crossings of a random geodesic spherical drawing of K_n is equal to

$$\frac{1}{16} \binom{n}{2} \binom{n-2}{2} = H(n) - O(n^3);$$

which explains why the bound of Theorem 5 matches the bound of [9].

Let $H(n, r)$ be the number of crossings in Harborth's drawing for $\text{cr}(K_n^r)$. Due to the complexity of the formula, we use the following approximation to $H(n, r)$ instead.

Lemma 6 *If n is a multiple of r , then*

$$H(n, r) \leq \frac{1}{16} \left(\frac{r-1}{r} \right)^2 \left(\frac{n^4}{4} - 2n^3 \right) + O(n^2).$$

Let D be as in Theorem 5; note that by Lemma 6, it holds that

$$E(\text{cr}(D)) - H(n, r) \leq \frac{1}{32} \left(\frac{r-1}{r} \right)^2 n^3 + O(n^2) = O(n^3).$$

Thus, the random embedding gives an upper bound on $\text{cr}(D)$ that matches the conjectured value up to the leading term, but it is a little worse in the lower terms.

We now upper bound $\overline{\text{cr}}(K_n^r)$, with this technique.

Theorem 7 *Let r be a positive integer and n a multiple of r . Let \overline{D} be a random embedding of K_n^r into an optimal rectilinear drawing of K_n . Then*

$$\begin{aligned} \overline{\text{cr}}(K_n^r) &\leq E(\text{cr}(\overline{D})) \\ &\leq \frac{\overline{q}}{4!} \left(\frac{r-1}{r} \right)^2 n^4 + o(n^4) \\ &< 0.015852 \left(\frac{r-1}{r} \right)^2 n^4 + o(n^4). \end{aligned}$$

For a lower bound we have the following.

Theorem 8 *Let r be a positive integer and n a multiple of r . Then*

$$\overline{\text{cr}}(K_n^r) \geq \overline{\text{cr}}(K_r) \left(\frac{n}{r} \right)^4.$$




Figure 2: A drawing of K_8^4 with 6 crossings (left) and K_9^4 with 15 crossings (right).

Theorems 7 and 8 imply the following.

Corollary 9 *Let $r = r(n)$ be a monotone function of n such that $r \rightarrow \infty$ as $n \rightarrow \infty$. Then*

$$\lim_{n \rightarrow \infty} \frac{\overline{\text{cr}}(K_n^r)}{\binom{n}{4}} = \bar{q}.$$

In both [8] and [9], it is conjectured that

$$\text{cr}(K_n^3) = \overline{\text{cr}}(K_n^3).$$

Using the order type database [11], we have verified that

$$\overline{\text{cr}}(K_8^4) = 8 \text{ and } \overline{\text{cr}}(K_9^4) = 15.$$

On the other hand

$$\text{cr}(K_8^4) \leq H(8, 4) = 6 \text{ and } \text{cr}(K_9^4) \leq H(9, 4) = 15.$$

See Figure 2 for an example. From the above results we conjecture the following.

Conjecture 10 *There exists a natural number $n_0 > 9$ such that for all $n \geq n_0$,*

$$\text{cr}(K_n^4) < \overline{\text{cr}}(K_n^4).$$

Layered Graphs

Using the random embedding technique into Hill's drawing of K_n , we obtain the following upper bound for $\text{cr}(L_n^r)$.

Theorem 11

$$\text{cr}(L_n^r) \leq \frac{(r-1)^2}{16r^4} n^4 + O(n^3).$$

We improve this upper bound in Section 3.

3 Planted Rectilinear Drawings

Let D be a rectilinear drawing of a graph G . For every vertex v of G , let ℓ_v be a directed straight line passing through v and no other vertex of D , such that to left of ℓ_v there are $\lfloor d(v)/2 \rfloor$ neighbors of v and to the right of ℓ_v there are the remaining $\lceil d(v)/2 \rceil$ neighbors of v . Let G^s be the graph whose vertex set is equal to

$$\{(v, i) : i = 1, \dots, s \text{ and } v \in V(G)\},$$

and in which (v, i) is adjacent to (w, j) whenever vw is an edge of G . We say that the set $\{(v, 1), \dots, (v, s)\}$ is the *cluster* of v . Let D^s be the rectilinear drawing of G^s in which for every vertex v of G , the vertices its cluster are placed arbitrarily close to ℓ_v and arbitrarily close to v (in D). We say that D^s is a *planted drawing* of G^s with *seed* D .

Lemma 12

$$\text{cr}(D^s) = \text{cr}(D)s^4 + \sum_{v \in V(G)} \left(\binom{\lfloor d(v)/2 \rfloor}{2} + \binom{\lceil d(v)/2 \rceil}{2} \right) \frac{s^3(s-1)}{2} + \|G\| \frac{s^2(s-1)^2}{4}.$$

Seeds and planted drawings were first used by Ábrego and Fernández-Merchant [12]¹ to upper bound the rectilinear crossing number of K_n . The current best upper bound on $\overline{\text{cr}}(K_n)$ is obtained via a seed of 2643 vertices and 771218714414 crossings.

Complete r -partite Graphs

Note that if we use K_{tr}^r as a seed for a planted drawing of K_n^r , we have that $s = \frac{n}{tr}$. Thus, from Lemma 12 we obtain the following.

Corollary 13 *Let D be a rectilinear drawing of K_{tr}^r . Then using D as a seed we obtain a planted drawing of K_n^r with*

$$\left(\frac{\text{cr}(D) + \frac{rt}{2} \left(\binom{\lfloor (r-1)t/2 \rfloor}{2} + \binom{\lceil (r-1)t/2 \rceil}{2} \right) + \frac{r(r-1)t^2}{8}}{(rt)^4} \right) n^4 - O(n^3)$$

crossings.

Using the seeds in Figure 3, we obtain planted rectilinear drawings of K_n^2 and K_n^3 , with the conjectured minimum number of crossings.

Using the random embedding technique and Theorem 5 we obtain a rectilinear drawing of K_n^4 with at most

$$0.0089676n^4 + o(n^4) \tag{1}$$

crossings; and since $\bar{q} > 0.379972$, the best we can hope to achieve with the random embedding technique is a rectilinear drawing of K_n^4 with

$$0.0089055n^4 + o(n^4) \tag{2}$$

crossings.

Using a planar drawing of K_4 as a seed, we obtain a rectilinear planted drawing of K_n^r with

$$\left(\frac{2 \left(\binom{1}{2} + \binom{2}{2} \right) + \frac{3}{2}}{4^4} \right) n^4 - O(n^3) = \frac{7}{29} n^4 - O(n^3) = 0.013671875n^4 - O(n^3)$$

¹They do it in a different way as presented here; first they duplicate each vertex along halving lines; then they choose halving lines for the original and new vertices and duplicate a new. They iterate this process.




Figure 3: The seeds for the planted drawings of K_n^2 and K_n^3

crossings. Using a rectilinear drawing of K_8^4 with 8 crossings, we obtain a planted rectilinear drawing with

$$\left(\frac{8 + 4 \left(\binom{3}{2} + \binom{3}{2} \right) + 6}{8^4} \right) n^4 - O(n^3) = \frac{38}{8^4} n^4 - O(n^3) = 0.009277344n^4 - O(n^3)$$

crossings.

Fabila-Monroy and López [13] used an heuristic of randomly moving vertices to obtain a rectilinear drawing of K_{75} with 45049 crossings. This was used as a seed for a previous best upper bound on \bar{q} . In [14] Duque, Fabila-Monroy, Hernández-Vélez and Hidalgo-Toscano gave an $O(n^2 \log n)$ time algorithm to compute the crossing number of a rectilinear drawing of a graph on n vertices. Using a similar heuristic as in [13] and the algorithm of [14], we obtained a rectilinear drawing of K_{24}^4 with 2033 crossings. Using this as a seed we obtain a planted rectilinear drawing of K_n^4 with

$$\left(\frac{2033 + 12 \left(\binom{9}{2} + \binom{9}{2} \right) + 54}{24^4} \right) n^4 - O(n^3) = \frac{2951}{24^4} n^4 - O(n^3) = 0.0088946n^4 - O(n^3)$$

crossings. This is better than the best possible upper bound obtainable with the random embedding technique. However, for $r \geq 5$, we have not found seeds that provide planted drawings with less crossings than the drawings obtained from the random embedding technique.

Layered Graphs

We now show a rectilinear planar drawing D_r of L_{2r}^r . For $i = 1, \dots, r$, let $\{u_i, v_i\}$ be the two vertices on layer i of L_{2r}^r . Place u_i and v_i at the points p_i and q_i , respectively; where

$$p_i := \begin{cases} (i, 0) & \text{if } i \text{ is odd,} \\ (0, i) & \text{if } i \text{ is even,} \end{cases} \quad \text{and} \quad q_i := \begin{cases} (-i, 0) & \text{if } i \text{ is odd,} \\ (0, -i) & \text{if } i \text{ is even.} \end{cases}$$

See Figure 4 for the drawing of L_{12}^6 .

Using this drawing as a seed for a planted drawing of L_n^r , we obtain a rectilinear drawing with

$$\begin{aligned} & \sum_{v \in V(D_r)} \left(\binom{\lfloor d(v)/2 \rfloor}{2} + \binom{\lceil d(v)/2 \rceil}{2} \right) \frac{s^4}{2} + \|D_r\| \frac{s^4}{4} - O(s^3) \\ &= (2(r-2) \cdot 2) \frac{n^4}{2 \cdot (2r)^4} + 4 \cdot (r-1) \frac{n^4}{4 \cdot (2r)^4} - O(n^3) \\ &= \frac{3r-5}{16r^4} n^4 - O(n^3) \end{aligned}$$




Figure 4: The rectilinear D_6 drawing of L_{12}^6

crossings. For $r \geq 4$, this is better than the upper bound obtained with the random embedding technique.

For $i = 2, \dots, r-1$, let H_i be the subgraph of L_n^r induced by the vertices in layers $i-1, i$ and $i+1$. Note that this graph is isomorphic to $K_{n/r, 2n/r}$. Thus, assuming that Zarankiewicz's conjecture holds, in every drawing of L_n^r , H_i produces at least $Z(n/r, 2n/r)$ crossings. Each of these crossings is produced by at most two such H_i 's. Therefore, assuming that Zarankiewicz's conjecture is true, we have that

$$\text{cr}(L_n^r) \geq \frac{(r-2)}{2} Z(n/r, 2n/r) = \frac{2r-4}{16r^4} n^4 - O(n^3).$$

References

- [1] R. Fabila-Monroy, R. Paul, J. Viafara-Chanchi, and A. Weinberger, "On the rectilinear crossing number of complete balanced multipartite graphs and layered graphs," in *XX Encuentros de Geometria Computacional (EGC'23)*, Santiago de Compostela, Spain, pp. 33–36, 2023.
- [2] M. Schaefer, *Crossing numbers of graphs*. CRC Press, 2018.
- [3] F. Harary and A. Hill, "On the number of crossings in a complete graph," *Proceedings of the Edinburgh Mathematical Society*, vol. 13, no. 4, pp. 333–338, 1963.
- [4] R. K. Guy, "A combinatorial problem," *Nabla (Bulletin of the Malayan Mathematical Society)*, vol. 7, pp. 68–72, 1960.
- [5] K. Zarankiewicz, "On a problem of P. Turan concerning graphs," *Fund. Math.*, vol. 41, pp. 137–145, 1954.
- [6] B. M. Ábrego, S. Fernández-Merchant, J. Leaños, and G. Salazar, "A central approach to bound the number of crossings in a generalized configuration," in *The IV Latin-American Algorithms, Graphs, and Optimization Symposium*, vol. 30 of *Electron. Notes Discrete Math.*, pp. 273–278, Elsevier Sci. B. V., Amsterdam, 2008.
- [7] O. Aichholzer, F. Duque, R. Fabila Monroy, O. E. García-Quintero, and C. Hidalgo-Toscano, "An ongoing project to improve the rectilinear and the pseudolinear crossing constants." Preprint.
- [8] H. Harborth, "Über die Kreuzungszahl vollständiger, n-geteilter Graphen," *Mathematische Nachrichten*, vol. 48, no. 1-6, pp. 179–188, 1971.
- [9] E. Gethner, L. Hogben, B. Lidický, F. Pfender, A. Ruiz, and M. Young, "On crossing numbers of complete tripartite and balanced complete multipartite graphs," *Journal of Graph Theory*, vol. 4, no. 84, pp. 552–565, 2017.

- [10] J. W. Moon, “On the distribution of crossings in random complete graphs,” *J. Soc. Indust. Appl. Math.*, vol. 13, pp. 506–510, 1965.
- [11] O. Aichholzer, F. Aurenhammer, and H. Krasser, “Enumerating order types for small point sets with applications,” *Order*, vol. 19, no. 3, pp. 265–281, 2002.
- [12] B. M. Ábrego and S. Fernández-Merchant, “Geometric drawings of K_n with few crossings,” *J. Combin. Theory Ser. A*, vol. 114, no. 2, pp. 373–379, 2007.
- [13] R. Fabila-Monroy and J. López, “Computational search of small point sets with small rectilinear crossing number,” *Journal of Graph Algorithms and Applications*, vol. 18, no. 3, pp. 393–399, 2014.
- [14] F. Duque, R. Fabila-Monroy, C. Hernández-Vélez, and C. Hidalgo-Toscano, “Counting the number of crossings in geometric graphs,” *Inform. Process. Lett.*, vol. 165, pp. Paper No. 106028, 5, 2021.

4 Appendix

We continuously use that:

- if x is an even integer, then $\lfloor (x-1)/2 \rfloor = (x-2)/2 = x/2 - 1$;
- and if x is an odd integer, then $\lfloor x/2 \rfloor = (x-1)/2$.

Lemma 14

$$\frac{H(n)}{3\binom{n}{4}} \leq \frac{1}{8} \left(1 - \frac{2}{n}\right).$$

Proof. If n is even, then

$$\begin{aligned} \frac{H(n)}{3\binom{n}{4}} &= \frac{2}{n(n-1)(n-2)(n-3)} \left\lfloor \frac{n}{2} \right\rfloor \left\lfloor \frac{n-1}{2} \right\rfloor \left\lfloor \frac{n-2}{2} \right\rfloor \left\lfloor \frac{n-3}{2} \right\rfloor \\ &= \frac{2}{n(n-1)(n-2)(n-3)} \binom{n}{2} \binom{n-2}{2} \binom{n-2}{2} \binom{n-4}{2} \\ &= \frac{1}{(n-1)(n-3)} \left(\frac{1}{2}\right) \binom{n-2}{2} \binom{n-4}{2} \\ &= \frac{1}{8(n-1)(n-3)} ((n-2)(n-4)) \\ &= \frac{n-2}{8} \left(\frac{(n-4)}{(n-1)(n-3)} \right) \\ &= \frac{n-2}{8} \left(\frac{(n-4)}{n^2-4n+3} \right) \\ &= \frac{n-2}{8} \left(\frac{(n-4)n}{(n^2-4n+3)n} \right) \\ &= \frac{n-2}{8} \left(\frac{n^2-4n}{(n^2-4n+3)n} \right) \\ &< \frac{n-2}{8} \left(\frac{1}{n} \right) \\ &= \frac{1}{8} \left(1 - \frac{2}{n}\right). \end{aligned}$$

If n is odd, then

$$\begin{aligned}
\frac{H(n)}{3\binom{n}{4}} &= \frac{2}{n(n-1)(n-2)(n-3)} \left\lfloor \frac{n}{2} \right\rfloor \left\lfloor \frac{n-1}{2} \right\rfloor \left\lfloor \frac{n-2}{2} \right\rfloor \left\lfloor \frac{n-3}{2} \right\rfloor \\
&= \frac{2}{n(n-1)(n-2)(n-3)} \left(\frac{n-1}{2} \right) \left(\frac{n-1}{2} \right) \left(\frac{n-3}{2} \right) \left(\frac{n-3}{2} \right) \\
&= \frac{1}{8n} \left(\frac{(n-1)(n-3)}{n-2} \right) \\
&= \frac{n-2}{8n} \left(\frac{(n-1)(n-3)}{(n-2)(n-2)} \right) \\
&= \frac{n-2}{8n} \left(\frac{n^2-4n+3}{n^2-4n+4} \right) \\
&< \frac{1}{8} \left(\frac{n-2}{n} \right) \\
&= \frac{1}{8} \left(1 - \frac{2}{n} \right).
\end{aligned}$$

□

According to Harborth [8], if n is a multiple of r , then

$$H(n, r) \leq \frac{3}{8} \binom{r}{4} \frac{n^4}{r^4} + r \left\lfloor \frac{n/r}{2} \right\rfloor \left\lfloor \frac{n/r-1}{2} \right\rfloor \left\lfloor \frac{n-n/r}{2} \right\rfloor \left\lfloor \frac{n-n/r-1}{2} \right\rfloor - \binom{r}{2} \left(\left\lfloor \frac{n/r}{2} \right\rfloor^2 \right) \left(\left\lfloor \frac{n/r-1}{2} \right\rfloor^2 \right) + O(n^2).$$

We bound this formula to obtain Lemma 6.

Proof. [Lemma 6] Let

$$\begin{aligned}
A &:= \frac{3}{8} \binom{r}{4} \frac{n^4}{r^4} = \frac{1}{64} \cdot \frac{(r-1)(r-2)(r-3)}{r^3} n^4, \\
B &:= r \left\lfloor \frac{n/r}{2} \right\rfloor \left\lfloor \frac{n/r-1}{2} \right\rfloor \left\lfloor \frac{n-n/r}{2} \right\rfloor \left\lfloor \frac{n-n/r-1}{2} \right\rfloor,
\end{aligned}$$

and

$$C := \binom{r}{2} \left\lfloor \frac{n/r}{2} \right\rfloor^2 \left\lfloor \frac{n/r-1}{2} \right\rfloor^2.$$

If n/r is even, then

$$\begin{aligned}
B &= r \left(\frac{n}{2r} \right) \left(\frac{n}{2r} - 1 \right) \left(\frac{n}{2} - \frac{n}{2r} \right) \left(\frac{n}{2} - \frac{n}{2r} - 1 \right) \\
&= \frac{n}{2} \left(\frac{n}{2r} - 1 \right) \left(\frac{n}{2} \left(\frac{r-1}{r} \right) \right) \left(\frac{n}{2} \left(\frac{r-1}{r} \right) - 1 \right) \\
&= \left(\frac{n^2}{4r} - \frac{n}{2} \right) \left(\frac{n^2}{4} \left(\frac{r-1}{r} \right)^2 - \frac{n}{2} \left(\frac{r-1}{r} \right) \right) \\
&= \frac{n^4}{16r} \left(\frac{r-1}{r} \right)^2 - \frac{n^3}{8r} \left(\frac{r-1}{r} \right) - \frac{n^3}{8} \left(\frac{r-1}{r} \right)^2 + O(n^2);
\end{aligned}$$

If n/r is odd, and n is even, then

$$\begin{aligned}
B &= r \left\lfloor \frac{n/r}{2} \right\rfloor \left\lfloor \frac{n/r-1}{2} \right\rfloor \left\lfloor \frac{n-n/r}{2} \right\rfloor \left\lfloor \frac{n-n/r-1}{2} \right\rfloor \\
&= r \left(\frac{n/r-1}{2} \right) \left(\frac{n/r-1}{2} \right) \left(\frac{n-n/r-1}{2} \right) \left(\frac{n-n/r-1}{2} \right) \\
&= \frac{r}{16} \left(\frac{n^2}{r^2} - \frac{2n}{r} + 1 \right) \left(\left(\frac{r-1}{r} \right)^2 n^2 - 2 \left(\frac{r-1}{r} \right) n + 1 \right) \\
&= \frac{n^4}{16r} \left(\frac{r-1}{r} \right)^2 - \frac{n^3}{8r} \left(\frac{r-1}{r} \right) - \frac{n^3}{8} \left(\frac{r-1}{r} \right)^2 + O(n^2)
\end{aligned}$$

If n/r is odd and n is odd, then

$$\begin{aligned}
B &= r \left\lfloor \frac{n/r}{2} \right\rfloor \left\lfloor \frac{n/r-1}{2} \right\rfloor \left\lfloor \frac{n-n/r}{2} \right\rfloor \left\lfloor \frac{n-n/r-1}{2} \right\rfloor \\
&= r \left(\frac{n/r-1}{2} \right) \left(\frac{n/r-1}{2} \right) \left(\frac{n-n/r}{2} \right) \left(\frac{n-n/r-2}{2} \right) \\
&= \frac{r}{16} \left(\frac{n^2}{r^2} - \frac{2n}{r} + 1 \right) \left(\left(\frac{r-1}{r} \right)^2 n^2 - 2 \left(\frac{r-1}{r} \right) n \right) \\
&= \frac{n^4}{16r} \left(\frac{r-1}{r} \right)^2 - \frac{n^3}{8r} \left(\frac{r-1}{r} \right) - \frac{n^3}{8} \left(\frac{r-1}{r} \right)^2 + O(n^2)
\end{aligned}$$

If n is even, then

$$\begin{aligned}
C &= \binom{r}{2} \left\lfloor \frac{n/r}{2} \right\rfloor^2 \left\lfloor \frac{n/r-1}{2} \right\rfloor^2 \\
&= \frac{r(r-1)}{2} \cdot \frac{n^2}{4r^2} \left(\frac{n}{2r} - 1 \right)^2 \\
&= \frac{r-1}{32r^3} n^4 - \frac{r-1}{8r^2} n^3 + O(n^2);
\end{aligned}$$

if n is odd, then

$$\begin{aligned}
C &= \binom{r}{2} \left\lfloor \frac{n/r}{2} \right\rfloor^2 \left\lfloor \frac{n/r-1}{2} \right\rfloor^2 \\
&= \frac{r(r-1)}{2} \left(\frac{n/r-1}{2} \right)^2 \left(\frac{n/r-1}{2} \right)^2 \\
&= \frac{r(r-1)}{32} \left(\frac{n^4}{r^4} - \frac{4n^3}{r^3} + O(n^2) \right) \\
&= \frac{r-1}{32r^3} n^4 - \frac{r-1}{8r^2} n^3 + O(n^2).
\end{aligned}$$

Therefore,

$$\begin{aligned}
H(n, r) &\leq A + B - C + O(n^2) \\
&= \frac{1}{16} \left(\frac{(r-1)(r-2)(r-3)}{4r^3} + \frac{(r-1)^2}{r^3} - \frac{r-1}{2r^3} \right) n^4 \\
&\quad + \frac{1}{8} \left(-\frac{r-1}{r^2} - \frac{(r-1)^2}{r^2} + \frac{r-1}{r^2} \right) n^3 \\
&\quad + O(n^2) \\
&= \frac{1}{16} \left(\frac{r^3 - 2r + r}{4r^3} \right) n^4 + n^3 + O(n^2) - \frac{1}{16} \left(\frac{r-1}{r} \right)^2 2n^3 + O(n^2) \\
&= \frac{1}{16} \left(\frac{r-1}{r} \right)^2 \left(\frac{n^4}{4} - 2n^3 \right) + O(n^2).
\end{aligned}$$

□

Lemma 15 *If n is a multiple of r , then*

$$\left(\binom{\|K_n^r\|}{2} - \sum_{v \in V(K_n^r)} \binom{d(v)}{2} \right) = \frac{1}{2} \left(\frac{r-1}{r} \right)^2 \left(\frac{n^4}{4} - n^3 \right) + O(n^2).$$

Proof. Every set in the partition has n/r vertices. Thus, the number of edges between two different sets is equal to n^2/r^2 . Therefore,

$$\|K_n^r\| = \frac{n^2}{r^2} \binom{r}{2} = \frac{n^2}{2} \cdot \frac{r-1}{r},$$

and

$$\binom{\|K_n^r\|}{2} = \frac{n^4}{8} \left(\frac{r-1}{r} \right)^2 - \frac{n^2}{4} \left(\frac{r-1}{r} \right).$$

For every vertex v of K_n^r , it holds that

$$d(v) = \frac{r-1}{r} n.$$

Thus,

$$\sum_{v \in V(K_n^r)} \binom{d(v)}{2} = \frac{n^3}{2} \left(\frac{r-1}{r} \right)^2 - \frac{r-1}{r} \cdot \frac{n^2}{2}.$$

It follows that

$$\begin{aligned}
\left(\binom{\|K_n^r\|}{2} - \sum_{v \in V(K_n^r)} \binom{d(v)}{2} \right) &= \frac{n^4}{8} \left(\frac{r-1}{r} \right)^2 - \frac{n^2}{4} \left(\frac{r-1}{r} \right) - \frac{n^3}{2} \left(\frac{r-1}{r} \right)^2 + \frac{n^2}{2} \left(\frac{r-1}{r} \right) \\
&= \frac{1}{2} \left(\frac{r-1}{r} \right) \left(\frac{n^4}{4} \left(\frac{r-1}{r} \right) - \frac{n^2}{2} - n^3 \left(\frac{r-1}{r} \right) + n^2 \right) \\
&= \frac{1}{2} \left(\frac{r-1}{r} \right)^2 \left(\frac{n^4}{4} - n^3 \right) + O(n^2).
\end{aligned}$$

□

Using Lemmas 4, 14 and 15, we can prove Theorem 5.

Proof. [Theorem 5] By Lemma 4, it holds that

$$E(\text{cr}(D)) = \frac{H(n)}{3\binom{n}{4}} \left(\binom{\|K_n^r\|}{2} - \sum_{v \in V(K_n^r)} \binom{d(v)}{2} \right).$$

Applying Lemmas 14 and 15 on the equality above yields

$$\begin{aligned} E(\text{cr}(D)) &\leq \frac{1}{8} \left(1 - \frac{2}{n} \right) \left(\frac{1}{2} \left(\frac{r-1}{r} \right)^2 \left(\frac{n^4}{4} - n^3 \right) + O(n^2) \right) \\ &\leq \frac{1}{16} \left(\frac{r-1}{r} \right)^2 \left(\frac{n^4}{4} - \frac{3n^3}{2} \right) + O(n^2). \end{aligned}$$

□

Proof. [Theorem 7] From Lemma 4 and the best upper bound known for $\overline{\text{cr}}(K_n)$, it follows that

$$\begin{aligned} E(\overline{\text{cr}}(\overline{D})) &= \frac{\overline{\text{cr}}(K_n)}{3\binom{n}{4}} \left(\binom{\|K_n^r\|}{2} - \sum_{v \in V(K_n^r)} \binom{d(v)}{2} \right) \\ &= \frac{\overline{q}\binom{n}{4} + o(n^4)}{3\binom{n}{4}} \left(\frac{1}{2} \left(\frac{r-1}{r} \right)^2 \left(\frac{n^4}{4} - n^3 \right) + O(n^2) \right) \\ &\leq \frac{\overline{q}}{4!} \left(\frac{r-1}{r} \right)^2 n^4 + o(n^4). \end{aligned}$$

□

Proof. [Theorem 8] Let D be a rectilinear drawing of K_n^r . Let D' be a rectilinear drawing of K_r obtained by choosing one point from each color class of D . There are $(n/r)^r$ such choices; and each choice provides at least $\overline{\text{cr}}(K_r)$ crossings. Each such crossing is counted exactly $(n/r)^{r-4}$ times. □

Proof. [Corollary 9] We have that

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\overline{\text{cr}}(K_n^r)}{\binom{n}{4}} &\geq \lim_{n \rightarrow \infty} \overline{\text{cr}}(K_r) \cdot \left(\frac{n}{r} \right)^4 \cdot \frac{4!}{n(n-1)(n-2)(n-3)} \\ &\geq \lim_{r \rightarrow \infty} \frac{\overline{\text{cr}}(K_r)}{\binom{r^4}{4!}} \\ &= \overline{q}. \end{aligned}$$

By Theorem 7, $\lim_{n \rightarrow \infty} \frac{\overline{\text{cr}}(K_n^r)}{\binom{n}{4}} \leq \overline{q} \left(\frac{r-1}{r} \right)^2$.

As $\left(\frac{r-1}{r} \right)^2 < 1$, it follows that

$$\lim_{n \rightarrow \infty} \frac{\overline{\text{cr}}(K_n^r)}{\binom{n}{4}} = \overline{q}.$$

□

To prove Theorem 11, we use the following proposition.

Proposition 16 *Let r be a positive integer and let n be a multiple of r . Then*

$$\left(\binom{\|L_n^r\|}{2} - \sum_{v \in V(L_n^r)} \binom{d(v)}{2} \right) = \frac{(r-1)^2}{2r^4} n^4 - \frac{2r-3}{r^3} n^3 + \frac{r-1}{r^2} n^2.$$

Proof. Note that

$$\|L_n^r\| = (r-1) \left(\frac{n}{r}\right)^2;$$

and

$$\binom{\|L_n^r\|}{2} = \frac{(r-1)^2}{2r^4} n^4 - \frac{r-1}{2r^2} n^2.$$

We have that

$$\begin{aligned} \sum_{v \in V(L_n^r)} \binom{d(v)}{2} &= \frac{2n}{r} \binom{n/r}{2} + \frac{(r-2)n}{r} \binom{2n/r}{2} \\ &= \frac{2n}{r} \left(\frac{n^2}{2r^2} - \frac{n}{2r} \right) + \frac{(r-2)n}{r} \left(\frac{2n^2}{r^2} - \frac{n}{r} \right) \\ &= \frac{2r-3}{r^3} n^3 - \frac{r-1}{r^2} n^2. \end{aligned}$$

Thus,

$$\left(\binom{\|L_n^r\|}{2} - \sum_{v \in V(L_n^r)} \binom{d(v)}{2} \right) = \frac{(r-1)^2}{2r^4} n^4 - \frac{2r-3}{r^3} n^3 + \frac{r-1}{r^2} n^2.$$

□

Combining Proposition 16 with Lemmas 4 and 14, we obtain Theorem 11.

Proof. [Theorem 11]

$$\begin{aligned} \text{cr}(L_n^r) \leq E(\text{cr}(D)) &= \frac{H(n)}{3 \binom{n}{4}} \left(\binom{\|L_n^r\|}{2} - \sum_{v \in V(L_n^r)} \binom{d(v)}{2} \right) \leq \\ &\frac{1}{8} \left(1 - \frac{2}{n} \right) \left(\frac{(r-1)^2}{2r^4} n^4 + O(n^3) \right) \leq \frac{(r-1)^2}{16r^4} n^4 + O(n^3). \end{aligned}$$

□

Proof. [Lemma 12] We classify the crossings of D^s depending on the number of different clusters in which the endpoints of the edges defining the crossing appear. Let e_1 and e_2 be a pair of edges of D^s that cross.

Suppose that the endpoints of e_1 and e_2 appear in four different clusters. We have that $e_1 = (u, i)(v, j)$ and $e_2 = (w, k)(x, l)$ for some four distinct vertices u, v, w, l of D and indices $1 \leq i, j, k, l \leq s$. Thus, uv, wx is a pair of crossing edges in D ; and for each pair of crossing edges in D we obtain s^4 pairs of crossing edges of D^s , such that its endpoints lie in four different clusters. Therefore, the number of crossings of D^s generated by pairs of edges whose endpoints lie in four different clusters is equal to

$$\text{cr}(D)s^4.$$

Suppose that the endpoints of e_1 and e_2 lie in three different clusters. Without loss of generality $e_1 = (u, i)(v, j)$ and $e_2 = (u, k)(w, l)$ for some three distinct vertices u, v, w of D and indices $1 \leq i, j, k, l \leq s$. Thus, v and w lie on the same side of ℓ_u ; and for every pair of vertices of D lying on the same side of ℓ_u we obtain $\binom{s}{2}s^2$ crossings in D^s generated by pairs of edges whose endpoints lie in three different clusters. Therefore, the number of crossings of D^s generated by pairs of edges whose endpoints lie in three different clusters is equal to

$$\sum_{v \in V(G)} \left(\binom{\lfloor d(v)/2 \rfloor}{2} + \binom{\lceil d(v)/2 \rceil}{2} \right) \frac{s^3(s-1)}{2}.$$

Suppose that the endpoints of e_1 and e_2 lie in two different clusters. We have that $e_1 = (u, i)(v, j)$ and $e_2 = (u, k)(v, l)$ for some edge uv of D and indices $1 \leq i, j, k, l \leq s$; and for every edge of D we obtain $\binom{s}{2} \binom{s}{2}$ crossings in D^s generated by pairs of edges whose endpoints lie in two different clusters. Therefore, the number of crossings of D^s generated by pairs of edges whose endpoints lie in two different clusters is equal to

$$||G|| \frac{s^2(s-1)^2}{4}.$$

□

We now give the coordinates of the rectilinear drawing D of K_{24}^4 with 2033 crossings. The colors are 0, 1, 2 and 3. We have appended the color of each point as a third coordinate.

$$\begin{aligned} V(D) = \{ & (-59260959, 44970123, 0), (261261347, -43693014, 0), (158829052, -28658158, 0), \\ & (-20273112, -23913465, 0), (20602644, -8343316, 0), (-8148611, -63519416, 0), \\ & (30209164, 4850528, 1), (12317574, -161508817, 1), (46649346, -344926319, 1), \\ & (-11015825, -47872739, 1), (-26347789, 22655563, 1), (-46729617, 35472331, 1), \\ & (-74136586, 66127255, 2), (-278900322, 316137789, 2), (14791528, -20163276, 2), \\ & (-140757971, 147565111, 2), (14081248, -20874215, 2), (9903931, -24183515, 2), \\ & (-38516867, 27953341, 3), (-60922797, 47350463, 3), (8267623, -135305393, 3), \\ & (-15043716, -39580158, 3), (41831995, 797354, 3), (181333931, -34086725, 3)\}. \end{aligned}$$

The vertices of this drawing can be seen in Figure 5.




Figure 5: The vertices of a rectilinear drawing of K_{24}^4