

A HILBERT-SCHMIDT INTEGRAL OPERATOR AND THE WEIL DISTRIBUTION

XIAN-JIN LI

ABSTRACT. In this paper, a positive operator is given. It is shown that the product of this positive operator and the convolution operator is a trace class Hilbert-Schmidt integral operator and has nonnegative eigenvalues. A formula is given for the trace of this product operator. It seems that this product operator is the closest trace class integral operator which has nonnegative eigenvalues and is related to the Weil distribution. A relation is given between the trace of the product operator and the Weil distribution.

1. INTRODUCTION

We first introduce some notations. Let $C_c^\infty(0, \infty)$ be the space of compactly supported smooth functions on $(0, \infty)$. Because of [1, Theorem 1, p. 191], we assume that g is a real-valued function in $C_c^\infty(0, \infty)$. Define

$$(1.1) \quad h(x) = \int_0^\infty g(xy)g(y)dy.$$

Then $h \in C_c^\infty(0, \infty)$. Since g belongs to $C_c^\infty(0, \infty)$, there exists a positive number $\mu < 1$ such that the support of g is contained in $(\sqrt{\mu}, \mu^{-1/2})$. Thus, $h(x) = 0$ for all $x \notin (\mu, \mu^{-1})$.

Let $\Lambda > 1$ be any fixed number, and let

- (1) $S = \{\infty, \text{all primes } p < \mu^{-1}\}$, $S' = S - \{\infty\}$, $Q = \text{field of rational numbers}$, $Q^* = Q - \{0\}$,
- (2) $\mathbb{A}_S = \mathbb{R} \times \prod_{p \in S'} Q_p$, $J_S = \mathbb{R} \times \prod_{p \in S'} Q_p^*$, $C_S = J_S / O_S^*$, $Q_p = p\text{-adic completion of } Q$,
- (3) $O_S^* = \{\xi \in Q^* : |\xi|_p = 1, p \notin S\}$, $I_S = \mathbb{R}_+ \times \prod_{p \in S'} O_p^*$, $O_p^* = \{x \in Q_p : |x|_p = 1\}$,
- (4) $E_S(f)(x) = \sqrt{|x|} \sum_{\xi \in O_S^*} f(\xi x)$, $e_S(f)(x) = \sum_{\xi \in O_S^*} f(\xi x)$,
- (5) S_Λ is the multiplication operator multiplied by the function which equals 0 when $|x| \leq \Lambda^{-1}$ and equals 1 when $|x| > \Lambda^{-1}$,
- (6) P_Λ is the multiplication by the function which is 0 for $|x| \geq \Lambda$ and is 1 for $|x| < \Lambda$,
- (7) \mathbb{N}_S is the set of all positive integers which are products of primes in S' ,
- (8) $h(x) := h(|x|)$ for $x \in \mathbb{A}_S$ or C_S , $|x| := |x|_S$.

1991 *Mathematics Subject Classification.* Primary 46C05, 47G10, 11M06.

Key words and phrases. Explicit formula, eigenvalues, positivity, trace of operator.

Let ψ_p be the additive character on Q_p given as in [10, Theorem 2.2.1, p. 309]. Let $\Psi_S = \prod_{p \in S} \psi_p$. For $f = \prod_{p \in S} f_p \in L^2(\mathbb{A}_S)$, we define

$$\mathfrak{F}_S f(\beta) = \int_{\mathbb{A}_S} f(\alpha) \Psi_S(-\alpha\beta) d\alpha.$$

Then $\mathfrak{F}_S f = \prod_{p \in S} \mathfrak{F}_p f_p$. As the Schwartz-Bruhat space $S(\mathbb{A}_S)$ is dense in $L^2(\mathbb{A}_S)$, the definition of \mathfrak{F}_S can be extended to all functions in $L^2(\mathbb{A}_S)$. The set of all functions f in $S(\mathbb{A}_S)$ satisfying $f(0) = 0$ and $\mathfrak{F}_S f(0) = 0$ is denoted by $S(\mathbb{A}_S)_0$.

The convolution operator $V(h)$ is defined by

$$V(h)F(x) = \int_{C_S} h(\lambda^{-1}x) \sqrt{|\lambda^{-1}x|} F(\lambda) d^\times \lambda$$

for $F \in L^2(C_S)$. The Weil distribution $\Delta(h)$ [13, p. 18] is given by

$$\Delta(h) = \widehat{h}(0) + \widehat{h}(1) - \sum_p \int'_{Q_p^*} \frac{h(|u|_p^{-1})}{|1-u|_p} d^* u,$$

where the sum on p is over all primes of Q including the infinity prime, and for $p \neq \infty$

$$\int'_{Q_p^*} \frac{h(|u|_p^{-1})}{|1-u|_p} d^* u = \sum_{m=1}^{\infty} \log p [h(p^m) + p^{-m} h(p^{-m})].$$

In particular,

$$\int'_{Q_p^*} \frac{h(|u|_p^{-1})}{|1-u|_p} d^* u = 0$$

for $p \notin S$ by our choice of S .

Similarly as function field case [6] we always assume that $\widehat{h}(1) = \widehat{h}(0) = 0$; see [5, Theorem 1.3].

The paper is organized as follows: Let

$$T = S_\Lambda (S_\Lambda - E_S \mathfrak{F}_S^t P_\Lambda \mathfrak{F}_S E_S^{-1}) S_\Lambda.$$

We show in Section 2 that T is a positive operator on $L^2(C_S)$ and that all eigenvalues of $V(h)T$ are nonnegative, namely the following two theorems.

Theorem 1.1. *Let $Z_\Lambda = \mathfrak{F}_S^t P_\Lambda \mathfrak{F}_S$ and*

$$T = S_\Lambda (S_\Lambda - E_S Z_\Lambda E_S^{-1}) S_\Lambda.$$

Then T is a positive operator on $L^2(C_S)$.

Theorem 1.2. *All eigenvalues of $V(h)T$ on $L^2(C_S)$ are nonnegative.*

In Section 3, we prove that $V(h)T$ is a trace class operator on $L^2(C_S)$.

Theorem 1.3. *$V(h)T$ is of trace class on $L^2(C_S)$.*

The trace of the operator $V(h)T$ is computed in Section 4. In particular, we obtained the following theorem.

Theorem 1.4. *We have*

$$\begin{aligned} \text{trace}_{L^2(C_S)} \{V(h)T\} &= \int_{\mathbb{A}_S, |v|>1} \Psi_S(v) \log |v| dv \int_{\mathbb{A}_S, |u|\geq 1} h(u) \Psi_S(-uv) du \\ &\quad + \int_{|v|>1} \Psi_S(v) \log |v| dv \int_{|u|>1} h(u) \Psi_S(-uv) du. \end{aligned}$$

By Theorem 1.2, all eigenvalues of $V(h)T$ on $L^2(C_S)$ are nonnegative. Theorem 1.3 says that $V(h)T$ is of trace class on $L^2(C_S)$. The Lidskii theorem (see Lemma 5.2) asserts that spectral and functional traces of a trace class operator are the same. So, the trace of $V(h)T$ is nonnegative. The following corollary then follows from Theorem 1.4; see Section 5.

Corollary 1.5. *Let h be any function given as in (1.1). Then*

$$\sum_{k_1, k_2, l_1, l_2 \in \mathbb{N}_S} \frac{\mu(k_1)\mu(k_2)}{k_1 k_2} \int_1^\infty \log v \cos \frac{2\pi l_1 v}{k_1} dv \int_1^\infty h(u) \cos \frac{2\pi l_2 uv}{k_2} du \geq 0.$$

We derive in Section 6 the following relation between the trace of the operator $V(h)T$ and the Weil distribution $\Delta(h)$.

Theorem 1.6. *We have*

$$\begin{aligned} \text{trace}_{L^2(C_S)} \{V(h)T\} &= \Delta(h) - \int_{|v|<1} \log |v| \Psi_S(-v) dv \int_{|u|\geq 1} h(u) \Psi_S(uv) du \\ &\quad - \int_{|v|<1} \log |v| \Psi_S(-v) dv \int_{|u|>1} h(u) \Psi_S(uv) du. \end{aligned}$$

2. NONNEGATIVITY OF EIGENVALUES OF $V(h)T$

In this section we show that all eigenvalues of $V(h)T$ are nonnegative.

Lemma 2.1. ([5, Lemma 3.3 and Lemma 3.4, p. 2467]) $E_S \mathfrak{F}_S E_S^{-1}$ is unitary on $L^2(C_S)$, and its inverse is $E_S \mathfrak{F}_S^t E_S^{-1}$.

Proof of Theorem 1.1. Since $S_\Lambda S_\Lambda = S_\Lambda$, we have

$$T = S_\Lambda (1 - E_S Z_\Lambda E_S^{-1}) S_\Lambda.$$

By Lemma 2.1, $E_S \mathfrak{F}_S E_S^{-1}$ is a unitary operator on $L^2(C_S)$. This implies that $E_S Z_\Lambda E_S^{-1}$ is an orthogonal projection on $L^2(C_S)$ as it is a nonzero self-adjoint idempotent. So is $1 - E_S Z_\Lambda E_S^{-1}$. Thus,

$$\langle (1 - E_S Z_\Lambda E_S^{-1}) F, F \rangle \geq 0$$

for $F \in L^2(C_S)$. As S_Λ is bounded linear operators on $L^2(C_S)$, we have

$$\langle (1 - E_S Z_\Lambda E_S^{-1}) S_\Lambda F, S_\Lambda F \rangle \geq 0$$

for $F \in L^2(C_S)$. It follows that

$$\begin{aligned} \langle TF, F \rangle &= \langle S_\Lambda (1 - E_S Z_\Lambda E_S^{-1}) S_\Lambda F, F \rangle \\ &= \langle (1 - E_S Z_\Lambda E_S^{-1}) S_\Lambda F, S_\Lambda F \rangle \geq 0 \end{aligned}$$

for every element $F \in L^2(C_S)$. Hence, T is a positive operator on $L^2(C_S)$.

This completes the proof of Theorem 1.1. \square

Lemma 2.2. *$V(h)$ is a positive operator on $L^2(C_S)$.*

Proof. Assume that F is any element in $L^2(C_S)$ with compact support. By definition,

$$V(h)F(x) = \int_0^\infty F(\lambda) \sqrt{|x/\lambda|} d^\times \lambda \int_0^\infty g(|x/\lambda|y) g(y) dy.$$

If we make the change of variables $y \rightarrow |\lambda|y$ then

$$\begin{aligned} &\int_{C_S} V(h)F(x) \bar{F}(x) d^\times x \\ &= \int_{C_S} \bar{F}(x) d^\times x \int_{C_S} F(\lambda) \sqrt{|x/\lambda|} d^\times \lambda \int_0^\infty g(|x/\lambda|y) g(y) dy \\ &= \int_{C_S} \bar{F}(x) \sqrt{|x|} d^\times x \int_{C_S} F(\lambda) \sqrt{|\lambda|} d^\times \lambda \int_0^\infty g(|x|y) g(|\lambda|y) dy. \end{aligned}$$

Since the above triple integral is absolute integrable as F and g are compactly supported, we can change order of integration to derive

$$\int_{C_S} V(h)F(x) \bar{F}(x) d^\times x = \int_0^\infty \overline{\left(\int_{C_S} F(x) g(|x|y) \sqrt{|x|} d^\times x \right)} \left(\int_{C_S} F(\lambda) g(|\lambda|y) \sqrt{|\lambda|} d^\times \lambda \right) dy \geq 0.$$

Since S is a finite set, compactly supported functions in $L^2(C_S)$ are dense in $L^2(C_S)$. As $V(h)$ is bounded, we have

$$\langle V(h)F, F \rangle \geq 0$$

for all $F \in L^2(C_S)$. That is, $V(h)$ is a positive operator on $L^2(C_S)$.

This completes the proof of the lemma. \square

Proof of Theorem 1.2. Since T is a positive operator on the complex Hilbert space $L^2(C_S)$ by Lemma 2.2, [8, Theorem VI.9 on p. 196 and 4th paragraph on p. 195] implies that there exists a positive bounded operator B on $L^2(C_S)$, which is self-adjoint, satisfying that $T = B^2$.

Let λ be any eigenvalue of $V(h)T$ on $L^2(C_S)$. Then a nonzero element $F \in L^2(C_S)$ exists such that $V(h)T(F) = \lambda F$. If $B(F) = 0$, then $\lambda F = V(h)TF = V(h)B \cdot B(F) = 0$. Hence $\lambda = 0$ as $F \neq 0$. Thus it suffices to consider the case when $B(F) \neq 0$.

As B is bounded, we have $B(F) \in L^2(C_S)$. Assume that $B(F) \neq 0$. Since $V(h)T(F) = \lambda F$, we have $BV(h)T(F) = \lambda B(F)$. Thus

$$\lambda \langle B(F), B(F) \rangle_{L^2(C_S)} = \langle \lambda B(F), B(F) \rangle = \langle BV(h)T(F), B(F) \rangle.$$

Since B is self-adjoint and $T = B^2$, we have

$$\langle BV(h)T(F), B(F) \rangle = \langle V(h)T(F), B^2(F) \rangle = \langle V(h)T(F), T(F) \rangle.$$

By Lemma 2.2, $V(h)$ is a positive operator on the space $L^2(C_S)$. Hence

$$\langle V(h)T(F), T(F) \rangle_{L^2(C_S)} \geq 0.$$

Therefore

$$\lambda \langle B(F), B(F) \rangle_{L^2(C_S)} \geq 0.$$

Since $B(F) \neq 0$, we must have $\langle B(F), B(F) \rangle_{L^2(C_S)} > 0$ as $L^2(C_S)$ is a Hilbert space. It follows that $\lambda \geq 0$.

This completes the proof of Theorem 1.2. \square

3. TRACEABILITY OF $V(h)T$

In this section we show that $V(h)T$ is of trace class on $L^2(C_S)$.

Lemma 3.1. ([5, (3.6), p. 2470]) *We have*

$$V(g)V(g^*) = V(h) = V(g^*)V(g)$$

on $L^2(C_S)$, where $g^*(x) = x^{-1}g(x^{-1})$.

We denote $[A, B] = AB - BA$.

Lemma 3.2. *Let T be given as in Theorem 1.1. Then*

$$V(h)T = -V(g)[S_\Lambda, V(g^*)]S_\Lambda + V(g)[S_\Lambda, V(g^*)S_\Lambda E_S \mathfrak{F}_S^t J E_S^{-1}]E_S J \mathfrak{F}_S E_S^{-1} S_\Lambda.$$

where $Jf(x) = |x|^{-1}f(x^{-1})$.

Proof. Since $JS_\Lambda J = P_\Lambda$ and

$$T = S_\Lambda(S_\Lambda - E_S \mathfrak{F}_S^t P_\Lambda \mathfrak{F}_S E_S^{-1})S_\Lambda,$$

we have

$$V(h)T = V(h)S_\Lambda(S_\Lambda - E_S \mathfrak{F}_S^t JS_\Lambda J \mathfrak{F}_S E_S^{-1})S_\Lambda.$$

By Lemma 2.1, $E_S \mathfrak{F}_S^t \mathfrak{F}_S E_S^{-1}$ is the identity map on $L^2(C_S)$. Since $E_S^{-1} S_\Lambda E_S = S_\Lambda$, from Lemma 3.1 we derive that

$$\begin{aligned} & V(h)S_\Lambda(S_\Lambda - E_S \mathfrak{F}_S^t JS_\Lambda J \mathfrak{F}_S E_S^{-1})S_\Lambda \\ &= -V(g)[S_\Lambda, V(g^*)]S_\Lambda + V(g)[S_\Lambda, V(g^*)S_\Lambda E_S \mathfrak{F}_S^t J E_S^{-1}]E_S J \mathfrak{F}_S E_S^{-1} S_\Lambda. \end{aligned}$$

Hence,

$$V(h)T = -V(g)[S_\Lambda, V(g^*)]S_\Lambda + V(g)[S_\Lambda, V(g^*)S_\Lambda E_S \mathfrak{F}_S^t J E_S^{-1}]E_S J \mathfrak{F}_S E_S^{-1} S_\Lambda.$$

This completes the proof of the lemma. \square

Lemma 3.3. ([8, Theorem VI.22(h), p. 210]) A linear operator on a Hilbert space is of trace class if and only if it is a product of two Hilbert-Schmidt operators.

Lemma 3.4. ([8, Theorem VI.19, p. 207]) Let A, B be bounded linear operators on a Hilbert space \mathcal{H} . If A is of trace class on \mathcal{H} , so are AB and BA . If A, B are of trace class, so is $A + B$.

Lemma 3.5. ([8, Theorem VI.23, p. 210]) Let $\langle M, \mu \rangle$ be a measure space. A bounded linear operator A on $L^2(M, d\mu)$ is Hilbert-Schmidt if and only if there exists a function $k(x, y) \in L^2(M \times M, d\mu \otimes d\mu)$ such that

$$(Af)(x) = \int_M k(x, y)f(y)d\mu(y)$$

for all $f \in L^2(M, d\mu)$.

Lemma 3.6. We have

$$V_S(g^*)S_\Lambda E_S \mathfrak{F}_S^t J E_S^{-1} F(x) = \int_{C_S} \left(\sqrt{|x/y|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda x) \Psi_S(\lambda x/y) d\lambda \right) F(y) d^\times y$$

for $F \in L^2(C_S)$.

Proof. Let $F = E_S(f)$ with $f \in S(\mathbb{A}_S)_0$. We denote $J_0 F(x) = F(1/x)$. Since

$$(3.1) \quad E_S^{-1} J_0 F(x) = E_S^{-1} \left[\sum_{\xi \in O_S^*} \frac{1}{\sqrt{|\xi x|}} f\left(\frac{1}{\xi x}\right) \right] = \frac{1}{|x|} f\left(\frac{1}{x}\right) = (J E_S^{-1} F)(x),$$

by Lemma 2.1 we have

$$\begin{aligned} V_S(g^*)S_\Lambda E_S \mathfrak{F}_S^t J E_S^{-1} F(x) &= V_S(g^*)S_\Lambda E_S \mathfrak{F}_S^t E_S^{-1} J_0 F(x) \\ &= \int_{C_S} g^*(|x/\lambda|) \sqrt{|x/\lambda|} S_\Lambda(\lambda) [(E_S \mathfrak{F}_S^t E_S^{-1}) J_0 F](\lambda) d^\times \lambda \\ &= \int_{C_S} \sqrt{\left|\frac{\lambda}{x}\right|} g\left(\left|\frac{\lambda}{x}\right|\right) S_\Lambda(\lambda) [(E_S \mathfrak{F}_S^t E_S^{-1}) J_0 F](\lambda) d^\times \lambda \\ &= \int_{C_S} \overline{(E_S \mathfrak{F}_S E_S^{-1})_\lambda \left(\sqrt{\left|\frac{\lambda}{x}\right|} g\left(\left|\frac{\lambda}{x}\right|\right) S_\Lambda(\lambda) \right)}(t) J_0 F(t) d^\times t \end{aligned}$$

where we used the fact that g and S_Λ are real-valued functions.

By [5, Lemma 3.4 and Remark, p. 2467],

$$\begin{aligned} (e_S \mathfrak{F}_S e_S^{-1})_\lambda \left[\frac{1}{\sqrt{|x|}} g\left(\left|\frac{\lambda}{x}\right|\right) S_\Lambda(\lambda) \right](t) &= (\mathfrak{F}_S)_\lambda \left[\frac{1}{\sqrt{|x|}} g\left(\left|\frac{\lambda}{x}\right|\right) S_\Lambda(\lambda) \right](t) \\ &= \int_{\mathbb{A}_S} \frac{1}{\sqrt{|x|}} g\left(\left|\frac{\lambda}{x}\right|\right) S_\Lambda(\lambda) \Psi_S(-\lambda t) d\lambda = \sqrt{|x|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda x) \Psi_S(\lambda tx) d\lambda. \end{aligned}$$

It follows that

$$(E_S \mathfrak{F}_S E_S^{-1})_\lambda \left\{ \sqrt{\left|\frac{\lambda}{x}\right|} g\left(\left|\frac{\lambda}{x}\right|\right) S_\Lambda(\lambda) \right\}(t) = \sqrt{|tx|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda x) \Psi_S(\lambda tx) d\lambda.$$

Thus,

$$\begin{aligned} V_S(g^*)S_\Lambda E_S \mathfrak{F}_S^t J E_S^{-1} F(x) &= \int_{C_S} J_0 F(t) d^\times t \sqrt{|tx|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda x) \Psi_S(\lambda tx) d\lambda \\ &= \int_{C_S} \left(\sqrt{|x/t|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda x) \Psi_S(\lambda x/t) d\lambda \right) F(t) d^\times t. \end{aligned}$$

Since $E_S(S(\mathbb{A}_S)_0)$ is dense in $L^2(C_S)$ and since $V_S(g^*)S_\Lambda E_S \mathfrak{F}_S^t J E_S^{-1}$ is bounded on $L^2(C_S)$, this identity holds for all $F \in L^2(C_S)$.

This completes the proof of the lemma. \square

Lemma 3.7. *For each $c \in (0, 1)$ we have*

$$\left| \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda \frac{z}{y}) d\lambda \right| \ll_S \left| \frac{z}{y} \right|^{-c}.$$

Proof. Since $|\gamma|_S = 1$ for $\gamma \in O_S^*$ and since the complement of J_S in \mathbb{A}_S is negligible as S is a finite set, by changing $\lambda \rightarrow \lambda|z/y|y/z$ also with $|z/y| := (|z/y|, 1, \dots, 1)$ depending on context we can write

$$\begin{aligned} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda \frac{z}{y}) d\lambda &= \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda |\frac{z}{y}|) d\lambda \\ &= \sum_{\gamma \in O_S^*} \int_{I_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\gamma \lambda |\frac{z}{y}|) d\lambda \\ &= \sum_{\gamma \in O_S^*} \varpi(\gamma) \int_0^\infty g(u) S_\Lambda(uz) e^{-2\pi i \gamma u |z/y|} du \\ &= 2\rho_S \sum_{k \in \mathbb{N}_S} \frac{\mu(k)}{\prod_{p|k} (p-1)} \sum_{l \in \mathbb{N}_S, (l,k)=1} \int_0^\infty g(u) S_\Lambda(uz) \cos \frac{2\pi l u |z/y|}{k} du \end{aligned}$$

where $\rho_S = \prod_{p \in S'} (1 - p^{-1})$ and

$$(3.2) \quad \varpi(\gamma) = \prod_{p \in S'} \begin{cases} 1 - p^{-1} & \text{if } |\gamma|_p \leq 1, \\ -p^{-1} & \text{if } |\gamma|_p = p, \\ 0 & \text{if } |\gamma|_p > p. \end{cases}$$

By [11, Example 10, p. 162],

$$\int_0^\infty t^{s-1} \cos t dt = \Gamma(s) \cos \frac{\pi s}{2}$$

for $0 < \Re s < 1$. Since $g(u) S_\Lambda(uz)$ as a function of u has a compact support in $(0, \infty)$, we have

$$\begin{aligned} \int_{\frac{N2lu\pi|z/y|}{k}}^\infty t^{s-1} \cos t dt &= -\left(\frac{N2lu\pi|z/y|}{k}\right)^{s-1} \sin \frac{N2lu\pi|z/y|}{k} - (s-1) \int_{\frac{N2lu\pi|z/y|}{k}}^\infty t^{s-2} \sin t dt \\ &\leq \left(\frac{N2lu\pi|z/y|}{k}\right)^{\Re s-1} \left(1 + \frac{|s-1|}{1-\Re s}\right) \rightarrow 0 \end{aligned}$$

uniformly for u in the support of $g(u)S_\Lambda(uz)$ as $N \rightarrow \infty$ when $0 < \Re s < 1$. This implies that for $0 < \Re s < 1$

$$\begin{aligned} & \int_0^\infty t^{s-1} dt \int_0^\infty g(u)S_\Lambda(uz) \cos \frac{2lu\pi t|z/y|}{k} du \\ &= \lim_{N \rightarrow \infty} \int_{\sqrt{\mu}}^{\frac{1}{\sqrt{\mu}}} g(u)S_\Lambda(uz) du \int_0^N t^{s-1} \cos \frac{2lu\pi t|z/y|}{k} dt \\ &= \lim_{N \rightarrow \infty} \int_{\sqrt{\mu}}^{\frac{1}{\sqrt{\mu}}} \left(\frac{2lu\pi|z/y|}{k}\right)^{-s} g(u)S_\Lambda(uz) du \int_0^{\frac{2lu\pi N|z/y|}{k}} t^{s-1} \cos t dt \\ &= k^s (2l\pi|z/y|)^{-s} \int_0^\infty g(u)S_\Lambda(uz) u^{-s} du \int_0^\infty t^{s-1} \cos t dt \\ &= k^s (2l\pi|z/y|)^{-s} g(u) \widehat{S_\Lambda}(uz) (1-s)\Gamma(s) \cos \frac{\pi s}{2} \end{aligned}$$

where the Mellin transform $g(u) \widehat{S_\Lambda}(uz) (1-s) = \int_0^\infty g(u)S_\Lambda(uz)u^{-s}du$.

By Mellin's inversion formula [12, Theorem 28, p. 46],

$$\int_0^\infty g(u)S_\Lambda(uz) \cos \frac{2\pi lut|\frac{z}{y}|}{k} du = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} t^{-s} k^s (2l\pi|\frac{z}{y}|)^{-s} g(u) \widehat{S_\Lambda}(uz) (1-s)\Gamma(s) \cos \frac{\pi s}{2} ds$$

for $0 < c < 1$. Because $\pm\gamma \in O_S^*$ for every $\gamma \in O_S^*$, choosing $t = 1$ we get

$$\begin{aligned} & \int_{\mathbb{A}_S} g(\lambda)S_\Lambda(\lambda z) \Psi_S(\lambda \frac{z}{y}) d\lambda \\ &= \frac{\rho_S}{\pi i} \sum_{k \in \mathbb{N}_S} \frac{\mu(k)}{\prod_{p|k} (p-1)} \sum_{l \in \mathbb{N}_S, (l,k)=1} \int_{c-i\infty}^{c+i\infty} k^s (2l\pi|\frac{z}{y}|)^{-s} g(u) \widehat{S_\Lambda}(uz) (1-s)\Gamma(s) \cos \frac{\pi s}{2} ds \\ &= \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} |z/y|^{-s} g(u) \widehat{S_\Lambda}(uz) (1-s)\chi(1-s) \prod_{p \in S'} \frac{1-p^{s-1}}{1-p^{-s}} ds \end{aligned}$$

where $\chi(s) = 2^s \pi^{s-1} \Gamma(1-s) \sin \frac{\pi s}{2}$. By partial integration,

$$\begin{aligned} g(u) \widehat{S_\Lambda}(uz) (1-s) &= \int_{\max(1, \frac{1}{\Lambda|z|})}^{\frac{1}{\mu}} g(u) u^{-s} du = \frac{1}{s-1} \{ g(\max(1, \frac{1}{\Lambda|z|})) (\max(1, \frac{1}{\Lambda|z|}))^{1-s} \\ &\quad - g(\frac{1}{\mu}) \mu^{s-1} - \int_0^\infty g'(u) S_\Lambda(uz) u^{1-s} du \} \ll \frac{1}{|s-1|}. \end{aligned}$$

Stirling's formula (cf. [11, line 13, p.151]) reads that for any fixed value of σ

$$|\Gamma(\sigma + it)| \sim \sqrt{2\pi} e^{-\frac{\pi}{2}|t|} |t|^{\sigma - \frac{1}{2}}$$

as $|t| \rightarrow \infty$. Thus, we can move line of integration so that

$$\left| \int_{\mathbb{A}_S} g(\lambda)S_\Lambda(\lambda z) \Psi_S(\lambda \frac{z}{y}) d\lambda \right| \ll_S |z/y|^{-c}$$

for each $0 < c < 1$.

This completes the proof of the lemma. \square

Proof of Theorem 1.3. By Lemma 3.2,

$$(3.3) \quad V(h)T = -V(g)[S_\Lambda, V(g^*)]S_\Lambda + V(g)[S_\Lambda, V(g^*)S_\Lambda E_S \mathfrak{F}_S^t J E_S^{-1}]E_S J \mathfrak{F}_S E_S^{-1} S_\Lambda.$$

Since $E_S J E_S^{-1} F(x) = F(1/x)$ by (3.1) and since $E_S J \mathfrak{F}_S E_S^{-1} = E_S J E_S^{-1} \cdot E_S \mathfrak{F}_S E_S^{-1}$, both S_Λ and $E_S J \mathfrak{F}_S E_S^{-1}$ are bounded operators on $L^2(C_S)$. By Lemma 3.4, in order to show that $V(h)T$ is of trace class on $L^2(C_S)$ it suffices to show that $V(g)[S_\Lambda, V(g^*)]$ and

$$V(g)[S_\Lambda, V(g^*)S_\Lambda E_S \mathfrak{F}_S^t J E_S^{-1}]$$

are of trace class.

We can write

$$(3.4) \quad \begin{aligned} & V_S(g)[S_\Lambda, V_S(g^*)]F(x) \\ &= \int_{C_S} \sqrt{|x/z|} g(|x/z|) d^\times z \int_{C_S} (S_\Lambda(|z|) - S_\Lambda(|y|)) g^*(|z/y|) \sqrt{|z/y|} F(y) d^\times y \\ &= \int_{C_S} \frac{g(|\frac{x}{z}|) \sqrt{|\frac{x}{z}|}}{1 + |\log|z||} d^\times z \int_{C_S} \frac{[S_\Lambda(|z|) - S_\Lambda(|y|)] g^*(|\frac{z}{y}|) \sqrt{|\frac{z}{y}|}}{(1 + |\log|z||)^{-1}} F(y) d^\times y = ABF(x) \end{aligned}$$

for $F \in L^2(C_S)$, where the operators A and B are defined by

$$BF(z) = \int_{C_S} \frac{[S_\Lambda(|z|) - S_\Lambda(|y|)] g^*(|\frac{z}{y}|) \sqrt{|\frac{z}{y}|}}{(1 + |\log|z||)^{-1}} F(y) d^\times y$$

and

$$AG(x) = \int_{C_S} \frac{g(|\frac{x}{z}|) \sqrt{|\frac{x}{z}|}}{1 + |\log|z||} G(z) d^\times z.$$

We find that

$$\begin{aligned} & \int_{C_S \times C_S} \left| \frac{g(|\frac{x}{z}|) \sqrt{|\frac{x}{z}|}}{1 + |\log|z||} \right|^2 d^\times x d^\times z \\ &= \int_{C_S} \frac{d^\times z}{(1 + |\log|z||)^2} \int_{C_S} \left| g\left(\left|\frac{x}{z}\right|\right) \left|\frac{x}{z}\right|^{\frac{1}{2}} \right|^2 d^\times x \\ &= \int_0^\infty \frac{d^\times |z|}{(1 + |\log|z||)^2} \int_0^\infty |g(|x|)| \sqrt{|x|}^2 d^\times |x| < \infty \end{aligned}$$

as $g(x)$ has a compact support. By Lemma 3.5, A is a Hilbert-Schmidt operator.

If

$$[S_\Lambda(|y|) - S_\Lambda(|z|)] g^*(|\frac{z}{y}|) \neq 0,$$

then we must have $\sqrt{\mu} \leq |z/y| \leq \mu^{-1/2}$ and either $|y| > \Lambda^{-1}$, $|z| \leq \Lambda^{-1}$ or $|z| > \Lambda^{-1}$, $|y| \leq \Lambda^{-1}$.

If $|y| > \Lambda^{-1}$ and $|z| \leq \Lambda^{-1}$, then

$$\Lambda^{-1} < |y| \leq |z|/\sqrt{\mu} \leq \frac{1}{\sqrt{\mu}\Lambda}$$

and

$$\Lambda^{-1} \geq |z| \geq |y|\sqrt{\mu} > \frac{\sqrt{\mu}}{\Lambda}.$$

If $|z| > \Lambda^{-1}$, $|y| \leq \Lambda^{-1}$, then

$$\Lambda^{-1} < |z| \leq |y|/\sqrt{\mu} \leq \frac{1}{\sqrt{\mu}\Lambda}$$

and

$$\Lambda^{-1} \geq |y| \geq \sqrt{\mu}|z| > \frac{\sqrt{\mu}}{\Lambda}.$$

Therefore, $[S_\Lambda(|y|) - S_\Lambda(|z|)]g^*(|\frac{z}{y}|) \neq 0$ only if either

$$(3.5) \quad |z| \in (\frac{\sqrt{\mu}}{\Lambda}, \frac{1}{\Lambda}], \quad |y| \in (\frac{1}{\Lambda}, \frac{1}{\Lambda\sqrt{\mu}}] \quad \text{or} \quad |z| \in (\frac{1}{\Lambda}, \frac{1}{\Lambda\sqrt{\mu}}], \quad |y| \in (\frac{\sqrt{\mu}}{\Lambda}, \frac{1}{\Lambda}].$$

Hence

$$\begin{aligned} & \int_{C_S \times C_S} \left| (1 + |\log|z||)[S_\Lambda(|z|) - S_\Lambda(|y|)]g^*(|\frac{z}{y}|) \sqrt{|\frac{z}{y}|} \right|^2 d^\times y d^\times z \\ & \leq \int_{\frac{\sqrt{\mu}}{\Lambda}}^{\frac{1}{\Lambda\sqrt{\mu}}} (1 + |\log|z||)^2 d^\times z \int_{C_S} |g^*(|z/y|)|^2 |z/y| d^\times y \\ & = \int_{\frac{\sqrt{\mu}}{\Lambda}}^{\frac{1}{\Lambda\sqrt{\mu}}} (1 + |\log|z||)^2 d^\times z \int_{C_S} |g^*(|y|)|^2 |y| d^\times y < \infty. \end{aligned}$$

It follows from Lemma 3.5 that B is a Hilbert-Schmidt operator.

Thus $V(g)[S_\Lambda, V(g^*)]$ is a product of two Hilbert-Schmidt operators A, B on $L^2(C_S)$, and hence is of trace class on $L^2(C_S)$ by Lemma 3.3.

By Lemma 3.6, we can write

$$(3.6) \quad \begin{aligned} & V(g)[S_\Lambda, V(g^*)S_\Lambda E_S \mathfrak{F}_S^t J E_S^{-1}]F(x) = \int_{C_S} \frac{g(|\frac{x}{z}|)\sqrt{|\frac{x}{z}|}}{1 + |\log|z||} d^\times z \\ & \times \int_{C_S} \frac{S_\Lambda(|z|) - S_\Lambda(|y|)}{(1 + |\log|z||)^{-1}} \left(\sqrt{|\frac{z}{y}|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda \frac{z}{y}) d\lambda \right) F(y) d^\times y \end{aligned}$$

for $F \in L^2(C_S)$. We have already shown above that

$$\int_{C_S \times C_S} \left| \frac{g(|\frac{x}{z}|)\sqrt{|\frac{x}{z}|}}{1 + |\log|z||} \right|^2 d^\times x d^\times z < \infty.$$

Since $S_\Lambda(|z|) - S_\Lambda(|y|) \neq 0$ only if either $\Lambda|z| > 1$, $\Lambda|y| \leq 1$ or $\Lambda|y| > 1$, $\Lambda|z| \leq 1$. That is,

$$(3.7) \quad |\log|z/y|| = |\log|\Lambda z/\Lambda y|| = |\log|\Lambda z| - \log|\Lambda y|| = |\log|\Lambda z|| + |\log|\Lambda y||.$$

By Lemma 3.7

$$\sqrt{|\frac{z}{y}|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda \frac{z}{y}) d\lambda \ll_S \left| \frac{\Lambda z}{\Lambda y} \right|^{\frac{1}{2}-c}.$$

If we choose $c = 1/4$ when $|\Lambda z/\Lambda y| < 1$ and choose $c = 3/4$ when $|\Lambda z/\Lambda y| > 1$, by (3.7)

$$\sqrt{|\frac{z}{y}|} \left| \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda \frac{z}{y}) d\lambda \right| \ll_S \frac{1}{(2 + |\log|z/y||)^3} = \frac{1}{(1 + |\log|\Lambda z|| + 1 + |\log|\Lambda y||)^3}$$

for all $y, z \in C_S$ satisfying $S_\Lambda(|z|) - S_\Lambda(|y|) \neq 0$. It follows that

$$(3.8) \quad \begin{aligned} & \int_{C_S \times C_S} \left| \frac{S_\Lambda(|z|) - S_\Lambda(|y|)}{(1 + |\log|z||)^{-1}} \left(\sqrt{|\frac{z}{y}|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda \frac{z}{y}) d\lambda \right) \right|^2 d^\times y d^\times z \\ & \ll_S \int_{C_S} \frac{(1 + |\log|z||)^2}{(1 + |\log|\Lambda z||)^4} d^\times z \int_{C_S} \frac{1}{(1 + |\log|\Lambda y||)^2} d^\times y < \infty. \end{aligned}$$

Thus $V(g)[S_\Lambda, V(g^*)S_\Lambda E_S \mathfrak{F}_S^t J E_S^{-1}]$ is a product of two Hilbert-Schmidt operators by Lemma 3.5, and hence is of trace class on $L^2(C_S)$ by Lemma 3.3.

This completes the proof of Theorem 1.3. \square

4. TRACE OF $V(h)T$

In this section we compute the trace of $V(h)T$ on $L^2(C_S)$. The S -local trace formula is first given in [3, Theorem 4, p. 56] for the operator $\mathfrak{F}_S P_\Lambda \mathfrak{F}_S^t P_\Lambda$. The author's variant of it in [5, Theorem 3.16, p. 2481] is due to Meyer [7, (19), p. 549] (although the author discovered it independently without knowing its existence already in [7]).

Lemma 4.1. ([2, Corollary 3.2, p. 237]) *Let μ be a σ -finite Borel measure on a second countable space X , and let A be a trace class Hilbert-Schmidt integral operator on $L^2(X, d\mu)$. If the kernel $k(x, y)$ is continuous at (x, x) for almost every x , then*

$$\text{trace}(A) = \int_X k(x, x) d\mu(x).$$

Lemma 4.2. *$V(h)T$ is a Hilbert-Schmidt integral operator on $L^2(C_S)$ with the kernel*

$$\begin{aligned} k(x, y) = & - \int_{C_S} \frac{g(|\frac{x}{z}|) \sqrt{|\frac{x}{z}|}}{1 + |\log|z||} \cdot \frac{[S_\Lambda(|z|) - S_\Lambda(|y|)]g^*(|\frac{z}{y}|) \sqrt{|\frac{z}{y}|}}{(1 + |\log|z||)^{-1}} S_\Lambda(y) d^\times z + \int_{C_S} g(|\frac{x}{z}|) \sqrt{|\frac{x}{z}|} \\ & \times (E_S \mathfrak{F}_S E_S^{-1})_u \left[[S_\Lambda(z) - S_\Lambda(\frac{1}{|u|})] \sqrt{|zu|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda \right] (y) S_\Lambda(y) d^\times z. \end{aligned}$$

Proof. By (3.3) and (3.4),

$$(4.1) \quad V(h)T = -V(g)[S_\Lambda, V(g^*)]S_\Lambda + V(g)[S_\Lambda, V(g^*)S_\Lambda E_S \mathfrak{F}_S^t J E_S^{-1}]E_S J \mathfrak{F}_S E_S^{-1} S_\Lambda$$

and

$$(4.2) \quad \begin{aligned} & V(g)[S_\Lambda, V(g^*)]S_\Lambda F(x) \\ &= \int_{C_S} \frac{g(|\frac{x}{z}|)\sqrt{|\frac{x}{z}|}}{1+|\log|z||} d^\times z \int_{C_S} \frac{[S_\Lambda(|z|)-S_\Lambda(|y|)]g^*(|\frac{z}{y}|)\sqrt{|\frac{z}{y}|}}{(1+|\log|z||)^{-1}} S_\Lambda(y)F(y)d^\times y. \end{aligned}$$

Since

$$E_S J f(y) = \sqrt{|y|} \sum_{\xi \in O_S^*} \frac{1}{|\xi y|} f(\frac{1}{\xi y}) = \frac{1}{\sqrt{|y|}} \sum_{\gamma \in O_S^*} f(\frac{\gamma}{y}) = E_S(f)(\frac{1}{y}),$$

by (3.6) we have

$$\begin{aligned} & V(g)[S_\Lambda, V(g^*)S_\Lambda E_S \mathfrak{F}_S^t J E_S^{-1}]E_S J \mathfrak{F}_S E_S^{-1} S_\Lambda F(x) = \int_{C_S} g(|\frac{x}{z}|)\sqrt{|\frac{x}{z}|}d^\times z \\ & \times \int_{C_S} [S_\Lambda(|z|)-S_\Lambda(|y|)]\{\sqrt{|\frac{z}{y}|} \int_{\mathbb{A}_S} g(\lambda)S_\Lambda(\lambda z)\Psi_S(\lambda \frac{z}{y})d\lambda\} (E_S J \mathfrak{F}_S E_S^{-1} S_\Lambda F)(y)d^\times y \\ &= \int_{C_S} g(|\frac{x}{z}|)\sqrt{|\frac{x}{z}|}d^\times z \int_{C_S} [S_\Lambda(|z|)-S_\Lambda(|y|)] \\ & \times \{\sqrt{|\frac{z}{y}|} \int_{\mathbb{A}_S} g(\lambda)S_\Lambda(\lambda z)\Psi_S(\lambda \frac{z}{y})d\lambda\} (E_S \mathfrak{F}_S E_S^{-1} S_\Lambda F)(\frac{1}{y})d^\times y \\ &= \int_{C_S} g(|\frac{x}{z}|)\sqrt{|\frac{x}{z}|}d^\times z \int_{C_S} [S_\Lambda(|z|)-S_\Lambda(\frac{1}{|y|})] \\ & \times \{\sqrt{|zy|} \int_{\mathbb{A}_S} g(\lambda)S_\Lambda(\lambda z)\Psi_S(\lambda zy)d\lambda\} (E_S \mathfrak{F}_S E_S^{-1} S_\Lambda F)(y)d^\times y \end{aligned}$$

for $F \in L^2(C_S)$.

Since

$$[S_\Lambda(|z|)-S_\Lambda(\frac{1}{|y|})]\{\sqrt{|\frac{z}{y}|} \int_{\mathbb{A}_S} g(\lambda)S_\Lambda(\lambda z)\Psi_S(\lambda \frac{z}{y})d\lambda\} \in L^2(C_S)$$

for fixed z by (3.8), we have

$$[S_\Lambda(|z|)-S_\Lambda(\frac{1}{|y|})]\{\sqrt{|zy|} \int_{\mathbb{A}_S} g(\lambda)S_\Lambda(\lambda z)\Psi_S(\lambda zy)d\lambda\} \in L^2(C_S)$$

as a function of y . By Lemma 2.1, $E_S \mathfrak{F}_S E_S^{-1}$ is a unitary operator on $L^2(C_S)$. This implies that

$$\langle E_S \mathfrak{F}_S E_S^{-1} F, G \rangle_{L^2(C_S)} = \langle F, E_S \mathfrak{F}_S^t E_S^{-1} G \rangle_{L^2(C_S)}$$

for $F, G \in L^2(C_S)$. From this identity we derive that

$$\begin{aligned} & \int_{C_S} [S_\Lambda(|z|) - S_\Lambda(\frac{1}{|y|})] \{ \sqrt{|zy|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zy) d\lambda \} (E_S \mathfrak{F}_S^t E_S^{-1} S_\Lambda F)(y) d^\times y \\ &= \int_{C_S} (E_S \mathfrak{F}_S E_S^{-1})_u \left[[S_\Lambda(z) - S_\Lambda(\frac{1}{|u|})] \sqrt{|zu|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda \right] (y) S_\Lambda F(y) d^\times y. \end{aligned}$$

Hence,

$$\begin{aligned} (4.3) \quad & V(g)[S_\Lambda, V(g^*) S_\Lambda E_S \mathfrak{F}_S^t J E_S^{-1}] E_S J \mathfrak{F}_S E_S^{-1} S_\Lambda F(x) = \int_{C_S} g(|\frac{x}{z}|) \sqrt{|\frac{x}{z}|} d^\times z \\ & \times \int_{C_S} (E_S \mathfrak{F}_S E_S^{-1})_u \left[[S_\Lambda(z) - S_\Lambda(\frac{1}{|u|})] \sqrt{|zu|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda \right] (y) S_\Lambda F(y) d^\times y \end{aligned}$$

for $F \in L^2(C_S)$.

Let

$$\begin{aligned} k(x, y) = & - \int_{C_S} \frac{g(|\frac{x}{z}|) \sqrt{|\frac{x}{z}|}}{1 + |\log|z||} \cdot \frac{[S_\Lambda(|z|) - S_\Lambda(|y|)] g^*(|\frac{z}{y}|) \sqrt{|\frac{z}{y}|}}{(1 + |\log|z||)^{-1}} S_\Lambda(y) d^\times z + \int_{C_S} g(|\frac{x}{z}|) \sqrt{|\frac{x}{z}|} \\ & \times (E_S \mathfrak{F}_S E_S^{-1})_u \left[[S_\Lambda(z) - S_\Lambda(\frac{1}{|u|})] \sqrt{|zu|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda \right] (y) S_\Lambda(y) d^\times z. \end{aligned}$$

Then

$$V(h)TF(x) = \int_{C_S} k(x, y) F(y) d^\times y,$$

for $F \in L^2(C_S)$, where the change order of integration is permissible as the involved integrals are absolutely integrable by (4.1), (4.2), (4.3). More precisely, by (3.5) the double integral in (4.2) is absolute integrable as $g(t) = 0$ when $t \notin (\sqrt{\mu}, \mu^{-1/2})$. By Schwarz inequality, Lemma 2.1, and (3.8), we have

$$\begin{aligned} & \int_{C_S} |(E_S \mathfrak{F}_S E_S^{-1})_u \left[[S_\Lambda(z) - S_\Lambda(\frac{1}{|u|})] \sqrt{|zu|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda \right] (y) S_\Lambda F(y)| d^\times y \\ & \leq \left(\int_{C_S} |[S_\Lambda(z) - S_\Lambda(y)] \sqrt{|\frac{z}{y}|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda \frac{z}{y}) d\lambda|^2 d^\times y \right)^{\frac{1}{2}} \left(\int_{C_S} |S_\Lambda F(y)|^2 d^\times y \right)^{\frac{1}{2}} < \infty. \end{aligned}$$

This implies that the double integral in (4.3) is absolutely integrable.

Next we show that $V(h)T$ is a Hilbert-Schmidt integral operator on $L^2(C_S)$, i.e., $k(x, y) \in L^2(C_S \times C_S)$. Notice that

$$\begin{aligned} & \int_{C_S \times C_S} |k(x, y)|^2 d^\times x d^\times y \leq 2 \int_{C_S \times C_S} \left| \frac{g(|\frac{x}{z}|) \sqrt{|\frac{x}{z}|}}{1 + |\log|z||} \right|^2 d^\times z d^\times x \\ & \times \left\{ \int_{C_S \times C_S} \left| \frac{[S_\Lambda(|z|) - S_\Lambda(|y|)] g^*(|\frac{z}{y}|) \sqrt{|\frac{z}{y}|}}{(1 + |\log|z||)^{-1}} S_\Lambda(y) \right|^2 d^\times z d^\times y + \int_{C_S \times C_S} \right. \\ & \left. \left| \frac{(E_S \mathfrak{F}_S E_S^{-1})_u \left[[S_\Lambda(z) - S_\Lambda(\frac{1}{|u|})] \sqrt{|zu|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda \right] (y) S_\Lambda(y)}{(1 + |\log|z||)^{-1}} \right|^2 d^\times z d^\times y \right\}. \end{aligned}$$

By Lemma 2.1, $E_S \mathfrak{F}_S^t E_S^{-1}$ is unitary on $L^2(C_S)$. Hence,

$$\begin{aligned} & \int_{C_S} \left| (E_S \mathfrak{F}_S E_S^{-1})_u \left[[S_\Lambda(z) - S_\Lambda(\frac{1}{|u|})] \sqrt{|zu|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda \right] (y) S_\Lambda(y) \right|^2 d^\times y \\ & \leq \int_{C_S} \left| (E_S \mathfrak{F}_S E_S^{-1})_u \left[[S_\Lambda(z) - S_\Lambda(\frac{1}{|u|})] \sqrt{|zu|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda \right] (y) \right|^2 d^\times y \\ & = \int_{C_S} \left| [S_\Lambda(z) - S_\Lambda(\frac{1}{|y|})] \sqrt{|zy|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zy) d\lambda \right|^2 d^\times y \\ & = \int_{C_S} \left| [S_\Lambda(z) - S_\Lambda(|y|)] \sqrt{|z/y|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda z/y) d\lambda \right|^2 d^\times y. \end{aligned}$$

By (3.5) and (3.8),

$$\begin{aligned} & \int_{C_S \times C_S} |k(x, y)|^2 d^\times x d^\times y \leq 2 \int_{C_S \times C_S} \left| \frac{g(|\frac{x}{z}|) \sqrt{|\frac{x}{z}|}}{1 + |\log|z||} \right|^2 d^\times z d^\times x \\ & \quad \times \left\{ \int_{C_S \times C_S} \left| \frac{[S_\Lambda(z) - S_\Lambda(y)] g^*(|\frac{z}{y}|) \sqrt{|\frac{z}{y}|}}{(1 + |\log|z||)^{-1}} S_\Lambda(y) \right|^2 d^\times z d^\times y + \int_{C_S \times C_S} \right. \\ & \quad \left. \left| \frac{[S_\Lambda(z) - S_\Lambda(y)] \sqrt{|z/y|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda z/y) d\lambda}{(1 + |\log|z||)^{-1}} \right|^2 d^\times z d^\times y \right\} < \infty. \end{aligned}$$

Therefore $k(x, y) \in L^2(C_S \times C_S)$, and hence $V(h)T$ is a Hilbert-Schmidt operator on $L^2(C_S)$.

This completes the proof of the lemma. \square

Lemma 4.3. *Let $O_p = \{x \in Q_p : |x|_p \leq 1\}$ and $B = \prod_{p \in S'} p^{-1} O_p$. If $F(y) = f(|y|)$ for some function f , then*

$$\int_{\mathbb{A}_S} F(y) \Psi_S(yx) dy = \int_{\mathbb{R} \times B} F(y) \Psi_S(y|x|) dy.$$

Proof. Since the complement of J_S in \mathbb{A}_S is negligible, as $|\xi| = 1$ for $\xi \in O_S^*$ we can write

$$\int_{\mathbb{A}_S} F(y) \Psi_S(yx) dy = \sum_{\xi \in O_S^*} \int_{I_S} F(y) \Psi_S(yx\xi) dy.$$

Since $F(y) = f(|y|)$, if we change variables $y \rightarrow y|x|/x$ with $|x| = (|x|, 1, \dots, 1)$ then

$$\int_{\mathbb{A}_S} F(y) \Psi_S(yx) dy = \sum_{\xi \in O_S^*} \int_{I_S} f(|y|) \Psi_S(y|x|\xi) dy = \sum_{\xi \in O_S^*} \varpi(\xi) \int_0^\infty f(t) e^{-2\pi i t|x|\xi} dt,$$

where $\varpi(\xi)$ is given as in (3.2).

If $\xi \in O_S^*$ satisfies $\varpi(\xi) \neq 0$, then $\xi = l/k$ for some integers $k, l \in \mathbb{N}_S$ with square free k . Thus, $\xi \prod_{p \in S'} O_p^* \subset B$. Hence,

$$\bigcup_{\xi \in O_S^*, \varpi(\xi) \neq 0} \xi I_S \subset \mathbb{R}^* \times B.$$

Conversely, if $x = (x_p) \in \mathbb{R}^* \times B$ then there exists a $\xi \in O_S^*$ such that $\xi^{-1}x \in I_S$, i.e., $|\xi|_p = |x|_p \leq p$ for $p \in S'$. By (3.2), $\varpi(\xi) \neq 0$. Hence, $x \in \bigcup_{\xi \in O_S^*, \varpi(\xi) \neq 0} \xi I_S$. Thus,

$$\mathbb{R}^* \times B \subset \bigcup_{\xi \in O_S^*, \varpi(\xi) \neq 0} \xi I_S.$$

Also if ξ, γ are distinct elements in O_S^* , then ξO_S^* and γO_S^* are disjoint sets. Therefore,

$$\mathbb{R}^* \times B = \bigcup_{\xi \in O_S^*, \varpi(\xi) \neq 0} \xi I_S, \text{ a disjoint union.}$$

Therefore,

$$\sum_{\xi \in O_S^*, \varpi(\xi) \neq 0} \int_{I_S} F(y) \Psi_S(y|x|\xi) dy = \int_{\mathbb{R} \times B} F(y) \Psi_S(y|x|\xi) dy.$$

That is,

$$\int_{\mathbb{A}_S} F(y) \Psi_S(yx) dy = \int_{\mathbb{R} \times B} F(y) \Psi_S(y|x|) dy.$$

This completes the proof of the lemma. \square

Lemma 4.4. *We have*

$$(E_S \mathfrak{F}_S E_S^{-1})_u \left[\sqrt{|u|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda \right] (y) = \frac{\sqrt{|y|}}{|z|} g\left(\frac{|y|}{z}\right) S_\Lambda(y).$$

That is,

$$\int_{\mathbb{A}_S} \Psi_S(-uy) du \int_{\mathbb{A}_S} \frac{1}{|z|} g\left(\frac{\lambda}{z}\right) S_\Lambda(\lambda) \Psi_S(\lambda u) d\lambda = \frac{1}{|z|} g\left(\frac{y}{z}\right) S_\Lambda(y).$$

Proof. By [5, Lemma 3.4 and Remark, p. 2467],

$$(e_S \mathfrak{F}_S e_S^{-1})_u \left[\int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda \right] (y) = (\mathfrak{F}_S)_u \left[\int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda \right] (y).$$

It follows that

$$\begin{aligned} & (E_S \mathfrak{F}_S E_S^{-1})_u \left[\sqrt{|u|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda \right] (y) \\ &= \sqrt{|y|} (\mathfrak{F}_S)_u \left[\int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda \right] (y) = \sqrt{|y|} (\mathfrak{F}_S)_u \left[\int_{\mathbb{A}_S} \frac{1}{|z|} g\left(\frac{\lambda}{z}\right) S_\Lambda(\lambda) \Psi_S(\lambda u) d\lambda \right] (y) \\ &= \frac{\sqrt{|y|}}{|z|} \int_{\mathbb{A}_S} \Psi_S(-uy) du \int_{\mathbb{A}_S} g\left(\frac{\lambda}{z}\right) S_\Lambda(\lambda) \Psi_S(\lambda u) d\lambda. \end{aligned}$$

Hence, the second stated identity follows from the first one.

By Lemma 4.3,

$$\begin{aligned} & \int_{\mathbb{A}_S} \Psi_S(-uy) du \int_{\mathbb{A}_S} g\left(\frac{\lambda}{z}\right) S_\Lambda(\lambda) \Psi_S(\lambda u) d\lambda \\ &= \int_{\mathbb{A}_S} \Psi_S(-uy) du \int_{\mathbb{R} \times B} g\left(\frac{\lambda}{z}\right) S_\Lambda(\lambda) \Psi_S(\lambda|u|) d\lambda \\ &= \int_{\mathbb{R} \times B} \Psi_S(-uy) du \int_{\mathbb{R} \times B} g\left(\frac{\lambda}{z}\right) S_\Lambda(\lambda) \Psi_S(\lambda u) d\lambda. \end{aligned}$$

Denote

$$(4.4) \quad x = (x_r, x_b) \text{ with } x_r \in \mathbb{R}, x_b \in B.$$

We can write

$$\begin{aligned} & \int_{\mathbb{A}_S} \Psi_S(-uy) du \int_{\mathbb{A}_S} g\left(\frac{\lambda}{z}\right) S_\Lambda(\lambda) \Psi_S(\lambda u) d\lambda \\ &= \int_{-\infty}^{\infty} e^{2\pi i u_r y_r} du_r \int_{\mathbb{A}_{S'}} \Psi_{S'}(-u_b y_b) du_b \int_{\mathbb{A}_{S'}} \phi(\lambda_b) \Psi_{S'}(\lambda_b u_b) d\lambda_b \end{aligned}$$

where

$$\phi(\lambda_b) = \begin{cases} \int_{-\infty}^{\infty} g\left(\frac{|\lambda_r||\lambda_b|}{|z|}\right) S_\Lambda(|\lambda_r||\lambda_b|) e^{-2\pi i \lambda_r u_r} d\lambda_r & \text{if } \lambda_b \in B \\ 0 & \text{if } \lambda_b \notin B. \end{cases}$$

Since $\phi(\lambda_b)$ is locally constant as a function of $\lambda_b \in \mathbb{A}_{S'}$ and is supported on the compact set B , the condition of [10, Theorem 2.2.2, p. 310] is satisfied by this function. By the Fourier inversion formula (Note that [10, Theorem 2.2.2, p. 310] is still true if we replace k_p^+ there by $\mathbb{A}_{S'}$), we have

$$\begin{aligned} & \int_{\mathbb{A}_{S'}} \Psi_{S'}(-u_b y_b) du_b \int_{\mathbb{A}_{S'}} \phi(\lambda_b) \Psi_{S'}(\lambda_b u_b) d\lambda_b = \phi(y_b) \\ &= \int_{-\infty}^{\infty} g\left(\frac{|\lambda_r||y_b|}{|z|}\right) S_\Lambda(|\lambda_r||y_b|) e^{-2\pi i \lambda_r u_r} d\lambda_r. \end{aligned}$$

Since $g\left(\frac{t|y_b|}{|z|}\right) S_\Lambda(t|y_b|)$ is a continuous and compactly supported function of $t \in (0, \infty)$, $t \neq 1/\Lambda|y_b|$ and is of bounded variation in an interval including y_r , by Fourier's single-integral formula [12, Theorem 12, p. 25] we have

$$\begin{aligned} & \int_{\mathbb{A}_S} \Psi_S(-uy) du \int_{\mathbb{A}_S} g\left(\frac{\lambda}{z}\right) S_\Lambda(\lambda) \Psi_S(\lambda u) d\lambda \\ &= \int_{-\infty}^{\infty} e^{2\pi i u_r y_r} du_r \int_{-\infty}^{\infty} g\left(\frac{|\lambda_r||y_b|}{|z|}\right) S_\Lambda(|\lambda_r||y_b|) e^{-2\pi i \lambda_r u_r} d\lambda_r \\ &= \int_{-\infty}^{\infty} e^{2\pi i u_r y_r} du_r \int_{\frac{|z|\sqrt{\mu}}{|y_b|}}^{\frac{|z|}{\sqrt{\mu}|y_b|}} g\left(\frac{|\lambda_r||y_b|}{|z|}\right) S_\Lambda(|\lambda_r||y_b|) e^{-2\pi i \lambda_r u_r} d\lambda_r \\ &= \lim_{X \rightarrow \infty} \int_{\frac{|z|\sqrt{\mu}}{|y_b|}}^{\frac{|z|}{\sqrt{\mu}|y_b|}} g\left(\frac{|\lambda_r||y_b|}{|z|}\right) S_\Lambda(|\lambda_r||y_b|) d\lambda_r \int_{-X}^X e^{2\pi i u_r (y_r - \lambda_r)} du_r \\ &= \lim_{X \rightarrow \infty} \frac{1}{\pi} \int_{\frac{|z|\sqrt{\mu}}{|y_b|}}^{\frac{|z|}{\sqrt{\mu}|y_b|}} g\left(\frac{|\lambda_r||y_b|}{|z|}\right) S_\Lambda(|\lambda_r||y_b|) \frac{\sin 2\pi X(y_r - \lambda_r)}{y_r - \lambda_r} d\lambda_r = g(|y/z|) S_\Lambda(y). \end{aligned}$$

Therefore,

$$(E_S \mathfrak{F}_S E_S^{-1})_u [\sqrt{|u|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda](y) = \frac{\sqrt{|y|}}{|z|} g(|\frac{y}{z}|) S_\Lambda(y).$$

This completes the proof of the lemma. \square

Lemma 4.5. *Let $k(x, y)$ be given as in Lemma 4.2. Then*

$$k(x, x) = S_\Lambda(x) \int_{\mathbb{A}_S, |u| \geq \Lambda|x|} \Psi_S(-u) du \int_{\mathbb{A}_S} h(\lambda) S_\Lambda(\lambda x) \Psi_S(\lambda u) d\lambda.$$

Proof. By Lemma 4.2,

$$\begin{aligned} k(x, y) &= S_\Lambda(y) \sqrt{|x|} \left\{ -\frac{1}{\sqrt{|y|}} \int_{C_S} g\left(\left|\frac{x}{z}\right|\right) [S_\Lambda(|z|) - S_\Lambda(|y|)] g^*\left(\left|\frac{z}{y}\right|\right) d^\times z \right. \\ &\quad \left. + \int_{C_S} g\left(\left|\frac{x}{z}\right|\right) (E_S \mathfrak{F}_S E_S^{-1})_u \left[[S_\Lambda(z) - S_\Lambda(\frac{1}{|u|})] \sqrt{|u|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda \right] (y) d^\times z \right\}. \end{aligned}$$

Notice that

$$\begin{aligned} \int_{C_S} g\left(\left|\frac{x}{z}\right|\right) g^*\left(\left|\frac{z}{y}\right|\right) d^\times z &= \int_0^\infty g(|xz|) |y| g(|yz|) d|z| \\ &= \int_0^\infty g\left(\left|\frac{x}{y}z\right|\right) g(|z|) d|z| = h(x/y). \end{aligned}$$

By Lemma 4.4,

$$(E_S \mathfrak{F}_S E_S^{-1})_u \left[\sqrt{|u|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda \right] (y) = \frac{\sqrt{|y|}}{|z|} g(|y/z|) S_\Lambda(y)$$

for $y \in C_S$. Hence,

$$\begin{aligned} &\int_{C_S} g\left(\left|\frac{x}{z}\right|\right) (E_S \mathfrak{F}_S E_S^{-1})_u \left[S_\Lambda(z) \sqrt{|u|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda \right] (y) d^\times z \\ &= \sqrt{|y|} S_\Lambda(y) \int_{C_S} g\left(\left|\frac{x}{z}\right|\right) S_\Lambda(z) \frac{1}{|z|} g\left(\left|\frac{y}{z}\right|\right) d^\times z \\ &= \frac{S_\Lambda(y)}{\sqrt{|y|}} \int_{C_S} g\left(\left|\frac{x}{z}\right|\right) S_\Lambda(z) g^*\left(\left|\frac{z}{y}\right|\right) d^\times z. \end{aligned}$$

As $S_\Lambda(y) S_\Lambda(y) = S_\Lambda(y)$, we obtain that

$$\begin{aligned} k(x, y) &= S_\Lambda(y) \left\{ \sqrt{\frac{|x|}{|y|}} h\left(\frac{x}{y}\right) \right. \\ &\quad \left. - \sqrt{|x|} \int_{C_S} g\left(\left|\frac{x}{z}\right|\right) (E_S \mathfrak{F}_S E_S^{-1})_u \left[S_\Lambda\left(\frac{1}{|u|}\right) \sqrt{|u|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda \right] (y) d^\times z \right\}. \end{aligned}$$

By [5, Lemma 3.4 and Remark, p. 2467],

$$\begin{aligned} & (e_S \mathfrak{F}_S e_S^{-1})_u [S_\Lambda(\frac{1}{|u|}) \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda](y) \\ &= (\mathfrak{F}_S)_u [S_\Lambda(\frac{1}{|u|}) \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda](y). \end{aligned}$$

It follows that

$$\begin{aligned} & (E_S \mathfrak{F}_S E_S^{-1})_u [S_\Lambda(\frac{1}{|u|}) \sqrt{|u|} \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda](y) \\ &= \sqrt{|y|} (\mathfrak{F}_S)_u [S_\Lambda(\frac{1}{|u|}) \int_{\mathbb{A}_S} g(\lambda) S_\Lambda(\lambda z) \Psi_S(\lambda zu) d\lambda](y) \\ &= \sqrt{|y|} (\mathfrak{F}_S)_u [S_\Lambda(\frac{1}{|u|}) \int_{\mathbb{A}_S} \frac{1}{|z|} g(\frac{\lambda}{z}) S_\Lambda(\lambda) \Psi_S(\lambda u) d\lambda](y). \end{aligned}$$

Hence, by the 2nd stated identity in Lemma 4.4 we have

$$\begin{aligned} k(x, y) &= S_\Lambda(y) \left\{ \sqrt{\frac{|x|}{|y|}} h\left(\frac{x}{y}\right) \right. \\ &\quad \left. - \sqrt{|xy|} \int_{C_S} g\left(\frac{|x|}{z}\right) d^\times z \int_{\mathbb{A}_S, |u| < \Lambda} \Psi_S(-uy) du \int_{\mathbb{A}_S} \frac{1}{|z|} g\left(\frac{\lambda}{z}\right) S_\Lambda(\lambda) \Psi_S(\lambda u) d\lambda \right\}. \end{aligned}$$

Since

$$\begin{aligned} & \int_{C_S} g\left(\frac{|x|}{z}\right) d^\times z \int_{\mathbb{A}_S, |u| < \Lambda} \Psi_S(-uy) du \int_{\mathbb{A}_S} \frac{1}{|z|} g\left(\frac{\lambda}{z}\right) S_\Lambda(\lambda) \Psi_S(\lambda u) d\lambda \\ &= \int_{\mathbb{A}_S, |u| < \Lambda} \Psi_S(-uy) du \int_{C_S} g\left(\frac{|x|}{z}\right) d^\times z \int_{\mathbb{A}_S} \frac{1}{|z|} g\left(\frac{\lambda}{z}\right) S_\Lambda(\lambda) \Psi_S(\lambda u) d\lambda \\ &= \int_{\mathbb{A}_S, |u| < \Lambda} \Psi_S(-uy) du \int_{\mathbb{A}_S} \frac{1}{|x|} h\left(\frac{\lambda}{x}\right) S_\Lambda(\lambda) \Psi_S(\lambda u) d\lambda \end{aligned}$$

where the changing of the order of integration is permissible as

$$\begin{aligned} & \int_{C_S} g\left(\frac{|x|}{z}\right) d^\times z \int_{\mathbb{A}_S} \frac{1}{|z|} g\left(\frac{\lambda}{z}\right) S_\Lambda(\lambda) \Psi_S(\lambda u) d\lambda \\ &= \int_{\sqrt{\mu}|x|}^{\frac{|x|}{\sqrt{\mu}}} g\left(\frac{|x|}{|z|}\right) d^\times |z| \int_{\mathbb{A}_S} \frac{1}{|z|} g\left(\frac{\lambda}{|z|}\right) S_\Lambda(\lambda) \Psi_S(\lambda u) d\lambda. \end{aligned}$$

we have

$$k(x, x) = h(1) S_\Lambda(x) - S_\Lambda(x) |x| \int_{\mathbb{A}_S, |u| < \Lambda} \Psi_S(-ux) du \int_{\mathbb{A}_S} \frac{1}{|x|} h\left(\frac{\lambda}{x}\right) S_\Lambda(\lambda) \Psi_S(\lambda u) d\lambda.$$

That is,

$$\begin{aligned} k(x, x) &= h(1) S_\Lambda(x) - S_\Lambda(x) \int_{\mathbb{A}_S, |u| < \Lambda|x|} \Psi_S(-u) du \int_{\mathbb{A}_S} h(\lambda) S_\Lambda(\lambda x) \Psi_S(\lambda u) d\lambda \\ &= S_\Lambda(x) \int_{\mathbb{A}_S, |u| \geq \Lambda|x|} \Psi_S(-u) du \int_{\mathbb{A}_S} h(\lambda) S_\Lambda(\lambda x) \Psi_S(\lambda u) d\lambda. \end{aligned}$$

This completes the proof of the lemma. \square

Proof of Theorem 1.4. By Lemma 4.5,

$$\begin{aligned} \text{trace}_{L^2(C_S)} \{V(h)T\} &= \int_{C_S} d^\times x \int_{\mathbb{A}_S, |u| \geq \Lambda|x|} \Psi_S(u) du \int_{\mathbb{A}_S} h(\lambda) S_\Lambda(x) S_\Lambda(\lambda x) \Psi_S(\lambda u) d\lambda \\ &= \int_{\mathbb{A}_S} \Psi_S(u) du \int_{C_S, |x| \leq \frac{|u|}{\Lambda}} d^\times x \int_{\mathbb{A}_S} h(\lambda) S_\Lambda(x) S_\Lambda(\lambda x) \Psi_S(\lambda u) d\lambda. \end{aligned}$$

Since the above integrand is nonzero if and only if $\max(\frac{1}{\Lambda}, \frac{1}{\Lambda|\lambda|}) < |x| \leq \frac{|u|}{\Lambda}$, we can write

$$\begin{aligned} &\int_{C_S, |x| \leq \frac{|u|}{\Lambda}} d^\times x \int_{\mathbb{A}_S} h(\lambda) S_\Lambda(x) S_\Lambda(\lambda x) \Psi_S(\lambda u) d\lambda \\ &= \int_{\mathbb{A}_S} h(\lambda) \Psi_S(\lambda u) d\lambda \int_{C_S, \max(\frac{1}{\Lambda}, \frac{1}{\Lambda|\lambda|}) < |x| \leq \frac{|u|}{\Lambda}} S_\Lambda(x) S_\Lambda(\lambda x) d^\times x \\ &= \int_{\mathbb{A}_S} h(\lambda) \Psi_S(\lambda u) \log^+ \frac{|u|}{\max(1, \frac{1}{|\lambda|})} d\lambda. \end{aligned}$$

Hence,

$$\text{trace}_{L^2(C_S)} \{V(h)T\} = \int_{\mathbb{A}_S} \Psi_S(u) du \int_{\mathbb{A}_S} h(\lambda) \Psi_S(\lambda u) \log^+ \frac{|u|}{\max(1, \frac{1}{|\lambda|})} d\lambda.$$

Notice that

$$\begin{aligned} &\int_{\mathbb{A}_S} \Psi_S(v) dv \int_{\mathbb{A}_S} h(u) \Psi_S(uv) \log^+ \frac{|v|}{\max(1, \frac{1}{|u|})} du \\ &= \int_{\mathbb{A}_S, |v| \geq 1} \Psi_S(v) \log |v| dv \int_{\mathbb{A}_S, |u| \geq 1} h(u) \Psi_S(-uv) du \\ &\quad + \int_{\mathbb{A}_S} \Psi_S(v) dv \int_{\mathbb{A}_S, |uv| \geq 1, |u| < 1} h(u) \Psi_S(-uv) \log |uv| du. \end{aligned}$$

We can write

$$\begin{aligned} &\int_{\mathbb{A}_S} \Psi_S(v) dv \int_{\mathbb{A}_S, |uv| \geq 1, |u| < 1} h(u) \Psi_S(-uv) \log |uv| du \\ &= \int_{\mathbb{A}_S} \Psi_S(v) dv \int_{\mathbb{A}_S, |u| \geq 1, |u/v| < 1} \frac{1}{|v|} h(u/v) \Psi_S(-u) \log |u| du \\ &= \int_{\mathbb{A}_S} \Psi_S(vu) dv \int_{\mathbb{A}_S, |u| \geq 1, |1/v| < 1} \frac{1}{|v|} h(1/v) \Psi_S(-u) \log |u| du \\ &= \int_{|u| \geq 1} \Psi_S(-u) \log |u| du \int_{|v| > 1} h(v) \Psi_S(vu) dv \\ &= \int_{|v| \geq 1} \Psi_S(v) \log |v| dv \int_{|u| > 1} h(u) \Psi_S(-uv) du. \end{aligned}$$

Hence,

$$\begin{aligned} \text{trace}_{L^2(C_S)} \{V(h)T\} &= \int_{\mathbb{A}_S, |v|>1} \Psi_S(v) \log |v| dv \int_{\mathbb{A}_S, |u|\geq 1} h(u) \Psi_S(-uv) du \\ &\quad + \int_{|v|>1} \Psi_S(v) \log |v| dv \int_{|u|>1} h(u) \Psi_S(-uv) du. \end{aligned}$$

This completes the proof of Theorem 1.4. \square

5. PROOF OF COROLLARY 1.5

Lemma 5.1. *We can write*

$$\begin{aligned} &\int_{\mathbb{A}_S, |v|>1} \Psi_S(v) \log |v| dv \int_{\mathbb{A}_S, |u|\geq 1} h(u) \Psi_S(-uv) du \\ &= 4 \sum_{k_1, k_2, l_1, l_2 \in \mathbb{N}_S} \frac{\mu(k_1)\mu(k_2)}{k_1 k_2} \int_1^\infty \log v \cos \frac{2\pi l_1 v}{k_1} dv \int_1^\infty h(u) \cos \frac{2\pi l_2 uv}{k_2} du. \end{aligned}$$

Proof. As $|\beta| = 1$, $|\gamma| = 1$ for $\beta, \gamma \in O_S^*$,

$$\begin{aligned} &\int_{\mathbb{A}_S, |v|>1} \Psi_S(v) \log |v| dv \int_{\mathbb{A}_S, |u|\geq 1} h(u) \Psi_S(-uv) du \\ &= \sum_{\beta \in O_S^*} \int_{\beta I_S, |v|>1} \Psi_S(v) \log |v| dv \sum_{\gamma \in O_S^*} \int_{\gamma I_S, |u|\geq 1} h(u) \Psi_S(-uv) du \\ &= \sum_{\beta \in O_S^*} \varpi(\beta) \int_1^\infty e^{-2\pi i v \beta} \log |v| dv \sum_{\gamma \in O_S^*} \varpi(\gamma) \int_1^\infty h(u) e^{2\pi \gamma u v} du \end{aligned}$$

where

$$\varpi(\beta) = \prod_{p \in S'} \begin{cases} 1 - p^{-1} & \text{if } |\beta|_p \leq 1, \\ -p^{-1} & \text{if } |\beta|_p = p, \\ 0 & \text{if } |\beta|_p > p. \end{cases}$$

That is,

$$\begin{aligned} &\int_{\mathbb{A}_S, |v|>1} \Psi_S(v) \log |v| dv \int_{\mathbb{A}_S, |u|\geq 1} h(u) \Psi_S(-uv) du \\ (5.1) \quad &= 4\rho_S^2 \sum_{k_1, l_1 \in \mathbb{N}_S, (k_1, l_1)=1} \frac{\mu(k_1)}{\prod_{p|k_1} (p-1)} \int_1^\infty \log v \cos \frac{2\pi l_1 v}{k_1} dv \\ &\times \sum_{k_2, l_2 \in \mathbb{N}_S, (k_2, l_2)=1} \frac{\mu(k_2)}{\prod_{p|k_2} (p-1)} \int_1^\infty h(u) \cos \frac{2\pi l_2 uv}{k_2} du \end{aligned}$$

By partial integration,

$$\begin{aligned}
& \int_1^\infty h(u) \cos \frac{2\pi l_2 u v}{k_2} du \\
(5.2) \quad &= \frac{k_2}{2\pi l_2 v} \left(-h(1) \sin \frac{2\pi l_2 v}{k_2} - \int_1^\infty h'(u) \sin \frac{2\pi l_2 u v}{k_2} du \right) dv \\
&= -h(1) \frac{k_2}{2\pi l_2 v} \sin \frac{2\pi l_2 v}{k_2} - h'(1) \left(\frac{k_2}{2\pi l_2 v} \right)^2 \cos \frac{2\pi l_2 v}{k_2} - \left(\frac{k_2}{2\pi l_2 v} \right)^2 \int_1^\infty h''(u) \cos \frac{2\pi l_2 u v}{k_2} du
\end{aligned}$$

and

$$\begin{aligned}
(5.3) \quad & \int_1^\infty \log v \cos \frac{2\pi l_1 v}{k_1} \left(\sum_{\substack{k_2, l_2 \in \mathbb{N}_S, (k_2, l_2)=1 \\ k_1 l_2 \neq k_2 l_1}} \frac{\mu(k_2)}{\prod_{p|k_2} (p-1)} h(1) \frac{-k_2}{2\pi l_2 v} \sin \frac{2\pi l_2 v}{k_2} \right) dv \\
&= \int_1^\infty \frac{\log v}{v} \left(\sum_{\substack{k_2, l_2 \in \mathbb{N}_S, (k_2, l_2)=1 \\ k_1 l_2 \neq k_2 l_1}} \frac{\mu(k_2)}{\prod_{p|k_2} (p-1)} h(1) \frac{-k_2}{4\pi l_2} \left[\sin 2\pi v \left(\frac{l_2}{k_2} + \frac{l_1}{k_1} \right) + \sin 2\pi v \left(\frac{l_2}{k_2} - \frac{l_1}{k_1} \right) \right] \right) dv \\
&= \int_1^\infty \frac{\log v - 1}{v^2} \sum_{\substack{k_2, l_2 \in \mathbb{N}_S, (k_2, l_2)=1 \\ k_1 l_2 \neq k_2 l_1}} \frac{\mu(k_2)}{\prod_{p|k_2} (p-1)} \frac{h(1) k_2}{4\pi l_2} \left\{ \frac{\cos 2\pi v \left(\frac{l_2}{k_2} - \frac{l_1}{k_1} \right)}{2\pi \left(\frac{l_2}{k_2} - \frac{l_1}{k_1} \right)} + \frac{\cos 2\pi v \left(\frac{l_2}{k_2} + \frac{l_1}{k_1} \right)}{2\pi \left(\frac{l_2}{k_2} + \frac{l_1}{k_1} \right)} \right\} dv \\
&= \sum_{\substack{k_2, l_2 \in \mathbb{N}_S, (k_2, l_2)=1 \\ k_1 l_2 \neq k_2 l_1}} \frac{\mu(k_2)}{\prod_{p|k_2} (p-1)} \frac{h(1) k_2}{4\pi l_2} \int_1^\infty \left\{ \frac{\cos 2\pi v \left(\frac{l_2}{k_2} - \frac{l_1}{k_1} \right)}{2\pi \left(\frac{l_2}{k_2} - \frac{l_1}{k_1} \right)} + \frac{\cos 2\pi v \left(\frac{l_2}{k_2} + \frac{l_1}{k_1} \right)}{2\pi \left(\frac{l_2}{k_2} + \frac{l_1}{k_1} \right)} \right\} \frac{\log v - 1}{v^2} dv \\
&= \sum_{\substack{k_2, l_2 \in \mathbb{N}_S, (k_2, l_2)=1 \\ k_1 l_2 \neq k_2 l_1}} \frac{\mu(k_2)}{\prod_{p|k_2} (p-1)} h(1) \frac{-k_2}{4\pi l_2} \int_1^\infty \frac{\log v}{v} \cos \frac{2\pi l_1 v}{k_1} \sin \frac{2\pi l_2 v}{k_2} dv.
\end{aligned}$$

Note that

$$\begin{aligned}
& \int_1^\infty \log v \cos \frac{2\pi l_1 v}{k_1} h(1) \frac{-k_2}{2\pi l_2 v} \sin \frac{2\pi l_2 v}{k_2} dv \\
(5.4) \quad &= \begin{cases} \frac{h(1) k_2}{4\pi l_2} \int_1^\infty \left\{ \frac{\cos 2\pi v \left(\frac{l_2}{k_2} - \frac{l_1}{k_1} \right)}{2\pi \left(\frac{l_2}{k_2} - \frac{l_1}{k_1} \right)} + \frac{\cos 2\pi v \left(\frac{l_2}{k_2} + \frac{l_1}{k_1} \right)}{2\pi \left(\frac{l_2}{k_2} + \frac{l_1}{k_1} \right)} \right\} \frac{\log v - 1}{v^2} dv & \text{if } k_2 l_1 \neq k_1 l_2 \\ \frac{h(1) k_2}{4\pi l_2} \int_1^\infty \frac{\cos 2\pi v \left(\frac{l_2}{k_2} + \frac{l_1}{k_1} \right)}{2\pi \left(\frac{l_2}{k_2} + \frac{l_1}{k_1} \right)} \frac{\log v - 1}{v^2} dv & \text{if } k_2 l_1 = k_1 l_2. \end{cases}
\end{aligned}$$

Since \mathbb{N}_S contains only finitely many square free integers k_1, k_2 , there are only finitely many integers l_1, l_2 in \mathbb{N}_S such that $k_1 l_2 - k_2 l_1 = 0$ for some square free integers k_1, k_2 in \mathbb{N}_S . It follows from (5.3) that

$$\begin{aligned}
(5.5) \quad & \int_1^\infty \log v \cos \frac{2\pi l_1 v}{k_1} \left(\sum_{k_2, l_2 \in \mathbb{N}_S, (k_2, l_2)=1} \frac{\mu(k_2)}{\prod_{p|k_2} (p-1)} h(1) \frac{-k_2}{2\pi l_2 v} \sin \frac{2\pi l_2 v}{k_2} \right) dv \\
&= \sum_{k_2, l_2 \in \mathbb{N}_S, (k_2, l_2)=1} \frac{\mu(k_2)}{\prod_{p|k_2} (p-1)} h(1) \frac{-k_2}{4\pi l_2} \int_1^\infty \frac{\log v}{v} \cos \frac{2\pi l_1 v}{k_1} \sin \frac{2\pi l_2 v}{k_2} dv.
\end{aligned}$$

From (5.2) and (5.5) we deduce that

$$(5.6) \quad \begin{aligned} & \int_1^\infty \log v \cos \frac{2\pi l_1 v}{k_1} dv \sum_{k_2, l_2 \in \mathbb{N}_S, (k_2, l_2)=1} \frac{\mu(k_2)}{\prod_{p|k_2}(p-1)} \int_1^\infty h(u) \cos \frac{2\pi l_2 uv}{k_2} du \\ &= \sum_{k_2, l_2 \in \mathbb{N}_S, (k_2, l_2)=1} \frac{\mu(k_2)}{\prod_{p|k_2}(p-1)} \int_1^\infty \log v \cos \frac{2\pi l_1 v}{k_1} dv \int_1^\infty h(u) \cos \frac{2\pi l_2 uv}{k_2} du. \end{aligned}$$

By (5.1) and (5.6),

$$(5.7) \quad \begin{aligned} & \int_{\mathbb{A}_S, |v|>1} \Psi_S(v) \log |v| dv \int_{\mathbb{A}_S, |u|\geq 1} h(u) \Psi_S(-uv du \\ &= 4\rho_S^2 \sum_{k_1, l_1 \in \mathbb{N}_S, (k_1, l_1)=1} \frac{\mu(k_1)}{\prod_{p|k_1}(p-1)} \sum_{k_2, l_2 \in \mathbb{N}_S, (k_2, l_2)=1} \frac{\mu(k_2)}{\prod_{p|k_2}(p-1)} \\ & \quad \times \int_1^\infty \log v \cos \frac{2\pi l_1 v}{k_1} dv \int_1^\infty h(u) \cos \frac{2\pi l_2 uv}{k_2} du \end{aligned}$$

By [11, Example 10, p. 162],

$$\int_0^\infty t^{s-1} \cos t dt = \Gamma(s) \cos \frac{\pi s}{2}$$

for $0 < \Re s < 1$. Since $h(u) = 0$ for $u \notin (\mu, \mu^{-1})$, we can change the order of integration and derive that for $0 < \Re s < 1$

$$\begin{aligned} & \int_0^\infty v^{s-1} dt \int_1^\infty h(u) \cos \frac{2\pi l_2 uv}{k_2} du = \int_1^\infty h(u) du \int_0^\infty v^{s-1} \cos \frac{2\pi l_2 uv}{k_2} dv \\ &= k_2^s (2l_2 \pi)^{-s} \int_1^\infty h(u) u^{-s} du \int_0^\infty v^{s-1} \cos v dv \\ &= k_2^s (2l_2 \pi)^{-s} \Gamma(s) \cos \frac{\pi s}{2} \int_1^\infty h(u) u^{-s} du. \end{aligned}$$

Since

$$k_2^s (2l_2 \pi)^{-s} \Gamma(s) \cos \frac{\pi s}{2} \int_1^\infty h(u) u^{-s} du$$

is analytic for $0 < \Re s < 1$ and is $\ll |s|^{\Re s - 3/2}$ in this vertical strip when $|s| \rightarrow \infty$. By Mellin's inversion formula [12, Theorem 28, p. 46],

$$\begin{aligned} \int_1^\infty h(u) \cos \frac{2\pi l_2 uv}{k_2} du &= \frac{1}{2\pi i} \int_{\Re s=c}^{c+i\infty} k_2^s (2l_2 \pi)^{-s} \Gamma(s) \cos \frac{\pi s}{2} \left(\int_1^\infty h(u) u^{-s} du \right) v^{-s} ds \\ &= \frac{1}{4\pi i} \int_{c-i\infty}^{c+i\infty} v^{-s} \chi(1-s) \left(\frac{k_2}{l_2} \right)^s \left(\int_1^\infty h(u) u^{-s} du \right) ds \end{aligned}$$

for $0 < c < 1$, where $\chi(s) = 2^s \pi^{s-1} \Gamma(1-s) \sin \frac{\pi s}{2}$.

Similarly, by (5.2) and (5.4) we find that

$$(5.8) \quad \begin{aligned} & \int_1^\infty \log v \cos \frac{2\pi l_1 v}{k_1} dv \int_1^\infty h(u) \cos \frac{2\pi l_1 u v}{k_1} du = \frac{1}{4\pi i} \int_{c-i\infty}^{c+i\infty} \chi(1-z) \left(\frac{k_1}{l_1}\right)^z \\ & \times \left(\int_1^\infty \log v \left\{ \frac{1}{4\pi i} \int_{c-i\infty}^{c+i\infty} v^{-s} \chi(1-s) \left(\frac{k_2}{l_2}\right)^s \left(\int_1^\infty h(u) u^{-s} du \right) ds \right\} v^{-z} dv \right) dz \end{aligned}$$

for $0 < c < 1$.

Since \mathbb{N}_S contains only finitely many primes and since $|\Gamma(x+iy)| \sim \sqrt{2\pi}|y|^{x-1/2}e^{-\pi|y|/2}$ for fixed x when $|y| \rightarrow \infty$, the double series and the double integral

$$\begin{aligned} & 4\rho_S^2 \sum_{k_1, l_1 \in \mathbb{N}_S, (k_1, l_1)=1} \frac{\mu(k_1)}{\prod_{p|k_1} (p-1)} \sum_{k_2, l_2 \in \mathbb{N}_S, (k_2, l_2)=1} \frac{\mu(k_2)}{\prod_{p|k_2} (p-1)} \frac{1}{4\pi i} \int_{c-i\infty}^{c+i\infty} \chi(1-z) \left(\frac{k_1}{l_1}\right)^z \\ & \times \left(\int_1^\infty \log v \left\{ \frac{1}{4\pi i} \int_{c-i\infty}^{c+i\infty} v^{-s} \chi(1-s) \left(\frac{k_2}{l_2}\right)^s \left(\int_1^\infty h(u) u^{-s} du \right) ds \right\} v^{-z} dv \right) dz \end{aligned}$$

are absolute convergent for $0 < c < 1/2$. Thus, by (5.7) and (5.8)

$$\begin{aligned} & \int_{\mathbb{A}_S, |v|>1} \Psi_S(v) \log |v| dv \int_{\mathbb{A}_S, |u|\geq 1} h(u) \Psi_S(-uv du) \\ & = 4\rho_S^2 \frac{1}{4\pi i} \int_{c-i\infty}^{c+i\infty} \chi(1-z) \left[\sum_{k_1, l_1 \in \mathbb{N}_S, (k_1, l_1)=1} \frac{\mu(k_1)}{\prod_{p|k_1} (p-1)} \left(\frac{k_1}{l_1}\right)^z \right] \int_1^\infty \log v \\ & \times \left\{ \frac{1}{4\pi i} \int_{c-i\infty}^{c+i\infty} v^{-s} \chi(1-s) \left(\sum_{k_2, l_2 \in \mathbb{N}_S, (k_2, l_2)=1} \frac{\mu(k_2)}{\prod_{p|k_2} (p-1)} \left(\frac{k_2}{l_2}\right)^s \right) \left(\int_1^\infty h(u) u^{-s} du \right) ds \right\} v^{-z} dv \} dz \end{aligned}$$

for $0 < c < 1/2$.

As

$$\begin{aligned} \rho_S & \sum_{k_1, l_1 \in \mathbb{N}_S, (k_1, l_1)=1} \frac{\mu(k_1)}{\prod_{p|k_1} (p-1)} \left(\frac{k_1}{l_1}\right)^z = \sum_{k_1 \in \mathbb{N}_S} \prod_{p|k_1} \frac{-1}{p^{1-z}} \cdot \prod_{p \nmid k_1} \frac{1-p^{-1}}{1-p^{-z}} \\ & = \prod_{p \in S'} \frac{1-p^{-1}}{1-p^{-z}} \sum_{k_1 \in \mathbb{N}_S} \prod_{p|k_1} \frac{-(1-p^{-z})}{p^{1-z}(1-p^{-1})} \\ & = \prod_{p \in S'} \frac{1-p^{-1}}{1-p^{-z}} \sum_{p \in S'} \left(1 + \frac{-(1-p^{-z})}{p^{1-z}(1-p^{-1})} \right) \\ & = \prod_{p \in S'} \frac{1-p^{-1}}{1-p^{-z}} \sum_{p \in S'} \frac{1-p^{z-1}}{1-p^{-1}} = \prod_{p \in S'} \frac{1-p^{z-1}}{1-p^{-z}}, \end{aligned}$$

we have

$$\begin{aligned}
& \int_{\mathbb{A}_S, |v| > 1} \Psi_S(v) \log |v| dv \int_{\mathbb{A}_S, |u| \geq 1} h(u) \Psi_S(-uv du) \\
&= \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \chi(1-z) \prod_{p \in S'} \frac{1-p^{z-1}}{1-p^{-z}} \left[\int_1^\infty \log v \right. \\
&\quad \times \left. \left\{ \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} v^{-s} \chi(1-s) \prod_{p \in S'} \frac{1-p^{s-1}}{1-p^{-s}} \left(\int_1^\infty h(u) u^{-s} du \right) ds \right\} v^{-z} dv \right] dz \\
&= \sum_{k_1, k_2, l_1, l_2 \in \mathbb{N}_S} \frac{\mu(k_1)\mu(k_2)}{k_1 k_2} \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \chi(1-z) \left(\frac{k_1}{l_1} \right)^z \int_1^\infty \log v \\
&\quad \times \left\{ \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} v^{-s} \chi(1-s) \left(\frac{k_2}{l_2} \right)^s \left(\int_1^\infty h(u) u^{-s} du \right) ds \right\} v^{-z} dv \} dz \\
&= 4 \sum_{k_1, k_2, l_1, l_2 \in \mathbb{N}_S} \frac{\mu(k_1)\mu(k_2)}{k_1 k_2} \int_1^\infty \log v \cos \frac{2\pi l_1 v}{k_1} dv \int_1^\infty h(u) \cos \frac{2\pi l_2 u v}{k_2} du.
\end{aligned}$$

This completes the proof of the lemma. \square

Lemma 5.2. ([4, Theorem 8.4, p.101]) If A is of trace class on a Hilbert space, then the functional trace of A coincides with its spectral trace.

Proof of Corollary 1.5. By Theorem 2.4, all eigenvalues of $V(h)T$ on $L^2(C_S)$ are nonnegative. Also, by Theorem 3.8 the operator $V(h)T$ is of trace class on the Hilbert space $L^2(C_S)$. It follows from Lemma 5.2 that the functional trace of $V(h)T$ is nonnegative.

As the functional trace of $V(h)T$ on $L^2(C_S)$ is equal to

$$\begin{aligned}
& \int_{\mathbb{A}_S, |v| > 1} \Psi_S(v) \log |v| dv \int_{\mathbb{A}_S, |u| \geq 1} h(u) \Psi_S(-uv du) \\
&+ \int_{|v| > 1} \Psi_S(v) \log |v| dv \int_{|u| > 1} h(u) \Psi_S(-u(|v|, 1, \dots, 1)) du
\end{aligned}$$

by Theorem 4.6, from Lemma 5.1 we derive that

$$\sum_{k_1, k_2, l_1, l_2 \in \mathbb{N}_S} \frac{\mu(k_1)\mu(k_2)}{k_1 k_2} \int_1^\infty \log v \cos \frac{2\pi l_1 v}{k_1} dv \int_1^\infty h(u) \cos \frac{2\pi l_2 u v}{k_2} du \geq 0$$

for all $g \in C_c^\infty(0, \infty)$.

This completes the proof of Corollary 1.5. \square

6. A RELATION TO THE WEIL DISTRIBUTION

Lemma 6.1. ([5, line 1-3 from bottom on p. 2483 and (3.15) on p. 2484]) We have

$$\int_{\mathbb{A}_S} \mathfrak{F}_S h(y) \Psi_S(y) \log |y| dy = - \sum_p \int'_{Q_p^*} \frac{h(u^{-1})}{|1-u|_p} d^* u$$

where the sum on p is over all primes of Q including the infinity prime.

We denote $x = (x_r, x_b)$ with $x_r \in \mathbb{R}$ and $x_b \in \mathbb{A}_{S'}$.

Lemma 6.2. *Let $f(\lambda) = h(\lambda) \log \max(1, \frac{1}{|\lambda|})$. Then*

$$\int_{\mathbb{A}_S} \Psi_S(-uy) du \int_{\mathbb{A}_S} f(\lambda) \Psi_S(\lambda u) d\lambda = f(y).$$

Proof. By [5, Lemma 3.4 and Remark, p. 2467],

$$(e_S \mathfrak{F}_S e_S^{-1})_u \left[\int_{\mathbb{A}_S} f(\lambda) \Psi_S(\lambda u) d\lambda \right] (y) = (\mathfrak{F}_S)_u \left[\int_{\mathbb{A}_S} f(\lambda) \Psi_S(\lambda u) d\lambda \right] (y).$$

That is,

$$\begin{aligned} & (E_S \mathfrak{F}_S E_S^{-1})_u \left[\sqrt{|u|} \int_{\mathbb{A}_S} f(\lambda) \Psi_S(\lambda u) d\lambda \right] (y) \\ &= \sqrt{|y|} (\mathfrak{F}_S)_u \left[\int_{\mathbb{A}_S} f(\lambda) \Psi_S(\lambda u) d\lambda \right] (y) \\ &= \sqrt{|y|} \int_{\mathbb{A}_S} \Psi_S(-uy) du \int_{\mathbb{A}_S} f(\lambda) \Psi_S(\lambda u) d\lambda. \end{aligned}$$

Since

$$\begin{aligned} \int_{\mathbb{A}_S} f(\lambda) \Psi_S(\lambda u) d\lambda &= \sum_{\gamma \in O_S^*} \int_{\gamma I_S} f(\lambda) \Psi_S(\lambda u) d\lambda \\ &= \sum_{\gamma \in O_S^*} \int_{I_S} f(\lambda) \Psi_S(\lambda \gamma u) d\lambda, \end{aligned}$$

we have

$$E_S^{-1} \left\{ \sqrt{|u|} \int_{\mathbb{A}_S} f(\lambda) \Psi_S(\lambda u) d\lambda \right\} = \int_{I_S} f(\lambda) \Psi_S(\lambda u) d\lambda.$$

Hence,

$$\begin{aligned} & (\mathfrak{F}_S E_S^{-1})_u \left\{ \sqrt{|u|} \int_{\mathbb{A}_S} f(\lambda) \Psi_S(\lambda u) d\lambda \right\} (y) \\ &= \int_{\mathbb{A}_S} \Psi_S(-uy) du \int_{I_S} f(\lambda) \Psi_S(\lambda u) d\lambda \\ &= \int_{-\infty}^{\infty} e^{2\pi i u_r y_r} du_r \int_{\mathbb{A}_{S'}} \Psi_{S'}(-u_b y_b) du_b \int_{\mathbb{A}_{S'}} \phi(\lambda_b) \Psi_{S'}(\lambda_b u_b) d\lambda_b \end{aligned}$$

where

$$\phi(\lambda_b) = \begin{cases} \int_{-\infty}^{\infty} f(|\lambda_r||\lambda_b|) e^{-2\pi i \lambda_r u_r} d\lambda_r & \text{if } \lambda_b \in \prod_{p \in S'} O_p^* \\ 0 & \text{if } \lambda_b \notin \prod_{p \in S'} O_p^*. \end{cases}$$

Also,

$$\int_{I_S} f(\lambda) \Psi_S(\lambda u) d\lambda = \varpi(u) \int_0^{\infty} f(t) e^{-2\pi i t u_r} dt.$$

This implies that

$$\int_{\mathbb{A}_{S'}} \phi(\lambda_b) \Psi_{S'}(\lambda_b u_b) d\lambda_b$$

as a function of u_b is supported on the compact set $B = \prod_{p \in S'} p^{-1} O_p$ with $O_p = \{x \in Q_p : |x|_p \leq 1\}$. Since $\phi(\lambda_b)$ is locally constant as a function of $\lambda_b \in \mathbb{A}_{S'}$ and is supported on the compact set $\prod_{p \in S'} O_p^*$, the condition of [10, Theorem 2.2.2, p. 310] is satisfied by this function. By the Fourier inversion formula (Note that [10, Theorem 2.2.2, p. 310] is still true if we replace k_p^+ there by $\mathbb{A}_{S'}$), we have

$$\int_{\mathbb{A}_{S'}} \Psi_{S'}(-u_b y_b) du_b \int_{\mathbb{A}_{S'}} \phi(\lambda_b) \Psi_{S'}(\lambda_b u_b) d\lambda_b = \phi(y_b)$$

where $\phi(y_b) = 0$ if $y_b \notin \prod_{p \in S'} O_p^*$.

Since $f(t|y_b|)$ is a continuous and compactly supported function of $t \in (0, \infty)$, $t \neq 1/\Lambda|y_b|$ and is of bounded variation in an interval including y_r , by Fourier's single-integral formula [12, Theorem 12, p. 25] we have for $y_b \in \prod_{p \in S'} O_p^*$

$$\begin{aligned} & (\mathfrak{F}_S E_S^{-1})_u \left\{ \sqrt{|u|} \int_{\mathbb{A}_S} f(\lambda) \Psi_S(\lambda u) d\lambda \right\} (y) \\ &= \int_{-\infty}^{\infty} e^{2\pi i u_r y_r} du_r \int_{-\infty}^{\infty} f(|\lambda_r| |y_b|) e^{-2\pi i \lambda_r u_r} d\lambda_r \\ &= \lim_{X \rightarrow \infty} \int_{-\infty}^{\infty} f(|\lambda_r| |y_b|) d\lambda_r \int_{-X}^X e^{2\pi i u_r (y_r - \lambda_r)} du_r \\ &= \lim_{X \rightarrow \infty} \frac{1}{\pi} \int_{-\infty}^{\infty} f(|\lambda_r| |y_b|) \frac{\sin 2\pi X (y_r - \lambda_r)}{y_r - \lambda_r} d\lambda_r = f(y) \end{aligned}$$

where the 2nd equality holds because f has a compact support on $(0, \infty)$. It follows that

$$\begin{aligned} & (E_S \mathfrak{F}_S E_S^{-1})_u \left\{ \sqrt{|u|} \int_{\mathbb{A}_S} f(\lambda) \Psi_S(\lambda u) d\lambda \right\} (y) \\ &= \sqrt{|y|} \sum_{\xi \in O_S^*} \begin{cases} f(\xi y) & \text{if } \xi y \in I_S, \\ 0 & \text{if } \xi y \notin I_S \end{cases} = \sqrt{|y|} f(y) \end{aligned}$$

because for each $y \in \mathbb{A}_S$ with $|y| \neq 0$ there exists exactly one $\xi \in O_S^*$ such that $\xi y \in I_S$. Therefore,

$$(E_S \mathfrak{F}_S E_S^{-1})_u [\sqrt{|u|} \int_{\mathbb{A}_S} f(\lambda) \Psi_S(\lambda z u) d\lambda] (y) = \sqrt{|y|} f(y).$$

This completes the proof of the lemma. \square

Corollary 6.3. *We have*

$$\int_{\mathbb{A}_S} \Psi_S(-v) dv \int_{\mathbb{A}_S} h(y) \log \max(1, \frac{1}{|y|}) \Psi_S(yv) dy = 0.$$

Proof. By Lemma 6.2,

$$\int_{\mathbb{A}_S} \Psi_S(-vx) dv \int_{\mathbb{A}_S} h(y) \log \max(1, \frac{1}{|y|}) \Psi_S(yv) dy = h(x) \log \max(1, \frac{1}{|x|}).$$

Choosing $x = 1$ in the above identity gives the stated identity.

This completes the proof of the lemma. \square

We denote $\log^+ x = \log x$ for $x > 1$, $\log^+ x = 0$ for $0 < x \leq 1$ and $\log^- x = \log x$ if $0 < x \leq 1$, $\log^- x = 0$ if $x > 1$.

Proof of Theorem 1.6. Since

$$\int_{\mathbb{A}_S} \Psi_S(-v) dv \int_{\mathbb{A}_S} h(u) \log \max(1, \frac{1}{|u|}) \Psi_S(uv) du = 0,$$

by Lemma 6.1 and Corollary 6.3 we can write

$$\begin{aligned} \Delta(h) &= \int_{\mathbb{A}_S} \Psi_S(-v) dv \int_{\mathbb{A}_S} h(u) \log |v| \Psi_S(uv) du \\ &\quad - \int_{\mathbb{A}_S} \Psi_S(-v) dv \int_{\mathbb{A}_S} h(u) \log \max(1, \frac{1}{|u|}) \Psi_S(uv) du \\ &= \int_{\mathbb{A}_S} \Psi_S(-v) dv \int_{\mathbb{A}_S} h(u) \log \frac{|v|}{\max(1, \frac{1}{|u|})} \Psi_S(uv) du \\ &= \int_{\mathbb{A}_S} \Psi_S(-v) dv \int_{\mathbb{A}_S} h(u) \left(\log^+ \frac{|v|}{\max(1, \frac{1}{|u|})} + \log^- \frac{|v|}{\max(1, \frac{1}{|u|})} \right) \Psi_S(uv) du \\ &= \int_{|v|>1} \log |v| \Psi_S(-v) dv \int_{|u|\geq 1} h(u) \Psi_S(uv) du + \int_{\mathbb{A}_S} \Psi_S(-v) dv \int_{|vu|>1, |u|<1} h(u) \log |vu| \Psi_S(uv) du \\ &\quad + \int_{|v|<1} \log |v| \Psi_S(-v) dv \int_{|u|\geq 1} h(u) \Psi_S(uv) du + \int_{\mathbb{A}_S} \Psi_S(-v) dv \int_{|vu|<1, |u|<1} h(u) \log |vu| \Psi_S(uv) du. \end{aligned}$$

As

$$\begin{aligned} &\int_{\mathbb{A}_S} \Psi_S(-v) dv \int_{|vu|>1, |u|<1} h(u) \log |vu| \Psi_S(uv) du \\ &= \int_{\mathbb{A}_S} \Psi_S(-v) \frac{dv}{|v|} \int_{|u|>1, |u|<|v|} h(\frac{u}{v}) \log |u| \Psi_S(u) du \\ &= \int_{|u|>1} \log |u| \Psi_S(u) du \int_{|u|<|v|} \frac{1}{|v|} h(\frac{u}{v}) \Psi_S(-v) dv \\ &= \int_{|u|>1} \log |u| \Psi_S(u) du \int_{|v|>1} h(v) \Psi_S(-vu) dv \end{aligned}$$

where we used the fact that $|v|^{-1}h(v^{-1}) = h(v)$. Similarly we have

$$\begin{aligned} &\int_{\mathbb{A}_S} \Psi_S(-v) dv \int_{|vu|<1, |u|<1} h(u) \log |vu| \Psi_S(uv) du \\ &= \int_{|u|<1} \log |u| \Psi_S(u) du \int_{|v|>1} h(v) \Psi_S(-vu) dv. \end{aligned}$$

Therefore,

$$\begin{aligned} \Delta(h) &= \int_{|v|>1} \log |v| \Psi_S(-v) dv \int_{|u|\geq 1} h(u) \Psi_S(uv) du + \int_{|v|>1} \log |v| \Psi_S(-v) dv \int_{|u|>1} h(u) \Psi_S(uv) du \\ &+ \int_{|v|<1} \log |v| \Psi_S(-v) dv \int_{|u|\geq 1} h(u) \Psi_S(uv) du + \int_{|v|<1} \log |v| \Psi_S(-v) dv \int_{|u|>1} h(u) \Psi_S(uv) du. \end{aligned}$$

By Theorem 1.4,

$$\begin{aligned} \text{trace}_{L^2(C_S)} \{V(h)T\} &= \int_{\mathbb{A}_S, |v|>1} \Psi_S(v) \log |v| dv \int_{\mathbb{A}_S, |u|\geq 1} h(u) \Psi_S(-uv) du \\ &+ \int_{|v|>1} \Psi_S(v) \log |v| dv \int_{|u|>1} h(u) \Psi_S(-uv) du. \end{aligned}$$

The stated identity follows from the above two identities.

This completes the proof of Theorem 1.6. \square

REFERENCES

1. E. Bombieri, *Remarks on Weil's quadratic functional in the theory of prime numbers, I*, Rend. Mat. Acc. Lincei, s. 9, v. 11 (2000), 183–233.
2. C. Brislawn, *Traceable integral kernels on countably generalized measure spaces*, Pacific J. Math. **150** (1991), 229–240.
3. A. Connes, *Trace formula in noncommutative geometry and the zeros of the Riemann zeta function*, Selecta Math. **5** (1999), 29–106.
4. I. C. Gohberg and M. G. Krein, *Introduction to the Theory of Linear Nonselfadjoint Operators in Hilbert Space*, Translations of Mathematical Monographs, Vol. 18, Amer. Math. Soc., Providence, R. I., 1969.
5. Xian-Jin Li, *On the explicit formula related to Riemann's zeta-function*, Int. J. Number Theory **11** (2015), 2451–2486.
6. A. Mattuck and J. Tate, *On the inequality of Castelnuovo-Severi*, Abh. Math. Sem. Univ. Hamburg **22** (1958), 295–299.
7. R. Meyer, *On a representation of the idele class group related to primes and zeros of L-functions*, Duke Math. J. **127** (2005), 519–595.
8. M. Reed and B. Simon, *Methods of Modern Mathematical Physics. I: Functional Analysis*, Academic Press, New York, 1980.
9. B. Riemann, *Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse*, in “Bernhard Riemann, Mathematische Werke”, Dover, New York, 1953, 145–153.
10. J. T. Tate, *Fourier analysis in number fields and Hecke's zeta-functions*, in “Algebraic Number Theory,” Edited by J.W.S. Cassels and A. Fröhlich, Academic Press, New York, 1967, 305–347.
11. E. C. Titchmarsh, *The Theory of Functions*, Second Edition, Oxford, 1939.
12. E. C. Titchmarsh, *Introduction to the Theory of Fourier Integrals*, Second Edition, Oxford, 1967.
13. A. Weil, *Sur les formules explicites de la théorie des nombres*, Izv. Akad. Nauk SSSR Ser. Mat. **36** (1972), 3–18.

DEPARTMENT OF MATHEMATICS, BRIGHAM YOUNG UNIVERSITY, PROVO, UTAH 84602, USA
E-mail address: xianjin@mathematics.byu.edu