Characterization of Isometric Words based on Swap and Mismatch Distance

M. Anselmo ¹	G. Castiglione ²	M. Flores ²
D. Giammarresi ³	M. Madonia ⁴	S. Mantaci ²

¹ Dipartimento di Informatica, Università di Salerno, Italy. {manselmo}@unisa.it

² Dipartimento di Matematica e Informatica, Università di Palermo, Italy

³ Dipartimento di Matematica. Università Roma "Tor Vergata" Italy.

giammarr@mat.uniroma2.it

⁴Dipartimento di Matematica eInformatica, Università di Catania, Italy. madonia@dmi.unict.it

Abstract

In this paper we consider an edit distance with swap and mismatch operations, called *tilde-distance*, and introduce the corresponding definition of *tilde-isometric* word. Isometric words are classically defined with respect to Hamming distance and combine the notion of edit distance with the property that a word does not appear as factor in other words. A word f is said tilde-isometric if, for any pair of f-free words u and v, there exists a transformation from u to v via the related edit operations such that all the intermediate words are also f-free. This new setting is here studied giving a full characterization of the tilde-isometric words in terms of overlaps with errors.

Keywords: Swap and mismatch distance; Isometric words; Overlap with errors.

1 Introduction

The notion of edit distance plays a crucial role in defining combinatorial properties of families of words as well as in designing many classical string algorithms that find applications in natural language processing, bioinformatics and, in general, in information retrieval problems. The edit distance is a string metric that quantifies how much two strings differ from each other and it is based on counting the minimum number of edit operations required to transform one string into the other one.

Changing the set of the operations will give rise to a different definition of edit distance. The operations of insertion, deletion and replacement of a character in the string characterize the Levenshtein distance which is probably the most widely known (cf. [26]). On the other hand, the most basic one is the Hamming distance which applies only to pair of strings having equal length and counts the positions where they have a mismatch; this corresponds to the restriction of using only the replacement operation. The Hamming distance finds a direct application in detecting and correcting errors in

strings and it plays a main role in the algorithms for string matching with mismatches (see [21]).

The notion of isometric word combines the notion of edit distance with the property that a word does not appear as factor in other words. Note that this property is important in combinatorics as well as in the investigation on similarities of DNA sequences, where the avoided factor is referred to as an absent word [12, 14, 15, 18].

Isometric words based on Hamming distance were first introduced in [24] as special binary strings that never appear as factors in some string transformations. A string is f-free if it does not contain f as factor. A word f is isometric if for any pair of ffree words u and v, there exists a sequence of symbol replacement operations that transforms u into v, where all the intermediate words are also f-free. In this original definition, isometric words are strictly related to and motivated by the special isometric subgraphs of the hypercubes, called generalized Fibonacci cubes. The hypercube Q_n is a graph whose vertices are the (binary) words of length n, and two vertices are adjacent when the corresponding words differ in exactly one symbol. Therefore, the distance between two vertices in Q_n is the Hamming distance of the corresponding vertex-words. The generalized Fibonacci cube $Q_n(f)$ is the subgraph of Q_n which contains only vertices that are f-free where f is an isometric word. This implies that the distances of the vertices in $Q_n(f)$ are the same as calculated in the whole Q_n , that is $Q_n(f)$ is isometric to Q_n . Fibonacci cubes have been introduced by Hsu in [22] and correspond to the case with f = 11. In [24, 25, 29, 30, 31] the structure of non-isometric words for alphabets of size 2 and Hamming distance is completely characterized and related to particular properties on their overlaps. The more general case of alphabets of size greater than 2 and Lee distance is studied in [5, 7, 6, 9, 8]. Using these characterizations, in [11] some linear-time algorithms are given to check whether a binary word is Hamming isometric and, for quaternary words, if it is Lee isometric. These algorithms were extended to provide further information on nonisometric words, still keeping linear complexity in [6]. Binary Hamming isometric two-dimensional words have been also studied in [10].

Many challenging problems in correcting errors in strings come from computational biology. Among the chromosomal operations on DNA sequences, in gene mutations and duplication, it seems natural to consider the *swap* operation, consisting in exchanging two adjacent symbols. The Damerau-Levenshtein distance adds also the swap to all edit operations. In [28], Wagner proves that computing the edit distance with insertion, deletion, replacement, and swap, is polynomially solvable in some restriction of the problem. The swap-matching problem has been considered in [1, 20, 19, 4, 3, 13], and related algorithms are given in [2, 16].

In this paper, we study the notion of binary isometric word using the edit distance based on swaps and mismatches. This distance will be here called *tilde-distance* by using the \sim symbol that somehow evokes the swap operation. The tilde-distance dist_{\sim}(u, v) of equal-length words u and v is the minimum number of replacement and swap operations needed to transform u into v.

It turns out that the addition of the swap operation to the definition makes the situation more complex, although interesting for applications. It is not a mere generalization of the Hamming case since special situations arise. A swap operation in fact is equivalent to two replacements, but it counts as one when computing the tilde-distance. Moreover, there could be different ways to transform u into v since particular triples of consecutive symbols can be managed, from left to right, either by a swap and then a replacement or by a replacement and then a swap. Furthermore there is also the possibility of making two swaps in consecutive positions that are in fact equivalent to two replacement operations; in this case there is a position whose symbol is changed twice. This fact is the major difference with the Hamming distance where all the transformations from u to v may differ only in the order in which the replacement operations are applyed to the positions to be changed. This difference will spawn new scenario when dealing with tilde-isometricity.

The definition of *tilde-isometric* word comes in a very natural way. A word f is tilde-isometric if for any pair of equal-length words u and v that are f-free, there is a tilde-transformation from u to v that uses exactly dist_{\sim}(u, v) replacement and swap operations and such that all the intermediate words still avoid f. As expected, the notions is not a mere generalization of the classic case. In fact we exibit some examples of tilde-isometric words that are not Hamming isometric and vice versa.

The main result of this paper consists in a complete characterization of the words that are not tilde-isometric in terms of special configurations in their overlap. Note that, in order to prove that a given string f is not tilde-isometric one should exhibit a pair of f-free words $(\tilde{\alpha}, \tilde{\beta})$ such that any tilde-transformation from $\tilde{\alpha}$ to $\tilde{\beta}$ of length dist_~ $(\tilde{\alpha}, \tilde{\beta})$ comes through words that contain f. Such a pair is called pair of *tildewitnesses* for f. The proof or our main theorem gives also an explicit construction of the tilde-witnesses in all possible cases.

The paper is organized as follows. In a short preliminary section we first fix some basic notations on words used through the paper and then we report known results on (Hamming) isometric words. Then two main sections follow. The first one starts by introducing the tilde-distance together with examples and the notion of tilde-transformations. This leads to the formal definition of tilde-isometric words and some examples of them. Then, we study the overlap with tilde-errors and finally we state our main result that consists in the fully characterization of tilde-isometric words in terms of special properties of their overlaps. The last section is completely dedicated to the proof of our main characterization theorem. The two directions of the proofs are given in two separate subsections respectively and are preceded by some technical lemmas. Finally, a partial and preliminary version of the results in this paper can be found in [4].

2 Preliminaries

Let Σ be a finite alphabet. A word (or string) w of length |w| = n, is $w = a_1 a_2 \cdots a_n$, where a_1, a_2, \ldots, a_n are symbols in Σ . The set of all words over Σ is denoted Σ^* and the set of all words over Σ of length n is denoted Σ^n . Finally, ϵ denotes the *empty* word and $\Sigma^+ = \Sigma^* - \{\epsilon\}$. For any word $w = a_1 a_2 \cdots a_n$, the *reverse* of w is the word $w^{rev} = a_n a_{n-1} \cdots a_1$. If $x \in \{0, 1\}$, we denote by \overline{x} the opposite of x, i.e $\overline{x} = 1$ if x = 0 and vice versa. Then we define *complement* of w the word $\overline{w} = \overline{a_1}\overline{a_2} \cdots \overline{a_n}$.

Let w[i] denote the symbol of w in position i, i.e. $w[i] = a_i$. If $f = w[i..j] = a_i \cdots a_j$, for $1 \le i \le j \le n$, we say that f is a *factor* of w that occurs in the interval [i..j], or equivalently, at position i. The *prefix* (resp. *suffix*) of w of length l, with $1 \le \ell \le n - 1$ is $\operatorname{pre}_{\ell}(w) = w[1..\ell]$ (resp. $\operatorname{suf}_{\ell}(w) = w[n - \ell + 1..n]$). When $\operatorname{pre}_{\ell}(w) = \operatorname{suf}_{\ell}(w) = u$ then u is here referred to as an *overlap* of w of length ℓ ; it is also called border, or bifix. A word w is said f-free if w does not contain f as a factor.

An *edit operation* is a function $O : \Sigma^* \to \Sigma^*$ that transforms a word into another one. Among the most common edit operations there are the insertion, the deletion or the replacement of a character and the swap of two adjacent characters. Let \mathcal{O} be a *set* of edit operations. The edit distance of two words $u, v \in \Sigma^*$ is the minimum number of edit operations in \mathcal{O} needed to transform u into v.

A well-known edit distance is the *Hamming distance* which is defined over a binary alphabet and uses only the replacement operation. Based on the Hamming distance, the definition of *good* words is introduced in [23] and

put in connection with isometric subgraphs of the hypercubes in [25]. In this paper we refer to this definition of isometric as *Ham-isometric*. More specifically, a word f is *Ham-isometric* if for any pair of f-free words u and v, there exists a sequence of replacement operations that applied to u transform u into v keeping all the intermediate words f-free. In [30], it is given a characterization of Ham-isometric words based on a property of their overlaps as follows. Here, a word w has a 2-error overlap if there exists $1 < \ell < |w|$ such that $\operatorname{pre}_{\ell}(w)$ and $\operatorname{suf}_{\ell}(w)$ differ in exactly 2 positions (i.e., they have Hamming distance equal to 2).

Proposition 1 A word f is not Ham-isometric if and only if f has a 2-error overlap.

For example the binary word 111000 is not Ham-isometric (take the prefix 11 and the suffix 00) while it is easy to verify that the word 1010 does not have any 2-error overlap and therefore it is Ham-isometric.

3 Tilde-distance and Tilde-isometric words

In this section, we introduce the tilde-distance as the edit distance based on replacements and swaps. Then, using the tilde-distance, we define the tilde-isometric words as a generalization of the (classical) Ham-isometric words. Very interesting situations and examples arise since the possibility of applying swap operations makes very different scenarios. At the end of the section we state the main theorem of the paper that provides a complete characterization of tilde-isometric words in terms of special blocks contained in their overlaps.

3.1 Tilde-distance and tilde-transformations

We start by formally introducing the operations of replacements and swaps; they are the base for the definition of tilde-distance. Then, we point out some interesting facts regarding the tilde-transformations between words considered in the definition of distance.

Definition 2 Let Σ be a finite alphabet and $w = a_1 a_2 \dots a_n$ a word over Σ . The replacement operation (or replacement, for short) on w at position i with $x \in \Sigma$, $x \neq a_i$, is defined by

 $R_{i,x}(a_1a_2...a_{i-1}a_ia_{i+1}...a_n) = a_1a_2...a_{i-1}xa_{i+1}...a_n.$

The swap operation (or swap, for short) on w at position i consists in exchanging characters at positions i and i + 1, provided that they are different, $a_i \neq a_{i+1}$,

$$S_i(a_1a_2\ldots a_ia_{i+1}\ldots a_n) = a_1a_2\ldots a_{i+1}a_i\ldots a_n.$$

When the alphabet $\Sigma = \{0, 1\}$ there is only one possible replacement at a given position *i*, so we write $R_i(w)$ instead of $R_{i,x}(w)$. In the following we always assume $\Sigma = \{0, 1\}$.

Remark that a single swap operation is a sort of shortcut since it has the same effect of two replacements on two adjacent positions respectively. Moreover note that both replacement and swap operations do not change the length of the word on which they are applied.

The edit distance based on swap and replacement operations will be called *tilde-distance* and denoted $dist_{\sim}$.

Definition 3 Let $u, v \in \Sigma^n$. The tilde-distance $dist_{\sim}(u, v)$ between u and v is the minimum number of replacements and swaps needed to transform u into v.

In this paper, we focus on the sequences of operations that transform a word into another one. We give first a formal definition of a transformation in the setting of the tilde-distance.

Definition 4 Let $u, v \in \Sigma^n$ be words of equal length and $dist_{\sim}(u, v) = d$. A tildetransformation τ from u to v is a sequence of d + 1 words (w_0, w_1, \ldots, w_d) such that $w_0 = u$, $w_d = v$, and for any $k = 0, 1, \ldots, d - 1$, $dist_{\sim}(w_k, w_{k+1}) = 1$. Moreover, let $f \in \Sigma^+$, τ is f-free if for any $i = 0, 1, \ldots, d$, the word w_i is f-free.

A tilde-transformation (w_0, w_1, \ldots, w_d) from u to v is associated to a sequence of d operations $(O_{i_1}, O_{i_2}, \ldots, O_{i_d})$ such that, for any $k = 1, \ldots, d, O_{i_k} \in \{R_{i_k}, S_{i_k}\}$ and $w_k = O_{i_k}(w_{k-1})$; it can be represented as follows:

$$u = w_0 \xrightarrow{O_{i_1}} w_1 \xrightarrow{O_{i_2}} \cdots \xrightarrow{O_{i_d}} w_d = v.$$

Therefore, an equivalent way to refer to a tilde-transformation is to give the sequence $(O_{i_1}, O_{i_2}, \ldots, O_{i_d})$ of its operations. When considering a tilde-transformation of u into v, it is helpful to easily detect the positions where they differ. For this, it is useful to write u and v aligned one above the other. So, in what follows, $\binom{u}{v}$ will denote an alignment of two words u and v of equal length n, meaning that for all $1 \le k \le n$, u[k] is aligned on v[k]. When an alignment $\binom{u}{v}$ is considered, a factor of length m of the alignment, denoted by $\binom{x}{y}$, is called a *block* of $\binom{u}{v}$ and we say that it occurs in position i if x, y occur in i as factors of u and v, respectively. For instance, the block $\binom{100}{011}$ occurs at position 3 in $\binom{001001}{010110}$.

Observe that the number of the operations in a tilde-transformation should correspond to the distance between the two words, but we do not put any restrictions on the kind of operations involved. In particular we also allow a position i to be changed twice by a pair of swaps S_{i-1} and S_i , for some position i (note that this was not permitted in the previous paper [4]). Further, observe that a sequence of swaps at consecutive positions $S_i, S_{i+1}, \ldots, S_{i+k}$, with $k \ge 2$, cannot be part of a tilde-transformation since the same result can be obtained by two replacements R_i and R_{i+k+1} in a shorter sequence of operations.

Another important issue is that, given two words u and v, there are many tildetransformations between them; such transformations may differ in the type of operations we choose and also in the order in which we perform such operations. Pay attention to the fact that you cannot always change the order of swaps. This is a major difference with the case of Hamming distance where only replacements are allowed and therefore all possible transformations can be obtained by exchanging the order of the operation application. The following significant cases explain this fact. **Example 5** Let $\binom{u}{v} = \binom{101}{010}$. Then (S_1, R_3) , (R_3, S_1) , (R_1, S_2) and (S_2, R_1) are all the different tilde-transformations from u to v.

Example 6 Let $\binom{u}{v} = \binom{100}{001}$. Then (R_1, R_3) (R_3, R_1) and (S_1, S_2) are three tildetransformations from u to v. Note that the operations S_1 and S_2 do not commute, since S_2 cannot be executed before S_1 . Therefore the pair (u, v) admits three different tilde-transformations. Note also that tilde-transformation (S_1, S_2) flips twice the bit in position 2.

Define the sets of blocks $\mathcal{B}_0 = \{\binom{10}{01}, \binom{01}{10}\}$, $\mathcal{B}_1 = \{\binom{101}{010}, \binom{010}{101}\}$ and $\mathcal{B}_2 = \{\binom{100}{001}, \binom{110}{011}, \binom{001}{100}, \binom{011}{100}\}$, obtained by applying complement, reverse and exchange of rows to blocks $\binom{10}{01}$, $\binom{101}{010}$ and $\binom{100}{001}$, respectively. In what follows we will only consider one block in \mathcal{B}_0 , \mathcal{B}_1 or in \mathcal{B}_2 , respectively, as representative of its class, since the results on the other blocks can be easily inferred from them.

Remark 7 Let $u, v \in \Sigma^*$. As a generalization of Example 5, note that, if u[i] = u[i + 2], each time R_i and S_{i+1} belong to a tilde-transformation τ , the sequence obtained from τ by substituting R_i and S_{i+1} with S_i and R_{i+2} is still a tilde-transformation, and vice versa. Similarly, as a generalization of Example 6, the operations S_i and S_{i+1} can be substituted with R_i and R_{i+2} in a tilde-transformation; the vice versa is possible only when the block occurring in $\binom{u}{v}$ at position *i* belongs to \mathcal{B}_2 . In particular, when dist_{\sim}(u, v) = 2, exactly two tilde-transformations from u to v exist, except in those cases in which a block in \mathcal{B}_1 occurs in $\binom{u}{v}$ yielding four tilde-transformations from u to v (cf. Example 5), and when a block in \mathcal{B}_2 occurs in $\binom{u}{v}$ and produces three tilde-transformations from u to v (cf. Example 6).

3.2 Tilde-isometric words

Tilde-transformations between two words play an important role in the definition of tilde-isometric words, here defined in analogy with the Ham-isometric words, but on the base on the swap and mismatch distance. We give here the formal definition.

Definition 8 Let $f \in \Sigma^n$, with $n \ge 1$, f is tilde-isometric if for any pair of f-free words u and v of length $m \ge n$, there exists a tilde-transformation from u to v that is f-free. It is tilde-non-isometric if it is not tilde-isometric.

Note that, in order to prove that a word is tilde-non-isometric, it is sufficient to exhibit a pair (u, v) of words contradicting Definition 8, that we call *pair of tilde-witnesses* for f. More formally we give the following definition.

Definition 9 A pair (u, v) of words in Σ^m is a pair of tilde-witnesses for f if:

- 1. dist_~ $(u, v) \ge 2$
- 2. u and v are f-free
- 3. there exists no f-free tilde-transformation from u to v.

Example 10 Let f = 1010, then f is tilde-non-isometric because the pair (u, v) = (11000, 10110) is a pair of tilde-witnesses for f. In fact, observe that there are two tilde-transformations from u to v, namely:

 $11000 \xrightarrow{S_2} 10100 \xrightarrow{R_4} 10110 \qquad 11000 \xrightarrow{R_4} 11010 \xrightarrow{S_2} 10110.$

In both tilde-transformations 1010 appears as factor after the first step, as evidenced by the underlined characters, i.e. they are not f-free.

Example 11 Let f = 100011 and consider the words u and v given by $\binom{u}{v} = \binom{100101011}{100010011}$. Four tilde-transformations, exist from u to v, namely:

 $100101011 \xrightarrow{R_4} 100001011 \xrightarrow{S_5} 100010011; 100101011 \xrightarrow{S_5} 100110011 \xrightarrow{R_4} 100010011,$

 $100101011 \xrightarrow{S_4} \underline{100011}011 \xrightarrow{R_6} 100010011; 100101011 \xrightarrow{R_6} 100\underline{100011} \xrightarrow{S_4} 100010011.$

Note that the first and the second tilde-transformations are *f*-free whereas the third and the fourth are not. This proves that (u, v) is not a pair of tilde-witnesses. Note that $\binom{u}{v}$ contains the block $\binom{101}{010}$ at position 4.

As a matter of fact, a pair of tilde-witnesses for the word f = 100011 of previous example does not exist, i.e. f is tilde-isometric. This can be proved by exploiting the the main theorem of this paper that consists of a characterization of tilde-isometric words in terms of a property of their overlaps. From now on, we study isometric binary words starting with 1, in view of the following lemma whose proof can be easily inferred by the definition.

Lemma 12 Let $f \in \Sigma^n$. The following statements are equivalent:

- 1. f is tilde-isometric
- 2. f^{rev} is tilde-isometric
- 3. \overline{f} is tilde-isometric.

Although the tilde-distance is more general than the Hamming distance, the sets of Ham-isometric and tilde-isometric words are incomparable, as stated in the following proposition.

Proposition 13 There exists a word which is tilde-isometric but Ham-non-isometric, and a word which is tilde-non-isometric, but Ham-isometric.

Proof: The word f = 111000 is tilde-isometric (cf. Example 4), but f is Ham-non-isometric. In fact, f has a 2-tilde-error overlap (with respect to the Hamming distance) with shift 4 and then it is Ham-non-isometric by Proposition 1.

Conversely, f' = 1010 is tilde-non-isometric (see Example 10), but Ham-isometric by Proposition 1. \Box

3.3 Tilde-error overlaps

The characterization of Ham-isometric words given in [30] and here reported as Proposition 1, uses the notion of 2-error overlap. In this section we introduce the corresponding definition that refers to the tilde-distance. Tilde-error overlaps will have a main role in the characterization of tilde-isometric words but the presence of swap operations will force us to handle them with care.

Definition 14 Let $f \in \Sigma^n$. Then, f has a q-tilde-error overlap of length ℓ and shift $r = n - \ell$, with $1 \le \ell \le n - 1$ and $0 \le q \le \ell$, if $\operatorname{dist}_{\sim}(\operatorname{pre}_{\ell}(f), \operatorname{suf}_{\ell}(f)) = q$.

Example 15 The word f = 1101110101101 has a 2-tilde-error overlap of length 6 and shift 7. Indeed, $\text{pre}_6(f) = 110111$, $\text{suf}_6(f) = 101101$ and $\text{dist}_{\sim}(110111, 101101) = 2$.

For our proofs, given a word f with a q-tilde-error overlap of length ℓ , we will study the tilde-transformations τ from $\operatorname{pre}_{\ell}(f)$ to $\operatorname{suf}_{\ell}(f)$ with q operations. For this reason we need to refer to the alignment of the two strings $\operatorname{pre}_{\ell}(f)$ to $\operatorname{suf}_{\ell}(f)$. Furthermore, in some technical proofs it will sometimes be relevant to consider the bits adjacent to a tilde-error overlap of a word. For this reason we introduce the following notation. Let f be a word in $\{0,1\}^*$ and \$ be a symbol different from 0, 1, here used as delimiter of a word that "matches" any symbol of the word. Consider f with its delimiters \$f\$. A q-tilde-error overlap of length ℓ is denoted by $\binom{\$xa}{by\$}$ where xa, by are a prefix and a suffix, respectively, of f, $a, b \in \Sigma$, $x, y \in \Sigma^*$ with $|x| = |y| = \ell$, and $\operatorname{dist}_{\sim}(\$xa, by\$) = \operatorname{dist}_{\sim}(x, y) = q$. This notation makes evident the fact that in f the prefix x is followed by a and the suffix y is preceded by $\binom{\$1}{01}\binom{1011}{0110}\binom{10}{0\$}$ because $\operatorname{dist}_{\sim}(110111, 101101) = 2 = \operatorname{dist}_{\sim}(1011, 0110)$.



Figure 1: A word f and its 2-tilde-error overlap of shift r and length $\ell = n - r$, with tilde-transformation $(O_i, O_j) = (R_i, R_j)$ (left), and $(O_i, O_j) = (S_i, R_j)$ (right)

In the sequel, we will be interested in the specific case of 1-tilde-error overlap where the single error in the alignment is a swap and in all the cases of 2-tilde-error overlaps. Consider a 2-tilde-error overlap of f of shift r, length $\ell = n - r$, and let $(O_i, O_j), 1 \le i < j \le \ell$, be a tilde-transformation from $\operatorname{pre}_{\ell}(f)$ to $\operatorname{suf}_{\ell}(f)$. Observe that the positions in $\operatorname{pre}_{\ell}(f)$ modified by O_i are either i or both i and i + 1, following that O_i is a replacement or a swap. Hence, the positions modified by O_i and O_j may be 2, 3 or 4. Fig. 1 shows a word f with its 2-tilde-error overlap of shift r and length $\ell = n - r$. With our notation, the 2-tilde-error overlap is $\binom{\$w_21w1w_3a}{bw_20w0w_3\$}$ in the figure on the left and $\binom{\$w_210w1w_3a}{bw_201w0w_3\$}$ in the figure on the right, where a is the last letter of w_1 and b the first letter of w_4 . A tilde-transformation from $\operatorname{pre}_{\ell}(f)$ to $\operatorname{suf}_{\ell}(f)$ is given by $(O_i, O_j) = (R_i, R_j)$ in the first case and by $(O_i, O_j) = (S_i, R_j)$ in the second case. We say that a 2-tilde-error overlap has *non-adjacent errors* when there is at least one character interleaving the positions modified by O_i and those modified by O_j .

The 2-tilde-error overlap $\binom{\$x}{ax}\binom{100}{001}\binom{yb}{y\$}$ is also considered as having non-adjacent errors because it admits the tilde-transformation $(O_i, O_j) = (R_i, R_{i+2})$, despite it has also the other $(O_i, O_j) = (S_i, S_{i+1})$.

In all the other cases, we say that the 2-tilde-error overlap has adjacent errors.

Characterization of tilde-isometric words 4

Let us state the main result of the paper that consists in the characterization of tildeisometric words in terms of special configurations in their overlap. The proof of the theorem is quite complicate and is given separately in the next section.

Recall that in this paper a block in \mathcal{B}_0 , \mathcal{B}_1 or \mathcal{B}_2 is taken as representative of its class, since the results on the other blocks can easily be inferred from it.

Theorem 16 A word $f \in \Sigma^n$ is tilde-non-isometric if and only if one of the following cases occurs (up to complement, reverse and inversion of rows):

- (C0) f has a 1-tilde-error overlap $\binom{\$x}{ax}\binom{01}{10}\binom{yb}{y\$}$ with $x, y \in \Sigma^*$, $a, b \in \Sigma$
- (C1) f has a 2-tilde-error overlap with non-adjacent errors, different from $\binom{\$x}{ax}\binom{000}{101}\binom{yb}{y\$}$ with $x, y \in \Sigma^+$, $a, b \in \Sigma$
- (C2) f has a 2-tilde-error overlap $\binom{\$x}{ax}\binom{0101}{1010}\binom{yb}{y\$}$ or $\binom{\$x}{ax}\binom{0110}{1001}\binom{yb}{y\$}$ with $x, y \in \Sigma^*$, $a, b \in \Sigma$
- (C3) f has a 2-tilde-error overlap $\binom{\$x}{ax}\binom{010}{101}\binom{yb}{y\$}$ with $x, y \in \Sigma^*$, $a, b \in \Sigma$
- (C4) f has a 2-tilde-error overlap $\binom{\$x}{ax}\binom{011}{100}\binom{0}{\$}$ with $x \in \Sigma^*$, $a \in \Sigma$
- (C5) f has a 2-tilde-error overlap $\binom{\$}{0}\binom{00}{11}\binom{1}{\$}$

As an application of Theorem 16 let us show the following example of a family of tilde-isometric words.

Example 17 All the words $f = 1^n 0^m$ for n, m > 2 are tilde-isometric. In fact, for $n, m > 2, f = 1^n 0^m$ has only two 2-tilde-error overlaps and none of them fall into a case in the statement of Theorem 16. The first one is the tilde-error overlap with shift 2, $\binom{\$1^{n-2}}{11^{n-2}}\binom{11}{00}\binom{0^{n-2}0}{0^{n-2}\$}$

the other one has shift n + m - 2, and it is $\binom{\$}{0}\binom{11}{00}\binom{1}{\$}$. Note that if n = m = 2, f = 1100 is tilde-non-isometric. In fact one can verify that the pair of words $\binom{u}{v} = \binom{110100}{101010}$ is a pair of tilde-witnesses for f. As expected from Theorem 16, f has only one 2-tilde-error overlap $\binom{\$}{1}\binom{11}{00}\binom{0}{\$}$, corresponding to case (C5). Note also that words $f = 1^n 0^m$ with $n, m \ge 2$ are Hamming non-isometric.

The following are two examples of words with a 2-tilde-error overlap with adjacent errors. The first one is tilde-isometric, the second one is tilde-non-isometric.

Example 18 The word f = 010110000 is tilde-isometric; indeed its unique 2-tildeerror overlap has shift 5 and length 4, $\binom{\$0}{10}\binom{101}{000}\binom{1}{\$}$. Note this is the case of nonadjacent errors but of the type prohibited by condition in (C1).

Example 19 The word f = 1011000 is tilde-non-isometric; indeed it has the 2-tildeerror overlap, of shift 4 and length 3, $\binom{\$}{1}\binom{101}{000}\binom{1}{\$}$, that verifies (C1). Note that the pair (u, v) = (10110011000, 10101001000) is a pair of tilde-witnesses for f.

5 The proof of the Characterization Theorem

This last section is very technical and contains the proof of Theorem 16. The cases (C0) up to (C5) list all the possible configurations we can find in an overlap of a tildenon-isometric word. The proof walks carefully through all these cases to show that they cover all possible situations. The two implications of the theorem are proved separately in two corresponding subsections and are preceded by some technical lemmas.

5.1 Properties of tilde-witnesses with minimal distance

When a word f is tilde-non-isometric then there exists a pair of tilde-witnesses. We focus our attention on a pair (u, v) of tilde-witnesses at minimal distance among all such pairs of a given length. With this constraint, any choice of the starting operation O_i in a tilde-transformation from u to v, will cause an occurrence of the factor f in $O_i(u)$. More specifically, this occurrence of f will cover at least one position modified by the operation O_i (i.e. position i and/or i + 1 in the case of a swap S_i).

We start with two technical lemmas.

Lemma 20 Let $f \in \Sigma^n$ be a tilde-non-isometric word and (u, v), with $u, v \in \Sigma^m$, be a pair of tilde-witnesses with minimal distance dist_~(u, v) among all pairs of tildewitnesses of length m. Let $\mathcal{O} = \{O_{i_1}, O_{i_2}, \ldots, O_{i_d}\}$ be the set of the operations in a tilde-transformation from u to v, that does not contain two swaps at two consecutive positions. Then,

- 1. for any $i_j \in \{i_1, i_2, \ldots, i_d\}$, f is a factor of $O_{i_j}(u)$. Let k_{i_j} denote the starting position of such occurrence of f in $O_{i_j}(u)$, and $I_{i_j} = [k_{i_j} \ldots k_i + |f| 1]$ the interval where f occurs in $O_{i_j}(u)$.
- 2. there exist $s, t \in \{i_1, i_2, ..., i_d\}$, with s < t, such that I_s and I_t share at least one position modified by O_s and at least one position modified by O_t . Moreover, without loss of generality, $k_s < k_t$.

Proof: Let $\mathcal{O} = \{O_{i_1}, O_{i_2}, \dots, O_{i_d}\}$ with $O_{i_j} \in \{R_{i_j}, S_{i_j}\}$ for any $j = 1, 2, \dots, d$, and let $1 \leq i_1 < i_2 < \dots < i_d \leq m$. Since the tilde-transformation does not contain two consecutive swaps, each O_{i_j} can be applied to u.

If, for some $j \in \{1, ..., m\}$, $O_{i_j}(u)$ were *f*-free, then the pair $(O_{i_j}(u), v)$ would still be a pair of tilde-witnesses of length *m*, with $dist_{\sim}(O_{i_j}(u), v) < d$, against the hypothesis that (u, v) are the tilde-witnesses of minimal distance. This proves statement (1).

Let I_{i_j} be the interval where f occurs in $O_{i_j}(u)$. This interval contains at least one position modified by O_{i_j} , because u is f-free; let o(j) denote the smallest position in I_{i_j} modified by O_{i_j} . Moreover, this occurrence of f must disappear in a tilde-transformation from u to v, because v is f-free. Hence, I_{i_j} contains a position modified by another operation in \mathcal{O} ; let p(j) denote the smallest such position. Overall, there must exist $s, t \in \{i_1, i_2, \ldots i_d\}$, such that I_s contains at least one position modified by O_t and I_t contains at least one position modified by O_s . To prove this, consider for any $j = 1, \ldots, d$, the interval I_{i_j} and the two positions o(j) and p(j) in I_{i_j} . Note that p(1) > o(1), whereas p(d) < o(d). Let i_k be the smallest position in $\{i_1, i_2, \ldots i_d\}$, such that p(k) < o(k); then p(k-1) > o(k-1). Since I_{i_k} contains o(k) and p(k), with p(k) < o(k), then I_{i_k} also contains o(k-1) because $p(k) \leq o(k-1)$. Then, I_{i_k} and $I_{i_{k-1}}$ both contain o(k-1) and o(k) and can play the role of I_s and I_t .



Figure 2: The representation of the three occurrences f^1 , f^2 , and f^3 of f in $O^1(u)$, $O^2(u)$ and $O^3(u)$, respectively, when u[s..s+2] = 101.

Finally, without loss of generality we can suppose that s < t and $k_s < k_t$. In fact, if this does not happen, exchange the roles of u and v.

Remark 21 Suppose that O satisfy the hypotheses of Lemma 20 and that contains a swap S_i . The swap operation S_i modifies two positions, i and i + 1, and interval I_i could contain just one of them. Then $i - |f| + 1 \le k_i \le i + 1$.

Lemma 22 Let f be a tilde-non-isometric word and let (u, v) be a pair of tildewitnesses for f of minimal distance dist $_{\sim}(u, v)$. Let O_s, O_t be as in Lemma 20, then

- If $(O_s = R_s \text{ and } O_t = S_{s+1})$ or $(O_s = S_s \text{ and } O_t = R_{s+2})$ then $\binom{u[s..s+2]}{v[s..s+2]} \neq \binom{101}{010};$
- If $O_s = R_s$ and $O_t = R_{s+2}$ then

$$\binom{u[s..s+2]}{v[s..s+2]} \neq \binom{100}{001}$$

Proof: First, consider the case that $O_s = R_s$, $O_t = S_{s+1}$. If $\binom{u[s..s+2]}{v[s..s+2]} = \binom{101}{010}$ we write $u = u_1 101u_2$ and $v = v_1 010v_2$, with $|u_1| = s - 1$.

Then we can obtain another tilde-transformation from u to v by replacing (R_s, S_{s+1}) with (S_s, R_{s+2}) (see Remark 7). By definition we have:

$$R_s(u_1\mathbf{101}u_2) = u_1\mathbf{001}u_2, \quad S_{s+1}(u_1\mathbf{101}u_2) = u_1\mathbf{110}u_2, \\ S_s(u_1\mathbf{101}u_2) = u_1\mathbf{011}u_2, \quad R_{s+2}(u_1\mathbf{101}u_2) = u_1\mathbf{100}u_2$$
(1)

The proof works considering only R_s , S_s and S_{s+1} . By Lemma 20, for each $O \in \{R_s, S_s, S_{s+1}\}$, O(u) has f as factor. Therefore, one can sort the operations in the set $\{R_s, S_s, S_{s+1}\}$ according to the increasing order of the positions where f occurs as factor in $R_s(u)$, $S_{s+1}(u)$ and $S_s(u)$, respectively. Hence, let O^1, O^2, O^3 be the sorted sequence of the operations, f^1, f^2, f^3 denote the relative occurrences of f at positions $k_1 < k_2 < k_3$ and I_1, I_2 and I_3 be the intervals of occurrence, respectively. Then, either $(O^3(u)[s+1] = a$ and $O^1(u)[s+1] = \overline{a})$ or $(O^3(u)[s+2] = a$ and $O^1(u)[s+2] = \overline{a}$, with $a \in \Sigma$. This fact is evident from Equation (1) where the bits s, s+1, s+2 are in bold. In what follows, without loss of generality, we can consider $k_1 = 1$ and denote $r = k_3 - 1$, $p = k_2 - 1$ and q = r - p.

We prove the statement in the case $O^1 = S_{s+1}$, $O^2 = R_s$ and $O^3 = S_s$. The proofs for all other cases apply the same technique. In such a case $O^3(u)[s+2] = 1$ and $O^1(u)[s+2] = 0$ (see Fig. 2). Note that I_2 contains s, s+1 and s+2. Indeed, by

Lemma 20, I_2 contains s and, at least, s + 1, but if I_2 does not contain s + 2 then I_1 does not contain s + 1 (because $k_1 < k_2$) and ends in s. This is not possible because u would contain f^1 , i.e. u is not f-free. For the same reason, $k_3 \le s + 1$ because u is f-free. Let i = s + 2 - r be the position of f^3 corresponding to position s + 2 of u (see Fig. 2); in other words, i - 2, i - 1 and i are the positions of f^3 corresponding to s, s + 1 and s + 2, respectively. Note that $f[i] = f^1[i] = f^2[i] = f^3[i] = 1$ and $f[r + i] = f^3[r + i] = f^1[r + i] = f^2[r + i] = 0$. Observe that $q \ne 2$, and $q \ne 1$, indeed, $f[i] = f^3[i] = 1$ and if either q = 1 or q = 0 then $f[i] = f^2[i] = 0$, a contradiction.

Since f^2 occurs in $R_s(u)$, f^2 matches the corresponding positions of u in I_2 , unless for the one where the replacement is applied, that is s. More precisely, for each $h \in I_2$ with $h \neq q + i - 2$ we have $u[p + h] = f^2[h]$.

It follows that $u[p+i] = f^2[i] = 1$, because $q \neq 2$.

Moreover, since f^1 occurs in $S_{s+1}(u)$, f^1 matches the corresponding positions of u in I_1 , except for the positions involved in the swap S_{s+1} , i.e. s + 1 and s + 2. More precisely, for each $h \in I_1$, with $h \neq r+i-1$ and $h \neq r+i$, we have $u[h] = f^1[h]$. It follows that $u[p+i] = f^1[p+i] = 1$, because q > 1.

Since f^3 occurs in $S_s(u)$, it matches the factor of u occurring at I_3 , except for the positions involved in the swap S_s , i.e. the ones corresponding to s and s + 1; in other words, $u[r+h] = f^3[h]$, for each $h \in I_3$, with $h \neq i-2$ and $h \neq i-1$.

It follows that $u[r + p + i] = f^3[p + i] = 1$ because p > 0. But $u[r + p + i] = f^2[r + i] = 0$ because p > 0. Then a contradiction follows.

The previous considerations can be extended to the other cases where $O^3(u)[s + 2] = a$ and $O^1(u)[s + 2] = \overline{a}$ getting the contradiction that u[r + p + i] = a and $u[r + p + i] = \overline{a}$, with i = s + 2 - r. On the other hand, if $O^3(u)[s + 1] = a$ and $O^1(u)[s + 1] = \overline{a}$ the contradiction that u[r + p + i] = a and $u[r + p + i] = \overline{a}$, can be similarly obtained for i = s + 1 - r.

Now, consider the case $O_s = R_s$ and $O_t = R_{s+2}$. If $\binom{u[s..s+2]}{v[s..s+2]} = \binom{100}{001}$ we write $u = u_1 100u_2$ and $v = v_1 001v_2$, with $|u_1| = s - 1$. Then we can can obtain another tilde-transformation from u to v by replacing (R_s, R_{s+2}) with (S_s, S_{s+1}) (see Remark 7). By definition:

$$R_s(u_1 \mathbf{100} u_2) = u_1 \mathbf{000} u_2, \quad R_{s+2}(u_1 \mathbf{100} u_2) = u_1 \mathbf{101} u_2, \\ S_s(u_1 \mathbf{100} u_2) = u_1 \mathbf{010} u_2,$$
(2)

One can sort the set of operations $\{R_s, R_{s+2}, S_s\}$ as before mentioned and denote by $k_1 < k_2 < k_3$ the start positions of f^1 , f^2 and f^3 , and I_1 , I_2 and I_3 the intervals of occurrence, respectively. From Equation (2), observe that either $(O^3(u)[s+1] = a$ and $O^1(u)[s+1] = \overline{a}$) or $(O^3(u)[s+2] = a$ and $O^1(u)[s+2] = \overline{a}$). Without loss of generality we can consider $k_1 = 1$, $r = k_3 - 1$, $p = k_2 - 1$ and q = r - p. We prove the statement for the order $O^1 = R_{s+2}$, $O^2 = S_s$ and $O^3 = R_s$ (see Fig.

We prove the statement for the order $O^1 = R_{s+2}$, $O^2 = S_s$ and $O^3 = R_s$ (see Fig. 3). Note that $O^3(u)[s+1] = 1$ and $O^1(u)[s+1] = 0$. By Lemma 20, both I_1 and I_3 contain both s and s+2. Therefore I_2 must contain s, s+1 and s+2. Let i = s+2-r, $f[i] = f^1[i] = 0 = f^2[i] = f^3[i]$ and $f[r+i] = f^3[r+i] = 1 = f^1[r+i] = f^2[r+i]$. First, let us remark that $q \neq 2$ and $q \neq 1$. If q = 2, one can easily verify that $f[h+1] = \overline{f[h]}$, for each h > i. By hypothesis $f^2[i+1] = 1$ then $f^3[i+3] = 1$ then $f^1[p+i+3] = 1$, but $f^1[p+i+2] = 1$ by hypothesis. A contradiction follows. If q = 1 then $f[i] = f^3[i] = 0$ and $f[i] = f^2[i] = 1$, a contradiction.

Since f^2 occurs in $S_s(u)$, f^2 matches the corresponding positions of u in I_2 except for the positions involved in the swap S_s , i.e. s and s + 1. Hence, for each $h \in I_2$ with



Figure 3: The representation of the three occurrences f^1 , f^2 , and f^3 of f in $O^1(u)$, $O^2(u)$ and $O^3(u)$, respectively, when u[s..s+2] = 100.

 $h \neq q + i - 2$ and $h \neq q + i - 1$ we have $u[p+h] = f^2[h]$.

It follows that $u[p+i] = f^2[i] = 1$, because $q \neq 2$ and $q \neq 1$. Moreover, since f^1 occurs in $R_{s+2}(u)$, f^1 matches the corresponding positions of u in I_1 unless for s + 2 where the replacement is applied. More formally, for each $h \in I_1$, with $h \neq i$ we have $u[h] = f^1[h]$. It follows that $u[p+i] = f^1[p+i] = 1$, because p > 0.

On the other hand $f^1[p+i] = f^3[p+i]$. Since f^3 occurs in $R_s(u)$, f^3 matches the corresponding positions of u in I_3 unless for s where the replacement is applied. In other words, for each $h \in I_3$ with $h \neq i-2$ we have $u[r+h] = f^3[h]$. It follows that $u[r+p+i] = f^3[p+i]$ because p > 0.

But $u[r+p+i] = f^2[r+i] = 0$ because p > 0. Then a contradiction follows.

5.2 The necessary condition

Once we have collected all the tools, we are now ready to prove the *only if* direction of Theorem 16.

Proposition 23 If $f \in \Sigma^n$ is tilde-non-isometric then one among (C0), (C1), (C2), (C3), (C4), (C5) cases of Theorem 16 occurs (up to complement, reverse and exchange of rows).

Proof: Let f be a tilde-non-isometric word, (u, v) be a pair of tilde-witnesses for f with minimal distance among the pairs of tilde-witnesses of minimal length. Consider a tilde-transformation from u to v that does not contain two swaps at two consecutive positions.

Recall that if a tilde-transformation contains swaps S_i and S_{i+1} for some position i in u then they can be substituted by replacements R_i and R_{i+2} still keeping the total number of operations. Throughout the proof let $s, t, k_s, k_t, O_s, O_t, I_s, I_t$, with $k_s < k_t$, be as in Lemma 20.

Further suppose that u[s] = c and u[t] = d for some $c, d \in \Sigma$.

First, consider the case where the characters modified by O_s and O_t are not adjacent.

If there are at least two characters of u interleaving the characters modified by O_s and the ones modified by O_t then f has a 2-tilde-error overlap with non-adjacent errors and any of its blocks containing all the error/ modified positions has length strictly greater than 3, thus falling in case (C1).

Suppose now that there is only one character of u interleaving the characters modified by O_s and the ones modified by O_t . Then, four situations may occur.

1. $O_s = R_s, O_t = R_t$ with t = s + 2 and u[s + 1] = v[s + 1]. Let u[s + 1] = eand u = wcedz with $w, z \in \Sigma^*$. Then, $O_s(u) = w\overline{a}cbz$ and $O_t(u) = wac\overline{b}z$ and, since I_s and I_t must share the positions s and t then f has a 2-tilde-error overlap with block $\binom{\overline{ced}}{ce\overline{d}}$. Applying Lemma 22, $\binom{u[s..s+2]}{v[s..s+2]} \neq \binom{100}{001}$ and then $\binom{\overline{ced}}{ce\overline{d}} \neq \binom{000}{101}$,

thus falling in case (C1).

- 2. $O_s = S_s, O_t = R_t$ with t = s + 3 and u[s + 2] = v[s + 2]. Let u[s + 2] = e and $u = wc\overline{c}edz$ with $w, z \in \Sigma^*$. Then, $O_s(u) = w\overline{c}cedz$ and $O_t(u) = wc\overline{c}edz$. If $I_s \cap I_t$ includes s, s + 1, s + 2, s + 3 then f has a 2-tilde-error overlap with non-adjacent error positions and any of its blocks containing the error positions has length strictly greater than 3, thus falling in case (C1). If $I_s \cap I_t$ includes s + 1, s + 2, s + 3 then f has a 2-tilde-error overlap with non-adjacent error positions with strictly greater than 3, thus falling in case (C1). If $I_s \cap I_t$ includes s + 1, s + 2, s + 3, but not s, then f has a 2-tilde-error overlap with non-adjacent error positions with block $\left(\frac{\overline{c}ed}{ced}\right)$ where $\overline{c}e\overline{d}$ is a prefix of f. If c = d then the block $\binom{000}{101}$, may occur in the 2-tilde-error overlap, but the 2-tilde-error overlap is not $\binom{8x}{ax}\binom{000}{101}\binom{yb}{yb}$ with $x \neq \varepsilon$, thus falling in case (C1).
- 3. $O_s = R_s, O_t = S_t$ with t = s + 2 and u[s + 1] = v[s + 1]. This case is the symmetric of the previous one. Indeed, f has a 2-tilde-error overlap with non-adjacent error positions where the block $\binom{000}{101}$ may occur, but in this case 000 is a suffix of f, thus falling again in case (C1).
- 4. $O_s = S_s, O_t = S_t$ with t = s + 3 and u[s + 2] = v[s + 2]. Similarly as in the two previous cases, if $I_s \cap I_t$ includes s + 1, s + 2, s + 3, but not s and not s + 4 then f has a 2-tilde-error overlap with non-adjacent error positions where the block $\binom{000}{101}$ may occur, but in this case 000 is a prefix of f and 101 is a suffix of f thus falling again in case (C1).

Suppose now that there is no character of u interleaving the characters modified by O_s and the ones modified by O_t .

Again, four situations may occur.

- 1. $O_s = R_s, O_t = R_t$ with t = s + 1. Then, u[s] = u[s + 1], otherwise R_s and R_t could be replaced in the tilde-transformation from u to v with a single swap S_s obtaining a tilde-transformation with a less number of operations. Hence, suppose u = w00z; then, $O_s(u) = R_s(u) = w10z$ and $O_{s+1}(u) = R_{s+1}(u) = w01z$, and since the intervals I_s and I_t must share the positions s and t, then f has a 1-tilde-error overlap $\binom{\$x}{ax}\binom{01}{y\$}\binom{ya}{y\$}$ with $x, y \in \Sigma^*$, $a \in \Sigma$, as in (CO).
- 2. $O_s = S_s, O_t = R_t$ with t = s + 2. In this case $u = wc\overline{c}dz$, with $w, z \in \Sigma^*$ and $c, d \in \Sigma$. Suppose c = 1 (then $\overline{c} = 0$). By Lemma 22, $d \neq 1$, because $\binom{u[s..s+2]}{v[s..s+2]} \neq \binom{101}{010}$. On the other hand, if d = 0 then $\binom{u[s..s+2]}{v[s..s+2]} = \binom{100}{011}$ and $O_s(u) = w010z$ and $O_t(u) = w101z$. If $I_s \cap I_t$ includes s, s + 1 and s + 2, then (C3) is verified. If $I_s \cap I_t$ includes only s + 1 and t = s + 2, then f has a 1-tilde-error overlap $\binom{\$x}{ax} \binom{01}{10} \binom{ya}{y\$}$, as in (C0).
- 3. $O_s = R_s, O_t = S_t = S_{s+1}$. The proof is equivalent to the previous case by considering the reverses of u and v in the above proof.
- 4. $O_s = S_s, O_t = S_t$ with t = s + 2. In this case $u = wc\overline{c}d\overline{d}z$. Suppose c = 1. Then two cases occur: d = 1 and d = 0.

If d = 1 then u = w1010z and $O_s(u) = S_s(u) = w0110z$ and $O_t(u) = w0110z$

 $S_t(u) = S_{s+2}(u) = w1001z$. If all positions s, s+1, t = s+2, t+1 = s+3 are included in $I_s \cap I_t$ then case (C2) is verified. Consider now the case where $s, s+1, s+2 \in I_s \cap I_t$ and $s+3 \notin I_s \cap I_t$. This means that I_s ends in position s+2, and f has a 2-tilde-error overlap $\binom{\$x}{ax}\binom{100}{011}\binom{1}{\$}$, as in case (C4), up to complement.

In a similar way, if s + 1, s + 2, $s + 3 \in I_s \cap I_t$ and $s \notin I_s \cap I_t$, then I_t starts in position s + 1 and f has a 2-tilde-error overlap $\binom{\$}{0}\binom{001}{110}\binom{xa}{x\$}$, as in case (C4), up to reverse and exchange of rows.

The last situation is when $s + 1, s + 2 \in I_s \cap I_t$ and $s, s + 3 \notin I_s \cap I_t$ then f starts with 001 and ends with 011, providing a 2-tilde-error overlap as in case (C5).

Consider now d = 0, then u = w1001z. Again, we have four cases. If $s, s + 1, s + 2, s + 3 \in I_s \cap I_t$ then case (C2) is verified.

If $s, s+1, s+2 \in I_s \cap I_t$ and $s+3 \notin I_s \cap I_t$, then f has a 2-tilde-error overlap $\binom{\$x}{ax}\binom{101}{010}\binom{0}{\$}$, as in case (C3), with $y = \varepsilon$ and b = 0 and up to complement.

If $s + 1, s + 2, s + 3 \in I_s \cap I_t$ and $s \notin I_s \cap I_t$, then f has a 2-tilde-error overlap $\binom{0}{0}\binom{010}{101}\binom{yb}{y\$}$, as in case (C3), with $x = \varepsilon$ and a = 0.

If s + 1, $s + 2 \in I_s \cap I_t$ and $s, s + 3 \notin I_s \cap I_t$, then f starts with 01 and ends with 10 corresponding to case (C0) with $x, y = \varepsilon$.

5.3 Construction of tilde-witnesses

As already discussed, in order to prove that a word is tilde-non-isometric it is sufficient to exhibit a pair of tilde-witnesses. As a preparation for the sufficient condition of the characterization theorem we introduce a technique to define a pair of tilde-witnesses for a word, starting from its tilde-error overlaps.

Definition 24 Let $f \in \Sigma^*$ have a 2-tilde-error overlap of shift r, length $\ell = n - r$, and let $(O_i, O_j), 1 \le i < j \le \ell$, be a tilde-transformation of $\operatorname{pre}_{\ell}(f)$ in $\operatorname{suf}_{\ell}(f)$, with $(O_i, O_j) \ne (S_i, S_{i+1})$.

 $If(O_i, O_j) \neq (R_i, S_{i+1}), (S_i, R_{i+2})$ then

$$\tilde{\alpha}_r(f) = \operatorname{pre}_r(f)O_i(f) \text{ and } \beta_r(f) = \operatorname{pre}_r(f)O_j(f)$$
(3)

else

$$\tilde{\alpha}_r(f) = \operatorname{pre}_r(f)S_i(f) \text{ and } \tilde{\beta}_r(f) = \operatorname{pre}_r(f)R_{i+2}(f)$$
 (4)

When there is no ambiguity, we omit f and we simply write $\tilde{\alpha}_r$ and β_r .

Lemma 25 If f has a 2-tilde-error overlap of shift r then $\tilde{\alpha}_r(f)$ is f-free.

Proof: Let f have a 2-tilde-error overlap of shift r and length $\ell = n - r$ and let $(O_i, O_j), 1 \leq i < j \leq \ell$, be a tilde-transformation of $\operatorname{pre}_{\ell}(f)$ in $\operatorname{suf}_{\ell}(f)$, with $(O_i, O_j) \neq (S_i, S_{i+1})$. If $(O_i, O_j) = (R_i, R_j)$ then $\tilde{\alpha}_r$ is f-free by Claim 1 of Lemma 2.2 in [29], also in the case of adjacent errors. If $(O_i, O_j) = (S_i, R_{i+2})$ then, by Equation (3), we have $\tilde{\alpha}_r = w_1 S_i(f)$ then $\tilde{\alpha}_r [r + k] = f[k]$, for any $1 \leq k \leq n$, with $k \neq i$ and $k \neq i + 1$. If f occurs in $\tilde{\alpha}_r$ in position $r_1 + 1$ we have that $1 < r_1 < r$ (if $r_1 = 1$ then f[i] = f[i + 1] and this is not possible since $O_i = S_i$) and $\tilde{\alpha}_r [r_1 + 1 \dots r_1 + n] = f[1 \dots n]$. Finally, we have that $\tilde{\alpha}_r[k] = f[k]$, for $k \neq r + j$.

In conclusion, we have that $f[i] = \tilde{\alpha}_r[r_1 + i] = f[r_1 + i]$ (trivially, $r_1 + i \neq r + j$). Furthermore $f[r_1 + i] = \tilde{\alpha}_r[r + r_1 + i]$ ($r_1 + i \neq i$ and $r_1 + i \neq i + 1$ because $r_1 > 1$). But $\tilde{\alpha}_r[r + r_1 + i] = f[r + i]$ then we have the contradiction that f[i] = f[r + i].

In all the other cases the proof is similar. For clarity, note that, also in the case of adjacent errors, supposing that f occurs in $\tilde{\alpha}_r$ leads to a contradiction in f[i] that is not influenced by $j.\square$

Note that while $\tilde{\alpha}_r$ is always *f*-free, $\tilde{\beta}_r$ is not. Indeed, the property of $\tilde{\beta}_r$ being not *f*-free is related to a condition on the overlap of *f*. We give the following definition.

Definition 26 Let $f \in \Sigma^*$ have a 2-tilde-error overlap of shift r, length $\ell = n - r$. Then, the 2-tilde-error overlap satisfies Condition[~] if any tilde-transformation $(O_i, O_j), 1 \leq i < j \leq \ell$, of $\operatorname{pre}_{\ell}(f)$ in $\operatorname{suf}_{\ell}(f)$, is $(O_i, O_j) = (R_i, R_j)$ or $(O_i, O_j) = (S_i, S_j)$ and

$$\begin{cases} r \text{ is even} \\ j-i=r/2 \\ f[i..(i+r/2-1)] = f[j..(j+r/2-1)] \end{cases}$$
 (Condition~)

Lemma 27 Let $f \in \Sigma^n$ have a 2-tilde-error overlap of shift r, then $\hat{\beta}_r(f)$ is not f-free iff the 2-tilde-error overlap satisfies Condition \sim .

Proof: Let us suppose f have a 2-tilde-error overlap of shift r and length $\ell = n - r$ and let $(O_i, O_j), 1 \le i < j \le \ell$, be a tilde-transformation of $\text{pre}_{\ell}(f)$ in $\text{suf}_{\ell}(f)$ that satisfies *Condition*~.

Now, if $(O_i, O_j) = (R_i, R_j)$ then the fact that $\tilde{\beta}_r(f)$ is not f-free can be shown as in the proof of Claim 2 of Lemma 2.2 in [29].

If $(O_i, O_j) = (S_i, S_j)$, then that proof must be suitably modified as follows. Let f[i] = f[j] = x, $f[i+r] = f[j+r] = \overline{x}$, $f[i+1] = f[j+1] = \overline{x}$ and f[i+1+r] = f[j+1+r] = x. Then, it is possible to show that, for some $k_1, k_2 \ge 0$, $f = \rho(uw)^{k_1}uwuw\overline{u}w\overline{u}(w\overline{u})^{k_2}\sigma$, where $u = x\overline{x}$, w = f[i+2..j-1] (*w* is empty, if j = i+2) and ρ and σ are, respectively, a suffix and a prefix of *w*. Then, $\tilde{\beta}_r(f) = \rho(uw)^{k_1+1}uwuw\overline{u}w\overline{u}(w\overline{u})^{k_2+1}\sigma$ and, hence, $\tilde{\beta}_r(f)$ is not *f*-free.

Assume now that $\hat{\beta}_r(f)$ is not f-free and suppose that a copy of f, say f', occurs in $\tilde{\beta}_r(f)$ at position $r_1 + 1$. A reasoning similar to the one used in the proof of Lemma 25, shows that $j - i = r_1$ and $j - i = r - r_1$. Hence $r = 2r_1$ is even and j - i = r/2. Therefore, f[i + t] = f'[i + t] = f[i + t + r/2] = f[j + t], for $0 \le t \le r/2$, i.e. f[i...(i + r/2 - 1)] = f[j...(j + r/2 - 1)]. Moreover, such conditions with the assumption that $\tilde{\beta}_r(f)$ is not f-free can be used to show that $(O_i, O_j) \ne (S_i, R_j)$ and $(O_i, O_j) \ne (R_i, S_j)$ and thus the 2-tilde-error overlap satisfies $Condition^{\sim}$. \Box

5.4 The sufficient condition

In this last subsection we conclude the proof of Theorem 16 by proving the *if* direction.

We distinguish the cases of 1-tilde-error overlap (case (C0)), 2-tilde-error overlap with non-adjacent errors (case (C1)) and, finally, 2-tilde-error overlap with adjacent errors (cases (C2) - (C5)). Non-adjacent errors can be dealt with the standard techniques used for the Hamming distance, while the case of adjacent ones may show new issues. Note that, for some error types, we need also to distinguish sub-cases related to the different characters adjacent to the overlap. For each case in the list, the proof consists in providing a pair of words which is proved to be a pair of tilde-witnesses for f by showing that it fulfills conditions 1., 2. and 3. in Definition 9.

Let us start with the case (C0) of a 1-tilde-error overlap.

Proposition 28 (C0) If f has a 1-tilde-error overlap $\binom{\$x}{ax}\binom{01}{10}\binom{yb}{y\$}$, with $x, y \in \Sigma^*$, $a, b \in \Sigma$, then f is tilde-non-isometric.

Proof: Let f have a 1-tilde-error overlap $\binom{\$x}{ax}\binom{01}{10}\binom{yb}{y\$}$ with shift r and block $\binom{01}{10}$ occurring at position i = |x| + 1. The pair (u, v) with:

$$u = \operatorname{pre}_r(f)R_i(f)$$
 $v = \operatorname{pre}_r(f)R_{i+1}(f)$

is a pair of tilde-witnesses for f. Indeed, first of all, note that (u, v) = (w11z, w00z)for some $w, z \in \Sigma^*$. Therefore, $dist_{\sim}(u, v) = 2$. Moreover, using techniques similar to the ones used in Lemma 25, one can prove that u and v are f-free. Finally, no f-free tilde-transformation from u to v exists. The only possible tilde-transformations from u to v are (R_{r+i}, R_{r+i+1}) and (R_{r+i+1}, R_{r+i}) and they both are not f-free since foccurs at position r + 1 in $R_{r+i}(u)$ and at position 1 in $R_{r+i+1}(u)$. \Box

Example 29 The word f = 101 has a 1-tilde-error overlap $\binom{\$}{1}\binom{10}{01}\binom{1}{\$}$ with block $\binom{10}{01}$ in position 1 therefore it is tilde-non-isometric. In fact, the pair (u, v) with u = 1001 and v = 1111 is a pair of tilde-witnesses.

Let us consider the case of non-adjacent errors and prove that f satisfies (C1) then it is tilde-non-isometric.

Proposition 30 (C1) If f has a 2-tilde-error overlap with non-adjacent errors, different from $\binom{\$x}{ax}\binom{000}{101}\binom{yb}{y\$}$ with $x, y \in \Sigma^+$, $a, b \in \Sigma$ then f is tilde-non-isometric.

Proof: Let f have a 2-tilde-error overlap with shift r and length $\ell = n - r$ with non-adjacent errors different from $\binom{\$x}{ax}\binom{000}{101}\binom{yb}{y\$}$. Let (O_i, O_j) , $1 \le i < j \le \ell$ be a tilde-transformation from $\operatorname{pre}_{\ell}(f)$ to $\operatorname{suf}_{\ell}(f)$. In case of the 2-tilde-error overlap $\binom{\$x}{ax}\binom{000}{001}\binom{yb}{y\$}$, consider $(O_i, O_j) = (R_i, R_{i+2})$. Then, if the 2-tilde-error overlap does not satisfy $Condition^{\sim}$ then the pair $(\tilde{\alpha}_r, \tilde{\beta}_r)$ as in Equation (3) is a pair of tilde-witnesses for f. Indeed, $dist_{\sim}(\tilde{\alpha}_r, \tilde{\beta}_r) = 2$ and from Lemma 25 and Lemma 27, $\tilde{\alpha}_r$ and $\tilde{\beta}_r$ are f-free. Finally, no f-free tilde-transformation from $\tilde{\alpha}_r$ to $\tilde{\beta}_r$ exists. In fact, the tilde-transformations from $\tilde{\alpha}_r$ to $\tilde{\beta}_r$, (O_{r+i}, O_{r+j}) and (O_{r+j}, O_{r+i}) , are not f-free because f occurs at position r + 1 in $O_{r+i}(\tilde{\alpha}_r)$ and at position 1 in $O_{r+j}(\tilde{\alpha}_r)$. And no other tilde-transformation is possible since $\binom{\tilde{\alpha}_r}{\tilde{\beta}_r}$ cannot contain either $\binom{100}{001}$ or $\binom{\$s}{101}\binom{900}{\frac{y\$}{\frac{9}{5}}}$ in the first case, against the hypotesis, and $\binom{\$x}{ax}\binom{101}{010}\binom{yb}{\frac{y\$}{\frac{9}{5}}}$, in the second case, that is not allowed since the errors would be adjacent. If, instead, the 2-tilde-error overlap satisfies $Condition^{\sim}$, then consider the pair $(\tilde{\eta}_r, \tilde{\gamma}_r)$

with $\tilde{\eta}_r = \operatorname{pre}_r(f)O_i(f)\operatorname{suf}_{r/2}(f)$, $\tilde{\gamma}_r = \operatorname{pre}_r(f)O_j(O_t(f))\operatorname{suf}_{r/2}(f)$, where t = j + r/2 and $O_t = R_t$, if $O_i = R_i$, and $O_t = S_t$, if $O_i = S_i$. Note that $O_j(O_t(f)) = O_t(O_j(f))$, since the 2-tilde-error overlap has non-adjacent errors. The pair $(\tilde{\eta}_r, \tilde{\gamma}_r)$ is a pair of tilde-witnesses for f. Indeed, $dist_{\sim}(\tilde{\gamma}_r, \tilde{\eta}_r) = 3$ (recall that the 2-tilde-error overlap has non-adjacent errors). Moreover, using techniques similar to the ones

used in Lemma 25, one can prove that $\tilde{\gamma}_r$ and $\tilde{\eta}_r$ are *f*-free. Finally, no *f*-free tildetransformation from $\tilde{\gamma}_r$ to $\tilde{\eta}_r$ exists. In fact, any possible tilde-transformation from $\tilde{\gamma}_r$ to $\tilde{\eta}_r$, is given by a permutation of $\{O_{r+i}, O_{r+j}, O_{r+t}\}$ and any of such tildetransformations let *f* occur at position 1, *r* or $3r/2.\Box$

Let us examine all the remaining cases (C2) up to (C5) that correspond to the different kinds of 2-tilde-error overlaps of f with adjacent errors and state the whole sufficient condition for a word to be non-tilde-isometric.

Proposition 31 [C2 - C5] Let $f \in \Sigma^n$. If one of the following cases occurs for f (up to complement, reverse, and exchange of rows) then f is tilde-non-isometric.

- (C2) f has a 2-tilde-error overlap $\binom{\$x}{ax}\binom{0101}{1010}\binom{yb}{y\$}$ or $\binom{\$x}{ax}\binom{0110}{1001}\binom{yb}{y\$}$ with $x, y \in \Sigma^*$, $a, b \in \Sigma$
- (C3) f has a 2-tilde-error overlap $\binom{\$x}{ax}\binom{010}{101}\binom{yb}{y\$}$ with $x, y \in \Sigma^*$, $a, b \in \Sigma$
- (C4) f has a 2-tilde-error overlap $\binom{\$x}{ax}\binom{011}{100}\binom{0}{\$}$ with $x \in \Sigma^*$, $a \in \Sigma$

(C5) f has a 2-tilde-error overlap $\binom{\$}{0}\binom{00}{11}\binom{1}{\$}$

Proof:

(C2).

Let f have a 2-tilde-error overlap $\binom{\$x}{ax}\binom{0101}{1010}\binom{yb}{y\$}$ or $\binom{\$x}{ax}\binom{0110}{1001}\binom{yb}{y\$}$ with $x, y \in \Sigma^*$, $a, b \in \Sigma$.

We distinguish the proof in two cases.

Case a. Suppose that f has a 2-tilde-error overlap $\binom{\$x}{ax}\binom{0101}{1010}\binom{yb}{y\$}$, $x, y \in \Sigma^*$, $a, b \in \Sigma$, with shift r and length $\ell = n - r$ and let (O_i, O_j) , be a tilde-transformation from $\operatorname{pre}_{\ell}(f)$ to $\operatorname{suf}_{\ell}(f)$. Note that it must be $(O_i, O_j) = (S_i, S_{i+2})$ or $(O_i, O_j) = (S_{i+2}, S_i)$ where i = |x| + 1.

If the 2-tilde-error overlap does not satisfy $Condition^{\sim}$ then the pair $(\tilde{\alpha}_r, \tilde{\beta}_r)$ as in Equation (3) is a pair of tilde-witnesses for f. Indeed, $dist_{\sim}(\tilde{\alpha}_r, \tilde{\beta}_r) = 2$ and from Lemma 25 and Lemma 27, $\tilde{\alpha}_r$ and $\tilde{\beta}_r$ are f-free. Finally, no f-free tilde-transformation from $\tilde{\alpha}_r$ to $\tilde{\beta}_r$ exists. In fact, the only possible tilde-transformations from $\tilde{\alpha}_r$ to $\tilde{\beta}_r$ are (S_{r+i}, S_{r+i+2}) and (S_{r+i+2}, S_{r+i}) . But they are not f-free because they let f occur at position 1 and at position r + 1 in $S_{r+i+2}(\tilde{\alpha}_r)$ and $S_{r+i}(\tilde{\alpha}_r)$, respectively.

If instead the 2-tilde-error overlap satisfies $Condition^{\sim}$, then consider the pair $(\tilde{\eta}_r, \tilde{\gamma}_r)$ with $\tilde{\eta}_r = \operatorname{pre}_r(f)S_i(f)\operatorname{suf}_{r/2}(f)$, $\tilde{\gamma}_r = \operatorname{pre}_r(f)S_{i+2}(S_{i+4}(f))\operatorname{suf}_{r/2}(f)$. Note that $S_{i+2}(S_{i+4}(f) = S_{i+4}(S_{i+2}(f))$.

The pair $(\tilde{\eta}_r, \tilde{\gamma}_r)$ is a pair of tilde-witnesses for f. Indeed, it is easy to see that $dist_{\sim}(\tilde{\gamma}_r, \tilde{\eta}_r) = 3$. Moreover, using techniques similar to the ones used in Lemma 25, one can prove that $\tilde{\gamma}_r$ and $\tilde{\eta}_r$ are f-free. Finally, no f-free tilde-transformation from $\tilde{\gamma}_r$ to $\tilde{\eta}_r$, consists. In fact, any possible tilde-transformations from $\tilde{\gamma}_r$ to $\tilde{\eta}_r$, consists of three swap operation and is given by a permutation of $\{S_{r+i}, S_{r+i+2}, S_{r+i+4}\}$. But any of such tilde-transformations let f occurs at position 1 or r + 1 or 3r/2 + 1.

Case b. Suppose that f has a 2-tilde-error overlap $\binom{\$x}{ax}\binom{0110}{1001}\binom{yb}{y\$}$, $x, y \in \Sigma^*$, $a, b \in \Sigma$, with shift r and length $\ell = n - r$ and let (O_i, O_j) , be a tilde-transformation from $\operatorname{pre}_{\ell}(f)$ to $\operatorname{suf}_{\ell}(f)$. Note that it must be $(O_i, O_j) = (S_i, S_{i+2})$ or $(O_i, O_j) = (S_{i+2}, S_i)$ where i = |x| + 1. Now, the pair $(\tilde{\alpha}_r, \tilde{\beta}_r)$ as in Equation (3) is a pair of tilde-witnesses for f. Note that, in this case, the 2-tilde-error overlap does not satisfy Condition[~] since j = i + 2 and f[i] is different from f[j].

(C3) Let f have a 2-tilde-error overlap $\binom{\$x}{ax}\binom{010}{101}\binom{yb}{y\$}$, $x, y \in \Sigma^*$, $a, b \in \Sigma$ with shift r and length $\ell = n - r$ and let (O_i, O_j) , be a tilde-transformation from $\operatorname{pre}_{\ell}(f)$ to $\operatorname{suf}_{\ell}(f)$. Note that it must be $(O_i, O_j) = (S_i, R_{i+2})$ or $(O_i, O_j) = (R_i, S_{i+1})$. Then, the 2-tilde-error overlap does not satisfy $Condition^\sim$. The pair $(\tilde{\alpha}_r, \tilde{\beta}_r)$ as in Equation (4) is a pair of tilde-witnesses for f. Indeed, it is easy to see that $\operatorname{dist}_{\sim}(\tilde{\alpha}_r, \tilde{\beta}_r) = 2$. Moreover, from Lemma 25 and Lemma 27, $\tilde{\alpha}_r$ and $\tilde{\beta}_r$ are f-free. Finally, no f-free tilde-transformation from $\tilde{\alpha}_r$ to $\tilde{\beta}_r$ exists. In fact, the tilde-transformations from $\tilde{\alpha}_r$ to $\tilde{\beta}_r$, (O_{r+i}, O_{r+j}) and (O_{r+j}, O_{r+i}) , are not f-free because they let f occur at position r + 1 and at position 1 in $O_{r+i}(\tilde{\alpha}_r)$ and $O_{r+j}(\tilde{\alpha}_r)$, respectively, and no other tilde-transformation is possible.

(C4) Let f have a 2-tilde-error overlap $\binom{\$x}{ax}\binom{011}{100}\binom{0}{\$}$, $x \in \Sigma^*$, $a \in \Sigma$ with shift r and length $\ell = n - r$ and let (O_i, O_j) , be a tilde-transformation from $\operatorname{pre}_{\ell}(f)$ to $\operatorname{suf}_{\ell}(f)$. Note that it must be $(O_i, O_j) = (S_i, R_{i+2})$ with i = |x| + 1. In this case we need a different technique to construct the pair of tilde-witnesses $(\tilde{\alpha}_r, \tilde{\delta}_r)$ for f. More exactly, we set $\tilde{\alpha}_r$ as in Equation (4) and $\tilde{\delta}_r = \text{pre}_r(f)S_{i+2}(f)$. It is easy to see that $dist_{\sim}(\tilde{\alpha}_r, \tilde{\delta}_r) = 2$. Moreover, from Lemma 25, $\tilde{\alpha}_r$ is f-free. Here we prove that $\tilde{\delta}_r$ is f-free. Indeed, suppose that f occurs in δ_r starting from position $r_1 + 1$. By the definition of δ_r , we have that $\delta_r[k] = f[k]$ for $1 \le k \le |f|$ and $k \ne r+i, r+i+1$. Moreover, $\tilde{\delta}_r[r+k] = f[k]$, for $1 \le k \le |f|$ and $k \ne j, j+1$. Now, we have 0 = f[i] and, since f occurs in $\tilde{\delta}_r$ starting from position $r_1 + 1$, $f[i] = \tilde{\delta}_r[r_1 + i]$. Since $\delta_r[r+i+1] = 1$, it holds $r_1 + i \neq r+i+1$. Moreover $r_1 + i \neq r+i$ and we can conclude that $\tilde{\delta}_r[r_1+i] = f[r_1+i]$ i.e. $0 = f[i] = \tilde{\delta}_r[r_1+i] = f[r_1+i]$. Note that it cannot be $r_1 + i = j + 1$, since $\tilde{\delta}_r[r + j + 1] = 1$. Now, if $r_1 + i \neq j$, we can say $f[r_1 + i] = \delta_r[r + r_1 + i]$. Since f occurs in $\tilde{\delta}_r$ starting from position $r_1 + 1$, we have $\tilde{\delta}_r[r+r_1+i] = f[r+i]$ Altogether, we have $0 = f[i] = \tilde{\delta}_r[r_1+i] = f[r_1+i] = f[r_1+i]$ $\tilde{\delta}_r[r+r_1+i] = f[r+i] = 1$, a contradiction. Therefore, the only possibility is that $r_1 + i = j$, i.e. $r_1 = 2$. Similar considerations on f[j] show that, if f occurs in δ_r starting from position $r_1 + 1$, then $r_1 = 1$. Therefore, if a copy of f occurs in δ_r starting from position $r_1 + 1$, then it should be $r_1 = 2$ and $r_1 = 1$ and this is impossible. Hence δ_r is *f*-free.

Moreover, no f-free tilde-transformation from $\tilde{\alpha}_r$ to $\tilde{\delta}_r$ exists. The only possible tilde-transformations are (S_{r+i}, S_{r+i+2}) and (S_{r+i+2}, S_{r+i}) and they are not f-free. Hence $(\tilde{\alpha}_r, \tilde{\delta}_r)$ is a pair of tilde-witnesses for f.

(C5) Let f have a 2-tilde-error overlap $\binom{\$}{0}\binom{00}{11}\binom{1}{1}$ with shift r and length 2 and let (O_i, O_j) , be a tilde-transformation from $\operatorname{pre}_2(f) = 00$ to $\operatorname{suf}_2(f) = 11$. Note that it must be $(O_i, O_j) = (R_i, R_{i+1})$ with i = 1. In this case we have f = w001 = 001z, for some $w, z \in \Sigma^*$. Consider the pair $(\tilde{\alpha}_r, \tilde{\psi}_r)$ with $\tilde{\alpha}_r$ as in Equation (3) and $\tilde{\psi}_r = \operatorname{pre}_{r-1}(f)1S_{i+1}(f)$. In other words $\tilde{\alpha}_r = w0101z$ and $\tilde{\psi}_r = w1010z$.

The pair $(\tilde{\alpha}_r, \psi_r)$ is a pair of tilde-witnesses for f. Indeed, it is easy to see that $dist_{\sim}(\tilde{\alpha}_r, \tilde{\psi}_r) = 2$. Moreover, from Lemma 25, $\tilde{\alpha}_r$ is f-free and, using techniques similar to the ones used in (C4) for $\tilde{\delta}_r$, one can prove that $\tilde{\psi}_r$ is f-free. Finally, no f-free tilde-transformation from $\tilde{\alpha}_r$ to $\tilde{\psi}_r$ exists. The only possible tilde-transformations are (S_{r-1}, S_{r+1}) and (S_{r+1}, S_{r-1}) and they both are not f-free. Remark that, in this case, the pair $(\tilde{\alpha}_r, \tilde{\beta}_r)$ of Equation 3 is not a pair of tilde-witnesses because $dist_{\sim}(\tilde{\alpha}_r, \tilde{\beta}_r) = 1$. \Box

The following is an example of construction of tilde-witnesses when f has a 2-tilde-error overlap with two adjacent swaps.

Example 32 The word f = 10010110 has a 2-tilde-error overlap $\binom{\$}{1}\binom{1001}{0110}\binom{0}{\$}$ of shift r = 4 and swaps in positions 1, 3. By Proposition 31 case (C2), the pair $(\tilde{\alpha}_4, \tilde{\beta}_4)$ with $\tilde{\alpha}_4 = 100101010110$ and $\tilde{\beta}_4 = 100110100110$ is a pair of tilde-witnesses. Then f is tilde-non-isometric. Note that f is Ham-isometric.

Propositions 28, 30 and 31 together, prove the sufficient condition for a word to be tilde-non-isometric. This concludes the proof of Theorem 16.

6 Conclusion

In this paper the problem of fully characterizing isometric words has been solved in the setting of the edit distance that allows swap and replacement operations. Compared with the setting of Hamming distance, the problem turned out to be much more complicated, since swap operation gives rise to transformations that differ from one another not only because of the execution order of operations, as in the case of Hamming, but also for the kind of operations executed. This fact makes things harder to manage and the characterization based on overlaps needs to specify some special cases that can occur.

Recall that the original reason for the introduction of isometric words was their connection with isometric subgraphs of the hypercube Q_n (cf.[24, 22, 25]). In fact, if all the nodes containing a given isometric word as factor are removed then the distances between the remaining nodes do not change. These subgraph of hypercubes are called generalized Fibonacci cubes and are subgraphs isometric to hypercubes in the graph-theoretic sense.

Over the years, many variations of the hypercube have been introduced in order to improve some of its features. For example, folded hypercubes (cf. [17]) and ehnanced hypercubes (cf. [27]) have been defined by adding some edges to the hypercube and present many advantages for some topological features. In [3], the definition of tilde-hypercube is introduced by exploiting the tilde-distance. Adding the swap operation increases the number of connections in the hypercube (since some nodes having distance 2 in the hypercube, using swap operation, have distance 1 in the tilde-hypercube). Tilde-isometric words then are used to introduce the generalized tilde-Fibonacci cubes that are isometric subgraphs of the tilde-hypercube and revealed important properties.

In conclusion, the swap and mismatch distance we adopted in this paper opens up new scenarios and presents interesting new situations that surely deserve further investigation. The definition and characterization of tilde-isometric words, besides having an important combinatorial value can serve as base for strings and graph algorithmic developments.

References

- A. Amir, R. Cole, R. Hariharan, M. Lewenstein and E. Porat, Overlap matching, *Inf. Comput.* 181(1) (2003) 57–74.
- [2] A. Amir, E. Eisenberg and E. Porat, Swap and mismatch edit distance, Algorithmica 45(1) (2006) 109–120.
- [3] M. Anselmo, G. Castiglione, M. Flores, D. Giammarresi, M. Madonia and S. Mantaci, Hypercubes and isometric words based on swap and mismatch dis-

tance, Descriptional Complexity of Formal Systems. DCFS 2023 Proceedings, Lecture Notes in Computer Science **13918**, (Springer, 2023), pp. 21–35.

- [4] M. Anselmo, G. Castiglione, M. Flores, D. Giammarresi, M. Madonia and S. Mantaci, Isometric words based on swap and mismatch distance, *Developments in Language Theory - 27th International Conference, DLT 2023 Proceedings, Lecture Notes in Computer Science* **13911**, (Springer, 2023), pp. 23–35.
- [5] M. Anselmo, M. Flores and M. Madonia, Quaternary *n*-cubes and isometric words, *Combinatorics on Words*, *Lect. Notes Comput. Sci.* **12842** (2021), pp. 27– 39.
- [6] M. Anselmo, M. Flores and M. Madonia, Fun slot machines and transformations of words avoiding factors, 11th International Conference on Fun with Algorithms, LIPIcs 226 (2022), pp. 4:1–4:15.
- [7] M. Anselmo, M. Flores and M. Madonia, On k-ary n-cubes and isometric words, *Theor. Comput. Sci.* 938 (2022) 50–64.
- [8] M. Anselmo, M. Flores and M. Madonia, Density of Ham- and Lee- nonisometric k-ary words, *ICTCS'23 Italian Conference on Theoretical Computer Science, CEUR Workshop Proceedings* 3587, (CEUR-WS.org, 2023), pp. 116– 128.
- [9] M. Anselmo, M. Flores and M. Madonia, Computing the index of non-isometric k-ary words with Hamming and Lee distance, *Computability*, *IOS* Pre-press (2024) 1–24.
- [10] M. Anselmo, D. Giammarresi, M. Madonia and C. Selmi, Bad pictures: Some structural properties related to overlaps, *DCFS 2020, Lect. Notes Comput. Sci.* 12442 (2020), pp. 13–25.
- [11] M. Béal and M. Crochemore, Checking whether a word is Hamming-isometric in linear time, *Theor. Comput. Sci.* 933 (2022) 55–59.
- [12] M. Béal, F. Mignosi and A. Restivo, Minimal forbidden words and symbolic dynamics, STACS 96, 13th Annual Symposium on Theoretical Aspects of Computer Science, Lecture Notes in Computer Science 1046 (1996), pp. 555–566.
- [13] G. Castiglione, M. Flores and D. Giammarresi, Isometric words and edit distance: Main notions and new variations, *Cellular Automata and Discrete Complex Systems. AUTOMATA 2023, Lect. Notes Comput. Sci.* 14152, (Springer, 2023), pp. 6–13.
- [14] G. Castiglione, S. Mantaci and A. Restivo, Some investigations on similarity measures based on absent words, *Fundam. Informaticae* 171(1-4) (2020) 97–112.
- [15] P. Charalampopoulos, M. Crochemore, G. Fici, R. Mercas and S. P. Pissis, Alignment-free sequence comparison using absent words, *Inf. Comput.* 262 (2018) 57–68.
- [16] Y. Dombb, O. Lipsky, B. Porat, E. Porat and A. Tsur, The approximate swap and mismatch edit distance, *Theor. Comput. Sci.* 411(43) (2010) 3814–3822.

- [17] A. El-Amawy and S. Latifi, Properties and performance of folded hypercubes, *IEEE Transactions on Parallel and Distributed Systems* **2**(1) (1991) 31–42.
- [18] C. Epifanio, A. Gabriele, F. Mignosi, A. Restivo and M. Sciortino, Languages with mismatches, *Theoretical Computer Science* 385(1) (2007) 152–166.
- [19] C. Epifanio, L. Forlizzi, F. Marzi, F. Mignosi, G. Placidi and M. Spezialetti, On the k-hamming and k-edit distances *CEUR Workshop Proceedings* **3587** (2023), p. 143 – 156.
- [20] S. Faro and A. Pavone, An efficient skip-search approach to swap matching, *Comput. J.* 61(9) (2018) 1351–1360.
- [21] Z. Galil and K. Park, An improved algorithm for approximate string matching., *SIAM J. Comput.* **19** (01 1990) 989–999.
- [22] W. Hsu, Fibonacci cubes-a new interconnection topology, *IEEE Transactions on Parallel and Distributed Systems* **4**(1) (1993) 3–12.
- [23] A. Ilić, S. Klavžar and Y. Rho, Generalized Fibonacci cubes, *Discrete Math.* 312(1) (2012) 2–11.
- [24] A. Ilić, S. Klavžar and Y. Rho, The index of a binary word, *Theor. Comput. Sci.* 452 (2012) 100–106.
- [25] S. Klavžar and S. V. Shpectorov, Asymptotic number of isometric generalized Fibonacci cubes, *Eur. J. Comb.* **33**(2) (2012) 220–226.
- [26] V. I. Levenshtein, Binary codes capable of correcting deletions, insertions, and reversals, *Cybern Control Theory* **10** (1966) 707–710.
- [27] N. Tzeng and S. Wei, Enhanced hypercubes, *IEEE Trans. Computers* **40**(3)(1991) 284–294.
- [28] R. A. Wagner, On the complexity of the extended string-to-string correction problem, *Proceedings of the 7th Annual ACM Symposium on Theory of Computing, May 5-7, 1975, Albuquerque, New Mexico, USA*, eds. W. C. Rounds, N. Martin, J. W. Carlyle and M. A. Harrison (1975), pp. 218–223.
- [29] J. Wei, The structures of bad words, Eur. J. Comb. 59 (2017) 204-214.
- [30] J. Wei, Y. Yang and X. Zhu, A characterization of non-isometric binary words, *Eur. J. Comb.* 78 (2019) 121–133.
- [31] J. Wei and H. Zhang, Proofs of two conjectures on generalized Fibonacci cubes, *Eur. J. Comb.* 51 (2016) 419 – 432.