

HOPF ALGEBRAS, STEINBERG MODULES, AND THE UNSTABLE COHOMOLOGY OF $\mathrm{SL}_n(\mathbb{Z})$ AND $\mathrm{GL}_n(\mathbb{Z})$

AVNER ASH, JEREMY MILLER, AND PETER PATZT

ABSTRACT. We prove that the direct sum of all homology groups of the integral general linear groups with Steinberg module coefficients form a commutative Hopf algebra, in particular a free graded commutative algebra. We use this to construct new infinite families of unstable cohomology classes of $\mathrm{SL}_n(\mathbb{Z})$.

1. INTRODUCTION

The goal of this paper is to prove a new structural result on the rational cohomology of general and special linear groups over the integers. Let $\mathrm{St}_n \mathbb{Q}$ denote the n th Steinberg module of the rationals (top degree reduced homology of the poset of subspaces of \mathbb{Q}^n) and \det denote the determinant representation (unless stated otherwise, all homology will be taken with coefficients in a field of characteristic zero). By work of Borel–Serre [BS73] (see also Putman–Studenmund [PS22]), the twisted homology groups

$$H_*(\mathrm{GL}_n \mathbb{Z}; \mathrm{St}_n \mathbb{Q}) \text{ and } H_*(\mathrm{GL}_n \mathbb{Z}; \mathrm{St}_n \mathbb{Q} \otimes \det)$$

determine the cohomology of $\mathrm{SL}_n(\mathbb{Z})$ and $\mathrm{GL}_n(\mathbb{Z})$:

- $H^{(n)}_{(2)} - k(\mathrm{SL}_n \mathbb{Z}) \cong H_k(\mathrm{GL}_n \mathbb{Z}; \mathrm{St}_n \mathbb{Q}) \oplus H_k(\mathrm{GL}_n \mathbb{Z}; \mathrm{St}_n \mathbb{Q} \otimes \det)$ for $n > 0$,
- $H^{(n)}_{(2)} - k(\mathrm{GL}_n \mathbb{Z}) \cong H_k(\mathrm{GL}_n \mathbb{Z}; \mathrm{St}_n \mathbb{Q})$ for n odd,
- $H^{(n)}_{(2)} - k(\mathrm{GL}_n \mathbb{Z}) \cong H_k(\mathrm{GL}_n \mathbb{Z}; \mathrm{St}_n \mathbb{Q} \otimes \det)$ for n even.

There is a well-known equivariant product structure on the Steinberg modules (see e.g. [MNP20] and [MPW]). In particular, there is a $\mathrm{GL}_n \mathbb{Z} \times \mathrm{GL}_m \mathbb{Z}$ -equivariant map

$$(1) \quad \mathrm{St}_n \mathbb{Q} \otimes \mathrm{St}_m \mathbb{Q} \longrightarrow \mathrm{St}_{n+m} \mathbb{Q}$$

which induces ring structures on

$$H := \bigoplus_{k, n \geq 0} H_k(\mathrm{GL}_n \mathbb{Z}; \mathrm{St}_n \mathbb{Q}) \quad \text{and} \quad H^{\det} := \bigoplus_{k, n \geq 0} H_k(\mathrm{GL}_n \mathbb{Z}; \mathrm{St}_n \mathbb{Q} \otimes \det).$$

We define coproduct structures on these rings which make them into Hopf algebras.

Theorem A. *H and H^{\det} have the structure of graded commutative Hopf algebras.*

Using a theorem of Leray, we deduce the following.

Corollary 1.1. *H and H^{\det} are free graded commutative algebras.*

Our freeness result lets us easily produce new nonzero classes in the cohomology of $\mathrm{SL}_n \mathbb{Z}$ and $\mathrm{GL}_n \mathbb{Z}$ simply by multiplying classes that were previously known to be nonzero. Moreover, using elementary results on Hopf algebras, one can check algebraic independence of cohomology classes using the coproduct. We apply our theory to the so-called wheel classes. Brown–Schnetz [BS, Theorem 4(i)] showed that there are nonzero classes

$$w_n \in H_n(\mathrm{GL}_n \mathbb{Z}; \mathrm{St}_n \mathbb{Q}) \cong H^{(n)}_{(2)} - n(\mathrm{GL}_n \mathbb{Z}) \subset H^{(n)}_{(2)} - n(\mathrm{SL}_n \mathbb{Z})$$

for every odd $n \geq 3$. For odd $n \geq 1$, let

$$t_n \in H^{(n)}_{(2)}(\mathrm{GL}_n \mathbb{Z}; \mathrm{St}_n \mathbb{Q}) \cong H^0(\mathrm{GL}_n \mathbb{Z})$$

Date: April 23, 2024.

Jeremy Miller was supported by NSF Grant DMS-2202943.

Peter Patzt was supported by a Simons Foundation Collaboration Grant.

be a generator of the one-dimensional vector space. (It turns out that t_3 and w_3 both generate the one-dimensional vector space $H_3(\mathrm{GL}_3 \mathbb{Z}; \mathrm{St}_3 \mathbb{Q})$.) And let, for even $n \geq 2$,

$$t_n \in H_{\binom{n}{2}}(\mathrm{GL}_n \mathbb{Z}; \mathrm{St}_n \mathbb{Q} \otimes \det) \cong H^0(\mathrm{GL}_n \mathbb{Z})$$

be a generator of the one-dimensional vector space.

Theorem B. *The free graded commutative algebra*

$$\bigwedge[t_1, t_5, t_9, \dots] \otimes \mathrm{Sym}[t_7, t_{11}, \dots] \otimes \mathrm{Sym}[w_3, w_5, \dots]$$

embeds as an algebra into H and the free graded commutative algebra

$$\bigwedge[t_2, t_6, t_{10}, \dots] \otimes \mathrm{Sym}[t_4, t_8, \dots]$$

embeds as an algebra into H^{\det} .

The cohomology $H^*(\mathrm{SL}_n(\mathbb{Z}))$ has been completely computed for $n \leq 7$ (see [Sou10, Hor14, EVGS13]). Using Theorem B, we describe bases of the vector spaces $H^k(\mathrm{SL}_n(\mathbb{Z}))$ for $n \leq 7$ in terms of the wheel classes, trivial classes, and the Borel classes $\sigma_5 \in H^5(\mathrm{SL}_6 \mathbb{Z})$, $\sigma_5 \in H^5(\mathrm{SL}_7 \mathbb{Z})$, and $\sigma_9 \in H^9(\mathrm{SL}_7 \mathbb{Z})$, which are known to be nonzero (see [Bor74, Fra08, GKT21, Bro]):




FIGURE 1. $H_k(\mathrm{SL}_n \mathbb{Z}; \mathrm{St}_n \mathbb{Q})$

Remark 1.2. Ash [Ash] recently proved that t_1 and w_3 generates a free graded commutative algebra inside of H . His paper inspired this collaboration and our coproducts on H and H^{\det} come from his product on cosharblies.

Remark 1.3. After a draft of this paper had been circulated, we learned of a forthcoming paper by F. Brown, M. Chan, S. Galatius, and S. Payne where they independently constructed a commutative Hopf algebra structure on H and use it to deduce many properties of the cohomology of $\mathrm{SL}_n(\mathbb{Z})$.

Remark 1.4. The bialgebra structure on H is the shadow of a bialgebra structure on the Steinberg modules themselves. Steinberg modules assemble to form a representation of the groupoid $\bigsqcup_n \mathrm{GL}_n(\mathbb{Z})$. This category of representations carries two monoidal structures, one often called the Day convolution product considered for example in [MNP20] and one based on parabolic induction considered for example by Nagpal [Nag19]. These monoidal structures are compatible in the sense that they form a duoidal category. The Steinberg modules form an algebra with respect to the first product, a coalgebra with respect to the second product, and bialgebra in a duoidal sense. For the definitions of a duoidal category and a bialgebra therein, see Böhm–Chen–Zhang [BCZ13]. To keep this paper less abstract, we will only work with classical Hopf algebras. However, we plan to revisit these ideas to study the unstable cohomology of congruence subgroups.

Remark 1.5. Free rings are determined by their indecomposables so it is natural to ask for a description of the indecomposables of H . It follows from [MPW, Remark 1.6] that these indecomposables are isomorphic to the reduced equivariant homology of Rognes’ common basis complex [Rog92].

Acknowledgements. We would like to thank Francis Brown, Benjamin Brück, Melody Chan, Søren Galatius, Rohit Nagpal, Sam Payne, Andrew Snowden, and Robin Sroka for helpful conversations.

2. A DIFFERENTIAL GRADED BIALGEBRA THAT COMPUTES STEINBERG HOMOLOGY

Let R be a PID and let F be its field of fractions. Further, let K be our coefficient ring, a field of characteristic zero.

The *Tits building* of F^n denoted by $\mathcal{T}_n(F)$ is the simplicial complex whose vertices are nonzero proper subspaces $0 \subsetneq V \subsetneq F^n$ and $\{V_0, \dots, V_p\}$ is a simplex if

$$V_0 \subset \dots \subset V_p.$$

The Solomon–Tits Theorem [Sol69] implies that $\mathcal{T}_n(F)$ is a wedge of $(n-2)$ -dimensional spheres and its only nonzero reduced homology group is called the Steinberg module

$$\mathrm{St}_n(F) := \tilde{H}_{n-2}(\mathcal{T}_n(F); K).$$

By convention, we define $\mathrm{St}_0(F)$ to be K .

Lee–Szczarba [LS76] gave a flat $K \mathrm{GL}_n(R)$ -resolution of $\mathrm{St}_n(F)$. We use a slight variant of it here, which can be established with the same proof. See also Ash–Gunnells–McConnell [AGM12]. Let V_n be the set of all nonzero vectors in $R^n \subset F^n$ and V_n^\pm be the quotient of V_n identifying negatives. By abuse of notation, we denote an element of V_n^\pm simply by its representative $v \in R^n$. Let X_n be the simplicial complex whose vertices are the elements of V_n^\pm and all (finite) sets of vertices are simplices. Let L_n the subcomplex of all simplices $\{v_0, \dots, v_p\}$ such that the F -span of v_0, \dots, v_p is not all of F^n . Clearly, X_n is contractible and using Lee–Szczarba’s argument it follows that L_n is homotopy equivalent to $\mathcal{T}_n(F)$, which implies that

$$H_i(X_n, L_n; K) \cong \tilde{H}_{i-1}(L_n; K) \cong \begin{cases} \mathrm{St}_n(F) & i = n-1 \\ 0 & i \neq n-1 \end{cases}.$$

Because $C_i(X_n, L_n; K) = 0$ for $i < n-1$, we get an exact sequence

$$\dots \longrightarrow C_n(X_n, L_n; K) \longrightarrow C_{n-1}(X_n, L_n; K) \longrightarrow \mathrm{St}_n(F) \longrightarrow 0.$$

[CP17, Lemma 3.2] implies that $C_i(X_n, L_n; K)$ is a flat $K \mathrm{GL}_n(R)$ -module.

We will use the notation

$$\text{Sh}_{n,k} = C_{n+k-1}(X_n, L_n; K)$$

and refer to $\text{Sh}_{n,*} \rightarrow \text{St}_n(F)$ as the *Sharbly resolution* of $\text{St}_n(F)$. Given nonzero vectors $v_1, \dots, v_{n+k} \in R^n$, we get a canonical element

$$[v_1, \dots, v_{n+k}] \in \text{Sh}_{n,k}$$

given by the corresponding simplex $\{v_1, \dots, v_{n+k}\}$ in X_n . These elements are nonzero if and only if the F -span of v_1, \dots, v_{n+k} is F^n . We call such an element then a *basic sharbly*.

Let $\chi: R^\times \rightarrow K^\times$ be a character (i.e. a group homomorphism), which we will view as a one-dimensional $K \text{GL}_n(R)$ -module that factors through the determinant. We will use the notation

$$[v_1, \dots, v_{n+k}]^\chi := [v_1, \dots, v_{n+k}] \otimes 1 \in \text{Sh}_{n,k}^\chi := \text{Sh}_{n,k} \otimes_K \chi.$$

We will further consider the $\text{GL}_n(R)$ -coinvariants of $\text{Sh}_{n,k}^\chi$ and call its elements *coinvariant sharblies*. In particular, we will denote the image of a basic sharbly $[v_1, \dots, v_{n+k}]^\chi \in \text{Sh}_{n,k}^\chi$ by

$$[v_1, \dots, v_{n+k}]_{\text{GL}_n(R)}^\chi \in (\text{Sh}_{n,k}^\chi)_{\text{GL}_n(R)}$$

and call it a *basic coinvariant sharbly*. Because $\text{GL}_n(R)$ acts on the set of all basic sharblies, which up to signs¹ form a basis of $\text{Sh}_{n,k}^\chi$, the set of nonzero basic coinvariant sharblies form a basis up to signs.

Let us denote by

$$B_{n,k}^\chi := (\text{Sh}_{n,k}^\chi)_{\text{GL}_n(R)} \quad \text{and} \quad B_m^\chi := \bigoplus_{n+k=m} B_{n,k}^\chi$$

the coinvariants of the sharblies and by

$$B^\chi := \bigoplus_{m \geq 0} B_m^\chi = \bigoplus_{n,k \geq 0} B_{n,k}^\chi$$

the (bi-)graded vector space of all coinvariant sharblies. It comes with a differential

$$\begin{aligned} \partial: B_{n,k}^\chi &= (\text{Sh}_{n,k}^\chi)_{\text{GL}_n(R)} \longrightarrow (\text{Sh}_{n,k-1}^\chi)_{\text{GL}_n(R)} = B_{n,k-1}^\chi \\ [v_1, \dots, v_{n+k}]_{\text{GL}_n(R)}^\chi &\longmapsto \sum_{i=1}^{n+k} (-1)^i [v_1, \dots, \hat{v}_i, \dots, v_{n+k}]_{\text{GL}_n(R)}^\chi \end{aligned}$$

of degree -1 , such that its homology is

$$H(B^\chi)_{n,k} \cong H_k(\text{GL}_n(R); \text{St}_n(F) \otimes_K \chi).$$

Definition 2.1. We define

$$\nabla: B_{n,k}^\chi \otimes B_{m,l}^\chi \longrightarrow B_{n+m,k+l}^\chi$$

by sending a tensor product of basic coinvariant sharblies $x = [v_1, \dots, v_{n+k}]_{\text{GL}_n(R)}^\chi$ and $y = [w_1, \dots, w_{m+l}]_{\text{GL}_m(R)}^\chi$ to

$$\nabla(x \otimes y) = xy = [\phi_{n,n+m}(v_1), \dots, \phi_{n,n+m}(v_{n+k}), \psi_{m,n+m}(w_1), \dots, \psi_{m,n+m}(w_{m+l})]_{\text{GL}_{n+m}(R)}^\chi,$$

where $\phi_{n,n+m}: R^n \rightarrow R^{n+m}$ sends the standard basis vector $e_i \in R^n$ to $e_i \in R^{n+m}$ and $\psi_{m,n+m}: R^m \rightarrow R^{n+m}$ sends $e_i \in R^m$ to $e_{n+i} \in R^{n+m}$.

The following lemma is easy to check.

Lemma 2.2. $\nabla: B^\chi \otimes B^\chi \rightarrow B^\chi$ defines a (bi-)graded algebra.

We call $\chi: R^\times \rightarrow K^\times$ *odd* if $\chi(-1) = -1$ and *even* if $\chi(-1) = 1$.

Lemma 2.3. If χ is odd and n is odd, then $B_{n,k}^\chi = 0$.

Proof. Let $x = [v_1, \dots, v_{n+k}]^\chi$ be a basic sharbly in $\text{Sh}_{n,k}^\chi$. Then

$$(-\text{id}) \cdot x = \chi(-1)^n \cdot [-v_1, \dots, -v_{n+k}]^\chi = -[v_1, \dots, v_{n+k}]^\chi = -x$$

and thus vanishes in the coinvariants $B_{n,k}^\chi$. \square

¹Up to signs” means that while $[v_1, \dots, v_{n+k}]^\chi$ is antisymmetric, if we choose a representative from each S_{n+k} -orbit of $(n+k)$ -tuples of vectors, we get a basis of $\text{Sh}_{n,k}^\chi$.

Lemma 2.4. B^χ is graded commutative, i.e. for $x \in B_{n,k}^\chi$ and $y \in B_{m,l}^\chi$, we get

$$xy = (-1)^{(n+k)(m+l)}yx.$$

Proof. Let $x = [v_1, \dots, v_{n+k}]_{GL_n(R)}^\chi$ and $y = [w_1, \dots, w_{m+l}]_{GL_m(R)}^\chi$ be basic coinvariant sharblies. We observe that

$$\begin{aligned} xy &= [\phi_{n,n+m}(v_1), \dots, \phi_{n,n+m}(v_{n+k}), \psi_{m,n+m}(w_1), \dots, \psi_{m,n+m}(w_{m+l})]_{GL_{n+m}(R)}^\chi \\ &= (-1)^{(n+k)(m+l)} [\psi_{m,n+m}(w_1), \dots, \psi_{m,n+m}(w_{m+l}), \phi_{n,n+m}(v_1), \dots, \phi_{n,n+m}(v_{n+k})]_{GL_{n+m}(R)}^\chi. \end{aligned}$$

Let $T_{n,m}$ be the permutation matrix sending e_i to e_{m+i} if $i \leq n$ and to e_{i-n} if $i > n$. We can use $T_{n,m} \in GL_{n+m}(R)$ to swap ϕ and ψ in the sense that

$$T_{n,m}\phi_{n,n+m} = \psi_{n,n+m} \quad \text{and} \quad T_{n,m}\psi_{m,n+m} = \phi_{m,n+m}.$$

The determinant of $T_{n,m}$ is $(-1)^{nm}$, so in particular 1 if n or m are even. We then get that

$$\begin{aligned} &[\psi_{m,n+m}(w_1), \dots, \psi_{m,n+m}(w_{m+l}), \phi_{n,n+m}(v_1), \dots, \phi_{n,n+m}(v_{n+k})]_{GL_{n+m}(R)}^\chi \\ &= T_{n,m}\chi(-1)^{nm}[\phi_{m,n+m}(w_1), \dots, \phi_{m,n+m}(w_{m+l}), \psi_{n,n+m}(v_1), \dots, \psi_{n,n+m}(v_{n+k})]_{GL_{n+m}(R)}^\chi \\ &= T_{n,m}\chi(-1)^{nm}yx. \end{aligned}$$

Therefore

$$xy = (-1)^{(n+k)(m+l)} \cdot \chi(-1)^{nm} \cdot yx$$

in the coinvariants $B_{n+m,k+l}^\chi$. The factor $\chi(-1)^{nm}$ is of course 1 if $\chi(-1) = 1$, and it is also 1 if $\chi(-1) = -1$ because then nm is even or $xy = yx = 0$. \square

Lemma 2.5. B^χ together with the differential ∂ is a differential graded algebra.

Proof. Given basic coinvariant sharblies $x = [v_1, \dots, v_{n+k}]_{GL_n(R)}^\chi$ and $y = [w_1, \dots, w_{m+l}]_{GL_m(R)}^\chi$, we need to prove that

$$\partial(xy) = (\partial x)y + (-1)^{n+k}x(\partial y).$$

This follows as

$$\begin{aligned} &\partial(xy) \\ &= \sum_{i=1}^{n+k} (-1)^i [\phi_{n,n+m}(v_1), \dots, \widehat{\phi_{n,n+m}(v_i)}, \dots, \phi_{n,n+m}(v_{n+k}), \psi_{m,n+m}(w_1), \dots, \psi_{m,n+m}(w_{m+l})]_{GL_{n+m}(R)}^\chi \\ &\quad + (-1)^{n+k} \sum_{i=1}^{m+l} (-1)^i [\phi_{n,n+m}(v_1), \dots, \phi_{n,n+m}(v_{n+k}), \psi_{m,n+m}(w_1), \dots, \widehat{\psi_{m,n+m}(w_i)}, \dots, \psi_{m,n+m}(w_{m+l})]_{GL_{n+m}(R)}^\chi \\ &= (\partial x)y + (-1)^{n+k}x(\partial y). \end{aligned} \quad \square$$

We now want to describe the coalgebra structure on B^χ . We will actually first describe it before taking coinvariants.

Definition 2.6. Let U be an a -dimensional subspace of F^n . Let $X(U)$ be the simplicial complex whose vertices are

$$V^\pm(U) := \{v \in V_n^\pm \mid v \in U\}$$

and every finite subset forms a face. Let $L(U)$ the subcomplex of $X(U)$ of all simplices $\{v_0, \dots, v_p\}$ such that the F -span of v_0, \dots, v_p is not all of U . We define

$$\text{Sh}_s^\chi(U) := C_{a+s-1}(X(U), L(U); K).$$

Note that a basis of the free R -module $U \cap R^n$ gives an isomorphism $\text{Sh}_*^\chi(U) \cong \text{Sh}_{a,*}^\chi$ of chain complexes.

Further, let W be a b -dimensional subspace of U . We define $\text{Sh}_s^\chi(U/W)$ as the quotient of $\text{Sh}_{s-b}^\chi(U)$ by identifying basic sharblies $[v_1, \dots, v_{a+(s-b)}]^\chi$ and $[w_1, \dots, w_{a+(s-b)}]^\chi$ in $\text{Sh}_{s-b}^\chi(U)$ if $v_i - w_i \in W$ for all i . We also identify a basic sharbly $[v_1, \dots, v_{a+(s-b)}]^\chi$ with zero if $v_i \in W$ for some i or if $v_i - v_j \in W$ for some $i \neq j$. Note that a basis of the free R -module $(U \cap R^n)/(W \cap R^n)$ gives an isomorphism $\text{Sh}_*^\chi(U/W) \cong \text{Sh}_{a-b,*}^\chi$ of chain complexes.

Definition 2.7. Let $W \subseteq U \subseteq F^n$ be subspaces of F^n . We define a map

$$\Delta: \text{Sh}_k^X(U/W) \longrightarrow \bigoplus_{\substack{W \subseteq V \subseteq U \\ p+q=k}} \text{Sh}_p^X(V/W) \otimes \text{Sh}_q^X(U/V)$$

in the following way.

Let $x = [v_1, \dots, v_{\dim U - \dim W + k}]^X \in \text{Sh}_k^X(U/W)$ be a basic sharply. We define \mathcal{U}_x to be the set of subspaces $V \subseteq U$ that are spanned by W and a subset of $\{v_1, \dots, v_{\dim U - \dim W + k}\}$. (W and U are necessarily included in \mathcal{U}_x .) Let $[m]$ denote the set $\{1, \dots, m\}$. For $V \in \mathcal{U}_x$, let

$$S(V) = \{i \in [\dim U - \dim W + k] \mid v_i \in V\}.$$

Let $s_1 < \dots < s_{\dim V - \dim W + p}$ be the elements of $S(V)$ and $t_1 < \dots < t_{\dim U - \dim V + q}$ the elements of $[a + k] \setminus S(V)$.

We set

$$x'_V = [v_{s_1}, \dots, v_{s_{\dim V - \dim W + p}}]^X \in \text{Sh}_p^X(V/W)$$

and

$$x''_V = [v_{t_1}, \dots, v_{t_{\dim U - \dim V + q}}]^X \in \text{Sh}_q^X(U/V).$$

We define

$$\Delta(x) = \sum_{V \in \mathcal{U}_x} (-1)^{\pi_{V/W \subset U/W}} x'_V \otimes x''_V,$$

where $\pi_{V/W \subset U/W}$ is the shuffle permutation sending i to s_i if $i \leq \dim V - \dim W + p$ and to $t_{i - \dim W - p}$ if $i > \dim V - \dim W + p$. If $U = F^n$ and $W = 0$, we will sometimes simply write π_V for $\pi_{V/W \subset F^n/W}$.

Lemma 2.8. *The map*

$$\Delta: \text{Sh}_{n,k}^X \longrightarrow \bigoplus_{\substack{U \subset F^n \\ p+q=k}} \text{Sh}_p^X(U) \otimes \text{Sh}_q^X(F^n/U)$$

is $\text{GL}_n(R)$ -equivariant and the $\text{GL}_n(R)$ -coinvariants of it give a map

$$\Delta: B_{n,k}^X \longrightarrow \bigoplus_{\substack{a+b=n \\ p+q=k}} B_{a,p}^X \otimes B_{b,q}^X.$$

Proof. For the equivariance it is important to note that $\text{GL}_n(R)$ permutes the summands by acting on the set of subspaces $U \subset F^n$.

The coinvariants on the left are $B_{n,k}^X$ by definition. On the right, we first note that $\text{GL}_n(R)$ acts transitively on the set of a -dimensional $U \subset F^n$. (Here we use that R is a PID.) Let $P_{a,b}$ denote the stabilizer of $F^a \subset F^n$ in $\text{GL}_n(R)$, where $b = n - a$.

There is a canonical isomorphism of $\text{Sh}_p^X(F^a) \rightarrow \text{Sh}_{a,p}^X$. There is also a canonical isomorphism of $\text{Sh}_q^X(F^n/F^a) \rightarrow \text{Sh}_{b,q}^X$ by only recording the last q coordinates of the representatives of vectors. $P_{a,b}$ acts on

$$\text{Sh}_p^X(F^a) \otimes \text{Sh}_q^X(F^n/F^a) \cong \text{Sh}_{a,p}^X \otimes \text{Sh}_{b,q}^X$$

via the projection onto its Levi factor: $P_{a,b} \rightarrow \text{GL}_a(R) \times \text{GL}_b(R)$. And that means that the coinvariants on the right-hand-side are isomorphic to

$$\bigoplus_{\substack{a+b=n \\ p+q=k}} B_{a,p}^X \otimes B_{b,q}^X. \quad \square$$

We get that B^X is a coalgebra by the following lemma.

Lemma 2.9. $\Delta: B^X \rightarrow B^X \otimes B^X$ defines a (bi-)graded coalgebra.

Proof. We will only check coassociativity here and leave all other properties to the reader. In fact, we check coassociativity before taking coinvariants.

Let $x = [v_1, \dots, v_{n+k}]^X \in \text{Sh}_{n,k}^X$ a basic sharply. We compute that

$$(\Delta \otimes \text{id}) \circ \Delta(x) = \sum_{\substack{W, U \in \mathcal{U}_x \\ W \subseteq U}} (-1)^{\pi_{U \subset F^n}} ((-1)^{\pi_{W \subset U}} (x'_U)'_W \otimes (x'_U)''_W) \otimes x''_U,$$

where $(x'_U)'_W \in \text{Sh}_a^X(W)$, $(x'_U)''_W \in \text{Sh}_b^X(U/W)$, and $x''_U \in \text{Sh}_c^X(F^n/U)$ for corresponding $a + b + c = k$.

We also compute that

$$(\text{id} \otimes \Delta) \circ \Delta(x) = \sum_{\substack{W, U \in \mathcal{U}_x \\ W \subseteq U}} (-1)^{\pi_{W \subset F^n}} x'_W \otimes ((-1)^{\pi_{U/W \subset F^n/W}} (x''_W)'_{U/W} \otimes (x''_W)''_{U/W}),$$

where $x'_W \in \text{Sh}_a^X(W)$, $(x''_W)'_U \in \text{Sh}_b^X(U/W)$, and $(x''_W)''_U \in \text{Sh}_c^X(F^n/U)$ for corresponding $a + b + c = k$.

It is now easy to see that

$$(x'_U)'_W = x'_W, \quad (x'_U)''_W = (x''_W)'_U, \quad x''_U = (x''_W)''_U, \text{ and } \pi_{W \subset U} \circ \pi_{U \subset F^n} = \pi_{U/W \subset F^n/W} \circ \pi_{W \subset F^n}. \quad \square$$

Let us check that it behaves well with the differential.

Lemma 2.10. *B^X together with the differential ∂ is a differential graded coalgebra.*

Proof. Given a basic sharply $x = [v_1, \dots, v_{n+k}]^X \in \text{Sh}_{n,k}^X$, we need to check that

$$\Delta(\partial x) = \sum_{U \in \mathcal{U}_x} (-1)^{\pi_U} (\partial x'_U \otimes x''_U + (-1)^{|S(U)|} x'_U \otimes \partial x''_U).$$

We want to show that both sides of this equation are equal to the sum

$$\sum_{\substack{U \in \mathcal{U}_x \\ i \in [n+k]}} (-1)^{\pi_U} \cdot (-1)^{\pi_U^{-1}(i)} d_i(x'_U \otimes x''_U),$$

where d_i omits the vector v_i from x'_U if $v_i \in U$ and from x''_U if $v_i \notin U$.

On the right-hand-side, we can index the summands by $U \in \mathcal{U}_x$ and the v_i that is being omitted. As for the sign, note that $\pi_U^{-1}(i)$ is the position of v_i in x'_U if $v_i \in U$ and the position of v_i in x''_U plus $|S(U)|$ if $v_i \notin U$.

For the left-hand-side, let us compute $\Delta(x_i)$ for every

$$x_i = [v_1, \dots, \hat{v}_i, \dots, v_{n+k}]^X.$$

This is

$$\Delta(x_i) = \sum_{U \in \mathcal{U}_{x_i}} (-1)^{\pi_{U,i}} (x'_i)_U \otimes (x''_i)_U,$$

where $\pi_{U,i}$ is the shuffle permutation obtained from π_U by removing the strand that comes from i . That means we get a summand $d_i(x'_U \otimes x''_U)$ for every pair (U, i) such that $U \in \mathcal{U}_{x_i}$. For these summands, the sign of $\pi_{U,i}$ is equal to $(-1)^{|i - \pi_U^{-1}(i)|}$.

We finish the proof by observing that if $U \in \mathcal{U}_x \setminus \mathcal{U}_{x_i}$, then $d_i(x'_U \otimes x''_U) = 0$. This follows because then $v_i \in U$ and the vectors v_s with $s \in S(U) \setminus \{i\}$ do not span U without v_i . \square

We conclude this section by proving that the product and the coproduct are compatible.

Lemma 2.11. *The diagram*

$$\begin{array}{ccccc} B^X \otimes B^X & \xrightarrow{\nabla} & B^X & \xrightarrow{\Delta} & B^X \otimes B^X \\ \Delta \otimes \Delta \downarrow & & & & \uparrow \nabla \otimes \nabla \\ B^X \otimes B^X \otimes B^X \otimes B^X & \xrightarrow{\text{id} \otimes \tau \otimes \text{id}} & B^X \otimes B^X \otimes B^X \otimes B^X & & \end{array}$$

commutes. Here $\tau: B_n^X \otimes B_m^X \rightarrow B_m^X \otimes B_n^X$ is given by

$$\tau(x \otimes y) = (-1)^{nm}(y \otimes x).$$

Proof. We evaluate both paths on $x \otimes y$ for basic sharblies $x = [v_1, \dots, v_{n+k}]^X \in \text{Sh}_{n,k}^X$ and $y = [w_1, \dots, w_{m+l}]^X \in \text{Sh}_{m,l}^X$.

For the lower path we get that $x \otimes y$ is sent to

$$\sum_{\substack{U \in \mathcal{U}_x \\ V \in \mathcal{U}_y}} (-1)^{pq} (-1)^{\pi_U} (-1)^{\pi_V} x'_U y'_V \otimes x''_U y''_V,$$

where p is the number of vectors in x''_U and q the number of vectors in y'_V . We note that we consider the product $x'_U y'_V$ to be in

$$\mathrm{Sh}_{a+c}^x(\phi(U) \oplus \psi(V)) \subset \mathrm{Sh}_{n+m, a+c}^x,$$

where $a = |S(U)| - \dim U$ and $c = |S(V)| - \dim V$. And we consider the product $x''_U y''_V$ to be in the quotient

$$\mathrm{Sh}_{b+d}^x(F^{n+m}/(\phi(U) \oplus \psi(V)))$$

of $\mathrm{Sh}_{n+m, b+d}^x$, where $b = k - a$ and $d = l - c$.

For the upper path, we get that $x \otimes y$ is sent to

$$\sum_{W \in \mathcal{U}_{xy}} (-1)^{\pi_W} (xy)'_W \otimes (xy)''_W.$$

To see that these two sums are the same, we first observe that every $W \in \mathcal{U}_{xy}$ uniquely decomposes into a direct sum $\phi(U) \oplus \psi(V) = W$ for pairs $(U, V) \in \mathcal{U}_x \times \mathcal{U}_y$.

It remains to show that the corresponding summands agree. For this note that

$$(xy)'_W = [\phi(v_{s_1}), \dots, \phi(v_{s_{\dim U+a}}), \psi(w_{\eta_1}), \dots, \psi(w_{\eta_{\dim V+c}})] \in \mathrm{Sh}_{a+c}^x(W),$$

where $S(U) = \{s_1 < \dots < s_{\dim U+a}\}$ and $S(V) = \{\eta_1 < \dots < \eta_{\dim V+c}\}$ which is the same as

$$x'_U y'_V = [\phi(v_{s_1}), \dots, \phi(v_{s_{\dim U+a}}), \psi(w_{\eta_1}), \dots, \psi(w_{\eta_{\dim V+c}})] \in \mathrm{Sh}_{a+c}^x(\phi(U) \oplus \psi(V)).$$

And

$$(xy)''_W = [\phi(v_{t_1}), \dots, \phi(v_{t_{n-\dim U+b}}), \psi(w_{\theta_1}), \dots, \psi(w_{\theta_{m-\dim V+b}})] \in \mathrm{Sh}_{b+d}^x(F^{n+m}/W),$$

where $[n+k] \setminus S(U) = \{t_1 < \dots < t_{n-\dim U+b}\}$ and $[m+l] \setminus S(V) = \{\theta_1 < \dots < \theta_{m-\dim V+b}\}$, which is the same as

$$x''_U y''_V = [\phi(v_{t_1}), \dots, \phi(v_{t_{n-\dim U+b}}), \psi(w_{\theta_1}), \dots, \psi(w_{\theta_{m-\dim V+b}})] \in \mathrm{Sh}_{b+d}^x(\phi(F^n/U) \oplus \psi(F^m/V)).$$

Finally, we need to compare the signs. Let us consider the two permutations π_U and π_V as one permutation $\pi_{U,V}$ on $n+k+m+l$ letters. We note that π_W is $\pi_{U,V}$ postcomposing with the permutation that swaps the number of vectors in x''_U , which we denoted by p , and the number of vectors in y'_V , which we denoted by q , with each other. Thus

$$(-1)^{\pi_W} = (-1)^{pq} (-1)^{\pi_U} (-1)^{\pi_V}. \quad \square$$

The result of this section is summarized in the following theorem.

Theorem 2.12. *For every PID R and every character $\chi: R^\times \rightarrow \mathbb{C}^\times$,*

$$B^\chi = \bigoplus_{n,k \geq 0} (\mathrm{Sh}_{n,k}^x \otimes_K \chi)_{\mathrm{GL}_n(R)}$$

is a differential graded commutative bialgebra.

We get the following corollary as an immediate consequence.

Corollary 2.13. *For every PID R and every character $\chi: R^\times \rightarrow \mathbb{C}^\times$,*

$$H(B^\chi) \cong \bigoplus_{n,k \geq 0} H_k(\mathrm{GL}_n(R); \mathrm{St}_n(F) \otimes_K \chi)$$

is a graded commutative bialgebra.

Remark 2.14. Let R be a PID with a finite group of units R^\times , for example $R = \mathbb{Z}$ or R a complex quadratic number ring which is a PID. And let K be a field of characteristic zero over which the R^\times -representations split into one-dimensional irreducibles, for example \mathbb{C} . Then for $n \geq 1$,

$$H_k(\mathrm{SL}_n(R); \mathrm{St}_n(F)) \cong H_k(\mathrm{GL}_n(R); \mathrm{St}_n(F) \otimes \chi_1) \oplus \dots \oplus H_k(\mathrm{GL}_n(R); \mathrm{St}_n(F) \otimes \chi_r),$$

where the regular representation $KR^\times \cong \chi_1 \oplus \dots \oplus \chi_r$.

In fact, Equation (1) induces an algebra structure on

$$\bigoplus_{n,k \geq 0} H_k(\mathrm{SL}_n(R); \mathrm{St}_n(F)),$$

but products between elements of summands corresponding to different characters are zero.

3. HOPF ALGEBRAS

In this section, we recall several definitions and facts about Hopf algebras and then prove a generalization of [Theorem A](#).

A graded bialgebra $B = \bigoplus_{n \geq 0} B_n$ is called *connected* if $B_0 = K$.

Usually, a Hopf algebra is defined to be a bialgebra that has an antipode. We will mainly cite Milnor–Moore [\[MM65\]](#) in this section, who do not require an antipode, i.e. their “Hopf algebras” are simply bialgebras. But it turns out that every connected bialgebra is automatically a Hopf algebra which follows from [\[MM65, Proposition 8.2\]](#). Together with [Corollary 2.13](#), this implies the following generalization of [Theorem A](#).

Theorem 3.1. *For every PID R and every character $\chi: R^\times \rightarrow \mathbb{C}^\times$,*

$$H(B^\chi) \cong \bigoplus_{n,k \geq 0} H_k(\mathrm{GL}_n(R); \mathrm{St}_n(F) \otimes_K \chi)$$

is a graded commutative Hopf algebra.

The *primitives* of a bialgebra B are given by

$$P(B) = \{x \in B \mid \Delta(x) = x \otimes 1 + 1 \otimes x\}.$$

For a connected bialgebra B , let B_+ denote the augmentation ideal $\bigoplus_{n \geq 1} B_n$. The *indecomposables* of B are

$$Q(B) = B_+ / (B_+)^2.$$

Because $P(B) \subset B_+$, there is a canonical map $P(B) \rightarrow Q(B)$.

Proposition 3.2 (Milnor–Moore [\[MM65, Proposition 4.17\]](#)). *A connected Hopf algebra H over a characteristic zero field is commutative if and only if $P(H) \rightarrow Q(H)$ is injective.*

Theorem 3.3 (Leray [\[MM65, Theorem 7.5\]](#)). *A connected graded commutative Hopf algebra H over a characteristic zero field is isomorphic to the free graded commutative algebra generated by $Q(H)$.*

Therefore we get the following generalization of [Corollary 1.1](#).

Corollary 3.4. *For every PID R and every character $\chi: R^\times \rightarrow \mathbb{C}^\times$,*

$$H(B^\chi) \cong \bigoplus_{n,k \geq 0} H_k(\mathrm{GL}_n(R); \mathrm{St}_n(F) \otimes_K \chi)$$

is a free graded commutative algebra.

4. NEW CLASSES

In this section, we take $R = \mathbb{Z}$, $F = \mathbb{Q}$, and $K = \mathbb{Q}$. Let $n \geq 3$. We call the basic coinvariant sharply

$$w_n = [e_1, \dots, e_n, e_1 - e_2, \dots, e_{n-1} - e_n, e_n - e_1]_{\mathrm{GL}_n(\mathbb{Z})} \in (\mathrm{Sh}_{n,n})_{\mathrm{GL}_n(\mathbb{Z})}$$

a *wheel sharply*. It is easy to check that $w_n = 0$ if n is even and that $\partial w_n = 0$ if n is odd. Recently, Brown–Schnetz [\[BS, Theorem 4\(i\)\]](#) proved² that for odd $n \geq 3$, w_n gives rise to a nonzero element in homology

$$H(B)_{n,n} \cong H_n(\mathrm{GL}_n(\mathbb{Z}); \mathrm{St}_n(\mathbb{Q})).$$

We want to prove the following two proposition about them and the trivial classes $t_n \in H^0(\mathrm{GL}_n(\mathbb{Z}))$.

Proposition 4.1. *w_n is primitive in B .*

Proposition 4.2. *t_n is primitive in H or H^{\det} , depending on the parity.*

²They show that a region τ_{W_n} in the space \mathcal{P}_n of positive definite $n \times n$ real matrices gives rise to a nonzero class in the locally finite homology $H_{2n}^{\mathrm{lf}}(\mathcal{P}_n / \mathrm{GL}_n(\mathbb{Z})) \cong H_n(\mathrm{GL}_n(\mathbb{Z}); \mathrm{St}_n(\mathbb{Q}))$. There is a quasi-isomorphism from $(\mathrm{Sh}_{n,*})_{\mathrm{GL}_n(\mathbb{Z})}$ to the locally finite chains of $\mathcal{P}_n / \mathrm{GL}_n(\mathbb{Z})$ defined by sending a basic coinvariant sharply $[v_1, \dots, v_{n+k}]_{\mathrm{GL}_n(\mathbb{Z})}$ to the region in \mathcal{P}_n which is the given by elements $c_1 \cdot v_1 v_1^T + \dots + c_{n+k} \cdot v_{n+k} v_{n+k}^T$ for all positive real numbers $c_1, \dots, c_{n+k} > 0$. Their discussion makes clear that our w_n is sent to their τ_{W_n} .

Because we know that w_n is nonzero in $H(B)$, it follows that it is a nonzero primitive in $H(B)$. Using [Proposition 3.2](#) and [Theorem 3.3](#), [Theorem B](#) follows as an immediate corollary.

Proof of [Proposition 4.1](#). For notational convenience, set $x = w_n$. Recall that we defined the coproduct using the formula

$$\Delta(x) = \sum_{U \in \mathcal{U}_x} (-1)^{\pi_U} x'_U \otimes x''_U.$$

We want to prove that if $U \in \mathcal{U}_x$ is a nonzero proper subspace of F^n , then $x''_U = 0$. Let β denote reduction modulo U .

First assume that none of the e_1, \dots, e_n are in U . Because U is nonzero, without loss of generality it contains $e_1 - e_2$. But that means that $\beta(e_1) = \beta(e_2)$ because $e_1 - e_2$ is in the kernel of β and thus $x''_U = 0$.

We can therefore assume without loss of generality that $e_1 \in U$. Let $k \geq 1$ be so that $e_1, \dots, e_k \in U$ and $e_{k+1} \notin U$. This implies that $e_k - e_{k+1} \notin U$, because $e_k \in U$. But $\beta(e_k - e_{k+1}) = -\beta(e_{k+1})$ because e_k is in the kernel of β . Therefore $x''_U = 0$.

This implies that x''_U is only nonzero if $U = 0$ or $U = F^n$ and we get that

$$\Delta(x) = 1 \otimes x + x \otimes 1. \quad \square$$

Proof of [Proposition 4.2](#). By Borel–Serre [\[BS73\]](#),

$$H_k(\mathrm{SL}_n(\mathbb{Z}); \mathrm{St}_n(\mathbb{Q})) \cong H_k(\mathrm{GL}_n(\mathbb{Z}); \mathrm{St}_n(\mathbb{Q})) \otimes H_k(\mathrm{GL}_n(\mathbb{Z}); \mathrm{St}_n(\mathbb{Q}) \otimes \det) \cong 0$$

for $k > \binom{n}{2}$. And

$$\binom{a}{2} + \binom{b}{2} \leq \binom{a+b}{2}$$

with equality only if a or b is zero. This means that the coproduct $\Delta(t_n)$ of $t_n \in H_{\binom{n}{2}}(\mathrm{GL}_n(\mathbb{Z}); \mathrm{St}_n(\mathbb{Q}) \otimes \chi)$, where χ is trivial if n is odd and the determinant representation if n is even, can only be $1 \otimes t_n + t_n \otimes 1$ for degree reasons. \square

REFERENCES

- [AGM12] Avner Ash, Paul E. Gunnells, and Mark McConnell, *Resolutions of the Steinberg module for $GL(n)$* , J. Algebra **349** (2012), 380–390. [3](#)
- [Ash] A. Ash, *On the cohomology of $\mathrm{SL}_n(\mathbb{Z})$* , Preprint, <https://arxiv.org/abs/2402.08840>. [3](#)
- [BCZ13] Gabriella Böhm, Yuanyuan Chen, and Liangyun Zhang, *On Hopf monoids in duoidal categories*, J. Algebra **394** (2013), 139–172. [3](#)
- [Bor74] A. Borel, *Stable real cohomology of arithmetic groups*, Ann. Sci. École Norm. Sup. (4) **7** (1974), 235–272 (1975). [2](#)
- [Bro] F. Brown, *Bordifications of the moduli spaces of tropical curves and abelian varieties, and unstable cohomology of $\mathrm{GL}_g(\mathbb{Z})$ and $\mathrm{SL}_g(\mathbb{Z})$* , Preprint, <https://arxiv.org/abs/2309.12753>. [2](#)
- [BS] F. Brown and O. Schnetz, *The wheel classes in the locally finite homology of $\mathrm{GL}_n(\mathbb{Z})$, canonical integrals and zeta values*, Preprint, <https://arxiv.org/abs/2402.06757>. [1, 9](#)
- [BS73] A. Borel and J.-P. Serre, *Corners and arithmetic groups*, Comment. Math. Helv. **48** (1973), 436–491, Avec un appendice: Arrondissement des variétés à coins, par A. Douady et L. Hérault. [1, 10](#)
- [CP17] T. Church and A. Putman, *The codimension-one cohomology of $\mathrm{SL}_n\mathbb{Z}$* , Geom. Topol. **21** (2017), no. 2, 999–1032. [3](#)
- [EVGS13] P. Elbaz-Vincent, H. Gangl, and C. Soulé, *Perfect forms, K-theory and the cohomology of modular groups*, Adv. Math. **245** (2013), 587–624. [2](#)
- [Fra08] Jens Franke, *A topological model for some summand of the Eisenstein cohomology of congruence subgroups*, Eisenstein series and applications, Progr. Math., vol. 258, Birkhäuser Boston, Boston, MA, 2008, pp. 27–85. [2](#)
- [GKT21] Jeffrey Giansiracusa, Alexander Kupers, and Bena Tshishiku, *Characteristic classes of bundles of $K3$ manifolds and the Nielsen realization problem*, Tunis. J. Math. **3** (2021), no. 1, 75–92. [2](#)
- [Hor14] I. Horozov, *Cohomology of $GL_4(\mathbb{Z})$ with nontrivial coefficients*, Math. Res. Lett. **21** (2014), no. 5, 1111–1136. [2](#)
- [LS76] R. Lee and R. H. Szczarba, *On the homology and cohomology of congruence subgroups*, Invent. Math. **33** (1976), no. 1, 15–53. [3](#)
- [MM65] John W. Milnor and John C. Moore, *On the structure of Hopf algebras*, Ann. of Math. (2) **81** (1965), 211–264. [9](#)
- [MNP20] J. Miller, R. Nagpal, and P. Patzt, *Stability in the high-dimensional cohomology of congruence subgroups*, Compos. Math. **156** (2020), no. 4, 822–861. [1, 3](#)

- [MPW] J. Miller, P. Patzt, and J. C. H. Wilson, *On rank filtrations of algebraic K-theory and Steinberg modules*, <https://arxiv.org/abs/2303.00245>. 1, 3
- [Nag19] Rohit Nagpal, *VI-modules in nondescribing characteristic, part I*, Algebra Number Theory **13** (2019), no. 9, 2151–2189. 3
- [PS22] Andrew Putman and Daniel Studenmund, *The dualizing module and top-dimensional cohomology group of $GL_n(\mathcal{O})$* , Math. Z. **300** (2022), no. 1, 1–31. 1
- [Rog92] John Rognes, *A spectrum level rank filtration in algebraic K-theory*, Topology **31** (1992), no. 4, 813–845. 3
- [Sol69] L. Solomon, *The Steinberg character of a finite group with BN-pair*, Theory of Finite Groups (Symposium, Harvard Univ., Cambridge, Mass., 1968), Benjamin, New York, 1969, pp. 213–221. 3
- [Sou10] C. Soulé, *Higher K-theory of algebraic integers and the cohomology of arithmetic groups*, Cohomology of groups and algebraic K-theory, Adv. Lect. Math. (ALM), vol. 12, Int. Press, Somerville, MA, 2010, Notes by Marco Varisco, pp. 503–517. 2

Email address: `avner.ash@bc.edu`

BOSTON COLLEGE, CHESTNUT HILL, MA 02445, USA

Email address: `jeremykmiller@purdue.edu`

PURDUE UNIVERSITY, WEST LAFAYETTE IN, 47907, USA

Email address: `ppatz@ou.edu`

UNIVERSITY OF OKLAHOMA, NORMAN OK, 73019, USA