

Transposed Poisson structures on quasi-filiform Lie algebras of maximum length*

Kobiljon Abdurasulov[†], Fatanah Deraman[‡], Azamat Saydaliyev[§] & Siti Hasana Sapar[¶]

Abstract: This article will discussing on $\frac{1}{2}$ -derivations of quasi-filiform Lie algebras of maximum length. The non-trivial transposed Poisson algebras with the quasi-filiform Lie algebras of maximum length are constructed by using $\frac{1}{2}$ -derivations of Lie algebras. We have established commutative associative multiplication to construct a transposed Poisson algebra with an associated given Lie algebra.

Keywords: Lie algebra, transposed Poisson algebra, $\frac{1}{2}$ -derivation.

MSC2020: 17A30, 17B40, 17B61, 17B63.

INTRODUCTION

Bai et.al. [3] introduced a dual notion of the Poisson algebra, which is called a *transposed Poisson algebra*, by changing the roles of the two multiplications in the Leibniz rule. Poisson algebra is introduced to commutative associative algebras by their derivation. Similarly, on a Lie algebra, the concept of a transposed Poisson algebra is defined by the $\frac{1}{2}$ -derivation. This way of defining not only shares some properties of a Poisson algebra, such as the closedness under tensor products and the Koszul self-duality as an operation, but also admits a rich class of identities [3, 5, 7, 9, 24–26].

The description of all Poisson algebras with a fixed Lie or associative part [10, 15, 30] is one of the natural tasks in the theory. In this paper, we present transposed Poisson algebras that demonstrate quasi-filiform Lie algebras of maximum length. Abdurasulov et.al. [1], obtained the description of all transposed Poisson algebra structures on solvable Lie algebras with filiform nilradical.

It is worth noting that any unital transposed Poisson algebra is a particular case of a “contact bracket” algebra and a quasi-Poisson algebra [5]. Every transposed Poisson algebra is a commutative Gelfand-Dorfman algebra [26], and is also an algebra of Jordan brackets [7]. Ferreira et. al. [6] established a relation between transposed Poisson algebras and $\frac{1}{2}$ -derivations of Lie algebras. These ideas played an active role in describing all transposed Poisson structures on Witt and Virasoro algebras in [6]. Twisted Heisenberg-Virasoro, Schrödinger-Virasoro and extended Schrödinger-Virasoro

*This work was supported by UIDB/MAT/00212/2020, UIDP/MAT/00212/2020, 2022.02474.PTDC and by grant F-FA-2021-423, Ministry of Higher Education, Science and Innovations of the Republic of Uzbekistan.

[†]CMA-UBI, Universidade da Beira Interior, Covilhã, Portugal; Institute of Mathematics Academy of Sciences of Uzbekistan, Tashkent, Uzbekistan; abdurashulov0505@mail.ru

[‡]Universiti Malaysia Perlis, Perlis, Malaysia; fatanah@unimap.edu.my

[§]Institute of Mathematics Academy of Sciences of Uzbekistan, Tashkent, Uzbekistan; National University of Uzbekistan, Tashkent, Uzbekistan; azamatsaydaliyev6@gmail.com

[¶]Universiti Putra Malaysia, Selangor, Malaysia; sitihas@upm.edu.my

algebras were studied in [32]. Meanwhile, Schrödinger algebra in $(n + 1)$ -dimensional space-time was discussed by [29]. Then, in [17] Witt type Lie algebras were discussed and [18] studied generalized Witt algebras. In the paper by [16, 18], they were studied on Block Lie algebras and [19] was studied on Lie algebra of upper triangular matrices and a transposed Poisson structures on Lie algebra of upper triangular matrices, while [20] was discussed on Lie incidence algebras. It was proved in [22] that any complex finite-dimensional solvable Lie algebra admit a non-trivial transposed Poisson structure. In [4] authors gave the algebraic and geometric classification of three-dimensional transposed Poisson algebras. The list of actual open questions on transposed Poisson algebras can be seen in [5]. Recently, all transposed Poisson algebra structures on oscillator Lie algebras, i.e., on one-dimensional solvable extensions of $(2n + 1)$ -dimensional Heisenberg algebra; on solvable Lie algebras with naturally graded filiform nilpotent radical; on $(n + 1)$ -dimensional solvable extensions of the $(2n + 1)$ -dimensional Heisenberg algebra; and on n -dimensional solvable extensions of n -dimensional algebra with the trivial multiplication were described [21]. This paper also gave an example of a finite-dimensional Lie algebra with non-trivial $\frac{1}{2}$ -derivations without non-trivial transposed Poisson algebra structures. Also, similar studies can be seen in [14, Section 7.3], and in the references given in that paper. In [4], they were obtained the algebraic and geometric classification of all complex 3-dimensional transposed Poisson algebras, and [31] was studied on the algebraic classification of all complex 3-dimensional transposed Poisson 3-Lie algebras.

The purpose of this article is to find all transposed Poisson algebras that demonstrate quasi-filiform Lie algebras of maximum length. In order to achieve our goal, we have organized this paper as follows: in Section 2, we described $\frac{1}{2}$ -derivations of quasi-filiform Lie algebras of maximum length and Section 3, will describe all non-trivial transposed Poisson algebras with quasi-filiform Lie algebras of maximum length. Next, by using descriptions of $\frac{1}{2}$ -derivations of Lie algebras, we established commutative associative multiplication to construct a transposed Poisson algebra associated with given Lie algebra.

1. PRELIMINARIES

This section will discuss the concepts and known results on algebras over the field \mathbb{C} unless otherwise stated.

We start with the definition of a Poisson algebra.

Definition 1. Let \mathfrak{L} be a vector space equipped with two bilinear operations

$$\cdot, [-, -] : \mathfrak{L} \otimes \mathfrak{L} \rightarrow \mathfrak{L}.$$

The triple $(\mathfrak{L}, \cdot, [-, -])$ is called a **Poisson algebra** if (\mathfrak{L}, \cdot) is a commutative associative algebra and $(\mathfrak{L}, [-, -])$ is a Lie algebra which satisfies the compatibility condition

$$[x, y \cdot z] = [x, y] \cdot z + y \cdot [x, z]. \quad (1)$$

Eq. (1) is called **Leibniz rule** since the adjoint operators of Lie algebra are derivations of the commutative associative algebra.

Definition 2. Let \mathfrak{L} be a vector space equipped with two bilinear operations

$$\cdot, [-, -] : \mathfrak{L} \otimes \mathfrak{L} \rightarrow \mathfrak{L}.$$

The triple $(\mathfrak{L}, \cdot, [-, -])$ is called a **transposed Poisson algebra** if (\mathfrak{L}, \cdot) is a commutative associative algebra and $(\mathfrak{L}, [-, -])$ is a Lie algebra which satisfies the following compatibility condition

$$2z \cdot [x, y] = [z \cdot x, y] + [x, z \cdot y]. \quad (2)$$

Eq. (2) is called **transposed Leibniz rule** because the roles of two binary operations in the Leibniz rule in a Poisson algebra are switched. Further, the resulting operation is rescaled by introducing a factor 2 on the left-hand side.

Transposed Poisson algebras were initially introduced by Bai et. al. [3]. A transposed Poisson structure \cdot on \mathfrak{L} is called **trivial**, if $x \cdot y = 0$ for all $x, y \in \mathfrak{L}$.

Definition 3. Let $(\mathfrak{L}, [-, -])$ be an algebra with a multiplication $[-, -]$, and φ be a bilinear map. Then φ is a $\frac{1}{2}$ -derivation if it satisfies:

$$\varphi([x, y]) = \frac{1}{2}([\varphi(x), y] + [x, \varphi(y)]). \quad (3)$$

Note that $\frac{1}{2}$ -derivations are a particular case of δ -derivations introduced by Filippov in [8] (see, [12, 13] and references therein). It is easy to see from Definition 3 that $[\mathfrak{L}, \mathfrak{L}]$ and $\text{Ann}(\mathfrak{L})$ are invariant under any $\frac{1}{2}$ -derivation of \mathfrak{L} . Definitions 2 and 3 immediately imply the following lemma.

Lemma 4. Let $(\mathfrak{L}, [-, -])$ be a Lie algebra and \cdot a new binary (bilinear) operation on \mathfrak{L} . Then $(\mathfrak{L}, \cdot, [-, -])$ is a transposed Poisson algebra if and only if \cdot is commutative and associative and for every $z \in \mathfrak{L}$ the multiplication by z in (\mathfrak{L}, \cdot) is a $\frac{1}{2}$ -derivation of $(\mathfrak{L}, [-, -])$.

Lower central series for a given Lie algebra \mathfrak{L} is defined as follows:

$$\mathfrak{L}^1 = \mathfrak{L}, \mathfrak{L}^{k+1} = [\mathfrak{L}^k, \mathfrak{L}], k \geq 1.$$

Definition 5. A Lie algebra \mathfrak{L} is said to be **nilpotent**, if there exists $k \in \mathbb{N}$ such that $\mathfrak{L}^k = \{0\}$. The minimal number k with such property is said to be the **index of nilpotency** of the algebra \mathfrak{L} .

Definition 6. An n -dimensional Lie algebra is called **quasi-filiform** if its index of nilpotency is equal to $n - 1$.

Now, let us define a maximum length for the nilpotent Lie algebras.

A Lie algebra \mathfrak{L} is \mathbb{Z} -graded, if $\mathfrak{L} = \bigoplus_{i \in \mathbb{Z}} V_i$, where $[V_i, V_j] \subseteq V_{i+j}$ for any $i, j \in \mathbb{Z}$ with a finite number of non-null spaces V_i . We say that a nilpotent Lie algebra \mathfrak{L} admits the **connected gradation** $\mathfrak{L} = V_{k_1} \oplus \cdots \oplus V_{k_t}$, if $V_{k_i} \neq \{0\}$ for any i ($1 \leq i \leq t$).

Definition 7. The number $l(\oplus \mathfrak{L}) = l(V_{k_1} \oplus \cdots \oplus V_{k_t}) = k_t - k_1 + 1$ is called **the length of gradation**. A gradation is called **of maximum length**, if $l(\oplus \mathfrak{L}) = \dim(\mathfrak{L})$.

We denote $l(\mathfrak{L}) = \max\{l(\oplus L) \text{ such that } L = V_{k_1} \oplus \cdots \oplus V_{k_t} \text{ is a connected gradation}\}$ of **the length of an algebra** \mathfrak{L} .

Definition 8. A Lie algebra \mathfrak{L} is called of **maximum length** if $l(\mathfrak{L}) = \dim(\mathfrak{L})$.

In the following theorem we present the classification of quasi-filiform Lie algebras of maximum length given in [11].

Theorem 9. Let L be an n -dimensional quasi-filiform Lie algebra of maximum length. Then the algebra L is isomorphic to one of the following pairwise non-isomorphic algebras:

$$\begin{aligned}
g_{(n,1)}^1 : & \begin{cases} [e_1, e_i] = e_{i+1}, & 2 \leq i \leq n-2, \\ [e_i, e_{n-i}] = (-1)^i e_n, & 2 \leq i \leq \frac{n-1}{2}, n \geq 5 \text{ and } n \text{ is odd}; \end{cases} \\
g_{(n,1)}^2 : & \begin{cases} [e_1, e_i] = e_{i+1}, & 2 \leq i \leq n-2, \\ [e_i, e_n] = e_{i+2}, & 2 \leq i \leq n-3, n \geq 5; \end{cases} \quad g_{(n,1)}^3 : \begin{cases} [e_1, e_i] = e_{i+1}, & 2 \leq i \leq n-2, \\ [e_i, e_n] = e_{i+2}, & 2 \leq i \leq n-3 \\ [e_2, e_i] = e_{i+3}, & 3 \leq i \leq n-4, n \geq 7; \end{cases} \\
g_7^1 : & \begin{cases} [e_1, e_i] = e_{i+1}, & 2 \leq i \leq 5, \\ [e_2, e_i] = e_{i+2}, & 3 \leq i \leq 4, \\ [e_i, e_{7-i}] = (-1)^i e_7, & 2 \leq i \leq 3; \end{cases} \quad g_9^2 : \begin{cases} [e_1, e_i] = e_{i+1}, & 2 \leq i \leq 7, \\ [e_2, e_i] = e_{i+2}, & 3 \leq i \leq 4, \\ [e_2, e_5] = 3e_7, & \\ [e_2, e_6] = 5e_8, & \\ [e_3, e_i] = -2e_{i+3}, & 4 \leq i \leq 5, \\ [e_i, e_{9-i}] = (-1)^i e_9, & 2 \leq i \leq 4; \end{cases} \\
g_{11}^3 : & \begin{cases} [e_1, e_i] = e_{i+1}, & 2 \leq i \leq 9, \\ [e_2, e_i] = e_{i+2}, & 3 \leq i \leq 4, \\ [e_2, e_i] = -e_{i+2}, & 6 \leq i \leq 7 \\ [e_3, e_7] = -e_{10}, & \\ [e_3, e_i] = e_{i+3}, & 4 \leq i \leq 5, \\ [e_4, e_i] = e_{i+4}, & 5 \leq i \leq 6, \\ [e_i, e_{11-i}] = (-1)^i e_{11}, & 2 \leq i \leq 5, \end{cases}
\end{aligned}$$

where $\{e_1, e_2, \dots, e_n\}$ is a basis of the algebra.

2. $\frac{1}{2}$ -DERIVATION OF QUASI-FILIFORM LIE ALGEBRAS OF MAXIMUM LENGTH

In this section, we will derive and calculate $\frac{1}{2}$ -derivation of quasi-filiform Lie algebras of maximum length. We start with the following theorem.

Theorem 10. Any $\frac{1}{2}$ -derivations of the algebra $g_{(n,1)}^1$ has the form

• for $n = 5$:

$$\begin{cases} \varphi(e_1) = \alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3 + \alpha_4 e_4 + \alpha_5 e_5, \\ \varphi(e_2) = \beta_1 e_1 + \beta_2 e_2 + \beta_3 e_3 + \beta_4 e_4 + \beta_5 e_5, \\ \varphi(e_3) = \frac{1}{2}(\alpha_1 + \beta_2)e_3 + \frac{1}{2}\beta_3 e_4 - \frac{1}{2}\alpha_3 e_5, \\ \varphi(e_4) = \frac{1}{4}(3\alpha_1 + \beta_2)e_4 + \frac{1}{2}\alpha_2 e_5, \\ \varphi(e_5) = \frac{1}{2}\beta_1 e_4 + \frac{1}{4}(\alpha_1 + 3\beta_2)e_5, \end{cases}$$

• for $n \geq 7$:

$$\begin{cases} \varphi(e_1) = \sum_{i=1}^n \alpha_i e_i, \\ \varphi(e_2) = \alpha_1 e_2 + \beta_{n-2} e_{n-2} + \beta_{n-1} e_{n-1} + \beta_n e_n, \\ \varphi(e_3) = \alpha_1 e_3 + \frac{1}{2}\beta_{n-2} e_{n-1} - \frac{1}{2}\alpha_{n-2} e_n, \\ \varphi(e_i) = \alpha_1 e_i + \frac{(-1)^i}{2} \alpha_{n-i+1} e_n, \quad 4 \leq i \leq n-1, \\ \varphi(e_n) = \alpha_1 e_n. \end{cases}$$

Proof. It is easy to see that $g_{(n,1)}^1$ has 2 generators. We use these generators to calculate of $\frac{1}{2}$ -derivation.

$$\varphi(e_1) = \sum_{i=1}^n \alpha_i e_i, \quad \varphi(e_2) = \sum_{i=1}^n \beta_i e_i.$$

Now consider the condition of $\frac{1}{2}$ -derivation for the elements e_1 and e_2 :

$$\begin{aligned} \varphi(e_3) &= \varphi([e_1, e_2]) = \frac{1}{2}([\varphi(e_1), e_2] + [e_1, \varphi(e_2)]) \\ &= \frac{1}{2}([\sum_{i=1}^n \alpha_i e_i, e_2] + [e_1, \sum_{i=1}^n \beta_i e_i]) = \frac{1}{2}((\alpha_1 + \beta_2)e_3 + \sum_{i=4}^{n-1} \beta_{i-1} e_i - \alpha_{n-2} e_n). \end{aligned}$$

We prove the following equality for $3 \leq i \leq n-1$ by induction:

$$\varphi(e_i) = \frac{1}{2^{i-2}} \left((2^{i-2} - 1)\alpha_1 + \beta_2 \right) e_i + \frac{1}{2^{i-2}} \sum_{t=i+1}^{n-1} \beta_{t-i+2} e_t + \frac{(-1)^i}{2} \alpha_{n-i+1} e_n.$$

If $i = 3$, the relationship holds according to the above equality. Now, we prove that it is true for i and $i+1$. By considering the condition of $\frac{1}{2}$ -derivation for the elements e_1, e_i we have

$$\begin{aligned} \varphi(e_{i+1}) &= \varphi([e_1, e_i]) = \frac{1}{2}([\varphi(e_1), e_i] + [e_1, \varphi(e_i)]) \\ &= \frac{1}{2} \left([\sum_{k=1}^n \alpha_k e_k, e_i] + [e_1, \frac{1}{2^{i-2}} \left((2^{i-2} - 1)\alpha_1 + \beta_2 \right) e_i + \frac{1}{2^{i-2}} \sum_{t=i+1}^{n-1} \beta_{t-i+2} e_t + \frac{(-1)^i}{2} \alpha_{n-i+1} e_n] \right) \\ &= \frac{1}{2} \left(\alpha_1 e_{i+1} + (-1)^{i+1} \alpha_{n-i} e_n + \frac{1}{2^{i-2}} \left((2^{i-2} - 1)\alpha_1 + \beta_2 \right) e_{i+1} + \frac{1}{2^{i-2}} \sum_{t=i+2}^{n-2} \beta_{t-i+2} e_{t+1} \right) \\ &= \frac{1}{2^{i-1}} \left((2^{i-1} - 1)\alpha_1 + \beta_2 \right) e_{i+1} + \frac{1}{2^{i-1}} \sum_{t=i+2}^{n-1} \beta_{t-i+1} e_t + \frac{(-1)^{i+1}}{2} \alpha_{n-i} e_n. \end{aligned}$$

Now, consider the condition of $\frac{1}{2}$ -derivation for the elements e_2, e_{n-2} :

$$\begin{aligned} \varphi(e_n) &= \varphi([e_2, e_{n-2}]) = \frac{1}{2}([\varphi(e_2), e_{n-2}] + [e_2, \varphi(e_{n-2})]) \\ &= \frac{1}{2}\left(\left[\sum_{i=1}^n \beta_i e_i, e_{n-2}\right] + [e_2, \frac{1}{2^{n-4}}\left((2^{n-4}-1)\alpha_1 + \beta_2\right)e_{n-2} + \frac{1}{2^{n-4}}\beta_3 e_{n-1} + \frac{(-1)^{n-2}}{2}\alpha_3 e_n]\right) \\ &= \frac{1}{2}\left(\beta_1 e_{n-1} + \beta_2 e_n + \frac{1}{2^{n-4}}\left((2^{n-4}-1)\alpha_1 + \beta_2\right)e_n\right) = \frac{1}{2}\beta_1 e_{n-1} + \frac{1}{2^{n-3}}\left((2^{n-4}-1)\alpha_1 + (2^{n-4}+1)\beta_2\right)e_n. \end{aligned}$$

Thus, we obtain that

$$\varphi(e_n) = \frac{1}{2}\beta_1 e_{n-1} + \frac{1}{2^{n-3}}\left((2^{n-4}-1)\alpha_1 + (2^{n-4}+1)\beta_2\right)e_n. \quad (4)$$

Now, we consider the condition of $\frac{1}{2}$ -derivation for the elements e_2 and e_i , $3 \leq i \leq n-3$:

$$\begin{aligned} 0 &= \varphi([e_2, e_i]) = \frac{1}{2}([\varphi(e_2), e_i] + [e_2, \varphi(e_i)]) \\ &= \frac{1}{2}\left(\left[\sum_{i=1}^n \beta_i e_i, e_i\right] + [e_2, \frac{1}{2^{i-2}}\left((2^{i-2}-1)\alpha_1 + \beta_2\right)e_i + \frac{1}{2^{i-2}}\sum_{t=i+1}^{n-1} \beta_{t-i+2} e_t + \frac{(-1)^i}{2}\alpha_{n-i+1} e_n]\right) \\ &= \frac{1}{2}\left(\beta_1 e_{i+1} - (-1)^i \beta_{n-i} e_n + \frac{1}{2^{i-2}}\beta_{n-i} e_n\right). \end{aligned}$$

By the comparison of coefficients at the basis elements we obtain that

$$\beta_1 = \beta_i = 0, \quad 3 \leq i \leq n-3.$$

By considering the condition of $\frac{1}{2}$ -derivation for the elements e_3 and e_{n-3} we have

$$\begin{aligned} \varphi([e_3, e_{n-3}]) &= \frac{1}{2}([\varphi(e_3), e_{n-3}] + [e_3, \varphi(e_{n-3})]) \\ &= \frac{1}{2}\left(\left[\frac{1}{2}((\alpha_1 + \beta_2)e_3 + \beta_{n-2}e_{n-1} - \alpha_{n-2}e_n), e_{n-3}\right] + [e_3, \frac{1}{2^{n-5}}\left((2^{n-5}-1)\alpha_1 + \beta_2\right)e_{n-3} + \frac{1}{2}\alpha_4 e_n]\right) \\ &= \frac{1}{2}\left(-\frac{1}{2}((\alpha_1 + \beta_2)e_n - \frac{1}{2^{n-5}}\left((2^{n-5}-1)\alpha_1 + \beta_2\right)e_n)\right). \end{aligned}$$

On the other hand,

$$\varphi([e_3, e_{n-3}]) = -\varphi(e_n) = -\left(\frac{1}{2^2}((\alpha_1 + \beta_2) + \frac{1}{2^{n-4}}\left((2^{n-5}-1)\alpha_1 + \beta_2\right))e_n\right). \quad (5)$$

By comparing the coefficients at the basis elements in expressions (4) and (5) we obtain that $\alpha_1 = \beta_2$. This completes the proof of the theorem. \square

Now we study the $\frac{1}{2}$ -derivation of the algebra $g_{(n,1)}^2$.

Theorem 11. Any $\frac{1}{2}$ -derivations of the algebra $g_{(n,1)}^2$ has the form

- for $n = 5$:

$$\begin{cases} \varphi(e_1) = \alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3 + \alpha_4 e_4 + \alpha_5 e_5, \\ \varphi(e_2) = \beta_2 e_2 + \beta_3 e_3 + \beta_4 e_4 + \beta_5 e_5, \\ \varphi(e_3) = \frac{1}{2}(\alpha_1 + \beta_2)e_3 + \frac{1}{2}(\beta_3 - \alpha_5)e_4, \\ \varphi(e_4) = \frac{1}{4}(3\alpha_1 + \beta_2)e_4, \\ \varphi(e_5) = -\alpha_2 e_3 + \gamma_4 e_4 + \frac{1}{2}(3\alpha_1 - \beta_2)e_5, \end{cases}$$

- for $n = 6$:

$$\begin{cases} \varphi(e_1) = \alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3 + \alpha_4 e_4 + \alpha_5 e_5 + \alpha_6 e_6, \\ \varphi(e_2) = \alpha_1 e_2 - 3\alpha_6 e_3 + \beta_4 e_4 + \beta_5 e_5, \\ \varphi(e_3) = \alpha_1 e_3 - 2\alpha_6 e_4 + \frac{1}{2}\beta_4 e_5, \\ \varphi(e_4) = \alpha_1 e_4 - \frac{3}{2}\alpha_6 e_5, \\ \varphi(e_5) = \alpha_1 e_5, \\ \varphi(e_6) = -\alpha_2 e_3 - \alpha_3 e_4 + \gamma_5 e_5 + \alpha_1 e_6, \end{cases}$$

- for $n \geq 7$:

$$\begin{cases} \varphi(e_1) = \sum_{i=1}^{n-1} \alpha_i e_i, \\ \varphi(e_2) = \alpha_1 e_2 + \beta_{n-2} e_{n-2} + \beta_{n-1} e_{n-1}, \\ \varphi(e_3) = \alpha_1 e_3 + \frac{1}{2}\beta_{n-2} e_{n-1}, \\ \varphi(e_i) = \alpha_1 e_i, \quad 4 \leq i \leq n-1, \\ \varphi(e_n) = -\sum_{i=3}^{n-2} \alpha_{i-1} e_i + \gamma_{n-1} e_{n-1} + \alpha_1 e_n. \end{cases}$$

Proof. From table multiplication of the algebra $g_{(n,1)}^2$ we conclude that e_1, e_2 and e_n are the generator basis elements of the algebra. We use these generators to calculate of $\frac{1}{2}$ -derivation:

$$\varphi(e_1) = \sum_{i=1}^n \alpha_i e_i, \quad \varphi(e_2) = \sum_{i=1}^n \beta_i e_i, \quad \varphi(x) = \sum_{i=1}^n \gamma_i e_i.$$

Now consider the condition of $\frac{1}{2}$ -derivation for the elements e_1 and e_2 :

$$\begin{aligned} \varphi(e_3) &= \varphi([e_1, e_2]) = \frac{1}{2}([\varphi(e_1), e_2] + [e_1, \varphi(e_2)]) \\ &= \frac{1}{2}([\sum_{i=1}^n \alpha_i e_i, e_2] + [e_1, \sum_{i=1}^n \beta_i e_i]) = \frac{1}{2}\left((\alpha_1 + \beta_2)e_3 + (\beta_3 - \alpha_n)e_4 + \sum_{i=5}^{n-1} \beta_{i-1} e_i\right). \end{aligned}$$

We prove the following equality for $3 \leq i \leq n-2$ by induction:

$$\varphi(e_i) = \frac{1}{2^{i-2}} \left((2^{i-2} - 1)\alpha_1 + \beta_2 \right) e_i + \frac{1}{2^{i-2}} \left(\beta_3 - (2^{i-2} - 1)\alpha_n \right) e_{i+1} + \frac{1}{2^{i-2}} \sum_{t=i+2}^{n-1} \beta_{t-i+2} e_t$$

and

$$\varphi(e_{n-1}) = \frac{1}{2^{n-3}} \left((2^{n-3} - 1)\alpha_1 + \beta_2 \right) e_{n-1}.$$

If $i = 3$, the relationship is fulfilled according to the above equality. Now, we prove that it is true for i and $i + 1$. By considering the condition of $\frac{1}{2}$ -derivation for the elements e_1, e_i , we have the following:

$$\begin{aligned} & \varphi(e_{i+1}) = \varphi([e_1, e_i]) = \frac{1}{2}([\varphi(e_1), e_i] + [e_1, \varphi(e_i)]) \\ &= \frac{1}{2} \left([\sum_{k=1}^n \alpha_k e_k, e_i] + [e_1, \frac{1}{2^{i-2}} \left((2^{i-2} - 1)\alpha_1 + \beta_2 \right) e_i] + \frac{1}{2^{i-2}} \left(\beta_3 - (2^{i-2} - 1)\alpha_n \right) e_{i+1} + \frac{1}{2^{i-2}} \sum_{t=i+2}^{n-1} \beta_{t-i+2} e_t \right) \\ &= \frac{1}{2} \left(\alpha_1 e_{i+1} - \alpha_n e_{i+2} + \frac{1}{2^{i-2}} \left((2^{i-2} - 1)\alpha_1 + \beta_2 \right) e_{i+1} + \frac{1}{2^{i-2}} \left(\beta_3 - (2^{i-2} - 1)\alpha_n \right) e_{i+2} \right. \\ &\quad \left. + \frac{1}{2^{i-2}} \sum_{t=i+2}^{n-2} \beta_{t-i+2} e_{t+1} \right) \\ &= \frac{1}{2^{i-1}} \left((2^{i-1} - 1)\alpha_1 + \beta_2 \right) e_{i+1} + \frac{1}{2^{i-1}} \left(\beta_3 - (2^{i-1} - 1)\alpha_n \right) e_{i+2} + \frac{1}{2^{i-1}} \sum_{t=i+3}^{n-1} \beta_{t-i+1} e_t. \end{aligned}$$

From

$$\begin{aligned} 0 &= \varphi([e_1, e_n]) = \frac{1}{2}([\varphi(e_1), e_n] + [e_1, \varphi(e_n)]) = \frac{1}{2} \left([\sum_{k=1}^n \alpha_k e_k, e_n] + [e_1, \sum_{k=1}^n \gamma_k e_k] \right) \\ &= \frac{1}{2} \left(\sum_{k=2}^{n-3} \alpha_k e_{k+2} + \sum_{k=2}^{n-2} \gamma_k e_{k+1} \right) = \frac{1}{2} \left(\sum_{k=4}^{n-1} \alpha_{k-2} e_k + \sum_{k=3}^{n-1} \gamma_{k-1} e_k \right), \end{aligned}$$

we obtain,

$$\gamma_2 = 0, \gamma_i = -\alpha_{i-1}, 3 \leq i \leq n-2.$$

Now consider the condition of $\frac{1}{2}$ -derivation for the elements e_2, e_n :

$$\begin{aligned} \varphi([e_2, e_n]) &= \frac{1}{2}([\varphi(e_2), e_n] + [e_2, \varphi(e_n)]) = \frac{1}{2} \left([\sum_{i=1}^n \beta_i e_i, e_n] + [e_2, \sum_{i=1}^n \gamma_i e_i] \right) \\ &= \frac{1}{2} \left(\sum_{i=2}^{n-3} \beta_i e_{i+2} - \gamma_1 e_3 + \gamma_n e_4 \right) = \frac{1}{2} \left(-\gamma_1 e_3 + (\beta_2 + \gamma_n) e_4 + \sum_{i=5}^{n-1} \beta_{i-2} e_i \right). \end{aligned}$$

Thus,

$$\varphi([e_2, e_n]) = \varphi(e_4).$$

By Comparing the coefficients at the basis elements, we obtain that

$$\gamma_1 = 0, 2\gamma_n = 3\alpha_1 - \beta_2; \text{ for } n \geq 6 \text{ we have } \beta_3 = -3\alpha_n, \beta_i = 0, 4 \leq i \leq n-3.$$

By using the property of $\frac{1}{2}$ -derivation for the products $[e_2, e_3] = 0$ and $[e_3, e_n] = e_5$, we have

$$\beta_1 = 0; \text{ and for } n \geq 6 \text{ we have } \beta_n = 0, \beta_2 = \alpha_1;$$

and for $n \geq 7$ we have $\beta_3 = \alpha_n = 0$.

This completes the proof of the theorem. \square

Now, we will study the $\frac{1}{2}$ -derivation of the algebra $g_{(n,1)}^3$.

Theorem 12. Any $\frac{1}{2}$ -derivations of the algebra $g_{(n,1)}^3$ has the form

- for $n = 7$:

$$\begin{cases} \varphi(e_1) = \alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3 + \alpha_4 e_4 + \alpha_5 e_5 + \alpha_6 e_6, \\ \varphi(e_2) = \alpha_1 e_2 + 4\alpha_2 e_4 + \beta_5 e_5 + \beta_6 e_6, \\ \varphi(e_3) = \alpha_1 e_3 + 2\alpha_2 e_5 + \frac{1}{2}(\beta_5 - \alpha_3) e_6, \\ \varphi(e_4) = \alpha_1 e_4 + \frac{3}{2}\alpha_2 e_6, \\ \varphi(e_5) = \alpha_1 e_5, \varphi(e_6) = \alpha_1 e_6, \\ \varphi(e_7) = -\alpha_2 e_3 - \alpha_3 e_4 - \alpha_4 e_5 + \gamma_6 e_6 + \alpha_1 e_7, \end{cases}$$

- for $n \geq 8$:

$$\begin{cases} \varphi(e_1) = \alpha_1 e_1 + \sum_{i=3}^{n-1} \alpha_i e_i, \\ \varphi(e_2) = \alpha_1 e_2 + \sum_{i=5}^{n-3} \alpha_{i-2} e_i + \beta_{n-2} e_{n-2} + \beta_{n-1} e_{n-1}, \\ \varphi(e_3) = \alpha_1 e_3 + \frac{1}{2}(\beta_{n-2} - \alpha_{n-4}) e_{n-1}, \\ \varphi(e_i) = \alpha_1 e_i, 4 \leq i \leq n-1, \\ \varphi(e_n) = -\sum_{i=4}^{n-2} \alpha_{i-1} e_i + \gamma_{n-1} e_{n-1} + \alpha_1 e_n. \end{cases}$$

Proof. The algebra $g_{(n,1)}^3$ has generators e_1, e_2 and e_n . Then, we have the following

$$\varphi(e_1) = \sum_{i=1}^n \alpha_i e_i, \quad \varphi(e_2) = \sum_{i=1}^n \beta_i e_i, \quad \varphi(e_n) = \sum_{i=1}^n \gamma_i e_i.$$

Now, consider the condition of $\frac{1}{2}$ -derivation for the elements e_2 and e_n we obtain

$$\begin{aligned} \varphi(e_3) &= \varphi([e_1, e_2]) = \frac{1}{2}([\varphi(e_1), e_2] + [e_1, \varphi(e_2)]) = \frac{1}{2}([\sum_{i=1}^n \alpha_i e_i, e_2] + [e_1, \sum_{i=1}^n \beta_i e_i]) \\ &= \frac{1}{2} \left(\alpha_1 e_3 - \sum_{i=3}^{n-4} \alpha_i e_{i+3} - \alpha_n e_4 + \sum_{i=2}^{n-2} \beta_i e_{i+1} \right) \\ &= \frac{1}{2} \left((\alpha_1 + \beta_2) e_3 + (\beta_3 - \alpha_n) e_4 + \beta_4 e_5 + \sum_{i=6}^{n-1} (\beta_{i-1} - \alpha_{i-3}) e_i \right). \end{aligned}$$

We can prove the following equalities by induction:

$$\begin{aligned}\varphi(e_i) &= \frac{1}{2^{i-2}} \left((2^{i-2} - 1)\alpha_1 + \beta_2 \right) e_i + \frac{1}{2^{i-2}} \left(\beta_3 - (2^{i-2} - 1)\alpha_n \right) e_{i+1} \\ &\quad + \frac{1}{2^{i-2}} \left((2^{i-2} - 2)\alpha_2 + \beta_4 \right) e_{i+2} + \frac{1}{2^{i-2}} \sum_{t=i+3}^{n-1} (\beta_{t-i+2} - \alpha_{t-i}) e_t\end{aligned}$$

for $3 \leq i \leq n-4$ and

$$\begin{aligned}\varphi(e_{n-3}) &= \frac{1}{2^{n-5}} \left((2^{n-5} - 1)\alpha_1 + \beta_2 \right) e_{n-3} + \frac{1}{2^{n-5}} \left(\beta_3 - (2^{n-5} - 1)\alpha_n \right) e_{n-2} \\ &\quad + \frac{1}{2^{n-5}} \left((2^{n-5} - 2)\alpha_2 + \beta_4 \right) e_{n-1}, \\ \varphi(e_{n-2}) &= \frac{1}{2^{n-4}} \left((2^{n-4} - 1)\alpha_1 + \beta_2 \right) e_{n-2} + \frac{1}{2^{n-4}} \left(\beta_3 - (2^{n-4} - 1)\alpha_n \right) e_{n-1}, \\ \varphi(e_{n-1}) &= \frac{1}{2^{n-3}} \left((2^{n-3} - 1)\alpha_1 + \beta_2 \right) e_{n-1}.\end{aligned}$$

From

$$\begin{aligned}0 = \varphi([e_1, e_n]) &= \frac{1}{2}([\varphi(e_1), e_n] + [e_1, \varphi(e_n)]) = \frac{1}{2} \left(\left[\sum_{k=1}^n \alpha_k e_k, e_n \right] + [e_1, \sum_{k=1}^n \gamma_k e_k] \right) \\ &= \frac{1}{2} \left(\sum_{k=2}^{n-3} \alpha_k e_{k+2} + \sum_{k=2}^{n-2} \gamma_k e_{k+1} \right) = \frac{1}{2} \left(\sum_{k=4}^{n-1} \alpha_{k-2} e_k + \sum_{k=3}^{n-1} \gamma_{k-1} e_k \right),\end{aligned}$$

we obtain that

$$\gamma_2 = 0, \gamma_i = -\alpha_{i-1}, 3 \leq i \leq n-2.$$

Now, we consider the condition of $\frac{1}{2}$ -derivation for the elements e_2, e_n :

$$\begin{aligned}\varphi([e_2, e_n]) &= \frac{1}{2}([\varphi(e_2), e_n] + [e_2, \varphi(e_n)]) = \frac{1}{2} \left(\left[\sum_{i=1}^n \beta_i e_i, e_n \right] + [e_2, \sum_{i=1}^n \gamma_i e_i] \right) \\ &= \frac{1}{2} \left(\sum_{i=2}^{n-3} \beta_i e_{i+2} - \gamma_1 e_3 + \sum_{i=3}^{n-4} \gamma_i e_{i+3} + \gamma_n e_4 \right) \\ &= \frac{1}{2} \left(-\gamma_1 e_3 + (\beta_2 + \gamma_n) e_4 + \beta_3 e_5 + \sum_{i=6}^{n-1} (\beta_{i-2} - \gamma_{i-3}) e_i \right).\end{aligned}$$

Thus, we have

$$\varphi([e_2, e_n]) = \varphi(e_4) = \frac{1}{4}(3\alpha_1 + \beta_2)e_4 + \frac{1}{4}(\beta_3 - 3\alpha_n)e_5 + \frac{1}{4}(2\alpha_2 + \beta_4)e_6 + \frac{1}{4} \sum_{t=7}^{n-1} (\beta_{t-2} - \alpha_{t-4}) e_t.$$

Compare the coefficients at the basis elements, we obtain that

$$\gamma_1 = 0, 2\gamma_n = 3\alpha_1 - \beta_2, \beta_3 = -3\alpha_n, \beta_4 = 4\alpha_2, \beta_i = \alpha_{i-2}, 5 \leq i \leq n-3.$$

By using the property of $\frac{1}{2}$ -derivation for the products $[e_2, e_3] = e_6$ and $[e_3, e_n] = e_5$, we have

$$\begin{aligned}\beta_1 &= 0, \quad \beta_n = 0, \quad \beta_2 = \alpha_1; \\ \beta_3 &= \alpha_n = 0, \text{ and for } n \geq 8 \text{ we have } \beta_4 = \alpha_2 = 0.\end{aligned}$$

This completes the proof of the theorem. \square

The following theorems, we present an analogous descriptions of $\frac{1}{2}$ -derivation for quasi-filiform Lie algebras of maximum length g_7^1, g_9^2 and g_{11}^3 .

Theorem 13. Any $\frac{1}{2}$ -derivations of the algebra g_7^1 has the form

$$\left\{ \begin{array}{l} \varphi(e_1) = \alpha_1 e_1 + \alpha_3 e_3 + \alpha_4 e_4 + \alpha_5 e_5 + \alpha_6 e_6 + \alpha_7 e_7, \\ \varphi(e_2) = \alpha_1 e_2 - \frac{1}{3} \alpha_3 e_4 + \beta_5 e_5 + \beta_6 e_6 + \beta_7 e_7, \\ \varphi(e_3) = \alpha_1 e_3 - \frac{2}{3} \alpha_3 e_5 + \frac{1}{2} (\beta_5 - \alpha_4) e_6 - \frac{1}{2} \alpha_5 e_7, \\ \varphi(e_4) = \alpha_1 e_4 - \frac{1}{3} \alpha_3 e_6 + \frac{1}{2} \alpha_4 e_7, \\ \varphi(e_5) = \alpha_1 e_5 - \frac{1}{2} \alpha_3 e_7, \\ \varphi(e_6) = \alpha_1 e_6, \\ \varphi(e_7) = \alpha_1 e_7. \end{array} \right.$$

Theorem 14. Any $\frac{1}{2}$ -derivations of the algebra g_9^2 has the form

$$\left\{ \begin{array}{l} \varphi(e_1) = \alpha_1 e_1 + \alpha_5 e_5 + \alpha_6 e_6 + \alpha_7 e_7 + \alpha_8 e_8 + \alpha_9 e_9, \\ \varphi(e_2) = \alpha_1 e_2 + \frac{1}{3} \alpha_5 e_6 + \beta_7 e_7 + \beta_8 e_8 + \beta_9 e_9, \\ \varphi(e_3) = \alpha_1 e_3 - \frac{4}{3} \alpha_5 e_7 + \frac{1}{2} (\beta_7 - 5\alpha_6) e_8 - \frac{1}{2} \alpha_7 e_9, \\ \varphi(e_4) = \alpha_1 e_4 + \frac{1}{3} \alpha_5 e_8 + \frac{1}{2} \alpha_6 e_9, \\ \varphi(e_5) = \alpha_1 e_5 - \frac{1}{2} \alpha_5 e_9, \\ \varphi(e_i) = \alpha_1 e_i, \quad 6 \leq i \leq 9. \end{array} \right.$$

Theorem 15. Any $\frac{1}{2}$ -derivations of the algebra g_{11}^3 has the form

$$\left\{ \begin{array}{l} \varphi(e_1) = \alpha_1 e_1 + \alpha_6 e_6 + \alpha_7 e_7 + \alpha_8 e_8 + \alpha_9 e_9 + \alpha_{10} e_{10} + \alpha_{11} e_{11}, \\ \varphi(e_2) = \alpha_1 e_2 - \alpha_6 e_7 - \alpha_7 e_8 + \beta_9 e_9 + \beta_{10} e_{10} + \beta_{11} e_{11}, \\ \varphi(e_3) = \alpha_1 e_3 + \frac{1}{2} \beta_9 e_{10} - \frac{1}{2} \alpha_9 e_{11}, \\ \varphi(e_4) = \alpha_1 e_4 + \frac{1}{2} \alpha_7 e_{10} + \frac{1}{2} \alpha_8 e_{11}, \\ \varphi(e_5) = \alpha_1 e_5 - \frac{1}{2} \alpha_6 e_{10} - \frac{1}{2} \alpha_7 e_{11}, \\ \varphi(e_6) = \alpha_1 e_6 + \frac{1}{2} \alpha_6 e_{11}, \\ \varphi(e_i) = \alpha_1 e_i, \quad 7 \leq i \leq 11. \end{array} \right.$$

3. TRANSPOSED POISSON STRUCTURE FOR QUASI-FILIFORM LIE ALGEBRAS OF MAXIMUM LENGTH

In this section, we will describe all transposed Poisson algebra structures on quasi-filiform Lie algebras of maximum length.

Theorem 16. Let $(g_{(5,1)}^1, \cdot, [-, -])$ be a transposed Poisson algebra structure defined on the Lie algebra $g_{(5,1)}^1$. Then the multiplication of $(g_{(5,1)}^1, \cdot)$ has the following form:

$$\begin{aligned} \mathbf{TP}_1(g_{(5,1)}^1) : & e_1 \cdot e_1 = \alpha_4 e_4 + \alpha_5 e_5, \quad e_1 \cdot e_2 = \beta_4 e_4 + \beta_5 e_5, \quad e_2 \cdot e_2 = \beta_9 e_4 + \beta_{10} e_5; \\ \mathbf{TP}_2(g_{(5,1)}^1) : & \begin{cases} e_1 \cdot e_1 = e_3 + \alpha_4 e_4 + \alpha_5 e_5, \quad e_1 \cdot e_2 = \beta_3 e_3 + \beta_4 e_4 + \beta_5 e_5, \quad e_1 \cdot e_3 = \frac{1}{2} \beta_3 e_4 - \frac{1}{2} e_5, \\ e_2 \cdot e_2 = \beta_3^2 e_3 + \beta_9 e_4 + \beta_{10} e_5, \quad e_2 \cdot e_3 = \frac{1}{2} \beta_3^2 e_4 - \frac{1}{2} \beta_3 e_5; \\ e_1 \cdot e_1 = \alpha_1 e_1 + e_2 + \alpha_3 e_3 + \alpha_4 e_4 + \alpha_5 e_5, \\ e_1 \cdot e_2 = -\frac{1}{4} (\alpha_1 - \beta_2)^2 e_1 + \beta_2 e_2 - \frac{1}{2} \alpha_3 (\alpha_1 - \beta_2) e_3 + \beta_4 e_4 + \beta_5 e_5, \\ e_1 \cdot e_3 = \frac{1}{2} (\alpha_1 + \beta_2) e_3 - \frac{1}{4} \alpha_3 (\alpha_1 - \beta_2) e_4 - \frac{1}{2} \alpha_3 e_5, \\ e_1 \cdot e_4 = \frac{1}{4} (3\alpha_1 + \beta_2) e_4 + \frac{1}{2} e_5, \\ e_1 \cdot e_5 = -\frac{1}{8} (\alpha_1 - \beta_2)^2 e_4 + \frac{1}{4} (\alpha_1 + 3\beta_2) e_5, \\ e_2 \cdot e_2 = -\frac{1}{4} (\alpha_1 - \beta_2)^2 \beta_2 e_1 + \frac{1}{4} (-\alpha_1^2 - 2\alpha_1\beta_2 + 3\beta_2^2) e_2 + \frac{1}{4} \alpha_3 (\alpha_1 - \beta_2)^2 e_3 + \\ + \frac{1}{8} (\alpha_1^2 (\alpha_5\beta_2 - \beta_5) - \beta_2 (2\alpha_4\beta_2 - \alpha_5\beta_2^2 - 10\beta_4 + \beta_2\beta_5) + \\ + 2\alpha_1 (\alpha_4\beta_2 - \alpha_5\beta_2^2 - \beta_4 + \beta_2\beta_5)) e_4 + \\ + \frac{1}{4} (2\beta_4 - 2\alpha_4\beta_2 - 3\alpha_5\beta_2^2 + 3\alpha_1 (\alpha_5\beta_2 - \beta_5) + 7\beta_2\beta_5) e_5, \\ e_2 \cdot e_3 = \frac{1}{4} (\beta_2^2 - \alpha_1^2) e_3 + \frac{1}{8} \alpha_3 (\alpha_1 - \beta_2)^2 e_4 + \frac{1}{4} \alpha_3 (\alpha_1 - \beta_2) e_5, \\ e_2 \cdot e_4 = \frac{1}{4} \alpha_1 (\beta_2 - \alpha_1) e_4 + \frac{1}{2} \beta_2 e_5, \\ e_2 \cdot e_5 = -\frac{1}{8} (\alpha_1 - \beta_2)^2 \beta_2 e_4 - \frac{1}{4} (\alpha_1^2 + \alpha_1\beta_2 - 2\beta_2^2) e_5, \\ e_3 \cdot e_3 = \frac{1}{8} (\beta_2^2 - \alpha_1^2) e_4 - \frac{1}{4} (\alpha_1 + \beta_2) e_5; \end{cases} \\ \mathbf{TP}_3(g_{(5,1)}^1) : & \end{aligned}$$

where the transposed Poisson algebra has its products with respect to the bracket $[-, -]$, and the remaining products are equal to zero.

Proof. Let $(g_{(5,1)}^1, \cdot, [-, -])$ be a transposed Poisson algebra structure defined on the Lie algebra $g_{(5,1)}^1$. Then for any element of $x \in g_{(5,1)}^1$, we have the operator of multiplication $\varphi_x(y) = x \cdot y$ is a $\frac{1}{2}$ -derivation. Hence, for $1 \leq i \leq 5$ we derive by Theorem 10

$$\begin{cases} \varphi_{e_i}(e_1) = \alpha_{i,1} e_1 + \alpha_{i,2} e_2 + \alpha_{i,3} e_3 + \alpha_{i,4} e_4 + \alpha_{i,5} e_5, \\ \varphi_{e_i}(e_2) = \beta_{i,1} e_1 + \beta_{i,2} e_2 + \beta_{i,3} e_3 + \beta_{i,4} e_4 + \beta_{i,5} e_5, \\ \varphi_{e_i}(e_3) = \frac{1}{2} (\alpha_{i,1} + \beta_{i,2}) e_3 + \frac{1}{2} \beta_{i,3} e_4 - \frac{1}{2} \alpha_{i,3} e_5, \\ \varphi_{e_i}(e_4) = \frac{1}{4} (3\alpha_{i,1} + \beta_{i,2}) e_4 + \frac{1}{2} \alpha_{i,2} e_5, \\ \varphi_{e_i}(e_5) = \frac{1}{2} \beta_{i,1} e_4 + \frac{1}{4} (\alpha_{i,1} + 3\beta_{i,2}) e_5. \end{cases}$$

By considering the property of $\varphi_{e_i}(e_j) = e_i \cdot e_j = e_j \cdot e_i = \varphi_{e_j}(e_i)$, we obtain the following restrictions:

$$\{e_1, e_2\}, \Rightarrow \alpha_{2,1} = \beta_{1,1}, \alpha_{2,2} = \beta_{1,2}, \alpha_{2,3} = \beta_{1,3}, \alpha_{2,4} = \beta_{1,4}, \alpha_{2,5} = \beta_{1,5},$$

$$\begin{aligned}
\{e_1, e_3\}, \quad &\Rightarrow \alpha_{3,1} = \alpha_{3,2} = 0, \alpha_{3,3} = \frac{1}{2}(\alpha_{1,1} + \beta_{1,2}), \alpha_{3,4} = \frac{1}{2}\beta_{1,3}, \alpha_{3,5} = -\frac{1}{2}\alpha_{1,3}, \\
\{e_1, e_4\}, \quad &\Rightarrow \alpha_{4,1} = \alpha_{4,2} = \alpha_{4,3} = 0, \alpha_{4,4} = \frac{1}{4}(3\alpha_{1,1} + \beta_{1,2}), \alpha_{4,5} = \frac{1}{2}\alpha_{1,2}, \\
\{e_1, e_5\}, \quad &\Rightarrow \alpha_{5,1} = \alpha_{5,2} = \alpha_{5,3} = 0, \alpha_{5,4} = \frac{1}{2}\beta_{1,1}, \alpha_{5,5} = \frac{1}{4}(\alpha_{1,1} + 3\beta_{1,2}), \\
\{e_2, e_3\}, \quad &\Rightarrow \beta_{3,1} = \beta_{3,2} = 0, \beta_{3,3} = \frac{1}{2}(\beta_{1,1} + \beta_{2,2}), \beta_{3,4} = \frac{1}{2}\beta_{2,3}, \beta_{3,5} = -\frac{1}{2}\beta_{1,3}, \\
\{e_2, e_4\}, \quad &\Rightarrow \beta_{4,1} = \beta_{4,2} = \beta_{4,3} = 0, \beta_{4,4} = \frac{1}{4}(3\beta_{1,1} + \beta_{2,2}), \beta_{4,5} = \frac{1}{2}\beta_{1,2}, \\
\{e_2, e_5\}, \quad &\Rightarrow \beta_{5,1} = \beta_{5,2} = \beta_{5,3} = 0, \beta_{5,4} = \frac{1}{2}\beta_{2,1}, \beta_{5,5} = \frac{1}{4}(\beta_{1,1} + 3\beta_{2,2}).
\end{aligned}$$

Thus, we have the following table of commutative multiplications of the transposed Poisson algebra structure defined on the Lie algebra $g_{(5,1)}^1$:

$$\left\{
\begin{aligned}
e_1 \cdot e_1 &= \alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3 + \alpha_4 e_4 + \alpha_5 e_5, \\
e_1 \cdot e_2 &= \beta_1 e_1 + \beta_2 e_2 + \beta_3 e_3 + \beta_4 e_4 + \beta_5 e_5, \\
e_1 \cdot e_3 &= \frac{1}{2}(\alpha_1 + \beta_2) e_3 + \frac{1}{2}\beta_3 e_4 - \frac{1}{2}\alpha_3 e_5, \\
e_1 \cdot e_4 &= \frac{1}{4}(3\alpha_1 + \beta_2) e_4 + \frac{1}{2}\alpha_2 e_5, \\
e_1 \cdot e_5 &= \frac{1}{2}\beta_1 e_4 + \frac{1}{4}(\alpha_1 + 3\beta_2) e_5, \\
e_2 \cdot e_2 &= \beta_6 e_1 + \beta_7 e_2 + \beta_8 e_3 + \beta_9 e_4 + \beta_{10} e_5, \\
e_2 \cdot e_3 &= \frac{1}{2}(\beta_1 + \beta_7) e_3 + \frac{1}{2}\beta_8 e_4 - \frac{1}{2}\beta_3 e_5, \\
e_2 \cdot e_4 &= \frac{1}{4}(3\beta_1 + \beta_7) e_4 + \frac{1}{2}\beta_2 e_5, \\
e_2 \cdot e_5 &= \frac{1}{2}\beta_6 e_4 + \frac{1}{4}(\beta_1 + 3\beta_7) e_5, \\
e_3 \cdot e_3 &= \frac{1}{4}(\beta_1 + \beta_7) e_4 - \frac{1}{4}(\alpha_1 + \beta_2) e_5,
\end{aligned}
\right.$$

By using the multiplication of Lie algebra $g_{(5,1)}^1$, we consider the general basis change and express the new basis elements $\{e'_1, e'_2, e'_3, e'_4, e'_5\}$ via the basis elements $\{e_1, e_2, e_3, e_4, e_5\}$ as follows:

$$\begin{aligned}
e'_1 &= \sum_{j=1}^5 A_j e_j, \quad e'_2 = \sum_{j=1}^5 B_j e_j, \quad e'_3 = \Delta_{1,2} e_3 + \Delta_{1,3} e_4 + \Delta_{2,3} e_5, \\
e'_4 &= \Delta_{1,2} (A_1 e_4 + A_2 e_5), \quad e'_5 = \Delta_{1,2} (B_1 e_4 + B_2 e_5),
\end{aligned}$$

where $\Delta_{i,j} = A_i B_j - A_j B_i$ and $\Delta_{1,2} \neq 0$.

Consider the multiplication $e'_1 \cdot e'_1 = \alpha'_1 e'_1 + \alpha'_2 e'_2 + \alpha'_3 e'_3 + \alpha'_4 e'_4 + \alpha'_5 e'_5$:

$$\begin{aligned}
e'_1 \cdot e'_1 &= \left(\sum_{j=1}^5 A_j e_j \right) \cdot \left(\sum_{j=1}^5 A_j e_j \right) \\
&= (A_1^2 \alpha_1 + 2A_1 A_2 \beta_1 + A_2^2 \beta_6) e_1 + (A_1^2 \alpha_2 + 2A_1 A_2 \beta_2 + A_2^2 \beta_7) e_2 + (*).
\end{aligned}$$

On the other hand, we have

$$\alpha'_1 e'_1 + \alpha'_2 e'_2 + \alpha'_3 e'_3 + \alpha'_4 e'_4 + \alpha'_5 e'_5 = \alpha'_1 \left(\sum_{j=1}^5 A_j e_j \right) + \alpha'_2 \left(\sum_{j=1}^5 B_j e_j \right) + (*)$$

$$= (A_1\alpha'_1 + B_1\alpha'_2)e_1 + (A_2\alpha'_1 + B_2\alpha'_2)e_2 + (*).$$

Now, we consider the multiplication $e'_1 \cdot e'_2 = \beta'_1 e'_1 + \beta'_2 e'_2 + \beta'_3 e'_3 + \beta'_4 e'_4 + \beta'_5 e'_5$:

$$e'_1 \cdot e'_2 = \left(\sum_{j=1}^5 A_j e_j \right) \cdot \left(\sum_{j=1}^5 B_j e_j \right) =$$

$$= (A_1 B_1 \alpha_1 + (A_1 B_2 + A_2 B_1) \beta_1 + A_2 B_2 \beta_6) e_1 + (A_1 B_1 \alpha_2 + (A_1 B_2 + A_2 B_1) \beta_2 + A_2 B_2 \beta_7) e_2 + (*).$$

On the other hand, we get

$$\begin{aligned} \beta'_1 e'_1 + \beta'_2 e'_2 + \beta'_3 e'_3 + \beta'_4 e'_4 + \beta'_5 e'_5 &= \beta'_1 \left(\sum_{j=1}^5 A_j e_j \right) + \beta'_2 \left(\sum_{j=1}^5 B_j e_j \right) + (*) \\ &= (A_1 \beta'_1 + B_1 \beta'_2) e_1 + (A_2 \beta'_1 + B_2 \beta'_2) e_2 + (*). \end{aligned}$$

Finally, we consider the multiplication $e'_2 \cdot e'_2 = \beta'_6 e'_1 + \beta'_7 e'_2 + \beta'_8 e'_3 + \beta'_9 e'_4 + \beta'_{10} e'_5$:

$$\begin{aligned} e'_2 \cdot e'_2 &= \left(\sum_{j=1}^5 B_j e_j \right) \cdot \left(\sum_{j=1}^5 B_j e_j \right) = \\ &= (B_1^2 \alpha_1 + 2B_1 B_2 \beta_1 + B_2^2 \beta_6) e_1 + (B_1^2 \alpha_2 + 2B_1 B_2 \beta_2 + B_2^2 \beta_7) e_2 + (*). \end{aligned}$$

Thus, we obtain

$$\begin{aligned} \beta'_6 e'_1 + \beta'_7 e'_2 + \beta'_8 e'_3 + \beta'_9 e'_4 + \beta'_{10} e'_5 &= \beta'_6 \left(\sum_{j=1}^5 A_j e_j \right) + \beta'_7 \left(\sum_{j=1}^5 B_j e_j \right) + (*) \\ &= (A_1 \beta'_6 + B_1 \beta'_7) e_1 + (A_2 \beta'_6 + B_2 \beta'_7) e_2 + (*). \end{aligned}$$

where $(*) \in \text{Span}\langle e_3, e_4, e_5 \rangle$. By verifying all the above multiplications in the basis elements e_1 and e_2 , we obtain the relations between parameters $x' = (\alpha'_1, \alpha'_2, \beta'_1, \beta'_2, \beta'_6, \beta'_7)$ and $x = (\alpha_1, \alpha_2, \beta_1, \beta_2, \beta_6, \beta_7)$:

$$\begin{cases} A_1 \alpha'_1 + B_1 \alpha'_2 = A_1^2 \alpha_1 + 2A_1 A_2 \beta_1 + A_2^2 \beta_6, \\ A_2 \alpha'_1 + B_2 \alpha'_2 = A_1^2 \alpha_2 + 2A_1 A_2 \beta_2 + A_2^2 \beta_7, \\ A_1 \beta'_1 + B_1 \beta'_2 = A_1 B_1 \alpha_1 + (A_1 B_2 + A_2 B_1) \beta_1 + A_2 B_2 \beta_6, \\ A_2 \beta'_1 + B_2 \beta'_2 = A_1 B_1 \alpha_2 + (A_1 B_2 + A_2 B_1) \beta_2 + A_2 B_2 \beta_7, \\ A_1 \beta'_6 + B_1 \beta'_7 = B_1^2 \alpha_1 + 2B_1 B_2 \beta_1 + B_2^2 \beta_6, \\ A_2 \beta'_6 + B_2 \beta'_7 = B_1^2 \alpha_2 + 2B_1 B_2 \beta_2 + B_2^2 \beta_7. \end{cases} \quad (6)$$

We can write the system (6) as follows:

$$x' \mathbb{A} = x \mathbb{B},$$

where $\det \mathbb{A} = \det \mathbb{B} = (\Delta_{1,2})^3$. This means that the system (6) has only a zero solution $x' = \mathbf{0}$ if and only if $x = \mathbf{0}$.

So we have the following possibility cases.

- (1) Let $x = \mathbf{0}$. Then, from the identity $e_1 \cdot (e_1 \cdot e_2) = (e_1 \cdot e_1) \cdot e_2$, we derive $\beta_3^2 - \alpha_3\beta_8 = 0$. Again, by considering the products of $e'_1 \cdot e'_1 = \alpha'_3 e'_3 + \alpha'_4 e'_4 + \alpha'_5 e'_5$, $e'_1 \cdot e'_2 = \beta'_3 e'_3 + \beta'_4 e'_4 + \beta'_5 e'_5$ and $e'_2 \cdot e'_2 = \beta'_8 e'_3 + \beta'_9 e'_4 + \beta'_{10} e'_5$, we have the following expressions:

$$\begin{aligned}\alpha'_3 &= \frac{A_1^2\alpha_3 + 2A_1A_2\beta_3 + A_2^2\beta_8}{\Delta_{1,2}}, \\ \beta'_3 &= \frac{A_1B_1\alpha_3 + (A_1B_2 + A_2B_1)\beta_3 + A_2B_2\beta_8}{\Delta_{1,2}}, \\ \beta'_8 &= \frac{B_1^2\alpha_3 + 2B_1B_2\beta_3 + B_2^2\beta_8}{\Delta_{1,2}}.\end{aligned}$$

From the above expressions we can conclude that $y = (\alpha_3, \beta_3, \beta_8)$ is zero invariant.

- (a) If $y = \mathbf{0}$, then we have the algebra $\mathbf{TP}_1(g_{(5,1)}^1)$.
- (b) If $y \neq \mathbf{0}$, then without loss of generality, we assume that $\alpha'_3 = 1$. Thus, we derive $\beta_8 = \beta_3^2$ and obtain the algebra $\mathbf{TP}_2(g_{(5,1)}^1)$.
- (2) Let $x \neq \mathbf{0}$. Then, the system (6) can be written as follows:

$$\begin{aligned}\alpha'_1 &= \frac{A_1^2B_2\alpha_1 + 2A_1A_2B_2\beta_1 + A_2^2B_2\beta_6 - A_1^2B_1\alpha_2 - 2A_1A_2B_1\beta_2 - A_2^2B_1\beta_7}{\Delta_{1,2}}, \\ \alpha'_2 &= \frac{A_1^2A_2\alpha_1 + 2A_1A_2^2\beta_1 + A_2^3\beta_6 - A_1^3\alpha_2 - 2A_1^2A_2\beta_2 - A_1A_2^2\beta_7}{-\Delta_{1,2}}, \\ \beta'_1 &= \frac{A_1B_1B_2\alpha_1 + (A_1B_2 + A_2B_1)B_2\beta_1 + A_2B_2^2\beta_6 - A_1B_1^2\alpha_2 - (A_1B_2 + A_2B_1)B_1\beta_2 - A_2B_1B_2\beta_7}{\Delta_{1,2}}, \\ \beta'_2 &= \frac{A_1A_2B_1\alpha_1 + (A_1B_2 + A_2B_1)A_2\beta_1 + A_2^2B_2\beta_6 - A_1^2B_1\alpha_2 - (A_1B_2 + A_2B_1)A_1\beta_2 - A_1A_2B_2\beta_7}{-\Delta_{1,2}}, \\ \beta'_6 &= \frac{B_1^2B_2\alpha_1 + 2B_1B_2^2\beta_1 + B_2^3\beta_6 - B_1^3\alpha_2 - 2B_1^2B_2\beta_2 - B_1B_2^2\beta_7}{\Delta_{1,2}}, \\ \beta'_7 &= \frac{A_2B_1^2\alpha_1 + 2A_2B_1B_2\beta_1 + A_2B_2^2\beta_6 - A_1B_1^2\alpha_2 - 2A_1B_1B_2\beta_2 - A_1B_2^2\beta_7}{-\Delta_{1,2}}.\end{aligned}$$

It can be shown that $\alpha'_2 = 1$ and by considering the associative identity $x \cdot (y \cdot z) = (x \cdot y) \cdot z$, we obtain the following restrictions on structure constants:

$$\begin{aligned}\{e_5, e_1, e_1\}, \quad &\Rightarrow \quad \beta_6 = \beta_1\beta_2, \quad \beta_7 = \frac{1}{4}(3\beta_2^2 - 2\alpha_1\beta_2 - \alpha_1^2), \\ \{e_1, e_1, e_3\}, \quad &\Rightarrow \quad \beta_1 = -\frac{1}{4}(\alpha_1 - \beta_2)^2, \quad \beta_3 = \frac{1}{2}\alpha_3(\beta_2 - \alpha_1), \quad \beta_8 = \frac{1}{4}\alpha_3(\alpha_1 - \beta_2)^2, \\ \{e_2, e_1, e_1\}, \quad &\Rightarrow \quad \beta_{10} = \frac{1}{4} \left(-2\alpha_4\beta_2 - 3\alpha_5\beta_2^2 + 2\beta_4 + 3\alpha_1(\alpha_5\beta_2 - \beta_5) + 7\beta_2\beta_5 \right), \\ &\quad \beta_9 = \frac{1}{8} \left(\alpha_1^2(\alpha_5\beta_2 - \beta_5) + \beta_2(-2\alpha_4\beta_2 + \alpha_5\beta_2^2 + 10\beta_4 - \beta_2\beta_5) + \right. \\ &\quad \left. + 2\alpha_1(\alpha_4\beta_2 - \alpha_5\beta_2^2 - \beta_4 + \beta_2\beta_5) \right).\end{aligned}$$

Hence, the algebra $\mathbf{TP}_3(g_{(5,1)}^1)$ is obtained. □

Theorem 17. Let $(g_{(n,1)}^1, \cdot, [-, -])$ be a transposed Poisson algebra structure defined on the Lie algebra $g_{(n,1)}^1$ and $n \geq 7$. Then, the multiplication of $(g_{(n,1)}^1, \cdot)$ has the following form:

$$\begin{aligned} \mathbf{TP}_1(g_{(n,1)}^1) : & \begin{cases} e_1 \cdot e_1 = \sum_{j=4}^n \alpha_j e_j, \quad e_1 \cdot e_2 = \beta_1 e_{n-2} + \beta_2 e_{n-1} + \beta_3 e_n, \\ e_1 \cdot e_3 = \frac{1}{2} \beta_1 e_{n-1} - \frac{1}{2} \alpha_{n-2} e_n, \quad e_1 \cdot e_j = \frac{(-1)^j}{2} \alpha_{n-j+1} e_n, \quad 4 \leq j \leq n-3, \\ e_2 \cdot e_2 = \beta_4 e_{n-2} + \beta_5 e_{n-1} + \beta_6 e_n, \quad e_2 \cdot e_3 = \frac{1}{2} \beta_4 e_{n-1} - \frac{1}{2} \beta_1 e_n; \end{cases} \\ \mathbf{TP}_2(g_{(n,1)}^1) : & \begin{cases} e_1 \cdot e_1 = e_3 + \sum_{j=4}^n \alpha_j e_j, \quad e_1 \cdot e_2 = \beta_1 e_{n-2} + \beta_2 e_{n-1} + \beta_3 e_n, \\ e_1 \cdot e_3 = \frac{1}{2} \beta_1 e_{n-1} - \frac{1}{2} \alpha_{n-2} e_n, \quad e_1 \cdot e_j = \frac{(-1)^j}{2} \alpha_{n-j+1} e_n, \quad 4 \leq j \leq n-3, \\ e_1 \cdot e_{n-2} = -\frac{1}{2} e_n, \quad e_2 \cdot e_2 = \beta_4 e_{n-1} + \beta_5 e_n, \quad e_2 \cdot e_3 = -\frac{1}{2} \beta_1 e_n; \end{cases} \\ \mathbf{TP}_3(g_{(n,1)}^1) : & \begin{cases} e_1 \cdot e_1 = e_2 + \sum_{j=3}^n \alpha_j e_j, \quad e_1 \cdot e_2 = 2\beta_1 e_{n-1} + \beta_2 e_n, \\ e_1 \cdot e_3 = -\frac{1}{2} \alpha_{n-2} e_n, \quad e_1 \cdot e_j = \frac{(-1)^j}{2} \alpha_{n-j+1} e_n, \quad 4 \leq j \leq n-2, \\ e_1 \cdot e_{n-1} = \frac{1}{2} e_n, \quad e_2 \cdot e_2 = \beta_1 e_n; \end{cases} \end{aligned}$$

where the transposed Poisson algebra has its products with respect to the bracket $[-, -]$, and the remaining products are equal to zero.

Proof. Let $(g_{(n,1)}^1, \cdot, [-, -])$ be a transposed Poisson algebra structure defined on the Lie algebra $g_{(n,1)}^1$. Then, for any element $x \in g_{(n,1)}^1$, we have the operator of multiplication $\varphi_x(y) = x \cdot y$ is a $\frac{1}{2}$ -derivation. Hence, for $1 \leq i \leq n$ we derive by Theorem 10:

$$\begin{cases} \varphi_{e_i}(e_1) = \sum_{j=1}^n \alpha_{i,j} e_j, \\ \varphi_{e_i}(e_2) = \alpha_{i,1} e_2 + \beta_{i,n-2} e_{n-2} + \beta_{i,n-1} e_{n-1} + \beta_{i,n} e_n, \\ \varphi_{e_i}(e_3) = \alpha_{i,1} e_3 + \frac{1}{2} \beta_{i,n-2} e_{n-1} - \frac{1}{2} \alpha_{i,n-2} e_n, \\ \varphi_{e_i}(e_j) = \alpha_{i,1} e_j + \frac{(-1)^j}{2} \alpha_{i,n-j+1} e_n, \quad 4 \leq j \leq n-1, \\ \varphi_{e_i}(e_n) = \alpha_{i,1} e_n. \end{cases}$$

By considering equality $\varphi_{e_i}(e_j) = e_i \cdot e_j = e_j \cdot e_i = \varphi_{e_j}(e_i)$, we have following restrictions:

$$\begin{aligned} \{e_1, e_2\}, \quad & \Rightarrow \alpha_{2,1} = 0, \quad \alpha_{2,2} = \alpha_{1,1}, \quad \alpha_{2,t} = 0, \quad 3 \leq t \leq n-3, \\ & \quad \alpha_{2,n-2} = \beta_{1,n-2}, \quad \alpha_{2,n-1} = \beta_{1,n-1}, \quad \alpha_{2,n} = \beta_{1,n}, \\ \{e_1, e_3\}, \quad & \Rightarrow \alpha_{3,1} = \alpha_{3,2} = 0, \quad \alpha_{3,3} = \alpha_{1,1}, \quad \alpha_{3,t} = 0, \quad 4 \leq t \leq n-2, \\ & \quad \alpha_{3,n-1} = \frac{1}{2} \beta_{1,n-2}, \quad \alpha_{3,n} = -\frac{1}{2} \beta_{1,n}, \\ \{e_1, e_j\}, \quad & \Rightarrow \alpha_{j,t} = 0, \quad 1 \leq t \leq j-1, \quad \alpha_{j,j} = \alpha_{1,1}, \\ 4 \leq j \leq n-1, \quad & \alpha_{j,n} = \frac{(-1)^j}{2} \alpha_{1,n-j+1}, \quad \alpha_{j,t} = 0, \quad j+1 \leq t \leq n-1, \\ \{e_1, e_n\}, \quad & \Rightarrow \alpha_{n,t} = 0, \quad 1 \leq t \leq n-1, \quad \alpha_{n,n} = \alpha_{1,1}, \\ \{e_2, e_3\}, \quad & \Rightarrow \beta_{3,n-2} = 0, \quad \beta_{3,n-1} = \frac{1}{2} \beta_{2,n-2}, \quad \beta_{3,n} = -\frac{1}{2} \beta_{1,n-2}, \\ \{e_2, e_j\}, \quad & \Rightarrow \beta_{j,n-2} = \beta_{j,n-1} = \beta_{j,n} = 0, \\ 4 \leq j \leq n-1, \quad & \\ \{e_2, e_n\}, \quad & \Rightarrow \beta_{n,n-2} = \beta_{n,n-1} = \beta_{n,n} = 0, \end{aligned}$$

$$\{e_3, e_j\}, \quad \Rightarrow \quad \alpha_{3,t} = 0, \quad 2 \leq t \leq n-3, \\ 4 \leq j \leq n-1.$$

Thus, we have the following table of commutative multiplications of the transposed Poisson algebra structure $(g_{(n,1)}^1, \cdot, [\cdot, \cdot])$:

$$\begin{cases} e_1 \cdot e_1 = \sum_{j=2}^n \alpha_j e_j, \\ e_1 \cdot e_2 = \beta_1 e_{n-2} + \beta_2 e_{n-1} + \beta_3 e_n, \\ e_1 \cdot e_3 = \frac{1}{2} \beta_1 e_{n-1} - \frac{1}{2} \alpha_{n-2} e_n, \\ e_1 \cdot e_j = \frac{(-1)^j}{2} \alpha_{n-j+1} e_n, \quad 4 \leq j \leq n-1, \\ e_2 \cdot e_2 = \beta_4 e_{n-2} + \beta_5 e_{n-1} + \beta_6 e_n, \\ e_2 \cdot e_3 = \frac{1}{2} \beta_4 e_{n-1} - \frac{1}{2} \beta_1 e_n. \end{cases} \quad (7)$$

Consider the associative identity $x \cdot (y \cdot z) = (x \cdot y) \cdot z$ in (7), we obtain the following restrictions on structure constants:

$$\begin{aligned} \{e_1, e_2, e_3\}, \quad &\Rightarrow \quad \alpha_2 \beta_4 = 0, \\ \{e_1, e_2, e_2\}, \quad &\Rightarrow \quad \alpha_3 \beta_4 - \alpha_2 \beta_5 = 0, \\ \{e_1, e_1, e_3\}, \quad &\Rightarrow \quad \alpha_2 \beta_1 = 0, \\ \{e_1, e_1, e_2\}, \quad &\Rightarrow \quad 2\alpha_2 \beta_5 + \alpha_3 \beta_4 = 0, \quad \alpha_2(2\beta_6 - \beta_2) = 0, \\ \{e_1, e_1, e_i\}, \quad &\Rightarrow \quad \alpha_2 \beta_4 = 0, \quad \alpha_3 \beta_4 = 0, \quad \alpha_2 \beta_5 = 0, \quad \alpha_2 \beta_1 = 0, \quad \alpha_2(2\beta_6 - \beta_2) = 0, \\ &4 \leq i \leq n-1. \end{aligned}$$

Now we consider the general change of basis:

$$e'_1 = \sum_{j=1}^n A_j e_j, \quad e'_2 = \sum_{j=1}^n B_j e_j, \quad e'_{i+1} = [e'_1, e'_i], \quad 2 \leq i \leq n-2, \quad e'_n = [e'_2, e'_{n-2}],$$

where $B_1 = 0$ and $A_1 B_2 \neq 0$. Then, from the multiplication $e_1 \cdot e_1 = \sum_{j=2}^n \alpha_j e_j$, we discover that the structure constant of the α_2 changes as follows:

$$\alpha'_2 = \frac{A_1^2}{B_2} \alpha_2.$$

Thus, we have the following cases.

- (1) If $\alpha_2 = 0$. Again, by using a change of basis and from the product $e_1 \cdot e_1 = \sum_{j=3}^n \alpha_j e_j$, we obtain the following relation:

$$\alpha'_3 = \frac{A_1}{B_2} \alpha_3.$$

Now, we consider the following subcases.

- (a) If $\alpha_3 = 0$. Then, we have the algebra $\mathbf{TP}_1(g_{(n,1)}^1)$.
(b) If $\alpha_3 \neq 0$, then, we have $\beta_4 = 0$, and via transformation

$$\phi(e_1) = e_1, \phi(e_i) = \alpha_3 e_i, 2 \leq i \leq n-1, \phi(e_n) = \alpha_3^2 e_n,$$

we get the algebra $\mathbf{TP}_2(g_{(n,1)}^1)$.

- (2) If $\alpha_2 \neq 0$, then we derive $\beta_4 = \beta_5 = \beta_1 = 0$, $\beta_2 = 2\beta_6$, and we via transformation

$$\phi(e_1) = e_1, \phi(e_i) = \alpha_2 e_i, 2 \leq i \leq n-1, \phi(e_n) = \alpha_2^2 e_n,$$

we obtain the algebra $\mathbf{TP}_3(g_{(n,1)}^1)$.

□

Theorem 18. Let $(g_{(5,1)}^2, \cdot, [-, -])$ be a transposed Poisson algebra structure defined on the Lie algebra $g_{(5,1)}^2$. Then the multiplication of $(g_{(5,1)}^2, \cdot)$ has the following form:

$$\begin{aligned} \mathbf{TP}_1(g_{(5,1)}^2) : & \begin{cases} e_1 \cdot e_1 = e_2 + \alpha_3 e_3 + \alpha_4 e_4 + \alpha_5 e_5, & e_1 \cdot e_2 = (\alpha_5(1 - 2\alpha_{16}) - 2\alpha_{15})e_3 + \alpha_8 e_4, \\ e_1 \cdot e_3 = -(\alpha_{15} + \alpha_5 \alpha_{16})e_4, & e_1 \cdot e_5 = -e_3 + \alpha_{10} e_4, \\ e_2 \cdot e_2 = (2(\alpha_{15} + \alpha_5 \alpha_{16})^2 - \alpha_5(2\alpha_{15} + \alpha_{16}))e_4, & e_2 \cdot e_5 = \alpha_{15} e_4, e_5 \cdot e_5 = \alpha_{16} e_4; \end{cases} \\ \mathbf{TP}_2(g_{(5,1)}^2) : & \begin{cases} e_1 \cdot e_1 = \alpha_3 e_3 + \alpha_4 e_4, & e_1 \cdot e_2 = \alpha_8 e_4, e_1 \cdot e_5 = \alpha_{10} e_4, \\ e_2 \cdot e_2 = \alpha_{13} e_4, & e_2 \cdot e_5 = \alpha_{15} e_4, e_5 \cdot e_5 = e_4; \end{cases} \\ \mathbf{TP}_3(g_{(5,1)}^2) : & \begin{cases} e_1 \cdot e_1 = \alpha_7^2 e_3 + \alpha_4 e_4, & e_1 \cdot e_2 = \alpha_7 e_3 + \alpha_8 e_4, e_1 \cdot e_3 = \frac{1}{2}\alpha_7 e_4, e_1 \cdot e_5 = \alpha_{10} e_4, \\ e_2 \cdot e_2 = e_3 + \alpha_{13} e_4, & e_2 \cdot e_3 = \frac{1}{2}e_4, e_2 \cdot e_5 = \alpha_{15} e_4, e_5 \cdot e_5 = e_4; \\ e_1 \cdot e_1 = \frac{(\alpha_7^2 - \alpha_5(\alpha_7 + 2\alpha_{15}))}{\alpha_{12}} e_3 + \alpha_4 e_4 + \alpha_5 e_5, & \\ e_1 \cdot e_2 = \alpha_7 e_3 + \alpha_8 e_4 + \alpha_9 e_5, & e_1 \cdot e_3 = \frac{1}{2}(\alpha_7 - \alpha_5)e_4, e_1 \cdot e_5 = \frac{1}{2}\alpha_5 \alpha_{12} e_4, \\ e_2 \cdot e_2 = \alpha_{12} e_3 + \alpha_{13} e_4 + e_5, & e_2 \cdot e_3 = \frac{1}{2}\alpha_{12} e_4, e_2 \cdot e_5 = \alpha_{15} e_4; \end{cases} \\ \mathbf{TP}_4(g_{(5,1)}^2) : & \begin{cases} e_1 \cdot e_1 = \alpha_3 e_3 + \alpha_4 e_4 + \alpha_5 e_5, & e_1 \cdot e_2 = \alpha_7 e_3 + \alpha_8 e_4, e_1 \cdot e_3 = \frac{1}{2}\alpha_7 e_4, e_1 \cdot e_5 = \frac{1}{2}\alpha_5 \alpha_{12} e_4, \\ e_2 \cdot e_2 = \alpha_{12} e_3 + \alpha_{13} e_4 + e_5, & e_2 \cdot e_3 = \frac{1}{2}\alpha_{12} e_4, e_2 \cdot e_5 = \alpha_{15} e_4; \end{cases} \\ \mathbf{TP}_5(g_{(5,1)}^2) : & \begin{cases} e_1 \cdot e_1 = \alpha_3 e_3 + \alpha_4 e_4 + \alpha_5 e_5, & e_1 \cdot e_2 = \alpha_7 e_3 + \alpha_8 e_4, \\ e_1 \cdot e_3 = \frac{1}{2}(\alpha_7 - \alpha_5)e_4, & e_2 \cdot e_2 = \alpha_{13} e_4 + e_5, e_2 \cdot e_5 = \frac{\alpha_7(\alpha_7 - \alpha_5)}{2\alpha_5} e_4; \end{cases} \\ \mathbf{TP}_6(g_{(5,1)}^2) : & \begin{cases} e_1 \cdot e_1 = \alpha_3 e_3 + \alpha_4 e_4, & e_1 \cdot e_2 = \alpha_8 e_4, e_2 \cdot e_2 = \alpha_{13} e_4 + e_5, e_2 \cdot e_5 = \alpha_{15} e_4; \\ e_1 \cdot e_1 = \alpha_3 e_3 + \alpha_4 e_4, & e_1 \cdot e_2 = 2\alpha_{15} e_3 + \alpha_8 e_4 + e_5, \end{cases} \\ \mathbf{TP}_7(g_{(5,1)}^2) : & \begin{cases} e_1 \cdot e_3 = \frac{1}{2}\alpha_{15} e_4, & e_1 \cdot e_5 = \frac{1}{2}(\alpha_3(\alpha_{12} - 1) - 4\alpha_{15}^2) e_4, \\ e_2 \cdot e_2 = \alpha_{12} e_3 + \alpha_{13} e_4, & e_2 \cdot e_3 = \frac{1}{2}(\alpha_{12} - 1)e_4, e_2 \cdot e_5 = \alpha_{15} e_4; \end{cases} \\ \mathbf{TP}_8(g_{(5,1)}^2) : & \begin{cases} e_1 \cdot e_1 = \alpha_7^2 e_3 + \alpha_4 e_4, & e_1 \cdot e_2 = \alpha_7 e_3 + \alpha_8 e_4, e_1 \cdot e_3 = \frac{1}{2}\alpha_7 e_4, \\ e_1 \cdot e_5 = \alpha_{10} e_4, & e_2 \cdot e_2 = e_3 + \alpha_{13} e_4, e_2 \cdot e_3 = \frac{1}{2}e_4, e_2 \cdot e_5 = \alpha_{15} e_4; \end{cases} \\ \mathbf{TP}_9(g_{(5,1)}^2) : & \begin{cases} e_1 \cdot e_1 = \alpha_3 e_3 + \alpha_4 e_4 + \alpha_5 e_5, & e_1 \cdot e_2 = \alpha_7 e_3 + \alpha_8 e_4, e_1 \cdot e_3 = \frac{1}{2}(\alpha_7 - \alpha_5)e_4, \\ e_1 \cdot e_5 = \alpha_{10} e_4, & e_2 \cdot e_2 = \alpha_{13} e_4, e_2 \cdot e_5 = \frac{\alpha_7(\alpha_7 - \alpha_5)}{2\alpha_5} e_4; \end{cases} \end{aligned}$$

TP₁₀($g_{(5,1)}^2$): $e_1 \cdot e_1 = \alpha_3 e_3 + \alpha_4 e_4$, $e_1 \cdot e_2 = \alpha_8 e_4$, $e_1 \cdot e_5 = \alpha_{10} e_4$, $e_2 \cdot e_2 = \alpha_{13} e_4$, $e_2 \cdot e_5 = \alpha_{15} e_4$; where the transposed Poisson algebra has its products with respect to the bracket $[-, -]$, and remaining products are equal to zero.

Proof. Let $(g_{(5,1)}^2, \cdot, [-, -])$ be a transposed Poisson algebra structure defined on Lie algebra $g_{(5,1)}^2$. Then, for any element of $x \in g_{(5,1)}^2$, we have operator of multiplication $\varphi_x(y) = x \cdot y$ is a $\frac{1}{2}$ -derivation. Hence, by using Theorem 11 1 $\leq i \leq n$, we derive

$$\begin{cases} \varphi_{e_i}(e_1) = \alpha_{i,1}e_1 + \alpha_{i,2}e_2 + \alpha_{i,3}e_3 + \alpha_{i,4}e_4 + \alpha_{i,5}e_5, \\ \varphi_{e_i}(e_2) = \beta_{i,2}e_2 + \beta_{i,3}e_3 + \beta_{i,4}e_4 + \beta_{i,5}e_5, \\ \varphi_{e_i}(e_3) = \frac{1}{2}(\alpha_{i,1} + \beta_{i,2})e_3 + \frac{1}{2}(\beta_{i,3} - \alpha_{i,5})e_4, \\ \varphi_{e_i}(e_4) = \frac{1}{4}(3\alpha_{i,1} + \beta_{i,2})e_4, \\ \varphi_{e_i}(e_5) = -\alpha_{i,2}e_3 + \gamma_{i,4}e_4 + \frac{1}{2}(3\alpha_{i,1} - \beta_{i,2})e_5, \end{cases}$$

By considering the property of $\varphi_{e_i}(e_j) = e_i \cdot e_j = e_j \cdot e_i = \varphi_{e_j}(e_i)$, we obtain the following restrictions:

$$\begin{aligned} \{e_1, e_2\}, &\Rightarrow \alpha_{2,1} = 0, \alpha_{2,2} = \beta_{1,2}, \alpha_{2,3} = \beta_{1,3}, \alpha_{2,4} = \beta_{1,4}, \alpha_{2,5} = \beta_{1,5}, \\ \{e_1, e_3\}, &\Rightarrow \alpha_{3,1} = \alpha_{3,2} = 0, \alpha_{3,3} = \frac{1}{2}(\alpha_{1,1} + \beta_{1,2}), \alpha_{3,4} = \frac{1}{2}(\beta_{1,3} - \alpha_{1,5}), \alpha_{3,5} = 0, \\ \{e_1, e_4\}, &\Rightarrow \alpha_{4,1} = \alpha_{4,2} = \alpha_{4,3} = 0, \alpha_{4,4} = \frac{1}{4}(3\alpha_{1,1} + \beta_{1,2}), \alpha_{4,5} = 0, \\ \{e_1, e_5\}, &\Rightarrow \alpha_{5,1} = \alpha_{5,2} = 0, \alpha_{5,3} = -\alpha_{1,2}, \alpha_{5,4} = \gamma_{1,4}, \alpha_{5,5} = \frac{1}{2}(3\alpha_{1,1} - \beta_{1,2}), \\ \{e_2, e_3\}, &\Rightarrow \beta_{3,2} = 0, \beta_{3,3} = \frac{1}{2}\beta_{2,2}, \beta_{3,4} = \frac{1}{2}(\beta_{2,3} - \beta_{1,3}), \beta_{3,5} = 0, \\ \{e_2, e_4\}, &\Rightarrow \beta_{4,2} = \beta_{4,3} = 0, \beta_{4,4} = \frac{1}{4}\beta_{2,2}, \beta_{4,5} = 0, \\ \{e_2, e_5\}, &\Rightarrow \beta_{5,2} = 0, \beta_{5,3} = -\beta_{1,2}, \beta_{5,4} = \gamma_{2,4}, \beta_{5,5} = -\frac{1}{2}\beta_{2,2}, \\ \{e_3, e_5\}, &\Rightarrow \gamma_{3,4} = -\frac{1}{4}(3\alpha_{1,1} + \beta_{1,2}), \\ \{e_4, e_5\}, &\Rightarrow \gamma_{4,4} = 0. \end{aligned}$$

Thus, we have the following table of commutative multiplications of the transposed Poisson algebra structure defined on $g_{(5,1)}^2$:

$$\begin{cases} e_1 \cdot e_1 = \alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3 + \alpha_4 e_4 + \alpha_5 e_5, e_1 \cdot e_2 = \alpha_6 e_2 + \alpha_7 e_3 + \alpha_8 e_4 + \alpha_9 e_5, \\ e_1 \cdot e_3 = \frac{1}{2}(\alpha_1 + \alpha_6)e_3 + \frac{1}{2}(\alpha_7 - \alpha_5)e_4, e_1 \cdot e_4 = \frac{1}{4}(3\alpha_1 + \alpha_6)e_4, \\ e_1 \cdot e_5 = -\alpha_2 e_3 + \alpha_{10} e_4 + \frac{1}{2}(3\alpha_1 - \alpha_6)e_5, e_2 \cdot e_2 = \alpha_{11} e_2 + \alpha_{12} e_3 + \alpha_{13} e_4 + \alpha_{14} e_5, \\ e_2 \cdot e_3 = \frac{1}{2}\alpha_{11} e_3 + \frac{1}{2}(\alpha_{12} - \alpha_9)e_4, e_2 \cdot e_4 = \frac{1}{4}\alpha_{11} e_4, e_2 \cdot e_5 = -\alpha_6 e_3 + \alpha_{15} e_4 + \frac{1}{2}\alpha_{11} e_5, \\ e_3 \cdot e_3 = \frac{1}{4}\alpha_{11} e_4, e_3 \cdot e_5 = -\frac{1}{4}(3\alpha_1 + \alpha_6)e_4, e_5 \cdot e_5 = \alpha_{16} e_4. \end{cases}$$

Now, consider the associative identity $x \cdot (y \cdot z) = (x \cdot y) \cdot z$, we obtain the following restrictions on structure constants:

$$\begin{aligned}\{e_4, e_2, e_2\}, &\Rightarrow \alpha_{11} = 0, \\ \{e_5, e_1, e_3\}, &\Rightarrow (\alpha_1 - \alpha_6)(3\alpha_1 + \alpha_6) = 0, \\ \{e_5, e_2, e_1\}, &\Rightarrow \alpha_6(\alpha_1 - \alpha_6) = 0, \\ \{e_5, e_5, e_2\}, &\Rightarrow \alpha_6(3\alpha_1 + \alpha_6) = 0.\end{aligned}$$

Hence, we can conclude that $\alpha_1 = \alpha_6 = 0$ and the table of multiplication of the algebra will have the following form:

$$\begin{cases} e_1 \cdot e_1 = \alpha_2 e_2 + \alpha_3 e_3 + \alpha_4 e_4 + \alpha_5 e_5, & e_1 \cdot e_2 = \alpha_7 e_3 + \alpha_8 e_4 + \alpha_9 e_5, \\ e_1 \cdot e_3 = \frac{1}{2}(\alpha_7 - \alpha_5)e_4, & e_1 \cdot e_5 = -\alpha_2 e_3 + \alpha_{10} e_4, & e_2 \cdot e_2 = \alpha_{12} e_3 + \alpha_{13} e_4 + \alpha_{14} e_5, \\ e_2 \cdot e_3 = \frac{1}{2}(\alpha_{12} - \alpha_9)e_4, & e_2 \cdot e_5 = \alpha_{15} e_4, & e_5 \cdot e_5 = \alpha_{16} e_4. \end{cases}$$

By considering the general change of the generators of basis

$$e'_1 = \sum_{t=1}^5 A_t e_t, \quad e'_2 = \sum_{t=1}^5 B_t e_t, \quad e'_5 = \sum_{t=1}^5 C_t e_t,$$

and from the product $e'_1 \cdot e'_1 = \alpha'_2 e'_2 + \alpha'_3 e'_3 + \alpha'_4 e'_4 + \alpha'_5 e'_5$, we derive

$$\alpha'_2 = \frac{A_1^2}{B_2} \alpha_2.$$

We have the following possible cases.

- (1) Let $\alpha_2 \neq 0$. Then by choosing $B_2 = A_1^2 \alpha_2$, we get $\alpha'_2 = 1$. By considering the associative identity $x \cdot (y \cdot z) = (x \cdot y) \cdot z$, we obtain the restrictions on structure constants as follows:

$$\begin{aligned}\{e_1, e_1, e_3\}, &\Rightarrow \alpha_{12} = \alpha_9, \\ \{e_1, e_1, e_2\}, &\Rightarrow \alpha_9 = \alpha_{14} = 0, \\ \{e_1, e_1, e_5\}, &\Rightarrow \alpha_7 = \alpha_5(1 - 2\alpha_{16}) - 2\alpha_{15}, \\ \{e_2, e_1, e_1\}, &\Rightarrow \alpha_{13} = 2\alpha_{15}^2 + 2\alpha_5 \alpha_{15}(-1 + 2\alpha_{16}) + \alpha_5^2 \alpha_{16}(-1 + 2\alpha_{16}).\end{aligned}$$

In this case, we have the algebra $\mathbf{TP}_1(g_{(5,1)}^2)$.

- (2) Let $\alpha_2 = 0$. Then, from the product $e'_5 \cdot e'_5 = \alpha'_{16} e'_4$, we obtain the following expression:

$$\alpha'_{16} = \frac{A_1^2}{B_2} \alpha_{16}.$$

- (a) If $\alpha_{16} \neq 0$, then by choosing $B_2 = A_1^2 \alpha_{16}$, we obtain $\alpha'_{16} = 1$. Considering the associative identity, we obtain the following restrictions:

$$\begin{cases} \{e_1, e_1, e_5\}, &\Rightarrow \alpha_5 = 0, \\ \{e_5, e_2, e_2\}, &\Rightarrow \alpha_{14} = 0, \end{cases} \quad \begin{cases} \{e_1, e_2, e_5\}, &\Rightarrow \alpha_9 = 0, \\ \{e_2, e_1, e_1\}, &\Rightarrow \alpha_7^2 = \alpha_3 \alpha_{12}. \end{cases}$$

Again, using the change of basis for the product $e'_2 \cdot e'_2 = \alpha'_{12}e'_3 + \alpha'_{13}e'_4$, we will get the following expression:

$$\alpha'_{12} = A_1\alpha_{12}.$$

If $\alpha_{12} = 0$, then we get $\alpha_7 = 0$ and obtain the algebra $\text{TP}_2(g_{(5,1)}^2)$.

If $\alpha_{12} \neq 0$, then by choosing $A_1 = \frac{1}{\alpha_{12}}$, we can assume $\alpha'_{12} = 1$, $\alpha_3 = \alpha_7^2$. Therefore, the algebra $\text{TP}_3(g_{(5,1)}^2)$ is obtained.

(b) If $\alpha_{16} = 0$, then apply the change of basis we get

$$\alpha'_{14} = \frac{B_2^2}{A_1^2}\alpha_{14}, \quad \alpha'_9 = \frac{B_2(A_1\alpha_9 + A_2\alpha_{14})}{A_1^2}, \quad \alpha'_{12} = \frac{B_2(A_1\alpha_{12} + A_2\alpha_{14})}{A_1^2}.$$

(i) If $\alpha_{14} \neq 0$ then by choosing $A_1^2 = B_2^2\alpha_{14}$, $A_2 = -\frac{A_1\alpha_9}{\alpha_{14}}$, we obtain $\alpha'_9 = 0$, $\alpha'_{14} = 1$. Again, using the change of basis elements of algebra, we have the following expression:

$$\alpha'_5 = \alpha_5, \quad \alpha'_{12} = \frac{B_2}{A_1}\alpha_{12}, \quad B_2^2 = A_1^2.$$

By considering the associative identity for the triples $\{e_1, e_1, e_2\}$ va $\{e_1, e_2, e_2\}$, we get

$$2\alpha_{10} = \alpha_5\alpha_{12}, \quad \alpha_5\alpha_7 - \alpha_7^2 + \alpha_3\alpha_{12} + 2\alpha_5\alpha_{15} = 0,$$

respectively. we consider following cases.

$\alpha_{12} \neq 0$, then we have the algebra $\text{TP}_4(g_{(5,1)}^2)$.

$\alpha_{12} = 0$ and $\alpha_5 \neq 0$, then we have the algebra $\text{TP}_5(g_{(5,1)}^2)$.

$\alpha_{12} = \alpha_5 = 0$, then we have the algebra $\text{TP}_6(g_{(5,1)}^2)$.

(ii) If $\alpha_{14} = 0$ then we derive

$$\alpha'_5 = \alpha_5 + \frac{2A_2}{A_1}\alpha_9, \quad \alpha'_9 = \frac{B_2}{A_1}\alpha_9, \quad \alpha'_{12} = \frac{B_2}{A_1}\alpha_{12}.$$

By using the associative identity for the triples $\{e_1, e_1, e_2\}$ and $\{e_1, e_2, e_2\}$, we obtain

$$2\alpha_7\alpha_9 = \alpha_5\alpha_{12} + 2\alpha_9\alpha_{15},$$

$$\alpha_5\alpha_7 - \alpha_7^2 - \alpha_3\alpha_9 - 2\alpha_9\alpha_{10} + \alpha_3\alpha_{12} + 2\alpha_5\alpha_{15} = 0.$$

Now we consider the following distinct cases:

$\alpha_9 \neq 0$, then by choosing $A_1 = B_2\alpha_9$, $A_2 = -\frac{B_2\alpha_5}{2}$, we can assume $\alpha'_5 = 0$, $\alpha'_9 = 1$ and obtain the algebra $\text{TP}_7(g_{(5,1)}^2)$.

$\alpha_9 = 0$, $\alpha_{12} \neq 0$, then by selecting $A_1 = B_2\alpha_{12}$, we get $\alpha'_{12} = 1$ and the algebra $\text{TP}_8(g_{(5,1)}^2)$.

$\alpha_9 = 0$, $\alpha_{12} = 0$, $\alpha_5 \neq 0$, then we have the algebra $\text{TP}_9(g_{(5,1)}^2)$.

$\alpha_9 = \alpha_{12} = \alpha_5 = 0$, then we have the algebra $\text{TP}_{10}(g_{(5,1)}^2)$.

□

Theorem 19. Let $(g_{(6,1)}^2, \cdot, [-, -])$ be a transposed Poisson algebra structure defined on the Lie algebra $g_{(6,1)}^2$. Then the multiplication of $(g_{(6,1)}^2, \cdot)$ has the following form:

$$\begin{aligned} \mathbf{TP}_1(g_{(6,1)}^2) : & \begin{cases} e_1 \cdot e_1 = e_2 + \alpha_2 e_3 + \alpha_3 e_4 + \alpha_4 e_5, & e_1 \cdot e_2 = -2\alpha_{11} e_4 + \alpha_7 e_5, & e_1 \cdot e_3 = -\alpha_{11} e_5, \\ e_1 \cdot e_6 = -e_3 - \alpha_2 e_4 + \alpha_8 e_5, & e_2 \cdot e_6 = \alpha_{11} e_5, & e_6 \cdot e_6 = e_4 + \alpha_{12} e_5; \end{cases} \\ \mathbf{TP}_2(g_{(6,1)}^2) : & \begin{cases} e_1 \cdot e_1 = e_3 + \alpha_3 e_4 + \alpha_4 e_5, & e_1 \cdot e_2 = \alpha_6 e_4 + \alpha_7 e_5, & e_1 \cdot e_3 = \frac{1}{2}\alpha_6 e_5, \\ e_1 \cdot e_6 = -e_4 + \alpha_8 e_5, & e_2 \cdot e_2 = \alpha_{10} e_5, & e_2 \cdot e_6 = \alpha_{11} e_5, & e_6 \cdot e_6 = \alpha_{12} e_5; \end{cases} \\ \mathbf{TP}_3(g_{(6,1)}^2) : & \begin{cases} e_1 \cdot e_1 = \alpha_3 e_4 + \alpha_4 e_5, & e_1 \cdot e_2 = \alpha_6 e_4 + \alpha_7 e_5, & e_1 \cdot e_3 = \frac{1}{2}\alpha_6 e_5, & e_1 \cdot e_6 = \alpha_8 e_5, \\ e_2 \cdot e_2 = \alpha_9 e_4 + \alpha_{10} e_5, & e_2 \cdot e_3 = \frac{1}{2}\alpha_9 e_5, & e_2 \cdot e_6 = \alpha_{11} e_5, & e_6 \cdot e_6 = \alpha_{12} e_5; \end{cases} \end{aligned}$$

where the transposed Poisson algebra has its products with respect to the bracket $[-, -]$, and remaining products are equal to zero.

Proof. Let $(g_{(6,1)}^2, \cdot, [-, -])$ be a transposed Poisson algebra structure defined on Lie algebra $g_{(6,1)}^2$. Then, for any element of $x \in g_{(6,1)}^2$, we have that operator of multiplication $\varphi_x(y) = x \cdot y$ is a $\frac{1}{2}$ -derivation. Hence, by using Theorem 11 for $1 \leq i \leq 6$, we derive the following:

$$\begin{cases} \varphi_{e_i}(e_1) = \alpha_{i,1} e_1 + \alpha_{i,2} e_2 + \alpha_{i,3} e_3 + \alpha_{i,4} e_4 + \alpha_{i,5} e_5 + \alpha_{i,6} e_6, \\ \varphi_{e_i}(e_2) = \alpha_{i,1} e_2 - 3\alpha_{i,6} e_3 + \beta_{i,4} e_4 + \beta_{i,5} e_5, \\ \varphi_{e_i}(e_3) = \alpha_{i,1} e_3 - 2\alpha_{i,6} e_4 + \frac{1}{2}\beta_{i,4} e_5, \\ \varphi_{e_i}(e_4) = \alpha_{i,1} e_4 - \frac{3}{2}\alpha_{i,6} e_5, \\ \varphi_{e_i}(e_5) = \alpha_{i,1} e_5, \\ \varphi_{e_i}(e_6) = -\alpha_{i,2} e_3 - \alpha_{i,3} e_4 + \gamma_{i,5} e_5 + \alpha_{i,1} e_6, \end{cases}$$

By considering the equality $\varphi_{e_i}(e_j) = e_i \cdot e_j = e_j \cdot e_i = \varphi_{e_j}(e_i)$ for the some elements e_i and e_j , we have the following restrictions:

$$\begin{aligned} \{e_1, e_2\}, & \Rightarrow \alpha_{2,1} = 0, \alpha_{2,2} = \alpha_{1,1}, \alpha_{2,3} = -3\alpha_{1,6}, \alpha_{2,4} = \beta_{1,4}, \alpha_{2,5} = \beta_{1,5}, \alpha_{2,6} = 0, \\ \{e_1, e_3\}, & \Rightarrow \alpha_{3,1} = \alpha_{3,2} = 0, \alpha_{3,3} = \alpha_{1,1}, \alpha_{3,4} = -2\alpha_{1,6}, \alpha_{3,5} = \frac{1}{2}\beta_{1,4}, \alpha_{3,6} = 0, \\ \{e_1, e_4\}, & \Rightarrow \alpha_{4,1} = \alpha_{4,2} = \alpha_{4,3} = 0, \alpha_{4,4} = \alpha_{1,1}, \alpha_{4,5} = -\frac{3}{2}\alpha_{1,6}, \alpha_{4,6} = 0, \\ \{e_1, e_5\}, & \Rightarrow \alpha_{5,1} = \alpha_{5,2} = \alpha_{5,3} = \alpha_{5,4} = 0, \alpha_{5,5} = \alpha_{1,1}, \alpha_{5,6} = 0, \\ \{e_1, e_6\}, & \Rightarrow \alpha_{6,1} = \alpha_{6,2} = 0, \alpha_{6,3} = -\alpha_{1,2}, \alpha_{6,4} = -\alpha_{1,3}, \alpha_{6,5} = \gamma_{1,5}, \alpha_{6,6} = \alpha_{1,1}, \\ \{e_2, e_3\}, & \Rightarrow \beta_{3,4} = 0, \beta_{3,5} = \frac{1}{2}\beta_{2,4}, \\ \{e_2, e_4\}, & \Rightarrow \beta_{4,4} = \beta_{4,5} = 0, \\ \{e_2, e_5\}, & \Rightarrow \beta_{5,4} = \beta_{5,5} = 0, \\ \{e_2, e_6\}, & \Rightarrow \alpha_{1,1} = 0, \beta_{6,4} = 3\alpha_{1,6}, \beta_{6,5} = \gamma_{2,5}, \\ \{e_3, e_6\}, & \Rightarrow \gamma_{3,5} = \frac{3}{2}\alpha_{1,6}, \\ \{e_4, e_6\}, & \Rightarrow \gamma_{4,5} = 0, \\ \{e_5, e_6\}, & \Rightarrow \gamma_{5,5} = 0. \end{aligned}$$

Thus, we have the following table of commutative multiplications of the transposed Poisson algebra structure defined on $g_{(6,1)}^2$:

$$\begin{cases} e_1 \cdot e_1 = \alpha_1 e_2 + \alpha_2 e_3 + \alpha_3 e_4 + \alpha_4 e_5 + \alpha_5 e_6, & e_1 \cdot e_2 = -3\alpha_5 e_3 + \alpha_6 e_4 + \alpha_7 e_5, \\ e_1 \cdot e_3 = -2\alpha_5 e_4 + \frac{1}{2}\alpha_6 e_5, & e_1 \cdot e_4 = -\frac{3}{2}\alpha_5 e_5, & e_1 \cdot e_6 = -\alpha_1 e_3 - \alpha_2 e_4 + \alpha_8 e_5, \\ e_2 \cdot e_2 = \alpha_9 e_4 + \alpha_{10} e_5, & e_2 \cdot e_3 = \frac{1}{2}\alpha_9 e_5, & e_2 \cdot e_6 = 3\alpha_5 e_4 + \alpha_{11} e_5, \\ e_3 \cdot e_6 = \frac{3}{2}\alpha_5 e_5, & e_6 \cdot e_6 = \alpha_1 e_4 + \alpha_{12} e_5. \end{cases}$$

By using the multiplication of Lie algebra $g_{(6,1)}^2$, we consider the general basis change:

$$e'_1 = \sum_{t=1}^6 A_t e_t, \quad e'_2 = \sum_{t=1}^6 B_t e_t, \quad e'_6 = \sum_{t=1}^6 C_t e_t.$$

Then the equality $e'_1 \cdot e'_1 = \alpha'_1 e'_2 + \alpha'_2 e'_3 + \alpha'_3 e'_4 + \alpha'_4 e'_5 + \alpha'_5 e'_6$ derives

$$\alpha'_1 = \frac{A_1^2}{B_2} \alpha_1.$$

- (1) If $\alpha_1 \neq 0$. Then by choosing $B_2 = A_1^2 \alpha_1$ we can assume $\alpha'_1 = 1$. Further, considering the associative identity for the triples $\{e_1, e_1, e_6\}$ and $\{e_1, e_1, e_2\}$, we obtain the following restrictions, respectively:

$$\alpha_5 = 0, \quad \alpha_6 = -2\alpha_{11}, \quad \alpha_9 = \alpha_{10} = 0.$$

In this case we obtain the algebra $\mathbf{TP}_1(g_{(6,1)}^2)$.

- (2) If $\alpha_1 = 0$. Using the associative identity for the triple $\{e_1, e_1, e_3\}$ we obtain $\alpha_5 = 0$. Again, using the change of basis we get the expression

$$\alpha'_2 = \frac{A_1}{B_2} \alpha_2.$$

- (a) If $\alpha_2 \neq 0$, then putting $B_2 = A_1 \alpha_2$ we get $\alpha'_2 = 1$. Furthermore, from the identity $e_1 \cdot (e_1 \cdot e_3) = (e_1 \cdot e_1) \cdot e_3$ we obtain $\alpha_9 = 0$. Hence, we have the algebra $\mathbf{TP}_2(g_{(6,1)}^2)$.

- (b) If $\alpha_2 = 0$, then we have the algebras $\mathbf{TP}_3(g_{(6,1)}^2)$.

□

Theorem 20. Let $(g_{(n,1)}^2, \cdot, [-, -])$ be a transposed Poisson algebra structure defined on Lie algebra $g_{(n,1)}^2$ and $n \geq 7$. Then, the multiplication of $(g_{(n,1)}^2, \cdot)$ has the following form:

$$\mathbf{TP}_1(g_{(n,1)}^2) : \begin{cases} e_1 \cdot e_1 = e_2 + \sum_{i=3}^{n-1} \alpha_i e_i, & e_1 \cdot e_2 = 2\alpha_1 e_{n-2} + \alpha_n e_{n-1}, & e_1 \cdot e_3 = \alpha_1 e_{n-1}, \\ e_1 \cdot e_n = -e_3 - \sum_{i=4}^{n-2} \alpha_{i-1} e_i + \alpha_{n+1} e_{n-1}, & e_2 \cdot e_n = -\alpha_1 \alpha_3 e_{n-1}, \\ e_n \cdot e_n = \sum_{i=4}^{n-2} \alpha_{i-1} e_i + \alpha_{n+2} e_{n-1}; \end{cases}$$

$$\begin{aligned} \mathbf{TP}_2(g_{(n,1)}^2) : & \left\{ \begin{array}{l} e_1 \cdot e_1 = e_3 + \sum_{i=4}^{n-1} \alpha_i e_i, \quad e_1 \cdot e_2 = \alpha_n e_{n-1}, \quad e_1 \cdot e_n = -e_4 - \sum_{i=5}^{n-2} \alpha_{i-1} e_i + \alpha_{n+1} e_{n-1}, \\ e_2 \cdot e_2 = \alpha_{n+3} e_{n-1}, \quad e_2 \cdot e_n = \alpha_{n+4} e_{n-1}, \quad e_n \cdot e_n = e_4 + \sum_{i=5}^{n-2} \alpha_{i-1} e_i + \alpha_{n+5} e_{n-1}; \end{array} \right. \\ \mathbf{TP}_3(g_{(n,1)}^2) : & \left\{ \begin{array}{l} e_1 \cdot e_1 = \sum_{i=4}^{n-1} \alpha_i e_i, \quad e_1 \cdot e_2 = 2\alpha_1 e_{n-2} + \alpha_n e_{n-1}, \quad e_1 \cdot e_3 = \alpha_1 e_{n-1}, \\ e_1 \cdot e_n = -\sum_{i=5}^{n-2} \alpha_{i-1} e_i + \alpha_{n+1} e_{n-1}, \quad e_2 \cdot e_2 = 2\alpha_{n+2} e_{n-2} + \alpha_{n+3} e_{n-1}, \\ e_2 \cdot e_3 = \alpha_{n+2} e_{n-1}, \quad e_2 \cdot e_n = \alpha_{n+4} e_{n-1}, \quad e_n \cdot e_n = \sum_{i=5}^{n-2} \alpha_{i-1} e_i + \alpha_{n+5} e_{n-1}; \end{array} \right. \end{aligned}$$

where the transposed Poisson algebra has its products with respect to the bracket $[-, -]$, and the remaining products are equal to zero.

Proof. Let $(g_{(n,1)}^2, \cdot, [-, -])$ be a transposed Poisson algebra structure defined on Lie algebra $g_{(n,1)}^2$. Then, for any element $x \in g_{(n,1)}^2$, we have operator of multiplication $\varphi_x(y) = x \cdot y$ is a $\frac{1}{2}$ -derivation. Hence, for $1 \leq i \leq n$, we derive

$$\begin{cases} \varphi_{e_i}(e_1) = \sum_{j=1}^{n-1} \alpha_{i,j} e_j, \\ \varphi_{e_i}(e_2) = \alpha_{i,1} e_2 + \beta_{i,n-2} e_{n-2} + \beta_{i,n-1} e_{n-1}, \\ \varphi_{e_i}(e_3) = \alpha_{i,1} e_3 + \frac{1}{2} \beta_{i,n-2} e_{n-1}, \\ \varphi_{e_i}(e_j) = \alpha_{i,1} e_j, \quad 4 \leq j \leq n-1, \\ \varphi_{e_i}(e_n) = -\sum_{j=3}^{n-2} \alpha_{i,j-1} e_j + \gamma_{i,n-1} e_{n-1} + \alpha_{i,1} e_n. \end{cases}$$

It is known that $\varphi_{e_i}(e_j) = e_i \cdot e_j = e_j \cdot e_i = \varphi_{e_j}(e_i)$. Then for some pair elements, we derive the following restrictions:

$$\begin{aligned} \{e_1, e_2\}, & \Rightarrow \alpha_{2,1} = 0, \quad \alpha_{2,2} = \alpha_{1,1}, \quad \alpha_{2,j} = 0, \quad 3 \leq j \leq n-3, \\ & \quad \alpha_{2,n-2} = \beta_{1,n-2}, \quad \alpha_{2,n-1} = \beta_{1,n-1}, \\ \{e_1, e_3\}, & \Rightarrow \alpha_{3,1} = \alpha_{3,2} = 0, \quad \alpha_{3,3} = \alpha_{1,1}, \quad \alpha_{3,j} = 0, \quad 4 \leq j \leq n-2, \\ & \quad \alpha_{3,n-1} = \frac{1}{2} \beta_{1,n-2}, \\ \{e_1, e_j\}, & \Rightarrow \alpha_{j,1} = \alpha_{1,1}, \quad \alpha_{j,t} = 0, \quad 1 \leq t \neq j \leq n-1, \\ & \quad 4 \leq j \leq n-1, \\ \{e_1, e_n\}, & \Rightarrow \alpha_{n,1} = \alpha_{n,2} = 0, \quad \alpha_{n,t} = -\alpha_{1,t-1}, \quad 3 \leq t \leq n-2, \\ & \quad \alpha_{n,n-1} = \gamma_{1,n-1}, \quad \alpha_{1,1} = 0, \\ \{e_2, e_3\}, & \Rightarrow \beta_{3,n-2} = 0, \quad \beta_{3,n-1} = \frac{1}{2} \beta_{2,n-2}, \\ \{e_2, e_j\}, & \Rightarrow \beta_{j,n-2} = \beta_{j,n-1} = 0, \\ & \quad 4 \leq j \leq n-1, \\ \{e_2, e_n\}, & \Rightarrow \beta_{n,n-2} = 0, \quad \beta_{n,n-1} = \gamma_{2,n-1}, \\ \{e_i, e_n\}, & \Rightarrow \gamma_{i,n-1} = 0, \quad 3 \leq i \leq n-1. \end{aligned}$$

Thus, we have the following table of commutative multiplications of the transposed Poisson algebra structure defined on $g_{(n,1)}^2$:

$$\begin{cases} e_1 \cdot e_1 = \sum_{i=2}^{n-1} \alpha_i e_i, \quad e_1 \cdot e_2 = 2\alpha_1 e_{n-2} + \alpha_n e_{n-1}, \quad e_1 \cdot e_3 = \alpha_1 e_{n-1}, \\ e_1 \cdot e_n = - \sum_{i=3}^{n-2} \alpha_{i-1} e_i + \alpha_{n+1} e_{n-1}, \quad e_2 \cdot e_2 = 2\alpha_{n+2} e_{n-2} + \alpha_{n+3} e_{n-1}, \\ e_2 \cdot e_3 = \alpha_{n+2} e_{n-1}, \quad e_2 \cdot e_n = \alpha_{n+4} e_{n-1}, \quad e_n \cdot e_n = \sum_{i=4}^{n-2} \alpha_{i-1} e_i + \alpha_{n+5} e_{n-1}. \end{cases} \quad (8)$$

Considering the associative identity $x \cdot (y \cdot z) = (x \cdot y) \cdot z$ in (8), we obtain the following restrictions on structure constants:

$$\begin{aligned} \{e_1, e_1, e_2\}, \quad & \Rightarrow \quad \alpha_2 \alpha_{n+2} = 0, \quad \alpha_2 \alpha_{n+3} + \alpha_3 \alpha_{n+2} = 0, \\ \{e_1, e_1, e_n\}, \quad & \Rightarrow \quad \alpha_1 \alpha_3 + \alpha_2 \alpha_{n+4} = 0. \end{aligned}$$

Similarly, by using the multiplication of Lie algebra $g_{(n,1)}^2$ we consider the general basis change:

$$e'_1 = \sum_{t=1}^n A_t e_t, \quad e'_2 = \sum_{t=1}^n B_t e_t, \quad e'_n = \sum_{t=1}^n C_t e_t.$$

Then the product $e'_1 \cdot e'_1 = \sum_{i=2}^{n-1} \alpha'_i e'_i$ gives

$$\alpha'_2 = \frac{A_1^2}{B_2} \alpha_2.$$

We have the following cases.

- (1) Let $\alpha_2 \neq 0$. Then by choosing $B_2 = A_1^2 \alpha_2$, we put $\alpha'_2 = 1$. In this case we obtain the algebra $\mathbf{TP}_1(g_{(n,1)}^2)$.
- (2) Let $\alpha_2 = 0$. Then from above restrictions we have $\alpha_3 \alpha_{n+2} = \alpha_1 \alpha_3 = 0$. If we apply a general change of basis, then we have

$$\alpha'_3 = \frac{A_1}{B_2} \alpha_3.$$

(a) If $\alpha_3 \neq 0$, then $\alpha'_3 = 1$ and we have the algebra $\mathbf{TP}_2(g_{(n,1)}^2)$.

(b) If $\alpha_3 = 0$, then we have the algebra $\mathbf{TP}_3(g_{(n,1)}^2)$.

□

Theorem 21. Let $(g_{(7,1)}^3, \cdot, [-, -])$ be a transposed Poisson algebra structure defined on Lie algebra $g_{(7,1)}^3$. Then, the multiplication of $(g_{(7,1)}^3, \cdot)$ has the following form:

$$\mathbf{TP}_1(g_{(7,1)}^3) : \begin{cases} e_1 \cdot e_1 = \alpha_4 e_4 + \alpha_5 e_5, \quad e_1 \cdot e_2 = 2\alpha_6 e_5 + \alpha_7 e_6, \quad e_1 \cdot e_3 = \alpha_6 e_6, \\ e_1 \cdot e_7 = -\alpha_4 e_5 + \alpha_8 e_6, \quad e_2 \cdot e_2 = 2\alpha_1 e_5 + \alpha_9 e_6, \quad e_2 \cdot e_3 = \alpha_1 e_6, \\ e_2 \cdot e_7 = \alpha_{10} e_6, \quad e_7 \cdot e_7 = \alpha_{11} e_6; \end{cases}$$

$$\mathbf{TP}_2(g_{(7,1)}^3) : \begin{cases} e_1 \cdot e_1 = 2e_3 + \alpha_4 e_4 + \alpha_5 e_5, e_1 \cdot e_2 = 2\alpha_6 e_5 + \alpha_7 e_6, e_1 \cdot e_3 = (\alpha_6 - 1)e_6, \\ e_1 \cdot e_7 = -2e_4 - \alpha_4 e_5 + \alpha_8 e_6, e_2 \cdot e_2 = \alpha_9 e_6, e_2 \cdot e_7 = \alpha_{10} e_6, \\ e_7 \cdot e_7 = 2e_5 + \alpha_{11} e_6; \end{cases}$$

where the transposed Poisson algebra has its products with respect to the bracket $[-, -]$, and remaining products are equal to zero.

Proof. Let $(g_{(7,1)}^3, \cdot, [-, -])$ be a transposed Poisson algebra structure defined on Lie algebra $g_{(7,1)}^3$. Then for any element $x \in g_{(7,1)}^3$, we have that operator of multiplication $\varphi_x(y) = x \cdot y$ is a $\frac{1}{2}$ -derivation. Hence, for $1 \leq i \leq 7$ by Theorem 12 we derive the following:

$$\begin{cases} \varphi_{e_i}(e_1) = \alpha_{i,1}e_1 + \alpha_{i,2}e_2 + \alpha_{i,3}e_3 + \alpha_{i,4}e_4 + \alpha_{i,5}e_5 + \alpha_{i,6}e_6, \\ \varphi_{e_i}(e_2) = \alpha_{i,1}e_2 + 4\alpha_{i,2}e_4 + \beta_{i,5}e_5 + \beta_{i,6}e_6, \\ \varphi_{e_i}(e_3) = \alpha_{i,1}e_3 + 2\alpha_{i,2}e_5 + \frac{1}{2}(\beta_{i,5} - \alpha_{i,3})e_6, \\ \varphi_{e_i}(e_4) = \alpha_{i,1}e_4 + \frac{3}{2}\alpha_{i,2}e_6, \\ \varphi_{e_i}(e_5) = \alpha_{i,1}e_5, \\ \varphi_{e_i}(e_6) = \alpha_{i,1}e_6, \\ \varphi_{e_i}(e_7) = -\alpha_{i,2}e_3 - \alpha_{i,3}e_4 - \alpha_{i,4}e_5 + \gamma_{i,6}e_6 + \alpha_{i,1}e_7, \end{cases}$$

Considering the property $\varphi_{e_i}(e_j) = e_i \cdot e_j = e_j \cdot e_i = \varphi_{e_j}(e_i)$, we obtain the following restrictions:

$$\begin{aligned} \{e_1, e_2\}, &\Rightarrow \alpha_{2,1} = 0, \alpha_{2,2} = \alpha_{1,1}, \alpha_{2,3} = 0, \alpha_{2,4} = 4\alpha_{1,2}, \alpha_{2,5} = \beta_{1,5}, \alpha_{2,6} = \beta_{1,6}, \\ \{e_1, e_3\}, &\Rightarrow \alpha_{3,1} = \alpha_{3,2} = 0, \alpha_{3,3} = \alpha_{1,1}, \alpha_{3,4} = -2\alpha_{1,2}, \alpha_{3,5} = 2\alpha_{1,2}, \alpha_{3,6} = \frac{1}{2}(\beta_{1,5} - \alpha_{1,3}), \\ \{e_1, e_4\}, &\Rightarrow \alpha_{4,1} = \alpha_{4,2} = \alpha_{4,3} = 0, \alpha_{4,4} = \alpha_{1,1}, \alpha_{4,5} = 0, \alpha_{4,6} = \frac{3}{2}\alpha_{1,2}, \\ \{e_1, e_5\}, &\Rightarrow \alpha_{5,1} = \alpha_{5,2} = \alpha_{5,3} = \alpha_{5,4} = 0, \alpha_{5,5} = \alpha_{1,1}, \alpha_{5,6} = 0, \\ \{e_1, e_6\}, &\Rightarrow \alpha_{6,1} = \alpha_{6,2} = \alpha_{6,3} = \alpha_{6,4} = \alpha_{6,5} = 0, \alpha_{6,6} = \alpha_{1,1}, \\ \{e_1, e_7\}, &\Rightarrow \alpha_{1,1} = 0, \alpha_{7,1} = \alpha_{7,2} = 0, \alpha_{7,3} = -\alpha_{1,2}, \alpha_{7,4} = -\alpha_{1,3}, \alpha_{7,5} = -\alpha_{1,4}, \alpha_{7,6} = \gamma_{1,6}, \\ \{e_2, e_3\}, &\Rightarrow \beta_{3,5} = 0, \beta_{3,6} = \frac{1}{2}\beta_{2,5}, \\ \{e_2, e_4\}, &\Rightarrow \beta_{4,5} = \beta_{4,6} = 0, \\ \{e_2, e_5\}, &\Rightarrow \beta_{5,5} = \beta_{5,6} = 0, \\ \{e_2, e_6\}, &\Rightarrow \beta_{6,5} = \beta_{6,6} = 0, \\ \{e_2, e_7\}, &\Rightarrow \beta_{7,5} = -4\alpha_{1,2}, \beta_{7,6} = \gamma_{2,6}, \\ \{e_3, e_7\}, &\Rightarrow \alpha_{1,6} = 0, \gamma_{3,6} = -\frac{3}{2}\alpha_{1,2}, \\ \{e_4, e_7\}, &\Rightarrow \gamma_{4,6} = 0, \\ \{e_5, e_7\}, &\Rightarrow \gamma_{5,6} = 0, \\ \{e_6, e_7\}, &\Rightarrow \gamma_{6,6} = 0. \end{aligned}$$

Thus, we have the following table of commutative multiplications of the transposed Poisson algebra structure $(g_{(7,1)}^3, \cdot, \cdot, [-, -])$:

$$\begin{cases} e_1 \cdot e_1 = 2\alpha_2 e_2 + 2\alpha_3 e_3 + \alpha_4 e_4 + \alpha_5 e_5, & e_1 \cdot e_2 = 8\alpha_2 e_4 + 2\alpha_6 e_5 + \alpha_7 e_6, \\ e_1 \cdot e_3 = 4\alpha_2 e_5 + (\alpha_6 - \alpha_3) e_6, & e_1 \cdot e_4 = 3\alpha_2 e_6, \\ e_2 \cdot e_2 = 2\alpha_1 e_5 + \alpha_9 e_6, & e_2 \cdot e_3 = \alpha_1 e_6, \\ e_3 \cdot e_7 = -3\alpha_2 e_6, & e_2 \cdot e_7 = -8\alpha_2 e_5 + \alpha_{10} e_6, \\ e_7 \cdot e_7 = 2\alpha_2 e_4 + 2\alpha_3 e_5 + \alpha_{11} e_6. \end{cases}$$

Considering the associative identity for the triples $\{e_1, e_1, e_7\}$ and $\{e_1, e_1, e_2\}$, we obtain the following restrictions, respectively:

$$\alpha_2 = 0, \alpha_1 \alpha_3 = 0.$$

Using multiplication of Lie algebra $g_{(7,1)}^3$ and multiplication $e_1 \cdot e_1 = 2\alpha_3 e_3 + \alpha_4 e_4 + \alpha_5 e_5$, and also consider a general change of basis, we can show that α_3 is an invariant.

If $\alpha_3 = 0$. Then, we have the algebra $\mathbf{TP}_1(g_{(7,1)}^3)$.

If $\alpha_3 \neq 0$. Then, we obtain $\alpha_1 = 0$. By choosing change of basis, we can assume that $\alpha_3 = 1$ and obtain the algebra $\mathbf{TP}_2(g_{(7,1)}^3)$. \square

Theorem 22. Let $(g_{(n,1)}^3, \cdot, [-, -])$ be a transposed Poisson algebra structure defined on Lie algebra $g_{(n,1)}^3$ and $n \geq 8$. Then, the multiplication of $(g_{(n,1)}^3, \cdot)$ has the following form:

$$\begin{aligned} \mathbf{TP}(g_{(8,1)}^3) : & \begin{cases} e_1 \cdot e_1 = \sum_{t=4}^7 \alpha_t e_t, & e_1 \cdot e_2 = \beta_1 e_6 + \beta_2 e_7, & e_1 \cdot e_3 = \frac{1}{2}(\beta_1 - \alpha_4) e_7, \\ e_1 \cdot e_8 = -\alpha_4 e_5 - \alpha_5 e_6 + \gamma_1 e_7, & e_2 \cdot e_2 = \beta_3 e_6 + \beta_4 e_7, \\ e_2 \cdot e_3 = \frac{1}{2}\beta_3 e_7, & e_2 \cdot e_8 = \gamma_2 e_7, & e_8 \cdot e_8 = \alpha_4 e_6 + \gamma_3 e_7; \end{cases} \\ \mathbf{TP}_1(g_{(9,1)}^3) : & \begin{cases} e_1 \cdot e_1 = \sum_{t=4}^8 \alpha_t e_t, & e_1 \cdot e_2 = \beta_1 e_7 + \beta_2 e_8, & e_1 \cdot e_3 = \frac{1}{2}(\beta_1 - \alpha_5) e_8, \\ e_1 \cdot e_9 = -\sum_{t=5}^7 \alpha_{t-1} e_t + \gamma_1 e_8, & e_2 \cdot e_2 = \beta_3 e_7 + \beta_4 e_8, & e_2 \cdot e_3 = \frac{1}{2}\beta_3 e_8, \\ e_2 \cdot e_9 = -\alpha_4 e_7 + \gamma_2 e_8, & e_3 \cdot e_9 = \frac{1}{2}\alpha_4 e_8, & e_9 \cdot e_9 = \sum_{t=6}^7 \alpha_{t-2} e_t + \gamma_3 e_8; \end{cases} \\ \mathbf{TP}_2(g_{(9,1)}^3) : & \begin{cases} e_1 \cdot e_1 = e_3 + \sum_{t=5}^8 \alpha_t e_t, & e_1 \cdot e_2 = e_5 + \beta_1 e_7 + \beta_2 e_8, \\ e_1 \cdot e_3 = \frac{1}{2}(\beta_1 - \alpha_5) e_8, & e_1 \cdot e_9 = -e_4 - \sum_{t=6}^7 \alpha_{t-1} e_t + \gamma_1 e_8, \\ e_2 \cdot e_2 = e_7 + \beta_4 e_8, & e_2 \cdot e_9 = -e_6 + \gamma_2 e_8, & e_9 \cdot e_9 = e_5 + \alpha_5 e_7 + \gamma_3 e_8; \end{cases} \end{aligned}$$

$$\begin{aligned} \mathbf{TP}_1(g_{(n,1)}^3) : & \left\{ \begin{array}{l} e_1 \cdot e_1 = \sum_{t=4}^{n-1} \alpha_t e_t, \quad e_1 \cdot e_2 = \sum_{t=6}^{n-3} \alpha_{t-2} e_t + \beta_1 e_{n-2} + \beta_2 e_{n-1}, \\ e_1 \cdot e_3 = \frac{1}{2}(\beta_1 - \alpha_{n-4}) e_{n-1}, \quad e_1 \cdot e_n = - \sum_{t=5}^{n-2} \alpha_{t-1} e_t + \gamma_1 e_{n-1}, \\ e_2 \cdot e_2 = \sum_{t=8}^{n-3} \alpha_{t-4} e_t + \beta_3 e_{n-2} + \beta_4 e_{n-1}, \quad e_2 \cdot e_3 = \frac{1}{2}(\beta_3 - \alpha_{n-6}) e_{n-1}, \\ e_2 \cdot e_n = - \sum_{t=7}^{n-2} \alpha_{t-3} e_t + \gamma_2 e_{n-1}, \quad e_3 \cdot e_n = \frac{1}{2} \alpha_{n-5} e_{n-1}, \quad e_n \cdot e_n = \sum_{t=6}^{n-2} \alpha_{t-2} e_t + \gamma_3 e_{n-1}; \end{array} \right. \\ \mathbf{TP}_2(g_{(n,1)}^3) : & \left\{ \begin{array}{l} e_1 \cdot e_1 = e_3 + \sum_{t=4}^{n-6} \alpha_t e_t + \sum_{t=n-4}^{n-1} \alpha_t e_t, \quad e_1 \cdot e_2 = e_5 + \sum_{t=5}^{n-4} \alpha_{t-2} e_t + \beta_1 e_{n-2} + \beta_2 e_{n-1}, \\ e_1 \cdot e_3 = \frac{1}{2}(\beta_1 - \alpha_{n-4}) e_{n-1}, \quad e_1 \cdot e_n = -e_4 - \sum_{t=5}^{n-5} \alpha_{t-1} e_t - \sum_{t=n-3}^{n-2} \alpha_{t-1} e_t + \gamma_1 e_{n-1}, \\ e_2 \cdot e_2 = e_7 + \sum_{t=8}^{n-2} \alpha_{t-4} e_t + \beta_4 e_{n-1}, \quad e_2 \cdot e_n = -e_6 - \sum_{t=7}^{n-3} \alpha_{t-3} e_t + \gamma_2 e_{n-1}, \\ e_n \cdot e_n = e_5 + \sum_{t=6}^{n-4} \alpha_{t-2} e_t + \alpha_{n-4} e_{n-2} + \gamma_3 e_{n-1}; \end{array} \right. \end{aligned}$$

where the transposed Poisson algebra has its products with respect to the bracket $[-, -]$, and remaining products are equal to zero.

Proof. Let $(g_{(n,1)}^3, \cdot, [-, -])$ be a transposed Poisson algebra structure defined on Lie algebra $g_{(n,1)}^3$. Then for any element $x \in g_{(n,1)}^3$ we have that, operator of multiplication $\varphi_x(y) = x \cdot y$ is a $\frac{1}{2}$ -derivation. Hence, for $1 \leq i \leq n$ we derive

$$\begin{cases} \varphi_{e_i}(e_1) = \alpha_{i,1} e_1 + \sum_{t=3}^{n-1} \alpha_{i,t} e_t, \\ \varphi_{e_i}(e_2) = \alpha_{i,1} e_2 + \sum_{t=5}^{n-3} \alpha_{i,t-2} e_t + \beta_{i,n-2} e_{n-2} + \beta_{i,n-1} e_{n-1}, \\ \varphi_{e_i}(e_3) = \alpha_{i,1} e_3 + \frac{1}{2}(\beta_{i,n-2} - \alpha_{i,n-4}) e_{n-1}, \\ \varphi_{e_i}(e_j) = \alpha_{i,1} e_j, \quad 4 \leq j \leq n-1, \\ \varphi_{e_i}(e_n) = - \sum_{t=4}^{n-2} \alpha_{i,t-1} e_t + \gamma_{i,n-1} e_{n-1} + \alpha_{i,1} e_n. \end{cases}$$

It is known that $\varphi_{e_i}(e_j) = e_i \cdot e_j = e_j \cdot e_i = \varphi_{e_j}(e_i)$. Then for some pair elements, we derive the following restrictions:

$$\begin{aligned} \{e_1, e_2\}, & \Rightarrow \alpha_{1,1} = \alpha_{2,1} = \alpha_{2,3} = \alpha_{2,4} = 0, \quad \alpha_{2,t} = \alpha_{1,t-2}, \quad 5 \leq t \leq n-3, \\ & \quad \alpha_{2,n-2} = \beta_{1,n-2}, \quad \alpha_{2,n-1} = \beta_{1,n-1}, \\ \{e_1, e_3\}, & \Rightarrow \alpha_{3,1} = 0, \quad \alpha_{3,t} = 0, \quad 3 \leq t \leq n-2, \quad \alpha_{3,n-1} = \frac{1}{2}(\beta_{1,n-2} - \alpha_{1,n-4}), \\ \{e_1, e_j\}, & \Rightarrow \alpha_{j,1} = 0, \quad \alpha_{j,t} = 0, \quad 3 \leq t \leq n-1, \\ 4 \leq j \leq n-1, & \\ \{e_1, e_n\}, & \Rightarrow \alpha_{n,1} = \alpha_{n,3} = 0, \quad \alpha_{n,t} = -\alpha_{1,t-1}, \quad 4 \leq t \leq n-2, \quad \alpha_{n,n-1} = \gamma_{1,n-1}, \end{aligned}$$

$$\begin{aligned}
\{e_2, e_3\}, \quad & \Rightarrow \beta_{3,n-2} = 0, \beta_{3,n-1} = \frac{1}{2}(\beta_{2,n-2} - \alpha_{1,n-6}), \\
\{e_2, e_j\}, \quad & \Rightarrow \beta_{j,n-2} = \beta_{j,n-1} = 0, \\
4 \leq j \leq n-1, \quad & \\
\{e_2, e_n\}, \quad & \Rightarrow \beta_{n,n-2} = -\alpha_{1,n-5}, \beta_{n,n-1} = \gamma_{2,n-1}, \\
\{e_3, e_n\}, \quad & \Rightarrow \gamma_{3,n-1} = \frac{1}{2}\alpha_{1,n-5}, \\
\{e_i, e_n\}, \quad & \Rightarrow \gamma_{i,n-1} = 0, \\
4 \leq i \leq n-1. \quad &
\end{aligned}$$

So, we have the following table of commutative multiplications of the transposed Poisson algebra structure $(g_{(n,1)}^3, \cdot, \cdot, [-, -])$

For $n = 8$:

$$\left\{
\begin{aligned}
e_1 \cdot e_1 &= \sum_{t=3}^7 \alpha_t e_t, \quad e_1 \cdot e_2 = \alpha_3 e_5 + \beta_1 e_6 + \beta_2 e_7, \quad e_1 \cdot e_3 = \frac{1}{2}(\beta_1 - \alpha_4) e_7, \\
e_1 \cdot e_8 &= -\sum_{t=4}^6 \alpha_{t-1} e_t + \gamma_1 e_7, \quad e_2 \cdot e_2 = \beta_3 e_6 + \beta_4 e_7, \quad e_2 \cdot e_3 = \frac{1}{2}\beta_3 e_7, \\
e_2 \cdot e_8 &= -\alpha_3 e_6 + \gamma_2 e_7, \quad e_3 \cdot e_8 = \frac{1}{2}\alpha_3 e_7, \quad e_8 \cdot e_8 = \sum_{t=5}^6 \alpha_{t-2} e_t + \gamma_3 e_7;
\end{aligned}
\right.$$

For $n \geq 9$:

$$\left\{
\begin{aligned}
e_1 \cdot e_1 &= \sum_{t=3}^{n-1} \alpha_t e_t, \quad e_1 \cdot e_2 = \sum_{t=5}^{n-3} \alpha_{t-2} e_t + \beta_1 e_{n-2} + \beta_2 e_{n-1}, \quad e_1 \cdot e_3 = \frac{1}{2}(\beta_1 - \alpha_{n-4}) e_{n-1}, \\
e_1 \cdot e_n &= -\sum_{t=4}^{n-2} \alpha_{t-1} e_t + \gamma_1 e_{n-1}, \quad e_2 \cdot e_2 = \sum_{t=7}^{n-3} \alpha_{t-4} e_t + \beta_3 e_{n-2} + \beta_4 e_{n-1}, \\
e_2 \cdot e_3 &= \frac{1}{2}(\beta_3 - \alpha_{n-6}) e_{n-1}, \quad e_2 \cdot e_n = -\sum_{t=6}^{n-2} \alpha_{t-3} e_t + \gamma_2 e_{n-1}, \\
e_3 \cdot e_n &= \frac{1}{2}\alpha_{n-5} e_{n-1}, \quad e_n \cdot e_n = \sum_{t=5}^{n-2} \alpha_{t-2} e_t + \gamma_3 e_{n-1}.
\end{aligned}
\right.$$

Considering the associative identity for the triples $\{e_1, e_1, e_2\}$ and $\{e_1, e_1, e_n\}$, we obtain the following restrictions, respectively:

$$\alpha_3(\beta_3 - \alpha_{n-6}) = 0, \quad \alpha_3\alpha_{n-5} = 0. \quad (9)$$

We have the following cases.

(1) Let $n = 8$. Then we get $\alpha_3 = 0$ and we have the algebra $\mathbf{TP}(g_{(8,1)}^3)$.

(2) Let $n \geq 9$. Then by using the multiplication of the Lie algebra $g_{(n,1)}^3$ we consider the general basis change:

$$e'_1 = \sum_{t=1}^n A_t e_t, \quad e'_2 = \sum_{t=1}^n B_t e_t, \quad e'_n = \sum_{t=1}^n C_t e_t.$$

The product $e'_1 \cdot e'_1 = \sum_{i=3}^{n-1} \alpha'_i e'_i$ gives

$$\alpha'_3 = \frac{A_1}{B_2} \alpha_3.$$

- (a) If $\alpha_3 = 0$. Then for $n = 9$ we derive the algebra $\mathbf{TP}_1(g_{(9,1)}^3)$ and for the case when $n \geq 10$ we obtain $\mathbf{TP}_1(g_{(n,1)}^3)$.
- (b) If $\alpha_3 \neq 0$. Then by choosing $B_2 = A_1 \alpha_3$ we can assume $\alpha'_3 = 1$ and from the restrictions (9) we get $\alpha_{n-5} = 0, \beta'_3 = \alpha_{n-6}$. In this case for the $n = 9$ we obtain $\mathbf{TP}_2(g_{(9,1)}^3)$ and for $n \geq 10$ we have the algebra $\mathbf{TP}_2(g_{(n,1)}^3)$.

□

Theorem 23. Let $(g_7^1, \cdot, [-, -])$ be a transposed Poisson algebra structure defined on Lie algebra g_7^1 . Then, the multiplication of (g_7^1, \cdot) has the following form:

$$\begin{aligned} \mathbf{TP}_1(g_7^1) : & \begin{cases} e_1 \cdot e_1 = \alpha_4 e_4 + \alpha_5 e_5 + \alpha_6 e_6 + \alpha_7 e_7, & e_1 \cdot e_2 = \alpha_1 e_5 + \alpha_2 e_6 + \alpha_8 e_7, \\ e_1 \cdot e_3 = \frac{1}{2}(\alpha_1 - \alpha_4)e_6 - \frac{1}{2}\alpha_5 e_7, & e_1 \cdot e_4 = \frac{1}{2}\alpha_4 e_7, \\ e_2 \cdot e_2 = \alpha_9 e_5 + \alpha_{10} e_6 + \alpha_{11} e_7, & e_2 \cdot e_3 = \frac{1}{2}\alpha_9 e_6 - \frac{1}{2}\alpha_1 e_7; \\ e_1 \cdot e_1 = 6e_3 + \alpha_4 e_4 + \alpha_5 e_5 + \alpha_6 e_6 + \alpha_7 e_7, & e_1 \cdot e_2 = -2e_4 + \alpha_1 e_5 + \alpha_2 e_6 + \alpha_8 e_7, \\ e_1 \cdot e_3 = -4e_5 + \frac{1}{2}(\alpha_1 - \alpha_4)e_6 - \frac{1}{2}\alpha_5 e_7, & e_1 \cdot e_4 = -2e_6 + \frac{1}{2}\alpha_4 e_7, e_1 \cdot e_5 = -3e_7, \\ e_2 \cdot e_2 = -\frac{2}{3}e_5 + \alpha_{10} e_6 + \alpha_{11} e_7, & e_2 \cdot e_3 = \frac{2}{3}e_6 - \frac{1}{2}\alpha_1 e_7, e_2 \cdot e_4 = -e_7, e_3 \cdot e_3 = 2e_7; \end{cases} \\ \mathbf{TP}_2(g_7^1) : & \begin{cases} e_1 \cdot e_1 = \alpha_4 e_4 + \alpha_5 e_5 + \alpha_6 e_6 + \alpha_7 e_7, & e_1 \cdot e_2 = \alpha_1 e_5 + \alpha_2 e_6 + \alpha_8 e_7, \\ e_1 \cdot e_3 = \alpha_9 e_5 + \alpha_{10} e_6 + \alpha_{11} e_7, & e_1 \cdot e_4 = \alpha_4 e_7, e_1 \cdot e_5 = -3e_7, \\ e_2 \cdot e_2 = -\frac{2}{3}e_5 + \alpha_{10} e_6 + \alpha_{11} e_7, & e_2 \cdot e_3 = \frac{2}{3}e_6 - \frac{1}{2}\alpha_1 e_7, e_2 \cdot e_4 = -e_7, e_3 \cdot e_3 = 2e_7; \end{cases} \end{aligned}$$

where the transposed Poisson algebra has its products with respect to the bracket $[-, -]$, and remaining products are equal to zero.

Proof. Let $(g_7^1, \cdot, [-, -])$ be a transposed Poisson algebra structure defined on the Lie algebra $g_{(n,1)}^3$. Then for any element of $x \in g_7^1$, we have that operator of multiplication $\varphi_x(y) = x \cdot y$ is a $\frac{1}{2}$ -derivation. Hence, for $1 \leq i \leq 7$ by using Theorem 13 we derive

$$\begin{cases} \varphi_{e_i}(e_1) = \alpha_{i,1} e_1 + \alpha_{i,3} e_3 + \alpha_{i,4} e_4 + \alpha_{i,5} e_5 + \alpha_{i,6} e_6 + \alpha_{i,7} e_7, \\ \varphi_{e_i}(e_2) = \alpha_{i,1} e_2 - \frac{1}{3}\alpha_{i,3} e_4 + \beta_{i,5} e_5 + \beta_{i,6} e_6 + \beta_{i,7} e_7, \\ \varphi_{e_i}(e_3) = \alpha_{i,1} e_3 - \frac{2}{3}\alpha_{i,3} e_5 + \frac{1}{2}(\beta_{i,5} - \alpha_{i,4}) e_6 - \frac{1}{2}\alpha_{i,5} e_7, \\ \varphi_{e_i}(e_4) = \alpha_{i,1} e_4 - \frac{1}{3}\alpha_{i,3} e_6 + \frac{1}{2}\alpha_{i,4} e_7, \varphi_{e_i}(e_5) = \alpha_{i,1} e_5 - \frac{1}{2}\alpha_{i,3} e_7, \\ \varphi_{e_i}(e_6) = \alpha_{i,1} e_6, \varphi_{e_i}(e_7) = \alpha_{i,1} e_7. \end{cases}$$

By considering equality $\varphi_{e_i}(e_j) = e_i \cdot e_j = e_j \cdot e_i = \varphi_{e_j}(e_i)$ for the some elements e_i and e_j , we have the following restrictions:

$$\begin{aligned}
\{e_1, e_2\}, \Rightarrow \alpha_{1,1} = \alpha_{2,1} = \alpha_{2,3} = 0, \alpha_{2,4} = -\frac{1}{3}\alpha_{1,3}, \alpha_{2,5} = \beta_{1,5}, \alpha_{2,6} = \beta_{1,6}, \alpha_{2,7} = \beta_{1,7}, \\
\{e_1, e_3\}, \Rightarrow \alpha_{3,1} = \alpha_{3,3} = \alpha_{3,4} = 0, \alpha_{3,5} = -\frac{2}{3}\alpha_{1,3}, \alpha_{3,6} = \frac{1}{2}(\beta_{1,5} - \alpha_{1,4}), \alpha_{3,7} = -\frac{1}{2}\alpha_{1,5}, \\
\{e_1, e_4\}, \Rightarrow \alpha_{4,1} = \alpha_{4,3} = \alpha_{4,4} = \alpha_{4,5} = 0, \alpha_{4,6} = -\frac{1}{3}\alpha_{1,3}, \alpha_{4,7} = \frac{1}{2}\alpha_{1,4}, \\
\{e_1, e_5\}, \Rightarrow \alpha_{5,1} = \alpha_{5,3} = \alpha_{5,4} = \alpha_{5,5} = \alpha_{5,6} = 0, \alpha_{5,7} = -\frac{1}{2}\alpha_{1,3}, \\
\{e_1, e_6\}, \Rightarrow \alpha_{6,1} = \alpha_{6,3} = \alpha_{6,4} = \alpha_{6,5} = \alpha_{6,6} = \alpha_{6,7} = 0, \\
\{e_1, e_7\}, \Rightarrow \alpha_{7,1} = \alpha_{7,3} = \alpha_{7,4} = \alpha_{7,5} = \alpha_{7,6} = \alpha_{7,7} = 0, \\
\{e_2, e_3\}, \Rightarrow \beta_{3,5} = 0, \beta_{3,6} = \frac{1}{6}(3\beta_{2,5} + \alpha_{1,3}), \beta_{3,7} = -\frac{1}{2}\beta_{1,5}, \\
\{e_2, e_4\}, \Rightarrow \beta_{4,5} = \beta_{4,6} = 0, \beta_{4,7} = -\frac{1}{6}\alpha_{1,3}, \\
\{e_2, e_5\}, \Rightarrow \beta_{5,5} = \beta_{5,6} = \beta_{5,7} = 0, \\
\{e_2, e_6\}, \Rightarrow \beta_{6,5} = \beta_{6,6} = \beta_{6,7} = 0, \\
\{e_2, e_7\}, \Rightarrow \beta_{7,5} = \beta_{7,6} = \beta_{7,7} = 0.
\end{aligned}$$

So, we have the following table of commutative multiplications of the transposed Poisson algebra structure $(g_{(7)}^1, \cdot, \cdot, [-, -])$

$$\left\{
\begin{aligned}
e_1 \cdot e_1 &= \alpha_3 e_3 + \alpha_4 e_4 + \alpha_5 e_5 + \alpha_6 e_6 + \alpha_7 e_7, \quad e_1 \cdot e_2 = -\frac{1}{3}\alpha_3 e_4 + \alpha_1 e_5 + \alpha_2 e_6 + \alpha_8 e_7, \\
e_1 \cdot e_3 &= -\frac{2}{3}\alpha_3 e_5 + \frac{1}{2}(\alpha_1 - \alpha_4) e_6 - \frac{1}{2}\alpha_5 e_7, \quad e_1 \cdot e_4 = -\frac{1}{3}\alpha_3 e_6 + \frac{1}{2}\alpha_4 e_7, \quad e_1 \cdot e_5 = -\frac{1}{2}\alpha_3 e_7, \\
e_2 \cdot e_2 &= \alpha_9 e_5 + \alpha_{10} e_6 + \alpha_{11} e_7, \quad e_2 \cdot e_3 = \frac{1}{6}(3\alpha_9 + \alpha_3) e_6 - \frac{1}{2}\alpha_1 e_7, \\
e_2 \cdot e_4 &= -\frac{1}{6}\alpha_3 e_7, \quad e_3 \cdot e_3 = \frac{1}{3}\alpha_3 e_7,
\end{aligned}
\right.$$

The equality $e_1 \cdot (e_1 \cdot e_2) = (e_1 \cdot e_1) \cdot e_2$ gives $\alpha_3 (\alpha_3 + 9\alpha_9) = 0$. Similarly above by using a general change of basis, we can show that α_3 is an invariant.

- (1) If $\alpha_3 = 0$, then we have the algebra $\text{TP}_1(g_7^1)$;
- (2) If $\alpha_3 \neq 0$, then we get the algebra $\text{TP}_2(g_7^1)$.

□

Theorem 24. Let $(g_9^2, \cdot, [-, -])$ be a transposed Poisson algebra structure defined on the Lie algebra g_9^2 . Then the multiplication of (g_9^2, \cdot) has the following form:

$$\text{TP}(g_9^2) : \left\{
\begin{aligned}
e_1 \cdot e_1 &= \alpha_1 e_5 + \alpha_2 e_6 + \alpha_3 e_7 + \alpha_4 e_8 + \alpha_5 e_9, \quad e_1 \cdot e_2 = \frac{1}{3}\alpha_1 e_6 + \alpha_6 e_7 + \alpha_7 e_8 + \alpha_8 e_9, \\
e_1 \cdot e_3 &= -\frac{4}{3}\alpha_1 e_7 + \frac{1}{2}(\alpha_6 - 5\alpha_2) e_8 - \frac{1}{2}\alpha_3 e_9, \quad e_1 \cdot e_4 = \frac{1}{3}\alpha_1 e_8 + \frac{1}{2}\alpha_2 e_9, \\
e_1 \cdot e_5 &= -\frac{1}{2}\alpha_1 e_9, \quad e_2 \cdot e_2 = \alpha_9 e_7 + \alpha_{10} e_8 + \alpha_{11} e_9, \\
e_2 \cdot e_3 &= \frac{1}{6}(3\alpha_9 - 5\alpha_1) e_8 - \frac{1}{2}\alpha_6 e_9, \quad e_2 \cdot e_4 = \frac{1}{6}\alpha_1 e_9, \quad e_3 \cdot e_3 = \frac{2}{3}\alpha_1 e_9,
\end{aligned}
\right.$$

where the transposed Poisson algebra has its products with respect to the bracket $[-, -]$, and remaining products are equal to zero.

Proof. Let $(g_9^2, \cdot, [-, -])$ be a transposed Poisson algebra structure defined on Lie algebra g_9^2 . Then for any element $x \in g_9^2$, we have that operator of multiplication $\varphi_x(y) = x \cdot y$ is a $\frac{1}{2}$ -derivation. Hence, for $1 \leq i \leq 9$ we derive

$$\begin{cases} \varphi_{e_i}(e_1) = \alpha_{i,1}e_1 + \alpha_{i,5}e_5 + \alpha_{i,6}e_6 + \alpha_{i,7}e_7 + \alpha_{i,8}e_8 + \alpha_{i,9}e_9, \\ \varphi_{e_i}(e_2) = \alpha_{i,1}e_2 + \frac{1}{3}\alpha_{i,5}e_6 + \beta_{i,7}e_7 + \beta_{i,8}e_8 + \beta_{i,9}e_9, \\ \varphi_{e_i}(e_3) = \alpha_{i,1}e_3 - \frac{4}{3}\alpha_{i,5}e_7 + \frac{1}{2}(\beta_{i,7} - 5\alpha_{i,6})e_8 - \frac{1}{2}\alpha_{i,7}e_9, \\ \varphi_{e_i}(e_4) = \alpha_{i,1}e_4 + \frac{1}{3}\alpha_{i,5}e_8 + \frac{1}{2}\alpha_{i,6}e_9, \quad \varphi_{e_i}(e_5) = \alpha_{i,1}e_5 - \frac{1}{2}\alpha_{i,5}e_9, \quad \varphi_{e_i}(e_j) = \alpha_{i,1}e_j, \quad 6 \leq j \leq 9. \end{cases}$$

It is known that $\varphi_{e_i}(e_j) = e_i \cdot e_j = e_j \cdot e_i = \varphi_{e_j}(e_i)$. Then for some pair elements, we derive the following restrictions:

$$\begin{aligned} \{e_1, e_2\}, \quad &\Rightarrow \alpha_{1,1} = \alpha_{2,1} = \alpha_{2,5} = 0, \quad \alpha_{2,6} = \frac{1}{3}\alpha_{1,5}, \quad \alpha_{2,7} = \beta_{1,7}, \quad \alpha_{2,8} = \beta_{1,8}, \quad \alpha_{2,9} = \beta_{1,9}, \\ \{e_1, e_3\}, \quad &\Rightarrow \alpha_{3,1} = \alpha_{3,5} = \alpha_{3,6} = 0, \quad \alpha_{3,7} = -\frac{4}{3}\alpha_{1,5}, \quad \alpha_{3,8} = \frac{1}{2}(\beta_{1,7} - 5\alpha_{1,6}), \quad \alpha_{3,9} = -\frac{1}{2}\alpha_{1,7}, \\ \{e_1, e_4\}, \quad &\Rightarrow \alpha_{4,1} = \alpha_{4,5} = \alpha_{4,6} = \alpha_{4,7} = 0, \quad \alpha_{4,8} = \frac{1}{3}\alpha_{1,5}, \quad \alpha_{4,9} = \frac{1}{2}\alpha_{1,6}, \\ \{e_1, e_5\}, \quad &\Rightarrow \alpha_{5,1} = \alpha_{5,5} = \alpha_{5,6} = \alpha_{5,7} = \alpha_{5,8} = 0, \quad \alpha_{5,9} = -\frac{1}{2}\alpha_{1,5}, \\ \{e_1, e_j\}, \quad &\Rightarrow \alpha_{j,1} = \alpha_{j,5} = \alpha_{j,6} = \alpha_{j,7} = \alpha_{j,8} = \alpha_{j,9}, \quad 6 \leq j \leq 9, \\ \{e_2, e_3\}, \quad &\Rightarrow \beta_{3,7} = 0, \quad \beta_{3,8} = \frac{1}{6}(3\beta_{2,7} - 5\alpha_{1,5}), \quad \beta_{3,9} = -\frac{1}{2}\beta_{1,7}, \\ \{e_2, e_4\}, \quad &\Rightarrow \beta_{4,7} = \beta_{4,8} = 0, \quad \beta_{4,9} = \frac{1}{6}\alpha_{1,3}, \\ \{e_2, e_j\}, \quad &\Rightarrow \beta_{j,7} = \beta_{j,8} = \beta_{j,9} = 0, \quad 5 \leq j \leq 9. \end{aligned}$$

So, we have the following table of multiplications of the transposed Poisson algebra structure $(g_9^2, \cdot, [-, -])$. Note the associative identity for any triple is hold.

$$\begin{cases} e_1 \cdot e_1 = \alpha_1e_5 + \alpha_2e_6 + \alpha_3e_7 + \alpha_4e_8 + \alpha_5e_9, \quad e_1 \cdot e_2 = \frac{1}{3}\alpha_1e_6 + \alpha_6e_7 + \alpha_7e_8 + \alpha_8e_9, \\ e_1 \cdot e_3 = -\frac{4}{3}\alpha_1e_7 + \frac{1}{2}(\alpha_6 - 5\alpha_2)e_8 - \frac{1}{2}\alpha_3e_9, \quad e_1 \cdot e_4 = \frac{1}{3}\alpha_1e_8 + \frac{1}{2}\alpha_2e_9, \\ e_1 \cdot e_5 = -\frac{1}{2}\alpha_1e_9, \quad e_2 \cdot e_2 = \alpha_9e_7 + \alpha_{10}e_8 + \alpha_{11}e_9, \\ e_2 \cdot e_3 = \frac{1}{6}(3\alpha_9 - 5\alpha_1)e_8 - \frac{1}{2}\alpha_6e_9, \quad e_2 \cdot e_4 = \frac{1}{6}\alpha_1e_9, \quad e_3 \cdot e_3 = \frac{2}{3}\alpha_1e_9. \end{cases}$$

□

Theorem 25. Let $(g_{11}^3, \cdot, [-, -])$ be a transposed Poisson algebra structure defined on the Lie algebra g_{11}^3 . Then the multiplication of (g_{11}^3, \cdot) has the following form:

$$\text{TP}(g_{11}^3) : \begin{cases} e_1 \cdot e_1 = \alpha_6e_6 + \alpha_7e_7 + \alpha_8e_8 + \alpha_9e_9 + \alpha_{10}e_{10} + \alpha_{11}e_{11}, \\ e_1 \cdot e_2 = -\alpha_6e_7 - \alpha_7e_8 + \alpha_1e_9 + \alpha_2e_{10} + \alpha_3e_{11}, \quad e_1 \cdot e_3 = \frac{1}{2}\alpha_1e_{10} - \frac{1}{2}\alpha_9e_{11}, \\ e_1 \cdot e_4 = \frac{1}{2}\alpha_7e_{10} + \frac{1}{2}\alpha_8e_{11}, \quad e_1 \cdot e_5 = -\frac{1}{2}\alpha_6e_{10} - \frac{1}{2}\alpha_7e_{11}, \quad e_1 \cdot e_6 = \frac{1}{2}\alpha_6e_{11}, \\ e_2 \cdot e_2 = \alpha_6e_8 + \alpha_4e_9 + \alpha_5e_{10} + \alpha_{12}e_{11}, \quad e_2 \cdot e_3 = \frac{1}{2}\alpha_4e_{10} - \frac{1}{2}\alpha_1e_{11}, \\ e_2 \cdot e_4 = -\frac{1}{2}\alpha_6e_{10} - \frac{1}{2}\alpha_7e_{11}, \quad e_2 \cdot e_5 = \frac{1}{2}\alpha_6e_{11}, \end{cases}$$

where it is taken into account that the transposed Poisson algebra has its products with respect to the bracket $[-, -]$, and the remaining products are equal to zero.

Proof. Let $(g_{11}^3, \cdot, [-, -])$ be a transposed Poisson algebra structure defined on Lie algebra g_{11}^3 . Then for any element of $x \in g_{11}^3$ we have that, operator of multiplication $\varphi_x(y) = x \cdot y$ is a $\frac{1}{2}$ -derivation. Further, using Theorem 15 for $1 \leq i \leq 11$ we derive

$$\left\{ \begin{array}{l} \varphi_{e_i}(e_1) = \alpha_{i,1}e_1 + \alpha_{i,6}e_6 + \alpha_{i,7}e_7 + \alpha_{i,8}e_8 + \alpha_{i,9}e_9 + \alpha_{i,10}e_{10} + \alpha_{i,11}e_{11}, \\ \varphi_{e_i}(e_2) = \alpha_{i,1}e_2 - \alpha_{i,6}e_7 - \alpha_{i,7}e_8 + \beta_{i,9}e_9 + \beta_{i,10}e_{10} + \beta_{i,11}e_{11}, \\ \varphi_{e_i}(e_3) = \alpha_{i,1}e_3 + \frac{1}{2}\beta_{i,9}e_{10} - \frac{1}{2}\alpha_{i,9}e_{11}, \\ \varphi_{e_i}(e_4) = \alpha_{i,1}e_4 + \frac{1}{2}\alpha_{i,7}e_{10} + \frac{1}{2}\alpha_{i,8}e_{11}, \\ \varphi_{e_i}(e_5) = \alpha_{i,1}e_5 - \frac{1}{2}\alpha_{i,6}e_{10} - \frac{1}{2}\alpha_{i,7}e_{11}, \\ \varphi_{e_i}(e_6) = \alpha_{i,1}e_6 + \frac{1}{2}\alpha_{i,6}e_{11}, \\ \varphi_{e_i}(e_j) = \alpha_{i,1}e_j, \quad 7 \leq j \leq 11. \end{array} \right.$$

Considering the property $\varphi_{e_i}(e_j) = e_i \cdot e_j = e_j \cdot e_i = \varphi_{e_j}(e_i)$, we obtain the following restrictions:

$$\begin{aligned} \{e_1, e_2\}, \quad & \Rightarrow \quad \alpha_{1,1} = \alpha_{2,1} = \alpha_{2,6} = 0, \\ & \quad \alpha_{2,7} = -\alpha_{1,6}, \quad \alpha_{2,8} = -\alpha_{1,7}, \quad \alpha_{2,9} = \beta_{1,9}, \quad \alpha_{2,10} = \beta_{1,10}, \quad \alpha_{2,11} = \beta_{1,11}, \\ \{e_1, e_3\}, \quad & \Rightarrow \quad \alpha_{3,1} = \alpha_{3,6} = \alpha_{3,7} = \alpha_{3,8} = \alpha_{3,9} = 0, \quad \alpha_{3,10} = \frac{1}{2}\beta_{1,9}, \quad \alpha_{3,11} = -\frac{1}{2}\alpha_{1,9}, \\ \{e_1, e_4\}, \quad & \Rightarrow \quad \alpha_{4,1} = \alpha_{4,6} = \alpha_{4,7} = \alpha_{4,8} = \alpha_{4,9} = 0, \quad \alpha_{4,10} = \frac{1}{2}\alpha_{1,7}, \quad \alpha_{4,11} = \frac{1}{2}\alpha_{1,8}, \\ \{e_1, e_5\}, \quad & \Rightarrow \quad \alpha_{5,1} = \alpha_{5,6} = \alpha_{5,7} = \alpha_{5,8} = \alpha_{5,9} = 0, \quad \alpha_{5,10} = -\frac{1}{2}\alpha_{1,6}, \quad \alpha_{5,11} = -\frac{1}{2}\alpha_{1,7}, \\ \{e_1, e_6\}, \quad & \Rightarrow \quad \alpha_{6,1} = \alpha_{6,6} = \alpha_{6,7} = \alpha_{6,8} = \alpha_{6,9} = \alpha_{6,10} = 0, \quad \alpha_{6,11} = \frac{1}{2}\alpha_{1,6}, \\ \{e_1, e_j\}, \quad & \Rightarrow \quad \alpha_{j,1} = \alpha_{j,6} = \alpha_{j,7} = \alpha_{j,8} = \alpha_{j,9} = \alpha_{j,10} = \alpha_{j,11} = 0, \quad 7 \leq j \leq 9, \\ \{e_2, e_3\}, \quad & \Rightarrow \quad \beta_{3,9} = 0, \quad \beta_{3,10} = \frac{1}{2}\beta_{2,9}, \quad \beta_{3,11} = -\frac{1}{2}\beta_{1,9}, \\ \{e_2, e_4\}, \quad & \Rightarrow \quad \beta_{4,9} = 0, \quad \beta_{4,10} = -\frac{1}{2}\alpha_{1,6}, \quad \beta_{4,11} = -\frac{1}{2}\alpha_{1,7}, \\ \{e_2, e_5\}, \quad & \Rightarrow \quad \beta_{5,9} = \beta_{5,10} = 0, \quad \beta_{5,11} = \frac{1}{2}\alpha_{1,6}, \\ \{e_2, e_j\}, \quad & \Rightarrow \quad \beta_{j,9} = \beta_{j,10} = \beta_{j,11} = 0, \quad 6 \leq j \leq 11. \end{aligned}$$

Thus, from the above restrictions we obtain the transposed Poisson algebra structure $\mathbf{TP}(g_{11}^3)$ defined on Lie algebra g_{11}^3 . \square

REFERENCES

- [1] Abdurasulov K., Adashev J., Eshmeteva S. *Transposed Poisson structures on solvable Lie algebras with filiform nilradical*, arXiv:2401.04443.
- [2] Ancochea Bermúdez J. M., Campoamor-Stursberg R., García Vergnolle L. *Solvable Lie algebras with naturally graded nilradicals and their invariants*, J. Phys. A, 39(6) (2006), 1339-1355.
- [3] Bai C., Bai R., Guo L., Wu Y., *Transposed Poisson algebras, Novikov-Poisson algebras, and 3-Lie algebras*, Journal of Algebra, 632 (2023), 535-566.

- [4] Beites P., Fernández Ouardi A., Kaygorodov I., *The algebraic and geometric classification of transposed Poisson algebras*, Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas, 117 (2023), 2, 55.
- [5] Beites P., Ferreira B. L. M., Kaygorodov I., *Transposed Poisson structures*, Results in Mathematics, 79 (2024), 93, 29.
- [6] Ferreira B. L. M., Kaygorodov I., Lopatkin V., $\frac{1}{2}$ -derivations of Lie algebras and transposed Poisson algebras, Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas, 115 (2021), 3, 142.
- [7] Fernández Ouardi A., *On the simple transposed Poisson algebras and Jordan superalgebras*, Journal of Algebra, 641 (2024), 173-198.
- [8] Filippov V., δ -Derivations of Lie algebras, Siberian Mathematical Journal, 39 (1998), 6, 1218-1230.
- [9] Huang J., Chen X., Chen Z., Ding M., *On a conjecture of transposed Poisson n-Lie algebras*, AIMS Mathematics, 9 (2024), 3, 6709-6733.
- [10] Jaworska-Pastuszak A., Pogorzały Z., *Poisson structures for canonical algebras*, Journal of Geometry and Physics, 148 (2020), 103564.
- [11] Gómez J.R. , Jiménez-Merchán A. , Reyes J., *Quasi-filiform Lie algebras of maximum length*. Linear Algebra and its Applications, 335, (2001), 119-135.
- [12] Kaygorodov I., δ -derivations of simple finite-dimensional Jordan superalgebras, Algebra and Logic, 46 (2007), 5, 318-329.
- [13] Kaygorodov I., δ -superderivations of simple finite-dimensional Jordan and Lie superalgebra, Algebra and Logic, 49 (2010), 2, 130-144.
- [14] Kaygorodov I., *Non-associative algebraic structures: classification and structure*, Communications in Mathematics, 32 (3) (2024), 1-62
- [15] Kaygorodov I., Khrypchenko M., *Poisson structures on finitary incidence algebras*, Journal of Algebra, 578 (2021), 402-420.
- [16] Kaygorodov I., Khrypchenko M., *Transposed Poisson structures on Block Lie algebras and superalgebras*, Linear Algebra and Its Applications, 656 (2023), 167-197.
- [17] Kaygorodov I., Khrypchenko M., *Transposed Poisson structures on Witt type algebras*, Linear Algebra and its Applications, 665 (2023), 196-210.
- [18] Kaygorodov I., Khrypchenko M., *Transposed Poisson structures on generalized Witt algebras and Block Lie algebras*, Results in Mathematics, 78 (2023), 5, 186.
- [19] Kaygorodov I., Khrypchenko M., *Transposed Poisson structures on the Lie algebra of upper triangular matrices*, Portugaliae Mathematica, 81 (2024), 1-2, 135-149.
- [20] Kaygorodov I., Khrypchenko M., *Transposed Poisson structures on Lie incidence algebras*, Journal of Algebra, 647 (2024), 458-491.
- [21] Kaygorodov I., Khudoyberdiyev A., *Transposed Poisson structures on solvable and perfect Lie algebras*, Journal of Physics A: Mathematical and Theoretical, 57 (2024), 035205.
- [22] Kaygorodov I., Lopatkin V., Zhang Z., *Transposed Poisson structures on Galilean and solvable Lie algebras*, Journal of Geometry and Physics, 187 (2023), 104781.
- [23] Khudoyberdiyev A., Sheraliyeva S., *Extensions of solvable Lie algebras with naturally graded filiform nilradical*, Journal of Algebra and Its Applications, doi.org/10.1142/S0219498824501615.
- [24] Liu G., Bai C., *A bialgebra theory for transposed Poisson algebras via anti-pre-Lie bialgebras and anti-pre-Lie-Poisson bialgebras*, Communications in Contemporary Mathematics, (2023), 2350050.
- [25] Liu G., Bai C., *New splittings of operations of Poisson algebras and transposed Poisson algebras and related algebraic structures*, In: Hounkonnou, M.N., Mitrović, M., Abbas, M., Khan, M. (eds) Algebra without Borders – Classical and Constructive Nonassociative Algebraic Structures. STEAM-H: Science, Technology, Engineering, Agriculture, Mathematics & Health. Springer, Cham., 2023, 49-96.

- [26] Sartayev B., *Some generalizations of the variety of transposed Poisson algebras*, Communications in Mathematics, 32 (2024), 2, 55-62.
- [27] Šnobl L. Winternitz P., *A class of solvable Lie algebras and their Casimir invariants*, Journal of Physics A, 38 (2005), 2687-2700.
- [28] Vergne M. *Cohomologie des algèbres de Lie nilpotentes. Application à l'étude de la variété des algèbres de Lie nilpotentes*. Bull. Soc. Math. France, 98 (1970), 81-116.
- [29] Yang Ya., Tang X., Khudoyberdiyev A., *Transposed Poisson structures on Schrodinger algebra in $(n + 1)$ -dimensional space-time*, arXiv:2303.08180.
- [30] Yao Y., Ye Y., Zhang P., *Quiver Poisson algebras*, Journal of Algebra, 312 (2007), 2, 570-589.
- [31] Yaxi J., Chuangchuang K., Jiafeng L., *Classification of Transposed Poisson 3-Lie algebras of dimension 3*, arXiv:2401.02593.
- [32] Yuan L., Hua Q., $\frac{1}{2}$ -(bi)derivations and transposed Poisson algebra structures on Lie algebras, Linear and Multilinear Algebra, 70 (2022), 22, 7672-7701.