Volumes of components of Lelong upper level sets II

Shuang Su and Duc-Viet Vu

April 23, 2024

Abstract

Let *X* be a compact Kähler manifold of dimension *n*, and let *T* be a closed positive (1,1)-current in a nef cohomology class on *X*. We establish an optimal upper bound for the volume of components of Lelong upper level sets of *T* in terms of cohomology classes of non-pluripolar self-products of *T*.

Keywords: relative non-pluripolar product, density current, Lelong number.

Mathematics Subject Classification 2010: 32U15, 32Q15.

1 Introduction

The aim of this paper is to investigate the singularities of closed positive currents on compact Kähler manifolds. Let X be a compact Kähler manifold of dimension n, and let T be a closed positive (1, 1)-current on X. We are interested in understanding the set of points where T has a strictly positive Lelong numbers. By the celebrated upper semicontinuity of Lelong numbers by Siu [12], we know that this set is a countable union of proper analytic subsets on X. Our goal is to estimate the size of this upper level set. We note that by [13], for every closed positive current R on X, there always exists a closed positive (1, 1)-current whose Lelong numbers coincide with those of R. Thus, to study questions about Lelong numbers in this paper, it suffices to consider only currents of bi-degree (1, 1). To delve into details, let us first introduce some necessary notions.

Let ω be a fixed smooth Kähler form on X. We equip X with the Riemannian metric induced by ω . Let S be a closed positive (p, p)-current for some $0 \le p \le n$. We define the mass ||S|| of S to be equal to $\int_X S \wedge \omega^{n-p}$. For an analytic set V of pure dimension l in X, we recall that

$$\operatorname{vol}(V) = \frac{1}{l!} \int_{\operatorname{Reg}V} \omega^l$$

where RegV is the regular locus of V. Given a closed positive (p, p)-current R, we denote by $\{R\} \in H^{p,p}(X, \mathbb{R})$ the cohomology class of R. We say $\alpha \in H^{p,p}(X, \mathbb{R})$ is *pseudoeffective* if $\alpha = \{R\}$ for some closed positive (p, p)-current R. Let $\alpha, \beta \in H^{p,p}(X, \mathbb{R})$. We say $\alpha \ge \beta$ if $\alpha - \beta$ is a pseudoeffective class. For every $x \in X$, let $\nu(T, x)$ denote the Lelong number of T at x. For every irreducible analytic subset V in X, we recall that the generic Lelong number $\nu(T, V)$ of T along V is defined as $\inf_{x \in V} \{\nu(T, x)\}$. The Lelong number is a notion measuring the singularities of T. We refer to [4] for the basics of Lelong numbers. For every constant c > 0, we denote $E_c(T) := \{x \in X | \nu(T, x) \ge c\}$ and $E_+ := \{x \in X | \nu(T, x) > 0\}$. By [12], $E_c(T)$ is a proper analytic subset in X, and $E_+(T) = \bigcup_{m \in \mathbb{N}^*} E_{1/m}(T)$ is a countable union of analytic sets.

Let W be an irreducible analytic subset of dimension m in X. We denote by $E^W_+(T) := \{x \in W | \nu(T, x) > \nu(T, W)\}$ the Lelong upper level set of T on W, which is also a countable union of proper analytic subsets in W. Let $V \subset E^W_+(T)$ be an irreducible analytic set. We say that V is *maximal* if there is no irreducible analytic subset V' of $E^W_+(T)$ such that V is a proper subset of V'. We call V a component of the Lelong upper level set of T along W, and let $\mathscr{V}_{T,W}$ be the set of such components V. Observe that $\mathscr{V}_{T,W}$ has at most countably many elements. For $0 \leq l \leq m$, we denote by $\mathscr{V}_{l,T,W}$ the set of $V \in \mathscr{V}_{T,W}$ such that dim V = l.

Write $T = dd^c u$ locally, where u is a plurisubharmonic (psh in short) function. We define $T|_{\text{Reg}W}$ to be $dd^c(u|_{\text{Reg}W})$ if $u \neq -\infty$ on RegW, and $T|_{\text{Reg}W} := 0$ otherwise. One sees that this definition is independent of the choice of u. Thus, $T|_{\text{Reg}W}$ is a current on RegW. Here is our main result in the paper.

Theorem 1.1. Let α be a nef (1,1)-class and let W be an irreducible analytic subset of dimension m in X. Let T be a closed positive current in α such that $\nu(T,W) = 0$. Let $1 \le m' \le m$ be an integer. Then, we have

$$\sum_{V \in \mathscr{V}_{m-m',T,W}} \nu(T,V)^{m'} \operatorname{vol}(V) \le \frac{1}{(m-m')!} \int_{\operatorname{Reg}W} \left(\alpha^{m'} - \langle (T|_{\operatorname{Reg}W})^{m'} \rangle \right) \wedge \omega^{m-m'}, \quad (1.1)$$

where in the integral, we identify α with a smooth closed form in α .

We have some comments on (1.1). To see why the term

$$I := \int_{\operatorname{Reg}W} \left(\alpha^{m'} - \langle (T|_{\operatorname{Reg}W})^{m'} \rangle \right) \wedge \omega^{m-m'}$$

is non-negative, one can consider the case where W is smooth. Then, by a monotonicity of non-pluripolar products (see Theorem 3.3 below), the cohomology class $(\alpha|_W)^{m'} - \{\langle (T|_W)^{m'} \rangle\}$ is pseudoeffective. Hence the integral in the right-hand side of (1.1) is nonnegative. In the general case where W is singular, one can use either a desingularisation of W or interpret I as the mass of some non-pluripolar product relative to [W] (the current of integration along W); see Lemma 3.2 below. We underline however that in order to prove Theorem 1.1, it is not possible to use desingularisation of W to reduce to the case where W is smooth. The reason is that in the process of desingularisation, one has to blow up submanifolds of W which in general could be some of components of the Lelong upper level sets of T on W.

In [8], a less precise upper bound of volume of components of the Lelong upper level set was given in terms of the volume of W and the mass of T; see also Theorem 3.7 for a more general statement. The estimate (1.1) is indeed optimal. To see this, consider n = m = m' = 1 and T is a sum of Dirac masses. In this case, (1.1) becomes an equality.

If we consider W = X, then the generic Lelong number of T along W is zero. Thus, by Theorem 1.1, we have the following result.

Corollary 1.2. Let α be a nef (1,1)-class, and T be a closed positive current in α . For $0 \leq l \leq n$, let $\mathscr{V}_{l,T}$ be the set of $V \in \mathscr{V}_{T,X}$ such that $\dim V = l$. Let $1 \leq m' \leq n$ be an integer. Then, we have

$$\sum_{V \in \mathscr{V}_{n-m',T}} \nu(T,V)^{m'} \operatorname{vol}(V) \le \frac{1}{(n-m')!} \int_X \left(\alpha^{m'} - \{ \langle T^{m'} \rangle \} \right) \wedge \omega^{n-m'}.$$

Corollary 1.2 generalizes [5, Corollary 7.6] by Demailly in which it was assumed additionally that the components of the upper Lelong level set of T are only of dimension 0 (hence the cohomology class of T is necessarily nef, see [5, Lemma 6.3]). The feature of Corollary 1.2 is that it holds for any current in a nef class.

This paper refines and substitutes [16]. The proof of Theorem 1.1 requires both the theory of density currents in [7] and relative non-pluripolar products in [17] (see also [1, 2]). One of the keys is Theorem 3.6 below following from a general comparison of Lelong numbers for density currents (see Corollary 2.6).

The organization of this paper is as follows. In Section 2, we recall basic properties of density currents from [7]. In Section 3, we discuss the connection between the non-pluripolar product and density currents, and prove Theorem 1.1.

Acknowledgments. This research is partially funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)-Projektnummer 500055552 and by the ANR-DFG grant QuaSiDy, grant no ANR-21-CE40-0016.

2 Density currents

We first recall some basic properties of density currents introduced by Dinh-Sibony in [7].

Let X be a complex Kähler manifold of dimension n, and V a smooth complex submanifold of X of dimension l. Let T be a closed positive (p, p)-current on X, where $0 \le p \le n$. Denote by $\pi : E \to V$ the normal bundle of V in X and $\overline{E} := \mathbb{P}(E \oplus \mathbb{C})$ the projective compactification of E. By abuse of notation, we also use π to denote the natural projection from \overline{E} to V.

Let U be an open subset of X with $U \cap V \neq \emptyset$. Let τ be a smooth diffeomorphism from U to an open neighborhood of $V \cap U$ in E which is identity on $V \cap U$ such that the restriction of its differential $d\tau$ to $E|_{V \cap U}$ is identity. Such a map is called *an admissible map*. Note that in [7], to define an admissible map, it is required furthermore that $d\tau$ is \mathbb{C} -linear at every point of V. This difference doesn't affect what follows. When U is a small enough tubular neighborhood of V, there always exists an admissible map τ by [7, Lemma 4.2]. In general, τ is not holomorphic. When U is a small enough local chart, we can choose a holomorphic admissible map by using suitable holomorphic coordinates on U. For $\lambda \in \mathbb{C}^*$, let $A_{\lambda} : E \to E$ be the multiplication by λ on fibers of E. Here is the first fundamental result for density currents. **Theorem 2.1.** ([7, Theorem 4.6]) Let τ be an admissible map defined on a tubular neighborhood of V. Then, the family $(A_{\lambda})_*\tau_*T$ is of mass uniformly bounded in λ on compact subsets in E, and if S is a limit current of the last family as $\lambda \to \infty$, then S is a current on E which can be extended trivially through $\overline{E} \setminus E$ to be a closed positive current on \overline{E} such that the cohomology class $\{S\}$ of S in \overline{E} is independent of the choice of S, and $\{S\}|_V = \{T\}|_V$, and $\|S\| \leq C\|T\|$ for some constant C independent of S and T, where $\{S\}|_V$ denotes the pull-back of $\{S\}$ under the natural inclusion map from V to \overline{E} .

We call *S* a tangent current to *T* along *V*, and its cohomology class is called *the total* tangent class of *T* along *V* and is denoted by $\kappa^V(T)$. Tangent currents are not unique in general. By [7, Theorem 4.6] again, if

$$S = \lim_{k \to \infty} (A_{\lambda_k})_* \tau_* T$$

for some sequence $(\lambda_k)_k$ converging to ∞ , then for every open subset U' of X and every admissible map $\tau': U' \to E$, we also have

$$S = \lim_{k \to \infty} (A_{\lambda_k})_* \tau'_* T.$$

This is equivalent to saying that tangent currents are independent of the choice of the admissible map τ .

Definition 2.2. ([7, Definition 3.1]) Let F be a complex manifold and $\pi_F : F \to V$ a holomorphic submersion. Let S be a positive current of bi-degree (p, p) on F. The h-dimension of S with respect to π_F is the biggest integer q such that $S \wedge \pi_F^* \theta^q \neq 0$ for some Hermitian metric θ on V.

By a bi-degree reason, the h-dimension of *S* is in $[\max\{l - p, 0\}, \min\{\dim F - p, l\}]$. We have the following description of currents with minimal h-dimension.

Lemma 2.3. ([7, Lemma 3.4]) Let $\pi_F : F \to V$ be a submersion. Let S be a closed positive current of bi-degree (p, p) on F of h-dimension (l-p) with respect to π_F . Then $S = \pi^*S'$ for some closed positive current S' on V.

By [7, Lemma 3.8], the h-dimensions of tangent currents to T along V are the same and this number is called *the tangential h-dimension of* T along V.

Let $m \ge 2$ be an integer. Let T_j be a closed positive current of bi-degree (p_j, p_j) for $1 \le j \le m$ on X and let $T_1 \otimes \cdots \otimes T_m$ be the tensor current of T_1, \ldots, T_m which is a current on X^m . A *density current* associated to T_1, \ldots, T_m is a tangent current to $\otimes_{j=1}^m T_j$ along the diagonal Δ_m of X^m . Let $\pi_m : E_m \to \Delta$ be the normal bundle of Δ_m in X^m . Denote by [V] the current of integration along V. When m = 2 and $T_2 = [V]$, the density currents of T_1 and T_2 are naturally identified with the tangent currents to T_1 along V (see [14, Lemma 2.3]).

The unique cohomology class of density currents associated to T_1, \ldots, T_m is called *the total density class of* T_1, \ldots, T_m . We denote the last class by $\kappa(T_1, \ldots, T_m)$. The tangential h-dimension of $T_1 \otimes \cdots \otimes T_m$ along Δ_m is called *the density h-dimension* of T_1, \ldots, T_m . **Lemma 2.4.** ([7, Section 5]) Let T_j be a closed positive current of bi-degree (p_j, p_j) on X for $1 \le j \le m$ such that $\sum_{j=1}^m p_j \le n$. Assume that the density h-dimension of T_1, \ldots, T_m is minimal, i.e, equals to $n - \sum_{j=1}^m p_j$. Then the total density class of T_1, \ldots, T_m is equal to $\pi_m^*(\wedge_{j=1}^m \{T_j\})$.

Let $h_{\overline{E}}$ be the Chern class of the dual of the tautological line bundle of \overline{E} . By [7, Page 535], we have

$$\kappa^{V}(T) = \sum_{j=\max\{0,l-p\}}^{\min\{l,n-p-1\}} \pi^{*} \left(\kappa_{j}^{V}(T)\right) \wedge h_{\overline{E}}^{p-(l-j)},$$
(2.1)

where $\pi : \overline{E} \to V$ is the natural projection and $\kappa_j^V(T) \in H^{2l-2j}(V,\mathbb{R})$. The tangential *h*-dimension of *T* along *V* is exactly equal to the maximal *j* such that $\kappa_j^V(T) \neq 0$, and it was known that the class $\kappa_i^V(T)$ is pseudoeffective ([7, Lemma 3.15]).

Theorem 2.5. ([7, Proposition 4.13]) Let V' be a submanifold of V and let T be a closed positive current on X. Let T_{∞} be a tangent current to T along V. Let s be the tangential h-dimension of T_{∞} along V'. Then, the tangential h-dimension of T along V' is at most s, and we have

$$\kappa_s^{V'}(T) \le \kappa_s^{V'}(T_\infty).$$

As a consequence, we obtain the following result generalizing the well-known lower bound of Lelong numbers of intersection of (1, 1)-currents due to Demailly [6, Page 169] in the compact setting. It is probably the first result dealing with comparison of Lelong numbers for intersection of currents of arbitrary bi-degree.

Corollary 2.6. Let T_j be a closed positive current on X for $1 \le j \le m$. Then, for every $x \in X$ and for every density current S associated to T_1, \ldots, T_m , we have

$$\nu(S, x) \ge \nu(T_1, x) \cdots \nu(T_m, x). \tag{2.2}$$

Proof. Let $x \in X$. Let Δ_m be the diagonal of X^m . Let $\pi : E \to \Delta_m$ be the natural projection from the normal bundle of the diagonal Δ_m of X^m in X^m . We identify x with a point in Δ_m via the natural identification Δ_m with X. Put $T := \bigotimes_{j=1}^m T_j$ and $V' := \{x\}$. By [11, Lemma 2.4], we have $\nu(T, x) \ge \nu(T_1, x) \cdots \nu(T_m, x)$. By [7, Proposition 5.6], we have

$$\kappa_0^{V'}(S) = \nu(S, x)\delta_x, \quad \kappa_0^{V'}(T) = \nu(T, x)\delta_x,$$

where δ_x is the Dirac measure on x (notice here dim V' = 0). This combined with Theorem 2.5 applied to X^m , $T := \bigotimes_{j=1}^m T_j$, Δ_m the diagonal of X^m and $V' := \{x\}$ implies

$$\nu(S, x) \ge \nu(T, x) \ge \nu(T_1, x) \cdots \nu(T_m, x).$$

Hence, the desired inequality follows. The proof is finished.

5

3 Relative non-pluripolar products

We first recall basic facts about the relative non-pluripolar product of currents and discuss its connection with the density current.

Let X be a compact Kähler manifold of dimension n. Let T_1, \ldots, T_m be closed positive (1, 1)-currents on X, and T be a closed positive (p, p)-current. By [17], the T-relative non-pluripolar product $\langle \wedge_{j=1}^m T_j \dot{\wedge} T \rangle$ is defined in a way similar to that of the usual non-pluripolar products in [1, 2]. For readers' convenience, we explain briefly how to do it.

Write $T_j = dd^c u_j + \theta_j$, where θ_j is a smooth form and u_j is a θ_j -psh function. Put

$$R_k := \mathbf{1}_{\bigcap_{i=1}^m \{u_i > -k\}} \wedge_{i=1}^m \left(dd^c \max\{u_j, -k\} + \theta_j \right) \wedge T$$

for $k \in \mathbb{N}$. By the strong quasi-continuity of bounded psh functions ([17, Theorems 2.4 and 2.9]), we have

$$R_{k} = \mathbf{1}_{\bigcap_{i=1}^{m} \{u_{i} > -k\}} \wedge_{i=1}^{m} (dd^{c} \max\{u_{j}, -l\} + \theta_{j}) \wedge T$$

for every $l \ge k \ge 1$. One can check that R_k is positive (see [17, Lemma 3.2]).

As in [2], we have that R_k is of mass bounded uniformly in k and $(R_k)_k$ converges to a closed positive current as $k \to \infty$. This limit is denoted by $\langle \wedge_{j=1}^m T_j \dot{\wedge} T \rangle$. We refer to [17, Proposition 3.5] for properties of relative non-pluripolar products.

For every closed positive (1,1)-current P, we denote by I_P the set of $x \in X$ so that local potentials of P are equal to $-\infty$ at x. Note that I_P is a complete pluripolar set. It is clear from the definition that $\langle \wedge_{j=1}^m T_j \wedge T \rangle$ has no mass on $\bigcup_{j=1}^m I_{T_j}$. Furthermore, for every locally complete pluripolar set A (*i.e.*, A is locally equal to $\{\psi = -\infty\}$ for some psh function ψ), if T has no mass on A, then so does $\langle \wedge_{j=1}^m T_j \wedge T \rangle$. The following is deduced from [17, Proposition 3.5].

Proposition 3.1. (i) For $R := \langle \wedge_{j=l+1}^m T_j \dot{\wedge} T \rangle$, we have $\langle \wedge_{j=1}^m T_j \dot{\wedge} T \rangle = \langle \wedge_{j=1}^l T_j \dot{\wedge} R \rangle$.

(ii) For every complete pluripolar set A, we have

$$\mathbf{1}_{X\setminus A}\langle T_1 \wedge T_2 \wedge \cdots \wedge T_m \dot{\wedge} T \rangle = \langle T_1 \wedge T_2 \wedge \cdots \wedge T_m \dot{\wedge} (\mathbf{1}_{X\setminus A} T) \rangle.$$

In particular, the equality

$$\langle \wedge_{j=1}^m T_j \dot{\wedge} T \rangle = \langle \wedge_{j=1}^m T_j \dot{\wedge} T' \rangle$$

holds, where $T' := \mathbf{1}_{X \setminus \bigcup_{i=1}^m I_{T_i}} T$.

Let V be an irreducible analytic set in X. For the case T = [V], we have the following lemma.

Lemma 3.2. ([18, Lemma 2.3]) Let T_1, \ldots, T_m be closed positive (1, 1)-currents on X. Then the following properties hold:

(i) If V is contained in $\bigcup_{j=1}^{m} I_{T_j}$, then $\langle T_1 \wedge \cdots \wedge T_m \wedge [V] \rangle = 0$ and there is $1 \leq j_0 \leq m$ so that $V \subset I_{T_{j_0}}$.

(ii) If V is not contained in $\cup_{i=1}^{m} I_{T_i}$, then

$$\langle \wedge_{j=1}^m T_j \dot{\wedge} [V] \rangle = i_* \langle T_{1,V} \wedge \dots \wedge T_{m,V} \rangle,$$

where $i: \operatorname{Reg}(V) \to X$ is the natural inclusion, and $T_{j,V} := dd^c(u_j|_{\operatorname{Reg}(V)})$ if $dd^c u_j = T_j$ locally.

Let T, T' be closed positive (1, 1)-currents in the same cohomology class on X. T' is said to be less singular than T if for every local potentials u of T and u' of T', $u \le u' + O(1)$. Here is a crucial property of relative non-pluripolar products.

Theorem 3.3. ([17, Theorem 1.1]) Let T'_j be closed positive (1, 1)-current in the cohomology class of T_j on X such that T'_j is less singular than T_j for $1 \le j \le m$. Then we have

$$\{\langle T_1 \wedge \dots \wedge T_m \dot{\wedge} T \rangle\} \leq \{\langle T'_1 \wedge \dots \wedge T'_m \dot{\wedge} T \rangle\}.$$

Weaker versions of the above result were proved in [2, 3, 19]. Let $\alpha_1, \ldots, \alpha_m$ be big (1, 1)-classed of X. A current $T_{j,\min} \in \alpha_j$ is said to have minimal singularities if it is less singular than any closed positive current in α_j .

By Theorem 3.3, the class $\{\langle T_{1,\min} \wedge \cdots \wedge T_{m,\min} \dot{\wedge} T \rangle\}$ is a well-defined pseudoeffective class which is independent of the choice of $T_{j,\min}$. We denote the last class by $\{\langle \alpha_1 \wedge \cdots \wedge \alpha_m \dot{\wedge} T \rangle\}$. When $T \equiv 1$, we simply write $\langle \alpha_1 \wedge \cdots \wedge \alpha_m \rangle$ for $\{\langle \alpha_1 \wedge \cdots \wedge \alpha_m \dot{\wedge} T \rangle\}$. In this case, the product $\langle \alpha_1 \wedge \cdots \wedge \alpha_m \rangle$ was introduced in [2].

Regarding the relation between relative non-pluripolar products and density currents, the following fact was proved in [15, Theorem 3.5], see also [9, 10].

Theorem 3.4. Let R_{∞} be a density current associated to T_1, \ldots, T_m, T . Then we have

$$\pi_{m+1}^* \langle \wedge_{j=1}^m T_j \dot{\wedge} T \rangle \le R_{\infty}, \tag{3.1}$$

where π_{m+1} is the natural projection from the normal bundle of the diagonal Δ of X^{m+1} to Δ , and as usual we identified Δ with X.

We will need the following to estimate the density h-dimension of currents.

Proposition 3.5. ([15, Proposition 3.6]) Let T_1, \ldots, T_m, T be as above. Let A be a Borel subset of X. Assume that for every $J \subset \{1, \ldots, m\}$ and for every density current R_J associated to $(T_j)_{j \in J}, T$ with $m_J := |J| < m$, we have that R_J has no mass on the set

$$\pi_{m_J+1}^*(\bigcap_{j \notin J} \{x \in A : \nu(T_j, x) > 0\}).$$

Then for every density current S associated to T_1, \ldots, T_m, T , the h-dimension of the current $\mathbf{1}_A S$ is equal to n - p - m, where $\mathbf{1}_A$ denotes the characteristic function of A.

For every pseudoeffective (p, p)-class γ on X, we put $\|\gamma\| := \int_X \Theta \wedge \omega^{n-p}$, where Θ is any closed smooth form in γ . This definition is independent of the choice of Θ and is nonnegative because of the pseudoeffectivity of γ .

Theorem 3.6. Let P and T be closed positive currents of bi-degree (1, 1) and (p, p) respectively on X, where $1 \le p \le n - 1$. Assume that T has no mass on I_P . Then, the cohomology class

$$\gamma := \{P\} \land \{T\} - \{\langle P \dot{\land} T \rangle\}$$

is pseudoeffective and we have

$$\|\gamma\| \ge \sum_{V} \nu(P, V)\nu(T, V)n_{V}! \operatorname{vol}(V),$$
(3.2)

where the sum is taken over every irreducible subset V of dimension at least n - p - 1 in X, and $n_V := \dim V$.

Proof. Let \mathscr{V} be the set of irreducible analytic subsets V of dimension at least n - p - 1 in X such that $\nu(T, V) > 0$ and $\nu(P, V) > 0$. We note that in (3.2), it is enough to consider $V \in \mathscr{V}$. We will see below that \mathscr{V} has at most countable elements.

Observe that if $\nu(P, x) > 0$, then $x \in I_P$. Hence, by hypothesis, the trace measure of T has no mass on the set $\{x \in X : \nu(P, x) > 0\}$. This allows us to apply Proposition 3.5 to P and T to obtain that the density h-dimension of P and T is minimal. Using this and Lemma 2.4 gives

$$\kappa(P,T) = \pi^*(\{P\} \land \{T\}), \tag{3.3}$$

where π is the natural projection from the normal bundle of the diagonal Δ of X^2 to Δ .

Let S be a density current associated to P and T. Since the h-dimension of S is minimal, using Lemma 2.3, we get that there exists a current S' on X such that $S = \pi^* S'$ (recall Δ is identified with X). Since the relative non-pluripolar product is dominated by density currents (Theorem 3.4), the current $S' - \langle P \wedge T \rangle$ is closed and positive. Moreover, by (3.3), the cohomology class of the last current is equal to γ . It follows that γ is pseudoeffective.

It remains to prove (3.2). Let $V \in \mathscr{V}$. By definition, the generic Lelong number of T along V is positive. Since T is of bi-degree (p, p), the dimension of V must be at most n - p. Hence, we have two possibilities: either dim V = n - p - 1 or dim V = n - p. The latter case cannot happen. If T has no mass on V, then $\nu(T, V) = 0$, which leads to a contradiction. If T has mass on V, which is contained in I_P (for $\nu(P, V) > 0$), then this contradicts the hypothesis that T has no mass on I_P .

Let $V \in \mathscr{V}$. Since the Lelong numbers are preserved by submersion maps ([11, Proposition 2.3]), by applying Corollary 2.6 to P, T and generic $x \in V$, we obtain

$$\nu(S',V) = \nu(S,V) \ge \nu(P,V)\nu(T,V).$$

This combined with the fact that dim V = n - p - 1 implies $S' \ge \nu(P, V)\nu(T, V) [V]$. We deduce that

$$S' \ge \langle P \dot{\wedge} T \rangle + \mathbf{1}_{I_P} S'$$
$$\ge \langle P \dot{\wedge} T \rangle + \sum_{V \in \mathscr{V}} \nu(P, V) \nu(T, V) [V].$$

The second inequality comes from Siu's decomposition theorem ([6, 2.18]), and this also shows that \mathscr{V} has at most countable elements. The desired assertion follows and the proof is finished.

We now prove Theorem 1.1.

Proof of Theorem 1.1. It suffices to consider the case where α is Kähler by using $\alpha + \epsilon \{\omega\}, T + \epsilon \omega$ instead of α, T and letting $\epsilon \to 0$. Hence we assume from now on that α is Kähler. By abuse of notation, we also denote by α a smooth Kähler form in α . By Lemma 3.2, the right hand-hand side of (1.1) can be written as

$$\frac{1}{(m-m')!} \int_{\operatorname{Reg} W} \left(\alpha^{m'} - \langle (T|_{\operatorname{Reg} W})^{m'} \rangle \right) \wedge \omega^{m-m'} = \frac{1}{(m-m')!} \big\| \langle \alpha^{m'} \dot{\wedge} [W] \rangle - \langle T^{m'} \dot{\wedge} [W] \rangle \big\|.$$

Step 1. We first focus on the case that *T* has analytic singularities. Set $S = \langle T^{m'-1} \dot{\wedge} [W] \rangle$. Since α is Kähler, by the monotonicity of non-pluripolar product (Theorem 3.3) and Proposition 3.1 (i), we get

$$\left\| \langle \alpha^{m'} \dot{\wedge} [W] \rangle - \langle T^{m'} \dot{\wedge} [W] \rangle \right\| \ge \left\| \langle \alpha \wedge T^{m'-1} \dot{\wedge} [W] \rangle - \langle T^{m'} \dot{\wedge} [W] \rangle \right\|$$

$$= \left\| \alpha \wedge S - \langle T \dot{\wedge} S \rangle \right\|$$

$$(3.4)$$

We now show that S has no mass on I_T . For m' > 1, this directly follows from the definition of non-pluripolar product. For m' = 1, the current S is just [W]. Since we assume that T has analytic singularities, the polar locus I_T is an analytic subset and it doesn't contain W. Hence, [W] also has no mass on I_T . Therefore, we can apply Lemma 3.6 to T, S, and get

$$\|\alpha \wedge \{S\} - \{\langle T \dot{\wedge} S \rangle\}\| \ge (m - m')! \sum_{V \in \mathscr{V}_{m-m',T,W}} \nu(T,V)\nu(S,V) \operatorname{vol}(V)$$
(3.5)

Let $V \in \mathscr{V}_{m-m',T,W}$ and let $\operatorname{Sing}(I_T \cap W)$ be the singular locus of the analytic set $I_T \cap W$. Put $X' := X \setminus \operatorname{Sing}(I_T \cap W)$ which is an open subset in X. Now, we claim that the classical intersection $T^{m'-1} \wedge [W]$ is well-defined on some open neighborhood U of $V \setminus \operatorname{Sing}(I_T \cap W)$ in X', in the sense given in [4, Chapter III]. Since T has analytic singularities, the Lelong number $\nu(T, x)$ is strictly positive if and only if x belongs to I_T . This implies that V is contained in $I_T \cap W$. Let K_1, \ldots, K_s be the irreducible components of $I_T \cap W$. Since V is a maximal irreducible proper analytic set such that $\nu(T, V) > 0$, V must be one of the irreducible components. By rearranging the index, we may assume $V = K_1$. Let U be an open neighborhood of $V \setminus \operatorname{Sing}(I_T \cap W)$ such that

$$U \cap K_j = \emptyset$$
, for $j = 2, \ldots, s$.

Notice that $V \setminus \text{Sing}(I_T \cap W)$ is contained in Reg(V), and is of dimension m - m'. Consequently, for $0 \le j' \le m' - 1$,

$$\mathscr{H}_{2m-2j'+1}(L(T)|_U \cap W) = \mathscr{H}_{2m-2j'+1}(I_T \cap W \cap U)$$
$$= \mathscr{H}_{2m-2j'+1}(V \setminus \operatorname{Sing}(I_T \cap W))$$
$$= 0,$$

where L(T) is the set of $x \in X$ such that the local potential of T is unbounded on any neighborhood of x. This allows us to apply [4, Chapter III, Theorem 4.5] and get the well-definedness of $T^{m'-1} \wedge [W]$ on U. By [17, Proposition 3.6], we obtain

$$\langle T^{m'-1}\dot{\wedge}[W]\rangle = \mathbf{1}_{U\setminus I_T}T^{m'-1}\wedge [W].$$
(3.6)

Actually, the equality also holds on $U \cap I_T$. To show this, we need to check that $T^{m'-1} \wedge [W]$ has no mass on $U \cap I_T$. Since $\dim(U \cap I_T \cap W) = m - m'$ and $T^{m'-1} \wedge [W]$ is of bi-dimension (m - m' + 1, m - m' + 1), the current $T^{m'-1} \wedge [W]$ must have no mass on $U \cap I_T \cap W$. Also, by the fact $\operatorname{Supp}(T^{m'-1} \wedge [W]) \subset W$, the current $T^{m'-1} \wedge [W]$ also has no mass on $(U \cap I_T) \setminus W$. Therefore, the equality (3.6) extends to U. This implies that the Lelong number $\nu(S, V)$ equals $\nu(T^{m'-1} \wedge [W], V \setminus \operatorname{Sing}(I_T \cap W))$ (remember that we consider the current $T^{m'-1} \wedge [W]$ on U, and $V \setminus \operatorname{Sing}(I_T \cap W)$ is an analytic subset in U), and we then have

$$\nu(S,V) = \nu(T^{m'-1} \wedge [W], V \setminus \operatorname{Sing}(I_T \cap W))$$

$$\geq \nu(T, V \setminus \operatorname{Sing}(I_T \cap W))^{m'-1} \nu([W], V \setminus \operatorname{Sing}(I_T \cap W))$$

$$\geq \nu(T, V)^{m'-1}.$$
(3.7)

By (3.4), (3.5) and (3.7), the desired inequality follows in the case where T has analytic singularities.

Step 2. Now, we remove the assumption that *T* has analytic singularities. First, we write $T = dd^c u + \theta$, where θ is a closed smooth (1, 1)- form and $u \in PSH(X, \theta)$. Demailly's analytic approximation theorem (see [6, Corollary 14.13]) allows us to construct a sequence $u_k^D \in PSH(X, \theta + \epsilon_k \omega)$, where ϵ_k decreases to 0, such that

- (1) $u_k^D \ge u$ and u_k^D converges to u in L^1 .
- (2) u_k^D has analytic singularities.
- (3) $\nu(T_k, x)$ converges to $\nu(T, x)$ uniformly on X, where $T_k = dd^c u_k^D + (\theta + \epsilon_k \omega)$.

By the monotonicity property of non-pluripolar product (Theorem 3.3), we have

$$\begin{aligned} \left\| \langle \alpha^{m'} \dot{\wedge}[W] \rangle - \langle T^{m'} \dot{\wedge}[W] \rangle \right\| &= \lim_{k \to \infty} \left\| \langle (\alpha + \epsilon_k \omega)^{m'} \dot{\wedge}[W] \rangle - \langle (T + \epsilon_k \omega)^{m'} \dot{\wedge}[W] \rangle \right\| \\ &\geq \limsup_{k \to \infty} \left\| \langle (\alpha + \epsilon_k \omega)^{m'} \dot{\wedge}[W] \rangle - \langle T_k^{m'} \dot{\wedge}[W] \rangle \right\| \end{aligned}$$
(3.8)

For every constant r > 0, set $A_r := \{V \in \mathscr{V}_{m-m',T,W} | \nu(T,V) \ge r\}$. Observe that A_r increases to $\mathscr{V}_{m-m',T,W}$ as $r \to 0$. Since $\nu(T_k, x)$ converges to $\nu(T, x)$ uniformly and T_k is less singular than T, for every fixed r > 0 we have

$$A_r \subset \mathscr{V}_{m-m',T_k,W}$$

when k is large enough. By Step 1, we therefore have

$$\begin{aligned} \left\| \langle (\alpha + \epsilon_k \omega)^{m'} \dot{\wedge} [W] \rangle - \langle T_k^{m'} \dot{\wedge} [W] \rangle \right\| &\geq (m - m')! \sum_{V \in \mathscr{V}_{m - m', T_k, W}} \nu(T_k, V)^{m'} \operatorname{vol}(V) \\ &\geq (m - m')! \sum_{V \in A_r} \nu(T_k, V)^{m'} \operatorname{vol}(V). \end{aligned}$$

Letting $k \to \infty$ and using (3.8) give

$$\begin{aligned} \left\| \langle \alpha^{m'} \dot{\wedge} [W] \rangle - \langle T^{m'} \dot{\wedge} [W] \rangle \right\| &\geq (m - m')! \limsup_{k \to \infty} \sum_{V \in A_r} \nu(T_k, V)^{m'} \operatorname{vol}(V) \\ &= (m - m')! \sum_{V \in A_r} \nu(T, V)^{m'} \operatorname{vol}(V), \end{aligned}$$

for every constant r > 0. Letting $r \to 0$, we obtain the desired estimate.

For the general case where $\nu(T, W) > 0$. We couldn't directly compare the volume of Lelong upper level sets of T on W and the mass of $\{\langle \alpha^{m'} \dot{\wedge} [W] \rangle\} - \{\langle T^{m'} \dot{\wedge} [W] \rangle\}\}$. In this case, we have the following modified inequality which is stronger than [8, Theorem 1.1].

Theorem 3.7. Let α be a nef (1, 1)-class. Let W be an irreducible analytic subset in X. Let T be a closed positive current in α . Let $1 \le m' \le m$ be an integer. Then we have

$$(m-m')! \sum_{V \in \mathscr{V}_{m-m',T,W}} \left(\nu(T,V) - \nu(T,W) \right)^{m'} \operatorname{vol}(V) \leq \left\| \left(\alpha + c\{\omega\} \right)^{m'} \wedge \{[W]\} - \left\{ \langle (T+c\omega)^{m'} \dot{\wedge}[W] \rangle \right\} \right\|$$
(3.9)

where $c = c_1 \cdot \nu(T, W)$ and $c_1 > 0$ is a constant independent of α, T, W . In particular, there is a constant $c_2 > 0$ independent of α, T, W such that

$$\sum_{V \in \mathscr{V}_{m-m',T,W}} \left(\nu(T,V) - \nu(T,W) \right)^{m'} \operatorname{vol}(V) \le c_2 \operatorname{vol}(W) \|T\|^{m'}.$$
(3.10)

Proof. The desired inequality (3.10) follows directly from (3.9). We prove now (3.9). For convenience, set $c_3 := \nu(T, W) > 0$. The regularization theorem of currents introduced in [5] allows us to cut down the Lelong upper level set $\{x \in X | \nu(T, V) \ge c_3\}$ from *T*. More precisely, by [5, Theorem 1.1], there exists a sequence of almost positive closed (1, 1)-currents $T_{c_3,k}$ in α such that

- (1) $T_{c_3,k} \ge -(c_1 \cdot c_3 + \epsilon_k)\omega$, where $\lim_{k\to\infty} \epsilon_k = 0$ and $c_1 > 0$ is a constant independent of α, T and W.
- (2) The global potentials of $T_{c_3,k}$ decreases to the global potential of T.
- (3) $\nu(T_{c_3,k},x) = \max\{\nu(T,x) c_3, 0\}.$

Set $\widetilde{T}_{c_3,k} = T_{c_3,k} + (c_1 \cdot c_3 + \epsilon_k)\omega$, which is a closed positive (1, 1)-current. By Theorem 3.3, we have

$$\left\| (\alpha + c_1 \cdot c_3\{\omega\})^{m'} \wedge \{[W]\} - \langle (T + c_1 \cdot c_3\omega)^{m'} \dot{\wedge}[W] \rangle \right\|$$

$$\geq \limsup_{k \to \infty} \left\| (\alpha + (c_1 \cdot c_3 + \epsilon_k)\{\omega\})^{m'} \wedge \{[W]\} - \langle \widetilde{T}_{c_3,k}^{m'} \dot{\wedge}[W] \rangle \right\|$$

Since $\nu(\widetilde{T}_{c_3,k},W) = 0$, we can apply Theorem 1.1 to the right-hand side of the above inequality and get

$$\left\| \left(\alpha + (c_1 \cdot c_3 + \epsilon_k) \{\omega\} \right)^{m'} \wedge \{[W]\} - \langle \widetilde{T}_{c_3,k}^{m'} \dot{\wedge}[W] \rangle \right\|$$

$$\geq (m - m')! \sum_{V \in \mathscr{V}_{m-m', \tilde{T}_{c_3,k}, W}} \nu(\widetilde{T}_{c_3,k}, V)^{m'} \operatorname{vol}(V), \qquad (3.11)$$

By the above properties of $T_{c_3,k}$, we have

$$\mathscr{V}_{m-m',\tilde{T}_{c_3,k},W}=\mathscr{V}_{m-m',T,W}.$$

Therefore, the right-hand side of (3.11) is equal to

$$(m-m')! \sum_{V \in \mathscr{V}_{m-m',T,W}} \nu(\widetilde{T}_{c_2,k},V)^{m'} \operatorname{vol}(V)$$

= $(m-m')! \sum_{V \in \mathscr{V}_{m-m',T,W}} (\nu(T,V) - \nu(T,W))^{m'} \operatorname{vol}(V).$

This completes the proof.

References

- [1] E. BEDFORD AND B. A. TAYLOR, *Fine topology, Šilov boundary, and* $(dd^c)^n$, J. Funct. Anal., 72 (1987), pp. 225–251.
- [2] S. BOUCKSOM, P. EYSSIDIEUX, V. GUEDJ, AND A. ZERIAHI, Monge-Ampère equations in big cohomology classes, Acta Math., 205 (2010), pp. 199–262.
- [3] T. DARVAS, E. DI NEZZA, AND C. H. LU, Monotonicity of nonpluripolar products and complex Monge-Ampère equations with prescribed singularity, Anal. PDE, 11 (2018), pp. 2049–2087.
- [4] J.-P. DEMAILLY, Complex analytic and differential geometry. http://www.fourier.ujf-grenoble.fr/~demailly.
- [5] —, Regularization of closed positive currents and intersection theory, J. Algebraic Geom., 1 (1992), pp. 361–409.
- [6] —, *Analytic methods in algebraic geometry*, vol. 1 of Surveys of Modern Mathematics, International Press, Somerville, MA; Higher Education Press, Beijing, 2012.
- [7] T.-C. DINH AND N. SIBONY, Density of positive closed currents, a theory of nongeneric intersections, J. Algebraic Geom., 27 (2018), pp. 497–551.
- [8] D. T. DO AND D.-V. VU, Volume of components of Lelong upper-level sets, J. Geom. Anal., 33 (2023), pp. Paper No. 303, 14.

- [9] D. T. HUYNH, L. KAUFMANN, AND D.-V. VU, Intersection of (1,1)-currents and the domain of definition of the Monge-Ampère operator, Indiana Univ. Math. J., 72 (2023), pp. 239–261.
- [10] L. KAUFMANN AND D.-V. VU, Density and intersection of (1,1)-currents, J. Funct. Anal., 277 (2019), pp. 392–417.
- [11] M. MEO, Inégalités d'auto-intersection pour les courants positifs fermés définis dans les variétés projectives, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 26 (1998), pp. 161– 184.
- [12] Y. T. SIU, Analyticity of sets associated to Lelong numbers and the extension of closed positive currents, Invent. Math., 27 (1974), pp. 53–156.
- [13] G. VIGNY, Lelong-Skoda transform for compact Kähler manifolds and self-intersection inequalities, J. Geom. Anal., 19 (2009), pp. 433–451.
- [14] D.-V. VU, Densities of currents on non-Kähler manifolds. https://doi.org/10.1093/imrn/rnz270. Int. Math. Res. Not. IMRN.
- [15] —, Density currents and relative non-pluripolar products, Bull. Lond. Math. Soc., 53 (2021), pp. 548–559.
- [16] _____, Loss of mass of non-pluripolar products. arXiv:2101.05483, 2021.
- [17] —, Relative non-pluripolar product of currents, Ann. Global Anal. Geom., 60 (2021), pp. 269–311.
- [18] —, Derivative of volumes of big cohomology classes. arXiv:2307.15909, 2023.
- [19] D. WITT NYSTRÖM, Monotonicity of non-pluripolar Monge-Ampère masses, Indiana Univ. Math. J., 68 (2019), pp. 579–591.

DUC-VIET VU, UNIVERSITY OF COLOGNE, DIVISION OF MATHEMATICS, DEPARTMENT OF MATHEMATICS AND

COMPUTER SCIENCE, WEYERTAL 86-90, 50931, KÖLN, GERMANY *E-mail address*: dvu@uni-koeln.de

SHUANG SU, UNIVERSITY OF COLOGNE, DIVISION OF MATHEMATICS, DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, WEYERTAL 86-90, 50931, KÖLN, GERMANY

E-mail address: ssu1@uni-koeln.de