arXiv:2404.14125v1 [math.GR] 22 Apr 2024

WEIGHTS FOR π -PARTIAL CHARACTERS OF π -SEPARABLE GROUPS

XUEWU CHANG AND PING JIN

ABSTRACT. The aim of this paper is to confirm an inequality predicted by Isaacs and Navarro in 1995, which asserts that for any π' -subgroup Q of a π -separable group G, the number of π' -weights of G with Q as the first component always exceeds that of irreducible π -partial characters of G with Q as their vertex. We also give some sufficient condition to guarantee that these two numbers are equal, and thereby strengthen their main theorem on the π -version of the Alperin weight conjecture.

1. INTRODUCTION

Let *p* be a prime, and let *G* be a finite group. Recall that a *p*-weight of *G* is a pair (Q, τ) , where *Q* is a *p*-subgroup of *G* and $\tau \in Irr(\mathbf{N}_G(Q)/Q)$ has *p*-defect zero, i.e., whose degree has the same *p*-part as $|\mathbf{N}_G(Q) : Q|$. The Alperin weight conjecture (often abbreviated AWC) asserts that the number of *G*-conjugacy classes of *p*-weights coincides with the number of conjugacy classes of *p*-weights coincides with the number of *G*.

The first complete published proof of the AWC for *p*-solvable groups was given by Isaacs and Navarro [4] in 1995 by introducing many new techniques to the π -theory of characters of π separable groups for a set π of primes, and in fact, they established a π -version of this conjecture. Specifically, they defined a π' -weight of a π -separable group *G* to be an ordered pair (Q, τ) , where *Q* is a π' -subgroup of *G* and $\tau \in Irr(N_G(Q)/Q)$ has *p*-defect zero for every prime $p \in \pi'$, and then proved the following result, which appears in [4] as Theorem A and reduces to the *p*-solvable case of AWC by taking $\pi = p'$, the complement of the prime *p*.

Theorem 1.1 (Isaacs-Navarro). Let G be a π -separable group, and assume that the Hall π complements of G are nilpotent. Then the number of G-conjugacy classes of π' -weights is equal to
the number of conjugacy classes of π -elements of G.

Furthermore, they defined the **vertices** for every irreducible π -partial character of a π -separable group G, and showed that the set of vertices for an irreducible π -partial character of G forms a single conjugacy class of π' -subgroups of G. We will review the relevant definitions and properties in the next section. Now let Q be a π' -subgroup of G, and write $I_{\pi}(G|Q)$ for the set of irreducible π -partial characters of G with vertex Q. It is easy to show that $|I_{\pi}(\mathbf{N}_G(Q)|Q)|$ is exactly the number of π' -weights of G with first component Q; see Lemma 6.28 of [3] or Lemma 2.4 below.

Key words and phrases. weight; vertex; π -separable group; π -partial character.

Using the notion of vertices, Isaacs and Navarro obtained a strengthened version of Theorem 1.1, and they also weakened the nilpotency condition on Q; see Theorem C and Theorem 6.3 of [4].

Theorem 1.2 (Isaacs-Navarro). Let G be π -separable, and let $Q \leq G$ be a solvable π' -subgroup with the property that whenever $Q \leq X < Y \leq G$, where Y is a π' -subgroup, we have $X < \mathbf{N}_Y(X)$. Then $|I_{\pi}(G|Q)| = |I_{\pi}(\mathbf{N}_G(Q)|Q)|$.

The main purpose of this paper is to confirm an inequality predicted by Isaacs and Navarro at the end of [4], which asserts that for any π' -subgroup Q of a π -separable group G, the number of π' -weights of G with Q as the first component always exceeds that of irreducible π -partial characters of G with Q as their vertex or, equivalently, that $|I_{\pi}(N_G(Q)|Q)| \ge |I_{\pi}(G|Q)|$.

Theorem A. Let G be a π -separable group, and let Q be a π '-subgroup of G. Then

 $|\mathbf{I}_{\pi}(G|Q)| \leq |\mathbf{I}_{\pi}(\mathbf{N}_G(Q)|Q)|.$

Our proof of Theorem A was inspired by the techniques developed by Navarro and Sambale in [7]. Actually, they explored nilpotent π' -weights in π -separable groups and gave a generalization of Theorem 1.1.

Using the same argument as in the proof of Theorem A, we obtain a strong form of Theorem 1.2.

Theorem B. Let G be a π -separable group, and suppose that $Q \leq G$ is a π' -subgroup with the property that $Q < \mathbf{N}_X(Q)$ for every π' -subgroup X of G that properly contains Q. Then

 $|\mathbf{I}_{\pi}(G|Q)| = |\mathbf{I}_{\pi}(\mathbf{N}_G(Q)|Q)|.$

The following is an immediate consequence of Theorem B in which $X_{\pi}(G)$ denotes the set of π -special characters of *G* (for the definition, see Section 2 below).

Corollary C. Let G be a π -separable group and let Q be a Hall π -complement of G. Then

- (a) $|I_{\pi}(G|Q)| = |I_{\pi}(N_G(Q)|Q)|$, and
- (b) $|X_{\pi}(G)| = |\operatorname{Irr}(\mathbf{N}_{G}(Q)/Q)|.$

We mention that part (b) in the above corollary is a deep result due to Wolf, which appears in [9] as Corollary 1.16.

All groups considered in this paper are finite, and the notation and terminology are mostly taken from Isaacs' books [1] and [3].

2. Preliminaries

In this section, we will review some necessary materials and most of them are taken from [3].

2.1. π -Partial characters and vertices. Let *G* be a π -separable group for a set π of primes, and let G^0 be the set of π -elements of *G*. For any complex character χ of *G*, we say that the restriction χ^0 of χ on G^0 is a π -partial character of *G*. If a π -partial character φ of *G* cannot be written as a sum of two π -partial characters of *G*, then we say that φ is an **irreducible** π -partial character of *G*. We use $I_{\pi}(G)$ to denote the set of irreducible π -partial characters of *G*. Note that if $\pi = p'$, the complement of a single prime *p*, then $I_{\pi}(G) = IBr_p(G)$ by the Fong-Swan theorem for *p*-solvable groups, and if π is the set of all primes, then $I_{\pi}(G) = Irr(G)$.

Recall that an irreducible complex character $\chi \in Irr(G)$ is π -special if $\chi(1)$ is a π -number and the determinantal order $o(\theta)$ is a π -number, where θ is any irreducible constituent of χ_S for every subnormal subgroup *S* of *G*. Denote by $X_{\pi}(G)$ for the set of all π -special characters of *G*.

The following is essential in the study of π -partial characters.

Lemma 2.1 ([3, Theorem 3.14]). Let G be π -separable. Then the map $\chi \mapsto \chi^0$ defines a bijection from $X_{\pi}(G)$ onto the set of those irreducible π -partial characters of G whose degree is a π -number.

Now let Q be a π' -subgroup of G. We say that Q is a **vertex** for an irreducible π -partial character $\varphi \in I_{\pi}(G)$ if there exists a subgroup U of G and $\theta \in I_{\pi}(U)$, such that $\varphi = \theta^{G}$, $\theta(1)$ is a π -number and Q is a Hall π' -subgroup of U. As mentioned in the introduction, we write $I_{\pi}(G|Q)$ for the set of irreducible π -partial characters of G with vertex Q.

Lemma 2.2 ([3, Corollary 5.18]). Let G be π -separable and let $\varphi \in I_{\pi}(G)$. If $\varphi \in I_{\pi}(G|Q)$, then $\varphi(1)_{\pi'}|Q| = |G|_{\pi'}$. In particular, if Q is a Hall π' -subgroup of G, then $\varphi \in I_{\pi}(G|Q)$ if and only if $\varphi(1)$ is a π -number.

Also, we say that $\chi \in \operatorname{Irr}(G)$ has π' -defect zero if $\chi(1)_{\pi'} = |G|_{\pi'}$ or equivalently, if χ has *p*-defect zero for every $p \in \pi'$. Write dz(G) for the set of irreducible complex characters of *G* having π' -defect zero, so a pair (Q, τ) is a π' -weight of *G* if *Q* is a π' -subgroup of *G* and $\tau \in dz(\mathbf{N}_G(Q)/Q)$.

We need two basic facts on vertices and weights.

Lemma 2.3 ([3, Theorem 5.17]). Let G be a π -separable group and let $\varphi \in I_{\pi}(G)$. Then all vertices for φ is a single conjugacy class of π' -subgroups of G.

Lemma 2.4 ([3, Lemma 6.28]). Let G be a π -separable group and let Q be a π '-subgroup of G. Then the number of π '-weights of G with first component Q is equal to $|I_{\pi}(\mathbf{N}_G(Q)|Q)|$. Actually, $\tau \mapsto \tau^0$ defines a canonical bijection $dz(\mathbf{N}_G(Q)/Q) \to I_{\pi}(\mathbf{N}_G(Q)|Q)$.

2.2. The Clifford correspondence for π -partial characters. Let $K \leq G$ and $\theta \in I_{\pi}(K)$. We say that $\varphi \in I_{\pi}(G)$ lies over θ or equivalently, that θ lies under φ if θ is an irreducible constituent of

 φ_K . We use $I_{\pi}(G|\theta)$ to denote the set of all irreducible π -partial characters of G lying over θ . It is natural to define θ^G by the usual formula for induced characters but applied only to π -elements of G, so that θ^G is also a π -partial character of G.

Furthermore, if $K \triangleleft G$, then $\theta^g \in I_{\pi}(K)$ for all $g \in G$, where θ^g is defined by $\theta^g(x^g) = \theta(x)$ for $x \in K^0$. Then *G* acts on the set $I_{\pi}(K)$ via conjugation, and we write G_{θ} for the stabilizer of θ in *G*. Now we can state the Clifford correspondence for π -partial characters.

Lemma 2.5 ([3, Theorem 5.11]). Suppose that $\theta \in I_{\pi}(N)$, where $N \triangleleft G$ and G is π -separable. Then induction $\alpha \mapsto \alpha^G$ defines a bijection $I_{\pi}(G_{\theta}|\theta) \rightarrow I_{\pi}(G|\theta)$. Also, if $\varphi = \alpha^G$, then α is the unique irreducible π -partial character of G_{θ} that lies under φ and over θ , and in this case, we often write $\alpha = \varphi_{\theta}$ and call it the Clifford correspondent of φ over θ .

Following Navarro-Sambale [7], we write $I_{\pi}(G|Q, \tau) = I_{\pi}(G|Q) \cap I_{\pi}(G|\tau)$, where Q is a π' -subgroup of a π -separable group G and $\tau \in I_{\pi}(N)$ for some $N \triangleleft G$.

Lemma 2.6 ([3, Lemma 6.33]). Let G be a π -separable group and let $\varphi \in I_{\pi}(G|Q)$, where Q is a π' -subgroup of G, and let $K \triangleleft G$.

- (a) There exists a unique $N_G(Q)$ -orbit of some $\theta \in I_{\pi}(K)$ such that $\varphi_{\theta} \in I_{\pi}(G_{\theta}|Q)$.
- (b) If $\tau \in I_{\pi}(K)$ is *Q*-invariant and $G = G_{\tau} \mathbf{N}_{G}(Q)$, then induction of π -partial characters defines a bijection $I_{\pi}(G_{\tau}|Q,\tau) \to I_{\pi}(G|Q,\tau)$.

The following corollary tells us how to count the quantity $|I_{\pi}(G|Q)|$ "over a normal subgroup".

Corollary 2.7. Let G be a π -separable group with $K \triangleleft G$, and suppose that Q is a π' -subgroup of G. Let \mathcal{A} be a set of representatives for the $N_G(Q)$ -orbits of Q-invariant members of $I_{\pi}(K)$. Then

$$|\mathbf{I}_{\pi}(G|Q)| = \sum_{\tau \in \mathscr{A}} |\mathbf{I}_{\pi}(G_{\tau}|Q,\tau)|.$$

Sometimes we need to work in quotient groups modulo normal π' -subgroups when studying π -partial characters and vertices.

Lemma 2.8 ([3, Theorem 3.10 and Lemma 5.31]). Let *G* be a π -separable group and let $N \triangleleft G$ be a π '-subgroup. Write $\overline{G} = G/N$. Then

- (a) There is a bijection $\varphi \mapsto \overline{\varphi}$ from $I_{\pi}(G)$ onto $I_{\pi}(\overline{G})$, where $\overline{\varphi}(\overline{x}) = \varphi(x)$ for all $x \in G^0$.
- (b) If $\varphi \in I_{\pi}(G|Q)$, where Q is a π' -subgroup of G, then $N \leq Q$ and $\bar{\varphi} \in I_{\pi}(\bar{G}|\bar{Q})$. In particular, we have $\mathbf{O}_{\pi'}(G) \leq Q$.

Lemma 2.9 ([3, Theorem 6.31]). Let G be a π -separable group and let $K \triangleleft G$ be a π -subgroup. Also, let $Q \leq G$ be a π' -subgroup, and let $L \triangleleft G$, where $L \leq Q$. Let $\theta \in \text{Irr}(K)$ be G-invariant, and write $\overline{G} = G/L$. Then $|I_{\pi}(G|Q, \theta)| = |I_{\pi}(\overline{G}|\overline{Q}, \overline{\theta})|$, where $\overline{\theta} \in \text{Irr}(\overline{K})$ corresponds to θ via the natural isomorphism $K \to KL/L$. 2.3. Vertices and character-triple isomorphisms. By a character triple (G, N, θ) , we mean that *G* is a group, $N \triangleleft G$ and $\theta \in Irr(N)$ is *G*-invariant. For the definition and properties of character-triple isomorphisms, see Chapter 1 of [3] or Chapter 11 of [1].

Lemma 2.10 ([3, Lemma 3.11]). Let (G, N, θ) be a character triple, where G is a group and N is a π -subgroup. Then there exists a character triple (G^*, N^*, θ^*) such that $N^* \leq \mathbb{Z}(G^*)$ is also a π -subgroup and (G^*, N^*, θ^*) is isomorphic to (G, N, θ) .

The irreducible π -partial characters behave very well under the isomorphism of character triples.

Lemma 2.11 ([3, Lemmas 6.21 and 6.32]). Let (G, N, θ) and (G^*, N^*, θ^*) be isomorphic character triples, where G and G^* are π -separable and N and N^* are π -groups. Then

- (a) There exists a unique bijection $\varphi \mapsto \varphi^*$ from $I_{\pi}(G|\theta)$ onto $I_{\pi}(G^*|\theta^*)$ such that if χ is a lift of φ in $Irr(G|\theta)$ and $\chi^* \in Irr(G^*|\theta^*)$ corresponds to χ under the character-triple isomorphism, then χ^* is a lift of φ^* . In particular, we have $\varphi(1)_{\pi'} = \varphi^*(1)_{\pi'}$.
- (b) Let $Q \leq G$ and $Q^* \leq G^*$ be π' -groups, and assume that the intermediate subgroups NQand N^*Q^* correspond via the character-triple isomorphism. Then $\varphi \in I_{\pi}(G|Q,\theta)$ if and only if $\varphi^* \in I_{\pi}(G^*|Q^*,\theta^*)$. In particular, we have $|I_{\pi}(G|Q,\theta)| = |I_{\pi}(G^*|Q^*,\theta^*)|$.

2.4. Above the Glauberman-Isaacs correspondence. The following deep theorem was proved by Dade and Puig in the case where Q is solvable, which is thoroughly explained in [8] as mentioned at the end of page 2532 of [7]. If Q is not solvable, then it must have even order by the Feit-Thompson theorem, and thus K has odd order. In this case, Lewis [5] showed that the result also holds.

Lemma 2.12 (Dade-Puig-Lewis). Let G be an arbitrary group with $K \triangleleft G$ and $Q \leq G$, and assume that $KQ \triangleleft G$ and that (|K|, |Q|) = 1. Let $\theta \in Irr(K)$ be Q-invariant, and write $\theta^* \in Irr(\mathbf{C}_K(Q))$ for its Glauberman-Isaacs correspondent. Then (G_{θ}, K, θ) and $(\mathbf{N}_G(Q)_{\theta^*}, \mathbf{C}_K(Q), \theta^*)$ are isomorphic character triples.

2.5. Weights and vertices. One of essential ingredients in our proofs of Theorems A and B is Theorem 2.14 below, whose proof relies on the following lemma.

Lemma 2.13 ([3, Lemma 6.30]). Let $K \triangleleft G$, where G is π -separable and K is a π -subgroup, and suppose $Q \leq G$ is a π' -subgroup such that $KQ \triangleleft G$. Let $\theta \in Irr(K)$ be Q-invariant and $\varphi \in I_{\pi}(G|\theta)$. Then the following hold.

- (a) There exists a vertex P of φ such that $Q \leq P \leq \mathbf{N}_G(Q)$. In particular, Q is a vertex for φ if and only if $\varphi(1)_{\pi'} = |G:Q|_{\pi'}$.
- (b) Furthermore, any two of such vertices P for φ are $N_G(Q)$ -conjugate.

Theorem 2.14. Let $K \triangleleft G$, where G is π -separable and K is a π -subgroup, and suppose $Q \leq G$ is a π' -subgroup such that $KQ \triangleleft G$. For any Q-invariant character $\tau \in Irr(K)$, we have

$$|\mathbf{I}_{\pi}(G|Q,\tau)| = |\mathbf{I}_{\pi}(\mathbf{N}_G(Q)|Q,\tau^*)|$$

where τ^* is the image of τ under the Glauberman-Isaacs correspondence $\operatorname{Irr}_Q(K) \to \operatorname{Irr}(\mathbb{C}_K(Q))$.

Proof. By the Frattini argument, we have $G = KN_G(Q) = G_\tau N_G(Q)$, and by Lemma 2.6(b), we see that $|I_{\pi}(G|Q,\tau)| = |I_{\pi}(G_{\tau}|Q,\tau)|$. Also, since $N_G(Q)_{\tau^*} = N_G(Q)_{\tau}$, the same reason yields

$$|I_{\pi}(\mathbf{N}_{G}(Q)|Q,\tau^{*})| = |I_{\pi}(\mathbf{N}_{G}(Q)_{\tau}|Q,\tau^{*})| = |I_{\pi}(\mathbf{N}_{G_{\tau}}(Q)|Q,\tau^{*})|.$$

To complete the proof, therefore, we can assume without loss that τ is *G*-invariant. Then, by Lemma 2.12, the character triples (G, K, τ) and $(\mathbf{N}_G(Q), \mathbf{C}_K(Q), \tau^*)$ are isomorphic, and by Lemma 2.11(a), the map $\varphi \mapsto \varphi^*$ defines a bijection $I_{\pi}(G|\tau) \to I_{\pi}(\mathbf{N}_G(Q)|\tau^*)$ such that $\varphi(1)_{\pi'} = \varphi^*(1)_{\pi'}$. In this case, Lemma 2.13(a) tells us that Q is a vertex for φ if and only if

$$\varphi(1)_{\pi'} = |G:Q|_{\pi'} = |\mathbf{N}_G(Q):Q|_{\pi'},$$

and this is equivalent to saying that $\varphi^*(1)_{\pi'} = |\mathbf{N}_G(Q) : Q|_{\pi'}$. Thus $\varphi \in I_{\pi}(G|Q, \tau)$ if and only if $\varphi^* \in I_{\pi}(\mathbf{N}_G(Q)|Q, \tau^*)$, which implies that the above map $\varphi \mapsto \varphi^*$ is also a bijection from $I_{\pi}(G|Q, \tau)$ onto $I_{\pi}(\mathbf{N}_G(Q)|Q, \tau^*)$, and the result follows.

3. Proof of Theorems A and B

In this section, we will establish Theorems A and B simultaneously from the introduction. As usual, we need to prove the following "relative version", which clearly includes Theorems A and B (by taking Z = 1).

Theorem 3.1. Let G be a π -separable group, and let Q be a π '-subgroup of G. Suppose that Z is a normal π -subgroup of G and that $\lambda \in Irr(Z)$ is invariant in G. Then the following hold.

- (a) $|I_{\pi}(G|Q,\lambda)| \leq |I_{\pi}(\mathbf{N}_G(QZ)|Q,\lambda)|.$
- (b) Equality holds in (a) if Q has the property that whenever Q < X, where X is a π' -subgroup of G, then $Q < \mathbf{N}_X(Q)$.

Proof. We proceed by induction on |G:Z|.

Step 1. We may assume that $Z \leq \mathbf{Z}(G)$, so that $\mathbf{N}_G(QZ) = \mathbf{N}_G(Q)$.

By Lemma 2.10, we can choose a character triple (G^*, Z^*, λ^*) isomorphic to (G, Z, λ) , where $Z^* \leq \mathbb{Z}(G^*)$ is a π -subgroup. Since $G^*/Z^* \cong G/Z$ and G is π -separable, we see that G^* is also π -separable.

For each π' -subgroup X of G, we know that the intermediate subgroup XZ corresponds to $(XZ)^*$ via the isomorphism $XZ/Z \cong (XZ)^*/Z^*$, and since $XZ/Z \cong X$ is a π' -group and Z^* is central

in $(XZ)^*$, there exists a unique π' -subgroup X^* of G^* such that $(XZ)^* = X^* \times Z^*$. It follows that $X \mapsto X^*$ defines a surjective map from the set of π' -subgroups of G onto that of G^* .

For (a), we conclude by Lemma 2.11(b) that $|I_{\pi}(G|Q,\lambda)| = |I_{\pi}(G^*|Q^*,\lambda^*)|$ and

$$|\mathbf{I}_{\pi}(\mathbf{N}_{G}(QZ)|Q,\lambda)| = |\mathbf{I}_{\pi}(\mathbf{N}_{G^{*}}(Q^{*}Z^{*})|Q^{*},\lambda^{*})|.$$

So, it suffices to prove (a) in G^* .

For (b), if Q < X, then QZ < XZ, which implies that $Q^*Z^* = (QZ)^* < (XZ)^* = X^*Z^*$ and hence $Q^* < X^*$. Using the natural isomorphism $X \cong XZ/Z$, it is easy to see that $Q < \mathbf{N}_X(Q)$ if and only if $QZ/Z < \mathbf{N}_{XZ/Z}(QZ/Z)$, and this is equivalent to saying that

$$(QZ)^*/Z^* < \mathbf{N}_{(XZ)^*/Z^*}((QZ)^*/Z^*) = \mathbf{N}_{X^*Z^*/Z^*}(Q^*Z^*/Z^*),$$

which again happens if and only if $Q^* < \mathbf{N}_{X^*}(Q^*)$. So, we may also work in G^* to establish (b). To complete the proof, therefore, we may assume without loss that $Z \leq \mathbf{Z}(G)$.

Step 2. We may assume that $\mathbf{O}_{\pi'}(G) = 1$ *.*

Let $L = \mathbf{O}_{\pi'}(G)$ and write $\overline{G} = G/L$. Suppose first that $L \nleq Q$. Then Lemma 2.8(b) tells us that $I_{\pi}(G|Q,\lambda)$ is empty, and (a) follows. For (b), we need to prove that $I_{\pi}(N|Q,\lambda)$ is also empty. To do this, write $N = \mathbf{N}_G(Q)$, and since $L \cap N \leq \mathbf{O}_{\pi'}(N)$, it suffices to show that $L \cap N \nleq Q$. Note that Q < LQ, and by the assumption on Q, we have

$$Q < \mathbf{N}_{LQ}(Q) = \mathbf{N}_{L}(Q)Q = (L \cap N)Q$$

and thus $L \cap N \not\leq Q$. This proves the theorem in the case where $L \not\leq Q$.

Now suppose that $L \leq Q$. By Lemma 2.9, we can deduce that

$$|I_{\pi}(G|Q,\lambda)| = |I_{\pi}(\bar{G}|\bar{Q},\bar{\lambda})| \text{ and } |I_{\pi}(\mathbf{N}_{G}(Q)|Q,\lambda)| = |I_{\pi}(\mathbf{N}_{\bar{G}}(\bar{Q})|\bar{Q},\bar{\lambda})|.$$

So, it is no loss to work in $\overline{G} = G/L$, and thus we may assume that L = 1 (note that $O_{\pi'}(G/L) = 1$).

Step 3. We may assume that |G:K| < |G:Z|, where $K = \mathbf{O}_{\pi}(G)$.

Note that $Z \leq K$, so we need to show that Z < K. Suppose that Z = K, so that K is central in G by Step 1. Since $\mathbf{O}_{\pi'}(G) = 1$ by Step 2, the Hall-Higman Lemma 1.2.3 (see Theorem 3.21 of [2] for example) implies that $G = \mathbf{C}_G(K) \leq K$. It follows that G is a π -group, and in this case, the result is trivial. Thus we may assume that Z < K.

Step 4. Complete the proof.

Let \mathscr{A} be a set of representatives of $N_G(Q)$ -orbits in the set of Q-invariant characters in $Irr(K|\lambda)$. Since λ is G-invariant, it follows by Corollary 2.7 that

$$|I_{\pi}(G|Q,\lambda)| = \sum_{\substack{\tau \in \mathscr{A} \\ \gamma}} |I_{\pi}(G_{\tau}|Q,\tau)|.$$

Furthermore, let \mathscr{A}^* be the image of \mathscr{A} under the Glauberman-Isaacs correspondence from $\operatorname{Irr}_{\mathcal{Q}}(K)$ to $\operatorname{Irr}(\mathbf{C}_K(Q))$. Then \mathscr{A}^* is clearly a set of representatives of $\mathbf{N}_G(Q)$ -orbits in $\operatorname{Irr}(\mathbf{C}_K(Q)|\lambda)$. As before, we have

$$|\operatorname{I}_{\pi}(\mathbf{N}_{G}(Q)|Q,\lambda)| = \sum_{\tau \in \mathcal{A}} |\operatorname{I}_{\pi}(\mathbf{N}_{G}(Q)_{\tau}|Q,\tau^{*})|.$$

Note that $K N_G(Q)_{\tau} = N_{G_{\tau}}(KQ)$, and by Theorem 2.14, we have

$$|\mathbf{I}_{\pi}(\mathbf{N}_{G_{\tau}}(KQ)|Q,\tau)| = |\mathbf{I}_{\pi}(\mathbf{N}_{G}(Q)_{\tau}|Q,\tau^{*})|,$$

for all $\tau \in \mathscr{A}$. To complete the proof, therefore, it suffices to show that

$$|\operatorname{I}_{\pi}(G_{\tau}|Q,\tau)| \leq |\operatorname{I}_{\pi}(\operatorname{N}_{G_{\tau}}(KQ)|Q,\tau)|$$

with equality if Q has the property described in (b). This follows by the inductive hypothesis applied in G_{τ} since then $|G_{\tau}: K| < |G: Z|$ by Step 4, and the proof is complete.

The following is Corollary C in the introduction, which we restate here for convenience.

Corollary 3.2. Let G be a π -separable group, and let Q be a Hall π -complement of G. Then

- (a) $|I_{\pi}(G|Q)| = |I_{\pi}(N_G(Q)|Q)|$ and
- (b) $|X_{\pi}(G)| = |\operatorname{Irr}(\mathbf{N}_{G}(Q)/Q)|.$

Proof. (a) is clear by Theorem B. To prove (b), we use Lemmas 2.1 and 2.2 to deduce that

$$|X_{\pi}(G)| = |I_{\pi}(G|Q)|$$
 and $|X_{\pi}(N_{G}(Q))| = |I_{\pi}(N_{G}(Q)|Q)|$

and by (a), we obtain $|X_{\pi}(G)| = |X_{\pi}(N_G(Q))|$. By definition, it is easy to see that all normal π' subgroups of a π -separable group are contained in the kernel of each π -special character, and thus
we may identify $X_{\pi}(N_G(Q))$ with $X_{\pi}(N_G(Q)/Q)$. But $N_G(Q)/Q$ is a π -group, so all irreducible
characters of $N_G(Q)/Q$ are π -special, that is, $Irr(N_G(Q)/Q) = X_{\pi}(N_G(Q)/Q)$. It follows that $|X_{\pi}(G)| = |Irr(N_G(Q)/Q)|$, as desired.

Acknowledgements

This work was supported by the NSF of China (12171289) and by Fundamental Research Programs of Shanxi Province (20210302123429 and 20210302124077).

References

- 1. I.M. Isaacs, Character Theory of Finite Groups, Academic Press, New York, 1976.
- 2. I.M. Isaacs, Finite Group Theory, Amer. Math. Soc., Providence, RI, 2008.
- 3. I.M. Isaacs, Characters of Solvable Groups, Amer. Math. Soc., Providence, RI, 2018.
- 4. I.M. Isaacs and G. Navarro, Weights and vertices for characters of π-separable groups, J. Algebra 177 (1995) 339-366.
- 5. M. Lewis, Characters, coprime actions, and operator groups. Arch. Math. 69 (1997) 455-460.
- 6. G. Navarro, Vertices for characters of p-solvable groups, Trans. AMS. 365 (2002) 2759-2773.

- 7. G. Navarro and B. Sambale, Weights and nilpotent subgroups, Int. Math. Res. Not. 2021 (2021) 2526-2538.
- 8. A. Turull, Above the Glauberman correspondence, Adv. Math. 217 (2008) 2170-2205.
- 9. T. Wolf, Variations on McKay's character degree conjecture, J. Algebra 135 (1990) 123-138.

School of Mathematical Sciences, Shanxi University, Taiyuan, 030006, China Email address: changxuewu@sxu.edu.cn Email address: jinping@sxu.edu.cn