arXiv:2404.14685v1 [math.FA] 23 Apr 2024

HILBERT SPACE VALUED GAUSSIAN PROCESSES, THEIR KERNELS, FACTORIZATIONS, AND COVARIANCE STRUCTURE

PALLE E.T. JORGENSEN AND JAMES TIAN

ABSTRACT. Motivated by applications, we introduce a general and new framework for operator valued positive definite kernels. We further give applications both to operator theory and to stochastic processes. The first one yields several dilation constructions in operator theory, and the second to general classes of stochastic processes. For the latter, we apply our operator valued kernelresults in order to build new Hilbert space-valued Gaussian processes, and to analyze their structures of covariance configurations.

Contents

1. Introduction	1
2. Operator valued p.d. kernels	2
2.1. Examples	5
3. Schwartz' constructions	7
4. <i>H</i> -valued Gaussian processes	8
References	10

1. INTRODUCTION

Our present paper deals with two closely related issues, (i) a number of questions from operator theory, factorizations, and dilations; and from (ii) non-commutative stochastic analysis. The first one addresses questions around classes of Hilbert space dilations; and the second (ii) deals with a general framework for Hilbert space valued Gaussian processes. While, on the face of it, it is not obvious that the two questions are closely related, but as we show, (i) and (ii) together suggest the need for an extension of both the traditional framework of positive definite (scalar valued) kernels and their associated reproducing kernel Hilbert spaces (RKHSs); and of the framework for non-commutative stochastic analysis. Hence our present starting point will be a fixed separable Hilbert space H, and we then study the notion of positive definite kernels K which take values in the algebra of all bounded operators on H, here denoted B(H).

For the theory or reproducing kernels, the literature is extensive, both pure and recent applications, we refer to the following, including current, [AS56, Aro48, AJ21, PR16, AFMP94, LYA23, CY17].

²⁰⁰⁰ Mathematics Subject Classification. 46E22, 47A20, 47B32, 60G15.

Key words and phrases. Positive definite functions, kernels, Gaussian processes, covariance, dilation, POVMs.

Notations. Throughout the paper, we continue with the physics convention: inner products linear in the second variable.

For a p.d. kernel $K : S \times S \to \mathbb{C}$, we denote by H_K the corresponding reproducing kernel Hilbert space. It is the Hilbert completion of

$$span \{K_y(\cdot) := K(\cdot, y) \mid y \in S\}$$

$$(1.1)$$

with respect to the inner product

$$\left\langle \sum_{i} c_{i} K\left(\cdot, x_{i}\right), \sum_{i} d_{j} K\left(\cdot, y_{j}\right) \right\rangle_{H_{K}} := \sum_{i} \sum_{j} \overline{c}_{i} d_{j} K\left(x_{i}, x_{j}\right).$$
(1.2)

The reproducing property is as follows:

$$\varphi(x) = \langle K(\cdot, x), \varphi \rangle_{H_K}, \forall x \in S, \forall \varphi \in H_K.$$
(1.3)

For any orthonormal basis (ONB) (φ_i) in H_K , one has

$$K(x,y) = \sum_{i} \varphi_i(x) \overline{\varphi_i(y)}, \quad \forall x, y \in S.$$
(1.4)

A B(H)-valued kernel $K: S \times S \to B(H)$ is positive definite (p.d.) if

$$\sum_{i=1}^{n} \left\langle a_i, K\left(s_i, s_j\right) a_j \right\rangle_H \ge 0 \tag{1.5}$$

for all $(a_i)_1^n$ in H, and all $n \in \mathbb{N}$.

2. Operator valued p.d. kernels

In this section, we introduce the general framework for operator valued positive definite kernels. Our analysis is applied to several dilation constructions, both in operator theory, and in stochastic processes. The latter will be taken up in Section 4 again where we apply the results here to build Hilbert space valued Gaussian processes, and examine their properties.

Let S be a set, and let $K: S \times S \to B(H)$ be positive definite. Set $X = S \times H$, and define $\tilde{K}: X \times X \to \mathbb{C}$ by

$$\tilde{K}\left(\left(s,a\right),\left(t,b\right)\right) = \left\langle a,K\left(s,t\right)b\right\rangle_{H},\tag{2.1}$$

for all $a, b \in H$, and all $s, t \in S$. Then \tilde{K} is a scalar-valued p.d. kernel on $X \times X$. Let $H_{\tilde{K}}$ be the corresponding RKHS.

It will follow from our subsequent results that this $H_{\tilde{K}}$ realization has properties that makes it universal. Moreover, one notes that in different but related p.d. settings, analogous $H_{\tilde{K}}$ -constructions serve as a generators of dilation Hilbert spaces, and representations. It has counterparts for such related constructions as GNS (Gelfand-Naimark-Segal), and Stinespring (complete positivity), see e.g., [DM20, PTD23, Ver22].

Theorem 2.1. The family of operators $\{V_s\}_{s \in S}$, where $V_s : H \to H_{\tilde{K}}$, defined as

$$V_s a = \tilde{K}(\cdot, (s, a)) : X \to \mathbb{C}, \quad a \in H,$$

$$(2.2)$$

satisfies the following:

(1) For all $s \in S$, and all $a \in H$,

$$\left\|V_{s}a\right\|_{H_{\tilde{K}}}^{2} = \left\langle a, K\left(s, s\right)a\right\rangle_{H}.$$
(2.3)

(2) The adjoint $V_s^*: H_{\tilde{K}} \to H$ is determined by

$$V_s^* \tilde{K}(\cdot, (t, b)) = K(s, t) b.$$
(2.4)

(3) $V_s V_s^* : H_{\tilde{K}} \to H_{\tilde{K}}$ is given by

$$V_s V_s^* \tilde{K}(\cdot, (t, b)) = \tilde{K}(\cdot, (s, K(s, t) b)).$$

$$(2.5)$$

(4) For all s_1, s_2, \dots, s_n in S,

$$(V_{s_1}V_{s_1}^*) \cdots (V_{s_n}V_{s_n}^*) \tilde{K}(\cdot, (t, b)) = \tilde{K}(\cdot, (s_1, K(s_1, s_2) \cdots K(s_{n-1}, s_n) K(s_n, t) b)).$$
 (2.6)

(5) For all $s, s' \in S$, and $b \in H$,

$$V_{s'}V_s^*\tilde{K}\left(\cdot,(t,b)\right) = \tilde{K}\left(\cdot,(s',K\left(s,t\right)b\right)\right).$$
(2.7)

(6) $V_s^* V_t b = K(s,t) b$, and

$$\left(V_{s_1}^* V_{t_1}\right) \cdots \left(V_{s_n}^* V_{t_n}\right) b = K\left(s_1, t_1\right) \cdots K\left(s_{n-1}, t_{n-1}\right) K\left(s_n, t_n\right) b.$$
(2.8)

If, in addition,

$$K(s,s) = I, \quad \forall s \in S$$

then the operators $V_s: H \to H_{\tilde{K}}$ are isometric, and $V_s V_s^*$ are (selfadjoint) projections in $H_{\tilde{K}}$.

Proof. (1) By definition,

$$\left\|V_{s}a\right\|_{H_{\tilde{K}}}^{2} = \left\langle \tilde{K}\left(\cdot, (s, a)\right), \tilde{K}\left(\cdot, (s, a)\right) \right\rangle_{H_{\tilde{K}}} = \left\langle a, K\left(s, t\right)b \right\rangle_{H}.$$

(2) For all $a, b \in H$, and all $s, t \in S$, since

$$\left\langle V_{s}a, \tilde{K}\left(\cdot, (t, b)\right)\right\rangle_{H_{\tilde{K}}} = \left\langle \tilde{K}\left(\cdot, (s, a)\right), \tilde{K}\left(\cdot, (t, b)\right)\right\rangle_{H_{\tilde{K}}} = \left\langle a, K\left(s, t\right)b\right\rangle_{H_{\tilde{K}}}$$

- it follows that $V_{s}^{*}\tilde{K}(\cdot,(t,b)) = K(s,t)b$, as stated.
 - (3) A direct calculation shows that

$$V_{s}V_{s}^{*}\tilde{K}\left(\cdot,\left(t,b\right)\right)=V_{s}K\left(s,t\right)b=\tilde{K}\left(\cdot,\left(s,K\left(s,t\right)b\right)\right).$$

(4)-(6) Immediate.

The above leads to a characterization of operator-valued p.d. kernels.

Corollary 2.2. Let $K : S \times S \to B(H)$. Then K is p.d. if and only if there exists a family of operators $\{V_s\}_{s \in S}$ from H into another Hilbert space L, such that K factors as

$$K(s,t) = V_s^* V_t, \quad s,t \in S.$$

$$(2.9)$$

If K(s,s) = I for all $s \in S$, then $V_s : H \to L$ are isometric. Furthermore, if $L = \overline{span} \{V_s a : s \in S, a \in H\}$, then $L \simeq H_{\tilde{K}}$.

Proof. The "only if" part is Theorem 2.1 (6).

3

Conversely, assume $K(s,t) = V_s^* V_t$ as stated. For all $(a_i)_{i=1}^n$ in H, and $(s_i)_{i=1}^n$ in S,

$$\sum_{i,j} \langle a_i, K(s_i, s_j) a_j \rangle_H = \sum_{i,j} \langle a_i, V_{s_i}^* V_{s_j} a_j \rangle_H$$
$$= \sum_{i,j} \langle V_{s_i} a_i, V_{s_j} a_j \rangle_L = \left\| \sum_i V_{s_i} a_i \right\|_L^2 \ge 0$$

which shows that K is positive definite.

Under the assumption $L = \overline{span} \{ V_s a : s \in S, a \in H \}$, the map

$$\tilde{K}(\cdot, (s, a)) \longmapsto V_s a$$

extends to a unique isometric isomorphism from $H_{\tilde{K}}$ onto L, and so any such space L is isomorphic to the RKHS $H_{\tilde{K}}$.

Corollary 2.3. Any function $F \in H_{\tilde{K}}$ satisfies that

$$F(s,a) = \langle a, V_s^* F \rangle_{H_{\tilde{K}}}, \quad (s,a) \in S \times H.$$
(2.10)

In particular, F is conjugate liner in the second variable, and

$$F(s,0) = 0, \quad \forall s \in S.$$

$$(2.11)$$

Proof. This follows from the reproducing property (see (1.3)) of $H_{\tilde{K}}$:

$$F(s,a) = \left\langle \tilde{K}(\cdot, (s,a)), F \right\rangle_{H_{\tilde{K}}} = \left\langle V_s a, F \right\rangle_{H_{\tilde{K}}} = \left\langle a, V_s^* F \right\rangle_H.$$

Corollary 2.4. For any ONB $(\varphi_i)_{i \in \mathbb{N}}$ in $H_{\tilde{K}}$, the following operator identity holds:

$$K(s,t) = \sum_{i \in \mathbb{N}} |V_s^* \varphi_i\rangle \langle V_t^* \varphi_i|.$$
(2.12)

Here $|a\rangle\langle b|$ is Dirac's notation for rank-1 operators, i.e., $c \mapsto a\langle b, c \rangle$.

Proof. Using the identity

$$I_{H_{\tilde{K}}} = \sum_{i} |\varphi_{i}\rangle \langle \varphi_{i}|$$

in $H_{\tilde{K}}$, it follows that

$$K(s,t) = V_s^* V_t = V_s^* I_{H_{\tilde{K}}} V_t = \sum_i |V_s^* \varphi_i\rangle \langle V_t^* \varphi_i|.$$

Equivalently,

$$\tilde{K}((s,a),(t,b)) = \sum_{i \in \mathbb{N}} \varphi(s,a) \overline{\varphi(t,b)} \\
= \sum_{i \in \mathbb{N}} \langle V_s a, \varphi_i \rangle \langle \varphi_i, V_t b \rangle \\
= \sum_{i \in \mathbb{N}} \langle a, V_s^* \varphi_i \rangle \langle V_t^* \varphi_i, b \rangle.$$

2.1. **Examples.** The current setting of operator-valued p.d. kernels contains a wide range of well-known constructions. A classical example is when

$$K(s,t) = \rho\left(s^{-1}t\right), \quad s,t \in G$$

where $\rho: G \to B(H)$ is a representation of the group G. Note that, in this case, $K(s,s) = \rho(e) = I_H$, the identity operator on H, for all $s \in G$.

Example 2.5. As an application, we reformulate the well-known power dilation of a single contraction $A \in B(H)$. Here, it is assumed that $||A|| \leq 1$. Consider

$$A^{(n)} := \begin{cases} A^n & n \ge 0\\ (A^*)^n & n < 0 \end{cases}$$

and set $K : \mathbb{Z} \times \mathbb{Z} \to B(H)$ by

$$K(m,n) = A^{(n-m)}.$$

This K is positive definite, i.e., $\langle h, Th \rangle \ge 0$, for all $h = (h_1, \dots, h_n)$ with $h_i \in H$ (see (1.5)); or, equivalently, the matrix below is p.d.:

$$T = \begin{bmatrix} 1 & A & A^2 & \cdots \\ A^* & 1 & A & \ddots \\ A^{*2} & A^* & 1 & \ddots \\ \vdots & \ddots & \ddots & \ddots \end{bmatrix}_{n \times n}$$
(2.13)

(See Lemma 2.6 for details.)

Then, let $X = \mathbb{Z} \times H$ and define $\tilde{K} : X \times X \to \mathbb{C}$ by

$$\tilde{K}\left(\left(m,h\right),\left(n,k\right)\right) = \left\langle h,K\left(m,n\right)k\right\rangle_{H} = \left\langle h,A^{\left(n-m\right)}k\right\rangle_{H}.$$

Denote by $H_{\tilde{K}}$ the RKHS of \tilde{K} . Set $V: H \to H_{\tilde{K}}$ by

$$Vh = \tilde{K}(\cdot, (0, h)).$$

This is an isometric embedding. Set $U: H_{\tilde{K}} \to H_{\tilde{K}}$ by

$$U\tilde{K}(\cdot, (m,h)) = \tilde{K}(\cdot, (m+1,h)).$$

Note, U is unitary. It follows from Theorem 2.1 that

$$V^*U^nVh = V^*U^n\tilde{K}\left(\cdot, (0, h)\right) = V^*\tilde{K}\left(\cdot, (n, h)\right) = A^nh,$$

that is,

$$A^n = V^* U^n V.$$

Lemma 2.6. The operator matrix (2.13) is positive definite.

Proof. An elementary approach to verify this is to rewrite T as

$$T = \begin{bmatrix} 1 & A & A^{2} & \cdots \\ A^{*} & 0 & 0 & \ddots \\ A^{*2} & 0 & 0 & \ddots \\ \vdots & \ddots & \ddots & \ddots \end{bmatrix} + \begin{bmatrix} 0 & 0 & 0 & \cdots \\ 0 & 1 & A & \cdots \\ 0 & A^{*} & 1 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$
$$= \begin{bmatrix} 1 \\ A^{*} \\ \vdots \\ A^{*n} \end{bmatrix} \begin{bmatrix} 1 & A & \cdots & A^{n} \end{bmatrix} - \begin{bmatrix} 0 \\ A^{*} \\ \vdots \\ A^{*n} \end{bmatrix} \begin{bmatrix} 0 & A & \cdots & A^{n} \end{bmatrix} + \begin{bmatrix} 0 & 0 & 0 & \cdots \\ 0 & 1 & A & \cdots \\ 0 & A^{*} & 1 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

Therefore,

$$\langle h, Th \rangle = \|h_1 + Ah_2 + \dots + A^n h_n\|^2 - \|Ah_2 + \dots + A^n h_n\|^2 + \|h_2 + Ah_3 + \dots + A^{n-1}h_n\|^2 - \|Ah_3 + \dots + A^{n-1}h_n\|^2 + \|h_3 + Ah_4 + \dots + A^{n-2}h_n\|^2 - \|Ah_4 + \dots + A^{n-2}h_n\|^2 \vdots + \|h_{n-1} + Ah_n\|^2 - \|Ah_n\|^2 + \|h_n\|^2 \ge \|h_1 + Ah_2 + \dots + A^n h_n\|^2 + (1 - \|A\|^2) \left[\|h_2 + Ah_3 + \dots + A^{n-1}h_n\|^2 + \dots + \|h_n\|^2 \right] \ge 0.$$

Another notable example that falls in the general setting of Theorem 2.1 is the dilation of positive operator valued measures (POVMs), namely, any POVM $Q: \mathscr{B}_S \to B(H)$ is the compression of a projection valued measure (PVM) P: $\mathscr{B}_S \to B(L)$ for some Hilbert space $L \supset H$, where \mathscr{B}_S denotes the Borel σ -algebra of S. See, e.g., [FLZG24, Hol22, RMC21, HCW21].

Example 2.7. Given $Q : \mathscr{B}_S \to B(H)$ as above, one may define $K : \mathscr{B}_S \times \mathscr{B}_S \to B(H)$ by

$$K(A,B) = Q(A \cap B), \quad \forall A, B \in \mathscr{B}_S,$$

which is positive definite. Then set

$$\tilde{K} : (\mathscr{B}_S, H) \times (\mathscr{B}_S, H) \to \mathbb{C},$$
$$\tilde{K} ((A, a), (B, b)) = \langle a, Q (A \cap B) b \rangle_H$$

for all $A, B \in \mathscr{B}_S$, and all $a, b \in H$.

Let $V: H \to H_{\tilde{K}}$ be the isometric embedding,

$$Va = \tilde{K}\left(\cdot, (S, a)\right)$$

where $H_{\tilde{K}}$ denotes the RKHS of $\tilde{K}.$ It follows that

$$V^*K\left(\cdot, (A, a)\right) = Q\left(A\right)a$$

$$P(A)\tilde{K}(\cdot, (B, a)) = \tilde{K}(\cdot, (A \cap B, a)).$$

Then $P(A)^* = P(A) = P(A)^2$, i.e., a selfadjoint projection in $H_{\tilde{K}}$, for all $A \in \mathscr{B}_S$. Finally,

$$\begin{split} V^*P\left(A\right)Va &= V^*P\left(A\right)\tilde{K}\left(\cdot,(S,a)\right) \\ &= V^*\tilde{K}\left(\cdot,(A\cap S,a)\right) = V^*\tilde{K}\left(\cdot,(A,a)\right) = Q\left(A\right)a. \end{split}$$

That is, $Q = V^* P V$, where P is a PVM acting in $H_{\tilde{K}}$.

3. Schwartz' constructions

Motivated by applications, in the present section we extend the framework for operator valued positive definite kernels to spaces more general than Hilbert spaces.

We explore an approach to operator theory that extends beyond the usual settings, moving into the realm of topological vector spaces L and their duals L^* . This idea was introduced to the first named author by Professor D. Alpay, who may have been influenced by L. Schwartz and others.

Here, we look at positive definite (p.d.) kernels K defined on $S \times S$ with values in continuous linear operators from L to L^* . This setup includes, but is not limited to, our previously discussed B(H)-valued kernels. By extending to these more general vector spaces, such as Banach spaces, we aim to broaden the application of these kernels. We will show how our earlier models using K can be adapted to this extended context, particularly by transforming these operator-valued kernels into scalar-valued kernels \tilde{K} on the product space $X = S \times L$. This transition illustrates the versatility and potential of our approach in various mathematical and practical scenarios.

Let S be a set, L a topological vector space, and L^* its *conjugate* dual space. We consider a positive definite (p.d.) kernel $K : S \times S \to \mathcal{L}(L, L^*)$, where $\mathcal{L}(L, L^*)$ denotes the space of continuous linear operators from L to L^* .

Specifically, the p.d. assumption on K is that

$$\sum_{i,j=1}^{n} \left\langle a_i, K\left(s_i, s_j\right) a_j \right\rangle \ge 0$$

for all $(s_i)_{i=1}^n$ in S, $(a_i)_{i=1}^n$ in L, and all $n \in \mathbb{N}$. Here, $\langle \cdot, \cdot \rangle : L \times L^* \to \mathbb{C}$ is the canonical dual pairing, i.e., $\langle \ell, \ell' \rangle := \ell'(\ell)$, for all $\ell \in L$, and $\ell' \in L^*$.

Define $X = S \times L$, and

$$K\left(\left(s,a\right),\left(t,b\right)\right) := \left\langle a,K\left(s,t\right)b\right\rangle.$$

Then \tilde{K} is a scalar-valued p.d. kernel on $X \times X$; indeed, we have

$$\sum_{i,j=1}^{n} \overline{c_i} c_j \tilde{K}\left(\left(s_i, a_i\right), \left(s_j, a_j\right)\right) = \sum_{i,j=1}^{n} \overline{c_i} c_j \left\langle a_i, K\left(s_i, s_j\right) a_j \right\rangle$$
$$= \sum_{i,j=1}^{n} \left\langle c_i a_i, K\left(s_i, s_j\right) c_j a_j \right\rangle \ge 0.$$

Let $H_{\tilde{K}}$ be the corresponding RKHS.

Then we get the following variant of Theorem 2.1:

Theorem 3.1. The family of operators $\{V_s\}_{s \in S}$, where $V_s : L \to H_{\tilde{K}}$, defined as

$$V_s a = \tilde{K}(\cdot, (s, a)) : X = S \times L \to \mathbb{C}, \quad a \in L,$$
(3.1)

satisfies the following:

(1) For all $s \in S$, and all $a \in L$,

$$\left\|V_{s}a\right\|_{H_{\tilde{K}}}^{2} = \left\langle a, K\left(s, s\right)a\right\rangle.$$

$$(3.2)$$

(2) The adjoint $V_s^*: H_{\tilde{K}} \to L^*$ is determined by

$$V_s^* \tilde{K}(\cdot, (t, b)) = K(s, t) b.$$
(3.3)

- (3) $V_s V_s^* : H_{\tilde{K}} \to H_{\tilde{K}} \text{ is given by}$ $V_s V_s^* \tilde{K} (\cdot, (t, b)) = \tilde{K} (\cdot, (s, K(s, t) b)).$ (3.4)
- (4) For all s_1, s_2, \dots, s_n in S,

$$(V_{s_1}V_{s_1}^*) \cdots (V_{s_n}V_{s_n}^*) \tilde{K}(\cdot, (t, b)) = \tilde{K}(\cdot, (s_1, K(s_1, s_2) \cdots K(s_{n-1}, s_n) K(s_n, t) b)).$$
(3.5)

(5) For all $s, s' \in S$, and $b \in L$,

$$V_{s'}V_s^*\tilde{K}\left(\cdot,(t,b)\right) = \tilde{K}\left(\cdot,(s',K\left(s,t\right)b\right)\right).$$
(3.6)

(6)
$$V_s^* V_t b = K(s,t) b$$
, and
 $\left(V_{s_1}^* V_{t_1}\right) \cdots \left(V_{s_n}^* V_{t_n}\right) b = K(s_1,t_1) \cdots K(s_{n-1},t_{n-1}) K(s_n,t_n) b.$ (3.7)

4. H-VALUED GAUSSIAN PROCESSES

Since the early developments of Ito, the stochastic analysis of Gaussian processes has been grounded in the use of Hilbert space and related operators, see e.g., [AJ15, AJ21, Sch52] and the papers, and early literature, cited there. Here we want to call attention to the following distinction: (i) the case of scalar valued Gaussian processes which may be indexed by a Hilbert space, often via an Itoisometry construction; and (ii) the case when the Gaussian processes itself takes values in Hilbert space, see e.g., [Lop23, Gue22, KM20, FS20, Kat19]. The latter is our present focus. And the setting of (ii) has the advantage of allowing a much more flexible encoding of more covariance structures arising in big data-sets. We now turn to the construction. It is based on the results in Section 2 above, and it makes use of both B(H)-valued positive definite kernels, and their induced scalar valued dilations, see Theorem 2.1 and Corollary 2.2 above.

Fix $S \times S \xrightarrow{K} B(H)$, and introduce $\tilde{K} : (S \times H) \times (S \times H) \to \mathbb{C}$, given by

$$\tilde{K}\left(\left(s,a\right),\left(t,b\right)\right) = \left\langle a,K\left(s,t\right)b\right\rangle_{H}, \quad \forall s,t\in S, \; \forall a,b\in H,$$

$$(4.1)$$

defined on $S \times H$. Let $H_{\tilde{K}}$ be the corresponding RKHS (consisting of scalar-valued functions on $S \times H$).

Lemma 4.1. For all $\varphi \in H_{\tilde{K}}$, $s \in S$, and $a, b \in H$, we have

$$\varphi(s, a+b) = \varphi(s, a) + \varphi(s, b).$$
(4.2)

(See also Corollary 2.3.)

Proof. It suffices to verify (4.2) for the generating functions $\varphi(t, b) = \tilde{K}(\cdot, (t, b))$ in $H_{\tilde{K}}$, which are functions defined on $S \times H$. By definition,

$$\varphi(t, b_1 + b_2)(s, a) = \left\langle \tilde{K}(s, a), \tilde{K}(t, b_1 + b_2) \right\rangle_{H_{\tilde{K}}}$$

$$= \left\langle a, K(s, t) (b_1 + b_2) \right\rangle_{H}$$

$$= \left\langle a, K(s, t) b_1 \right\rangle_{H} + \left\langle a, K(s, t) b_2 \right\rangle_{H}$$
since $K(s, t) \in B(H)$

$$= \varphi(t, b_1)(s, a) + \varphi(t, b_2)(s, a).$$

Corollary 4.2. For any ONB $\{e_i\}$ in H, we have the following identity for functions $\varphi \in H_{\tilde{K}}$:

$$\sum_{i} \varphi(s, e_{i}) \langle e_{i}, a \rangle_{H} = \varphi(s, a), \quad \forall (s, a) \in S \times H.$$
(4.3)

Proof. Since $K(s,t) \in B(H)$ for all $(s,t) \in S \times S$, (4.3) follows from the ONB representation in H, i.e.,

$$\sum_{i} e_i \langle e_i, a \rangle_H = a, \quad \forall a \in H,$$
(4.4)

and

$$\sum |\langle e_i, a \rangle_H|^2 = ||a||_H^2.$$
(4.5)

Similarly,

$$\sum_{i} \langle a, e_i \rangle_H \langle e_i, b \rangle_H = \langle a, b \rangle_H, \quad \forall a, b \in H.$$
(4.6)

Let K, \tilde{K} , S, H be as above, and let $\{Z_i\}$ be i.i.d. N(0,1) random variables, realized on a probability space (Ω, \mathbb{P}) .

Introduce the operators $(V_s)_{s \in S}$ from Theorem 2.1. Recall that

$$V_s^* V_t = K(s,t), \quad \forall s, t \in S.$$

$$(4.7)$$

The argument for the identity (4.7) relies on the pair of operators, where

$$V_t a = \tilde{K}_{(t,a)}\left(\cdot\right) := \tilde{K}\left(\cdot, (t,a)\right)$$

and

$$V_s^*\left(\tilde{K}_{(t,a)}\right) = K\left(s,t\right)a.$$

$$H_{\tilde{K}}$$

$$V_t \checkmark V_s^*$$

$$H \xrightarrow{V_s^* V_t} H$$

We verify that $V_{s}^{*}V_{t} = K(s,t)$, or equivalently,

$$\langle V_{s}a, V_{t}b \rangle_{H_{\tilde{K}}} = \langle a, K(s, t) t \rangle_{H}$$

for all $s, t \in S$, and all $a, b \in H$.

Definition 4.3. Consider the *H*-valued Gaussian process $W_s : \Omega \to H$,

$$W_t = \sum_i \left(V_t^* \varphi_i \right) Z_i, \tag{4.8}$$

where (φ_i) is an ONB in $H_{\tilde{K}}$.

Theorem 4.4. We have

$$\mathbb{E}\left[\left\langle a, W_s \right\rangle_H \left\langle W_t, b \right\rangle_H\right] = \left\langle a, K\left(s, t\right) b \right\rangle_H.$$
(4.9)

Proof. Note that $\mathbb{E}[Z_i Z_j] = \delta_{i,j}$. From this, we get

LHS_(4.9) =
$$\mathbb{E} [\langle a, W_s \rangle_H \langle W_t, b \rangle_H]$$

= $\sum_i \sum_j \langle a, V_s^* \varphi_i \rangle \langle V_t^* \varphi_j, b \rangle \mathbb{E} [Z_i Z_j]$
= $\sum_i \langle V_s a, \varphi_i \rangle_{H_{\tilde{K}}} \langle \varphi_i, V_t b \rangle_{H_{\tilde{K}}}$
= $\langle V_s a, V_t b \rangle_{H_{\tilde{K}}} = \langle a, V_s^* V_t b \rangle_H = \langle a, K(s, t) b \rangle_H.$

Remark 4.5. The fact that $\{W_t\}_{t\in S}$ in (4.8) is well defined, and a Gaussian process with the stated properties, is an application of the central limit theorem to the choice $\{Z_i\}$ of i.i.d. N(0,1) Gaussians on the right-side of (4.8).

Different choices of ONBs (φ_i) and i.i.d. N(0,1) Gaussian (Z_i) would yield different Gaussian processes $\{W_t\}_{t\in S}$, but they will all satisfy the covariance condition (4.9). Also recall that if (φ_i) , (Z_i) are fixed, then the corresponding Gaussian process $\{W_t\}_{t\in S}$ is unique, since Gaussian processes are determined by the first two moments.

References

- [AB97] D. Alpay and V. Bolotnikov, On tangential interpolation in reproducing kernel Hilbert modules and applications, Topics in interpolation theory (Leipzig, 1994), Oper. Theory Adv. Appl., vol. 95, Birkhäuser, Basel, 1997, pp. 37–68. MR 1473250
- [AD93] Daniel Alpay and Harry Dym, On a new class of structured reproducing kernel spaces, J. Funct. Anal. 111 (1993), no. 1, 1–28. MR 1200633
- [AD06] D. Alpay and C. Dubi, Some remarks on the smoothing problem in a reproducing kernel Hilbert space, J. Anal. Appl. 4 (2006), no. 2, 119–132. MR 2223568
- [AFMP94] Gregory T. Adams, John Froelich, Paul J. McGuire, and Vern I. Paulsen, Analytic reproducing kernels and factorization, Indiana Univ. Math. J. 43 (1994), no. 3, 839– 856. MR 1305949
- [AJ15] Daniel Alpay and Palle Jorgensen, Spectral theory for Gaussian processes: reproducing kernels, boundaries, and L²-wavelet generators with fractional scales, Numer. Funct. Anal. Optim. 36 (2015), no. 10, 1239–1285. MR 3402823
- [AJ21] Daniel Alpay and Palle E. T. Jorgensen, New characterizations of reproducing kernel Hilbert spaces and applications to metric geometry, Opuscula Math. 41 (2021), no. 3, 283–300. MR 4302453
- [AL95] Daniel Alpay and Juliette Leblond, Traces of Hardy functions and reproducing kernel Hilbert spaces, Arch. Math. (Basel) 64 (1995), no. 6, 490–499. MR 1329822
- [Alp91] Daniel Alpay, Some reproducing kernel spaces of continuous functions, J. Math. Anal. Appl. 160 (1991), no. 2, 424–433. MR 1126127
- [Aro48] N. Aronszajn, Reproducing and pseudo-reproducing kernels and their application to the partial differential equations of physics, Harvard University, Graduate School of Engineering, Cambridge, MA, 1948, Studies in partial differential equations. Technical report 5, preliminary note. MR 31663

- [AS56] N. Aronszajn and K. T. Smith, Functional spaces and functional completion, Ann. Inst. Fourier (Grenoble) 6 (1955/56), 125–185. MR 80878
- [CLPL19] Wen-Sheng Chen, Jingmin Liu, Binbin Pan, and Yugao Li, Block kernel nonnegative matrix factorization for face recognition, Int. J. Wavelets Multiresolut. Inf. Process. 17 (2019), no. 1, 1850059, 28. MR 3911880
- [CY17] Rupak Chatterjee and Ting Yu, Generalized coherent states, reproducing kernels, and quantum support vector machines, Quantum Inf. Comput. 17 (2017), no. 15-16, 1292– 1306. MR 3700848
- [DM20] Nicolò Drago and Valter Moretti, The notion of observable and the moment problem for *-algebras and their GNS representations, Lett. Math. Phys. 110 (2020), no. 7, 1711–1758. MR 4114386
- [FLZG24] Lingxuan Feng, Shunlong Luo, Yan Zhao, and Zhihua Guo, Equivorlapping measurements as extensions of symmetric informationally complete positive operator valued measures, Phys. Rev. A 109 (2024), no. 1, Paper No. 012218, 10. MR 4705975
- [FS20] Martin Forde and Benjamin Smith, The conditional law of the Bacry-Muzy and Riemann-Liouville log correlated Gaussian fields and their GMC, via Gaussian Hilbert and fractional Sobolev spaces, Statist. Probab. Lett. 161 (2020), 108732, 11. MR 4073554
- [Gue22] J. C. Guella, On Gaussian kernels on Hilbert spaces and kernels on hyperbolic spaces, J. Approx. Theory 279 (2022), Paper No. 105765, 36. MR 4414825
- [HCW21] Shan Huang, Zeng-Bing Chen, and Shengjun Wu, Entropic uncertainty relations for general symmetric informationally complete positive operator-valued measures and mutually unbiased measurements, Phys. Rev. A 103 (2021), no. 4, Paper No. 042205, 9. MR 4256782
- [Hol22] A. S. Holevo, On optimization problem for positive operator-valued measures, Lobachevskii J. Math. 43 (2022), no. 7, 1646–1650. MR 4509531
- [JYZ21] Hao Jiang, Ming Yi, and Shihua Zhang, A kernel non-negative matrix factorization framework for single cell clustering, Appl. Math. Model. 90 (2021), 875–888. MR 4160499
- [Kat19] Victor Katsnelson, On the completeness of Gaussians in a Hilbert functional space, Complex Anal. Oper. Theory 13 (2019), no. 3, 637–658. MR 3940383
- [KM20] Vít Kubelka and Bohdan Maslowski, Filtering of Gaussian processes in Hilbert spaces, Stoch. Dyn. 20 (2020), no. 3, 2050020, 22. MR 4101087
- [Lop23] Oleh Lopushansky, Bernstein-Jackson inequalities on Gaussian Hilbert spaces, J.
 Fourier Anal. Appl. 29 (2023), no. 5, Paper No. 58, 21. MR 4641110
- [LYA23] Binglin Li, Changwon Yoon, and Jeongyoun Ahn, Reproducing kernels and new approaches in compositional data analysis, J. Mach. Learn. Res. 24 (2023), Paper No. [327], 34. MR 4690276
- [PPPR20] Satish K. Pandey, Vern I. Paulsen, Jitendra Prakash, and Mizanur Rahaman, Entanglement breaking rank and the existence of SIC POVMs, J. Math. Phys. 61 (2020), no. 4, 042203, 14. MR 4085703
- [PR16] Vern I. Paulsen and Mrinal Raghupathi, An introduction to the theory of reproducing kernel Hilbert spaces, Cambridge Studies in Advanced Mathematics, vol. 152, Cambridge University Press, Cambridge, 2016. MR 3526117
- [PTD23] J. E. Pascoe and Ryan Tully-Doyle, Induced Stinespring factorization and the Wittstock support theorem, Results Math. 78 (2023), no. 4, Paper No. 135, 15. MR 4581177
- [PTG15] Andrés Esteban Páez-Torres and Fabio A. González, Online kernel matrix factorization, Progress in pattern recognition, image analysis, computer vision, and applications, Lecture Notes in Comput. Sci., vol. 9423, Springer, Cham, 2015, pp. 651–658. MR 3487535
- [QL17] Likuan Qin and Vadim Linetsky, Long-term factorization of affine pricing kernels, Math. Financ. Econ. 11 (2017), no. 4, 479–498. MR 3709384
- [RMC21] Benjamin Robinson, Bill Moran, and Doug Cochran, Positive operator-valued measures and densely defined operator-valued frames, Rocky Mountain J. Math. 51 (2021), no. 1, 265–272. MR 4280112
- [Sch52] Laurent Schwartz, Théorie des noyaux, Proceedings of the International Congress of Mathematicians, Cambridge, Mass., 1950, vol. 1, Amer. Math. Soc., Providence, RI, 1952, pp. 220–230. MR 45307

- [Sch64] _____, Sous-espaces hilbertiens d'espaces vectoriels topologiques et noyaux associés (noyaux reproduisants), J. Analyse Math. 13 (1964), 115–256. MR 179587
- [Shi05] Serguei Shimorin, Commutant lifting and factorization of reproducing kernels, J. Funct. Anal. 224 (2005), no. 1, 134–159. MR 2139107
- [Ver22] Dominic Verdon, A covariant Stinespring theorem, J. Math. Phys. 63 (2022), no. 9, Paper No. 091705, 49. MR 4482713

(Palle E.T. Jorgensen) Department of Mathematics, The University of Iowa, Iowa City, IA 52242-1419, U.S.A.

Email address: palle-jorgensen@uiowa.edu

(James F. Tian) Mathematical Reviews, 416 4th Street Ann Arbor, MI 48103-4816, U.S.A.

Email address: jft@ams.org