

TIGHT CONTACT STRUCTURES ON HYPERBOLIC HOMOLOGY 3-SPHERES

MAHAN MJ, BALARKA SEN

ABSTRACT. We produce a large class of hyperbolic homology 3-spheres admitting arbitrarily many distinct tight contact structures. We also produce a sub-class admitting arbitrarily many distinct tight contact structures within the same homotopy class of oriented plane distributions. We also introduce a notion of geometric limits of contact structures compatible with geometric limits of hyperbolic manifolds. We study the behavior of the tight contact structures we construct under geometric limits.

CONTENTS

1. Introduction	1
2. Construction of hyperbolic pseudo-Anosov braids	4
2.1. Braid closures	4
2.2. An application of Thurston's hyperbolization	5
2.3. Genus one surfaces	7
2.4. Increasing genus	9
3. Proof of Theorem 1.2	11
3.1. Construction of the manifolds	11
3.2. Construction of the contact structures	12
3.3. Distinguishing contact structures	14
3.4. Homotopy-indistinguishability of plane fields	15
4. Further examples: rational surgery on links	16
5. Geometric limit of contact structures	18
5.1. Preliminary definitions	18
5.2. Basic slice and stabilization	19
5.3. Geometric convergence of rational surgeries	22
References	26

1. INTRODUCTION

A contact structure on a 3-manifold M is a rank-2 distribution $\xi \subset TM$ such that ξ is nowhere integrable. Contact structures on 3-manifolds come in two distinct flavors: tight and overtwisted. A result due to Eliashberg [Eli89] states that overtwisted contact structures

2010 *Mathematics Subject Classification.* 57R17, 57M50, 53D35.

Key words and phrases. hyperbolic 3-manifold, tight contact structure, geometric limit.

Both authors are supported in part by the Department of Atomic Energy, Government of India, under project no.12-R&D-TFR-5.01-0500, and by an endowment of the Infosys Foundation. MM is supported in part by the Fields Institute, Toronto during the special semester on "Randomness and Geometry", and by a DST JC Bose Fellowship.

are completely classified by the homotopy class of their underlying rank-2 distributions. Consequently, all oriented closed 3-manifolds admit overtwisted contact structures. In [Eli92, 8.2.], Eliashberg asked: which 3-manifolds admit tight contact structures? Etnyre and Honda [EH01] have demonstrated that the reducible manifold $P \# \bar{P}$ does not admit a tight contact structure, where P denotes the Poincaré homology sphere. Nevertheless, the following question remains open:

Question 1.1 (Question 8.2.2. in [Eli92] and Problem 4.142 in [kir97]). Let M be an irreducible 3-manifold. Does it admit a tight contact structure? Does it admit a fillable contact structure?

The problem of existence and classification of tight contact structures on lens spaces [Hon00a] and Seifert fibered spaces [Gom98, Hon00b, LS09] is completely settled. By the work of Agol [Ago13, Theorem 9.2] and Wise [Wis21], every closed hyperbolic 3-manifold admits a finite-sheeted cover which is fibered over the circle. Combining this with the work of Eliashberg-Thurston [ET98] on confoliations, one deduces that every closed hyperbolic 3-manifold admits a finite-sheeted cover possessing a tight contact structure.

However, not much is known beyond this with regard to Question 1.1 for the class of hyperbolic 3-manifolds. In this direction, there is a complete classification of tight contact structures for surgeries on the figure-eight knot, due to Conway-Min [CM20]. Moreover, there are examples of hyperbolic 3-manifolds which do not admit fillable contact structures, due to Li-Liu [LL19]. For more examples of tight contact structures on hyperbolic 3-manifolds, see [HKM03], [AS17], [Etg12].

Gabai [Gab83] observed that any oriented closed irreducible 3-manifold M with $b_1(M) > 0$ admits a taut foliation. Using the theory of confoliations, one can therefore deduce that such manifolds also admit tight contact structures. Thus, the attention can be narrowed down to hyperbolic rational homology 3-spheres. The main theorem of this article provides examples of tight contact structures on a large class of integer homology 3-spheres.

Theorem 1.2. *There exists an infinite family of sequences $\{M_{i,1}, M_{i,2}, M_{i,3}, \dots\}$ of hyperbolic integer homology 3-spheres, such that*

- (1) $M_{i,n} \rightarrow M_{i,\infty}$, i.e. $M_{i,n}$ converges geometrically to the finite volume complete hyperbolic manifold $M_{i,\infty}$,
- (2) $M_{i,\infty}, M_{j,\infty}$ are distinct for $i \neq j$.

such that the following holds: For any such sequence $\{M_{i,1}, M_{i,2}, M_{i,3}, \dots\}$, there is a sequence of natural numbers $(a_n)_{n \geq 1}$ with $a_n \rightarrow \infty$ such that

- (1) $M_{i,n}$ admits at least a_n contact structures ξ_1, \dots, ξ_{a_n} that are distinct up to contactomorphism,
- (2) ξ_1, \dots, ξ_{a_n} are all Weinstein-fillable.
- (3) ξ_1, \dots, ξ_{a_n} all belong to the same homotopy class of rank-2 distributions on M_n .

Gromov [Gro85] and Eliashberg [Eli90a] observed that Weinstein-fillable contact structures are tight. Colin-Giroux-Honda [CGH09] established that for any closed oriented atoroidal 3-manifold (e.g., hyperbolic 3-manifold) M , the number of tight contact structures that M admits, hereby denoted as $\# \text{Tight}(M)$, is finite. We immediately have the following corollary of Theorem 1.2:

Corollary 1.3. *For any sequence $M_{i,n} \rightarrow M_{i,\infty}$ as in Theorem 1.2, $\#\text{Tight}(M_n) \rightarrow \infty$ as $n \rightarrow \infty$.*

A word about the history of this problem is in order:

- (1) In [LM97], Lisca-Matic proved an analogous result for certain Seifert-fibered spaces that are rational homology 3-spheres, given by $1/n$ -Dehn surgery on the trefoil knot.
- (2) In [BM24, Theorem A.7], Barthelme, Mann and Bowden prove the existence of closed hyperbolic 3-manifolds M_n with at least n non-contactomorphic Anosov contact structures. However, their examples all have positive first Betti number, i.e., $b_1(M_n) > 0$. In fact, the examples in [BM24, Theorem A.7] satisfy $b_1(M_n) \rightarrow \infty$ as $n \rightarrow \infty$.

Outline of the paper:

The examples in Theorem 1.2 arise as $1/n$ -Dehn surgeries on certain hyperbolic knots K as $n \rightarrow \infty$. The knots K in question satisfy the following:

- (1) They are closures $\widehat{\beta}$ of pseudo-Anosov braids β ,
- (2) The braid β has Haken genus greater than one (see Definition 2.1).
- (3) Suppose β has m strands. Let c_+ and c_- respectively denote the number of positive and negative crossings of β . Then

$$c_+ - 2c_- - m \geq 1.$$

Conditions (1) and (2) above guarantee that the closure $\widehat{\beta}$ is, in fact, a hyperbolic knot. This is established in Section 2 and provides a reasonably elementary sufficient condition guaranteeing hyperbolicity of $\widehat{\beta}$. A more sophisticated sufficient condition may be found in [Ito11]. Condition (3) ensures the existence of a Legendrian representative of K with an appropriate framing (see Section 3.2). In the beginning of Section 3, we assume in addition that for the braid β above,

- (4) $c_+ + c_- \equiv m + 1 \pmod{2}$.

This is only logically necessary for the third conclusion of Theorem 1.2 regarding the fact that the underlying rank-2 distributions of the contact structures are in the same homotopy class. This part of the argument is somewhat independent of the arguments needed for the rest of the conclusion of Theorem 1.2, and is discussed in Section 3.4. We refer the reader to Remark 2.12 and Section 3.1 for a discussion on sufficient conditions that guarantee conditions (1), (2), (3) and (4).

While Section 3 is devoted to a proof of Theorem 1.2, we generalize it in two directions in Section 4 (see Theorem 4.4):

- (1) First, we allow links L in place of K . Such links L satisfy conditions (1) and (2) above. Further, each component of L satisfies condition (3).
- (2) The $1/n$ -surgeries are replaced by p/q -surgeries (on each component of L). The surgery coefficients p/q are required to satisfy the property that $-q/p$ has a *terminating negative continued fraction expansion* $[a_0, \dots, a_n]$ with each $a_i \leq -2$ (see Definition 4.1).

This allows us to construct rational homology spheres M with arbitrarily many tight contact structures by rational Dehn filling on links L as above (see Corollary 4.5).

Section 5 provides a notion of geometric convergence of hyperbolic 3-manifolds *equipped with contact structures*. Tripp [Tri06] studied contact structures on non-compact 3-manifolds. He showed that there exist uncountably many non-contactomorphic tight contact structures on an irreducible open 3-manifold with an end of positive genus. We use the technology of bypasses developed by Honda [Hon00a, Hon00b] together with the notion of slope at infinity from [Tri06] to show that there exist uncountably many non-contactomorphic tight contact structures on the complement of a hyperbolic knot K where

- (1) $S^3 \setminus K$ is a geometric limits of closed hyperbolic 3-manifolds *equipped with contact structures*,
- (2) The knots K satisfy conditions (1), (2) and (3) above.

See Theorem 5.10 and Corollary 5.15.

Acknowledgments: We thank Marc Kegel for comments on an earlier draft. The second author thanks Dishant Pancholi and Mike Miller Eisemeier for useful conversations.

2. CONSTRUCTION OF HYPERBOLIC PSEUDO-ANOSOV BRAIDS

2.1. Braid closures. Let B_m denote the braid group on m strands, and let $\beta \in B_m$ be a braid. Let $\widehat{\beta} \subset S^3$ denote the link obtained from taking the closure of β . Fix m points on a disk, hereby denoted as $p_1, \dots, p_m \in D^2$. Let $\text{MCG}_{\partial}(D^2, m)$ denote the mapping class group of the disk with p_1, \dots, p_m as marked points. Then, $B_m \cong \text{MCG}_{\partial}(D^2, m)$. Therefore, β gives rise to a mapping class ϕ . Let

$$T(\phi) = (I \times D^2)/(x, 0) \sim (\phi(x), 1),$$

denote the mapping torus of ϕ . Fix a basepoint $z_0 \in \partial D^2$. The image of $I \times \{z_0\}$ under the quotient map $I \times D^2 \rightarrow T(\phi)$ gives an embedded curve $c \subset \partial T(\phi)$. Choose a genus 1 Heegaard decomposition $S^3 = (S^1 \times D^2) \cup (D^2 \times S^1)$, where we suppose coordinates on either torus are chosen in a way that $S^1 \times \{z_0\}$ in the first solid torus bounds a meridian disk $D^2 \times \{z_0\}$ in the second solid torus. Let us also choose an identification $T(\phi) \cong S^1 \times D^2$ given by sending c to $S^1 \times \{z_0\}$. Consider the composition,

$$D^2 \times I \rightarrow T(\phi) \cong S^1 \times D^2 \hookrightarrow S^3,$$

where the first map is the quotient map to the mapping torus, the middle identification is the one chosen above, and the final embedding is the inclusion of the first solid torus in the chosen Heegaard decomposition of S^3 . The image of $\bigsqcup_i \{p_i\} \times [0, 1]$ under the above map defines the link $\widehat{\beta} \subset S^3$, contained within the solid torus $S^1 \times D^2$. We call $S^1 \times D^2$ the *braid torus*.

We will say that β is *pseudo-Anosov* if the mapping class ϕ is pseudo-Anosov. In this case, the interior of $(S^1 \times D^2) \setminus \widehat{\beta}$ admits the structure of a complete cusped hyperbolic 3-manifold by Thurston's work on hyperbolization of fibered 3-manifolds [Thu86, Ota01].

Definition 2.1. Let $\beta \in B_m$ be a pseudo-Anosov braid. We define an *essential surface* for β to be a properly embedded compact surface $\Sigma \subset (S^1 \times D^2) \setminus \widehat{\beta}$ with boundary $\partial \Sigma = \bigsqcup_i S^1 \times \{x_i\}$ (where $x_i \in \partial D^2$), such that Σ is incompressible, ∂ -nonparallel and

cuspidal-nonparallel. We define the *Haken genus* of β as

$$h(\beta) := \min\{g(\Sigma) : \Sigma \subset (S^1 \times D^2) \setminus \widehat{\beta}\} \cup \{\infty\},$$

where $g(\Sigma)$ denotes the genus of Σ , and Σ varies over all essential surfaces for β .

Essential surfaces with minimal intersection with a Heegaard torus.

Next, let $\Sigma \subset S^3 \setminus \widehat{\beta}$ be an embedded incompressible surface. Note that

$$S^3 \setminus \widehat{\beta} = (S^1 \times D^2 \setminus \widehat{\beta}) \cup (D^2 \times S^1).$$

We refer to this decomposition as the Heegaard decomposition of $S^3 \setminus \widehat{\beta}$ and $\mathcal{T} = \partial D^2 \times S^1$ as the *toral Heegaard surface*, or simply the Heegaard torus. As in Definition 2.1, we refer to $(S^1 \times D^2) \setminus \widehat{\beta}$ as the *braid-torus*. We also refer to $D^2 \times S^1$ as the *solid Heegaard torus*.

Assume that Σ intersects \mathcal{T} transversely and minimally, i.e. $\Sigma \cap \mathcal{T}$ has the minimum number of components possible.

If $S^3 \setminus \widehat{\beta}$ is irreducible, then no component σ of $\Sigma \cap \mathcal{T}$ is homotopically trivial. Indeed, otherwise σ bounds disks D_Σ and $D_\mathcal{T}$ on Σ, \mathcal{T} respectively. By irreducibility of $S^3 \setminus \widehat{\beta}$, $D_\Sigma \cup D_\mathcal{T}$ bounds a ball, and we can isotope \mathcal{T} to reduce the number of components of $\Sigma \cap \mathcal{T}$, contradicting minimality of the intersection. Hence, $\Sigma \cap \mathcal{T}$ consists of finitely many disjoint freely homotopic essential curves $\gamma_1, \dots, \gamma_k$ in \mathcal{T} . Let m/n denote the slope of each γ_i . Here, we use the convention that a curve has slope m/n if it represents m -times a longitude plus n -times a meridian of the *solid Heegaard torus*.

Lemma 2.2. *Suppose $k > 0$. Then each γ_i has slope zero.*

Proof. If not, then $\Sigma \cap \mathcal{T} = \gamma_1 \cup \dots \cup \gamma_n$ where each γ_i has slope m/n with $m \neq 0$. Hence, γ_i is homotopically non-trivial in the solid Heegaard torus $D^2 \times S^1$. Since $\Sigma \cap (D^2 \times S^1)$ is minimal, it follows that each component of $\Sigma \cap (D^2 \times S^1)$ is an annulus \mathcal{A} where the boundary curves have slope m/n on \mathcal{T} . Since any two such curves bound an annulus on \mathcal{T} isotopic rel. boundary to \mathcal{A} , we can isotope Σ so that each such \mathcal{A} is isotoped rel. boundary into \mathcal{T} ; and subsequently into the braid torus. Since $\Sigma \cap \mathcal{T}$ is minimal, it follows that $\Sigma \cap \mathcal{T} = \emptyset$, contradicting $k > 0$. \square

The following is immediate from Lemma 2.2.

Corollary 2.3. *Let Σ be an embedded incompressible surface in an irreducible $M = S^3 \setminus \widehat{\beta}$ having minimal intersection with \mathcal{T} . Then*

- (1) *either Σ is contained in the braid torus,*
- (2) *or each γ_i bounds a disk in the solid Heegaard torus.*

2.2. An application of Thurston's hyperbolization. Recall that a compact 3-manifold M (possibly with boundary) is *atoroidal* if any essential torus in M is homotopic into ∂M . Also, M is *Haken* if M is irreducible and admits an embedded incompressible surface of positive genus.

Remark 2.4. The above definition of atoroidal is sometimes referred to as *geometrically atoroidal*, where the (homotopically) essential torus is, by definition, embedded. However, there are a few examples of irreducible geometrically atoroidal 3-manifolds M that satisfy $\mathbb{Z}^2 \subset \pi_1(M)$, where the \mathbb{Z}^2 is carried by an immersed torus not homotopic into ∂M , i.e.

M is not homotopically atoroidal. We discuss the relevant examples now. In the following, we shall be interested in link complements, where ∂M consists of tori. If M is atoroidal, this forces M to be a Seifert-fibered space over a base-orbifold of type (p, q, ∞) with $p, q \in \mathbb{N} \cup \{\infty\}$, and $p \leq q$. If $p = q = \infty$, M is necessarily homeomorphic to $S_{0,3} \times S^1$, where $S_{0,3}$ is a 3-holed sphere. In this case, M is the complement of a trivial 3-component link.

If $q = \infty$, and $p \in \mathbb{N}$, then $\pi_1(M)$ is $\mathbb{Z} \times (\mathbb{Z} * \mathbb{Z}/p)$. M is the complement in S^3 of a 2-component link L of the form $L = K_1 \cup K_2$, where K_1 is the core curve of a solid torus \mathbb{T} embedded in a standard way in S^3 , and K_2 is a (p_1, p) torus knot on the boundary $\partial\mathbb{T}$ (with any $p_1 \geq 1$). We shall refer to such links L as 2-component $T^2 \times I$ -links, as the components of L may be assumed to lie on the boundary components of $T^2 \times \{0, 1\}$ of $T^2 \times I$.

Finally, let $p \leq q < \infty$ and $(p, q) = 1$. Then M is the complement of a (p, q) -torus knot. It is indeed possible that such an M can be the complement in S^3 of the closure of a pseudo-Anosov braid. In fact, consider the braid group B_3 for braids on 3 strands, with standard generators σ_1, σ_2 . The braid $\beta = \sigma_1^3 \sigma_2^{-1}$ satisfies

- (1) β is pseudo-Anosov. Note that this means that its complement in the braid torus is hyperbolic.
- (2) $\widehat{\beta}$ is a (p, q) -torus knot in S^3 . Note that this implies that its complement in S^3 is not hyperbolic.

This unusual phenomenon accounts for the hypothesis in Corollary 2.6 excluding such examples.

More generally, if $dp_1 = p \leq q = dq_1 < \infty$ and $(p_1, q_1) = 1$, then M is the complement of a link L consisting of d parallel copies of a (p_1, q_1) -torus knot in S^3 . We shall refer to M in this case as a (p, q) -torus link. But in this case, L is a satellite of a (p_1, q_1) -torus knot; in fact it consists of d parallel copies of a (p_1, q_1) -torus knot K . Let $\nu(K)$ denote a solid torus neighborhood of K . Then $\partial\nu(K)$ is an embedded incompressible torus in the complement of L .

We start with Thurston's geometrization theorem:

Theorem 2.5. [Ota98, Kap01] *Let M be a compact atoroidal Haken 3-manifold (possibly with boundary). Then the interior of M admits a complete hyperbolic structure.*

Corollary 2.6. *Let $\beta \in B_m$ be any pseudo-Anosov braid with Haken genus $h(\beta) > 1$. Then the closure $\widehat{\beta} \subset S^3$ is a hyperbolic link, provided $\widehat{\beta}$ is not one of the following (see Remark 2.4):*

- (1) a torus knot,
- (2) a 2-component $T^2 \times I$ -link,
- (3) a trivial 3-component link.

Proof. By the Definition of Haken genus, Remark 2.4, and Theorem 2.5, it suffices to show that $S^3 \setminus \widehat{\beta}$ is irreducible and atoroidal. Recall that $S^3 \setminus \widehat{\beta} = (S^1 \times D^2 \setminus \widehat{\beta}) \cup (D^2 \times S^1)$ is a Heegaard decomposition of $S^3 \setminus \widehat{\beta}$ and $\mathcal{T} = \partial D^2 \times S^1$ is the toral Heegaard surface.

If $S^3 \setminus \widehat{\beta}$ is reducible, then there exists an embedded essential sphere Σ_0 in $S^3 \setminus \widehat{\beta}$. We assume that Σ_0 intersects \mathcal{T} minimally. In particular, $\Sigma_0 \cap \mathcal{T}$ consists of finitely

many disjoint freely homotopic essential curves in \mathcal{T} . Further, $(S^1 \times D^2 \setminus \widehat{\beta}) \cap \Sigma_0$ must then consist of finitely many annuli. But the existence of any such annulus would force the automorphism ϕ associated to $\widehat{\beta}$ to be reducible, violating the hypothesis that ϕ is pseudo-Anosov.

Next, if $S^3 \setminus \widehat{\beta}$ is not atoroidal, then there exists an embedded essential torus Σ_1 in $S^3 \setminus \widehat{\beta}$ that is not parallel to a boundary component. We assume that Σ_1 intersects \mathcal{T} minimally so that $\Sigma_1 \cap \mathcal{T}$ consists of finitely many disjoint freely homotopic essential curves in \mathcal{T} . Further, $(D^2 \times S^1) \cap \Sigma_1$ must then

- (1) either consist of finitely many embedded essential annuli,
- (2) or finitely many embedded meridinal disks.

In the former case, each of these annuli can be homotoped into \mathcal{T} and subsequently into $(S^1 \times D^2 \setminus \widehat{\beta})$. Thus, minimality of $\Sigma_1 \cap \mathcal{T}$ forces this intersection to be empty. Finally, hyperbolicity of $(S^1 \times D^2 \setminus \widehat{\beta})$ forces Σ_1 to be homotopic to a boundary component of $(S^1 \times D^2 \setminus \widehat{\beta})$. If this boundary component is \mathcal{T} , then it is compressible in $S^3 \setminus \widehat{\beta}$ violating essentiality of Σ_1 . Else it is homotopic to a boundary component of a neighborhood of $\widehat{\beta}$ in S^3 . But this contradicts the choice of Σ_1 which was chosen not to be parallel to a boundary component of $S^3 \setminus \widehat{\beta}$. In the latter case, $h(\beta) = 1$.

Note that the above argument also rules out the case that $\widehat{\beta}$ is a (p, q) -torus link in the sense of Remark 2.4. Indeed $(p, q) > 1$ forces the existence of an incompressible Σ_1 (with boundary) of genus equal to one.

The above facts force $S^3 \setminus \widehat{\beta}$ to be irreducible and atoroidal. Theorem 2.5 now furnishes the conclusion. \square

2.3. Genus one surfaces. This subsection is aimed at understanding understanding genus one essential surfaces Σ_1 in the braid torus. We assume further that β is pseudo-Anosov. By Corollary 2.3, either such a Σ_1 is embedded in the braid torus or each component of $\Sigma_1 \cap \mathcal{T}$ bounds a disk in the solid Heegaard torus. Since β is pseudo-Anosov, the first alternative forces Σ_1 to be parallel to a boundary component of the braid torus, i.e. Σ_1 is either ∂ -parallel or cusp-parallel in the sense of Definition 2.1. We therefore assume henceforth that each component of $\Sigma_1 \cap \mathcal{T}$ bounds a disk in the solid Heegaard torus. In particular, the genus of Σ_1 equals that of $\Sigma_1 \cap (S^1 \times D^2 \setminus \widehat{\beta})$.

Next, we cut the braid torus $(S^1 \times D^2) \setminus \widehat{\beta}$ along $(\{p\} \times D^2) \setminus \widehat{\beta}$ for some $p \in S^1$ so as to get the braid complement $\mathcal{C} = ([0, 1] \times D^2) \setminus \beta$. Further, from the above description of $\Sigma_1 \cap \mathcal{T}$, we can assume that

- (1) Σ_1 is transverse to $(\{p\} \times D^2) \setminus \widehat{\beta}$ in the braid torus $(S^1 \times D^2) \setminus \widehat{\beta}$.
- (2) Let Σ_1^c denote $\Sigma_1 \cap (S^1 \times D^2) \setminus \widehat{\beta}$ cut open along $(\{p\} \times D^2) \setminus \widehat{\beta}$ (and then completed in \mathcal{C}). Then Σ_1^c intersects the *vertical boundary* $\partial_v \mathcal{C} = [0, 1] \times \partial D^2$ in vertical lines of the form $[0, 1] \times \{z_i\}$, $i = 1, \dots, 2l$,
- (3) Σ_1^c intersects the *horizontal boundary* $\partial_h \mathcal{C} = \{0, 1\} \times D^2$ in finitely many pairs of disjoint chords $\alpha_1 \times \{0, 1\}, \dots, \alpha_l \times \{0, 1\}$ (the same for the top and bottom components of $\partial_h \mathcal{C}$, where $\alpha_1, \dots, \alpha_l$ are chords in the disk avoiding the initial points of the braid β).
- (4) The standard height function $H : \mathcal{C} \rightarrow [0, 1]$ given by the projection onto the first coordinate of \mathcal{C} is a Morse function on Σ_1^c .

Let $\mathcal{S}_i := \alpha_i \times [0, 1] \subset D^2 \times [0, 1]$ denote *vertical rectangles* corresponding to chords α_i . Then $\partial \Sigma_1^c = \bigcup_i \partial \mathcal{S}_i$, i.e. the boundary of Σ_1^c consists of the disjoint union of the boundary of the vertical rectangles \mathcal{S}_i . Further, there exist

- (1) finitely many disks $\mathcal{D}_1, \dots, \mathcal{D}_j$ contained in the interior of $\bigcup_i \mathcal{S}_i$, and
- (2) finitely many orientable surfaces $\mathcal{A}_1, \dots, \mathcal{A}_k$, where each \mathcal{A}_i is homeomorphic to a genus g surface, $g \geq 0$ with exactly *two* boundary components. We refer to the \mathcal{A}_i as *generalized annuli*.

such that

$$\Sigma = (\bigcup_i \mathcal{S}_i \setminus (\bigcup_r \mathcal{D}_r)) \bigcup (\bigcup_s \mathcal{A}_s),$$

i.e. Σ_1^c is obtained from $\bigcup_i \mathcal{S}_i$ by removing the interiors of the disks \mathcal{D}_r and attaching the generalized annuli \mathcal{A}_s along the boundaries of the removed disks. We construct an auxiliary (undirected) graph $\mathcal{G} := \mathcal{G}(\mathcal{S}, \mathcal{A})$ as follows:

- (1) the vertex set $\mathcal{V}(\mathcal{G})$ equals $\{\mathcal{S}_i\}$,
- (2) the edge set $\mathcal{E}(\mathcal{G})$ equals $\{\mathcal{A}_s\}$,
- (3) the edge \mathcal{A}_s connects vertices $\mathcal{S}_a, \mathcal{S}_b$ if the boundary circles of \mathcal{A}_s lie on $\mathcal{S}_a, \mathcal{S}_b$. (Note that we allow $\mathcal{S}_a = \mathcal{S}_b$.)

Lemma 2.7. *Let Σ_1 be an essential genus 1 surface in a braid link. Then, with notation as above, the generalized annuli \mathcal{A}_s used in constructing Σ_1^c all have $g = 0$, i.e. each \mathcal{A}_s is an annulus. Further, \mathcal{G} has exactly one edge-loop, i.e. $\pi_1(\mathcal{G}) = \mathbb{Z}$.*

Proof. First, observe that \mathcal{G} encodes a graph of spaces description of Σ_1^c , and hence a graph of groups description of $\pi_1(\Sigma_1^c)$, where

- (1) the vertex groups are $\pi_1(\mathcal{S}_i \setminus (\bigcup_r \mathcal{D}_r))$,
- (2) the edge groups are $\pi_1(\mathcal{A}_i)$.

Since Σ_1^c has genus one, we can assume without loss of generality that each \mathcal{A}_i is an annulus, so that $\pi_1(\mathcal{A}_i) = \mathbb{Z}$. (Note that the only other possibility necessitates that \mathcal{A}_1 is a twice-punctured torus connecting two squares $\mathcal{S}_1, \mathcal{S}_2$. This is equivalent to having two disjoint annuli between \mathcal{S}_1 and \mathcal{S}_2 . The associated graph of spaces has two vertices corresponding to \mathcal{S}_1 and \mathcal{S}_2 and two edges between them. We assume henceforth that such a twice-punctured torus is replaced by two annuli.)

Next, we glue the top and bottom of \mathcal{C} together to get back the braid torus. Gluing the horizontal boundary chords of Σ_1^c back together, we also get back $\Sigma_1 \cap (S^1 \times D^2 \setminus \widehat{\beta})$. Note that each $\mathcal{S}_i \setminus (\bigcup_r \mathcal{D}_r)$ now glues up to give an annulus $\widehat{\mathcal{S}}_i \setminus (\bigcup_r \mathcal{D}_r)$ with holes corresponding to \mathcal{D}_r . Thus, $\Sigma_1 \cap (S^1 \times D^2 \setminus \widehat{\beta})$ admits a graph of spaces description, with the same underlying graph \mathcal{G} , where

- (1) the vertex spaces are $\widehat{\mathcal{S}}_i \setminus (\bigcup_r \mathcal{D}_r)$,
- (2) the edge spaces are the annuli \mathcal{A}_i .

We next note that the genus of $\Sigma_1 \cap (S^1 \times D^2 \setminus \widehat{\beta})$ equals the genus of \mathcal{G} , where the genus of \mathcal{G} equals the rank of its (free) fundamental group. Since the genus of $\Sigma_1 \cap (S^1 \times D^2 \setminus \widehat{\beta})$ equals one, the conclusion follows. \square

2.4. Increasing genus. Theorem 2.9 below is the main output of this section. We shall need some basic terminology. Let $\mathcal{C} = (I \times D^2) \setminus \beta$ denote the braid complement as before.

Definition 2.8. A surface $\mathcal{F} \subset \mathcal{C}$ is called an *elementary saddle* if there exist 4 cyclically ordered distinct points $p_1, p_2, p_3, p_4 \in \partial D^2$, and a quadrilateral with sides $[p_1, p_2]$, $[p_2, p_3]$, $[p_3, p_4]$, $[p_4, p_1] \subset D^2$ such that

- (1) \mathcal{F} is homeomorphic to a closed disk.
- (2) the boundary $\partial \mathcal{F}$ consists of
 - the vertical arcs $\cup_i I \times \{p_i\}$,
 - the horizontal arcs $[p_1, p_2] \times \{0\}$, $[p_3, p_4] \times \{0\}$ on $\{0\} \times D^2$,
 - the horizontal arcs $[p_2, p_3] \times \{1\}$, $[p_4, p_1] \times \{1\}$ on $\{1\} \times D^2$.
- (3) the height function $H : \mathcal{C} \rightarrow [0, 1]$ given by projection to the first coordinate is a Morse function with a single saddle singularity.

A surface $\mathcal{F} \subset \mathcal{C}$ is called a *trivial rectangle* if there exists a pair of distinct points p, q and an arc $[p, q] \subset D^2$ such that \mathcal{F} is isotopic rel. boundary to $I \times [p, q]$.

Theorem 2.9. For any pseudo-Anosov braid β , $h(\beta^k) > 1$ for $k > 4$.

Proof. We first note that the second paragraph of the proof of Corollary 2.6 in fact shows that $M = S^3 \setminus \widehat{\beta}$ is irreducible. (The hypothesis $h(\beta) > 1$ in Corollary 2.6 was not used for the proof of this fact.) Hence $h(\beta) \neq 0$.

We observe that the complement M_k in S^3 of the closure $\widehat{\beta^k}$ is obtained as follows. Let $M = S^3 \setminus \widehat{\beta}$ as before. Let \mathcal{H} denote the solid Heegaard torus, and let $c \subset \mathcal{H}$ denote the core curve of \mathcal{H} . Then M_k is a k -fold branched cover of M branched over c .

For concreteness, we show first that an essential embedded torus in M pulls back to an essential surface of genus greater than one in M_k . (Strictly speaking, the argument in this paragraph is not logically necessary for the rest of the proof, but it provides an easy first case.) Let Σ_1 denote the essential genus one surface in M . Recall the structure of Σ_1 by Lemma 2.7. Recall also that Σ_1 intersects the solid Heegaard torus of M in finitely many disks $\mathcal{D}_1, \dots, \mathcal{D}_j$. Let Σ_1^k denote its pullback to M_k . Then Σ_1^k intersects the solid Heegaard torus of M_k in finitely many disks $\mathcal{D}'_1, \dots, \mathcal{D}'_j$, where \mathcal{D}'_r is a k -fold branched cover of \mathcal{D}_r over the origin. Then Σ_1^k has genus greater than one. If possible, suppose that Σ_1^k is compressible, and $\gamma \subset \Sigma_1^k$ is an essential closed curve that is contractible in M_k . Then γ can be isotoped slightly to be disjoint from the branch locus, forcing the image of γ in M to be contractible. This contradicts the fact that Σ_1 is essential.

It remains to show that there is no other essential surface in M_k of genus zero or one. If Σ_0 is an essential sphere in M_k , its image, under the branched cover map Π to M , would give an immersed essential sphere $\Pi(\Sigma_0)$ in M . Hence by the sphere theorem, M contains an essential embedded sphere, forcing $h(\beta) = 0$ and contradicting that β is pseudo-Anosov as in the proof of Corollary 2.6.

If Σ'_0 is an essential torus in M_k , its image $\Pi(\Sigma'_0)$ is an immersed essential torus in M . Let $\mathcal{B} = (S^1 \times D^2) \setminus \widehat{\beta} \subset M$ denote the complement of the braid β within the braid torus. Let \mathcal{B}_k denote its lift to M_k , so that \mathcal{B}_k is a k -fold regular cover of \mathcal{B} . Note that \mathcal{B}_k is the complement of the braid β^k within the braid torus of β^k . We assume Σ'_0 intersects the boundary of the braid torus $\partial \mathcal{B}_k$ (i.e., the Heegaard torus) transversely and minimally.

By Corollary 2.3, either Σ'_0 is contained entirely inside the braid torus or intersects the Heegaard torus in a collection of circles which bound meridinal disks in the solid Heegaard torus. The first scenario is impossible, as \mathcal{B}_k is hyperbolic and therefore atoroidal, since β is pseudo-Anosov.

Let $\mathcal{C}_k = ([0, k] \times D^2) \setminus \widehat{\beta^k}$ denote \mathcal{B}_k cut open along a disk. Let $H_k : \mathcal{C}_k \rightarrow [0, k]$ denote the associated height function. Let $\Sigma_0^{cc} \subset \mathcal{C}_k$ denote $\Sigma'_0 \cap \mathcal{B}_k$ cut along the disk. We assume that

- (1) H_k is a Morse function when restricted to Σ_0^{cc} ,
- (2) the total number of critical points with respect to H_k is the minimum possible in the isotopy class of Σ_0^{cc} rel. boundary. In particular, H_k has no maxima or minima.

Since \mathcal{B}_k is a k -fold regular cover of \mathcal{B} , we assume that \mathcal{C}_k is a vertical concatenation of k copies of \mathcal{C} . Let $\mathcal{Q}_i := (D^2 \times [i-1, i]) \setminus \beta$ denote the i -th block of \mathcal{C} in \mathcal{C}_k , $i = 1, \dots, k$. Let $\mathcal{F}_i = \Sigma_0^{cc} \cap \mathcal{Q}_i$. By the discussion in Section 2.3 we can assume that there exists $l > 0$ and points p_1, \dots, p_{2l} such that

- (1) the vertical boundary of \mathcal{F}_i equals $[i-1, i] \times \{p_1, \dots, p_{2l}\}$ for all i ,
- (2) the horizontal boundary of \mathcal{F}_i restricted to $\{i\} \times D^2$ consists of l disjoint arcs $\alpha_{i1}, \dots, \alpha_{i2l}$ in $\{i\} \times D^2$ pairing points in $\{i\} \times \{p_1, \dots, p_{2l}\}$.

We next observe that \mathcal{F}_i cannot contain a trivial rectangle, else the trivial rectangle would separate strands of the braid β into two non-empty clusters, forcing $\widehat{\beta}$ to be a split link, violating the hypothesis that β is a pseudo-Anosov braid. Hence, $\alpha_{i1}, \dots, \alpha_{i2l}$ contribute at least $l/2$ elementary saddles in \mathcal{F}_i , i.e. \mathcal{F}_i has at least $l/2$ critical points of index 1. Recall that H_k restricted to \mathcal{F}_i has no maxima or minima.

Hence, the total number of index 1 critical points on Σ_0^{cc} is at least $k \cdot l/2$. Hence, gluing Σ_0^{cc} back along its top and bottom boundaries, we see $\Sigma'_0 \cap \mathcal{B}_k$ has a circle-valued Morse function with at least $k \cdot l/2$ index 1 critical points, and no index 0 or index 2 critical points. Moreover, the circle-valued Morse function is identity restricted to the $2l$ boundary circles of $\Sigma'_0 \cap \mathcal{B}_k$. Therefore, by the Poincare-Hopf index theorem,

$$\chi(\Sigma'_0 \cap \mathcal{B}_k) \leq -k \cdot \frac{l}{2}$$

Note that Σ'_0 is obtained from $\Sigma'_0 \cap \mathcal{B}_k$ by capping off the $2l$ boundary circles by disks. Thus, $\chi(\Sigma'_0) = \chi(\Sigma'_0 \cap \mathcal{B}_k) + 2l$. Combining with the above, we get,

$$\chi(\Sigma'_0) \leq 2l - k \cdot \frac{l}{2}$$

Hence, for $k > 4$, we must have $\chi(\Sigma'_0) < 0$. Therefore, Σ'_0 has genus greater than 1. Hence β^k has Haken genus greater than 1. This proves the theorem. \square

Remark 2.10. We summarize some conditions to ensure that a braid has hyperbolic braid closure. From Corollary 2.6 and Theorem 2.9 we have the following. Let β be a pseudo-Anosov braid such that its closure is not one of the following:

- (1) a torus knot,
- (2) a 2-component $T^2 \times I$ -link,
- (3) a trivial 3-component link.

Then $\widehat{\beta^k}$ is hyperbolic for $k > 4$.

We refer to [Ito11] for the notion of Dehornoy floor $[\beta]_D$ of a braid β . In [Ito11, Theorem 2], Ito shows the following:

Theorem 2.11. [Ito11] *Let $\beta \in B_m$ be a braid whose closure is a knot satisfying the condition that its Dehornoy floor $[\beta]_D$ satisfies $[\beta]_D \geq 3$. Then $\widehat{\beta}$ is hyperbolic if and only if β is pseudo-Anosov.*

Remark 2.12 (Genericity). In [Car17, CW17], Caruso and Wiest proved the genericity of pseudo-Anosov elements of braid groups. More precisely, equip the braid group B_m ($m \geq 3$) with Garside's generating set, and let Γ be the resulting Cayley graph. Then it is shown in [CW17] that the proportion of pseudo-Anosov braids in the ball of radius r in Γ tends to 1 exponentially fast as $r \rightarrow \infty$.

Remark 2.10 and Theorem 2.11 now furnish two different ways of upgrading a pseudo-Anosov braid β to a hyperbolic link in S^3 .

3. PROOF OF THEOREM 1.2

3.1. Construction of the manifolds. We begin by constructing the required 3-manifolds. Let $\beta \in B_m$ be a braid on m strands. Let c_+ and c_- respectively denote the number of positive and negative crossings of β .

Hypothesis 3.1. Let us suppose $\beta \in B_m$ is a braid such that,

- (1) The closure $\widehat{\beta}$ is a hyperbolic knot,
- (2) $c_+ - 2c_- - m \geq 1$,
- (3) $c_+ + c_- \equiv m + 1 \pmod{2}$.

Remark 3.2. The third condition of Hypothesis 3.1 is only logically necessary in Section 3.4 to guarantee that the non-contactomorphic tight contact structures on M_n we will construct in Section 3.2 are homotopic as 2-plane distributions. In particular, the first two conditions of Hypothesis 3.1 are sufficient to ensure all but the third conclusion in Theorem 1.2.

Satisfying Hypothesis 3.1:

Theorem 2.9 and Corollary 2.6 together furnish a large family of examples of $\beta \in B_m$ such that $\widehat{\beta}$ is a hyperbolic knot using the notion of Haken genus (see Remark 2.10). Theorem 2.11 uses the notion of Dehornoy floor to give a different sufficient condition. Hence, braids satisfying the first condition above are furnished by the discussion at the end of Section 2. In particular, Remark 2.12 shows that such $\widehat{\beta}$ may be obtained from generic braids.

The second condition is much easier to satisfy, once a pseudo-Anosov braid β has been fixed. Let Δ denote the Garside element in the braid group B_m given by

$$\Delta := (\sigma_1 \sigma_2 \cdots \sigma_{n-1})(\sigma_1 \sigma_2 \cdots \sigma_{n-2}) \cdots (\sigma_1 \sigma_2)(\sigma_1).$$

Then Δ^2 generates the infinite cyclic center of B_m and has only *positive crossings*. If β is pseudo-Anosov, then so is $\Delta^{2\ell} \beta$ for all $\ell \in \mathbb{Z}$. Next, let $\widehat{\Delta^{2\ell} \beta}$ denote the closure of $\Delta^{2\ell} \beta$ in S^3 . Then $S^3 \setminus \widehat{\Delta^{2\ell} \beta}$ is obtained from $(S^1 \times D^2) \setminus \widehat{\beta}$ by attaching a disk to the $(1, \ell)$ -curve on the boundary $S^1 \times S^1$ of $S^1 \times D^2$. Choosing $\ell > 0$ large enough, the second condition is also satisfied by Thurston's hyperbolic Dehn surgery [Thu22, Chapter 4].

For the third condition, we observe that $c_+ + c_-$ is the total number of crossings (or equivalently, length of the braid word) of β . This can be ensured, for instance, (together with the first and third conditions) as follows: let β be a braid on m strands such that m is odd, and β^2 is a knot. That is to say, the permutation corresponding to β squares to an m -cycle. Then the total number of crossings of β^2 is even. Hence, it agrees with the parity of $m + 1$. Thus, $\Delta^{2\ell}\beta^2$ satisfies all three conditions provided $\ell > 0$ is sufficiently large, as in the previous paragraph.

Constructing hyperbolic homology spheres:

Let $M_n := S_{1/n}(\widehat{\beta})$ denote the $1/n$ -Dehn filling of $S^3 \setminus \widehat{\beta}$. Here, we use the convention that $1/n$ -Dehn filling attaches a disk to $\mu + n\lambda$, where μ is the meridian of the knot $\widehat{\beta}$, and λ is a longitude given by pushing the knot off along a Seifert surface. This is an integer homology 3-sphere, and by Thurston's hyperbolic Dehn surgery theorem [Thu22, Chapter 4], M_n is a closed hyperbolic 3-manifold for $n > 0$ sufficiently large.

We now construct a smooth 4-dimensional handlebody $(X_n, \partial X_n)$ such that $\partial X_n \cong M_n$. Let $K := \widehat{\beta} \subset S^3$ and let $U \subset S^3$ denote the meridian of K . By the slam-dunk operation [GS99, Section 5.3., p. 163], we may rewrite the surgery description of M_n as a surgery on the link $K \cup U \subset S^3$ with surgery coefficients 0 and $-n$ on K and U respectively. By treating this integral surgery diagram as a Kirby diagram, we obtain a 4-dimensional handlebody $(X_n, \partial X_n)$ by attaching a pair of 2-handles h_K^2, h_U^2 to D^4 along K and U , respectively, with framing as in the surgery coefficients.

3.2. Construction of the contact structures. The aim of this section is to prove:

Theorem 3.3. *Let $(X_n, \partial X_n)$ be the 4-dimensional handlebodies with boundary constructed in Section 3.1. Then $(X_n, \partial X_n)$ admit Weinstein structures.*

Below, tb denotes the *Thurston-Bennequin number*. We shall use the following theorem:

Theorem 3.4. [Eli90b, Gom98] *Let $(W, \partial W)$ be a smooth 4-manifold-with-boundary, with a handlebody decomposition $W = D^4 \cup_i h_i^2$ consisting of only 2-handles. Suppose that there exists a Legendrian link $L = \cup_i K_i \subset \partial D^4$ in the framed isotopy class of the union of the attaching circles of the 2-handles h_i^2 such that the framing on K_i is $tb(K_i) - 1$ for all i . Then W admits the structure of a Weinstein domain.*

Proof of Theorem 3.3. First, we draw a knot diagram for the braid closure $K = \widehat{\beta}$ (cf. Section 3.1) by drawing a horizontal planar diagram of the braid β , followed by a collection of arcs $\alpha_1, \dots, \alpha_m$ closing the endpoints of β . Note that m is the number of strands of β . We orient K so that the strands of β traverse “left-to-right” while the arcs α_i traverse “right-to-left”. Each arc α_i gives rise to two vertical tangencies of the knot diagram, which we replace by a pair of cusps of appropriate orientation. (See Figure 1.)

We shall draw every left-handed (or negative) crossing of β as in one of the two pictures in Figure 2, so that the overcrossing arc has slope less than that of the undercrossing arc. More precisely, let us fix an ordering on all the negative crossings. If c_- is even, we draw the negative crossings alternately as in the two pictures in Figure 2. If c_- is odd, we draw




FIGURE 1. Braid closure $\hat{\beta}$ obtained by joining the end-points of braid β by arcs $\alpha_1, \dots, \alpha_m$ (left); Legendrian realization of $\hat{\beta}$ (right).

the first negative crossing as in the right picture in Figure 2, and alternately use the two pictures for the rest.




FIGURE 2. Two Legendrian realizations of a left-handed crossing by introduction of two cusps of the same orientation.

These modifications lead to a Legendrian front projection for a Legendrian knot K' in the smooth isotopy class of $K \subset S^3$. We note the following regarding the front diagram for K' that we have constructed:

- (1) The number of positive and negative crossings are the same as those for the braid β , i.e. c_+ and c_- , respectively.
- (2) Each arc α_i as well as each negative crossing of K contributes to a pair of cusps. Therefore, the total number of cusps for the front diagram of K' is $2(m + c_-)$. The pair of cusps corresponding to each arc α_i are of opposite orientation, while the pair of cusps corresponding to each crossing are of the same orientation.

We compute the Thurston-Bennequin number for the Legendrian realization:

$$\text{tb}(K') = c_+ - c_- - \frac{1}{2} \cdot 2(m + c_-) = c_+ - 2c_- - m.$$

Moreover, by construction,

$$\text{rot}(K') = \begin{cases} 0, & \text{if } c_- \text{ is even,} \\ 1, & \text{if } c_- \text{ is odd.} \end{cases}$$

By Condition (2) of Hypothesis 3.1 on β , $\text{tb}(K') \geq 1$. By Condition (3) of Hypothesis 3.1 on β , $\text{tb}(K')$ has the same parity as c_- . Therefore, we may stabilize K' precisely $\text{tb}(K') - 1$ times, and choose the sign of the stabilizations so that $\text{rot}(K') = 0$.

Let K_{Leg} denote the resulting Legendrian knot obtained by stabilizing K' exactly $\text{tb}(K') - 1$ times in this manner. Then K_{Leg} is a Legendrian knot which is still contained in the smooth isotopy class of $K \subset S^3$. Moreover, we have $\text{tb}(K_{\text{Leg}}) = 1$ and $\text{rot}(K_{\text{Leg}}) = 0$.

For any $1 \leq k \leq n - 1$, there is a Legendrian realization U_{Leg} of the unknot with $\text{tb}(U_{\text{Leg}}) = -n + 1$ and $\text{rot}(U_{\text{Leg}}) = 2k - n$. We use such a Legendrian realization for the meridian of K_{Leg} .

Thus, the framed link $K_{\text{Leg}} \cup U_{\text{Leg}} \subset S^3$ with framing 0 and $-n$ respectively, satisfies the hypothesis of Theorem 3.4. Since the smooth 4-dimensional handlebody corresponding to this framed link is the manifold X_n constructed in Section 3.1, this produces a Weinstein structure on X_n . Note that *a priori* the ambiguity of the choice of U_{Leg} could lead to different Weinstein structures on X_n . We therefore denote the Weinstein domains obtained in this way as $W_{n,k}$, $1 \leq k \leq n - 1$. Note that the underlying smooth manifold for $W_{n,k}$ is X_n . Thus, the Weinstein structures induce contact structures $\xi_{n,k}$ ($1 \leq k \leq n - 1$) on the boundary $M_n = \partial X_n$. Since $\xi_{n,k}$ are Weinstein-fillable contact structures, they must be tight [Gro85, Eli90a]. \square

3.3. Distinguishing contact structures. To distinguish the contact structures $\xi_{n,k}$ ($1 \leq k \leq n - 1$) on M_n up to isotopy, we follow the overall strategy of Lisca-Matić [LM97]. We shall use the relative Seiberg-Witten invariants constructed by Kronheimer-Mrowka [KM97]. In particular, we need the following theorem:

Theorem 3.5. [KM97, Theorem 1.2.] *For a pair (X, ξ) consisting of a compact, connected, oriented 4-manifold X with nonempty boundary and a contact structure ξ on ∂X , there exists a monopole invariant*

$$\text{SW} : \text{Spin}^c(X, \xi) \rightarrow \mathbb{Z},$$

defined on the space of Spin^c -structures on X compatible with ξ . If X admits an exact symplectic form ω compatible with ξ , then $\text{SW}(\mathfrak{s}) \neq 0$ if and only if $\mathfrak{s} = \mathfrak{s}_0$ is the canonical Spin^c -structure corresponding to ω .

Proposition 3.6. *The contact structures $\xi_{n,k}$, $1 \leq k \leq n - 1$, on M_n constructed in Section 3.2 are distinct up to isotopy.*

Proof. Suppose $\xi_{n,k}$ and $\xi_{n,l}$ are isotopic contact structures on $M_n = \partial X_n$. We choose an isotopy between them, and extend the isotopy into a symplectized collar of ∂X_n in X_n . Thus, we obtain two different Weinstein structures on X_n , Weinstein-isotopic to those of $W_{n,k}$ and $W_{n,l}$ respectively, inducing the same contact structure ξ on M_n . Let ω_k, ω_l denote the corresponding exact symplectic forms on X_n . Let \mathfrak{s}_k and \mathfrak{s}_l denote the canonical Spin^c structures on X_n corresponding to the symplectic forms ω_k and ω_l , respectively. Explicitly, we may think of \mathfrak{s}_k (resp. \mathfrak{s}_l) as induced from the choice of an almost complex structure J_k (resp. J_l) compatible with ω_k (resp. ω_l).

By Theorem 3.5, $\text{SW}(\mathfrak{s}_k)$ and $\text{SW}(\mathfrak{s}_l)$ are both nonzero, and hence isomorphic. Indeed, they are respectively equal to the canonical Spin^c structures corresponding to possibly different exact symplectic forms ω_k and ω_l on X_n . However, Theorem 3.5 says that there can only be one Spin^c structure on which SW takes nonzero value. In particular, for the

two different symplectic forms ω_k, ω_l , the canonical Spin^c structures they give rise to are the same.

Consequently, J_k, J_l are isomorphic almost-complex structures on X_n . In particular, $c_1(W_{n,k}, J_k) = c_1(W_{n,l}, J_l)$ as cohomology classes in $H^2(X_n; \mathbb{Z})$. Let $[h_U^2] \in H_2(X_n; \mathbb{Z})$ denote the homology class corresponding to the 2-handle h_U^2 in X_n . As the rotation number of the attaching Legendrian knot of a Weinstein 2-handle records the first Chern class of the Weinstein handlebody, we have

$$\begin{aligned}\langle c_1(W_{n,k}, J_k), [h_U^2] \rangle &= 2k - n, \\ \langle c_1(W_{n,l}, J_l), [h_U^2] \rangle &= 2l - n.\end{aligned}$$

Therefore, $k = l$. □

Proposition 3.7. *There exists a natural number $c > 0$ depending only on the braid β chosen in Section 3.1, such that for all $n > 0$ sufficiently large, $\{\xi_{n,k} : 1 \leq k \leq n-1\}$ contains a subset of size at least n/c such that all elements of the subset are pairwise distinct up to contactomorphism.*

Proof. By the hypothesis that $n > 0$ is sufficiently large, M_n is hyperbolic by Thurston's Dehn surgery theorem. Suppose $\xi_{n,k}$ and $\xi_{n,l}$ are contactomorphic. As they are not isotopic by Proposition 3.6, any contactomorphism between them must be a nontrivial smooth mapping class of M_n , i.e., it must not be diffeotopic to identity. By Mostow rigidity and [GMT03], the smooth mapping class group of M_n is isomorphic to the isometry group of M_n . If $n > 0$ is chosen to be sufficiently large, then M_n admits a unique shortest geodesic given by the core of the solid torus used in Dehn filling [Thu22]. Any isometry of M_n must preserve this curve. Deleting this curve results in an isometry of the manifold $S^3 \setminus K$. Therefore, $\text{Isom}(M_n) \cong \text{Isom}(S^3 \setminus K)$. Let $c := |\text{Isom}(S^3 \setminus K)|$ denote the order of this group. This is a finite number by Mostow rigidity, and only depends on the braid β whose closure gives the knot K . Thus, the action of $\text{MCG}(M_n)$ on the set $\{\xi_{n,1}, \dots, \xi_{n,n-1}\}$ has all orbits of size at most c . Therefore, there exist at least n/c contact structures, one from each orbit, that are pairwise distinct up to contactomorphism. □

3.4. Homotopy-indistinguishability of plane fields. In contrast to Proposition 3.6 and 3.7, we shall show that the contact structures $\xi_{n,k}$ ($1 \leq k \leq n-1$) are all homotopic as oriented rank-2 distributions. We will use Gompf's θ -invariant (also known as the d_3 -invariant in some of the more recent literature) to accomplish this.

Definition 3.8. [Gom98, Definition 4.2.] Let M^3 be an oriented 3-manifold and ξ be a co-oriented plane field on M . Let (X^4, J) be an almost-complex 4-manifold with $\partial X = M$ such that $\xi = TM \cap JTM$. The θ -invariant of (M, ξ) is defined to be

$$\theta(M, \xi) := \langle c_1(X, J)^2, [X, \partial X] \rangle - 2\chi(X) - 3\sigma(X),$$

where $c_1(X, J)$ is the first Chern class of (X, J) , $\chi(X)$ is the Euler characteristic of X and $\sigma(X)$ is the signature of $(X, \partial X)$. This is an invariant of (M^3, ξ) independent of the choice of the bounding almost-complex manifold (X, J) .

The following theorem shows that the θ -invariant is a complete invariant of oriented plane fields up to homotopy on integer homology 3-spheres:

Theorem 3.9. [Gom98][LM97, Proposition 2.2.] *If M^3 is an oriented integer homology 3-sphere and ξ_1, ξ_2 are co-oriented plane fields on M , then ξ_1 is homotopic to ξ_2 if and only if $\theta(\xi_1) = \theta(\xi_2)$.*

Proposition 3.10. *The contact structures $\xi_{n,k}$, $1 \leq k \leq n-1$, belong to the same homotopy class of oriented rank-2 distributions.*

Proof. We show that the θ -invariants of $\xi_{n,k}$ are the same for all $1 \leq k \leq n-1$. We choose the bounding almost-complex 4-manifold to be $(W_{n,k}, J_k)$. Since the smooth manifold underlying $W_{n,k}$ is just X_n , the Euler characteristic and signatures agree for all $1 \leq k \leq n-1$. The homology group $H_2(W_{n,k}; \mathbb{Z})$ is freely generated by the fundamental classes $[h_K^2], [h_U^2]$ of the 2-handles.

(1) In Proposition 3.6, we had already computed

$$\langle c_1(W_{n,k}, J_k), [h_U^2] \rangle = 2k - n$$

(2) By the construction of K_{Leg} in Section 3.2, the rotation number of K_{Leg} is zero. This is the only place where Condition (3) of Hypothesis 3.1 is used. Therefore,

$$\langle c_1(W_{n,k}, J_k), [h_K^2] \rangle = 0$$

Thus, $c_1(W_{n,k}) = (2k - n)\text{PD}[h_K^2]$. Hence, $c_1(W_{n,k})^2 = 0$ for all $1 \leq k \leq n-1$ as K has framing 0. This concludes the proof. \square

4. FURTHER EXAMPLES: RATIONAL SURGERY ON LINKS

In Section 3, we constructed tight contact structures on hyperbolic manifolds obtained from choosing appropriate braids β with closure $\widehat{\beta}$ a hyperbolic knot, and performing $1/n$ -surgery on the closure. In this section, we generalize the results of Section 3 in two related directions:

- (1) By considering surgery coefficients of the form p/q instead of $1/n$,
- (2) By allowing $\widehat{\beta}$ to be a hyperbolic link instead of a hyperbolic knot.

First, we set up some notation and preliminary definitions.

Definition 4.1. Let r be a real number. A *negative continued fraction expansion* for r , denoted $[a_0, a_1, a_2, \dots]$, is a (possibly non-terminating) expression of the form

$$r = a_0 - \frac{1}{a_1 - \frac{1}{a_2 - \dots}}$$

such that $a_i \leq -2$. Every real number $r < -1$ admits a unique negative continued fraction expansion. For a terminating continued fraction expansion $r = [a_0, a_1, \dots, a_k]$, we denote

$$\Phi(r) = |a_0 + 1| |a_1 + 1| \cdots |a_k + 1|$$

Remark 4.2. Any real number with a terminating negative continued fraction expansion is rational, but the converse is not true. For instance, $-3 = [-2, -2, -2, \dots]$.

Let $\beta \in B_m$ be a braid. Let $\sigma : B_m \rightarrow S_m$ denote the natural epimorphism from the braid group to the permutation group. The components of the link $\widehat{\beta}$ are in one-to-one correspondence with the cycle decomposition of the permutation $\sigma(\beta) \in S_m$. Suppose $\sigma(\beta)$ has cycle type (m_1, \dots, m_ℓ) . Then, we may write $\widehat{\beta}$ as a union of its components, each of which is a braid knot:

$$\widehat{\beta} = \widehat{\beta}_1 \cup \dots \cup \widehat{\beta}_\ell,$$

where $\beta_i \in B_{m_i}$ is a braid on m_i strands, $1 \leq i \leq \ell$. We denote $c_{i,+}, c_{i,-}$ to be the number of positive and negative crossings of β_i , respectively.

Definition 4.3. Let $\beta \in B_m$ be a braid, with its closure $\widehat{\beta} = \widehat{\beta}_1 \cup \dots \cup \widehat{\beta}_\ell$ as above. Define a *slope vector*

$$\mathbf{v} = (p_1/q_1, \dots, p_\ell/q_\ell),$$

to be an ordered tuple of rational numbers. Let $M(\beta, \mathbf{v})$ be the 3-manifold obtained by Dehn filling $S^3 \setminus \widehat{\beta}$ with surgery coefficient p_i/q_i on the component $\widehat{\beta}_i$, for all $1 \leq i \leq \ell$.

Let $\mathbf{v} = (p_1/q_1, \dots, p_\ell/q_\ell)$ be a slope vector such that $-q_i/p_i$ ($1 \leq i \leq \ell$) admits a terminating negative continued fraction expansion. Define

$$\Phi(\mathbf{v}) = \Phi(-q_1/p_1)\Phi(-q_2/p_2) \cdots \Phi(-q_\ell/p_\ell),$$

where $\Phi(r)$ is defined in 4.1.

Theorem 4.4. Let $\beta \in B_m$ be a braid with closure $\widehat{\beta} = \widehat{\beta}_1 \cup \dots \cup \widehat{\beta}_\ell$ such that $\widehat{\beta}_i$ satisfies the condition $c_{i,+} - c_{i,-} - 2m_i \geq 1$ (cf. Section 3.1) for all $1 \leq i \leq \ell$. Suppose that the slope vector $\mathbf{v} = (p_1/q_1, \dots, p_\ell/q_\ell)$ has the property that $-q_i/p_i$ admits a terminating negative continued fraction expansion for all $1 \leq i \leq \ell$. Then $M(\beta, \mathbf{v})$ admits at least $\Phi(\mathbf{v})$ distinct tight contact structures up to isotopy.

Proof. The proof proceeds along the general strategy described in Section 3. Here we indicate the necessary modifications we have to make.

- (1) As in Section 3.1, we construct a 4-manifold-with-boundary X such that $\partial X = M(\beta, \mathbf{v})$. This is accomplished by first writing the surgery diagram for $M(\beta, \mathbf{v})$ given by:

$$\widehat{\beta} = \widehat{\beta}_1 \cup \dots \cup \widehat{\beta}_\ell,$$

with surgery coefficients as the slope vector \mathbf{v} . We write $p_i/q_i = 0 - q_i/p_i$ and apply the slam-dunk operation to change surgery coefficients of $\widehat{\beta}_i$ to 0, at the cost of introducing a meridian unknot for $\widehat{\beta}_i$ with surgery coefficient $-q_i/p_i$, for all $1 \leq i \leq \ell$. Then we use the terminating negative continued expansion of $-q_i/p_i$ to further modify the surgery diagram so that all the coefficients are integral, by iterated applications of slam-dunk. This replaces the meridian of each component by a chain of Hopf links, with surgery coefficients given by the continued fraction expansion. The final surgery diagram defines a Kirby diagram for the desired 4-manifold X .

- (2) For each $1 \leq i \leq \ell$, the conditions $c_{+,i} - c_{-,i} - 2m_i \geq 1$ together with the conditions $a_i \leq -2$ (cf. Definition 4.1) for the negative continued fraction expansion for $-q_i/p_i$ allow us to modify the Kirby diagram obtained in Step (1) into a Weinstein-Kirby diagram. This is accomplished by choosing a Legendrian realization of all the components (stabilizing them if necessary) and applying Theorem 3.4. Thus, we equip X with the

structure of a Weinstein manifold, which in turn induces a tight contact structure on $\partial X = M(\beta, \mathbf{v})$ just as in Section 3.2.

- (3) While choosing a Legendrian realization in Step (2), there may be ambiguities as to the choice of the Legendrian realization of the unknots occurring in the chain of Hopf links described in Step (1). For each unknot with framing f , the Thurston-Bennequin number is $f + 1$. The possible rotation numbers of a Legendrian realization of the unknot with Thurston-Bennequin number $f + 1$ are $2k - f$, where $1 \leq k \leq -f - 1$. Thus, there are $|f + 1|$ such choices. As these choices can be made independently on each component of the chain of Hopf links appearing in the surgery diagram described in Step (1), there are $\Phi(\mathbf{v})$ many such choices in total. The proof of Proposition 3.6 now shows that these $\Phi(\mathbf{v})$ many contact structures are all distinct up to isotopy, since the tuple of rotation numbers of all components disagree for any two such contact structures. \square

Let $r < -1$ be an irrational real number. Let $r = [a_0, a_1, \dots]$ be a continued fraction expansion and let

$$-q_n/p_n = [a_0, a_1, \dots, a_n]$$

be the n -th convergent. Then, we observe $|p_n| + |q_n| \rightarrow \infty$, and for infinitely many $i \geq 0$, $a_i \leq -3$. The last assertion follows using Remark 4.2, as otherwise r would be rational. Therefore, $\Phi(-q_n/p_n) \rightarrow \infty$ as $n \rightarrow \infty$. Using this, we deduce the following corollary of Theorem 4.4:

Corollary 4.5. *Let $\beta \in B_m$ be a braid with closure $\widehat{\beta} = \widehat{\beta}_1 \cup \dots \cup \widehat{\beta}_\ell$ a hyperbolic link. Suppose that $\widehat{\beta}_i$ satisfies the condition $c_{i,+} - c_{i,-} - 2m_i \geq 1$ for all $1 \leq i \leq \ell$. Let $r_1, \dots, r_\ell < -1$ be irrational numbers, and let $-q_{i,n}/p_{i,n}$ denote the n -th convergent of the negative continued fraction expansion of r_i . Let $\mathbf{v}_n = (p_{1,n}/q_{1,n}, \dots, p_{\ell,n}/q_{\ell,n})$. Then for sufficiently large $n \geq 0$, $M(\beta, \mathbf{v}_n)$ is a hyperbolic manifold and $\#\text{Tight}(M(\beta, \mathbf{v}_n)) \rightarrow \infty$ as $n \rightarrow \infty$.*

Proof. Since $\widehat{\beta}$ is a hyperbolic link and the surgery coefficients on each component satisfy the condition $|p_{i,n}| + |q_{i,n}| \rightarrow \infty$ as $n \rightarrow \infty$ for all $1 \leq i \leq \ell$, $M(\beta, \mathbf{v}_n)$ is hyperbolic for sufficiently large values of n , by Thurston's hyperbolic Dehn surgery theorem. We have established in Theorem 4.4 that $M(\beta, \mathbf{v}_n)$ has at least $\Phi(\mathbf{v}_n)$ non-isotopic contact structures. Since $\Phi(\mathbf{v}_n) \rightarrow \infty$, the proof of Proposition 3.7 shows $M(\beta, \mathbf{v}_n)$ admits an arbitrarily large number of contact structures distinct up to contactomorphism, as $n \rightarrow \infty$. \square

5. GEOMETRIC LIMIT OF CONTACT STRUCTURES

5.1. Preliminary definitions. Recall the following definition from hyperbolic geometry:

Definition 5.1. Let M_n ($n \geq 1$) and M be finite-volume hyperbolic 3-manifolds. The sequence of manifolds $(M_n)_{n \geq 1}$ is said to *geometrically converge* to the manifold M if there exists a collection of points $p_n \in M_n$ and $p \in M$, a collection of radii $r_n > 0$, a collection of Lipschitz constants $\lambda_n > 0$ and a collection of λ_n -Lipschitz diffeomorphisms $\phi_n : B(p_n, r_n) \rightarrow B(p, r_n)$ such that $r_n \rightarrow \infty$ and $\lambda_n \rightarrow 1$ as $n \rightarrow \infty$.

We define a notion of geometric convergence for a sequence of hyperbolic 3-manifolds equipped with a contact structure.

Definition 5.2. Let (M_n, ξ_n) ($n \geq 1$) and (M, ξ) be finite-volume hyperbolic 3-manifolds equipped with contact structures. We shall say the sequence $(M_n, \xi_n)_{n \geq 1}$ *geometrically converges* to (M, ξ) if,

- (1) M_n converges to M geometrically (cf. Definition 5.1),
- (2) We use the notation as in Definition 5.1. For each $n \geq 1$, there exists a diffeotopy $\{f_t\}_{t \in [0,1]}$ of $\text{Op}(\partial B_n(p_n, r_n))$ with $f_0 = \text{id}$, such that

$$f_1^* \xi_n = \phi_n^* \xi,$$

and $\{f_t\}$ extends to a diffeotopy $\{\tilde{f}_t\}_{t \in [0,1]}$ of $B_n(p_n, r_n)$ satisfying $\tilde{f}_0 = \text{id}$, $\tilde{f}_1^* \xi_n = \phi_n^* \xi$. (Here, we are using Gromov's notation of $\text{Op}(Q)$ from [Gro86] to denote a germ of an open neighborhood of Q .)

We remark that geometric limits of contact structures in the sense of Definition 5.2 are not necessarily unique. However, the possible ambiguities occur only from contact topological information disappearing to infinity under geometric limits.

In this section, we investigate geometric convergence of the contact structures constructed in Section 4. To do this, we shall require some amount of convex surface theory (see, [Hon00a]). We spend some time on this in the next subsection.

5.2. Basic slice and stabilization. Let us fix an integral basis of $H_1(T^2; \mathbb{Z})$. We recall the following definition due to Honda [Hon00a]:

Definition 5.3. [Hon00a, Section 4.3] A *basic slice* is a contact structure ξ^b on $T^2 \times [0, 1]$ such that,

- (1) $(T^2 \times [0, 1], \xi^b)$ is tight,
- (2) $T^2 \times \{i\}$, $i = 0, 1$ are convex surfaces, and the number of dividing curves on each is 2,
- (3) Let s_i be the slopes of the dividing curves on $T^2 \times \{i\}$, for $i = 0, 1$. If $s_i = p_i/q_i$, $i = 0, 1$ denotes the reduced fraction representing the slopes, then $(p_0, q_0), (p_1, q_1)$ forms an integral basis of \mathbb{Z}^2 .
- (4) ξ^b is minimally twisting, i.e., the dividing curves of every ∂ -parallel convex torus in $T^2 \times [0, 1]$ have slope between s_0 and s_1 .

Theorem 5.4. [Hon00a, Proposition 4.7] *There are exactly two basic slices $(T^2 \times I, \xi^b)$ with $s_0 = 0$ and $s_1 = -1$, up to isotopy relative to boundary. They are distinguished by their relative Euler classes which are $(1, 0)$ and $(-1, 0)$ in $H^1(T^2; \mathbb{Z})$, respectively.*

Remark 5.5. We shall say that the *sign* of a basic slice is positive (resp. negative), if the relative Euler class referred to in Theorem 5.4 is $(1, 0)$ (resp. $(-1, 0)$).

Definition 5.6. [Hon00a, Section 4.4.5] Fix $m \geq 1$. A *continued fraction block of length m* is a contact structure ξ_m^{cf} on $T^2 \times [1, m+1]$ such that there exists a decomposition,

$$T^2 \times I = B_1 \cup B_2 \cup \cdots \cup B_m,$$

where $B_i = T^2 \times [i, i+1]$ for all $1 \leq i \leq m$ and $(B_i, \xi_m^{cf}|_{B_i})$ is a basic slice and the dividing slopes on the boundary tori $T^2 \times \{i\}$ of the basic slices are given by $s_i = -i$ for all $1 \leq i \leq m+1$.

We shall give a description of basic slices in terms of stabilization of Legendrian knots. Although this description is well-known to experts (see [Hon00a] for instance), we give a proof for completeness. First we set up some notation.

Let $K \subset (S^3, \xi_{\text{std}})$ be a Legendrian knot in the sphere with the standard contact structure. Suppose $\text{tb}(K) = -n < 0$. Then K has a tubular neighborhood $N(K)$ contactomorphic to $(S^1 \times D^2, \xi)$ where

$$\xi = \ker(\sin(2n\pi z)dx + \cos(2n\pi z)dy)$$

Here, x, y are coordinates of D^2 and z is the coordinate on $S^1 = \mathbb{R}/\mathbb{Z}$. We will call $N(K)$ the *standard neighborhood* of K . The boundary torus $\partial N(K) \cong T^2$ is a convex surface. The characteristic foliation on $\partial N(K)$ consists of two lines of singularities, called the *Legendrian divides*, and is otherwise foliated by the lines $\{y = \text{const.}\}$, called the *ruling curves*. Let μ be a meridian curve $\{pt\} \times \partial D^2$ and λ be one of the ruling curves on $\partial N(K)$. Let us declare (a little unconventionally) that μ has slope 0 and λ has slope ∞ . Then the characteristic foliation has two dividing curves parallel to the Legendrian divides, each of slope $-1/n$. Notice that each ruling curve intersects each dividing curve n times. (See Figure 3.)




FIGURE 3. Characteristic foliation of boundary of standard neighborhood of a Legendrian knot. Ruling curves are in green. Legendrian divides are in red.

Definition 5.7. [Gei08, p. 318] Let $\mathcal{L} \subset (\mathbb{R}^3, \xi)$ be a Legendrian knot in \mathbb{R}^3 equipped with the standard contact structure ξ . Let $\pi : \mathbb{R}_{xyz}^3 \rightarrow \mathbb{R}_{xz}^2$ denote the Legendrian front projection. The positive (resp. negative) *stabilization* of \mathcal{L} is obtained from \mathcal{L} by

- (1) first choosing a smooth embedded arc α without crossings on $\pi(\mathcal{L})$,
- (2) modifying the arc α in a compactly supported manner by adding two cusps of positive (resp. negative) sign,
- (3) and finally lifting the modified front projection back to a Legendrian knot in (\mathbb{R}^3, ξ) .

Proposition 5.8. Let $K \subset (S^3, \xi_{\text{std}})$ be a Legendrian knot with $\text{tb}(K) = -n < 0$ and K' denote a stabilization of K . We isotope K' to be contained in a standard neighborhood

$N(K)$ of K . Let $N(K')$ be a standard neighborhood of K' contained in $N(K)$. Let $T = N(K) \setminus N(K')$. Then $(T, \xi_{\text{std}}|_T)$ is a basic slice with sign the same as that of the stabilization performed on K to yield K' .

Proof. Certainly, $(T, \xi_{\text{std}}|_T)$ is tight as it is contactly embedded in the tight contact sphere (S^3, ξ_{std}) . Moreover, $\partial T = \partial N(K) \cup \partial N(K')$ is a convex surface, and each torus component has precisely two dividing curves. Note that the reduced fraction representations $-1/n, -1/(n+1)$ of the slopes of the dividing curves of $\partial N(K), \partial N(K')$ (respectively) form an integral basis of \mathbb{Z}^2 . Therefore, $(T, \xi_{\text{std}}|_T)$ satisfies conditions (1), (2) and (3) in Theorem 5.4. It remains to show $(T, \xi_{\text{std}}|_T)$ satisfies condition (4), i.e., that it is minimally twisting.

Let $A \subset T$ be an annulus transverse to ∂T , such that $\partial A = c \cup c'$ where $c \subset \partial N(K)$ and $c' \subset \partial N(K')$ are a pair of ruling curves. Then $\partial A \subset T$ is Legendrian. Let fr denote the framing on ∂A induced from A . This agrees with the surface framing induced from the tori $\partial N(K)$ and $\partial N(K')$, respectively, as A is transverse to ∂T . Since c (resp. c') intersects the dividing multi-curve of $\partial N(K)$ (resp. $\partial N(K')$) at $2n$ (resp. $2n+2$) points, we obtain

$$\begin{aligned} \text{tw}(c, fr) &= -n \\ \text{tw}(c', fr) &= -(n+1) \end{aligned}$$

As $\text{tw}(c, fr), \text{tw}(c', fr) < 0$, we may perturb A relative to ∂A to be convex. Since $\text{tw}(c', fr) < \text{tw}(c, fr) < 0$, the Imbalance Principle [Hon00a, Proposition 3.17] implies A contains a bypass attached along the boundary component $c' \subset \partial A$.

Therefore, $\partial N(K)$ is obtained from attaching a bypass to $\partial N(K')$ along a ruling curve. By a change of basis we may assume that the dividing curves of $\partial N(K), \partial N(K')$ have slopes -1 and 0 , respectively. In the new basis, the ruling curve $c' \subset \partial N(K')$ along which the bypass is attached has slope $-(n+1) \in (-\infty, -1)$. Therefore, $(T, \xi_{\text{std}}|_T)$ is minimally twisting [Hon00a] (see also, [Etn04, Theorem 5.11]). This concludes the proof that $(T, \xi_{\text{std}}|_T)$ is a basic slice.

To determine the sign of the basic slice $(T, \xi_{\text{std}}|_T)$ (in the sense of Theorem 5.4), we compute the relative Euler class $e(\xi_{\text{std}}|_T)$. Recall the convex annulus $A \subset T$ constructed above. Then,

$$\langle e(\xi_{\text{std}}|_T), [A] \rangle = \chi(R_+) - \chi(R_-),$$

where R_{\pm} are the (\pm) -ve regions of A in the complement of the dividing arcs of A . Such a region is either an annulus, or a disk. If it is an annulus, the Euler characteristic is zero. Each disk region is necessarily bounded by a boundary-parallel arc in A , therefore giving rise to a bypass. Since A contains precisely one bypass, we conclude $\langle e(\xi_{\text{std}}|_T), [A] \rangle = \pm 1$. Whether it is $+1$ or -1 depends on whether the unique disk region is a $(+)$ -ve region or a $(-)$ -ve region. Equivalently, it depends on the sign of the bypass. This is precisely the sign of the stabilization, given by $\text{rot}(K) - \text{rot}(K')$. \square

Remark 5.9. In Definition 5.6, there is an ambiguity in the description of the contact structure on the continued fraction block $(T^2 \times [1, m+1], \xi_m^{cf})$ since each basic slice B_i in the decomposition may have positive or negative sign. However, one can always re-order (or *shuffle*) the basic slices in a continued fraction block by contact isotopies relative

to boundary so that B_1, \dots, B_k are the positive basic slices and B_{k+1}, \dots, B_n are the negative ones. Indeed, this follows from our description of basic slices in Proposition 5.8, as order of positive and negative stabilizations on a Legendrian knot is irrelevant up to isotopy. Therefore, the ambiguity in the description of Definition 5.6 is simply the number $k \in \{0, \dots, m\}$ of positive basic slices in the block.

5.3. Geometric convergence of rational surgeries. In Section 4, we constructed examples of tight contact structures on hyperbolic manifolds obtained from rational surgery on certain hyperbolic braid links. We shall show that appropriate sequences of these geometrically converge to tight contact structures on the link complement, in the sense of Definition 5.2. For ease of exposition, we will only consider braid knots.

5.3.1. The manifolds. Let $\beta \in B_m$ be a braid such that the closure $K := \widehat{\beta}$ is a hyperbolic knot. Assume β satisfies the condition $c_+ - 2c_- - m \geq 1$ (cf. Section 3.1). Let $r < -1$ be a real number, and $r = [a_0, a_1, a_2, \dots]$ denote a negative continued fraction expansion (cf. Definition 4.1). Let $-q_n/p_n$ denote the n -th convergent of r . Let M_n denote the p_n/q_n -surgery on K . Suppose $|p_n| + |q_n| \rightarrow \infty$. For instance, this is the case if r is irrational. Then M_n is a hyperbolic manifold for $n \gg 0$ sufficiently large.

5.3.2. The contact structures on the manifolds. In Theorem 4.4, we constructed

$$\Phi(-q_n/p_n) = |a_0 + 1| |a_1 + 1| \cdots |a_n + 1|$$

nonisotopic contact structures on M_n . Recall that these contact structures are indexed by tuples (k_0, k_1, \dots, k_n) where $1 \leq k_i \leq -a_i - 1$ for all $0 \leq i \leq n$. M_n equipped with such a contact structure admits a Weinstein filling W_n , given by the Weinstein-Kirby surgery presentation

$$K \cup U_0 \cup U_1 \cup \cdots \cup U_n,$$

- (1) K is a Legendrian realization of the closure of β with Thurston-Bennequin number 1,
- (2) U_i is a Legendrian unknot with Thurston-Bennequin number $a_i + 1$ and rotation number $2k_i - a_i$,
- (3) U_0 is the meridian of K ,
- (4) U_i is the meridian of U_{i-1} for all $0 < i \leq n$.

Let $\mathbf{k} = (k_0, k_1, \dots)$ be an infinite tuple such that $1 \leq k_i \leq |a_i + 1|$. We denote $\mathbf{k}(n) = (k_0, k_1, \dots, k_n)$. We denote the contact structure on M_n corresponding to the tuple $\mathbf{k}(n)$ constructed above as $\xi_{r, \mathbf{k}(n)}$.

5.3.3. The contact structures on the limit manifold. Let $M_\infty := S^3 \setminus K$. Let $N(K) \subset S^3$ denote a Legendrian standard neighborhood of K . Let μ, λ denote the meridian and longitude of K , respectively. Let us fix an integral basis of $H_1(T^2; \mathbb{Z})$. We write:

$$M_\infty = S^3 \setminus N(K) \cup T^2 \times [-1, 0] \cup T^2 \times [0, 1] \cup T^2 \times [1, 2] \cup \cdots,$$

where for all $i \geq 0$, $T^2 \times [i, i+1]$ is glued to $T^2 \times [i-1, i]$ along $T^2 \times \{i\}$ by the linear diffeomorphism,

$$T^2 \times [i, i+1] \supset T^2 \times \{i\} \xrightarrow{\phi_i} T^2 \times \{i\} \subset T^2 \times [i-1, i]$$

$$\phi_i = \begin{pmatrix} 0 & -1 \\ 1 & -a_i \end{pmatrix},$$

and $T^2 \times \{-1\}$ is glued to $\partial N(K)$ by sending $(1, 0)$ to $-\lambda$ and $(0, 1)$ to μ . We define a contact structure $\xi_{r, \mathbf{k}}$ on M_∞ such that

- (1) $\xi_{r, \mathbf{k}}$ agrees with ξ_{std} on $S^3 \setminus N(K)$, where ξ_{std} is the standard contact structure on S^3 ,
- (2) For all $i \geq 0$, $T^2 \times [i-1, i]$ is a continued fraction block of length $|a_i + 2|$ (cf. Definition 5.6), consisting of $k_i - 1$ positive basic slices appearing in any order (cf. Remark 5.9).

This can be achieved since the slopes of the dividing curves of consecutive continued fraction blocks match upon applying the diffeomorphism φ_i . Hence, the contact structures glue consistently.

5.3.4. Geometric convergence. We are now prepared to state the main result:

Theorem 5.10. *Let $r < -1$ be a real number with a negative continued fraction expansion $r = [a_0, a_1, a_2, \dots]$, and let $-q_n/p_n$ denote the convergents. Suppose $|p_n| + |q_n| \rightarrow \infty$ as $n \rightarrow \infty$. Let $\mathbf{k} = (k_0, k_1, k_2, \dots)$ be an infinite tuple such that $1 \leq k_i \leq |a_i + 1|$. Let $\mathbf{k}(\mathbf{n}) = (k_0, k_1, k_2, \dots, k_n)$, and $\xi_{r, \mathbf{k}(n)}$ denote the contact structure on M_n constructed in Section 5.3.2. Then, $(M_n, \xi_{r, \mathbf{k}(n)})$ geometrically converges to $(M_\infty, \xi_{r, \mathbf{k}})$.*

Proof. Recall that M_n is obtained by a p_n/q_n -Dehn surgery on the hyperbolic knot $K := \widehat{\beta}$, and $M_\infty = S^3 \setminus K$ is the knot complement. Let $\varepsilon > 0$ be sufficiently small. Then for all n sufficiently large, M_n admits a ε -thick-thin decomposition,

$$M_n = \text{Thick}_\varepsilon(M_n) \cup \text{Thin}_\varepsilon(M_n),$$

such that $\text{Thin}_\varepsilon(M_n)$ is a tubular neighborhood of the systole of M_n given by core of the surgery solid torus, i.e., a ε -thin Margulis tube. For any given $\varepsilon > 0$, there exists $N(\varepsilon) \gg 0$ such that for all $n \geq N(\varepsilon)$, there exists a $(1 + o(\varepsilon))$ -Lipschitz diffeomorphism,

$$\phi_{n, \varepsilon} : \text{Thick}_\varepsilon(M_n) \rightarrow \text{Thick}_\varepsilon(M_\infty).$$

Thus, the sequence of hyperbolic manifolds M_n geometrically converges to M_∞ .

For convenience of the reader we recall some relevant terminology. A *Weinstein 2-handle* is the standard $D^2 \times D^2 \subset \mathbb{R}^4$ equipped with

- (1) the standard symplectic form $\omega_{\text{std}} = dx_1 \wedge dy_1 + dx_2 \wedge dy_2$ induced from \mathbb{R}^4 ,
- (2) the Liouville vector field

$$X = -x_1 \frac{\partial}{\partial x_1} - x_2 \frac{\partial}{\partial x_2} + 2y_1 \frac{\partial}{\partial y_1} + 2y_2 \frac{\partial}{\partial y_2}.$$

The regions $S^1 \times D^2$ and $D^2 \times S^1$ are called the *attaching* and *co-attaching regions* respectively. The circles $S^1 \times \{0\}$ and $\{0\} \times S^1$ are called the *core circle* and *belt circle*

respectively. Since $(D^2 \times D^2, \omega_{std}, X)$ is a Weinstein manifold-with-corners, the attaching and co-attaching regions $S^1 \times D^2$ and $D^2 \times S^1$ inherit canonical induced contact structures.

Recall the Weinstein filling W_n of M_n described in Section 5.3.2, given by the Weinstein-Kirby presentation $K \cup U_0 \cup U_1 \cup \cdots \cup U_n$. Let h_K^2 and $h_{U_i}^2$ ($0 \leq i \leq n$) denote the Weinstein 2-handles in W_n corresponding to the components of this link. We smoothly slide the 2-handle $h_{U_n}^2$ over the boundary $h_{U_{n-1}}^2$ so that the attaching circle of $h_{U_n}^2$ is the belt circle of $h_{U_{n-1}}^2$. The effect on the boundary of the 4-manifold is the same as the slam-dunk isotopy [GS99, Section 5.3., p. 163] pushing U_n inside the surgery solid torus of U_{n-1} . Similarly, we inductively slide $h_{U_i}^2$ over $h_{U_{i-1}}^2$ for all $1 \leq i \leq n$, and finally slide $h_{U_0}^2$ over h_K^2 . Taking boundary, we obtain the decomposition:

$$(5.3.1) \quad M_n \cong (S^3 \setminus N(K)) \cup T^2 \times [-1, 0] \cup T^2 \times [0, 1] \cup \cdots \cup T^2 \times [n-1, n] \cup S^1 \times D^2$$

Here,

- (1) $N(K)$ is a fixed standard neighborhood of the Legendrian knot $K \subset (S^3, \xi_{std})$,
- (2) $T^2 \times [-1, 0]$ is the co-attaching region of h_K^2 with a neighborhood of the belt circle removed. Or equivalently, the surgery solid torus for K with a neighborhood of the core circle removed.
- (3) $T^2 \times [i, i+1]$, $0 \leq i \leq n-1$ is the co-attaching region of $h_{U_i}^2$ after the slide, with a neighborhood of the belt circle removed. Or equivalently, the surgery solid torus of U_i after the slam-dunk isotopy, with a neighborhood of the core circle removed.
- (4) $S^1 \times D^2$ is the co-attaching region of $h_{U_n}^2$ after the slide. Or equivalently, the surgery solid torus of U_n after the slam-dunk isotopy.

We may imagine the hyperbolic structure on M_n as the hyperbolic manifold-with-boundary $S^3 \setminus N(K)$ and a Margulis tube given by the solid torus in Item (4), connected by a “long, narrow neck” $T^2 \times [-1, n]$. Observe that pulling back $\xi_{r, k(n)}$ by the handleslide/slam-dunk isotopies, we get a contact structure which agrees with the contact structure on $S^3 \setminus N(K)$ induced from (S^3, ξ_{std}) . Furthermore, the contact structure on the solid torus in Item (4) is the canonical contact structure on the co-attaching boundary of a Weinstein 2-handle. Thus, convergence of the contact structures $\xi_{r, k(n)}$ on M_n need only be verified on the neck region.

We proceed to carry this out more precisely. Let $M_\infty = S^3 \setminus K$ be hyperbolic, so that $M_\infty = \mathbb{H}^3/\Gamma$. Then an embedded incompressible torus in M_∞ will be called a *horotorus* if it is the quotient of a horosphere by a $\mathbb{Z} \oplus \mathbb{Z}$ stabilizing it. With respect to the complete hyperbolic structure on $M_\infty = S^3 \setminus K$, fix now a horotorus $T_{-1} \subset M_\infty$. Let $T_n \subset M_\infty$ be a horotorus such that the unbounded component of $M_\infty \setminus T_n$ is exactly the $1/n$ -thin part of M_∞ . Let T_0, \dots, T_{n-1} be such that T_{-1}, T_0, \dots, T_n are a collection of equidistant, parallel horotori. Since (M_n) converges to M_∞ , after possibly passing to a subsequence of (M_n) we can find a $(1 + o(1/n))$ -Lipschitz diffeomorphism

$$\varphi_n := \phi_{n, 1/n} : \text{Thick}_{1/n}(M_n) \rightarrow \text{Thick}_{1/n}(M_\infty)$$

We isotope the contact structure $\xi_{r, k(n)}$ on M_n so that in Equation 5.3.1, the tori $T^2 \times \{i\}$ are precisely $\varphi_n^{-1}(T_i)$, for $-1 \leq i \leq n$.

We equip M_∞ with the contact structure $\xi_{r, k}$ constructed in Section 5.3.3, where the domains bounded by the horotori T_{i-1}, T_i ($i \geq 0$) are continued fraction blocks of length

$|a_i + 2|$ containing $k_i - 1$ positive basic slices. By Proposition 5.8, φ_n is a contactomorphism. Therefore, the diffeomorphisms φ_n certify the geometric convergence of $(M_n, \xi_{r, \mathbf{k}(n)})$ to $(M_\infty, \xi_{r, \mathbf{k}})$. \square

5.3.5. Uncountably many tight contact limits. In Tripp [Tri06], it is shown that there are uncountably many tight contact structures on an irreducible open three-manifold which are pairwise distinct up to contactomorphism, provided the manifold possesses an end of nonzero genus. In this section we exhibit a family of such examples which also naturally arise as geometric limits of contact structures on closed three-manifolds. We begin with the following notion due to [Tri06]:

Definition 5.11. Two contact structures ξ, ξ' on a three-manifold M are said to be *properly isotopic* if there exists an isotopy of M taking ξ to ξ' . More precisely, there exists a smooth homotopy of diffeomorphisms $\{f_t : M \rightarrow M\}$ such that $f_0 = \text{id}$, $f_1^* \xi' = \xi$.

As Gray's stability theorem fails on noncompact manifolds, this is distinct from the notion of an isotopy of contact plane fields in general. Next, we recall the notion of *slope at infinity* introduced in [Tri06], which is an invariant of contact structures on open three-manifolds up to proper isotopy. We specialize the definition to the situation of a manifold with toric end, for simplicity.

Definition 5.12. [Tri06, Section 3] Let (M, ξ) be a contact three-manifold, where M is the interior of a compact three-manifold \overline{M} with torus boundary $\partial \overline{M} \cong T^2$. Fix an integral basis of $H_1(\partial M; \mathbb{Z})$. We say an embedded convex torus $T \subset M$ is *end-parallel* if $T \subset \overline{M}$ is isotopic to $\partial \overline{M}$. Let $T \times [0, \infty) \subset M$ be the unbounded component bounded by such a torus. We call the contact manifold $E = (T \times [0, \infty), \xi)$ together with its embedding in M , a *contact end* of M . The set of contact ends denoted $\text{Ends}(M, \xi)$ is a directed set, directed by reverse inclusion.

Fix an integral basis of $H_1(\partial M; \mathbb{Z})$. For any end-parallel convex torus T , we obtain a corresponding integral basis of $H_1(T; \mathbb{Z})$. We denote the slope of the dividing curves of T measured against this basis as $\text{slope}(T)$. For any contact end E of M , we define the slope of E as

$$\text{slope}(E) = \sup_T \text{slope}(T),$$

where T varies over all end-parallel convex tori contained in E . Then $\text{slope} : \text{Ends}(M, \xi) \rightarrow \mathbb{R} \cup \{\infty\}$ defines a net. If the net is convergent, we define the *slope at infinity* of M to be the limit.

Definition 5.13. Let $r < -1$ be a real number with negative continued fraction expansion $r = [a_0, a_1, a_2, \dots]$. Let $\mathbf{k} = (k_0, k_1, k_2, \dots)$ be an infinite tuple such that $1 \leq k_i \leq |a_i + 1|$. We shall say the *sign* of \mathbf{k} is $+$ (resp. $-$) if for all $i \geq 0$ sufficiently large, $k_i = |a_i + 1|$ (resp. $k_i = 1$). Otherwise, we say sign of \mathbf{k} is \pm .

We have the following corollary of Theorem 5.10:

Corollary 5.14. Let $\xi_{r, \mathbf{k}}$ denote the contact structures on M_∞ constructed in Section 5.3.3. Then,

- (1) $\xi_{r, \mathbf{k}}$ is a tight contact structure on M_∞ ,

- (2) $\xi_{r,\mathbf{k}}$ and $\xi_{r',\mathbf{k}'}$ are properly isotopic if and only if $r = r'$ and \mathbf{k}, \mathbf{k}' are of the same sign.

Proof.

- (1) Suppose $(M_\infty, \xi_{r,\mathbf{k}})$ was overtwisted. Then it would contain an overtwisted disk D_{ot} . By compactness of the disk, there exists $n \gg 0$ large such that D_{ot} is entirely contained inside $\text{Thick}_{1/n}(M_\infty)$. Recall the diffeomorphism $\varphi_n : \text{Thick}_{1/n}(M_n) \rightarrow \text{Thick}_{1/n}(M_\infty)$ from the Proof of Theorem 5.10. Thus, $\varphi_n^{-1}(D_{ot}) \subset \text{Thick}_{1/n}(M_n)$ is an overtwisted disk in $\varphi_n^* \xi_{r,\mathbf{k}}$. As $\varphi_n^* \xi_{r,\mathbf{k}}$ is isotopic to $\xi_{r,\mathbf{k}(n)}$, we conclude $(M_n, \xi_{r,\mathbf{k}(n)})$ is also overtwisted. This is a contradiction.
- (2) As $\xi_{r,\mathbf{k}}$ is tight, by [Tri06, Proposition 4.1] the slope at infinity exists. By taking the cofinal sequence of ends bounded by $T^2 \times \{i\}$ in Section 5.3.3, we see that the slope at infinity is r (after choosing an appropriate reference homology basis for the toric end). Finally, observe that if \mathbf{k}, \mathbf{k}' are of the same sign, then an isotopy between $\xi_{r,\mathbf{k}}, \xi_{r,\mathbf{k}'}$ can be constructed by shuffling the positive and negative basic blocks (cf. Remark 5.9) \square

The following Corollary shows that the notion of geometric convergence of contact structures in Definition 5.2 is compatible with Tripp's work in [Tri06].

Corollary 5.15. *M_∞ admits uncountably many tight contact structures up to contactomorphism arising as geometric limits of tight contact structures on M_n .*

Proof. By Corollary 5.14, M_∞ admits uncountably many tight contact structures up to proper isotopy. Note that $M_\infty = S^3 \setminus K$ is a knot complement. Observe that the smooth mapping class group of the open manifold M_∞ agrees with the smooth mapping class group (not relative to the boundary) of the thickened knot complement $S^3 \setminus \nu(K)$. Indeed, any diffeomorphism of M_∞ must send an end-parallel torus $T^2 \subset M_\infty$ to an isotopic copy of itself. After isotopy, we may ensure the diffeomorphism preserves this torus. We cut along this torus and observe that the smooth mapping class group $\text{MCG}_\partial(T^2 \times [0, \infty))$, relative to $T^2 \times \{0\}$, is trivial.

Since the thickened knot complement is oriented Haken, a theorem of Waldhausen (see, for instance, Hempel [Hem04, Corollary 13.7]) implies the smooth mapping class group $\text{MCG}(S^3 \setminus \nu(K))$ is isomorphic to the group of self-homotopy equivalences $\text{hAut}(S^3 \setminus \nu(K))$. Since K is hyperbolic, the latter is a finite group by the Mostow rigidity theorem. Therefore, $\text{MCG}(M_\infty)$ is finite.

Thus, we must have uncountably many mapping class orbits of tight contact structures, as desired. \square

REFERENCES

- [Ago13] Ian Agol. The virtual Haken conjecture. *Doc. Math.*, 18:1045–1087, 2013. With an appendix by Agol, Daniel Groves, and Jason Manning. [2](#)
- [AS17] M. Firat Arikan and Merve Seçgin. Tight contact structures on hyperbolic three-manifolds. *Topology Appl.*, 231:345–352, 2017. [2](#)
- [BM24] Thomas Barthelmé and Kathryn Mann. Orbit equivalences of \mathbb{R} -covered Anosov flows and hyperbolic-like actions on the line. *Geom. Topol.*, 28(2):867–899, 2024. [3](#)

- [Car17] Sandrine Caruso. On the genericity of pseudo-Anosov braids I: rigid braids. *Groups Geom. Dyn.*, 11(2):533–547, 2017. [11](#)
- [CGH09] Vincent Colin, Emmanuel Giroux, and Ko Honda. Finitude homotopique et isotopique des structures de contact tendues. *Publ. Math. Inst. Hautes Études Sci.*, (109):245–293, 2009. [2](#)
- [CM20] James Conway and Hyunki Min. Classification of tight contact structures on surgeries on the figure-eight knot. *Geom. Topol.*, 24(3):1457–1517, 2020. [2](#)
- [CW17] Sandrine Caruso and Bert Wiest. On the genericity of pseudo-Anosov braids II: conjugations to rigid braids. *Groups Geom. Dyn.*, 11(2):549–565, 2017. [11](#)
- [EH01] John B. Etnyre and Ko Honda. On the nonexistence of tight contact structures. *Ann. of Math. (2)*, 153(3):749–766, 2001. [2](#)
- [Eli89] Y. Eliashberg. Classification of overtwisted contact structures on 3-manifolds. *Invent. Math.*, 98(3):623–637, 1989. [1](#)
- [Eli90a] Yakov Eliashberg. Filling by holomorphic discs and its applications. In *Geometry of low-dimensional manifolds, 2 (Durham, 1989)*, volume 151 of *London Math. Soc. Lecture Note Ser.*, pages 45–67. Cambridge Univ. Press, Cambridge, 1990. [2](#), [14](#)
- [Eli90b] Yakov Eliashberg. Topological characterization of Stein manifolds of dimension > 2 . *Internat. J. Math.*, 1(1):29–46, 1990. [12](#)
- [Eli92] Yakov Eliashberg. Contact 3-manifolds twenty years since J. Martinet’s work. *Ann. Inst. Fourier (Grenoble)*, 42(1-2):165–192, 1992. [2](#)
- [ET98] Yakov M. Eliashberg and William P. Thurston. *Confoliations*, volume 13 of *University Lecture Series*. American Mathematical Society, Providence, RI, 1998. [2](#)
- [Etg12] Tolga Etgü. Tight contact structures on laminar free hyperbolic three-manifolds. *Int. Math. Res. Not. IMRN*, (20):4775–4784, 2012. [2](#)
- [Etn04] John B. Etnyre. Convex surfaces in contact geometry: class notes. 2004. Available at <https://etnyre.math.gatech.edu/preprints/papers/surfaces.pdf>. [21](#)
- [Gab83] David Gabai. Foliations and the topology of 3-manifolds. *J. Differential Geom.*, 18(3):445–503, 1983. [2](#)
- [Gei08] Hansjörg Geiges. *An introduction to contact topology*, volume 109 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2008. [20](#)
- [GMT03] David Gabai, G. Robert Meyerhoff, and Nathaniel Thurston. Homotopy hyperbolic 3-manifolds are hyperbolic. *Ann. of Math. (2)*, 157(2):335–431, 2003. [15](#)
- [Gom98] Robert E. Gompf. Handlebody construction of Stein surfaces. *Ann. of Math. (2)*, 148(2):619–693, 1998. [2](#), [12](#), [15](#), [16](#)
- [Gro85] M. Gromov. Pseudo holomorphic curves in symplectic manifolds. *Invent. Math.*, 82(2):307–347, 1985. [2](#), [14](#)
- [Gro86] Mikhael Gromov. *Partial differential relations*, volume 9 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1986. [19](#)
- [GS99] Robert E. Gompf and András I. Stipsicz. *4-manifolds and Kirby calculus*, volume 20 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1999. [12](#), [24](#)
- [Hem04] John Hempel. *3-manifolds*. AMS Chelsea Publishing, Providence, RI, 2004. Reprint of the 1976 original. [26](#)
- [HKM03] Ko Honda, William H. Kazez, and Gordana Matić. Tight contact structures on fibered hyperbolic 3-manifolds. *J. Differential Geom.*, 64(2):305–358, 2003. [2](#)
- [Hon00a] Ko Honda. On the classification of tight contact structures. I. *Geom. Topol.*, 4:309–368, 2000. [2](#), [4](#), [19](#), [20](#), [21](#)
- [Hon00b] Ko Honda. On the classification of tight contact structures. II. *J. Differential Geom.*, 55(1):83–143, 2000. [2](#), [4](#)
- [Ito11] Tetsuya Ito. Braid ordering and the geometry of closed braid. *Geom. Topol.*, 15(1):473–498, 2011. [3](#), [11](#)
- [Kap01] M. Kapovich. Hyperbolic Manifolds and Discrete Groups. *Progress in Mathematics, vol. 183*. Birkhauser, 2001. [6](#)
- [kir97] Problems in low-dimensional topology. In Rob Kirby, editor, *Geometric topology (Athens, GA, 1993)*, volume 2.2 of *AMS/IP Stud. Adv. Math.*, pages 35–473. Amer. Math. Soc., Providence, RI, 1997. [2](#)

- [KM97] P. B. Kronheimer and T. S. Mrowka. Monopoles and contact structures. *Invent. Math.*, 130(2):209–255, 1997. [14](#)
- [LL19] Youlin Li and Yajing Liu. Hyperbolic 3-manifolds admitting no fillable contact structures. *Proc. Amer. Math. Soc.*, 147(1):351–360, 2019. [2](#)
- [LM97] P. Lisca and G. Matić. Tight contact structures and Seiberg-Witten invariants. *Invent. Math.*, 129(3):509–525, 1997. [3](#), [14](#), [16](#)
- [LS09] Paolo Lisca and András I. Stipsicz. On the existence of tight contact structures on Seifert fibered 3-manifolds. *Duke Math. J.*, 148(2):175–209, 2009. [2](#)
- [Ota98] J. P. Otal. 'Thurston's hyperbolization of Haken manifolds. ' *Surveys in differential geometry, Vol. III Cambridge, MA, Int. Press, Boston, MA*, pages 77–194, 1998. [6](#)
- [Ota01] Jean-Pierre Otal. *The hyperbolization theorem for fibered 3-manifolds*, volume 7 of *SMF/AMS Texts and Monographs*. American Mathematical Society, Providence, RI; Société Mathématique de France, Paris, 2001. Translated from the 1996 French original by Leslie D. Kay. [4](#)
- [Thu86] William P. Thurston. Hyperbolic structures on 3-manifolds, ii: Surface groups and 3-manifolds which fiber over the circle. *arXiv preprint*, 1986. arXiv:1410.5093. [4](#)
- [Thu22] William P. Thurston. *The geometry and topology of three-manifolds. Vol. IV*. American Mathematical Society, Providence, RI, [2022] ©2022. Edited and with a preface by Steven P. Kerckhoff and a chapter by J. W. Milnor. [11](#), [12](#), [15](#)
- [Tri06] James J. Tripp. Contact structures on open 3-manifolds. *J. Symplectic Geom.*, 4(1):93–116, 2006. [4](#), [25](#), [26](#)
- [Wis21] Daniel T. Wise. *The structure of groups with a quasiconvex hierarchy*, volume 209 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, [2021] ©2021. [2](#)