

α_i -Metric Graphs: Hyperbolicity ^{*}

Feodor F. Dragan¹ and Guillaume Ducoffe²

¹ Computer Science Department, Kent State University, Kent, USA dragan@cs.kent.edu

² National Institute for Research and Development in Informatics and
University of Bucharest, București, România guillaume.ducoffe@ici.ro

Abstract. A graph is called α_i -metric ($i \in \mathcal{N}$) if it satisfies the following α_i -metric property for every vertices u, w, v and x : if a shortest path between u and w and a shortest path between x and v share a terminal edge vw , then $d(u, x) \geq d(u, v) + d(v, x) - i$. The latter is a discrete relaxation of the property that in Euclidean spaces the union of two geodesics sharing a terminal segment must be also a geodesic. Recently in (Dragan & Ducoffe, *WG'23*) we initiated the study of the algorithmic applications of α_i -metric graphs. Our results in this prior work were very similar to those established in (Chepoi et al., *SoCG'08*) and (Chepoi et al., *COCOA'18*) for graphs with bounded *hyperbolicity*. The latter is a heavily studied metric tree-likeness parameter first introduced by Gromov. In this paper, we clarify the relationship between hyperbolicity and the α_i -metric property, proving that α_i -metric graphs are $f(i)$ -hyperbolic for some function f linear in i . We give different proofs of this result, using various equivalent definitions to graph hyperbolicity. By contrast, we give simple constructions of 1-hyperbolic graphs that are not α_i -metric for any constant i . Finally, in the special case of $i = 1$, we prove that α_1 -metric graphs are 1-hyperbolic, and the bound is sharp. By doing so, we can answer some questions left open in (Dragan & Ducoffe, *WG'23*).

Keywords: metric graph classes; α_i -metric; hyperbolicity.

1 Introduction

For any undefined graph terminology, see [7]. In what follows, we only consider graphs $G = (V, E)$ that are finite, undirected, unweighted, simple and connected. The distance $d_G(u, v)$ between two vertices $u, v \in V$ is the minimum length (number of edges) of a path between u and v in G . The interval $I_G(u, v)$ between u and v contains every vertex on a shortest (u, v) -path, *i.e.*, $I_G(u, v) = \{w \in V : d_G(u, v) = d_G(u, w) + d_G(w, v)\}$. We will omit the subscript if G is clear from the context. In this paper, we compare two families of metric properties on graphs, defined on quadruples of vertices u, v, w, x as follows for every $i \in \mathcal{N}$:

α_i -metric property: if $v \in I(u, w)$ and $w \in I(v, x)$ are adjacent, then

$$d(u, x) \geq d(u, v) + d(v, x) - i = d(u, v) + 1 + d(w, x) - i;$$

β_i -metric property: if $d(u, v) + d(w, x) \leq d(u, w) + d(v, x) \leq d(u, x) + d(v, w)$, then

$$d(u, x) + d(v, w) - d(u, w) - d(v, x) \leq i.$$

A graph is called α_i -metric if it satisfies the α_i -metric property for every four vertices u, v, w, x . It is called δ -hyperbolic if it satisfies the $\beta_{2\delta}$ -metric property for every four vertices u, v, w, x . The hyperbolicity of G is the smallest half-integer δ such that it is δ -hyperbolic. Roughly, in an α_i -metric graph G , the union of two shortest paths with a common terminal edge results in an “almost shortest” path with defect at most i . By comparison, the less intuitive parameter

^{*} This work was supported by a grant of the Romanian Ministry of Research, Innovation and Digitalization, CCCDI - UEFISCDI, project number PN-III-P2-2.1-PED-2021-2142, within PNCDI III.

hyperbolicity represents the smallest possible distortion for embedding any four vertices of G in a weighted tree [27]. Both the α_0 -metric property and the β_0 -metric property are satisfied by trees. In [3], it was observed that if there exists an embedding of a graph G in some weighted tree T with distortion at most η , then G must be $\alpha_{\lfloor 3\eta \rfloor}$ -metric and $\frac{\lfloor 4\eta \rfloor}{2}$ -hyperbolic. This makes graphs satisfying β_i -metric property as well as graphs satisfying α_i -metric property important in Metric Graph Theory.

Related work. The α_1 -metric property was first considered in [14,15] for chordal graphs. Then, the general α_i -metric property ($i \in \mathcal{N}$) was introduced and studied in [45]. It was proved that all chordal graphs [14] and all plane triangulations with inner vertices of degree at least seven [25] are α_1 -metric. All distance-hereditary graphs [45], and even more strongly, all HHD-free graphs [16], are α_2 -metric. The α_0 -metric graphs are exactly the ptolemaic graphs, *i.e.* the chordal distance-hereditary graphs [33]. Furthermore, a characterization of α_1 -metric graphs was given in [45]: they are exactly the graphs with convex disks and one forbidden isometric subgraph W_6^{++} (see Fig. 6). More recently, additional properties of α_1 -metric graphs and α_i -metric graphs ($i \in \mathcal{N}$) were reported in [22,24,25]. In [22], we presented the first algorithmic applications of α_i -metric graphs to classical distance problems such as diameter, radius and all eccentricities computations. More specifically, all vertex eccentricities in an α_i -metric graph can be approximated in linear time up to some additive term in $\mathcal{O}(i)$. Furthermore, there exists a subquadratic-time algorithm for exact computation of the radius of an α_1 -metric graph.

Hyperbolicity was introduced by Gromov in his study on automatic groups [31]. Since then, the study of δ -hyperbolic graphs has become an important topic in Metric Graph Theory [4]. This parameter has attracted further attention in Network Science, both as a way to better classify complex networks [1,36] and to explain some of their properties such as core congestion [20]. Many real-world networks have small hyperbolicity [1,36,38]. See also [2,6,26,35] for other related results on the hyperbolicity. Many different approaches have been proposed in order to upper bound the hyperbolicity in some graph classes [23,37,44]. In particular, chordal graphs are 1-hyperbolic, and the chordal graphs with hyperbolicity strictly less than one can be characterized with two forbidden isometric subgraphs [8]. The 0-hyperbolic graphs are exactly the block graphs, *i.e.* the graphs such that every biconnected component is a clique [34]. Characterizations of $\frac{1}{2}$ -hyperbolic graphs were given in [3,21]. Furthermore, the algorithmic applications of δ -hyperbolic graphs have been studied much earlier than for α_i -metric graphs [17,18,19]. In [18,19] it was proved that all vertex eccentricities in a δ -hyperbolic graph can be approximated in linear time up to some additive term in $\mathcal{O}(\delta)$. Therefore, it can be argued that α_i -metric graphs and δ -hyperbolic graphs have very similar algorithmic properties.

Little is known about the relationships between α_i -metric properties and hyperbolicity. On one hand, the authors of [3] observed that every 0-hyperbolic graph must be α_0 -metric, and every $\frac{1}{2}$ -hyperbolic graph must be α_1 -metric. However, for any positive integer i , there exists a 1-hyperbolic graph that is not α_i -metric (we give a simple construction in Section 3). On the other hand, the authors of [9] briefly discussed the hyperbolicity and the α_i -metric property for geodesic metric spaces. They observed that Euclidean spaces must be α_0 -metric (*i.e.*, because the union of two geodesics that overlap on one terminal segment must be also a geodesic) whereas they have unbounded hyperbolicity. However, they also noted that “for graphs, the links between δ -hyperbolic graphs and graphs with α_i -metrics are less clear”.

Our Contributions. The main result in the paper is that α_i -metric graphs are $f(i)$ -hyperbolic, for some function f that only depends on i (Theorem 2). Throughout Section 3 we give different proofs of this result, one direct proof in Section 3.4 and other indirect proofs in Section 3.1- 3.3 where we compare the α_i -metric property with some graph parameters that are different, but functionally equivalent, to hyperbolicity. By doing so, we obtain arguably simpler proofs, and sharper relations between the α_i -metric property and these other parameters than by using their known relationships with hyperbolicity. Our current best estimate on function f is linear, but it is probably not tight. We conjecture the right upper bound to be $f(i) = \frac{i+1}{2}$, that would be sharp. In Section 4, we prove this conjecture if $i \leq 1$. Some consequences of our results are further discussed. Some preliminary results are recalled in Section 2. We conclude this paper in Section 5, where we introduce an intriguing generalization of both α_i -metric graphs and δ -hyperbolic graphs.

2 Preliminaries

In what follows, for every two vertices u and v in a graph $G = (V, E)$, we write $u \sim v$ if and only if u and v are adjacent. For every vertex v in a graph $G = (V, E)$, its open and closed neighbourhoods are defined as $N(v) = \{u \in V : u \sim v\}$ and $N[v] = N(v) \cup \{v\}$. The disk of center v and radius r is defined as $D(v, r) = \{u \in V : d(u, v) \leq r\}$. The following variation of Lemma 4 in [22] is used in our proofs:

Lemma 1. *Let $D(u, r_u), D(v, r_v), D(w, r_w)$ be pairwise intersecting disks of G . If G is an α_i -metric graph, then there exists a vertex x such that $d(u, x) \leq r_u$, $d(v, x) \leq r_v$ and $d(w, x) \leq r_w + i$.*

Proof. Let $x \in D(u, r_u) \cap D(v, r_v)$ be such that $d(w, x)$ is minimized. Suppose, by way of contradiction, that $d(x, w) > r_w + i$, and let $y \in N(x) \cap I(x, w)$ be arbitrary. By minimality of $d(x, w)$, we must have $d(u, y) > r_u$ or $d(v, y) > r_v$. By symmetry, we may assume that $d(u, y) > r_u$. It implies $d(u, w) \geq d(u, x) + 1 + d(y, w) - i \geq r_u + 1 + r_w > d(u, w)$, giving a contradiction. \square

Let u and v be arbitrary vertices in G . Recall that $I(u, v)$ contains every vertex on a shortest (u, v) -path. Let also $I_G^o(u, v) = I_G(u, v) \setminus \{u, v\}$. For every integer k such that $0 \leq k \leq d(u, v)$, we define the slice $S_k(u, v) = \{x \in I(u, v) : d(u, x) = k\}$. If graph G is not clear from the context, we will write $S_k(u, v, G)$. Let $\kappa(u, v)$ denote the maximum diameter of a slice between u and v : $\kappa(u, v) = \max_{0 \leq k \leq d(u, v)} \max\{d(x, y) : x, y \in S_k(u, v)\}$. We define the interval thinness of G as $\kappa(G) = \max\{\kappa(u, v) : u, v \in V\}$.

Lemma 2 ([22]). *Let G be an α_i -metric graph, and let u, v, x, y be its vertices such that $x \in I(u, v)$, $d(u, x) = d(u, y)$, and $d(v, y) \leq d(v, x) + k$. Then, $d(x, y) \leq k + i + 2$.*

Lemma 3 ([22]). *If $G = (V, E)$ is an α_i -metric graph, then its interval thinness is at most $i + 1$.*

A geodesic triangle $\Delta(u, v, w) = P(u, v) \cup P(v, w) \cup P(w, u)$ is the union of a shortest (u, v) -path $P(u, v)$, a shortest (v, w) -path $P(v, w)$ and a shortest (w, u) -path $P(w, u)$. Note that $P(u, v), P(v, w), P(w, u)$ are called the sides of the triangle, and they may not be disjoint.

A metric triangle is a triple uvw such that $I^o(u, v), I^o(v, w)$ and $I^o(w, u)$ are pairwise disjoint. Assuming further $d(u, v) \leq d(v, w) \leq d(w, u)$, the type of this triangle is the triple $(d(u, v), d(v, w), d(w, u))$. Let also $d(w, u)$ be the maximum side-length of the triangle. Finally, given a triple $u, v, w \in V$, a quasi-median is a metric triangle $u'v'w'$ such that we have:

- $d(u, v) = d(u, u') + d(u', v') + d(v', v)$;
- $d(v, w) = d(v, v') + d(v', w') + d(w', w)$;
- $d(w, u) = d(w, w') + d(w', u') + d(u', u)$.

Every triple has at least one quasi median (*e.g.*, see [18]).

Other notations and terminology are locally defined at appropriate places throughout the paper.

3 Hyperbolicity of α_i -metric graphs for arbitrary $i \geq 0$

3.1 Using Interval Thinness

It is easy to prove that every δ -hyperbolic graph has interval thinness at most 2δ . Conversely, odd cycles are examples of graphs with interval thinness equal to zero (they are so-called “geodetic graphs”) but unbounded hyperbolicity. However, let the 1-subdivision graph $\Sigma(G)$ of a graph G be obtained by replacing all its edges $e = uv$ by internally vertex-disjoint paths $[u, e, v]$ of length two. Papasoglu [41] proved that the hyperbolicity of G is at most doubly exponential in the interval thinness of $\Sigma(G)$.

By Lemma 3, the interval thinness of α_i -metric graphs is at most $i + 1$. Therefore, in order to prove that α_i -metric graphs are $f(i)$ -hyperbolic, for some arbitrary f , it would be sufficient to prove that their 1-subdivision graphs are $\alpha_{g(i)}$ -metric, for some g . Unfortunately, this is false, even for $i = 1$, as shown in Fig. 1. Our counter-example is a triangular $(2, n)$ -grid graph G . This graph is α_1 -metric (*e.g.*, see Theorem 3). However, $2n - 1 = d_{\Sigma(G)}(u, e) < d_{\Sigma(G)}(u, w)$ and $2n - 2 = d_{\Sigma(G)}(x, w) < d_{\Sigma(G)}(x, e)$ imply that the smallest i such that $\Sigma(G)$ is α_i -metric must satisfy $2 \geq 4n - 2 - i$, and so, $i \geq 4n - 4$.




Fig. 1. The triangular $(2, n)$ -grid is α_1 -metric, but its 1-subdivision graph is not α_i -metric for any $i \leq 4n - 5$.

We now give a direct proof that 1-subdivision graphs of α_i -metric graphs have interval thinness at most linear in i .

Lemma 4. *If $G = (V, E)$ is an α_i -metric graph, then $\kappa(\Sigma(G)) \leq 2i + 12$.*

Proof. Let $s, t \in V \cup E$ be arbitrary. Since $\Sigma(G)$ is bipartite with partite sets V and E , for every two consecutive slices $S_k(s, t)$ and $S_{k+1}(s, t)$, there is one slice fully in V and one slice fully in E . In particular, $\kappa(s, t)$ is at most two units more than the maximum diameter of a slice $S_k(s, t) \subseteq V$. If $s, t \in V$, by Lemma 3, $\kappa(s, t) \leq 2 + 2(i + 1) = 2i + 4$. Otherwise, we assume, without loss of generality, that $s = uv \in E$. Let $p, q \in S_k(s, t) \subseteq V$, for some $k \leq d_{\Sigma(G)}(s, t)$ such that $d_{\Sigma(G)}(p, q)$ is maximized. Consider a shortest (s, t) -path P such that $p \in V(P)$. We may assume, without loss of generality, that $P \cap V$ is a shortest (u, w) -path in G for some $w \in t$ such that $d_G(u, w) \leq d_G(v, w)$ (if t is a vertex then $w = t$, else w is one of the ends of edge t). By now considering a shortest (s, t) -path Q such that $q \in V(Q)$, we obtain that q lies on some (u, w) -path (not necessarily shortest) in G of length at most $d_G(u, w) + 2$. Furthermore, $d_G(u, p) \leq d_G(u, q) \leq d_G(u, w) + 2$. By Lemma 2, $d_G(p, q) \leq (i + 2 + 2) + 1 = i + 5$. Therefore, $\kappa(s, t) \leq d_{\Sigma(G)}(p, q) + 2 \leq 2i + 12$. \square

Corollary 1. *If $G = (V, E)$ is α_i -metric, then it is $f(i)$ -hyperbolic for some doubly exponential function f .*

3.2 Using Cop-Robber Games

An n -vertex graph G , with $n > 1$, is *dismantlable* if its vertices can be ordered v_1, \dots, v_n so that for each vertex v_k , $1 \leq k < n$, there exists another vertex v_ℓ with $\ell > k$, such that $N[v_k] \cap V_k \subseteq N[v_\ell]$, where $V_k := \{v_k, v_{k+1}, \dots, v_n\}$. Such a vertex ordering is called a *dismantling ordering*. It is known [40,42] that the dismantlable graphs are exactly those graphs where in the classical cop and robber game the cop has a winning strategy. In this classical cop and robber game, two players, the cop \mathcal{C} and the robber \mathcal{R} , move alternatively along edges of a graph G . The cop captures the robber if both players are on the same vertex at the same moment of time. A graph G is called *cop-win* if the cop always captures the robber after a finite number of steps. Nowakowski, Winkler [40] and Quilliot [42] characterized the cop-win graphs as graphs admitting a dismantling ordering.

First we show that every $(i + 1)^{st}$ power of an α_i -metric graph G is dismantlable. Recall that the λ^{th} power of a graph $G = (V, E)$ is graph $G^\lambda = (V, E')$ such that $uv \in E'$ if and only if $0 < d(u, v) \leq \lambda$.

Lemma 5. *If G is an α_i -metric graph, then G^{i+1} is dismantlable. In particular, every BFS-ordering of G is a dismantling ordering of G^{i+1} .*

Proof. Let v_1, \dots, v_n with $v_n := u$ be a $BFS(u)$ -ordering of G started at an arbitrary vertex u . We will show that for every vertex v_k there is a vertex v_ℓ with $\ell > k$ such that for every vertex $x \in \{v_{k+1}, \dots, v_n\}$, $d(x, v_k) \leq i + 1$ implies $d(x, v_\ell) \leq i + 1$. As a vertex v_ℓ we will choose any neighbor of v_k in $I(v_k, v_n)$. Assume that for some $x \in \{v_{k+1}, \dots, v_n\}$ with $d(x, v_k) \leq i + 1$, $d(x, v_\ell) > i + 1$ holds. Then, necessarily, $d(x, v_\ell) = i + 2$ and $d(x, v_k) = i + 1$. By α_i -metric property applied to v_n, v_ℓ, v_k, x , $d(x, v_n) \geq d(v_n, v_k) + d(v_k, x) - i = d(v_n, v_k) + 1$ must hold, contradicting with the fact that in any $BFS(u)$ -ordering, $d(u, v_k) \geq d(u, x)$ for every $x \in \{v_{k+1}, \dots, v_n\}$. \square

For the case when $i = 1$, this result follows also from [11,45].

Recently, several other variants of the classical cop and robber game were introduced and investigated [12,13,29,30,39]. In a most general extension of the game, the cop \mathcal{C} and the robber \mathcal{R} move at speeds $s' \geq 1$ and $s \geq 1$, respectively. This game was introduced and thoroughly

investigated in [12,13]. It generalizes the cop and fast robber game from [29,30,39]. The unique difference between this “ (s, s') -cop and robber game” and the classical cop and robber game is that at each step, \mathcal{C} can move along a path of length at most s' and \mathcal{R} can move along a path of length at most s not traversing the position occupied by the cop. In [12], the class of cop-win graphs for this game was denoted by $\mathcal{CWFR}(s, s')$.

Similar to the characterization of classical cop-win graphs given in [40,42], the (s, s') -cop-win graphs have been characterized in [12] via a special (s, s') -dismantling scheme. A graph $G = (V, E)$ is called (s, s') -*dismantlable* if its vertices can be ordered v_1, \dots, v_n so that for each vertex v_k , $1 \leq k < n$, there exists another vertex v_ℓ with $\ell > k$, such that $D_{G-v_\ell}(v_k, s) \cap V_k \subseteq D_G(v_\ell, s')$, where $V_k := \{v_k, v_{k+1}, \dots, v_n\}$ and $D_{G-v_\ell}(v_k, s)$ is the disk of radius s and with center v_k considered in the graph $G \setminus \{v_\ell\}$. Such a vertex ordering is called an (s, s') -*dismantling ordering*. It was proven in [12] that G belongs to the class $\mathcal{CWFR}(s, s')$, $s' \leq s$, if and only if G is (s, s') -dismantlable¹. Furthermore, any δ -hyperbolic graph belongs to the class $\mathcal{CWFR}(2r, r + 2\delta)$ for any $r > 0$, and the graphs in $\mathcal{CWFR}(s, s')$ are $(s - 1)$ -hyperbolic for any $s \geq 2s'$ [12]. In the follow-up paper [13], those results were complemented by proving that if $s' < s$, then any graph of $\mathcal{CWFR}(s, s')$ is δ -hyperbolic with $\delta = O(s^2)$ (unfortunately, a sufficiently large constant is hidden under the big-O). Authors of [13] showed also that the dependency between δ and s is linear if $s - s' = \Omega(s)$ and G satisfies a slightly stronger (s, s') -dismantling condition. A graph $G = (V, E)$ is called (s, s') -*dismantlable* if its vertices can be ordered v_1, \dots, v_n so that for each vertex v_k , $1 \leq k < n$, there exists another vertex v_ℓ with $\ell > k$, such that $D(v_k, s) \cap V_k \subseteq D(v_\ell, s')$, where $V_k := \{v_k, v_{k+1}, \dots, v_n\}$ and both disks are considered in G . Such a vertex ordering is called an (s, s') -*dismantling ordering*. Paper [13] demonstrated also that weakly modular graphs from $\mathcal{CWFR}(s, s')$ with $s' < s$ are $184s$ -hyperbolic.

Next, we show that α_i -metric graphs with $n > 1$ vertices belong to the class $\mathcal{CWFR}(r, \lceil r/2 \rceil + 2i + 1)$ for any $r > 0$. In fact, we will prove that they are $(r, \lceil r/2 \rceil + 2i + 1)$ -dismantlable. For that, we will need Lemma 3.

Lemma 6. *If $G = (V, E)$ is an α_i -metric graph with $n > 1$ vertices, then every BFS-ordering of G is its $(r, \lceil r/2 \rceil + 2i + 1)$ -dismantling ordering for every $r > 0$.*

Proof. We may assume that $r \geq 2$. Let v_1, \dots, v_n with $v_n := u$ be a BFS(u)-ordering of G started at an arbitrary vertex u . We will show that for every vertex v_k there is a vertex v_ℓ with $\ell > k$ such that for every vertex $x \in \{v_{k+1}, \dots, v_n\}$, $d(x, v_k) \leq r$ implies $d(x, v_\ell) \leq \lceil r/2 \rceil + 2i + 1$. As a vertex v_ℓ we will choose any vertex from $S_q(v_k, v_n)$, where $q = \lceil r/2 \rceil$ (if $d(v_k, v_n) < \lceil r/2 \rceil$, then set $v_\ell := v_n$). Let x be an arbitrary vertex from $\{v_{k+1}, \dots, v_n\}$ with $d(x, v_k) \leq r$, x' be a vertex of $S_q(v_k, v_n)$ closest to x and x'' be a neighbor of x' on a shortest path from x' to x . By the choice of x' , $d(x'', v_k) > d(x', v_k)$ or $d(x'', v_n) > d(x', v_n)$. In the former case, by the α_i -metric property applied to v_k, x', x'', x , we get $d(v_k, x) \geq d(v_k, x') + d(x', x) - i$. That is, $d(x', x) \leq r - \lceil r/2 \rceil + i = \lceil r/2 \rceil + i$. In the latter case, by the α_i -metric property applied to v_n, x', x'', x , we get $d(v_n, x) \geq d(v_n, x') + d(x', x) - i$. We also know, by the property of a BFS(u)-ordering, that $d(v_n, v_k) \geq d(v_n, x)$. Hence, $d(x', x) \leq d(v_n, x) - d(v_n, x') + i \leq d(v_n, v_k) - d(v_n, x') + i \leq d(x', v_k) + i \leq \lceil r/2 \rceil + i$.

Since in either case $d(x', x) \leq \lceil r/2 \rceil + i$ holds and, by Lemma 3, $d(x', v_\ell) \leq i + 1$, we get $d(x, v_\ell) \leq \lceil r/2 \rceil + 2i + 1$. \square

From Lemma 6 and [13, Corollary 3], we conclude:

¹ Note that if $s' > s$, then the cop can always capture the robber by strictly decreasing at each move his distance to the robber.

Corollary 2. *Every α_i -metric graph is δ -hyperbolic for $\delta = O(i)$.*

Proof. By Lemma 6, any α_i -metric graph G with $n > 1$ vertices has an $(r, \lceil r/2 \rceil + 2i + 1)^*$ -dismantling ordering for every $r > 0$. It is known [13, Corollary 3] that if a graph G is $(s, s')^*$ -dismantlable with $s - s' \geq ks$ for some constant $k > 0$, then G is $64s/k$ -hyperbolic. For our α_i -metric graph G , we can pick $k := 1/4$ and $r := 4(2i + 1)$. Then, G is $(s, s')^*$ -dismantlable with $s = 4(2i + 1)$, $s' = 3(2i + 1)$, and $s = s' + (1/4)s$. Therefore, G must be δ -hyperbolic for $\delta = 2^6 s/k = 2^{10}(2i + 1)$. \square

3.3 Using Geodesic Triangles

For every vertices x, y, z in a graph $G = (V, E)$, the Gromov product of x, y with respect to z is defined as $(x | y)_z = \frac{1}{2}(d(x, z) + d(y, z) - d(x, y))$. Let $\Delta(x, y, z) = P(x, y) \cup P(y, z) \cup P(z, x)$ be some geodesic triangle. There is a canonical distance-preserving embedding of x, y, z in some weighted star $T(x, y, z)$: the leaves are x, y, z , and the respective weights of their incident edges are $\alpha_x = (y | z)_x, \alpha_y = (z | x)_y, \alpha_z = (x | y)_z$. Furthermore, if we replace each edge of $T(x, y, z)$ by some continuous segment, then one can define a unique mapping $\varphi : \Delta(x, y, z) \mapsto T(x, y, z)$ such that the restriction of φ on either side $P(x, y), P(y, z)$ or $P(z, x)$ is an isometry. The thinness of $\Delta(x, y, z)$ equals the smallest δ such that, for every $u, v \in \Delta(x, y, z)$ such that $\varphi(u) = \varphi(v)$, $d_G(u, v) \leq \delta$. See Fig. 2 taken from [10] for an illustration.




Fig. 2. Thinness in graphs.

It is known [10, Proposition 3.1] that in δ -hyperbolic graphs all geodesic triangles are 4δ -thin. Conversely [20, Lemma 3.1], if all geodesic triangles in a graph are δ -thin, then it is δ -hyperbolic. Next we show that the thinness of geodesic triangles in α_i -metric graphs, and so, their hyperbolicity, is at most $3(i + 1)$.

Theorem 1. *All geodesic triangles in an α_i -metric graph G are $3(i + 1)$ -thin.*

Proof. Consider an arbitrary geodesic triangle formed by vertices x, y, z . By [10, Lemma 3.2] and by symmetry, it is sufficient to show that any two vertices $z' \in S_k(y, x)$ and $x' \in S_k(y, z)$, where $k = \lfloor (x|z)_y \rfloor$, are at distance at most $3(i + 1)$. To do so, we will show that a vertex z^* exists in $S_k(y, x)$ whose distance to any vertex $x' \in S_k(y, z)$ is at most $2(i + 1)$. Then, by Lemma 3, it will follow that $d(z', x') \leq 3(i + 1)$ for every vertices $z' \in S_k(y, x)$ and $x' \in S_k(y, z)$.

Consider three disks $D(y, r_y), D(x, r_x), D(z, r_z)$, where $r_y = k = \lfloor (x|z)_y \rfloor$, $r_x = \lceil (z|y)_x \rceil$ and $r_z = \lceil (x|y)_z \rceil$. Since $\lfloor (x|z)_y \rfloor + \lceil (z|y)_x \rceil = d(x, y)$, $\lfloor (x|z)_y \rfloor + \lceil (x|y)_z \rceil = d(y, z)$ and $\lceil (z|y)_x \rceil + \lceil (x|y)_z \rceil \geq d(x, z)$, those disks pairwise intersect. By Lemma 1, there is a vertex $z^* \in S_k(y, x)$ whose distance to z is at most $r_z + i = \lceil (x|y)_z \rceil + i$. Applying now Lemma 2 to z, y, z^* and any vertex $x' \in S_k(y, z)$, we get $d(z^*, x') \leq i + i + 2 = 2i + 2$. \square

It was further proved in [18, Proposition 10] that if the interval thinness of a graph G is at most μ , and the maximum side-length of its metric triangles is at most ν , then G must be $(16\mu + 4\nu)$ -hyperbolic. By Lemma 3, $\mu \leq i + 1$ if G is α_i -metric. Furthermore, if $i = 1$, then it was proved in [3] that $\nu \leq 2$ (see also Lemma 14 in the next section). However, we conclude this section by proving that for every $i \geq 2$, the maximum side-length of metric triangles in α_i -metric graphs is unbounded.

Lemma 7. *For every integer $p \geq 2$, there exists an α_2 -metric graph G_p with $4p$ vertices and a metric triangle $x_0 y_{p-1} z_{p-1}$ such that $d(x_0, y_{p-1}) = d(x_0, z_{p-1}) = p \geq 2 = d(y_{p-1}, z_{p-1})$. In particular, there is no constant upper bound on the side-length of metric triangles in an α_2 -metric graph.*

Proof. The vertex set of G_p ($p \geq 1$) consists of four disjoint sets: $W = \{w_0, w_1, \dots, w_{p-1}\}$; $X = \{x_0, x_1, \dots, x_{p-1}\}$; $Y = \{y_0, y_1, \dots, y_{p-1}\}$; and $Z = \{z_0, z_1, \dots, z_{p-1}\}$. Its edge set is as follows:

- for every i such that $0 \leq i < p - 1$, $y_i \sim y_{i+1}$ and $z_i \sim z_{i+1}$;
- for every i such that $0 \leq i < p$, x_i, y_i, w_i, z_i induce a C_4 ;
- for every i such that $0 \leq i < p - 1$, $w_i \sim y_{i+1}, z_{i+1}$;
- for every i such that $0 < i \leq p - 1$, $x_i \sim y_{i-1}, z_{i-1}$.




Fig. 3. Graph G_3 in the proof of Lemma 7.

See Fig. 3 for an illustration. We can compute from the above construction the following distances (by induction on p):

1. $d(y_i, y_j) = d(z_i, z_j) = |j - i|$;
2. $d(y_i, z_i) = 2$, and $d(y_i, z_j) = |j - i| + 1$ if $i \neq j$;
3. $d(x_i, y_j) = d(x_i, z_j) = 1 + j - i$ if $j \geq i$, and $d(x_i, y_j) = d(x_i, z_j) = i - j$ if $j < i$;
4. $d(w_i, y_j) = d(w_i, z_j) = 1 + i - j$ if $j \leq i$, and $d(w_i, y_j) = d(w_i, z_j) = j - i$ if $j > i$;

5. $d(x_i, x_j) = d(w_i, w_j) = |j - i| + 1$ if $i \neq j$;
6. $d(x_i, w_j) = j - i + 2$ if $i \leq j$, $d(x_i, w_{i-1}) = 2$, and $d(x_i, w_j) = i - j$ if $i > j + 1$.

In particular, for $p \geq 2$, we can directly deduce from the above distance formulas that $x_0 y_{p-1} z_{p-1}$ is a metric triangle because the open intervals $I^o(x_0, y_{p-1}) = Y \setminus \{y_{p-1}\}$, $I^o(x_0, z_{p-1}) = Z \setminus \{z_{p-1}\}$, $I^o(y_{p-1}, z_{p-1}) = \{w_{p-2}, x_{p-1}, w_{p-1}\}$ are pairwise disjoint. Furthermore, $d(x_0, y_{p-1}) = d(x_0, z_{p-1}) = p$, and $d(y_{p-1}, z_{p-1}) = 2$.

It remains to prove that G_p is an α_2 -metric graph. For $p = 1$, this is true, because $G_1 = C_4$. From now on, we assume $p > 1$ and G_{p-1} is an α_2 -metric graph. Then, $H_1 = G_p \setminus \{w_{p-1}, x_{p-1}, y_{p-1}, z_{p-1}\}$ is an isometric subgraph of G_p that is isomorphic to G_{p-1} . Furthermore, let us consider the automorphism φ of G_p such that, for every i such that $0 \leq i < p$, we have $\varphi(w_i) = x_{p-1-i}$, $\varphi(x_i) = w_{p-1-i}$, $\varphi(y_i) = y_{p-1-i}$ and $\varphi(z_i) = z_{p-1-i}$. We obtain that $H_2 = G_p \setminus \{w_0, x_0, y_0, z_0\} = \varphi(H_1)$ is also an isometric copy of G_{p-1} . We next observe that every edge of G_p must be covered by at least one of H_1, H_2 . Hence, due to the aforementioned isomorphism between H_1 and H_2 , it is sufficient to prove the α_2 -metric property in G_p for every edge uv of H_1 . For that, let us pick arbitrary s and t such that $d(s, u) < d(s, v)$ and $d(t, v) < d(t, u)$. By induction, H_1 is α_2 -metric. Therefore, we may assume that $s \notin V(H_1)$.

Suppose first $t \notin V(H_1)$. If $s = y_{p-1}$, $t = z_{p-1}$, then $u \in Y$ and $v \in Z$, which is impossible because there are no edges between Y and Z . Otherwise, up to reverting the respective roles of s and t , we may assume (see the distance formulas above) that $d(s, r) \leq d(t, r)$ for every $r \in V(H_1)$. In particular, $d(s, u) < d(s, v) \leq d(t, v) < d(t, u)$. This implies $d(s, u) = d(t, u) - 2$, $d(s, v) = d(t, v)$, and so $s = x_{p-1}, t = w_{p-1}$. However, $|d(x_{p-1}, r) - d(w_{p-1}, r)| \leq 1$ for every $r \in V(H_1)$, which holds in particular for $r = u$, a contradiction. As a result, $t \in V(H_1)$.

Assume furthermore $s \sim u$. Then, $d(s, u) + d(u, t) - 2 = d(u, t) - 1 \leq d(s, t) + 1 - 1 = d(s, t)$, and we are done. Hence, from now on, we assume $s \not\sim u$ and, similarly, $t \not\sim v$.

- *Case $s = x_{p-1}$.* For every $r \in V(H_1) \setminus \{w_{p-2}\}$, we have $d(r, x_{p-1}) = d(r, w_{p-2})$. It implies $t \neq w_{p-2}$ and $u \neq w_{p-2}$. Since we also assume that $u \not\sim s, v \neq w_{p-2}$ holds. Then, $d(s, t) = d(w_{p-2}, t) \geq d(w_{p-2}, u) + d(u, t) - 2 = d(s, u) + d(u, t) - 2$.
- *Case $s = w_{p-1}$.* For every $r \in V(H_1) \setminus \{w_{p-2}\}$, we have $d(r, w_{p-1}) = d(r, w_{p-2}) + 1$. It implies $t \neq w_{p-2}$. Furthermore, since $d(w_{p-1}, w_{p-2}) = 2$, we get that w_{p-1} is equidistant to w_{p-2} and any of its neighbours in H_1 . In particular, $w_{p-2} \neq u, v$. Then, $d(s, t) = d(w_{p-2}, t) + 1 \geq (d(w_{p-2}, u) + d(u, t) - 2) + 1 = (d(w_{p-2}, u) + 1) + d(u, t) - 2 = d(s, u) + d(u, t) - 2$.
- *Case $s = y_{p-1}$ or $s = z_{p-1}$.* By symmetry, we only consider case $s = y_{p-1}$.

Let us first consider the subcase $\{u, v\} \cap \{y_{p-2}, z_{p-2}, w_{p-2}\} \neq \emptyset$. Since $s \not\sim u$, edge uv must be incident to z_{p-2} and another non-neighbour of s . Since furthermore we must have $d(s, u) < d(s, v)$, $u = z_{p-2}$ and $v = z_{p-3}$ is the only possibility. Let $H'_1 = H_1 \setminus \{w_{p-2}, x_{p-2}, y_{p-2}, z_{p-2}\}$. Note that H'_1 is an isometric copy of G_{p-2} , and that $t \in V(H'_1)$ because we assumed $d(t, v) < d(t, u)$. Assume $t \in Y$. We must have $t = y_j$ for some $j \leq p-4$. Then, $d(y_{p-1}, y_j) = p-1-j$ and $d(y_{p-1}, z_{p-2}) + d(z_{p-2}, y_j) - 2 = p-1-j$. Assume now $t \notin Y$. Then, $d(t, y_{p-1}) = d(t, z_{p-2}) + 1$ and $d(t, z_{p-2}) + d(z_{p-2}, y_{p-1}) - 2 = d(t, z_{p-2}) < d(t, y_{p-1})$.

From now on, $\{u, v\} \cap \{y_{p-2}, z_{p-2}, w_{p-2}\} = \emptyset$. For every $r \in V(H_1) \setminus \{y_{p-2}, z_{p-2}, w_{p-2}\}$, we have $d(r, y_{p-1}) = d(r, y_{p-2}) + 1$. It implies $y_{p-2} \neq t$. Then, $d(t, y_{p-1}) = d(t, y_{p-2}) + 1 \geq (d(t, u) + d(u, y_{p-2}) - 2) + 1 = d(t, u) + d(u, y_{p-1}) - 2$.

Overall, this above case analysis implies that edge uv satisfies the α_2 -metric property in G_p . \square

3.4 Using Injective Hulls

A graph is called Helly if every family of pairwise intersecting disks has a nonempty common intersection. It is known (*e.g.*, see [20]) that every graph G can be isometrically embedded in a (unique) smallest Helly graph $\mathcal{H}(G)$, which is sometimes called the injective hull of G . In what follows, we will show that using some recent results on injective hulls of graphs, a better bound on the hyperbolicity of α_i -metric graphs can be obtained. First note that, unfortunately, the injective hull of an α_i -metric graph may not necessarily be α_i -metric. This complicates some considerations but the difficulties can be circumvented. In Fig.4 we give an example of an α_1 -metric graph whose injective hull is not α_1 -metric. To have other such examples, one needs to note that the bound of Lemma 3 is sharp. In particular, that means that there are α_1 -metric graphs G whose injective hulls $\mathcal{H}(G)$ are not α_1 -metric, because every α_1 -metric graph has convex disks [45], however the interval thinness of a Helly graph with convex disks must be at most 1 [28].




Fig. 4. (a) An α_1 -metric graph G with interval thinness 2. (b) The injective hull of G which is not α_1 -metric (contains a C_4).

We recall the following properties of injective hulls.

Lemma 8 ([32]). *A graph G is δ -hyperbolic if and only if its injective hull $\mathcal{H}(G)$ is δ -hyperbolic.*

Lemma 9 ([23]). *If H is Helly and its interval thinness is at most τ , then it is $\lceil \frac{\tau}{2} \rceil$ -hyperbolic.*

Lemma 10 ([24]). *If H is the injective hull of G and $x, y \in V(H)$ are arbitrary, then there exist $x', y' \in V(G)$ such that any shortest (x, y) -path in H is contained in a shortest (x', y') -path in H .*

We complement these above properties with the following simple observation about distances in the injective hull of a graph:

Lemma 11. *If H is the injective hull of G and x, y are vertices of H such that $d(x, v) \leq d(y, v) + \lambda$ for every $v \in V(G)$, then $d(x, y) \leq \lambda$.*

Proof. By Lemma 10, there exist $u, v \in V(G)$ such that any shortest (x, y) -path in H is contained in a shortest (u, v) -path in H . Without loss of generality, $d(v, y) \leq d(v, x)$. Since $d(v, x) \leq d(v, y) + \lambda$, $d(x, y) = d(v, x) - d(v, y) \leq \lambda$.

We are now ready to prove our main result in this section.

Theorem 2. *If $G = (V, E)$ is α_i -metric, then it is δ -hyperbolic for some $\delta \leq i + \lceil \frac{i+1}{2} \rceil \leq \frac{3}{2} \cdot (i+1)$.*

Proof. Set $\delta = i + \lceil \frac{i+1}{2} \rceil$. By Lemma 8, G and its injective hull $\mathcal{H}(G)$ have the same hyperbolicity constant. Therefore, we are left proving that $\mathcal{H}(G)$ is δ -hyperbolic. For that, by Lemma 9, it is sufficient to prove that the interval thinness of $\mathcal{H}(G)$ is at most 2δ . Furthermore in order to bound the interval thinness of $\mathcal{H}(G)$, by Lemma 10, it is sufficient to consider the shortest-paths between vertices $u, v \in V$ (i.e., we discard the pairs of vertices with at least one vertex in $\mathcal{H}(G) \setminus V$).

Thus, from now on, let $u, v \in V$ be fixed, and let $S_k(u, v, \mathcal{H}(G))$ be some fixed slice of $I_{\mathcal{H}(G)}(u, v)$. By Lemma 3, the disks $D(x, \lceil \frac{i+1}{2} \rceil)$, for every $x \in S_k(u, v, G)$, pairwise intersect. Therefore, by the Helly property, there exists a vertex c of $\mathcal{H}(G)$ such that $S_k(u, v, G) \subseteq D(c, \lceil \frac{i+1}{2} \rceil)$. Then, in order to prove that every two vertices of $S_k(u, v, \mathcal{H}(G))$ are pairwise at distance at most 2δ , it suffices to prove that $S_k(u, v, \mathcal{H}(G)) \subseteq D(c, \delta)$. For that, let $y \in S_k(u, v, \mathcal{H}(G)) \setminus V$ be arbitrary. Let us define $r_y(w) = d(y, w)$ for every $w \in V \setminus \{u, v\}$. By Lemma 1, there is a $x \in S_k(u, v, G)$ such that $d(x, w) \leq r_y(w) + i$. It implies $d(c, w) \leq \lceil \frac{i+1}{2} \rceil + r_y(w) + i = r_y(w) + \delta$. By Lemma 11, $d(y, c) \leq \delta$. \square

Finally, note that, although the hyperbolicity of an α_i -metric graph is upper-bounded by $i + \lceil \frac{i+1}{2} \rceil$, there are n -vertex 1-hyperbolic graphs which satisfy α_i -metric only for $i = \Omega(n)$. Consider, for example, a ladder with height ℓ (see Fig. 5). It is a 1-hyperbolic graph and satisfies α_i -metric only for $i = 2\ell$.




Fig. 5. A ladder with height ℓ . It is a 1-hyperbolic graph but satisfies α_i -metric only for $i = 2\ell$ as $v \in I(u, w), w \in I(v, x)$ and $1 = d(u, x) = d(u, v) + d(v, x) - 2\ell$.

4 Graphs with α_1 -metric

By Theorem 2, every α_0 -metric graph must be 1-hyperbolic, and similarly every α_1 -metric graph must be 2-hyperbolic. In fact [44, Corollary 30], the hyperbolicity of an α_0 -metric graph is at most $\frac{1}{2}$. In this section, we improve the bound on the hyperbolicity of α_1 -metric graphs from 2 to 1 (Theorem 4). The bound is sharp even for the subclass of chordal graphs [8].

4.1 Intermediate Results

First, we recall an elegant characterization of α_1 -metric graphs.

Theorem 3 ([45]). G is an α_1 -metric graph if and only if all disks $D(v, k)$ ($v \in V, k \geq 1$) of G are convex and G does not contain the graph W_6^{++} from Fig. 6 as an isometric subgraph.

Lemma 12 ([43]). All disks $D(v, k)$ ($v \in V, k \geq 1$) of a graph G are convex if and only if G does not contain isometric cycles of length $\ell > 5$, and for any two vertices x, y of G the slice $S_1(x, y)$ is a clique.




Fig. 6. Forbidden isometric subgraph W_6^{++} .

The next lemma characterizes the situation in which, in an α_1 -metric graph, the union of two shortest paths with one common terminal edge is not a shortest path.

Lemma 13 ([3]). *Let G be an α_1 -metric graph. Let x, y, v, u be vertices of G such that $v \in I(x, y)$, $x \in I(v, u)$, and x and v are adjacent. Then $d(u, y) = d(u, x) + d(v, y)$ holds if and only if there exist a neighbor x' of x in $I(x, u)$ and a neighbor v' of v in $I(v, y)$ with $d(x', v') = 2$; in particular, x' and v' lie on a common shortest path of G between u and y .*

We will also need in our proofs the following properties of metric triangles in α_1 -metric graphs.

Lemma 14 ([3]). *In an α_1 -metric graph G , every metric triangle is of type $(1, 1, 1)$, $(1, 2, 2)$, $(2, 1, 2)$, $(2, 2, 1)$ or $(2, 2, 2)$.*

Finally, we conclude this subsection with two more technical results. We note that they could be also deduced from [11].

Lemma 15 ([22]). *Let G be an α_1 -metric graph. Then, for every edge $xy \in E$ and a vertex $u \in V$ with $d(u, x) = d(u, y) = k$, either there is a common neighbor u' of x and y at distance $k - 1$ from u or there exists a vertex u' at distance 2 from x and y and at distance $k - 2$ from u such that, for every $z \in N(x) \cap N(u')$ and $w \in N(y) \cap N(u')$, the sequence (x, z, u', w, y) forms an induced C_5 in G .*

Corollary 3. *Let $x, y \in S_k(u, v)$ be adjacent in an α_1 -metric graph G . Then, x, y have common neighbours in both $S_{k-1}(u, v)$ and $S_{k+1}(u, v)$.*

Proof. By symmetry, it suffices to prove that x, y have a common neighbour in $S_{k-1}(u, v)$. Suppose, for the sake of contradiction, that it is not the case. By Lemma 15, there exists a vertex w such that $d(x, w) = d(y, w) = 2$ and $d(u, w) = k - 2$. Note that in this situation, $w \in S_{k-2}(u, v)$ and $x, y \in S_2(w, v)$. Let $x' \in N(x) \cap N(w)$ and let $y' \in N(y) \cap N(w)$. Again, by Lemma 15, the sequence (x, x', w, y', y) induces a C_5 . It implies that $x', y' \in S_1(w, v)$ are nonadjacent, thus implying that not all the disks of G are convex by Lemma 12. However, the latter contradicts Theorem 3. \square

4.2 Hyperbolicity

Our strategy in order to prove Theorem 4 is essentially the same as what we did for Theorem 2: we upper bound the interval thinness of the injective hull. For that, we heavily use structural and metric properties of α_1 -metric graphs.

We first improve Lemma 1, but only for the α_1 -metric graphs.

Lemma 16. *Let $D(u, r_u), D(v, r_v), D(w, r_w)$ be pairwise intersecting disks of G such that $d(u, v) = r_u + r_v$. If G is an α_1 -metric graph, then every vertex $x \in S_{r_u}(u, v) (= S_{r_v}(v, u))$ satisfies $d(w, x) \leq r_w + 2$.*

Then, $a_u \in I(x, y) \setminus D(v, r_v)$, contradicting Theorem 3. Therefore, $ya_u \notin E$, and we can prove similarly $ya_v \notin E$. Doing so, we obtain that the C_6 with vertex-set $\{x, a_v, b_v, y, b_u, a_u\}$ is induced. Next, we prove that z is adjacent to every vertex of $\{x, a_v, b_v, y, b_u, a_u\}$. For x, y that follows from the definition of z and for a_u, a_v that follows from Lemma 12 because $z, a_u \in S_1(x, u')$ and $z, a_v \in S_1(x, v')$. Now, there is a C_4 with vertex-set $\{z, a_u, b_u, y\}$. Since we already proved that $ya_u \notin E$, by Lemma 12, we obtain $zb_u \in E$. In the same way, by considering a C_4 with vertex-set $\{z, a_v, b_v, y\}$, we obtain that $zb_v \in E$. Doing so, we get that $\{x, a_v, b_v, y, b_u, a_u, z\}$ induces a copy of the wheel W_6 . By the choice of c_v , its only neighbours in X are a_v, b_v . Last, since $u' \in S_{r_u-1}(u, v)$ and $d(u', x) = 3$, we obtain that the only neighbours of u' in X are b_u, y . So, we proved as claimed that X induces a W_6^{++} . \diamond

It remains to prove that $G[X]$ is an isometric subgraph, thus contradicting Theorem 3. The only pairs of vertices that are at distance 3 in $G[X]$ are: (u', x) , (u', a_v) , (u', c_v) , (a_u, c_v) , (b_u, c_v) . Note that we already proved that $d(u', x) = 3$. Furthermore, $d(u', a_v) > 2$ because otherwise $a_v \notin D(u, r_u)$ would be on a shortest (u', x) -path, thus contradicting Theorem 3. Finally, $\min\{d(u', c_v), d(a_u, c_v), d(b_u, c_v)\} \geq 3$ because $u', a_u, b_u \in S_{r_u-1}(u, v)$ and $c_v \in S_{r_u+2}(u, v)$. \square

We need one more technical lemma.

Lemma 17. *Let u, v, a, b be vertices of a graph $G = (V, E)$ such that:*

- $d(u, v) = r_u + r_v$;
- $d(u, a) \leq r_u + r_a$, $d(v, a) \leq r_v + r_a$;
- $d(u, b) \leq r_u + r_b$, $d(v, b) \leq r_v + r_b$.

If G is an α_1 -metric graph, then $d(a, b) \leq r_a + r_b + 2$.

Proof. Suppose, by way of contradiction, that $d(a, b) > r_a + r_b + 2$. The reader can follow our construction using Fig. 8. By Lemma 1, there exists a vertex $x \in S_{r_u}(u, v)$ such that $d(a, x) \leq r_a + 1$. By Lemma 16, $d(x, b) \leq r_b + 2$. As a result, $d(a, b) = r_a + r_b + 3$. Furthermore, $x \in I(a, b)$.

Let $y \in S_1(x, b)$ be arbitrary. If $d(u, y) > r_u$ then, since G is an α_1 -metric graph, we obtain $d(u, b) \geq d(u, x) + d(y, b) = r_u + r_b + 1$, and a contradiction with the assumption of the lemma arises. In the same way, if $d(v, y) > r_v$, then we obtain $d(v, b) \geq r_v + r_b + 1$, giving a contradiction again. As a result, $y \in S_{r_u}(u, v)$.

Let $z \in S_1(y, b)$ be arbitrary. We claim that $z \notin S_{r_u}(u, v)$. Indeed, otherwise, by Lemma 16, $d(z, a) \leq r_a + 2$, and so $d(a, b) \leq d(a, z) + d(z, b) \leq r_a + r_b + 2$. By symmetry, let $d(u, z) = r_u + 1$. Since G is an α_1 -metric graph, $d(u, b) \geq d(u, y) + d(z, b) = r_u + r_b$. This together with the assumption $d(u, b) \leq r_u + r_b$ implies $d(u, b) = d(u, y) + d(z, b) = r_u + r_b$. By Lemma 13, there exist $w_u \in S_1(y, u)$ and $w_b \in S_1(z, b)$ satisfying $d(w_u, w_b) = 2$.

Let $c \in N(x) \cap N(y) \cap S_{r_u-1}(u, v)$, whose existence follows by Corollary 3. Note that $d(c, w_b) > 1$ (else, $d(u, b) \leq d(u, c) + 1 + d(w_b, b) = r_u - 1 + 1 + r_b - 1 < r_u + r_b$). If $d(c, w_b) = 2$, then we replace w_u by c . Doing so, we claim that $N(w_u) \cap N(w_b) \subseteq S_{r_b}(b, a)$. For that, let $w_{ub} \in N(w_u) \cap N(w_b)$ be arbitrary. Since $c, w_u \in S_1(y, u)$, by Lemma 12, either $c = w_u$ or $cw_u \in E$. We consider both cases separately.

- *Case $c = w_u$.* Recall that $d(b, w_u) = r_b + 1$. Furthermore, $d(a, w_u) \leq 1 + d(a, x) = r_a + 2$. Therefore, $w_u, w_{ub} \in I(w_b, a) \subseteq I(b, a)$. It implies $w_{ub}, z \in S_1(w_b, a)$. By Lemma 12, $w_{ub}z \in E$.
- *Case $cw_u \in E$.* In particular, $d(w_b, c) > 2$. But then $d(c, w_b) = 3$ and $z, w_{ub} \in S_1(w_b, c)$. By Lemma 12, we obtain $w_{ub}z \in E$.




Fig. 8. To the proof of Lemma 17.

Summarizing, in both cases we get $w_{ub}z \in E$. Hence, we obtain that there is a C_4 with vertex-set $\{w_{ub}, z, y, w_u\}$. Since $w_uz \notin E$ (else, $d(u, z) \leq r_u$), by Lemma 12, we have $w_{ub}y \in E$. As a result, $w_{ub} \in S_1(y, b) \subseteq S_{r_b}(b, a)$.

By construction, $d(u, w_{ub}) \leq r_u$. Since $w_{ub} \notin S_{r_u}(u, v)$, $d(v, w_{ub}) = r_v + 1$. We replace z by w_{ub} in the above, and we invert the respective roles of u and v . Doing so, we obtain the following information: $d(v, b) = r_v + r_b$; and there exist adjacent vertices $w_v, w_{vb} \in N(y)$ such that $d(v, w_{vb}) = d(v, w_v) + 1 = r_v$ and $d(w_{vb}, b) = r_b$.

We invert the respective roles of a and b in the above. Doing so, we obtain the following additional information:

- $d(u, a) = r_u + r_a$; $d(v, a) = r_v + r_a$;
- there exist adjacent vertices $w'_u, w'_{ua} \in N(x)$ such that $d(u, w'_{ua}) = d(u, w'_u) + 1 = r_u$ and $d(w'_{ua}, a) = r_a$;
- there exist adjacent vertices $w'_v, w'_{va} \in N(x)$ such that $d(v, w'_{va}) = d(v, w'_v) + 1 = r_v$ and $d(w'_{va}, a) = r_a$.

By Lemma 12, we have that $w_{ub}, w_{vb} \in S_1(y, b)$ are adjacent, and similarly $w'_{ua}, w'_{va} \in S_1(x, a)$ are adjacent. Furthermore, $w_u \neq w'_u$ (otherwise, $d(a, b) \leq d(a, w'_u) + d(w_u, b) = r_a + r_b + 2$) and, similarly, $w_v \neq w'_v$. In what follows, let us assume $d(w_u, w'_u) + d(w_v, w'_v)$ to be minimized. By Lemma 3, $1 \leq d(w_u, w'_u), d(w_v, w'_v) \leq 2$. We need to distinguish between several cases.

Case $d(w_u, w'_u) = 2$ or $d(w_v, w'_v) = 2$. See Fig. 9. By symmetry, let $d(w_u, w'_u) = 2$. By Theorem 3, $w_u x, w'_u y \notin E$ (else, $D(u, r_u - 1)$ would not be convex). Let $c \in N(x) \cap N(y) \cap S_{r_u-1}(u, v)$, that exists by Corollary 3. Since we have $w_u, c \in S_1(y, u)$ and $w'_u, c \in S_1(x, u)$, by Lemma 12, $w_u c, w'_u c \in E$. Let $s \in N(w_u) \cap N(c) \cap S_{r_u-2}(u, v)$ and $t \in N(w'_u) \cap N(c) \cap S_{r_u-2}(u, v)$, that exist by Corollary 3. Since $c \in N(s) \cap N(t)$, $c \notin D(u, r_u - 2)$, by Theorem 3, we get $st \in E$. Furthermore, $sw'_u, tw_u \notin E$ (else, $D(v, r_v + 1)$ would not be convex). Then, $\{x, y, w_u, s, t, w'_u\}$ induces a C_6 and $\{c, x, y, w_u, s, t, w'_u\}$ induces a wheel W_6 .




Fig. 9. Case $d(w_u, w'_u) = 2$.

Since we have $d(w'_{ua}, w_{ub}) = 3$, one of the edges $w'_{ua}c, w_{ub}c$ must be missing. By symmetry, let us assume w'_{ua}, c are non-adjacent.

Claim. $d(w'_{ua}, w_u) = 3$.

Suppose $d(w'_{ua}, w_u) < 3$. Then, $d(w'_{ua}, w_u) = 2$ (else, $d(a, b) \leq d(a, w'_{ua}) + 1 + d(w_u, b) = r_a + r_b + 2$). Let $w' \in N(w'_{ua}) \cap N(w_u)$ be arbitrary. Observe that $(w'_{ua}, w', w_u, w_{ub})$ and (w'_{ua}, x, y, w_{ub}) are shortest paths. Therefore, by Lemma 12, $xw' \in E$. Note that in this situation, there are non-adjacent vertices $x, w_u \in N(w') \cap N(c)$, and therefore also by Lemma 12, $w'c \in E$. But then, there are non-adjacent vertices $w'_{ua}, c \in N(w') \cap N(w'_u)$, and so again, by Lemma 12, $w'_u w' \in E$. Since $D(u, r_u - 1)$ must be convex (Theorem 3) and $w' \in N(w_u) \cap N(w'_u)$, $d(u, w') = r_u - 1$. We replace w'_u by w' , thus contradicting the minimality of $d(w_u, w'_u) + d(w_v, w'_v)$. \diamond

Claim. $d(w'_{ua}, s) = 3$.

Consider the edge yw_u . We have $d(w'_{ua}, y) < d(w'_{ua}, w_u)$ and $d(s, w_u) < d(s, y)$. Since G is α_1 -metric, $d(w'_{ua}, s) \geq d(w'_{ua}, y) + d(w_u, s) = 2 + 1 = 3$. \diamond

Let $w_{st} \in N(s) \cap N(t) \cap S_{r_u-3}(u, v)$, that exists by Corollary 3. It follows from the above that the subgraph induced by $\{c, x, y, w_u, s, t, w'_u\} \cup \{w'_{ua}, w_{st}\}$ is an isometric W_6^{++} , thus contradicting Theorem 3.

Case $d(w_u, w'_u) = d(w_v, w'_v) = 1$. See Fig. 10. Since there is a C_4 with vertex-set $\{x, w'_u, w_u, y\}$ (respectively, $\{x, w'_v, w_v, y\}$), by Lemma 12, one of the edges xw_u, yw'_u (respectively, xw_v, yw'_v) must be present. Furthermore, since by Theorem 3 the disk $D(a, r_a + 1)$ (respectively, $D(b, r_b + 1)$)

must be convex, one of the edges $w'_u y, w'_v y$ (respectively, $w_u x, w_v x$) must be missing. As a result, by symmetry, we only need to consider the subcase when $w'_u y, w_v x \in E$ but $w_u x, w'_v y \notin E$. Then, $\{x, w'_u, w_u, w_{ub}, w_{vb}, w_v\}$ induces a cycle C_6 and $Y = \{x, w'_u, w_u, w_{ub}, w_{vb}, w_v\} \cup \{y\}$ induces a wheel W_6 .




Fig. 10. Case $d(w_u, w'_u) = d(w_v, w'_v) = 1$.

Let $\alpha \in N(w_u) \cap N(w'_u) \cap S_{r_u-2}(u, v)$ and $\beta \in N(w_{ub}) \cap N(w_{vb}) \cap S_{r_b-1}(b, a)$, that both exist by Corollary 3. By the choice of α, β , we have $N(\alpha) \cap Y = \{w_u, w'_u\}$ and $N(\beta) \cap Y = \{w_{ub}, w_{vb}\}$. Therefore, $X = \{\alpha, \beta\} \cup Y$ induces a W_6^{++} .

Moreover,

- $d(\alpha, \beta) \geq d(u, b) - d(u, \alpha) - d(b, \beta) = r_u + r_b - (r_u - 2) - (r_b - 1) = 3$;
- $d(\alpha, w_{vb}) \geq d(u, w_{vb}) - d(u, \alpha) = r_u + 1 - (r_u - 2) = 3$;
- $d(\alpha, w_v) \geq d(u, v) - d(u, \alpha) - d(v, w_v) = r_u + r_v - (r_u - 2) - (r_v - 1) = 3$;
- $d(\beta, x) \geq d(a, b) - d(a, x) - d(b, \beta) = r_a + r_b + 3 - (r_a + 1) - (r_b - 1) = 3$;
- $d(\beta, w'_u) \geq d(a, b) - d(a, w'_u) - d(b, \beta) = r_a + r_b + 3 - (r_a + 1) - (r_b - 1) = 3$.

As a result, $G[X]$ is an isometric subgraph, thus contradicting Theorem 3. \square

We are now ready to prove the main result in this section.

Theorem 4. *If $G = (V, E)$ is an α_1 -metric graph, then it is 1-hyperbolic.*

Proof. As we already discussed in the proof of Theorem 2, it is sufficient to prove that in $\mathcal{H}(G)$, for every $u, v \in V$ and for every $k \leq d(u, v)$, if $x, y \in S_k(u, v, \mathcal{H}(G))$ then we have $d(x, y) \leq 2$. For that, we need to consider three different cases.

- *Case $x, y \in V$.* The result follows from Lemma 3.
- *Case $x \in V, y \notin V$* (the case $x \notin V, y \in V$ is symmetric to this one). For every $w \in V$, let $r_y(w) = d(y, w)$. By Lemma 16, we have $d(x, w) \leq r_y(w) + 2$. By Lemma 11, $d(x, y) \leq 2$.
- *Case $x, y \notin V$.* By Lemma 10, there exist $x', y' \in V$ such that any shortest (x, y) -path in $\mathcal{H}(G)$ is contained in a shortest (x', y') -path in $\mathcal{H}(G)$. Let $r_x = d(x, x')$ and $r_y = d(y, y')$. By Lemma 17, $d(x', y') \leq r_x + r_y + 2$. Therefore, $d(x, y) \leq 2$.

The above proves, as claimed, that the interval thinness of $\mathcal{H}(G)$ is at most two, and therefore that G is 1-hyperbolic. \square

Next, we present two consequences of this result (in Section 4.3 and Section 4.4, respectively).

4.3 A new characterization of α_1 -metric graphs

The following characterization of 1/2-hyperbolic graphs was presented in [3].

Theorem 5 ([3]). *A graph G is 1/2-hyperbolic if and only if G is α_1 -metric and none of the graphs of Fig. 11 occur as isometric subgraphs.*




Fig. 11. Forbidden isometric subgraphs for 1/2-hyperbolic graphs.

Thus, it follows from Theorem 4 and Theorem 5 that the class of α_1 -metric graphs is a superclass of 1/2-hyperbolic graphs and a subclass of 1-hyperbolic graphs. Theorem 5 describes the place of 1/2-hyperbolic graphs within the class of α_1 -metric graphs in terms of forbidden isometric subgraphs. Using Theorem 3, Theorem 4 and a recent characterization of graphs with convex disks [11] (see Theorem 6), we can describe also the place of α_1 -metric graphs within the class of 1-hyperbolic graphs in terms of forbidden subgraphs.

Theorem 6 ([11]). *All disks of a graph G are convex if and only if G has no isometric C_k for every $k \geq 4$ such that $k \neq 5$, no isometric PT , every induced subgraph isomorphic to PP_1 has diameter at most 3 and every induced subgraph isomorphic to PP_2 has diameter 2 (see Fig. 12).*




Fig. 12. The graphs PT , PP_1 and PP_2 .

Corollary 4. *A graph G is α_1 -metric if and only if G is 1-hyperbolic, has no isometric C_4 , C_6 , C_7 , W_6^{++} or PT , every induced subgraph isomorphic to PP_1 has diameter at most 3 and every induced subgraph isomorphic to PP_2 has diameter 2 (see Fig. 6 and Fig. 12).*

Proof. It follows from Theorem 3, Theorem 4, Theorem 6, and the fact that a 1-hyperbolic graph cannot contain any isometric C_k for every $k \geq 8$. \square

4.4 Algorithmic Applications

Recall that the *eccentricity* $e(v)$ of a vertex v in G is defined by $\max_{u \in V} d(u, v)$, i.e., it is the distance to a most distant vertex. The *diameter* of a graph is the maximum over the eccentricities of all vertices: $\text{diam}(G) = \max_{u \in V} e(u) = \max_{u, v \in V} d(u, v)$. It was left as an open question in [22] whether in α_1 -metric graphs the eccentricity of any vertex furthest from an arbitrary vertex is at least $\text{diam}(G) - 2$. Our Theorem 4 provides an affirmative answer to that question. Indeed, it is known that in any δ -hyperbolic graph the eccentricity of any vertex furthest from an arbitrary vertex is at least $\text{diam}(G) - 2\delta$ [18]. So, we have the following corollary from Theorem 4.

Corollary 5. *In every α_1 -metric graph G , the eccentricity of any vertex furthest from an arbitrary vertex is at least $\text{diam}(G) - 2$. Hence, an additive 2-approximation of the diameter of G can be computed in linear time by a Breadth-First-Search.*

Note that computing the exact diameter in subquadratic time, even in chordal graphs (a proper subclass of α_1 -metric graphs), is impossible unless the well known Strong Exponential Time Hypothesis (SETH) is false [5].

5 Conclusion

Our main result in the paper is that every α_i -metric graph must be $f(i)$ -hyperbolic, for some function f such that $\frac{i+1}{2} \leq f(i) \leq 3 \cdot \frac{i+1}{2}$. It would be interesting to close the gap between the upper bound and the lower bound. We only managed to do so for α_1 -metric graphs. By contrast, general α_i -metric graphs ($i \in \mathcal{N}$) seem much less structured. For instance, they can have metric triangles with side-length unbounded (Lemma 7). Furthermore, unlike δ -hyperbolic graphs, the α_i -metric graphs are not well-behaved under important operations such as 1-subdivision (see Fig. 1) and injective hull (see Fig. 4).

Nevertheless, there is a natural generalization of an α_i -metric, which we call a (λ, μ) -bow metric: namely, if two shortest paths $P(u, w)$ and $P(v, x)$ share a common shortest subpath $P(v, w)$ of length more than λ (that is, they overlap by more than λ), then the distance between u and x is at least $d(u, v) + d(v, w) + d(w, x) - \mu$. Clearly, α_i -metric graphs satisfy $(0, i)$ -bow metric. However, this generalization is more robust to some graph operations. For instance, the 1-subdivision of an (λ, μ) -bow metric graph must satisfy $(2\lambda + 2, 2\mu)$ -bow metric. This notion of (λ, μ) -bow metric can also be considered for all geodesic metric spaces. Furthermore, in what follows, we show that every δ -hyperbolic graph (in fact, every δ -hyperbolic geodesic metric space) satisfies $(\delta, 2\delta)$ -bow metric.

Proposition 1. *Every δ -hyperbolic graph and, generally, every δ -hyperbolic geodesic metric space satisfies $(\delta, 2\delta)$ -bow metric.*

Proof. Consider arbitrary four vertices u, v, w, x such that $v \in I(u, w), w \in I(v, x)$ and $d(v, w) > \delta$. We need to show that $d(x, u) \geq d(u, v) + d(v, w) + d(w, x) - 2\delta$. Consider the three distance sums: (A) $d(u, w) + d(x, v)$, (B) $d(x, u) + d(w, v)$, and (C) $d(x, w) + d(u, v)$. Clearly, (A) is strictly larger than (C). If (B) is strictly larger than (A), then $d(x, u) + d(w, v) > d(u, w) + d(x, v) = d(u, v) + 2d(v, w) + d(w, x)$. On the other hand, $d(x, u) + d(w, v) \leq d(u, v) + d(v, w) + d(w, x) + d(w, v) = d(u, v) + 2d(v, w) + d(w, x)$. The contradiction obtained shows that (A) is at least (B). So, (A) is largest out of the three sums. If (C) is second largest sum then, by δ -hyperbolicity, $2\delta \geq (A) - (C) = d(u, w) + d(x, v) - d(x, w) - d(u, v) = 2d(v, w) > 2\delta$, which is impossible. Hence,

(B) is the second largest sum and, by δ -hyperbolicity, $2\delta \geq (A) - (B) = d(u, w) + d(x, v) - d(x, u) - d(w, v) = d(u, v) + 2d(v, w) + d(w, x) - d(x, u) - d(v, w) = d(u, v) + d(v, w) + d(w, x) - d(x, u)$, i.e., $d(x, u) \geq d(u, v) + d(v, w) + d(w, x) - 2\delta$. \square

We believe that the study of (λ, μ) -bow metrics could help in deriving new properties of α_i -metric graphs and δ -hyperbolic graphs.

Data Availability Statement

Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.

References

1. M. Abu-Ata, & F.F. Dragan. Metric tree-like structures in real-world networks: an empirical study, *Networks* 67(1) (2016), 49-68.
2. R. Albert, B. DasGupta, & N. Mobasher. Topological implications of negative curvature for biological and social networks, *Physical Review E* 89(3) (2014), 032811.
3. H.-J. Bandelt, V. Chepoi, 1-Hyperbolic graphs, *SIAM J. Discr. Math.* 16 (2003), 323–334.
4. H.-J. Bandelt, V. Chepoi, Metric graph theory and geometry: a survey, *Contemporary Mathematics* 453 (2008), 49-86.
5. M. Borassi, P. Crescenzi, and M. Habib, Into the square: On the complexity of some quadratic-time solvable problems, *Electronic Notes in TCS*, 322(2016), 51–67.
6. M. Borassi, A. Chessa, & G. Caldarelli. Hyperbolicity measures democracy in real-world networks, *Physical Review E* 92(3) (2015), 032812.
7. J.A. Bondy, U.S.R. Murty, Graph Theory, *Graduate Texts in Mathematics*, 244 (2008).
8. G. Brinkmann, J.H. Koolen, & V. Moulton. On the hyperbolicity of chordal graphs, *Annals of Combinatorics* 5(1) (2001), 61-69.
9. J. Chalopin, M. Changat, V. Chepoi, & J. Jacob. First-order logic axiomatization of metric graph theory, *arXiv:2203.01070*, 2022.
10. J. Chalopin, V. Chepoi, F.F. Dragan, G. Ducoffe, A. Mohammed, Y. Vaxès, Fast Approximation and Exact Computation of Negative Curvature Parameters of Graphs, *Discret. Comput. Geom.*, 65(2021), 856–892.
11. J. Chalopin and V. Chepoi, U. Giocanti, Graphs with convex balls, *Geom. Dedicata* (2023) <https://doi.org/10.21203/rs.3.rs-2293810/v1>.
12. J. Chalopin, V. Chepoi, N. Nisse, Y. Vaxès, Cop and robber games when the robber can hide and ride, *SIAM J. Discrete Math.* 25 (2011) 333–359.
13. J. Chalopin, V. Chepoi, P. Papasoglu, T. Pecatte, Cop and robber game and hyperbolicity, *SIAM J. Discrete Mathematics* 28 (2014), 1987–2007.
14. V. Chepoi, Some d -convexity properties in triangulated graphs, in *Mathematical Research*, vol. 87, Ştiinţa, Chişinău, 1986, pp. 164–177 (Russian).
15. V. Chepoi, Centers of triangulated graphs, *Math. Notes* 43 (1988), 143–151.
16. V. Chepoi and F. F. Dragan, Finding a central vertex in an HHD-free graph, *DAM*, 131(1) (2003), 93–111.
17. V. Chepoi, F.F. Dragan, B. Estellon, M. Habib, Y. Vaxés, & Y. Xiang. Additive spanners and distance and routing labeling schemes for hyperbolic graphs, *Algorithmica* 62 (2012), 713-732.
18. V.D. Chepoi, F.F. Dragan, B. Estellon, M. Habib and Y. Vaxès, Diameters, centers, and approximating trees of δ -hyperbolic geodesic spaces and graphs, Proceedings of the 24th Annual ACM Symposium on Computational Geometry (SoCG 2008), June 9-11, 2008, College Park, Maryland, USA, pp. 59-68.
19. V. Chepoi, F.F. Dragan, M. Habib, Y. Vaxès, and H. Alrasheed, Fast approximation of eccentricities and distances in hyperbolic graphs, *Journal of Graph Algorithms and Applications*, 23(2019), 393–433.
20. V.D. Chepoi, F.F. Dragan, & Y. Vaxés. Core congestion is inherent in hyperbolic networks, in *Proceedings of the Twenty-Eighth Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, Society for Industrial and Applied Mathematics (2018), pp. 2264-2279.
21. D. Coudert, G. Ducoffe. Recognition of C_4 -free and $1/2$ -hyperbolic graphs, *SIAM Journal on Discrete Mathematics* 28(3) (2014), 1601-1617.

22. F.F. Dragan, and G. Ducoffe, α_i -Metric Graphs: Radius, Diameter and all Eccentricities, WG 2023, Lecture Notes in Computer Science 14093, 276–290.
23. F.F. Dragan, and H.M. Guarnera, Obstructions to a small hyperbolicity in Helly graphs, in *Discrete Mathematics*, vol. 342.2, 2019, pp. 326-338.
24. F.F. Dragan, and H.M. Guarnera, Helly-gap of a graph and vertex eccentricities, in *Theoretical Computer Science*, vol. 867, 2021, pp. 68-84.
25. F.F. Dragan, E. Köhler, H. Alrasheed, Eccentricity Approximating Trees, *Discrete Applied Mathematics*, 232 (2017), 142–156.
26. F.F. Dragan, A. Mohammed, Slimness of graphs, *Discret. Math. Theor. Comput. Sci.* 21(3) (2019).
27. A.W.M. Dress, Trees, tight extensions of metric spaces, and the cohomological dimension of certain groups: A note on combinatorial properties of metric spaces, *Adv. Math.*, 53 (1984), pp. 321–402.
28. G. Ducoffe and F.F. Dragan, A story of diameter, radius, and (almost) Helly property, *Networks*, 77(2021), 435-453.
29. F.V. Fomin, P. Golovach, and J. Kratochvíl, On tractability of cops and robbers game, in *5th Ifip International Conference On Theoretical Computer Science*, vol. 273 (2008), pp. 171-185, Springer, Heidelberg.
30. F.V. Fomin, P. Golovach, J. Kratochvíl, N. Nisse, and K. Suchan, Pursuing a fast robber on a graph, *Theor. Comput. Sci.* 411 (2010), 1167–1181.
31. M. Gromov, Hyperbolic Groups, pages 75–263. Springer, New York, NY, 1987.
32. H.M. Guarnera, F.F. Dragan, and A. Leitert. Injective hulls of various graph classes, in *Graphs and Combinatorics*, vol. 38.4, 2022, 112.
33. E. Howorka, A characterization of distance-hereditary graphs, *Quart. J. Math. Oxford Ser.* 28 (1977), 417–420.
34. E. Howorka, On metric properties of certain clique graphs, *Journal of Combinatorial Theory, Series B* 27(1) (1979), 67–74.
35. E. Jonckheere, P. Lohsoonthorn. Geometry of network security, in *Proceedings of the American Control Conference*, IEEE, Vol. 2 (2004), pp. 976-981.
36. W. S. Kennedy, I. Saniee, & O. Narayan. On the hyperbolicity of large-scale networks and its estimation, in *IEEE International Conference on Big Data (Big Data)*, IEEE (2016), pp. 3344-3351.
37. J.H. Koolen, V. Moulton. Hyperbolic bridged graphs, *European Journal of Combinatorics* 23(6) (2002), 683-699.
38. F. de Montgolfier, M. Soto, L. Viennot, Treewidth and Hyperbolicity of the Internet. NCA 2011: 25–32.
39. N. Nisse and K. Suchan, Fast robber in planar graphs, in *Proceedings of the 34th International Workshop on Graph-Theoretic Concepts in Computer Science (WG)*, Springer LNCS vol. 5344 (2008), pp. 312–323.
40. R.J. Nowakowski and P. Winkler, Vertex-to-vertex pursuit in a graph, *Discr. Math.* 43 (1983), 235–239.
41. P. Papasoglu. Strongly geodesically automatic groups are hyperbolic, *Inventiones Math.* 121 (1995), 323– 334.
42. A. Quilliot, A short note about pursuit games played on a graph with a given genus, *J. Comb. Theory, Ser. B* 38 (1985), 89–92.
43. V.P. Soltan, V.D. Chepoi, Conditions for invariance of set diameters under d -convexification in a graph, *Cybernetics* 19 (1983), 750–756 (Russian, English transl.).
44. Y. Wu, C. Zhang. Hyperbolicity and chordality of a graph, *The Electronic Journal of Combinatorics* 18(1) (2001), P43.
45. S.V. Yushmanov, V. Chepoi, A general method of investigation of metric graph properties related to the eccentricity, in *Mathematical Problems in Cybernetics*, vol. 3, Nauka, Moscow, 1991, pp. 217–232 (Russian).